Efficient Generation of Humanized Cloned Cattle via Embryonic Stem Cell-Mediated

Dual-Round Site-Specific Editing

Zhiqiang Feng[1, #], Jinying Zhang[1, #], Shenao Tao[1, #], Peiru Lv[2, #], Xiaowei Zhang[1], Shunxin Wang[1],

Minglei Zhi[1], He Zhang[1], Chang Zeng[3], Xin Li[4], Zengyuan Zhao[4], Yanhua Li[5], Tianzhi Chen[1],

Yadi Teng[1], Yuhan Yang[1], Zimo Zhao[1], Hongkuan Ruan[1], Zhu Ma[5], Suying Cao[3, *], Shujing li[4, *],

Jianyong Han[1, *]

[1] State Key Laboratory of Animal Biotech Breeding, Frontiers Science Center for Molecular

Design Breeding (MOE), College of Biological Sciences, China Agricultural University, Beijing,

China.

[2] Henan Chuangyuan Biotechnology Co., Ltd.

[3] Animal Science and Technology College, Beijing University of Agriculture, Beijing, China.

[4] Beijing Dairy Cattle Center, Beijing 100192, People's Republic of China

[5] Shijiazhuang Tianquan Elite Dairy Co., Ltd. Shijiazhuang, Hebei 050200, People's Republic of

China

[#] These authors contributed equally: Zhiqiang Feng, Jinying Zhang, Shenao Tao, Peiru Lv.

[*] Correspondence should be addressed to:

Suying Cao, 20137602@bua.edu.cn

Shujing Li, embryochina@163. com

Jianyong Han, hanjy@cau.edu.cn

Abstract

Introduction: transgenic cloning has enabled genetic modification in cattle, but its

translational potential is constrained by low cloning efficiency and limited

gene-editing capacity, due largely to the restricted proliferative ability of somatic

donor cells. Pluripotent stem cells (PSCs), with their unlimited self-renewal capacity,

offer a promising alternative to overcome these bottlenecks.

Results: in this study, we first established stable bovine epiblast stem cell lines

(bEpiSCs) and achieved site-specific integration of the HCSN2-HLF-GFP cassette

into the β-casein locus (transfection efficiency: 72.07% $\pm$ 2.25%; correct integration:

80.48% $\pm$ 6.43%). The selectable marker was excised via Cre-LoxP, yielding

marker-free HCSN2-HLF bEpiSCs. This cell line retained pluripotency markers,

tri-lineage differentiation potential, and supported target protein expression in

immortalized bovine mammary epithelial cells (IBMECs). As nuclear donors,

HCSN2-HLF bEpiSCs produced blastocysts with quality comparable to IVF embryos.

Twelve cloned calves were generated, exhibiting significantly higher birth and

survival rates (30.71% $\pm$ 5.68%; 16.74% $\pm$ 1.40%) than the fibroblast group.

PCR-based genotyping of the edited loci and microsatellite kinship analysis

confirmed that all cloned calves were genetically derived from the HCSN2-HLF

bEpiSCs.

Conclusion: this study has successfully established an efficient technical platform

based on embryonic stem cell-targeted strategy, enabling the rapid generation of

multigene-edited cloned cattle with considerable industrial translation prospects.

Transgenic cloning enables genetic modification in livestock [1-3], with dairy cattle serving as

effective bioreactors for the production of functional recombinant proteins in milk [4], thereby

enhancing both biological activity and commercial value of dairy products. Humanized dairy

models have been obtained to express key human proteins including lactoferrin [5], serum albumin [6],

and lysozyme [7] in bovine milk, significantly improving its nutritional and functional properties.

While translational potential of the current strategy is dual constrained by low cloning efficiency

and limited gene editing capability due to the proliferative ability of donor somatic cells, which

cannot tolerate continuous rounds of complex gene editing. Specifically, the survival rate of

cloned animals typically remains as low as 1–5% [5, 8], which severely impedes the industrial-scale

application of this technology. Endowed with inherent unlimited proliferative capacity, pluripotent

stem cells tolerate continuous multiple rounds gene editing [9]. This may effectively overcome the

predicaments encountered in somatic cells. Here, we established a technical system to efficiently

obtained cloned cattle from stable embryonic stem cells with at least three milk protein genes

precise edits.

We selected human β-casein (HCSN2) and human lactoferrin (HLF), which are two key

functional proteins in human milk, for transgene candidates. As a major nutritional structural

protein, HCSN2 directly participates in regulating digestion, absorption, and mineral transport;

while, HLF serves as a core immunodefensive protein and plays a central role in human

milk-mediated immune protection. To perform dual-round site-specific editing, we use stable

bovine epiblast stem cells (bEpiSCs) generated from early embryos for this study. We have

designed a gene manipulation strategy (Fig. 1A) and successfully achieved site-specific

integration of the HCSN2-HLF fragment into the third exon of the bovine β-casein gene by

tandemly linking the *HCSN2* and *HLF* genes and adopting a two-step strategy assisted by the GFP

fluorescent reporter system. Subsequently, the edited HCSN2-HLF bEpiSCs were used as donors

for cell nuclear transfer.

We first successfully established three stable bEpiSC lines, which can be passaged for over

50 times without any signs of differentiation, from the epiblasts of three E7-stage blastocysts of

high-yield dairy cows (Fig S1A). These cells consistently maintained a typical dome-shaped

colony morphology (Fig S1B), positive alkaline phosphatase (AP) staining (Fig S1C), and a

normal karyotype of 60 chromosomes (Fig S1D) during long-term passage. Immunofluorescence

results demonstrated that the cells stably expressed the core pluripotency transcription factors such

as OCT4, NANOG, and SOX2 (Fig S1E). Further sex identification was performed via specific

PCR detection of the SRY gene, and the results showed that two lines were male and one was

female (Fig S1F, G). The female cell line was selected as the candidate for subsequent gene

editing in this study.

Initially, the HCSN2-HLF-GFP plasmid was transfected into bEpiSCs at passage 13 (Fig 1B).

At 48 hours post-transfection, robust GFP fluorescence was observed in most colonies (Fig 1B).

GFP-positive cells were then enriched via flow cytometry, achieving a sorting efficiency of 72.07%

± 2.25% (Fig 1C, D). The sorted GFP-positive cells were subjected to clonal expansion, and

individual colonies were randomly selected for PCR-based genotyping (Fig. S2C). Amplification

of fragments spanning the 5′ homology arm (5′HA: 6.1 kb), 3′ homology arm (3′HA: 4.1 kb),

HCSN2 (360 bp), and *HLF* (360 bp) confirmed that the proportion of correctly targeted colonies,

termed as HCSN2-HLF-GFP bEpiSCs, was 80.48% ± 6.43% (Fig. S2E).

To ensure that cloned animals exclusively express the target protein and eliminate the

potential risks associated with the reporter gene, site-specific excision was mediated by Cre

recombinase transfection following the establishment of passage 17 HCSN2-HLF-GFP bEpiSCs

(Fig. 1E). GFP-negative cell populations were isolated via flow cytometry, with an excision

efficiency of 69.07% ± 2.00% (Fig. 1F, G). After clonal expansion of these GFP-negative cells,

individual colonies were randomly selected for systematic genotyping, which included PCR

validation of long homologous arms (5′HA: 6.1 kb; 3′HA: 2.1 kb) and target genes (*HCSN2*: 360

bp; *HLF*: 360 bp) (Fig. S2D). The results demonstrated that all tested colonies exhibited precise

integration and stably harbored the intact *HCSN2* and *HLF* genes (Fig. S2F), culminating in the

successful construction of a marker-free HCSN2-HLF bEpiSCs. Notably, the efficiency of

large-fragment precise integration in stem cells is significantly higher than that in somatic cells,

and the operation is convenient [5, 10].

Next, we analyzed the pluripotency characteristics of HCSN2-HLF bEpiSCs. We found that

the cells retained canonical pluripotent morphological and molecular features after genetic editing.

They maintained a dome-shaped colony morphology and positive AP staining (Fig. S3A), with

sustained and stable nuclear expression of the core pluripotent transcription factors such as OCT4,

SOX2, and NANOG (Fig. 1H, Fig. S3B). In addition, *in vitro* embryoid body (EB) formation

assays showed that the cells were capable of self-assembling into EBs with intact structures and distinct outlines (Fig. S3C). Following spontaneous differentiation, cells derived from these EBs expressed markers specific to the three germ layers, such as ectodermal marker β-III-tubulin, mesodermal marker α-SMA, and endodermal marker GATA6, confirming their robust differentiation potential (Fig. 1I, Fig. S3D).

To validate the feasibility of the gene-editing strategy developed in this study, we first confirmed the suitability of immortalized bovine mammary epithelial cells (IBMECs) for subsequent functional assays via identification of mammary gland-specific markers (Fig. S4A). Subsequently, the editing strategy previously optimized in bovine embryonic stem cells (bEpiSCs) was applied in parallel to IBMECs. PCR amplification spanning homology arms and the target genes enabled the isolation of cell lines with site-specific integration of the HCSN2-HLF cassette (Fig. S4B). Furthermore, *in vitro* lactation induction assays combined with enzyme-linked immunosorbent assay (ELISA) analyses of both the supernatant and cell pellets revealed that endogenous bovine β-casein expression was significantly downregulated in edited cells, whereas the expression levels of exogenous human β-casein and human lactoferrin were markedly elevated (Fig. S4E, H, K). These findings demonstrate the efficacy and functionality of our gene-editing strategy at both the protein secretion and intracellular storage levels.

Subsequently, to systematically evaluate the developmental potential of HCSN2-HLF bEpiSCs as donor cells for nuclear transfer, we examined the early developmental efficiency of cloned embryos derived from this cell line. The results demonstrated that morphologically normal blastocysts could be successfully obtained using HCSN2-HLF bEpiSCs as donors (Fig. 1J). To further clarify the characteristics of their developmental capacity, we performed a horizontal comparison of the early embryonic developmental efficiency among the groups of *in vitro* fertilization (IVF), HCSN2-HLF bEpiSCs nuclear transfer, and fibroblast nuclear transfer. Interestingly, the early developmental capacity of HCSN2-HLF bEpiSCs-derived cloned embryos is significantly superior to that of the fibroblast cloning group and closer to that of IVF embryos (Fig. 1K, Table S1). Furthermore, cell counting analysis at the blastocyst stage showed that both the total cell number and the inner cell mass (ICM) cell number of blastocysts in the HCSN2-HLF bEpiSCs nuclear transfer group were significantly higher than those in the fibroblast group (Fig. S5A, B, C), indicating better embryonic developmental quality and further confirming that

HCSN2-HLF bEpiSCs possess the potential to serve as high-quality donor cells for nuclear transfer.

Finally, 105 cloned embryos were generated using HCSN2-HLF bEpiSCs as nuclear donors and transplanted into 71 surrogate cows in three separate batches (Table S2), leading to the successful birth of 12 healthy cloned calves (Fig. 1L, Fig. S6A, Table S3). Quantitative analysis revealed that both the birth rate (30.71% ± 5.68%) and survival rate (16.74% ± 1.40%) in the HCSN2-HLF bEpiSCs group were significantly higher than those in the fibroblast group (birth rate: 7.14% ± 4.40%; survival rate: 3.91% ± 1.68%) (Fig. 1M, N, Table S2). Subsequent genotypic characterization of the 12 calves via cross-homologous arm PCR and target gene detection confirmed that all calves contained the expected integrated fragments (5′ HA: 6.1 kb; 3′ HA: 2.1 kb) along with intact *HCSN2* and *HLF* genes (Fig. 1O, Fig. S6B, C). Moreover, kinship analysis based on fragment length polymorphism of 20 short tandem repeats (STR) loci demonstrated that all cloned calves were genetically consistent with the parental bEpiSCs and HCSN2-HLF bEpiSCs, while showing no genetic association with the 11 surrogate cows (Fig. 1P, Fig. S6D, E, Fig. S7A, B, C). Collectively, these results confirm that all cloned calves were derived from HCSN2-HLF bEpiSCs, thereby validating the reliability and efficiency of this cell line as a nuclear transfer donor.

Our study establishes an efficient and precise gene-editing cloning platform mediated by embryonic stem cells, enabling the rapid generation of multiplex gene-edited cloned cattle. It provides a scalable, integrated technological solution for large-scale pharmaceutical protein production via mammary gland bioreactors and the construction of humanized large-animal disease models.

Fig. 1: Generation of Cloned Cattle for Human β-Casein and Lactoferrin Production via

bEpiSCs

A Schematic diagram of cloned cattle editing mediated by bEpiSCs.

B Pre-transfection and post-transfection cellular status of bEpiSCs following transfection with the

HCSN2-HLF-GFP plasmid. Scale bar, 50 μm.

C The proportion of GFP-positive cells in bEpiSCs pre-transfection and post-transfection, with

trapezoidal boxes indicating GFP-positive cells.

D Bar plots showing the proportion of positive cells in bEpiSCs following transfection with the

HCSN2-HLF-GFP plasmid as shown in C. Data are presented as mean ± SD, n = 3 replicates.

****P*<0.001.

E Pre- and post-Cre transfection cellular status of HCSN2-HLF-GFP bEpiSCs. Scale bar, 50 μm.

F The proportion of GFP-negative cells in HCSN2-HLF-GFP bEpiSCs pre- and post-Cre

transfection, with trapezoidal boxes indicating GFP-negative cells.

G Bar plots showing the proportion of negative cells in HCSN2-HLF-GFP bEpiSCs transfection

with Cre as shown in F. Data are presented as mean ± SD, n = 3 replicates. *****P*<0.0001.

H Cellular status and immunofluorescence staining of core pluripotency marker proteins OCT4

NANOG and SOX2 in HCSN2-HLF bEpiSCs. Scale bar, 50 μm.

I *In vitro* embryoid body (EB) formation of HCSN2-HLF bEpiSCs and the capacity of EB for

spontaneous differentiation into the three germ layers, along with immunofluorescence staining of

the ectodermal neural-specific marker β-□-tubulin, mesodermal muscle-specific marker α-SMA,

and endodermal-specific marker GATA6. Scale bar, 50 μm.

J Schematic diagram of E7 blastocysts derived from *in vitro* fertilization (IVF), HCSN2-HLF

bEpiSCs, and fibroblasts. Scale bar, 200 μm.

K The line graph showing the early developmental capacity of different types of embryos. Data

are presented as the mean ± SD., n = 3 replicates.

L Healthy calves derived from HCSN2-HLF bEpiSCs via nuclear transfer technology, with the

photograph taken on the 45th day of the calves' survival. Clone#1 refers to cloned calf No.1, and

so forth.

M Bar plots showing a comparison of the birth rates of cloned calves derived from HCSN2-HLF

bEpiSCs and fibroblasts. Data are presented as the mean ±SD., n = 3 replicates. ***P*<0.01.

N Bar plots showing a comparison of the survival rates of cloned calves derived from
HCSN2-HLF bEpiSCs and fibroblasts. Data are presented as the mean ±SD., n = 3 replicates.
****P<0.0001.
O Detection of the 5' homology arm, 3' homology arm, and target gene in cloned calves derived
from HCSN2-HLF bEpiSCs, with bEpiSCs serving as the negative control and HCSN2-HLF
bEpiSCs as the positive control. SC#1 denotes surrogate cow No.1, and so forth. Clone#1 was
derived from SC#1; Clone#2 and Clone#3 were derived from SC#2; Clone#3 was derived from
SC#3.
P Calf parentage testing. Through microsatellite analysis, the allele fragment lengths at 20 loci
were compared among bEpiSCs, HCSN2-HLF bEpiSCs, cloned calves, and surrogate cows;
fragments consistent with those of bEpiSCs were labeled in orange, whereas inconsistent ones
were labeled in blue.

References

1 Cibelli JB, Stice SL, Golueke PJ, Kane JJ, Jerry J *et al.* Cloned transgenic calves produced

from nonquiescent fetal fibroblasts. *Science* **280**, 1256-1258 (1998).

2 Kato Y, Tani T, Sotomaru Y, Kurokawa K, Kato J *et al.* Eight calves cloned from somatic cells

of a single adult. *Science* **282**, 2095-2098 (1998).

3 Kubota C, Yamakuchi H, Todoroki J, Mizoshita K, Tabara N *et al.* Six cloned calves

produced from adult fibroblast cells after long-term culture. *Proc Natl Acad Sci U S A* **97**,

990-995 (2000).

4 Yu S, Luo J, Song Z, Ding F, Dai Y, Li N. Highly efficient modification of beta-lactoglobulin

(BLG) gene via zinc-finger nucleases in cattle. *Cell Res* **21**, 1638-1640 (2011).

5 Yang P, Wang J, Gong G, Sun X, Zhang R *et al.* Cattle mammary bioreactor generated by a

novel procedure of transgenic cloning for large-scale production of functional human

lactoferrin. *PLoS One* **3**, e3453 (2008).

6 Luo Y, Wang Y, Liu J, Cui C, Wu Y *et al.* Generation of TALE nickase-mediated

gene-targeted cows expressing human serum albumin in mammary glands. *Sci Rep* **6**, 20657

(2016).

7 Lu D, Liu S, Ding F, Wang H, Li J *et al.* Large-scale production of functional human lysozyme

from marker-free transgenic cloned cows. *Sci Rep* **6**, 22947 (2016).

8 Wu H, Wang Y, Zhang Y, Yang M, Lv J *et al.* TALE nickase-mediated SP110 knockin

endows cattle with increased resistance to tuberculosis. *Proc Natl Acad Sci U S A* **112**,

E1530-1539 (2015).

9 Zhi M, Zhang J, Tang Q, Yu D, Gao S *et al.* Generation and characterization of stable pig

pregastrulation epiblast stem cell lines. *Cell Res* **32,** 383-400 (2022).

10 Yuan M, Zhang J, Gao Y, Yuan Z, Zhu Z *et al.* HMEJ-based safe-harbor genome editing

enables efficient generation of cattle with increased resistance to tuberculosis. *J Biol Chem*

296, 100497 (2021).

239