A revised genome annotation of the model cyanobacterium *Synechocystis* based on start and stop codon-enriched ribosome profiling and proteogenomics

Lydia Hadjeras[1#], Vanessa Krauspe[2#], Rick Gelhausen[3#], Benjamin Heiniger[4,5], Philipp Spät[6], Viktoria Reimann[2], Garance Jaques[4], Paul Minges[2], Raphael Bilger[2], Maren Gerstner[2], Boris Maček[6], Christian H. Ahrens[4,*], Rolf Backofen[3,7], Cynthia M. Sharma[1] and Wolfgang R. Hess[2,*]

[1]University of Würzburg, Institute of Molecular Infection Biology (IMIB), University of Würzburg, Germany;

[2]Genetics and Experimental Bioinformatics, Faculty of Biology, University of Freiburg, Germany;

[3]Bioinformatics Group, Department of Computer Science, Technical Faculty, University of Freiburg, Germany;

[4]Molecular Ecology, Agroscope and SIB Swiss Institute of Bioinformatics, 8046 Zurich, Switzerland;

[5]PhD Program in Systems Biology, Zurich Life Science Graduate School

[6]Department of Quantitative Proteomics, Interfaculty Institute for Cell Biology, University of Tübingen, D-72076 Tübingen, Germany;

[7]Signalling Research Centre CIBSS, University of Freiburg, Germany

[#]Co-sharing first authors

*Corresponding authors: Christian H. Ahrens: christian.ahrens@agroscope.admin.ch, Wolfgang R. Hess: wolfgang.hess@biologie.uni-freiburg.de

Lead contact: Wolfgang R. Hess, wolfgang.hess@biologie.uni-freiburg.de

Running title: TIS- and TTS-Ribo-seq in *Synechocystis* sp. PCC 6803

ABSTRACT

Cyanobacteria are important primary producers and are used in biotechnology as microbial cell factories due to their ability to use solar light for oxygenic photosynthesis. *Synechocystis* sp. PCC 6803 is a popular model cyanobacterium, yet there are ambiguities in the precise coding regions of many genes, and numerous genes encoding small proteins have remained undetected. Here we present the results of a ribosome profiling (Ribo-seq) analysis involving inhibitors that stall ribosomes at translation initiation and termination sites (TIS- and TTS-Ribo-seq), combined with a proteogenomic reevaluation and reannotation of its entire genome. We report evidence for the translation of 3,050 annotated genes based on proteogenomics (83%), of 3,492 based on Ribo-seq (95.2%), and of 3,009 supported by both methods (82%). The data suggested both novel protein-coding genes and corrections for annotated ones. We validated 15 novel small proteins translated from antisense RNAs, from intergenic and intragenic regions and identified 69 novel, mostly small proteins based on proteogenomics. With *slr0489, slr1079* and *slr1082* we identified three genes with ~300 nt long intragenic out-of-frame coding regions and show that both the internal and host reading frames are translated. The resulting proteins interact with each other, resembling certain defense or toxin-antitoxin systems. Our data illustrate the enormous value of consolidating genome annotations in the context of integrated experimental data and suggest that genome annotations in general need to be extended and revised. All of our data can be accessed via an intuitive and interactive genome browser platform at https://www.bioinf.uni-freiburg.de/~ribobase/.

INTRODUCTION

Oxygenic photosynthesis powers photoautotrophic organisms such as cyanobacteria enabling them to build complex organic molecules from inorganic carbon. As primary producers, cyanobacteria have a pivotal ecological significance[1] and considerable value in biotechnological applications[2–4]. Among cyanobacteria, *Synechocystis* sp. PCC 6803 (from here: *Synechocystis* 6803), the third organism to have its complete genome sequenced[5], is considered one of the most advanced models. However, despite careful annotation attempts, severe annotation ambiguities still exist.

Cyanobacteria, like other prokaryotes, harbor a higher share of genes encoding small
proteins compared to eukaryotes due to their more compact genome architecture[6].
Importantly, small protein-coding genes are frequently missed during standard
genome annotation which aims to minimize the over-prediction of spurious short
ORFs. Moreover, standard mass spectrometry-based approaches fail to
comprehensively detect this class of proteins[7], making small proteins a systematically
underestimated component of the total proteome. Small proteins (here defined as ≤70
amino acids (aa)) are often considered too small to exert enzymatic activity.
Nevertheless, many small proteins were demonstrated to interact with larger proteins
or multi-protein complexes and have important functions[8]. Examples are the One-Helix
Proteins in the assembly or stabilization of photosynthetic pigment-protein complexes
in plant chloroplasts and cyanobacteria[9,10], small structural proteins in photosystem
(PS) I[11], from which PsaJ (40 aa) is essential for the stable assembly and structural
integrity of the entire complex[12]. Due to the extensive previous work on the
photosynthetic apparatus, cyanobacteria provide a paradigm for small protein
functions: Nineteen well-characterized proteins smaller than 50 aa are associated with
the photosynthetic machinery alone[13] . Globally, small proteins operate as modulators
of larger proteins' enzymatic activities[8,14–16], such as the 48 aa protein AtpΘ, which
inhibits the futile reverse reaction of ATP synthase under unfavorable conditions[17].
Lastly, many small proteins are toxic, including those that are part of type I toxin–
antitoxin systems[18], or antimicrobial peptides (AMPs)[19].

Ribosome profiling (Ribo-seq) detects translated mRNA segments on a genome-wide
level[20]. Following cellular lysis and nuclease treatment, intact RNA associated with
proteins – including the mRNA engaged with ribosomes during translation, is
separated from cell debris by sucrose gradient ultracentrifugation followed by recovery
of the associated RNA, cDNA synthesis and sequencing[21]. The analysis of RNA
associated with the bacterial 70S ribosomes yields ribosome footprints with a length
of 15–40 nt, peaking around 24 nt for *E. coli*[22]. Through comparison with an RNA-seq
library generated in parallel from total RNA harvested from the same culture and
fragmented into similar size, the coverage by sequenced ribosome footprints then
allows to infer ribosome occupancy, open reading frames (ORFs) boundaries, and
translational efficiency. While Ribo-seq was initially developed to study translational
regulation[23–25], the discovery of extensive, previously unrecognized, coding potential

in genomes of diverse organisms, such as yeast[26], vertebrates including humans[27,28],

plants[29], bacteria[30–32], and viruses[33] was another major benefit. The identification of

potential short ORF encoded proteins (SEPs) catalogued by Ribo-seq in diverse

organisms is gaining momentum. After the adaptation of the Ribo-seq workflow, this

has only recently been extended to a range of bacteria and archaea[32,34–36].

The antibiotic retapamulin, an inhibitor of the initiation of translation in bacteria, has

revolutionized the view on the small proteome through Ribo-seq. Retapamulin (a

pleuromutilin antibiotic) stalls ribosomes in treated cells at the translation initiation site

(TIS), while elongating ribosomes finish the translation and run off the mRNA releasing

the nascent peptide. Thus, retapamulin-assisted TIS-Ribo-seq results in defined

peaks for translated proteins at their respective translational start sites, which allows

to detect both unknown internal proteins, alternative versions of known proteins and

unannotated, novel small proteins[30,37]. Further improvements have been obtained by

the introduction of Apidaecin, a proline-rich antimicrobial peptide, which arrests

translation close to the stop codon[38] and leads to the accumulation of ribosome

footprints close to the 3′ ends of coding regions[39], and thus can be used for global

mapping of stop codons via translation termination site (TTS)-Ribo-seq[32,40].

Combining TIS- and TTS-Ribo-seq data can thus yield the precise borders of a

translated sequence from start to stop[32,40]. These recent methodological

breakthroughs applied alone have already yielded unprecedented insight not only into

the small proteome of bacteria, but also suggested an unexpectedly high number of

proteins translated from regions within annotated larger genes and even in antisense

orientation[15,30–32,34,35,40,41]. However, thus far these approaches have only been

applied to a small number of model species, and even more rarely as powerful

combination of TIS-Ribo-seq and TTS-Ribo-seq. Given the large diversity among

prokaryotes, substantial insight can be expected from describing the small and

alternative proteome of species from diverse taxonomic ranks.

Previously collected Ribo-seq data for wild type and carbon-starved cells of

Synechocystis 6803 showed allocation of ribosomes in carbon depletion[42] and helped

to characterize the protein turnover at different time points of the light-dark cycle[43]. To

our knowledge, there are no TIS- or TTS-Ribo-seq datasets at all and no Ribo-seq

datasets for any other cyanobacterium available.

Altogether, 165 genes encoding small proteins ≤70 aa are annotated (Kazusa (KZS)
annotation) in the *Synechocystis* 6803 genome (4.5% of all genes). We have recently
characterized five additional small proteins NblD, AtpΘ, AcnSP, SliP4 and NirP1 that
are functionally relevant in the energy and nutrient metabolism of *Synechocystis*
6803[9,17,44–46] demonstrating the immense value of fully unraveling the composition and
functional relevance of the small and alternative proteome.

Here, we report the results of TIS-, TTS-Ribo-seq and proteogenomic analyses in the
model cyanobacterium *Synechocystis* 6803. We present evidence for the translation
of 3,492 and 3,050 annotated genes based on Ribo-seq and proteogenomics, and of
1,069 and 58 previously unknown ORFs, including 935 SEPs of 70 aa or less. We
detected an instance of ribosome frameshifting and small proteins translated from
intragenic out-of-frame alternative reading frames and from antisense RNAs
(asRNAs). We integrated our Ribo-seq data with the annotations from several
reference databases, performed a reanalysis of previously generated proteomic
datasets[47], and *ab initio* gene predictions. This integrated view of several annotations
overlaid with experimental evidence allowed to consolidate many of the genome
annotation differences. The proteogenomic approach, based on a search database
that captures these differences and novel predictions[48] also validated some Ribo-Seq
candidates and identified additional small and larger novel proteins. The consolidated
data provide a significantly updated and extended annotation of the *Synechocystis*
6803 genome that is available for the community through an easily accessible
interactive web-based genome-browser at https://www.bioinf.uni-
freiburg.de/~ribobase/. The here developed generic approach provides an optimal
basis to better exploit the genome sequence information; we expect it to be highly
beneficial for many other model organisms and research communities.

RESULTS

Ribo-seq and retapamulin-based TIS-Ribo-seq captures the translatome of *Synechocystis* 6803

To globally measure translation of annotated ORFs and to discover potential new short ORF encoded proteins (SEPs), we applied Ribo-seq detecting actively translated mRNA segments to *Synechocystis* 6803. In parallel to this standard Ribo-seq approach, we added retapamulin to stall ribosomes on the start codon positions (set-up in **Fig. 1A**) for TIS-Ribo-seq. Cells were rapidly harvested to halt ribosomes along ORFs and preserve polysomes, and lysed, followed by converting polysomes into monosomes and trimming of ribosome footprints by MNase. Polysome profiles generated by sucrose density gradient centrifugation showed successful capture of translating ribosomes as well as their controlled digestion into monosomes (**Fig. 1A**). Following the results of our pre-experiments (**Fig. S1**), we used a concentration of 10 µg/mL retapamulin in 100 mL *Synechocystis* 6803 wild type cultures for TIS-Ribo-seq. We found that retapamulin treatment collapsed polysomes to monosomes even without micrococcal nuclease (MNase) digestion (**Fig. 1A**), as reported for *E. coli*[30], validating our conditions for TIS profiling.

The mapped reads from the standard Ribo-seq analysis revealed the transcriptome-wide ribosome occupancy. Comparison to the RNA-seq coverage inferred in parallel allowed us to calculate the translation efficiency (TE) for a particular transcript (ratio Ribo-seq/RNA-seq coverage). The comparison of TEs between different annotated gene classes showed a much higher mean TE for protein-coding sequences than for sRNAs or 5′ and 3′ UTR sequences (**Fig. 1B**).

Typical examples for the obtained read profiles are shown for gene *psaM* encoding the small PSI subunit XII protein (**Fig. 1C**) and the tricistronic *cpcBAC2* genes encoding the phycobilisome subunits CpcB, CpcA and the rod linker polypeptide CpcC2 (**Fig. 1D**). The profiles show transcriptome coverage starting at the transcriptional initiation sites (TSS) and ending within the 3′UTR downstream the coding region. In contrast, the Ribo-seq read coverage is confined to the coding sections. As reported for MNase-generated Ribo-seq libraries in other bacteria[22,32], the mRNA fragments protected by ribosomes showed a much cleaner, more defined cutoff at the 3′ ends, where the ribosomes' footprints ended compared to the 5′ ends,

where the footprints started (**Fig. S2**). As examples, we show the sharp 3′ ends
observed 19 nt downstream the respective annotated start codons of *psaM* (**Fig. 1C**)
and the *cpcB, cpcA* and *cpcC2* genes (**Fig. 1D**). To generalize these observations, we
performed a meta-analysis of footprints enriched in TIS-Ribo-seq near annotated start
codons (ATG, TTG, GTG). This revealed an average length of the footprints of
retapamulin-stalled ribosomes of 32 nt, with the vast majority of their 3′ ends 19 nt
downstream the annotated start codons (**Fig. 1E**). A parallel metagene analysis
focusing on short ORFs (sORFs; ≤50 nt) was conducted to confirm that translation
initiation patterns in these shorter sequences did not differ substantially from those
observed in longer ORFs (**Fig. S2**).

Minor peaks for the footprint 3′ ends were observed 2 nt and 4 nt downstream the
annotated start codons. As an example, we have zoomed in the region around the
start codon of *rpl34*, a house-keeping gene encoding the large ribosomal subunit
protein L34. For control, the 3′ ends of cDNA fragments generated from standard Ribo-
Seq (in the absence of retapamulin), are plotted for all three examples (**Fig. 1C, D, E**).

The nature and conservation of possible ribosome binding sites for the initiation of
translation in cyanobacteria is not known. Therefore, we searched for potentially
conserved nucleotides 30 nt upstream and 10 nt downstream of the start codons,
focusing on those genes contributing to the read density at the specific offsets
(+2,+4,+19) and filtered them using a 10 read cutoff. This revealed a clear preference
for ATG over GTG as start codon, and a slight preference for 2 preceding pyrimidine
residues. A purine-rich motif that would indicate a likely ribosome binding site was
found in only a subset of genes (**Fig. 1F**).

Overall, metagene analysis and single-gene inspection suggested successful set-up
of TIS-Ribo-seq profiling in *Synechocystis* 6803, an average footprint length of
retapamulin-stalled ribosomes of 32 nt, with their 3′ ends mapping with an offset of 2
nt and 4 nt, but mainly 19 nt downstream the annotated start codons. A purine-rich
motif suitable as ribosome binding site was observed only for a subset of genes (**Fig.**
1F).

Establishing TTS-Ribo-seq in a cyanobacterium

While the location of stop codons of annotated protein-coding genes is uncontroversial, their comprehensive analysis and precise localization is highly valuable for accurately calling previously overlooked translated reading frames. Translation termination site (TTS)-Ribo-seq is a method to infer stop codon positions based on the accumulation of ribosome footprints in the presence of inhibitors of termination[32,39,40]. This method has not been established yet for cyanobacteria. To facilitate TTS-Ribo-seq, we used the inhibitor Api137 that traps the release factors during translation termination, impeding the natural process of peptide liberation[38]. The experimental set-up is shown in **Fig. 2A**. In contrast to the polysome depletion in retapamulin-treated cultures (**Fig. 1A**), the gradient profile of Api-treated cells did not change much in terms of polysome abundance (**Fig. 2A**). Remaining disome peaks were also observed quite frequently in samples digested with MNase, even without Api137. We reasoned that this was most probably due to missing space between two adjacent ribosomes which would be necessary for the digestion of mRNAs, an effect that appeared pronounced in the presence of Api-137 stalling ribosomes at termination codons.

The 3′ ends of sequenced footprints from treated samples compared to the Ribo-seq control were mapped for the TTS-Ribo-seq analysis. As examples, we show the sharp 3′ ends observed 14 nt downstream of the respective stop codons of *norf1* (**Fig. 2B**), and the dicistronic *slr1648* and *ssr2754* genes encoding the carotenoid oxygenase SynDiox2[49], and a YefM-type antitoxin[50] (**Fig. 2C**). Note that the inclusion of TIS-Ribo-seq data allowed the precise definition of the *norf1* reading frame (**Fig. 2B**) that had not been annotated in any of the used reference genome annotations (KZS, GBK, RefSeq; see Methods). This gene has meanwhile been identified to encode the 48 aa ATP synthase inhibitor AtpΘ[17].

Meta-analysis of footprints enriched in TTS-Ribo-seq from isolated monosomes revealed that the majority of their 3′ ends mapped 14 nt downstream of the respective stop codons (**Fig. 2D**). For illustration, we show the stop codon region of *petB* encoding the cytochrome b6 protein involved in photosynthetic electron transport (**Fig. 2D**). The analysis of cDNA 3′ ends enriched in TTS-Ribo-seq from prepared disomes

revealed peaks mapping 14 nt downstream of the respective stop codons as well (**Fig.**

2E), exemplified for *psbN* encoding the 43 aa photosystem II protein PsbN.

To summarize, the 3′ ends of the footprints obtained from Api137-stalled ribosomes

were observed on average 14 nt downstream of the respective stop codons. Metagene

analysis and single-gene inspection suggested successful set-up of TTS-Ribo-seq

profiling in *Synechocystis* 6803.

Validation by proteogenomic analysis and database visualization

The combination and filtering of all TIS- TTS- and classical Ribo-seq data yielded

2,711 instances of potentially novel protein-coding regions or suggested corrections

of annotated genes (1,069 unannotated, 1,190 Internal-Out of-Frame (IOF), 305

Internal-In-Frame (IIF), 88 longer and 59 shorter proteoforms). The majority of these

instances were short protein-coding genes (**Fig. S3, Table S1, S2**). After manual

inspection of 808 instances from this list (30%) and based on the Ribo-seq coverage,

we considered 148 of those to be correct (18%). Among these 148 candidates, we

found 80 unannotated CDS, 46 IOF, 9 IIF, 6 N-terminal extensions and 7 possible N-

terminal truncations. 114/148 (77%) encode proteins of 70 or less aa.

Because manual curation is subjective, we attempted to also leverage the large

amount of proteomics data acquired for *Synechocystis* 6803[47]. For this, we created

integrated proteogenomics databases (iPtgxDBs) based on the *Synechocystis* 6803

chromosome and plasmid sequences and searched publicly available mass

spectrometry datasets[47] against them (**Fig. S4**). For the initial Ribo-seq data mapping,

we used an in-house curated genome sequence based on the KZS annotation that

contained all previously mapped sRNAs, transcriptional units, 5′ and 3′ UTRs that were

defined in previous comparative differential RNA-seq analyses over ten different

growth conditions[51]. Moreover, to distinguish differences in gene calling and

annotation, we integrated the annotations according to NCBI RefSeq 2022 (ID:

GCF_000009725.1) (2022) and GenBank (2016). Despite being based on the same

genome sequence, these differed substantially (**Fig. S5**). Furthermore, to capture the

protein-coding potential comprehensively[48], we also included two *ab initio* gene

predictions (Prodigal 2.6.3[52] and ChemGenome 2.1[53]), and an *in-silico* ORF prediction

that considers the main alternative start codons. These were hierarchically integrated
into a standard iPtgxDB[48] and a smaller custom iPtgxDB (see Methods) that are
accessible via the platform at https://www.bioinf.uni-freiburg.de/~ribobase/.

Altogether, from the list of 165 KZS-annotated small proteins (**Table S3**), 138 (83.6%)
were considered expressed in our dataset based on the Ribo-seq data (RIBO-WT-
avg_TE was ≥0.288), of which 97 had support from TIS-Ribo-seq. In addition, peptide
evidence was observed in the proteogenomic searches for 63 of the 165 SEPs
(38.2%) (**Table S3**), all of which had some form of Ribo-seq support. Previously, Baers
et al. had validated 47 small proteins[54]. Our proteogenomic dataset missed 6 of those
(*ssl0312, ssr0854, ssl0511, hliD, ssl5096* and *MYO_RS19000*), which however, were
all covered by the Ribo-seq data. Thus, based on our combined data, 138 annotated
SEPs are translated, almost tripling the previously available experimental evidence for
47 annotated and validated SEPs. Among the 27 remaining sORFs lacking evidence
for translation, 12 are pseudogenes, mainly fragments of transposase genes.

To identify novel proteins or those unique to one of the annotations, we performed
searches against the standard and custom iPtgxDB. Overall, 11 proteins were
identified that were annotated in KZS and/or GenBank, but not in the latest RefSeq
annotation, i.e., examples of CDS that got wrongly removed from the genome
annotation (**Fig. S6**). Searches against both iPtgxDBs identified 7 of these 11 proteins
as novel (compared to RefSeq) that were annotated by KZS and GBK: 4 of these were
classified as pseudogenes in RefSeq and 3 were removed (one being 4 amino acids
shorter in KZS compared to GBK). Both iPtgxDBs uniquely identified one KZS-only
annotated protein missing from RefSeq, while the custom iPtgxDB identified 3 more
GBK-only annotated proteins, two of which were annotated as pseudogenes in
RefSeq. The other 58 proteins identified through searches of the iPtgxDBs were
entirely new (**Fig. S6**). The custom iPtgxDB uniquely uncovered 22 proteins with Ribo-
seq evidence, while the standard iPtgxDB uniquely found 27 *in-silico* ORFs (not
contained in the custom iPtgxDB), and 6 were predicted by both. Finally, two
predictions from ChemGenome and one from Prodigal were only found using the much
larger standard iPtgxDB, again illustrating the value of integrating several
annotations[48].

TIS- and TTS-Ribo-seq permit start codon re-annotation and the detection of ribosome frameshifting

We observed that the annotation of reading frames and their respective start codons frequently differs between the considered annotation files, although these were based on identical genome sequences. This can cause substantial problems if proteins of interest are expressed only in a truncated version or contain unannotated N-terminal extensions. These problems can be resolved by considering the TIS- and TTS-Ribo-seq data and consolidating the different genome annotations on a case-by-case basis.

We illustrate this here for the gene *sll0553* encoding an alpha/beta fold hydrolase, that is widely conserved in cyanobacteria. The annotated longest gene model in GBK is 1,038 nt, encoding a 345 aa homolog with a GTG start codon (WP_010873707.1), while the corresponding gene model in the KZS annotation suggests a 306 aa protein. In contrast to both annotations, the Ribo-seq coverage and TIS-Ribo-seq mapping data indicated a reading frame encoding a 298 aa protein. DNA and protein sequence alignments show that the here reported start codon is conserved among different *Synechocystis* strains and matches the annotations of homologs in *Synechocystis* strains PCC 6714 (hereafter *Synechocystis* 6714) and 7339 (**Fig. 3A**). We further noticed that the protein length of 298 aa is supported by 19 different peptides from the proteogenomic analysis, while no peptides were found that support the longer N termini annotated in the GBK and KZS annotations. Hence, based on the combined evidence, the gene model for *sll0553* should be truncated by 47 codons.

As a second example, we show the gene *sll0872,* annotated by both GBK and KZS to encode a 130 aa protein (BAL28778.1). However, the very closely related homolog D082_22070 in *Synechocystis* 6714 with its annotated 68 aa is much shorter. Here, our Ribo-seq data indicate that likely the annotations in both strains are incorrect. Ribo-seq coverage and a TIS signal 19 nt downstream of another suitable start codon suggest that Sll0782 is a 101 aa protein in *Synechocystis* 6803 (**Fig. 3B**), consistent with 9 detected peptides, while none was found matching the annotated 29 codons of the N-terminus before the here defined start. Sequence alignments show that a start codon at a matching position exists also in *Synechocystis* 6714, for which consequently the gene model for D082_22070 should be extended by 33 aa, yielding

101 aa proteins in both strains (**Fig. 3B**). Thus, the data suggest that the *sll0872* gene
model should be truncated by 29 codons.

The predictive power of TIS- and TTS-Ribo-seq profiling is furthermore evident in the
region around genes *slr1220* and *slr1222* encoding unknown proteins. TIS-Ribo-seq
in combination with Ribo-seq coverage supported an N-terminal extension of Slr1222
by 116 aa and initiation from a rare TTG codon, yielding a 321 aa protein **(Fig. S7A)**.

In many different bacteria, translation of *prfB* encoding release factor 2 (RF2 or PrfB)
includes ribosomal frameshifting[55–57]. This *prfB* ribosomal frameshifting has not yet
been demonstrated in cyanobacteria. However, the identification of two peptides by
mass spectrometry provided proteogenomic support for the translation of an 26 aa
SEP located upstream of *prfB*[47], while the annotated *prfB* gene (*sll1865*) appears
truncated and starts from a rare TTG codon (**Fig. 3C**). Here, we noticed abundant
signals in both the Ribo-seq and RNA-seq coverage exactly upstream of the likely
pausing site (**Fig. 3C**). The encoded release factor 2 mediates termination at UAA and
UGA codons. Almost all described frameshifting instances in other bacteria involve
CUU UGA as the shift site/stop codon. A cytosine base 3′ of the UGA stop codon was
shown in *E. coli* to make UGA a relatively weak termination codon, contributing to the
frameshifting[58]. These bases – CUU UGA C – are also found in the C-terminal region
of the 26 aa sORF here. In the presence of high PrfB concentrations, most ribosomes
terminate at the UGA stop codon and only the 26 aa peptide is formed. If the
concentration of PrfB is low, the CUU codon can detach from the anticodon of peptidyl-
tRNA$_{Leu}$, mediating a shift from the +1 to the +2 frame[57]. Further described elements
involved in the frameshifting are a short spacer upstream of the shift site (often UAU;[57],
here CUAU), upstream of which an internal Shine-Dalgarno sequence is located,
consistent with the TIS signal at the pausing site. All of these elements are also present
here **(Fig. 3C)**. However, direct proof for the ribosomal frameshifting comes from an
identified peptide (LGQTQDYLDLPG) that directly spans the frameshifting site **(Fig.**
3C). This data corroborates the presence of this naturally occurring frameshift in
cyanobacteria and provides a view of how such a site appears in ribosome profiling
data. The abundant Ribo-seq coverage and TIS signals at the shift site provide direct
evidence for ribosome stalling upstream of the termination codon and re-initiation at
the internal SD element as evidenced from the TIS signal.

We conclude that the generated Ribo-seq data are useful to spot ambiguities
concerning functional start codons and support gene models that should be truncated
or N-terminally extended. They furthermore illustrate an instance of ribosomal
frameshifting in cyanobacteria.

Ribosome profiling aids the discovery of novel toxin-antitoxin candidates

Our ribosome profiling data allowed us to identify several novel protein-coding genes.
Names were assigned conferring to their location in non-coding areas according to the
location of the forward or reverse strand (*ncr* or *ncl*), their transcriptional unit (TU), or
their location within an annotated gene. Two previously identified transcripts, Ncl1450
and Ncr1470, were thought to be two non-coding sRNAs complementary to each
other[51]. Their origin from the same locus in forward and reverse orientation, their
upregulation under low carbon[51], as well as the arrangement of *ncl1450* resembling
an excludon[59] with regard to adjacent genes, prompted further investigation.
Intriguingly, Ribo-seq coverage, TIS, and TTS signals indicated the translation of both
Ncl1450 and Ncr1470 (**Fig. 4A**). Fusions of the reading frames with the P*petE* promoter
led to their expression detected in western blots (**Fig. 4B**). Ncl1450 homologs are
found in several other cyanobacteria (**Fig. 4C**), including *Pleurocapsa*, *Stanieria* (both
53% aa identity) and *Cyanothece* (50% aa identity), while *ncr1470* is only found in
Synechocystis strains (**Fig. 4E**). Ncl1450 is a 36 aa protein that was predicted as
containing a single transmembrane domain by DeepTMHMM[60]. Corresponding
fractionation experiments into soluble and membrane fractions and comparison with
the membrane protein PetC supported a membrane localization for Ncl1450 (**Fig. 4D**).
The combination of a gene encoding a single transmembrane protein with an asRNA
resembles the architecture of type I toxin-antitoxin systems[61]. We conclude that
Ncl1450 and Ncr1470, previously considered as two non-coding sRNAs
complementary to each other, encode SEPs of 36 and 40 aa and appear as a
derivative of a type I toxin-antitoxin system.

An even more compelling case is provided by *ncr1610* previously considered as an
sRNA[51]. Both TIS- and TTS-Ribo-seq data suggested that the *ncr1610* gene contains
two sORFs encoding small proteins of 71 and 34 aa (**Fig. 4F**). The coding potential of
sORF1 was verified by the detection of 4 peptides from the search against the iPtgxDB

and immunologically by FLAG tagging and western blot detection (**Fig. 5A**). Homologs to *ncr1610-sORF1* exist throughout the cyanobacterial phylum, in Pseudomonadota, Chloroflexota and other bacteria (**Fig. S7B**), while no conservation was found for *ncr1610-sORF2.* The *ncr1610-sORF1* start codon overlaps the stop codon of the preceding gene *slr1421* indicating a tight genetic association between the two genes. Searches using Foldseek[62] suggested that Slr1421, a thus far uncharacterized DUF4258 protein, is a structural homolog of MqsR (**Fig. 4G**), an mRNA interferase toxin[63]. Therefore, the Ncr1610-sORF1 protein is suggested as the corresponding antitoxin.

Toxin-antitoxin proteins are often small, which makes the detection of the corresponding loci particularly difficult. Ribosome profiling aids in the discovery of novel elements of such systems, or entirely new ones. This is illustrated here by the antitoxin candidate Ncr1610-sORF1, and the *ncl1450/ncr1470* locus.

ORFBounder identification of unannotated sORFs

Existing tools for sORF identification from ribosome profiling data have significant limitations for TIS-Ribo-seq and TTS-Ribo-seq analysis. While HRIBO[64] incorporates DeepRibo for ORF detection, this tool is designed for conventional Ribo-seq data and not optimized for TIS or TTS datasets. The only analysis software available at the time for TIS-Ribo-seq data was RETscript[30], which had been applied to TIS peak calling in *E. coli*, but was difficult to adapt to other organisms and lacked any support for TTS data. To address these limitations, we developed ORFBounder, a customizable and efficient tool specifically designed for calling TIS and TTS peaks from the respective Ribo-seq data types, with a particular focus on identifying small ORFs. ORFBounder identifies translation sites based on ribosomal footprint enrichment following retapamulin treatment near potential start codons for TIS data. Using optimal read lengths and P-site offsets determined by metagene analysis (**Fig. 1E**), start codons were detected by TIS-Ribo-seq for 100/165 of the previously annotated sORFs (KZS annotation), which overlapped with the 98 ORFs that had significant Ribo-seq coverage (TE ≥0.288, **Table S3)**. All predictions made by ORFBounder were corroborated by HRIBO and confirmed through manual curation.

The power of combining the Ribo-seq and ORFBounder analysis with proteogenomic
searches was well illustrated in the assignment of start codons. There was a single
ATT start codon annotated (KZS), for gene *ureF/slr1899* encoding urease accessory
protein F, while in the GBK and RefSeq annotations, *ureF* was annotated to start from
a GTG. However, ORFBounder specified yet another, more distal TTG as start with a
very good score, leading to a 243 aa protein, instead of the 217 aa (GBK and RefSeq)
and 180 aa (KZS) forms. This N-terminally extended form was directly supported by a
peptide MDDNLLPLLSGSAQLK identified in the standard iPtgxDB search with a
spectral count of 5. It serves as direct proof for the N terminus, as the genomically
encoded sequence would have been LDDNLLPLLSGSAQLK.

Validation of SEPs

Based on the DeepRibo and ORFBounder predictions, we selected 18 sORFs for
experimental validation in addition to the Ncl1450/Ncr1470 pair and Ncr1610-sORF1
discussed above. Six of these sORFs were intergenic, three were short internal out-
of-frame ORFs (IOFs), three were asRNA-located sORFs, and three were intragenic
IIFs in the same reading frame as the main gene. All sORFs were fused to C-terminal
3xFLAG tags and if these could not be detected, we repeated the experiment using a
SPA tag. Using this strategy, 15 of the 18 candidates were validated by western
blotting (**Fig. 5A**, **Table 1**).

Validated intergenically encoded SEPs

The genes *sml0012* and *norf2* were suggested before to encode hypothetical
proteins[13,51,65], but have never been validated. The gene *sml0012* encodes a 45 aa
polypeptide and is located upstream of *aroQ*, the 3-dehydroquinate dehydratase
involved in the shikimate/chorismate pathway, essential for the synthesis of aromatic
amino acids. We validated it by western blotting (**Fig. 5A**) and detection of a single
peptide in the iPtgxDB search (**Fig. S8A**). Homologs of Sml0012 are widespread in
the cyanobacterial phylum, sharing 38 to 88% aa identity (**Fig. S8A**). Sml0012 might
function as an attenuator peptide for *aroQ* translation, as the closely spaced
arrangement of both genes in the same operon is conserved. Translation of the 67 aa
protein Norf2 was well supported by western blotting, TIS- and TTS-Ribo-seq data,

and the detection of 4 peptides in the iPtgxDB search (**Fig. S8B**). There is no homolog

for *norf2* in any other cyanobacterium, but some closely related proteins exist in

gamma proteobacteria (**Fig. S8B**). The region surrounding the *norf2* gene in the

Synechocystis 6803 genome has been characterized as a genomic island[66],

consistent with its paraphyly. The *norf2* gene overlaps by three codons with the

downstream located gene *slr1062* encoding a PemK-type toxin[50] suggesting Norf2 is

possibly the cognate antitoxin.

We also found SEPs on the ~100 kb plasmid pSYSA encoding the three distinct

CRISPR-Cas systems of *Synechocystis* 6803[67]. There are substantial numbers of

accessory proteins encoded with type III CRISPR-Cas gene cassettes[68], and more

such genes might have been overlooked. Indeed, there is an sORF close to the 3′ end

of the CRISPR3 array encoding a 46 aa SEP, here called Ssr70935, supported by

strong Ribo-seq, TIS- and TTS-Ribo-seq coverage (**Fig. S8C**) and by western blotting

(**Fig. 5A**). Ssr70935 is a very basic protein due to the presence of 9 lysine and 8

glutamine residues (isoelectric point (pI) of 12.04), a hallmark of nucleic acid-binding

proteins.

We validated the translation of three previously defined sRNAs, Ncl1350, Ncr1420 and

NsiR7[51]. Ncl1350 and Ncr1420 encode a 57 and a 52 aa SEP, respectively. Homologs

of similar size exist in many different cyanobacteria (**Fig. S9** and **S10**). The transcript

level of *ncl1350* was previously found to be high and almost constitutive under ten

different conditions. The transcription of *ncl1350* is linked with that of the downstream

gene *sll0376* encoding a core protein of the cyanoglobule lipid droplet[69], to which it

may be functionally related. Translation of Ncl1420 was validated by six different

peptides identified in the iPtgxDB. Transcription of *ncl1420* is highest during

exponential growth and very low under adverse conditions suggesting involvement in

vegetative growth. In contrast, NsiR7 is upregulated under nitrogen starvation[70] and

encodes a 51 aa SEP. Putative homologs of NsiR7 were identified only in a small

number of other *Synechocystis* strains, which share the presence of cysteine residues

at conserved positions (**Fig. S10B)** suggesting redox sensitivity.

SEPs translated from asRNAs

Three asRNA-located SEPs were chosen for validation. One of them, *rsbU*-asORF,

was found on an asRNA to *slr2031* encoding a homolog to the protein phosphatase

RsbU[71]. The asRNA to *slr2031* was detected earlier and divergent transcript
accumulation with the *slr2031* mRNA was observed under high light compared to
standard conditions and cultivation in darkness[65]. In the Δ*slr2031* mutant, pigment
contents (Chl and phycocyanin) were significantly higher compared to wild type[71]. This
finding was consistent with reports that spontaneous mutations can be observed in
this gene[72] and that deletion of a 154 bp DNA segment encoding part of the N-terminal
domain of Slr2031 led to a glucose-resistant substrain[73] that also displayed higher
pigment contents and showed an increased high-light tolerance[74]. The RNA-seq and
Ribo-seq signals are stronger for *rsbU*-asORF than for the gene *slr2031* in sense
orientation (**Fig. 5B**). Within this asRNA, we found two sORFs in-frame with each
other, encoding a 62 or a 38 aa peptide. Support from TTS-Ribo-Seq is excellent for
both, since they end with the same stop codon. TIS-Ribo-Seq support is very clear
and at the optimal distance of 19 nt for the shorter sORF. However, a peptide was
found in the custom iPtgxDB search, directly supporting the long variant (**Fig. 5B**) and
expression of the FLAG tagged protein was confirmed by western blotting (**Fig. 5A**).
Therefore, it is also possible that the 38 aa variant is an IOF to the longer sORF.
Corresponding contiguous ORFs exist in some other cyanobacteria including possible
start codons at the corresponding positions for the 62 and 38 aa sORFs in
Synechocystis 6803 (**Fig. 5B**). RsbU-type phosphatases such as Slr2031 commonly
belong to a partner switching system that consists of four modules. These are RsbU,
a phosphorylatable anti-sigma antagonist, a kinase anti-sigma factor and an
alternative sigma factor[75]. This already complex system is made even more complex
by the asRNA and the herein encoded SEP.

We also validated the coding potential in two more asRNAs, in as_csx18 to gene
slr7091, encoding a CRISPR-associated Csx18 protein **(Fig. S11),** and in as_psbC to
gene *sll0851,* encoding the photosystem II protein PsbC/CP43 **(Fig. S12)**. For
as_csx18 and its cognate gene in sense orientation, *slr7091,* both RNA-seq, Ribo-
seq, and TIS-Ribo-seq coverage supported the expression of both DNA strands. The
coding potential within *as_csx18* was well supported by the characteristic TIS-Ribo-
seq signals 3′ of the start codon at position +19 nt. Our TIS-Ribo-seq signals also
indicated that the *slr7091* ORF encodes a 96 aa protein, which is shorter than the
annotation as a 122 aa protein in the KZS, RefSeq and GBK databases, but consistent
with the shorter Prodigal prediction (**Fig. S11A**). The translation of the *as_csx18* sORF

was validated by western blot, detecting the SPA-tagged asCsx18 protein (**Fig. S11B**).
Homologs of the asCsx18 protein could be encoded on asRNAs to the respective
csx18 gene in *Synechocystis* 6714, and possibly further cyanobacteria (**Fig. S11C**).
The asRNA overlapping the 3′ end of the *psbC* coding region originates from a TSS at
position 1,347,388 on the forward strand of the chromosome[51]. Thus, this TSS is
located 86 nt downstream of the *psbC* stop codon, while the asRNA overlaps >300 nt
of the *psbC* coding region, suggesting a regulatory function. Indeed, both transcripts
are divergently regulated. While the *as_psbC* transcript is maximum under adverse
conditions, *psbC* mRNA accumulation is maximum under optimal growth conditions
and exponential growth[51]. Translation of the *as_psbC*-encoded 28 aa SEP was
supported by Ribo-Seq, TIS-Ribo-seq, TTS-Ribo-seq data and western blotting **(Fig.**
S12A, B). The asPsbC protein in *Synechocystis* 6803 and several possible homologs
in other cyanobacteria are very basic (pI of ~12) and very rich in alanine **(Fig. S12C)**.

Translation of SEPs from alternative start sites within coding regions

In addition to SEPs encoded in intergenic regions or within asRNAs, we also found
evidence for the translation of SEPs from alternative start sites within protein-coding
genes Internal-Out of-Frame (IOF). Three examples were selected for validation, *der*-
IOF, *crtR*-IOF and *slr1923*-IOF. The *der*-IOF encodes an only 25 aa SEP, located
within the *der* gene (*slr1974*) for translation initiation factor IF-3 (**Fig. 5C**). Although
translation of the 42 aa SEP originating from the IOF within the *crtR* gene and of the
SEP originating from the IOF within *cvrA* was well supported (**Fig. S13**), their reading
frames appear not as widely conserved. Lastly, we picked three examples of IIF-
encoded SEPs (within *sll0871, slr0601* and *slr1397*), but these could not be validated.

To summarize, we validated 15 of 18 selected sORFs in addition to the Ribo-seq
support by western blots, and five were further supported by the proteogenomic
detection of peptides (**Table 1**). From these 15 validated instances, one was annotated
as a hypothetical protein (*sml0012*), one was a previously predicted protein-coding
gene based on transcriptomic evidence (*norf2*), while the others were known only as
sRNAs or asRNAs, or are antisense intragenic IOF-encoded SEPs. These results
indicated the protein-coding potential of widely different types of loci.

A family of genes encoding two proteins by internal out-of-frame translation

Thus far we have reported on SEPs encoded in intergenic regions, antisense transcripts, or within coding regions. However, the Ribo-seq data also uncovered the existence of a hitherto uncharacterized family of genes that encoded a second, relatively long protein IOF. In **Fig. 6** we show that a 112 aa protein is encoded within the coding sequence for the 151 aa protein Slr0489. From here on, we call these proteins Slr0489L for the longer, annotated gene product, and Slr0489S for the shorter protein, encoded IOF. Reads from the standard Ribo-seq analysis cover the entire annotated *slr0489* gene, but the coverage is visibly higher for the internal segment encoding Slr0489S (**Fig. 6A**). There are TIS-Ribo-seq signals for both reading frames at the characteristic position 19 nt 3′ of the respective start codons, and TTS-Ribo-seq signals mapped 14 nt 3′ of the respective stop codons. These signals are strikingly higher for the internal segment. Ultimate proof for the simultaneous expression of both proteins was obtained by the proteogenomic detection of 6 different peptides for Slr0489L and 5 for Slr0489S (**Fig. 6B, C**).

A search for related genes yielded two additional loci in *Synechocystis* 6803, *slr1079* and *slr1082*. These genes harbor IOF reading frames as well, and Ribo-seq data supported their expression (**Fig. S14**) and the detection of peptides in the iPtgxDB search directly supported the presence of Slr1079L, Slr1079S and Slr1082S (**Fig. S14D, E**). Notably, the internally encoded proteins Slr0489S, Slr1079S and Slr1082S are closely related to each other, as are Slr0489L, Slr1079L and Slr1082L (**Fig. 6B, C**). However, the respective longer and shorter proteins encoded within each of these loci are not related in sequence or structure. While Slr0489S has a slightly acidic pI of 5.56, Slr0489L has a pI of 9.38. Moreover, Slr0489L shares the presence of a signal peptide and a transmembrane domain with Slr1079L and Slr1082L (**Fig. 6B**). The presence of these three genes does not follow phylogeny — the closely related strain *Synechocystis* 6714 lacks a related gene, while in the only distantly related *Synechococcus* sp. PCC 7003, a close homolog exists (75% identical and 87% conserved aa with Slr0489, sequence ID WP_065714473.1).

To search for interacting proteins, we expressed a FLAG-tagged version of Slr0489L from a plasmid vector and performed coimmunoprecipitation (co-IP) analysis compared to a separate co-IP for sfGFP-3xFLAG as control. The most enriched

protein was Slr0489S, suggesting that the internally encoded smaller and the longer
2TM proteins interact with each other (log$_2$-fold enrichment of 10.6 for Slr0489L and
8.8 for Slr0489S, **Fig. 6D**). However, the overexpression of *slr0489* inherently leads
to a higher level of both, Slr0489L and Slr0489S, as they are both encoded on the
same mRNA. Therefore, we constructed a cell line separately overexpressing
Slr0489S fused to a C-terminal 3xFLAG tag (Slr0489S-3xFLAG) and performed co-
IP. In this case, a log$_2$-fold enrichment of 8.3 was observed for Slr0489S and of 6.4 for
Slr0489L (**Fig. 6E**). We noticed that in both cases two other highly enriched proteins
were the CRISPR3 type III-Bv system Cmr5 protein Sll7086[76], and the minor pilin
Slr2018[77] suggesting a connection to complexes close to the cell membrane. We
conclude that the two different proteins produced from the *slr0489* locus interact with
each other. In parallel, we performed the same type of co-IP experiments with Slr1079
(**Fig. S14**). The co-IP using Slr1079L as bait yielded Slr1079S as the most enriched
protein (log$_2$-fold enrichment of 12.1 for Slr1079L and 6.7 for Slr1079S, **Fig. S14B**),
while the co-IP using Slr1079S as bait yielded Slr1079L as the most enriched protein
(log$_2$-fold enrichment of 10.1 for Slr1079S and of 8.8 for Slr1079L, **Fig. S14C**).

To measure possible physiological effects of the Slr0489L/S system, we constructed
a *Synechocystis* 6803 Δ*slr0489* deletion mutant. In this mutant, we then
overexpressed the native *slr0489* gene with intact reading frames for both Slr0489L
and Slr0489S. Alternatively, we expressed only Slr0489S, or only Slr0489L. For the
latter, we introduced a *slr0489* gene with a single point mutation substituting the ATG
start codon by an ACG codon for threonine, thus preventing the translation of
Slr0489S. In a growth experiment lacking the inducer rhamnose, all gene variants had
no or only negligible growth effects compared to wild type (**Fig. 6F**). In this line only,
clear growth retardation was observed in the presence of the inducer rhamnose, while
all other strains grew similarly over a 5-day time course (**Fig. 6F**). We conclude that
the sole expression of Slr0489L had a toxic effect on cell growth, which was
neutralized in the presence of Slr0489S.

DISCUSSION

Genome consolidation effort in *Synechocystis* 6803

Synechocystis 6803 is an important model organism, which is reflected by the fact that its genome was the overall third to be fully sequenced 30 years ago[5]. Nevertheless, similar to other model systems, precise determination of coding regions has remained challenging up to this date, visible in differences in the available annotations of the identical genome sequence, such as the ones used here, deriving from NCBI RefSeq (2022), GenBank (2016) and the KZS 2009 annotation. These annotations differ both in terms of the overall numbers of coding sequences predicted and their precise start sites (**Fig. S5, S6**). As we show in this study, several reading frames should be extended or truncated (**Fig. 3**), which resolves ambiguities concerning functional start codons. Nevertheless, we are aware that ultimate proteomics proof requires in many instances specialized N-terminomics approaches[78], beyond the scope of this work. We have integrated both the different reference annotations and complemented them by *ab initio* gene predictions using Prodigal 2.6.3[52], ChemGenome 2.1[53], and an *in-silico* ORF prediction integrated into a standard and a custom integrated proteogenomic search database (iPtgxDB), which are both accessible through a single platform. The generation of a database comprising different annotations is generic; it enables researchers to jointly consolidate the genome annotation of their favorite model organism in the context of overlaid experimental data. The addition of data from multiple conditions and sources forms the basis to more comprehensively discover the full coding potential of genome sequences. To facilitate access to the entire set of data, we developed a JBrowse 2 instance[79], a web-accessible, fully interactive genome browser provided at https://www.bioinf.uni-freiburg.de/~ribobase/.

While SEPs mediate multiple important functions in all domains of life, they frequently have escaped annotation. Here, the combination of TIS- and TTS-Ribo-seq approaches with a proteogenomic analysis identified hundreds of candidate genes, of which several were experimentally validated to be present (**Table 1, Fig.s 4, 5, S7–S13**). Our discovery of many new SEPs is in line with the recent prediction of extensive microprotein families in the Enterobacteriaceae[16]. More than 140 novel small proteins (in this case even ≤50 aa) have been manually validated over the span of roughly ten years in *E. coli*[80], recent work in *Synechocystis* 6803 provided evidence for 516 SEPs

<100 aa[81], both consistent with our findings. We have previously characterized the

Synechocystis 6803 SEPs AtpΘ as a regulator of ATP synthase[17], NblD as a factor

involved in the nitrogen starvation-induced degradation of the phycobilisomes[45],

AcnSP as a regulator of aconitase activity[44] and NirP1 as regulating nitrite reductase[46]

demonstrating the physiological relevance of SEPs. In extension, many of the newly

identified candidates in this study warrant further consideration. These include the

slr1421-ncr1610-sORF1 locus (**Fig. 4F and G**), or the *ncl1450/ncr1470* locus, which

are related to toxin-antitoxin systems. Our data indicate that a SEP evolved in the

asRNA Ncr1470, which can be detected in some but not in the majority of related loci

in other cyanobacteria possessing homologs of *ncl1450* (**Fig. 4C, E**). The *ncl1450*

gene and its overlapping *ncr1470* are located in an arrangement between *slr0841* and

slr0842 resembling an excludon[59]. The Ncl1450/Ncr1470 pair might constitute an

associated type I toxin-antitoxin system, possibly with a stabilizing role[82]. Secondarily,

the Ncr1470 sORF may have evolved in an originally non-coding asRNA.

CRISPR-associated hidden genes

We present two instances of previously unknown CRISPR-associated sORFs and

corrected the ORF for gene *slr7091/csx18* **(Fig. S8C, S11A)**. The here defined gene

ssr70935 encodes a 46 aa protein. It is located directly between the CRISPR array of

a type III-Bv system[76], and the genes *slr7094* and *slr7095* with which it is cotranscribed

as part of a tricistronic operon. Homologs of *ssr70935* in other cyanobacteria are also

located adjacent to CRISPR-Cas loci, indicating a genetic association. There are

several examples for the relevance of small proteins encoded in atypical sORFs

associated with CRISPR-Cas systems. In the archaeal CRISPR-Cas type I-B system

of *Haloferax volcanii*, the Cas11b is encoded by an internal in-frame translation within

the *cas11* gene[83]. In a study on the type I-D system of the here investigated

Synechocystis 6803, cryo-EM structures of the purified complex revealed an additional

small subunit Cas11d[84]. Initially, no gene could be assigned to this protein until it was

discovered that it originated from an IIF translation within the gene of the large subunit

cas10d[84]. Therefore, the here identified asCsx18 protein translated from an asRNA

further illustrates the coding capacity of CRISPR-Cas systems to be extended by

translation of ORFs from unorthodox locations. While currently neither the precise

function of Csx18 nor of the asCsx18 protein is known, we noticed that Csx18 proteins

are generally associated with type III-D and III-B CRISPR systems and likely

adaptation-associated[68]. This would be consistent with the location of *slr7091* and its

asRNA just upstream of the *cas1* and *cas2* adaptation module and next to the

CRISPR3 effector genes on plasmid pSYSA in *Synechocystis* 6803[76].

Genes-in-genes

Our work not only provided evidence for several novel sORFs, but also uncovered the

existence of a hitherto uncharacterized, novel family of 3 genes encoding a second,

relatively long protein out-of-frame, which may be considered as genes-in-genes (**Fig.**

6, Fig. S14). This genetic arrangement resembles some bacteriophage loci encoding

two-component spanins. Spanins are lysis proteins produced by phages that are

required to disrupt the outer membrane during lysis of gram-negative host cells. Two-

component spanins consist of an integral inner membrane i-spanin, and an outer

membrane o-spanin, which is a lipoprotein[85]. In bacteriophage lambda, the reading

frame Rz1 encoding the o-spanin is embedded within the reading frame Rz encoding

the i-spanin[85]. The lambda phage i-spanin possesses a transmembrane domain and

is located in the inner membrane. The C-terminal domain of the i-spanin, which

extends into the periplasm, consists of alpha helical segments. In contrast, the o-

spanin possesses a proline-rich region, is connected to the outer membrane, lacks a

transmembrane domain and any other detectable structure[86]. Therefore, there are

striking parallels to the here described *slr0489, slr1079* and *slr1082* loci. In addition to

the genetic arrangement, the three proteins Slr0489L, Slr1079L and Slr1082L share

the presence of a signal peptide and a transmembrane domain (**Fig. 6B**), while

Slr0489S, Slr1079S and Slr1082S encoded by the embedded ORFs possess a

conserved proline-rich motif at positions 6–8 (**Fig. 6C**), recognized as a membrane

fusion motif in o-spanins[85].

However, there are also important differences. These three loci do not belong to any

bacteriophage or prophage genome and their overexpression did not cause the cells

to lyse. However, we did observe that the expression of Slr0489L without Slr0489S

led to growth retardation (**Fig. 6F**). Thus, the tight genetic connectivity between

Slr0489L and Slr0489S first had to be disentangled to observe this effect, resembling

a toxin-antitoxin effect in which binding of Slr0489S negates the deleterious effect of
Slr0489L on our cultures. This resembles rather a defense-associated toxin–antitoxin
relationship, as in the Panoptes defense system[87,88], providing an exciting topic for
further research.

Outlook and limitations of this work

There are two previously published instances of alternative proteins produced from
single genes in *Synechocystis* 6803. These are the IIF-encoded CRISPR-Cas subunit
Cas11d[84] and the two alternative forms of ferredoxin:NADP oxidoreductase[89].
Translation of the corresponding genes *slr7011* and *slr1643/petH* can be seen in the
Ribo-seq data, but not the initiation of translation at the internal sites. This points at a
sensitivity limit and that we rather under- than overestimate the number of alternative
ORFs. Part of this may relate to the fact that we have established TIS- and TTS-Ribo-
seq and performed that analysis under one condition only. The shorter
ferredoxin:NADP oxidoreductase proteoform is mainly made under heterotrophic
conditions[89], which were not included here. Nevertheless, we provide information
about 2,711 instances of potentially novel protein-coding sections or suggested
corrections of annotated genes. This number includes more than 1,000 candidates
each for SEPs from loci considered non-coding, or IOF-produced SEPs and we
validated several instances related to the CRISPR-Cas systems, genes encoding core
proteins of photosystem II, or regulatory factors. The functional analysis of these SEPs
as well as of the instances of longer genes-in-genes is an exciting topic for further
research.

METHODS

Strains and growth conditions

Synechocystis 6803 cultures were grown in TES-buffered BG11[90]. One hundred mL
cultures were grown in 250 mL Erlenmeyer flasks with shaking under an illumination
of 40 µmol photons m^{-2} s^{-1} and a temperature of 30°C until an OD_{750} of 0.6-0.8 was
reached. For TIS profiling, 10 µg/mL retapamulin final concentration (approx. 10-fold
MIC) solved in DMSO was added to the culture and incubated for 15 min with

continued cultivation. For control, an equal volume of DMSO was added (mock-treated
cultures). For TTS profiling, the cultures were treated with 10 µM final concentration
(approx. 10-fold MIC) of the proline-rich antimicrobial peptide Apidaecin 137 (Api,
solved in ddH$_2$O, NovoPro BioSciences, Shanghai) for 15 min. Mock-treated controls
without the antibiotic were grown in parallel. Two independent biological replicates
were prepared and processed for Ribo-seq, TIS and TTS profiling experiments.
Immediately before cell harvest, aliquots were taken from the untreated cultures for
total RNA analysis, mixed with 0.2 vol stop mix (5% buffer-saturated phenol (Roth) in
95% ethanol), and snap-frozen in liquid N$_2$.

MIC assays for TIS and TTS profiling

In *E. coli* TIS profiling, mutants lacking the gene for efflux transporter *tolC* were used
because translation stalling effects were limited due to retapamulin secretion via the
transporter in the wild type strain[30]. A homologous transporter is also present in
Synechocystis 6803[91,92] and might be responsible for the efficient efflux of
retapamulin. This is unfavorable for TIS profiling, since the appropriate antibiotic levels
need to be maintained in the cell to stall initiating ribosomes. Minimum inhibitory
concentrations (MIC) of retapamulin (Sigma CDS023386) were determined in liquid
cultures for the wild type and the Δ*tolC,* Δ*acrA* and Δ*acrB* transporter mutants. Added
retapamulin inhibited all three mutants, while the wild type resisted retapamulin
concentrations up to 1 µg/mL (**Fig. S1A, B**). Without antibiotics, all strains grew
steadily from an OD$_{750}$ of 0.2-0.4 to > 2 within 3 days. To avoid possible
unphysiological effects of working with mutant strains, we chose the wild type for
retapamulin-assisted TIS profiling, applying a ~10x MIC of 10 µg/mL retapamulin in
100 mL cultures. To establish proper conditions for TTS-Ribo-seq, MIC were
determined using the proline-rich antimicrobial peptide (prAMP) Oncocin 112
(Onc112, Novopro) in small scale liquid cultures. Onc112 is similar in chemical nature
(sequence   VDKPPYLPRPRPPRrIYNr-NH$_2$)   compared   to   Api137   (Gu-
ONNRPVYIPRPRPPHPRL-OH   (Gu=N,N,N',N'-tetramethylguanidine   and   O=L-
ornithine))[37], but is less costly. Both prAMPs are supposed to be taken up by a SbmA-
like transporter[93]. A MIC of 0.8 µM Onc112 was determined for *Synechocystis* 6803
wild type (**Fig. S1C**). Consequently, a 12.5-fold MIC of 10 µM Api137 was applied in
the TTS approach.

Ribosome Profiling

Cells were collected by fast filtration through hydrophilic polyethersulfone filters (Pall Supor®-800, 0.8 µm). The cells were scratched off the filters, quickly frozen in liquid nitrogen and stored at -80°C. Cells were then disrupted by cryogenic grinding (Retsch M-400) using prechilled grinding ball and chamber in the presence of 1 mL frozen lysis buffer (50 mM HEPES pH 8, 100 mM NaCl, 10 mM $MgCl_2$, 5 mM $CaCl_2$, 0.4 % Triton X-100, 2 mM DTT, 500 U RiboLock). In contrast to other approaches, chloramphenicol has been omitted in all buffers to avoid bias[22]. The grinding (15 Hz, 3 min) was repeated four more times. The grinding jars were cooled with liquid nitrogen between cycles. Frozen powder was collected and then centrifuged (4°C, 8,000 × g, 10 min) to separate cell debris and lysate. The lysate was kept on ice for further processing, or stored at - 80°C. For the digestion, an aliquot containing ~100 µg nucleic acid (quantified with nanodrop) was filled up to a volume of 500 µL with digestion buffer (50 mM HEPES pH 8, 100 mM NaCl, 10 mM $MgCl_2$, 10 mM $CaCl_2$, 0.4% Triton X-100, 2 U/mL TURBO DNase, 100 U/mL RiboLock) and digested with 300 U MNase (NEB) for 45-60 min at 25°C, 450 rpm. A negative control was treated the same way, but MNase was omitted. To stop the digestion, 5 mM EGTA (final concentration) was added. The lysate was immediately layered on a sucrose gradient with 10%-55% sucrose solved in gradient buffer (100 mM NH_4Cl, 20 mM Tris-HCl pH 8, 5 mM $CaCl_2$, 10 mM $MgCl_2$, 2 mM DTT). Gradients were formed by Gradient Master (Biocomp) in 1:50 min, with 80° angle and 21 rpm, at 4°C, and then centrifuged for 2.5 h, 217k × g at 4°C in a Beckman Coulter Optima L-80 XP ultracentrifuge and SW40 Ti rotor), the gradients were fractionated and the absorbance at 260 nm was measured with Gradient Fractionator (Biocomp) integrated TRIAX flow cell. For TIS profiling, the monosomes (70S) fractions were collected for both the untreated (Ribo-seq) and Retapamulin-treated samples (TIS). For TTS profiling, fractions corresponding to monosomes, disomes as well as trisomes were collected for both untreated (Ribo-seq) and Api-treated samples (TTS). The collected fractions were immediately frozen in liquid N_2 until further analysis.

Isolation of ribosome footprints

Ribosome footprints were prepared from monosomes, disomes, and trisomes. Total RNA was extracted from fractions or cell pellets using hot phenol-chloroform-isoamyl

alcohol (25:24:1, Roth) or hot phenol (Roth), respectively[94]. Ribosomal RNA (rRNA)
was depleted from 5 µg of TURBO DNase-digested total RNA by subtractive
hybridization with the Pan-Prokaryotes riboPOOLs (siTOOLs, Germany) according to
the manufacturer's protocol with Dynabeads MyOne Streptavidin T1 beads
(Invitrogen). Total RNA was fragmented with an RNA fragmentation reagent (Ambion)
following the manufacturer's instructions. Monosomes RNA and fragmented total RNA
were size selected (26–34 nt) on 15% polyacrylamide (PAA), 7 M urea gels, as
previously described[95] using RNA oligonucleotides NI-19 and NI-20 as ladder. In
parallel, RNA from disomes (50-80 nt) and trisomes (80-120 nt) from both untreated
and Api-treated samples were size selected on the same gel electrophoresis as
indicated above using Ultra-low RNA ladder (RNA Decade Markers, #AM7778,
ThermoFischer). RNA was extracted from the gel, cleaned and concentrated by
isopropanol precipitation with 15 µg of GlycoBlue (Ambion) and dissolved in H_2O.
cDNA libraries were prepared by Vertis Biotechnologie AG (Freising, Germany) using
the adapter ligation protocol without fragmentation. First, an oligonucleotide adapter
was ligated to the 3′ end of the RNA molecules. First-strand cDNA synthesis was
performed using M-MLV reverse transcriptase and the 3′ adapter as the primer. The
first strand of cDNA was purified, and the 5′ Illumina TruSeq sequencing adapter was
ligated to the 3′ end of the antisense cDNA. The resulting cDNA was PCR-amplified
to approximately 10–20 ng/µL using a high-fidelity DNA polymerase. The DNA was
purified using the Agencourt AMPure XP kit (Beckman Coulter Genomics) and
analyzed by capillary electrophoresis. The primers used for PCR amplification were
designed for TruSeq sequencing according to the instructions of Illumina. The
following adapter sequences flank the cDNA inserts: TruSeq_Sense_primer:
(NNNNNNNN = i5 Barcode for multiplexing) 5′-
AATGATACGGCGACCACCGAGATCTACAC-NNNNNNNN-ACACTCTTTCCCTACA
CGACGCTCTTCCGATCT-3′; TruSeq_Antisense_primer: (NNNNNNNN = i7 Barcode
for multiplexing) 5′-CAAGCAGAAGACGGCATACGAGAT-NNNNNNNN-
GTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT-3′. cDNA libraries were pooled
on an Illumina NextSeq 500 high-output flow cell and sequenced in single-end mode
(75 cycles, with 20 million reads per library) at the Core Unit SysMed at the University
of Würzburg.

Ribo-seq, TIS and TTS profiling data analysis

Ribo-seq and TIS/TTS datasets were processed using the HRIBO workflow (version 1.7.0)[64], which has previously been used for the analysis of other prokaryotic Ribo-seq data[34,35,96]. Sequencing read files were processed with a Snakemake[97] workflow that downloads all required tools from bioconda[98] and automatically determines the necessary processing steps. Adapters were trimmed from the reads with cutadapt (version 2.1)[99] and then mapped against the *Synechocystis* genome with segemehl (version 0.3.4)[100]. Reads corresponding to rRNA and other multiple mapping reads were removed using SAMtools (version 1.9)[101]. Quality control was performed by creating read count statistics for each processing step and RNA class with Subread featureCounts (1.6.3)[102]. All processing steps were analyzed with FastQC (version 0.11.8)[103], and the results were aggregated with MultiQC (version 1.7)[104].

Read coverage files were generated with HRIBO using different full-read mapping approaches (global or centered) and single-nucleotide mapping strategies (5′ or 3′ end). Read coverage files using two different normalization methods were created (mil and min). For the mil normalization, read counts were normalized by the total number of mapped reads within the sample and scaled by a per-million factor. For the min normalization, the read counts were normalized by the total number of mapped reads within the sample and scaled by the minimum number of mapped reads among all analyzed samples. The coverage files generated using the min normalization and the global mapping (full read) approach were used for genome browser visualization.

Metagene analysis of ribosome occupancy

A genome wide analysis of ribosome occupancy was conducted using TIS-Ribo-seq and TTS-Ribo-seq. To this end, all annotated coding sequences (CDS) were collected, and windows consisting of upstream and downstream regions around each annotated start codon (or stop codon, respectively) were defined. For each position in the analyzed window, reads occupying that position were counted for each annotated CDS. The total number of reads per position was subsequently reported and plotted in a metagene graph. To generate a high-confidence set of genes, multiple filters were applied. These filters removed genes in close-proximity ($\leq$50 nt apart), genes below a coverage threshold of 10 RPKM and genes that fell below a minimum length cutoff (i.e., genes shorter than the downstream window for start codons or shorter than the

upstream windows for stop codons). However, we performed a parallel metagene
analysis focusing on 39 short ORFs (≤50 aa) that confirmed that translation initiation
patterns in these shorter sequences did not differ substantially from those observed in
longer ORFs (compare **Fig. 1E** and **Fig. S2**).

To assess sequence motifs in the 5'-UTR regions of annotated start codons, all
annotated CDS contributing to positions with the highest ribosome occupancy (at
offset of +2, +4, and +19) were collected, and sequence windows were extracted. Motif
plots were then generated using weblogo (3.7.9)[105].

All scripts used for metagene analysis are available in our Github repository
(https://github.com/RickGelhausen/Synechocystis_pub_scripts_2025/).

Ribo-seq ORF predictions using HRIBO modules

ORFs were called with an adapted variant of REPARATION[106] using blast instead of
usearch (see https://github.com/RickGelhausen/REPARATION_blast) and
DeepRibo[107]. Generic feature format (GFF) track files with this information, plus
potential start and stop codons and ribosome binding site information were created for
in-depth manual genome browser inspection. Summary statistics for KZS and
GenBank annotated and merged novel ORFs detected by REPARATION and
DeepRibo were computed in a tabularized form, including, among other values,
translation efficiency (TE), RPKM (reads per kilobase of transcript per million mapped
reads) normalized read counts, codon counts, and nucleotide and aa sequences (see
Table S1 for annotated genes and **Table S2** for newly predicted candidates or known
genes with suggested corrections). Annotated sORFs were classified as translated if
they fulfilled an empirical mean TE cut-off of ≥ 0.288 and RNA-seq and Ribo-seq
RPKM of ≥ 10 (cut-offs chosen based on the lowest TE and RPKM values associated
with housekeeping genes [i.e., ribosomal protein genes] and the genes detected by
proteomics). To identify robust novel sORF candidates based on Ribo-seq coverage,
we inspected HRIBO ORF predictions from DeepRibo and REPARATION. As
DeepRibo is prone to a high rate of false positives[108], we first generated a reasonable
set of potential novel sORFs by applying the following expression cut-off filters: mean
TE of ≥0.288 and RNA-seq and Ribo-seq RPKM of ≥40 (in one of the replicates). In
addition, novel translated sORF candidates were required to have a DeepRibo
prediction score[107] of ≥-3.009 that allows for ORF candidate ranking. The filtered

sORFs were then subjected to manual curation as described[108]. This parallel
inspection of Ribo-seq and RNA-seq read coverage files in a genome browser allowed
for asserting the translation status for the filtered novel predicted sORFs. Briefly, the
Ribo-seq and RNA-seq read coverage files were loaded in the Integrated Genome
Browser (IGB) along the sequence of the reference genome and the Kazusa 2009
annotation (KZS; with annotated 5'- and 3'-UTRs and further manual curation from the
Hess laboratory). The Ribo-seq and RNA-seq read coverage files (normalized to the
lowest number of reads between the two) were visually inspected with similar scales.
To assess translation of the predicted novel sORFs, we used the following criteria: 1)
Ribo-seq read coverage within ORF boundaries with the detection of ribosome
footprints in the UTRs (15-16 nt) near the start and stop codons resulting from initiating
and terminating ribosomes; 2) exclusion of Ribo-seq read coverage from the residual
5'- and 3'-UTRs; 3) the shape of the Ribo-seq read coverage; here the evenness of
the read coverage was considered and predicted sORFs with uneven read coverage
(exhibiting peaks with plateau, which resulted from either RNA structures or cDNA
library preparation artifacts) were not taken into account; 4) the Ribo-seq read signal
was generally required to be comparable to or higher than the transcriptome signal
from the RNA-seq library. For visualization, we developed an interactive web-based
genome browser using JBrowse 2 (https://www.bioinf.uni-freiburg.de/~ribobase/)[109],
where the coverage files for the Ribo-seq replicates, the different genome annotations,
results of proteogenomic mappings and the predicted sORF can be accessed
interactively.

TIS and TTS peak calling, ORF predictions

To define a set of potentially novel ORFs, the three different datasets resulting from
(1) classical riboprofiling (Ribo-seq), (2) TIS-Ribo-seq and (3) TTS-Ribo-seq were
filtered separately. Then, the filtered results were combined; for an ORF to be called,
it had to be detected by Ribo-seq and at least one of the other two methods (TIS or
TTS-Ribo-seq). The three datasets were filtered as follows:

(1) Ribo-seq data were analyzed using the workflow HRIBO (version 1.4.3)[64]. ORFs
were initially called with two algorithms, REPARATION[106], but using blast instead of
usearch [34,35], and with DeepRibo[107]. Summary statistics included TE, RPKM
normalized read-counts and codon counts. The nt and aa sequences for all available

annotated and merged novel ORFs detected by REPARATION and DeepRibo were
computed in a tabular format (**Tables S1** and **S2**). For benchmarking, we used a
dataset of 15 independently validated small and alternative ORFs (Ncl1450, Ncr1470,
NbID, NirP1, NsiR7, Norf4, SliP4, as_rsbU, AcnSP, AtpΘ, IOF slr0978, IOF sll1954,
IOF slr1082, IOF slr0489 and IOF slr1079). IOF indicates ORFs located
"Internal_OutofFrame". (IOF). The worst averaged translational efficiency (aTE) for
these ORFs was detected for NsiR7 with 0.32. However, one of the annotated house-
keeping genes, *rpl36*, encoding ribosomal protein L36 had an even lower TE of 0.288
in one of the replicates. Therefore, we set the minimum threshold for aTE to ≥0.288.
Similarly, we set the DeepRibo prediction score at ≥-3.009 according to the score of -
3.0091 for gene *psbF* (*smr0006*) encoding the cytochrome b_{559} beta chain
(photosystem II protein VI). The other filter criteria were that the RPKM was ≥40 in
either of the 2 replicates, the number of codons ≥12 including the stop codon and that
none of these ORFs was annotated at the time of analysis.

(2) TIS-Ribo-seq were analyzed using OrfBounder[108]. According to the results of our
metagene analysis, we required the three different distances from the mapped 3' end
of the peak to be located at 19, 4 and 2 nt from the respective start codon.

The other filter criteria were that the RPKM was ≥40 in either of the 2 replicates for the
TIS as well as the associated Ribo-seq read mapping, a TIS/Ribo-seq $\log_2$FC of ≥1 in
either of the 2 replicates, a number of codons ≥12, an aTE of ≥0.282 and that none of
these ORFs was annotated.

(3) The TTS-Ribo-seq filter criteria were an RPKM value ≥10 in either of the 2
replicates for the TTS as well as the associated Ribo-seq read mapping, a TTS/Ribo-
seq $\log_2$FC of ≥1 in either of the 2 replicates, a codon count ≥12, an aTE of ≥0.15 and
that none of these ORFs was annotated. These criteria were applied separately to the
"TTS-next" and the "TTS-furthest" dataset.

Conservation and protein domain analysis

The identification of novel small protein homologues was performed using Blastp and
TBlastn searches in bacteria at the National Center for Biotechnology Information
website (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The protein sequences for novel
protein candidates identified in this study were used as query sequences. For TBlastn,

the following parameters were used: the filter for low complexity regions off, a word

size of 6, 60% coverage of the query sequence with at least 40% identity, an E value

≤100 to capture all potential orthologs, and an E value <0.1 for high-confidence hits[34].

Moreover, novel small protein candidates were further analyzed for signal peptides

and transmembrane domains using deepTMHMM[60].

Integrated proteogenomics search databases

Integrated proteogenomics databases (iPtgxDBs) were generated based on the

Synechocystis 6803 genome sequence (RefSeq ID: GCF_000009725.1). Annotations

from reference genome annotation centers (NCBI RefSeq (2022), GenBank (2016;

from here GBK) and the KZS annotation, two *ab initio* gene predictions (Prodigal

2.6.3[52] and ChemGenome 2.1[53]), and an *in-silico* ORF prediction considering

alternative start codons were hierarchically integrated into a standard iPtgxDB (**Table**

S5A) as previously described[48]. A minimum length cutoff of 10 aa was chosen for the

in-silico ORFs (closely matching the minimal length of Ribo-seq identified novel CDS

candidates of 11 aa) and the iPtgxDBs were generated using the strict trypsin

cleavage pattern (cleavage also before P). Since ChemGenome can only process

fasta files containing a single contig, ChemGenome and Prodigal predictions were

generated separately for the chromosome and each plasmid and then combined. To

improve search statistics, we also created a much smaller custom iPtgxDB (**Table**

S5B) as described[34]. For this, we included the 2,708 top candidates implicated by TIS

and TTS Ribo-seq data instead of the 5,071 ChemGenome entries and 107,626 *in-*

silico ORF predictions included in the much larger standard iPtgxDB (**Tables S5A and**

B).

Analysis of available mass spectrometry data and post-processing

Public proteomics data for *Synechocystis* 6803[47] were downloaded from 2 large

datasets deposited in PRIDE (PXD033243, 628 files; PXD033290, 367 files) using

pridepy 0.0.2[110] and converted to mzML format with pyrawr 0.1.0. The files contained

six different data types (unlabeled, dimethyl labeled with and without

phosphoproteomics, TMT-6 labeled, and TMT-10 labeled with quantification at the

MS2 or MS3 level). For each data type, three searches were performed (**Fig. S4**): A

regular search against an NCBI RefSeq database (November 2022) was carried out

to establish the coverage of the annotated proteome. Searches against the larger

standard iPtgxDB and a smaller custom iPtgxDB enabled the discovery of proteomic evidence for the expression of several novel SEPs. FragPipe 21.1 was used to search the data with MSFragger 4.0[111] and validate peptides with PeptideProphet (Philosopher 5.1.0[112]). Apart from specifying the labels for the TMT and dimethyl searches, standard parameters were used for the corresponding workflows. The search results in PEPxml format were processed using custom Java tools as described previously[48]: The results of all 6 searches were combined for each of the three databases and filtered based on the PeptideProphet probability using a false discovery rate (FDR) of 0.0065% at the peptide-spectrum match (PSM) level, leading to protein level FDRs of 0.83% for the searches against the standard iPtgxDB and 0.98% for the searches against the custom iPtgxDB. To obtain a list of novel proteins identified with higher confidence, the identified proteins were further filtered to have a minimum number of PSMs depending on the origin of the annotation (≥2 PSMs if annotation came from RefSeq, GBK, KZS, or Ribo-Seq; ≥3 PSMs if annotation was based on ab initio prediction by Prodigal or ChemGenome; ≥4 PSMs if based on a prediction of in-silico ORFs). Based on the iPtgxDB identifier, proteins were then categorized into annotated proteins, novel proteins, longer and shorter proteoforms of annotated proteins, and expressed pseudogenes. Importantly, all peptides were also classified with the PeptideClassifier tool[113], adapted for prokaryotic proteogenomics[48]. This allowed to filter the most information-rich peptides that either unambiguously identify one unique novel proteoform (class 1a), a subset or all proteoforms encoded by a gene model (class 2a, class 2b) and de-select those peptides shared by proteoforms encoded by different gene models (class 3b). For more information, see (https://iptgxdb.expasy.org/creating_iptgxdbs/).

Consolidation of genome annotations

A consolidated version of major reference annotation sources included in the standard iPtgxDB (see above) was generated by post-processing the GFF file produced when creating the iPtgxDB using a custom Python script. A comparison of RefSeq, GBK, KZS and Prodigal illustrates that substantial differences exist (**Fig. S5**). Notably, the Hess laboratory has been using their manually curated and extended KZS genome annotation as basis, while we used the RefSeq2022 annotation as main reference for the proteogenomics approach. This can nevertheless be resolved; it in fact allowed us to identify evidence for annotations from Genbank 2016 that were removed in RefSeq,

but still represent valid protein-coding genes as judged by the protein expression
evidence we detected. We in fact assume that many research communities could
benefit from using such an integrative approach. The consolidated GFF file was
constructed such that for each annotation cluster (annotations from different sources
that share the same stop codon position), the longest annotation was used to
represent the cluster. A respective cluster was then colored according to the
conservation among annotations: clusters represented in all reference annotation
sources were colored grey if all annotations were identical and yellow if any of the
annotations differed in length, while clusters only appearing in a subset of annotations
were colored red. Coloring entries in JBrowse2 by the color attribute in the GFF file
was achieved by writing a simple no-build plugin.

Genetic Engineering and protein verification

Genes of interest were cloned into the self-replicating expression plasmid pVZ322
flanked by the copper-inducible P_{petE} promoter at the 5' end and a coding segment for
either a 3xFLAG tag (DYKDHDGDYKDHDIDYKDDDDK) or a SPA tag[114]. This was
started by subcloning the coding sequences into plasmid pUC19. Therefore, a part of
pUC19 (primers P11-45 + P11-46; **Table S4**) was amplified to generate the backbone.
For asPsbC inserts, the P_{petE} promoter and the *asPsbC* coding sequence were
amplified from *Synechocystis* 6803 genomic DNA using primers G3-64 + VR278 and
VR279 + VR280. The SPA tag, STOP codon, *ncl1450* 3´UTR and the *oop* terminator
sequences were PCR-amplified from plasmid pVZ322::P_{petE}_*ncr1470*_SPA using
primers VR281 + G3-65. For asCsx18 inserts, the P_{petE} promoter and *asCsx18* coding
sequence were amplified using primers G3-64 + VR285 and VR286 + VR287, while
the SPA tag, STOP codon, *ncl1450* 3´UTR and *oop* terminator sequences were
amplified from plasmid pVZ322_P_{petE}::*ncr1470*-SPA using primers VR288 + G3-65.
The PCR-amplified pUC19 backbone and the respective insert fragments each were
assembled using AQUA cloning[115] and subcloned into *E. coli* Top10F' (Thermo Fisher
Scientific). The isolated plasmids were used as a PCR template with primers VR277
+ VR282 to amplify the fragment P_{petE}_*asPsbC* or *asCsx18*_SPA_STOP_*ncl1450*
3´UTR_*oop* with overhangs to the self-replicating expression plasmid pVZ322. The
vector was therefore cut with *Sal*I and *Pst*I (NEB) and treated with FAST AP (Thermo
Fisher). The linearized plasmid and the insert fragments were assembled using AQUA
cloning[115] and subcloned into *E. coli* Top10F' cells. All other genes of interest were

cloned following the same strategy but using the respective primers listed in **Table S4.**

All PCR reactions were performed with PCRBIO HiFi Polymerase (PCR Biosystems).

All PCRs using a plasmid as template were treated with FastDigest *Dpn*I (Thermo

Fisher Scientific). All PCR products were gel-purified (NucleoSpin Gel and PCR Clean-

up, Macherey-Nagel). Sanger Sequencing Economy Run (Microsynth) was used to

verify the correct sequence of all plasmids. The constructs were then conjugated into

the *Synechocystis* 6803 wild type[67]. Conjugated clones were selected with 50 µg/mL

kanamycin and 5 µg/mL gentamicin in BG11 medium[90], grown to exponential phase

(OD_{750} ~ 0.6-0.8) and protein expression was stimulated by the addition of 1.25 or 2

µM $CuSO_4$. Bortezomib (20 µM) was used to inhibit the Clp protease system during

the expression phase for candidates that lacked Western blot signals. After 2 h of

incubation, induced and control wild type cells were harvested by centrifugation (4000

× g, 4°C, 10 min) and lysed in phosphate buffered saline (137 mM sodium chloride,

2.7 mM potassium chloride, 10 mM disodium phosphate, 1.8 mM potassium

dihydrogen phosphate, pH 7.4) supplemented with cOmplete protease inhibitor

cocktail (Roche). Cell disruption was performed mechanically in a prechilled Precellys

24 (Bertin Instruments) with three rounds of shaking at 6,000 rpm (10 s) followed by a

pause (5 s) per cycle, for a total of 5 cycles and using glass beads with a 0.1–0.25 mm

diameter. Glass beads were removed by centrifugation (1,000 x g, 1 min, 4°C).

Obtained cell lysates were mixed with denaturing and reducing loading dye (5×

concentrated: 250 mM Tris-HCl pH 6.8, 25% glycerol, 10% SDS, 500 mM DTT, and

0.05% bromophenol blue G-250). The samples were subsequently separated by SDS-

Tricine-PAGE with Precision Plus Protein Dual Xtra molecular weight marker (Bio-

Rad) as standard followed by western blotting[13].

Several vectors were constructed for the expression of the different versions of

Slr0489 and Slr1079, including the presence or absence of a sequence encoding a C-

terminal 3xFLAG tag. The genes *slr0489* and *slr1079* were amplified from

Synechocystis 6803 genomic DNA using primer pairs RB1 + RB2 and RB3 + RB4,

respectively. For the nested genes, *slr0489S* and *slr1079S*, we also amplified the 5´

UTR with primer pairs RB5 + RB6 and RB7 +RB8, respectively. The amplicons were

cloned into the reverse-amplified pUC19P_{rha}YFP3×FLAG vector with or without the

FLAG tag[116]. For the wild-type version, pUC19P_{rha}YFP3×FLAG was reverse amplified

with RB9 + RB10 for the tagged version, or RB10 + RB11 for the untagged version.

Similarly, *slr0489S* and *slr1079S* were amplified with RB12 and RB9 or RB11 for the

untagged version. The inserts of interest were amplified and cloned into *Xmn*I-digested

pVZ322s as described previously[117]. For the co-immunoprecipitation assay, the

3xFLAG constructs were transformed into wild-type *Synechocystis* 6803 by

electroporation as described[46].

To assess possible physiological effects of expressing either the short or the long

Slr0489 version on *Synechocystis*, first the native version, encoding both the large and

the small protein, was amplified with primer pairs RB13 + RB14 from a previous

construct. The fragment was cloned into a shorter version of a modified pACYC-Duet-

1 vector (with Km^R instead of Cm^R), after amplification with primers RB15 + RB16, via

AQUA cloning[118]. From this new construct (pACYC-*slr0489*WT-6×His), a single point

mutation was inserted into *slr0489*WT to change the start codon of the small gene into

a threonine codon, creating a construct expressing only the long protein (Slr0489L).

The construct carrying the native *slr0489* gene was reverse amplified with primers

RB17 + RB18 and RB18, carrying 5´overhangs for AQUA cloning. For expression in

Synechocystis, *slr0489*L was amplified with the primer pair RB1 + RB19 from pACYC-

slr0489L to generate overhangs to the linearized pUC19-Rha-YFP-3×FLAG vector

and later cloned into pVZ322s[116] without any tag sequence. The final pVZ322s-P*rha*-

slr0489L was then conjugated into *Synechocystis* Δ*slr0489* via tri-parental mating. The

constructs pVZ322s-P*rha-slr0489* and pVZ322s-P*rha-slr0489S* were also conjugated

into Δ*slr0489*. The strains were then cultivated in BG11 under standard conditions.

The expression was induced with 0.6 mg/mL L-rhamnose, while uninduced cultures

served as control. The growth was monitored by measuring the OD_{750} of 150 μL

samples in a 96-well plate for 5 d. Each strain was cultivated in biological duplicates

for each condition.

The Δ*slr0489* deletion mutant was generated by replacing the *slr0489* gene by

homologous recombination with a streptomycin resistance cassette (*aadA*) amplified

from plasmid pRL692, and using pUC19 as a vector for subcloning. For this, flanking

regions present up- and downstream of *slr0489* were amplified from *Synechocystis*

6803 genomic DNA using primer pairs 117/118 and 123/124. Transcription of *slr0489*

occurs from a transcriptional start site at position 2,622,526 on the fw strand matching

the first nt of the start codon (leaderless expression), but leading to a joint

transcriptional unit TU2770 with the downstream genes *slr0491–slr0492*[51]. To allow
expression of the downstream-located genes, the promoter region was amplified
separately using primers121/122 and inserted at the end of the *aadA* resistance
cassette. Note, that the *slr0489* 3′ end and the *slr0491* 5′ end overlap by 15 nt, which
were kept intact behind the inserted *aadA*-promoter construct. The construct for gene
replacement was generated by AQUA cloning[118]. The corresponding plasmid was
used to transform *Synechocystis* 6803 cells by electroporation according to Kraus et
al.[46]. Transformants were selected on 20 µg/mL streptomycin. The segregation of
Δ*slr0489*/*aadA* from wild type alleles was followed by colony PCR using the primers
133/134.

1143

Separation of soluble and insoluble fractions

For localization of the Ncl1450 protein, wild type *Synechocystis* 6803, P$_{petE}$::*ncl1450*-
3×FLAG and P$_{ncl1450}$::*ncl1450*-3×FLAG ectopic overexpression mutants were grown
until exponential phase, then the first sample was taken. The P$_{petE}$ promoter was
induced by 2 µM CuSO$_4$. Carbon-depletion for the native promoter was induced by
washing the pellet three times in carbon-depleted BG11 medium. 24 hours after
induction, samples were taken and centrifuged (4,000 × g, 10 min, 4°C) to collect cells.
The pellet was resuspended in membrane buffer (25 mM HEPES-NaOH, pH 7.0, 15
mM CaCl$_2$, 5 mM MgCl$_2$, 15% v/v glycerol) with protease inhibitor and then frozen in
liquid nitrogen and stored at -80°C. Cells were disrupted in a Precellys homogenizer
as described above. Whole cell lysate was centrifuged (21,000 × g, 20 min, 4°C) for
separation of soluble and insoluble fractions, then supernatant (soluble fraction) and
pellet were separated. The pellet was resuspended in membrane buffer. SDS-PAGE
and anti-FLAG western blot were performed as described[13].

Co-IP, LC-MS/MS analyses and data processing of nested gene products

Tagged   proteins   (Slr0489L-3xFLAG;   Slr0489S-3xFLAG;   Slr1079L-3xFLAG;
Slr1079S-3xFLAG; sfGFP-3xFLAG) were expressed in *Synechocystis* 6803 from
plasmid pVZ322s[119] under control of the P$_{rha}$ promoter[120] after induction by adding 0.6
mg/mL L-rhamnose at an OD$_{750}$ of 0.5. After 24 h, cells from 50 mL batch cultures
were harvested by centrifugation (3,750 × g, 4°C, 20 min), resuspended in 700 µL
FLAG buffer (50 mM HEPES-NaOH pH 7; 5 mM MgCl$_2$; 25 mM CaCl$_2$, 150 mM NaCl;

10% glycerol; 0.1% Tween-20) supplemented with Protease Inhibitor (cOmplete,

Roche) and mixed with 100 µL of glass beads (Ø 0.1–0.25 mm, Retsch). Cell

disruption was performed in a prechilled Precellys 24. Crude cell debris and glass

beads were removed by centrifugation (1,000 × g, 1 min, 4°C) and supernatants were

subsequently incubated at gentle shaking in presence of 2% w/v n-dodecyl- ß -D-

maltoside for 60 min at 4°C in the dark. Remaining cell debris was removed by

centrifugation (21,300 × g, 4°C, 30 min). The cleared total cell lysate was subjected to

co-IP using anti-FLAG M2 affinity magnetic beads (Sigma–Aldrich) rotating for 2 h at

4°C. The beads were washed 5 times with 600 µL FLAG buffer supplemented with

protease inhibitors for 2 min, 4°C, rotating. Bound proteins were eluted by incubation

with 150 µL FLAG peptide (Sigma–Aldrich, 100 ng/µL) in TBS buffer (150 mM NaCl in

20 mM Tris/HCl, pH 7.6) for 30 min at 4°C in two subsequent steps and subsequently

combined. Three independent replicates were prepared for each Co-IP. Fractions (20

µL) of each eluate protein were analyzed by SDS-PAGE and anti-FLAG

immunoblotting for quality control. Remaining eluates were lyophilized and

resolubilized in denaturation buffer (6 M urea, 2 M thiourea in 100 mM Tris/HCl; pH

8.0) for LC-MS/MS analysis. Protein disulfide bonds were reduced with dithiothreitol

at a final concentration of 1 mM for 45 min and the resulting thiol groups were alkylated

with iodoacetamide at a final concentration of 5.5 mM for 45 min in the dark. Proteins

were pre-digested with Lys-C (Santa Cruz Biotechnology) for 3 h, diluted with 4

volumes of 20 mM ammonium bicarbonate buffer, pH 8, and overnight digested with

trypsin (sequencing grade, Promega) at room temperature with a protease/protein

ratio of 1/100 for both proteases. Resulting peptide solutions were acidified with

trifluoroacetic acid to pH 2.5, and purified by Stage Tips[121]. For LC-MS/MS-based

protein analysis, 1/20 of each sample was loaded onto an in-house-packed reverse-

phase (RP) column (20 cm; 75 µm ID, packed with ReproSil-Pur 1.9 µm C18 from Dr.

Maisch, Germany) and separated by RP chromatography on an EASY-nLC 1200

(Thermo Fisher Scientific, USA) using linear 36 min gradients. Eluting peptides were

directly ionized by electro spray ionization and analyzed on a Q-Exactive HF mass

spectrometer (Thermo Fisher Scientific, USA) in the data dependent acquisition mode.

MS1 resolution was set to 60,000 at a scan range of 300 – 1650 m/z. The 7 most

abundant ions with charge states 2-5 were subjected to fragmentation by higher-

energy collision induced fragmentation. Fragment ions were analyzed at an MS2

resolution of 60,000 and a scan range of 200-2000 m/z. Dynamic exclusion of
previously sequenced ions was set to 30 s.

Raw spectra of each Co-IP experiment (Slr0489L; Slr0489S; Slr1079L and Slr1079S)
were analyzed separately including all replicates and processed with the common
control (sfGFP-FLAG) with the MaxQuant software (version 1.6.8.0)[122] at default
settings. Label-free quantification (LFQ), matching between runs and iBAQ options
were enabled. Peak lists were searched against Co-IP specific modified target-decoy
database of *Synechocystis* 6803[47] with 3,959 protein entries, including the tagged bait
protein sequences truncated for the indifferentiable 3xFLAG sequence. False
discovery rates were limited to 1% at peptide and protein levels. Missing LFQ protein
intensities were imputed to increase number of quantifiable proteins. Therefore,
random values the low intensity range were given in case of missing values for each
replicate separately within a width of 10% of the standard deviation of the data and a
downshift corresponding to the 2-fold standard deviation using the Perseus software
suite (version 1.6.5.0)[123] (**Fig.s S16–S19**). Co-IP enriched proteins were analyzed by
Student's *t*-test relative to the control (sfGFP-FLAG IP) with a permutation-based FDR
of 0.05 (**Tables S6–S9**).

1215

DATA AVAILABILITY

Ribo-seq, TIS and TTS profiling datasets were deposited in the Gene Expression
Omnibus (GEO) Database, with accession number GSE246336 (direct access:
https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE246336). The mass
spectrometry proteomics data generated in this study have been deposited at the
ProteomeXchange Consortium via the PRIDE[124] partner repository with the dataset
identifier PXD060427. The datasets described in this study can be viewed with an
interactive online JBrowse 2 instance at https://www.bioinf.uni-freiburg.de/~ribobase/.

1224

ACKNOWLEDGEMENTS

This work was supported by the German Research Foundation (DFG) priority program
SPP2002 "Small Proteins in Prokaryotes, an Unexplored World" (grants HE2544/12-

1/2 to WRH, SH580/7-1/2 to CMS, BA 2168/21-1/2 to ROB), by grants HE 2544/22-1

to WRH and DFG_MA 49184-1 to BM and by Germany's DFG-funded Excellence

Strategy (CIBSS - EXC-2189 - Project ID 390939984) to ROB. CHA acknowledges

funding from the Swiss National Science Foundation for BH and GJ (grant 197391).

The computational resources were supported by the de.NBI Cloud within the German

Network   for   Bioinformatics   Infrastructure   (de.NBI)   and   ELIXIR-DE

(Forschungszentrum Jülich and W-de.NBI-001, W-de.NBI-004, W-de.NBI-008, W-

de.NBI-010, W-de.NBI-013, W-de.NBI-014, W-de.NBI-016, W-de.NBI-022). We thank

Ann-Janine Imsiecke, Stephanie Färber and Philipp Kible (Würzburg, Germany) for

the technical assistance in establishing the Ribo-seq workflow for *Synechocystis* 6803,

Paulo Oliveira (Porto, Portugal) for the *tolC* transporter mutants and Joëlle Sasse

Schläpfer (Zurich, Switzerland) for critical reading.

1240

1241 **CONFLICT OF INTEREST STATEMENT**

1242 The authors declare that they have no conflicts of interest.

1243

AUTHOR CONTRIBUTIONS

WRH, CMS and ROB designed and supervised the Ribo-seq part of the project and

secured funding. CHA supervised the proteogenomics part of the project. VK and PM

constructed and cultivated strains. LH and VK designed and established Ribo-seq and

TIS/TTS methods for *Synechocystis* 6803. RG performed bioinformatic processing of

the Ribo-seq, TIS and TTS data, developed custom analysis tools and scripts, and

created a JBrowse 2 instance for the project. VK, LH and WRH performed the analysis

of Ribo-seq, TIS and TTS data including the manual labeling of the implied novel ORFs

and sORFs and curation of gene annotations. PS and BM performed mass

spectrometry, BH, GJ and CHA performed the proteogenomic analysis to identify

known and novel ORFs/sORFs including iPtgxDB-based protein searches informed

by Ribo-seq and TIS/TTS data, and contributed the generic approach for genome

annotation consolidation. VK, VR and MG constructed strains and performed western

blot analyses. RAB performed cultivation experiments with *slr0489* and *slr1079*. VK,

LH, CHA and WRH wrote the manuscript with input from all authors. All authors read

and agreed on the manuscript.

1260

REFERENCES

1. Flombaum, P. *et al.* Present and future global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc Natl Acad Sci USA* **110**, 9824–9829 (2013).

2. Hagemann, M. & Hess, W. R. Systems and synthetic biology for the biotechnological application of cyanobacteria. *Curr Opin Biotechnol* **49**, 94–99 (2018).

3. Vijay, D., Akhtar, M. K. & Hess, W. R. Genetic and metabolic advances in the engineering of cyanobacteria. *Curr. Opin. Biotechnol.* **59**, 150–156 (2019).

4. Klähn, S., Opel, F. & Hess, W. R. Customized molecular tools to strengthen metabolic engineering of cyanobacteria. *Green Carbon* 2, 149-163 (2024).

5. Kaneko, T. *et al.* Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* **3**, 109–136 (1996).

6. Zhang, J. Protein-length distributions for the three domains of life. *Trends Genet. TIG* **16**, 107–109 (2000).

7. Ahrens, C. H., Wade, J. T., Champion, M. M. & Langer, J. D. A practical guide to small protein discovery and characterization using mass spectrometry. *J. Bacteriol.* **204**, e0035321 (2022).

8. Burton, A. T., Zeinert, R. & Storz, G. Large roles of small proteins. *Annu. Rev. Microbiol.* **78**, 1–22 (2024).

9. Alvarenga-Lucius, L. V. *et al.* The high-light-induced protein SliP4 binds to NDH1 and photosystems facilitating cyclic electron transport and state transition in *Synechocystis* sp. PCC 6803. *New Phytol.* **239**, 1083–1097 (2023).

10. Komenda, J. & Sobotka, R. Cyanobacterial high-light-inducible proteins--

Protectors of chlorophyll-protein synthesis and assembly. *Biochim. Biophys. Acta*

1857, 288–295 (2016).

11. Fromme, P., Melkozernov, A., Jordan, P. & Krauss, N. Structure and function of

photosystem I: interaction with its soluble electron carriers and external antenna

systems. *FEBS Lett.* **555**, 40–44 (2003).

12. Xu, W., Tang, H., Wang, Y. & Chitnis, P. R. Proteins of the cyanobacterial

photosystem I. *Biochim. Biophys. Acta BBA - Bioenerg.* **1507**, 32–40 (2001).

13. Baumgartner, D., Kopf, M., Klähn, S., Steglich, C. & Hess, W. R. Small proteins in

cyanobacteria provide a paradigm for the functional analysis of the bacterial micro-

proteome. *BMC Microbiol.* **16**, 285 (2016).

14. Brandenburg, F. & Klähn, S. Small but smart: On the diverse role of small proteins

in the regulation of cyanobacterial metabolism. *Life* **10**, 322 (2020).

15. Gray, T., Storz, G. & Papenfort, K. Small proteins; Big questions. *J. Bacteriol.* **204**,

e0034121 (2022).

16. Fesenko, I. *et al.* The hidden bacterial microproteome. *Mol. Cell* **85**, 1024–1041

(2025).

17. Song, K. *et al.* AtpΘ is an inhibitor of F0F1 ATP synthase to arrest ATP hydrolysis

during low-energy conditions in cyanobacteria. *Curr. Biol.* **32**, 136-148.e5 (2022).

18. Jurėnas, D., Fraikin, N., Goormaghtigh, F. & Van Melderen, L. Biology and

evolution of bacterial toxin–antitoxin systems. *Nat. Rev. Microbiol.* **20**, 335–350

(2022).

19. Santos-Júnior, C. D. *et al.* Discovery of antimicrobial peptides in the global

microbiome with machine learning. *Cell* **187**, 3761-3778.e16 (2024).

20. Ingolia, N. T., Ghaemmaghami, S., Newman, J. R. S. & Weissman, J. S. Genome-

wide analysis in vivo of translation with nucleotide resolution using ribosome

profiling. *Science* **324**, 218–223 (2009).

21. Ingolia, N. T., Hussmann, J. A. & Weissman, J. S. Ribosome profiling: global views

of translation. *Cold Spring Harb. Perspect. Biol.* **11**, a032698 (2019).

22. Mohammad, F., Green, R. & Buskirk, A. R. A systematically-revised ribosome

profiling method for bacteria reveals pauses at single-codon resolution. *eLife* **8**,

e42591 (2019).

23. Gerashchenko, M. V., Lobanov, A. V. & Gladyshev, V. N. Genome-wide ribosome

profiling reveals complex translational regulation in response to oxidative stress.

Proc. Natl. Acad. Sci. USA **109**, 17394–17399 (2012).

24. Zupanic, A. *et al.* Detecting translational regulation by change point analysis of

ribosome profiling data sets. *RNA* **20**, 1507–1518 (2014).

25. Chotewutmontri, P. & Barkan, A. Ribosome profiling elucidates differential gene

expression in bundle sheath and mesophyll cells in maize. *Plant Physiol.* **187**, 59–

72 (2021).

26. Xiao, Z. *et al.* De novo annotation and characterization of the translatome with

ribosome profiling data. *Nucleic Acids Res.* **46**, e61 (2018).

27. Ingolia, N. T., Lareau, L. F. & Weissman, J. S. Ribosome profiling of mouse

embryonic stem cells reveals the complexity and dynamics of mammalian

proteomes. *Cell* **147**, 789–802 (2011).

28. Bazzini, A. A. *et al.* Identification of small ORFs in vertebrates using ribosome

footprinting and evolutionary conservation. *EMBO J.* **33**, 981–993 (2014).

29. Hsu, P. Y. *et al.* Super-resolution ribosome profiling reveals unannotated

translation events in Arabidopsis. *Proc. Natl. Acad. Sci. USA* **113**, E7126–E7135

(2016).

30. Meydan, S. *et al.* Retapamulin-assisted ribosome profiling reveals the alternative

bacterial proteome. *Mol. Cell* **74**, 481-493.e6 (2019).

31. Orr, M. W., Mao, Y., Storz, G. & Qian, S.-B. Alternative ORFs and small ORFs:

shedding light on the dark proteome. *Nucleic Acids Res.* **48**, 1029–1042 (2020).

32. Froschauer, K. *et al.* Complementary Ribo-seq approaches map the translatome

and provide a small protein census in the foodborne pathogen *Campylobacter*

jejuni. Nat. Commun. **16**, 3078 (2025).

33. Weingarten-Gabbay, S. *et al.* Pan-viral ORFs discovery using massively parallel

ribosome profiling. *Science* **388**, 1218–1224 (2025).

34. Hadjeras, L. *et al.* Unraveling the small proteome of the plant symbiont

Sinorhizobium meliloti by ribosome profiling and proteogenomics. *microLife* **4**,

uqad012 (2023).

35. Hadjeras, L. *et al.* Revealing the small proteome of *Haloferax volcanii* by combining

ribosome profiling and small-protein optimized mass spectrometry. *microLife* **4**,

uqad001 (2023).

36. Tufail, M. A. *et al.* Uncovering the small proteome of *Methanosarcina mazei* using

Ribo-seq and peptidomics under different nitrogen conditions. *Nat. Commun.* **15**,

8659 (2024).

37. Weaver, J., Mohammad, F., Buskirk, A. R. & Storz, G. Identifying small proteins by

ribosome profiling with stalled initiation complexes. *mBio* **10**, (2019).

38. Florin, T. *et al.* An antimicrobial peptide that inhibits translation by trapping release

factors on the ribosome. *Nat. Struct. Mol. Biol.* **24**, 752–757 (2017).

39. Mangano, K. *et al.* Genome-wide effects of the antimicrobial peptide apidaecin on

translation termination in bacteria. *eLife* **9**, e62655 (2020).

40. Stringer, A., Smith, C., Mangano, K. & Wade, J. T. Identification of novel translated

small open reading frames in *Escherichia coli* using complementary ribosome

profiling approaches. *J. Bacteriol.* **204**, e00352-21 (2022).

41. Vazquez-Laslop, N., Sharma, C. M., Mankin, A. & Buskirk, A. R. Identifying small

open reading frames in prokaryotes with ribosome profiling. *J. Bacteriol.* **204**,

e00294-21 (2020).

42. Karlsen, J., Asplund-Samuelsson, J., Thomas, Q., Jahn, M. & Hudson, E. P.

Ribosome profiling of *Synechocystis* reveals altered ribosome allocation at carbon

starvation. *Msystems* **3**, e00126-18 (2018).

43. Karlsen, J., Asplund-Samuelsson, J., Jahn, M., Vitay, D. & Hudson, E. P. Slow

protein turnover explains limited protein-level response to diurnal transcriptional

oscillations in cyanobacteria. *Front. Microbiol.* **12**, 657379 (2021).

44. de Alvarenga, L. V., Hess, W. R., Hagemann, M. & Hagemann, M. AcnSP - a novel

small protein regulator of aconitase activity in the cyanobacterium *Synechocystis*

sp. PCC 6803. *Front. Microbiol.* **11**, 1445 (2020).

45. Krauspe, V. *et al.* Discovery of a novel small protein factor involved in the

coordinated degradation of phycobilisomes in cyanobacteria. *Proc. Natl. Acad. Sci.*

USA **118**, e2012277118 (2021).

46. Kraus, A. *et al.* Protein NirP1 regulates nitrite reductase and nitrite excretion in

cyanobacteria. *Nat. Commun.* **15**, 1911 (2024).

47. Spät, P., Krauspe, V., Hess, W. R., Maček, B. & Nalpas, N. Deep proteogenomics

of a photosynthetic cyanobacterium. *J. Proteome Res.* **22**, 1969–19 (2023).

48. Omasits, U. *et al.* An integrative strategy to identify the entire protein coding

potential of prokaryotic genomes by proteogenomics. *Genome Res.* **27**, 2083–

2095 (2017).

49. Chen, Q. *et al.* Deletion of *sll1541* in *Synechocystis* sp. strain PCC 6803 allows

formation of a far-red-shifted holo-proteorhodopsin in vivo. *Appl. Environ.*

Microbiol. **84**, e02435-17 (2018).

50. Kopfmann, S., Roesch, S. K. & Hess, W. R. Type II toxin-antitoxin systems in the

unicellular cyanobacterium *Synechocystis* sp. PCC 6803. *Toxins* **8**, 228.1-228.23

(2016).

51. Kopf, M. *et al.* Comparative analysis of the primary transcriptome of *Synechocystis*

sp. PCC 6803. *DNA Res.* **21**, 527–539 (2014).

52. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

53. Mishra, A., Siwach, P., Singhal, P. & Jayaram, B. ChemGenome2.1: An ab initio

gene prediction software. in *Gene Prediction: Methods and Protocols* (ed. Kollmar,

M.) 121–138 (Springer, New York, NY, 2019). doi:10.1007/978-1-4939-9173-0_7.

54. Baers, L. L. *et al.* Proteome mapping of a cyanobacterium reveals distinct

compartment organization and cell-dispersed metabolism. *Plant Physiol.* **181**,

1721–1738 (2019).

55. Craigen, W. J., Cook, R. G., Tate, W. P. & Caskey, C. T. Bacterial peptide chain

release factors: conserved primary structure and possible frameshift regulation of

release factor 2. *Proc. Natl. Acad. Sci. USA* **82**, 3616–3620 (1985).

56. Craigen, W. J. & Caskey, C. T. The function, structure and regulation of E. coli

peptide chain release factors. *Biochimie* **69**, 1031–1041 (1987).

57. Baranov, P. V., Gesteland, R. F. & Atkins, J. F. Release factor 2 frameshifting sites

in different bacteria. *EMBO Rep.* **3**, 373–377 (2002).

58. Tate, W. P. *et al.* Translational termination efficiency in both bacteria and mammals

is regulated by the base following the stop codon. *Biochem. Cell Biol.* **73**, 1095–

1103 (1995).

59. Sesto, N., Wurtzel, O., Archambaud, C., Sorek, R. & Cossart, P. The excludon: a

new concept in bacterial antisense RNA-mediated gene regulation. *Nat. Rev.*

Microbiol. **11**, 75–82 (2013).

60. Hallgren, J. *et al.* DeepTMHMM predicts alpha and beta transmembrane proteins

using   deep   neural   networks.   2022.04.08.487609   Preprint   at

https://doi.org/10.1101/2022.04.08.487609 (2022).

61. Shore, S. F. H., Leinberger, F. H., Fozo, E. M. & Berghoff, B. A. Type I toxin-

antitoxin systems in bacteria: from regulation to biological functions. *EcoSal Plus*

12, eesp-0025-2022 (2024).

62. van Kempen, M. *et al.* Fast and accurate protein structure search with Foldseek.

Nat. Biotechnol. **42**, 243–246 (2024).

63. Yu, V., Ronzone, E., Lord, D., Peti, W. & Page, R. MqsR is a noncanonical

microbial RNase toxin that is inhibited by antitoxin MqsA via steric blockage of

substrate binding. *J. Biol. Chem.* **298**, 102535 (2022).

64. Gelhausen, R. *et al.* HRIBO: high-throughput analysis of bacterial ribosome

profiling data. *Bioinformatics* **37**, 2061–2063 (2021).

65. Mitschke, J. *et al.* An experimentally anchored map of transcriptional start sites in

the model cyanobacterium *Synechocystis* sp. PCC6803. *Proc. Natl. Acad. Sci.*

USA **108**, 2124–2129 (2011).

66. Kopf, M. *et al.* Comparative genome analysis of the closely related *Synechocystis* strains PCC 6714 and PCC 6803. *DNA Res.* **21**, 255–266 (2014).

67. Scholz, I., Lange, S. J., Hein, S., Hess, W. R. & Backofen, R. CRISPR-Cas systems in the cyanobacterium *Synechocystis* sp. PCC6803 exhibit distinct processing pathways involving at least two Cas6 and a Cmr2 protein. *PLoS ONE* **8**, e56470 (2013).

68. Shah, S. A. *et al.* Comprehensive search for accessory proteins encoded with archaeal and bacterial type III CRISPR-cas gene cassettes reveals 39 new cas gene families. *RNA Biol.* **16**, 530–542 (2019).

69. Susanto, F. A., Jones, A. D., Donnelly, S. L. & Lundquist, P. K. Cyanoglobule lipid droplets are a stress-responsive metabolic compartment of cyanobacteria and the progenitor of plant plastoglobules. *Plant Cell* **37**, koaf127 (2025).

70. Kopf, M. & Hess, W. R. Regulatory RNAs in photosynthetic cyanobacteria. *FEMS Microbiol. Rev.* **39**, 301–315 (2015).

71. Huckauf, J., Nomura, C., Forchhammer, K. & Hagemann, M. Stress responses of *Synechocystis* sp. strain PCC 6803 mutants impaired in genes encoding putative alternative sigma factors. *Microbiology* **146**, 2877–2889 (2000).

72. Trautmann, D., Voß, B., Wilde, A., Al-Babili, S. & Hess, W. R. Microevolution in cyanobacteria: re-sequencing a motile substrain of *Synechocystis* sp. PCC 6803. *DNA Res.* **19**, 435–448 (2012).

73. Katoh, A., Sonoda, M. & Ogawa, T. A possible role of 154-base pair nucleotides located upstream of ORF440 on CO2 transport of *Synechocystis* sp. PCC 6803. in *Photosynthesis: From Light to Biosphere* vol. 3 481–484 (Kluwer Academic Publishers, Dordrecht, The Netherlands, 1995).

74. Kamei, A., Ogawa, T. & Ikeuchi, M. Identification of a novel gene (*slr2031*) involved

in high-light resistance in the cyanobacterium *Synechocystis* sp. PCC 6803. in

Photosynthesis: Mechanisms and Effects: Volume I–V: Proceedings of the XIth

International Congress on Photosynthesis, Budapest, Hungary, August 17–22,

1998 (ed. Garab, G.) 2901–2904 (Springer Netherlands, Dordrecht, 1998).

doi:10.1007/978-94-011-3953-3_680.

75. Bouillet, S., Arabet, D., Jourlin-Castelli, C., Méjean, V. & Iobbi-Nivol, C. Regulation

of σ factors by conserved partner switches controlled by divergent signalling

systems. *Environ. Microbiol. Rep.* **10**, 127–139 (2018).

76. Behler, J. *et al.* The host-encoded RNase E endonuclease as the crRNA

maturation enzyme in a CRISPR–Cas subtype III-Bv system. *Nat. Microbiol.* **3**,

367–377 (2018).

77. Oeser, S. *et al.* Minor pilins are involved in motility and natural competence in the

cyanobacterium *Synechocystis* sp. PCC 6803. *Mol. Microbiol.* **116**, 743–765

(2021).

78. Kaushal, P. & Lee, C. N-terminomics - its past and recent advancements. *J.*

Proteomics **233**, 104089 (2021).

79. Diesh, C. *et al.* JBrowse 2: a modular genome browser with views of synteny and

structural variation. *Genome Biol.* **24**, 74 (2023).

80. Hemm, M. R., Weaver, J. & Storz, G. *Escherichia coli* small proteome. *EcoSal Plus*

9, (2020).

81. Peng, Z., Xiao, Q. & Wan, C. Identification and validation of SmORF-encoded

peptides by genomics and proteomics in five cyanobacteria. *J. Proteome Res.* **24**,

5818−5829 (2025).

82. Jurėnas, D., Fraikin, N., Goormaghtigh, F. & Van Melderen, L. Biology and

evolution of bacterial toxin–antitoxin systems. *Nat. Rev. Microbiol.* **20**, 335–350

(2022).

83. Sailer, A.-L. *et al.* Internal in-frame translation generates Cas11b, which is

important for effective interference in an archaeal CRISPR-Cas system. *Front.*

Microbiol. **16**, 1543464 (2025).

84. McBride, T. M. *et al.* Diverse CRISPR-Cas complexes require independent

translation of small and large subunits from a single gene. *Mol. Cell* **80**, 971–979

(2020).

85. Kongari, R. *et al.* Phage spanins: diversity, topological dynamics and gene

convergence. *BMC Bioinformatics* **19**, 326 (2018).

86. Cahill, J. *et al.* Genetic analysis of the lambda spanins Rz and Rz1: identification

of functional domains. *G3 GenesGenomesGenetics* **7**, 741–753 (2017).

87. Doherty, E. E. *et al.* A miniature CRISPR–Cas10 enzyme confers immunity by

inhibitory signalling. *Nature* **647**, 997–1004 (2025).

88. Sullivan, A. E. *et al.* The Panoptes system uses decoy cyclic nucleotides to defend

against phage. *Nature* **647**, 988-996 (2025).

89. Thomas, J.-C., Ughy, B., Lagoutte, B. & Ajlani, G. A second isoform of the

ferredoxin:NADP oxidoreductase generated by an in-frame initiation of translation.

Proc. Natl. Acad. Sci. USA **103**, 18368–18373 (2006).

90. Rippka, R., Deruelles, J., Waterbury, J. B., Herdman, M. & Stanier, R. Y. Generic

assignments, strain histories and properties of pure cultures of cyanobacteria.

Microbiology **111**, 1–61 (1979).

91. Oliveira, P. *et al.* The versatile TolC-like Slr1270 in the cyanobacterium

Synechocystis sp. PCC 6803. *Environ. Microbiol.* **18**, 486–502 (2016).

92. Gonçalves, C. F., Pacheco, C. C., Tamagnini, P. & Oliveira, P. Identification of

inner membrane translocase components of TolC-mediated secretion in the

cyanobacterium *Synechocystis* sp. PCC 6803. *Environ. Microbiol.* **20**, 2354–2369

(2018).

93. Mattiuzzo, M. *et al.* Role of the Escherichia coli SbmA in the antimicrobial activity

of proline-rich peptides. *Mol. Microbiol.* **66**, 151–163 (2007).

94. Sharma, C. M., Darfeuille, F., Plantinga, T. H. & Vogel, J. A small RNA regulates

multiple ABC transporter mRNAs by targeting C/A-rich elements inside and

upstream of ribosome-binding sites. *Genes Dev.* **21**, 2804–2817 (2007).

95. Ingolia, N. T., Brar, G. A., Rouskin, S., McGeachy, A. M. & Weissman, J. S. The

ribosome profiling strategy for monitoring translation in vivo by deep sequencing

of ribosome-protected mRNA fragments. *Nat. Protoc.* **7**, 1534–1550 (2012).

96. Venturini, E. *et al.* A global data-driven census of *Salmonella* small proteins and

their potential functions in bacterial virulence. *microLife* **1**, uqaa002 (2020).

97. Köster, J. & Rahmann, S. Snakemake-a scalable bioinformatics workflow engine.

Bioinforma. Oxf. Engl. **34**, 3600 (2018).

98. Grüning, B. *et al.* Bioconda: sustainable and comprehensive software distribution

for the life sciences. *Nat. Methods* **15**, 475–476 (2018).

99. Martin, M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* **17**, 10 (2011).

100. Otto, C., Stadler, P. F. & Hoffmann, S. Lacking alignments? The next-

generation sequencing mapper segemehl revisited. *Bioinforma. Oxf. Engl.* **30**,

1837–1843 (2014).

101. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**,

giab008 (2021).

102. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics* **30**,

923–930 (2014).

103. Wingett, S. W. & Andrews, S. FastQ Screen: A tool for multi-genome mapping

and quality control. *F1000Research* **7**, 1338 (2018).

104. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis

results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–

3048 (2016).

105. Crooks, G. E., Hon, G., Chandonia, J.-M. & Brenner, S. E. WebLogo: A

sequence logo generator. *Genome Res.* **14**, 1188–1190 (2004).

106. Ndah, E. *et al.* REPARATION: ribosome profiling assisted (re-) annotation of

bacterial genomes. *Nucleic Acids Res.* **45**, e168–e168 (2017).

107. Clauwaert, J., Menschaert, G. & Waegeman, W. DeepRibo: a neural network

for precise gene annotation of prokaryotes by combining ribosome profiling signal

and binding site patterns. *Nucleic Acids Res.* **47**, e36–e36 (2019).

108. Gelhausen, R. *et al.* RiboReport-benchmarking tools for ribosome profiling-

based identification of open reading frames in bacteria. *Brief. Bioinform.* **23**,

bbab549 (2022).

109. Buels, R. *et al.* JBrowse: a dynamic web platform for genome visualization and

analysis. *Genome Biol.* **17**, 66 (2016).

110. Kamatchinathan, S. *et al.* pridepy: A Python package to download and search

data from PRIDE database. *J. Open Source Softw.* **10**, 7563 (2025).

111. Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. &

Nesvizhskii, A. I. MSFragger: ultrafast and comprehensive peptide identification in

mass spectrometry–based proteomics. *Nat. Methods* **14**, 513–520 (2017).

112.   da Veiga Leprevost, F. *et al.* Philosopher: a versatile toolkit for shotgun

proteomics data analysis. *Nat. Methods* **17**, 869–870 (2020).

113.   Qeli, E. & Ahrens, C. H. PeptideClassifier for protein inference and targeted

quantitative proteomics. *Nat. Biotechnol.* **28**, 647–650 (2010).

114.   Zeghouf, M. *et al.* Sequential Peptide Affinity (SPA) system for the identification

of mammalian and bacterial protein complexes. *J. Proteome Res.* **3**, 463–468

(2004).

115.   Beyer, H. M. *et al.* AQUA cloning: a versatile and simple enzyme-free cloning

approach. *PLOS ONE* **10**, e0137652 (2015).

116.   Hemm, L. *et al.* Interactors and effects of overexpressing YlxR/RnpM, a

conserved RNA binding protein in cyanobacteria. *RNA Biol.* **21**, 1–19 (2024).

117.   Kaltenbrunner, A. *et al.* Regulation of pSYSA defense plasmid copy number in

Synechocystis through RNase E and a highly transcribed asRNA. *Front. Microbiol.*

14, 1112307 (2023).

118.   Beyer, H. M. *et al.* AQUA cloning: a versatile and simple enzyme-free cloning

approach. *PLOS ONE* **10**, e0137652 (2015).

119.   Kaltenbrunner, A. *et al.* Regulation of pSYSA defense plasmid copy number in

Synechocystis through RNase E and a highly transcribed asRNA. *Front. Microbiol.*

14, 1112307 (2023).

120.   Kelly, C. L., Taylor, G. M., Hitchcock, A., Torres-Méndez, A. & Heap, J. T. A

rhamnose-inducible system for precise and temporal control of gene expression in

cyanobacteria. *ACS Synth. Biol.* **7**, 1056–1066 (2018).

121.   Ishihama, Y., Rappsilber, J. & Mann, M. Modular stop and go extraction tips

with stacked disks for parallel and multidimensional peptide fractionation in

proteomics. *J. Proteome Res.* **5**, 988–994 (2006).

122. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein

quantification. *Nat. Biotechnol.* **26**, 1367–1372 (2008).

123. Tyanova, S. *et al.* The Perseus computational platform for comprehensive

analysis of (prote)omics data. *Nat. Methods* **13**, 731–740 (2016).

124. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass

spectrometry-based proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552

(2022).

125. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 26 (2011).

126. Darty, K., Denise, A. & Ponty, Y. VARNA: Interactive drawing and editing of the

RNA secondary structure. *Bioinformatics* **25**, 1974–1975 (2009).

1589

TABLES

Table 1. SEPs validated in this study by tagging and western blotting. The SEP
ID is given (ID), followed by the coordinates and orientation (O). The SEPs are sorted
by length (L), and the sequences are given. If proteogenomic support was found, the
number of identified peptides is given in column P. All sORFs are located on the
chromosome (acc. NC_000911), or, if indicated, on plasmid pSYSA (AP004311).

ID	Location	O	L	Sequence	Class	P	Fig.
infC-IOF,	343,330-343,407	(+)	25	MNTTIKCASIRPNASSKLGIKSRQR	IOF		Fig. 5A, C
asPsbC	1,347,490 - 1,347,576	(+)	28	MNNVGSVSRSIPFSNPAAAARARPACQR	as		Fig. S6
Ncl1450	2,854,576 - 2,854,686	(-)	36	MTLIDTNSLHNAAFVLGTVLFVGILVWSVWFAFHMD	Ig/as		Fig. 4
Ncr1470	2,854,583 - 2,854,705	(+)	40	MWNANHTDQTRIPTKRTVPNTKAALWRLLVSMSVIQSRRC	Ig/as		Fig. 4,
crtR-IOF	982,131-982,259	(-)	42	MTLKMTQTILFPPAVPSSSLPPGSSTMRSFSLNGGYGANMSY	IOF		Fig. 5A
Sml0012	903,627 - 903,764	(-)	45	MDYLEKLLDKLRELAQKLIEGLLGPQGEPEPELIPIPVNDRRSRR	uORF	1	Fig. 5A, Fig. S8A
Ssr70935 (TU7087)	93,076 - 93,216 pSYSA	(+)	46	MQLKHWQSQKQEQKKSPGAPKKVRRQKARLQALRKKLHSSWAGNQP	ig		Fig. 5A, Fig. S8C
slr1923-IOF	2,211,455 – 2,211,595	(+)	46	MAGRENWVKKMRFILVFTKKCSPPRKKNLSPGPSGQALSAPLVVKC	IOF		Fig. 5A
NsiR7	450,353 - 450,505	(+)	49	MKPTHFIGHCPLLEILSVVQYRIADNANTKTICACISPNCGPMVSNVTDS	ig		Fig. 5A, Fig. S10B
Ncr1420	2,741,413 - 2,741,571	(+)	52	VISVKISSQPLRPSPYKAYRDPDTGQWRIAKQNREEQNNNSPNATDQDQDNG	ig	6	Fig. 5A, Fig. S10A
Ncl1350	2,706,192 - 2,706,365	(-)	57	MFWKLRQVSRFWFCYLNQPIFSRESKSTWRLSHFWSMYQISLLESCWQKSPIKETSL	ig		Fig. 5A, Fig. S9
rsbU-as	782,439 - 782,627	(-)	62	MGGVAIGKYARFGVIFRIKQGDKAMGIFQVGIHYPVEVVKNLGWTMAVQHFCPQHPPSHGHN	as	1	Fig. 5A, B
Norf2	365,077 - 365,283	(+)	68	MYAIEFEADLQDGVLTIPDHYKSLKNQHVRVVIMLDGHGNELELRALSDHSATLIDEWHDPSEDDVWI	ig	4	Fig. 5A, Fig. S8B
asCsx18	88,276 - 88,488 pSYSA	(-)	70	MKVEQATNRVLIITAAKPKGAISKITKVIRPLAMAKTRLRAKRSREGEYMLRATVRKVAEILTPPTTKIV	as		Fig. S11
Ncr1610 sORF1	3,274,013 - 3,274,231	(+)	72	MNTRTQTKIIHEGDYMAEVQVELTYTDHDWSPYLSLTEAQKLDQLRLALRRNDVKTASSLARVYHLTPVVVA	ig	3	Fig. 4F, 5A, Fig. S7B

1596

FIGURES

Fig. 1 | Establishment of TIS-Ribo-seq for the cyanobacterium *Synechocystis* 6803. **A.** Experimental set-up. Ribosomes were captured in the presence or absence of retapamulin on the translated mRNAs. In parallel, total RNA was prepared from aliquots of the same cultures. After sucrose gradient ultracentrifugation, absorption profiles were taken at 260 nm comparing samples from cultures treated with retapamulin (10 µg/mL) for 15 min and digested with MNase (Ret+(TIS)_MNase+; dark blue) or not (Ret+(TIS)_MNase-; black). The middle panel shows samples without retapamulin treatment, either after MNase digestion (Ribo_MNase+, light blue), or without (Ribo_MNase-, black). The MNase treatment depleted polysomes between 40 and 60 mm, enriching the 70S monosome peak at 30 mm. Retapamulin treatment stalled ribosomes at translation initiation sites while elongating ribosomes finished translation, resulting in an enrichment of 70S monosomes. Approximately 30 nt long footprints protected by and co-purified with 70S ribosomes were then subjected to cDNA library preparation and deep sequencing to identify the translatome and start codons, respectively. **B.** Translational efficiency (TE) for different RNA categories. The scatter plots show global TEs (TE = Ribo-seq/RNA-seq) computed from Ribo-seq replicates for all annotated coding sequences (CDS), 5′- and 3′-UTRs, tRNAs,

1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615

annotated small RNAs (sRNAs), and annotated sORFs encoding proteins of ≤70 aa.

The purple lines indicate the mean TE of the respective transcript category. **C.** Upper

2 panels, read coverage from mapped 3′ ends from TIS-Ribo-seq (dark blue) and the

Ribo-seq control (light blue), and from Ribo-seq and RNA-seq libraries (lower 2

panels) for the gene *psaM* encoding the photosystem I subunit XII. A sharp 3′ end was

observed 19 nt downstream the annotated start codon. The Ribo-seq read coverage

is enriched for the coding part, in contrast to the RNA-seq coverage, which starts at

the position of the transcription start site (bent arrow). **D.** Read coverage for the

cpcBAC2 genes encoding the phycobilisome subunits CpcB, A and the rod linker

polypeptide C2. **E.** Meta-analysis of footprints enriched in TIS-Ribo-seq. A sharp 3′

end was observed in 12,979 reads 19 nt downstream the annotated start codons of

582 genes (upper panel), exemplified in the lower panel for gene *rpl34* encoding the

large ribosomal subunit protein L34. The 3′ ends of cDNA reads are depicted for

ribosome-associated fractions after retapamulin treatment (dark blue) or without the

drug (light blue lines). **F.** Enrichment of nucleotides associated with TIS-Ribo-seq 3′

end signals with an offset of 2 nt (29 genes), 4 nt (28 genes), or 19 nt (407 genes)

from the annotated start codons after filtering for ≥10 reads. Sequence logos of the 30

nt preceding and ten nt following the start codons are shown. The preference for ATG

over GTG codons is visible, as well as a slight preference for preceding pyrimidine

residues. An enrichment for purine residues was only seen for the TIS-Ribo-seq

signals with an offset of 2 nt in the region from -10 to -15 nt, or with an offset of 4 nt

for the region from -7 to -14.

1638

1639

1640

Fig. 2 | Establishment of TTS-Ribo-seq for *Synechocystis* 6803. **A.** Experimental
set-up. Gradient profiles comparing samples from cultures treated with 10 µM Api137
for 15 min (right panel) digested with MNase (TTS (Api+)_MNase +; red) or without
(TTS (Api+)_MNase -; black) compared to samples without Api137 treatment (panel
in the middle), after MNase digestion (Ribo_MNase +, pink), or without (Ribo_MNase

-, black). After RNase treatment, approximately 30 nt long footprints protected by and

co-purified with 70S ribosomes were obtained, subjected to cDNA library preparation

and deep sequencing to identify the translatome and stop codons, respectively. In

parallel, transcriptome samples were prepared and analyzed. **B.** TTS and TIS

ribosome profiling for the gene *atpT* encoding the ATP synthase inhibitor AtpΘ[17]. Read

coverage from RNA-seq and Ribo-seq libraries (upper 2 panels), mapped 3′ ends from

Ribo-seq as control and TIS-Ribo-seq (middle panels) and from Ribo-seq control and

TTS-Ribo-seq prepared from monosomes and disomes (lower 3 panels). The Ribo-

seq read coverage is enriched for the coding part, in contrast to the RNA-seq

coverage, which starts at the position of the transcription start site (bent arrow). The

TIS-Ribo-seq 3′ ends map 19 nt downstream the start codon and the TTS-Riboseq 3′

ends 14 downstream the stop codons. **C.** TTS and TIS ribosome profiling for the genes

slr1648 and *ssr2754*. Ribosomes jam into blocked termination sites resulting in visible

polysome fractions in Api-treated samples prepared from disomes as well as

monosomes (TTS-seq, upper 2 panels). The gene *slr1648* encodes the carotenoid

oxygenase SynDiox2 involved in the biosynthesis of retinal[49], *ssr2754* encodes a

YefM-type antitoxin that forms an antitoxin-toxin gene pair with *ssr2755*[50]. **D.** Meta-

analysis of footprints enriched in TTS-Ribo-seq from prepared monosomes (replicate

2, 32 nt reads, counts per million (CPM)-normalized). A sharp 3′ end was observed 14

nt downstream of the annotated stop codons across 1,055 genes (left panel),

exemplified for gene *petB* (right panel). The 3′ ends of cDNA reads are depicted for

ribosome-associated fractions after Api137 treatment (red) or without the drug (pink

lines). **E.** Metagene analysis of footprints enriched in TTS-Ribo-seq from prepared

disomes for 358 genes (replicate 2, 56 nt reads, CPM-normalized) (left panel),

exemplified for the *psbN* gene (right panel).

1671

1672

Fig. 3 | Start codon re-annotation and detection of ribosomal frameshifting. A. Re-annotation of the start codon of *sll0553* based on TIS profiling data. The annotated longest gene model in GBK is 1038 nt, encoding a 345 aa homolog with a GTG start codon (WP_010873707.1), while the KZS annotation suggests a 306 aa protein. However, the Ribo-seq coverage and TIS-Ribo-seq mapping indicate a reading frame encoding a 298 aa protein. Blue: current annotations (345 and 306 aa). Grey: Novel annotation based on TIS profiling data (298 aa). The enriched TIS peak associated with the correct start codon is indicated with a black arrow in the TIS library and indicates an N-terminal truncation for the *sll0553*-encoded protein. The lower panel shows DNA and protein alignments of the translation initiation regions from different *Synechocystis* strains. Thus, the TIS data supports a shorter form of *sll0553* in *Synechocystis* 6803, consistent with the annotations in *Synechocystis* strains PCC

6714 and 7339. The annotated and new start codons are indicated. The corrected protein length of 298 aa is supported by 19 different peptides from the proteogenomics analysis, while no peptides were found for the elongated N termini based on the GBK and KZS annotations. **B.** Correction of Sll0872. It is annotated as a 393 nt ORF in *Synechocystis* 6803 (GBK and KZS annotations) yielding a 130 aa protein (S.6803L; BAL28778.1), while the reading frame of the homolog D082_22070 in *Synechocystis* 6714 with 69 codons is much shorter (S.6714S). The Ribo-seq data suggest that both variants are incorrect. The predicted new start is depicted in red in both strains. **C.** Ribosome frameshifting. The *prfB* gene (*sll1865*) encodes peptide chain release factor 2. RNA-seq coverage (top, black) indicates contiguous transcription from a TSS approximately 140 nt upstream the annotated reading frame beginning with a TTG start codon. Ribo-seq coverage in the 2nd panel from top indicates translation in the upstream region, supported by a TIS-Ribo-seq signal 19 nt downstream of a suitable start codon (blue line +19 in the 4th panel from top), pointing at a 26-codon reading frame finishing with a TGA stop codon. However, both datasets show an unusual, abundant read accumulation in the same subregion (red arrows) and another TIS-Ribo-seq signal, which is not associated with a start codon. This pattern is consistent with ribosomal pausing and +1 ribosomal frameshifting, which is strongly supported by mapped peptides (black horizontal lines under the aa sequence). Two (MITELTDLK and MITELTDLKR) map to the upstream 26 aa frame, including the start amino acid, while a third (LGQTQDYLDLPG) directly spans the frameshifting site (lines connected by a dashed line). Elements involved in the frameshifting are annotated alongside the RNA secondary structure with a minimum free energy of -18.60 kcal/mol. The shift site is preceded by a short spacer and an internal Shine-Dalgarno (SD) sequence. A second UAA stop site downstream is indicated that is a conserved feature of these elements to suppress translation from re-initiation at the UUG. The RNA secondary structure was predicted by RNAfold[125] with default parameters and visualized with VARNA version 3.93[126].

1714

Fig. 4 | Identification of novel sORFs encoding toxin-antitoxin candidates. A.
Ncr140 and Ncl1450 were annotated as a pair of overlapping sRNAs[51], the genomic
arrangement is shown in the center of panel A. Ribo-seq, TIS- and TTS-Ribo-seq
coverage indicate the translated segments in each transcript (orange arrows). **B.** Both
coding sequences were fused to the copper-regulated P$_{petE}$ promoter and a segment
encoding a C-terminal 3xFLAG tag. A sequence encoding superfolder GFP (sfGFP)
was engineered as control. The time course of protein induction after induction is

shown. The lower panel shows the corresponding ponceau red-stained membrane. **C.**
Alignment of 6 identical Ncl1450 homologs in *Synechocystis* strains 6803, 6714, 7338,
7339, IPPAS B-1465 and CACIAM 05 (6 Syn), with homologs from *Cyanothece* sp.
PCC 7822 and *Pleurocapsa* sp. PCC 7327. A predicted transmembrane domain is
indicated (red line, TMR). **D.** Fractionation of total cellular protein (C) into soluble (S)
and membrane (M) fractions yielded a similar localization for Ncl1450 in the membrane
fraction as the known membrane protein PetC. Mouse monoclonal anti-FLAG serum
coupled to horseradish peroxidase (A8592, Sigma) was used at a titer of 1:5000, anti-
PetC antiserum (AS08 330, Agrisera) was used at a titer of 1:2,000 and developed
with horseradish peroxidase-conjugated anti rabbit secondary antiserum (A8275,
Sigma-Aldrich) at a titer of 1:10,000. **E.** Ncr1470 is only found in *Synechocystis* strains.
The alignment shows all putative homologs that could be identified. Three sequence
variants were found, with identical sequences each in *Synechocystis* 6803 and IPPAS
B-1465 (top), in strains CACIAM 05 and PCC 6714 (center), and in strains PCC 7339
and PCC 7338 (bottom). None of the respective genes is annotated. **F.** Coding
potential around the *slr1421* locus. TIS- and TTS-Ribo-seq data suggest two sORFs
encoding small proteins of 71 and 34 aa within the previously defined sRNA *ncr1610*[51].
The coding potential of sORF1 was verified immunologically (**Fig. 5A**) and by the
detection of 3 peptides (horizontal tabs). Its start codon overlaps the stop codon of
slr1421. Bent arrows indicate transcription start sites in panels A and F. **G.** Alphafold-
predicted structure of Slr1421 (blue) and overlay with the best-matching structure
identified by Foldseek (yellow), the protein MqsR (TM score: 0.6003).

1744

1745

Fig. 5 | Validation of sORF candidates. A. Western Blot verification using anti-FLAG serum showing signals for 7 3xFLAG-tagged sORFs (upper panel) and 5 dual SPA-tagged proteins (lower panel) expressed from plasmid pVZ322 under control of the P*petE* promoter induced by copper addition. *Synechocystis* wild type (WT) samples served as negative controls. **B.** Coding potential in an asRNA to *slr2031* (*rsbU*). Both DNA strands are expressed, supported by RNA-seq (black), Ribo-Seq (turquoise), TIS-Ribo-seq (dark blue) and TTS-Ribo-seq (red) datasets. The sequence alignment at the bottom shows possible homologs of the 62 aa protein encoded by the *rsbU*-asORF in *Synechocystis* strains 6714 and 6312, and in *Leptolyngbya* sp. BL0902 identified by TBlastN in antisense orientation to their respective *rsbU* genes. However,

1756   it is unknown if corresponding asRNAs exist in these strains and the respective ORFs

1757   are longer than the shown sections. **C.** Evidence for an IOF encoding a 25 aa peptide

1758   within *slr0974* encoding the ribosome initiation factor IF-3. Bent arrows indicate

1759   transcription start sites.

1760

Fig. 6 | A family of genes encoding proteins in long internal out-of-frame reading frames. A. A protein Slr0489S is encoded within the *slr0489* reading frame (blue arrow) by a long IOF reading frame (orange arrow). Read coverage from Ribo-seq and RNA-seq libraries is shown in the lower 2 panels, the Ribo-seq read coverage is higher for the IOF segment. Mapped dual signals from TIS-Ribo-seq at the characteristic position 19 nt 3′ of the respective start codons (two central panels including control), and 14 nt 3′ of the respective stop codons from TTS-Ribo-seq (two upper panels including control) are shown. **B.** Sequence comparison of the proteins encoded by *slr0489* and two closely related loci, *slr1079* and *slr1082*. The proteins expressed from the annotated genes are labeled "L" for long. Signal peptides and transmembrane segments predicted by DeepTMHMM[60] are boxed. Black horizontal lines indicate some of the proteogenomically detected peptides for Slr0489L. Altogether, 6 peptides

were identified in the custom and 5 in the standard iPtgxDB. **C.** Sequence comparison of the proteins (labeled "S" for short) encoded IOF within *slr0489*, *slr1079* and *slr1082*. Black horizontal lines indicate some of the proteogenomically detected peptides for Slr0489S. **D.** Co-IP analysis of Slr0489L-3xFLAG in three replicates. The volcano plot shows the protein interaction partners of Slr0489L-3xFLAG identified by mass spectrometry compared to sfGFP-3xFLAG. The two most enriched proteins, Slr0489L and Slr0489S are colored red, the locus IDs are given for co-enriched proteins in grey. For the complete list of 512 detected proteins and further details, see **Table S6**. **E.** Co-IP analysis of Slr0489S-3xFLAG. Details as in panel D. For the complete list of 586 detected proteins, see **Table S7.** The results of an analogous co-IP experiment for *slr1079* encoding Slr1079L/S are shown in **Fig. S14**. **F.** Effects of overexpressing *slr0489* (without FLAG tag) in a *Synechocystis* 6803 Δ*slr0489* deletion mutant, either with both intact reading frames (Δ*slr0489*+*slr0489* +R), of Slr0489S separately (Δ*slr0489*+*slr0489*S +R), or of Slr0489L from a variant in which a single point mutation prevented the expression of Slr0489S (Δ*slr0489*+*slr0489*L +R). The overexpression was triggered by the addition of rhamnose (+R), which was omitted from the respective controls (-R). The wild type (WT) and Δ*slr0489* deletion mutant served as further controls. The lines show the averaged increase in OD_{750} over 120 h. Cultures were measured in technical duplicates of two independent biological replicates.