Title: Insecticide alters the evolution of glyphosate resistance in *Ipomoea purpurea*

Authors: Grace M. Zhang[1] and Regina S. Baucom[1]

[1]Ecology and Evolutionary Biology Department, University of Michigan

Corresponding author: Grace M. Zhang (gmzhang@umich.edu), Ecology and Evolutionary

Biology Department, 4034 Biological Sciences Building, University of Michigan, Ann Arbor,

MI 48109

Data availability: Data will be made publicly available upon manuscript acceptance.

Author contributions: GMZ and RSB designed the field experiment. GMZ performed the

experiment and analyzed the data. GMZ and RSB wrote the manuscript.

Funding: This work is supported by the AFRI Predoctoral Fellowships Program, project award

no. 2024-67011-42912, from the U.S. Department of Agriculture's National Institute of Food

and Agriculture. This work was also supported by funding from the University of Michigan.

Conflicts of interest: None declared.

Acknowledgements: We thank N. Hammond, G. Lu, and M. Vaughan-Coston for assisting with

the field experiment and data processing, M. Palmer and the Matthaei Botanical Gardens for

field support, and V. Iriart for providing feedback for the manuscript.

ABSTRACT

The evolution of plant resistance naturally occurs in complex, multifaceted environments that

consist of multiple simultaneous stressors. Understanding how shifting environmental contexts

may shape resistance evolution requires empirical studies that consider the combined effects of

interacting stressors on fitness and selection. Here, we examined how exposure to an insecticide

impacts the evolution of resistance to the herbicide glyphosate in *Ipomoea purpurea* (common

morning glory). Through a factorial field experiment, we manipulated glyphosate and an

insecticide to estimate selection on glyphosate and herbivory resistance. We found that

glyphosate acted as the primary agent of selection, favoring higher levels of glyphosate

resistance. In the presence of glyphosate alone, positive correlational selection favored a

combination of higher glyphosate and herbivory resistance, supporting prior work that suggested

these traits may be linked. Importantly, insecticide exposure modified both glyphosate resistance

and the strength of selection acting upon the trait by increasing resistance and weakening

selection. Together, our results indicate that the evolution of herbicide resistance is context-

dependent and that secondary stressors like insecticide can alter the evolutionary trajectories of

plant defense.

INTRODUCTION

Plants experience natural and human-altered environments characterized by multiple, shifting

stressors. How organisms adapt to such complex environments remains a central question in

evolutionary biology. The evolution of plant resistance to damage—whether from natural

enemies, abiotic stressors, or anthropogenic agents—provides some of the clearest

demonstrations of adaptation in action. Classic studies of resistance have yielded insights into the

genetic basis of adaptation, the costs of defense, and the trade-offs that constrain evolutionary

trajectories (Warwick 1991; Bergelson and Purrington 1996; Strauss et al. 1999). For example,

the framework of assessing fitness costs and benefits emerged from studies of plant defense

against herbivory (Simms and Rausher 1987, 1989), and has since been extended to other natural

biotic stressors such as fungal pathogens (Simms and Rausher 1993; Tian et al. 2003) and

competition (Chaney and Baucom 2014), as well as abiotic stressors like herbicide (Baucom and

Mauricio 2008; Debban et al. 2015). Increasingly, studies recognize that organisms experience

multiple, co-occurring stressors in nature (Siemens et al. 2003; Gassmann and Futuyma 2005;

Wise and Rausher 2016), where both the benefits and costs of resistance can depend strongly on

environmental context and shift across ecological gradients (Koricheva 2002; Cipollini et al.

2017). However, more empirical work involving multiple stressors is needed to predict plant

evolutionary trajectories when selective pressures are indirect, nonlinear, or causally linked

(Wise 2023).

Recent work has shown that overlapping biotic and abiotic pressures characterizing a complex

environment may interact to shape both the expression and fitness consequences of plant defense

(Côté et al. 2016; Desaint et al. 2021; Zandalinas et al. 2024). Anthropogenic chemicals such as

herbicide and insecticide provide powerful systems for studying contemporary evolution in

agricultural landscapes. Herbicides like glyphosate (the active ingredient of Roundup) are highly

effective at weed control and impose strong selection on weed populations, driving the evolution

of resistance (Culpepper 2006; Baucom 2019; Kreiner et al. 2022). Both the application of

glyphosate and the evolution of glyphosate resistance may have cascading ecological and

evolutionary consequences for plants and their associated communities, such as altering the

patterns of insect herbivory (Zhang and Baucom 2024). At the same time, insecticides commonly

co-occur with herbicides in agricultural contexts (Kwon and Penner 1995), potentially modifying

the selective landscape beyond their direct effects on herbivores. Indeed, combinations of

herbicide and insecticide with diverse modes of action may differentially affect plant species

exposed to these chemicals (Daramola et al. 2023). However, while ecological interactions

between herbicide and insecticide have been documented, there have been few empirical tests of

whether a secondary chemical can alter the strength, direction, and causal pathways of selection

on resistance traits.

We address this gap by examining how plant resistance evolves in environments mediated by

anthropogenic chemicals: the herbicide glyphosate, which exerts strong selective pressure on

plants, and the insecticide spinosad, which reduces insect herbivory. To test whether exposure to

an insecticide influences the evolution of glyphosate resistance, we conducted a factorial

common garden experiment examining the effects of each treatment environment on resistance

and fitness in *Ipomoea purpurea* (common morning glory). Previous work in this species shows

that glyphosate resistance confers a fitness benefit in the presence of glyphosate, but no

detectable fitness cost measured as seed production in glyphosate's absence (Baucom and

Mauricio 2008; Van Etten et al. 2016). The addition of insecticide could modify these outcomes

if it interacts with glyphosate synergistically or antagonistically (Figure 1). For example, the

fitness benefit of glyphosate resistance in the presence of glyphosate may be strengthened or

weakened by the application of the insecticide. Conversely, in the absence of glyphosate but in

the presence of insecticide, glyphosate resistance may incur a cost, or alternatively confer a

previously undetected fitness benefit.

In this study, we explore how interacting stressors—herbicide and insecticide—may influence

plant adaptation by directly impacting fitness and indirectly *via* trait-mediated pathways by

asking: 1) Is there a fitness benefit of glyphosate resistance (Figure 1 green box), but no evidence

of a cost in terms of seed production (Figure 1 blue box), in line with previous work? 2) Does

insecticide exposure impact glyphosate resistance directly and/or alter the patterns of selection

acting on this trait? If so, the direction and magnitude of selection acting on glyphosate

resistance in the presence of insecticide (Figure 1 yellow and white boxes) would differ from the

patterns of selection detected in spinosad's absence (Figure 1 green and blue boxes). And 3) do

glyphosate and insecticide application directly and/or indirectly impact the plant evolutionary

trajectory *via* resistance to glyphosate or herbivory? Untangling direct selection from indirect

selection is a key component of studying multivariate evolution, and a factorial experimental

design allows for the examination of chains of selective causality in complex, multi-stressor

environments.

MATERIALS AND METHODS

Study system

Ipomoea purpurea (L.) Roth (Convolvulaceae), the common morning glory, is an annual weed

frequently found in agricultural fields and disturbed soil beds in the United States. The species

begins germination in the early spring and continues through the growing season. It grows

rapidly, capable of producing over 8,000 seeds in a season before being killed by the first winter

frost (Chaney and Baucom 2014). In Michigan, *I. purpurea* is fed upon by generalist coleopteran

and lepidopteran herbivores such as *Popillia japonica* (Japanese beetle) and *Trichoplusia ni*

(cabbage looper) (Zhang and Baucom 2024), and prior work shows that herbivory resistance is

under positive selection in this species (Rausher and Simms 1989). Populations of the species are

also naturally resistant to the herbicide glyphosate, likely through detoxification and cell

transport mechanisms (Baucom and Mauricio 2004; Kuester et al. 2015; Gupta et al. 2023).

Field experiment

On June 9, 2022, we planted replicate *Ipomoea purpurea* seeds from 45 family lines in a

common garden field plot at the Matthaei Botanical Gardens in Ann Arbor, Michigan. These

lines were generated from crosses between individuals (*i.e.*, maternal lines) sampled as seed from

natural populations found in the US Southeast that exhibited variable levels of glyphosate

resistance (Kuester et al. 2015). Individuals were randomly chosen from populations, selfed

once, and then crossed in a haphazard design to randomize the genetic background. To generate

experimental individuals, we selfed between one to five randomly selected half-sib families from

19 paternal lines to generate sufficient sample sizes. Four replicate seeds from each half-sib

family were then randomly planted in each of four treatment environments in a factorial design

of herbicide (glyphosate) and insecticide (spinosad, details below)—1) glyphosate-present,

spinosad-absent, 2) glyphosate-absent, spinosad-absent, 3) glyphosate-present, spinosad-present,

and 4) glyphosate-absent, spinosad-present. These combinations were repeated in two blocks, for

a total of 1440 individual seeds. We watered daily and used a combination of weed fabric and

hand-weeding to control weeds. Germination started a few days after planting and was complete

within ten days.

Two weeks after the seeds were planted, we started applying a field dose of 0.30 kg ai/L of the

insecticide spinosad (Garden Insect Spray, Monterey Lawn and Garden Products, Inc., Fresno,

CA, USA) to the plants every other week for a total of five applications throughout the length of

the experiment. Spinosad, an broad-spectrum organic insecticide derived from a naturally

occurring soil bacterium *Saccharopolyspora spinosa* (Bacteria: Actinobacteridae) (Mertz and

Yao), is commonly used to protect vegetable crops and stored grain against insect herbivores by

triggering uncontrollable muscle activity in targeted insects that leads to death (Mertz and Yao

1990; Bunch et al. 2014; Zhang et al. 2021). We conducted a preliminary growth room

experiment to ensure that spinosad would not affect plant biomass or act synergistically with

glyphosate (see Figure S1 for details). We counted the number of leaves on all plants five weeks

after planting in order to estimate plant size (Debban et al. 2015). Six weeks after planting, we

applied the herbicide glyphosate (Roundup WeatherMax, Bayer Corp., Whippany, NJ, USA) at

the sublethal dose of 1.0 kg ai/ha to the glyphosate-present treatment plots using a handheld

CO_2-pressurized sprayer (Spraying Systems Co., Wheaton, IL, USA) at a rate of 187 L/ha at 30

psi with a stride pace of 90 paces/min (Kuester et al. 2015). Because it was necessary to measure

individual-level fitness later in the experiment, we chose a dose lower than the recommended

field dose for *Ipomoea spp.* of 1.54 kg ai/ha (Monsanto 2017). Our chosen dose of 1.0 kg ai/ha

was in line with past studies that aimed to merely stress the plants rather than eradicate them and

is well within the range of recommended field doses of glyphosate for weed control (Duke et al.

2012; Vázquez et al. 2021). We scored damage two weeks after glyphosate application by

dividing the number of damaged (*i.e.,* yellowed, browned, and/or shriveled) leaves on each plant

by the total number of leaves in order to calculate the proportion of damaged leaves (per

(Baucom and Mauricio 2008)). We also took note of which plants had died due to glyphosate

application.

Five weeks after the glyphosate application, we randomly collected four leaves from each plant

for herbivory damage estimates. Leaves were sandwiched between paper towels and kept cool on

ice to be brought back to the lab, where they were scanned with a Canon CanoScan LIDE 400

flatbed scanner (Canon USA, Melville, NY, USA) to generate high-resolution images at 600 dpi.

Herbivory damage was estimated as the percentage of tissue missing per leaf using the iOS app

LeafByte (Getman-Pickering et al. 2020), then averaged across the four leaves for each plant. In

line with our previous work, we elected to be conservative with our herbivory damage estimates

and excluded ambiguous damage that could be attributed to glyphosate from our estimates

(Zhang and Baucom 2024). Additionally, we noticed that many plants had leaves exhibiting

patches of rust fungus; thus we scored the presence/absence of the fungus on each leaf as a

binomial (1 - present, 0 - absent) and averaged those values across the four leaves for each plant.

Finally at the end of the field season, we collected seeds from all plants for two rounds of

collection in order to estimate fitness using seed number.

Statistical analyses

We operationally defined resistance as 1-p, where p is the amount of damage experienced by an

individual plant. Thus, when calculating glyphosate resistance, p refers to the proportion of

leaves damaged by glyphosate per plant (Baucom and Mauricio 2008). Likewise, when

calculating herbivory resistance, p is the average percentage of leaf tissue missing due to

herbivory for each plant (Rausher and Simms 1989).

All analyses were conducted using R 4.3.2 (R Core Team 2023). For the field experiment, we

excluded any individual plants that did not germinate (N = 229), as well as plants in the

glyphosate-absent treatments that were inadvertently damaged by glyphosate (N = 43). We also

removed any plants that died before herbivory data and/or seed number could be recorded.

Ultimately, we had 258 plants in the glyphosate-present, spinosad-absent treatment, 233 plants in

the glyphosate-absent, spinosad-absent treatment, 262 plants in the glyphosate-present, spinosad-

present treatment, and 255 plants in the glyphosate-absent, spinosad-present treatment, for a total

of 1008 plants.

Treatment effects of glyphosate and spinosad

We wanted to confirm the efficacy of treatment regimes. First, we built linear mixed models

using the package *lme4* (Bates et al. 2015) to determine if spinosad reduced herbivory as

expected. Using all experimental plants, we constructed the following model:

herbivory damage ~ spinosad + glyphosate + leaf count + rust + 1|family + 1|block

We transformed the response variable of herbivory damage using rank normalization in order to

ensure normality after using the package *bestNormalize* (Peterson 2021) to identify the

appropriate transformation. Spinosad and glyphosate were included as fixed effects, while leaf

count (as a proxy for plant size) and the proportion of collected leaves showing rust fungus were

included as covariates in all models. The Satterthwaite approximation was used to determine the

degrees of freedom of the fixed-effect variables. Family line and block were included as random

effects unless stated otherwise. To test the significance of random effects, we removed each one

and used the χ^2 test of difference (df = 1) to compare the simplified model to the full model.

Next, we wanted to determine if glyphosate influences herbivory damage (based on prior results

(Zhang and Baucom 2024)) or alternatively, if spinosad altered glyphosate resistance. We first

tested for an effect of glyphosate application on herbivory damage in the spinosad-present and -

absent environments separately using the following model, with herbivory damage transformed

using rank normalization in both environments:

herbivory damage ~ glyphosate + leaf count + rust + 1|family + 1|block

To examine whether spinosad application may have had an effect on glyphosate resistance, we

built the following linear mixed model using just the plants treated with glyphosate, with

glyphosate resistance transformed using rank normalization:

glyphosate resistance ~ spinosad + leaf count + rust + (1 + spinosad|family) + 1|block

We specifically tested for an interaction between spinosad and family line while letting the

slopes and intercepts vary because we were interested in whether the effect of spinosad

application on glyphosate resistance showed genetic variation.

Genetic variation and among-family variance in glyphosate and herbivory resistance

We assessed if there was evidence for genetic variation for glyphosate and herbivory resistance

by including family line as a fixed effect in the following linear mixed model:

$$\text{resistance} \sim \text{family} + \text{leaf count} + \text{rust} + 1|\text{block},$$

where the resistance was either glyphosate or herbivory resistance. For either resistance, we

tested for genetic variation first among all plants that had been exposed to that stressor to

determine overall variation in the trait, then in both the presence and the absence of the other

stressor to account for any treatment effects of the second stressor. Glyphosate resistance as a

response variable in the absence of spinosad was transformed using a log transformation to

ensure normality, while both glyphosate and herbivory resistance in all other environments were

transformed using rank normalization.

Using the model we previously built to assess the effect of spinosad on glyphosate resistance

(glyphosate resistance ~ spinosad + leaf count + rust + (1 + spinosad|family) + 1|block), we also

determined whether the presence of spinosad affected the partitioning of variance in glyphosate

resistance attributable to family line among glyphosate-treated plants. This model allowed for

among-family variance to be partitioned separately depending on the presence of spinosad, with

variance in the intercept reflecting among-family variance in the spinosad-absent environment,

and variance in the slope reflecting additional among-family variance attributed to differential

family responses to spinosad. We then used parametric bootstrapping (n = 1000 simulations) to

generate 95% confidence intervals on intraclass correlation coefficients (ICCs) within spinosad

environments. ICCs were calculated in each environment as the ratio of among-family variance

to total variance (*i.e.*, among-family + block + residual variance).

Similarly, we tested if the presence of glyphosate influenced the variance partitioning in

herbivory resistance attributable to family line among plants in the spinosad-absent environment

using a similar model: herbivory resistance ~ glyphosate + leaf count + rust + (1 +

glyphosate|family) + 1|block. ICCs and 95% confidence intervals were calculated in the presence

and absence of glyphosate as previously described.

Selection analyses on resistance traits

We conducted selection analyses on the two forms of resistance in each of the four treatment

environments to determine if either spinosad or glyphosate would affect the patterns of selection

acting on resistance to glyphosate and herbivory. We also assessed if there were fitness costs and

benefits of each form of resistance depending on the treatment environment. If there was positive

selection for a form of resistance in the presence of the corresponding stressor, then we would

conclude there was a fitness benefit associated with that form of resistance. The converse was

also examined—negative selection for a form of resistance in the absence of the stressor would

suggest there was a fitness cost associated with resistance to that stressor.

In accordance with past studies, we used a modified form of Lande and Arnold's method of

detecting selection using partial regression coefficients (Lande and Arnold 1983; Rausher 1992;

Mauricio and Mojonnier 1997). Instead of performing selection analyses on phenotypic trait

values, we used family line means of trait values to conduct genotypic selection analyses in order

to eliminate environmental biases (Rausher 1992; Stinchcombe et al. 2002). Additionally, we

used the residuals of untransformed values of relative fitness and resistance after removing the

effects of block to further remove environmental effects from the analyses. Relative fitness was

calculated within each environment by dividing the fitness of each plant by the average fitness of

all plants in that environment. After calculating family means of relative fitness and resistance

trait values, we standardized the resistance traits in each environment such that they had an

average of zero and a standard deviation of one. Relative fitness was then regressed on each form

of resistance within each environment to determine the pattern of selection for that particular

trait. Directional selection was estimated using models with only linear terms, while nonlinear

(*i.e.*, stabilizing/disruptive) selection was estimated using models containing both linear and

quadratic terms. All selection gradients for nonlinear selection were reported as twice the

quadratic regression coefficient.

In the glyphosate-present, spinosad-absent environment (Figure 1 green box), we assessed the

fitness benefits of glyphosate and herbivory resistance. The color in parentheses next to each trait

indicates the environment from which mean trait values were obtained, in line with Figure 1:

relative fitness (green) ~ glyphosate resistance (green)

relative fitness (green) ~ herbivory resistance (green)

In the glyphosate-absent, spinosad-absent environment (blue box), we tested for the cost of

glyphosate resistance and the benefit of herbivory resistance:

relative fitness (blue) ~ glyphosate resistance (green)

relative fitness (blue) ~ herbivory resistance (blue)

In the glyphosate-present, spinosad-present environment (yellow box), we assessed the benefit of

glyphosate resistance and cost of herbivory resistance:

relative fitness (yellow) ~ glyphosate resistance (yellow)

relative fitness (yellow) ~ herbivory resistance (green)

Lastly, in the glyphosate-absent, spinosad-present environment (white box), we tested for the

costs of glyphosate and herbivory resistance:

relative fitness (white) ~ glyphosate resistance (yellow)

relative fitness (white) ~ herbivory resistance (blue)

To detect correlational selection between the two forms of resistance in each environment, we regressed relative fitness on the interaction term between the two resistance traits, as well as all linear and quadratic terms of the resistance traits. We then used the *fields* package (Nychka et al. 2021) to generate a fitness landscape fitted with thin-plate splines. We further tested for genetic correlations between the two forms of resistance in each environment by calculating Pearson's correlation coefficients.

Since we were interested in whether spinosad might alter patterns of selection (if any) that were acting on glyphosate resistance, we built the following model to compare the selection gradients between the two glyphosate-present environments:

relative fitness ~ glyphosate resistance × spinosad + glyphosate resistance + spinosad

Relative fitness and the standardization of glyphosate resistance were calculated within environments, as described above. A significant interaction between glyphosate resistance and spinosad would suggest that the selection gradients were altered by the presence of spinosad.

Direct and indirect effects on fitness

We assessed the direct and indirect effects of glyphosate and spinosad on fitness—how do the dual treatments of herbicide and insecticide affect resistance, and how does that impact fitness? To answer these questions, we constructed an SEM using all experimental plants, specifically testing the effects that glyphosate and spinosad have on both glyphosate resistance and herbivory

resistance, as well as seed number. We also tested for relationships between the resistances and

fitness. With family line and block included as random effects, the following relationships were

modeled using the package *piecewiseSEM* (Lefcheck 2016), and the Fisher's C statistic was used

to determine model fit:

glyphosate resistance ~ glyphosate + spinosad + 1|family + 1|block

herbivory resistance ~ glyphosate + spinosad + 1|family + 1|block

seed count ~ glyphosate + spinosad + glyphosate resistance + herbivory resistance

353   + 1|family + 1|block

We did not include a relationship between glyphosate resistance and herbivory resistance in the

final model. Doing so would have resulted in a saturated model, which would not have allowed

us to assess model fit. Additionally, the saturated model reported that the relationship between

the two resistances was not significant, and including the relationship resulted in a model with a

higher AIC score (saturated: 6408.696 vs. current model: 6402.301), which suggests a worse fit.

Thus, we proceeded with the SEM that omitted this relationship.

RESULTS

Treatment effects of glyphosate and spinosad

Glyphosate was an effective stressor of *Ipomoea purpurea*. While only ten plants died from

herbicide application, all but one of the 520 glyphosate-treated plants exhibited leaf damage from

glyphosate (on average, 37.7% of leaves per plant were damaged) and reduced total seed

production (mean seed number in glyphosate-treated plants: 290 vs. untreated plants: 1723).

There was no effect of glyphosate on chewing herbivory damage overall ($F_{1, 979} = 2.14$, p =

0.144, Figure 2a, Table S1), nor was there difference in herbivory damage between glyphosate

treatments for plants in the spinosad-absent environment ($F_{1, 486} = 0.21$, p = 0.648, Table S2).

However, glyphosate marginally increased the amount of herbivory damage experienced by

plants in the presence of spinosad ($F_{1, 487} = 3.18$, p = 0.075, Table S2).

Strikingly, plants that were exposed to both glyphosate and spinosad showed higher levels of

resistance to glyphosate than plants that were just exposed to glyphosate (spinosad: 69.43%

leaves showed no glyphosate damage, no spinosad: 55.14% leaves showed no glyphosate

damage; $F_{1, 233} = 71.83$, p < 0.001; Figure 2b, Table S3). There was no interaction between

spinosad treatment and family line, however, indicating that families generally experienced

similar levels of increased glyphosate resistance in the presence of the spinosad ($\chi^2_1 = 0.81$, p =

0.847; Table S3). As expected, spinosad application significantly reduced herbivory overall

(untreated: 0.53%, treated: 0.15%; $F_{1, 975} = 174.06$, p < 0.001; Figure 2c, Table S1), but did not

eliminate herbivory.

Genetic variation and among-family variance in resistance to glyphosate and herbivory

We investigated whether there was genetic variation for the two forms of resistance and found

evidence of genetic variation for glyphosate resistance across all plants in glyphosate-present

environments regardless of spinosad treatment ($F_{44, 472} = 1.46$, $p = 0.033$; Table S4), as well as

glyphosate-treated plants that were not treated with spinosad ($F_{44, 210} = 1.67$, $p = 0.009$; Table

S4). There was no genetic variation for glyphosate resistance detected among plants that were

treated with both glyphosate and spinosad ($F_{44, 214} = 0.99$, $p = 0.492$; Table S4). Among-family

variance in glyphosate resistance in the presence of spinosad (ICC = 0.079, 95% CI: 0.006 -

0.178; Table S5) was comparable to in spinosad's absence (ICC = 0.036, 95% CI: 0.001 - 0.121;

Table S5) given the wide, overlapping confidence intervals.

In contrast to glyphosate resistance, there was no evidence of genetic variation underlying

herbivory resistance among plants that were not treated with spinosad ($F_{44, 443} = 0.61$, $p = 0.977$;

Table S6). Nor was there evidence of genetic variation for herbivory resistance in either the

glyphosate-present or glyphosate-absent environments (glyphosate-present: $F_{44, 211} = 0.70$, $p =$

0.924; glyphosate-absent: $F_{44, 185} = 0.89$, $p = 0.664$; Table S6). Likewise, there was no difference

in among-family variance in herbivory resistance in the presence and absence of glyphosate

(glyphosate-present ICC = 0.002, 95% CI: 0.000 - 0.091; glyphosate-absent ICC = 0.002, 95%

CI: 0.000 - 0.089; Table S7).

Glyphosate resistance is under positive selection

We tested for selection acting upon family line averages of glyphosate and herbivory resistance

in each of the four treatment environments in order to determine if the presence or absence of

412 either glyphosate or spinosad would alter the patterns of selection, as well as whether there were

413 fitness costs and benefits associated with either form of resistance. There were two instances in

414 which selection was detected. In the presence of glyphosate and but the absence of spinosad (*i.e.*,

415 herbivores were present), there was positive selection for glyphosate resistance, meaning that

416 there is a fitness benefit associated with that trait ($\beta = 0.46$, $F_{1,43} = 20.21$, $p < 0.001$; Figure 3a;

417 Table 1). In contrast, in the presence of both glyphosate and spinosad, there was marginal

418 positive selection for glyphosate resistance ($\beta = 0.18$, $F_{1, 43} = 3.28$, $p = 0.077$; Figure 3c). The

419 strength of selection acting on glyphosate resistance was contingent on the presence of spinosad

420 ($F_{1, 86} = 3.96$, $p = 0.050$; Table S8); selection was stronger in the spinosad-absent environment.

421 No other evidence of selection of either linear or quadratic on glyphosate or herbivory resistance

422 was detected (Table 1, Table S9).

424 Interestingly, in the environment with only glyphosate, there was correlational selection acting

425 on resistance to glyphosate and herbivory, favoring high levels of both forms of resistance

426 (correlational selection gradient $= 0.40$, $F_{1, 39} = 3.99$, $p = 0.053$; Figure 4; Table S9). However,

427 there was no selection acting on herbivory resistance in this environment ($\beta = 0.13$, $F_{1, 43} = 1.10$,

428 $p = 0.299$; Figure 3a; Table 1), and the two resistance traits were not genetically correlated ($r^2 =$

429 0.13, $p = 0.383$). There were no signs of correlational selection in any other environment (Table

430 S9).

432 **Direct and indirect effects of treatments on resistance and seed production**

We assessed the direct and indirect effects of glyphosate and spinosad treatments on glyphosate

resistance, herbivory resistance, and fitness by constructing a well-fit structural equation model

(SEM) that had a Fisher's C statistic of 0.007 (p = 0.996; Figure 5). As expected, glyphosate

treatment negatively impacted glyphosate resistance (standardized β = -0.75, df = 984, p <

0.001), which is necessarily the case because unexposed plants would not have experienced

glyphosate damage and therefore would have perfect resistance. In turn, glyphosate resistance

was positively associated with seed production, meaning that more resistant plants produced

more seeds (standardized β = 0.28, df = 725, p < 0.001). In contrast, the impact of glyphosate on

seed production was negative (total effect = -0.76), but split into the primary direct effect

(standardized β = -0.55, df = 923, p < 0.001) and an indirect effect that was mediated by

glyphosate resistance (-0.75 * 0.28 = -0.21). Glyphosate had no effect on herbivory resistance

(standardized β = -0.02, df = 923, p = 0.423).

Spinosad treatment predictably led to increased herbivory resistance (*i.e.*, decreased herbivory;

standardized β = 0.38, df = 978, p < 0.001)—plants exposed to spinosad are by necessity more

resistant to herbivory because they experience less herbivory damage. Strikingly, spinosad also

increased glyphosate resistance (standardized β = 0.16, df = 981, p < 0.001), thereby indirectly

and weakly increasing seed production (total effect (0.04) = indirect effect through glyphosate

resistance (0.16 * 0.28 = 0.04)). The insecticide had no direct effect on seed production

(standardized β = -0.01, df = 965, p = 0.727), nor did herbivory resistance impact fitness

(standardized β = -0.03, df = 985, p = 0.149). Thus, the negative effect of glyphosate and the

positive effect of spinosad on fitness were both mediated through resistance to glyphosate to

some degree.

DISCUSSION

DISCUSSION

We conducted a factorial field experiment to examine how exposure to an insecticide (spinosad)

shapes the evolution of resistance to an herbicide (glyphosate) in *Ipomoea purpurea*. Two main

patterns emerge. First, glyphosate is the dominant selective agent, consistently reducing fitness

and favoring higher resistance in glyphosate-present environments. Second, spinosad

unexpectedly alters glyphosate resistance and its evolutionary dynamics, increasing resistance

and modifying the strength of selection. Together, these results demonstrate that resistance

evolution is strongly context-dependent, and additional stressors can reshape both the expression

of resistance traits and the selection acting upon them.

Glyphosate resistance is under positive selection

Glyphosate application, even at a sublethal dose, imposed strong fitness consequences, delaying

development and substantially reducing seed production. Consistent with previous work in this

system (Baucom and Mauricio 2008), glyphosate resistance in the presence of glyphosate

experienced significant positive selection in the spinosad-absent environment, indicating a clear

fitness benefit of increased glyphosate resistance. Importantly, we detected genetic variation for

glyphosate resistance across plants exposed to glyphosate, suggesting continued evolutionary

potential (Zhang and Baucom 2024). The structural equation model (SEM) reinforces this

conclusion—glyphosate reduced fitness primarily through a strong direct effect, with an

additional indirect effect mediated by glyphosate resistance. These results highlight glyphosate

as the principal driver of evolutionary change in this study. However, despite strong selection,

we detected no fitness costs of glyphosate resistance measured as seed production, regardless of

the presence of spinosad. This absence of detectable costs is consistent with prior studies that

identified costs in seed viability, germination, or plant size (Debban et al. 2015; Van Etten et al.

2016) but not seed production. Although glyphosate resistance clearly enhances fitness in the

presence of herbicide, its associated costs may be conditional on dose, genetic background, and

experimental environment (Neve et al. 2014). Further, while our experimental design is only

suitable to explicitly detect constitutive but not induced resistance, previous work has shown that

glyphosate resistance in *Ipomoea purpurea* is likely to be constitutive (Gupta et al. 2023). If the

mechanism of resistance included induced resistance, however, we would not expect to identify

fitness costs of resistance in the absence of glyphosate.

Although there was no univariate selection on herbivory resistance, we did find correlational

selection favoring jointly higher levels of glyphosate and herbivory resistance in the presence of

glyphosate and absence of spinosad (*i.e.*, herbivores were present). This pattern suggests that as

glyphosate resistance evolves, herbivory resistance may also increase—even in the absence of

detectable genetic variation for herbivory resistance itself. This interpretation aligns with our

previous SEM-based work, which inferred a causal relationship between glyphosate and

herbivory resistance (Zhang and Baucom 2024).

One explanation is that the two resistance traits may share underlying genetic mechanisms.

Glyphosate inhibits 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), a key enzyme

upstream of metabolic pathways that synthesize aromatic amino acids—the precursors of many

secondary metabolites involved in chemical defense against herbivores (Amrhein et al. 1980;

Mithöfer and Boland 2012; Fuchs et al. 2021). The likely mechanism of glyphosate resistance in

Ipomoea purpurea—metabolic detoxification coupled with enhanced transport (Gupta et al.

2023)—could affect herbivory resistance in two ways. First, detoxification may reduce the

number of glyphosate molecules that reach and inhibit EPSPS (Baek et al. 2021), thereby

mitigating glyphosate's indirect suppression of anti-herbivore defense compound biosynthesis.

Second, loci underlying detoxification pathways (*e.g.*, cytochrome P450s, glycosyltransferases,

and ABC and phosphate transporters) may have pleiotropic effects on herbivory resistance. For

example, uridine diphosphate glucosyltransferases (UGTs) regulate the glycosylation of

phytohormones central to herbivory defense such as abscisic acid, jasmonates, and salicylic acid

(Dimunová et al. 2022; Gharabli et al. 2023). If glyphosate and herbivory resistance are linked

due to pleiotropy, the relationship between the traits will persist, and correlational selection will

likely shape the evolution of those traits concurrently over time (Svensson et al. 2021).

Spinosad alters glyphosate resistance and selection

One of the more interesting findings of this study was that spinosad application increased

glyphosate resistance and indirectly increased fitness through this effect. Plants exposed to

spinosad exhibited higher glyphosate resistance, as well as weaker evidence (and strength) of

positive selection for that form of resistance. While there was reduced genetic variation for

glyphosate resistance in the presence of spinosad, the proportion of variance attributable to

family line was not affected by spinosad, suggesting that the loss of significance in the spinosad-

present environment is due to reduced statistical power to detect family line variation.

Nonetheless, the results suggest that spinosad modifies the expression and evolutionary

dynamics of glyphosate resistance. Our design does not allow us to fully disentangle the effects

of spinosad from reduced herbivory, but the SEM suggests that spinosad acts directly on

glyphosate resistance rather than indirectly through herbivory. This illustrates how additional

stressors can alter not only trait expression, but also the strength of selection.

A plausible explanation for spinosad's effect on glyphosate resistance is physiological priming

via shared detoxification pathways. Resistance to spinosad in insects can be conferred *via*

metabolic detoxification, particularly by upregulating enzymes such as monooxygenases,

cytochrome P450s, and UGTs (Sparks et al. 2012; Bastarache et al. 2023; Wang et al. 2024).

Notably, plants have metabolic detoxification processes that trigger the same classes of enzymes

(Edwards et al. 2011; Gharabli et al. 2023), and importantly, the same processes likely confer

resistance to glyphosate in *Ipomoea purpurea* (Gupta et al. 2023). Thus, it is possible that the

application of spinosad may prime plants to become more resistant to glyphosate by activating

the same detoxification pathways. Moreover, while spinosad is generally thought to have low

systemic activity in plants (National Organic Standards Board Technical Advisory Panel 2002),

one study found that applying the insecticide to the roots of *Solanum lycopersicum* (tomato) was

enough to control the herbivore *Tetranychus urticae* (spider mites) on its leaves (Van Leeuwen et

al. 2005). Additionally, spinosad was detected in both the aboveground and belowground tissue

of *Allium cepa* (onion) that had been treated with the insecticide as seeds for up to 60 days post

planting (Moretti et al. 2021). It seems evident that spinosad has the potential to translocate in

plant tissue. If *I. purpurea* can similarly take up spinosad, the insecticide may activate

detoxification pathways, priming the plants to respond quickly to the herbicide and better protect

from damage.

The potential of insecticides to modify herbicide efficacy has long been considered, given the

potential for antagonizing chemicals to reduce weed control. However, such studies generally

focus on herbicide performance in terms of plant death or reduction in biomass, without

considering further ecological or evolutionary implications (Pankey et al. 2004; Daramola et al.

2023). Our work explicitly explores the consequences of insecticide application on the evolution

of herbicide resistance in an agricultural weed, showing that spinosad enhances glyphosate

resistance and thereby increases fitness, but also reduces the strength of selection acting on this

trait. Interestingly, spinosad does not affect the lack of a fitness cost of glyphosate resistance.

These results imply that with time, plants exposed to both glyphosate and spinosad will be more

resistant to glyphosate and therefore more fit than plants that have been exposed to glyphosate

alone due to the plastic effects of spinosad. Additionally, while these effects are likely species-

specific, to our knowledge our study is the first to show that spinosad interacts antagonistically

with glyphosate to reduce herbicide efficacy in a weed (Sparks et al. 2004; Scroggs et al. 2005).

Understanding how the interactions between herbicide and insecticide impacts the evolution of

herbicide resistance is essential for predicting the persistence of resistance in heterogeneous,

complex environments such as agricultural landscapes.

Context-dependent ecological interactions

Much of the work examining interactions between insecticides and herbicides have focused on

the impact of insecticide on herbicide efficacy. However, herbicide may influence insecticide

efficacy as well. This relationship has primarily been studied in lab settings, with an emphasis on

insect mortality ((Lichtenstein et al. 1973; Choung et al. 2013), but see (Finlayson et al. 1975)).

Our study offers a glimpse at the potentially antagonistic effect of glyphosate on spinosad

efficacy: plants treated with glyphosate exhibited marginally higher levels of herbivory damage,

but only in the presence of spinosad. While this result somewhat contrasts with our previous

work showing that glyphosate application results in increased herbivory damage in the absence

of any insecticide (Zhang and Baucom 2024), the discrepancy may be due to a difference in

glyphosate dosage between experiments. Nonetheless, our finding here is largely in line with

other studies demonstrating an increase of herbivores on herbicide-treated plants (Dewar et al.

2000; Johnson and Baucom 2022), and suggests that spinosad may mediate this phenomenon.

Given that glyphosate can alter herbivory damage patterns (*e.g.*, hole-feeding vs. margin-

feeding) in this system (Zhang and Baucom 2024), further work is necessary to determine

whether spinosad may affect the community composition of insect herbivores.

In conclusion, our study examined how plant resistance evolves in different environmental

contexts as mediated by glyphosate and spinosad, showing that spinosad modified the expression

and genetic variation of glyphosate resistance, as well as the strength of selection acting on the

trait. By increasing glyphosate resistance, spinosad indirectly improves the fitness of exposed

plants, thereby influencing their evolutionary trajectory and demonstrating the context-

dependency of resistance evolution. As plants in nature rarely face stressors in isolation, it is

important to examine the effects of multiple stressors on resistance traits to predict how plants

may adapt in real time. Using factorial experiments to evaluate fitness effects across multiple

environmental contexts and stressors offers a powerful way to understand how plants respond to

shifting, multifaceted environments.

LITERATURE CITED

Amrhein, N., Brigitte Deus, Peter Gehrke, and Hans Christian Steinrücken. 1980. The site of the

inhibition of the shikimate pathway by glyphosate: II. Interference of glyphosate with

chorismate formation *in vivo* and *in vitro*. Plant Physiol. 66:830–834. American Society of

Plant Biologists (ASPB).

Baek, Y., L. K. Bobadilla, D. A. Giacomini, J. S. Montgomery, B. P. Murphy, and P. J. Tranel.

2021. Evolution of glyphosate-resistant weeds. Rev. Environ. Contam. Toxicol. 255:93–

128. Springer International Publishing.

Bastarache, P., R. Bouafoura, E. Omakele, C. E. Moffat, J. L. Vickruck, and P. Morin Jr. 2023.

Spinosad-associated modulation of select cytochrome P450s and glutathione S-transferases

in the Colorado potato beetle, *Leptinotarsa decemlineata*. Arch. Insect Biochem. Physiol.

112:e21993. Wiley.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using

lme4. J. Stat. Softw. 67:1–48.

Baucom, R. S. 2019. Evolutionary and ecological insights from herbicide-resistant weeds: what

have we learned about plant adaptation, and what is left to uncover? New Phytol. 223:68–

82.

Baucom, R. S., and R. Mauricio. 2008. Constraints on the evolution of tolerance to herbicide in

the common morning glory: resistance and tolerance are mutually exclusive. Evolution

62:2842–2854.

Baucom, R. S., and R. Mauricio. 2004. Fitness costs and benefits of novel herbicide tolerance in

a noxious weed. Proc. Natl. Acad. Sci. U. S. A. 101:13386–13390.

Bergelson, J., and C. B. Purrington. 1996. Surveying patterns in the cost of resistance in plants.

Am. Nat. 148:536–558. [University of Chicago Press, American Society of Naturalists].

Bunch, T. R., C. Bond, K. Buhl, and D. Stone. 2014. *Spinosad General Fact Sheet*.

Chaney, L., and R. S. Baucom. 2014. The costs and benefits of tolerance to competition in

Ipomoea purpurea, the common morning glory. Evolution 68:1698–1709.

Choung, C. B., R. V. Hyne, M. M. Stevens, and G. C. Hose. 2013. The ecological effects of a

herbicide-insecticide mixture on an experimental freshwater ecosystem. Environ. Pollut.

172:264–274. Elsevier BV.

Cipollini, D., D. Walters, and C. Voelckel. 2017. Costs of resistance in plants: From theory to

evidence. John Wiley & Sons, Ltd, Chichester, UK.

Côté, I. M., E. S. Darling, and C. J. Brown. 2016. Interactions among ecosystem stressors and

their importance in conservation. Proc. Biol. Sci. 283:20152592. The Royal Society.

Culpepper, A. S. 2006. Glyphosate-Induced Weed Shifts. Weed Technol. 20:277–281.

[Cambridge University Press, Weed Science Society of America].

Daramola, O. S., G. E. MacDonald, R. G. Kanissery, and P. Devkota. 2023. Effects of co-applied

agrochemicals on herbicide performance: A review. Crop Prot. 174:106396. Elsevier BV.

Debban, C. L., S. Okum, K. E. Pieper, A. Wilson, and R. S. Baucom. 2015. An examination of

fitness costs of glyphosate resistance in the common morning glory, *Ipomoea purpurea*.

Ecol. Evol. 5:5284–5294.

Desaint, H., N. Aoun, L. Deslandes, F. Vailleau, F. Roux, and R. Berthomé. 2021. Fight hard or

die trying: when plants face pathogens under heat stress. New Phytol. 229:712–734. Wiley.

Dewar, A. M., L. A. Haylock, K. M. Bean, and M. J. May. 2000. Delayed control of weeds in

glyphosate☐tolerant sugar beet and the consequences on aphid infestation and yield. Pest

Manag. Sci. 56:345–350. Wiley.

Dimunová, D., P. Matoušková, R. Podlipná, I. Boušová, and L. Skálová. 2022. The role of UDP-

glycosyltransferases in xenobiotic resistance. Drug Metab. Rev. 54:282–298.

Duke, S. O., J. Lydon, W. C. Koskinen, T. B. Moorman, R. L. Chaney, and R. Hammerschmidt.

2012. Glyphosate effects on plant mineral nutrition, crop rhizosphere microbiota, and plant

disease in glyphosate-resistant crops. J. Agric. Food Chem. 60:10375–10397.

Edwards, R., D. P. Dixon, I. Cummins, M. Brazier-Hicks, and M. Skipsey. 2011. New

perspectives on the metabolism and detoxification of synthetic compounds in plants. Pp.

125–148 in P. Schröder and C. D. Collins, eds. Organic Xenobiotics and Plants: From Mode

of Action to Ecophysiology. Springer Netherlands, Dordrecht.

Finlayson, D. G., C. J. Campbell, and H. A. Roberts. 1975. Herbicides and insecticides: their

compatibility and effects on weeds, insects and earthworms in the minicauliflower crop.

Ann. Appl. Biol. 79:95–108. Wiley.

Fuchs, B., K. Saikkonen, and M. Helander. 2021. Glyphosate-modulated biosynthesis driving

plant defense and species interactions. Trends Plant Sci. 26:312–323.

Gassmann, A. J., and D. J. Futuyma. 2005. Consequence of herbivory for the fitness cost of

herbicide resistance: photosynthetic variation in the context of plant-herbivore interactions.

J. Evol. Biol. 18:447–454.

Getman-Pickering, Z. L., A. Campbell, N. Aflitto, A. Grele, J. K. Davis, and T. A. Ugine. 2020.

LeafByte: A mobile application that measures leaf area and herbivory quickly and

accurately. Methods Ecol. Evol. 11:215–221. Wiley.

Gharabli, H., V. Della Gala, and D. H. Welner. 2023. The function of UDP-glycosyltransferases

in plants and their possible use in crop protection. Biotechnol. Adv. 67:108182.

Gupta, S., A. Harkess, A. Soble, M. Van Etten, J. Leebens-Mack, and R. S. Baucom. 2023.

Interchromosomal linkage disequilibrium and linked fitness cost loci associated with

selection for herbicide resistance. New Phytol. 238:1263–1277.

Johnson, N. M., and R. S. Baucom. 2022. Dicamba drift alters plant–herbivore interactions at the

agro☐ecological interface. Ecosphere 13:e4274. Wiley.

Koricheva, J. 2002. Meta-analysis of sources of variation in fitness costs of plant antiherbivore

defenses. Ecology 83:176–190. Wiley.

Kreiner, J. M., S. M. Latorre, H. A. Burbano, J. R. Stinchcombe, S. P. Otto, D. Weigel, and S. I.

Wright. 2022. Rapid weed adaptation and range expansion in response to agriculture over

the past two centuries. Science 378:1079–1085.

Kuester, A., S.-M. Chang, and R. S. Baucom. 2015. The geographic mosaic of herbicide

resistance evolution in the common morning glory, *Ipomoea purpurea*: Evidence for

resistance hotspots and low genetic differentiation across the landscape. Evol. Appl. 8:821–

833.

Kwon, C. S., and D. Penner. 1995. The interaction of insecticides with herbicide activity. Weed

Technol. 9:119–124. [Cambridge University Press, Weed Science Society of America].

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.

Evolution 37:1210–1226. [Society for the Study of Evolution, Wiley].

Lefcheck, J. S. 2016. piecewiseSEM : Piecewise structural equation modelling in r for ecology,

evolution, and systematics. Methods Ecol. Evol. 7:573–579. Wiley.

Lichtenstein, E. P., T. T. Liang, and B. N. Anderegg. 1973. Synergism of insecticides by

herbicides. Science 181:847–849. American Association for the Advancement of Science

(AAAS).

Mauricio, R., and L. E. Mojonnier. 1997. Reducing bias in the measurement of selection. Trends

Ecol. Evol. 12:433–436.

Mertz, F. P., and R. C. Yao. 1990. *Saccharopolyspora spinosa* sp. nov. isolated from soil

collected in a sugar mill rum still. Int. J. Syst. Bacteriol. 40:34–39. Microbiology Society.

Mithöfer, A., and W. Boland. 2012. Plant defense against herbivores: chemical aspects. Annu.

Rev. Plant Biol. 63:431–450.

Monsanto. 2017. Roundup Weathermax Complete Directions for Use.

Moretti, E. A., A. G. Taylor, K. Wickings, and B. A. Nault. 2021. Insights into how spinosad

seed treatment protects onion from onion maggot (Diptera: Anthomyiidae). J. Econ.

Entomol. 114:694–701. Oxford University Press (OUP).

National Organic Standards Board Technical Advisory Panel. 2002. Spinosad. USDA.

Neve, P., R. Busi, M. Renton, and M. M. Vila-Aiub. 2014. Expanding the eco-evolutionary

context of herbicide resistance research. Pest Manag. Sci. 70:1385–1393.

Nychka, D., R. Furrer, J. Paige, and S. Sain. 2021. fields: Tools for spatial data. University

Corporation for Atmospheric Research, Boulder, CO, USA.

Pankey, J. H., J. L. Griffin, B. R. Leonard, D. K. Miller, R. G. Downer, and R. W. Costello.

2004. Glyphosate–insecticide combination effects on weed and insect control in cotton.

Weed Technol. 18:698–703. Cambridge University Press (CUP).

Peterson, R. 2021. Finding optimal normalizing transformations via bestNormalize. R J. 13:310.

708  The R Foundation.

709 Rausher, M. D. 1992. The measurement of selection on quantitative traits: biases due to

710  environmental covariances between traits and fitness. Evolution 46:616–626. [Society for

711  the Study of Evolution, Wiley].

712 Rausher, M. D., and E. L. Simms. 1989. The evolution of resistance to herbivory in *Ipomoea*

713  *purpurea*. I. Attempts to detect selection. Evolution 43:563–572.

714 R Core Team. 2023. R: A Language and Environment for Statistical Computing. R Foundation

715  for Statistical Computing, Vienna, Austria.

716 Scroggs, D. M., D. K. Miller, J. L. Griffin, J. Geaghan, P. Vidrine, and A. M. Stewart. 2005.

717  Glyphosate efficacy on selected weed species is unaffected by chemical Coapplication.

718  Weed Technol. 19:1012–1016. [Cambridge University Press, Weed Science Society of

719  America].

720 Siemens, D. H., H. Lischke, N. Maggiulli, S. Schürch, and B. A. Roy. 2003. Cost of resistance

721  and tolerance under competition: the defense-stress benefit hypothesis. Evolutionary

722  Ecology 17:247–263. Springer.

723 Simms, E. L., and M. D. Rausher. 1987. Costs and benefits of plant resistance to herbivory. Am.

724  Nat. 130:570–581. [University of Chicago Press, American Society of Naturalists].

725 Simms, E. L., and M. D. Rausher. 1993. Patterns of selection on phytophage resistance in

726  *Ipomoea purpurea*. Evolution 47:970–976. [Society for the Study of Evolution, Wiley].

727 Simms, E. L., and M. D. Rausher. 1989. The evolution of resistance to herbivory in *Ipomoea*

728  *purpurea*. II. Natural selection by insects and costs of resistance. Evolution 43:573–585.

729 Sparks, O. C., J. L. Barrentine, N. R. Burgos, and M. R. McClelland. 2004. The effect of

730  glyphosate and insecticide tank mixtures on cotton bollworm (*Helicoverpa zea*) and

selected weed species. Pp. 159–166 *in* D. M. Oosterhuis, ed. Summaries of Arkansas Cotton

Research 2003. University of Arkansas, Fayetteville, Fayetteville, AR.

Sparks, T. C., J. E. Dripps, G. B. Watson, and D. Paroonagian. 2012. Resistance and cross-

resistance to the spinosyns – A review and analysis. Pestic. Biochem. Physiol. 102:1–10.

Elsevier BV.

Stinchcombe, J. R., M. T. Rutter, D. S. Burdick, P. Tiffin, M. D. Rausher, and R. Mauricio.

2002. Testing for environmentally induced bias in phenotypic estimates of natural selection:

theory and practice. Am. Nat. 160:511–523.

Strauss, S. Y., D. H. Siemens, M. B. Decher, and T. Mitchell-Olds. 1999. Ecological costs of

plant resistance to herbivores in the currency of pollination. Evolution 53:1105–1113.

Oxford University Press (OUP).

Svensson, E. I., S. J. Arnold, R. Bürger, K. Csilléry, J. Draghi, J. M. Henshaw, A. G. Jones, S.

De Lisle, D. A. Marques, K. McGuigan, M. N. Simon, and A. Runemark. 2021.

Correlational selection in the age of genomics. Nat. Ecol. Evol. 5:562–573. Springer

Science and Business Media LLC.

Tian, D., M. B. Traw, J. Q. Chen, M. Kreitman, and J. Bergelson. 2003. Fitness costs of R-gene-

mediated resistance in *Arabidopsis thaliana*. Nature 423:74–77. Springer Science and

Business Media LLC.

Van Etten, M. L., A. Kuester, S.-M. Chang, and R. S. Baucom. 2016. Fitness costs of herbicide

resistance across natural populations of the common morning glory, I*pomoea purpurea*.

Evolution 70:2199–2210.

Van Leeuwen, T., W. Dermauw, M. van de Veire, and L. Tirry. 2005. Systemic use of spinosad

to control the two-spotted spider mite (Acari: Tetranychidae) on tomatoes grown in

rockwool. Exp. Appl. Acarol. 37:93–105. Springer Science and Business Media LLC.

Vázquez, M. B., M. V. Moreno, M. R. Amodeo, and M. V. Bianchinotti. 2021. Effects of

glyphosate on soil fungal communities: A field study. Rev. Argent. Microbiol. 53:349–358.

Elsevier BV.

Wang, J., Y. Wan, Y. Zhang, J. Yuan, X. Zheng, H. Cao, K. Qian, J. Feng, Y. Tang, S. Chen, Y.

Zhang, X. Zhou, P. Liang, and Q. Wu. 2024. Uridine diphosphate glucosyltransferases are

involved in spinosad resistance in western flower thrips *Frankliniella occidentalis*

(Pergande). J. Hazard. Mater. 466:133575. Elsevier BV.

Warwick, S. 1991. Herbicide resistance in weedy plants: Physiology and population biology.

Annual Review of Ecology, Evolution, and Systematics 22:95–114. Annual Reviews.

Wise, M. J. 2023. Why fitness impacts of different herbivores may combine nonadditively, and

why it matters to the ecology and evolution of plant-herbivore communities. Plant Ecol.

Evol. 156:13–28. Agentschap Plantentuin Meise.

Wise, M. J., and M. D. Rausher. 2016. Costs of resistance and correlational selection in the

multiple-herbivore community of Solanum carolinense. Evolution 70:2411–2420.

Zandalinas, S. I., M. Á. Peláez-Vico, R. Sinha, L. S. Pascual, and R. Mittler. 2024. The impact of

multifactorial stress combination on plants, crops, and ecosystems: how should we prepare

for what comes next? Plant J. 117:1800–1814. Wiley.

Zhang, G. M., and R. S. Baucom. 2024. Herbicidal interference: glyphosate drives both the

ecology and evolution of plant-herbivore interactions. New Phytol. 245:807–817. John

Wiley & Sons, Ltd.

Zhang, Y., X. Liu, T. Yin, Q. Li, Q. Zou, K. Huang, D. Guo, and X. Zhang. 2021. Comparative

transcriptomic analysis of two *Saccharopolyspora spinosa* strains reveals the relationships

between primary metabolism and spinosad production. Sci. Rep. 11:14779. Springer

Science and Business Media LLC.

FIGURES AND TABLES

Figure 1. Factorial design of the common garden study, showing predictions of the fitness costs and benefits of glyphosate resistance (gold lines) in the form of selection analyses in the four treatment environments: glyphosate-present, spinosad-absent (green box), glyphosate-absent, spinosad-absent (blue box), glyphosate-present, spinosad-present (yellow box), and glyphosate-absent, spinosad-present (white box). Based on past studies, there is a fitness benefit (green box), but not a fitness cost (blue box) in terms of seed production. We examine whether the addition of spinosad, an agriculturally relevant and commonly used insecticide, alters those selection patterns (as represented by the dashed lines). The benefit of glyphosate resistance in the presence of glyphosate (yellow box) may be enhanced by spinosad (a), unaffected (b), or dampened (c).

Likewise, the cost of glyphosate resistance in the absence of glyphosate (white box) may be outweighed by the benefit (d), remain undetectable (e), or manifest (f). Spinosad as a secondary stressor may potentially interact with glyphosate to shape plant evolutionary trajectories.

Figure 2. Treatment effects of glyphosate and spinosad application. Circles show unadjusted treatment means with standard error bars. (a) Glyphosate treatment did not have a significant effect on the amount of herbivory damage overall ($F_1 = 2.14$, $p = 0.144$), nor did it impact herbivory in the absence of spinosad ($F_1 = 0.21$, $p = 0.648$). In the presence of spinosad, however, plants treated with glyphosate showed marginally more herbivory damage in comparison to untreated plants ($F_1 = 3.18$, $p = 0.075$). (b) Spinosad-treated plants had higher levels of glyphosate resistance ($F_1 = 71.83$, $p < 0.001$). (c) Spinosad-treated plants experienced less herbivory damage ($F_1 = 174.06$, $p < 0.001$).

Figure 3. Family means of fitness in relation to resistance to glyphosate (orange) and herbivory (blue) in the four treatment environments. Each point is a family-line mean for that particular trait. (a) In the glyphosate-present, spinosad-absent environment, there was positive selection for glyphosate resistance ($\beta = 0.46$, $F_1 = 20.21$, $p < 0.001$) as denoted by the solid line. (b) In the glyphosate-absent, spinosad-absent environment, there was no selection acting on either glyphosate or herbivory resistance. (c) In the glyphosate-present, spinosad-present environment, there was marginal positive selection for glyphosate resistance ($\beta = 0.18$, $F_1 = 3.28$, $p = 0.077$) as denoted by the dashed line. (d) In the glyphosate-absent, spinosad-present environment, there was no selection acting on either form of resistance.

Figure 4. In the glyphosate-present, spinosad-absent environment (*i.e.*, herbivores are present), there is correlational selection that favors high levels of both glyphosate and herbivory resistance (correlational selection gradient = 0.04, F_1 = 3.99, p = 0.053). The colored bar on the side shows relative fitness, with higher values being warmer and lower values being cooler.

Figure 5. Structural equation model (SEM) of the relationships between glyphosate treatment, spinosad treatment, glyphosate resistance, herbivory resistance, and seed production. All models in the SEM are linear mixed models, with family line and block included as random effects. Boxes represent the measured variables and arrows show the unidirectional relationships between variables. Gray arrows indicate positive relationships and red arrows represent negative relationships. The thickness of the arrows reflect the magnitude of the standardized β coefficients, which are shown along the arrows. All solid arrows show statistically significant paths (p < 0.001), while dotted arrows show nonsignificant paths. Conditional R^2 values are included within the box of each response variable. This model fit well, with a Fisher's C score of 0.007 (p = 0.996).

Table 1. Selection analyses for resistance to glyphosate and herbivory in each of the four treatment environments. Models for each type of resistance were conducted separately in each environment by regressing relative fitness on the standardized form of each resistance. Bolded p-values denote statistical significance, while italicized p-values show marginal statistical significance.

relative fitness ~ standardized resistance

Trait	β	SE	F	df	p
glyphosate-present, spinosad-absent					
glyphosate resistance	0.46	0.10	20.21	1	**<0.001**
herbivory resistance	0.13	0.12	1.10	1	0.299
glyphosate-absent, spinosad-absent					
glyphosate resistance	0.06	0.03	2.68	1	0.109
herbivory resistance	-0.04	0.03	1.14	1	0.292
glyphosate-present, spinosad-present					
glyphosate resistance	0.18	0.10	3.28	1	*0.077*
herbivory resistance	0.03	0.10	0.07	1	0.797
glyphosate-absent, spinosad-present					
glyphosate resistance	0.01	0.03	0.03	1	0.858
herbivory resistance	<0.001	0.06	0.00	1	0.996