Longitudinal Neurocognitive Trajectories in a Large Cohort of Youth Who Use Cannabis:
Combining Self-Report and Toxicology

Natasha E. Wade[1], Ryan M. Sullivan[1], Alexander L. Wallace[1], Rachel Visontay[2], Veronica Szpak[1], Krista M. Lisdahl[3], Marilyn A. Huestis[4], Priscila Dib Gonçalves[5], Hollie Byrne[2], Louise Mewton[2], Joanna Jacobus[1], Susan F. Tapert[1]

[1] Department of Psychiatry, University of California, San Diego, La Jolla CA
[2] The Matilda Centre for Research in Mental Health and Substance Use, The University of Sydney, NSW, Australia
[3] Department of Psychological & Brain Sciences, University of Wisconsin-Milwaukee
[4] Huestis & Smith Toxicology, LLC, Fredericksburg, VA
[5] Department of Psychiatry, Columbia University

*Correspondence and reprint requests to:
Natasha E. Wade, Ph.D.
Assistant Professor, Department of Psychiatry
University of California, San Diego
9500 Gilman Drive, MC 0405
La Jolla, CA 92093
email: nwade@ucsd.edu

Abstract

Adolescents experience extensive neurocognitive development, which cannabis use potentially impacting developmental trajectories. Here, we comprehensively assess the influence of adolescent cannabis use onset on neurocognitive trajectories and consider how recent delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) may influence neurocognition. We use the large, diverse longitudinal Adolescent Brain Cognitive Development (ABCD) Study dataset, combining self-reported substance use with objective toxicological tests (hair, urine, breath, oral fluid). Longitudinal mixed methods of the full cohort (n=11,036, ages 9-17) investigate time-varying cannabis onset on neurocognitive performance. Secondarily, in participants with repeat toxicological hair testing (n=645) at ages 12-16, we consider the influence of THC v. CBD v. Controls. Covariates include sociodemographics, family history of substance use disorder, prenatal substance exposure, early psychopathology, other substance use, and nesting for participant ID, study site, and family ID. Cannabis group interacted with age to show altered neurocognitive trajectories across domains (immediate recall and delayed memory, processing speed, inhibitory control, visuospatial processing, language, and working memory; βs=-0.11 - -0.52). Secondary models indicated hair-identified THC exposure predicted worse episodic memory than in Controls (β=-0.60, *p*=.007), with no difference between CBD exposed and Controls. Data suggest those who use cannabis show a slight cognitive advantage during late childhood, with demonstrate diminished trajectories over time. These flattened neurocognitive trajectories in youth (ages 9-17) who initiate cannabis use were demonstrated after accounting for within-person change and numerous known confounds and improving accuracy in cannabis groupings through incorporating toxicological measures. Continued monitoring of this cohort will clarify cannabinoid-cognition relationships into young adulthood, including the impact of timing of cannabis use initiation.

Introduction

Over the past several decades, high rates of cannabis use in individuals ages 13-24 remained relatively stable[1], but the legal status of cannabis changed in many US states, the potency of the primary psychoactive cannabinoid delta-9-tetrahydrocannabinol (THC) increased[2], vaped inhalation increased, and public perception of risk decreased[3]. Critically, adolescents experience extensive brain and cognitive development[4-6], making adolescents particularly vulnerable to the influence of cannabis[7]. This includes within the endocannabinoid system, a neuromodulatory system critical to brain development (e.g., neural progenitor proliferation, neurogenesis)[7,8] and to which THC binds. Early longitudinal studies suggest neurocognitive deficits in select domains are associated with adolescent cannabis onset and escalation, such as poorer verbal learning and memory[9-15], visuospatial functioning[10,11,13,15], working memory[10,11,15,16], inhibition and attention[11,14,15], and psychomotor and processing speed[9,11,13,14]. However, these studies often do not fully account for important pre-substance use confounds (e.g., other substance use, family history, family factors, sociodemographics), include small sample sizes, or do not clearly identify the onset of cannabis use[16-18].

Studies of adolescent cannabis use can also be limited by the accuracy of self-report, the primary method of identifying cannabis use in most studies. Recent studies comparing objective, toxicological measures with self-report in adolescents and young adults underestimated substance use prevalence by up to 60%[19], or mis-identified up to 9% of participants as "controls" when they actually used substances[20]. Therefore, research on the sensitive topic of substance use in teens may at times present with bias in self-report, which impedes identifying substance use consequences. Incorporating toxicological measures, including hair cannabinoids, may improve accuracy in characterizing cannabis-neurocognition relationships[21,22].

Accurate assessment of cannabis use histories is further complicated by the complexity of the plant that contains over 120 cannabinoids[23], with the most consistent evidence demonstrating repeated THC exposure is linked with changes in CB1 density and downstream cognitive functioning[24]. Adult evidence demonstrates that cannabidiol (CBD), a CB1 cannabinoid receptor allosteric modulator, may offer mitigating influence on cognitive performance[25,26]. Accordingly, objective evidence of cannabinoid analyte concentration exposures may be more informative than relying on self-reported cannabis use episodes or potency, especially given individual differences in processing of cannabis due to genetics, rates of exposure, and other biological mechanisms[27-30], in addition to inconsistencies in self-reported use[20]. The literature on specific cannabinoid constituents as related to cognitive functioning in the developing brain is exceptionally limited[31,32], yet different cannabinoids may moderate clinical outcomes. Thus, combining toxicological cannabinoid information with self-reported use may more accurately and thoroughly assess the impact of cannabis use onset on cognition and further investigate the influence of specific cannabinoid constituents (i.e., THC, CBD, the most studied cannabinoids) as potential explanatory mechanisms.

Here, we comprehensively assess the influence of adolescent cannabis use onset on neurocognitive trajectories. We probe potential mechanisms by investigating specific cannabinoid content through toxicological hair testing to examine whether use of certain cannabinoids may predict differential trajectories. Leveraging data from the Adolescent Brain Cognitive Development (ABCD) Study – a large and richly-characterized, multisite US cohort— we first identify those who initiated cannabis use through comprehensive substance use assessment, including self-report and multiple toxicological tests. We then conduct longitudinal mixed effect models to assess trajectories of neurocognitive performance in those who initiated

cannabis use relative to those who did not. Secondarily, follow-up analyses utilize robust cannabinoid content from hair to examine whether recent use of specific cannabinoid constituents (THC v. CBD v. Controls) differentially alter neurocognitive trajectories. Importantly, models covary for core confounds and include within-person nesting to account for individual trajectories and neurocognition prior to cannabis use onset.

Methods

Participants

Participants were recruited at ages 9-10 in 2016-2018 via propensity-based school recruitment at 21 sites across the United States, resulting in a cohort of 11,868[33,34], and followed for 10+ years. ABCD data release 6.0 contains data collected from September 1, 2016 to January 15, 2024, which includes complete annual data from Waves 0-5 and half of Wave 6. **Figure 1** displays sample sizes for the present analyses. Eligible participants and their caregivers presented at their local site for enrollment and a Baseline session, with mid-year brief assessment and annual in-person visits each year thereafter. Annual sessions included biosample collection, self- and parent-report surveys, clinical mental health interview, and a semi-structured substance use interview[35-40]. On even years (Baseline, Wave 2, Wave 4, etc.) participants also completed a full neurocognitive battery. Starting Wave 3, all youth were told to remain substance-free 24 hours prior to the session. Participants and their caregivers completed written informed assent and consent, respectively. All study protocols were completed consistent with the Declaration of Helsinki and approved through a single IRB at the University of California, San Diego.

Figure 1. Longitudinal Sample Size by Neurocognitive Task

Notes: Chart describing sample sizes and number of complete administrations required to be included in analyses. N=Sample size by task; Nobs=number of observations included in the models. * indicates tasks that were included in secondary hair cannabinoid analyses, as these have data available at Waves 2, 4, and 6 (to match waves of available hair data); only data from half the cohort was released for use in the present analyses.

Measures

 Substance Use Measures.

 Self-Reported Substance Use and Timeline Followback Interview. Prior to querying

substance use, research assistants (RAs) reminded participants of confidentiality, including that

caregivers will not be informed of their responses to the substance use questions unless they

report clear, imminent threat or harm or other state-mandated reporting requirements. Participants annually completed a substance use interview covering lifetime at Baseline and the past year (or since last visit) at annual follow-ups. When participants endorsed full standard substance use (1+ standard alcohol drink or any other substance use), RAs administered a Timeline Followback (TLFB) interview to collect detailed substance use patterns since their last visit (approximately past year) by each substance class[37,41]. A mid-year assessment was given at approximately the mid-point between study visits (~6 months) covering substance use since their last visit (binary reported yes/no).

Medicinal CBD Use. Beginning at Wave 2, caregivers were asked, "Has your son/daughter used any CBD (or cannabidiol) products with your or your doctor's permission, such as Epidiolex or other over the counter CBD product, in the past year?". Youth participants and their caregivers were also asked whether they had used CBD within the past 24 hours.

Oral Fluid Toxicology. An oral fluid drug screen was administered to ~5% randomly selected participants at Baseline and increased to all participants at Wave 6. Drug classes tested include amphetamines, benzodiazepines, methamphetamines, cocaine, cannabis (THC), and opioids. Oral fluid tests sensitively identify past 12-48 hours of drug use, depending on the substance[39].

Urine Toxicology. At Waves 4-6, participants completed a full drug urinalysis screening for amphetamines, barbiturates, buprenorphine, benzodiazepines, cannabinoids, cocaine, MDMA, methamphetamine, methadone, opioids, oxycodone, PCP, propoxyphene, tricyclics, and cannabis (THC). Depending on drug class and frequency/amount of drug used, urine can test positive up to a month after last use.

Hair Toxicology. Hair was collected from all participants annually who were willing to give a sample and did not have a hair style that would be disrupted by collection (e.g., too short of hair; braids). Approximately 100mg hair 1.5cm from the root were collected from 3-4 regions around the crown of the participant's head. Hair was securely packaged and stored until selected for testing, based on funding and prioritization of sample selection[20,42], with testing processes and cut-offs described in the Supplement. Hair samples provide on average a 3-month window of detection for substance use[43]. Hair testing for cannabis is sensitive for detecting moderate-to-heavy use[44] (using a minimum of twice a month on average[21]) and demonstrates excellent specificity in adolescents and young adults[21].

Predictors of interest include: (Aim 1) <u>Any Lifetime Cannabis Use</u>, combining all cannabis data (self-report, toxicological) into a binary variable, in which all participants are coded "Controls" until any use is identified via self-reported use exceeding a puff/taste or any positive cannabis toxicology, then subsequently coded "yes" for lifetime cannabis use. Mid-year interview substance use (e.g., Wave 4.5) was applied to the following study visit (e.g., Wave 5). In models, indicators of cannabis use were entered as time-varying predictors with a binary yes/no variable representing cannabis use at the wave with first-identified use and then carried forward for each subsequent wave.

(Aim 2) <u>Hair Cannabinoids</u> coded as "Controls" (no cannabinoids detected in hair), "THC Only" (THC/THCCOOH/Δ8-THC detected only in hair), and "CBD+" (CBD detected in hair, regardless of THC/THCCOOH/Δ8-THC status) to assess the influence of recent, specific cannabinoids on neurocognition, consistent with prior work[26]; in these models, hair cannabinoid groups were time-varying at each wave.

Neurocognitive Outcomes.

Tasks selected for the present analyses include those given on even study years and administered at least three times between Baseline and Wave 6 (see **Figure 1** for task administration time). Analyses were restricted to participants who completed at least two administrations of the same task across waves. For List Sort Working Memory and Rey Auditory Verbal Learning Test (RAVLT), data are only available for less than half of participants at Wave 6 and there were only three possible administrations across time. Therefore, all three administrations were required given it was not possible for a majority of participants to complete the task.

NIH Toolbox[45]. Inhibitory control was assessed via the Flanker Inhibitory Control and Attention task, where participants were to press an arrow key matching the direction a stimulus arrow was pointing. For receptive language (Picture Vocabulary), participants listened to a word and matched it to one of four pictures. For oral reading, participants read and pronounced a list of words. In the List Sort Working Memory, participants ordered by size a list of animals and produce. Pattern Comparison Processing Speed asked participants to quickly identify whether two pictures were identical. Picture Sequencing Memory test assessed episodic memory via showing the participants a series of pictures, then asking them to sort the pictures in the same order as demonstrated. Primary outcomes included uncorrected standard scores for each subtest, consistent with prior studies[46].

Little Man Task[47]. Participants were shown a "little man" holding a briefcase in various orientations to assess visuospatial processing and mental rotation. As the figure shifts (e.g., upside down; faces away from the participant), the participant must select whether the briefcase is in the man's left or right hand. Total correct responses were used as the primary visuospatial performance outcome.

RAVLT[48]. Verbal recall and memory were assessed through the RAVLT. Participants were read a list of words five times, followed by a distractor list, the original list, and then a long delay. Here, <u>immediate verbal recall</u> was assessed with number of correct Trial 1 responses and memory was determined by number of correct responses after the <u>long delay</u>.

Covariates. Covariates included those previously associated with cannabis and behavioral outcomes[49] and variables are described in more detail in the Supplement. Covariate use within models is described in the Statistical Analyses.

<u>Time-Invariant</u>. *Sociodemographic covariates*. Baseline sociodemographic characteristics, as reported by the caregiver, were selected as covariates given their relevance to cognition and substance use: highest level of parental education attainment[50], biological sex of the participant as assigned at birth[51], and race and ethnicity[36] (due to potential differences in rates of hair collection[52]). *Family History of Substance Use.* Caregivers reported whether any of the participant's biological relatives had alcohol use problems (AUD) or, separately, substance use problems (SUD)[53]. *Prenatal Substance Exposure*. Caregivers reported whether the participant was exposed to any drug classes in utero, both before and after knowledge of the pregnancy[36]. Each drug class was coded as its own exposure (1/0, regardless of whether using before/after knowledge of pregnancy; see Supplement for details). *Baseline Psychopathology*. Caregivers completed the Childhood Behavior Checklist (CBCL), with age- and sex-normed t-scores subscales for Externalizing and Internalizing symptoms included as covariates, given youth psychopathology is often identified as a predictor of substance initiation[54].

<u>Time-Varying</u>. Youth age at each Wave was included. *Other Substance Use*. Polysubstance use is common and may influence cognition[55,56]. Using the same identification method of lifetime substance exposure as described for cannabis use, binary (yes/no) time-

varying lifetime alcohol use, nicotine use, and any other substance use variables were computed across all observations.

Analytic Plan

ABCD Data Release 6.0 was used (https://doi.org/10.82525/jy7n-g441). Analyses were conducted in R.

Aim 1: Longitudinal Influence of Cannabis Onset. Linear mixed effect models were run using *lme4* package in R[57]. Neurocognitive task performance over time was predicted by lifetime cannabis use group status (CU group) interacting with age.

Covariates of interest, often time-varying, were included in the full linear mixed effect models: lifetime alcohol use, lifetime nicotine use, lifetime other substance use, sex, and age. Additionally, for variables shown to be linked to cannabis group status and cognition which are more static in nature or potential confounds of no interest (i.e., prenatal exposure, family history of substance use disorder, baseline psychopathology, sociodemographic factors), an inverse propensity score was created to account for covariates while also reducing variable dimensionality (see Supplement for specific variables). To estimate the inverse propensity score variable, logistic regression was run, predicting lifetime cannabis exposure by all listed factors, clustering on site and family ID. The inverse of logistic regression values (their individual probability of exposure) were saved for each participant and included as a covariate in analyses.

Study site, family ID, and subject ID (to account for within-person change) were included as nested effects. FDR corrections were applied within models[58]. Standardized β were calculated using the *MuMin* package[59]. Estimated marginal means by age were extracted via the *emmeans* package[60].

Aim 2: Longitudinal Influence of Hair Cannabinoids. To specifically assess potential recent THC or CBD influence relative to no use, a subsample of the cohort was included in secondary longitudinal analyses. All participants with hair tested at Waves 2, 4, and 6 were included (n=648). Time-varying groups by wave were then defined by hair cannabinoid content: no cannabinoids present (Controls n=546); THC (identified by THC or THCCOOH; THC Only Group n=81); and CBD (identified by confirmed CBD, regardless of whether THC/THCCOOH were present; CBD+ group n=21), consistent with prior research[26]. Notably, n=13 in the CBD+ group also had THC present in their hair sample and n<10 participants with hair cannabinoids were both in the CBD and THC groups on altering years, though descriptively are included within the CBD group (**Table S2**). Mean performance by hair cannabinoid group and age are presented in **Table S3**. Linear mixed effects models were run to predict cognitive performance by hair group*age. Analyses included tasks that were administered at the same waves as selected hair samples. Given limited prior research using novel repeat hair testing results and cognition and due to the smaller cell sizes, use of covariates were constrained to more sensitively assess potential cannabinoid relationships: sex assigned at birth and any other substance use (i.e., alcohol, nicotine, or any other substance combined) positive in hair were included as covariates. Models were nested for family ID and subject ID; study site was included as a fixed effect due to convergence issues. Interpretation of significance was set at $p<.01$.

Results

Participant Sociodemographic and Substance Use Characteristics. Participants included the full ABCD cohort (see **Table 1** for descriptives). Participants who used cannabis by the Wave 6 follow-up differed from those who did not in sex ($\chi^2=7.3$, $p=.007$), race ($\chi^2=62.9$, $p<.0001$), and parental educational attainment ($\chi^2=147.3$, $p<.0001$). Onset of cannabis use was

most commonly identified by self-report (67%), followed by hair testing (22%), urine toxicology

(21%), oral fluid testing (5%), and self-/parent-reported medicinal CBD use (3%; percentages

not mutually exclusive).

Table 1. Sociodemographics of groups

Characteristic	Category	Controls N=9,664	Cannabis Users N=2,204	
Sex				*
	Female	47%	51%	
Ethnicity				
	Hispanic	20%	22%	
Race				*
	White	67%	53%	
	Black	16%	19%	
	Asian	3%	<1%	
	American Indian/ Alaska Native	<1%	1%	
	Native Hawaiian or Other Pacific Islander	<1%	<1%	
	More than one race	10%	13%	
	Other (unknown or not reported)	4%	4%	
Parental Education				*
	<High School	5%	7%	
	High School Diploma/GED	9%	12%	
	Some College	24%	33%	
	Bachelor's	26%	22%	
	Graduate	36%	26%	

Notes: Data reflects Baseline sociodemographic characteristics of participants; * indicates p<.05
between group differences.

For secondary models, hair groups (THC, CBD, or Controls) differed by sex (χ^2=12.4,

p=.002), parental education (χ^2=64.8, p<.0001), race (χ^2=46.9, p<.0001), and ethnicity (χ^2=13.1,

p=.001), with descriptives included in **Table S4**.

Aim 1: Primary Neurocognitive Models.

As aims were to investigate longitudinal trajectories of cognition, only the interaction between CU status and age are reported, though main effects were found across tasks (FDR-ps<.05). Mean performances by group and age are in **Table S2**.

Working Memory: Working memory performance exhibited a significant CU*age interaction (β=-0.52, p<.0001, FDR-p<.0001), such that CU participants did not demonstrate the same improvement in working memory with age as Controls. Review of estimated marginal means indicates groups differed from ages 9-14 with CU group demonstrating better performance (ps<.0001), whereas at 15-16 groups did not differ (**Figure 2a**). At age 17, CU exhibited toward worse performance (p=.0001).

Inhibitory Control: Inhibitory control exhibited a significant CU*age interaction (β=-0.21, p<.0001, FDR-p<.0001; **Figure 2b**). The CU group performed better at ages 9-10 (ps<.01) and 11 (p=.03), with the relationship flipping starting at age 14 (p=.03) through 15-17 (ps<.0001).

Episodic Memory: CU group interacted with age (β=-0.32, p<.0001, FDR-p<.0001; **Figure 2c**) in predicting episodic memory. At ages 9-13, CU demonstrated better episodic memory performance (ps<.01), no difference at ages 13-14, and worse performance than Controls at ages 15-17 (ps<.001).

Receptive Language: There was an interaction of CU*age in receptive language (β=-0.14, p<.0001, FDR-p<.0001), with the CU group demonstrating less change over time (**Figure 2d**). At ages 9-10, CU demonstrated better performance (ps<.01), with worse performance at ages 14 (p=.03) and 15+ (ps<.0001).

Figure 2. Cannabis Group Status *Age on Neurocognitive Performance

Notes: Cannabis use group status interactions with age by neurocognitive domain; a-e y-axes reflect uncorrected NIH Toolbox performance by domain [working memory n=4,371; picture memory n=11,021; vocabulary n=11,036; processing speed n=10,091]; f is the Little Man Task number correct scores [n=10,941]; g and h are Rey Auditory Verbal Learning Test (RAVLT) immediate recall (Trial 1) and memory (delayed recall) raw scores, respectively [n=4,487]. Total sum N of those who used cannabis are in blue (n=9,664); those who did not use cannabis are in red (n=2,204). Groups were time-varying with onset identified over time.

Processing Speed: The interaction of CU*age (β=-0.24, *p*<.0001, FDR-*p*<.0001) significantly predicted processing speed, with Controls demonstrating greater improvement with age (**Figure 2e**). At ages 9-12, CU demonstrated better processing speed (*p*s<.05), with no difference ages 13-14, and CU demonstrating worse performance than Controls at ages 15-17 (*p*s<.0001).

Oral Reading: For oral reading, there was an interaction of CU*age (β=-0.11, *p*=.0001, FDR-*p*=.0003; **Figure 2f**). CU group performed at ages 9-11 (*p*s<.05) and worse than Controls at ages 15-17 (*p*s<.01).

Visuospatial: For predicting number correct on visuospatial performance, site was included as a fixed effect due to convergence issues. Results indicate a significant interaction between CU*age (β=-0.16, *p*=.001, FDR-*p*=.003), such that those in the CU group demonstrated lower number correct with age (**Figure 2g**). While CU exhibited more correct responses than Controls at age 9 (*p*=.04) and no difference at ages 10-13, Controls had more correct responses at ages 15-17 (*p*<.0001).

Verbal Recall and Memory: CU group*age significantly interacted to predict immediate recall (β=-0.21, *p*=.02, FDR-*p*=.03; **Figure 2h**). Groups did not differ from ages 9-14, though in ages 15-17 the CU group demonstrated worse performance than Controls (*p*s<.05). For long delay memory, there was also a significant interaction between CU*age (β=-0.21, *p*=.008, FDR-

p=.01; **Figure 2i**), such that those in the CU group demonstrated less improvement over time. Groups differed from ages 9-11 with CU group demonstrating better performance (ps<.05), whereas at 12-16 there was no difference between groups, and at age 17 CU demonstrated worse performance (p=.02).

Aim 2: Secondary Hair Analyses

After controlling for sex, other positive hair toxicology results, and site, and nesting for family ID and subject ID, THC Only group status significantly predicted <u>episodic memory</u> performance relative to Controls (β=0.55, p=.007; n=645, $n_{observations}$=1,885). There was also a THC*age interaction relative to Controls (β=-0.60, p=.007), such that those in the THC Only group demonstrated worse performance with age (**Figure 3**). Estimated marginal means revealed that at ages 15-17, THC demonstrated worse episodic memory than Controls (ps=.04-.001) and worse performance than CBD+ (ps=.02-008). CBD+ group and Controls were not significantly different. For all other neurocognitive tasks (receptive language, processing speed, inhibitory control, and oral reading), no significant relationships were found between hair cannabinoid groups and performance.

Figure 3. Hair Cannabinoid Group*Age on Picture Memory

Figure 3. Hair cannabinoid use group identified by whether there was THC (without CBD) in hair, CBD in hair (regardless of THC), or no hair cannabinoids. Hair testing and episodic memory (NIH Toolbox Picture Memory) testing occurred every other year (approximately ages 11-12, 13-14, and 15-16). Total sum N of those without hair cannabinoids (Controls) are in red (n=546); those with CBD in hair are in green (n=21); those with THC are in blue (n=81). Groups were time-varying with onset identified over time.

Discussion

Findings support and extend prior work suggesting aberrant neurocognitive developmental trajectories in youth ages 9-17 who use cannabis, significantly extending the existing literature by including a comprehensive assessment of self-report and objective toxicological measures to accurately categorize cannabis use onset. Utilizing a prospective longitudinal design that considered pre-substance use confounding factors (i.e., sociodemographics, prenatal substance use history, family history of AUD/SUD, internalizing and externalizing symptoms) and controlled for comorbid substance use onset, we found that those who use cannabis actually show a slight cognitive advantage during late childhood, but demonstrate diminished trajectories in immediate recall and delayed memory, processing speed, inhibitory control, visuospatial processing, language, and working memory over adolescence.

Follow-up analyses considered a potential mechanism of altered trajectories through unique contributions of THC and CBD in analyzed in objective hair toxicology. Those with THC use exhibited declining trajectories in episodic memory relative to those without hair cannabinoids, whereas those with CBD did not differ from controls.

In the present study the cannabis use group demonstrated either restricted improvement or plateaued performance relative to non-using youth across cognitive domains. Longitudinal studies previously identified changes in those who initiated cannabis use in learning and memory[9-15], working memory[10,11,15,16], inhibition and attention[11,14,15] visuospatial functioning[10,11,13,15], and processing speed[9,11,13,14]. Here, those who use cannabis exhibited potential neurocognitive advantage in earlier years with a diminished positive slope over time. This is despite the fact that youth (including within the ABCD Study) are expected to show improvement on tasks over time[46]. Other prior work within this cohort describes higher cognition being associated with substance use onset and cannabis specifically[61,62]; it may be that early maturation increases risk of using, and that these youth then demonstrate a diminished cognitive trajectory. Following this cohort into young adulthood can determine how variations in use trajectories, including escalation of use and cessation, influence cognition over time.

Secondary analyses indicate that youth who have objective evidence of recent THC, but no CBD, exposure in hair samples are likely to demonstrate worse memory performance over time than those with no cannabinoids in their hair. Findings align with prior hair cannabinoid results in adults[26], and are generally consistent with cross-sectional analyses of CBD hair cannabinoids in this cohort that did not reveal CBD-cognition relationships[22]. These data indicate a potential mechanism (THC) driving the displayed neurocognitive trajectories observed in the larger CU analyses, consistent with theorized processes[24,63]. THC, then, is likely particularly

important for otherwise healthy adolescents to avoid or delay use. On balance, other data within this cohort indicated that even those who use CBD for medicinal reasons may be unintentionally exposed to THC[64], and product mislabeling is a known concern even in regulated markets[65-68]. Therefore, while CBD may be considered a harm reduction option for those regularly using cannabis in the context of mitigating THC-related consequences or cessation, the exact impact of CBD and the potential for contamination of CBD (including with THC) remain significant concerns for adolescent health.

Small-to-medium effect sizes were observed, comparable to prior research[69]. Importantly, given the sensitive sociodevelopmental period of adolescence, even modest effects may translate to clinically significant alterations[70,71]. For instance, relative decrements in learning and memory can impact standardized testing, grade advancement, and higher education opportunities in a highly competitive contemporary environment. Similarly, visuospatial, processing speed, and working memory skills are paramount to driving, an area where teens may already be at a disadvantage[72,73]. Taken together, the impact of these altered neurocognitive trajectories may have substantial potential in individual lives, particularly when individuals initiate use before age 16[74], though further investigation to the real-world impact of such cannabis use is needed.

Findings are considered in light of strengths and limitations. Results here expand on the prior literature through several notable methodological improvements: (1) use of subjective (self-report) and objective (toxicological) cannabis use variables; (2) wide consideration of potential confounds, including prenatal history, family factors, sociodemographics, other substance use, and within-person change; and (3) a large cohort that was almost exclusively substance naïve at Baseline (ages 9-10)[75], with neurocognitive measures that predate cannabis exposure. On balance, secondary analyses of THC/CBD consisted of a smaller subsample, including a

relatively small number of those with CBD in their hair. Oral fluid and urinalysis tests were not confirmed, though both demonstrate adequate specificity[21,30]; inclusion of toxicological measures here are therefore more likely to contribute to more accurate grouping rather than mis-labeling someone as using cannabis when they do not. Participants included here are largely in the middle of adolescence and will still undergo more neurocognitive development; early cannabis use (e.g., before age 16) is known to be particularly risky for adolescents[74]. Additional follow-up to consider trajectories of those who initiate cannabis use or continue to use in later adolescence are needed.

Conclusions

Adolescence marks a period of substantial development, but cannabis use may constrain one's full cognitive developmental performance. Data here indicate novel flattened neurocognitive trajectories in youth (ages 9-17) who initiate cannabis use, after accounting for within-person change and numerous known confounds and improving accuracy in cannabis groupings through incorporating toxicological measures. Unique analyses harnessing objective toxicology demonstrate that THC use may specifically relate to episodic memory performance over time, and CBD was not linked with cognition during this developmental period. Prevention efforts might inform adolescents of the risk of early cannabis use on cognitive performance. Continued monitoring of this cohort will clarify cannabinoid-cognition relationships into young adulthood, including the impact of timing of cannabis use initiation.

References

1. Schulenberg JE, Johnston LD, O'Malley PM, Bachman JG, Miech RA, Patrick ME. Monitoring the Future national survey results on drug use, 1975-2017: Volume II, college students and adults ages 19-55. 2018:454.

2. ElSohly MA, Chandra S, Radwan M, Majumdar CG, Church JC. A Comprehensive Review of Cannabis Potency in the United States in the Last Decade. *Biol Psychiatry Cogn Neurosci Neuroimaging.* 2021;6(6):603-606.

3. Nguyen N, Holmes LM, Pravosud V, Cohen BE, Ling PM. Changes in perceived harms of tobacco and cannabis and their correlations with use: A panel study of young adults 2014-2020. *Addict Behav.* 2023;144:107758.

4. Giedd JN, Raznahan A, Alexander-Bloch A, Schmitt E, Gogtay N, Rapoport JL. Child psychiatry branch of the National Institute of Mental Health longitudinal structural magnetic resonance imaging study of human brain development. *Neuropsychopharmacology.* 2015;40(1):43-49.

5. Gogtay N, Giedd JN, Lusk L, et al. Dynamic mapping of human cortical development during childhood through early adulthood. *PNAS.* 2004;101(21):8174-8179.

6. Shaw P, Kabani NJ, Lerch JP, et al. Neurodevelopmental trajectories of the human cerebral cortex. *J Neurosci.* 2008;28(14):3586-3594.

7. Schneider M. Puberty as a highly vulnerable developmental period for the consequences of cannabis exposure. *Addict Biol.* 2008;13(2):253-263.

8. Harkany T, Guzman M, Galve-Roperh I, Berghuis P, Devi LA, Mackie K. The emerging functions of endocannabinoid signaling during CNS development. *Trends Pharmacol Sci.* 2007;28(2):83-92.

9. Nguyen-Louie TT, Castro N, Matt GE, Squeglia LM, Brumback T, Tapert SF. Effects of emerging alcohol and marijuana use behaviors on adolescents' neuropsychological functioning over four years. *J Stud Alcohol Drugs.* 2015;76:738-748.

10. Hanson KL, Medina KL, Padula CB, Tapert SF, Brown SA. Impact of Adolescent Alcohol and Drug Use on Neuropsychological Functioning in Young Adulthood: 10-Year Outcomes. *J Child Adolesc Subst Abuse.* 2011;20(2):135-154.

11. Jacobus J, Squeglia LM, Infante MA, et al. Neuropsychological performance in adolescent marijuana users with co-occurring alcohol use: A three-year longitudinal study. *Neuropsychology.* 2015;29(6):829-843.

12. Becker MP, Collins PF, Schultz A, Urosevic S, Schmaling B, Luciana M. Longitudinal changes in cognition in young adult cannabis users. *J Clin Exp Neuropsychol.* 2018;40(6):529-543.

13. Meier MH, Caspi A, Ambler A, et al. Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proc Natl Acad Sci U S A.* 2012;109(40):E2657-2664.

14. Meier MH, Caspi A, A RK, et al. Long-Term Cannabis Use and Cognitive Reserves and Hippocampal Volume in Midlife. *Am J Psychiatry.* 2022;179(5):362-374.

15.   Morin JG, Afzali MH, Bourque J, et al. A Population-Based Analysis of the Relationship Between Substance Use and Adolescent Cognitive Development. *Am J Psychiatry.* 2019;176(2):98-106.

16.   Meier MH, Caspi A, Danese A, et al. Associations between adolescent cannabis use and neuropsychological decline: a longitudinal co-twin control study. *Addiction.* 2018;113(2):257-265.

17.   Rogeberg O. Correlations between cannabis use and IQ change in the Dunedin cohort are consistent with confounding from socioeconomic status. *Proc Natl Acad Sci U S A.* 2013;110(11):4251-4254.

18.   Conrod PJ. Cannabis and Brain Health: What Is Next for Developmental Cohort Studies? *Am J Psychiatry.* 2022;179(5):317-318.

19.   Steinhoff A, Shanahan L, Bechtiger L, et al. When Substance Use Is Underreported: Comparing Self-Reports and Hair Toxicology in an Urban Cohort of Young Adults. *J Am Acad Child Adolesc Psychiatry.* 2023;62(7):791-804.

20.   Wade NE, Sullivan RM, Tapert SF, et al. Concordance between substance use self-report and hair analysis in community-based adolescents. *Am J Drug Alcohol Abuse.* 2023;49(1):76-84.

21.   Wade NE, Wallace AL, Baca R, et al. Cannabinoid Profiling Across Toxicology Samples in Adolescents and Young Adults by Route of Administration and in Relation to Depression Symptoms. *Journal of Analytical Toxicology.* 2025:bkaf051.

22.   Wade NE, Wallace AL, Huestis MA, Lisdahl KM, Sullivan RM, Tapert SF. Cannabis use and neurocognitive performance at 13-14 Years-Old: Optimizing assessment with hair toxicology in the Adolescent brain cognitive development (ABCD) study. *Addict Behav.* 2024;150:107930.

23.   ElSohly MA, Radwan MM, Gul W, Chandra S, Galal A. Phytochemistry of Cannabis sativa L. In: Kinghorn AD, Falk H, Gibbons S, Kobayashi Ji, eds. *Phytocannabinoids: Unraveling the Complex Chemistry and Pharmacology of Cannabis sativa.* Cham: Springer International Publishing; 2017:1-36.

24.   Mizrahi R, Watts JJ, Tseng KY. Mechanisms contributing to cognitive deficits in cannabis users. *Neuropharmacology.* 2017;124:84-88.

25.   Colizzi M, Bhattacharyya S. Does Cannabis Composition Matter? Differential Effects of Delta-9-tetrahydrocannabinol and Cannabidiol on Human Cognition. *Curr Addict Rep.* 2017;4(2):62-74.

26.   Morgan CJ, Gardener C, Schafer G, et al. Sub-chronic impact of cannabinoids in street cannabis on cognition, psychotic-like symptoms and psychological well-being. *Psychol Med.* 2012;42(2):391-400.

27.   Hryhorowicz S, Walczak M, Zakerska-Banaszak O, Slomski R, Skrzypczak-Zielinska M. Pharmacogenetics of Cannabinoids. *Eur J Drug Metab Pharmacokinet.* 2018;43(1):1-12.

28.   Stout SM, Cimino NM. Exogenous cannabinoids as substrates, inhibitors, and inducers of human drug metabolizing enzymes: a systematic review. *Drug Metab Rev.* 2014;46(1):86-95.

29.   Sharma P, Murthy P, Bharath MMS. Chemistry, metabolism, and toxicology of cannabis: Clinical implications. *Iran J Psychiatry.* 2012;7(4):149-156.

30. Huestis MA, Smith ML. Cannabinoid Markers in Biological Fluids and Tissues: Revealing Intake. *Trends Mol Med.* 2018;24(2):156-172.
31. Mokrysz C, Freeman TP, Korkki S, Griffiths K, Curran HV. Are adolescents more vulnerable to the harmful effects of cannabis than adults? A placebo-controlled study in human males. *Transl Psychiatry.* 2016;6(11):e961.
32. Lawn W, Trinci K, Mokrysz C, et al. The acute effects of cannabis with and without cannabidiol in adults and adolescents: A randomised, double-blind, placebo-controlled, crossover experiment. *Addiction.* 2023;118(7):1282-1294.
33. Garavan H, Bartsch H, Conway K, et al. Recruiting the ABCD sample: Design considerations and procedures. *Dev Cogn Neurosci.* 2018;32:16-22.
34. Jernigan TL, Brown SA, Coordinators AC. Introduction. *Dev Cogn Neurosci.* 2018;32:1-3.
35. Casey BJ, Cannonier T, Conley MI, et al. The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Dev Cogn Neurosci.* 2018;32:43-54.
36. Barch DM, Albaugh MD, Avenevoli S, et al. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Dev Cogn Neurosci.* 2018;32:55-66.
37. Lisdahl KM, Sher KJ, Conway KP, et al. Adolescent brain cognitive development (ABCD) study: Overview of substance use assessment methods. *Dev Cogn Neurosci.* 2018;32:80-96.
38. Luciana M, Bjork JM, Nagel BJ, et al. Adolescent neurocognitive development and impacts of substance use: Overview of the adolescent brain cognitive development (ABCD) baseline neurocognition battery. *Dev Cogn Neurosci.* 2018;32:67-79.
39. Uban KA, Horton MK, Jacobus J, et al. Biospecimens and the ABCD study: Rationale, methods of collection, measurement and early data. *Dev Cogn Neurosci.* 2018;32:97-106.
40. Zucker RA, Gonzalez R, Feldstein Ewing SW, et al. Assessment of culture and environment in the Adolescent Brain and Cognitive Development Study: Rationale, description of measures, and early data. *Dev Cogn Neurosci.* 2018;32:107-120.
41. Robinson SM, Sobell LC, Sobell MB, Leo GI. Reliability of the Timeline Followback for cocaine, cannabis, and cigarette use. *Psychol Addict Behav.* 2014;28(1):154-162.
42. Wade NE, Tapert SF, Lisdahl KM, Huestis MA, Haist F. Substance use onset in high-risk 9-13 year-olds in the ABCD study. *Neurotoxicol Teratol.* 2022;91:107090.
43. Cooper GA, Kronstrand R, Kintz P, Society of Hair T. Society of Hair Testing guidelines for drug testing in hair. *Forensic Sci Int.* 2012;218(1-3):20-24.
44. Taylor M, Collin SM, Munafo MR, MacLeod J, Hickman M, Heron J. Patterns of cannabis use during adolescence and their association with harmful substance use behaviour: findings from a UK birth cohort. *J Epidemiol Community Health.* 2017;71(8):764-770.

45. Akshoomoff N, Newman E, Thompson WK, et al. The NIH Toolbox Cognition Battery: results from a large normative developmental sample (PING). *Neuropsychology.* 2014;28(1):1-10.
46. Anokhin AP, Luciana M, Banich M, et al. Age-related changes and longitudinal stability of individual differences in ABCD Neurocognition measures. *Dev Cogn Neurosci.* 2022;54:101078.
47. Nixon SJ, Prather R, Lewis B. Sex differences in alcohol-related neurobehavioral consequences. *Handb Clin Neurol.* 2014;125:253-272.
48. Schmidt M. Rey Auditory and Verbal Learning Test: A Handbook. 1996, Los Angeles.
49. Halladay J, Kershaw S, Devine EK, et al. Covariates in studies examining longitudinal relationships between substance use and mental health problems among youth: A meta-epidemiologic review. *Drug Alcohol Depend.* 2025;271:112665.
50. Sheridan MA, Peverill M, Finn AS, McLaughlin KA. Dimensions of childhood adversity have distinct associations with neural systems underlying executive functioning. *Dev Psychopathol.* 2017;29(5):1777-1794.
51. Crane NA, Schuster RM, Mermelstein RJ, Gonzalez R. Neuropsychological sex differences associated with age of initiated use among young adult cannabis users. *J Clin Exp Neuropsychol.* 2015;37(4):389-401.
52. Wade NE, Si Y, Tapert SF, et al. Prevalence of Biochemically-Verified Substance Use in Healthy Adolescents Across the United States: Hair Toxicology Results in the ABCD Study. *medRxiv [Preprint].* 2025:2025.2009.2019.25336190.
53. Rice JP, Reich T, Bucholz KK, et al. Comparison of direct interview and family history diagnoses of alcohol dependence. *Alcohol Clin Exp Res.* 1995;19:1018-1023.
54. Sullivan RM, Wade NE, Wallace AL, et al. Substance use patterns in 9 to 13-year-olds: Longitudinal findings from the Adolescent Brain Cognitive Development (ABCD) study. *Drug Alcohol Depend Rep.* 2022;5.
55. Wade NE, Bagot KS, Tapert SF, Gruber SA, Filbey FM, Lisdahl KM. Cognitive Functioning Related to Binge Alcohol and Cannabis Co-Use in Abstinent Adolescents and Young Adults. *J Stud Alcohol Drugs.* 2020;81:479-483.
56. Hernandez Mejia M, Wade NE, Baca R, Diaz VG, Jacobus J. The Influence of Cannabis and Nicotine Co-use on Neuromaturation: A Systematic Review of Adolescent and Young Adult Studies. *Biol Psychiatry.* 2021;89(2):162-171.
57. Bates D, Mächler M, Bolker B, Walker S, 2015 Y. Fitting linear mixed-effects models using lme4. *J Stat Softw.* 2015;67:1-48.
58. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society.* 1995;57(1):289-300.
59. Bartoń K. MuMIn: Multi-Model Inference. *R package version 14811.* 2025.
60. Lenth R, Piaskowski J. emmeans: Estimated Marginal Means, aka Least-Squares Means. *emmeans: Estimated Marginal Means, aka Least-Squares Means.* 2025.
61. Green R, Wolf BJ, Chen A, et al. Predictors of Substance Use Initiation by Early Adolescence. *Am J Psychiatry.* 2024;181(5):423-433.

62. Miller AP, Baranger DAA, Paul SE, et al. Characteristics Associated With Cannabis Use Initiation by Late Childhood and Early Adolescence in the Adolescent Brain Cognitive Development (ABCD) Study. *JAMA Pediatr.* 2023;177:861-863.

63. Prini P, Zamberletti E, Manenti C, Gabaglio M, Parolaro D, Rubino T. Neurobiological mechanisms underlying cannabis-induced memory impairment. *Eur Neuropsychopharmacol.* 2020;36:181-190.

64. Wade NE, Nguyen-Louie TT, Wallace AL, Sullivan RM, Tapert SF. Commercial Cannabidiol for Community-Based Young Adolescents: Predicting Medicinal Use. *Cannabis Cannabinoid Res.* 2025;10:e353-e361.

65. Johnson E, Kilgore M, Babalonis S. Label accuracy of unregulated cannabidiol (CBD) products: measured concentration vs. label claim. *J Cannabis Res.* 2022;4(1):28.

66. Johnson E, Kilgore M, Babalonis S. Cannabidiol (CBD) product contamination: Quantitative analysis of Delta(9)-tetrahydrocannabinol (Delta(9)-THC) concentrations found in commercially available CBD products. *Drug Alcohol Depend.* 2022;237:109522.

67. Miller OS, Elder EJ, Jr., Jones KJ, Gidal BE. Analysis of cannabidiol (CBD) and THC in nonprescription consumer products: Implications for patients and practitioners. *Epilepsy Behav.* 2022;127:108514.

68. Gidal BE, Vandrey R, Wallin C, et al. Product labeling accuracy and contamination analysis of commercially available cannabidiol product samples. *Front Pharmacol.* 2024;15:1335441.

69. Scott JC, Slomiak ST, Jones JD, Rosen AFG, Moore TM, Gur RC. Association of Cannabis With Cognitive Functioning in Adolescents and Young Adults: A Systematic Review and Meta-analysis. *JAMA Psychiatry.* 2018;75(6):585-595.

70. Matthay EC, Hagan E, Gottlieb LM, et al. Powering population health research: Considerations for plausible and actionable effect sizes. *SSM Popul Health.* 2021;14:100789.

71. Durlak JA. How to select, calculate, and interpret effect sizes. *J Pediatr Psychol.* 2009;34(9):917-928.

72. Banz BC, Hersey D, Vaca FE. Coupling neuroscience and driving simulation: A systematic review of studies on crash-risk behaviors in young drivers. *Traffic Inj Prev.* 2021;22(1):90-95.

73. Walshe EA, Ward McIntosh C, Romer D, Winston FK. Executive Function Capacities, Negative Driving Behavior and Crashes in Young Drivers. *Int J Environ Res Public Health.* 2017;14(11).

74. Lisdahl KM, Gilbart ER, Wright NE, Shollenbarger S. Dare to delay? The impacts of adolescent alcohol and marijuana use onset on cognition, brain structure, and function. *Front Psychiatry.* 2013;4:53.

75. Lisdahl KM, Tapert S, Sher KJ, et al. Substance use patterns in 9-10 year olds: Baseline findings from the adolescent brain cognitive development (ABCD) study. *Drug Alcohol Depend.* 2021;227:108946.

Acknowledgements: Authors would like to thank the families who participate in the ABCD Study and the staff who collect the data. Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development™ (ABCD) Study, held in the NIH Brain Development Cohorts Data Sharing Platform. This is a multisite, longitudinal study designed to recruit more than 10,000 children aged 9–10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the **National Institutes of Health** and additional federal partners under award numbers: U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147.

A full list of supporters is available at Federal Partners – ABCD Study. ABCD Consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD Consortium investigators. The ABCD data repository grows and changes over time. The ABCD data used in this report came from https://doi.org/10.82525/jy7n-g441. Additional support for this work was made possible from NIEHS R01-ES032295 and R01-ES031074.

Author Contributions:
NEW conceptualized, conducted statistical analyses, interpreted fndings, drafted the original manuscript, and completed critical revisions. RMS and ALW assisted in conceptualization, data analysis, interpretation, and critical revisions. RV assisted in design, stastisical methods, and critical revisions. VS, KML, MAH, PDB, HB, LM, and JJ all contributed to conceptualization, interpretation, and critical revisions. SFT contributed to conceptualization and design, interpretation, and critical revisions. All authors contributed to the writing and revision of the manuscript and have reviewed and approved the final version of the manuscript.

Funding: N.E. Wade was supported by National Institute on Drug Abuse (DA050779, PI: Wade). This work was also supported by DA064409 (PI: Sullivan), DA062011 (PI: Wallace), T32 AA013525 (PI: Riley/Spadoni to Szpak), R01DA062432 (MPI Lisdahl, Hillard), 2U01DA041025 (MPI Lisdahl, Larson), K01DA057389 (PI: Gonçalves), and NARSAD/Brain Behavior Research Foundation (Gonçalves).

Competing Interests: Authors have no conflicts of interest to disclose.