Developmental candidate GHP-88310/EIDD-3608 with high tolerability and oral efficacy in measles and respiratory paramyxovirus models

Carolin M Lieber[1], Josef D Wolf[1], Mugunthan Govindarajan[2], Jeong-Joong Yoon[1], Zachary M Sticher[2], Claire E Ruckel[1], Alexander I Leach[1], Lauren A Harrison[1], Dariia Vyshenska[3], Amalia Anne Cruz[2], Meghan K. Andrews[2], Rebecca E. Krueger[2], Robert M Cox[1], George R Painter[2], Alexander L Greninger[3], Michael G Natchus[2], Richard K Plemper[1]*

[1]Center for Translational Antiviral Research, Georgia State University Institute for Biomedical Sciences, Atlanta, GA, 30303, USA.

[2]Emory Institute for Drug Development, Emory University, Atlanta, GA 30322, USA.

[3]Virology Division, Department of Laboratory Medicine and Pathology, University of Washington Medical Center, Seattle, Washington, USA

*Corresponding author: rplemper@gsu.edu

Abstract

Orthoparamyxoviruses such as human parainfluenza virus type-3 (HPIV3), measles virus (MeV), and

emerging henipaviruses pose a substantial threat to human health. We discovered an orally efficacious

broad-spectrum non-nucleoside inhibitor of orthoparamyxovirus polymerases. However, here we found

that tolerability in higher mammals was limited to 50 mg/kg b.i.d., close to the lowest efficacious dose.

We report development of a clinical candidate analog, GHP-88310 (EIDD-3608), which combines

improved oral efficacy with favorable tolerability in non-rodent species (ferrets and dogs), greatly

expanding the therapeutic window. GHP-88310 had potent activity against the primary indication,

HPIV3 and related respiroviruses, and the secondary indication, MeV and related morbilliviruses. In 7-

day multi-dose tolerability studies, daily doses of 2,000 mg/kg were well-tolerated. Pharmacokinetic

analysis revealed an altered plasma exposure profile after oral delivery of GHP-88310 compared to the

original hit. In the HPIV3 cotton rat model, GHP-88310 significantly reduced viral load in the upper and

lower respiratory tract when administered orally at 50 mg/kg twice daily. Therapeutic administration to

ferrets infected with canine distemper virus (CDV), causing lethal measles-like disease, resulted in

complete survival, significant reduction of primary viremia and shed viral load, and alleviated

lymphocytopenia. Once-daily GHP-88310 was efficacious at respective 50 mg/kg and 150 mg/kg in the

CDV ferret and HPIV3 cotton rat models. The compound was sterilizing against HPIV3 at

pharmacokinetics-informed dynamic concentrations in disease-relevant human airway epithelium

organoids. These results identify GHP-88310 as a candidate for urgently needed improved

orthoparamyxovirus disease management.

Introduction

The orthoparamyxovirus subfamily contains major emerging and reemerging pathogens, such as

measles virus (MeV), the human parainfluenzaviruses type 1 and 3 (HPIV1 and HPIV3), and the highly

pathogenic Nipah and Hendra viruses (NiV and HeV) (*1, 2*). These viruses circulate in the human

population or can readily transition species barriers as exemplified by spillover of NiV from *Pteropus* fruit

bats to domestic animals and humans (*2*).

Although a live-attenuated measles vaccine has been available for decades, endemic transmission

of MeV continues in large geographical areas including parts of Europe. Largely fueled by reports of an

alleged link between the MMR vaccine and the incidence of autism (*3, 4*), measles vaccination rates have

declined in the last decade in the US. Further exacerbated by the COVID-19 pandemic (*5, 6*), low herd

immunity in 2025 resulted in the largest measles outbreak in North America in 3 decades with over 1,500

confirmed cases around the epicenter in West Texas, three deaths, and thousands of additional cases

in neighboring states, Mexico, and Canada(*7-9*). Approximately 20% of patients were hospitalized with

severe disease (*10*). However, no approved antivirals for improved measles management beyond

supportive care exist (*11-13*). A major obstacle that has hampered drug development against MeV is the

challenge of conducting clinical efficacy trials in a predominantly pediatric patient population (*14-16*). In

contrast, related HPIV3 can reinfect people throughout life. Older adults and the immunocompromised

are at greatest risk of HPIV3 spread to the small airways, causing lower respiratory tract infection (LRTI)

that presents with bronchiolitis and pneumonia (*17, 18*). Adult hematopoietic stem cell transplant

recipients are among the most vulnerable groups with mortality rates from HPIV3 LRI of up to 75% (*19,*

20), and no vaccine prophylaxis or therapeutic is currently available to manage the disease.

We have identified HPIV3 as the primary indication for an anti-orthoparamyxovirus drug development

program based on the rationale that the HPIVs constitute an unmet medical need with an estimated 3

million medically attended cases in the United States annually (*21, 22*); that progression to severe HPIV3

disease in at-risk adults is relatively slow with a median of 3 days between upper respiratory symptoms

and onset of pneumonia, opening a window for intervention (*20*); and that precedence for clinical

evaluation of HPIV3 inhibitors in transplant patients has been established (*19, 20*).

In previous work, we identified a novel non-nucleoside inhibitor class of the paramyxovirus RNA-

dependent RNA polymerase (RdRP) complex through an anti-HPIV3 high-throughput drug screen (*23,*

24). The hit compound, GHP-88309, showed unusually broad-spectrum sub-micromolar activity against

orthoparamyxoviruses of the respirovirus (HPIV1, HPIV3, Sendai virus (SeV)), morbillivirus (MeV and

canine distemper virus (CDV)), and henipavirus (Ghana henipavirus and Cedar virus) genera (*23-25*).

Mechanistic characterization revealed that GHP-88309 binding prevents initiation of the viral RdRP

complex at the promoter, most likely by sterically blocking structural reorganization of the polymerase

from initiation into elongation configuration (*23, 24, 26*). For proof-of-concept, we established oral

efficacy in the SeV-mouse model as a lethal substitute assay for human respirovirus disease (*23, 24*).

In parallel, we validated broad anti-orthoparamyxovirus efficacy in CDV-infected ferrets, which develop

measles-like disease that presents with high fever, rash, and lymphocytopenia (*25*). The infection

recapitulates severe measles cases, since CDV pathogenicity in ferrets is high and all infected animals

succumb within 10-12 days of infection (*25*). A course of oral GHP-88309 started after the onset of

clinical signs but before peak viremia changed outcome and all treated animals survived (*23, 24*). Thus,

GHP-88309 met three fundamental requirements of the target product profile that we require of an anti-

orthoparamyxovirus candidate: orally bioavailable to facilitate rapid self-administration; active against

different high-consequence orthoparamyxoviruses including an indication with a clear path to clinical

testing; and shelf-stable under ambient conditions for stockpiling and rapid distribution. However, the

envisioned clinical use in pediatric patients and the immunocompromised demands high tolerability with

wide safety margins.

In this study, we report an unexpected tolerability limit of GHP-88309 in higher mammals of 50

mg/kg. In search of a clinical candidate, we launched a synthetic lead development program that

yielded a novel chemotype class with favorable tolerability in rodent and non-rodent (ferrets and dogs)

toxicology species. We report enhanced oral efficacy of this candidate against the primary HPIV3

indication in a relevant animal model, show a broadened therapeutic window against the secondary

measles indication, and establish dosing paradigms against HPIV3 in disease-relevant human airway

epithelium organoid cultures based on PK-informed physiologically achievable dynamic concentrations.

Results

Prolonged 17-day treatment of CDV disease in ferrets with the first-generation lead GHP-88309

was highly effective at a minimally efficacious oral dose of 50 mg/kg b.i.d. and all animals survived the

infection (*25*). In mice, multi-dose administration of GHP-88309 at 150 mg/kg (highest dose tested)

b.i.d. was well tolerated without adverse events (*23*).

Synthetic lead development to overcome tolerability limitations of GHP-88309

To estimate the therapeutic index (TI) of GHP-88309 in a non-rodent species, we increased oral

dose levels to 150 mg/kg in a multi-dose PK study in search of tolerability limits in ferrets (Fig. 1A).

Unexpectedly, ferrets in the 150 mg/kg group reached endpoint within hours following the second dose

(Fig. 1B, Fig. S1A,B). Complete blood count (CBC) analysis revealed a drastic drop in platelet counts in

animals of this group after compound administration (Fig. S1C). Plasma PK analysis of the 50 mg/kg

group on day 7 (after 14 doses) showed high plasma concentration peaks (C_{max} 43.5 μM) and an

overall exposure of 270 μM $\times$ hr (Fig. 1C). A single ascending dose study confirmed that a dose level of

50 mg/kg was well tolerated, whereas animals receiving 150 mg/kg showed transient paralysis and a

drop in body temperature, and ferrets dosed at 500 mg/kg died within 4 hours of dosing (Fig. 1D-F).

Building on our initial understanding of the structure-activity relationship (SAR) of the scaffold (*23*),

we launched a synthetic lead development program with the primary objective to improve tolerability in

non-rodent species. Six azaquinoline lead candidates emerged from this exercise (Table S1), each

featuring a substituted heterocyclic ring (Fig. 1G). None showed appreciable cytotoxicity in cell culture

(Fig. 1H) and all maintained antiviral potency *in vitro* within a <2-fold range against the HPIV3 primary

indication (Fig. 1I) and respective <8-fold and <3-fold range against the morbillivirus (Fig. 1J) and

henipavirus (Fig. 1K) secondary indications.

Single-dose PK prescreening of these six candidates in mice at an oral dose level of 20 mg/kg

revealed highest plasma exposure for, in descending order, GHP-88382, GHP-88310, and GHP-88370.

However, none of these analogs reached the high exposure levels of GHP-88309 (Fig. 1L). Based on

PK performance, we selected these three analogs for proof-of-concept efficacy testing against the

primary respirovirus indication.

GHP-88310 efficacy against the primary respirovirus indication

Using the lethal SeV 129X1/SvJ mouse model as a surrogate for human respirovirus disease, we

screened the three developmental candidates in comparison with GHP-88309, each dosed orally at 150

mg/kg twice daily (b.i.d.), initiated 24 hours after infection (Fig. 2A). All animals receiving GHP-88309

and GHP-88310 survived without developing clinical signs, whereas GHP-88382 mediated only partial

survival and GHP-88370 provided no therapeutic benefit (Fig. 2B, Fig. S2A,B). Viral load in the upper

(Fig. 2C) and lower (Fig. 2D) respiratory tract of GHP-88310-treated animals was significantly reduced

compared to vehicle-treated mice. Effect size was equivalent to that achieved by GHP-88309. In

contrast, GHP-88382 reduced viral burden only in the lower respiratory tract and GHP-88370 had no

significant effect on viral load.

Accordingly, we selected GHP-88310 for a dose-to-failure study against the primary indication in

the HPIV3 cotton rat model, measuring viral load in the respiratory tract 3.5 days post-infection (dpi;

Fig. 2E). In this model, the minimal dose of original hit GHP-88309 that significantly reduced viral load

in both trachea and lungs was 300 mg/kg b.i.d. (Fig. 2F,G; Fig. S3A,B). GHP-88310 was substantially

more potent, showing lowest efficacious doses of 50 mg/kg b.i.d. and 16.5 mg/kg b.i.d. in the upper and

lower respiratory tract, respectively. Plasma exposure of GHP-88310 in cotton rats was approximately

one-half that of GHP-88309 after a single oral dose of 50 mg/kg, despite greater efficacy of GHP-88310

(Fig. 2H,I). Corresponding variations in tissue distribution were greatest in brain, heart, and kidneys,

smallest in liver and lung (Fig. 2J). However, none of these differences were statistically significant.

A 7-day ascending multi-dose study with GHP-88310 culminating in a 2,400 mg/kg daily dose (Fig.

2K) showed that the compound was well tolerated at all dose levels without changes in bodyweight,

temperature, or other adverse signs (Fig. 2L; Fig. S4A,B). Plasma exposure built up to nearly 2,000 μM

$\times$ hr at the highest dose (Fig. 2M). Trough plasma concentrations at the lowest dose tested exceeded

19 μM, corresponding to 5-times the cell-culture EC_{90} concentration of GHP-88310 against HPIV3.

Based on these high exposure levels, we examined anti-HPIV3 efficacy in a once-daily (q.d.)

therapeutic dosing regimen, starting treatment of cotton rats 12 hpi (Fig. 2N). Oral GHP-88310 at 150

mg/kg q.d. or higher significantly reduced viral load in trachea and the lower respiratory tract (Fig.

2O,P) without changes in body weight or temperature (Fig. S5A,B).

GHP-88310 tolerability in non-rodent toxicology species

Having validated oral efficacy of GHP-88310 against the primary indication in cotton rats, we

explored PK properties of the compound in ferrets in a single ascending dose oral PK study (Fig 3A). A

dose level of 500 mg/kg GHP-88310 was well tolerated and plasma exposure increased dose-

proportionally between 150 and 500 mg/kg doses (Fig. 3B). A subsequent escalating 7-day ascending

multi-dose tolerability study in ferrets demonstrated that administering GHP-88310 at a daily dose of up

to 2,000 mg/ml (Fig. 3C) had no negative effect on bodyweight, body temperature, and CBC

parameters including platelet counts (Fig. S6A-E). All animals survived the full 7-day course without

signs of adverse effects, in contrast to reference animals dosed with GHP-88309 at 150 mg/kg, which

succumbed within hours of receiving the second dose (Fig. 3D). Plasma concentrations examined

once-daily 4 hours after dosing (expected C_{max}) plateaued dose-dependently after administration of four

(500 mg/kg) to eight (1,000 mg/kg) doses (Fig. 3E). Overall exposure after 7 days (14 doses) increased

dose-dependently from 336 $\mu M \times$ hr (50 mg/kg b.i.d.) to 2,140 $\mu M \times$ hr (500 mg/kg b.i.d) (Fig. 3F). At a

dose of 50 mg/kg b.i.d., total plasma exposure of GHP-88310 was slightly higher than that of GHP-

88309 with 336 $\mu M \times$ hours versus 270 $\mu M \times$ hours at study end (Fig. 1C vs Fig. 3F).

CBC parameters of GHP-88310-treated animals did not deviate from the normal range over the

course of the study (Fig. 3G) and whole blood chemistry panels taken on days 2 and 7 were likewise

unremarkable (Fig. S7). In contrast, a reference animal that had received two doses of GHP-88309 at

150 mg/kg showed changes in three of four liver enzymes examined – alkaline phosphatase (ALP),

alanine transaminase (ALT), and aspartate aminotransferase (AST) – by more than 750% one day

post-dosing and prior to death, suggesting GHP-88309-induced hepatotoxicity. Due to rapid death after

repeated administration of GHP-88309 at this dose level, a comprehensive blood chemistry analysis of

all animals in the reference group was not possible.

Based on greatly improved tolerability of GHP-88310 compared to GHP-88309, we advanced the

compound to single ascending-dose PK testing in beagle dogs, the predominant non-rodent toxicology

species used to inform investigational new drug (IND)-enabling safety studies (Fig. 3H). Dogs were

dosed orally with GHP-88310 at 50, 150, or 500 mg/kg, each followed by assessment of plasma

exposure over a 24-hour period (Fig. 3H). All dose levels were well tolerated without adverse signs or

weight loss of the animals (Fig. 3I). Plasma exposure ranged from 135 to 2,150 $\mu M \times hr$ (Fig. 3J),

demonstrating approximate dose-proportionality and exceeding at high-dose that observed in ferrets.

These results demonstrated that GHP-88310 plasma exposure levels exceeding those at which GHP-

88309 was lethal in ferrets were well tolerated in both ferrets and dogs (Fig. 3B and 3J).

GHP-88310 efficacy against measles-like disease in ferrets

Measles has emerged as an urgent secondary indication for GHP class paramyxovirus inhibitors

due to growing vaccination hesitancy in many high-income countries and endemic MeV transmission in

large geographical regions in sub-Saharan Africa and Southeast Asia (*12, 27, 28*). We have

demonstrated potent oral efficacy of GHP-88309 against the CDV/ferret surrogate model of human

measles (*25*). GHP-88310 and GHP-88309 had virtually identical antiviral activity against both MeV

(Fig. 1J) and CDV (Fig. 4A) in cell culture, returning active (EC_{50}) concentrations within a 2-fold range

of each other.

In a proof-of-concept oral efficacy study, we infected ferrets with a lethal inoculum of the wild type

CDV-5804p strain and initiated treatment at 3 dpi, when PBMC-associated viremia and fever become

first detectable in this model (Fig. 4B and Fig. S8A,B). Exploring two distinct treatment regimens,

animals received GHP-88310 either at 50 and 150 mg/kg b.i.d., or 50 and 100 mg/kg q.d., the latter

corresponding to the lowest efficacious daily dose of 100 mg/kg of GHP-88309 (*25*) and half that dose.

Treatment was continued in all groups until 10 dpi (150 mg/kg b.i.d. group) or 21 dpi (50 mg/kg b.i.d.

and q.d. groups). All treated animals survived the infection (Fig. 4C) and clinical signs were fully

mitigated (Fig. S8A,B), whereas all animals in a vehicle-treated reference group succumbed to CDV

disease with a median survival of 11 dpi. Cell-associated viremia was first detectable 3 dpi, when

therapeutic treatment was initiated, and rapidly progressed in vehicle-treated animals to peak viremia

titers of approximately 1.3×10^5 $TCID_{50}$ units per 10^6 PBMCs at 7 dpi (Fig. 4D). Treatment at 50 or 100

mg/kg q.d. or 50 mg/kg b.i.d. lowered peak titers by ~1 order of magnitude and resolved viremia with

similar kinetics by 15-16 dpi. Treatment at 150 mg/kg reduced peak viremia by 2 orders of magnitude

and was sterilizing at 10 dpi. The effect of treatment on shed virus load in nasal lavages closely

mirrored that on viremia titers (Fig. 4E). Virus shedding from animals receiving GHP-88310 at 150

mg/kg b.i.d. peaked 2 orders of magnitude lower than in the vehicle group and shedding from treated

animals ceased 6 dpi. Low b.i.d. dose animals and animals of both q.d. groups showed biphasic

shedding kinetics, characterized by an initial decline after treatment start followed by a second

shedding peak at approximately 10 dpi. Infectious virus was undetectable in all treated animals 17 dpi.

Lymphocytopenia, a hallmark of morbillivirus disease, was efficiently mitigated by 50 mg/kg b.i.d. or 100

mg/kg q.d. GHP-88310 and fully suppressed at the 150 mg/kg b.i.d. dose level (Fig. 4F). Animals in the

50 mg/kg q.d. treatment group experienced vehicle-like lymphocytopenia until 9 dpi. Subsequently,

lymphocyte counts rapidly stabilized and returned to normal range by 27 dpi. Consistent with transient

lymphocytopenia, the emergence of neutralizing antibodies was delayed in animals of the 50 mg/kg q.d.

cohort compared to the other treatment groups (Fig. 4G). Whole genome sequencing of treatment-

experienced virus populations recovered from animals 3, 5, 8,13, and 14 dpi did not detect any

mutations in polymerase proteins (supplementary dataset file S1), indicating a high barrier against viral

escape from GHP-88310 *in vivo*.

Molecular interactions between GHP-88310 and the viral L protein

We have identified the central cavity of the viral P-L polymerase complex as target site for GHP-

88309-class inhibitors through photo-affinity crosslinking and reduced susceptibility profiling (*23*).

Several high-resolution structural models of the MeV polymerase complex were recently solved (*29,*

30), setting the stage for extraction of the GHP-88310 docking pose. Activity testing of GHP-88310

against panels of recombinant HPIV3 (Fig. 5A), recombinant SeV (Fig. 5B), and MeV minigenome (Fig.

5C) harboring mutations mediating escape from GHP-88309 demonstrated, with few exceptions, high

consistency of resistance sites, suggesting conserved docking poses of both compounds. We defined

substitutions that caused a >5-fold increase in EC_{50} and/or EC_{90} values as major drivers of reduced

viral susceptibility (highlighted in Fig. 5A-C). Exceptions were the HPIV3 L-T1010A variant, which

remained fully sensitive to GHP-88310, and SeV L-Y942H, which mediated robust resistance to GHP-

88310 but moderate escape from GHP-88309.

Informed by these conserved mediators of resistance and photoaffinity mapping results for GHP-

88309, we docked both compounds *in silico* into a native structural model of the MeV polymerase

(PDB: 9DUS). Top scoring poses consistently placed both inhibitors into the same defined pocket at the

interface between the L protein RdRP and PRNTase domains (Fig. 5D and Fig. S9). However, distinct

predicted docking poses emphasized a high-affinity interaction between the GHP-88310 diazo quinoline

ring and residue A866 of the RdRP domain that was not present in the GHP-88309 pose (Fig. S10A,B).

Major resistance sites lined the binding pocket, predicting direct contact between the ligand and L-

protein residues A866, Y942, and T1010. Additional hot-spot sites were located in close proximity,

suggesting that viral escape from either inhibitor is predominantly due to primary, rather than long-

range secondary, resistance.

GHP-88310 efficacy in primary human cells and organoids

To assess whether antiviral potency of GHP-88310 supports realistic human dose concentrations,

we determined efficacy in physiologically relevant primary human airway epithelium cells and well-

differentiated human airway epithelium organoids. Wash-in and wash-out studies assessed intracellular

exposure levels during 2-hour periods in undifferentiated human bronchial tracheal epithelial cells

(HBTECs) in, respectively, the presence of 80 μM extracellular GHP-88310 (wash-in) or after removal

of extracellular compound (wash-out) (Fig. S11). At wash-in, a maximal intracellular exposure plateau

of approximately 10^5 pmol/10^6 cells was reached in 15 minutes (earliest sampling time point), indicating

high plasma-membrane permeability, and was sustained for the duration of the experiment. After

removal of extracellular GHP-88310, intracellular drug levels declined by approximately 2 orders of

magnitude within the 2-hour monitoring window.

Efficacy testing of GHP-88310 against the primary HPIV3 indication in undifferentiated HBTECs

showed sub-micromolar potency and a steep Hill slope, resulting in a fully sterilizing antiviral effect at

concentrations of 10 μM and greater (Fig. 6A). After generation of well-differentiated airway epithelium

organoids grown at air-liquid interface (Figure 6B), we exposed cultures to ascending concentrations of

GHP-88310 in the basolateral chamber of the transwell system for 48 hours each and measured

corresponding transepithelial electrical resistance (TEER) as a well-established marker of cytotoxicity in

the system(*31-33*). Electrical resistance remained unchanged at approximately 600 ohm $\times$ cm^2 before

and after exposure (Fig. 6C), indicating that GHP-88310 was well-tolerated at the highest

physiologically achievable concentrations without disturbing tissue integrity.

Dose-response assays using static concentrations of basolateral GHP-88310 after apical infection

of organoids with 5,000 plaque forming units (pfu) HPIV3 returned nanomolar EC$_{50}$ values with

favorable hill slope (EC$_{90}$ 0.13 μM), based on titers of apically released progeny virions (Fig. 6D).

Sterilizing conditions were reached at 1.85 μM GHP-88310 in the basolateral chamber, approximately

equivalent to trough plasma concentrations in the multi-dose ferret PK studies when compound was

administered at 50 mg/kg. To verify enhanced antiviral potency of HPIV3 in the organoid models

compared to immortalized cell lines, we imaged HPIV3-infected, fixed, and stained organoid sections

after incubation in the presence of 0.5 or 10 μM compound. Titration of progeny virions prior to imaging

demonstrated that 0.5 μM basolateral GHP-88310 reduced virus titers by approximately 2 orders of

magnitude (Fig. 6E), which was consistent with the dose-response assay results. Anti-ZO-1 staining

revealed an unperturbed tight junction network in the presence of both concentrations examined,

similar to mock-treated uninfected cultures (Fig. 6F). Only sporadic viral antigen was detectable after

incubation of infected cells in the presence of 0.5 μM GHP-88310, and none after incubation with 10

μM compound, whereas HPIV3 proteins were abundant in organoids that had received vehicle volume-

equivalents. Overlay of viral and beta-actin stains revealed that viral replication was concentrated in

ciliated cells (Fig. S12), consistent with previous reports for HPIV3(*34, 35*).

To assess GHP-88310 efficacy in human organoids at physiological, dynamic concentrations, we

recapitulated ferret plasma PK curves in the basolateral chambers of the airway epithelium organoids

infected with HPIV3. Two dose levels, 20 mg/kg and 50 mg/kg, each represented by b.i.d. and q.d.

dosing regimens, were simulated over a 48-hour period each (Fig. 6G). At study end, we determined

progeny virus yields released from the apical surface. Validating our finding of high anti-HPIV3 potency

of static GHP-88310 in the organoids, both high-dose regimens were sterilizing (Fig. 6H). Also

simulating a low-dose b.i.d. treatment fully suppressed production of infectious progeny virions (Fig. 6I),

although plasma PK profiles called for peak GHP-88310 concentrations of only ~15 μM and no

compound was present in the basolateral chamber for 6 hours in every 12-hour interval (Fig. 6H). Even

recapitulating a 20 mg/kg q.d. plasma exposure profile in the basolateral chamber, which comprised

compound-free incubation periods of 18 hours in every 24-hour interval, greatly inhibited virus

replication. Progeny virus yield was statistically significantly reduced by over two orders of magnitude

compared to organoid cultures that had been incubated in the presence of basolateral vehicle volume-

equivalents (Fig. 6I). These results support high developmental potential of GHP-88310 for human

therapy, demonstrating that potent antiviral efficacy is achieved at physiological concentrations in

disease-relevant human tissue models without cytotoxicity limitations.

Discussion

This study identified the broad-spectrum anti-orthoparamyxovirus clinical candidate GHP-88310,

which showed favorable tolerability properties in rodent species and higher mammals including dogs. De-

risking GHP-88309 illuminated a sharp tolerability limit in higher mammals at concentrations exceeding

100 mg/kg daily dose. We addressed this issue synthetically through development of GHP-88310, which

occupies novel chemical space and was tolerated without adverse effects by ferrets and dogs at very

high dose levels that far exceeded the lethal dose of GHP-88309 in ferrets. Differential responses

between rodents and non-rodents to drug exposure are not unusual and often attributable to the high

metabolic rate of rodents(*36-38*), an expanded rodent cytochrome P450 system that results in fast

breakdown of drugs with frequently distinct metabolites (*39-41*), and/or the high liver to body weight ratio

of rodents that increases resistance to chemical stress (*42*). In the case of GHP-88309, plasma exposure

was high after oral dosing in both rodents and non-rodents, but only non-rodents experienced tolerability

limits. At equal dose levels, GHP-88310 exposure trailed that of GHP-88309. Our ascending dose PK

studies demonstrated, however, that improved tolerability of GHP-88310 was not a trivial consequence

of lower exposure compared to GHP-88309, since plasma exposure in ferrets and dogs after high-dose

GHP-88310 far exceeded lethal levels of GHP-88309 in ferrets. We propose that toxicity of high-dose

GHP-88309 in non-rodents results most likely from accumulation of an unidentified metabolite, which is

not produced in rodents due to their distinct cytochrome P450 composition, and absent from GHP-88310

breakdown products due to its distinct chemical structure.

A hallmark of viral escape from GHP-88309 is that resistance emerged readily in cell culture, but all

escape variants were apathogenic and no escape mutations appeared *in vivo* (*23*). Susceptibility

testing of developmental candidate GHP-88310 against recombinant strains of three viral targets – the

primary indication HPIV3, surrogate respirovirus SeV, and secondary indication MeV – that each

harbored confirmed GHP-88309 escape mutations (*23*) revealed overlapping susceptibility profiles. All

mutations were located in the viral L protein, lining the interface between the RdRP and PRNTase

domains in the central polymerase cavity. However, tolerance to GHP-88310 was, in most cases, less

robust than escape from GHP-88309. Conserved susceptibility profiles spotlight a common molecular

target site of both compounds, whereas reduced robustness of escape from GHP-88310 suggests

altered docking poses, which was substantiated by *in silico* QSAR models. Importantly, this hallmark

feature of GHP-88309 resistance appears to equally apply to GHP-88310, since whole genome

sequencing of heavily treatment-experienced virus populations recovered from ferrets did not reveal

substitutions at known escape hot-spots or any other polymerase residues. Although future work must

address whether additional specific GHP-88310 escape sites exist, our results indicate a high barrier

against viral escape from GHP compounds without concomitant loss in pathogenicity.

It was the primary focus of this synthetic lead identification program to improve tolerability in higher

mammals without compromising antiviral potency, and GHP-88309 and GHP-88310 indeed have equal

activity in cell culture. We therefore did not anticipate also achieving a substantial *in vivo* potency boost

in both rodent and non-rodent host species. Quantitation of drug tissue levels of both compounds in

cotton rats excluded improved lung tissue distribution of GHP-88310 as a possible explanation for this

approximately 6-fold increase in antiviral efficacy. Whereas the lowest efficacious dose of GHP-88309

and GHP-88310 in the CDV ferret model was the same (50 mg/kg), plasma exposure of GHP-88310

was lower than that achieved by GHP-88309 after equal doses. The conclusion of an *in vivo* potency

boost in both models was further supported by our observation that GHP-88310 was efficacious also in

a q.d. regimen, lowering the minimally required efficacious daily dose by 50%. We propose that

breakdown of GHP-88310 in both rodents and non-rodents may generate a bioactive, but non-toxic,

metabolite that carries part of the antiviral efficacy, whereas GHP-88309 metabolism produces a toxic

by-product. Both effects, increased tolerability and enhanced antiviral efficacy of GHP-88310, have

substantially expanded the therapeutic window of the compound against the primary and secondary

indication in relevant animal models, underscoring high clinical potential.

Ex vivo, potency of GHP-88309 against HPIV3 in established cell lines and primary HBTECs was

virtually identical. However, we noted an approximately 5-fold higher antiviral potency in disease-

relevant, well-differentiated human airway epithelium organoids versus the undifferentiated HBTECs.

This effect may be due to a combination of the diverse cell types present in the organoids, their

complex 3D-architecture, and the more natural nutrient concentration gradient compared to simple cell

monolayers (*43-46*). Indeed, confocal microscopy identified, consistent with previous reports (*34, 35*),

ciliated cells as the primary host cell type for HPIV3 replication in the airway epithelium cultures. To test

the correlation between oral dose level and antiviral effect, we recapitulated ferret PK-informed dynamic

compound concentrations in the basolateral chamber of the transwell cultures, simulating repeat oral

dosing ranging from 20 mg/kg q.d. to 50 mg/kg b.i.d. We first implemented this approach when

developing molnupiravir (*47*). Simulating plasma concentrations of a 20 mg/kg q.d. GHP-88310 dosing

regimen in the basolateral chamber significantly reduced progeny virus yields by several orders of

magnitude, which translates under consideration of established species conversion factors (*48*) to a

feasible predicted oral human dose target as low as 200 mg daily. Corroborating these results, higher

dose levels were fully sterilizing.

Simulation of 20 mg/kg b.i.d. or 20 mg/kg q.d. regimens entailed two 6-hour or one 18-hour

window, respectively, in every 24-hour period in which no compound was present in the basolateral

chamber. Effective viral inhibition despite prolonged incubation in the absence of inhibitor is likely due

to inhibition of both replicase and transcriptase activity of the viral polymerase complex by GHP class

compounds (*26, 49*). This concerted action may arrest the viral replication cycle in the primary

transcription stage (*49*), preventing exponential expansion of viral protein expression and suppressing

production of immune-modulatory viral V and C proteins that counteract the type I host interferon

response (*50-53*), which exposes the virus to host cells in an unmitigated antiviral state. Both the

HPIV3 cotton rat and CDV ferret models confirmed that effective virus inhibition by only intermittently

present GHP-88310 was not an artifact of the *ex vivo* organoid system, since q.d. administration of

GHP-88310 at the lowest efficacious dose likewise resulted in either model in prolonged periods with

undetectable plasma drug levels. These results thus establish proof-of-concept that efficacious

orthoparamyxovirus therapy with a non-nucleoside polymerase inhibitor does not depend on continuous

drug plasma concentrations above the minimum inhibitory level, illuminating an overproportional

beneficial effect of targeting the mononegavirus polymerase complex.

Naturally, human organoid systems can only partially simulate the complexity of human infection.

Whereas first-in-human dose levels will be determined based on toxicology results, the organoids

provide valuable insight into defining desirable dosing objectives. Other limitations of our study arise

from incomplete recapitulation of all aspects of human disease presentation by animal models,

currently undefined highest tolerated dose levels of GHP-88310, and limited insight into a possible

effect of co-morbidities and/or low patient immune competence on antiviral efficacy. However, ferrets

are a natural host for CDV and hallmarks of human morbillivirus disease are fully preserved, supporting

confidence that treatment paradigms established in this model will provide valuable guidance for

performance in human therapy. In summary, this study identified a much-needed novel developmental

candidate for treatment of orthoparamyxovirus infections that combines attractive drug-like properties

with potent antiviral efficacy and unusually broad indication spectrum, opening a viable path to clinical

testing.

Material and Methods

Study design

Cells, normal human bronchial/tracheal epithelial (NHBE) cells, well-differentiated air liquid

interface (ALI) human airway epithelial (HAE) cells, ferrets, cotton rats and mice were used as models

to examine the efficacy of GHP-88310 and other GHP-88309 analogs against HPIV3, SeV, and CDV.

Viruses were administered through intranasal inoculation and virus load was monitored continuously in

nasal lavages (ferrets only), and in respiratory tissues of cotton rats and mice extracted when indicated

after infection. Virus titers were determined through $TCID_{50}$-titration/ plaque assays.

Cells, viruses, and compound synthesis

Vero E6, Vero E6 TMPRSS2, Vero cells stably expressing canine signaling lymphocyte activation

molecule (Vero c-SLAM, VDS), Vero-hSLAM, and BHK-T7 cells (BHK cells stably expressing T7) were

cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 7.5% heat-inactivated

fetal bovine serum (FBS) at 37°C and 5% CO_2. Cells were tested regularly for mycoplasma

contamination prior to usage. Recombinant SeV-GFP, rCDV 5804p, HPIV3 (clinical isolate 13J5,

HPIV3/Seattle/USA/13J5/2016; KY629774.1), rPIV3-nano (JS strain), rMeV (Edmonston strain), and

rCedar virus (CedV)-nano were propagated on Vero E6 TMPRSS2 or VDS cells using DMEM with

7.5% serum, scraped when necessary, clarified, aliquoted, and stored at −80°C. The generation of all

GHP-88309 resistant recombinant viruses used in these studies has been previously described (*23*).

Virus stock titers were determined through $TCID_{50}$-titration (rSeV, rCDV, rMeV, and rCedV) or plaque

assay (PIV3), and stocks were stored in aliquots at -80°C. All virus stocks were sequence confirmed

prior to usage. GHP-88310 and other GHP-88309 analogs were authenticated through elemental

analysis (Table S1) and stored as dry powder (supplementary information. Working aliquots were

dissolved in DMSO or 1% methylcellulose for *in vitro* and *in vivo* studies, respectively.

Primary cells

Undifferentiated cells: Normal human bronchial/tracheal epithelial cells (Lonza CC-2540S, lot

number 519670, Caucasian, 42-year-old female donor "F3", unknown cause of death, smoker) were

amplified and expanded in lifeline cell culture media (LIFELINE Cell Technology) for no more than 3

passages. Cells were seeded in 24-well plates for experiments.

Differentiated ALI HAE cells (MatTek): EpiAirways (AIR-100; 23-year-old healthy male, Caucasian;

grown at air-liquid interface for 13 days) were equilibrated after arrival and further differentiated in

hanging-top plates (following manufacturers protocol) for 14-21 days prior to use in experiments. ALI

HAE cells were fed every two days 6 ml maintenance media (MatTek), rinsed weekly and before a

procedure (MatTek, TEER buffer). Transepithelial/transendothelial electrical resistance (TEER) was

measured on selected inserts using an EVOM3 (World Precision Instruments) following manufacturers

protocol.

Virus titration

TCID$_{50}$: Samples were serially diluted (10-fold starting at 1:10 initial dilution) in serum-free DMEM.

Serial dilutions were added to either VDS (rCDV), Vero hSLAM (rMeV), BHKT7 (rCedV), or Vero E6

TMPRSS2 (rSeV) cells seeded in 96-well plates at 1×10^5 cells/ml, 14 hours before infection. Infected

plates were incubated for 72-96 hours at 37°C with 5% CO_2, and titers were determined (Reed and

Muench).

Plaque assays: HPIV3 virus samples were serially diluted (10-fold starting at 1:10 initial dilution) in

serum-free DMEM. Serial dilutions were added to Vero E6 TMPRSS2 cells seeded in 12-well plates at

2×10^5 cells per well, 20 hours before infection. Ninety minutes post infection, cells were overlaid with 1

ml of 2× DMEM and avicel (1:1 ratio). Infected plates were incubated for 84 hours at 37°C with 5% CO_2.

Upon incubation, cells were washed twice with PBS, stained with 1% crystal violet (in 20% EtOH), and

plaques were enumerated.

Dose-response and minigenome antiviral assays and cytotoxicity assessment

Compound stocks were prepared in dimethyl sulfoxide (DMSO) and diluted in cell culture media

(final DMSO concentration of less than 0.1% in all wells). For luciferase-based dose-response assays,

cells were seeded in white-walled 96-well plates one day prior to the experiment to achieve 50%

confluency. Threefold serial dilutions of compounds were prepared in triplicate using an automated

Nimbus liquid handler (Hamilton) and transferred to the cells. Immediately after the addition of

compound, cells were infected with virus as previously described (*54*). Each plate contained four wells

of positive and negative control (media containing 100 μM cycloheximide or vehicle (DMSO),

respectively). Luciferase activities were determined 48 hours after infection using Nano-GLO buffer

(Promega) and an H1 Synergy plate reader (Biotek). Normalized luciferase activities were analyzed

with the following formula: % inhibition = (Signal$_{sample}$ – Signal$_{min}$)/(Signal$_{max}$ – Signal$_{min}$) × 100, and

dose-response curves were further analyzed by normalized nonlinear regression with variable hill slope

to determine EC$_{50}$ and EC$_{90}$.

BSR-T7/5 cells (5×10^4 per well in a 96-well plate format) were transfected with plasmids encoding

for P-ICB (0.01 µg), N-ICB (0.01 µg), L-ICB (0.0075 µg) and L-ICB mutant variants (0.0125 µg), and the

MeV luciferase replicon reporter (0.02 µg) using GeneJuice transfection reagent. Cells were lysed 42

hours after transfection with OneGlo substrate (Promega), which was directly added to the cells, and

bioluminescence was quantified using the above-mentioned formula.

Virus yield reduction assays with CDV and SeV: Cells were seeded in 24-well plates. Once

reaching 90% confluency, cells were overlaid with compound dilutions (three-fold dilutions, in triplicate)

and cells were infected at an MOI of 0.1. CDV: After 28 hours of incubation, the supernatant was

removed, and cells were scraped in 300 µl media, subjected to two freeze-thaw cycles, clarified, and

the resulting supernatant was collected for titration. SeV: Supernatant was collected after 52 hours and

clarified. Samples were titrated and scored using the $TCID_{50}$ method after 4 days.

Virus yield reduction assays in undifferentiated NHBE cells: Cells were seeded at 60% confluency

in a 24-well plate. Upon reaching 80% confluency, cells were overlaid with compound dilutions (50 µM,

three-fold dilutions, in triplicate) and were infected at an MOI of 0.1 with HPIV3 (clinical isolate 13J5).

Supernatant was collected after 48 hours and clarified. Samples were titrated by plaque assay on Vero

E6 TMPRSS2 cells and quantified after 4 days.

Virus yield reduction assays in EpiAirways: Inserts were rinsed using DPBS and infected on the

apical side with 10,000 PFU of HPIV3 (clinical isolate 13J5) per well in 100 µl PBS. Sixty minutes after

infection, virus was aspirated to allow for air-liquid exchange. Compound dilutions (50 µM, three-fold

dilutions, in triplicate) were added on the basolateral side at the time of infection and ALI HAE cells

were incubated for 48 hours (after 24 hours, the basolateral media was exchanged - same static

compound concentration). After 20 hours of initial incubation, the apical surface was washed once to

remove accumulated mucus. Apical shed was harvested in 300 µl PBS that had been incubated at

37°C for 30 min. Clarified samples were titrated by plaque assay on Vero E6 TMPRSS2 cells and

measured after 4 days.

To assess the impact of the GHP-compound class on cell metabolic activity, Vero E6 cells were

seeded at 6,000 cells per well in a 96-well plate and incubated at 37°C (5% CO_2) following the addition

of a 3-fold serial dilution of each compound, starting at 150 µM. After 72 hours of incubation, cells were

incubated with 10 µl per well of PrestoBlue (ThermoFisher Scientific) for 1 hour at 37°C and

fluorescence was measured using an H1 synergy plate reader (Biotek) with excitation and emission

wavelengths set at 560 nm and 590 nm, respectively. Cellular metabolic activity as a surrogate for cell

viability was expressed as a percentage relative to the media control.

To determine the effect of GHP-88310 on cell metabolic activity in primary cells, three to four ALI

HAE cultures were exposed to increasing concentrations of GHP-88310 (25 µM - 200 µM) in

maintenance media (MatTek). To assess barrier integrity, TEER was measured after 48 hours of

incubation at 37°C with 5% CO_2 and immediately prior to compound addition to the basolateral

chamber of a 12-well hanging insert plate (MatTek). The measurements are displayed in Ohm $\times$ cm^2.

Confocal microscopy

PIV3- and mock-infected ALI HAE cells were infected and treated with GHP-88310 at either 0.5

µM or 10 µM as above. Forty-eight hours after infection the apical shed was harvested, and the cells

were prepared for confocal microscopy. All incubation steps of immunostaining were performed at room

temperature and in the absence of light as previously described (*47*). Briefly, the inserts were fixed for

40 min with 4% paraformaldehyde-PBS, washed once with PBS, cut, and mounted on a chamber slide

(Fisher, cat# 154461PK). Cells were permeabilized with 500 µl of 0.5% Triton X-100 in PBS for 2 hours,

then washed, and blocked with 500 µl per insert of 5% BSA in PBS containing 0.05% Tween-20 for 1

hour. Inserts were washed and stained with the primary antibody in an antibody-binding solution (2%

BSA-PBS) for 1 hour (PIV3: goat anti-HPIV3 HN, Abcam ab28584; TJ: mouse anti-ZO-1 BD,

Biosciences 610966; Mucin/Goblet cells: mouse anti-MUC5AC, ThermoFisher MA5-12175). Cells were

washed three times and incubated with a conjugated antibody for 45 min (donkey anti-goat Alexa Fluor

568 ThermoFisher A-11057; goat anti-mouse IgG (H+L) highly cross-adsorbed secondary antibody,

Alexa Fluor 488, Invitrogen, A-11029). Cells were washed an additional three times in PBS and then

incubated for 5 min with Hoechst 34580. Following three additional washes, cells were mounted with

coverslips using a drop of ProLong™ Diamond Antifade Reagent (Thermo Scientific, Cat# P36970) and

stored overnight.

Confocal Imaging was performed using the Zeiss Axio Observer Z.1 and an LSM 800 confocal

microscope with AiryScan, controlled by the Zeiss Zen 3.1 Blue software (Windows 10). This software

as well as Adobe Photoshop (25.9.0) were employed for image analysis.

Single-dose pharmacokinetic property profiling in mice, cotton rats, ferrets, and dogs

Mice: Eight-week-old female Balb/cJ mice (The Jackson Laboratory) were acclimated for at least

three days prior to study initiation. Mice were weighed and administered 20 mg/kg of GHP-class

compounds by oral gavage in 1% methylcellulose (200 µl total volume). Blood samples (150 µl) were

collected retro-orbitally at designated time points post dosing, centrifuged at 2,000 rpm for 5 minutes at

4°C, and the resulting plasma was stored at -80°C.

Cotton rats: Female cotton rats (Envigo), aged 2-3 months, were rested for at least three days prior

to study start. Animals were weighed and dosed via oral gavage with 50 mg/kg of GHP-88309 or GHP-

88310, formulated in 1% methylcellulose (500 µl total volume). Blood samples (150 µl) were collected

at 30 minutes, and at 1, 2, 3, 4 and 8 hours post dosing. Plasma was processed as described above. At

8 hours post-dosing, organs (lung, brain, spleen, heart, kidney, small intestine, large intestine, and liver)

were harvested and immediately frozen in liquid nitrogen.

Ferrets: Female ferrets (Triple F Farm), 4-8 months of age, were acclimated for a minimum of three

days prior to dosing. Animals were weighed and administered GHP-88309 or GHP-88310 via oral

gavage at doses of 150 mg/kg (GHP-88309) and 150 or 500 mg/kg (GHP-88310). Both compounds

were prepared in 1% methylcellulose, with a final gavage volume of 2 ml. Blood samples (200 µl) were

collected at 0.5, 1, 2, 3, 4, 6, 8, 12, and 24 hours post-dose and processed as described previously.

Dogs: Male and female beagles, aged 10-12 months old, were weighed and administered GHP-

88310 orally at 50, 150, or 500 mg/kg, formulated in 1% methylcellulose (study conducted by SRI

Biosciences™). Blood samples (300 µl) were collected at pre-dose, 0.5, 1, 2, 4, 6, 8, 12, and 24 hours

post-dosing for analysis. Clinical observations were conducted at pre-dose, immediately after dosing, 2-

4 hours post-dose, and again at 24 hours. The same four animals were used across all dose levels,

with a minimum washout period of seven days between each dose escalation.

For analysis, aliquots of animal plasma were mixed with 70% acetonitrile in water that included an

internal standard. Samples were then vortexed for one minute and collected at 15,000 rpm for 5

minutes. The resulting supernatants were transferred to HPLC vials for analysis by LC-MS/MS at 4°C.

HPLC separation was performed on an Agilent 1260 system (Agilent Technologies, Santa Clara, CA,

USA) equipped with an autosampler, column oven, UV lamp, and binary pump.

For the mouse PK study screening, a Zorbax SB-Phenyl (50x4.6mm, 3.5 µm) column (Waters) was

used. Mobile phase A consisted of 20 mM ammonium formate and mobile phase B consisted of

acetonitrile. A 2-minute isocratic HPLC method was used at 40% mobile phase A. For the remaining PK

studies, an XTerra MS C18 (50x2.1mm, 5 µm) column (Waters) was used. Mobile phase A consisted of

25 mM ammonium bicarbonate in HPLC grade water pH adjusted to 9.8 and mobile phase B consisted

of acetonitrile. A 2-minute isocratic HPLC method was used at 75 or 80% mobile phase A. MS analysis

was performed on a QTrap 7500 or Triple 7500 Mass Spectrometer (Sciex, Framingham, MA, USA) or

using positive electrospray ionization (ESI) in multiple reaction monitoring (MRM) mode. Data analysis

was performed using SciexOS Software (Sciex). PK parameters were calculated using Phoenix

WinNonLin 8.4 (Build 8.4.2.6172; Certara, Princeton, NJ) using the non-compartmental analysis tool.

Samples of frozen animal tissue were homogenized with 4°C 70% acetonitrile in water that

included an internal standard. Homogenate were collected for 5 minutes at 10,000 rpm and

supernatants subjected to a second clearance step (5 minutes, 15,000 rpm). The remaining

supernatant was analyzed via LC/MS/MS, maintaining samples at 4°C. HPLC separation was

performed on an Agilent 1260 system (Agilent Technologies) equipped with an autosampler, column

oven, UV lamp, and binary pump, using an Xterra MS C18 (50 x 2.1 mm, 5 µm) column (Waters).

Mobile phase A consisted of 25 mM ammonium bicarbonate buffer in HPLC grade water, pH 9.8, and

mobile phase B consisted of acetonitrile. A 4.5-minute gradient HPLC method was performed to

separate the analytes beginning with a 1-minute isocratic hold at 10% B, followed by a 2-minute

gradient to 30% B, and a return to starting conditions for 1.5 minutes. MS analysis was performed on a

ZenoTOF 7600 Mass Spectrometer (Sciex) using positive ESI in multiple reaction monitoring (MRM)

mode. Data analysis was performed using SciexOS Software (Sciex).

Single-dose GHP-88309 tolerability in ferrets

Female ferrets (Triple F Farm), between 6-8 months old, were acclimated for at least three days.

Ferrets were weighed and dosed with GHP-88309 once via oral gavage at doses ranging from 50-

1,000mg/kg. The compound was formulated in 1% methylcellulose and a final gavage volume of 2 ml

was administered. Rectal temperature and clinical scores were measured before dosing and at 30

minutes, as well as 1, 2, 3, 4, 5, 6, 8, 12, and 24 hours post-dose.

Clinical scoring evaluated: i) temperature to monitor hypothermia and ii) neurological status to

detect signs of acute drug toxicity. Temperature scoring was assigned as follows: 0, >38°C; 1, 37.5-

37.9°C; 2, 37-37.4°C; 3, <37°C. To assess neurological status, each ferret was individually placed in a

single-ventilated cage at the time of evaluation to observe its response to a novel environment.

Neurological scoring was partially adapted from the Merck Veterinary Manual (*55*) and assigned as

follows: 0, explores objects, weasel war dance(*56*), intact reflexes (pinch test); 1, decreased interest in

exploring objects; 2, lethargic, and/or altered gait, and/or decreased reflexes; 3, obtunded, and/or

seizures, and/or paralysis in one or more limbs. Upon reaching the predetermined endpoint, a clinical

score of 3 in either category, ferrets were euthanized.

Multidose tolerability studies in cotton rats and ferrets

Cotton rats: Female cotton rats (Envigo), aged 2-3 months were habituated for at least three days

prior to the beginning of the study. Cotton rats were weighed and administered GHP-88310 twice daily

by oral gavage at doses of 300, 600 and 1,200 mg/kg. The compound was formulated in 0.5%

methylcellulose, 5 mM sodium citrate, and 0.25% Tween80, with a final gavage volume of 500 µl. Each

animal received a total of 14 doses and was monitored daily for bodyweight and temperature using

implantable electronic ID transponders (TP-1000, BMDS). Following the final dose, blood samples (150

µl) were collected at designated time points, centrifuged at 2,000 rpm for 5 minutes at 4°C, and the

plasma was stored at -80°C.

Ferrets: Female ferrets (Triple F Farm), between 6-8 months old, were allowed to rest for at least

three days. Ferrets were weighed and dosed with GHP-88309 or GHP-88310 twice daily via oral

gavage at doses ranging from 50-1,000 mg/kg. The compounds were formulated in 1% methylcellulose,

with a final gavage volume of 2 ml. Animals were weighed daily, and body temperature was measured

using a rectal probe. Ferrets received a total of 14 doses under a twice-daily dosing regimen or 7 doses

under a once-daily regimen. Blood samples were collected on days 0, 2, 4, 5, 6 and 7 post-dosing for

complete blood cell (CBC) analysis, and on days 1 or 2 and 7 for blood chemistry analysis. CBC

analysis was performed using a Vetscan HM5 hematology analyzer (Abaxis), following the

manufacturer's protocol. For the 500 mg/kg and 1,000 mg/kg dosing groups, as well as the multi-dose

once-daily dosing study, maximum plasma concentration (C_{max}) at 5 hours post-dose was measured

after doses 1, 2, 4, 6, 8, 10, 12, and 14, or daily (once-daily regimen only). Blood samples (150 µl) were

collected for plasma preparation as previously described.

***In vivo* infections with SeV, HPIV3 and CDV**

Efficacy studies in mice: Female 129x1/SvJ mice (The Jackson Laboratory), 6 weeks of age, were

acclimated for at least three days before being randomly assigned to study groups and housed under

ABSL-2 conditions for infections with sequence-validated rSeV. Bodyweight and rectal temperature

were measured once daily. For infection, mice were anesthetized with isoflurane and intranasally

inoculated with 5×10^6 $TCID_{50}$ units/ml of rSeV (*23*) in a total volume of 50 µl PBS (25 µl per nare).

Treatment was initiated 24 hours post-infection and administered twice-daily via oral gavage at a dose

of 150 mg/kg in 200 µl of 1% methylcellulose, continuing through study day 8. A subset of animals was

euthanized either five days post-infection or upon reaching a predetermined endpoint, and tracheas

and lungs were collected for analysis.

Efficacy studies in cotton rats: Female cotton rats (Envigo), 3-4 weeks of age, were allowed to rest

for at least three days prior to being randomly assigned to study groups and single-housed under

ABSL-2 conditions for infections with sequence-validated HPIV3 clinical isolate 13J5. Bodyweight and

body temperature, using an implanted electronic ID transponder (TP-1000, BMDS), were measured

once daily. For infection, cotton rats were anesthetized with isoflurane and intranasally inoculated with

1×10^7 pfu/ml of HPIV3 in a total volume of 100 µl PBS (50 µl per nare). For therapeutic dosing, animals

were anesthetized with isoflurane 12 hours post-infection and administered GHP-88309 or GHP-88310

in 500 µl of 1% methylcellulose via oral gavage, either once or twice-daily, through study day 3.

Efficacy studies in ferrets: CDV unvaccinated female ferrets (Triple F Farms), 8 months of age,

were rested for one week prior to random assignment into study groups and housed in groups of three

animals under ABSL-2 condition. Bodyweight and rectal temperature were recorded once daily. Ferrets

were anesthetized with dexmedetomidine/ketamine and intranasally inoculated with 2×10^5 $TCID_{50}$ units

of CDV 5804p in a total volume of 200 µl PBS (100 µl per nare) (*25*). Daily, nasal lavages were

performed using 1 ml of PBS supplemented with $2\times$ antibiotics-antimycotics (Gibco). Peripheral blood

mononuclear cells (PBMCs) were collected twice weekly for titration of cell-associated viremia and CBC

analysis. On days 17, 21, and 35 post-infection blood samples were collected to assess neutralizing

antibody titers. For therapeutic treatments, ferrets were administered GHP-88310 by oral gavage

following either a once or twice-daily regimen (50-300 mg/kg) in 2 ml of 1% methylcellulose, beginning

at 3-days post-infection. Ferrets were euthanized either upon reaching a predetermined study endpoint

or at the conclusion of the study on day 35 post-infection.

Virus titration of tissue samples (mice and cotton rats) and nasal lavages (ferrets only)

Organs were weighed and homogenized in 300 µl PBS using a Bead Blaster 24R (Benchmark) set

to three 30 second cycles at 4°C, with one minute rest intervals between cycles. Homogenates and

ferret lavage samples were clarified by centrifugation at $13,000 \times g$ for 10 minutes at 4°C, then

subsequently titrated. Viral titers were expressed as $TCID_{50}$ units/plaques per gram input tissue or per

ml.

Virus titration of PBMCs (ferrets only)

Blood samples were collected at specified time points twice per week. As previously described

(*25*), viremia was assessed by isolating PBMCs from 1 ml of whole blood. Red blood cells were lysed

using ACK buffer (150 mM NH_4Cl, 10 mM $KHCO_3$, 10 µM EDTA pH 7.4) for 7 minutes, followed by two

PBS washes and resuspension in DMEM. PBMC titers were determined by co-culturing serial dilutions

of purified PBMCs with VDS cells and expressed as $TCID_{50}$ per 10^6 PBMCs.

Neutralizing antibody assay (ferrets only)

Blood samples (300 µl) were collected on days 17 and 21 post-CDV infection (and on day 35 for

the ferret CDV QD study only) to obtain plasma, which was subjected to heat inactivation (56°C, 30

minutes) and clearance centrifugation (4,000 $\times$ g, 5 minutes). Heat-inactivated and clarified plasma was

diluted two-fold in serum free DMEM, with a final volume of 50 µl per well. Fifty microliters of rCDV

5804p (2×10^3 TCID$_{50}$ units) was added to the plasma dilutions, followed by incubation of the plasma-

virus mixture at 37°C for 75 minutes. Subsequently, the mixture was added in duplicate to VDS cells in

96-well plates, incubated for 3 days, and cytopathic effect was assessed by light microscopy.

Whole genome sequencing of virus stocks and ferret nasal lavage samples

Whole genome sequencing (WGS) was performed using metagenomic next-generation

sequencing (mNGS) as described previously (*57*). Briefly, extracted RNA was treated with a TURBO

DNA-free Kit (Thermo Fisher, product #AM1907) to remove DNA. RNA was then reverse transcribed

using random hexamers (Thermo Fisher, product #N8080127) and Superscript IV (Thermo Fisher,

product #18090010), following the manufacturer's protocol. Double-stranded cDNA synthesis was

performed using Sequenase v2.0 (Thermo Fisher, product #70775Z1000UN), and the resulting cDNA

was purified using 1.8$\times$ AMPure XP magnetic beads (Beckman Coulter, product # A63882).

Metagenomic viral WGS libraries were created from purified double-stranded cDNA with

tagmentation reagents from the Illumina DNA Prep with Enrichment Kit (Illumina, product #20025524),

followed by 14 cycles of dual-indexed PCR for SeV and 18 cycles for HPIV3 and CDV isolates.

Amplified libraries were cleaned with 0.8$\times$ AMPure XP magnetic beads. Viral isolate libraries were not

subjected to enrichment and proceeded directly to sequencing. Libraries were sequenced 2$\times$150 bp on

NextSeq 2000 (CDV, HPIV3) or NovaSeq 6000 (SeV).

Due to its high viral titer, the sample "CDV ferret study - BID treatment – 50 mg/kg - ferret 2 – D8pI"

was sequenced using metagenomic WGS on NextSeq 2000 with 1$\times$100 bp read format.

Viral WGS was also performed on all ferret nasal lavage samples with detectable PFU titers,

except "CDV ferret study - BID treatment – 50 mg/kg - ferret 2 – D8pI". These samples were processed

using QIAseq xHYB Microbial Hyb&Lib Kit A (Qiagen, product #334525) and a custom Qiagen

hybridization panel containing probes designed from CDV strain 5804 (AY386315.1) containing eGFP

reporter (Qiagen, product # 334586). Following the manufacturer's protocol, extracted RNA was

depleted of ribosomal RNA and converted to double-stranded cDNA. Double-stranded cDNA was then

enzymatically fragmented, end-repaired, indexed, purified with 0.9× and 1.1× bead cleanup, and

amplified with 14 cycles of PCR, followed by a final bead purification, according to the manufacturer's

protocol.

Pre-capture sequencing libraries were pooled based on PFU titer, with one to four samples per

pool, and incubated with the custom capture panel. After overnight hybridization with biotinylated

probes, probe-target hybrids were bound to streptavidin-coated magnetic beads and washed to remove

unbound library fragments. The enriched libraries were amplified with 20 cycles of post-hybridization

PCR and purified via 1.1× bead clean-up. Post-amplification library fragment sizes were estimated with

the TapeStation 4200 D1000 (Agilent), and concentrations were measured with Qubit 4 Fluorometer

and Qubit dsDNA HS Assay Kit. Libraries were sequenced on Illumina NextSeq 2000 instruments using

a 1×100 bp read format.

Data analysis: Raw reads were trimmed and quality filtered using fastp (v0.23.4)(58) with the

following parameters: --cut_mean_quality 20 --cut_front --cut_tail --length_required 20 --

low_complexity_filter --trim_poly_g --trim_poly_x. The consensus sequence of "recCDV_5804p WT"

was generated by majority voting with Geneious Prime 2025.0, using the CDV plasmid sequence

(AY386315.1 background) containing eGFP reporter as a reference.

Variants were identified using RAVA workflow with default parameters

(https://github.com/greninger-lab/RAVA_Pipeline/tree/publication_GSU_2025_03_01)(59), (60) .

References used in RAVA analysis: for "recCDV_5804p WT" we used plasmid map as a reference

(AY386315.1 background), for the "HPIV3 clinical isolate 13J5" – HPIV3/Seattle/USA/13J5/2016

(KY629774.1), for "recSeV_GFP" – Respirovirus muris strain 52 (MH557085.1), and, finally, for CDV

ferret nasal lavage samples – "recCDV_5804p WT" consensus sequence. Sample "CDV ferret study -

BID treatment – 50 mg/kg - ferret 2 – D8pI" had insufficient coverage (mean coverage of ~49x with

some CDS regions having 0x coverage) and was therefore excluded from the RAVA analysis.

Complete lists of identified non-synonymous and complex SNVs (allelic frequency ≥ 5%, sequencing

depth of the nucleotide position ≥ 30) can be found in the supplementary data file S1. Data availability:

raw sequencing data is publicly available in NCBI BioProject PRJNA1217449.

***In silico* docking**

Docking studies were performed with MOE (version 2024.06). A structure of MeV L (Protein Data

Bank (PDB) 9dus) was used for the docking studies. After protonation and energy minimization, the

triangle matcher method was used for initial compound placement, followed by refinement using an

induced-fit protocol to dock GHP-88310 or GHP-88309 in the L structure based on resistance data

information. For MeV L, residues Y942, A866, and I1009 were selected to identify the target site for

docking and resulting docking poses from each polymerase target were compared. The top scoring

docking poses conserved between GHP-88309 and GHP-88310 were chosen for further analysis.

Cellular uptake and clearance in undifferentiated NHBE cells

Primary normal HBTECs (donor "F3") were seeded in 48 well plates and utilized for a wash-

in/wash-out study once confluency was achieved.

Wash-in study: An 80 µM dilution of GHP-88310 was prepared in pre-warmed Lifeline media.

Following media removal from the cells, 0.5 ml of either the 80 µM compound or vehicle (volume

equivalent DMSO)-media mixture was added to the wells. Plates were incubated at 37°C, 5% CO_2 and

extracted in triplicate at the indicated time points. Cells were washed twice with cold DPBS, followed by

the addition of 250 µl of 70% methanol/30% water spiked with the internal standard to each well treated

with GHP-88310. A mock-treated blank plate (16 well) was extracted with 250 µl of 70% methanol/30%

water (no internal standard) per well at one-hour post-dosing. All samples were mixed by pipetting and

transferred to 1.5 ml microcentrifuge tubes. After centrifugation at 16,000 $\times$ g for 10 minutes at 4°C, the

supernatants were transferred to 2 ml tubes and stored at -80°C until LC/MS/MS analysis.

Wash-out study: An 80 µM dilution of GHP-88310 was prepared in pre-warmed lifeline media. After

media removal, 0.5 ml of the 80 µM compound-media mixture was added to the wells. Plates were

incubated at 37°C, 5% CO_2 for one hour, washed twice with 0.5 ml warm DPBS, and compound free

pre-warmed Lifeline media was added. Plates were further incubated as specified, followed by washing

(twice with 0.5 ml cold DPBS) and extraction, processing, and analysis as described above.

Recapitulation of ferret plasma PK profile in EpiAirways

Air liquid interface HAE cells were exposed basolaterally to dynamic GHP-88310 concentrations

reflected ferret plasma levels after either 50 mg/kg oral twice-daily dosing on day 7 or a single oral dose

of 20 mg/kg. All ALI HAE cell cultures (n=4 per condition: vehicle, 50 or 20 mg/kg bid, and 50 or 20

mg/kg qd) were infected with 1×10^4 PFU per well of the HPIV3 clinical isolate 13J5 one hour prior to

treatment. Infected cultures underwent either four 12-hour treatment cycles (bid) or two 24-hour

treatment cycles (qd). Twenty hours post infection the apical side was rinsed once with 400 µl PBS to

reduce mucus accumulation. After 48 hours post-infection, apically shed virus was collected by

incubating the cultures with 300 µl of DPBS at 37°C for 30 minutes. Viral titers were then determined by

plaque assay on Vero E6 TMPRSS2 cells.

Statistical analysis and data preparation

For statistical analysis of studies comparing more than two study groups, a one-way analysis of

variance (ANOVA) or two-way ANOVA with multiple comparison *post hoc* tests as specified were used

to assess statistical difference between samples. An unpaired, two-tailed t-test was used to compare

compound concentrations across the analyzed organs. Statistical analyses were performed using

GraphPad Prism (v10.1.0). All figures were assembled in GraphPad Prism and Adobe

Illustrator/Photoshop. Study schematics were created using Adobe Illustrator, with animal illustrations

generated via the vector tool. Objects in figures 5 and 6 were drawn in Adobe Illustrator and adapted

from BioRender. The number of individual biological replicates (n values) and exact P values are shown

in the figures. The threshold of statistical significance (α) was set to 0.05. Source data and statistical

analysis are shown in the source data (supplementary data file 2) and statistical analysis

(supplementary data file 3) files, respectively**.**

Ethical compliance

All animal work was performed in compliance with the *Guide for the Care and Use of Laboratory*

Animals of the National Institutes of Health and the Animal Welfare Act Code of Federal Regulations.

Experiments involving mice, cotton rats and ferrets were approved by the Georgia State University

Institutional Animal Care and Use Committee (IACUC) under protocols A23011, A24022 and A22035,

respectively. All experiments using infectious material were approved by the Georgia State University

Institutional Biosafety Committee (IBCs) and performed in BSL-2/ABSL-2 containment facilities,

respectively.

Materials & Correspondence

Correspondence and material requests should be addressed to R.K. Plemper, rplemper@gsu.edu

List of Supplementary Materials

Table S1. Elemental analysis of selected GHP-88309 analogs

Figure S1. Multi-dose tolerability of GHP-88309 in ferrets

Figure S2. Clinical parameters of SeV infection in mice

Figure S3. Clinical parameters and viral titers of cotton rats infected with HPIV3

Figure S4. Multi-dose tolerability in cotton rats

Figure S5. Once daily administration of GHP-88310 against HPIV3 in cotton rats

Figure S6. Multi-dose tolerability and CBC parameters in ferrets

Figure S7. Clinical parameters of multi-dose GHP-88309 and GHP-88310 tolerability

Figure S8. Evaluation of once- and twice-daily GHP-88310 dosing for CDV infection in ferrets

Figure S9. *In silico* docking of GHP-88309

Figure S10. 2D docking poses

Figure S11. Cellular uptake and retention of GHP-88310

Figure S12. Confocal imaging of GHP-88310 antiviral activity in human airway epithelium organoids

References and Notes
1.   R. K. Plemper, R. A. Lamb, in *Fields Virology,* D. M. Knipe, P. M. Howley, S. Whelan, Eds.
(Wolters Kluwer/Lippincott Williams & Wilkins, Philadelphia, 2020), vol. 1, chap. 33, pp. 504-558.
2.   B. Lee, C. C. Broder, L. Wang, in *Fields Virology,* D. M. Knipe, P. M. Howley, S. Whelan, Eds.
(Wolters Kluwer/Lippincott Williams & Wilkins, Philadelphia, 2200), vol. 1, chap. 34, pp. 559-595.
3.   A. J. Wakefield *et al.*, RETRACTED: Ileal-lymphoid-nodular hyperplasia, non-specific colitis, and
pervasive developmental disorder in children. *The Lancet* **351**, 637-641 (1998).
4.   L. Eggertson, Lancet retracts 12-year-old article linking autism to MMR vaccines. *Cmaj* **182**,
E199-200 (2010).
5.   H. J. Larson, E. Gakidou, C. J. L. Murray, The Vaccine-Hesitant Moment. *N Engl J Med* **387**, 58-
65 (2022).
6.   L. A. H. Do, K. Mulholland, Measles 2025. *N Engl J Med*, (2025).
7.   CDC. (2025), vol. 2025.
8.   W. H. Organization, in *Disease Outbreak News*. (2025), vol. 2025.

9.   G. o. Canada, in *Weekly surveillance reports and monitoring maps for measles and rubella.*
756   (2025), vol. 2025.
757   10.   NMA Statement on Measles Outbreak – Children's Lives in Danger in West Texas.
758   https://nmanet.org/press-release/nma-statement-on-measles-outbreak-childrens-lives-in-
759   danger-in-west-texas/ (accessed 02/27/2025). (2025).
760   11.   R. K. Plemper, Measles Resurgence and Drug Development. *Curr. Opin. Virol.* **41**, 8-17 (2020).
761   12.   D. V. Parums, A Review of the Resurgence of Measles, a Vaccine-Preventable Disease, as
762   Current Concerns Contrast with Past Hopes for Measles Elimination. *Med Sci Monit* **30**, e944436
763   (2024).
764   13.   S. S. Chen. (emedicine.medscape, 2025), vol. 2025.
765   14.   S. E. Kern, Challenges in conducting clinical trials in children: approaches for improving
766   performance. *Expert Rev Clin Pharmacol* **2**, 609-617 (2009).
767   15.   V. Bencheva, N. K. Mann, T. Rombey, D. Pieper, S. Schmiedl, Barriers and facilitators to
768   enrollment in pediatric clinical trials: an overview of systematic reviews. *Syst Rev* **13**, 283 (2024).
769   16.   C. Domingues, I. Jarak, F. Veiga, M. Dourado, A. Figueiras, Pediatric Drug Development:
770   Reviewing Challenges and Opportunities by Tracking Innovative Therapies. *Pharmaceutics.* 2023
771   (10.3390/pharmaceutics15102431).
772   17.   R. H. Parrott *et al.*, Clinical features of infection with hemadsorption viruses. *N. Engl. J. Med.* **260**,
773   731-738 (1959).
774   18.   R. C. Welliver, D. T. Wong, M. Sun, N. McCarthy, Parainfluenza virus bronchiolitis. Epidemiology
775   and pathogenesis. *Am. J. Dis. Child.* **140**, 34-40 (1986).
776   19.   A. J. Peck *et al.*, Respiratory virus infection among hematopoietic cell transplant recipients:
777   evidence for asymptomatic parainfluenza virus infection. *Blood* **110**, 1681-1688 (2007).
778   20.   W. G. Nichols, L. Corey, T. Gooley, C. Davis, M. Boeckh, Parainfluenza virus infections after
779   hematopoietic stem cell transplantation: risk factors, response to antiviral therapy, and effect on
780   transplant outcome. *Blood* **98**, 573-578 (2001).
781   21.   M. S. Lee, R. E. Walker, P. M. Mendelman, Medical burden of respiratory syncytial virus and
782   parainfluenza virus type 3 infection among US children. Implications for design of vaccine trials.
783   *Hum Vaccin* **1**, 6-11 (2005).
784   22.   G. R. Abedi *et al.*, Estimates of Parainfluenza Virus-Associated Hospitalizations and Cost Among
785   Children Aged Less Than 5 Years in the United States, 1998-2010. *J Pediatric Infect Dis Soc* **5**,
786   7-13 (2016).
787   23.   R. M. Cox *et al.*, Orally efficacious broad-spectrum allosteric inhibitor of paramyxovirus
788   polymerase. *Nat Microbiol* **5**, 1232-1246 (2020).
789   24.   G. D. Haas *et al.*, Tetracistronic Minigenomes Elucidate a Functional Promoter for Ghana Virus
790   and Unveils Cedar Virus Replicase Promiscuity for all Henipaviruses. *bioRxiv*, (2024).
791   25.   R. M. Cox *et al.*, Therapeutic mitigation of measles-like immune amnesia and exacerbated
792   disease after prior respiratory virus infections in ferrets. *Nat Commun* **15**, 1189 (2024).
793   26.   J. D. Wolf, M. R. Sirrine, R. M. Cox, R. K. Plemper, Structural basis of paramyxo- and
794   pneumovirus polymerase inhibition by non-nucleoside small-molecule antivirals. *Antimicrob*
795   *Agents Chemother* **68**, e0080024 (2024).
796   27.   P. A. Rota *et al.*, Measles. *Nature Reviews Disease Primers* **2**, 16049 (2016).
797   28.   A. T. Truong *et al.*, Genetic analysis of Asian measles virus strains – new endemic genotype in
798   Nepal. *Virus Research* **76**, 71-78 (2001).
799   29.   Y. Wang *et al.*, Structures of the measles virus polymerase complex with non-nucleoside inhibitors
800   and mechanism of inhibition. *Cell* **188**, 4913-4923.e4913 (2025).
801   30.   D. Wang, G. Yang, B. Liu, Structure of the measles virus ternary polymerase complex. *Nature*
802   *Communications* **16**, 3819 (2025).
803   31.   B. Srinivasan *et al.*, TEER measurement techniques for in vitro barrier model systems. *J Lab*
804   *Autom* **20**, 107-126 (2015).
805   32.   M. Malik, S. A. Steele, D. Mitra, C. J. Long, J. J. Hickman, Trans-epithelial/endothelial electrical
806   resistance (TEER): Current state of integrated TEER measurements in organ-on-a-chip devices.
807   *Current Opinion in Biomedical Engineering* **34**, 100588 (2025).

33. A. R. Calabro, R. Konsoula, F. A. Barile, Evaluation of in vitro cytotoxicity and paracellular
permeability of intact monolayers with mouse embryonic stem cells. *Toxicology in Vitro* **22**, 1273-
1284 (2008).
34. L. C. Rijsbergen *et al.*, Modeling Infection and Tropism of Human Parainfluenza Virus Type 3 in
Ferrets. *mBio* **13**, e0383121 (2021).
35. L. Zhang *et al.*, Infection of ciliated cells by human parainfluenza virus type 3 in an in vitro model
of human airway epithelium. *J Virol* **79**, 1113-1124 (2005).
36. B. Martin, S. Ji, S. Maudsley, M. P. Mattson, "Control" laboratory rodents are metabolically morbid:
why it matters. *Proc Natl Acad Sci U S A* **107**, 6127-6133 (2010).
37. C. Moro, C. Magnan, Revisited guidelines for metabolic tolerance tests in mice. *Lab Animal* **54**,
16-23 (2025).
38. J. K. Corrigan *et al.*, A big-data approach to understanding metabolic rate and response to obesity
in laboratory mice. *eLife* **9**, e53560 (2020).
39. M. Martignoni, G. M. Groothuis, R. de Kanter, Species differences between mouse, rat, dog,
monkey and human CYP-mediated drug metabolism, inhibition and induction. *Expert Opin Drug
Metab Toxicol* **2**, 875-894 (2006).
40. T. Nakajima *et al.*, Cytochrome P450-related differences between rats and mice in the metabolism
of benzene, toluene and trichloroethylene in liver microsomes. *Biochem Pharmacol* **45**, 1079-
1085 (1993).
41. U. M. Zanger, M. Schwab, Cytochrome P450 enzymes in drug metabolism: Regulation of gene
expression, enzyme activities, and impact of genetic variation. *Pharmacology & Therapeutics* **138**,
103-141 (2013).
42. R. Mezencev *et al.*, The association between histopathologic effects and liver weight changes
induced in mice and rats by chemical exposures: an analysis of the data from Toxicity Reference
Database (ToxRefDB). *Toxicol Sci* **200**, 404-413 (2024).
43. S. P. Jiang *et al.*, Airway Organoid Models as Pivotal Tools for Unraveling Molecular Mechanisms
and Therapeutic Targets in Respiratory Diseases: A Literature Review. *Ther Clin Risk Manag* **21**,
975-986 (2025).
44. H. Joo, S. Min, S. W. Cho, Advanced lung organoids for respiratory system and pulmonary
disease modeling. *J Tissue Eng* **15**, 20417314241232502 (2024).
45. J. H. Lee *et al.*, Development of robust antiviral assays using relevant apical-out human airway
organoids. *bioRxiv*, (2024).
46. C. T. Ekanger *et al.*, Comparison of air-liquid interface transwell and airway organoid models for
human respiratory virus infection studies. *Front Immunol* **16**, 1532144 (2025).
47. M. Toots *et al.*, Characterization of orally efficacious influenza drug with high resistance barrier in
ferrets and human airway epithelia. *Sci. Transl. Med.* **11**, (2019).
48. A. Nair, M. A. Morsy, S. Jacob, Dose translation between laboratory animals and human in
preclinical and clinical phases of drug development. *Drug Dev Res* **79**, 373-382 (2018).
49. S. Plumet, W. P. Duprex, D. Gerlier, Dynamics of viral RNA synthesis during measles virus
infection. *J. Virol.* **79**, 6900-6908 (2005).
50. T. Komatsu, K. Takeuchi, J. Yokoo, B. Gotoh, C and V proteins of Sendai virus target signaling
pathways leading to IRF-3 activation for the negative regulation of interferon-β production.
Virology **325**, 137-148 (2004).
51. A. Kato *et al.*, Y2, the smallest of the Sendai virus C proteins, is fully capable of both counteracting
the antiviral action of interferons and inhibiting viral RNA synthesis. *J Virol* **75**, 3802-3810 (2001).
52. B. Gotoh *et al.*, Knockout of the Sendai virus C gene eliminates the viral ability to prevent the
interferon-alpha/beta-mediated responses. *FEBS Lett* **459**, 205-210 (1999).
53. D. Garcin, P. Latorre, D. Kolakofsky, Sendai virus C proteins counteract the interferon-mediated
induction of an antiviral state. *J Virol* **73**, 6559-6565 (1999).
54. J. Sourimant *et al.*, 4'-Fluorouridine is an oral antiviral that blocks respiratory syncytial virus and
SARS-CoV-2 replication. *Science* **375**, 161-167 (2022).
55. W. B. Thomas. (2023), vol. 2025.
56. M. J. Goodfellow *et al.*, Behavioral assessment of well-being in the naïve laboratory ferret
(Mustela putorius furo). *Sci Rep* **14**, 30119 (2024).

57.   A. L. Greninger *et al.*, Rapid Metagenomic Next-Generation Sequencing during an Investigation
863   of Hospital-Acquired Human Parainfluenza Virus 3 Infections. *J. Clin. Microbiol.* **55**, 177-182
864   (2017).
865   58.   S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
866   *Bioinformatics* **34**, i884-i890 (2018).
867   59.   P. Di Tommaso *et al.*, Nextflow enables reproducible computational workflows. *Nat Biotechnol*
868   **35**, 316-319 (2017).
869   60.   M. J. Lin, R. C. Shean, N. Makhsous, A. L. Greninger, LAVA: a streamlined visualization tool for
870   longitudinal analysis of viral alleles. *bioRxiv*, 2019.2012.2017.879320 (2019).
871

Acknowledgements

We thank the Georgia State University Department of Animal Resources for expert assistance and

M. Sirrine for assistance. The recHPIV3 and recCDV-5804p genomic plasmids and Vero-canine SLAM

cells were kind gifts of B. Lee, V. von Messling, and Y. Yanagi, respectively. Blood chemistry analysis

was provided by the Pathology Core of Emory University. This study was supported, in part, by public

health service grants AI071002 (to R.K.P.), AI171403 project 3 (to R.K.P.), AI171403 Scientific Core B

(to M.G.N), AI171403 Scientific Core C, and AI171403 Scientific Core D (to A.L.G.) from the NIH/NIAID.

Competing interests

CML, MG, RMC, MGN, and RKP are co-inventors on a patent filing covering composition and method

of use of GHP-88310/EIDD-3608 for antiviral therapy. This study could affect their personal financial

status. RKP reports contract testing to GSU from Enanta Pharmaceuticals and Gilead Sciences, outside

of the described work. ALG reports contract testing to UW from Abbott, Cepheid, Novavax, Pfizer,

Janssen and Hologic, research support from Gilead, and personal fees from Arisan Therapeutics, outside

of the described work. All other authors declare that they have no competing interests to report.

Figures

Figure 1. SAR development of broad-spectrum GHP-class polymerase inhibitors. A) Study

schematic of multi-dose GHP-88309 tolerability and PK assessment. Ferrets received 14 oral doses of

GHP-88309 at two different concentrations, administered twice daily. **B)** Survival curves of animals

from (A). **C)** PK profile of GHP-88309 (50 mg/kg) in ferret plasma following 14 oral doses administered

b.i.d. Exposure is specified. **D-F)** Survival curves (D), body temperature (E), and clinical scores (F) of

animals after single oral dose of GHP-88309. Body temperature scoring: 0, >38°C; 1, 37.5-37.9°C; 2,

37-37.4°C; 3, <37°C. Neurologic scoring: 0, explores objects, weasel war dance, intact reflexes; 1,

decreased interest in exploring objects; 2, lethargic, and/or altered gait, and/or decreased reflexes; 3,

obtunded, and/or seizures, and/or paralysis in one or more limbs. **G)** Top-ranking GHP-88309 analogs.

Dashed circles mark regions that QSAR models (*23*) have identified as most amenable to synthetic

elaboration. **H)** Evaluation of cellular viability of GHP compounds on Vero-E6 cells. **I-K)** Recombinant

virus reporter-based dose response inhibition of HPIV3 (I), MeV (J), and CedV (K), n $\geq$3. **L)** Study

schematic of single-dose PK assessment in BALB/c mice. Mice received an oral dose of GHP-class (20

mg/kg) and were monitored over 24 hours. Blood was collected from each sub-group of mice twice,

following euthanasia. **M)** Single-dose PK profile in mouse plasma. Exposure is given as AUC_{0-8hrs} with

95% confidence intervals in parenthesis. Group sizes (n) specified in study schematics (A,L). Log-rank

(Mantel-Cox) test with median survival shown (B,D). Symbols represent geometric mean $\pm$ geometric

standard deviation (SD) (C,M). Variable slope 4-parameter non-linear regression modeling (H-K), n $\geq$3.

Symbols represent means $\pm$ SD. LLOQ, lower limit of quantitation. AUC, area under the curve.

Figure 2. PK and efficacy evaluation of GHP-88309 analogs in rodents. A) Study schematic of

compound efficacy testing in 129x1/SvJ mice. Mice were infected intranasally with 1.5×10^5 TCID$_{50}$ units

of recombinant SeV. Compounds were administered orally b.i.d., starting one day after infection and

continuing through 8.5 dpi. **B-D)** Survival curves (B), trachea (C) and lung (D) viral titers determined 5

dpi of animals shown in (A). **E)** Study overview. A direct comparison of GHP-88309 and GHP-88310

efficacy at varying concentrations was conducted in cotton rats. Animals were intranasally infected with

1×10^6 TCID$_{50}$ units of HPIV3 clinical isolate 13J5. Oral dosing was initiated 12 hours post infection and

continued b.i.d. **F,G)** Trachea (F) and combined left and right cranial lungs (G) were collected 3.5 days

after infection and viral titers determined by plaque assay. **H)** Study outline of single-dose PK and tissue

distribution analysis in cotton rats, comparing 50 mg/kg doses of GHP-88309 and GHP-88310. **I,J)** PK

profiles of GHP-88309 and GHP-88310 in cotton rat plasma (I) and **t**issue concentrations of GHP-

88309 and GHP-88310 measured 8 hours post dosing (J). Unpaired, two-tailed t-test. Bars indicate the

geometric mean $\pm$ geometric SD. **K)** Overview of a multi-dose GHP-88310 tolerability and PK study.

Cotton rats received 14 oral doses of varying concentrations, administered twice daily. **L,M)** Survival

curves (L) and PK profile (M) after 14 doses of GHP-88310. **N)** Summary overview of an efficacy

evaluation of GHP-88310 administered once daily at different concentrations in cotton rats. Animals

were intranasally infected with 1×10^6 TCID$_{50}$ units of HPIV3, and oral dosing started 12 hours after

infection. **O,P)** Trachea (O) and combined left and right cranial lung lobes (P) were collected 3.5 dpi

and viral titers determined by plaque assay. In (B,L), log-rank (Mantel-Cox) test with median survival. In

(C,D,F,G,O,P), bars indicate the geometric mean $\pm$ geometric SD, symbols represent individual

biological repeats. One-way ANOVA with Tukey's (between group comparisons) or Dunnet's

(comparisons to vehicle) post-hoc multiple comparisons test. In (I,M), symbols represent geometric

means $\pm$ geometric SD.

Figure 3. Single and multi-dose tolerability and PK of twice daily GHP-88310 in ferrets. A)

Overview of single oral dose PK study of GHP-88309 and GHP-88310 in ferrets. **B)** PK profile of GHP-

88309 and GHP-88310 in ferret plasma. **C)** Schematic of multi-dose tolerability and PK study of GHP-

88309 and GHP-88310 at different concentrations. Ferrets received 14 oral doses, administered b.i.d.

Maximal plasma concentration (C_{max}) was measured daily in ferrets administered 500 and 1,000 mg/kg

GHP-88310. **D)** Survival curves; data for the 150 mg/kg GHP-88309 group, originally shown in (Fig. 1A-

B, are included for direct comparison. Log-rank (Mantel-Cox) test with median survival. **E)** C_{max} of GHP-

88310 to monitor plasma build-up after doses 1, 2, 4, 6, 8, 10, 12, and 14. **F)** PK profile of GHP-88310

in ferret plasma after 14 oral doses. **G)** CBC analysis of ferret blood samples quantifying white blood

cells (left), red blood cells (middle), and red blood cell width (right). Normal range for each parameter is

shown in green shading. (Left, middle), symbols indicate the geometric mean $\pm$ geometric SD, (right)

symbols represent means $\pm$ SD. Two-way ANOVA followed by Dunnet's (comparisons to mock) post-

hoc multiple comparisons test. **H-J)** Single dose PK study of GHP-88310 in dogs. Shown are study

outline (H), neurologic status scores (I) (0, explores objects, intact reflexes; 1, decreased interest in

exploring objects; 2, lethargic, and/or altered gait, and/or decreased reflexes; 3, obtunded, and/or

seizures, and/or paralysis in one or more limbs), and PK profile of GHP-88310 in dog plasma (J). In (B,

E-F, J), symbols indicate geometric means $\pm$ geometric SD.

Figure 4. Therapeutic effect of GHP-88310 on CDV infection in ferrets. A) Virus yield reduction of

recCDV-5804p. Variable slope 4-parameter non-linear regression modeling; n ≥3. Symbols indicate the

geometric mean ± geometric SD. **B)** Overview of assessment of q.d. and b.i.d. GHP-88310 therapeutic

potential. Ferrets were infected intranasally with $2{\times}10^5$ $TCID_{50}$ units of recCDV-5804p. Oral

administration of GHP-88310 began 3 days after infection and continued until 10 (150 mg/kg) or 21 (50

mg/kg) dpi. **C)** Survival curves. Log-rank (Mantel-Cox) test with median survival. **D-F)** PBMC-

associated viremia (D), nasal lavage viral titers quantified by $TCID_{50}$ assay (E), and lymphocyte counts

(F) are shown. Normal range of lymphocytes is depicted by green shading. **G)** Neutralizing antibody

titers. In (D-F), symbols indicate individual animals and lines represent geometric means. Two-way

ANOVA followed by Dunnet's (comparisons to vehicle) post-hoc multiple comparisons test.

Figure 5. Functional and molecular profiling of GHP-88310. A-C) Dose-response assays with

recombinant virus variants or in a minigenome harboring mutations that mediate reduced susceptibility

to GHP-88309. Shown are 4-parameter variable slope non-linear regression curves for HPIV3 (A), SeV

(B), and in a MeV minigenome (C). Fifty and 90% inhibitory concentrations against each compound are

provided. Yellow highlights mark mutations that result in a $\geq$5-fold increase in EC_{50} and/or EC_{90}

compared to the unmodified genetic parent virus. **D)** *In silico* docking of GHP-88310. Ribbon

representation of the MeV L-P structure (PDBID: 9dus), highlighting the predicted docking location

(black spheres in red dashed box) of GHP-88310 (left). The RdRP and capping domains are colored in

cyan and green, respectively. Ribbon representation of the top scoring docking pose conserved

between GHP-88309 and GHP-88310 (middle). GHP-88310 docking pose is shown as grey sticks.

Residues that have been shown to confer resistance to GHP-88309 are highlighted in red. Surface

representation of the docking site of GHP-88310 showing the location of the docking pose between the

capping and RdRP domains (right). In (A-C), curves show 4-parameter variable slope non-linear

regression models; n $\geq$3. Symbols represent means $\pm$ SD (A,C) or the geometric mean $\pm$ geometric SD

(B).

Figure 6. GHP-88310 efficacy in well-differentiated air liquid interface human airway epithelial

cultures. A) Virus yield reduction of HPIV3 (clinical isolate 13J5) in undifferentiated NHBE cells (F3,

n=3). **B)** Schematic of well-differentiated air-liquid interface human airway epithelial organoids. **C)**

Assessment of GHP-88310 cytotoxicity in organoids. TEER was measured before and after 48 hours of

exposure to increasing concentrations of GHP-88310 in the basolateral chamber. Symbols represent

individual biological measurements, and lines show medians. **D)** Virus yield reduction assay of HPIV3

in organoids. Shed progeny virus was collected from the apical side of the transwells. Curves show 4-

parameter variable slope non-linear regression models, symbols represent the geometric mean $\pm$

geometric SD. **E)** HPIV3 titers in infected organoids used for confocal imaging. **F)** Confocal microscopy

of organoids infected with HPIV3 at 2dpi. Cells were stained with anti-HPIV3 HN (red), anti-ZO-1 (tight

junctions; green), and Hoechst 34580 (nuclei; blue). Z-stacks of 100-150 0.22 µm slices with 63× oil

objective. Scale bar, 10 µm. **G)** *Ex vivo* PK-guided assessment of GHP-88310 efficacy against HPIV3

in organoids. The simulation recapitulates the PK profile of GHP-88310 following 14 doses of 50 mg/kg

b.i.d. (top) or the PK profile after 20 mg/kg q.d. Solid lines represent the experimentally measured PK

profiles in ferrets. Corresponding compound concentrations of GHP-88310 applied at each time point

are shown as bars. **H,I)** HPIV3 titers from the *ex vivo* simulation study. In (E,H,I), bars indicate the

geometric mean $\pm$ geometric SD and symbols represent individual biological repeats. One-way ANOVA

with Dunnet's (comparisons to vehicle) post-hoc multiple comparisons test. Group sizes (n) are

indicated in each study schematic.