The role of type IV pilus in the interaction of *Neisseria gonorrhoeae* with a corneal epithelium tissue model

Lina Kafuri[1], Nicola Knetzger[2], Christian Lotz[2, 3], Vignesh Parasuraman[1], Thomas Rudel[1], Vera Kozjak-Pavlovic[1,*]

[1] Department of Microbiology, Biocenter, Julius-Maximilian University Würzburg, Würzburg, Germany

[2] Translational Center Regenerative Therapies TLC-RT, Fraunhofer Institute for Silicate Research ISC, Würzburg, Germany

[3] Department for Functional Materials in Medicine and Dentistry, University Hospital Würzburg, Würzburg, Germany

[*] Corresponding author: Vera Kozjak-Pavlovic (vera.kozjak@uni-wuerzburg.de)

Short title: Cornea models for *Neisseria gonorrhoeae* infection

Keywords: cornea, *Neisseria gonorrhoeae*, tissue models, type IV pilus, *ophthalmia neonatorum*

Abstract

Neisseria gonorrhoeae, the causative agent of the sexually transmitted disease gonorrhea, can infect intact corneal epithelium, leading to inflammation, cornea perforation, and blindness. Models for studying this type of gonococcal infection are few and limited. We have tested cornea models based on an immortalized hTCEpi corneal epithelium cell line for infection research using four derivatives of the *N. gonorrhoeae* MS11 strain that differ in pilus expression. We show that bacterial adherence depends on the functional type IV pilus in the early stages of infection, and that the formation of bacterial microcolonies and biofilms on the model surface leads to tissue destruction. The infection induces a specific cytokine response characterized by an increase in the secretion of IL-8 and TNF-α, but not of IL-6. The testing of trifluoperazine, a drug that induces pilus retraction, on infected corneal tissue models showed that the drug strongly diminished the number of adherent gonococci only when applied simultaneously with bacteria, and not when bacteria were allowed to form microcolonies on the tissue surface. Our work describes, for the first time, hTCEpi-based corneal epithelium tissue models as a useful tool for investigating *N. gonorrhoeae* infection, with potential for application in high-throughput studies and drug screening.

Introduction

Neisseria gonorrhoeae is a gram-negative diplococcus that causes the sexually transmitted disease gonorrhea. The infection in women is usually asymptomatic, and the risk of transmission to the newborn during birth is considerably high for undiagnosed pregnant women, which is often the case in countries where the screening for *N. gonorrhoeae* is lacking (1). Gonococcus (GC) can adhere and colonize the intact corneal epithelium, and is the main cause of *ophthalmia neonatorum*, an aggressive eye infection characterized by the inflammation of the eyelids and the presence of a yellow discharge, which can lead to cornea ulceration, perforation, and blindness (2). Adults are also susceptible to gonococcal eye infections. In recent years, the number of reported cases of eye infections caused by multi-antibiotic-resistant GC has increased due to misdiagnosis of the causing pathogen, delaying proper treatment and leading to permanent cornea injuries (1).

N. gonorrhoeae uses different virulence factors to adhere to epithelial cells, including the opacity proteins (Opa) and the type IV pilus. This complex protein machinery consists of many subunits, which together make dynamic filaments with the possibility of retraction and extrusion (3). The main subunit, pilin or PilE, has a high intra-strain structural variation dependent on the homologous recombination of the large family of PilS silent genes, mediated by the recombinase A (RecA) (4). This high variation not only allows *N. gonorrhoeae* to escape the immune system but also affects how they adhere to epithelial cells. However, PilE is not the only determining factor of the adherence behavior of GC (5). PilC, encoded by two different genes, pilC1 and pilC2, is located on the tip of the pilus where it acts as an adhesin, thus allowing the binding of the bacterium not only to the target tissue (6) but also to other bacteria, which facilitates the formation of microcolonies.

N. gonorrhoeae infection research, especially in the genitourinary tract, has been carried out using various models, including humanized hormone-treated mice, *in vitro* models using 2D cell monolayers, and 3D tissue models and organoids. This has allowed a better understanding of the key factors in gonococcal adherence and invasion (7). Previous infection studies focusing on corneal tissue mostly used human corneal explants (8) or the culture of human corneal epithelial cells on a bovine cornea-derived scaffold (1). Of the different human corneal tissue models established so far, most have been made as an alternative to the Draize irritation test (9), but not specifically for the infection studies. The accessibility and high variability of the patient-derived primary human corneal epithelial cells can make these models difficult to make and standardize. As an alternative, different human corneal epithelial cell lines can be used, such as a telomerase-immortalized human corneal epithelial cell line (hTCEpi), which can form stratified epithelium expressing corneal differentiation markers like ZO-1 and cytokeratin 3 when cultured under air-lift conditions (10).

In this work, we used an hTCEpi-based corneal model to study GC infection. We
assessed the role of the type IV pilus, using derivatives of the MS11 strain with
differences in pilus expression and adherence: MS11 F3 Pil- (Pil-, Opa+, PilC+,
recA+), MS11 F3 Pil+ (Pil+, Opa+, PilC+, recA+), MS11 N159 (Pil+, Opa+, PilC+,
recA-), and MS11 N191 (Pil+, Opa+, PilC-, recA+). We analyzed the effect of GC
infection on tissue permeability, cytotoxicity, and the production of key cytokines, as
well as demonstrated the importance of microcolony formation for the damage of the
corneal tissue, which is a symptom often observed during GC infection in patients.
Lastly, we tested the effect of trifluoperazine, a drug that can induce pilus retraction
in *Neisseria meningitidis*. We assessed the pilus retraction effect of the drug in terms
of bacterial adherence and IL-8 secretion. We also analyzed the bactericidal activity
of trifluoperazine and the induction of GC microcolony disaggregation. In sum, the
cornea model used in this study is a suitable and reproducible tool not only for
studying basic aspects of the GC eye infection but also for the testing of
antimicrobials.

Materials and methods

Generation of corneal epithelium tissue models

Human corneal epithelium tissue models were derived from telomerase-immortalized
human corneal epithelial cell line (hTCEpi) (Evercyte, Vienna, Austria, #CHT-045-
0237) (10). These cells were maintained in E1 medium (EpiLife medium) (Thermo
Fisher Scientific, Massachusetts, USA, #MEPI500CA) supplemented with 1 %
human keratinocyte growth supplement (HKGS) (Thermo Fisher Scientific,
Massachusetts, USA, #S0015) and 1 % penicillin/streptomycin (P/S) (Sigma-Aldrich,
Darmstadt, Germany, #P4333). Cells were seeded into culture flasks at a density of
4–6 cells/cm² and grown to 80–90 % confluence under standard conditions (37 °C, 5
% CO_2, 95 % humidity). The medium was exchanged every 2–3 days.

For passaging, cells were detached with trypsin and centrifuged at 200xg for 5 min.
The pellet was resuspended in E2 medium (EpiLife medium supplemented with 1 %
HKGS, 1 % P/S, and 0.48 % 300 mM $CaCl_2$).

Human corneal epithelium models were cultured in BRAND cell culture inserts
(BRAND, Wertheim, Germany, #782700) with polycarbonate membranes (0.4 µm
pore size and 0.59 cm² growth area). For seeding, 3x10☐ cells in 300 µl E2 medium
were added to the apical compartment of each insert and incubated for 2 h under
standard conditions (37 °C, 95 % humidity, 5 % CO_2). Subsequently, 1.4 ml E2
medium was added to the basolateral compartment. After 24 h of submerged culture,
the medium on the apical side of the inserts was removed, while the medium on the
basolateral side was replaced with 1.4 ml E3 medium (EpiLife medium
supplemented with 1 % HKGS, 1 % penicillin/streptomycin, 0.48 % $CaCl_2$, 73 µg/ml
ascorbyl-2-phosphate and 10 ng/ml KGF) to initiate air–liquid interface culture. The
medium was exchanged every 2–3 days thereafter.

Histology

Corneal epithelium models were fixed for 2 h with 4 % paraformaldehyde (Morphisto, Offenbach am Main, Germany, #11762.01000) and, after washing three times with 1xPBS, they were disassembled using an 8 mm biopsy puncher. After paraffin embedding, the samples were sectioned into 7 µm-thick slides. After a deparaffinization process including xylene and rehydration with a gradient of different concentrated solutions of ethanol, hematoxylin and eosin staining was done according to standard protocols (11). The stained tissue slides were analyzed with a light microscope (Leica Microsystems, Wetzlar, Germany).

Neisseria gonorrhoeae **strains**

All the *N. gonorrhoeae* strains used in this study are MS11 derivatives, and their whole genomes were sequenced using Oxford Nanopore (Microsynth, Balgach, Switzerland). The sequences described in this publication are available at NCBI's BioProject database under the accession number PRJNA1356842.

N. gonorrhoeae F3 Pil- (Pil$^-$, PilC$^+$, Opa$^+$, recA$^+$), F3 Pil+ (Pil$^+$, PilC$^+$, Opa$^+$, recA$^+$), N159 (Pil$^+$, PilC$^+$, Opa$^+$, recA$^-$), and N191 (Pil$^+$, PilC2, Opa$^+$, recA$^+$) were grown for 24 h at 37 °C and 5 % CO_2 on plates with Oxoid GC agar base (Thermo Fischer Scientific, Massachusetts, USA, #CM0367) supplemented with 1 % (v/v) vitamin mix. The day after, approximately 10 single colonies were selected, re-streaked onto a new plate, and grown for 16 h before being used for infection.

Infection of corneal epithelium models

Human corneal epithelium models were infected with the different *N. gonorrhoeae* strains on day 7 of the development of the models. The bacteria from the plate were taken with a sterile cotton swab and resuspended in 1 ml of DMEM F12 medium (Gibco, Thermo Fisher Scientific, Massachusetts, USA, #31331093). The OD_{550} was measured to calculate the bacterial number, and, taking into account the number of epithelial cells, the models were infected from the apical side for 2 h at an MOI of 20 in DMEM F12 medium. The models were then washed three times with 1xDPBS and incubated under airlift conditions until further evaluation at different time points (2 h, 24 h, and 72 h).

Immunoblotting

PilE (pilin) expression was analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and western blot. Briefly, the samples were dissolved in Laemmli buffer (62.5 mM Tris pH 6.8, 2 % SDS, 10 % glycerol, 5 % β-mercaptoethanol, and 0.002 % Bromophenol Blue) and separated on a 4-12 % NuPAGE Bis-Tris Precast gel (Thermo Fisher Scientific, Massachusetts, USA, # NP0322BOX). The separated proteins were then transferred onto a polyvinylidene fluoride (PVDF) membrane, which was blocked for 1 h with 5 % skimmed milk in PBS, followed by the overnight incubation with the pilin SM1 mouse monoclonal

156 antibody (a kind gift from Magdalene Yh So, University of Arizona), Omp85 (custom-
157 made in rabbits against the full-length protein; Davids Biotechnology, Regensburg,
158 Germany), or opa (in-house made in rabbits) at 4 °C. For detection, the membrane
159 was incubated with an HRP-coupled secondary antibody (Biozol, Eching, Germany),
160 and the images were processed with FIJI (12).

Barrier integrity of corneal epithelium models

To test the tissue integrity of the human cornea models upon infection, we measured
the permeability to 4 kDa fluorescein isothiocyanate (FITC)-Dextran (Sigma Aldrich,
Darmstadt, Germany, #46944). For this, 0.25 g/ml FITC-Dextran was dissolved in
Dulbecco's Modified Eagle Medium (DMEM) (Sigma Aldrich, Darmstadt, Germany,
#D6429) and sterile filtered. The medium on the basolateral side of the insert was
removed and replaced with 1 ml of DMEM, while 300 µl of the FITC-dextran solution
was added to the apical side. The models were incubated for 30 min (37 °C and 5 %
CO_2), and then 200 µl from the basolateral side was transferred to a black clear-
bottom 96-well plate (Corning, New York, USA, #3603). The fluorescence was
analyzed using a TECAN reader (490 nm absorption and 525 nm emission), and the
results were normalized to the values measured from an empty cell culture insert.

Cytotoxicity

174 Cytotoxicity was assessed by measuring the lactate dehydrogenase (LDH) activity in
175 the cell supernatant using the Cytotoxicity Detection KitPlus (Roche, Basel,
176 Switzerland, #4744926001). The assay was performed according to the
177 manufacturer's instructions, using the medium from the basolateral side of the
178 models.

Bacterial adherence

After the different time points, the medium was removed from the basolateral side of
the cell culture insert, and using an 8 mm biopsy puncher, the membrane was
carefully removed and placed into an Eppendorf tube containing 300 µl of 1 % (w/v)
saponin solution (Sigma Aldrich, Darmstadt, Germany, #47036), and the tubes were
vortexed for 1 min. After 30 min of incubation, serial dilutions were made and plated
on GC agar plates. The CFU/ml was determined after 48 h of incubation at 37 °C
with 5 % CO_2.

Cytokine measurements

72 h after infection, the medium was collected from the basolateral side of the cell
culture insert and analyzed using a personalized Legendplex cytokine panel assay
kit (BioLegend, San Diego, USA) to detect and measure the concentration of IL-1β,
TNF-α, IL-6, IL-8, IL-10, and MCP-1. The assay was performed according to the
manufacturer's instructions, and the results were analyzed using the BioLegend
software.

For the drug treatment experiments, the concentration of secreted IL-8 was
measured in the medium collected from the basolateral compartment of the models
infected with *N. gonorrhoeae* MS11 N159 and simultaneously treated with 40 µM
trifluoperazine (SimTr). For this, the Human IL-8 DuoSet ELISA kit (R&D Systems,
Minneapolis, USA, #DY208) was used, and the analysis performed according to the
manufacturer's instructions.

Immunofluorescence staining and imaging

At different time points, the medium was removed from the basolateral side of the
cell culture inserts, and the infected corneal epithelium models were fixed with 4 %
paraformaldehyde (Morphisto, #11462.01000) for 2 h at room temperature, followed
by washing 3 times with PBS and blocking with 3 % (w/v) bovine serum albumin
(BSA) (Carl Roth, Karlsruhe, Germany, #8076.3) and 0.01 % Triton X (Carl Roth,
Karlsruhe, Germany, #3051.4) in PBS for 1 h. The models were then incubated with
primary antibody against *N. gonorrhoeae* (United States Biological, Salem, USA,
#N0600-02) at 4 °C overnight. This was followed by incubation with a fluorophore-
coupled secondary antibody (Alexa Fluor 488, Thermo Fisher Scientific,
Massachusetts, USA, #A-11008), Phalloidin-Alexa Fluor 555 (Invitrogen/ Thermo
Fisher Scientific, Massachusetts, USA, #A34055), and DAPI (Sigma Aldrich,
Darmstadt, Germany, #D9542). The samples were mounted using Mowiol® (Carl
Roth, Karlsruhe, Germany, #0713.2). For imaging, Z-stacks were made using a
Leica STELLARIS 5 confocal imaging platform (Leica Microsystems, Wetzlar,
Germany), and the images were processed using FIJI (12).

Scanning Electron Microscopy (SEM)

After 72 hours of infection, the membranes of the cell culture inserts were carefully
removed with an 8 mm biopsy puncher and were fixed with 6.5 % glutaraldehyde at
4 °C overnight. Subsequent sample preparation was done as previously described
(13) and imaging was performed using a JEOL JSM-7500F microscope.

Drug assays

We applied two different approaches to study the effects of trifluoperazine on pilus
retraction. For the first approach, post infection treatment (PostTr), the corneal
epithelium models were infected with *N. gonorrhoeae* MS11 N159 for 1 h, as
described previously in this study. The models were washed 3 times with DPBS to
remove the non-adhered bacteria and were then treated with 40 µM trifluoperazine
(Sigma Aldrich, Darmstadt, Germany, #T8516) for 1 h. After this time, the models
were washed 3 times with DPBS and were left under the airlift condition for another
72 h. For the second approach, the simultaneous addition of bacteria and drug
(SimTr), the models were infected with *N. gonorrhoeae* MS11 N159 as described
and treated at the same time with 40 µM trifluoperazine. 1 h later, the models were
washed 3 times with DPBS and left under the airlift condition for further 72 h.
Bacterial adherence was then measured as previously described in this study.

Bacterial growth curves

Bacterial growth curves

The *N. gonorrhoeae* strains were grown for 24 h on GC agar plates. After this time,
they were re-streaked onto a fresh GC agar plate for 16 h before the experiment.
The bacteria were then taken up with a sterile cotton swab and diluted in 1 ml PPM+
medium (15 g/l proteose peptone (Becton Dickinson, New Jersey, USA, #211693), 5
g/l sodium chloride (VWR, Radnor, USA, #27810.364), 0.5 g/l soluble starch (Sigma-
Aldrich, Darmstadt, Germany, #33615), 1 g/l potassium dihydrogen phosphate (Carl
Roth, Karlsruhe, Germany, #3904.1), and 4 g/l dipotassium hydrogen phosphate
(Carl Roth, Karlsruhe, Germany, #P749.3), supplemented with 1 % (v/v) vitamin mix
and 0.5 % (v/v) sodium hydrogen carbonate (VWR, Radnor, USA, #27775.293). The
OD_{550} was measured and adjusted to 0.3 in 20 mL PPM+ medium and grown at 37
°C with constant shaking (130 rpm), until they reached an OD_{550} of at least 0.4
(exponential phase). Then they were diluted to an OD_{550} of 0.1, with 40 µM
trifluoperazine added to the corresponding cultures. The OD_{550} was measured every
hour for 5 h.

Bactericidal assay

To test the bactericidal effect of different concentrations of trifluoperazine in the two
approaches (PostTr and SimTr), *N. gonorrhoeae* MS11 N159 was grown on GC agar
plates for 24 h and then re-streaked onto a fresh plate, where it grew for
approximately 16 h. The bacteria were then taken up with a sterile cotton swab,
resuspended in 1 ml DMEM F12 medium, and the OD_{550} was measured. The same
volume needed to infect the corneal epithelium models with an MOI 20 was
calculated, and the solutions were prepared accordingly, with the different
concentrations of trifluoperazine (10, 20, 30, 40, and 50 µM) and gentamicin (150
µg/ml) (Gibco, Thermo Fisher Scientific, Massachusetts, USA, # 15710-049). 100 µl
of each solution was added to a flat-bottom 96-well plate.

For the first approach (PostTr), the plate containing just bacteria was incubated for 1
h at 37 °C with 5 % CO_2; after this time, trifluoperazine and gentamicin were added
to the corresponding wells, and the bacteria were treated for 1 h.

For the second approach (SimTr), the plate containing the bacterial suspension with
different concentrations of trifluoperazine and gentamicin was incubated for 1 h at 37
°C with 5 % CO_2.

After incubation, serial dilutions were made and plated onto GC agar plates.
Colonies were counted after 48 h to calculate the CFU/ml.

Aggregation assay

To analyze the effect of trifluoperazine on the disaggregation of microcolonies, *N.*
gonorrhoeae MS11 N159 was grown for 24 h on a GC agar plate, re-streaked onto a
fresh GC agar plate, and grown for approximately 16 h. The bacteria were collected
with a sterile cotton swab, diluted in 1 ml PPM+ medium, and the OD_{550} was

measured. The OD was adjusted to 0.3 in 15 ml PPM+ medium, and bacteria were
grown at 37 °C with constant shaking (130 rpm) until they reached an OD_{550} of
approximately 0.4. The bacterial culture was diluted with the DMEM F12 medium to
an OD of 0.1. 500 µl of this suspension was distributed into a 24-well plate and
incubated (37 °C, 5 % CO_2) for 2 h. After this time, the bacteria were treated with
increasing concentrations of trifluoperazine (10, 20, 30, 40, and 50 µM) and with
gentamicin (150 µg/ml) for 1 h. Before and after the treatment, three images per well
were taken using a Leica DMIRB microscope (Leica Microsystems, Wetzlar,
Germany), and the area of the microcolonies was calculated using FIJI (12).

Statistical methods

283 Statistical analyses were performed on at least three independent biological
284 replicates using 2-way ANOVA and Tukey's multiple comparison tests, with the help
285 of the GraphPad Prism software version 10.4.2.

Results

Type IV pilus plays a crucial role in the adherence of *N. gonorrhoeae* to the corneal epithelium

To study the role of the type IV pilus of *N. gonorrhoeae* in the adherence to the intact
corneal epithelium, we used a multilayer model made of immortalized human derived
corneal epithelial cell lines (hTCEpi) seeded onto polycarbonate membrane cell
culture inserts. These models show corneal differentiation traits, and their
architecture resembles the native tissue (Fig. 1A) (10). We allowed bacteria to
interact with the tissue for 2 h before washing and removing the ones that did not
adhere. Further analyses of the models were then performed at different infection
times, starting directly after the wash (2 h time point) and finishing 72 h later (Fig.
1B).

We infected the models with four different *N. gonorrhoeae* strains derived from
MS11, which exhibit different pilus phenotypes (Fig. 2A). To assess these
differences, we first performed whole genome sequencing of each strain and
analyzed the regions encoding for important proteins like pilE, pilC, opacity-
associated (opa) proteins, and recA (Table 1). We observed that one of the two pilE
genes in the F3 Pil- strain was not present and the other one was partially deleted,
resulting in no expression of pilin, which was confirmed by western blot of bacteria
before and after 2 h of infection (Fig. 2A, Supp. Fig. 1). This gene was intact in F3
Pil+, N159, and N191; however, there was a noticeably higher expression of the pilus
in N159 (Fig. 2A). In addition, recA gene in N159 is inactivated due to a large
insertion, which drastically reduces phase variation of the pilus. N191, although
possessing an intact pilE gene, has mutations in both pilC genes, which strongly
diminished the adherence of these bacteria (Table 1).

We assessed bacterial adherence, cytotoxicity, and tissue permeability during
infection. N159 strain showed the highest adherence at all time points (2 h, 24 h, and

72 h). The adherence capacity of F3 Pil+ was the second highest (Fig. 2B). This
strain resembles the wildtype MS11 genotype and contains the intact recA gene, so
the pilus undergoes phase-variation at a much higher rate compared to N159; this
was reflected in the fact that the pilus expression in the bacteria collected after 2 h of
infection was reduced (Fig. 2A), indicating that nonadherent bacteria were less
piliated. F3 Pil- strain showed a somewhat reduced adherence capacity, as
expected, due to the lack of PilE expression (Fig. 2A, B), especially at 2 h. Lastly, the
N191 strain showed the lowest adherence to the corneal epithelium, emphasizing
the importance of PilC in the adhesion to the target tissue (6) (Fig. 2B).

The mature models at the time of infection have a relative permeability of 8 %, as
measured by the FITC-dextran assay. The permeability increased over time to up to
15 % in both uninfected and infected models, indicating that this was probably
caused by the aging of the tissue models and not by infection (Fig. 2C).

The cytotoxicity upon infection was measured in relation to the concentration of
lactate dehydrogenase (LDH) released into the medium of the basolateral
compartment. This is connected to cell lysis, usually associated with apoptosis or
necrosis. We observed an increase in cytotoxicity with time, showing the highest
percentage in the models infected with the N159 strain for 72 h (17 %). Interestingly,
the F3 Pil- strain showed the second highest cytotoxicity, followed by F3 Pil+ and
N191, for which the values were comparable to those obtained for the non-infected
models (Fig. 2D).

In conclusion, adherence and cytotoxicity mostly correspond to the ability of bacteria
to adhere to the corneal tissue models with the help of the pilus. Relatively high
adherence and cytotoxicity of F3 Pil- strain indicate, however, that bacteria can use
other mechanisms besides pili to efficiently adhere to the corneal tissue.

***N. gonorrhoeae* forms microcolonies on the surface of corneal tissue models,**
depending on the presence of the pilus

Another important role of the type IV pilus in *N. gonorrhoeae* is the formation of
microcolonies. They allow GC to survive longer in the target tissue in the presence of
some antibiotics, due to their limited diffusion within the microcolony (14).

We observed large microcolonies already after 2 h of infection for the piliated strains
F3 Pil+ and N159. These microcolonies were distributed over the whole surface of
the corneal epithelium models. In contrast, F3 Pil- and N191 bacteria were present
on the models as single bacteria or in much smaller aggregates, visible only after 72
h of infection (Fig. 3A).

Scanning electron microscopy (SEM) of the infected tissue models 72 h post
infection revealed that F3 Pil+ and N159 microcolonies led to tissue damage, with
pits forming underneath bacterial aggregates attached and connected through long
filaments, most likely representing pili. Such filaments were absent in F3 Pil-, though
we observed occasional accumulations of these bacteria resembling a biofilm. N191

were hard to detect in SEM images and were found only as single diplococci
attached to the cell surface. Interestingly, pili filaments of N159 appeared much
longer and more abundant than those of F3 Pil+ (Fig. 3B).

Microscopy shows, therefore, that only piliated bacteria form large microcolonies,
which lead to tissue damage. Consequently, the high cytotoxicity observed for the
N159 strain corresponds to the most pronounced tissue damage visible on the SEM
images.

360 Corneal epithelial cells produce different cytokines and chemokines upon
361 infection with *N. gonorrhoeae*

Previous studies have shown that bacteria induce the secretion of different cytokines
by the mucosal epithelial cells, which are the first responders to infection. The
secreted cytokines regulate other cell types, recruiting immune cells to the site of
infection. The degree of this response is dependent on the cell type, as well as on
the pathogen and its virulence factors (15).

We evaluated the secretion of different cytokines and chemokines after 72 h of
infection with the four strains we used. Interleukin (IL)-8 was the most secreted
cytokine, followed by Monocyte Chemoattractant Protein (MCP)-1, IL-1β, and Tumor
Necrosis Factor (TNF)-α, whereas IL-6 and IL-10 were not detectable, or present in
very low amounts (below 3 pg/ml) (Fig. 4A). The main function of IL-8 is the
recruitment of neutrophils in response to tissue damage caused by infection or injury
(16). Whereas infection generally led to a significant increase in IL-8 production, the
models infected with the piliated strains F3 Pil+ and N159 showed the highest
secretion of IL-8, while the non-adherent N191 showed the lowest. The differences in
IL-8 production between the models infected with F3 Pil-, F3 Pil+, or N159 were not
statistically significant (Fig. 4A, B). The high secretion of IL-8 correlates in part with
the observed adherence and tissue damage (Fig. 2B, D).

The secretion of MCP-1, another chemokine responsible for the recruitment of
monocytes and macrophages, was also observed in all infected models, although
without statistically significant differences among them and non-infected models.
However, similar to the IL-8 secretion, F3 Pil-, F3 Pil+, and N159 strains showed the
highest values, while N191 showed the lowest (Fig. 4C). Regarding IL-1β and TNF-
α, we detected amounts less than 10 pg/ml in the infected tissue models with F3
Pil+, N159, and F3 Pil-. For the N191 strain, values were under the detection limit
(Fig. 4D, E).

The infection, therefore, induces the secretion of high amounts of IL-8 that correlate
with the ability of bacteria to adhere to the tissue, and are the highest for the strains
that express a functional pilus.

390 Trifluoperazine decreases bacterial adherence and secretion of IL-8

We next explored whether our multilayer corneal epithelium infection models are
suitable for drug testing. For this, we used trifluoperazine, a small molecule that
belongs to the group of phenothiazines and is currently used as an antipsychotic
agent. Previously, trifluoperazine was shown to have antimicrobial properties,
especially against bacteria that express the type IV pilus (17). In previous studies,
trifluoperazine disrupted pilus-dependent function, such as twitching mobility,
aggregation, and adherence to inert surfaces or cell surfaces in *N. meningitidis* (18).
In our study, we assessed the adherence to the corneal epithelium tissue models
and the IL-8 secretion after the treatment with trifluoperazine, using highly piliated
N159 *N. gonorrhoeae* MS11 derivative.

In the first approach, we infected the tissue models with the N159 strain for 1 h,
administered trifluoperazine (40 µM) for another hour, and then removed the medium
and the drug, allowing the infection to proceed for another 72 h (post-treatment
(PostTr)). In the second approach, bacteria and the drug were simultaneously added
to the models for 1 h, after which they were removed by washing, followed by
additional 72 h of infection (simultaneous treatment (SimTr)) (Fig. 5A).

To control for the bactericidal effects of trifluoperazine, we monitored bacterial growth
and survival in the presence of the drug. Starting from the second hour,
trifluoperazine strongly affected the growth of N159 (Fig. 4B). However, analysis of
bacterial survival after 1 h of treatment with different trifluoperazine concentrations in
the PostTr and SimTr setups revealed interesting differences. The drug had no
significant effect on bacterial survival when bacteria were first allowed to form
microcolonies for 1 h (Supp. Fig. 2A). On the other hand, we observed a
concentration-dependent decrease in the number of surviving bacteria when the
drug was added directly to the bacterial suspension for 1 h, without previous
incubation that would let bacteria aggregate (Supp. Fig. 2B). In both cases,
treatment with gentamicin completely eradicated the bacteria.

Additionally, we studied the effect that trifluoperazine has on *N. gonorrhoeae*
microcolonies. We incubated the suspension of N159 bacteria in a tissue culture
plate for 2 h, which was sufficient for microcolony formation, and then added
trifluoperazine in increasing concentrations, measuring the surface area of the
microcolonies before and after 1 h of drug treatment. The drug significantly
diminished the size of the microcolonies (Supp. Fig. 2C), though this did not
correlate with the bactericidal effect (Supp. Fig. 2A). Gentamicin, on the other hand,
did not affect bacterial microcolony size even though no bacteria survived the
treatment (Supp. Fig. 2C).

In agreement with these observations, we observed no significant difference in N159
adherence to the corneal tissue models in PostTr setup, whereas the simultaneous
addition of trifluoperazine significantly diminished the number of adherent bacteria
(Fig. 5C). Accordingly, the drug treatment led to a reduction in IL-8 secretion in the
infected corneal epithelium models (Fig. 5D). In addition, the SEM showed that, after

addition of trifluoperazine, N159 microcolonies on the surface of the tissue models
were reduced to only a few bacteria, with no visible pilus present (Fig. 5E).

Corneal epithelium tissue models, therefore, represent a reliable tool to test drugs
that affect bacterial pilus formation and adherence. One of these drugs is
trifluoperazine, which can reduce gonococcal infectivity by affecting pilus formation.

Discussion

In this work, we show the application of a multilayer corneal epithelium tissue model
for the study of the *N. gonorrhoeae* infection. These models were developed using
an immortalized human corneal epithelial cell line (hTCEpi) grown on a
polycarbonate membrane scaffold. hTCEpi expresses a human telomerase reverse
transcriptase (hTERT), leading to the activation of telomerase, which prevents the
shortening of telomeres and the telomere-dependent senescence in the cells without
altering their differentiation capacity (19). Under airlift conditions, these models
exhibit differentiation markers of the corneal epithelium and natural desquamation
comparable to the native cornea tissue (10). However, for infection studies,
additional parameters like the tissue permeability and integrity are also important.
Our multilayer corneal epithelium models have a permeability of 8 % and
transepithelial electrical resistance (TEER) values of 327 Ω^*cm^2 on day 7 after
introducing airlift conditions, which correlates with values measured for human
cornea (200 Ω^*cm^2) (20), making them suitable to be used as infection models.

One of the most important roles of the type IV pilus of *N. gonorrhoeae* is to enable
bacteria to adhere to tissues and form microcolonies (21, 22). In our experiments
using hTCEpi-based corneal tissue models, we observed that the adherence of the
piliated F3 Pil+ and N159 strains was indeed the highest, as expected, compared to
lower adherence of F3 Pil- and especially N191 (Fig. 2B). N191 strain expresses
pilus and has an intact recA gene but has both PilC genes mutated or deleted. These
mutations in PilC, which is the subunit located at the tip of the pilus and described as
the adhesin that allows the contact of the bacteria with the target tissue and other
bacteria, are responsible for the reduced adherence capacity of this strain, even
though PilE is expressed (Fig. 2A) (6). In our infection model, we could faithfully
reproduce the effect of the PilC defect and consistently observe low adherence of
N191 bacteria.

The differences in adhesion between piliated and non-piliated/non-adherent bacteria
were particularly noticeable at the earliest time point of 2 h. At later time points, the
number of adherent bacteria increased for all strains (Fig. 2B), probably due to the
multiplication of already-attached GC (Fig. 3A). Interestingly, F3 Pil- showed better
adherence than the N191 strain, indicating that the presence of a non-functional
pilus might interfere with the tissue contact. Both strains still interacted with the
tissue, however, possibly through opa proteins (23), since we did not specifically
select opa-negative bacterial colonies for infection (Fig. 2A). An earlier report
showed that non-piliated bacteria did not adhere to human corneal explants, and that

the adherence was independent of the opa status (8). However, in that study,
bacteria were incubated with the tissue for 1 h, whereas in our experiments, the
incubation time was 2 h. It is possible that longer incubation allowed bacteria to
interact with the tissue using opa proteins, and that the differences in adherence
between piliated and non-piliated bacteria would be more apparent at shorter
incubation times.

The same earlier study reported thinning and exfoliation of the human cornea
explants upon infection (8). *N. gonorrhoeae* was also shown to disrupt the apical
junctions of polarized epithelial cells (24, 25) and induce epithelial cell exfoliation in
cervical tissue explants (26). Contrary to this, we did not see major differences in
tissue permeability changes over time between infected and non-infected hTCEpi
cornea models that would be indicative of tissue thinning or exfoliation (Fig. 2C). The
cytotoxicity also did not seem to strictly depend on the presence of the pilus but
mostly correlated with the number of adherent bacteria (Fig. 2D). This might be due
to the differences between tissue explants and models, with models showing greater
robustness in infection scenario.

The formation of microcolonies is important for the survival of GC in the host
because it facilitates invasion and reduces the effectiveness of antibiotics, as they
are less able to diffuse within the microcolony. In our study, we observed tissue
damage and formation of pits under the microcolonies formed by highly piliated
N159. This is reminiscent of the observations made using human cornea explants
(8). Tissue damage caused by GC attachment could be related to cornea perforation,
one of the most severe complications in patients with untreated eye infections
caused by *N. gonorrhoeae* (27, 28). Likewise, SEM images at 72 h revealed that F3
Pil- forms occasional biofilm-like patches on the surface of the models. No pilus was
visible, but there was an apparent tissue destruction beneath these patches (Fig.
3B), which could explain the high cytotoxicity measured for models infected with this
strain (Fig. 2D). Interestingly, however, the lowest percentage of cytotoxicity was
seen after F3 Pil+ infection and not after N191 infection, suggesting that the cell
death caused by GC in the corneal epithelium tissue is not exclusively pilus-
dependent, and the role of other virulence factors is also relevant.

We also measured the concentration of different cytokines and chemokines
secreted upon infection by the corneal epithelial cells. The pattern of cytokines
depends on the tissue in question. In T-cells, the pilus-dependent adherence of *N.*
gonorrhoeae induces the activation and proliferation of these cells and the
production of anti-inflammatory IL-10 (29). In mature macrophages incubated with
piliated GC, the infection causes an increased secretion of the pro-inflammatory
cytokines IL-6, TNF-α, GRO-α, MIP-1α, and RANTES, with no effect on IL-8 (30). In
the male urethra (31), there is an increased expression of IL-8, IL-6, and TNF-α in
urine before the start of symptoms after GC infection, and in fallopian tubes, IL-1α,
IL-1β, and TNF-α are upregulated (32). Our previous work using 3D models that
mimicked endometrial and urethral tissue showed that urethral tissue models

secreted high amounts of IL-8, IL-6, and TNF-α, whereas endometrial tissue models
secreted significantly less of the same cytokines (11). There is little known, however,
about the cytokine secretion of corneal epithelium upon GC infection. We observed
an increase in IL-8 and TNF-α secretion in infected cornea models, which correlated
with the number of adherent bacteria. MCP-1 secretion was also increased, but this
did not seem to be related to the infection. Only small amounts of IL-1β, and no IL-6
or IL-10, were produced upon infection (Fig. 4). These results point to a specific,
pro-inflammatory response of corneal tissue to GC infection, mostly directed at the
recruitment of neutrophils, which corresponds to the clinical picture of the disease.

Finally, we tested the effects of trifluoperazine on pilus-mediated adherence and
infection using corneal tissue models. Trifluoperazine reduces bacterial piliation by
affecting the establishment of the sodium gradient through alteration of the activity of
the Na$^+$-pumping NADH-ubiquinone oxidoreductase complex (18). This complex
creates a Na$^+$ gradient across the cell membrane using energy released by the
oxidation of NADH and the reduction of quinone (33). Trifluoperazine was described
to induce pilus retraction and dispersal of microcolonies in *N. meningitidis* (18). We
demonstrated a similar effect on *N. gonorrhoeae* in liquid culture (Supp. Fig. 2).
However, in tissue models, we observed that when we allowed piliated GC to adhere
and form microcolonies on the tissue models for 1 h before the addition of
trifluoperazine, the drug was not as efficient as when added simultaneously with
bacteria (Fig. 5C). This is in agreement with the role of the pilus only in the initial
attachment to the tissue, with adherence to host cells being mediated by other
adhesins as the infection progresses.

Our study demonstrates for the first time the usefulness of hTCEpi-based tissue
models for infection research, shedding light on several important aspects of the
infection of the cornea with *N. gonorrhoeae* and the role of type IV pilus in this
process. In comparison to the primary tissue in the form of explants, corneal
epithelial tissue models are more accessible and show lower variability, while still
faithfully mimicking the original tissue, making them suitable for high-throughput
research and drug screening.

Acknowledgments

546 We thank Leonie Speth for technical assistance. This study was supported by the
547 Deutsche Forschungsgemeinschaft (DFG, German Research Society) through a
548 GRK2157/1+2 grant to VK-P, which included the salary of LK. The funders had no
549 role in study design, data collection and analysis, decision to publish, or preparation
550 of the manuscript.

Competing Interests

552 The authors have declared that no competing interests exist.

Data Availability

The sequencing data related to this study is available and can be accessed from BioProject: PRJNA1356842.

References

1. Churchward CP, Snyder LAS. 2019. In Vitro Models of Eye Infection with *Neisseria gonorrhoeae*. Methods Mol Biol 1997:363-376.
2. Churchward CP, Alany RG, Kirk RS, Walker AJ, Snyder LAS. 2017. Prevention of Ophthalmia Neonatorum Caused by *Neisseria gonorrhoeae* Using a Fatty Acid-Based Formulation. mBio 8:e00534-17.
3. Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional consequences. Nat Rev Microbiol 17:429-440.
4. Rudel T, van Putten JP, Gibbs CP, Haas R, Meyer TF. 1992. Interaction of two variable proteins (PilE and PilC) required for pilus-mediated adherence of *Neisseria gonorrhoeae* to human epithelial cells. Mol Microbiol 6:3439-50.
5. Rudel T, Boxberger HJ, Meyer TF. 1995. Pilus biogenesis and epithelial cell adherence of *Neisseria gonorrhoeae* pilC double knock-out mutants. Mol Microbiol 17:1057-71.
6. Rudel T, Scheuerpflug I, Meyer TF. 1995. Neisseria PilC protein identified as type-4 pilus tip-located adhesin. Nature 373:357-359.
7. Heydarian M, Rühl E, Rawal R, Kozjak-Pavlovic V. 2022. Tissue Models for *Neisseria gonorrhoeae* Research—From 2D to 3D. Front Cell Infect Microbiol 12.
8. Tjia KF, van Putten JP, Pels E, Zanen HC. 1988. The interaction between *Neisseria gonorrhoeae* and the human cornea in organ culture. An electron microscopic study. Graefes Arch Clin Exp Ophthalmol 226:341-5.
9. Lotz C, Kiesewetter L, Schmid FF, Hansmann J, Walles H, Groeber-Becker F. 2018. Replacing the Draize eye test: Impedance spectroscopy as a 3R method to discriminate between all GHS categories for eye irritation. Sci Rep 8.
10. Robertson DM, Li L, Fisher S, Pearce VP, Shay JW, Wright WE, Cavanagh HD, Jester JV. 2005. Characterization of Growth and Differentiation in a Telomerase-Immortalized Human Corneal Epithelial Cell Line. Invest Ophthalmol Vis Sci 46:470.
11. Heydarian M, Yang T, Schweinlin M, Steinke M, Walles H, Rudel T, Kozjak-Pavlovic V. 2019. Biomimetic Human Tissue Model for Long-Term Study of *Neisseria gonorrhoeae* Infection. Front Microbiol 10.
12. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. Nat Methods 9:676-682.
13. Spiliotis M, Lechner S, Tappe D, Scheller C, Krohne G, Brehm K. 2008. Transient transfection of *Echinococcus multilocularis* primary cells and complete in vitro regeneration of metacestode vesicles. Int J Parasitol 38:1025-1039.
14. Tseng BS, Zhang W, Harrison JJ, Quach TP, Song JL, Penterman J, Singh PK, Chopp DL, Packman AI, Parsek MR. 2013. The extracellular matrix protects *Pseudomonas aeruginosa* biofilms by limiting the penetration of tobramycin. Environ Microbiol 15:2865-2878.

15. Hedges SR, Agace WW, Svanborg C. 1995. Epithelial cytokine responses and
mucosal cytokine networks. Trends Microbiol 3:266-270.
16. Kobayashi Y. 2008. The role of chemokines in neutrophil biology. Front Biosci
13:2400-7.
17. Duménil G. 2019. Type IV Pili as a Therapeutic Target. Trends Microbiol
27:658-661.
18. Denis K, Le Bris M, Le Guennec L, Barnier J-P, Faure C, Gouge A, Bouzinba-
Ségard H, Jamet A, Euphrasie D, Durel B, Barois N, Pelissier P, Morand PC,
Coureuil M, Lafont F, Join-Lambert O, Nassif X, Bourdoulous S. 2019.
Targeting Type IV pili as an antivirulence strategy against invasive
meningococcal disease. Nat Microbiol 4:972-984.
19. Vaziri H, Benchimol S. 1998. Reconstitution of telomerase activity in normal
human cells leads to elongation of telomeres and extended replicative life
span. Curr Biol 8:279-282.
20. Knetzger N, Regensburger AK, Goy A, Han H, Theuersbacher J, Tarau IS,
Cleve C, Königer L, Finger T, Eibichova S, Haider MS, Schwarz T, Hansmann
J, Hillenkamp J, Kampik D, Lotz C. 2025. From injury to recovery:
investigating wound healing in a 3D tissue-engineered cornea equivalent.
Biomaterials 323:123409.
21. Jonsson AB, Ilver D, Falk P, Pepose J, Normark S. 1994. Sequence changes
in the pilus subunit lead to tropism variation of *Neisseria gonorrhoeae* to
human tissue. Mol Microbiol 13:403-416.
22. Zöllner R, Cronenberg T, Kouzel N, Welker A, Koomey M, Maier B. 2019.
Type IV Pilin Post-Translational Modifications Modulate Material Properties of
Bacterial Colonies. Biophys J 116:938-947.
23. Sadarangani M, Pollard AJ, Gray-Owen SD. 2011. Opa proteins and
CEACAMs: pathways of immune engagement for pathogenic *Neisseria*.
FEMS Microbiol Rev 35:498-514.
24. Edwards VL, Wang L-C, Dawson V, Stein DC, Song W. 2013. *Neisseria
gonorrhoeae* breaches the apical junction of polarized epithelial cells for
transmigration by activating EGFR. Cell Microbiol 15:1042-1057.
25. Rodriguez-Tirado C, Maisey K, Rodriguez FE, Reyes-Cerpa S, Reyes-Lopez
FE, Imarai M. 2012. *Neisseria gonorrhoeae* induced disruption of cell junction
complexes in epithelial cells of the human genital tract. Microbes Infect
14:290-300.
26. Wang L-C, Yu Q, Edwards V, Lin B, Qiu J, Turner JR, Stein DC, Song W.
2017. *Neisseria gonorrhoeae* infects the human endocervix by activating non-
muscle myosin II-mediated epithelial exfoliation. PLOS Pathog 13:e1006269.
27. Das S, D'Souza S, Gorimanipalli B, Shetty R, Ghosh A, Deshpande V. 2022.
Ocular Surface Infection Mediated Molecular Stress Responses: A Review. Int
J Mol Sci 23:3111.
28. Jongkhajornpong P, Nakamura T, Sotozono C, Inatomi T, Kinoshita S. 2015.
Phenotypic Investigation of Regenerated Epithelial Cells After Gonococcal
Corneal Perforation: A Clinical, Histological, and Immunohistochemical Study.
Cornea 34:1508-1512.
29. Plant LJ, Jonsson A-B. 2006. Type IV Pili of *Neisseria gonorrhoeae* Influence
the Activation of Human CD4[+] T Cells. Infect Immun 74:442-448.
30. Makepeace BL, Watt PJ, Heckels JE, Christodoulides M. 2001. Interactions of
Neisseria gonorrhoeae with Mature Human Macrophage Opacity Proteins

Influence Production of Proinflammatory Cytokines. Infect Immun 69:1909-
1913.
31.   Ramsey KH, Schneider H, Cross AS, Boslego JW, Hoover DL, Staley TL,
653   Kuschner RA, Deal CD. 1995. Inflammatory cytokines produced in response
654   to experimental human gonorrhea. J Infect Dis 172:186-91.
655   32.   Maisey K, Nardocci G, Imarai M, Cardenas H, Rios M, Croxatto HB, Heckels
656   JE, Christodoulides M, Velasquez LA. 2003. Expression of proinflammatory
657   cytokines and receptors by human fallopian tubes in organ culture following
658   challenge with *Neisseria gonorrhoeae*. Infect Immun 71:527-532.
659   33.   Juárez O, Barquera B. 2012. Insights into the mechanism of electron transfer
660   and   sodium   translocation   of   the   Na(+)-pumping   NADH:quinone
661   oxidoreductase. Biochim Biophys Acta 1817:1823-32.

Figure legends

Figure 1. Schematic representation of the corneal epithelium structure and study
design. (A) The scheme of the cornea structure (on the left side) in comparison to
the corneal epithelium model used in this study. The corneal tissue model was fixed,
embedded in paraffin, and sectioned into 7 µm-thick sections, stained using
hematoxylin and eosin (scale bar 50 µm). (B) Corneal tissue models were infected
from the apical side with the different *N. gonorrhoeae* MS11 derivative strains for 2 h,
followed by washing with PBS and further incubation under airlift conditions. Different
parameters were analyzed at different time points.

Figure 2. Impact of pilus expression on bacterial adherence, corneal tissue model
barrier integrity, and cytotoxicity. (A) The expression of Pilin in different *N.*
gonorrhoeae MS11 derivative strains was analyzed by immunoblotting before and
after 2 h of infection using antibodies against Pilin, Omp85 (Outer membrane protein
assembly factor BamA) and opa. (B) To determine the number of adherent bacteria,
corneal tissue models were solubilized with 1 % saponin for 30 min. Serial dilutions
were plated on GC agar to calculate the CFU/ml. (C) The corneal epithelium barrier
integrity was assessed before (0 h) and at different time points of infection by
measuring the diffusion of a FITC-dextran solution from the apical to the basolateral
compartment of the models. The values were normalized to the values of an empty
cell insert. (D) Cytotoxicity during different time points of infection was measured as
the percentage of lactate dehydrogenase LDH released to the medium collected
from the basolateral side of the corneal epithelium models. All graphs represent
mean values ± SD from three independent replicates. Statistical analysis was done
using 2-way ANOVA and Tukey's multiple comparison tests. * - $p \leq 0.05$, ** - $p \leq 0.01$,
*** - $p \leq 0.001$, and **** - $p \leq 0.0001$.

Figure 3. Pilus-expressing *N. gonorrhoeae* MS11 derivative strains form
microcolonies on the surface of corneal tissue models. (A) The models were fixed 2
h and 72 h post-infection and analyzed by immunofluorescence and confocal
microscopy, using DAPI (blue), Phalloidin-Alexa Fluor 555 (red), and the primary

antibody against *N. gonorrhoeae*/Alexa Fluor 488-coupled secondary antibody
(green). The images were made using the Leica Stellaris 5 confocal imaging
platform. Scale bar is 50 μm. (B) Corneal tissue models infected with different *N.*
gonorrhoeae MS11 derivative strains were fixed after 72 h of infection and analyzed
by scanning electron microscopy (SEM). Scale bars are 10 μm for low magnification
(Low Mag.) and 1 μm for other images.

Figure 4. Infection of corneal epithelium models *N. gonorrhoeae* MS11 leads to an
increase in IL-8 and TNF-α secretion. (A-E) Different cytokines and chemokines
secreted by the corneal epithelial cells were measured in the medium collected from
the basolateral side of the tissue models after 72 hours of infection using a
personalized Legendplex cytokine panel to assess the secretion of interleukin (IL)-8
(B), monocyte chemotactic protein 1 (MCP-1) (C), IL-1β (D), and TNF-α (E). The
graphs represent the mean values ± SD from three independent replicates.
Statistical analysis was performed using 2-way ANOVA and Tukey's multiple
comparison tests. * - $p \leq 0.05$, ** - $p \leq 0.01$, *** - $p \leq 0.001$, and **** - $p \leq 0.0001$.

Figure 5. Trifluoperazine decreases bacterial adherence to the corneal epithelium
models in the early stages of infection. (A) Schematic representation of two different
approaches to test the effect of trifluoperazine. In the PostTr approach, models were
first infected with N159 bacteria for 1 h, washed, and treated with 40 μM
trifluoperazine for 1 h, followed by another wash and incubation for 72 h under airlift
conditions. For the SimTr approach, the bacteria were treated with 40 μM
trifluoperazine simultaneously with infection for 1 h, after which the models were
washed and kept for 72 h under airlift conditions. (B) To measure the growth of N159
bacteria without and with 40 μM trifluoperazine, bacteria were grown in the PPM+
medium until they reached an $OD_{550} = 0.4$, diluted to an $OD_{550} = 0.1$, and 40 μM
trifluoperazine was added to one culture. OD_{550} was measured every hour for 5 h.
The graph represents mean values ± SD of at least three independent replicates. (C)
Models were prepared as described in (A) and solubilized with 1 % saponin solution.
The CFU/ml was calculated from the number of colonies growing after plating of
serial dilutions of the suspension on GC agar plates. The results were normalized to
the non-treated models. The graph represents mean values ± SD of at least three
independent replicates. (D) The concentration of IL-8 was measured in the
basolateral medium of the non-infected (NI) and N159-infected corneal tissue
models, with and without 40 μM trifluoperazine, using ELISA. The graph shows the
mean value of at least three independent replicates. Statistical analysis was
performed using 2-way ANOVA and Tukey's multiple comparison tests. * - $p \leq 0.05$, **
- $p \leq 0.01$. (E) Scanning electron microscopy images of the corneal epithelium models
infected with N159 with and without the 1 h treatment with 40 μM trifluoperazine.
Scale bar is 1 μm.

Supplementary figure legends

Supplementary Figure 1. Pilin protein in F3 Pil- strain has several deletions. Amino
acid sequences were translated using ExPasy according to the nucleotide
sequences obtained from whole genome sequencing and aligned to the reference
MS11 sequence (NCBI (CP003909)) using Clustal Omega.

Supplementary Figure 2. Trifluoperazine has a bactericidal effect as well as
causing disaggregation of GC microcolonies. (A-B) The same number of N159
bacteria used for the infection of corneal epithelium models (6×10^6) was treated with
increasing concentrations of trifluoperazine. For the PostTr approach, trifluoperazine
was added after 1 h of incubation (37 °C, 5 % CO_2) for 1 h. For the SimTr approach,
the bacteria were treated simultaneously with infection for 1 h. Serial dilutions were
plated to calculate the CFU/ml. 150 µg/ml of gentamicin was used as a positive
control. (C) N159 bacteria were grown in PPM+ medium until they reached an OD_{550}
of 0.4, diluted to an OD_{550} of 0.1 in DMEM F12, transferred into a 24-well plate, and
incubated for 2 h (37 °C, 5 % CO_2). After this time, the bacteria were treated with
increasing concentrations of trifluoperazine or with gentamicin for 1 h. Three images
of each of the triplicates were made before and after the treatment using a phase-
contrast microscope, and the area index of GC microcolonies was calculated using
FIJI. The graph represents the mean values ±SD of three independent replicates.
Statistical analysis was performed using 2-way ANOVA and Tukey's multiple
comparison tests. * - $p \leq 0.05$, ** - $p \leq 0.01$, and **** - $p \leq 0.0001$.

Table 1.

Gene	Gene ID	Function	Strain			
			MS11 F3 Pil-	**MS11 F3 Pil+**	**MS11 N159**	**MS11 N191**
PilE	NGFG_01821	Major pilin subunit	Deletions	NC	NC	NC
	NGFG_01819		Not present	NC	NC	NC
PilS	NGFG_02405		NC	NC	NC	NC
	NGFG_02431	Antigenic variation of PilE	NC	NC	NC	NC
	NGFG_02481		NC	NC	NC	NC
	NGFG_02482		NC	NC	NC	NC
	NGFG_02484		NC	NC	NC	NC
PilX	NGFG_00609	GC aggregation	NC	NC	NC	NC
PilG	NGFG_02119	Pilus assembly	NC	NC	NC	NC
PilC1	NGFG_01977	Pilus tip adhesin	NC	NC	NC	1158 bp insertion
PilC2	NGFG_00188		NC	NC	NC	Not present
Opa	NGFG_02238		NC	NC	NC	NC
	NGFG_02258		NC	NC	NC	NC
	NGFG_02259		NC	NC	NC	NC
	NGFG_02330	Mediate adhesion and invasion (epithelial cells, neutrophils, T and B cells)	NC	NC	NC	NC
	NGFG_02337		NC	NC	NC	NC
	NGFG_02351		NC	NC	NC	NC
	NGFG_02406		NC	shorter sequence	shorter sequence	shorter sequence
	NGFG_02432		NC	NC	NC	NC
	NGFG_02435		NC	NC	NC	NC
	NGFG_02467		NC	NC	NC	NC
	NGFG_0248		NC	Not	Not	Not

	3			present	present	present
RecA	NGFG_01045	Pilus antigenic variation	NC	NC	518 bp insertion	NC
PorB_IB	NGFG_01725	Outer membrane porin	NC	NC	NC	NC

754 NC: No changes in the sequence compared to the reference genome sequence

756 **Table 1.** Comparison of relevant genes involved in bacterial adherence and invasion
757 between the different *N. gonorrhoeae* MS11 derivative strains and the reference
758 genome of *N. gonorrhoeae* MS11 (NCBI, CP003909) according to the information
759 obtained by whole genome sequencing.

Figure 1

A

Epithelium

Bowman's layer

Stroma

Descemet's membrane

Endothelium

B

2 h

Infection
and adherence

Washing and
medium removal

2 h
24 h
72 h

Bacterial adherence
SEM
Confocal microscopy
Cell viability
Cytokine measurement

Figure 2

Figure 3

Figure 4

Figure 5

Figure legends

Figure 1. Schematic representation of the corneal epithelium structure and study design. (A) The scheme of the cornea structure (on the left side) in comparison to the corneal epithelium model used in this study. The corneal tissue model was fixed, embedded in paraffin, and sectioned into 7 µm-thick sections, stained using hematoxylin and eosin (scale bar 50 µm). (B) Corneal tissue models were infected from the apical side with the different *N. gonorrhoeae* MS11 derivative strains for 2 h, followed by washing with PBS and further incubation under airlift conditions. Different parameters were analyzed at different time points.

Figure 2. Impact of pilus expression on bacterial adherence, corneal tissue model barrier integrity, and cytotoxicity. (A) The expression of Pilin in different *N. gonorrhoeae* MS11 derivative strains was analyzed by immunoblotting before and after 2 h of infection using antibodies against Pilin, Omp85 (Outer membrane protein assembly factor BamA) and opa. (B) To determine the number of adherent bacteria, corneal tissue models were solubilized with 1 % saponin for 30 min. Serial dilutions were plated on GC agar to calculate the CFU/ml. (C) The corneal epithelium barrier integrity was assessed before (0 h) and at different time points of infection by measuring the diffusion of a FITC-dextran solution from the apical to the basolateral compartment of the models. The values were normalized to the values of an empty cell insert. (D) Cytotoxicity during different time points of infection was measured as the percentage of lactate dehydrogenase LDH released to the medium collected from the basolateral side of the corneal epithelium models. All graphs represent mean values ± SD from three independent replicates. Statistical analysis was done using 2-way ANOVA and Tukey's multiple comparison tests. * - $p \leq 0.05$, ** - $p \leq 0.01$, *** - $p \leq 0.001$, and **** - $p \leq 0.0001$.

Figure 3. Pilus-expressing *N. gonorrhoeae* MS11 derivative strains form microcolonies on the surface of corneal tissue models. (A) The models were fixed 2 h and 72 h post-infection and analyzed by immunofluorescence and confocal microscopy, using DAPI (blue), Phalloidin-Alexa Fluor 555 (red), and the primary antibody against *N. gonorrhoeae*/Alexa Fluor 488-coupled secondary antibody (green). The images were made using the Leica Stellaris 5 confocal imaging platform. Scale bar is 50 µm. (B) Corneal tissue models infected with different *N. gonorrhoeae* MS11 derivative strains were fixed after 72 h of infection and analyzed by scanning electron microscopy (SEM). Scale bars are 10 µm for low magnification (Low Mag.) and 1 µm for other images.

Figure 4. Infection of corneal epithelium models *N. gonorrhoeae* MS11 leads to an increase in IL-8 and TNF-α secretion. (A-E) Different cytokines and chemokines secreted by the corneal epithelial cells were measured in the medium collected from the basolateral side of the tissue models after 72 hours of infection using a personalized Legendplex cytokine panel to assess the secretion of interleukin (IL)-8 (B), monocyte chemotactic protein 1 (MCP-1) (C), IL-1β (D), and TNF-α (E). The graphs represent the mean values ± SD from three independent replicates. Statistical analysis was performed using 2-way ANOVA and Tukey's multiple comparison tests. * - $p \leq 0.05$, ** - $p \leq 0.01$, *** - $p \leq 0.001$, and **** - $p \leq 0.0001$.

Figure 5. Trifluoperazine decreases bacterial adherence to the corneal epithelium models in the early stages of infection. (A) Schematic representation of two different approaches to test the effect of trifluoperazine. In the PostTr approach, models were first infected with N159 bacteria for 1 h, washed, and treated with 40 μM trifluoperazine for 1 h, followed by another wash and incubation for 72 h under airlift conditions. For the SimTr approach, the bacteria were treated with 40 μM trifluoperazine simultaneously with infection for 1 h, after which the models were washed and kept for 72 h under airlift conditions. (B) To measure the growth of N159 bacteria without and with 40 μM trifluoperazine, bacteria were grown in the PPM+ medium until they reached an $OD_{550} = 0.4$, diluted to an $OD_{550} = 0.1$, and 40 μM trifluoperazine was added to one culture. OD_{550} was measured every hour for 5 h. The graph represents mean values ± SD of at least three independent replicates. (C) Models were prepared as described in (A) and solubilized with 1 % saponin solution. The CFU/ml was calculated from the number of colonies growing after plating of serial dilutions of the suspension on GC agar plates. The results were normalized to the non-treated models. The graph represents mean values ± SD of at least three independent replicates. (D) The concentration of IL-8 was measured in the basolateral medium of the non-infected (NI) and N159-infected corneal tissue models, with and without 40 μM trifluoperazine, using ELISA. The graph shows the mean value of at least three independent replicates. Statistical analysis was performed using 2-way ANOVA and Tukey's multiple comparison tests. * - $p≤0.05$, ** - $p≤0.01$. (E) Scanning electron microscopy images of the corneal epithelium models infected with N159 with and without the 1 h treatment with 40 μM trifluoperazine. Scale bar is 1 μm.