Compromised ESCRT signalling is sufficient for resistance to the Target of Rapamycin Complex inhibitor Torin1 in fission yeast

John P. Alao[1,] Rowshan A. Islam[1], Yusuke Toyoda[2], Shigeaki Saitoh[2] and Charalampos Rallis[1]*

[1] Research Centre of Molecular Cell Biology, Research Centre of Evolutionary and Functional Genomics, School of Biological and behavioural Sciences, Queen Mary University of London, Mile End Road, London, E1 4NS, United Kingdom

[2] Institute of Life Science, Kurume University, Asahi-machi 67, Kurume, Fukuoka 830-0011, Japan

*Correspondence: c.rallis@qmul.ac.uk

Abstract

Background

Fission yeast cells defective in Golgi-endosomal sorting display high resistance to Torin1, a pan-Target of Rapamycin (TOR) inhibitor. TOR complexes regulate the ESCRT system to integrate nutrient availability with cell division. TOR activity is frequently deregulated in cancer, making it an attractive therapeutic target. Deregulated ESCRT activity has also been associated with cancer, but its role in mediating drug resistance is not fully understood. Herein, we have investigated the role of the ESCRT system in regulating sensitivity to Torin1.

Methods

Growth assays were used to monitor the growth of yeast cells. The effect of Torin1 on protein

expression was monitored by immunoblotting. Fluorescence microscopy was used to

investigate the action of Torin1 on protein localization.

Results

The ESCRT system mediates Torin1-induced degradation of amino acid and glucose

transporters. The expression of these transporters at the plasma membrane is not abolished

in ESCRT mutants. Mutants unable to effectively ubiquitylate these transporters are also

resistant to Torin1. Impaired ESCRT-mediated protein degradation is associated with strong

resistance to Torin1.

Conclusions

Mutations in genes encoding ESCRT components have been reported in cancer. We present

evidence that compromised ESCRT signalling is sufficient for resistance to Torin1. Cells

defective in ESCRT signalling or ubiquitin homeostasis are highly resistant to Torin1. Our

studies demonstrate that defective ESCRT-mediated proteolysis can suppress sensitivity to

Torin1.

Keywords: TORC1**,** TORC2, ESCRT, Gsk3, fission yeast, nutrient transport, cell division,

cancer, caffeine, rapamycin, Torin1, drug resistance

Background

The Target of Rapamycin Complexes, TORC1 and TORC2, integrate nutrient availability and

environmental stress conditions with cell metabolism, growth and division. They, thus, play

important roles in cancer development and drug resistance, and dual complex inhibitors are

in clinical development. Crosstalk between the two complexes following ligand-receptor

activation complicates the effective use of TORC inhibitors as anti-cancer agents. As with all

anti-cancer therapeutics, *de novo* or acquired resistance limits the efficacy of pan-TORC

inhibitors (1) (2). We previously reported that *Schizosaccharomyces pombe* (*S. pombe*) fission

yeast mutants defective in Endosomal Sorting Complex Required for Transport (ESCRT)

signalling display strong resistance to the pan-TORC inhibitor Torin1 (3). Herein, we have

investigated the links between deregulated ESCRT signalling and Torin1 resistance in *S.*

pombe.

The ESCRT pathway is an ancient, evolutionarily conserved pathway consisting of

multicomponent complexes that execute various cellular functions, such as membrane

modification, endosomal sorting, cytokinesis, membrane repair, nuclear pore sealing and viral

budding in all organisms (4,5). Deregulated ESCRT activity has also been linked to genomic

instability and resistance to Epidermal Growth Factor Receptor (EGFR) inhibitors and

Oxaliplatin in cancer cells (6–8). In fission yeast, the TORC1 and TORC2 complexes regulate

nutrient transporter localization and stability (9–11). Additionally, these complexes also

regulate amino acid transporter expression by Isp7 a member of the 2-oxoglutarate-Fe(II)-

dependent oxygenase gene family (12). Ubiquitylation of the Aat1 transporter by the HECT-

type Pub1 E3 ligase is required for its localization to the Golgi-endosome. In the absence of

Pub1, Aat1 is localized to the plasma membrane (PM). Arrestin-related molecular adaptors

such as Any1 (Arn1) and Aly3 are required for Pub1-mediated ubiquitylation of nutrient

transporters. When TORC1 is inhibited, or *pub1* is deleted, Aat1 localizes to the PM and is

then degraded via the ESCRT pathway (9,10). Aly3 similarly regulates ubiquitylation of the

Ght5 high-affinity hexose transporter. The inhibition of TORC2 abolishes Gad8-dependent

phosphorylation of Aly3. This permits ubiquitylation of Ght5 by Pub1 and results in the

transport of Ght5 to the endosomes for degradation in lysosomes (10,11). Inhibition of TORC2

also results in the activation of the downstream Gsk3 kinase, which is itself implicated in

cancer development and progression. Gsk3 mutants are also highly resistant to Torin1.

Phosphorylation of Pub1 by Gsk3 has been reported to regulate its activity (3,13). Interestingly,

Pub1 also regulates Cdc25 stability, targeting it for ubiquitin-dependent proteasomal

degradation. Cdc25 is a phosphatase that regulates cell cycle progression by counteracting

the inhibitory phosphorylation of Cdc2 by Wee1(14). Cdc25 degradation is required for

cytokinesis and proper cell division following mitosis (15). Wee1 has been reported to delay

cell cycle progression in G2 following glucose withdrawal, and *wee1* mutants lose viability

under these conditions (16). Inhibition of Cdc2 activity leads to arrest in the G1, S or G2 phases

of the cell cycle in a context-dependent manner (14).

The expression of PM nutrient transporters is regulated by TORC1 and TORC2 according to

nutrient availability. In the absence of nitrogen, cells rapidly divide and exit the cell cycle or

undergo mating. In contrast, glucose limitation induces a transient cell cycle arrest in G2 as

Ght5 expression is upregulated (11,16,17). Exposure of fission yeast cells to caffeine or

rapamycin inhibits TORC1 but not TORC2, driving entry into mitosis without subsequent cell

cycle arrest. In contrast, exposure to Torin1 inhibits TORC1 and TORC2, which drives cells

into mitosis and cell cycle arrest in G2 (17,18). TORC1 and TORC2, together with Gsk3 and

Pub1, thus regulate Cdc25 activity, integrating nutrient availability with cell cycle regulation.

Furthermore, Torin1 induces Cdc25 degradation and cell cycle arrest due to the inhibition of

Cdc2 activity by Wee1 (17,19). We hypothesized that Torin1 induces the global degradation

of amino acid and glucose transporters, leading to cell cycle arrest due to Cdc2 inhibition.

Mutations that confer resistance to Torin1 would need to rewire cell signalling pathways to

reactivate or prevent the inactivation of Cdc2.

We report that Torin1 treatment results in the degradation of the Aat1 and Cat1 amino acid

transporters as well as the Ght5 glucose transporter in an ESCRT-dependent manner.

Furthermore, Pub1 is required for the degradation of these transporters following exposure to

Torin1. In *pub1Δ* and ESCRT mutants, Aat1, Cat1 and Ght5 expression is not abolished in

response to Torin1. Interestingly, suppression of global protein ubiquitylation also confers

resistance to Torin1. We propose that fission yeast strains defective in ESCRT-mediated

protein degradation maintain the expression of nutrient transporters at the PM following

exposure to Torin1. Deregulated ESCRT activity can thus contribute to Torin1 resistance by

preventing cell cycle exit.

Methods

Strains, media, and reagents

Strains are listed in Additional file 1. Cells were grown in yeast extract plus supplements medium (YES). For experiments in the absence of nitrogen, we used EMM2-N medium (Formedium, Hunstanton, United Kingdom). Stock solutions of caffeine (Sigma Aldrich, Gillingham, United Kingdom) (100 mM) were prepared in water stored at -20°C. Rapamycin (Tocris, Abingdon, United Kingdom) was dissolved in DMSO (500μg/mL) and stored at -20°C. Torin1 (Tocris) was dissolved in DMSO (3.3mM) and stored at -20°C.

Microscopy

Fluorescent microscopy of GFP-expressing strains was performed using an EVOS M5000 and Leica DMRA2 epi-fluorescent microscope attached to a monochrome Orca-ER camera from Hamamatsu Photonics.

Immunoblotting

Antibodies directed against GFP (11814460001 Roche) were purchased from Sigma-Aldrich (Gillingham, United Kingdom). Secondary antibodies directed against mouse (ab205719) IgG were purchased from Abcam (Abcam, Cambridge, United Kingdom). Cell pellets were prepared for SDS-PAGE and treated as previously reported (Alao et al., 2025). Where deemed appropriate, images were quantified using Image J software and Student's t-test for statistical analyses.

Results

Torin1 and nitrogen withdrawal induce Ght5 degradation

We previously identified genes encoding ESCRT pathway components in screens for Torin1 resistance (3,21). As part of this study, we also identified *hse1* (ESCRT 0) and *sst6* (*vps23*)

(ESCRT I) deletion mutants as being resistant to the drug. We also identified *atg6* and *vps38*

(required for the activation of ESCRT signalling) but not *atg36* (regulates the activation of

autophagy) mutants in our screens, indicating that the ESCRT pathway confers sensitivity to

Torin1 (Fig. 1A (Light green and dark green ovals, respectively) and Fig. 1B). *S. pombe* cells

cannot efficiently degrade GFP, which accumulates as Ght5 is degraded (22,23). We thus

investigated the effect of Torin1 on transmembrane nutrient transporters using strains

expressing Ght5-GFP (11). Exposure to 5 µM Torin1 but not 10 mM caffeine for 3h induced

Ght5-GFP degradation independently of Tor1. We noted, however, that Ght5-GFP degradation

was impaired in the *tor1Δ* mutant compared to *wt* cells (See discussion) (Fig. 1C and 1D).

Exposure to 100 ng/ mL rapamycin slightly induced Ght5-GFP degradation, as indicated by

the accumulation of GFP. However, exposure to rapamycin induced Ght5-GFP accumulation.

Exposure to caffeine and rapamycin induced Ght5-GFP degradation to an extent like that of

Torin1. It is unclear if this effect is due to the induction of autophagy rather than Tor1 inhibition

and degradation *via* ESCRT signalling. Caffeine also counteracted the effect of rapamycin on

Ght5-GFP accumulation (Fig. 1E). While we identified *atg6* as the only autophagy-regulating

gene in our screen for Torin1 resistance, exposure to the drug also induces autophagy as

detected by GFP-Atg8 cleavage (Additional file 1A). In contrast to glucose deprivation,

exposure to Torin1 thus mimics the effects of nitrogen withdrawal on Ght5-GFP (Fig. 1F).

Microscopic analyses demonstrated that exposure to 5 µM Torin1 induced Ght5-GFP vacuolar

accumulation after 1h, followed by degradation (1h vs 3h) (Fig. 1C and 1G). Similar results

were observed only when caffeine was combined with rapamycin. Caffeine and rapamycin

also exerted differential effects on Ght5 localization, which was more diffuse than the latter.

This is consistent with their mechanisms of TORC1 inhibition. We also noted that in contrast

to cells exposed to Torin1, cells exposed to caffeine with or without rapamycin were still

undergoing division after 3h (Fig. 1G). Since TORC2 prevents Ght5-GFP ubiquitylation, we

conclude that caffeine combined with rapamycin induces its degradation independently of the

ESCRT system. In contrast, Torin1 induces the degradation of Ght5-GFP via the ESCRT

pathways by inhibiting TORC2. Pub1 and Pub3 have been identified as Ght5 E3 ligases (10).

The deletion of *pub1* clearly suppressed the degradation of Ght5-GFP, confirming our previous

findings. Ght5-GFP degradation was not abolished in this mutant, likely due to residual Pub3

activity (Additional file 1C). Torin1 thus arrests cell division in G2, in part by preventing glucose

import via the Ght5 transporter at the PM (10,17).

Defective ESCRT signaling attenuates Torin1-mediated Ght5 degradation

Next, we investigated the effect of Torin1 on Ght5-GFP expression in ESCRT mutants. In

contrast to wt cells, Torin1 failed to induce Ght5-GFP degradation in *hse1Δ* mutants. We noted

that this mutation did not abolish Ght5-GFP degradation as GFP still accumulated. Deletion of

sst4 did not prevent Ght5-GFP degradation, although basal levels were higher, and we did not

observe the accumulation of GFP in this mutant (Fig. 2A to 2C). Exposure to Torin1 did not

induce Ght5-GFP degradation in *vps25Δ* mutants. GFP still accumulated in these mutants,

suggesting the degradation of Ght5-GFP is not completely abolished (Fig. 2D and 2E).

Inhibition of Tor1 or Gad8 activates Gsk3, which regulates Pub1 and Pub3 activity (ref) (Fig.

2F). We deleted *gsk3* in the Ght5-GFP mutant to investigate its role in regulating Ght5 stability.

However, deletion of *gsk3* did not affect Ght5-GFP expression or following exposure to Torin1

(Fig. 2G to 2I, Additional file 1B). The resistance of *gsk3* mutants to Torin1, thus, appears to

be unrelated to its regulation of Pub1/ Pub3 expression and Ght5-GFP ubiquitylation/

degradation (ref, ref). Our results identify the ESCRT system as being required for Torin1-

induced Ght5 degradation.

ESCRT mutants maintain Ght5 expression at the PM following exposure to Torin1

We next compared the expression of Ght5-GFP in wt and ESCRT pathway mutants following

exposure to 5 µM Torin1. In untreated cells, Ght5-GFP was expressed at the PM in bright

patches with little expression in the cytoplasm. Exposure to Torin1 induced the rapid import

of Ght5-GFP in a speckled pattern and subsequent degradation within the vacuole after 3h

(Fig. 1 and 3A). ESCRT mutants also expressed Ght5 at the PM, but the expression pattern

differed in different mutants. Ght5-GFP was visible in a granular cytoplasmic pattern even in

untreated *hse1Δ*, *vps25Δ* mutants and strongly at the cell membrane in *sst4Δ,* and *sst6Δ* cells

(Fig. 3B to 3E). Defective TORC2 and ESCRT-mediated protein degradation thus result in the

accumulation of Ght5 (ref). We noted by immunoblotting that basal Ght5-GFP levels were not

identical in these mutants (not shown). Following exposure to Torin1, Ght5-GFP was still

expressed at the PM in *hse1Δ*, *vps25Δ*, *sst4Δ,* and *sst6Δ* mutants. Torin1 exposure did not

affect the expression of Ght5 in *sst4Δ* mutants (Fig. 3D). Exposure to Torin1 appeared to

increase the expression of Ght5 at the PM in *sst6Δ* mutants (Fig. 3E). We concluded that,

unlike wt cells, *hse1Δ*, *vps25Δ*, *sst4Δ,* and *sst6Δ* mutants continue to express Ght5-GFP at

the PM following exposure to Torin1 (Fig. 3A to 3E). Glucose is required for cell division, and

low concentrations strongly induce the expression of Ght5 at the PM (Fig. 1F) (11). We

investigated whether growth on 0.03 % glucose induces resistance to Torin1. The Torin1

sensitivity of the *wt*, *tor1Δ* and ESCRT mutants rather increased under these conditions

(Additional file 1B). Thus, overexpression of Ght5 alone is insufficient to suppress Torin1

sensitivity. Torin1 induces Ght5 degradation, and its expression is maintained at the cell

membrane of ESCRT mutants. These findings partly explain the different effects of caffeine,

rapamycin and Torin1 on cell cycle progression in fission yeast (17,24).

Deletion of *pub1* abolishes Torin1- induced Aat1 degradation

TORC1-mediated ubiquitylation of the amino acid transporter Aat1 by Pub1 is required for its

retention at the Golgi/ endosomes (ref). We investigated the effect of Torin1 on Aat1 in wt and

pub1Δ mutants. As observed with Ght5-GFP, exposure to 5 µM Torin1 and nitrogen withdrawal

induced the degradation of Aat1-GFP. The effect was not observed following exposure to

caffeine and rapamycin, which induced Aat1-GFP degradation (Fig. 4A). This effect of Torin1

on Aat1 was abolished when *pub1* was deleted (Fig. 4B and 4D). Unlike Ght5-GFP, exposure

to caffeine and rapamycin did not induce Aat1-GFP degradation (Fig. 1C and 4A). Interestingly,

exposure to rapamycin induced the accumulation of Aat1-GFP in *pub1Δ* mutants. TORC1

negatively regulates Aat1 via Isp7 at the transcriptional level (12). Inhibition of TORC1 with

rapamycin may induce *aat1* expression and accumulation of Aat1 in *pub1* mutants, as it cannot

be degraded. (Fig. 4B). These findings also highlight significant differences between caffeine

and rapamycin in terms of TORC1 inhibition (19,24). The deletion of *gsk3* did not abolish the

Torin1-induced degradation of Aat1-GFP. Our observations suggest, however, that it does

affect the dynamics of Aat1-GFP degradation following exposure to Torin1 (Fig. 4E and 4F).

We next examined the effect of Torin1 on Aat1 expression by fluorescence microscopy. In *wt*

cells, exposure to Torin1 induced the transport of Aat1-GFP at the Golgi/ endosomes and

subsequent degradation (Fig. 4G). Deletion of *pub1* resulted in the constitutive expression of

Aat1-GFP at the PM as previously reported. Aat1-GFP also accumulated at the septa of

dividing cells following exposure to Torin1. Pub1 has been reported to be partly localized to

the septum in *S. pombe* (25). These findings demonstrate that Pub1 is required for the Torin1-

induced degradation of Aat1-GFP (Fig. 4G). It was also observed that *pub1* mutants continued

cell division, albeit asynchronously, following exposure to Torin1 (Fig. 4G). The larger cell size

of *pub1*Δ mutants has been previously reported (25,26). Studies using Aat1-GFP *gsk3*Δ

mutants produced variable results. In our experiments, the localization of Aat1-GFP in *gsk3*Δ

mutants mimicked that observed in *pub1*Δ mutants (Fig. 4H). We noted, however, that Aat1-

GFP exhibits a differential localization pattern in *gsk3*Δ mutants (Fig. 4G, 4H and I). These

findings are consistent with previous reports linking TORC2 and Gsk3 to Pub1 stability and

regulation of Aat1 expression (13). The inhibition of TORC2 induces Aat1 degradation.

Furthermore, Pub1 but not Gsk3 is required for the Torin1-induced degradation of Aat1 in

fission yeast. Both Pub1 and Gsk3 regulate Aat1 expression.

Deletion of *pub1* abolishes Torin1- induced Cat1 degradation

As the PM amino acid transporter Cat1 is regulated like Aat1, we next investigated the effect

of Torin1 on its expression. Exposure to 5 µM Torin1 induced Cat1-GFP degradation as

observed in Aat1-GFP and Ght5-GFP expressing strains (Fig. 1, 2 and 5A). Cat1-GFP

appeared to be more sensitive to exposure to 10 mM caffeine alone or with 100 ng/ mL

rapamycin than Aat1-GFP (Fig. 5B). Deletion of *pub1Δ* abolished the effect of Torin1 on Cat1-

GFP expression. We noted, however, that GFP still accumulated, indicating Pub1-independent

alternate Cat1-GFP degradation mechanisms. Pub3 may thus substitute for Pub1 in mediating

Cat1 but not Aat1 degradation (Fig. 5C). As observed with Aat1-GFP, deletion of *gsk3Δ*

delayed but did not abolish Torin1-induced Cat1-GFP degradation in this strain (Fig. 5C and

5D). This observation resembles that observed with the *aat1-GFP* strain (Fig. 4E and 4F).

Fluorescence microscopy revealed that exposure to Torin1 induced the relocation of Cat1-

GFP from the Golgi/ endosomes to vacuoles. In untreated wt cells, Cat1 was localized to the

Golgi/ endosomes. Exposure to Torin1 resulted in a diffuse appearance in the cytoplasm with

bright speckles (Fig. 5Ea and 5Eb). Deletion of *pub1* resulted in the dispersion of Cat1-GFP

from the Golgi/ endosomes to vacuoles to the cytoplasm, PM and septum of dividing cells (Fig.

5Ec and Ed). Deletion of *pub1Δ* prevented the degradation of Cat1-GFP following exposure

to Torin1. While exposure to Torin1 in wt cells resulted in a speckled appearance, this was not

observed in *pub1Δ* mutants, which fail to degrade Cat1 (Fig. 5B). Furthermore, *pub1* mutants

continued to express Cat1-GFP at the PM following exposure to Torin1 (Fig. 5Ec and 5Ed).

We next examined the effect of deleting *gsk3* in the *aat1-GFP, cat1-GFP, and Ght5-GFP*

strains. The deletion of *gsk3* did not affect the localization of *cat1* -GFP as observed in pub1Δ

mutants (Figure 5E and 5F). The deletion of *gsk3*, similarly, did not affect the expression of

Ght5-GFP in untreated cells. Together, our findings suggest that while Pub1 regulates the PM

expression of both Aat1 and Cat1, Gsk3 regulates only the localization of the former (Fig. 5F).

This observation is consistent with previous reports linking Gsk3 to Pub1 activity and Aat1

expression. Pub1 is, however, required for the degradation of Aat1, Cat1 and Ght5 following

exposure to Torin1. In this model, the deletion of *gsk3* would limit the availability of Pub1 and

suppress sensitivity to Torin1 by maintaining nutrient transporter expression (13). Fission yeast

cells can thus selectively fine-tune the expression of nutrient transporters under various

conditions in a TORC1 and TORC2-dependent manner.

The ubiquitin system is required for Torin1 sensitivity and links TORC1 and TORC2

signalling with cell division

Torin1 induces the ubiquitin-dependent degradation of nutrient transporters and Cdc25 by

Pub1 (Fig. 1, 2, 3, 4 and 5) (19). E3 ligases such as Pub1 regulate multiple targets and

pathways. Loss of Pub1 expression would prevent the degradation of Cdc25 and numerous

nutrient transporters (9,9,10,19). The deletion of *pub1* did not confer resistance to caffeine

combined with rapamycin or Torin1 but rather increased sensitivity to these drugs (Fig. 6A).

Mutants lacking Pub1 are sensitive to various stresses and exhibit cell cycle defects, which

may override its effect on Torin1 sensitivity (27). Gsk3 activity is elevated in *gad8Δ* mutants.

These cells displayed increased sensitivity to caffeine combined with rapamycin or Torin1,

consistent with TORC2 regulating sensitivity to these agents. As previously reported, *gsk3Δ*

mutants are highly resistant to Torin1 (3). The deletion of *pub2* did not affect sensitivity to either

caffeine plus rapamycin or Torin1. When both *pub2* and *pub3* were deleted, the cells exhibited

resistance to Torin1 but not caffeine combined with rapamycin. A WD-repeat protein Lub1, the

loss of which was found to show resistance to Torin1, is required for adequate ubiquitin

expression in fission yeast, and *lub1Δ* mutants exhibit low global ubiquitylated protein levels

(28). These mutants are highly resistant to Torin1, presumably due to their inability to

sufficiently ubiquitylate target proteins (Fig. 6A). These findings identify a novel ubiquitin-

dependent, ESCRT-regulated degradation system required for sensitivity to Torin1. We

deleted *gsk3* in a strain expressing a Cdc25-GFP variant that lacks negative phosphorylation

sites (29). In the Cdc25(12A)-GFP *gsk3Δ* mutant, expression of the phosphatase was

extremely low and cleaved GFP was not detectable. Furthermore, exposure to Torin induced

Cdc25(12A) degradation in wt but not *gsk3Δ* cells. Surprisingly, Torin1 induced Cdc25

accumulation in the absence of Gsk3 (Fig. 6B). This ESCRT-dependent system thus

integrates nutrient transport with cell division through dual regulation of PM nutrient transporter

and Cdc25 stability dependent on Pub1 (Fig. 6C).

Discussion

We have identified an ESCRT-dependent protein degradation system that selectively

regulates sensitivity to Torin1 in fission yeast. E3 ligases target nutrient transporters for

degradation in lysosomes via the ESCRT pathway, which differs mechanistically from

autophagy (4,30). TOR inhibitors have drawn considerable attention as anticancer agents, due

to their central role in cellular metabolism, autophagy and cell division (1,2). Their use as

anticancer agents has been limited by modest anti-tumor activity and various drug resistance

mechanisms. Current studies investigate the impact of cell metabolism on the inhibitory

activities of these drugs and the mechanisms that underlie resistance to them (2). We present

evidence that suppression of Torin1-induced degradation of nutrient transporters is sufficient

for resistance to this drug. To our knowledge, such a resistance mechanism has not been

previously reported for TOR inhibitors. Clinically, deregulated ESCRT signalling has been

proposed as a potential mechanism of drug resistance. Compromised EGFR sorting to

lysosomes via the ESCRT pathway could potentially result in resistance to EGF receptor

signalling. Thus, drug resistance need not be necessarily linked to mutations altering the drug

affinity of a target protein or the activity of downstream effectors. Our findings in an

evolutionarily conserved model system confirm the need for further investigations into

deregulated ESCRT signalling as a mechanism of TORC inhibitor drug resistance (6,31).

Crucially, this system does not mediate Torin1 resistance due to a single deregulated protein.

Rather, compromised individual proteins at any stage of the ESCRT-mediated degradation

pathway, including low ubiquitin levels, confer resistance to Torin1. We identified only *atg6* and

vps38 mutants as exhibiting Torin1 resistance in the study by Rodríguez-López *et al*., (27).

Atg6 and Vps38 direct protein degradation via the ESCRT pathway, while Atg38 regulates

autophagy. We also noted that the majority of *atg* gene deletions strongly enhanced sensitivity

to Torin1 (21). Thus, defective ESCRT signalling but not autophagy induces resistance to

Torin1 (Fig. 1A).

TORC1 and 2 regulate the expression of nutrient transporters through the ESCRT system,

which is dependent on the E3 ligases Pub1 and Pub3 (10,11,13,32). Cells ensure the

adequate import of nutrients by selectively adjusting the expression levels of nutrient

transporters, including Aat1, Cat1 and Ght5. On nitrogen withdrawal, membrane transporters

relocate to the PM and/ or are directly degraded via the ESCRT system. Glucose deprivation

induces   G2 arrest, and both conditions lead to cell cycle exit and induction of autophagy

(11,32–34).   TORC2 positively regulates protein expression levels by preventing their

degradation by the ESCRT system. Exposure to Torin1 but not caffeine or rapamycin induces

cell cycle arrest. Thus, Torin1 inhibits fission yeast proliferation by inhibiting TORC2. We have

demonstrated that exposure to Torin1 mimics nitrogen withdrawal by inducing the degradation

of amino acid and glucose transporters. This degradation requires the prior arrestin-related

adaptor-dependent ubiquitylation of substrate proteins by Pub1 and Pub3 (10,11,25). We have

shown that mutants defective in ESCRT-mediated degradation display strong resistance to

and fail to degrade Ght5 in response to Torin1. Hence, Torin1 sensitivity correlates strongly

with uncompromised ESCRT signalling. ESCRT mutants such as *hse1Δ* and *vps25Δ* fail to

degrade Ght5 following exposure to Torin1 and are highly resistant to the drug. Aat1, Cat1 and

Ght5 mutants lacking *pub1* fail to degrade these transporters following Torin1 exposure.

Indeed, the degradation of Aat1 does not occur in response to nitrogen withdrawal in *pub1Δ*

mutants. In contrast to *wt* cells, therefore, ESCRT and *pub1Δ* mutants continue to express

nutrient transporters at the PM. This may permit the sufficient importation of nutrients to

prevent cell cycle exit. The deletion of *pub*1 did not confer resistance to Torin1. This may reflect

its dual role in negatively regulating Cdc25. We noted that Torin1 failed to arrest but induced

asynchronous cell division in these mutants. Interestingly, low endogenous ubiquitin levels are

also associated with Torin1 resistance. In these mutants, insufficient ubiquitin availability

prevents the degradation of target proteins. Mutants that fail to de-ubiquitylate target proteins

are also resistant to this agent, likely due to ubiquitin sequestration by target proteins, which

accumulate within the cell or on the PM (3,7). Our study identifies an ESCRT system that

includes proteins maintaining ubiquitin homeostasis, arrestin-related adaptors, Pub1 and

Pub3, and vacuolar biosynthesis whose expression is required for Torin1 sensitivity. TORC1

and 2 integrate the availability of nutrients with cell division through this system by co-

regulation of nutrient transporters and Cdc25 (ref, ref) (Fig. 6C)

While caffeine in combination with rapamycin does slow cell division, our studies suggest that

the underlying mechanism differs substantially from that of Torin1. Torin1 directs Ght5 for

degradation through the ESCRT system. In contrast, caffeine plus rapamycin induces the

degradation of Cat1 and Ght5 but not Aat1. Significantly, rapamycin induced Aat1

accumulation in *pub1Δ* mutants. We noted that cleavage of Aat1-, Cat1- and Ght5-GFP still

occurs in *pub1* mutants. Furthermore, ESCRT mutants activate autophagy in the presence of

Torin1. Caffeine combined with rapamycin may target Cat1 and Ght5 for degradation via

autophagy. The co-deletion of *pub2* and *pub3* conferred resistance to Torin1 but not caffeine

and rapamycin. Low ubiquitin expression in *lub1* mutants presumably induces Torin1

resistance by suppressing Pub1 and Pub3 activity (28). These mutants exhibited resistance

to caffeine plus rapamycin and Torin1, but there is no correlation in drug resistance between

them. The ESCRT system thus mediates sensitivity to Torin1 but not caffeine when combined

with rapamycin. Caffeine has been shown to indirectly inhibit TORC1 by inducing the

activation of AMPK. These findings suggest that caffeine, a well-characterised drug, can

suppress resistance to TOR inhibitors. Caffeine in combination with rapamycin may trigger

nutrient transporter degradation independently of ESCRT signalling. We previously reported

that mutants unable to transfer substrate proteins from the Golgi/ endosomes to lysosomes

are also resistant to Torin1. Future studies will investigate the role of the ESCRT system in

mediating sensitivity to Rapalink-1 (3,35).

Conclusions

We have identified an ESCRT-dependent pathway required for sensitivity to Torin1. A role for

this pathway in mediating resistance to TOR inhibitors has not been previously reported.

Mutants unable to degrade PM nutrient transporters continue to undergo cell division in the

presence of Torin1. Our findings partly explain why Torin1, but not caffeine, combined with

Rapamycin induces cell cycle arrest. Crucially, ubiquitin homeostasis is similarly required for

sensitivity to Torin1.

377 List of abbreviations

EGFR- Epidermal Growth Factor Receptor

ESCRT- Endosomal Sorting Complex Required for Transport

GFP- Green Fluorescent Protein

PM- Plasma Membrane

TORC1- Target of Rapamycin Complex 1

TORC2- Target of Rapamycin Complex 2

384 Declarations

385 Consent for publication

386 All authors consented to the publication of the manuscript.

387 Availability of data and materials

388 Data and yeast strains are available on request, subject to permission of the creators.

389 Competing interests

390 The authors declare no competing interests.

391 Funding

392 C.R. acknowledges funding from the Royal Society (Research Award grant number

393 RGS\R1\201348), BBSRC (grant numbers BB/V006916/1, BB/V006916/2) and MRC (grant

394 number MR/W001462/1). Y.T. acknowledges funding from Grants-in-Aid for Scientific

395 Research (C) from the Japan Society for the Promotion of Science (23K05758).

Authors' contributions

J.P.A. and C.R. designed the study. J.P.A. and R.A.I. carried out experiments. Y.T. prepared the yeast mutants. J.P.A., Y.T. and S.S. and C.R. analyzed the data. J.P.A. wrote the manuscript. J.P.A., R.A.I., C.R., Y.T. and S.S. revised and edited the manuscript

Acknowledgements

We are grateful to the Matsumoto and Nakashima laboratories and the Yeast Genetic Resource Centre (YGRC) Japan for mutant *S. pombe* strains.

References

1.   Eyre TA, Collins GP, Goldstone AH, Cwynarski K. Time now to TORC the TORC? New developments in mTOR pathway inhibition in lymphoid malignancies. Br J Haematol. 2014 Aug;166(3):336–51.

2.   Faes S, Demartines N, Dormond O. Resistance to mTORC1 Inhibitors in Cancer Therapy: From Kinase Mutations to Intratumoral Heterogeneity of Kinase Activity. Cai Z, editor. Oxid Med Cell Longev. 2017 Jan;2017(1):1726078.

3.   Rodríguez-López M, Gonzalez S, Hillson O, Tunnacliffe E, Codlin S, Tallada VA, et al. The GATA Transcription Factor Gaf1 Represses tRNAs, Inhibits Growth, and Extends Chronological Lifespan Downstream of Fission Yeast TORC1. Cell Rep. 2020 Mar;30(10):3240-3249.e4.

4.   Hurley JH, Coyne AN, Miączyńska M, Stenmark H. The expanding repertoire of ESCRT functions in cell biology and disease. Nature. 2025 June 26;642(8069):877–88.

5.   Remec Pavlin M, Hurley JH. The ESCRTs – converging on mechanism. J Cell Sci. 2020 Sept 15;133(18):jcs240333.

6.   Abdelrazik NM, Patel A, Conn A, Sutton CW, Kantamneni S, Shnyder SD. A Novel

Modulator of Resistance for Oxaliplatin-Based Therapy for Colorectal Cancer: The

ESCRT Family Member VPS4A. Cells. 2025 June 19;14(12):929.

7.   Baldys A, Raymond JR. Critical Role of ESCRT Machinery in EGFR Recycling.

Biochemistry. 2009 Oct 13;48(40):9321–3.

8.   Singh S, Yeat NY, Wang YT, Lin SY, Kuo IY, Wu KP, et al. PTPN23 ubiquitination by

WDR4 suppresses EGFR and c-MET degradation to define a lung cancer therapeutic

target. Cell Death Dis. 2023 Oct 11;14(10):671.

9.   Nakashima A, Kamada S, Tamanoi F, Kikkawa U. Fission yeast arrestin-related

trafficking adaptor, Arn1/Any1, is ubiquitinated by Pub1 E3 ligase and regulates

endocytosis of Cat1 amino acid transporter. Biol Open. 2014 June 15;3(6):542–52.

10.   Toyoda Y, Masuda F, Saitoh S. Phosphorylation of the Aly3 C-terminus impedes

aberrant endocytosis of *Schizosaccharomyces pombe* hexose transporter Ght5. J

Cell Sci. 2025 Apr 1;138(7):jcs263572.

11.   Toyoda Y, Soejima S, Masuda F, Saitoh S. TORC2 inhibition of α-arrestin Aly3

mediates cell surface persistence of *S. pombe* Ght5 glucose transporter in low

glucose. J Cell Sci. 2021 May 15;134(10):jcs257485.

12.   Laor D, Cohen A, Pasmanik-Chor M, Oron-Karni V, Kupiec M, Weisman R. Isp7 Is a

Novel Regulator of Amino Acid Uptake in the TOR Signaling Pathway. Mol Cell Biol.

2014 Mar 1;34(5):794–806.

13.   Wang T, Woodman P, Humphrey SJ, Petersen J. Environmental control of Pub1

(NEDD4 family E3 ligase) in *Schizosaccharomyces pombe* is regulated by TORC2

and Gsk3. Life Sci Alliance. 2022 May;5(5):e202101082.

14.   Nefsky B, Beach D. Pub1 acts as an E6-AP-like protein ubiquitiin ligase in the

degradation of cdc25. EMBO J. 1996 Mar 15;15(6):1301–12.

15.   Esteban V, Sacristán M, Andrés S, Bueno A. The Flp1/Clp1 phosphatase cooperates

with HECT-type Pub1/2 protein-ubiquitin ligases in Schizosaccharomyces pombe.

Cell Cycle Georget Tex. 2008 May 1;7(9):1269–76.

16.   Masuda F, Ishii M, Mori A, Uehara L, Yanagida M, Takeda K, et al. Glucose restriction

450   induces transient G2 cell cycle arrest extending cellular chronological lifespan. Sci

451   Rep. 2016 Jan 25;6(1):19629.

452   17.   Atkin J, Halova L, Ferguson J, Hitchin JR, Lichawska-Cieslar A, Jordan AM, et al.

453   Torin1-mediated TOR kinase inhibition reduces Wee1 levels and advances mitotic

454   commitment in fission yeast and HeLa cells. J Cell Sci. 2014 Mar 15;127(Pt 6):1346–

455   56.

456   18.   Ikai N, Nakazawa N, Hayashi T, Yanagida M. The reverse, but coordinated, roles of

457   Tor2 (TORC1) and Tor1 (TORC2) kinases for growth, cell cycle and separase-

458   mediated mitosis in *Schizosaccharomyces pombe*. Open Biol. 2011 Nov;1(3):110007.

459   19.   Alao JP, Kumar J, Stamataki D, Rallis C. Dissecting the cell cycle regulation, DNA

460   damage sensitivity and lifespan effects of caffeine in fission yeast. Microb Cell. 2025

461   June 24;12:141–56.

462   20.   Alao JP, Sjölander JJ, Baar J, Özbaki‐Yagan N, Kakoschky B, Sunnerhagen P.

463   Caffeine stabilizes C$_{DC}$25 independently of R$_{AD}$3 in *S chizosaccharomyces pombe*

464   contributing to checkpoint override. Mol Microbiol. 2014 May;92(4):777–96.

465   21.   Lie S, Banks P, Lawless C, Lydall D, Petersen J. The contribution of non-essential

466   *Schizosaccharomyces pombe* genes to fitness in response to altered nutrient supply

467   and target of rapamycin activity. Open Biol. 2018 May;8(5):180015.

468   22.   Arines FM, Li M. An optimized protocol to analyze membrane protein degradation in

469   yeast using quantitative western blot and flow cytometry. STAR Protoc. 2022

470   June;3(2):101274.

471   23.   Vijayasimha K, Tran MV, Leestemaker-Palmer AL, Dolan BP. Direct Conjugation of

472   NEDD8 to the N-Terminus of a Model Protein Can Induce Degradation. Cells. 2021

473   Apr 9;10(4):854.

24. Rallis C, Codlin S, Bähler J. TORC 1 signaling inhibition by rapamycin and caffeine

affect lifespan, global gene expression, and cell proliferation of fission yeast. Aging

Cell. 2013 Aug;12(4):563–73.

25. Nakase Y, Nakase M, Kashiwazaki J, Murai T, Otsubo Y, Mabuchi I, et al. Fission

yeast Any1, β-arrestin-like protein, is involved in TSC-Rheb signaling and the

regulation of amino acid transporters. J Cell Sci. 2013 Jan 1;jcs.128355.

26. Graml V, Studera X, Lawson JLD, Chessel A, Geymonat M, Bortfeld-Miller M, et al. A

Genomic Multiprocess Survey of Machineries that Control and Link Cell Shape,

Microtubule Organization, and Cell-Cycle Progression. Dev Cell. 2014

Oct;31(2):227–39.

27. Rodríguez-López M, Bordin N, Lees J, Scholes H, Hassan S, Saintain Q, et al. Broad

functional profiling of fission yeast proteins using phenomics and machine learning.

eLife. 2023 Oct 3;12:RP88229.

28. Ogiso Y, Sugiura R, Kamo T, Yanagiya S, Lu Y, Okazaki K, et al. Lub1 Participates in

Ubiquitin Homeostasis and Stress Response via Maintenance of Cellular Ubiquitin

Contents in Fission Yeast. Mol Cell Biol. 2004 Mar 1;24(6):2324–31.

29. Frazer C, Young PG. Carboxy-terminal phosphorylation sites in Cdc25 contribute to

enforcement of the DNA damage and replication checkpoints in fission yeast. Curr

Genet. 2012 Aug;58(4):217–34.

30. Nachmias D, Frohn BP, Sachse C, Mizrahi I, Elia N. ESCRTs – a multi-purpose

membrane remodeling device encoded in all life forms. Trends Microbiol. 2025

Feb;S0966842X25000083.

31. Hasegawa J, Jebri I, Yamamoto H, Tsujita K, Tokuda E, Shibata H, et al. SH3YL1

cooperates with ESCRT-I in the sorting and degradation of the EGF receptor. J Cell

Sci. 2019 Oct 1;132(19):jcs229179.

32. Nakase M, Nakase Y, Chardwiriyapreecha S, Kakinuma Y, Matsumoto T, Takegawa

K. Intracellular trafficking and ubiquitination of the Schizosaccharomyces pombe

amino acid permease Aat1p. Microbiology. 2012 Mar 1;158(3):659–73.

33. Aspuria PJ, Tamanoi F. The Tsc/Rheb signaling pathway controls basic amino acid
uptake via the Cat1 permease in fission yeast. Mol Genet Genomics. 2008
May;279(5):441–50.
34. Pérez-Díaz AJ, Vázquez-Marín B, Vicente-Soler J, Prieto-Ruiz F, Soto T, Franco A, et
al. cAMP-Protein kinase A and stress-activated MAP kinase signaling mediate
transcriptional control of autophagy in fission yeast during glucose limitation or
starvation. Autophagy. 2023 Apr 3;19(4):1311–31.
35. Kumar J, Ng K, Rallis C. Rapalink-1 reveals TOR-dependent genes and an
agmatinergic axis-based metabolic feedback regulating TOR activity and lifespan in
fission yeast. Commun Biol. 2025 Sept 29;8(1):1364.

Figure legends

Fig. 1 Torin1 induces Ght5 degradation. **A** Mutations in ESCRT complex protein subunits

induce resistance to Torin1. Subunits in dark green were identified independently of our drug

screening. The resistance of *atg6* and *vps38*, but not *atg38* mutants links ESCRT signalling to

Torin1 sensitivity. **B** Wt, *hse1Δ*, *vps25Δ* and *vps28Δ* mutants were grown to stationary phase

and then cultured in fresh YES media for at least 2h. Cells were adjusted to an $O.D._{600}$ of 0.3

to 0.5 and plated on YES agar with or without 5µM Torin1 and incubated at 32°C for 2-3 days.

C Cells expressing Ght5-GFP were grown to log phase and exposed to 5µM Torin1 for 3h or

as indicated. Cell pellets were stored at -80° C and analysed by SDS-PAGE. GFP expression

was monitored by immunoblot analysis using monoclonal antibodies directed against GFP.

Full-length Ght5-GFP and cleaved GFP is indicated. Ponceau S was used to monitor gel

loading. **D** Quantification of Torin1-induced Ght5-GFP degradation. Experiments were carried

out as in **C** (*n*= 3). **E** Cells were grown to log phase and incubated with 10mM caffeine,

100ng/mL Rapamycin alone or in combination, 150mM NaCl or 5µM Torin1 for 3h. **F** Cells

were grown to log phase and transferred into fresh media with 3% glucose, 0.03% glucose or

EMM2-N for 3h. Samples were analysed as in D. **G** Cells expressing Ght5-GFP were grown

to log phase, exposed to 5μM Torin1 for 1 and 3h and examined by fluorescent microscopy.

White arrowheads indicate septa. Scale bars indicate 10μm.

Fig. 2 Torin1 induces ESCRT-dependent Ght5 degradation. **A** *Wt* cells expressing Ght5-GFP

were exposed to 5μM Torin1 for 3h and samples analysed by immunoblot analysis. Numbers

indicate relative expression levels. **B** *hse1Δ* and *sst4Δ* mutants were exposed to 5μM Torin1

for 3h and analysed as in A. **C** Quantification of Torin1-induced Ght5 degradation in wt and

hse1Δ cells (*n*=3). **D** Wt and *vps25Δ* mutants were treated as in A. **E** Quantification of Torin1-

induced Ght5 degradation in wt and *vpsΔ* cells (*n*=3). **F** Regulation of nutrient transporter

stability by TORC2. Torin1 inhibits TORC2, blocks Gad8 activity and induces Gsk3 activation.

Gsk3 stabilizes Pub1, which targets Cdc25 for ubiquitin-dependent proteasomal degradation.

Inhibition of Gad8 facilitates arrestin-related adaptor-mediated ubiquitylation of Aat1, Cat and

Ght5 by Pub1 and Pub3. TORC2 also regulates Aat1 and Cat1 expression by regulating the

activity of Isp7. **G** *Wt* and *gsk3Δ* mutants were exposed to 5μM Torin1 as indicated. Samples

were processed as in A. **H** Quantification of Torin1-induced Ght5 degradation in *wt* and *gsk3Δ*

mutants (*n*=1). **I** *Wt* and *gsk3Δ* mutants expressing Ght5-GFP were exposed to 5μM Torin1 for

3h and examined by fluorescent microscopy. Scale bars indicate 10μm.

Fig 3. ESCRT mutants maintain Ght5 expression at the PM following exposure to Torin1. **A**

Wt **B** *hse1Δ*, **C** *vps25Δ*, **D** *sst4Δ* and **E** *sst6Δ* strains were examined by fluorescent microscopy.

Cells were exposed to 5μM Torin1 for 3h. White arrowheads indicate Ght5 expression at the

PM. Scale bars indicate 10μm.

Fig 4. Torin1 induces Aat1 degradation in an ESCRT-dependent manner. **A** Cells expressing

Aat1-GFP were exposed to 10mM caffeine, 100ng/mL Rapamycin alone or in combination,

5μM Torin1 or EMM2-N for 3h. Samples were analysed by immunoblotting. Full-length Aat1-

GFP and cleaved GFP are indicated. **B** *pub1Δ* mutants expressing Aat1-GFP were treated as

in A. **C** Wt cells expressing Aat1-GFP were exposed to 5μM Torin1 for the indicated timepoints

and treated as in A. **D** *Wt* and *pub1Δ* mutants expressing Aat1-GFP were treated as in C. **E**

Wt and *gsk3Δ* mutants expressing Aat1-GFP were treated as in C. **F** Quantification of Aat1-

GFP expression in *wt* and *gsk3Δ* mutants exposed to 5μM Torin1 for 3h (n=3). **G** *Wt* and *pub1Δ*

mutants expressing Aat1-GFP were exposed to 5μM Torin1 for 1 to 3h and examined by

fluorescent microscopy. Scale bar indicates 10μm. **H** *Wt* and *gsk3Δ* mutants expressing Aat1-

GFP were examined by fluorescent microscopy.   Scale bar indicates 10μm. **I** *Wt, gsk3Δ* and

pub1Δ mutants expressing Aat1-GFP were examined by fluorescent microscopy.   White

arrowheads indicate septa. Scale bar indicates 10μm.

Fig 5. Torin1 induces Cat1 degradation in a Pub1-dependent manner. **A** Cells expressing

Cat1-GFP were exposed to 5μM Torin1 for the indicated time points. Alternatively, the cells

were exposed to 10mM caffeine, 100ng/mL Rapamycin alone or in combination, 5μM Torin1

or EMM2-N for 3h. Samples were analysed by immunoblotting. Full-length Cat1-GFP and

cleaved GFP are indicated. **B** *Wt* and *pub1Δ* mutants expressing Cat1-GFP were exposed to

5μM Torin1 for the indicated timepoints. Samples were treated as in A. **C** *Wt* and *gsk3Δ*

mutants expressing Cat1-GFP were exposed to 5μM Torin1 for the indicated timepoints.

Samples were treated as in A. **D** Quantification of Aat1-GFP expression in *wt* and *gsk3Δ*

mutants exposed to 5μM Torin1 for the indicated timepoints (n=3). **E** *Wt* and *pub1Δ* mutants

expressing Cat1-GFP were exposed to 5μM Torin1 for 3h and examined by fluorescent

microscopy.   White arrowheads indicate expression of Cat1 at the PM. Scale bar indicates

10μm.

Fig 6. TORC2 and Pub1 integrate nutrient availability with cell division. **A** *Wt*, *gad8Δ*, *gsk3Δ*,

pub1Δ, *pub2Δ*, *pub2Δ pub3Δ* strains were grown to stationary phase, diluted into fresh media

at an O.D.$_{600}$ of 0.3 and incubated for 2h. Cells were serially diluted and plated on YES media

with 10mM caffeine and 100ng/mL rapamycin or 5μM Torin1 and incubated for 3-5 days. **B**

Wt and *gsk3Δ Cdc25(12A)-GFP* strains were incubated with 5μM Torin1 for the indicated time

points. Cell lysates were analyzed by immunoblotting. Full-length Cdc25(12A)-GFP and

cleaved GFP are indicated. **C** TORC1 and TORC2 adjust nutrient transporter expression to

cell cycle regulation. Environmental and nutrient stress inhibit TORC1-mediated Gad8

inhibition, resulting in Gsk3 activation. Gsk3 stabilises Pub1 which is required for the

ubiquitylation and degradation of PM nutrient transporters. Gad8 inhibits the arrestin-related-

mediated ubiquitylation of Ght5 by Pub1. Low endogenous ubiquitin levels prevent optimal

target protein degradation. TORC2 also positively regulates Cdc25 stability and is required for

cell cycle re-entry following activation of DNA damage checkpoints. TORC1 and TORC2 also

interact with Ssp1 and Ssp2 which also regulate cell cycle kinetics.

Additional file 1 A Wt, *bro1Δ*, *sst2Δ* and *vps8Δ* strains expressing GFP-Atg8 were grown to

log phase and exposed to 5μM Torin1 for 2h. Cell pellets were resolved by SDS-PAGE and

blotted with monoclonal antibodies directed against GFP. **B** Wt, *bro1Δ*, *did2Δ*, *sst2Δ* and *vps8Δ*

strains were grown to stationary phase, diluted into fresh media and incubated for a further

2h. Cells were serially diluted and plated on YES media with 3% or 0.03% glucose with or

without 5μM Torin1 and incubated for 3-5 days.

Additional Table 1 *S. pombe* strains used in this study.

Fig. 1

A

B

wt
hse1Δ
vps25Δ
vps28Δ

C

WT | *tor1Δ* | Torin1

Control | Caffeine | Torin1 | Control | Caffeine | Torin1 | 0 | 1 | 2 | 3 | 4h

Ght5- GFP

GFP

Ponceau S

D

Control | Torin1

p= 0.003

E

Control | Caffeine | Rapamycin | Caffeine+ rapamycin | NaCl | Torin1

Ght5- GFP

GFP

Ponceau S

F

Control | 0.03 % glucose | - Nitrogen

Ght5- GFP

GFP

Ponceau S

G

Control | Torin1 (1h) | Torin1 (3h)

Caffeine | Rapamycin | Caffeine+ Rapamycin

Ght5- GFP

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

A

B

C

Additional file 1

A

wt   bro1Δ   sst2Δ   vps8Δ

Torin1:   −   +   −   +   −   +   −   +

GFP-Atg8
GFP

Ponceau S

B

3% Glucose   3% Glucose+ torin1   0.03% Glucose   0.03% Glucose+ torin1

wt
bro1Δ
did2Δ
sst2Δ
vps8Δ
wt
tor1Δ

C

Torin1   Torin1

0   1   2   3   0   1   2   3h

Ght5- GFP

GFP

Ponceau S

wt   pub1Δ

Additional Table 1. List of fission yeast strains used in this study.

Strain	Genotype	Source/Reference
h⁻ L972	*h-*	Laboratory stocks
tor1Δ	*h- tor1::kanMX6*	Laboratory stocks/(Rallis *et al*., 2013)
gad8Δ	*h- gad8:: ura4 ade6- M216 leu1 ura4- D18*	YGRC
gsk3Δ	*h− gsk3::kanMX6*	Laboratory stocks/ (Rallis *et al*., 2017)
lub1Δ	*h+ lub1::KanMX ade6-M216 ura4-D18 leu1-32*	Laboratory stocks
ght5-GFP	*h- gad8ᵀˢ ght5-GFP::KanR*	Laboratory stocks/ (Saitoh *et al*., 2015)
ght5-GFP gad8ᵀˢ gsk3Δ	*h- gad8ᵀˢ gsk3::natMX ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2021)
ght5-GFP gad8ᵀˢ pub1Δ	*h- gad8ᵀˢ pub1::hphR ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2025)
ght5-GFP gad8ᵀˢ hse1Δ	*h- gad8ᵀˢ hse1::hphR ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2021)
ght5-GFP gad8ᵀˢ vps25Δ	*h- gad8ᵀˢ vps25::hphR ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2021)
ght5-GFP gad8ᵀˢ vps28Δ	*h- gad8ᵀˢ vps28::hphR ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2021)
ght5-GFP gad8ᵀˢ sst4Δ	*h- gad8ᵀˢ sst4::hphR ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2021)
ght5-GFP gad8ᵀˢ sst6Δ	*h- gad8ᵀˢ sst6::hphR ght5-GFP::KanR*	Laboratory stocks/ (Toyoda *et al*., 2021)
aat1-GFP	*h- aat1-GFP::KanR*	Matsumoto Laboratory
aat1-GFP gsk3Δ	*h- aat1-GFP<LEU2 leu1-32 gsk3::KanMx6*	This study
aat1-GFP pub1Δ	*h- pub1:: ura4+ ura4-D18 aat1-GFP<<LEU2 leu1-32*	Matsumoto Laboratory
cat1-GFP	*h- cat1+-GFP::kanR*	Nakashima laboratory

cat1-GFP gsk3Δ	*h- gsk3::natMX cat1+-GFP-kanMX*	This study
cat1-GFP pub1Δ	*h- pub1::hphMX cat1+-GFP-kanMX*	Nakashima laboratory
cdc25 (12A)- GFP	*h- cdc25(12A)-GFPint cdc25:: kanMX6 ura4-D18 leu1-32*	Young laboratory
cdc25 (12A)- GFP *gsk3Δ*	*h- cdc25(12A)-GFPint cdc25:: kanMX6 gsk3::natMX6 ura4-D18 leu1-32*	This study
pub1Δ	*h-* pub1*::*ura4*+* leu1*-32* ura4*-D18*	YGRC
pub2Δ	*h+* pub2*::*ura4*+* ura4*-D18*	YGRC
pub2Δ pub3Δ	*h90* pub2*::*ura4*+* pub3*::*ura4*+* ade6*-M216* leu1*-32* ura4*-D18*	YGRC
GFP-atg8		Laboratory stocks
GFP-atg8 bro1Δ		Laboratory stocks
GFP-atg8 sst2Δ		Laboratory stocks
GFP-atg8 vps8Δ		Laboratory stocks

*Yeast Genetic Resource Center (YGRC) Japan