Persist or Give up? Fire ants motivated to search for a high-quality food source even if they don't know how to find it

Chinmay Hemant Joshi [1], and Anna Dornhaus [1]

[1] Department of Ecology & Evolutionary Biology, University of Arizona, Tucson, Arizona

Abstract

Finding resources for the colony is one of the most difficult and risky tasks for a social insect worker. A worker on a foraging trip can face a number of challenges, including interference from other individuals, her own errors, and environmental disturbances. Collectively, colonies may use a variety of strategies to minimize the impact of such perturbations on the foraging process. Here, we investigated how individual *Solenopsis xyloni* ant workers react to perturbation of an established pheromone trail. We trained foragers from colonies in the field to either a low or high concentration sucrose solution in a feeder on a T-maze setup, then replaced a section of floor covering, removing a section of the pheromone trail previously laid. We found that while ants made correct choices on the T-maze when the trail was intact, their choices did not differ from chance when the trail was absent, indicating strong reliance on a pheromone trail (and not, for example, memory) to return to the resource. Moreover, when the trail was absent, we found that a majority of ants abandoned the resource, and that even the ants that were able to reach the resource did not repair the perturbed trail. However, with a high-quality resource, more ants persisted in attempting to reach it (instead of abandoning). We interpret these responses in the framework of robustness mechanisms discussed in systems biology. Our study thus links individual and collective responses to perturbations, and provides an empirical example of how information use interacts with system robustness.

Keywords: Collective foraging, ants, robustness, and behavioral ecology.

Introduction

Social insects are complex adaptive systems that perform various tasks, such as foraging cooperatively (Bonabeau 1998; Garnier et al. 2007). Worker behavior has evolved to maximize the collective foraging performance of the colony (Wilson 1971; Garnier et al. 2007; Doering et al. 2024). Individual workers in a colony can use a variety of ways to communicate information about food sources to their nestmates, often leading to recruitment towards the food source (Frisch 1967; Hölldobler and Wilson 1990; Nieh 1998; Czaczkes et al. 2015; Leonhardt et al. 2016; Ganeshaiah 2023). These communication systems differ in several aspects, including signal modality, information content, how signals are reinforced to sustain recruitment, and how they are adapted to the local environmental conditions (e.g. Lanan 2014; Leonhardt et al. 2016). However, communication systems can be affected by sudden changes (i.e., perturbations) in the environment. Perturbations such as fluctuations in air pressure (Sujimoto et al. 2020), rain

(Farji-Brener et al. 2018), surface temperature (van Oudenhove et al. 2011), and wind (Alma et al. 2016) can affect communication among workers and thereby collective foraging performance. Specifically, these perturbations can severely affect workers directly (Alma et al. 2016; Farré-Armengol et al. 2016; Farji-Brener et al. 2018), affect their communication signals (van Oudenhove et al. 2011, 2012; Alma et al. 2022) or ability to amplify these signals to sustain recruitment (van Oudenhove et al. 2012; Zhang et al. 2020), or affect the food source itself (Farji-Brener et al. 2018).

Robustness is an emergent, system-level property that allows a system, such as an ant colony, to maintain its performance when subjected to different perturbations (Kitano 2004; Masel and Siegal 2009). Robustness arises from how its components interact with each other and with the environment around them (Kitano 2004, 2007; Wagner et al. 2007; Whitacre 2010; Nijhout et al. 2019). Kitano (2004), in his seminal conceptual review, proposed the following main mechanism categories, which are thought to apply across a variety of complex systems: (1) *feedback loops*, (2) *buffering,* (3) *fail-safes,* and (4) *modularity*. *Feedback loops* (or 'system control') allow the system to return to its original state by counteracting the perturbation (negative feedback loop) or by moving the system to another, more suitable state given the new environment (positive feedback loops). *Buffers* (or 'decoupling') entail having more capacity in existing functional units than normally needed, so that the system can withstand a perturbation. *Fail-safes* (or 'redundancy and diversity') involve the existence of additional functional units that can take over for any failed components. Finally, *modularity* entails having separate (yet interconnected) modules that allow the system to restrict negative effects to a particular module to prevent their spread to other modules. These mechanisms have been identified and studied primarily in genetic networks and biochemical pathways across different organisms (Alon et al. 1999; Freeman 2000; Conant and Wagner 2004; Fritsche-Guenther et al. 2011; Holme 2011), within engineered systems (Tell et al. 2022; Kiakojouri et al. 2023), and in designed (human) organizational frameworks (Xie et al. 2022; Duvald et al. 2025).

Both molecular systems and social insect colonies contain individual components whose actions and behaviors have been optimized at the collective level (Seeley 2002; Kitano 2004; Garnier et al. 2007). Therefore, the systems-level thinking developed in molecular systems can potentially be applied to understand how social insects respond to different perturbations (and vice versa to study molecular systems). Social insects use various strategies to maintain foraging performance that fall under one or more of the above categories. Workers may use inhibitory signals or modulate positive communication to steer nestmates away from exhausted and/or suboptimal food sources (Evison et al. 2008; Czaczkes and Heinze 2015; Alma and Buteler 2019) - both examples of feedback loops maintaining a kind of homeostasis in foraging. Alternatively, workers can flexibly alter their behavior to continue foraging amidst perturbations (strong wind gust, for instance) (Gordon et al., 2013; Sujimoto et al., 2020; Alma et al., 2022), an example of decoupling. Sampling multiple sources of the same signal (Tanner and Visscher 2008) or combining information from multiple information sources (Czaczkes et al., 2011, 2013; Letendre & Moses, 2013) could likewise be seen as an example of buffering (more capacity

available than typically needed). If insects on the other hand switch between information sources when one becomes unreliable, this could be considered a fail-safe mechanism: such as when bees switch to using their memory when social signals are unreliable (Grüter et al. 2011; Dunlap et al. 2016; Jones et al. 2019).

Our aim in this study was to identify mechanisms used by social insects to maintain their collective performance in the face of environmental perturbations. Specifically, we focus on how ants deal with the disruption of communication during foraging, and which of the categories of robustness mechanisms identified in systems research broadly may be most common in social insects. We used southern fire ants (*Solenopsis xyloni*) as a model system for this study, and disrupted their foraging by removing a section of their pheromone trail to a food source. We then examine the responses of individual ants. We test the following specific hypotheses, which are matched to the types of robustness mechanisms proposed by Kitano 2004: (1) Ants use a fail-safe mechanism, i.e., an alternative mechanism to locate the food source, such as memory, (2) Ants employ a buffer mechanism in the face of pheromone trail disappearance, such as continuing to walk in the same direction as the original trail to relocate the food source, (3) A disrupted trail causes negative feedback (i.e., compensatory behavior), such as increased deposition of trail pheromone on the path to food, (4) Ants abandon the perturbed trail to redirect towards another food source nearby, and (5) Finally, it is also possible that ants show little robustness to disruption of their trail communication, and that foraging performance is severely impacted. We use a classic T-maze setup to examine these hypotheses, and a food source of low or high value to test whether increased food source value results in different strategies used by ants.

Methods

Study area and Study species

The southern fire ant (*Solenopsis xyloni*, McCook (Myrmicinae)) is one of the most common ant species across the Southwestern U.S (MacKay and Mackay 2002). *Solenopsis xyloni* nests in soil or under stones in open areas and near the edges of lawns in urban and semi-urban areas (Trager 1991; MacKay and Mackay 2002). *Solenopsis xyloni* relies upon trail pheromones to recruit nestmates to food sources (Barlin et al. 1976; Tschinkel 2013a). This pheromone is produced in the Dufour's gland and extruded via their sting, which is both connected to the Dufour's and venom glands (Barlin et al., 1976). Individual fire ants press the stinger against the ground and drag it on their way to the nest from a food source (Hangartner, 1969, and personal observations in *S. xyloni*). We conducted this study at the University of Arizona campus (Tucson, Arizona, 32.23ºN, 110.95ºS). In total, we used 18 ant colonies across four years (July-November, 2022-2025).

Experimental setup

We used a laser-cut plexiglass T-maze (arm length = 12.6 cm, stem length = 8.6 cm, thickness = 0.32 cm) supported by cardboard columns (diameter = 0.80 cm) attached to small Petri dishes (diameter=3.3 cm) (Fig. 1a). To ensure that ants only entered and exited through one point in the setup, we applied Fluon (Company name: byFormica) on the underside of the T-maze and the ramp, and filled the petri dishes with water. The removable ramp allowed us to restrict the entry of ants during the experimental trials (such that we could isolate a single focal ant, see below). We cut printer paper to the dimensions of the 'T' of the maze, and cut it into sections so it could be selectively removed (Fig. 1b, and see below for more details). We attached these paper sections as surface covering using washable glue (Elmer's Repositionable Washable Glue Stick). Lines were lightly drawn on the paper with black pencil to mark key measurement locations (Fig. 2a). We used an Eppendorf tube (1.5 ml) as a feeder by punching small holes on the side near the cap, and attaching it upside down to the maze with a small lump of putty to stabilize the feeder (Fig. 1b). The feeder was filled with a sucrose in water solution (either 1.5 M or 3 M sucrose).

Experimental procedure

We conducted all experiments between 9:30 am and 12 pm and 3:00 pm and 7:00 pm (July-November, 2022-25), when the ants were most active. We used two experiment types that only differed in the sucrose concentration used in the feeder (1.5 M (n = 116 trials) and 3 M (n = 86 trials); hereafter referred to as low sucrose and high sucrose, respectively). Both experiment types included trials in the control and experimental treatment (see 'Test phase' below). Each trial consisted of a training phase followed by a test phase.

Training phase

We kept the T-maze ~1m away from the colony's nest entrance (the exact location near the nest and its position randomly chosen before every trial, Fig. 1a). We kept the feeder (filled with low or high sucrose solution) on the ground close to the colony entrance and allowed the ants to feed for 5 min. We then moved the feeder onto the T-maze (with ramp access) for ants to access the feeder for another 5 min. Following that, we moved the feeder one last time to either of the T-maze arms randomly and allowed the colony to forage for another 10 min. This allowed the colony to lay a pheromone trail on the paper overlay towards the arm containing the feeder.

Test phase

Following the 20-minute training phase, we removed the ramp and gently brushed off all the ants from the setup (including the feeder). Using nitrile gloves, we then removed the middle part of the T-shaped paper overlay (Length = 5.1 cm, Width = 4 cm) and replaced it in the same position (control treatment) or with a fresh paper section (experimental treatment). Thus, the control trials contained the same pheromone trail as laid by the ants in the training phase, but in experimental trials, there was a gap in the trail where the paper was replaced. We pointed the camera (Nikon D7000 DSLR) at a constant distance perpendicular to the setup so that it recorded the side-view

of the majority of the T-maze stem (including the specific area where the paper had been replaced). Then, using the ramp, we allowed only a single ant to enter the setup. For every individual trial, once the focal ant entered the setup, we removed the ramp. We started the video recording once the ant crossed the 'line of commitment' (Fig. 2a). Once the ant returned and crossed the 'line of commitment' again, we stopped the recording. For the experimental treatment across both experiment types, we tested at most three ants in succession using the same pheromone trail. This was possible because we put a fresh paper section on the T-maze for each experimental trial. However, for the control treatment, since we replaced the same paper section (with the trail on the paper), we tested just one ant per pheromone trail laid during the training phase. For both treatments, once the trial ended, we gently removed the ant using a paintbrush and placed it slightly away (>1m) from the colony to prevent it from attempting to return to the T-maze immediately.

Data collection

We defined four relevant 'lines', drawn on the T-maze with pencil, to operationally assess the decisions made by foraging ants (Fig. 2a). The section of the paper, and thus the pheromone trail, removed in experimental trials was the one between the 'line of disturbance' and the two 'choice' lines. We defined an ant as having decided to forage once it crossed the 'line of commitment' and walked onto the maze (at which point we started recordings). We defined an ant as having made a choice when it crossed either the 'line of choice' towards the left or right arm of the T-maze. For each individual ant, we recorded (1) the first choice made (crossed either of the line of choices), (2) whether it abandoned the setup (turned back between the 'line of choice' and 'line of disturbance', i.e., never crossed a 'line of choice'), and (3) the angle at which it held its gaster while walking between the 'line of choice' and the 'line of disturbance' on the way back from the top of the T-maze (extracted from the video). In ant species that use pheromone trail communication, workers typically bend/curve their gaster in a species-stereotypical fashion when laying a pheromone trail (Hangartner 1969; Hölldobler and Wilson 1990; Beckers et al. 1992; Jackson and Châline 2007). During this bending, the angle between the gaster and the vertical decreases compared to when the ant is not laying a trail (Wilson 1962; Beckers et al. 1992). Fire ant workers typically drag their gaster continuously on the ground (to lay pheromone trail) when they are returning to their nest from a food source (Wilson 1962; Hangartner 1969). We therefore used our measure of 'gaster angle' as a proxy for trail-laying behavior. We used Adobe Premiere Pro to extract all video frames in which the ant was between the line of choice and the line of disturbance on the T-maze, on the way back from the food source. Then we used ImageJ to quantify the gaster angle on every third frame. Fig. 1c illustrates the entire experimental protocol, and the Supplementary File S1 contains a more detailed explanation on the video analysis pipeline.

Statistical analysis

We used Chi-square tests to compare the number of choices for either maze arm and the number of ants abandoning vs. staying on the setup between treatments. We used a binomial test to compare the number of ants choosing the left and right maze arms, and either abandoning or staying on the trail (i.e., making either choice) to a null expectation of random (50%) choice, within each treatment. We averaged the gaster angles across all frames measured for one ant (Range: 1-58 frames) for each trial. We then ran Kruskal-Wallis, Dunn posthoc, and Mann-Whitney U-Tests to compare these average gaster angles across ants making correct versus wrong choices between the control and experimental treatments. All data were analyzed in R (v. 4.5.2, R Core Team, 2024). Detailed sample sizes for each experiment are given in Table 1. The original data and the R script are made available in the Supplementary section and this GitHub repository: https://github.com/chjoshi24/Ch1-robustness-mechanisms-paper-2025

Fig. 1 Experimental setup and procedure. (a) Camera and T-maze setup in the field, with measuring tape to determine the distance of the camera. (b) T-maze with feeder, ramp, and petri dish 'moats'. (c) Schematic of the procedure. In the training phase, we allowed the whole colony to feed on the sucrose solution for 20 minutes in total (5 minutes near the colony entrance, 5 minutes on the T-maze stem, and 10 minutes on either arm of the T-maze that was selected

randomly). We then brushed off the ants from the setup and feeder, and either applied the control treatment (removing and replacing the same paper section, highlighted with orange) or the experimental treatment (removal and replacement with fresh paper of the same section). Following that, we allowed one ant at a time to enter the setup, started a video recording, and recorded whether the ant made a choice or abandoned it. The video recording was stopped when the ant exited the setup (crossed the 'line of commitment'). The wiggly lines represent the pheromone trail. All the objects here (T-maze, ants, camera, brush, and pheromone trail) are schematics and not drawn as per the relative sizes of the objects in reality.

Fig. 2 (a) Schematic of the pencil-drawn lines on the T-shaped paper cutout with measurements for each section. (b) Schematic of the measurements done during the experiments. We quantified three things for every ant: The first choice (left or right, i.e. correct or incorrect depending on feeder placement) or whether the ant abandoned the food source (turned back without crossing a choice line), and the average gaster angle on the return trip from a choice (while within orange area). The wiggly line represents the pheromone trail as present in the experimental treatment. Objects in Fig. 2b are not drawn to scale.

Results

In total, we tested 202 ants from 18 colonies across four field seasons (2022-2025).

Ant choices

Individually tested ants strongly preferred the arm of the T-maze that contained the food source, i.e. made a 'correct' choice (90% of ants in low sucrose treatment, 82% of ants in high sucrose treatment; Binomial tests: low sucrose - p<0.001, n = 40; high sucrose - p<0.001, n = 33; Fig. 3). However, when the pheromone trail was removed, the ants did not significantly prefer either direction (i.e., made 'correct' decisions at chance levels: 50% in low, 52% in high sucrose treatment; Binomial tests: low sucrose - p = 1, n = 16; high sucrose - p = 1, n = 21). In each condition (low vs high sucrose), the proportion of correct choices was higher in the control than the experimental (pheromone trail perturbed) treatment (Chi-square tests: low sucrose - n=56, χ^2 = 10.86, df = 1, p<0.001; high sucrose - n = 54, χ^2 = 5.33, df = 1, p = 0.02). We did not find an effect of food quality (sucrose concentration) on the proportion of correct choices for either control (Chi-square test: n = 73, χ^2=1.02, df = 1, p = 0.31) or experimental treatments (Chi-square test: n = 37, χ^2 = 0.02, df=1, p = 0.89).

Ants giving up the food source

In both low and high sucrose experiments, we found a significant relationship between the treatment and the proportion of ants that abandoned the resource, with more ants abandoning when the trail was perturbed (low sucrose: 10% in control, 77% in experiment; high sucrose: 6% in control, 59% in experiment; Chi-square tests: low sucrose - n = 116, χ^2 = 48.57, df = 1, p<0.0001; high sucrose - n = 86, χ^2 = 25.06, df = 1, p<0.0001; Fig. 4). There was no difference between low and high sucrose experiments in the control treatment (Chi-square test: n = 80, χ^2 = 0.718, df = 1, p = 0.40). However, we found that abandonment was significantly lower in the high sucrose compared to the low sucrose experiments when the trail was removed (i.e., in the experimental treatment, Chi-square test: n = 122, $\chi2$ = 4.88, df = 1, p = 0.03).

Average gaster angle

Across both experiments, there were some ants that made a correct choice but did not feed on the sucrose, and some ants that first made a wrong choice but still fed on the sucrose (by moving across the other arm of the T maze to the feeder arm before exiting). We excluded both types of ants, only including ants that indeed reached the food after a correct choice and fed, and those that, if they made an initial wrong choice, did not reach the food (and did not feed), in the gaster angle analysis.

We found that ants that made a correct choice had a significantly lower mean gaster angle compared to the ones that made a wrong choice in the high sucrose experiment (Kruskal-Wallis test: n=31, χ^2 = 13.47, df = 2, p = 0.004; see Fig. 5 for the pairwise differences, Dunn's post hoc test). This indicates that our gaster angle measure indeed quantifies trail laying, and that ants

who reached a high-quality food source were more likely to lay a trail than those who did not. In the low-sucrose experiment, this was not the case (correctness of choice did not significantly affect gaster angle; however, we did not have any cases of ants making an incorrect choice in the control treatment for low sucrose; Fig. 5, Kruskal-Wallis test: n=36, $\chi2 = 2.21$, df = 2, p = 0.331).

For the ants that made correct choices within control treatments across both experiments, the gaster angle was significantly lower in the high sucrose experiment compared to the low sucrose (difference between categories indicated by asterisk in Fig. 5, Mann-Whitney-U test: n=46, W= 364, p = 0.004). However, the analogous difference was not significant for the experimental treatments (i.e. when the trail was removed, Mann-Whitney-U test: n=12, W = 20, p = 0.570; however, note the much lower sample size).

Fig. 3 Proportion of ants making correct vs. wrong choices in control and experimental treatments across two experiment types. Ants made more correct choices in the control treatments (i.e., when the trail was intact). The proportion of correct choices in either control or experimental treatments did not differ depending on food quality. The dotted line, at 50%, is the expectation if ants show no preference for either choice.

Fig. 4 Proportions of ants that abandoned vs. stayed on the trail (correct and wrong choices combined). Significantly more ants abandoned the food source in the experimental treatment compared to the control across both experiments. There was a significant reduction in rate of abandonment in the high sucrose experiment compared to the low sucrose one in the experimental treatment, but not in the control.

Fig. 5 Gaster angle (averaged, for each ant, across video frames) in different treatments across low and high sucrose experiments. Only ants that made a correct choice and did feed on the feeder (red boxes), and ants that made a wrong choice and did not feed on the feeder (blue boxes), are included here. The year of the experiment is indicated by the color of the data points. We did not see a significant effect of treatment (control, with trail intact, or experiment, with trail removed). We also did not find a difference in gaster angles between ants that made correct or wrong choices in the low sucrose experiment. However, in the high sucrose experiment, ants that made a correct choice had a significantly lower gaster angle, i.e. were more likely to lay a new recruitment trail, compared to the ants that made a wrong choice. For the high sucrose experiment, the boxes that do not share the same letter are significantly different from each other (significant Kruskal-Wallis test followed by Dunn's post hoc test). Additionally, in the control treatments across both experiments, the gaster angle was significantly lower for the high sucrose experiment compared to the low sucrose, but no such difference was found for the experimental treatments (Mann Whitney U-tests, differences indicated by the asterisks).

Discussion

We found that when confronted with an apparent dead-end in a pheromone trail, individual *S. xyloni* workers primarily reacted by turning back (Fig. 4). Those ants that continued towards the food source made random choices on the T-maze bifurcation in the absence of pheromone trail information (Fig. 3), suggesting that they were unable to use backup information sources such as their own memory or long-distance cues like vision or odor to locate the food source. When ants foraged on a more valuable food source, they remained more committed and did not turn around as often even in the absence of a pheromone trail; this was despite the fact that they still appeared to make random choices on the T-maze regardless of food source quality (Fig. 3). Our measure of gaster angle was lower when foragers were returning from high quality food rather than from an incorrect choice (and thus an unsuccessful trip) (Fig. 5). This confirms that the measure of gaster angle indeed indicates trail-laying. However, foragers equally reinforced the perturbed trail across both food qualities overall, and did not reinforce the trail more if it had disappeared (in control vs experiment).

Overall, our study thus showed a surprising lack of robustness to pheromone trail perturbation in fire ants. This failure indicates that individual *S. xyloni* may be highly reliant on the pheromone trail for navigating their way towards food sources, instead of using trails only to recruit naive ants (and then using their own memory to return to known food sources). Given that the pheromone trail serves as an externalized memory of the food source location, there may be a low selection pressure for every forager to be successful at remembering the food source location (Feinerman and Traniello 2016). Studies from other fire ant species have shown that foragers can in principle use landmarks to navigate and memorize food locations, and pheromone trail presence can speed up this process (Stratton and Coleman 1973; Vander Meer et al. 2023). However, one of these studies shows that *Solenopsis saevissima* foragers took significantly longer to reach a food source in a multi-turn maze without a pheromone trail compared to when the trail was present, indicating that the pheromone trail is important for orientation during foraging (Stratton and Coleman 1973). It is also possible that *S. xyloni* foragers need more time to memorize the food source locations. All of our training phases only lasted 20 minutes. In the study mentioned above, authors also found that the *S. saevissima* foragers become less error-prone over 15-21 days in navigating their way towards the food source without using trails (Stratton and Coleman 1973).

Solenopsis xyloni and many other mass recruiting species (including other species from the *Solenopsis* genus) typically have big colony sizes with aggressive, but individually small workers (Valone and Kaspari 2005; Cerdá et al. 2013; Tschinkel 2013a). This can enable colonies to monopolize a food source even against competing ants (Detrain and Deneubourg 2008; Lanan 2014; Price et al. 2016). It may be that in a foraging system like this, food resources are only valuable to the colony when mass recruitment and strong symmetry breaking is possible (Lanan et al. 2012; Cerdá et al. 2013; Price et al. 2016), and that colonies thus invest little in exploration. If that is the case, this would explain why foragers are not adapted to recover a food source individually when no trail can be or has been formed.

As noted above, the frequent abandonment of the food source by workers when the trail was perturbed might be interpreted as a lack of individual or collective robustness to this type of disturbance. In the robustness mechanism framework of Kitano 2004, we expected to find perhaps feedback (trail reinforcement or otherwise strengthened recruitment) or individual-level fail-safes (alternative information sources) in response to trail perturbation, but did not find either of these mechanisms in fire ants in our experiment. However, we could argue that an alternative interpretation to 'absence of robustness' is one of modularity or buffering. Ant colonies may maintain several concurrent, independent pheromone trails to different food sources, thereby constituting a modular system (Lanan, 2014; and personal observations in *S. xyloni*). Disturbance of one of these trails likely does not affect foraging on other trails, thus containing the perturbation to one 'module' and maintaining (some) food intake by the colony. On the other hand, ants that 'abandoned' our T-maze may also not have abandoned foraging altogether; they may have switched to another available trail or resource. If individual workers have a choice between multiple maintained or discoverable resources, this would constitute a kind of 'buffering' mechanism in Kitano's framework, where individual performance is only moderately impacted by the perturbation because of extra capacity. Future experiments are needed that involve tracking ants' decisions to further understand the downstream consequences of individual-level and food-source-specific abandonment on colony-level foraging performance (in ants and in general across social insects).

Despite an overall high reliance on trail pheromone, we did find that even an apparently complete lack of information about which way the food was located did not prevent about 41% of foragers from attempting to find it when the resource they expected was of high quality. This illustrates a tradeoff between the risks of exploration (searching for a food source that may or may not be found) and the potential gains of exploitation (sucrose reward when food is found).

Overall, we found an apparent lack of ability to compensate for a disturbed pheromone trail in fire ants in our study, indicating that mass recruiting ant species like *S. xyloni* may not prioritize robustness in this context. Possible reasons are a high reliance on strength in numbers and resulting low levels of individual reliability. This indicates a possible tradeoff between robustness, efficiency, and cost (Latty et al. 2011; Cook et al. 2014; Cabanes et al. 2015; Middleton and Latty 2016). Applying a general framework to categorizing robustness mechanisms allowed the identification of different possible approaches social insect colonies could take in dealing with perturbations of collective activity such as trail disruption. Application of such a framework may therefore be fruitful in a comparative context, by examining both different species (e.g., ones that differ in reliance on individual vs. mass foraging, or memory vs. communication) and examining different types of collective activity (e.g. foraging vs. task allocation vs. colony defense). Such endeavors may allow us to understand when and how robustness is prioritized and realized by collective systems of different types. Such studies will contribute to identifying the rules of life that are common to different scales of organization.

References

Alma AM, Buteler M (2019) Plant odors trigger clearing behavior in foraging trails- do they represent olfactory obstacles? Behav Processes 169:103989. https://doi.org/10.1016/j.beproc.2019.103989

Alma AM, Buteler M, Martinez A, Corley J (2022) Wind disrupts trail pheromone communication in the leaf-cutting ant Acromyrmex lobicornis. Anim Behav 192:39–49. https://doi.org/10.1016/j.anbehav.2022.07.007

Alma AM, Farji-Brener AG, Elizalde L (2016) Collective Response of Leaf-Cutting Ants to the Effects of Wind on Foraging Activity. Am Nat 188:576–581. https://doi.org/10.1086/688419

Alon U, Surette MG, Barkai N, Leibler S (1999) Robustness in bacterial chemotaxis. Nature 397:168–171

Barlin MR, Blum MS, Brand MM (1976) Fire ant trail pheromones: Analysis of species specificity after gas chromatographic fractionation. J Insect Physiol 22:839–844. https://doi.org/10.1016/0022-1910(76)90253-5

Beckers R, Deneubourg JL, Goss S (1992) Trail laying behaviour during food recruitment in the antLasius niger (L.). Insectes Sociaux 39:59–72. https://doi.org/10.1007/BF01240531

Bonabeau E (1998) Social Insect Colonies as Complex Adaptive Systems. Ecosystems 1:437–443. https://doi.org/10.1007/s100219900038

Cabanes G, Van Wilgenburg E, Beekman M, Latty T (2015) Ants build transportation networks that optimize cost and efficiency at the expense of robustness. Behav Ecol 26:223–231. https://doi.org/10.1093/beheco/aru175

Cerdá X, Arnan X, Retana J (2013) Is competition a significant hallmark of ant (Hymenoptera: Formicidae) ecology. Myrmecol News 18:131–147

Conant GC, Wagner A (2004) Duplicate genes and robustness to transient gene knock-downs in *Caenorhabditis elegans*. Proc R Soc Lond B Biol Sci 271:89–96. https://doi.org/10.1098/rspb.2003.2560

Cook Z, Franks DW, Robinson EJH (2014) Efficiency and robustness of ant colony transportation networks. Behav Ecol Sociobiol 68:509–517. https://doi.org/10.1007/s00265-013-1665-8

Czaczkes TJ, Grüter C, Ratnieks FLW (2015) Trail Pheromones: An Integrative View of Their Role in Social Insect Colony Organization. Httpsdoiorg101146annurev-Ento-010814-020627 60:581–599. https://doi.org/10.1146/ANNUREV-ENTO-010814-020627

Czaczkes TJ, Heinze J (2015) Ants adjust their pheromone deposition to a changing environment and their probability of making errors. Proc R Soc B Biol Sci 282:.

https://doi.org/10.1098/rspb.2015.0679

Detrain C, Deneubourg J-L (2008) Collective Decision-Making and Foraging Patterns in Ants and Honeybees. In: Advances in Insect Physiology. Elsevier, pp 123–173

Doering GN, Prebus MM, Suresh S, et al (2024) Emergent collective behavior evolves more rapidly than individual behavior among acorn ant species. Proc Natl Acad Sci 121:e2420078121. https://doi.org/10.1073/pnas.2420078121

Dunlap AS, Nielsen ME, Dornhaus A, Papaj DR (2016) Foraging Bumble Bees Weigh the Reliability of Personal and Social Information. Curr Biol 26:1195–1199. https://doi.org/10.1016/J.CUB.2016.03.009

Duvald I, Obel B, Lohse C, et al (2025) Organizational resilience in emergency healthcare: Lessons from the COVID-19 pandemic in five Danish hospitals. Soc Sci Med 366:117728. https://doi.org/10.1016/j.socscimed.2025.117728

Evison SEF, Hart AG, Jackson DE (2008) Minor workers have a major role in the maintenance of leafcutter ant pheromone trails. Anim Behav 75:963–969. https://doi.org/10.1016/j.anbehav.2007.07.013

Farji-Brener AG, Dalton MC, Balza U, et al (2018) Working in the rain? Why leaf-cutting ants stop foraging when it's raining. Insectes Sociaux 65:233–239. https://doi.org/10.1007/s00040-018-0605-z

Farré-Armengol G, Peñuelas J, Li T, et al (2016) Ozone degrades floral scent and reduces pollinator attraction to flowers. New Phytol 209:152–160. https://doi.org/10.1111/nph.13620

Feinerman O, Traniello JFA (2016) Social complexity, diet, and brain evolution: modeling the effects of colony size, worker size, brain size, and foraging behavior on colony fitness in ants. Behav Ecol Sociobiol 70:1063–1074. https://doi.org/10.1007/s00265-015-2035-5

Freeman M (2000) Feedback control of intercellular signalling in development. Nature 408:313–319. https://doi.org/10.1038/35042500

Frisch K von (1967) Dance language and orientation of bees

Fritsche-Guenther R, Witzel F, Sieber A, et al (2011) Strong negative feedback from Erk to Raf confers robustness to MAPK signalling. Mol Syst Biol 7:489. https://doi.org/10.1038/msb.2011.27

Ganeshaiah KN (2023) Recruitment Strategies and Foraging Patterns of Ants: What Shapes Them and Why? J Indian Inst Sci 103:1129–1141. https://doi.org/10.1007/s41745-023-00403-7

Garnier S, Gautrais J, Theraulaz G (2007) The biological principles of swarm intelligence. Swarm Intell 1:3–31. https://doi.org/10.1007/s11721-007-0004-y

Gordon DM, Dektar KN, Pinter-Wollman N (2013) Harvester Ant Colony Variation in Foraging Activity and Response to Humidity. PLOS ONE 8:e63363. https://doi.org/10.1371/journal.pone.0063363

Grüter C, Czaczkes TJ, Ratnieks FLW (2011) Decision making in ant foragers (Lasius niger) facing conflicting private and social information. Behav Ecol Sociobiol 65:141–148. https://doi.org/10.1007/s00265-010-1020-2

Hangartner W (1969) Structure and variability of the individual odor trail in Solenopsis geminata Fabr. (Hymenoptera, Formicidae). Z F�r Vgl Physiol 62:111–120. https://doi.org/10.1007/BF00298046

Hölldobler B, Wilson EO (1990) The ants. Harvard University Press

Holme P (2011) Metabolic Robustness and Network Modularity: A Model Study. PLOS ONE 6:e16605. https://doi.org/10.1371/journal.pone.0016605

Jackson DE, Châline N (2007) Modulation of pheromone trail strength with food quality in Pharaoh's ant, Monomorium pharaonis. Anim Behav 74:463–470. https://doi.org/10.1016/j.anbehav.2006.11.027

Jones S, Czaczkes TJ, Gallager AJ, et al (2019) Copy when uncertain: lower light levels increase trail pheromone depositing and reliance on pheromone trails in ants. Anim Behav 156:87–95. https://doi.org/10.1016/j.anbehav.2019.08.007

Kiakojouri F, De Biagi V, Abbracciavento L (2023) Design for Robustness: Bio-Inspired Perspectives in Structural Engineering. Biomimetics 8:95. https://doi.org/10.3390/biomimetics8010095

Kitano H (2004) Biological robustness. Nat Rev Genet 5:826–837. https://doi.org/10.1038/nrg1471

Kitano H (2007) Towards a theory of biological robustness. Mol Syst Biol 3:. https://doi.org/10.1038/msb4100179

Lanan M (2014) Spatiotemporal resource distribution and foraging strategies of ants (Hymenoptera: Formicidae)

Lanan MC, Dornhaus A, Jones EI, et al (2012) The Trail Less Traveled: Individual Decision-Making and Its Effect on Group Behavior. PLOS ONE 7:e47976. https://doi.org/10.1371/journal.pone.0047976

Latty T, Ramsch K, Ito K, et al (2011) Structure and formation of ant transportation networks. J R Soc Interface 8:1298–1306. https://doi.org/10.1098/rsif.2010.0612

Leonhardt SD, Menzel F, Nehring V, Schmitt T (2016) Ecology and Evolution of Communication in Social Insects. Cell 164:1277–1287. https://doi.org/10.1016/j.cell.2016.01.035

MacKay WP, Mackay E (2002) The ants of New Mexico (Hymenoptera: Formicidae). Edwin Mellen Press Lewiston, NY

Masel J, Siegal ML (2009) Robustness: mechanisms and consequences. Trends Genet 25:395–403. https://doi.org/10.1016/j.tig.2009.07.005

Middleton EJT, Latty T (2016) Resilience in social insect infrastructure systems. J R Soc Interface 13:. https://doi.org/10.1098/rsif.2015.1022

Nieh JC (1998) The role of a scent beacon in the communication of food location by the stingless bee, Melipona panamica. Behav Ecol Sociobiol 43:47–58. https://doi.org/10.1007/s002650050465

Nijhout HF, Best JA, Reed MC (2019) Systems biology of robustness and homeostatic mechanisms. Wiley Interdiscip Rev Syst Biol Med 11:. https://doi.org/10.1002/wsbm.1440

Price RI, Grüter C, Hughes WOH, Evison SEF (2016) Symmetry breaking in mass-recruiting ants: extent of foraging biases depends on resource quality. Behav Ecol Sociobiol 70:1813–1820. https://doi.org/10.1007/s00265-016-2187-y

R Core Team (2024) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/Accessed 23 September, 2024. Found Stat Comput Vienna Austria

Seeley TD (2002) When Is Self-Organization Used in Biological Systems? Biol Bull 202:314–318. https://doi.org/10.2307/1543484

Stratton LO, Coleman WP (1973) Maze learning and orientation in the fire ant (Solenopsis saevissima). J Comp Physiol Psychol 83:7–12. https://doi.org/10.1037/h0034261

Sujimoto FR, Costa CM, Zitelli CHL, Bento JMS (2020) Foraging activity of leaf-cutter ants is affected by barometric pressure. Ethology 126:290–296. https://doi.org/10.1111/eth.12967

Tanner DA, Visscher PK (2008) Do honey bees average directions in the waggle dance to determine a flight direction? Behav Ecol Sociobiol 62:1891–1898. https://doi.org/10.1007/s00265-008-0619-z

Tell S, Andersson A, Najafi A, et al (2022) Real-time hybrid testing for efficiency assessment of magnetorheological dampers to mitigate train-induced vibrations in bridges. Int J Rail Transp 10:436–455. https://doi.org/10.1080/23248378.2021.1954560

Trager JC (1991) A revision of the fire ants, Solenopsis geminata group (Hymenoptera: Formicidae: Myrmicinae). J N Y Entomol Soc 141–198

Tschinkel WR (2013a) The fire ants. Harvard University Press

Tschinkel WR (2013b) The Morphometry of Solenopsis Fire Ants. PLoS ONE 8:79559. https://doi.org/10.1371/JOURNAL.PONE.0079559

Valone TJ, Kaspari M (2005) Interactions between granivorous and omnivorous ants in a desert grassland: results from a long‑term experiment. Ecol Entomol 30:116–121. https://doi.org/10.1111/j.0307-6946.2005.00670.x

van Oudenhove L, Billoir E, Boulay R, et al (2011) Temperature limits trail following behaviour through pheromone decay in ants. Naturwissenschaften 98:1009–1017. https://doi.org/10.1007/s00114-011-0852-6

van Oudenhove L, Boulay R, Lenoir A, et al (2012) Substrate Temperature Constrains Recruitment and Trail Following Behavior in Ants. J Chem Ecol 38:802–809. https://doi.org/10.1007/s10886-012-0130-x

Vander Meer RK, Alonso LE, Lofgren CS (2023) Light Affects the Homing Ability of Solenopsis invicta (Hymenoptera: Formicidae) Foraging Workers. Fla Entomol 106:. https://doi.org/10.1653/024.106.0304

Wagner GP, Pavlicev M, Cheverud JM (2007) The road to modularity. Nat Rev Genet 2007 812 8:921–931. https://doi.org/10.1038/nrg2267

Whitacre JM (2010) Degeneracy: a link between evolvability, robustness and complexity in biological systems

Wilson EO (1971) The insect societies. Cambridge, Mass. : Belknap Press of Harvard University Press, Cambridge, Mass.

Wilson EO (1962) Chemical communication among workers of the fire ant Solenopsis saevissima (Fr. Smith) 1. The Organization of Mass-Foraging. Anim Behav 10:134–147. https://doi.org/10.1016/0003-3472(62)90141-0

Xie Y, Desouza KC, Jabbari M (2022) On organizational robustness: A conceptual framework. J Contingencies Crisis Manag. https://doi.org/10.1111/1468-5973.12423

Zhang ZY, Li Z, Huang Q, et al (2020) Honeybees (Apis mellifera) modulate dance communication in response to pollution by imidacloprid. J Asia-Pac Entomol 23:477–482. https://doi.org/10.1016/j.aspen.2020.03.011

Acknowledgments

We thank Michele Lanan for help with ant identification, Cycle Havel, Hayley Hananman, Tiffany Sessions, and Tyler Reese for contributions to data collection and video analysis, and the Staff at the Catalyst Studio, Main Library, University of Arizona, for help with T-maze laser cutting. This work was supported by the Department of Ecology and Evolutionary Biology, University of Arizona, the National Science Foundation (Grant No. DBI 1564521 to A.D.), and a project Grant from the Graduate and Professional Student Council, University of Arizona, to C.H.J.

Statements and declarations

We do not have competing or financial interests

Data availability

The annotated R code and all the data files are available on github repository (https://github.com/chjoshi24/Ch1-robustness-mechanisms-paper-2025). The detailed breakdown of year wise experimental data, information on datasheet columns, and video analysis pipeline can be found in Supplementary file S1.