15-Hydroxyeicosatetraenoic Acid and GPR39 Together Orchestrate Coronary Autoregulation: A Comprehensive Metabolomic Analysis

D. Elizabeth Le, M.D.[1,2]
Masaki Kajimoto, M.D., Ph.D.[1]
Yan Zhao, M.D.[1]
Carmen Methner, Ph.D.[1]
Agostino Cianciulli, Ph.D.[4]
Teresa Semeraro, Ph.D.[4]
Iuni Margaret Laura Trist, Ph.D.[4]
Fabrizio Micheli, Ph.D.[4]
Sanjiv Kaul, M.D.[1,3]
(Orchid ID 0000-0003-0124-3920)

From the [1]**Knight Cardiovascular Institute,** [2]**Veterans Administration Portland Health Care System, Oregon Health & Science University,** [3]**Vasocardea, Inc. Portland, Oregon, USA, and the** [4]**Discovery Chemistry Unit, Aptuit (Evotec), Verona, Italy**

Running title: 15-HETE and GPR39 in coronary autoregulation

Word count: 6792

Address correspondence to:

Sanjiv Kaul, M.D.
Knight Cardiovascular Institute
Oregon Health & Science University
UHN62
3181 SW Sam Jackson Park Rd.
Portland, OR 97239

E: kauls@ohsu.edu
T: 503-494-3889
F: 503-494-8550

Abstract

Background: Coronary autoregulation is the ability of the normal heart to maintain constant coronary blood flow (CBF) over a wide range of coronary driving pressures (CDP). Despite being vital for survival, the mechanism of coronary autoregulation is unknown. We hypothesized that GPR39, present in vascular smooth muscle cells, together with its endogenous agonist 15-hydroxyeicosatetraenoic acid (15-HETE) is responsible for coronary autoregulation.

Methods: We created coronary stenoses of varying degrees in open-chest, anesthetized dogs where we measured CBF and CDP. In a subset of animals, coronary venous blood was sampled for eicosanoid, adenosine, endothelin-1, polyunsaturated fatty acids, and prostaglandins levels. Stenoses were recreated during intravenous administration of VC108, a specific GPR39 antagonist and systemic, pulmonary and coronary hemodynamics measured.

Results: 15-HETE correlated linearly with CDP over the autoregulatory range (r^2=0.47, p=0.0024); none of the other metabolites had any relation with CDP. Prior to administration of VC108, as expected, CBF did not change with CDP in the coronary autoregulatory range. VC108 had no effect of systemic and pulmonary hemodynamics but increased CBF (p=0.02 versus vehicle) by decreasing coronary microvascular resistance (p=0.01 versus vehicle), indicating that GPR39 participates in control of normal coronary vascular tone. With VC108, coronary autoregulation was abolished and CBF became CDP dependent (r^2=0.96, p=0.004).

Conclusion: GPR39 and its endogenous agonist 15-HETE together orchestrate coronary autoregulation. These novel findings provide a mechanism for coronary autoregulation and could direct pharmacological treatment of various coronary syndromes in humans.

Non-standard Abbreviations and Acronyms

14,15-EET   14,15-epoxyeicosatrienoic acid

15-HETE   15-hydroxyeicosatetraenoic acid

CBF   coronary blood flow

CDP   coronary driving pressure

COX-1   cyclooxygenase-1

DHET   Dihydroxyeicosatrienoic acid

DMSO   dimethyl sulfoxide

EDTA   Ethylenediaminetetraacetic acid

eNOS   endothelial NO synthetase

ERK   extracellular signal-regulated kinase

Gαq   G-protein αq

GPR39   G-protein coupled receptor

H_2O_2 hydrogen peroxide

LOX   lipoxygenase

MVR   microvascular resistance

NO   nitric oxide

PGE2   prostaglandin E2

PGD2   prostaglandin D2

PGF2α prostaglandin F2α

PLA_2 phospholipase A2

RA   right atrial

TBXA2   thromboxane A2

VSMC   vascular smooth muscle cell

Coronary autoregulation is the ability of the normal heart to maintain constant coronary

blood flow (CBF) over a wide range of coronary driving pressures (CDPs) provided cardiac

work remains constant: lowering CDP results in vasodilation of the resistance arterioles present

within the heart while raising CDP cause vasoconstriction of these vessels, thus maintaining

constant CBF.[1-4] Autoregulation is a teleologically protective mechanism that is meant to keep

blood flow constant to vital organs despite wide alterations in systemic pressures. This ensures

constant capillary hydrostatic pressure (CHP), necessary for cell survival.[1] In large animals and

humans, the CDP range where CBF remains constant is between approximately 45 to 120 mm

Hg.[2-5]

Despite the relevance of coronary autoregulation to survival its mechanism is unknown.[6]

The eicosanoid 20-hydroxyeicosatetraenoic acid (20-HETE) has been implicated in cerebral

autoregulation.[7] Eicosanoids form a wide variety of lipid-based signaling molecules present in

various forms and concentrations in different tissues.[8] Although some eicosanoids are

vasoactive, their role in coronary autoregulation has not been examined. Further, while single

mediators like adenosine have been investigated as a putative mechanism of coronary

autoregulation[9,10] a comprehensive metabolomic analysis of all possible vasoactive compounds

that could influence resting coronary vascular tone, and autoregulation has not been performed.

We reported the presence of a G-protein coupled receptor (GPR39) in the vascular

smooth muscle cells (VSMCs) of coronary arterioles in the mouse heart.[11] We showed that two

eicosanoids are endogenous ligands for GPR39: the vasoconstrictor 15-hydroxyeicosatetraenoic

acid (15-HETE) is the agonist, and the vasodilator 14,15-epoxyeicosatrienoic acid (14,15-EET)

is the antagonist. Both act through GPR39 to maintain coronary vascular tone.[11] Consequently,

we hypothesized that the levels of 15-HETE and 14,15-EET, acting on GPR39, will determine

the coronary vascular tone depending on the CDP. Thus, we postulated that the receptor and its

ligands together orchestrate coronary autoregulation.

117   **Methods**

An instrumented canine model was selected for the study because coronary physiology

can be reliably studied in these animals and because most of the classical papers on coronary

autoregulation were performed in this model. All data and materials will be made available upon

request. Results are reported according to ARRIVE guidelines. The study protocols were

approved by the Institutional Animal Care and Use Committee at the Oregon Health & Science

University and conformed to the National Institutes of Health guidelines for animal research.

Animal Preparations

Thirty-five adult male mongrel dogs (30-35 kg) were studied. Females were excluded

because of potential pregnancy. Dogs were sedated with intramuscular injection of

hydromorphone (0.1-0. 2 mg·Kg^{-1}) and then induced with intravenous administration of

ketamine (2-5 mg·Kg^{-1}) and midazolam (0.1-0.4 mg·Kg^{-1}). They were then intubated and

ventilated (Integra SL with AV-S ventilator, Avante Health Solutions, Louisville, KY,

USA) with isoflurane gas (0.5-1.0%) and medical-grade room air. Constant-rate infusion

anesthesia was maintained with intravenous administration of ketamine (0.3-0.5 mg·Kg^{-1} bolus

followed by an infusion of 0.3-1.5 mg·Kg^{-1}hr^{-1}), fentanyl (0.001-0.003 mg bolus followed by an

infusion of 0.05-0.07 µg·Kg^{-1}min^{-1}), and a very low dose lidocaine infusion (0.6-6.0 mg·Kg^{-1}hr^{-1}

IV). These infusion rates of the drugs were adjusted throughout the experiment to optimize

anesthesia without affecting coronary reactivity. The mean coronary hyperemic response after a

20 s occlusion was 3-fold, which is normal in open-chest dogs indicating that anesthesia did not

affect vasomotor tone.

Heart rate, blood pressure, O_2 saturation, end-tidal CO_2, and temperature were

continuously monitored (Advisor® Vital Signs Monitor, Surgivet, Norwell, MA, USA). A 6F

sheath was inserted in the right femoral artery for arterial pressure measurement. 7F catheters

were inserted into the right femoral, right atrium (RA), and peripheral veins for administration of

drugs and fluids and measurement of pressure. In a subset of animals, a 7F sheath was inserted

for placement of a balloon-tipped catheter (Edwards Lifesciences, Santa Ana, CA, USA) for

measurement of pulmonary artery (PA) pressures. In the same dogs a micromanometer-tipped

catheter (Millar Instruments, Inc., Houston, TX, USA) was inserted into the left ventricular

cavity to measure left ventricular $\Delta P/\Delta t$.

A left lateral thoracotomy was performed, and the heart was suspended in a pericardial

cradle. The left anterior descending (LAD) coronary artery was dissected free from surrounding

structures in its proximal and mid portions. An ultrasonic time-of-flight flow probe (Series SC,

Transonics, Ithaca, NY, USA) was placed on the proximal segment of the artery and connected

to a digital flow meter (Model T206, Transonics).   A 20-gauge polyethylene catheter was

inserted into the distal portion of the artery to measure pressure. A similar catheter was inserted

in the anterior cardiac vein for obtaining blood samples. CDP was calculated as the difference

between the distal LAD and RA pressures. Coronary microvascular resistance (MVR) was

derived by the CDP/CBF ratio. A screw occluder was positioned around the mid-portion of the

artery to create coronary stenoses of varying degrees.

Hemodynamic Measurements

All catheters and flow meters were interfaced with a multi-channel recorder (PowerLab,

AD Instruments, Inc., Colorado Springs, CO, USA).   Phasic and mean CBF and systemic

pressures were acquired digitally and displayed on-line on an iMac (Apple Inc., Cupertino, CA,

USA). All data were analyzed off-line using LabChart 6.

Mass Spectroscopy

The details for mass spectroscopy measurements of lipids and adenosine have been

published before.[12,13] For measurement of eicosanoids and prostaglandins blood was drawn in a

stop solution containing the antioxidants butylated hydroxytoluene and triphenylphosphonium

mixed with indomethacin in a methanol solution. The samples were centrifuged, placed on ice

and sent for analysis.

Measurements were made using MS/MS with a 5500 Q-TRAP hybrid/triple quadrupole

linear ion trap mass spectrometer (Applied Biosystems, Waltham, MA, USA) with electrospray

ionization (ESI) in negative mode. The mass spectrometer (MS/MS) was interfaced to a

Shimadzu (Columbia, MD, USA) SIL-20AC XR auto-sampler followed by 2 LC-20AD XR LC

pumps and analysis on an Applied Biosystems/SCIEX Q5500 instrument. The instrument was

operated with the following settings: source voltage -4000 kV, GS1 70, GS2 70, CUR 45, TEM

750 and CAD gas MED. The gradient mobile phase was delivered at a flow rate of 0.5 mL/min,

and consisted of two solvents, A: 0.05% acetic acid in water and B: acetonitrile. The LC column

used was a Kinetex Phenyl-Hexyl 50x2.1 mm 2.6 μM. The initial concentration of solvent B

used was 20%, this was held for 0.5 minutes, increased to 60% B over 10.5 minutes, increased to

95% B over 1 min, held at 95% B for 1 min, and then returned to initial conditions over 0.1 min,

followed by re-equilibration for 1.9 min for a total run time of 15 min. The column oven was set

to 50°C, autosampler to 15°C with needle rinse before and after each injection. Data was

acquired using Analyst 1.7.1 and analyzed with Multiquant 3.03.

For arachidonic acid and other polyunsatured fatty acids, analysis was performed using

an Agilent 7890B GC (Santa Clara, CA, USA) with a 5977A MS detector with an autosampler

and split/splitless injector operated in split mode. The column was an Agilent DB-FastFAME

column (30 m, 0.25 mm id, 0.25 μm film thickness). Helium was the carrier gas at a flow rate of

1 mL/min. The injection port and auxiliary heater were maintained at 245°C. A 1 μL sample was

injected in split mode (1:10) at an initial oven temperature of 150°C, held for 0.5 min and then

increased at 15°C/min to 200°C, followed by 25°C/min to 240°C, held for 2 min and then

returned to 50°C. The MS was operated at a source temperature of 230°C and a MS quad

temperature of 150°C in positive electron impact mode. The GC-MS data were acquired using

enhanced MassHunter software. Calibration curves for quantification were generated from peak

area ratios for authentic standard:internal standard for calibrator samples.

For adenosine blood was obtained in stop solution containing dipyridamole, sodium

EDTA, erythro-hydroxy-nonyladenine, α,β-methylene-adenosine 5'-diphosphatenucleotidase,

and heparin.[14] The sample was centrifuged, placed on ice, and sent for analysis. Plasma was

analyzed using MS/MS with a 5500 Q-TRAP PLUS hybrid/triple quadrupole linear ion trap

mass spectrometer (Applied Biosystems) with electrospray ionization (ESI) in positive mode.

The tandem MS/MS was interfaced to a Shimadzu SIL-20AC XR auto-sampler followed by 2

LC-20AD XR LC pumps. The MS instrument was operated with the following settings: source

voltage 5500 kV, GS1 60, GS2 560, CUR 35, TEM 750, and CAD gas MED. Compounds were

infused individually and MS/MS instrument parameters optimized for each MRM transition.

Those used for quantification were m/z=268→136 for adenosine with a dwell time of 100 msec,

DP= 75, EP=10, CXP=15 and CE=25 and m/z=273.2→136 with a dwell time of 100 msec, DP=

75, EP=10, CXP=15 and CE=25. Data was acquired using Analyst 1.7.3 and analyzed with

Multiquant 3.03.

ELISA for Endothelin-1

Plasma samples were analyzed for endothelin-1 using an ELISA kit (Catalog# ADI-900-

020A, Enzo, Farmingdale, NY, USA) as described in the manufacturer's manual. The

absorbance was measured at 450nm using a 96-well plate reader (Molecular Devices,

FlexStation 3, San Jose, CA, USA).

Measurement of VC108 blood levels

A 12-point calibration curve and quality control samples were prepared in blood diluted

1:4.8 in 0.1N Hepes buffer in micronic tubes., 20 µL of which was transferred into micronic

tubes containing 80µL 0.1N Hepes buffer. Calibration samples, quality control samples, and

analysis samples were extracted by adding 400µL CH3CN containing IS (Rolipram) 20 ng/mL.

After vortex and centrifugation (3000 rpm/10 min), samples supernatants were

transferred to 96 well plate and diluted (450µL CH3CN/aq=37/63 + 50µL supernatant). Samples

were injected using an Agilent 1290 Infinity II UPLC system (Santa Clara, CA, USA) and

separated on a Acquity UPLC BEH C18 (30x2.1mm, 1.7µm, Milford, MA, USA) connected

directly to the TurboIonSpray source of an AB Sciex 6500QTrap mass spectrometer

(Framingham, MA, USA). Mobile Phase A was 0.1% (v/v) formic acid in water and Mobile

Phase B was 0.1% (v/v) formic acid in acetonitrile. The flow rate was 1 mL/min and a generic

gradient of 5% B to 95% B in 1.3min was applied.

Study Protocols and their Rationale

We used 2 groups of dogs (Figure 1). Group 1 comprised 7 dogs in which coronary

stenoses of different degrees were created within the autoregulatory range and eicosanoid levels

from the anterior cardiac vein were measured at each stage. Our aim was to determine whether

eicosanoids levels changed during decreases in CDP, which would allow us to further explore

our hypothesis.

Based on the results from Group 1 dogs we proceeded with experiments in the 28 Group

2 dogs (Figure 1). By then we had access to a specific GPR39 inhibitor, VC108,[15] that had been

found to increase CBF in rodents.[16] We created stenoses of varying severities (including flow

limiting) in 22 of these dogs to demonstrate the presence of coronary autoregulation, and in 12 of

them we collected anterior cardiac venous blood at each stage for metabolomic analysis:

adenosine, endothelin-1, polyunsaturated fatty acids (including arachidonic acid and

prostaglandins. We then administered VC108 (n=14) or vehicle (n=8) in these animals and added

3 additional dogs to the vehicle group to make the total =11, and measured systemic, pulmonary

and coronary hemodynamics 30 min later. Vehicle comprising 10% DMSO was administered as

a 5-8 mL IV bolus over 30 s followed by a continuous infusion (120 $mL \cdot hr^{-1}$). VC108 was

suspended in 10% DMSO was administered as a 1.5 mg IV bolus over 30 s followed by a

continuous infusion (0.6 $mg \cdot Kg^{-1} hr^{-1}$).   This dose was calculated based on >90% receptor

occupancy. We then recreated stenoses of varying severities in the 14 dogs receiving VC108 to

demonstrate abolishment of coronary autoregulation. Finally, we used 3 additional dogs in which

blood levels of VC108 were measured every 10 min for an hour.

At the end of the experiment, with animals still under deep anesthesia, 1 mL/4.55 kg of

Euthasol (sodium pentobarbital 390 mg/mL and sodium phenytoin 50 mg/mL) was administered

intravenously in accordance with the June 2007 American veterinary medical association

guidelines on euthanasia that is approved by the Institutional Animal Care and Use Committee at

our institution.

Statistical Methods

Data were expressed as mean±1 standard error of the mean (SEM). Comparison of

hemodynamic data between vehicle and VC108 was performed using unpaired t test.

Comparisons of hemodynamic data between different stenosis stages prior to and during VC108

administration was performed using repeated measures ANOVA and subsequent comparison

between any 2 stages was performed using Dunn-Sidàak post hoc analysis. Data before and

during VC108 administration were compared using 2-way ANOVA. Aggregate CDP versus CBF

data in the 22 dogs were fit to a one phase decay function. Linear regression for performed for

the aggregate CDP versus CBF data in 14 dogs, and CDP versus all metabolites in 12 dogs.

Significance was determined at $P<0.05$.

Results

Table 1 lists the relation between CDP and all the eicosanoids in 7 Group 1 dogs during

17 stages. Only 15-HETE (vasoconstrictor) showed a significant relation with CDP: As CDP

levels decreased so did 15-HETE levels whereas the levels of 14,15-EET (vasodilator) did not

change (Figure 2A). Since the ratio of 15-HETE (vasoconstrictor) and 14,15-EET (vasodilator)

could influence coronary vascular tone, we determined its relationship with CDP (Figure 2B)

and, as expected based on our results, it correlated with CDP.

Since 15-HETE is the endogenous agonist for GPR39,[11] we elected to test the effect of a

specific GPR39 antagonist, VC108,[15] on coronary autoregulation. This drug increases CBF in

rodents,[16] however, its hemodynamic effects in large animals are unknown. We measured its

systemic, pulmonary, and coronary hemodynamics in addition to its effect on coronary

autoregulation. Because various vasoactive metabolites have previously been implicated in the

mechanism of coronary autoregulation,[9,10,17-22] we performed comprehensive metabolomics of

known coronary vasoactive compounds at different CDPs prior to VC108 administration.

We first created coronary stenoses (including where CDP declined below the

autoregulatory range) of the left anterior descending coronary artery in 22 dogs and measured

CDP and CBF during each stage (Figure 1). In 12 of these dogs, we sampled 2.5 mL of blood

from the anterior cardiac vein for measurement of adenosine, endothelin-1, polyunsaturated fatty

acids (including arachidonic acid), and prostaglandins. For prostaglandin and adenosine

measurements blood was collected in stop solutions. We then removed the stenosis and allowed

coronary hemodynamics to return to normal.

Table 2 lists the systemic and coronary hemodynamics at baseline and during different

stenoses stages in 22 dogs (total number of stages = 77). Systemic pressures did not change

between stages. In contrast, as expected, CDP decreased with increasing degrees of stenoses and

CBF remained constant for non-critical stenoses and declined when stenoses became severe

(flow limiting). Constant CBF over a wide range of CDP with non-critical stenosis demonstrates

that autoregulation was operative at baseline. MVR decreased progressively with increasing

stenoses severity and increased with severe stenosis, although these changes were not

statistically significant. We previously reported increased MVR with reduction in resting CBF

despite exhaustion of autoregulation caused by a decrease in myocardial blood volume.[23]

Figure 3A illustrates data from individual dogs in the presence of different degrees of

coronary stenosis. There is wide variation in baseline CBF between animals depending on the

size of the instrumented coronary artery. As a rule, CBF decreased only when CDP was low.

Aggregate results from the 22 animals are shown in Figure 3B where autoregulation is clearly

seen with mild and moderate degrees of stenosis, and CBF declined only when CDP was low

(severe stenosis). The data fit well to a one phase decay function.

Table 3 depicts the relation between CDP and different metabolites measured from the

anterior cardiac vein during different stenosis stages. Only 6-keto-PGF1α, a stable metabolite of

prostacyclin (PGI2) that is not vasoactive, showed a significant correlation with CDP. PGI2 itself

is quickly metabolized and therefore cannot be measured. Figure 4 illustrates the relation with

CDP and arachidonic acid and prostaglandins, while Figure 5 illustrates the relation between

CDP and adenosine and endothelin-1. These cardiac metabolites do not participate in coronary

autoregulation.

After these data were collected we initiated the administration of either VC108 (14

animals) or vehicle (8 animals). We added 3 more dogs to the vehicle group to obtain a

comparable number to the VC108 group. Thirty minutes later we measured systemic, pulmonary,

and coronary hemodynamics. Table 4 lists the baseline systemic and coronary hemodynamics

before administration of VC108 (n=13) or vehicle (n=11). RA pressure was not measured in an

additional dog receiving VC108 and data from it are not included in the table. At baseline, prior

to administration of either vehicle or VC108, CDP was lower in dogs that were assigned to

receive VC108 because of lower mean aortic and consequently distal coronary pressure.

Figure 6 illustrates the change in hemodynamics after VC108/vehicle. There was no

difference between VC108 and vehicle in terms of changes in systemic or pulmonary

hemodynamics (panels A to E). In comparison, there was a small, albeit non-significant

decrease in distal coronary artery pressure (CAP) caused by VC108 compared to vehicle (panel

F) accompanied by a significant increase in CBF (panel G) resulting in a significant reduction in

the calculated coronary microvascular resistance (MVR, panel H). These results indicate that at

rest inhibition of GPR39 causes reduction in coronary MVR without any change in systemic and

pulmonary vascular resistances. No change in cardiac work (product of HR, aortic pressure, left

ventricular $\Delta P/\Delta t$) was noted with either vehicle or drug.

We then recreated stenoses of varying severities in the 14 dogs receiving VC108 and

measured CDP and CBF at each stage. Table 5 shows results from the different stenoses stages.

Systemic pressures did not change between stages. Baseline CBF was higher (26±2 vs. 19±2

mL.min^{-1}, p=0.05) and MVR was lower (2.42±0.20 vs. 3.58±0.27 mmHg·mL^{-1}·min^{-1}, p=0.03)

during VC108 compared to prior to VC108 (Table 2). CBF declined with each level of non-

critical stenosis, indicating abolishment of coronary autoregulation. Because of coronary

vasodilation caused by VC108, MVR remained similar during all levels of non-critical coronary

stenosis but increased when the stenosis was critical. We previously showed that the majority

(>75%) of increase in MVR distal to critical stenosis during coronary vasodilation occurs from

capillary derecruitment distal to stenosis.[23]

Figure 3C illustrates results from the 14 individual dogs where different degrees of

stenoses were placed in the presence of intravenously administered VC108. As a rule, CBF

decreased with decreasing CDP, although the slopes of the decline and baseline CBF varied

between animals. Aggregate results of the 74 stages from 14 animals are shown in Figure 3D.

There is a linear relation between CDP and CBF implying that VC108 abolishes autoregulation.

Table 6 illustrates the results of 2-way ANOVA comparison of systemic and coronary

hemodynamics during various degrees of stenosis in the absence (Table 2) or presence (Table 5)

of VC108. Except for RA pressure, the interactions between each variable are highly significant,

indicating that the conditions were independent of each other.

In 3 additional naïve dogs we administered VC108 and measured drug levels every 10

min for one hour (Table 7). The elevated level at 10 min is from the bolus injection (see

methods), after which the drug level remains steady during VC108 infusion. The mean

blood/plasma ratio from 3 animals was 0.98. Of note, the ratio of free plasma concentration/IC50

of VC108, which in the dog is 31.6 nM, is well over unity, implying therapeutic effect.

Discussion

Using a comprehensive metabolomic analysis of vasoactive molecules that could have a

putative role in coronary autoregulation, we show for the first time that coronary venous levels of

15-HETE, the endogenous agonist of GPR39, is the only molecule that correlates to changes in

CDP, whereas other vasoactive metabolites do not. Since 15-HETE acts through GPR39, our

results indicate that GPR39 is the molecular switch that controls coronary vasomotion. This is

supported by the finding that after administration of VC108, a specific GPR39 inhibitor,

coronary MVR decreased. In the absence of VC108, CBF remained constant over a wide range

of CDPs induced by varying degrees of coronary stenosis indicating intact autoregulation.

VC108 abolished coronary autoregulation so that CBF became CDP dependent. Therefore,

GPR39 and its endogenous agonist 15-HETE together orchestrate coronary autoregulation.

In our previous study, where we showed that GPR39 was the receptor for the endogenous

ligands 15-HETE and 14,15-EET in the mouse, we stripped the heart of all large vessels and

demonstrated the presence of GPR39 on VSMCs in the coronary microvasculature using

immunohistochemistry: GPR39 was present in coronary resistance arterioles.[11] Using a mouse

Langendorf constant flow preparation we showed that 15-HETE increased CDP above the

normal range, an effect that was attenuated by co-administration of 14,15-EET and abolished in

GPR39 knockout mice. In these experiments, 14,15-EET alone had no effect on CDP. Our

current results extend these observations.

Zn^{++} serves as an allosteric modulator of GPR39.[24,25] In the presence of Zn^{++}, GPR39

activation leads to extracellular signal-regulated kinase (ERK) phosphorylation and intracellular

Ca^{++} mobilization,[11] suggesting activation of G-protein αq (Gαq). Increased intracellular Ca^{++}

results in VSMC contraction. The higher the level of 15-HETE, the greater the stimulation of

GPR39 and the greater the intracellular Ca^{++} and consequent vasoconstriction.[11] As 15-HETE

levels decrease the reverse occurs, leading to less intracellular Ca^{++} and less vasoconstriction (or

more vasorelaxation). In the intact animal changes in CDP can occur moment to moment

requiring a corresponding change in CBF. Therefore, from a systems perspective it seems

efficient to have one molecule act on one receptor to affect instant changes, akin to a thermostat

rather than an off/on switch. The very short plasma half-life of 15-HETE supports this notion.

Role of Arachidonic acid Metabolites

Prostaglandins, arachidonic acid derivatives, were examined previously where

administration of two inhibitors of prostaglandin synthesis (5,9,11,14-eicosatetraynoic acid and

indomethacin) abolished coronary autoregulation in the perfused isolated rabbit heart,[26] although

studies in larger animals did not show any effects of indomethacin on CBF.[27,28] The same

investigators who reported the effect of prostaglandin synthesis inhibition in the isolated rabbit

heart subsequently reported an 'active' substance, an unidentified albumin bound lipid molecule,

that counteracted the effects of indomethacin.[29] Although it cannot be definitively proven, our

results suggest that this 'active' substance could have been 15-HETE, a metabolite of

arachidonic acid, which the investigators did not measure. When human endothelial cells are

stimulated by histamine, they enhance synthesis of both prostaglandin and 15-HETE.[30] Both are

inhibited by indomethacin, which might explain why autoregulation was abolished with the drug

in the perfused isolated rabbit heart.[31] Our results indicate that prostaglandins, specifically

TBXA2, PDE2, PGE2, and PGF2$_\alpha$, do not participate in autoregulation. We saw a mild decline

in the prostacyclin metabolite 6-keto-PGF1α, which is not vasoactive.

Of the eicosanoids, another group of arachidonic acid metabolites, only 15-HETE

showed a significant correlation with CDP in our study. However, a borderline correlation

between CDP and 12-HETE was also noted. Both 15- and 12-HETE are produced by the same

enzyme (12/15LOX) and their ratio depends on the species. In our study the 12-HETE/15-HETE

ratio was approximately 20:1. The main receptor for 12-HETE is GPR31, which is thought not to

be present in VSMCs.[22] We previously reported that it had no effect on GPR39.[11] 12-HETE is a

competitive thromboxane A2 receptor antagonist, which would make it a vasodilator although its

action on VSMCs is complex and controversial.[22] Coronary venous plasma levels of arachidonic

acid itself remained unchanged at different CDPs, indicating that changes in coronary pressure

influenced arachidonic acid to 15-HETE conversion by 12/15 LOX probably by release of more

substrate from the lipid bilayer by phospholipase A_2.

Role of Other Pathways-Comparison with Previous Studies

Nitric oxide (NO) has been implicated in regulation of coronary autoregulation.[17] There

are several pathways through which GPR39, 15-HETE, and NO interact with each other to

explain these findings. First, increased intracellular Ca^{++} stimulates endothelial NO synthetase

(eNOS) resulting in increased NO production.[32] NO, in turn, stimulates sarcoplasmic

endoplasmic reticulum Ca^{++}ATPase, thereby increasing Ca^{++} reuptake into sarcoplasmic

endoplasmic reticulum and decreasing vascular tone.[33] GPCRs can undergo post-translation

modification induced by S-nitrosylation whereby NO could modify the response of GPR39 to 15-

HETE in vivo.[34] Thus, inhibition of eNOS by NG-nitro-L-arginine would interfere with

autoregulation as previously reported[17] by inhibiting signaling both at the level of GPR39 and

downstream to it.

Inhibition of α-adrenergic systemic has been shown to alter coronary autoregulation.[18] α-

adrenergic stimulation causes increased production of arachidonic acid through activation of

mitogen-activated protein kinases that are downstream to ERK, enhancing arachidonic acid

production from the nuclear envelope.[35] It also increases production of arachidonic acid and its

metabolites by increasing intracellular Ca^{++}, which stimulates cytosolic phospholipase (PLA_2)

production that is responsible for release of arachidonic acid from plasma membrane.[36] Thus, α-

adrenergic inhibition by drugs such as prazocin reported previously[18] could affect coronary

autoregulation through inhibition of arachidonic acid and 15-HETE production, mechanisms

proximal to GPR39.

Vasodilation requires VSMC hyperpolarization that is mediated through increased levels

of intracellular K^+ resulting from K^+ channel opening.[19] 15-HETE directly inhibits K^+ channel

activity, thus attenuation vasodilation.[37] Lower levels of 15-HETE seen with coronary arteriolar

vasodilation at lower CDPs are therefore likely associated with more active K^+ channels that are

open and allow intracellular K^+ flux under appropriate circumstances. Open K^+ channels-

induced hyperpolarization also decreases Ca^{++} sensitivity.[38] Other K^+ channel blockers have also

been associated with disruption of coronary autoregulation.[10,20,39] H_2O_2 opens Ca^{++} gated and

voltage-dependent K^+ channels allowing VSMC hyperpolarization.[39] Therefore, K^+ channel

blockers like tetraethyl ammonium and 4-aminopyridine block the effect of H_2O_2, thus

interfering with vasodilation.[20] A similar explanation can be provided for attenuation of coronary

autoregulation by glibenclamide, an ATP-sensitive K^+ channel blocker.[20] K^+ channel activity is

downstream to GPR39 activation.

Finally, adenosine was previously shown not to participate in coronary autoregulation[9,10]

and we confirmed the finding. Additionally, endothelin-1, a potent coronary vasoconstrictor[40]

also did not participate in coronary autoregulation.

Proposed Mechanism of Coronary Autoregulation

Based on our results we propose the following scenario for coronary autoregulation

shown in the central illustration. Changes in CDP are mechanosensed by ion channels[41,42]

present on VSMCs, which then either activate PLA_2 to release the substrate arachidonic acid

from plasma membrane to produce its metabolites or directly modulate 12/15 LOX activity

resulting in 15-HETE production. 15-HETE then activates GPR39, which in turn causes

intracellular Ca^{++} release resulting in VSMC contraction. Higher CDP causes higher 15-HETE

production and more vasoconstriction while lower CDP causes less 15-HETE production and

less vasoconstriction (or relative vasodilation). Counterintuitively, no vasodilators actively

participate in this process. Coronary vascular tone needs to be rapidly adjusted to myocardial

oxygen needs or CDP changes, implying that the molecular switch for coronary autoregulation

needs to act more like a thermostat than a lock and key, which is readily accomplished by a

single biomolecule acting on a single receptor.

Limitations of our Study

We did not study the upper range of autoregulation (100-120 mm CDP). Nonetheless,

based on our previous results using a constant flow Langendorff mouse model,[11] we predict that

increasing CDP above the normal range would increase 15-HETE levels and cause further

vasoconstriction to maintain constant CBF. In the dog, where CBF is not controlled,

intracoronary injection of 15-HETE causes CBF decline from increased MVR that is mitigated

by pre-treatment with VC108.[43] We did not see elevation of adenosine or endothelin-1 levels at

low CDP probably because ischemia was not produced due to adequate collateral flow to the

myocardial region supplied by the left anterior descending artery. Collateral blood flow is high

in the dog.[44]

We did not measure cardiac metabolite levels during VC108 administration because in

our early experiments we noted hemolysis on centrifugation of the coronary venous blood

samples. It was then determined that hemolysis was caused by DMSO.[45]

<u>Conclusion</u>

In this study we show for the first time that 15-HETE participates in the resting coronary

vascular tone in dogs. We also demonstrated that 15-HETE acting through GPR39 is responsible

for coronary autoregulation. Factors that have previously been described to effect coronary

autoregulation either modify GPR39 directly or act through signaling pathways upstream or

downstream of GPR39. Hence, GPR39 serves as the molecular switch responsible for coronary

autoregulation.

471   **Author Contributions**

472   Drs. Le and Kaul developed the concept and design for the studies. They also wrote the

473   manuscript. Drs. Kajimoto and Zhao participated in the studies and performed the surgery and

474   delivered anesthesia and drugs. Dr. Methner performed ELISA for endothelin-1 and prepared

475   VC108 bolus and infusions. Dr. Le analyzed the data and created the tables and figures. Drs.

476   Cianciulli, Semeraro, Trist and Micheli contributed to the development of VC108.

477   **Acknowledgements**

478   Metabolomic   analysis   was   performed   by   the Bioanalytical   Shared   Resource/

Pharmacokinetics Core Facility (Research Resource Identifier (RRID): SCR_009963), which is

supported in part by the University Shared Resource Program at Oregon Health and Sciences

University. Special thanks to Lisa Byle and Jenny Luo for the analysis. We are also grateful to

Joseph Sciurba DVM and his team in the department of comparative medicine for their

participation in the study and care of the animals.

Conflict of Interest

Oregon Health & Science University and Vasocardea, Inc. have a joint patent application

filed globally (see WO2021222858 and corresponding national filings) that encompasses drugs

that inhibit GPR39. Vasocardea, Inc. has an additional patent application filed globally (see

WO2023076219 and corresponding national filings) that encompasses VC108. Vasocardea, Inc.

has exclusive rights to develop and commercialize drugs under all of these patent applications.

Evotec has assigned all patent rights to Oregon Health & Science University and Vasocardea,

Inc.

Dr. Kaul is the founder and President of Vasocardea, Inc., a Delaware incorporated

company located in Portland, Oregon.

The information in this manuscript does not represent the views of the Veterans

Administration or the US government.

References

1.   Starling, EH. On the Absorption of Fluids from the Connective Tissue Spaces. *J*
Physiol. 1896;19:312-326.

2.   Mosher P, Ross J Jr., McFate PA, Shaw RF. Control of coronary blood flow by an
autoregulatory mechanism. *Circ Res* 1964;16:250-259.

3.   Rouleau J, Boerboom LE, Surjadhana J, Hoffman JIE. The role of autoregulation and
tissue diastolic pressures in the transmural distribution of left ventricular blood flow
in anesthetized dogs. *Circ. Res.*1979;45:804–815.

4.   Kanatsuka H, Lamping KG, Eastham CL, Marcus ML. Heterogeneous changes in
epimyocardial microvascular size during graded coronary stenosis. Evidence of the
microvascular site for autoregulation. *Circ Res*. 19900;66, 389–396.

5.   Dick GM, Namani R, Patel B, Kassab GS. Role of coronary myogenic response in
pressure-flow autoregulation in swine: A meta-analysis with coronary flow modeling.
Frontiers in Physiology 2018;10.3389/ fphys. 2018. 00580/full.

6.   Deussen A. Mechanisms underlying coronary autoregulation continue to await
clarification. *Basic. Res. Cardiol.* 2018;113:33-34.

7.   Gebremedhin D, Lange AR, Lowry TF, Taheri R, Birks EK, Hudetz AG, Narayanan
J, Falck JR, Okamoto H, Roman RJ, Nithipatikom K, Campbell WB, Harder DR.
Production of 20-HETE and Its Role in Autoregulation of Cerebral Blood Flow. *Circ.*
Res. 2000;87:60-65.

8.   Smith WL. The eicosanoids and their biochemical mechanisms of action. *Biochem. J.*
1989;259: 315-324.

9. Hanley FL, Grattan MT, Stevens MB, Hoffman JIE. Role of adenosine in coronary

autoregulation. *Am. J. Physiol.* 1986;250:H558-H566.

10. Stepp DW, Kroll K, Feigl EO. K$^+$ATP channels and adenosine are not necessary for

coronary autoregulation. *Am. J. Physiol*. 1997;273:H1299–H1308.

11. Alkayed NJ, Cao Z, Qian ZY, Nagarajan S, Liu X, Nelson J, Xie F, Li B, Fan W,

Xiao X, Barnes AP, Kaul S. Control of Coronary Vascular Resistance by Eicosanoids

via a Novel GPCR. *Am. J. Physiol.* 2022;322:C1011-C1021.

12. Dumlao DS, Buczynski MW, Norris PC, Harkewicz R and Dennis EA. High-

throughput lipidomic analysis of fatty acid derived eicosanoids and N-

acylethanolamines. *Biochim Biophys Acta* 2011;1811:724-736.

13. Sharma K, Singh R, Giri S, Rajagopal S and Mullangi R. Highly sensitive method for

the determination of adenosine by LC-MS/MS-ESI: method validation and scope of

application to a pharmacokinetic/pharmacodynamic study. *Biomed Chromatogr* 2012;

26:81-88

14. Shryock JC, Boukin MT, Hill JA, Belardinelli L. A new method of sampling blood

for measurement of plasma adenosine. *Am J Physiol* 1990;258:H1232-H1239.

15. Cianciulli A, Bernadelli A, Faedo S, Le E, Methner C, Pellacani A, Semeraro T, Trist

ILM, Kaul S, Micheli F. Identification of Novel and selective GPR39 receptor

antagonists for the treatment of Acute Myocardial Ischemia. Chem Med Chem 2025;

https://doi.org/10.1002/cmdc.202500662

16. Methner C, Le DE, Kaul S. Targeting GPR39 for no reflow after acute myocardial

infarction. *Circulation* 2023;148 (Suppl_1):18781.

17.   Ueeda M, Silvia SK, Olsson RA. Nitric oxide modulates autoregulation in guinea
543   pigs. *Circ. Res.* 1992;70:1296-1303.

544   18.   Jones CE, Liang IYS, Gwirtz PA. Effect of α-adrenergic blockage on coronary
545   autoregulation in dogs. *Am. J. Physiol.* 1987;253:H365-H372.

546   19.   Jackson WF. Potassium Channels in regulation of vascular smooth muscle
547   contraction and growth. *Adv. Pharmacol.* 2017;78:89-144

548   20.   Yada T, Shimokawa H, Hiramatsu O, Kajita T, Shigeto F, Goto M, Ogasawara Y,
549   Kajiya F. Hydrogen peroxide, an endogenous endothelium-derived hyperpolarizing
550   factor, plays an important role in coronary autoregulation in vivo. *Circulation.*
551   2003;107:1040-1045.

552   21.   Narishige T, Egashira K, Akatsuka Y, Katsuda Y, Numaguchi K, Sakata M,
553   Takeshita A. Glibenclamide, a putative ATP-Sensitive K^+ channel blocker, inhibits
554   coronary autoregulation in anesthetized dogs. *Circ. Res.* 1993;73:771-776

555   22.   Pascale JV, Lucchesi PA, Garcia V. Unraveling the Role of 12- and 20- HETE in
556   Cardiac Pathophysiology: G-Protein-Coupled Receptors, Pharmacological Inhibitors,
557   and Transgenic Approaches. *J Cardiovasc Pharmacol.* 2021;77:707-717.

558   23.   Jayaweera AR, Wei K, Coggins M, Bin JP, Goodman C, Kaul S. Role of capillaries
559   in determining coronary blood flow reserve: New insights using myocardial contrast
560   echocardiography. *Am J Physiol* 1999;277:H2363-H2372.

561   24.   Laitakari A, Liu L, Frimurer TM, Holst B. The Zinc-Sensing receptor GPR39 in
562   physiology and as a pharmacological target. *Int. J. Mol. Sci.* 2021;22:3872.

563   25.   Storjohann L, Holst B, Schwartz TW. Molecular mechanism of Zn^{2+} agonism in the
564   extracellular domain of GPR39, *FEBS Letters*, 2008;582.

26. Moretti RL, Abraham S, Ecker RR. The stimulation of cardiac prostaglandin

production by blood plasma and its relationship to the regulation of coronary flow in

isolated isovolumic rabbit hearts. *Circ. Res.* 1976;39:231-238.

27. Rubio R, Berne RM. Regulation of coronary blood flow. *Prog. Cardiovasc. Dis.*

1975;18:105-122.

28. Dai XJ, Bache RJ. Effect of indomethacin on coronary blood flow during graded

treadmill exercise in the dog *Am. J. Physiol.* 1984:247:H452-H458.

29. Moretti RL, Abraham S. Stimulation of microsomal prostaglandin synthesis by a

blood plasma constituent which augments autoregulation and maintenance of vascular

tone in isolated rabbit hearts. *Circ. Res.* 1978;42:317-323.

30. Revtyak GE, Hughes MJ, Johnson AR, Campbell WB. Histamine stimulation of

prostaglandin and HETE synthesis in human endothelial cells. *Am. J. Physiol.*

1988;255(2 Pt 1):C214-225.

31. Salari H, Braquet P, Borgeat P. Comparative effects of indomethacin, acetylenic

acids, 15-HETE, nordihydroguaiaretic acid and BW755C on the metabolism of

arachidonic acid in human leukocytes and platelets. *Prostaglandins Leukotrienes and*

Medicine 1984;13:53-60.

32. Fleming I, Busse R. Signal transduction of eNOS activation. *Cardiovasc. Res.*

1999;43:532–541.

33. Tong X, Evangelista A, Cohen RA. Targeting the redox regulation of SERCA in

vascular physiology and disease. *Curr. Opin. Pharmacol*. 2010;10:133-138.

34. Daaka Y. S-Nitrosylation-regulated GPCR signaling. *Biochim. Biophys. Acta*. 2012;

1820:743–751.

588 35. Burch RM, Luini A, Mais DE, Corda D, Vanderhoek JY, Kohn LD, Axelrod J. Alpha

589   1-adrenergic stimulation of arachidonic acid release and metabolism in a rat thyroid

590   cell line. Mediation of cell replication by prostaglandin E2. *J Biol. Chem.*

591   1986;261:11236-11241.

592 36. Nebigil C, Malik KU. Alpha adrenergic receptor subtypes involved in prostaglandin

593   synthesis are coupled to Ca^{++} channels through a pertussis toxin-sensitive guanine

594   nucleotide-binding protein. *J. Pharmacol. Exp. Ther.* 1993;266:1113-1124.

595 37. Li Y, Li Q, Wang Z, Liang D, Liang S, Tang X, Guo L, Zhang R, Zhu D. 15-HETE

596   suppresses K(+) channel activity and inhibits apoptosis in pulmonary artery smooth

597   muscle cells. *Apoptosis.* 2009;1:42-51.

598 38. Gutterman DD, Miura H, Liu Y. Redox modulation of vascular tone. Focus of

599   potassium channel mechanisms of dilation. *Arterioscl. Thromb. Vasc. Biol.*

600   2005;25:671–678.

601 39. Nacitarhan C, Bayram Z, Eksert B, Usta C, Golbasi I, Ozdem SS. The Effect of

602   hydrogen peroxide in human internal thoracic arteries: role of potassium channels,

603   nitric oxide and cyclooxygenase products. *Cardiovasc. Drugs. Ther.* 2007*;*21:257–

604   262

605 40. Teerlink JR, Breu V, Sprecher U, Clozel M, Clozel JP. Potent vasoconstriction

606   mediated by endothelin ETB receptors in canine coronary arteries. *Circ Res.* 1994;

607   74:105-114..

608 41. Hardie RC. TRP channels and lipids: from Drosophila to mammalian physiology. *J*

609   *Physiol* 2007;578:9-24.

42.   Munaron L. Shuffling the cards in signal transduction: Calcium, arachidonic acid and

mechanosensitivity. *World. J. Biol. Chem.* 2011;2:59-66.

43.   Le DE, Kajimoto M, Zhao Y, Methner C, Kaul S. VC108, a GPR39 blocker,

increases coronary blood flow and inhibits the vascular effects of 15-

hydroxyeicosatetraenoic acid. Circulation 152 (suppl_3):4357647, 2025.

44.   Schaper W, Binz K, Sass S, Winkler B. Influence of collateral blood flow and of

variations in MV̇O2 on tissue-ATP content in ischemic and infarcted myocardium. *J.*

Mol. Cell. Cardiol. 1987;19:19-37.

45.   Yi X, Liu M, Luo Q, Zhuo H, Cao H, Wang J, Han Y. Toxic effects of dimethyl

sulfoxide on red blood cells, platelets, and vascular endothelial cells *in vitro*. *FEBS*

Open Bio. 2017;7:485-494.

Table 1. Relation between Coronary Driving Pressure and Coronary Venous Eicosanoid Levels (n=7 dogs, 17 stages)

Eicosanoid Isomer	Equation	r^2	p value
5,6-EET	y = 0.9392x + 42.17	0.0758	0.3902
8,9-EET	y = 0.0273x + 40.01	0.0005	0.9338
11,12-EET	y = 0.0843x + 31.53	0.0206	0.3376
14,15-EET	y = 0.0911x + 48.96	0.0063	0.7626
5,6-DHET	y = -0.0957x + 37.02	0.0143	0.6481
8,9-DHET	y = -0.0230x + 51.04	0.0009	0.9106
11,12-DHET	y = -0.2599x + 132.6	0.0192	0.5955
14,15-DHET	y = -0.1528x + 104.6	0.0086	0.7240
11-HETE	y = 2.7060x + 15.70	0.0316	0.4950
12-HETE	y = 44.94x + 129.10	0.2308	0.0596
15-HETE	y = 3.0350x + 4.7770	0.4700	0.0024
20-HETE	y = -5.523x + 1964	0.0188	0.5994

Table 2. Systemic and Coronary Hemodynamics for 22 Protocol 3 Dogs at Baseline and Varying Degrees of
643
Coronary Stenosis (n=77 Stages) Prior to VC108 Administration (mean ± SEM)
644

Parameter	Baseline (n=22)	Mild Stenosis Gradient ≤10 mmHg (n=19)	Moderate Stenosis Gradient 11 – 34 mmHg (n=28)	Severe Stenosis Gradient ≥35 mmHg (n=8)	p value Anova
Heart rate (bpm)	97±3	93±3	96±3	104±3	0.3023
Mean blood pressure (mmHg)	74±3	71±2	70±1	64±4	0.2897
Right atrial blood pressure (mmHg)	12±1	12±1	12±1	13±1	0.7037
Mean distal coronary artery pressure (mmHg)	74±3*‡§	65±2‡§	55±2§	35±3	<0.0001
Coronary artery driving pressure (mmHg)	63±3*‡§	53±2†§	43±2§	22±3	<0.0001
Coronary artery blood flow (mL·min^{-1})	19±2#	17±18#	17±1#	9±1	0.0002
Coronary microvascular resistance (mmHg·mL·min^{-1})	3.58±0.27	3.52±0.31	2.88±0.28	3.14±0.45	0.2607

647   *$p < 0.05$ v. mild stenosis gradient
648   †$p < 0.05$ v. moderate stenosis gradient
649   ‡$p < 0.0001$ v. moderate stenosis gradient
650   #$p < 0.01$ v. severe stenosis gradient
651   §$p < 0.0001$ v. severe stenosis gradient

Table 3. Relation between Coronary Driving Pressure and Cardiac Venous Metabolite Levels During Baseline and Various Stenosis Stages Prior to VC108 Administration in 12 Dogs

Metabolite (pg/mL)	Equation	r^2	p value
Adenosine (n=49)	y = 0.2641x + 28.78	0.0201	0.3309
Endothelin (n=49)	y = -0.0066x + 9.94	0.0001	0.9340
Prostaglandins			
6-keto Prostaglandin (n=46)	y = 1.570x + 77.28	0.0992	0.0330
Thromboxane 2 (n=46)	y = 11.79x + 8.53	0.0502	0.1345
Prostaglandin E2 (n=48)	y = 0.2014x + 27.46	0.0195	0.3435
Prostaglandin D2 (n=48)	y = 0.1326x + 23.56	0.0047	0.6420
Prostaglandin F2$_\alpha$ (n=49)	y = 0.3365x + 41.22	0.0047	0.6390
Polyunsaturated Fatty Acid			
Arachidonic acid (n-49)	y = -0.1259x + 416.50	0.0008	0.8454
Oleic Acid (n=49)	y = 0.2577x + 146.90	0.0071	0.5652
Linolenic Acid (49)	y = 0.0051x + 0.0019	0.0019	0.7642
Gama Linoleic Acid (n=49)	y = 0.0209x + 5.10	0.0349	0.1988
Linoleic Acid (n=49)	y = 0.0418x + 478.20	0.0022	0.7491
Eicosapentaenoic Acid (n=49)	y = 0.0133x + 5.19	0.0212	0.3185
Docosahexanoic Acid (n=49)	y = -0.0369x + 29.64	0.0029	0.7132

Hemolyzed samples not analyzed

**Table 4. Baseline Systemic and Coronary Hemodynamics in Protocol 2 Dogs
Prior to Vehicle/Drug Administration (mean ± SEM)**

Parameter	Vehicle (n=11)	VC108 (n=13)	p value
Heart rate (bpm)	101±6	104±3	0.6092
Mean aortic pressure (mmHg)	80±5	72±2	0.1048
Mean right atrial pressure (mmHg)	11±2	12±1	0.5283
*Left ventricular $\Delta P/\Delta t$ (mmHg·s^{-1})	1082±185	868±127	0.3533
*Mean pulmonary artery pressure (mmHg)	23±5	16±1	0.1461
Distal coronary artery pressure (mmHg)	82±4	72±2	0.0508
Coronary artery driving pressure (mmHg)	71±4	60±2	0.0108
Coronary artery blood flow (mL·min^{-1})	21±2	22±2	0.9097
Coronary microvascular resistance (mmHg·mL^{-1}·min^{-1})	3.66±0.49	3.14±0.30	0.3747

*Measured in 6 dogs

710 **Table 5. Systemic and Coronary Hemodynamics for 14 Protocol 3 Dogs at Baseline and with Varying Degrees**
711 **of Coronary Stenoses (n=74 stages) During VC108 Administration (mean ± SEM)**
712
713

Parameter	Baseline (n=14)	Mild Stenosis Gradient ≤10 mmHg (n=10)	Moderate Stenosis Gradient 11 – 34 mmHg (n=32)	Severe Stenosis Gradient ≥35 mmHg (n=18)	p value Anova
Heart rate (bpm)	104±3	104±3	108±2	106±2	0.6373
Mean blood pressure (mmHg)	71±2	76±3	75±2	76±2	0.3918
Right atrial blood pressure (mmHg)	13±0	13±0	13±0	13±0	0.4900
Mean distal coronary artery pressure (mmHg)	71±2‡§	70±3‡§	56±1§	40±1	<0.0001
Coronary artery driving pressure (mmHg)	59±2‡§	56±3‡§	43±1§	27±2	<0.0001
Coronary artery blood flow (mL·min^{-1})	26±2§	22±2§	20±2§	8±1	<0.0001
Coronary microvascular resistance (mmHg·mL·min^{-1})	2.42±0.20¥	2.64±0.22♯	2.56±0.16§	4.63±0.59	<0.0001

714
*p < 0.05 v. mild stenosis gradient
†p < 0.05 v. moderate stenosis gradient
‡p < 0.0001 v. moderate stenosis gradient
♯p < 0.01 v. severe stenosis gradient
¥p < 0.001 v. severe stenosis gradient
§p < 0.0001 v. severe stenosis gradient
721
722

Table 6. 2-Way ANOVA interactions between variables in Tables 5 and 6 (Protocols 3 Dogs)

Parameter	F value	p value
Heart rate (bpm)	21.22	<0.0001
Mean blood pressure (mmHg)	52.22	<0.0001
Right atrial blood pressure (mmHg)	0.98	0.4011
Mean distal coronary artery pressure (mmHg)	25.25	<0.0001
Coronary artery driving pressure (mmHg)	23.65	<0.0001
Coronary artery blood flow (mL·min^{-1})	30.55	<0.0001
Coronary microvascular resistance (mmHg·mL·min^{-1})	86.41	<0.0001

**Table 7. Total and Free Blood Concentrations (nM) as well as 1C50/Free Blood Concentration of VC108
After a 1.5 mg Bolus Injection Followed by 0.6 mg/kg/h Infusion in 3 Protocol 1 Dogs**

Time After Administration (min)	Total Blood Concentration	Free Blood Concentration	Free Plasma Concentration	Free Plasma Concentration/IC50
10	14752	4661	4568	71
20	6824	2156	2113	33
30	6382	2017	1977	31
40	5876	1857	1820	28
50	6094	1926	1887	29
60	5902	1653	1620	25

Blood plasma ratio of VC108=0.98
IC50 of VC108=31.6 nM

Figure Legends

Figure 1: Breakdown of the 35 animals used in the study and protocols they underwent. Created

in https://Biorender.com

Figure 2: Relation between coronary driving pressure and anterior cardiac venous plasma levels

of A)15-HETE and 14,15-EET from 7 dogs undergoing different degrees of coronary stenosis

within the autoregulatory range, and B) 15-HETE/14,15-EET ratio.

Figure 3: Coronary driving pressure versus coronary artery blood flow results from protocols 2

and 3. A) Data from 22 individual dogs (77 unique stages) in the absence of VC108 showing flat

coronary artery blood flow over a wide range of coronary driving pressures, with decreasing

flow at low coronary driving pressures. B) Aggregate results from the same animals as in panel

A, showing classical coronary autoregulation over a wide range of coronary driving pressures

with a decline in coronary artery flow at low coronary driving pressures. Data were fit using one

phase decay least-squares analysis. C) Data from 14 of the 22 individual dogs (74 unique stages)

from protocol 2 in the presence of VC108 (protocol 3) showing declining coronary artery blood

flow with lowering of coronary driving pressure. D) Aggregate results from the same animals as

in panel C, showing abolishment of coronary autoregulation with coronary artery blood flow

becoming coronary driving pressure dependent. Data were fit using linear regression analysis.

Figure 4: Relation between coronary driving pressure and anterior coronary vein plasma

concentrations of A) adenosine and B) endothelin-1 from 12 dogs undergoing various degrees of

coronary stenoses at rest.

Figure 5: Relation between coronary driving pressure and anterior coronary vein plasma

concentrations of A) Arachidonic acid; B) 6-keto-PGF$_\alpha$, (an inactive metabolite of prostaglandin

I2; C) Thromboxane A2; D) Prostaglandin E2; E) Prostaglandin D2; and F) Prostaglandin F2$_\alpha$.

Figure 6: Percent change from baseline in systemic, pulmonary, and coronary hemodynamics

with administration of Vehicle (VEH, n=11) or VC108 (n=13). A) HR=heart rate; B) AoP=aortic

pressure; C) PAP=pulmonary artery pressure; D) RAP=right atrial pressure; E) LV dP/dt=rate of

change of left ventricular pressure; E) CAP=coronary artery pressure; F) CBF=coronary blood

flow; and H) MVR=coronary microvascular resistance. Comparisons between groups were made

using unpaired *t*-test.

Central Illustration Mechanism of Autoregulation: Proposed mechanism of intraluminal

coronary pressure sensing and GPR39 stimulation and vasoconstriction (see text for details).

Increased or decreased intraluminal pressure is sensed by ion channels or G-protein coupled

receptors that will either increase (high pressure) or decrease (low pressure) arachidonic acid

production leading to either vasoconstriction or vasodilation. Created in https://Biorender.com

Figure 1: Breakdown of the dogs into groups and protocols. See text for details. Created in

https\\Biorender.com

Figure 2: Relation between coronary driving pressure and anterior cardiac venous plasma levels

of A)15-HETE and 14,15-EET from 7 dogs undergoing different degrees of coronary stenosis

within the autoregulatory range, and B) 15-HETE/14,15-EET ratio.

Figure 3: Coronary driving pressure versus coronary artery blood flow results from protocols 2

and 3. A) Data from 22 individual dogs (77 unique stages) in the absence of VC108 showing flat

coronary artery blood flow over a wide range of coronary driving pressures, with decreasing

flow at low coronary driving pressures. B) Aggregate results from the same animals as in panel

A, showing classical coronary autoregulation over a wide range of coronary driving pressures

with a decline in coronary artery flow at low coronary driving pressures. Data were fit using one

phase decay least-squares analysis. C) Data from 14 of the 22 individual dogs (74 unique stages)

from protocol 2 in the presence of VC108 (protocol 3) showing declining coronary artery blood

flow with lowering of coronary driving pressure. D) Aggregate results from the same animals as

in panel C, showing abolishment of coronary autoregulation with coronary artery blood flow

becoming coronary driving pressure dependent. Data were fit using linear regression analysis.

Figure 4: Relation between coronary driving pressure and anterior coronary vein plasma

concentrations of A) adenosine and B) endothelin-1 from 12 dogs undergoing various degrees of

coronary stenoses at rest.

A.

B.

Figure 5: Relation between coronary driving pressure and anterior coronary vein plasma

concentrations of A) Arachidonic acid; B) 6-keto-PGF$_\alpha$, (an inactive metabolite of prostaglandin

I2; C) Thromboxane A2; D) Prostaglandin D2; E) Prostaglandin E2; and F) Prostaglandin F2$_\alpha$.

Figure 6: Percent change from baseline in systemic, pulmonary, and coronary hemodynamics

with administration of Vehicle (VEH, n=11) or VC108 (n=13). A) HR=heart rate; B) AoP=aortic

pressure; C) PAP=pulmonary artery pressure; D) RAP=right atrial pressure; E) LV dP/dt=rate of

change of left ventricular pressure; E) CAP=coronary artery pressure; F) CBF=coronary blood

flow; and H) MVR=coronary microvascular resistance. Comparisons between groups were made

using unpaired *t*-test.

829 **Central Illustration Mechanism of Autoregulation:** Proposed mechanism of intraluminal

830 coronary pressure sensing and GPR39 stimulation and vasoconstriction (see text for details).

831 Increased or decreased intraluminal pressure is sensed by ion channels or G-protein coupled

832 receptors that will either increase (high pressure) or decrease (low pressure) arachidonic acid

833 production leading to either vasoconstriction or vasodilation. Created in https://Biorender.com

834