Title: A meningeal circadian clock and brain state co-ordinately

enhance macromolecular clearance during wake

Authors: Talya Goble, Tanushree Kundu, Léticia S. Meyer, Dimitra Klinaki,

Thomas A. Hawkins, Jason Rihel*

Department of Cell and Developmental Biology, University College London, London,

United Kingdom

*corresponding author: j.rihel@ucl.ac.uk

Abstract

The removal of brain waste is thought to be regulated by the circadian clock and

sleep-wake cycle. A meningeal scavenger cell population, called mural Lymphatic

Endothelial Cells (muLECs), uptake macromolecules from the cerebrospinal fluid

(CSF) and are poised to clear soluble brain waste. Whether CSF clearance is

regulated across the 24-hour day is unknown. Using a quantitative, *in vivo* assay in

zebrafish, we show that the capacity of muLECs to remove macromolecules is

highest during the waking day. The muLEC protein uptake is regulated by a cell-

autonomous circadian oscillator that drives rhythmic protein levels of Mannose

Receptor 1a (Mrc1a); in contrast, lipoprotein uptake increases as a function of time

spent awake. Thus, circadian and brain state cues independently maximize muLECs'

function during wake. Consistently, muLECs are required for normal recovery after

the heightened metabolic demand of a seizure, suggesting that muLEC clearance

contributes to both normal and disrupted brain homeostasis.

Impact Statement:

28 A meningeal scavenger cell population called Mural Lymphatic Endothelial Cells

29 preferentially clear macromolecules from the CSF during the waking day, and this

30　uptake is regulated independently by brain state and circadian cues in a substrate-
31　specific manner.

Introduction

The mechanisms that maintain fluid homeostasis and clearance of waste
macromolecules from the brain remain incompletely understood. Current frameworks
propose that brain waste clearance involves the movement and subsequent
excretion of soluble waste via fluid exchange between the interstitial fluid (ISF) of the
brain and the cerebrospinal fluid (CSF) of the encasing leptomeninges along with
intracellular degradation of waste by resident parenchymal and meningeal cell
populations (Neumann et al, 2008; Liu et al, 2022; Zhan et al, 2025). In 2017, a new
scavenger cell type was discovered that resides in the meninges of zebrafish termed
Mural Lymphatic Endothelial Cells (muLECs; also termed brain LECs and
Fluorescent Granular Perithelial cells) (Bower et al, 2017; Galanternik et al, 2017;
van Lessen et al, 2017), and subsequently a similar cell type has been described in
mammals (Shibata-Germanos et al, 2020; Siret et al, 2022). Similar to other LECs,
muLECs arise from vascular endothelium and rely on lymphatic developmental
signals (Bower et al, 2017; Galanternik et al, 2017; van Lessen et al, 2017).
Uniquely, however, muLECs do not form lumenised vessels, instead forming a
loosely connected meshwork of cells that closely associate with meningeal blood
vessels covering the adult zebrafish brain (van Lessen et al, 2017). Although the
physiological functions of muLECs are yet unknown, these cells are poised as CSF
scavengers to clear metabolic waste from the brain. For example, muLECs readily
endocytose a variety of macromolecules spanning proteins, lipoproteins, liposomes,
and polysaccharides, with a higher uptake efficiency than brain-resident microglia
(Huisman et al, 2021). Recent work has demonstrated that muLEC development is
sensitive to, and regulated by, neuronal activity (Li et al, 2025), hinting at a functional
association between the brain and this meningeal cell population.

Over the past decade, numerous studies have reported that brain clearance
pathways are regulated by both behavioural state and circadian cues, although there
is some discrepancy as to whether this drives peak clearance preferentially during
the sleep or wake period. CSF oscillations in the ventricles that are dependent on

sleep stages have been observed in humans (Fultz et al, 2019) and birds (Ungurean et al, 2023) and are thought to drive fluid influx into the brain. In mice, CSF influx has been reported to be higher during sleep (Xie et al, 2013), potentially driven by sleep-associated changes in arterial pulsations (Bojarskaite et al, 2023) and circadian regulation of Aquaporin-4 water channels during the subjective rest period (Hablitz et al, 2020). Clearance of exogenous tracers from the brain has been shown to be elevated during sleep (Xie et al, 2013; Eide et al, 2021), whereas lymphatic drainage of CSF from the head cavity via meningeal lymphatic vessels was higher during the subjective active period (Hablitz et al, 2020). Microglia, the brain-resident phagocytes, show sleep- and circadian-regulation in morphology and uptake capacity of amyloid β (Liu et al, 2019; Clark et al, 2022; Parhizkar et al, 2023). However, recent work has reported preferential tracer clearance during wakefulness (Miao et al, 2024), challenging the notion that brain clearance is enhanced during sleep and may in fact be highest during wake. Nevertheless, these data collectively demonstrate brain clearance is dynamic across the day-night cycle. Given that muLECs are poised to remove waste from the brain, we hypothesized that their function may likewise show dynamic regulation over the 24-hour day-night cycle.

Here, we developed an *in vivo*, quantitative method to measure muLEC CSF solute clearance over the day-night cycle in zebrafish larvae and show that clearance of multiple substrate classes is highest during the waking day. Experimental disentanglement of light, circadian, and sleep-wake cues surprisingly revealed that while muLEC-uptake capacity for both protein and low-density lipoprotein (LDL) peaks during the day-time, these are independently regulated by a muLEC-autonomous circadian oscillator and sleep-wake state, respectively. Perturbing muLEC function slowed behavioural recovery following a seizure, suggesting that muLECs contribute to maintaining solute homeostasis during periods of high metabolic demand.

Results

muLEC morphology is dynamic over the day-night cycle

If muLEC function changes across the 24-hour day-night cycle, we reasoned that
this regulation may be detectable as alterations in cellular morphology during the day
versus night. To test this, we first raised *Tg(flt4:mCitrine)* embryos, which have
fluorescently labelled muLECs (van Impel et al, 2014) and other lymphatic
vasculature, to 6 days post fertilisation (6dpf) on opposing 14hr:10hr light-dark cycles
to yield developmentally matched larvae in different phases of their day-night cycle.
At 6 dpf, muLECs form bilateral loops in the developing meninges dorsal to the optic
tectum (Figure 1a) and harbour distinctive, round intracellular inclusions of varying
sizes and filopodial-like structures (Figure 1d; S1a; S1g). During the day, the total
volume (Figure 1b, p=0.008) and surface area (Figure S1c, p=0.0007) of muLECs
were significantly bigger compared to at night. Curiously, these daytime size
changes were also accompanied by a reorganization of muLEC interactivity, with
significantly more disconnections between muLECs during the day versus the night
(Figure 1c, p=0.049), although neither the width (Figure S1e, p=0.15) nor length
(Figure S1f, p=0.27) of remaining connections were altered (Figure S1b). In addition,
the muLECs had a trend towards more filopodia (Figure 1e) and significantly more
intracellular inclusions (Figure 1f, p=0.02) during the day than at night, although
there were no changes in either filopodial length (Figure S1d, p=0.45) or average
inclusion size (Figure 1g, p=0.24). As an increased number of intracellular inclusions
likely represent increased macromolecular internalization, these results suggest that
muLEC scavenger activity is higher during the waking day than at night.
117

muLEC-uptake of diverse macromolecules is higher during the day

To investigate whether diurnal changes in muLEC morphology relate to a difference
in uptake capacity, we developed a quantitative assay to measure muLEC-uptake
rates *in vivo*. Dorsally mounted *Tg(flt4:mCitrine)* larvae were co-injected into the
hindbrain ventricle with the pH-sensitive dye, pHrodo-Red avidin (hereafter termed
pHrodo-avidin), which is readily internalized by muLECs into acidified compartments,
and a 2MDa Alexa-fluor, 488 dextran (hereafter termed 2MDa-dextran), which is too
large for muLEC uptake (van Lessen et al, 2017) and serves as an injection
efficiency control. Following ventricular injection, larvae were imaged by confocal
microscopy for several hours to measure the increase of the pHrodo-avidin signal

within muLECs (Figure 2a). In a representative larva sampled at high frequency

(every 4 minutes), muLECs steadily accumulated pHrodo-avidin over the imaging

window in a sigmoidal pattern (Figure 2b). To enable sufficient experimental

throughput, we tested whether muLEC accumulation of pHrodo-avidin could be

accurately modelled if down sampled (once every 60 minutes) and found that the

data is well captured by a simple linear regression model with no apparent biases to

early, middle, or late imaging starting points post-injection (Figure S1a). Therefore, in

subsequent experiments we used the slope from a simple linear regression to

quantify muLEC-uptake of pHrodo-avidin.

Using the multiplexed muLEC-uptake assay, we then asked whether muLEC-uptake

rates of pHrodo-avidin varied over the day-night cycle. Developmentally matched

Tg(flt4:mCitrine) larvae raised on opposing light schedules were co-injected with

2MDa-dextran and pHrodo-avidin either in the middle of the day (ZT7, i.e., 7 hours

after lights on) or night (ZT19, i.e., 5 hours after lights off) (Figure 2c, d). Larvae were

imaged every 60 minutes for 5 hours post-injection with no difference in the

integrated intensity of 2MDa-dextran at the first imaging time point, indicating similar

injection volumes between groups (Figure S2e, p=0.95), and no time-sampling bias

(Figure S2g, p=0.95). The rate of muLEC-uptake of pHrodo-avidin was higher during

the late day compared to the middle of the night (Figure 2e,f; p=0.0044). This diurnal

change in muLEC-uptake capacity of pHrodo-avidin was not observed when larvae

were injected at the start of the day (ZT1) versus the start of the night (ZT14) (Figure

S2b, c; p=0.42).

As muLECs take up a wide variety of macromolecules (Huisman et al, 2020), we

next tested whether uptake of other cargo also varies across the day-night cycle. We

co-injected larvae during the day or night (Figure 2c) with 2MDa-dextran and pHrodo

conjugated to low-density lipoprotein (LDL; herein termed pHrodo-LDL) (Figure 2g)

or pHrodo conjugated to 10kDa dextran (dex; herein termed pHrodo-dex) (Figure 2j)

and imaged every 60 minutes for 6 hours. For both pHrodo-LDL (Figure S2h) and

pHrodo-dex (Figure S2k), the rates of uptake were best captured at low sampling

frequency by a one-phase association model. Similar to pHrodo-avidin (Figure 2e,f),

the uptake rates for both pHrodo-LDL (Figure 2h,i; p=0.0135) and pHrodo-dex

(Figure 2k,l; p=0.0089) were higher during the day compared to the night, with no

differences in initial injection volume (pHrodo-LDL, Figure S2i, p=0.95; pHrodo-dex,

Figure S2l, p=0.95) or in time-sampling (pHrodo-LDL, Figure S2j, p=0.95; pHrodo-

dex, Figure S2m, p=0.95) between groups. Thus, muLECs are more capable of

internalizing a variety of macromolecules during the waking day than at night.

168 Uptake of pHrodo-avidin is regulated by a muLEC-autonomous
169 circadian clock

The variation in muLEC uptake across the 24-hour day-night cycle could be

regulated by three co-varying, yet experimentally separable factors: the direct effects

of light exposure; the propensity to be awake (during the day) or asleep (at night); or

the timing of the molecular circadian clock (Figure 3a). We therefore designed a

series of pharmacological, genetic, and behavioural experiments to disentangle

which of these factors are regulating the daytime increase in muLEC uptake rates,

focusing on pHrodo-avidin.

First, to test whether the day-night difference in pHrodo-avidin uptake is directly

influenced by light, we tracked muLEC uptake in larvae that were entrained on

14hr:10hr light-dark cycles to 4-5dpf then switched into constant light (free-running

conditions) for 1 day. Although light levels are constant across the 24-hour day, the

phases of both the circadian clock (Kaneko and Cahill, 2005) and the diurnal timing

of wake and sleep states are preserved (Hurd and Cahill, 2002; Suppermpool et al,

2024). Developmentally matched *Tg(flt4:mCitrine)* larvae were then injected either

during the middle of the subjective day (circadian time 7, CT7) or night (CT19) as

entrained by the prior light-dark cycles (Figure 3b). Under these free-running

conditions, muLEC-uptake of pHrodo-avidin remained higher during the subjective

day than during the subjective night (Figure 3c, p=0.0035) with no difference in

loading injection volume between groups (Figure S4a, p=0.1) or time sampling

(Figure S4b, p=0.54). Since light exposure was the same at both timepoints, the

subjective day increase in muLEC uptake cannot be due to signalling events that

directly depend on a light-dark cycle alone.

Next, we assessed whether the 24-hour dynamic of pHrodo-avidin uptake is
modulated by brain state by acutely modulating sleep and wake behaviours
independently of the lighting conditions and time of day. To test whether acute
inductions of sleep modulated the rate of pHrodo-avidin uptake, larvae were injected
in the morning (ZT1) when locomotor activity is typically high and then transferred
into a solution containing either 1µM melatonin dissolved in 0.2% dimethyl sulfoxide
(DMSO) or a 0.2% DMSO control (Figure 3d). In zebrafish, melatonin is an acute
hypnotic that works downstream of the circadian clock to regulate sleep (Zhdanova
et al, 2001; Gandhi et al, 2015); melatonin-treated fish will therefore have a higher
propensity to sleep while remaining circadian phase-matched to their DMSO-treated
conspecifics (Suppermpool et al, 2024). muLEC-uptake of pHrodo-avidin remained
the same between melatonin and DMSO treated larvae (Figure 3e, p=0.098). In a
second sleep assay, we acutely but temporarily exposed larvae to the GABA-A
receptor antagonist, pentylenetetrazol (PTZ, 10mM), which induces a prolonged
increase in rebound sleep during the day with no effect on molecular circadian
rhythms (Reichert et al, 2019; Benoit et al, 2024), and then assessed muLEC uptake
capacity for pHrodo-avidin (Figure S3a). Similar to melatonin-treated fish, post-PTZ-
induced daytime sleep did not impact muLEC-uptake rates compared to fish-water
treated controls (Figure S3b, p=0.84).

Because inducing sleep during the day with drugs may not fully recapitulate natural
night-time sleep states, we also assessed whether promoting even more sleep
during the subjective night could impact muLEC rate measurements. To do this, we
used a gentle handling sleep deprivation protocol to keep larvae awake for 4hr at the
start of the night (ZT14-ZT18), which is followed by increased sleep consolidation
and duration (Suppermpool et al, 2024; Elias et al, 2025). We then injected pHrodo-
avidin immediately after sleep deprivation and tracked muLEC uptake for the
remainder of the night period (ZT19-24) (Figure 3f). Again, we observed no change
in uptake rates in this post-sleep deprivation period (Figure 3g, p=0.82). Taken
together, these results indicate that muLEC uptake of pHrodo-avidin is not
modulated by acute inductions of sleep, either during the day or night, suggesting
that the lower rate of uptake during the subjective night cannot be due to increased
sleep alone.

We then tested whether uptake of pHrodo-avidin could instead be boosted by
increased wakefulness at night. Larvae were injected in the middle of the night
(ZT19), immediately transferred into a solution containing 250µM caffeine to promote
wakefulness, and uptake was tracked for the remainder of the night (Figure 3h). This
caffeine-induced wakefulness did not affect muLEC uptake rates (Figure 3l, p=0.28).
To test the effects of natural waking states on uptake, larvae were injected with
pHrodo-avidin prior to the start (ZT13.5) of 4hr of gentle handling sleep deprivation,
and muLEC uptake rates were imaged immediately afterward at ZT18 (Figure 3j).
Again, no difference in uptake of pHrodo-avidin was observed during sleep
deprivation (Figure 3k, p=0.35), demonstrating that boosting wakefulness with either
drugs or gentle handling does not alter the muLEC's capacity to internalize pHrodo-
avidin.

Since neither acute light exposure nor sleep-wake state can explain the diurnal
variation in pHrodo-avidin uptake, we next examined whether the difference could be
due to non-behavioral outputs of the circadian clock. Zebrafish embryos raised from
fertilization under constant light and temperature conditions fail to develop coherent
24-hour circadian rhythms, as assessed by the loss of rhythmic clock gene
expression and behaviour (Whitmore et al, 2000; Prober et al, 2006; Wang et al,
2020; Suppermpool et al, 2024). We therefore tested the muLEC uptake of pHrodo-
avidin in larvae under these continuous light "clock-break" conditions compared to
developmentally matched larvae under entrained, free-running conditions (Figure
S3c). Larvae under free-running conditions had higher levels of pHrodo-avidin
accumulating in muLECs relative to "clock-broken" larvae when sampled in the
subjective day (CT7) but not during the subjective night (CT19) (Figure S3d; clock-
break vs CT7-13, p=0.0125; clock-break vs CT19-1 p=0.346). We conclude that day-
night differences in pHrodo-avidin uptake rely on intact circadian rhythms.

Zebrafish larvae, like many species, have endogenous molecular circadian
oscillators in most, if not all, cells (Whitmore et al, 2000; Mohawk et al, 2012; Moore
& Whitmore, 2014). Therefore, the clock-dependent diurnal variation in pHrodo-
avidin uptake could be due either to extrinsic circadian cues that signal onto muLECs
(non-cell autonomous), or to muLEC-intrinsic circadian rhythms (cell-autonomous)
that boost muLEC scavenger function during the subjective day. To test this, we

expressed in muLECs a dominant negative Clock (dnCLK), which can bind to its
partner BMAL but lacks the ability to bind DNA and direct transcription of key
circadian clock genes, thereby cell-autonomously disrupting the essential positive
arm of the transcription-translation feedback loop that underlies intrinsic circadian
timekeeping (Dekens & Whitmore, 2008; Livne et al, 2016; Kelu et al, 2020). Double
transgenic *Tg(flt4:Gal4);Tg(UAS:GFPp2AdnCLK)* larvae showed robust and uniform
expression of the dnCLK construct in muLECs at 5/6 dpf (Figure 3l), and this dnCLK
expression abolished the day-night difference in pHrodo-avidin uptake (Figure 3m, n;
p=0.7). Thus, a muLEC-intrinsic circadian oscillator regulates the daily rhythm of
pHrodo-avidin uptake capacity.

The muLEC-intrinsic clock regulates day-night uptake through the rhythmic expression of Mrc1a protein

An attractive hypothesis for how a muLEC-intrinsic circadian clock might dictate day-
night uptake rates of pHrodo-avidin is through timed regulation of the molecular
machinery required for uptake, via transcriptional or post-transcriptional
mechanisms. We therefore focused on the pattern-recognition receptor Mannose
receptor C-type lectin 1a (Mrc1a), which had previously been implicated in mediating
uptake in muLECs (van Lessen et al, 2017; Huisman et al, 2021). First, we
confirmed the requirement of *mrc1a* for uptake of avidin by injecting *Tg(flt4:mCitrine)*
embryos at the one-cell stage with Cas9 and guideRNAs targeting three separate
mrc1a exons to generate *mrc1a* F_0 knockouts (KO) (Figure 4a; S5a-d). We
confirmed that *mrc1a* KOs harboured a high-proportion of frameshift mutations
(Figure S5e) upstream of sequences encoding key Mrc1a protein domains (Figure
S5b), strongly indicating that *mrc1a* KOs are loss of function mutants. Unexpectedly,
compared to sibling controls injected with non-targeting gRNAs, *mrc1a* KOs
exhibited abnormal muLEC morphology (Figure 4b; S6a), with an increase in total
cell volume (Figure S6b) and mislocalisation of muLECs inside the tectal rings, an
ectopic distribution for this stage of development (Figure S6c), even though there
were no differences in the surrounding blood vasculature (Figure S6d-f) or microglia
(Figure S6g-k). Mrc1a must therefore play a key role in regulating either the
differentiation, cell division, or migration of muLECs at this stage as has been
previously described in macrophages (Sturge et al, 2007). Nevertheless, despite the

larger muLEC pool, and in line with previous work (Huisman et al, 2021), *mrc1a* loss-
of-function completely abolished muLEC uptake of pHrodo–avidin at all measured
time points post-injection (Figure 4c; 4d, p<0.0001). Interestingly, round and dynamic
pHrodo-avidin signal was still detected in the brain in *mrc1a* KOs suggesting that
Mrc1a is not required for pHrodo-avidin uptake by brain resident phagocytes (Figure
4b). This indicates that Mrc1a functions as a non-redundant receptor for avidin
internalization and muLECs have no compensatory pathways.
302
We then examined whether Mrc1a expression or availability changes in a circadian
clock-controlled manner by performing immunohistochemistry during the day or night
with a custom anti-Mrc1a antibody (Figure 4e; S5f) on muLEC-clock-intact
(*Tg(flt4:mCitrine))* or muLEC-clock-broken (*Tg(flt4:Gal4; Tg(UAS:GFPp2AdnCLK))*
larvae. In muLEC-clock intact larvae, Mrc1a protein expression was significantly
higher in the day compared to the night (Figure 4f, p=0.0048), but this diurnal
variation was abolished in muLEC-clock broken larvae (Figure 4g, p=0.135). These
results are consistent with a model in which the muLEC-intrinsic circadian clock
drives higher daytime Mrc1a protein levels that enhance muLEC capacity for the
internalization of pHrodo-avidin.
313

Diurnal variation of LDL uptake is regulated by sleep-wake state and not a muLEC-intrinsic clock

316
Unlike pHrodo-avidin, muLEC-uptake of both pHrodo–dex and pHrodo-LDL occur in
an Mrc1a-independent manner (Huisman et al, 2021). However, despite relying on
different receptor machinery, is the diurnal uptake of carbohydrate and lipoprotein
similarly controlled by a muLEC-autonomous circadian clock? To test this, larvae
expressing dnCLK in muLECs were injected with either pHrodo-dex or pHrodo-LDL
in the day and night, and the muLEC accumulation of these substrates was tracked
for several hours (Figure 5a). Despite interference with the muLEC-intrinsic clock,
uptake of both pHrodo-dex (Figure 5b, p=0.0039; Figure S7a) and pHrodo-LDL
(Figure 5c; p=0.03; Figure S7d) remained higher during the day than night, showing
that the muLEC-intrinsic clock is insufficient to drive diurnal changes in the uptake of
LDL and dextran.
328

We then asked whether pHrodo-dex and pHrodo-LDL uptake are instead governed
by non-circadian processes, such as brain state (e.g., sleep vs. wake). To test this,
Tg(flt4:mCitrine) larvae were injected with pHrodo-dex or pHrodo-LDL before the
start of the night (ZT13.5) and either kept awake by gentle handling for four hours
(ZT14-18) or left to sleep undisturbed (Figure 5d). Similar to pHrodo-avidin uptake
(Figure 3j), forced wakefulness did not impact muLEC-uptake rates of pHrodo-dex
(Figure 5e, p=0.216). Intriguingly, however, sleep deprivation significantly increased
the muLEC-uptake rates of pHrodo-LDL compared to undisturbed controls (Figure
5f, p=0.035). Furthermore, inducing night-time wakefulness either by sleep
deprivation or 250µM caffeine treatment significantly increased the size (Figure 5h,
p=0.0012; Figure S7k; Figure 5i p=0.0003; Figure S7o) but not the number of
muLEC inclusions (Figure S7h, p=0.156; S7l, p=0.13), while forced daytime sleep
with melatonin trended towards a decrease in inclusion size (Figure 5j, p=0.098;
Figure S7s) with no change in the average number of inclusions (Figure S7j, p>0.99)
or filopodia measurements (Figure S7i,j ; S7m,n ; S7q,r). Since LDL is trafficked into
larger, acidified late endosomes (Islam et al, 2022), these observations are
consistent with the heightened increase in LDL uptake seen after extended
wakefulness.

Thus, although muLECs have a higher capacity for all three macromolecular cargoes
during the waking day, this increase is accomplished by at least two, and possibly
three, independent mechanisms: avidin relies on a muLEC-autonomous clock; LDL
is attuned to the waking state of the brain; and dextran requires neither.

muLECs facilitate behavioural homeostasis post-seizure

Taken together, muLEC scavenging is modulated by circadian and behavioural state
to peak during the day when metabolic demand is greatest. Consequently, we
hypothesised that muLECs contribute to maintaining brain solute homeostasis during
metabolic challenges and that their disruption would impact physiological recovery
following a metabolically taxing insult. To test this, we used short multiphoton laser
pulses to ablate muLECs, which eliminated all muLECs while leaving the
surrounding blood vasculature intact (Figure 6a). Mock ablations were performed by

aiming the laser adjacent to muLECs. The mock- and muLEC-ablated larvae were
then treated with 10mM PTZ for 1 hour to induce seizures (Figure S3a) and
subsequently video-tracked for 48 hours in free-running conditions. Representative
traces show that both mock- and muLEC- ablated larvae have an initial multi-hour
period of extreme inactivity following PTZ treatment, but all larvae eventually resume
normal levels of locomotor activity (Figure 6b), with no difference in daytime (Figure
6d, p=0.329) or night-time activity levels between mock and ablated groups after
recovery (Figure 6e, p=0.592). However, muLEC-ablated larvae took significantly
longer (p<0.05, rank sum test) to recover normal locomotor activity (defined by
reaching activity levels equal to their next days' 50^{th} percentile) than mock-ablated
larvae (Figure 6c), with 50% of muLEC-ablated larvae recovering after 11.38 hours
compared to 7.75 hours for mock-ablated larvae. This delay in post-seizure recovery
was consistently found irrespective of the percentile cut-off used to define recovery
(Figure S8a). These data suggest that muLECs contribute to re-establishing solute
homeostasis after a metabolically taxing insult, which may also explain why muLEC
function is boosted to meet the higher metabolic demands of the waking day.
378

Discussion

Here, we demonstrate that at least two distinct mechanisms converge to boost CSF
solute clearance by a meningeal scavenger cell population during the waking day
and that this function is critical for recovery after a brain metabolic challenge. Given
that the muLECs are poised to participate in maintaining solute homeostasis of the
CSF, these results have important implications for the regulation and timing of brain
clearance.

The brain is metabolically demanding during the waking day, and this is well
reflected in the content of the CSF. The total protein content of the CSF undergoes
diurnal fluctuation and is higher in the latter portions of the waking day (Kakarla et al,
2023). The CSF abundance of a range of solutes spanning the neuromodulators
dopamine (Poceta et al, 2009), cortisol (Kalin et al, 1980), and hypocretin (Zeitzer et
al, 2003) as well as the waste products amyloid β (Bateman et al, 2007; Huang et al,
2012; Lucey et al, 2015) and tau (Holth et al, 2019) all peak during active periods as
a function of brain state or circadian cues. Additionally, peroxidated lipids build up

during the waking day and accumulate in glia as lipid droplets (Haynes et al, 2024),
while sleep disruption has been shown to alter the brain (Rorsman et al, 2025) and
CSF lipidome (Koethe et al, 2009; Saito et al, 2021; Dakterzada et al, 2023) as well
as the CSF distribution of lipid-associated apolipoproteins (Achariyar et al, 2016).
Together, these findings demonstrate that routes exist to connect the ISF of the brain
to the CSF within the surrounding meninges and further argue that wakefulness is a
period of active solute release into the CSF that creates a demand for the removal of
an array of macromolecular waste during behaviourally active periods.

Our data show that muLEC-uptake of proteins, lipoproteins, and carbohydrates are
all higher during the day, with uptake of specific proteins relying on the timed
expression of Mrc1a via a cell-autonomous clock. In contrast, lipoprotein uptake is
attuned to, and regulated by, wake-dependent cues, independently of the circadian
clock. Why use two distinct mechanisms to regulate protein versus lipoprotein
uptake, given that under normal conditions, both lead to a co-ordinated increase
during the waking day? One hypothesis is that each uptake pathway is regulated in
accordance with the clearance demands of the endogenous substrate/s that it
serves. Since the circadian clock anticipates the 24-hour day but does not typically
allow for more dynamic modulation, it could be that this cell-autonomous regulation
serves to clear predictably oscillating Mrc1a-dependent substrates from the CSF
such as glycosylated endo-lysosomal proteins (van der Zande et al, 2021;
Cummings, 2022; Orduña Dolado et al, 2024). Alternatively, muLECs might provide
an important barrier to prevent proteinaceous material entering the CSF/brain; such
risks might be more reliably likely to occur during active behavioural exploration. In
contrast, the regulation of lipoprotein uptake may require faster kinetics in response
to more acute events like sleep deprivation, when harmful material such as oxidized
lipids start to accumulate (Haynes et al., 2024). An open question for future work is
to what extent microglia and other endocytic cell populations also engage in this
dual-mode of regulation.

Another important question remains as to which wake-related signal(s) muLECs are
attuned, and how this signal(/s) is translated within muLECs to potentiate LDL
uptake. A recent study reported that muLEC development is dependent on, and
regulated by, neuronal activity via stimulation of radial glia that release

lymphangiogenic vascular endothelial growth factor C and D (Vegfc, Vegfd) into the
meninges (Li et al, 2025). It is not yet clear whether this neuron → glia → muLEC
signalling cascade is limited to controlling only muLEC development or could also be
used to link brain state to the muLEC scavenging functions that we have tested here.
The release of other VEGF signals has been shown to upregulate expression of the
lipid scavenger receptors Stabillin-1 and low-density lipoprotein receptor-related
protein 1 (LRP1) and to mediate lipid uptake and metabolism by macrophages and
liver sinusoidal cells in the periphery (Wary et al, 2003; Hagberg et al, 2010;
Falkevall et al, 2017; Shew et al, 2018; Tirronen et al, 2018; Ning et al, 2020; Luo et
al, 2022). Future work should examine whether lipid uptake by muLECs is also
regulated by VEGF-signalling and whether this provides a mechanism to link active
brain states to muLEC function.

Finally, we demonstrated that muLECs are required to restore behavioural
homeostasis following drug-induced seizure. It is well documented that seizures alter
the composition of CSF, leading to elevated total protein content, lactate, and tau
(Chatzikonstantinou et al, 2015; Tumani et al, 2015; Süße et al, 2019; Langenbruch
et al, 2021). Additionally, there are elevated CSF levels of somnogenic cytokines
such as interleukin-6 (Peltola et al, 1998; Vgontzas et al, 1999; Peltola et al, 2000;
Rohleder et al, 2012) and meningeal-derived prostaglandins that promote post-ictal
sleep and recovery (Kaushik et al, 2014; Mello & Oliveira, 2014). Therefore, it is
possible that muLEC scavenging is critical for refining the CSF milieu in response to
changing brain states, especially under insult. Future work should explore whether
CSF solute clearance by muLECs contributes to other disease states known to
impact CSF composition, such as in traumatic brain injury, chronic sleep disorders,
CNS infections, or neurodegenerative disease (Morganti-Kossman et al, 1997;
Vgontzas et al, 1997; Prasad et al, 2014; Carvalho et al, 2022).

To conclude, we demonstrate that CSF solute clearance by muLECs is differentially
regulated by state and circadian cues to peak during the waking day. We propose
that muLECs represent one node of many mechanisms involved in brain clearance
and homeostasis. Understanding how the wake-active muLECs operate alongside
other avenues of brain clearance proposed to undergo diurnal regulation will be

important for a fuller picture of how brain waste is regulated in both normal and
diseased brains.
463

Acknowledgements:

We thank Jeff Kelu and Simon Hughes for the UAS-dnCLK construct, Sandip Patel
for consultation on pH-sensitive dyes, the Payne lab for MiSeq, the UCL Fish Facility
for animal husbandry, and all members of the Rihel lab for constructive feedback on
all stages of the project. The work was supported by a Wellcome Trust Investigator
Award to JR (217150/Z/19/Z), a BBSRC grant to JR and TAH (BB/T001844/1), and a
Medical Research Council Doctoral Training Partnership (MRC-DTP) fellowship
awarded to TG.

Author Contributions:

TG, TK, JR and TAH conceived of the project. TK, with help from DK, imaged and
analysed muLEC morphology in Figure 1 and 5g. TG designed, performed and
analysed all uptake experiments (Figure 2-5) with help from LSM for Mrc1a
immunostaining (Figure 4e-g). TG, TK, and JR performed muLEC ablation
experiments (Figure 6). TG designed the figures and wrote the manuscript with input
from all authors. JR supervised the project.

Materials and Methods

Animals

Zebrafish (Danio rerio) husbandry and experiments were conducted according to
UCL Fish Facility standard protocols and under the project licenses PA8D4D0E5 and
PP6325955 awarded to JR under the auspices of the UK Animal Scientific
Procedures Act (1986). Embryos were kept in 10cm Petri dishes in E3 media (0.3
g/L Instant Ocean and 0.1% Methylene blue) in 14□hr-10□hr light–dark cycle
incubators at 28□°C, except of clock break experiments, when they were kept under
constant light from 0 dpf. Unless otherwise stated, all experiments were conducted
on larvae 5-6 days post fertilisation. At the end of all experiments, larvae were

494 euthanised using by anaesthetic overdose of MS222 (Sigma Aldrich, UK) or 2-
495 phenoxyethanol (ACROS Organics).

497 All transgenic lines were kept in a *nacre (mitfa)$^{w2/w2}$* mutant background. The
498 following lines were used for experiments:

Transgenic Line	Abbreviated Name	Reference
Tg(flt4^{BAC}:mCitrine)^{hu7135}	*Tg(flt4:mCitrine)*	van Impel et al, 2014
Tg(flt4^{BAC}:mCitrine)^{hu7135};Tg(kdr-l:HRAS-mCherry-CAAX)^{s916}	*Tg(flt4:mCitrine);Tg(kdrl:mCherry)*	van Impel et al, 2014; Hogan et al, 2009
Tg(flt4^{BAC}:mCitrine)^{hu7135} Tg(mpeg1:Gal4-FF)gl25; Tg(UAS:E1b:nfsB.mCherry)^{c264}	*Tg(flt4:mCitrine); Tg(mpeg:mCherry)*	van Impel et al, 2014; Voelz, Gratacap, & Wheeler, 2015
Tg(Ola_Flt4:GAL4FF)^{hu9236};Tg(UAS:EGFPp2AdnCLK)	*Tg(flt4:Gal4)Tg(UAS: GFPp2AdnCLK)*	van Impel et al, 2014;

500
501
Transgenesis

A plasmid encoding the pT2-UAS-EGFP-p2A-dnCLK-5xMyc construct was kindly
provided by Jeff Kulu and Yoav Gothlif. Approximately 1nl of plasmid (20ng/µl)
alongside Tol2 transposase mRNA (50ng/µl) was injected in *Tg(flt4:Gal4FF)^{hu9236}*
(van Impel et al., 2014) embryos at the one-cell stage as previously described (Clark
et al., 2011). F_0 larvae from a single clutch were screened at 4dpf for positive EGFP
expression and raised to adulthood. Founder adults were outcrossed to *nacre*
(mitfa)$^{w2/w2}$ larvae to screen for transgene germline integration. A Nikon fluorescent
SMZ1270i fluorescent stereo microscope was used to identify F_1 progeny with the
brightest EGFP fluorescence and further propagated. Experiments were performed
on F_1 or F_2 larvae from a single parental clutch incross.

514
515
516 **muLEC cellular and subcellular morphology**

517
518 To visualize and quantify muLEC morphology, 6dpf *Tg(flt4:mCitrine)* larvae were
519 culled by anesthetic overdose (MS222, approximately 420 mg/L) and fixed in 4%
520 paraformaldehyde (PFA). Following washing in 1x phosphate buffered saline (PBS)
521 and 1x PBS with Triton (PBST), larvae were ventrally mounted in 1% low-melt

agarose and imaged using a LSM 980 inverted confocal microscope with Airyscan 2 (Zeiss). For assessing muLEC volume, surface area, disconnectedness and width/length of muLEC connections a x20/1.0 NA objective was used and z-stacks were obtained at 1µm intervals of 4076 x 4084 px or 4060 x 4060 px and 16-bit using SR-8Y acquisition mode. For assessing muLEC inclusions and filopodia a x40/1.2 NA objective was used and z-stacks were obtained at 0.21 µm intervals of 4084 x 4084 px and 16-bit using CO-8Y acquisition mode. mCitrine was excited at 488nm.

Analysis of muLEC volume, surface area, disconnectedness were performed in Imaris. ROIs were constructed around all muLECs within one hemisphere and raw z-stacks were segmented and auto-thresholded to generate 3D surface reconstructions using the Imaris surface function. For quantification of the width/length of muLEC connections, 2D maximum projections were analysed in ImageJ (NIH) using a modified macro script from McDowell et al, 2020. For assessing muLEC inclusion sizes, 2D maximum projections were analysed in ImageJ with a custom-built macro (https://github.com/JRihel/muLEC-Circadian-Imaging). Filopodial lengths and numbers were manually analyzed using the segmented line tool in ImageJ.

Intraventricular injection assay for assessing muLEC uptake

Lighting schedules

Larvae were reared in incubators kept at 28°C on various lighting schedules as specified (hours light: hours dark): light:dark, 14hr:10hr; free-running, 14:hr:10hr up to 4dpf then transferred to constant light; clock-break, 24hr:0hr, constant light. When multiple ZT times or different light schedules were compared within an experiment, water bath incubators were set up to run in parallel. Temperature and light transitions of water bath incubators were recorded with a HOBO Pendant (Onset Data Loggers) immersed in the water bath. If larvae were collected in their dark period, larvae were removed from the water baths using an external red head light (Blackburn Local Bike Rear Light 15 Lumen, UK) and transported in light proof boxes to be taken through the intraventricular injection assay where ambient light exposure was limited as much as possible.

Drug treatment

For PTZ-treatment, Tg(*flt4:mCitrine*) larvae were transferred to a solution containing
10mM pentylenetetrazol (PTZ) (Sigma-Alrich, UK) dissolved in E3 media or the
same volume of E3 media for 1 hr (ZT1-2) as previously described (Reichert et al,
2019). PTZ-treated fish were assessed for behavioural hyperactivity and tachycardia
using a Nikon SMZ1500 bright-field microscope, subsequently washed out of
solution 3x in E3 media, anaesthetised in MS222 (42 mg/L) and intraventricularly
injected (see *injection regime*) and imaged for several hours (see *live imaging*). For
melatonin-treatment, Tg(*flt4:mCitrine*) larvae were intraventricularly injected at ZT1
and immediately transferred into solution containing 1µM melatonin (Sigma-Alrich,
UK) dissolved in 0.2% DMSO or 0.2% DMSO control for subsequent imaging. For
caffeine-treatment, Tg(*flt4:mCitrine*) larvae were intraventricularly injected at ZT19
and immediately transferred into solution containing 250µM caffeine (Sigma-Alrich,
UK) dissolved in E3 media or E3 media alone for subsequent imaging. For inclusion
and filopodia analysis, the treatment group was placed in melatonin solution (1 µM)
or caffeine solution (250µM) and control group in 0.2% DMSO or H20 respectively,
for 4 hrs (ZT1-5 for melatonin and ZT-19-23 for caffeine) before they were culled,
fixed in 4%PFA and imaged (see *muLEC cell and subcellular morphology)*.

Sleep deprivation assay

Sleep deprivation by gentle handing was performed as previously described
(Suppermpool et al., 2024). Prior to lights-off (ZT14), *Tg(flt4:mCitrine)* larvae were
transferred into two separate identical 24-well clear plates (ThermoFisher, UK).
Immediately after lights off (ZT14-18) both plates were placed in a dark room under
dim-red light (Blackburn Local Bike Rear Light 15 Lumen, UK). Larvae in one plate
were left undisturbed while larvae in the other plate were manually sleep deprived by
repeated gentle stimulation using a No. 1-2 paintbrush (Daler Rowney Graduate
Brush, UK) to prevent larvae from being immobile for longer than 1 minute. Placing
the paintbrush into the water of the well was usually stimulatory enough to elicit
movement, but if larvae remained immobile, they were gently touched until
movement was initiated. The 4-hr sleep deprivation was performed by two
experimenters in 2-hr shifts.

Injection regime

Intraventricular injections were conducted as previously described (Shibata-
Germanos et al, 2020) using a PM1000 Cell Microinjector (MicroData, Instrument,
Inc) with glass capillary needles (Harvard Apparatus, # GC100F-15) prepared with a
Micropipette Puller (Shutter Instruments). Larvae were anaesthetised in E3 media
with MS222 (42 mg/L) and mounted dorsally in 2% low-melt agarose (Sigma Aldrich,
UK) dissolved in ddH_2O on a 60mm plastic petri dish lid which was scratched with
forceps prior to mounting to reinforce agarose adherence. The following substrates
were injected into the rhombenchephalic ventricle with a total bolus volume 1nl at 20
psi pressure at the specified concentration: Dextran, Fluorescein 2MDa, 1mg/ml
(ThermoFisher, UK); pHrodo Red Avidin, 2mg/ml (ThermoFisher, UK); pHrodo Red
Dextran 10kDa, 0.5mg/ml (ThermoFisher, UK); pHrodo Red LDL, 1mg/ml
(ThermoFisher, UK). pHrodo-conjugated dyes were pre-mixed with Dextran,
Fluorescein 2MDa immediately prior to co-injection so that in each 1nl bolus, 0.5nl
constituted each respective dye.

Live imaging

All live imaging data were acquired using a Zeiss LSM 980 confocal microscope with
a 20x/1.0 NA water immersion objective. Z-stacks were obtained with sequential
acquisition settings of 1024 x 1024 pixels. Z-depth intervals ranged from 1-1.5µm
which were standardised within each assay. Dextran, Fluorescein 2MDa was excited
at 488nm and pHrodo Red Avidin, pHrodo Red Dextran and pHrodo Red LDL were
excited at 546nm. Multi-position time lapses were set up and larvae were
sequentially imaged every 60 minutes post-injection over the course of 6-7 hours,
unless otherwise stated.

Image Processing

Confocal stacks were processed and analysed in Imaris. Briefly, raw z-stacks were
segmented and thresholded to provide representative surfaces of a given object in
3D space. Segmentation and thresholding limits were standardised across all fish in
a given assay. Standardised regions of interest (ROIs) were constructed for muLECs
along the midline of both tectal rings as representative to reduce variability of data

from lateral muLECs at lower z-depths. For assessment of the initial injection bolus,

3D surfaces of Dextran, Fluorescein 2MDa signal were constructed using

standardised ROIs spanning the most anterior point of the telencephalic ventricle to

the posterior point of the diencephalic ventricle. All z-stacks are represented as

dorsal maximum projections unless otherwise stated.

Quantitative immunohistochemistry and imaging

Larvae were culled by anaesthetic overdose (MS222, approximately 420 mg/L) and

fixed overnight (12-16 hours) in 4% PFA supplemented with 4% sucrose at 4°C. The

following day, larvae were washed three times in 1x PBS and dissected. To remove

batch effects, day and night larvae were given distinctive oblique cuts on the tail for

later identification and transferred to the same vial for staining. Dissected larvae

were permeablised with Proteinase K (30 µg/mL) for 90 seconds, post-fixed for 20

minutes in 4% PFA at room temperature and subsequently washed three times in

PBST. Samples were blocked with blocking buffer (10% normal goat serum; 1%

DMSO; 0.5% Triton-X100 in PBS) for 1 hr at room temperature. Samples were then

incubated with a custom anti-Mrc1a antibody (1:200 dilution, rabbit polyclonal, Sino

Biological, UK) in blocking buffer overnight at room temperature. Primary antibody

was removed, and samples were rinsed 3x in PBST and washed 4x30mins in PBST.

Samples were placed in fresh block buffer with 1:500 anti-rabbit Alexa Fluor 568

(ThermoFisher, UK). Excess secondary antibodies were washed 4-6 times for 30

minutes and mounted dorsally in 2% LM agarose for imaging.

All imaging data were acquired using a Zeiss LSM 980 confocal microscope with a

20x/1.0NA water-immersion objective. Z-stacks were obtained at 1.5µm depth

intervals with sequential acquisition settings of 1024 x 1024 pixels. Endogenous

mCitrine and anti-rabbit Alexa Fluor 568 were excited at 488nm and 561nm,

respectively. Image analysis was conducted in Imaris. Raw z-stacks were

segmented and standardised thresholds were applied to provide representative 3D

surfaces of muLEC and anti-Mrc1a immunofluorescence.

Generation of *mrc1a* F$_0$ knockouts

crRNA Selection

F_0 knockout larvae were generated as described previously (Kroll et al., 2021; Kroll et al., 2025). All crRNAs used in F_0 knockout experiments were designed using CHOPCHOP (chopchop.cbu.uib.no) (Labun et al., 2019). Putative targets were checked for SNPs using SNPfisher (Butler et al., 2015). Sequences of the targets (and genomic positions) used to generate F_0 knockout are as follows with the protospacer adjacent motif (PAM) underlined (5'→3'): mrc1a_guide_1, TGAGGGTCAACGCTTTCGAT<u>GGG</u> (chr7:58939687); mrc1a_guide_2, ACTACGGAGGGAAGATCAGA<u>TGG</u> (chr7:58938982); mrc1a_guide_3, TGGGACAGTGATCCAGTGAC<u>TGG</u> (chr7:58936563). Scrambled, nonsense crRNAs were prepared identically and had the following sequences (5'→3'): scrambled_guide_1, CCCGACTAACATGGCGAGAC; scrambled_guide_2, GTATCCACCACGGGAGACAC; scrambled_guide_3, GTATATGGTTTGGGGCCACT; scrambled_guide_4, CGCAAGTTAAGGCCGACCCA. Genomic positions are for reference genome GRCz11 (danRer11).

Cas9/gRNA preparation

Cas9/gRNA were prepared as previously described (Kroll et al., 2021; Kroll et al., 2025). CRISPR-Cas9 RNPs were composed of: crRNA (Alt-R® CRISPR-Cas9 crRNA) and tracrRNA (Alt-R® CRISPR-Cas9 tracrRNA), Cas9 (Alt-R® S.p. Cas9 Nuclease V3) (Integrated DNA Technologies, UK). crRNA and tracrRNA were resuspended in Duplex buffer (IDT, UK) to form 200 µM stocks. Stocks of crRNA and tracrRNA were stored at −70˚C before use and Cas9 was stored at −20˚C before use. Each crRNA was annealed separately with the tracrRNA by mixing 1 µL crRNA 200 µM; 1 µL tracrRNA 200 µM; 1.28 µL Duplex buffer. The mix was heated to 95˚C for 5 min, then cooled on ice, to obtain a 61 µM gRNA solution. The gRNA solutions were then mixed in equal volumes with Cas9 (1 µL gRNA 61 µM; 1 µL Cas9 61 µM directly from the IDT vial), incubated at 37˚C for 5 min then cooled on ice, generating three 30.5 µM RNP solutions. The three RNP solutions were pooled; the final concentration of each RNP in the pool was thus 10.2 µM and the total RNP concentration 30.5 µM. The RNPs were usually kept overnight in a 4˚C fridge on ice

before injections the following day. Some experiments used RNPs stored at −70°C

for a few weeks.

Injections

Approximately 1nL of the three-RNP pool (~10.2fmol per RNP) was injected into the

yolk at the single-cell stage before cell inflation. Injected embryos were kept at room

temperature for approximately 45 minutes post-injection before transfer to a 28.5°C

incubator, as delaying the first cell divisions may increase mutagenesis and reduce

the diversity of alleles (Terzioglu et al., 2020).

Preparation of samples for Illumina MiSeq

Target loci were deep sequenced by Illumina MiSeq as previously described (Kroll et

al., 2021; Kroll et al., 2025). 8dpf *mrc1a* KO and scrambled control larvae were

culled by MS222 overdose and genomic DNA was extracted by HotSHOT as

previously described (Meeker et al., 2007). Individual larvae were transferred to a

96-well PCR plate where excess liquid was removed from each well before adding

50 µl of 1x base solution (25 mM KOH, 0.2 mM EDTA in water). Plates were sealed

and incubated at 95°C for 30 min then cooled to room temperature before the

addition of 50 µL of 1x neutralisation solution (40 mM Tris-HCl in water). Genomic

DNA was then stored at −20°C. PCR primers were designed for each target locus

using Primer-BLAST (NCBI) to amplify a window of 150–200 bp with at least 30 bp

between each primer binding site and the predicted double-strand break site.

Forward and reverse PCR primers were ordered with overhang sequences at the 5′-

end and 3' end of each primer to allow indexing (Forward 5' overhang:

TCGTCGGGCAGCGTCAGATGTGTATAAGAGACAG;   Reverse   3'   overhang:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG).

Each PCR well contained: 7.98 µL PCR mix (2 mM $MgCl_2$, 14 mM pH 8.4 Tris-HCl,

68 mM KCl, 0.14% gelatine in water, autoclaved for 20 min, cooled to room

temperature, chilled on ice, then added 1.8% 100 mg/mL BSA and 0.14% 100 mM

d[A, C, G, T]TP), 3 µL 5✕ Phusion HF buffer (New England Biolabs, UK), 2.7 µL

dH2O, 0.3 µL forward primer (100 µM), 0.3 µL reverse primer (100 µM), 0.12 µL

Phusion High-Fidelity DNA Polymerase (New England Biolabs, UK), 1.0 µL genomic

DNA; for a total of 15.4 µL. The PCR plate was sealed and placed into a

thermocycler. The PCR program was: 95°C – 5 min, then 40 cycles of: 95°C – 30

sec, 60°C – 30 sec, 72°C – 30 sec, then 72°C – 10 minutes, then cooled to 10°C until

collection. The PCR product's concentration was quantified with Qubit (dsDNA High

Sensitivity or Broad Range Assay) and its length was verified on a 1.5% agarose gel

with GelRed (Biotium, USA). Excess primers and dNTPs were removed by ExoSAP-

IT (ThermoFisher, UK) following the manufacturer's instructions. The samples were

then sent for Illumina MiSeq, which used MiSeq Reagent Nano Kit v2 (300 Cycles)

and analysed as in Kroll et al, 2025.

Morphological assessments of mrc1a F_0 knockouts

mrc1a KOs and scrambled controls were generated in the following lines:

Tg(*flt4:mCitrine*), *Tg*(*flt4:mCitrine);Tg*(*kdrl:mCherry*), and *Tg(flt4:mCitrine;*

Tg(mpeg:mCherry). At 5dpf, larvae were anaesthetised (MS222, 42 mg/L), mounted

dorsally in 2% LM agarose on a small petri dish lid. Imaging data were acquired

using a Zeiss LSM 980 confocal microscope with a 20x/1.0NA water immersion

objective. Z-stacks were obtained with sequential acquisition settings of 1024 x 1024

pixels with z interval depths of 1µm. mCitrine and mCherry were excited at 488nm

and 594nm, respectively.

Confocal stacks were processed and analysed in Imaris. For muLEC morphology

assessment, standardised ROIs were constructed bilaterally over the optic tectum

and standardised threshold and segmentation were applied to construct 3D surfaces

using the Imaris Surface function. For assessment of the vascular network, as shown

previously (Spangenberg et al, 2023), standardised ROI were constructed from the

dorsal midline junction to the most anterior point of the prosencephalic artery to

depth of 120µm from the skins surface as visible by autofluorescent signal. The

Imaris Filament function with loops based on local intensity contrast without the

detection of spines was performed to reconstruct the vasculature within the ROI as a

connected network in 3D space. For assessment of *mpeg*[+] cells, the Imaris Surfaces

and Filament (with tree autopath algorithm) function was used to reconstruct
mCherry positive signal within a standardised ROI covering the head region and
eyes at a depth of 120µm from the autofluorescent skin into 3D space.

muLEC ablation and seizure recovery

muLEC ablation

Tg(flt4:mCitrine); Tg(kdrl:mCherry) larvae were raised to 5dpf, anaesthetised in 42
mg/L MS222 solution and dorsally mounted in 2% LM agarose. mCitrine positive
cells within the bilateral muLEC tectal loops and along the midline were ablated
using a Zeiss LSM 980 confocal microscope with two-photon laser at 800nm (Mai
Tai HP laser 690-1040nm tuneable, Spectra Physics). Control ablations were
performed by directing an identical laser pulse just adjacent to mCitrine positive
muLECs in a fluorescent negative space. Larvae were imaged before and after
muLEC-ablation and mock-ablation to confirm successful ablations and intact
surrounding blood vasculature. Larvae were re-imaged 3 days post ablation to
assess for regeneration. Regenerated muLECs were quantified in Imaris using the
Imaris Surface function.

PTZ-induced seizures and locomotor tracking

Following ablations, larvae were unmounted from agarose and transferred to a
solution 10mM PTZ dissolved in E3 media or the same volume of E3 media alone for
1 hr at approximately ZT7. PTZ-treated fish were assessed for behavioural
hyperactivity and tachycardia using a Nikon SMZ1500 bright-field microscope,
subsequently washed out of solution 3x in E3 media and transferred into individual
wells of a 96-well plate (ThermoFisher, UK) containing approximately 600µL of fish
water for behavioural tracking.

Locomotor activity of larvae was monitored using an automated video tracking
system (Zebrabox, Viewpoint LifeSciences) in a temperature-regulated room
(26.5°C) and illuminated with constant white light. Larval movement was recorded
using the Videotrack 'quantization' mode with the following detection parameters:
detection threshold, 15; burst, 100; freeze, 3; bin size, 60s. Tracking was performed

at 25 frames per second for approximately 48 hours. Behavioural data was analysed
with custom Matlab (R2024a, the Mathworks) scripts (available at
https://github.com/JRihel/muLEC-Circadian-Imaging).
802
803
Data Analysis and Statistical tests

Imaging processing, analysis and presentation was performed in Imaris 9.82 - 10.0.0
(Imaris, UK) and Fiji-Image J version 2.14.0i. Graph construction and statistical
analyses were performed in Prism 9.2.0 (GraphPad software, USA). All data were
assessed for normality by a Shapiro-Wilk test. Parametric data were assessed by
One-way ANOVA (two-tailed), Mixed-Effects Model and log rank tests, and non-
parametric data were assessed by Mann-Whitney U tests (two tailed) as specified.
All tests were conducted with an alpha of 0.05. Figures and schematics were
constructed in Inkscape version 1.4.
814
Bibliography

Achariyar, T.M., Li, B., Peng, W., Verghese, P.B., Shi, Y., McConnell, E.,
Benraiss, A., Kasper, T., Song, W., Takano, T., Holtzman, D.M., Nedergaard, M. and
Deane, R. (2016). Glymphatic distribution of CSF-derived apoE into brain is isoform
specific and suppressed during sleep deprivation. *Molecular Neurodegeneration*,
[online] 11(1), p.74. doi:https://doi.org/10.1186/s13024-016-0138-8.

Agorastos, A., Hauger, R.L., Barkauskas, D.A., Moeller-Bertram, T., Clopton,
P.L., Haji, U., Lohr, J.B., Geracioti, T.D., Patel, P.M., Chrousos, G.P. and Baker,
D.G. (2014). Circadian rhythmicity, variability and correlation of interleukin-6 levels in
plasma and cerebrospinal fluid of healthy men. Psychoneuroendocrinology, 44,
pp.71–82. doi:https://doi.org/10.1016/j.psyneuen.2014.02.020.

Bateman, R.J., Wen, G., Morris, J.C. and Holtzman, D.M. (2007). Fluctuations
of CSF amyloid-β levels. Neurology, 68(9), pp.666–669.
doi:https://doi.org/10.1212/01.wnl.0000256043.50901.e3.

Benoit, E., Lyons, D.G. and Rihel, J. (2023). Noradrenergic tone is not
required for neuronal activity-induced rebound sleep in zebrafish. Journal of
Comparative Physiology B, 194(3), pp.279–298. doi:https://doi.org/10.1007/s00360-
023-01504-6.

Bojarskaite, L., Vallet, A., Bjørnstad, D.M., Gullestad Binder, K.M., Cunen, C.,
Heuser, K., Kuchta, M., Mardal, K.-A. and Enger, R. (2023). Sleep cycle-dependent
vascular dynamics in male mice and the predicted effects on perivascular
cerebrospinal fluid flow and solute transport. Nature Communications, [online] 14(1),
p.953. doi:https://doi.org/10.1038/s41467-023-36643-5.

Bower, N.I., Koltowska, K., Pichol-Thievend, C., Virshup, I., Paterson, S.,
Lagendijk, A.K., Wang, W., Lindsey, B.W., Bent, S.J., Baek, S., Rondon-Galeano,
M., Hurley, D.G., Mochizuki, N., Simons, C., Francois, M., Wells, C.A., Kaslin, J. and
Hogan, B.M. (2017). Mural lymphatic endothelial cells regulate meningeal
angiogenesis in the zebrafish. Nature Neuroscience, [online] 20(6), pp.774–783.
doi:https://doi.org/10.1038/nn.4558.

Butler, M.G., Iben, J.R., Marsden, K.C., Epstein, J.A., Granato, M. and
Weinstein, B.M. (2015). SNPfisher: tools for probing genetic variation in laboratory-
reared zebrafish. Development. doi:https://doi.org/10.1242/dev.118786.

Carvalho, D.Z., St, E.K., Przybelski, S.A., Morgenthaler, T.I., Machulda, M.M.,
Boeve, B.F., Petersen, R.C., Jack, C.R., Graff-Radford, J., Vemuri, P. and Mielke,
M.M. (2022). Sleepiness in Cognitively Unimpaired Older Adults Is Associated With
CSF Biomarkers of Inflammation and Axonal Integrity. Frontiers in Aging
Neuroscience, 14. doi:https://doi.org/10.3389/fnagi.2022.930315.

Chatzikonstantinou, A., Ebert, A.D. and Hennerici, M.G. (2015). Cerebrospinal
fluid findings after epileptic seizures. Epileptic Disorders, 17(4), pp.453–459.
doi:https://doi.org/10.1684/epd.2015.0779.

Clark, G.T., Yu, Y., Urban, C.A., Fu, G., Wang, C., Zhang, F., Linhardt, R.J.
and Hurley, J.M. (2022). Circadian control of heparan sulfate levels times
phagocytosis of amyloid beta aggregates. PLOS Genetics, 18(2), p.e1009994.
doi:https://doi.org/10.1371/journal.pgen.1009994.

Clark, K.J., Urban, M.D., Skuster, K.J. and Ekker, S.C. (2011). Transgenic
Zebrafish Using Transposable Elements. Methods in Cell Biology, pp.137–149.
doi:https://doi.org/10.1016/b978-0-12-374814-0.00008-2.

Cummings, R.D. (2022). The mannose receptor ligands and the macrophage
glycome. Current Opinion in Structural Biology, 75, pp.102394–102394.
doi:https://doi.org/10.1016/j.sbi.2022.102394.

Dakterzada, F., Benítez, I.D., Targa, A., Carnes, A., Pujol, M., Jové, M.,
Mínguez, O., Vaca, R., Sánchez-de-la-Torre, M., Barbé, F., Pamplona, R. and Piñol-
Ripoll, G. (2023). Cerebrospinal fluid lipidomic fingerprint of obstructive sleep apnoea
in Alzheimer's disease. Alzheimer's Research & Therapy, [online] 15(1).
doi:https://doi.org/10.1186/s13195-023-01278-7.

Dekens, M. and Whitmore, D. (2008). Autonomous onset of the circadian
clock in the zebrafish embryo. The EMBO Journal, 27(20), pp.2757–2765.
doi:https://doi.org/10.1038/emboj.2008.183.

Dolado, A.O., Stomrud, E., Ashton, N.J., Nilsson, J., Quijano-Rubio, C.,
Jethwa, A., Brum, W.S., Westman, A.B., Zetterberg, H., Blennow, K., Janelidze, S.
and Hansson, O. (2024). Effects of time of the day at sampling on CSF and plasma
levels of Alzheimer' disease biomarkers. Alzheimer s Research & Therapy, 16(1).
doi:https://doi.org/10.1186/s13195-024-01503-x.

Eide, P.K., Vinje, V., Pripp, A.H., Mardal, K.-A. and Ringstad, G. (2021). Sleep
deprivation impairs molecular clearance from the human brain. Brain, [online] 144(3).
doi:https://doi.org/10.1093/brain/awaa443.

Elias, L.J., Khoo, H., Kroll, F., Zhang, C., Hur, S.C., Rihel, J. and Blackshaw,
S. (2025). JACUZI-SD: An automated, high-throughput, minimally stressful approach
to sleep depriving larval zebrafish. bioRxiv (Cold Spring Harbor Laboratory).
doi:https://doi.org/10.1101/2025.04.03.647099.

Falkevall, A., Mehlem, A., Palombo, I., Heller Sahlgren, B., Ebarasi, L., He, L.,
Ytterberg, A.J., Olauson, H., Axelsson, J., Sundelin, B., Patrakka, J., Scotney, P.,
Nash, A. and Eriksson, U. (2017). Reducing VEGF-B Signaling Ameliorates Renal

Lipotoxicity and Protects against Diabetic Kidney Disease. Cell Metabolism, 25(3),
pp.713–726. doi:https://doi.org/10.1016/j.cmet.2017.01.004.

Fultz, N.E., Bonmassar, G., Setsompop, K., Stickgold, R.A., Rosen, B.R.,
Polimeni, J.R. and Lewis, L.D. (2019). Coupled electrophysiological, hemodynamic,
and cerebrospinal fluid oscillations in human sleep. Science, [online] 366(6465),
pp.628–631. doi:https://doi.org/10.1126/science.aax5440.

Galanternik, M.V., Castranova, D., Gore, A.V., Blewett, N.H., Jung, H.M.,
Stratman, A.N., Kirby, M.R., Iben, J., Miller, M.F., Kawakami, K., Maraia, R.J. and
Weinstein, B.M. (2017). A novel perivascular cell population in the zebrafish brain.
eLife, [online] 6, p.e24369. doi:https://doi.org/10.7554/eLife.24369.

Gandhi, Avni V., Mosser, E.A., Oikonomou, G. and Prober, David A. (2015).
Melatonin Is Required for the Circadian Regulation of Sleep. Neuron, 85(6),
pp.1193–1199. doi:https://doi.org/10.1016/j.neuron.2015.02.016.

Haberland, M.E., Mottino, G., Le, M. and Frank, J.S. (2001). Sequestration of
aggregated LDL by macrophages studied with freeze-etch electron microscopy.
Journal of Lipid Research, 42(4), pp.605–619. doi:https://doi.org/10.1016/s0022-
2275(20)31170-6.

Hablitz, L.M., Plá, V., Giannetto, M., Vinitsky, H.S., Stæger, F.F., Metcalfe, T.,
Nguyen, R., Benrais, A. and Nedergaard, M. (2020). Circadian control of brain
glymphatic and lymphatic fluid flow. Nature Communications, [online] 11(1), p.4411.
doi:https://doi.org/10.1038/s41467-020-18115-2.

Hagberg, C.E., Falkevall, A., Wang, X., Larsson, E., Huusko, J., Nilsson, I.,
van Meeteren, L.A., Samen, E., Lu, L., Vanwildemeersch, M., Klar, J., Genove, G.,
Pietras, K., Stone-Elander, S., Claesson-Welsh, L., Ylä-Herttuala, S., Lindahl, P. and
Eriksson, U. (2010). Vascular endothelial growth factor B controls endothelial fatty
acid   uptake.   Nature,   464(7290),   pp.917–921.
doi:https://doi.org/10.1038/nature08945.

Haynes, P.R., Pyfrom, E.S., Li, Y., Stein, C., Cuddapah, V.A., Jacobs, J.A.,
Yue, Z. and Sehgal, A. (2024). A neuron–glia lipid metabolic cycle couples daily

sleep to mitochondrial homeostasis. Nature Neuroscience, [online] 27(4), pp.666–
678. doi:https://doi.org/10.1038/s41593-023-01568-1.

Hogan, B.M., Bos, F.L., Bussmann, J., Witte, M., Chi, N.C., Duckers, H.J. and
Schulte-Merker, S. (2009). ccbe1 is required for embryonic lymphangiogenesis and
venous   sprouting.   Nature   Genetics,   41(4),   pp.396–398.
doi:https://doi.org/10.1038/ng.321.

Holth, J.K., Fritschi, S.K., Wang, C., Pedersen, N.P., Cirrito, J.R., Mahan,
T.E., Finn, M.B., Manis, M., Geerling, J.C., Fuller, P.M., Lucey, B.P. and Holtzman,
D.M. (2019). The sleep-wake cycle regulates brain interstitial fluid tau in mice and
CSF   tau   in   humans.   Science,   363(6429),   pp.880–884.
doi:https://doi.org/10.1126/science.aav2546.

Huang, Y. (2012). Effects of Age and Amyloid Deposition on Aβ Dynamics in
the Human Central Nervous System. Archives of Neurology, 69(1), p.51.
doi:https://doi.org/10.1001/archneurol.2011.235.

Huisman, Y., Uphoff, K., Berger, M., Dobrindt, U., Schelhaas, M., Zobel, T.,
Bussmann, J., Impel, A. and Schulte‑Merker, S. (2021). Meningeal lymphatic
endothelial cells fulfill scavenger endothelial cell function and cooperate with
microglia   in   waste   removal   from   the   brain.   Glia,   70(1),   pp.35–49.
doi:https://doi.org/10.1002/glia.24081.

Hurd, M.W. and Cahill, G.M. (2002). Entraining Signals Initiate Behavioral
Circadian Rhythmicity in Larval Zebrafish. Journal of Biological Rhythms, 17(4),
pp.307–314. doi:https://doi.org/10.1177/074873002129002618.

Islam, M.M., Hlushchenko, I. and Pfisterer, S.G. (2022). Low-Density
Lipoprotein Internalization, Degradation and Receptor Recycling Along Membrane
Contact   Sites.   Frontiers   in   Cell   and   Developmental   Biology,   10.
doi:https://doi.org/10.3389/fcell.2022.826379.

Kakarla, V., Letendre, S.L. and Ellis, R.J. (2023). Time of Day Influences
Concentrations of Total Protein and Albumin in Cerebrospinal Fluid in HIV.

International Journal of Molecular Sciences, 24(3), p.2832.
doi:https://doi.org/10.3390/ijms24032832.

Kalin, N.H., Cohen, R.M. and Murphy, D.L. (1980). Circadian variation in the
CSF cortisol concentration of the rhesus monkey. Life Sciences, 26(18), pp.1485–
1487. doi:https://doi.org/10.1016/0024-3205(80)90270-2.

Kaneko, M. and Cahill, G.M. (2005). Light-Dependent Development of
Circadian Gene Expression in Transgenic Zebrafish. PLoS Biology, 3(2), p.e34.
doi:https://doi.org/10.1371/journal.pbio.0030034.

Kaushik, M.K., Aritake, K., Kamauchi, S., Hayaishi, O., Huang, Z.-L., Lazarus,
M. and Urade, Y. (2014). Prostaglandin D2 is crucial for seizure suppression and
postictal sleep. Experimental Neurology, 253, pp.82–90.
doi:https://doi.org/10.1016/j.expneurol.2013.12.002.

Kelu, J.J., Pipalia, T.G. and Hughes, S.M. (2020). Circadian regulation of
muscle growth independent of locomotor activity. Proceedings of the National
Academy of Sciences, 117(49), pp.31208–31218.
doi:https://doi.org/10.1073/pnas.2012450117.

Koethe, D., Schreiber, D., Giuffrida, A., Mauss, C., Faulhaber, J.,
Heydenreich, B., Hellmich, M., Graf, R., Klosterkötter, J., Piomelli, D. and Leweke,
F.M. (2009). Sleep deprivation increases oleoylethanolamide in human cerebrospinal
fluid. Journal of Neural Transmission, 116(3), pp.301–305.
doi:https://doi.org/10.1007/s00702-008-0169-6.

Kroll, F., Donnelly, J., Özcan, G.G., Mackay, E. and Rihel, J. (2025).
Behavioural pharmacology predicts disrupted signalling pathways and candidate
therapeutics from zebrafish mutants of Alzheimer's disease risk genes. eLife, 13.
doi:https://doi.org/10.7554/elife.96839.3.

Kroll, F., Powell, G.T., Ghosh, M., Gestri, G., Antinucci, P., Hearn, T.J.,
Tunbak, H., Lim, S., Dennis, H.W., Fernandez, J.M., Whitmore, D., Dreosti, E.,
Wilson, S.W., Hoffman, E.J. and Rihel, J. (2021). A simple and effective F0 knockout

method for rapid screening of behaviour and other complex phenotypes. eLife,
[online] 10, p.e59683. doi:https://doi.org/10.7554/eLife.59683.

Labun, K., Montague, T.G., Krause, M., Torres Cleuren, Y.N., Tjeldnes, H.
and Valen, E. (2019). CHOPCHOP v3: Expanding the CRISPR Web Toolbox beyond
Genome   Editing.   Nucleic   Acids   Research,   47(W1),   pp.W171–W174.
doi:https://doi.org/10.1093/nar/gkz365.

Langenbruch, L., Wiendl, H., Groß, C. and Kovac, S. (2021). Diagnostic utility
of cerebrospinal fluid (CSF) findings in seizures and epilepsy with and without
autoimmune-associated   disease.   Seizure,   91,   pp.233–243.
doi:https://doi.org/10.1016/j.seizure.2021.06.030.

Li, J., Liu, M.-J., Du, W.-J., Peng, X.-L., Deng, H., Zi, H.-X., Shang, H.-B. and
Du, J.-L. (2025). Neural-activity-regulated and glia-mediated control of brain
lymphatic   development.   Cell,   [online]   12.
doi:https://doi.org/10.1016/j.cell.2025.04.008.

Li, R., Oteiza, A., Sørensen, K.K., McCourt, P., Olsen, R., Smedsrød, B. and
Svistounov, D. (2011). Role of liver sinusoidal endothelial cells and stabilins in
elimination of oxidized low-density lipoproteins. American Journal of Physiology-
Gastrointestinal   and   Liver   Physiology,   300(1),   pp.G71–G81.
doi:https://doi.org/10.1152/ajpgi.00215.2010.

Liu, J., Guo, Y., Zhang, C., Zeng, Y., Luo, Y. and Wang, G. (2022). Clearance
Systems in the Brain, From Structure to Function. Frontiers in Cellular Neuroscience,
15. doi:https://doi.org/10.3389/fncel.2021.729706.

Liu, Y.U., Ying, Y., Li, Y., Eyo, U.B., Chen, T., Zheng, J., Umpierre, A.D., Zhu,
J., Bosco, D.B., Dong, H. and Wu, L.-J. (2019). Neuronal network activity controls
microglial process surveillance in awake mice via norepinephrine signaling. Nature
Neuroscience, 22(11), pp.1771–1781. doi:https://doi.org/10.1038/s41593-019-0511-
3.

Livne, Z.B.-M., Alon, S., Vallone, D., Bayleyen, Y., Tovin, A., Shainer, I.,
Nisembaum, L.G., Aviram, I., Smadja-Storz, S., Fuentès, M., Falcón, J., Eisenberg,

E., Klein, D.C., Burgess, H.A., Foulkes, N.S. and Gothilf, Y. (2016). Genetically
Blocking the Zebrafish Pineal Clock Affects Circadian Behavior. PLOS Genetics,
12(11), pp.e1006445–e1006445. doi:https://doi.org/10.1371/journal.pgen.1006445.

Lucey, B.P., Gonzales, C., Das, U., Li, J., Siemers, E.R., Slemmon, J.R.,
Bateman, R.J., Huang, Y., Fox, G.B., Claassen, J.A.H.R., Slats, D., Verbeek, M.M.,
Tong, G., Soares, H., Savage, M.J., Kennedy, M., Forman, M., Sjögren, M.,
Margolin, R. and Chen, X. (2015). An integrated multi-study analysis of intra-subject
variability in cerebrospinal fluid amyloid-β concentrations collected by lumbar
puncture and indwelling lumbar catheter. Alzheimer's research & therapy, 7(1).
doi:https://doi.org/10.1186/s13195-015-0136-z.

Luo, B., Song, J., Zhang, J., Han, J., Zhou, X. and Chen, L. (2024). The
contribution of circadian clock to the biological processes. Frontiers in Molecular
Biosciences, 11. doi:https://doi.org/10.3389/fmolb.2024.1387576.

Luo, X., Li, R.-R., Li, Y.-Q., Yu, H.-P., Yu, H.-N., Jiang, W.-G. and Li, Y.-N.
(2022). Reducing VEGFB expression regulates the balance of glucose and lipid
metabolism in mice via VEGFR1. Molecular Medicine Reports, 26(3).
doi:https://doi.org/10.3892/mmr.2022.12801.

McDowell, K.P., Berthiaume, A.-A., Tieu, T., Hartmann, D.A. and Shih, A.Y.
(2020). VasoMetrics: unbiased spatiotemporal analysis of microvascular diameter in
multi-photon imaging applications. Quantitative Imaging in Medicine and Surgery,
11(3), pp.969–982. doi:https://doi.org/10.21037/qims-20-920.

Mello, C.F. and Oliveira, M.S. (2014). Commentary on Kaushik et al.:
Prostaglandin D2 is crucial for seizure suppression and postictal sleep. Novel
evidence supporting a role for prostanoid receptors in seizure control. Experimental
Neurology, 257, pp.157–161. doi:https://doi.org/10.1016/j.expneurol.2014.05.002.

Miao, A., Luo, T., Hsieh, B., Edge, C.J., Gridley, M., Wong, R.T.C.,
Constandinou, T.G., Wisden, W. and Franks, N.P. (2024). Brain clearance is
reduced during sleep and anesthesia. Nature Neuroscience, [online] (6).
doi:https://doi.org/10.1038/s41593-024-01638-y.

Mohawk, J.A., Green, C.B. and Takahashi, J.S. (2012). Central and
Peripheral Circadian Clocks in Mammals. Annual Review of Neuroscience, [online]
35(1), pp.445–462. doi:https://doi.org/10.1146/annurev-neuro-060909-153128.

Moore, H.A. and Whitmore, D. (2014). Circadian Rhythmicity and Light
Sensitivity of the Zebrafish Brain. PLoS ONE, 9(1), p.e86176.
doi:https://doi.org/10.1371/journal.pone.0086176.

Morganti-Kossman, M.C., Lenzlinger, P.M., Hans, V., Stahel, P., Csuka, E.,
Ammann, E., Stocker, R., Trentz, O. and Kossmann, T. (1997). Production of
cytokines following brain injury: beneficial and deleterious for the damaged tissue.
Molecular Psychiatry, 2(2), pp.133–136. doi:https://doi.org/10.1038/sj.mp.4000227.

Neumann, H., Kotter, M.R. and Franklin, R.J.M. (2008). Debris clearance by
microglia: an essential link between degeneration and regeneration. Brain, 132(2),
pp.288–295. doi:https://doi.org/10.1093/brain/awn109.

Ning, F.C., Jensen, N., Mi, J., Lindström, W., Balan, M., Muhl, L., Eriksson,
U., Nilsson, I. and Nyqvist, D. (2020). VEGF-B ablation in pancreatic β-cells
upregulates insulin expression without affecting glucose homeostasis or islet lipid
uptake. Scientific Reports, 10(1). doi:https://doi.org/10.1038/s41598-020-57599-2.

Parhizkar, S., Gent, G., Chen, Y., Rensing, N., Gratuze, M., Strout, G.W.,
Sviben, S., Tycksen, E., Zhang, Q., Gilmore, P., Sprung, R.W., Malone, J., Chen,
W., Serrano, J.R., Bao, X., Lee, C., Wang, C., Landsness, E.C., Fitzpatrick, J.A.J.
and Wong, M. (2023). Sleep deprivation exacerbates microglial reactivity and Aβ
deposition in a TREM2 -dependent manner in mice. Science Translational Medicine,
15(693). doi:https://doi.org/10.1126/scitranslmed.ade6285.

Peltola, J., Hurme, M., Miettinen, A. and Keränen, T. (1998). Elevated levels
of interleukin-6 may occur in cerebrospinal fluid from patients with recent epileptic
seizures. Epilepsy Research, 31(2), pp.129–133. doi:https://doi.org/10.1016/s0920-
1211(98)00024-2.

Peltola, J., Palmio, J., Korhonen, L., Suhonen, J., Miettinen, A., Hurme, M.,
Lindholm, D. and Keränen, T. (2000). Interleukin-6 and Interleukin-1 receptor

antagonist in cerebrospinal fluid from patients with recent tonic–clonic seizures. Epilepsy Research, 41(3), pp.205–211. doi:https://doi.org/10.1016/s0920-1211(00)00140-6.

Poceta, J.S., Parsons, L., Engelland, S. and Kripke, D.F. (2009). Circadian rhythm of CSF monoamines and hypocretin-1 in restless legs syndrome and Parkinson's disease. Sleep Medicine, 10(1), pp.129–133. doi:https://doi.org/10.1016/j.sleep.2007.11.002.

Prasad, R., Kapoor, R., Srivastava, R., Mishra, O.P. and Singh, T.B. (2014). Cerebrospinal Fluid TNF-α, IL-6, and IL-8 in Children With Bacterial Meningitis. Pediatric Neurology, 50(1), pp.60–65. doi:https://doi.org/10.1016/j.pediatrneurol.2013.08.016.

Prober, D.A., Rihel, J., Onah, A.A., Sung, R.-J. . and Schier, A.F. (2006). Hypocretin/Orexin Overexpression Induces An Insomnia-Like Phenotype in Zebrafish. Journal of Neuroscience, 26(51), pp.13400–13410. doi:https://doi.org/10.1523/jneurosci.4332-06.2006.

Reichert, S., Pavón Arocas, O. and Rihel, J. (2019). The Neuropeptide Galanin Is Required for Homeostatic Rebound Sleep following Increased Neuronal Activity. Neuron, [online] 104(2), pp.370-384.e5. doi:https://doi.org/10.1016/j.neuron.2019.08.010.

Rohleder, N., Aringer, M. and Boentert, M. (2012). Role of interleukin-6 in stress, sleep, and fatigue. Annals of the New York Academy of Sciences, 1261(1), pp.88–96. doi:https://doi.org/10.1111/j.1749-6632.2012.06634.x.

Rorsman, H.O., Müller, M.A., Liu, P.Z., Sanchez, L.G., Kempf, A., Gerbig, S., Spengler, B. and Gero Miesenböck (2025). Sleep pressure accumulates in a voltage-gated lipid peroxidation memory. Nature, [online] 641(8061). doi:https://doi.org/10.1038/s41586-025-08734-4.

Saito, K., Hattori, K., Hidese, S., Sasayama, D., Miyakawa, T., Matsumura, R., Tatsumi, M., Yokota, Y., Ota, M., Hori, H. and Kunugi, H. (2021). Profiling of

Cerebrospinal Fluid Lipids and Their Relationship with Plasma Lipids in Healthy Humans. Metabolites, 11(5), p.268. doi:https://doi.org/10.3390/metabo11050268.

Shew, T., Wolins, N.E. and Cifarelli, V. (2018). VEGFR-3 Signaling Regulates Triglyceride Retention and Absorption in the Intestine. Frontiers in Physiology, 9. doi:https://doi.org/10.3389/fphys.2018.01783.

Shibata-Germanos, S., Goodman, J.R., Grieg, A., Trivedi, C.A., Benson, B.C., Foti, S.C., Faro, A., Castellan, R.F.P., Correra, R.M., Barber, M., Ruhrberg, C., Weller, R.O., Lashley, T., Iliff, J.J., Hawkins, T.A. and Rihel, J. (2020). Structural and functional conservation of non-lumenized lymphatic endothelial cells in the mammalian leptomeninges. Acta Neuropathologica, [online] 139(2), pp.383–401. doi:https://doi.org/10.1007/s00401-019-02091-z.

Siret, C., Lessen, M. van, Bavais, J., Jeong, H.W., Reddy, K., Kapupara, K., Wang, S., Simic, M., de Fabritus, L., Tchoghandjian, A., Fallet, M., Huang, H., Sarrazin, S., Sieweke, M.H., Stumm, R., Sorokin, L., Adams, R.H., Schulte-Merker, S., Kiefer, F. and van de Pavert, S.A. (2022). Deciphering the heterogeneity of the Lyve1+ perivascular macrophages in the mouse brain. Nature Communications, 13(1), pp.7366–7366. doi:https://doi.org/10.1038/s41467-022-35166-9.

Sturge, J., Todd, S.K., Kogianni, G., McCarthy, A. and Isacke, C.M. (2007). Mannose receptor regulation of macrophage cell migration. Journal of Leukocyte Biology, 82(3), pp.585–593. doi:https://doi.org/10.1189/jlb.0107053.

Suppermpool, A., Lyons, D.G., Broom, E. and Rihel, J. (2024). Sleep pressure modulates single-neuron synapse number in zebrafish. Nature, [online] 629(8012), pp.1–7. doi:https://doi.org/10.1038/s41586-024-07367-3.

Süße, M., Saathoff, N., Hannich, M. and von Podewils, F. (2019). Cerebrospinal fluid changes following epileptic seizures unrelated to inflammation. European Journal of Neurology, 26(7), pp.1006–1012. doi:https://doi.org/10.1111/ene.13924.

Tang, M., Song, C., Zhang, Y., Xu, X., Wang, C., Zhang, Z. and Chen, T. (2021). Levels of pretreatment blood lipids are prognostic factors in advanced

NSCLC patients treated with anlotinib. Lipids in Health and Disease, 20(1), pp.165–
165. doi:https://doi.org/10.1186/s12944-021-01596-5.

Tumani, H., Jobs, C., Brettschneider, J., Hoppner, A.C., Kerling, F. and
Fauser, S. (2015). Effect of epileptic seizures on the cerebrospinal fluid – A
systematic retrospective analysis. Epilepsy Research, 114, pp.23–31.
doi:https://doi.org/10.1016/j.eplepsyres.2015.04.004.

Ungurean, G., Behroozi, M., Böger, L., Helluy, X., Libourel, P.-A., Güntürkün,
O. and Rattenborg, N.C. (2023). Wide-spread brain activation and reduced CSF flow
during avian REM sleep. Nature Communications, [online] 14(1), p.3259.
doi:https://doi.org/10.1038/s41467-023-38669-1.

van der Zande, H.J.P., Nitsche, D., Schlautmann, L., Guigas, B. and Burgdorf,
S. (2021). The Mannose Receptor: From Endocytic Receptor and Biomarker to
Regulator of (Meta)Inflammation. Frontiers in Immunology, 12.
doi:https://doi.org/10.3389/fimmu.2021.765034.

van Impel, A., Zhao, Z., Hermkens, D., Roukens, M.G., Fischer, J.C.,
Peterson-Maduro, J., Duckers, H.J., Ober, E.A., Ingham, P.W. and Schulte-Merker,
S. (2014). Divergence of zebrafish and mouse lymphatic cell fate specification
pathways. Development, 141(6), pp.1228–1238.
doi:https://doi.org/10.1242/dev.105031.

van Lessen, M., Shibata-Germanos, S., van Impel, A., Hawkins, T.A., Rihel, J.
and Schulte-Merker, S. (2017). Intracellular uptake of macromolecules by brain
lymphatic endothelial cells during zebrafish embryonic development. eLife, [online] 6,
p.e25932. doi:https://doi.org/10.7554/eLife.25932.

Vgontzas, A.N., Papanicolaou, D.A., Bixler, E.O., Kales, A., Tyson, K. and
Chrousos, G.P. (1997). Elevation of Plasma Cytokines in Disorders of Excessive
Daytime Sleepiness: Role of Sleep Disturbance and Obesity. The Journal of Clinical
Endocrinology & Metabolism, 82(5), pp.1313–1316.
doi:https://doi.org/10.1210/jcem.82.5.3950.

Vgontzas, A.N., Papanicolaou, D.A., Bixler, E.O., Lotsikas, A., Zachman, K.,
Kales, A., Prolo, P., Wong, M.-L., Licinio, J., Gold, P.W., Hermida, R.C., Mastorakos,
G. and Chrousos, G.P. (1999). Circadian Interleukin-6 Secretion and Quantity and
Depth of Sleep. The Journal of Clinical Endocrinology & Metabolism, [online] 84(8),
pp.2603–2607. doi:https://doi.org/10.1210/jcem.84.8.5894.

Voelz, K., Gratacap, R.L. and Wheeler, R.T. (2015). A zebrafish larval model
reveals early tissue-specific innate immune responses to Mucor circinelloides.
Disease   Models   &   Mechanisms,   8(11),   pp.1375–1388.
doi:https://doi.org/10.1242/dmm.019992.

Wadhwa, M., Kumari, P., Chauhan, G., Roy, K., Alam, S., Kishore, K., Ray, K.
and Panjwani, U. (2017). Sleep deprivation induces spatial memory impairment by
altered hippocampus neuroinflammatory responses and glial cells activation in rats.
Journal   of   Neuroimmunology,   312,   pp.38–48.
doi:https://doi.org/10.1016/j.jneuroim.2017.09.003.

Wang, H., Yang, Z., Li, X., Huang, D., Yu, S., He, J., Li, Y. and Yan, J. (2020).
Single-cell in vivo imaging of cellular circadian oscillators in zebrafish. PLOS Biology,
18(3), p.e3000435. doi:https://doi.org/10.1371/journal.pbio.3000435.

Wary, K.K., Thakker, G.D., Humtsoe, J.O. and Yang, J. (2003). Analysis of
VEGF-responsive Genes Involved in the activation of endothelial cells. Molecular
Cancer, 2(1), p.25. doi:https://doi.org/10.1186/1476-4598-2-25.

Whitmore, D., Foulkes, N.S. and Sassone-Corsi, P. (2000). Light acts directly
on organs and cells in culture to set the vertebrate circadian clock. Nature,
404(6773), pp.87–91. doi:https://doi.org/10.1038/35003589.

Xie, L., Kang, H., Xu, Q., Chen, M.J., Liao, Y., Thiyagarajan, M., O'Donnell, J.,
Christensen, D.J., Nicholson, C., Iliff, J.J., Takano, T., Deane, R. and Nedergaard,
M. (2013). Sleep Drives Metabolite Clearance from the Adult Brain. Science,
342(6156), pp.373–377. doi:https://doi.org/10.1126/science.1241224.

Zareba, J., Cattaneo, E.F., Villani, A., Othman, A., Streb, S. and Peri, F.
(2024). NPC1 links cholesterol trafficking to microglial morphology via the

gastrosome. Nature Communications, 15(1). doi:https://doi.org/10.1038/s41467-024-52874-6.

Zhdanova, I.V., Wang, S.Y., Leclair, O.U. and Danilova, N.P. (2001). Melatonin promotes sleep-like state in zebrafish. Brain Research, 903(1-2), pp.263–268. doi:https://doi.org/10.1016/s0006-8993(01)02444-1.

Figures:

Figure 1: muLEC morphology is dynamic over the day-night cycle. **a**) Representative maximum projection of a muLEC tectal loop (green) in 6dpf Tg*(flt4:mCitrine)* larvae during the day (ZT1, left) versus night (ZT19, right). Connected (arrow) and disconnected (arrowhead) muLECs are highlighted. **b**) Quantification of total muLEC volume within a tectal loop during the day (n=16) and night (n=16). **c**) Quantification of muLEC disconnectedness in the day (n=16) and night (n=16). **d**) High-resolution muLEC imaging reveals fluorescent negative inclusions (blue arrows) and filopodia-like protrusions (pink arrows). **e**) Average number of muLEC filopodial protrusions per cell per larva in the day (n=7) and night (n=7). (**f & g**) Average size (f) and number (g) of intracellular muLEC inclusions, day (n=7) and night (n=7). Data are mean ± SEM. Each dot represents one larva. Statistical differences were determined by one-way ANOVA (two-tailed, b-g**)**. ns $p > 0.05$; * $p < 0.05$; ** $p < 0.01$. Abbreviations: ZT, Zeitgeber Time. Scale bar = 100µm (a), 20µm (b).

Figure 2: muLEC-uptake of macromolecules fluctuates over the day-night cycle.
a) *Tg(flt4:mCitrine)* larva were intraventricularly injected with dyes and imaged for several hours to track muLEC macromolecular uptake. **b**) Time-course of pHrodo-avidin accumulation in muLECs, sampled every 4 minutes. **c**) Developmentally matched larvae on reverse LD cycles were injected and imaged in the middle of the day (ZT7) or night (ZT19). **d**) Representative maximum projection from a timeseries of a *Tg(flt4:mCitrine)* larva injected with 2MDa Alexa-fluor 488 dextran (Dex; blue) and pHrodo Red avidin (pHr-av; red). Signal was extracted from muLEC and ventricle masks removing any brain derived signal. **e**) The average integrated intensity ± SEM (60 min bins) of pHrodo-avidin uptake by muLECs is significantly higher in the day (ZT7-12, n=20) than night (ZT19-24, n=17). **f**) Slope of muLEC-uptake of pHrodo-avidin from a simple linear regression. **g**) Larva as in d) but injected with pHrodo Red LDL (pHr-LDL; magenta). **h, i**) muLEC uptake of pHrodo-LDL is higher during the day (ZT7-13, n=19) than at night (ZT19-1, n=18); the slope is from a one phase association regression. **j**) Larva as in d) but injected with pHrodo Red dextran (10kDa) (pHr-dex; cyan). **k**, **l**) muLEC-uptake of pHrodo-dextran is higher during the day (ZT7-13, n=23) than at night (ZT19-1, n=21). Data are mean ± SEM; each dot represents one larva (**f, i, l**). Statistical differences were determined by Mixed-Effects Model (**e, h, k**) or one-way ANOVA (two-tailed, **f, i, l**). ^{ns} $p > 0.05$;* $p < 0.05$; ** $p < 0.01$. Scale bar = 100μm.

Figure 3: A muLEC-intrinsic circadian clock regulates the diurnal variation of pHrodo-avidin uptake. a) Candidate physiological drivers of uptake rates covary across the day-night cycle. **b**) muLEC-uptake rates were measured under free-running constant light conditions following injection (black arrow) during the subjective day (CT7) or night (CT19). **c**) The slope of muLEC uptake of pHrodo-avidin is higher during the subjective day (CT7-13, n=13) than subjective night (CT19-24, n=16). **d**) Experimental design to test the effect of melatonin. **e**) muLEC uptake of pHrodo-avidin is not significantly different between melatonin (n=18) treated and DMSO controls (n=19). **f**) Experimental design to test the effect of recovery sleep after sleep deprivation. **g**) muLEC uptake of pHrodo-avidin is not significantly affected by recovery sleep (n=14) vs controls (n=12). **h**) Experimental design to test the effect of caffeine. **i**) muLEC uptake of pHrodo-avidin is not affected by caffeine (n=15) vs controls (n=17). **j**) Experimental design to test the effect of "gentle handling" sleep deprivation. **k**) muLEC uptake of pHrodo-avidin during 4 hours of sleep deprivation (n=16) is similar to undisturbed controls (n=18). **l**) Timeseries maximum projection of a *Tg(flt4:Gal4); Tg(UAS:GFPp2AdnCLK)* larva injected with 2MDa Alexa-fluor, 488 dextran (Dex; blue) and pHrodo Red avidin (pHr-av; red). Signal was extracted from muLEC and ventricle masks removing any brain derived signal. **m,n)** The average integrated intensity ± SEM (m) and slope of muLEC-uptake of pHrodo-avidin by simple linear regression (n) are not significantly different between the day (n=35) and night (n=27). Data are mean ± SEM and each dot represents one larva (**c,e,g,i,k,n**). Statistical differences were determined by: one-way ANOVA (two-tailed, **c,e,g,j**). ^{ns} p>0.05;** p<0.01. Scale bar = 100µm.

Figure 4: muLEC-intrinsic circadian regulation of Mrc1a protein regulates pHrodo-avidin uptake. a) The experimental paradigm. *Tg(flt4:mCitrine)* larvae were injected at the one-cell stage with Cas9 and *mrc1a*-targeting or scrambled control gRNAs, then injected with dyes at 5dpf. **b**) Representative maximum projection from a timeseries of a scrambled control larva (left) and *mrc1a* KO (right) with muLECs (green) post-intraventricular injection of 2MDa Alexa-fluor, 488 dextran (Dex; blue) and pHrodo Red avidin (pHr-av; red). White arrowheads indicate pHr-avidin uptake by brain-resident phagocytes. **c**) The average integrated intensity ± SEM of pHrodo-avidin accumulation in *mrc1a* KO larvae (n=8) and scrambled control siblings (n=10). **d**) The slope of muLEC-uptake of pHrodo-avidin. **e**) Representative maximum projection of muLECs (green) and anti-Mrc1a (magenta) immunofluorescence in 5dpf *Tg(flt4:mCitrine)* larvae. **f**) Integrated intensity of anti-Mrc1a immunofluorescence in muLECs is higher during the day (ZT7, n=19) than night (ZT19, n=17). **g**) Integrated intensity of anti-Mrc1a immunofluorescence in muLECs is similar in the day (n=22) and night (n=26) in muLEC-clock disrupted larvae. Data are mean ± SEM. Each dot represents one larva (c,d,f,g). Statistical differences were determined by a Mixed-Effects Model (c) or Mann-Whitney U tests (two-tailed, d,f,g).). ^{ns} p>0.05; ** p<0.01; ****p<0.0001. Scale bar = 100μm.

Circadian effects on uptake of pHr-dex and pHr-LDL

State effects on uptake of pHr-dex and pHr-LDL

State effects on muLEC inclusions

Figure 5: Diurnal variation in pHrodo-dex and pHrodo-LDL uptake are independent of the intrinsic clock. a) The experimental paradigm to measure day (ZT7) vs. night (ZT19) muLEC uptake rates in *Tg(flt4:Gal4); Tg(UAS:GFPp2AdnCLK)* larvae. **b,c**) Slopes of muLEC-uptake of pHrodo-dex (b) pHrodo-LDL (c) from a one-phase association regression show that uptake differences between day (pHr-dex, n=22; pHr-LDL, n=33) versus night (pHr-dex, n=22; pHr-LDL, n=29) persist despite transgenic muLEC-clock disruption. **d**) Experimental design to test the effect of extended waking on muLEC uptake of pHrodo-dex and pHrodo-LDL. **e**) The integrated intensity of pHrodo-dex accumulation is similar during 4 hours of sleep deprivation (n=24) vs controls (n=26). **f**) The integrated intensity of pHrodo-LDL accumulation is significantly higher after sleep deprivation (n=24) vs controls (n=27). **g**) A muLEC from a *Tg(flt4:mCitrine)* larva following 4 hours of sleep deprivation (right) or undisturbed control (left). **h**) Average size of intracellular muLEC inclusions after sleep deprivation (n=10) vs controls (n=10). **i**) Average size of muLEC intracellular inclusions at ZT23 after 4 h of Caffeine exposure (250µM, n=13) or water control (n=13). **j**) Average size of muLEC intracellular inclusions at ZT5 after 4 h of Melatonin exposure (1µM, n=13) or DMSO control (n=13). Data are mean ± SEM; each dot represents one larva. Statistical differences were determined by Mann-Whitney U tests (two-tailed, c,f,h,j) or one-way ANOVA (two-tailed, b,e,i). ^{ns} p>0.05;* p<0.05; ** p<0.01; p<0.001. Scale bar:10µm.

Figure 6. muLECs facilitate behavioural recovery following drug-induced seizures. a) Maximum projections of confocal stacks pre- and post-mock (left) or pre- and post-targeted laser ablation (right) of muLECs (green) in a 5dpf *Tg(flt4:mCitrine);Tg(kdrl:mCherry)* larva. Blood vessels *(*magenta) are unaffected. **b**) Multi-day activity traces showing the recovery of baseline behavior following PTZ-induced seizures (1hr ending at time 0). A representative larva from mock- (left, grey) and muLEC-ablated (right, red) conditions are shown. Recovery is determined by the time the activity levels reach the 50th percentile of each larva's post-recovery baseline activity level (dashed navy line). **c**) The fraction of fish that have recovered by a given time post seizure. Mock ablated fish recover faster than muLEC-ablated fish (log rank test, $p < 0.05$). **d, e**) Average daytime (**d**) and nighttime (**e**) activity of ablated (n=21) and mock ablated (n=23) fish is similar after recovery. Statistical differences were determined by Mann-Whitney U tests (two-tailed, e); one-way ANOVA (two-tailed, d). ns $p > 0.05$; *** $p < 0.001$. Scale bar:100µm.

Figure S1: Additional morphological characteristics of muLECs over the day night cycle and analysis methodology. a) Surface area of the muLEC loop day (n=16) vs night (n=16). **b)** Length of muLEC filopodial protrusions in the day (n=7) and night (n=7). **c,d)** The average width (c) and length (d) of connections between muLECs between the day (n=13) and night (n=14). **e)** Confocal imaging of a single muLEC in a *Tg(flt4:mCitrine)* larva (grayscale) was processed and segmented in 2D to identify fluorescent-negative spaces as intracellular inclusions. **f)** Morphometric analysis of the muLEC tectal loop (grayscale) reveals connections (yellow) between muLECs. **g)** Distribution of inclusion size as a cumulative percentage frequency of total recorded inclusions in all cells in all larvae during the day versus night. **a-d**, data are mean ± SEM; each dot represents one larva. Statistical differences were determined by one-way ANOVA (two-tailed, **a-d**). ns p>0.05; *** p<0.001. Scale bar: 5µm (e); 20µm (f).

Figure S2: muLEC-uptake curve modelling and experimental controls. a) The rate of muLEC-uptake of pHrodo-avidin is well captured by a simple linear regression when data are sampled at low frequency (every 60 min), with the slope from the linear fit similar to 4 minute sampling (grey continuous line; data from Figure 2b) irrespective of early, middle or late starting points (T_0= 30, 60, 90 mins post-injection, respectively). **b)** The average integrated intensity ± SEM of pHrodo-avidin uptake by muLECs is similar between the early day (ZT1-6, n=15) and early night (ZT14-19, n=11). **c)** Slope of muLEC-uptake of pHrodo-avidin from a simple linear regression. **d,e)** Average integrated intensity at T_0 of 2MDa dextran across both telencephalic and diencephalic ventricles is the same across groups presented in b,c (d) or Figure 2e,f (e). **f,g)** Time-point T_{final} is the same between groups presented in b,c (f) or Figure 2e,f (g). **h)** Representative larva sampled every 10 minutes post-injection of pHrodo-LDL (grey line). One-phase association regression captures slopes when simulated at low frequency. **i,j)** There is no difference in injection bolus (**i**) or time-sampling (**j**) from the data presented in Figure 2h,i. **k)** Representative larva sampled every 8 minutes post-injection of pHrodo-dextran (grey line). One-phase association regression captures slopes when simulated at low frequency. **l,m)** There is no difference in injection bolus (l) or time-sampling (m) from the data presented in Figure 2k,l. Data are mean ± SEM and each dot represents one larva. Statistical differences were determined by one-way ANOVA (two-tailed, **b-e**, **g-k**). [ns] $p>0.05$.

Figure S3: Diurnal variation of muLEC-uptake of pHrodo-avidin is regulated by the clock and not brain state. a) Experimental design to test the effect of drug-induced rebound sleep. **b**) The slope of pHrodo-avidin uptake was not significantly different between PTZ treated (n=17) and controls (n=12). **c**) *Tg(flt4:mCitrine)* embryos were either (top) entrained on a normal light-dark cycle to 4dpf before switching to free-running conditions (as in Figure 2b) or (bottom) raised under constant light conditions to suppress circadian rhythm development, referred to as 'clock-break'. **d**) Normalised slope of muLEC-uptake of pHrodo-avidin from a simple linear regression revealed that entrained larvae show higher levels of muLEC-uptake relative to clock-broken fish (purple) when sampled in the middle of their subjective day (CT7, yellow) and not in the middle of their subjective night (CT19, grey). As per x-axis: Clock break, n=12; CT7, n=14; Clock break, n=12; CT19, n=10. Data are mean ± SEM; each dot represents one larva. Statistical differences were determined by one-way ANOVA (two-tailed). ^{ns} p>0.05; * p<0.05.

Figure S4: Injection dose and final sampling times for Figure 3 and Figure S3. Average integrated intensity of 2MDA dextran signal in both the telencephalic and diencephalic ventricles at T_0 is the same between groups corresponding to experiments presented in: **a**, Figure 3c; **c**, Figure 3e; **e**, Figure 3g; **g**, Figure S3b; **i**, Figure 3i; **k**, Figure 3m,n; **m,o**, Figure S3d. Final sampled time-points post-injection demonstrating no sampling bias between groups corresponding to experiments presented in: **b**, Figure 3c; **d**, Figure 3e; **f**, Figure 3g; **h**, Figure S3b; **j**, Figure 3i; **l**, Figure 3m,n; **n,p**, Figure S3d. All data are mean ± SEM; each dot represents one larva. Statistical differences were determined by Mann-Whitney U tests (two-tailed, a,b,d,f,j); one-way ANOVA (two-tailed, c,e,g,h,l,k-p). [ns] $p > 0.05$.

a

locus 3   locus 1

mrc1a

exon 6   exon 3   exon 1

locus 2

b

RICIN domain
37-108

FN2 domain
165-213

CLECT domain
235-343; 361-485
503-625; 656-771
794-914; 944-1072
1098-1205; 1221-1347

0   485   1221 1440

c

mrc1a   TGAGGGTCAACGCTTTCGATGGG
           58939709   58939687
guide 1   TGAGGGTCAACGCTTTCGAT**GGG**

mrc1a   ACTACGGAGGGAAGATCAGATGG
           58939004   58938982
guide 2   ACTACGGAGGGAAGATCAGA**TGG**

mrc1a   TGGGACAGTGATCCAGTGACTGG
           58936563   58936585
guide 3   TGGGACAGTGATCCAGTGAC**TGG**

d

mrc1b   TGAGGGTCACGAGCCACTGGCATTGTCATTT - - - TGGG
           31536497   31536462
guide 1   TGAGGGTCA- - - - - - -AC - -GC - - - - - - -TTTCGAT**GGG**

mrc1b   ACTGCGTGTTAATGAGAGGGGAGGTAAGACATTATTTGG
           31533766   31533728
guide 2   ACTACG - - - - - - - - - - -GAGGGAAGATCAGA - - - - - -**TGG**

mrc1b   TGGTACTGTCAGATCCACAAAGGTAAATGG
           31527075   31527046
guide 3   TGGGACAGT - -GATCCA - - - - -GTGAC**TGG**

e

Percentage reads (%)

mutated   frameshifted

KO1 KO2 KO3 KO4 KO5 scr1 scr2 scr3
locus 1

KO1 KO2 KO3 KO4 KO5 scr1 scr2 scr3
locus 2

KO1 KO2 KO3 KO4 KO5 scr1 scr2 scr3
locus 3

f

muLEC   anti-Mrc1a   merge

scrambled

mrc1a KO

Figure S5: Deep sequencing of *mrc1a* F$_0$ knockouts and validation of the anti-Mrc1a antibody. a) Schematic of the *mrc1a* gene in the 5'-3' genomic direction. Exons are displayed in dark grey; short exons are protein coding; tall exons are 5' or 3' UTRs. Orange lines show the 3 target loci in exons 2, 3 and 4 for CRISPR-mediated mutagenesis of *mrc1a*. Exons and introns are not to scale. **b**) Schematic of the Mrc1a protein with key conserved protein domains and their respective amino acid positions. Orange arrows indicate the predicted locations of truncations generated from the 3 target loci. **c**, **d**) Genomic sequences of *mrc1a* (c) and *mrc1b* (d) showing that the selected gRNAs targeted *mrc1a* and not *mrc1b*. Genomic bases complimentary to the guide have a grey background. The protospacer adjacent motif (PAM) where Cas9 binds is underlined in orange. **e**) Stacked plot of the percentage of total reads mutated (grey) and with a frameshift mutation (orange) as computed by ampliCan analysis following MiSeq deep sequencing of each targeted locus of *mrc1a* (white label, knockout, n=5) or in scrambled controls (grey label, scr, n=3). Each number represents the same larva sequenced at multiple loci. **f**) Dorsal maximum projection of anti-Mrc1a immunohistochemistry (magenta) and endogenous fluorescence of muLECs (green) in 5dpf *Tg(flt4:mCitrine)* larvae. Mrc1a immunofluorescence is present in scrambled controls (top) but lost in *mrc1a* KO larvae (bottom). Abbreviations: RICIN, ricin B, lectin domain; FN2, fibronectin type II; CLECT, C-type Lectin; Scale bar, 100μm.

Figure S6: *mrc1a* knockout affects muLEC morphology but not the surrounding vasculature or *mpeg+* macrophages. *mrc1a* KO or control larvae were generated in 3 transgenic lines that label **a**) muLECs (Tg(*flt4:mCitrine*)) in green; **d**) muLECs and surrounding vasculature in red (*Tg*(*flt4:mCitrine*);*Tg*(*kdrl:mCherry*)); and **g**) muLECs and *mpeg+* cells in magenta (*Tg*(*flt4:mCitrine*);*Tg(mpeg:mCherry)*). Representative scrambled-control larva (left) and *mrc1a* F_0 KO larva (right) are shown. **b,c**) muLEC morphology is quantified as volume of mCitrine$^+$ mass in the bilateral loops over the optic tectum (**b**) or populating the inner tectal rings (**c**) reveals altered muLEC morphology in *mrc1a* F_0 KOs (green, n=11) compared to scrambled-injected controls (grey, n=27). **e,f**) Vascular morphology is quantified in *mrc1a* KOs (red, n=13) or controls (grey, n=10) by (**e**) the cumulative length of the *kdrl*$^+$ vessels in diencephalic and telencephalic regions 120μm from the skin surface and (**f**) the total number of branching points. **h-k**) There were no differences between *mrc1a* KOs (magenta, n=8) and controls (grey, n=10) in total mpeg$^+$ cell volume in the head cavity (**h**), or the number (**i**), volume (**j**) or branches (**k**) of individual *mpeg*$^+$ cells. Data are mean ± SEM; each dot represents one larva. Statistical differences were determined by one-way ANOVA (two-tailed). ^{ns} p>0.05; ** p<0.01; *** p<0.001. Scale bar:100μm.

Figure S7: supplemental data relating to Figure 5. a, **d)** The average integrated intensity ± SEM of pHrodo-dex (a) or pHrodo-LDL (d) accumulated in muLECs in *Tg(flt4:Gal4);Tg(UAS:GFP2AdnCLK)* larvae day (ZT7) vs night (ZT19)*.* Data corresponding to Figure 5b (a) and Figure 5c (d). **b,e)** Average integrated intensity of 2MDa dextran signal in both telencephalic and diencephalic ventricles at T_0 is not different between groups, corresponding to experiments presented in Figure 5b (b); Figure 5c (e). **c,f)** Time T_{final} is similar between groups corresponding to experiments presented in: Figure 5b (c); Figure 5c (f).**h,l,p)** Average number of intracellular muLEC inclusions per larvae following: h, sleep deprivation (n=10) or undisturbed sleep (n=10); l, Caffeine-treatment (n=13) or fish water control (n=13); p, Melatonin-treatment (n=13) or DMSO controls (n=13). **i,j,m,n,q,r)** Average number and length of filopodia per cell per larva following: i,j, sleep-deprivation; m,n, Caffeine-treatment; q,r, Melatonin-treatment. **k,o,s)** Distribution of inclusion size as a cumulative percentage frequency of all recorded inclusions in all cells in all larvae following: k, sleep-deprivation; o, Caffeine-treatment; s, Melatonin-treatment. Data are mean ± SEM; each dot represents one larva (b,c,e,f,h-j,l-n,p-r). Statistical differences were determined by Mann-Whitney U tests (two-tailed, b,c,p,r), or one-way ANOVA (two-tailed, e,f,h,i,j,m,n,q). [ns] p>0.05.

Figure S8: Behavioural recovery post treatment and muLEC regeneration post-ablation. a) The delay in the recovery time (i.e., when 50% of larvae have recovered) in muLEC-ablated fish is observed across a wide range of recovery cutoff values (Activity Percentiles). The values derived from the curves in Figure 6c are noted by the vertical dashed line. **b**) Representative dorsal maximum projections of a mock-ablated (left) and muLEC-ablated (right) larva 3 days post intervention (dpi). muLECs (green) and blood vessels (magenta) in a (*Tg(flt4:mCitrine;Tg(kdrl:mCherry larva)*. **c**) Regenerative capacity of muLECs post-ablation as quantified by the volume of mCitrine[+] cells in bi-lateral tectal loops structures 3 days post- muLEC (n=10) or mock ablation (n=9). Data are mean ± SEM; each dot represents one larva. Statistical differences were determined by Mann-Whitney U tests (two-tailed, c). [ns] p>0.05; *** p<0.001.

Supplemental Video 1: Timelapse of a 5dpf *Tg(flt4:mCitrine)* larva injected with 2MDa Alexa-fluor, 488 dextran (Dex; blue) and pHrodo Red avidin (pHr-av; red) and imaged at high-time resolution as depicted in Figure 2a, S2a. Signal was extracted from muLEC and ventricle masks removing any brain derived signal. First frame captured 5 minutes post-injection and every subsequent frame +4 minutes for a total imaging window of 330 minutes. Scale bar, 100μm.