Title: Oestradiol and Progesterone Variability in Adolescent Females: Associations with Brain Structure and Mental Health Problems

Short title - Hormone Variability, Brain Structure and Mental Health Risk

Author details: Muskan Khetan, M.Tech[a,b], Ye Ella Tian, PhD[c], Nandita Vijayakumar, PhD[d,e,f], Megan M. Herting, PhD[e], Michele O' Connell, MD[d,g,h], Marc L Seal, PhD[d,g], Sarah Whittle, PhD[a,f,i]

[a]Centre for Youth Mental Health, The University of Melbourne, Parkville, VIC, Australia
[b]Orygen, The National Centre of Excellence in Youth Mental Health, Parkville, VIC, Australia
[c]Department of Psychiatry, University of Melbourne, Parkville, VIC, Australia
[d]Murdoch Children's Research Institute, Parkville VIC, Australia
[e]Department of Population and Public Health Sciences, University of Southern California, Los Angeles, California, USA
[f]School of Psychology, Deakin University, Burwood, VIC, Australia
[g]Department of Paediatrics, The University of Melbourne, Parkville, VIC, Australia
[h]Department of Endocrinology & Diabetes, The Royal Children's Hospital, Parkville, VIC, Australia
[i]Melbourne School of Psychological Sciences, The University of Melbourne, Parkville, VIC, Australia

Correspondence to: Muskan Khetan, M.Tech, Melbourne, Australia; email: m.khetan@unimelb.edu.au

ABSTRACT

Objective

Adolescent females are at increased risk for adverse mental health problems, and puberty is thought to be a critical factor. Pubertal onset involves substantial variability in oestradiol (E2) and progesterone (P4), yet the impact of these hormone dynamics remains unclear. We investigated whether within-individual E2 and P4 variability across one month relates to brain structure and mental health problems in adolescent females.

Methods

Participants were 147 females (11–17 years) from the community-based, cross-sectional Puberty and NeuroDevelopment in Adolescents (PANDA) study in Australia. Salivary E2 and P4 were collected weekly for one month, with variability indexed as within-subject standard deviation. Structural MRI was acquired on the final sampling day. Mental health problems were assessed via self- and parent-report. Linear regression and mediation analyses tested associations between hormone variability, brain structure, and mental health, adjusting for demographics.

Results

Greater P4 variability was associated with smaller left thalamus volume (Cohen's d = -0.26, pFDR = 0.031). Higher E2 variability was associated with fewer anxiety (d = -0.170, pFDR = 0.043) and depressive symptoms (d = -0.179, pFDR = 0.036), lower emotion dysregulation (d = -0.179, pFDR = 0.036), and smaller right hippocampus volume in pre-menarchal adolescents (d = -0.192, p = 0.02). Weekly reductions in E2 predicted lower positive affect (d = 0.084, p = 0.048).

Conclusion

These findings indicate that hormone variability may meaningfully shape brain and behavioural development in adolescent girls differently from hormone-related transitions in adulthood. Larger, replicable, and longer-term studies are needed to confirm these patterns.

Keywords: Female Adolescents, hormones, brain structure, mental illness

1. <u>Introduction</u>

Adolescent females have a higher vulnerability to experiencing emotional dysregulation and a range of mental health problems than adolescent males[1,2]. Pubertal factors may contribute to this risk by influencing brain maturation[3]. During female puberty, oestradiol (E2) and progesterone (P4) begin to cycle across the month, but this hormonal cycling is marked by high variability, i.e., variability in the hormonal cyclical patterns across the month, comparable to other transitional stages such as peri-menopause[4]. While research has examined how hormone variability during postpartum and perimenopause affects brain structure[5], far less is known about the adolescent period. Investigating how hormone variability influences adolescent brain structure could offer crucial insights into why females are at increased risk for many mental health problems during this stage of development.

During adolescence, female ovarian E2 and P4 rise and begin cycling during puberty, marking the onset of the reproductive stage[6]. Both E2 and P4 follow a distinct cyclical (or pulsatile) pattern throughout the menstrual cycle. E2 peaks twice, first in the late follicular phase (just before ovulation) and again alongside P4 during the mid-luteal phase (around seven days before menstruation). Both hormones decline during the premenstrual phase (or luteal phase)[7]. Notably, adolescent menstrual cycles are often irregular in this pattern, resulting in greater individual variability in hormonal patterns, i.e., a greater rise and fall of these hormones in a non-cyclical (non-pulsatile) manner, compared to adults[11].

Although limited, existing evidence suggests that adolescents are sensitive to hormone variability, and this may begin even before menarche. For instance, greater E2 variability is associated with increased moodiness within a day in premenarchal girls as young as 9–10

years old[12]. Similarly, weekly changes in estrone (a urinary metabolite of E2) have been associated with greater mood sensitivity in girls aged 11–14, with stronger effects observed in premenarchal girls[4]. Menstrual cycle-related mood disorders such as premenstrual syndrome (PMS) and premenstrual dysphoric disorder (PMDD) are also common in adolescent females[13], and are characterised by mood disturbances, anger/irritability, reduced interest in usual activities, and emotion dysregulation during the luteal phase of the menstrual cycle[14]. Similar behavioural patterns are observed during other hormonally dynamic life stages, such as post-pregnancy and perimenopause, where greater variability in E2 and P4 has been linked to higher levels of depressive symptoms[15,16]. Together, these findings suggest that hormonal variability may contribute to individual differences in emotion dysregulation and vulnerability to mental health difficulties across key transitional periods in the female lifespan. However, evidence during adolescence remains limited, and the underlying neural mechanisms are not yet well understood.

One possible mechanism by which hormonal variability is associated with vulnerability to mental illness symptoms and emotion dysregulation could involve the effects of hormones on brain structure during this critical period of neurodevelopment[17]. During puberty, E2 and P4 have key organisational effects on the brain via membrane-associated and intracellular receptors[18,19]. These hormone receptors are concentrated in regions that are involved in emotion regulation and are implicated in anxiety, depression and emotion dysregulation, such as the prefrontal cortex (PFC), amygdala, and hippocampus[20–22]. Prior adolescent studies have reported positive and negative associations between E2 levels and gray matter (GM) structure in PFC, parietal, middle temporal region and amygdala, and between P4 levels and GM structure within the default mode network that includes medial PFC, posterior cingulate cortex, medial and lateral temporal regions[23,24]. However, these studies have typically relied

on average hormone levels, which may mask the impact of intra-individual variability. Fluctuations in hormone levels (or non-cyclical changes in hormones) within an individual during this sensitive developmental window may uniquely exert some influence on brain structure.

To date, only one study, to our knowledge, has investigated links between E2 variability and brain structure and mental illness symptoms in adolescent females[25]. This study found a negative association between individual E2 variability and medial orbitofrontal cortical thickness; however, there was no association with internalising symptoms in this relatively small sample (n = 44). Notably, no adolescent study, to our knowledge, has explored the effects of P4 variability on brain structure or mental illness symptoms. In adult studies, women with menstrual cycle related mood disorders show structural alterations in brain regions rich in E2 and P4 receptors—such as the hippocampus, parahippocampus, and amygdala[26,27]. Further, alterations in amygdala structure were also negatively associated with depressive symptoms in this population[26]. These findings underscore the importance of understanding how individual differences in hormone variability may shape brain structure (particularly the regions involved in emotion regulation) and mental illness symptoms in a larger adolescent sample.

Thus, the current study aimed to investigate how individual variability in E2 and P4 levels across a month relates to GM structure, focusing on brain regions involved in emotion regulation—the PFC, hippocampus, and amygdala[28]. Given the widespread effects of average E2 levels across the brain[29], we also employed an exploratory whole-brain approach. We further aimed to investigate associations between hormone variability and internalising and externalising symptoms and emotion regulation difficulties. Links with internalising and

externalising symptoms were investigated, given higher rates of these symptoms in females with menstrual cycle related mood disorders[30]. Finally, we aimed to test whether individual differences in brain structure mediate links between E2/P4 variability and mental health problems (defined as mental illness symptoms and emotion dysregulation in this study).

We hypothesised that variability in E2 and P4 would be associated with the structure of brain regions involved in emotion regulation, given that these regions are dense in E2 and P4 receptors[31]. Further, we hypothesised that there would be a significant association between E2/P4 variability and mental health problems via these brain structures. Due to the lack of adolescent research, we do not specify directional hypotheses.

2. Materials and Methods:

Study hypotheses and analyses were preregistered (**10.17605/OSF.IO/RB84H**). Non-preregistered analyses are noted as exploratory.

2.1 Participants:

Female participants aged 11 to 17 years were drawn from the Puberty and NeuroDevelopment in Adolescents (PANDA) Study. The PANDA Study is a cross-sectional study that recruited participants from Melbourne, Australia. Ethics approval was granted through the Royal Children's Hospital Human Ethics Research Office (#1955580). Participants diagnosed with any developmental or endocrine disorders were excluded from this study. Additionally, those taking hormonal contraceptives were also excluded (for detailed inclusion and exclusion criteria, refer to the study protocol)[32]. A subset of individuals with clinical and MRI assessments, and weekly saliva samples over one month (i.e., minimum four saliva samples, more details in

SM), were used in the current study (n=145, Figure 1). Data quality control and outlier detection are provided in the Supplementary Material (SM).

Figure 1: Study overview. Each week, participants collected saliva samples every morning to assay oestradiol (E2) and progesterone (P4). Along with saliva samples, each participant recorded their mood in the past week using the Positive and Negative Affect Schedule (PANAS). Clinical measures, including mental health symptoms and emotion regulation, were collected during the week before the in-person MRI visit.

2.2 Hormone (E2 and P4) acquisition and processing:

Participants collected saliva samples immediately after waking (before eating, drinking, or brushing their teeth) once per week for four weeks before their MRI scan, plus an additional sample on the day of the scan (five samples total). Each sample (~2.5 mL) was collected via passive drool into 10 mL Techno Plas centrifuge tubes using a straw. After collection, samples were frozen in participants' homes within a sealed Bio-Bottle® container until the day of their MRI assessment and transported on ice to the assessment site. Upon arrival, samples were stored at –30°C until processing. Saliva was then thawed, centrifuged, and the supernatant was assayed for estradiol (E2) and progesterone (P4) by the Vaccine Immunology Group at MCRI

(Murdoch Children's Research Institute) using Salimetrics ELISA kits, with an E2 assay range of 1 – 32 pg/ml and a P4 assay range of 10 – 2430 pg/ml.

<u>E2 & P4 variability measures:</u>

We indexed intra-individual E2 and P4 variability using the standard deviation (SD) of each participant's hormone measurements, consistent with prior adolescent work[25]. Although alternative metrics such as the mean absolute successive difference (MASD) and mean squared successive difference (MSSD) have been proposed[4], these were highly correlated with SD in our sample (see SM). We therefore used SD as the primary variability measure.

2.3 Magnetic resonance imaging (MRI)

Data acquisition:

Neuroimaging data were acquired on the 3T Siemens MAGNETOM Prisma fit scanner – 32 channel head coils (Siemens, Erlangen, Germany) at the Royal Children's Hospital, Melbourne, Australia. Due to software upgrades, some sequence parameters were different in subsets of participants. However, the majority of the T1-weighted scans acquired (~80%) had the following parameters: repetition time $=$ 2500ms; echo time1 $=$ 1.72ms; echo time2 $=$ 3.45ms; echo time3 $=$ 5.18ms; echo time 4 $=$ 6.91ms), flip angle $=$ 8°; field of view $=$ 256 mm; 208 contiguous 0.9 mm thick slices (voxel dimensions $=$ 0.9mm^3). Details on the imaging parameters for other participants are in SM.

Data preprocessing:

T1-weighted images were processed using FreeSurfer v7.0.0 (http://surfer.nmr.mgh.harvard.edu/). Cortical reconstruction and segmentation were done with the recon-all function from FreeSurfer with 10 mm FWHM smoothing. This pipeline includes intensity normalisation, skull stripping, surface reconstruction, and segmentation, yielding vertex-wise cortical thickness and surface area measures[33,34]. Details on T1- weighted quality control procedures are in SM

Subcortical segmentation was based on the *aseg* atlas[35,] providing volumes of 40 regions (20 for each hemisphere), of which we included 14: thalamus, caudate, putamen, pallidum, hippocampus, amygdala, and nucleus accumbens for each hemisphere.

2.4 Mental illness symptoms and Emotion regulation:

Internalising symptoms were measured using the self-report Spence Children's Anxiety Scale-Short (SCAS-S)[36] and Children's Depression Inventory 2 (CDI-2)[37]. Externalising symptoms were measured using the parent-report Strengths and Difficulties Questionnaire (SDQ)[38]. Emotion regulation was measured using the self-report Difficulties in Emotion Regulation Scale – Short Form (DERS-SF)[39]. All scales demonstrated good internal consistency in the current sample (Cronbach's α = 0.72–0.90; see SM for details).

2.5 <u>PANAS:</u>

Positive and negative affect over the past week was measured using the Positive and Negative Affect Schedule (PANAS)[40]. The PANAS has demonstrated good validity and internal consistency, with Cronbach's $\alpha = 0.87$ for positive affect and Cronbach's $\alpha = 0.86$ for negative affect in our sample.

2.6 <u>Covariates:</u>

The current study included both pre- and post-menarchal adolescents. Menarche status was assessed using the self-report Pubertal Development Questionnaire (PDQ)[41], which asked participants whether they had begun menstruating and their age at first period. Given that we were interested in the effects of E2 and P4 variability over and above pubertal development, menarche status was included as a categorical covariate (yes/no) in all analyses.

Sociodemographic covariates included race (caregiver-reported), caregiver's education, and income-to-needs ratio (derived from after-tax household income and number of children). Given that the participants were scanned using different MRI sequences, we controlled for this as a dummy variable.

2.7 <u>Statistical analyses:</u>

There were some missing values for CDI (N = 4), DERS-SF (N = 10), SCAS (N = 1) and income-to-needs ratio data (N = 6). All missing values for each variable were handled using multiple imputation ("mice" package in R), with 50 iterations. We then

used the "merge_imputations" function[42] in R to impute the missing values with the mean imputed value across all the iterations.

Mental illness symptoms and emotion regulation scores were transformed using the "transformTukey" function (R package *rcompanion*) to address skewness in the raw data. Additionally, normality of residuals was checked for all models, and if there were any deviations from normality, robust linear regression ("lmrob") was used. Finally, influential outliers were checked using the Cook's distance.

Association between hormone variability and brain structure:

For the hypothesis-driven analysis, we used a PFC mask comprising cortical labels defined by the Desikan-Killiani atlas[34,35], which included merging the medial orbitofrontal cortex, lateral orbitofrontal cortex, superior frontal, rostral, and caudal middle frontal cortex, pars orbitalis, pars triangularis, pars opercularis, rostral and caudal anterior cingulate labels for the left and right hemispheres separately.

We performed both whole-brain and masked vertex-wise analyses using Freesurfer's general linear model analysis ("mri_glmfit"). In each model, E2 or P4 variability was entered as the predictor, with structural brain measures—cortical thickness and surface area at each vertex across the whole brain, and within the PFC mask—as the outcomes. To control for individual differences in brain size, surface area was scaled using estimated total intracranial volume (eTIV) during preprocessing. Vertex-wise results were corrected for multiple comparisons using Monte Carlo cluster-wise correction (cluster-forming threshold $p < 0.001$; FWE-corrected threshold $p < 0.05$)

For the subcortical volume analysis, the same linear regression model was fitted in R (using "glm" package), with volume at each subcortical region as an outcome and eTIV as an additional covariate in the models. To control for Type I error, false discovery rate (FDR) correction was applied across the 14 subcortical regions, with corrected p-values < 0.05 considered statistically significant.

Association between hormone variability and mental health/emotion regulation:

Linear regression models were run in R (using the "glm" function) to examine the relationship between hormone variability and the outcomes. Mental illness symptoms (SCAS, CDI, SDQ) and emotion dysregulation (DERS) were entered as separate outcome variables. For any significant associations, follow-up analyses were conducted on relevant subscales. To adjust for multiple comparisons, FDR correction (p < 0.05) was applied.

Mediation:

For any significant cortical clusters or subcortical volumes detected in the aforementioned analyses, mediation models were tested with mental health problem measures as outcomes, E2 or P4 variability as the predictor, and brain measures as the mediator. The mediation model was fitted using the R package "mediation" with a bootstrapping method (boot = 5000).

2.8 Exploratory analyses:

To test whether any significant findings were specific to hormone variability, in exploratory analyses, we tested mean hormone values as predictors. We also tested both menarchal status and cycle irregularity (pre-menarche, regular, and irregular cycling) as moderators. We additionally examined whether week-to-week changes in hormone levels were associated with weekly mood (positive and negative affect) changes. Refer to SM for further details on the classification of cycle irregularity.

3. Results

Demographic, pubertal and clinical characteristics (DERS-SF, CDI, SCAS, and SDQ) of the final sample are shown in Table 1.

Table 1: Descriptives of the final sample

	min – max[c]	Mean (sd)[c]	% in clinical range[g]
E2 variability[a] (pg/ml)	0.06 – 1.37	0.41 (0.21)	-
P4 variability[a] (pg/ml)	3.31 - 220.06	63.09 (44.22)	-
E2 mean[b] (pg/ml)	0.33 – 3.20	1.55 (0.65)	-
P4 mean[b] (pg/ml)	19.95 - 432.28	141.48 (99.38)	-

Age range (years)	11.11 – 16.90	14.04 (1.77)	-
Income-to-needs ratio[d]	0.00 – 44.654	2.81 (3.92)	-
	N (%)		
Pre-menarche	46 (31.72)	**-**	**-**
Post-menarche	99 (68.28)	**-**	**-**

Tanner stage for breast development

Stage 1	4 (2.76)	**-**	**-**
Stage 2	15 (10.35)	**-**	**-**
Stage 3	46 (31.72)	**-**	**-**
Stage 4	45 (31.03)	**-**	**-**
Stage 5	29 (20.00)	**-**	**-**
Missing	6 (4.14)	**-**	**-**

Tanner stage for pubic hair development

Stage 1	5 (3.45)	**-**	**-**
Stage 2	13 (8.97)	**-**	**-**
Stage 3	30 (20.69)	**-**	**-**
Stage 4	55 (37.93)	**-**	**-**
Stage 5	36 (24.83)	**-**	**-**
Missing	6 (4.14)	**-**	**-**

Child Race[e]

Mixed	21 (14.48)	**-**	**-**
White/Caucasian	96 (66.21)	**-**	**-**
Asian	28 (19.31)	**-**	**-**

Parent education

Partial high school/upper secondary school (Grade 7 to 12)	1 (0.69)	-	-
Completed high school/upper secondary school (Grade 12, typically age ~18)	2 (1.38)	-	-
TAFE (Technical and Further Education)/Vocational training course	10 (6.89)	-	-
Partial university/College	8 (5.52)	-	-
Graduated three-year University/College (Bachelor's) degree	40 (27.59)	-	-
Honours (four-year University/College degree)	27 (18.62)	-	-
Partial graduate school (Master's. PhD/doctorate, etc.)	9 (6.21)	-	-
Completed graduate school (Master's, PhD/doctorate, etc.)	48 (33.10)	-	-
	Min - Max[f]	Mean (SD)[f]	
SCAS	1-37	14.10 (7.75)	17.2 %
SDQ	0-12	2.80 (2.53)	0.7 %
DERS-SF	1-60	21.06 (11.29)	-
CDI	0-27	8.90 (6.49)	4.1 %
Positive affect	10 - 48	29.29 (7.27)	-
Negative affect	10 - 41	18.95 (6.48)	-

a. Variability in the hormones represents the calculated standard deviation using multiple hormone measurements for each individual

b. Mean in the hormones represents the calculated mean using multiple hormone measurements for each individual

c. The range (min and max) and mean (sd) is after calculating within-individual hormone mean and variability across the whole sample

d. Income-to-needs ratio was calculated as annual after-tax household income divided by the minimum income required for a family with a given number of children, based on the Australian poverty line 2023.

e. Some race categories were combined into broader groups: for example, 'Asian' includes Indian and other Asian backgrounds reported by the primary caregiver. 'Mixed' includes 'Other' or a combination of two races as reported by the primary caregiver. A detailed breakdown of the specific reported races is in the SM.

f. raw scores were used

g. T-scores were used; For SCAS & CDI = T-score > 65; For SDQ = T-score > 70

3.1 Association between hormone variability and brain structure:

There were no significant association between hormone variability and cortical structure using vertex-wise analysis (refer to uncorrected cortical images in the repository https://neurovault.org/collections/IVDQQFDM/)

Linear regression models showed a significant negative association between P4 variability and the volume of the left thalamus (Cohen's d = -0.26, pFDR = 0.031; refer to Figure 2). There were no other significant associations between hormone variability and subcortical structure (see Table 2 for subcortical results).

Table 2: Linear model between subcortical regions and hormone variability[a]

Subcortical regions (mm³)	P4 variability (pg/ml)		E2 variability (pg/ml)	
	Cohen's D	pFDR	Cohen's D	pFDR
Left Thalamus	-0.259	0.031[b]	-0.168	0.435
Right Thalamus	-0.133	0.775	-0.103	0.720
Left Caudate	0.022	0.856	0.079	0.720
Right Caudate	-0.032	0.856	0.060	0.720
Left Putamen	-0.024	0.856	-0.054	0.720
Right Putamen	0.010	0.903	-0.060	0.720
Left Pallidum	-0.046	0.814	-0.025	0.770
Right Pallidum	-0.077	0.814	-0.042	0.770
Left Hippocampus	-0.077	0.814	0.024	0.770
Right Hippocampus	-0.075	0.814	-0.065	0.720
Left Amygdala	-0.057	0.814	-0.103	0.720
Right Amygdala	-0.050	0.814	-0.157	0.435
Left Accumbens	-0.064	0.814	0.031	0.770
Right Accumbens	0.079	0.814	0.062	0.720

a. Linear models were adjusted for menarche status, age at the time of MRI, adolescent's race, primary caregiver's education, income-to-needs ratio, a dummy variable for different T1-w sequences, and ICV

b. Significant after FDR correction

Figure 2: The fitted dashed line with 95% CI (shaded area) shows a negative association between progesterone variability (pg/ml) and volume in the left thalamus (mm^3). Linear models were adjusted for menarche status, age at the time of MRI, adolescent's race, primary caregiver's education, income-to-needs ratio, a dummy variable for different T1-w sequences, and ICV

3.2 Association between hormone variability and mental illness symptoms/emotion regulation:

Hormone variability was not significantly associated with mental illness symptom/emotion regulation (p > 0.05), either before or after FDR correction. However, after checking for influential outliers using the Cook's distance method, two influential outliers were found for models testing associations between E2 variability and symptoms (see SM, Figure 1 for Cook's distance statistics and model output). After removing these outliers, there was a significant negative association between E2 variability and DERS-SF (d = -0.179; pFDR = 0.036), CDI (d = -0.179; pFDR = 0.036) and SCAS total (d = -0.170; pFDR = 0.043) scores

(as shown in Figure 3 and Table 3). Follow-up models tested associations between E2 variability and DERS-SF, CDI and SCAS sub-scales. The models showed significant negative associations between E2 variability and DERS 'strategies' (d = -0.243; pFDR = 0.010), CDI 'emotional problems' (d = -0.208; pFDR = 0.028), and SCAS 'separation' (d = -0.302; pFDR = 0.001) and 'social anxiety' (d = 0.226; pFDR = 0.007).

Figure 3: Negative association between E2 variability and A) DERS-SF score, B) CDI score, C) SCAS score. DERS-SF = Difficulties in Emotion Regulation Scale – Short Form; CDI = Children's Depression Inventory; SCAS = Spence Children's Anxiety Scale-Short. Dashed line represents the fitted linear regression with 95% CI (shaded area). Linear models were adjusted for menarche status, age at the time of MRI, adolescent's race, primary caregiver's education, and income-to-needs ratio

Table 3: Linear regression output for models testing the association between hormone variability and mental illness symptoms/emotion regulation[b]

	E2 variability (pg/ml)		P4 variability (pg/ml)	
Outcome[a]	Cohen's D	pFDR	Cohen's D	pFDR

CDI Total	-0.179	0.036[c]	0.025	0.645
SCAS Total	-0.170	0.043 [c]	-0.036	0.645
DERS-SF Total	-0.179	0.036 [c]	0.073	0.645
SDQ Total	-0.032	0.529	0.053	0.645

sub-scales[a]

DERS_strategies	-0.243	0.010 [c]	0.003	0.930
DERS_non-acceptance	-0.153	0.0847	0.0008	0.930
DERS_impulse	-0.102	0.195	0.094	0.752
DERS_goals	0.167	0.081	-0.014	0.930
DERS_awareness	0.040	0.824	0.058	0.752
DERS_clarity	-0.157	0.090	0.127	0.752
CDI_emotional problem	-0.208	0.028 [c]	0.008	0.975
CDI_functional problem	-0.118	0.322	-0.067	0.342
SCAS_separation	-0.302	0.001 [c]	-0.195	0.125
SCAS_social anxiety disorder	-0.226	0.007 [c]	-0.017	0.873
SCAS_panic disorder	-0.034	0.707	0.055	0.873
SCAS_phobias	-0.102	0.398	-0.025	0.873
SCAS_generalised anxiety disorder	-0.109	0.425	-0.072	0.873

a. All mental health scores were transformed before model fitting

b. Linear models were adjusted for menarche status, age at the time of MRI, adolescent's race, primary caregiver's education, and income-to-needs ratio

c. significant after FDR correction

CDI = Children's Depression Inventory; SCAS = Spence Children's Anxiety Scale-Short;

DERS-SF = Difficulties in Emotion Regulation Scale – Short Form; SDQ = Strengths and

Difficulties Questionnaire

3.3 Mediating role of Left Thalamus volume in the association between hormone variability and mental illness symptoms and emotion regulation:

There were no significant indirect effects of P4 variability on mental health symptoms (CDI = -0.0005 [-0.0075 0.01], SCAS = -0.0001 [-0.0003 0.00], SDQ = -0.00074 [-0.00208 0.00]) or emotion dysregulation (DERS-SF = -0.0007 [-0.0022 0.00]) via left thalamus volume.

3.4 Exploratory analyses

i) **Association between mean hormone levels and brain structure and mental illness symptoms/emotion regulation:**

Mean P4 levels were negatively associated with CDI total symptoms (Cohen's d = -0.192, p = 0.022), but this effect did not survive multiple comparison correction (pFDR = 0.088). No other significant associations were found between mean E2 or P4 levels and brain structure or mental illness symptoms/emotion regulation. Refer to the SM Table 5 for model output.

ii) **Menarche status or cycle irregularity as a moderator in the association between hormone variability and brain structure and mental illness symptoms/emotion regulation:**

Refer to SM for pre- versus post-menarche comparisons of hormone variability and correlation between Tanner stages and hormone variability

There was a significant moderating effect of menarche status on the associations between E2 variability and right hippocampus volume, DERS total and CDI total scores. Simple slope analysis showed significant effects for the pre-menarche group only, with greater E2 variability associated with reduced right hippocampal volume, depressive symptoms and emotion dysregulation. Upon removal of detected influential outliers (as discussed in section 3.2 – I.A), the moderation effects became non-significant, but the effects in the pre-menarche group remained significant (Refer to SM for Figure 2 and model statistics).

There were no significant moderating cycle irregularity effects on any of the mental illness symptoms, emotion dysregulation, or brain structure measures.

iii) **Association between weekly hormone changes and weekly positive and negative affect:**

Linear mixed effect models showed a significant positive association between the absolute change in P4 levels from one week to the next and weekly

negative affect scores in the corresponding week (d = 0.115, p = 0.012). That is, greater week-to-week fluctuations in P4 levels were linked to increased negative affect. Additionally, there was a significant absolute change*direction effect in the association between weekly E2 levels and weekly positive affect (d = 0.084, p = 0.048). A reduction in E2 levels from one week to the next was linked with a reduction in weekly positive affect (refer to SM).

4. Discussion

This study examined how hormone variability relates to brain structure and mental health problems in adolescent females. While we found that greater P4 variability was associated with lower left thalamus volume, we did not observe significant effects for other hypothesised regions. Additionally, we found that E2 variability was associated with lower depressive and anxiety symptoms, as well as emotion dysregulation. However, these associations were not mediated by brain structural changes as hypothesised.

Our finding that greater P4 variability was associated with smaller left thalamus volume may reflect a role of P4 in the maturation of the thalamus, a region with high expression of P4 receptors[43]. Prior longitudinal studies show that thalamic volume tends to decrease during early to mid-adolescence, to a greater extent in females compared to males[44]. Thus, our findings might suggest P4 variability as a biological mechanism partly responsible for thalamic development among females. Interestingly, structural changes in the thalamus have also been observed during other hormonal transitional stages, such as pregnancy and menopause[5], as well as in menstrual cycle related clinical conditions such as PMDD , PMS and Primary Dysmenorrhea (PD), with studies reporting reduced thalamic volume associated

with higher pain in women with these conditions[45]. Given the thalamus's role in emotion

regulation through its connections with the midbrain and cortex[46], the current findings

highlight it as a potential region through which P4 variability may influence emotional

functioning during hormonal transition periods and disorders. However, we did not find links

between P4 variability and mental health problems, directly or indirectly via thalamus

volume. This null finding may indicate that P4-related changes in thalamus volume influence

emotional (or other) functioning not measured in this study, or that effects on mental health

may unfold over time.

Menarche or cycle regularity did not significantly moderate this effect, suggesting that the

link between P4 variability and thalamic volume may be independent of menarche or the

development of stable cyclic patterns. Further research is needed to clarify whether P4-

related structural changes in the thalamus are specific to certain pubertal periods or persist

into adulthood, which would help clarify whether these patterns reflect temporary

developmental adaptations or stable individual differences in thalamic structure.

Contrary to our hypothesis, hormone variability was not associated with the structure of any

other subcortical or cortical regions. Similarly prior study[25] reported no vertex-wise

associations, although they did find a negative relation between E2 variability and thickness

in the right medial orbitofrontal cortex using an ROI approach. Notably, their study involved

a smaller sample and a higher proportion of pre-menarche participants. Interestingly, in our

exploratory subcortical ROI analyses, higher E2 variability was linked to smaller right

hippocampus volume only among the pre-menarche group. Although speculative, this may

reflect delayed brain maturation with higher E2 variability in this group, as these structures

typically increase during adolescence[47]. However, future longitudinal studies are required to

verify these patterns. Prior adolescent studies have linked average E2 or P4 levels to cortical density in prefrontal, parietal, and middle temporal cortices, and adult and animal studies have shown a reduced hippocampal volume during menstrual phases with average low E2 levels[23,24]. However, we found no associations for mean E2 or P4. This may partly reflect methodological differences, as our mean hormone estimates were based on four-five samples collected across a month, whereas prior studies used only two consecutive samples collected over two days. Additionally, previous work often included smaller female samples (<50 participants), did not account for menarchal status, and controlled for a specific cycle phase, all of which could contribute to inconsistent findings. Thus, future large-scale and replicable studies are needed to clarify hormone–brain associations.

In line with our hypothesised relationship between hormone variability and mental health problems, we found that increased E2 variability over a month was linked to reduced internalising symptoms and emotion regulation problems. To date, there are mixed findings regarding the direction of the relationship between E2 variability and mental health problems. For instance, a recent study has shown no association between E2 variability and depressive or anxiety symptoms in adolescent females, although their sample was relatively small and had limited variability in symptom levels[25]. In contrast, adult studies have shown that greater E2 variability during perimenopause was associated with heightened depressive symptoms[15,48]. Notably, our exploratory analysis indicated that the association between E2 variability and reduced mental health problems was stronger among pre-menarchal participants. This finding suggests that E2 variability during early adolescence may not have the same adverse implications observed during other hormonal transitions (e.g., postpartum, perimenopause). While speculative, greater E2 variability before menarche may reflect a normative feature of pubertal maturation or the gradual establishment of the biological

mechanism, facilitating further emotional maturational processes. However, because symptom measures were collected concurrently with hormone sampling, and given the limited number of adolescent studies examining E2 variability, these interpretations should be considered preliminary and warrant cautious interpretation.

Interestingly, our within-person exploratory results indicated that a reduction in E2 levels from one week to the next was associated with less weekly positive affect, aligning with prior adolescent studies showing heightened mood sensitivity to E2 withdrawal and affective disturbances during other E2 withdrawal stages (i.e., peripartum symptoms)[4,49]. In addition, greater week-to-week change in P4 levels (regardless of the direction of change) was associated with greater weekly negative affect, suggesting that P4 instability may heighten sensitivity to negative mood. While the mechanisms by which weekly changes in E2 and P4 contribute to affect remain unclear, increased stress sensitivity during periods of hormonal fluctuation may be one potential pathway[50]. Moreover, heightened stress sensitivity has also been an important risk factor for the development of psychopathology during adolescence[51,52]. Thus, changes in weekly E2 and P4 levels that coincide with changes in affect may serve as early markers of emerging emotional patterns or vulnerabilities

Notably, E2 changes showed a divergence in short- versus longer-term associations: while short-term reductions in E2 corresponded to lower positive mood, greater overall E2 variability across the month was associated with fewer mental health problems. This pattern suggests that transient hormonal declines may influence momentary mood states, whereas a more dynamic hormonal profile may reflect adaptive physiological responsivity. Further research is needed to clarify how greater hormonal variability, particularly when coupled

with chronic or cumulative stress exposure, may contribute to the transition from normative affective sensitivity to later psychopathology.

5. Limitations and future directions:

The current study is the first to examine how hormone variability relates to brain structure and mental health problems in adolescent females within a sample of this scale. However, several additional limitations should be considered when interpreting these findings.

Although the study captured weekly hormone data and self-reported menstrual cycle information, self-reports likely introduce error, particularly when adjusting for individual differences in menstrual cycle regularity among post-menarche girls. This limitation may have contributed to null associations when used as a moderator in exploratory analysis. Future research should aim to collect hormone samples at a higher frequency (e.g., every few days) and incorporate ovulation testing (e.g., using ovulation kits[53]) to more precisely map menstrual cycle phases and control for irregularity in cycle length.

Our study focused exclusively on E2 or P4. However, prior adolescent research has shown that other hormones, such as estrone (E1), demonstrate earlier increases during the pubertal transition compared with E2[4,54]. Future work should therefore examine a broader range of hormones to more fully characterise patterns of hormonal variability and their relevance for adolescent female development.

Further, Saliva was used as the collection medium because it is non-invasive and well-suited to frequent sampling in adolescents. Although plasma is the traditional gold standard for hormones like E2 and P4, saliva captures the free, biologically active fraction and may be

more relevant for behaviour[55]. However, salivary assays are influenced by rapid fluctuations in hormone levels, necessitating frequent sampling for reliable estimates. While all biospecimens are affected by temporal hormone variation, saliva reflects these shifts more immediately, making sample timing and density particularly important for valid measurement[56]. Despite using a sensitive Salimetrics ELISA kit (lower limit: 0.998 pg/ml), four participants had values below the detection limit, indicating that approaches such as Liquid chromatography- tandem mass spectrometry (LC-MS/MS) and denser sampling schedules may be needed to improve accuracy for low-concentration hormone measurements[57].

Finally, the current sample was recruited from Melbourne and comprised a community-based cohort, with a higher proportion of White/Caucasian participants (66.2%), and only a small number of individuals falling within the clinical range for mental illness symptoms. This limited diversity and restricted symptom range may have reduced the generalisability of our findings and contributed to the modest associations with mental illness symptoms (effect size ~ 0.18). Replication in more diverse and clinically enriched samples will be important for determining the broader and clinical relevance of hormone–brain–behaviour associations.

Conclusion

In conclusion, this study provides the first evidence of the links between E2 and P4 variability and brain structure and mental health problems in adolescent females. These findings lay an important foundation for future research examining how fluctuations in hormones contribute to brain development and risk for mental health difficulties across the pubertal transition. Moreover, the distinct associations observed between hormone variability,

affect, and behaviour highlight the importance of investigating these dynamics in future developmental studies.

CRediT authorship contribution statement

MK: Writing – original draft, Conceptualisation, Formal analysis, Visualisation. **YT:** Writing – review & editing, Conceptualisation, Supervision. **NV:** Conceptualisation, Supervision, Funding acquisition. **MH**: Writing – review & editing, Funding acquisition**. MC**: Writing – review & editing, Funding acquisition. **MS:** Writing – review & editing, Funding acquisition. **SW:** Conceptualisation, Methodology, Resources, Supervision, Project administration, Funding acquisition, Writing – review & editing.

References:

1. Campbell OLK, Bann D, Patalay P. The gender gap in adolescent mental health: A cross-national investigation of 566,829 adolescents across 73 countries. *SSM - Popul Health*. 2021;13:100742. doi:10.1016/j.ssmph.2021.100742

2. Yoon Y, Eisenstadt M, Lereya ST, Deighton J. Gender difference in the change of adolescents' mental health and subjective wellbeing trajectories. *Eur Child Adolesc Psychiatry*. 2023;32(9):1569-1578. doi:10.1007/s00787-022-01961-4

3. Juraska JM, Willing J. Pubertal onset as a critical transition for neural development and cognition. *Brain Res*. 2017;1654:87-94. doi:10.1016/j.brainres.2016.04.012

4. Andersen E, Fiacco S, Gordon J, et al. Methods for characterizing ovarian and adrenal hormone variability and mood relationships in peripubertal females. *Psychoneuroendocrinology*. 2022;141:105747. doi:10.1016/j.psyneuen.2022.105747

5. Catenaccio E, Mu W, Lipton ML. Estrogen- and progesterone-mediated structural neuroplasticity in women: evidence from neuroimaging. *Brain Struct Funct*. 2016;221(8):3845-3867. doi:10.1007/s00429-016-1197-x

6. Das PK, Mukherjee J, Banerjee D. Female Reproductive Physiology. In: Das PK, Sejian V, Mukherjee J, Banerjee D, eds. *Textbook of Veterinary Physiology*. Springer Nature; 2023:513-568. doi:10.1007/978-981-19-9410-4_22

7. Barbieri RL. The Endocrinology of the Menstrual Cycle. In: Rosenwaks Z, Wassarman PM, eds. *Human Fertility: Methods and Protocols*. Springer; 2014:145-169. doi:10.1007/978-1-4939-0659-8_7

8. Kessler RC, McGonagle KA, Swartz M, Blazer DG, Nelson CB. Sex and depression in the National Comorbidity Survey. I: Lifetime prevalence, chronicity and recurrence. *J Affect Disord*. 1993;29(2-3):85-96. doi:10.1016/0165-0327(93)90026-g

9. Deecher D, Andree TH, Sloan D, Schechter LE. From menarche to menopause: Exploring the underlying biology of depression in women experiencing hormonal changes. *Psychoneuroendocrinology*. 2008;33(1):3-17. doi:10.1016/j.psyneuen.2007.10.006

10. Joffe H, Cohen LS. Estrogen, serotonin, and mood disturbance: where is the therapeutic bridge? *Biol Psychiatry*. 1998;44(9):798-811. doi:10.1016/S0006-3223(98)00169-3

11. Treloar AE, Boynton RE, Behn BG, Brown BW. Variation of the human menstrual cycle through reproductive life. *Int J Fertil*. 1967;12(1 Pt 2):77-126.

12. Buchanan CM. *Hormone Concentrations and Variability: Associations with Self-Reported Moods and Energy in Early Adolescent Girls*.; 1989. Accessed March 24, 2025. https://eric.ed.gov/?id=ED305166

13. Itriyeva K. Premenstrual syndrome and premenstrual dysphoric disorder in adolescents. *Curr Probl Pediatr Adolesc Health Care*. 2022;52(5):101187. doi:10.1016/j.cppeds.2022.101187

14. Eggert L, Witthöft M, Hiller W, Kleinstäuber M. Emotion Regulation in Women with Premenstrual Syndrome (PMS): Explicit and Implicit Assessments. *Cogn Ther Res*. 2016;40(6):747-763. doi:10.1007/s10608-016-9788-7

15. Gordon JL, Peltier A, Grummisch JA, Sykes Tottenham L. Estradiol Fluctuation, Sensitivity to Stress, and Depressive Symptoms in the Menopause Transition: A Pilot Study. *Front Psychol*. 2019;10. doi:10.3389/fpsyg.2019.01319

16. Lozza-Fiacco S, Gordon JL, Andersen EH, et al. Baseline anxiety-sensitivity to estradiol fluctuations predicts anxiety symptom response to transdermal estradiol treatment in perimenopausal women – A randomized clinical trial. *Psychoneuroendocrinology*. 2022;143:105851. doi:10.1016/j.psyneuen.2022.105851

17. Schulz KM, Sisk CL. The organizing actions of adolescent gonadal steroid hormones on brain and behavioral development. *Neurosci Biobehav Rev*. 2016;70:148-158. doi:10.1016/j.neubiorev.2016.07.036

18. Kasubuchi M, Watanabe K, Hirano K, et al. Membrane progesterone receptor beta (mPRβ/Paqr8) promotes progesterone-dependent neurite outgrowth in PC12 neuronal cells via non-G protein-coupled receptor (GPCR) signaling. *Sci Rep*. 2017;7(1):5168. doi:10.1038/s41598-017-05423-9

19. Schumacher M, Mattern C, Ghoumari A, et al. Revisiting the roles of progesterone and allopregnanolone in the nervous system: Resurgence of the progesterone receptors. *Prog Neurobiol*. 2014;113:6-39. doi:10.1016/j.pneurobio.2013.09.004

20. Österlund MK, Gustafsson JÅ, Keller E, Hurd YL. Estrogen Receptor β (ERβ) Messenger Ribonucleic Acid (mRNA) Expression within the Human Forebrain: Distinct Distribution Pattern to ERα mRNA1. *J Clin Endocrinol Metab*. 2000;85(10):3840-3846. doi:10.1210/jcem.85.10.6913

21. Schmaal L, Hibar DP, Sämann PG, et al. Cortical abnormalities in adults and adolescents with major depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major Depressive Disorder Working Group. *Mol Psychiatry*. 2017;22(6):900-909. doi:10.1038/mp.2016.60

22. Vijayakumar N, Whittle S, Yücel M, Dennison M, Simmons J, Allen NB. Thinning of the lateral prefrontal cortex during adolescence predicts emotion regulation in females. *Soc Cogn Affect Neurosci*. 2014;9(11):1845-1854. doi:10.1093/scan/nst183

23. Curtis M, Flournoy JC, Kandala S, et al. Disentangling the unique contributions of age, pubertal stage, and pubertal hormones to brain structure in childhood and adolescence. *Dev Cogn Neurosci*. 2024;70:101473. doi:10.1016/j.dcn.2024.101473

24. Peper JS, Brouwer RM, Schnack HG, et al. Sex steroids and brain structure in pubertal boys and girls. *Psychoneuroendocrinology*. 2009;34(3):332-342. doi:10.1016/j.psyneuen.2008.09.012

25. Zwaan IS, Felmingham K, Vijayakumar N, et al. Estradiol variability is associated with brain structure in early adolescent females. *Psychoneuroendocrinology*. 2022;146:105943. doi:10.1016/j.psyneuen.2022.105943

26. Dubol M, Wikström J, Lanzenberger R, Epperson CN, Sundström-Poromaa I, Comasco E. Grey matter correlates of affective and somatic symptoms of premenstrual dysphoric disorder. *Sci Rep*. 2022;12(1):5996. doi:10.1038/s41598-022-07109-3

27. Jeong HG, Ham BJ, Yeo HB, Jung IK, Joe SH. Gray matter abnormalities in patients with premenstrual dysphoric disorder: An optimized voxel-based morphometry. *J Affect Disord*. 2012;140(3):260-267. doi:10.1016/j.jad.2012.02.010

28. Guyer AE, Silk JS, Nelson EE. The neurobiology of the emotional adolescent: From the inside out. *Neurosci Biobehav Rev*. 2016;70:74-85. doi:10.1016/j.neubiorev.2016.07.037

29. Vijayakumar N, Mills KL, Alexander-Bloch A, Tamnes CK, Whittle S. Structural brain development: A review of methodological approaches and best practices. *Dev Cogn Neurosci*. 2018;33:129-148. doi:10.1016/j.dcn.2017.11.008

30. Luo D, Dashti SG, Sawyer SM, Vijayakumar N. Pubertal hormones and mental health problems in children and adolescents: a systematic review of population-based studies. *eClinicalMedicine*. 2024;76. doi:10.1016/j.eclinm.2024.102828

31. Bixo M, Ba¨ckstro¨m T, Winblad B, Andersson A. Estradiol and testosterone in specific regions of the human female brain in different endocrine states. *J Steroid Biochem Mol Biol*. 1995;55(3):297-303. doi:10.1016/0960-0760(95)00179-4

32. Bray KO, Durbin O, Hartanto S, et al. Puberty and NeuroDevelopment in adolescents (PANDA): a study protocol. *BMC Pediatr*. 2024;24(1):768. doi:10.1186/s12887-024-05197-w

33. Dale AM, Fischl B, Sereno MI. Cortical Surface-Based Analysis: I. Segmentation and Surface Reconstruction. *NeuroImage*. 1999;9(2):179-194. doi:10.1006/nimg.1998.0395

34. Fischl B, Salat DH, Busa E, et al. Whole Brain Segmentation: Automated Labeling of Neuroanatomical Structures in the Human Brain. *Neuron*. 2002;33(3):341-355. doi:10.1016/S0896-6273(02)00569-X

35. Desikan RS, Ségonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*. 2006;31(3):968-980. doi:10.1016/j.neuroimage.2006.01.021

36. Ahlen J, Vigerland S, Ghaderi A. Development of the Spence Children's Anxiety Scale - Short Version (SCAS-S). *J Psychopathol Behav Assess*. 2018;40(2):288-304. doi:10.1007/s10862-017-9637-3

37. Kovacs M. Children's Depression Inventory (CDI and CDI 2). In: *The Encyclopedia of Clinical Psychology*. John Wiley & Sons, Ltd; 2015:1-5. doi:10.1002/9781118625392.wbecp419

38. Goodman R. The Strengths and Difficulties Questionnaire: a research note. *J Child Psychol Psychiatry*. 1997;38(5):581-586. doi:10.1111/j.1469-7610.1997.tb01545.x

39. A bifactor analysis of the Difficulties in Emotion Regulation Scale - Short Form (DERS-SF) in a sample of adolescents and adults | Current Psychology. Accessed December 8, 2025. https://link.springer.com/article/10.1007/s12144-019-00602-5

40. Dahl RE, Forbes EE. Pubertal Development and Behavior: Hormonal Activation of Social and Motivational Tendencies. *Brain Cogn*. 2010;72(1):66-72. doi:10.1016/j.bandc.2009.10.007

41. Petersen AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal status: Reliability, validity, and initial norms. *J Youth Adolesc*. 1988;17(2):117-133. doi:10.1007/BF01537962

42. Burns RA, Butterworth P, Kiely KM, et al. Multiple imputation was an efficient method for harmonizing the Mini-Mental State Examination with missing item-level data. *J Clin Epidemiol*. 2011;64(7):787-793. doi:10.1016/j.jclinepi.2010.10.011

43. Brinton RD, Thompson RF, Foy MR, et al. Progesterone receptors: form and function in brain. *Front Neuroendocrinol*. 2008;29(2):313-339. doi:10.1016/j.yfrne.2008.02.001

44. Dennison M, Whittle S, Yücel M, et al. Mapping subcortical brain maturation during adolescence: evidence of hemisphere- and sex-specific longitudinal changes. *Dev Sci*. 2013;16(5):772-791. doi:10.1111/desc.12057

45. Tu CH, Niddam DM, Chao HT, et al. Brain morphological changes associated with cyclic menstrual pain. *Pain*. 2010;150(3):462-468. doi:10.1016/j.pain.2010.05.026

46. Asami T, Yoshida H, Takaishi M, et al. Thalamic shape and volume abnormalities in female patients with panic disorder. *PLOS ONE*. 2018;13(12):e0208152. doi:10.1371/journal.pone.0208152

47. Krogsrud SK, Tamnes CK, Fjell AM, et al. Development of hippocampal subfield volumes from 4 to 22 years. *Hum Brain Mapp*. 2014;35(11):5646-5657. doi:10.1002/hbm.22576

48. Joffe H, de Wit A, Coborn J, et al. Impact of Estradiol Variability and Progesterone on Mood in Perimenopausal Women With Depressive Symptoms. *J Clin Endocrinol Metab*. 2020;105(3):e642-e650. doi:10.1210/clinem/dgz181

49. Bloch M, Schmidt PJ, Danaceau M, Murphy J, Nieman L, Rubinow DR. Effects of gonadal steroids in women with a history of postpartum depression. *Am J Psychiatry*. 2000;157(6):924-930. doi:10.1176/appi.ajp.157.6.924

50. Gordon JL, Girdler SS, Meltzer-Brody SE, et al. Ovarian hormone fluctuation, neurosteroids, and HPA axis dysregulation in perimenopausal depression: a novel heuristic model. *Am J Psychiatry*. 2015;172(3):227-236. doi:10.1176/appi.ajp.2014.14070918

51. Muñoz-Samons D, Tor J, Rodríguez-Pascual M, et al. Recent stressful life events and stress sensitivity in children and adolescents at clinical risk for psychosis. *Psychiatry Res*. 2021;303:114017. doi:10.1016/j.psychres.2021.114017

52. McLaughlin KA, Hatzenbuehler ML. Stressful Life Events, Anxiety Sensitivity, and Internalizing Symptoms in Adolescents. *J Abnorm Psychol*. 2009;118(3):659-669. doi:10.1037/a0016499

53. Ayoola AB, Slager D, Feenstra C, Zandee GL. A Feasibility Study of Women's Confidence and Comfort in Use of a Kit to Monitor Ovulation. *J Midwifery Womens Health*. 2015;60(5):604-609. doi:10.1111/jmwh.12347

54. Biro FM, Pinney SM, Huang B, Baker ER, Walt Chandler D, Dorn LD. Hormone changes in peripubertal girls. *J Clin Endocrinol Metab*. 2014;99(10):3829-3835. doi:10.1210/jc.2013-4528

55. Chen MC, Hamilton JP, Gotlib IH. Decreased Hippocampal Volume in Healthy Girls at Risk of Depression. *Arch Gen Psychiatry*. 2010;67(3):270-276. doi:10.1001/archgenpsychiatry.2009.202

56. Stern J, Arslan RC, Penke L. Stability and validity of steroid hormones in hair and saliva across two ovulatory cycles. *Compr Psychoneuroendocrinology*. 2022;9:100114. doi:10.1016/j.cpnec.2022.100114

57. Brouillard A, Davignon LM, Cernik R, et al. Comparing immunoassay and mass spectrometry techniques for salivary sex hormone analysis. *Psychoneuroendocrinology*. 2025;174:107379. doi:10.1016/j.psyneuen.2025.107379