Universal rapid RNA-based quantification of toxigenic *Alexandrium*

species (Dinophyceae) using quantitative recombinase polymerase

amplification

Ioannis Markopoulos[a], Isavella Papadopoulou[a], Christos Chantzaras[a], Katherine Hartle-

Mougiou[b,a], Frédéric Verret[a], <u>Electra Gizeli[b,a*]</u>, <u>Martha Valiadi[a]*</u>

[a] Foundation for Research and Technology Hellas, Institute of Molecular Biology and

Biotechnology, Heraklion, Crete, Greece

[b] Department of Biology, University of Crete, Heraklion, Crete, Greece

Underlined authors share senior authorship

*Corresponding authors: (martha_valiadi@imbb.forth.gr; gizeli@imbb.forth.gr)

Abstract

Harmful algal blooms caused by toxigenic *Alexandrium* species pose recurrent risks to coastal

ecosystems and public health, yet current monitoring approaches rely on microscopy and

laboratory-based toxin analysis with limited capacity for rapid, functional early warning. Here, we

present a universal quantitative reverse-transcriptase recombinase polymerase amplification (qRT-

RPA) assay targeting the *sxt*A4 transcript, an essential gene in saxitoxin biosynthesis. The RPA

chemistry is isothermal with a low running temperature, making it suitable for portable, on-site

testing. The assay was designed against a conserved *sxt*A4 region and validated using isolated

*sxt*A4 amplicons from multiple *Alexandrium* species, synthetic DNA and RNA templates, and total

RNA of *Alexandrium minutum* as a widespread reference species. The assay achieved uniform

amplification kinetics across species, a limit of detection below 10^3 synthetic RNA copies and 0.1

ng total RNA per reaction for *A. minutum*, and a runtime of less than 15 minutes. The assay

selectively detected *sxt*A4 transcripts from toxigenic *Alexandrium* strains and showed no cross-

reactivity with non-target phytoplankton. A limit of detection that is relevant to early warning for

Alexandrium species (under 20 cells per reaction) was retained in complex RNA matrices. Mock
samples prepared by spiking cultured cells into natural seawater also demonstrated detection at
field-relevant concentrations. These results establish qRT-RPA as a rapid, RNA-based,
functionally informative molecular tool that provides a foundation for portable, early-warning
monitoring of potentially toxigenic *Alexandrium* blooms.

Keywords

34 *Alexandrium*, dinoflagellate, harmful algal blooms, saxitoxin, *sxt*A4 RNA

1 Introduction

Harmful Algal Blooms (HABs) are posing increasing ecological and socioeconomic pressure in coastal areas worldwide. Their global expansion is attributed to anthropogenic nutrient enrichment and ocean warming (Dai et al., 2023; Gobler et al., 2017; Griffith et al., 2019; Hayashida et al., 2020; Rabalais et al., 2007; Yu et al., 2023). Warming temperatures can promote bloom initiation (Anderson et al., 2021a) accelerated cell growth rates and prolonged bloom seasons (Anderson et al., 2021a; Gobler et al., 2017; Griffith et al., 2019) especially in temperate regions (Dai et al., 2023), whilst also allowing their intrusion into previously unaffected polar areas (Anderson et al., 2021a). The consequences of HABs range from ecosystem-scale deoxygenation to the accumulation of potent toxins that threaten food security and human health (Anderson et al., 2021a; Anderson et al., 2021b). Despite progress in ocean observing systems, early-warning capacity often remains limited, particularly in regions where toxic events develop rapidly or surveillance infrastructure is sparse.

The dinoflagellate genus *Alexandrium* is responsible for HABs worldwide. Species of this genus produce an array of toxins as secondary metabolites, being most well-known for the production of saxitoxins. These cause Paralytic Shellfish Poisoning (PSP) in humans upon consumption of contaminated shellfish ((Anderson et al., 2012; Gribble et al., 2025; Lewis et al., 2018; Suleiman et al., 2017). Most regulatory monitoring programs still rely on a dual structure: (i) microscopic identification and enumeration of cells in water samples, and (ii) chemical toxin analysis in shellfish tissue (Parks et al., 2020; Turner et al., 2014; union, 2004). The first assesses putatively toxic cell abundance, while the second measures saxitoxin and congeners which are converted to toxicity equivalence factors (TEFs) (Commission Implementing Regulation (EU) 2021/1709, (Commission., 2021). Concerned countries like the UK follow this regime with practical thresholds of 400 STX eq. (Turner et al., 2014) in shellfish flesh or 40 - 100 *Alexandrium* cells L^{-1} of water triggering intensified monitoring frequency ((Parks et al., 2020; Senovilla-Herrero et al., 2025). In other locations the thresholds may be much higher if cells are less toxic; there is considerable variability in toxicity level regionally, interspecifically and intraspecifically (Geffroy et al., 2021; Sung et al., 2025). While these methods underpin long-established monitoring frameworks, they present significant operational constraints. Microscopy cannot reliably distinguish between toxic and non-toxic strains of morphologically similar taxa and provides no

functional information on active toxin production at the time. Meanwhile, toxin chemical analyses
by HPLC (Standard EN 14526) requires centralized laboratories, limiting sampling frequency and
increasing turnaround time to minimum of 3 days. Together, these limitations hinder early warning
and intervention, increasing the likelihood of food contamination and ecological damage.

Molecular amplification-based methods represent an alternative to microscopy analysis, offering
higher sensitivity, faster time-to-result, species-level resolution and quantification of specific
cellular functions (John et al., 2007; Kon et al., 2015; Murray et al., 2019). A wealth of DNA-
based qPCR assays have been developed to enumerate specific toxic species (Dai et al., 2020;
Erdner et al., 2010; Gao et al., 2015a; Gao et al., 2015b; Garneau et al., 2011; Hatfield et al., 2019;
Ruvindy et al., 2018; Toebe et al., 2013; Vandersea et al., 2017). Most existing assays target the
small or large ribosomal genes of specific *Alexandrium* species to estimate abundance, whereas
several qPCR assays targeting genes of the saxitoxin cluster enable direct assessment of the
population's toxigenic potential (Gao et al., 2015b; Han-Sol et al., 2024; Murray et al., 2019;
Murray et al., 2011; Penna et al., 2015; Savela et al., 2016; Stüken et al., 2013). Despite these
advances, the transition from traditional light microscopy to molecular monitoring for HAB
surveillance has been slow to materialize. While qPCR is the gold standard of molecular detection
due to its cost-effectiveness and high throughput, it is mainly applied in a laboratory setting due to
the requirement for high-power consuming thermal cycling.

HAB surveillance will significantly benefit from on-site or *in situ* monitoring offering real-time
detection. In contrast to qPCR, isothermal amplification can achieve molecular quantification in
remote field settings, using portable battery-powered analyzers (e.g. Pebble R by BIOPIX-T;
TwistAmp by TwistDx). In particular, recombinase polymerase amplification (RPA) operates at a
low, stable temperature (37 – 42 ºC), has high tolerance to impurities and a fast time-to-result in
less than 15 minutes (Tan et al., 2022). RPA chemistry can be used for qualitative detection using
lateral flow strips (Liu et al., 2023; Yao et al., 2024; Yu et al., 2025) and quantitative analysis by
incorporating fluorescent probes into the reaction (Tan et al., 2022). RNA can be amplified in one
step simply by adding reverse transcriptase in the RPA reaction. These features make RPA a
versatile and fast amplification technology that is suitable for portability, and can be applied to
quantify functional genes in field-monitoring applications.

Saxitoxin biosynthesis in dinoflagellates is linked to a family of genes homologous to the

cyanobacterial *sxt* cluster (Kellmann et al., 2008; Stüken et al., 2011). The *sxt*A4 domain is

recognized as essential for toxin production and its expression is strongly correlated with

intracellular toxin content (Geffroy et al., 2021; Murray et al., 2015). This makes the *sxt*A4 gene

domain an appropriate target for quantifying toxic populations rather than simply detecting genus

presence.

In this study, we use the HAB forming, saxitoxin-producing dinoflagellate genus *Alexandrium* as

a case-study, to demonstrate the utility of qRT-RPA for RNA-based quantification of

metabolically active toxigenic cells. We present optimization strategies for qRT-RPA to achieve

an equivalent performance across four *Alexandrium* species, despite minor sequence variation. We

the use *Alexandrium minutum* as a reference species to benchmark limits of detection and standard

curves for sample quantification. We demonstrate the assay performance in realistic conditions

using simulated samples. To our knowledge, this is the first report of an RNA-based quantitative

RPA assay for a harmful algal bloom species. We therefore provide a methodological foundation

for transitioning HAB molecular detection into portable-ready, rapid-response monitoring,

positioning RPA as a robust approach for HAB surveillance and early warning.

116   2   Materials and Methods

2.1 Culture maintenance and growth experiments

Culture isolates of dinoflagellates and other phytoplankton species were obtained from the Roscoff

Culture Collection (RCC, France), the Culture Collection of Algae and Protozoa (CCAP, UK), and

the Culture Collection of Harmful and Environmental Marine Microalgae (CCVIEO, Spain).

Strains were maintained in L1 culturing media (CCAP, UK) in polypropylene flasks (Starstedt),

at 18°C, with a light : dark cycle of 16 : 8 hours and an irradiance of approximately 100 μmol m^{-2}

s^{-1}. Cells were kept in exponential growth phase by frequent subculturing and monitoring the cell

concentration using a Sedgewick-Rafter counting chamber and a brightfield microscope (Olympus

CX-43).

2.2 Nucleic acid isolation

Cell material was collected for nucleic acid extraction by either pelleting by centrifugation at 4000g for 10 mins, or by vacuum filtration onto a 3.0 μm pore size polycarbonate membrane (Nuclepore, Whatman, USA). Samples were flash frozen in liquid nitrogen and stored at -80 C until further processing. Immediately prior to nucleic acid isolation, cells were disrupted using liquid nitrogen and grinding with a pestle. DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Netherlands). RNA was extracted using the GRS Total RNA kit – plant (GRiSP, Portugal), residual DNA was removed from RNA using Turbo DNAse treatment (Thermo Fisher Scientific, USA) and re-purified using the Monarch RNA cleanup kit (NEB, USA). DNA and RNA purity was assessed spectrophotometrically (Nanodrop One, Thermo Fisher Scientific, USA) and quantity was measured using a Qubit 4 instrument (Thermo Fisher Scientific, USA). Additionally, RNA integrity was also assessed using a Bioanalyzer 2100 with Plant RNA pico reagents (Agilent, USA) electrophoresis.

2.3 RPA primer and probe design

*Sxt*A4 gene sequences from all dinoflagellate species were retrieved from GenBank and aligned through NCBI Blastn. A consensus sequence was generated for each species using EMBOSS cons (https://www.ebi.ac.uk/jdispatcher/msa/emboss_cons). The most conserved regions for assay design covering all *Alexandrium* species were identified manually.

RPA primers (Eurofins, Germany and IDT, USA) were designed according to TwistDx's RPA assay design manual (GC content: 20–70%, length: 30–35 nt, Tm: 50–100 °C), using Primer-BLAST on *Alexandrium minutum sxt*A4 gene (GenBank accession number KM438016.1). Primer candidates were evaluated for 1) potential self- and cross-dimer formation using the Multiple Primer Analyzer (Thermo Fisher Scientific), 2) stable secondary structures using Mfold (unafold.org) and 3) cross-reactivity using Blastn. Where potential primer dimers were detected at the 5' or 3' end, the primer sequence was truncated by 1-2 nucleotides.

The RPA probe was designed according to TwistDx's RPA assay design manual and produced by
Biomers (Germany). The design criteria were expanded based on the literature to specifically
minimize the G-content of the probe and reduce the distance between fluorophore and quencher
to a minimum (Liu et al., 2019). A custom script identified regions of 47-nucleotides on both the
forward and reverse strands of the target sequence, ensuring a thymine was at position 31 as
recommended by the manufacturer. Probes predicted by the Multiple Primer Analyzer to form
dimers on the abasic site were excluded to prevent non-specific fluorescence emission. The final
selected primers and probe are shown in **Table 1**.

2.4 PCR amplification of *Sxt*A4 from gDNA of cultured isolates

To confirm the presence of the *sxt*A4 gene in *Alexandrium* cultured strains VGO0650, VGO0657,
CCAP 1119/48 and FE126, we used a touch-up PCR protocol utilizing *sxt*A4 specific primers
(Stüken et al., 2011) (**Table 1**). All PCRs were run at 25 µL reaction volumes containing 1x GoTaq
G2 green Master Mix (Promega, USA), 5% v/v DMSO, 0.4 µM Forward and Reverse primers.
PCR cycling was carried out as follows: 95°C for 3 minutes, 5 x (95°C for 30s, 58°C for 45s, and
72°C for 50s), 5 x (95°C for 30s, 60°C for 45s, and 72°C for 50s), 27 x (95°C for 30s, 62°C for
45s, and 72°C for 50s), 72°C for 5 minutes. Results were visualized by electrophoresis on a 1.5%
agarose gel.

Table 1. Primer and probe sequences used in this study.

Primers/Probe	Amplification	Sequence (5'-3')	Source
AlexU_SxtA4_3256F	RPA	AGACGGCMGCCAGGCTCTGCAAGAAGGG	This study
AlexU_SxtA4_3301F	RPA	TGAAGCACAACGACACGGAGCAGCTYGAG	This study
AlexU_SxtA4_3565R	RPA	CCCGGCGTAGATGATGTTGTTCTCCGCGT	This study
AlexU_SxtA4_3396Pr	RPA	GACGGAGAGCTCGCCGACTTGCCCGCCATA[dT-FAM][dSpacer][BMN-Q3535]TGCTTGTTTGAGGCC[Spacer C3]	This study
AlexU_SxtA4_3264F	qPCR	GCCAGGCTCTGCAAGAAGGG	This study

AlexU_SxtA4_3410R	qPCR	CACATATGGCGGGCAAGTCG	This study
sxt007	PCR	ATGCTCAACATGGGAGTCATCC	(Stüken et al., 2011)
sxt008	PCR	GGGTCCAGTAGATGTT GACGATG	(Stüken et al., 2011)

Where *sxt*A4 was detected, we repeated the PCR in triplicate with a high-fidelity polymerase to

recover the amplicons for sequencing in order to verify their identity against the reference sequence

of the strain on NCBI. All PCRs were run in 25 μL reaction volumes containing 1x KAPA HiFi

HotStart ReadyMix (Roche, Switzerland), 5% v/v DMSO, 0.4 μM Forward and Reverse primer

and water adjusted for the volume of the template. The PCR cycle was: 95°C for 3 minutes, 35 x

(95°C for 20s, 60-65°C for 15s, and 72°C for 50s), 72°C for 2 minutes. After gel electrophoresis

the bands were excised, purified using the NucleoSpin Gel and PCR Clean-up kit (Macherey-

Nagel, Germany), both strands were sequenced by Genewiz (Azenta Life Sciences, USA).

2.5 Recombinase Polymerase Amplification

Quantitative RPA (qRPA) and Reverse Transcriptase-RPA (qRT-RPA) reactions were performed

following the manufacturer's instructions for the TwistAmp Exo kit (TwistDx, Cambridge, UK).

Real-time, exonuclease-assisted RPA reactions were run at 40°C, at a final volume of 25 μL per

reaction and detected on a CFX Opus 96 Real-Time qPCR system (BioRad, USA) at 15 second

intervals. Initially, RPA reactions were setup following the manufacturer's instructions and further

optimized as shown in section 3.2. Briefly, for a 50 μL reaction which was then split in half, a

rehydration master mix was prepared containing 29.5 μL rehydration buffer, variable

concentrations of forward primer (240 – 600 nM), variable concentrations of reverse primer (240

– 600 nM), variable concentrations of probe (90 – 150 nM), 1 μL Superscript IV for RNA templates

(Thermo Fisher Scientific, USA) and water up to the required volume. The freeze-dried enzyme

pellet was reconstituted with rehydration master mix and split into two individual 25 μL reactions,

followed by template addition. MgOAc at variable concentrations (14 – 30 mM) were added to the

flat cap of each reaction. The introduction of MgOAc into each reaction and the subsequent mixing

were performed by simultaneously spinning down, vortexing and spinning down again all

reactions together, before being placed in the qPCR instrument. Measurements of Time to
Positivity (TTP) were calculated as the timepoint at which the fluorescent signal crossed the
fluorescence threshold, using OriginPro's (2024) Intersect Gadget. The threshold for each reaction
was set as the mean fluorescence signal plus three times the standard deviation of all the negative
(no template) control measurements.

2.6 Assay performance across multiple *Alexandrium* species

Firstly, the assay's performance was tested across four species *A. tamarense* (RCC292)*, A.*
ostenfeldii (CCAP 1119/53)*, A. pacificum* (CCAP 1119/52)*, A. minutum* (VGO0657). For this test, we
used *sxt*A4 amplicons generated from the cultured cells and amplified by qRPA (omitting reverse
transcriptase). This step reveals any difference in amplification based on the target sequence,
without requiring costly production of synthetic RNA for each representative species, with minor
sequence differences. The amplicon concentrations were determined spectofluorometrically
(Qubit, Invitrogen, USA) and serially diluted for testing over 4 orders of magnitude. A synthetic
DNA fragment of the target region representing *A. minutum* consensus sequence, plus 20 bp
flanking either side (IDT, USA) was also included for additional quantification accuracy.

2.7 Assay sensitivity, quantitative range and specificity

The assay was benchmarked for RNA-based amplification using *A. minutum* as a representative
species. To assess the limit of detection (LoD) and quantitative range, a synthetic RNA fragment
corresponding to the target region of the *A. minutum* consensus sequence (produced by Genewiz,
UK) was used as input for a dilution series spanning 6 orders of magnitude. The equivalent test
was then performed using total RNA of *A. minutum* VGO0657. Field-relevant RNA inputs were
tested over 4 orders of magnitude spanning 0.1 – 100 ng per reaction, without prior information
on the *sxt*A4 transcript content of the cells.

For specificity testing we included four strains of *Alexandrium minutum* (FE126, CCAP 1119/48,
VGO0657, VGO0650) as positive controls, another SXT producing dinoflagellate *Gymnodinium*
catenatum (VGO1486), and four representative non-dinoflagellate phytoplankton species. For

these non-dinoflagellate species, genomic DNA was tested directly. All tests were performed using

inputs of 10 ng reaction^{-1}.

2.8 Determination of *sxt*A4 transcripts per cell in *A. minutum* strains

We determined the number of *sxt*A4 transcripts per cell under nutrient replete condition in three

cultured strains with different cell volume and growth rate profiles (strains VGO0657, CCAP

1119/48 and FE126). Growth rates were calculated by bi-daily counting of triplicate culture flasks

in the middle of the light phase when cells were not dividing. The specific growth rate μ (cells day$^-$

^{1}) during the exponential phase was calculated using the formula $\mu = (\ln(N_1)-\ln(N_0)) / (t_1 - t_0)$

(Anderson et al., 1990), where N denotes cell concentration and t denotes day of culture between

two timepoints. Cells were then kept in exponential growth phase for at least 10 generations for

stable physiology. Cells were collected for RNA extraction in mid-exponential phase using 5 μm

pore size polycarbonate filters to exclude bacterial cells (Cytiva, US). Cell size of the 3 strains was

also measured using an Operetta CLS High-Content Analysis System (Revvity).

From these samples we determined total RNA cell^{-1} as well as *sxt*A4 transcripts cell^{-1} using both

our qRT-RPA assay as described above and qRT-PCR primers that we specifically designed for

this purpose (Table 1). The qRT-PCR primers uSxtA4_3264F - uSxtA4_3410R, overlap with the

RPA probe and reverse primer for comparability with the RPA assay, and amplify with 86%

efficiency based on a standard curve analysis. The 20 μL reactions volumes contained 1x GoTaq

qPCR Master Mix (Promega, UK), 1x GoScript™ RT Mix (Promega, UK), 200nM each primer

and 2μL template. The cycle was: 1x (42°C for 15 minutes), 1x (95°C for 10 minutes), 40x (95°C

for 30s, 64°C for 30s, and 72°C for 30s) on a CFX Opus 96 Real-Time qPCR system (BioRad,

USA).

2.9 Assay performance with in simulated samples

Validation of the assay involved the production of simulated samples, as a bloom did not occur in

our planned sampling area during the study period. We performed three tests: 1) Spiking target

RNA into a mixture of non-target RNA from cultured species, 2) Spiking target RNA into RNA

extracted from natural seawater, and c) spiking cultured cells into natural seawater and then
extracting the total community RNA.

The first test was performed for three *A. minutum* strains (FE126, CCAP 1119/48, VGO0657).
Spike RNA was added to 1 ng reaction^{-1} into a mixture of non-target RNA composed at 10 ng
reaction^{-1}. This background was made of 4 equal parts (2.5 ng reaction^{-1} each) of the green alga
Bathycoccus sp., the coccolithophore *Gephyrocapsa (Emiliania) huxleyi* and the diatoms
Phaeodactylum trichornutum and *Thalassiossira pseudonana*.

In the second test, a dilution series of *A. minutum* CCAP 1119/48 total RNA was performed using
RNA from a natural seawater sample. A 1.5 L water sample was collected from Karteros Beach,
10 km east of Heraklion, Crete, Greece on the 3rd of July 2025, containing non-target dinoflagellate
species including *Peridiniales*, *Gymnodinium*, *Tripos* spp*.*, *Prorocentrales*, *Protoperidinium* and
Gonyaulax, as well as other phytoplankton and microzooplankton. The water sample was filtered
through a 3.0 μm pore size polycarbonate membrane (Cytiva, US), snap frozen in liquid nitrogen
and stored at -80 C until the RNA extraction. Again, this background RNA matrix was kept at a
fixed concentration of 10 ng reaction^{-1}.

Finally, in the third test the assay's performance was validated with spiking *A. minutum* VGO0657
cells in sea water sample. We collected 2 L seawater from Ammoudara beach in Heraklion, Crete,
Greece on the 22nd of November 2025. The sample was rich in diatoms and small naked
dinoflagellates. Two mock samples were created by adding 1,000 and 10,000 cells into 0.4 L of
the natural sample. Samples were filtered through a 3.0 μm pore size polycarbonate membrane
(Cytiva, US) and RNA extraction, DNAse treatment and purification was performed as described
for the cultured isolates. The total RNA was eluted in 10 μL water and 6.5 μL were added to the
amplification reaction.

3 Results

In this study, we developed an RNA-based qRT-RPA assay to quantify expressed *sxt*A4 from *Alexandrium* species. The main steps of the assay development are shown in **Figure 1**. Following a series of optimization to achieve the most efficient amplification, the performance across four dinoflagellate species was evaluated using *sxt*A4 PCR amplicons. Specificity was tested with several non-target species representing different phytoplankton taxa. The remaining characterization focused on *Alexandrium minutum* RNA as a reference species that is relevant for HABs globally, and was physiologically stable in our culture conditions. Amplification range and sensitivity was determined using both total RNA, where *sxt*A4 copies per cell were unknown, as well as synthetic RNA to enable *sxt*A4 transcript quantification across different *A. minutum* strains. Analysis of three strains with differing size, total RNA content and *sxt*A4 transcript abundance provide an approximate relationship of *sxt*A4 transcripts to cell abundance, to demonstrate relevance to HAB thresholds. Finally, to demonstrate the field applicability of the assay, yet in the absence of a bloom during the study period, the assay performance was tested using simulated samples. These were prepared either by adding target RNA into a background RNA from natural samples, or adding target cells to natural seawater and treating it as a simulated field sample.

Figure 1. Key steps in qRT-RPA assay development. (a) Assay performance across *Alexandrium* species using *sxt*A4 amplicons. (b) Quantitative range of the assay using synthetic *sxt*A4 RNA and total *A. minutum* VGO0657 RNA. (c) Calculation of *sxt*A4 transcripts per cell in 3 different *A. minutum* strains using qRT-RPA assay. (d) Mock samples including (i) spiking a dilution series of *A. minutum* CCAP 1119/48 RNA into background RNA and (ii) spiking two concentrations of *A. minutum* VGO0657 cells in sea water sample. The image was created with BioRender.com.

3.1 Primer and probe design

The assay design was targeted to the first ~ 400 bp part of the *sxt*A4 open reading frame which contains a highly conserved region with the least nucleotide polymorphisms among *Alexandrium* species. This is key for a truly "universal" RPA probe design, as even a single mismatch can reduce or preclude target amplification. Mismatches are better tolerated in the primers than in the probe (Liu et al., 2019). Nevertheless, we avoided regions with polymorphisms at the 3' end of the primer or used degenerate bases to encompass possible sequence variation (**Figure 2**). We tested 2 forward primers, in combination with a single reverse and probe. By using amplicons of the *sxt*A4 region from four different species, we could compare amplification based solely on sequence variation, by conducting tests at a single template input concentration of 10^6 copies per reaction.

Despite mismatches in the 3' end of the forward primer AlexU_SxtA4_3301F, *sxt*A4 amplified from all *Alexandrium* species. However, TTP differed among species and those with no mismatches amplified faster. The primer and probe combination AlexU_SxtA4_3256F / 3565R / 3396Pr amplified a 309 bp region from all species with an equal TTP. This was selected as the Universal qRT-RPA assay and taken through further optimization.

Figure 2. qRT-RPA assay design and testing. a) Nucleotide alignment of *Alexandrium* and *Gymnodinium catenatum sxt*A4 sequences from NCBI, along with the tested universal *sxt*A4 primers and probes. Numbers in brackets denote the number of sequences used to generate the consensus sequence for each species. Multiple alleles are represented as degenerate bases. The probe fluorophore and quencher positions are highlighted in orange and blue, respectively; the dash represents the abasic site. b) qRT-RPA amplification curves using and AlexU_SxtA4_3256F / AlexU_SxtA4_3565R / 3396Pr. c) qRT-RPA amplification curves using AlexU_SxtA4_3301F / AlexU_SxtA4_3565R / 3396Pr. The notation Syn denotes synthetic RNA generated based on the *A. minutum* sequence. All amplification tests were done using amplicons of the target regions at 10^6 copies reaction^{-1}.

3.2 qRT-RPA assay optimization

To maximise assay sensitivity and speed, we performed optimization tests for the reaction temperature, primer and probe concentrations, and MgOAc levels, based on the ranges of tolerance

suggested by the manufacturer. These tests were performed using the synthetic DNA template
representing *Alexandrium minutum sxt*A4, prior to inclusion of the reverse transcriptase needed
for RNA amplification. The template input was invariable 10^4 copies per reaction, which was near
the initial limit of detection using RPA's default recommended parameters.

We performed optimization tests for 5 parameters individually using a sequential approach. At
each step we selected the optimal condition and used it in the next optimization step – the
conditions were tested in the order they are presented in **Figure 3a-f**. The final optimized assay
consisted of the highest recommended probe concentration for a maximum fluorescence signal,
the lowest possible primer concentrations that did not compromise amplification signal to
minimize primer-dimer formation, the default MgOAc concentration a temperature of 40 °C which
is favorable for the reverse transcription enzyme.

We selected the conditions yielding maximum signal intensity as the optimization strategy, rather
than the quickest TTP. This accounts for the need to maintain a detectable signal at low
concentrations for enhanced sensitivity. Also, it enables the assay to be run on portable devices
that have less sensitive fluorescence detectors than laboratory-based qPCR cyclers. The final
protocol for a 50 μL reaction was as follows: 29.5 μL rehydration buffer, 2.1 μL of 10 μM forward
primer (420 nM final concentration), 1.2 μL of 10 μM reverse primer (240 nM final concentration),
0.75 μL 10 μM probe (150 nM final concentration) and 1 μL Superscript IV for RNA templates
(Thermo Fisher Scientific, USA). These conditions resulted in the TTP becoming faster by 2.5
mins and a 3.3-fold increase in the maximum signal intensity (**Figure** 3**f**).

Figure 3. Assay optimization for different parameters using 10^4 copies of *Alexandrium minutum* synthetic DNA template. (a) Temperature (b) Forward primer concentration (c) Reverse primer concentration (d) Magnesium acetate concentration (e) Probe concentration (f) Comparison between initial conditions based on manufacturer's instructions and final optimized assay conditions. The coloured halo of each curve represents the SD from 3 technical replicates.

3.3 Amplification efficiency across multiple *Alexandrium* species

Assay optimization is normally initiated using gDNA as it is a more stable molecule and the type and amount of reverse transcriptase can be separately optimized. However, the *Alexandrium sxt*A4 does not amplify efficiently from gDNA, normally requiring addition of DMSO into PCR reaction. This not compatible with qRPA chemistry, making this approach unsuitable for *sxt*A4 quantification (**Figure S1**).

The performance of the optimized qRPA assay was evaluated using *sxt*A4 amplicons spanning a longer section of *sxt*A4 that contained the RPA assay target region. These were generated from four *Alexandrium* species: *A. tamarense* (RCC292), *A. ostenfeldii* (CCAP 1119/53), *A. pacificum* (CCAP 1119/52) and *A. minutum* (VGO0657) (**Figure 4**). This approach enabled direct comparison of amplification efficiency based solely on sequence variation, using accurately quantified inputs. Under optimized conditions, the assay showed uniform amplification kinetics across the four

species and over four orders of magnitude, retaining quantitative performance in serial dilutions.

The LoD was 10^3 copies reaction^{-1} with the exception of the *A. tamarense* amplicon which failed

to amplify below 10^4 copies reaction^{-1}. Overall, these results demonstrate robust, cross-species

quantitative performance over a wide dynamic range.

3.4 Sensitivity and quantitative range

Following the initial assay optimization using synthetic DNA and amplicon templates, the

remaining assay characterization was performed on RNA templates using *Alexandrium minutum*

as a reference strain. We assessed analytical sensitivity for RNA using both total RNA from *A.*

minutum VGO0657 and synthetic *sxt*A4 RNA as a quantification standard (**Figure 4**). The assay's

quantification range spans at least 4 orders of magnitude and the limit of detection is 10^3 synthetic

RNA copies and 0.1 ng of *A. minutum* VGO0657 total RNA. Furthermore, the assay run time is

fast, with the lowest concentrations of total RNA amplifying in under 12 minutes.

Figure 4. Characterization of the *sxt*A4 qRT-RPA universal assay. (a) Serial dilutions of isolated

amplicons from four different *Alexandrium* species (*A. tamarense* (RCC292), *A. ostenfeldii* (CCAP

1119/53), *A. pacificum* (CCAP 1119/52), *A. minutum* (VGO0657)). (b) Standard curves of *A.*

minutum synthetic *sxt*A4 RNA (black) and *A. minutum* VGO0657 total RNA (blue). Error bars

represent the standard deviation of 3 technical triplicates.

3.5 Assay specificity

We tested the RNA-based assay specificity using four strains of *A. minutum* as positive controls,

the non-target saxitoxin-producing dinoflagellate *Gymnodinium catenatum* (VGO1486), and other

non-target cosmopolitan microalgal species representing green algae, coccolithophores and

diatoms (**Table 2**). Amplification in other *Alexandrium* species was already shown in previous

tests using amplicons amplified from *sxt*A4 genomic DNA; active *sxt*A4 expression in these

cultures was not tested due to unsatisfactory physiological state in our growth conditions. At the

RNA level, the *sxt*A4 transcript was not detected in any non-target species.

Table 2. *Alexandrium sxt*A4 qRT-RPA assay specificity.

Species	Strain ID	Phylum	sxtA4 RNA
Alexandrium minutum	VGO0650	Dinoflagellata	+
	VGO0657		+
	CCAP119/48		+
	FE126		+
Gymonodinium catenatum	VGO1486	Dinoflagellata	-
Bathycoccus sp.	RCC4222	Chlorophyta	-
Emiliania huxleyi	RCC6241	Haptophyta	-
Phaeodactylum trichornutum	CCAP 1055/1	Heterokontophyta	-
Thalasiossira pseudonana	CCMP 1355	Heterokontophyta	-

3.6 Relationship of *sxt*A4 transcripts to cell abundance

Trigger limits for HAB monitoring programs generally refer to specific cell concentrations.

Therefore, we sought to equate our RNA-based assay results to equivalent cell number. For this

purpose, we determined the number of *sxt*A4 transcripts per cell for three *A. minutum* strains

VGO0657, CCAP 1119/48 and FE126 which had differing growth rates and cell size (**Table 3,**
Figure S2). Cells were collected during the exponential growth phase and samples of 100,000
cells were collected by filtration for each strain. We measured cell size using and *Operetta CLS*
High-Content Analysis System, total RNA cell^{-1} by scaling RNA yield to the cell input, and
transcript number cell^{-1} by qRT-RPA for defined total RNA amounts with their originating cell
numbers. Cell diameter and cell volume varied 1.8-fold and 3.4-fold, respectively. The total RNA
content, transcripts pg^{-1} RNA and transcripts cell^{-1}, all increased with increasing cell size.
Considering the qRT-RPA assay LoD of 0.1 ng reaction^{-1} *A. minutum* VGO0657 total RNA, this
equates to the detection of 6 cells reaction^{-1}.

Table 3. Cell size, total RNA content and *sxt*A4 transcript abundance analysis based on qRT-RPA
for three *A. minutum* strains. Standard deviation is based on 3 replicates.

Strain	Cell diameter (μm)	Cell volume (μm^3)	Growth rate (day^{-1})	Total RNA (pg cell^{-1})	Transcripts pg^{-1}	Transcripts cell^{-1}
FE126	14.7 ± 1.7	1701 ± 557	0.19	5.1 ± 0.68	3.7 ± 2.2	19 ± 1
CCAP1119/48	16.8 ± 2.4	2625 ± 1025	0.16	9.1 ± 0.52	6.8 ± 1.2	62 ± 11
VGO0657	22.0 ± 3.2	5718 ± 2226	0.15	15.9 ± 1.96	7.0 ± 0.7	112 ± 11

3.7 Assay performance in complex samples

We evaluated the influence of background RNA on *sxt*A4 RNA quantification by using
sequentially more complex mock samples (**Figure 5**). Spiking 1 ng RNA from three *A. minutum*
strains into 10 ng background RNA from a mixture of phytoplankton species (**Figure 5a**), showed
a small degree of inhibition delaying the assay reaction by 1.7 minutes, on average, for the 3
strains. Based on the equations of VGO0567 synthetic and total RNA standard curves (**Figure 4b**),
this would cause a quantitative underestimation of one order of magnitude.

To further test this in more realistic conditions, we used 10 ng total RNA from a natural sample,
spiked with *A. minutum* total RNA ranging 0.1 – 10 ng reaction^{-1}. For this test we used *A. minutum*

CCAP 1119/48, which showed healthier growth at the time of the experiment. Also, it is the

representative strain with average size and *sxt*A4 transcript content. In this test, no inhibitory effect

was observed near the assay's LoD at 0.1 ng, however, amplification was more variable resulting

in a higher standard deviation. The final LoD of the assay in complex samples was therefore set

as 0.1 ng reaction^{-1} of *A. minutum* CCAP 1119/48 total RNA. According to the information in

Table 3, this is equivalent to 11 *A. minutum* CCAP 1119/48 cells reaction^{-1}. This LoD is indicative

and expected to vary according to the strain *sxt*A4 transcript content, both intraspecifically and

depending on physiology.

Finally, we tested the assay performance using mock seawater samples spiked with 1,000 and

10,000 cells of *A. minutum* VGO0657 (**Figure 5c**). Detection of 1,000 cells was easily achieved

within the assay's 12 min run time. The TTPs for samples spiked with 1,000 cells and 10,000 were

9.75 and 6.75 mins respectively, which based on the equation shown in **Figure 5b** (blue, with

background RNA) equates to 0.5 ng and 12.9 ng total RNA per reaction respectively. This

approximates one order of magnitude difference in the cell inputs.

Figure 5. Assay performance in complex samples. (a) qRT-RPA of total RNA from *three A.*

minutum strains (CCAP1119/48, FE126, VGO0657) in the presence (blue) and absence (black) of

10 ng background total RNA (b) qRT-RPA of total RNA from *A. minutum* CCAP 1119/48 in the

presence (blue) and absence (black) of 10 ng total RNA isolated from a natural seawater sample.

(c) qRT-RPA of total RNA from 1,000 and 10,000 cells of *A. minutum* VGO0657 spiked in sea

water.

4 Discussion

4.1 Methodological advance

This study demonstrates the utility of qRT-RPA for quantification of active toxigenic *Alexandrium* species. Targeting RNA enables preferential detection of cells that are actively expressing genes required for toxin production, while also improving analytical sensitivity due to the potentially higher abundance of transcripts relative to genomic copies (Geffroy et al., 2021; Murray et al., 2015). Amplification is achieved very fast in less than 12 mins, compared to approximately one hour for qPCR and ddPCR. Importantly, amplification at a constant low temperature around 40 °C, makes qRT-RPA ideal to use with simple portable devices. Furthermore, the tolerance of qRT-RPA to minor interspecific sequence variation, resulted in quantification of *sxt*A4 amplicons representing multiple *Alexandrium* species with equal efficiency. This ensures applicability across different geographic locations, where the causative microorganism might vary or where multiple saxitoxin producing species may coexist. The speed, ease of assay design and energy efficiency makes qRT-RPA ideal for RNA-based testing for toxigenic *Alexandrium* species in the field, enabling functional gene monitoring for early-warning HAB surveillance.

4.2 Performance characteristics

The qRT-RPA assay exhibited a quantitative range spanning four orders of magnitude, with limits of detection of approximately 10^3 synthetic *sxt*A4 RNA copies reaction^{-1} and 0.1 ng total *Alexandrium minutum* RNA reaction^{-1}, corresponding to approximately 10-20 cell equivalents depending on strain. These sensitivities are comparable to those reported for DNA-based qPCR assays targeting the *sxt* gene, with limits of detection ranging from 2.5 to 150 cell equivalents reaction^{-1} (Gao et al., 2015b; Han-Sol et al., 2024; Murray et al., 2019), or <10 copies reaction^{-1} for purified DNA standards (Penna et al., 2015; Savela et al., 2016).

Cross-species testing using *sxt*A4 amplicons revealed uniform amplification kinetics across the *Alexandrium* species tested, despite some differences in the sequence. These results indicate that the assay design successfully minimizes sequence-dependent bias. Furthermore, the assay was highly specific to the target genus, with no amplification in SXT containing *G. catenatum*, other

phytoplankton species, or natural samples containing diverse phytoplankton collected from the
coast of Crete.

By validating assay performance using isolated *sxt*A4 amplicons rather than genomic DNA, we
were able to separate sequence-dependent amplification behavior from confounding effects related
to genome size and variable gene copy number, which are known to complicate molecular
ecological analyses of dinoflagellates (Ruvindy et al., 2023). This approach revealed that the probe
specificity was a key determinant for equivalent performance of qRT- RPA across different
species. This is consistent with previous studies on RPA probe mismatch sensitivity (Liu et al.,
2019).

Background non-target nucleic acids are known to have an inhibitory effect on RPA, most likely
due to non-specific primer binding at the low running temperature (Munawar, 2022; Rohrman and
Richards-Kortum, 2015). In our assay, we found moderate inhibition in the presence of complex
background RNA from cultured phytoplankton, which was expected as one of the primers is
degenerate (Liu et al., 2019). The quantitative underestimation equated to less than one order of
magnitude. However, this effect did not persist when a more complex natural sample was used as
the background RNA. Perhaps a larger diversity of sequences causes a dilution of potentially
interfering sequences. Our results highlight that, standard curves with natural background RNA
are more representative for future quantification of the target in unknown samples, and should be
adopted as a standard practice for qRT-RPA assays. Importantly, reliable detection at low target
concentrations was achieved.

Although translating *sxt*A4 RNA-based detection to cell equivalents is desirable for comparability
to established monitoring programme thresholds, this will vary considerably with strain and
physiological state. Nevertheless, detection of as little as 11 cell equivalents reaction^{-1} from cells
grown in optimal conditions, usually associated with low toxicity (Bui et al., 2024), suggests that
the assay is likely to be highly sensitive for actively toxigenic field populations. Furthermore, this
limit of detection is highly suitable for the target ranges of HAB monitoring programmes with the
strictest alert thresholds from 100 *Alexandrium* cells L^{-1}.

4.3 Comparison with other isothermal amplification approaches

Isothermal amplification strategies for HAB detection have been explored primarily aiming at field deployability. Most reported approaches rely on DNA-based detection with qualitative or endpoint readouts, including RPA or LAMP coupled to lateral flow dipsticks or CRISPR–Cas signal amplification for species identification rather than functional risk assessment (e.g. haptophytes, *Karenia mikimotoi*, *Pseudo-nitzschia multiseries* (Wang et al., 2023; Yao et al., 2024; Yu et al., 2025; Zhang et al., 2024). A notable exception is the use of RPA coupled to Nanopore sequencing, which enables portable, high-resolution taxonomic profiling but remains non-quantitative and DNA-based (Hatfield et al., 2020). Among isothermal methods, the only study to date targeting a toxin biosynthesis gene rather than a ribosomal marker is a LAMP assay targeting *dab*A in *Pseudo-nitzschia multistriata*, demonstrated on a custom-made field-deployable device; however, it is also DNA-based and lacks detailed quantitative and specificity characterization (Alrefaey et al., 2026). In contrast, the qRT-RPA assay presented here uniquely combines isothermal amplification, transcript-level detection and real-time quantification, making it well suited for early-warning applications where speed, functional relevance and portability are prioritized.

4.4 Conclusion

This study establishes the analytical performance of a universal qRT-RPA assay targeting the *sxt*A4 transcript, enabling rapid, cross-species detection of toxigenic *Alexandrium* with functional relevance and short time-to-result. Validation under controlled conditions, including sequence diversity and increasing matrix complexity, provides a solid methodological foundation for RNA-based isothermal monitoring. As with any emerging molecular approach, targeted field testing with natural bloom samples and direct comparison to qPCR, microscopy and toxin measurements will be required to define operational performance and establish informed risk thresholds. With full compatibility to low-power portable fluorescence readers and sensor-integrated platforms, qRT-RPA is well positioned as a rapid early-warning tool in HAB monitoring programmes.

Acknowledgements

The authors would like to thanks Dr. Nuria Lluch, Dr. Rosa Isabel Figueroa and Dr. Esther Garces at Insitut de Ciencias del Mar, Spain, for contributing *Alexandrium minutum* and *Gymnodinium*

562 *catenatum* cultures. This research was funded by the EU Horizon project AquaBioSens no.

563 101135432.

565 REFERENCES

Alrefaey, A.I., McQuillan, J.S., Schaap, A., Siracusa, F., Cardwell, C.L., Walk, J., Rogers, D.,
Forrester, R., Mowlem, M.C., Robidart, J.C., 2026. A lab-on-a-chip system integrating DNA
purification and loop-mediated isothermal amplification for the quantification of the toxic diatom
Pseudo-nitzschia multistriata. Sensors & Diagnostics.
Anderson, D.M., Alpermann, T.J., Cembella, A.D., Collos, Y., Masseret, E., Montresor, M., 2012.
The globally distributed genus Alexandrium: Multifaceted roles in marine ecosystems and impacts
on human health. Harmful Algae 14, 10-35.
Anderson, D.M., Fachon, E., Pickart, R.S., Lin B □, P., Fischer, A.D., Richlen, M.L., Uva, V.,
Brosnahan, M.L., McRaven, L., Bahr, F., Lefebvre, K., Grebmeier D □, J.M., Danielson, S.L.,
Lyu, Y., Fukai, Y., 2021a. Evidence for massive and recurrent toxic blooms of Alexandrium
catenella in the Alaskan Arctic. Proc Natl Acad Sci U S A 118(41):e2107387118.
Anderson, D.M., Fensin, E., Gobler, C.J., Hoeglund, A.E., Hubbard, K.A., Kulis, D.M.,
Landsberg, J.H., Lefebvre, K.A., Provoost, P., Richlen, M.L., Smith, J.L., Solow, A.R., Trainer,
V.L., 2021b. Marine harmful algal blooms (HABs) in the United States: History, current status
and future trends. Harmful Algae 102.
Anderson, D.M., Kulis, D.M., Hall, S., Lee, C., 1990. Dynamics and physiology of saxitoxin
production by the dinoflagellates Alexandrium spp, Marine Biology, pp. 511-524.
Bui, Q.T.N., Pradhan, B., Kim, H.-S., Ki, J.-S., 2024. Environmental Factors Modulate Saxitoxins
(STXs) Production in Toxic Dinoflagellate Alexandrium: An Updated Review of STXs and
Synthesis Gene Aspects. Toxins 16(5), 210.
Commission., E., 2021. Commission Implementing Regulation (EU) 2021/1709 of 23 September
2021 amending Implementing Regulation (EU) 2019/627. (Official Journal L 339, p. 84.
Dai, L., Geng, H.X., Yu, R.C., Liu, Y., Zhao, J.Y., Wang, J.X., Zhang, Q.C., Kong, F.Z., Zhou,
M.J., 2020. Distribution of Alexandrium pacificum cysts in the area adjacent to the Changjiang
River estuary, China. Marine Pollution Bulletin 156.
Dai, Y., Yang, S., Zhao, D., Hu, C., Xu, W., Anderson, D.M., Li, Y., Song, X.P., Boyce, D.G.,
Gibson, L., Zheng, C., Feng, L., 2023. Coastal phytoplankton blooms expand and intensify in the
21st century. Nature 615(7951), 280-284.
Erdner, D.L., Percy, L., Keafer, B., Lewis, J., Anderson, D.M., 2010. A quantitative real-time PCR
assay for the identification and enumeration of Alexandrium cysts in marine sediments. Deep-Sea
Research Part II: Topical Studies in Oceanography 57(3-4), 279-287.
Gao, Y., Yu, R.C., Chen, J.H., Zhang, Q.C., Kong, F.Z., Zhou, M.J., 2015a. Distribution of
Alexandrium fundyense and A. pacificum (Dinophyceae) in the Yellow Sea and Bohai Sea. Marine
Pollution Bulletin 96(1-2), 210-219.
Gao, Y., Yu, R.C., Murray, S.A., Chen, J.H., Kang, Z.J., Zhang, Q.C., Kong, F.Z., Zhou, M.J.,
2015b. High specificity of a quantitative PCR assay targeting a saxitoxin gene for monitoring toxic
algae associated with paralytic shellfish toxins in the Yellow Sea. Applied and Environmental
Microbiology 81(20), 6973-6981.
Garneau, M.È., Schnetzer, A., Countway, P.D., Jones, A.C., Seubert, E.L., Caron, D.A., 2011.
Examination of the seasonal dynamics of the toxic dinoflagellate Alexandrium catenella at

Redondo Beach, California, by quantitative PCR. Applied and Environmental Microbiology 77(21), 7669-7680.

Geffroy, S., Lechat, M.-M., Le Gac, M., Rovillon, G.-A., Marie, D., Bigeard, E., Malo, F., Amzil, Z., Guillou, L., Caruana, A.M.N., 2021. From the sxtA4 Gene to Saxitoxin Production: What Controls the Variability Among Alexandrium minutum and Alexandrium pacificum Strains? Frontiers in Microbiology Volume 12 - 2021.

Gobler, C.J., Doherty, O.M., Hattenrath-Lehmann, T.K., Griffith, A.W., Kang, Y., Litaker, R.W., 2017. Ocean warming since 1982 has expanded the niche of toxic algal blooms in the North Atlantic and North Pacific oceans, Proceedings of the National Academy of Sciences of the United States of America. National Academy of Sciences, pp. 4975-4980.

Gribble, M.O., Bennett, B.J., Liddie, J.M., Borchert, W., Pfluger, B.A., Segars, J.S., Keast, J.M., Hans, A., Kikkeri, N.S., Shin, C., Roland, H.B., Hoysala, S., Kohlhoff, J., Alahmad, B., Nagalli, S., Rushton, W., Enevoldsen, H., Bell, M., Fortenberry, D., Harley, J.R., Schneider, A.L.C., 2025. Global epidemiology of paralytic shellfish poisoning: a systematic search literature review. The Lancet Planetary Health, 101271-101271.

Griffith, A.W., Doherty, O.M., Gobler, C.J., 2019. Ocean warming along temperate western boundaries of the Northern Hemisphere promotes an expansion of Cochlodinium polykrikoides blooms. Proceedings of the Royal Society B: Biological Sciences 286(1904).

Han-Sol, K., Taehee, K., Jaeyeon, P., Tae Gyu, P., Jang-Seu, K., 2024. Development of saxitoxin biosynthesis gene sxtB-targeted qPCR assay for the quantification of toxic dinoflagellates Alexandrium catenella (group I) and A. pacificum (group IV) occurring in the Korean coast. Harmful Algae Volume 134,102603.

Hatfield, R.G., Batista, F.M., Bean, T.P., Fonseca, V.G., Santos, A., Turner, A.D., Lewis, A., Dean, K.J., Martinez-Urtaza, J., 2020. The Application of Nanopore Sequencing Technology to the Study of Dinoflagellates: A Proof of Concept Study for Rapid Sequence-Based Discrimination of Potentially Harmful Algae. Frontiers in Microbiology Volume 11 - 2020.

Hatfield, R.G., Bean, T., Turner, A.D., Lees, D.N., Lowther, J., Lewis, A., Baker-Austin, C., 2019. Development of a TaqMan qPCR assay for detection of Alexandrium spp and application to harmful algal bloom monitoring. Toxicon: X 2.

Hayashida, H., Matear, R.J., Strutton, P.G., 2020. Background nutrient concentration determines phytoplankton bloom response to marine heatwaves. Global Change Biology 26(9), 4800-4811.

John, D.E., Wang, Z.A., Liu, X., Byrne, R.H., Corredor, J.E., López, J.M., Cabrera, A., Bronk, D.A., Tabita, F.R., Paul, J.H., 2007. Phytoplankton carbon fixation gene (RuBisCO) transcripts and air-sea CO"2 flux in the Mississippi River plume. ISME Journal 1(6), 517-531.

Kellmann, R., Mihali, T.K., Young, J.J., Pickford, R., Pomati, F., Neilan, B.A., 2008. Biosynthetic intermediate analysis and functional homology reveal a saxitoxin gene cluster in cyanobacteria. Applied and Environmental Microbiology 74(13), 4044-4053.

Kon, N.F., Teng, S.T., Hii, K.S., Yek, L.H., Mujahid, A., Lim, H.C., Lim, P.T., Leaw, C.P., 2015. Spatial distribution of toxic Alexandrium tamiyavanichii (Dinophyceae) in the southeastern South China Sea-Sulu Sea: A molecular-based assessment using real-time quantitative PCR (qPCR) assay. Harmful Algae 50, 8-20.

Lewis, A.M., Coates, L.N., Turner, A.D., Percy, L., Lewis, J., 2018. A review of the global distribution of Alexandrium minutum (Dinophyceae) and comments on ecology and associated paralytic shellfish toxin profiles, with a focus on Northern Europe, Journal of Phycology. Blackwell Publishing Inc., pp. 581-598.

Liu, S., Yang, D.S., Wang, S., Luan, H., Sekine, Y., Model, J.B., Aranyosi, A.J., Ghaffari, R., Rogers, J.A., 2023. Soft, environmentally degradable microfluidic devices for measurement of sweat rate and total sweat loss and for colorimetric analysis of sweat biomarkers. EcoMat 5(1), e12270.

Liu, X., Yan, Q., Huang, J., Chen, J., Guo, Z., Liu, Z., Cai, L., Li, R., Wang, Y., Yang, G., Lan, Q., 2019. Influence of design probe and sequence mismatches on the efficiency of fluorescent RPA. World Journal of Microbiology and Biotechnology 35(6).

Munawar, M.A., 2022. Critical insight into recombinase polymerase amplification technology. Expert Review of Molecular Diagnostics 22(7), 725-737.

Murray, S.A., Diwan, R., Orr, R.J.S., Kohli, G.S., John, U., 2015. Gene duplication, loss and selection in the evolution of saxitoxin biosynthesis in alveolates. Molecular Phylogenetics and Evolution 92, 165-180.

Murray, S.A., Ruvindy, R., Kohli, G.S., Anderson, D.M., Brosnahan, M.L., 2019. Evaluation of sxtA and rDNA qPCR assays through monitoring of an inshore bloom of Alexandrium catenella Group 1. Scientific Reports 9(1).

Murray, S.A., Wiese, M., Stüken, A., Brett, S., Kellmann, R., Hallegraeff, G., Neilan, B.A., 2011. SxtA-based quantitative molecular assay to identify saxitoxin-producing harmful algal blooms in marine waters. Applied and Environmental Microbiology 77(19), 7050-7057.

Parks, R., Bear, E., Coates, L., Maskrey, B., Algoet, M., 2020. Annual report on the results of the Biotoxin and Phytoplankton Official Control Monitoring Programmes.

Penna, A., Perini, F., Dell'Aversano, C., Capellacci, S., Tartaglione, L., Giacobbe, M.G., Casabianca, S., Fraga, S., Ciminiello, P., Scardi, M., 2015. The sxt Gene and Paralytic Shellfish Poisoning Toxins as Markers for the Monitoring of Toxic Alexandrium Species Blooms. Environmental Science and Technology 49(24), 14230-14238.

Rabalais, N.N., Turner, R.E., Sen Gupta, B.K., Boesch, D.F., Chapman, P., Murrell, M.C., 2007. Hypoxia in the Northern Gulf of Mexico: Does the Science Support the Plan to Reduce, Mitigate, and Control Hypoxia?, pp. 753-772.

Rohrman, B., Richards-Kortum, R., 2015. Inhibition of Recombinase Polymerase Amplification by Background DNA: A Lateral Flow-Based Method for Enriching Target DNA. Analytical Chemistry 87(3), 1963-1967.

Ruvindy, R., Barua, A., Bolch, C.J.S., Sarowar, C., Savela, H., Murray, S.A., 2023. Genomic copy number variability at the genus, species and population levels impacts in situ ecological analyses of dinoflagellates and harmful algal blooms. ISME Communications 3(1).

Ruvindy, R., Bolch, C.J., MacKenzie, L., Smith, K.F., Murray, S.A., 2018. qPCR Assays for the Detection and Quantification of Multiple Paralytic Shellfish Toxin-Producing Species of Alexandrium. Frontiers in Microbiology 9.

Savela, H., Harju, K., Spoof, L., Lindehoff, E., Meriluoto, J., Vehniäinen, M., Kremp, A., 2016. Quantity of the dinoflagellate sxtA4 gene and cell density correlates with paralytic shellfish toxin production in Alexandrium ostenfeldii blooms. Harmful Algae 52, 1-10.

Senovilla-Herrero, D., Chen, Y., Liu, C., Suresh, S., Moore, H., McKinney, A., Service, M., Helyar, S., Mbadugha, L., Campbell, K., 2025. Environmental and Biological Factors of Relevance to Shellfish Production in Northern Ireland: Insights From 20 Years of Regional Monitoring Data. Reviews in Aquaculture 17(1), e12998.

Stüken, A., Dittami, S.M., Eikrem, W., McNamee, S., Campbell, K., Jakobsen, K.S., Edvardsen, B., 2013. Novel hydrolysis-probe based qPCR assay to detect saxitoxin transcripts of dinoflagellates in environmental samples. Harmful Algae 28, 108-117.

Stüken, A., Orr, R.J.S., Kellmann, R., Murray, S.A., Neilan, B.A., Jakobsen, K.S., 2011. Discovery
of nuclear-encoded genes for the neurotoxin saxitoxin in dinoflagellates. PLoS ONE 6(5).
Suleiman, M., Jelip, J., Rundi, C., Chua, T.H., 2017. Case report: Paralytic shellfish poisoning in
Sabah, Malaysia. American Journal of Tropical Medicine and Hygiene 97(6), 1731-1736.
Sung, J., Choi, D.H., Lee, Y., Kim, J.-H., Shin, H.H., Kim, Y.-E., Choi, J.H., Noh, J.H., Gobler,
C.J., Park, B.S., 2025. Temperature-Driven Intraspecific Diversity in Paralytic Shellfish Toxin
Profiles of the Dinoflagellate Alexandrium pacificum and Intragenic Variation in the Saxitoxin
Biosynthetic Gene, sxtA4. Microbial Ecology 88(1), 87.
Tan, M., Liao, C., Liang, L., Yi, X., Zhou, Z., Wei, G., 2022. Recent advances in recombinase
polymerase amplification: Principle, advantages, disadvantages and applications, Frontiers in
Cellular and Infection Microbiology. Frontiers Media S.A.
Toebe, K., Alpermann, T.J., Tillmann, U., Krock, B., Cembella, A., John, U., 2013. Molecular
discrimination of toxic and non-toxic Alexandrium species (Dinophyta) in natural phytoplankton
assemblages from the Scottish coast of the North Sea. European Journal of Phycology 48(1), 12-
26.
Turner, A.D., Dhanji-Rapkova, M., Baker, C., Algoet, M., 2014. Assessment of a semiquantitative
liquid chromatography-fluorescence detection method for the determination of paralytic shellfish
poisoning toxin levels in bivalve molluscs from Great Britain. Journal of AOAC International
97(2), 492-497.
union, E.p.a.t.c.o.t.e., 2004. Regulation (EC) No 853/2004 of the European Parliament and of the
Council of 29 April 2004 laying down specific hygiene rules for food of animal origin. Official
Journal of the European Communities.
Vandersea, M.W., Kibler, S.R., Van Sant, S.B., Tester, P.A., Sullivan, K., Eckert, G., Cammarata,
C., Reece, K., Scott, G., Place, A., Holderied, K., Hondolero, D., Litaker, R.W., 2017. QPCR
assays for Alexandrium fundyense and A. ostenfeldii (Dinophyceae) identified from Alaskan
waters and a review of species-specific Alexandrium molecular assays. Phycologia 56(3), 303-
320.
Wang, L., Chen, X., Pan, F., Yao, G., Chen, J., 2023. Development of a rapid detection method
for Karenia mikimotoi by using CRISPR-Cas12a. Frontiers in Microbiology Volume 14 - 2023.
Yao, Y., Luo, N., Zong, Y., Jia, M., Rao, Y., Huang, H., Jiang, H., 2024. Recombinase Polymerase
Amplification Combined with Lateral Flow Dipstick Assay for the Rapid and Sensitive Detection
of Pseudo-nitzschia multiseries. International Journal of Molecular Sciences 25(2), 1350.
Yu, J., Shen, Y., Zhang, Q., Luo, X., Zong, Y., Zhou, C., Huang, H., Jiang, H., 2025. Development
of an RPA-CRISPR/LbaCas12a-Lateral Flow Assay for the Visual Detection of Chrysotila dentata
(Haptophyta). Microorganisms 13(9), 2203.
Yu, Z., Tang, Y., Gobler, C.J., 2023. Harmful algal blooms in China: History, recent expansion,
current status, and future prospects, Harmful Algae. Elsevier B.V.
Zhang, J., Ryderheim, F., Selander, E., Wünsch, U., Kiørboe, T., 2024. Predator-induced defense
decreases growth rate and photoprotective capacity in a nitrogen-limited dinoflagellate,
Alexandrium minutum. Harmful Algae 139, 102734.