Universal rapid RNA-based quantification of toxigenic *Alexandrium* species (Dinophyceae) using quantitative recombinase polymerase amplification

Ioannis Markopoulos[a], Christos Chantzaras[a], Martha Valiadi[a]*, Electra Gizeli[a,b]*

[a] Foundation for Research and Technology Hellas, Institute of Molecular Biology and Biotechnology, Heraklion, Crete, Greece

[b] Department of Biology, University of Crete, Heraklion, Crete, Greece

Underlined authors share senior authorship

*Corresponding authors: (martha_valiadi@imbb.forth.gr; gizeli@imbb.forth.gr)

Abstract

Harmful algal blooms caused by toxigenic *Alexandrium* species pose recurrent risks to coastal ecosystems and public health, yet current monitoring approaches rely on microscopy and laboratory-based toxin analysis with limited capacity for rapid, functional early warning. Here, we present a universal quantitative reverse-transcriptase recombinase polymerase amplification (qRT-RPA) assay targeting the *sxtA4* transcript, an essential gene in saxitoxin biosynthesis. The RPA chemistry is isothermal with a low running temperature, making it suitable for portable, on-site testing. The assay was designed against a conserved *sxtA4* region and validated using synthetic templates, isolated amplicons and total RNA from multiple *Alexandrium* species. Using *Alexandrium minutum* as a reference, the assay achieved a limit of detection below 10^3 synthetic RNA copies and 0.1 ng total RNA per reaction, with uniform amplification kinetics across species and a runtime of less than 15 minutes. The assay selectively detected *sxtA4* transcripts from toxigenic *Alexandrium* strains, showed no cross-reactivity with non-target phytoplankton, and retained it's LoD in complex RNA matrices. These results establish qRT-RPA as a rapid, RNA-

26   based, functionally informative molecular tool that provides a foundation for portable, early-
27   warning monitoring of toxigenic *Alexandrium* blooms.

Keywords

30   *Alexandrium*, saxitoxin, *sxt*A4 RNA, recombinase polymerase amplification, harmful algal
31   blooms

1 Introduction

Harmful Algal Blooms (HABs) are posing increasing ecological and socioeconomic pressure in coastal areas worldwide. Their global expansion is attributed to anthropogenic nutrient enrichment and ocean warming (Dai et al., 2023; Gobler et al., 2017; Griffith et al., 2019; Hayashida et al., 2020; Rabalais et al., 2007; Yu et al., 2023). Warming temperatures can promote bloom initiation (Anderson et al., 2021a) accelerated cell growth rates and prolonged bloom seasons (Anderson et al., 2021a; Gobler et al., 2017; Griffith et al., 2019) especially in temperate regions (Dai et al., 2023), whilst also allowing their intrusion into previously unaffected polar areas (Anderson et al., 2021a). The consequences of HABs range from ecosystem-scale deoxygenation to the accumulation of potent toxins that threaten food security and human health (Anderson et al., 2021a; Anderson et al., 2021b). Despite progress in ocean observing systems, early-warning capacity often remains limited, particularly in regions where toxic events develop rapidly or surveillance infrastructure is sparse.

Most regulatory monitoring programs still rely on a dual structure: (i) microscopic identification and enumeration of phytoplankton in water samples, and (ii) chemical toxin analysis in shellfish tissue [EC 853/2004; Turner 2014; Parks 2020]. The first assesses putatively toxic cell abundance, while the second measures saxitoxin and congeners which are converted to toxicity equivalence factors (TEFs) (Commission Implementing Regulation (EU) 2021/1709, (Commission., 2021)). Concerned countries like the UK follow this regime with practical thresholds of 400 STX eq. (Turner et al., 2014) in shellfish flesh or 40 - 100 *Alexandrium* cells L^{-1} of water triggering intensified monitoring frequency (Parks et al., 2020; Senovilla-Herrero et al., 2025). While these methods underpin long-established monitoring frameworks, they present significant operational constraints. Microscopy cannot reliably distinguish between toxic and non-toxic strains of morphologically similar taxa and provides no functional information on toxin biosynthesis. Meanwhile, toxin chemical analyses by HPLC (Standard EN 14526) requires centralized laboratories, limiting sampling frequency and increasing turnaround time to minimum of 3 days. Together, these limitations hinder early warning and intervention, increasing the likelihood of food contamination and ecological damage.

Molecular amplification-based represent an alternative to microscopy analysis, offering higher
sensitivity, faster time-to-result, species-level resolution and quantification of specific cellular
functions (John et al., 2007; Kon et al., 2015; Murray et al., 2019). A wealth of DNA-based qPCR
assays to enumerate specific toxic species (Dai et al., 2020; Erdner et al., 2010; Gao et al., 2015;
Garneau et al., 2011; Hatfield et al., 2019; Ruvindy et al., 2018; Toebe et al., 2013; Vandersea et
al., 2017). Most assays target the small or large ribosomal genes of specific *Alexandrium* species
to provide abundance measurements, but are not able to assess the populations' toxigenic potential.
Several qPCR assays also target genes of the saxitoxin cluster, to directly assess toxigenic potential
(Gao et al., 2015; Han-Sol et al., 2024; Murray et al., 2019; Murray et al., 2011; Penna et al., 2015;
Savela et al., 2016; Stüken et al., 2013). Nevertheless, the transition from traditional light
microscopy to molecular monitoring for HAB surveillance has been slow to materialize. While
qPCR is the gold standard of molecular detection due to its cost-effectiveness and high throughput,
it is limited to a laboratory setting due to the requirement for high-power consuming thermal
cycling. Therefore, it does not overcome the need for centralized analytical facilities and
76
HAB surveillance will significantly benefit from on-site or *in situ* monitoring offering real-time
detection. In contrast to qPCR, isothermal amplification can achieve molecular quantification in
remote field settings, using portable battery-powered analyzers (e.g. Pebble R, BIOPIX-T;
TwistAmp, TwistDx). In particular, recombinase polymerase amplification (RPA) operates at a
low amplification at a low, stable temperature (37 – 42 ºC), has high tolerance to impurities and a
fast time-to-result in less than 15 minutes (Tan et al., 2022). RPA chemistry can be used for
qualitative detection using lateral flow strips (Liu et al., 2023; Yao et al., 2024; Yu et al., 2025)
and quantitative analysis by incorporating fluorescent probes into the reaction, with run times
below 15 minutes (Tan et al., 2022). Also, RNA can be amplified in one step simply by adding
reverse transcriptase in the RPA reaction. These features make RPA a versatile and fast
amplification technology that is suitable for portability, that can be applied to quantify functional
genes in field-monitoring applications.
89
Here, we use the HAB forming, saxitoxin-producing dinoflagellate genus *Alexandrium* as a case-
study, to develop and evaluate a qRPA assay for RNA-based quantification of metabolically active
toxigenic cells. *Alexandrium* species are globally distributed and responsible for recurrent paralytic

shellfish poisoning (PSP) outbreaks (Anderson et al., 2012; Gribble et al., 2025; Lewis et al., 2018;

Suleiman et al., 2017). They produce an array of toxins as secondary metabolites, being most well-

known for the production of saxitoxins that cause Paralytic Shellfish Poisoning (PSP). Saxitoxin

biosynthesis in dinoflagellates is linked to a family of genes homologous to the cyanobacterial *sxt*

cluster (Kellmann et al., 2008; Stüken et al., 2011). The *sxt*A4 domain is recognised as essential

for toxin production and strongly correlated with transcript abundance and intracellular toxin

content (Geffroy et al., 2021; Murray et al., 2015). This makes the *sxt*A4 gene domain an

appropriate target for quantifying toxic populations rather than simply detecting genus presence.

In this study, we present the development, optimization and performance testing of a universal

*Alexandrium sxt*A4 RNA-based qRT-RPA assay. To our knowledge, this is the first report of an

RNA-based quantitative RPA assay for a harmful algal bloom species. We present optimization

strategies for qRT-RPA and quantify analytical sensitivity using synthetic and total RNA

templates. We establish realistic limits of detection and standard curves for sample quantification

using three *A. minutum* strains of differing size, RNA content and *sxt*A4 transcript content. We

demonstrate the assay performance using mock samples, prepared by spiking known amounts of

target RNA into background RNA from natural samples. We therefore provide a methodological

foundation for transitioning HAB molecular detection into portable-ready, rapid-response

monitoring, positioning RPA as a robust approach for HAB surveillance and early warning.

112 2 Materials and Methods

113 2.1 Culture maintenance and growth experiments

114 Dinoflagellate species were maintained in L1 culturing media (CCAP, UK) in polypropylene

115 flasks (Starstedt), at 18°C, with a light : dark cycle of 16 : 8 hours and an irradiance of

116 approximately 100 µmol m^{-2} s^{-1}. Cells were kept in exponential growth phase by frequent

117 subculturing and monitoring the cell concentration using a Sedgewick-Rafter counting chamber

118 and a brightfield microscope (Olympus CX-43).

120 2.2 Nucleic acid isolation

Cell material was collected for nucleic acid extraction by either pelleting by centrifugation at 4000g for 10 mins, or by vaccum filtration onto a 3.0 μm pore size polycarbonate membrane (Nuclepore, Whatman, USA). Samples were flash frozen in liquid nitrogen and stored at -80 C until further processing. Immediately prior to nucleic acid isolation, cells were disrupted using liquid nitrogen and grinding with a pestle. DNA was extracted using the DNeasy Plant Mini Kit (Qiagen, Netherlands). RNA was extracted using the GRS Total RNA kit – plant (GRiSP, Portugal), residual DNA was removed from RNA using Turbo DNAse treatment (Thermo Fisher Scientific, USA) and re-purified using the Monarch RNA cleanup kit (NEB, USA). DNA and RNA purity was assessed spectrophotometrically (Nanodrop One, Thermo Fisher Scientific, USA) and quantity was measured using a Qubit 4 instrument (Thermo Fisher Scientific, USA). Additionally, RNA integrity was also assessed using a Bioanalyzer 2100 with Plant RNA pico reagents (Agilent, USA) electrophoresis.

2.3 RPA primer and probe design

*Sxt*A4 gene sequences from all dinoflagellate species were assembled and aligned through NCBI. For consensus sequence was generated for each species EMBOSS cons (https://www.ebi.ac.uk/jdispatcher/msa/emboss_cons). The most conserved regions for assay design covering all of *Alexandrium* species were initially identified manually.

RPA primers (Eurofins, Germany and IDT, USA) were designed according to TwistDx's RPA assay design manual (GC content: 20–70%, length: 30–35 nt, Tm: 50–100 °C), using Primer-BLAST on *Alexandrium minutum sxtA4* gene (accession number KM438016.1). Primer candidates were evaluated for 1) potential self- and cross-dimer formation using the Multiple Primer Analyzer (Thermo Fisher Scientific), 2) stable secondary structures using Mfold (unafold.org) and 3) cross-reactivity using Blastn. Where potential primer dimers were detected at the 5' or 3' end, the primer sequence was truncated by 1-2 nucleotides.

The RPA probe was designed according to TwistDx's RPA assay design manual and produced by Biomers (Germany). The design criteria were expanded based on the literature to specifically minimize the G-content of the probe and reduce the distance between fluorophore and quencher

150 to a minimum (Liu et al., 2019). A custom script identified regions of 47-nucleotides on both the
151 forward and reverse strands of the target sequence, ensuring a thymine was at position 31 as
152 recommended by the manufacturer. Probes predicted by the Multiple Primer Analyzer to form
153 dimers on the abasic site were excluded to prevent non-specific fluorescence emission. The final
154 selected primers and probe are shown in Table 1.

2.4 Identification and sequencing of *Sxt*A4 from cultured isolates

157 To confirm the presence of the *sxtA4* gene *Alexandrium* cultures, we used touch-up PCR protocol
158 utilizing *sxtA4* specific primers (Stüken et al., 2011) (**Table 1**). All PCRs were run at 25 µL
159 reaction volumes containing 1x GoTaq G2 green Master Mix (Promega, USA), 5% v/v DMSO,
160 0.4 µM Forward and Reverse primers. PCR cycling was carried out as follows: 95°C for 3 minutes,
161 5 x (95°C for 30s, 58°C for 45s, and 72°C for 50s), 5 x (95°C for 30s, 60°C for 45s, and 72°C for
162 50s), 27 x (95°C for 30s, 62°C for 45s, and 72°C for 50s), 72°C for 5 minutes. Results were
163 visualized by electrophoresis on a 1.5% agarose gel.

165 **Table 1.** Primer and probe sequences used in this study.

Primers/Probe	Amplification	Sequence (5'-3')	Source
AlexU_SxtA4_3256F	RPA	AGACGGCMGCCAGGCTCTGCAAGAAGGG	This study
AlexU_SxtA4_3301F	RPA	TGAAGCACAACGACACGGAGCAGCTYGAG	This study
AlexU_SxtA4_3565R	RPA	CCCGGCGTAGATGATGTTGTTCTCCGCGT	This study
AlexU_SxtA4_3355Pr	RPA	GGGCCGACATCACCTACGTGTGCGACGGCG[FAM-dT][dSpacer]T[BMN-Q3535]ACTCCACGGACGGAG[Spacer C3]	This study
AlexU_SxtA4_3396Pr	RPA	GACGGAGAGCTCGCCGACTTGCCCGCCATA[dT-FAM][dSpacer][BMN-Q3535]TGCTTGTTTGAGGCC[Spacer C3]	This study
AlexU_SxtA4_3264F	qPCR	GCCAGGCTCTGCAAGAAGGG	This study
AlexU_SxtA4_3410R	qPCR	CACATATGGCGGGCAAGTCG	This study

sxt007	PCR	ATGCTCAACATGGGAGTCATCC	(Stüken et al., 2011)
sxt008	PCR	GGGTCCAGTAGATGTT GACGATG	(Stüken et al., 2011)

Where *sxt*A4 was detected, we repeated the PCR in triplicate with a high-fidelity polymerase to

recover the amplicons for sequencing. All PCRs were run in 25 µL reaction volumes containing

1x KAPA HiFi HotStart ReadyMix (Roche, Switzerland), 5% v/v DMSO, 0.4 µM Forward and

Reverse primer and water adjusted for the volume of the template. The PCR cycle was: 95°C for

3 minutes, 35 x (95°C for 20s, 60-65°C for 15s, and 72°C for 50s), 72°C for 2 minutes. After gel

electrophoresis the bands were excised, purified using the NucleoSpin Gel and PCR Clean-up kit

(Macherey-Nagel, Germany), both strands were sequenced by Genewiz (Azenta Life Sciences,

USA) and the sequences were validated using NCBI's blastn.

176 2.5 Recombinase Polymerase Amplification

Quantitative RPA and Reverse Transcriptase-RPA (qRT-RPA) reactions were performed

following the manufacturer's instructions for the TwistAmp Exo kit (TwistDx, Cambridge, UK).

Real-time, exonuclease-assisted RPA reactions were run at 40°C, at a final volume of 25 µL per

reaction and detected on a CFX Opus 96 Real-Time qPCR system (BioRad, USA) at 15 second

intervals. Briefly, for the reaction setup, a rehydration master mix was prepared containing 29.5

µL rehydration buffer, 1.2 µL of 10 µM forward primer, 2.1 µL of 10 µM reverse primer, 0.75 µL

10 µM probe and 1 µL Superscript IV for RNA templates (Thermo Fisher Scientific, USA). The

freeze-dried enzyme pellet was reconstituted with rehydration master mix and split into two

individual 25 µL reactions. After template addition, 1.25 µL of 280mM MgOAc were added to the

flat cap of each reaction. The introduction of MgOAc into each reaction and the subsequent mixing

were performed by simultaneously spinning down, vortexing and spinning down again all

reactions together. Measurements of Time to Positivity (TTP) were calculated as the timepoint at

which the fluorescent signal crossed the fluorescence threshold, using OriginPro's (2024) Intersect

Gadget. The threshold for each reaction was set as the mean fluorescence signal plus three times

the standard deviation of all the negative control measurements.

2.6 Determination of *sxt*A4 transcripts per cell in *A. minutum* strains

We determined the number of *sxt*A4 transcripts per cell under nutrient replete condition in three
cultured strains with different cell volume and growth rate profiles (strains VGO 657, CCAP
1119/48 and FE126). Growth rates were calculated by bi-daily counting of triplicate culture flasks
in the middle of the light phase when cells were not dividing. The specific growth rate μ (cells day$^-$
^{1}) during the exponential phase was calculated using the formula $\mu = (\ln(N_1)-\ln(N_0)) / (t_1 - t_0)$
(Anderson et al., 1990), where N denotes cell concentration and t denotes day of culture between
two timepoints. Cells were then kept in exponential growth phase for at least 10 generations for
stable physiology. Cells were collected for RNA extraction in mid-exponential phase using 5 μm
polycarbonate filters to exclude bacterial cells (Cytiva, US). Cell size of the 3 strains was also
measured using an Operetta CLS High-Content Analysis System (Revvity).

From these samples we determined total RNA cell^{-1} as well as *sxt*A4 trancripts per cell using both
our qRT-RPA assay as described above and qRT-PCR primers that we specifically designed for
this purpose. The qRT-PCR primers uSxtA4_3264F - uSxtA4_3410R, overlap with the RPA probe
and reverse primer for comparability with the RPA assay, and amplify with 86% efficiency based
on a standard curve analysis. The 20 μL reactions volumes contained 1x GoTaq qPCR Master Mix
(Promega, UK), 1x GoScript™ RT Mix (Promega, UK), 200nM each primer and 2μL template.
The cycle was: 1x (42°C for 15 minutes), 1x (95°C for 10 minutes), 40x (95°C for 30s, 64°C for
30s, and 72°C for 30s) on a CFX Opus 96 Real-Time qPCR system (BioRad, USA).

2.7 Sensitivity and quantitative range across species

Firstly, the assay's performance was tested across species using the partial *sxt*A4 amplicons
generated initially four sequencing as well as synthetic DNA of the target region plus 20 bp
flanking either side (IDT, USA). The amplicon concentrations were determined
spectofluorometrically (Qubit, Invitrogen, USA) and serially diluted for testing over 5 orders of
magnitude. The assay's quantitative range was then tested using RNA of *A. minutum*, as a
representative *Alexandrium* species. Initially, this was done using dilution series of synthetic RNA

that corresponded to the region of the synthetic DNA (Genewiz), followed by total RNA extracted

from *Alexandrium minutum* VGO657.

2.8 Assay inclusivity and specificity

Inclusivity was tested using four strains of *Alexandrium minutum* and another 4 species of

Alexandrium that are known to produce toxins and one that is putatively non-toxic. Because

Alexandrium genomic DNA was not suitable for amplification by q-RPA, the sequenced served as

a template for testing with q-RPA. Specificity to the genus was tested against another SXT

producing dinoflagellate, two putative SXT producing cyanobacteria, and four representative non-

dinoflagellate phytoplankton species. For these non-dinoflagellate species, genomic DNA was

tested directly at a concentration of 10 ng reaction^{-1} .

2.9 Assay performance with increasing sample complexity

To test whether non-target RNA affected amplification behaviour, we created mock samples where

the target was spiked into a non-target background matrix. Initially, *A. minutum* total RNA from

strains FE126, CCAP1119/48, VGO657 at 1 ng reaction^{-1} was spiked into a mixture of non-target

RNA from cultured cells provided by colleagues (*Bathycoccus sp., Emiliania huxleyi,*

Phaeodactylum trichornutum and *Thalassiossira pseudonana*). The background matrix was

composed of equal amounts of these four species at a fixed concentration of 10 ng reaction^{-1}.

Finally, a dilution series of *A. minutum* CCAP 1119/48 total RNA was performed using RNA from

a natural seawater sample that did not contain *Alexandrium* cells. A 1.5 L water sample was

collected from Karteros Beach, 10km east of Heraklion, Crete, Greece and filtered with a 3.0μm

polycarbonate (Cytiva, US) and snap frozen with liquid nitrogen and stored at -80 C until the RNA

extraction. Again, the background matrix was kept at a fixed concentration of 10 ng reaction^{-1}.

3   Results

3.1 Primer and probe design

The assay design was targeted to the initial ~400bp part of the *sxt*A4 gene which contains a highly conserved region without nucleotide polymorphisms among *Alexandrium* species. This is key for a truly "universal" RPA probe design, as even a single mismatch can reduce or preclude target amplification. Mismatches are better tolerated in the primers than in the probe (Liu et al., 2019). Nevertheless, we avoided regions with polymorphisms at the 3' end of the primer or used degenerate bases to encompass possible sequence variation (**Figure 1**). We tested a total of 2 combinations of primer pairs and 2 probes. By using amplicons of the *sxt*A4 region from four different species, we could compare amplification based solely on sequence variation, by conducting tests at a single template input concentration of 10^6 copies per reaction.

Despite mismatches in the 3' end of the forward primer AlexU_SxtA4_3301F, *sxt*A4 amplified from all *Alexandrium* species. However, TTP differed among species and those with no mismatches amplified faster. Furthermore, Probe 1, did not produce any fluorescence signal in the amplification, possibly due to cross-reactions with the reverse primer. The primer and probe combination AlexU_SxtA4_3256F / 3565R / 3396Pr amplified a 309 bp region from all species with an equal TTP. This was selected as the Universal qRT-RPA assay and taken through further optimization.

Figure 1. qRT-RPA assay design and testing. a) Nucleotide alignment of *Alexandrium* and *Gymnodinium sxt*A4 sequences from NCBI, along with the tested universal *sxt*A4 primers and probes. Numbers in brackets denote the number of sequences used to generate the consensus sequence for each species. Multiple alleles are represented as degenerate bases. The probe fluorophore and quencher positions are highlighted in orange and blue, respectively; the dash represents the abasic site. b) qRT-RPA amplification curves using Probe 1 and AlexU_SxtA4_3256F / AlexU_SxtA4_3565R. c) qRT-RPA amplification curves using Probe 1 and AlexU_SxtA4_3330F / AlexU_SxtA4_3565R. The notation Syn denotes synthetic RNA generated based on the *A. minutum* sequence. All amplification tests were done using amplicons of the target regions at 10^6 copies reaction^{-1}.

3.2 qRT-RPA assay optimization

To maximise assay sensitivity and speed, we performed optimization tests for the reaction temperature, primer and probe concentrations, and MgOAc levels, based on the ranges of tolerance suggested by the manufacturer. The template for all tests was synthetic *Alexandrium minutum sxt*A4 DNA at a concentration of 10^4 copies per reaction, which was near the initial limit of detection using RPA's default recommended parameters.

We performed optimization tests for 5 parameters using a sequential approach instead of a multi-parameter matrix. At each step we selected the optimal condition and used it in the next optimization step – the conditions were tested in the order they are presented in **Figure 2**. The final optimized assay consisted of the highest recommended probe concentration for a maximum fluorescence signal, the lowest possible primer concentrations that did not compromise amplification signal to minimize primer-dimer formation, the default MgOAc concentration a temperature of 40 ℃ which is favorable for the reverse transcription enzyme.

We chose to use maximum signal intensity as the optimization strategy, rather than the quickest TTP. This accounts for the need to maintain a detectable signal at low concentrations for enhanced sensitivity. Also, it enables the assay to be run on portable devices that have less sensitive fluorescence detectors than laboratory-based qPCR cyclers (Figure 2 f or g). These changes resulted in the TTP becoming faster by 2.5 mins and a 3.3-fold increase in the maximum signal intensity.

Figure 2. Universal assay optimization using 10^4 copies of *Alexandrium minutum* synthetic DNA. (a) Temperature (b) Forward primer concentration (c) Reverse primer concentration (d) Magnesium acetate concentration (e) Probe concentration (f) Comparison between initial and optimized assay conditions. The coloured halo of each curve represents the SD from 3 technical triplicates.

3.3 Performance of the assay with different template types

We next evaluated the performance of the assay with different templates types to determine a) any potential effect of reverse transcription step in a one-pot reaction on assay speed and sensitivity, and b) whether the assay can be used for both genomic DNA as a direct cell abundance proxy despite the high GC-content of the gene (Figure S1). Genomic DNA showed very poor amplification in qRPA on genomic DNA, with a low signal intensity and resolution among different template concentrations. Normally, amplification of the *sxt*A4 gene by PCR or by basic, non-quantitative RPA, requires the addition of DMSO due to the high template GC-content. However, this is not with qRPA chemistry (exo RPA kit) and is the most probable reason for inefficient amplification. We therefore concluded that genomic DNA is an unsuitable target for *sxt*A4 qRT-RPA, and focused solely on RNA amplification.

3.4 Sensitivity and quantitative range across multiple species

The optimized assay was then evaluated using isolated *sxt*A4 amplicons from four *Alexandrium*
species (**Figure 3**). This approach enabled direct comparison of amplification efficiency based
solely on sequence variation. Under optimized conditions, the assay showed uniform amplification
kinetics across four species and over four orders of magnitude, retaining quantitative performance
in serial dilutions. The LoD was 10^3 copies per reaction with the exception of *A. tamarense* which
failed to amplify below 10^4 copies per reaction. Overall, these results demonstrate robust, cross-
species quantitative performance over a wide dynamic range.
Following the initial assay optimization using synthetic and amplicon DNA templates, the
remaining assay characterization was performed on RNA templates. We assessed analytical
sensitivity for RNA using two approaches: i) a 10-fold, 6-point dilution series (10^8-10^3 copies per
reaction) using synthetic *sxtA4* RNA and ii) a 10-fold, 4-point dilution series (0.1-100 ng per
reaction) of total RNA, isolated from *A. minutum* VGO 657 (**Figure 3**). The assay's quantification
range spans at least 4 orders of magnitude and the limit of detection is 10^3 synthetic RNA copies
and 0.1 ng of *A. minutum* VGO 657 total RNA. Furthermore, the assay run time is fast, with the
lowest concentrations of total RNA amplifying in under 12 minutes.
331

Figure 3. Characterization of the *sxt*A4 qRT-RPA universal assay. (a) Serial dilutions of isolated amplicons from four different *Alexandrium* species (*A. tamarense, A. ostenfeldii, A. pacificum, A. minutum*). (b) Standard curves of *A. minutum* synthetic sxtA4 RNA (black) and *A. minutum* VGO657 total RNA (blue). Error bars represent the standard deviation of 3 technical triplicates.

3.5 Inclusivity and specificity

To test inclusivity and specificity of the assay we used a panel of marine dinoflagellates of the target genus as well as a non-target saxitoxin-producing dinoflagellate, two putative saxitoxin-producing freshwater cyanobacteria, and other non-target cosmopolitan microalgal species (**Table 2**). The assay successfully detected *sxt*A4 in RNA from all 5 toxic *Alexandrium* strains, as well as *A. tamutum* which is generally regarded as non-toxic. We did not detect *sxt*A4 in toxic *Gymnodinium catenatum* neither by RT-qRPA nor by PCR; this species is generally regarded as toxic, but toxicity in this particular strain had not been assessed by the provider. Of the cyanobacteria species, *Planktothrix agardhii* is known to produce saxitoxins and the *Anabaena* genus contains saxitoxin-producing species, although the toxicity profile of the *A. cylindri*ca strain we used was not tested by the provider. The sxtA4 gene was not detected in neither of these strains, nor in any other non-target phytoplankton species.

353   **Table 2.** *Alexandrium sxt*A4 assay inclusivity and specificity.

Species	Strain ID	Phylum	sxtA4 DNA	sxtA4 RNA
Alexandrium andersonii	RCC10728	Dinoflagellata	-	n.a.
Alexandrium minutum	VGO657	Dinoflagellata	+	+
	CCAP119/48			
	FE126			
	VGO650			
Alexandrium ostenfeldii	CCAP 1119/53	Dinoflagellata	+	n.a.
Alexandrium pacificum	CCAP 1119/52	Dinoflagellata	+	n.a.
Alexandrium tamarense	RCC292	Dinoflagellata	+	n.a.
Alexandrium tamutum	RCC4860	Dinoflagellata	+	n.a.
Gymonodinium catenatum	VGO1486	Dinoflagellata	-	n.a.
Anabaena cylindrica	CCAP 1403/2A	Cyanobacteria	-	n.a.
Planktothrix agardii	CCAP 1459/21	Cyanobacteria	-	n.a.
Bathycoccus sp.	RCC4222	Chlorophyta	n.a.	-
Emiliania huxleyi	RCC6241	Haptophyta	n.a.	-
Phaeodactylum trichornutum	CCAP 1055/1	Heterokontophyta	n.a.	-
Thalasiossira pseudonana	CCMP 1355	Heterokontophyta	n.a.	-

3.6 Relationship of *sxt*A4 transcripts to cell abundance

As trigger limits for HAB monitoring programs generally refer to specific cell concentrations, we
equated our RNA-based assay results to cell concentrations by determining the number of *sxt*A4
transcripts per cell. We used *A. minutum* as a representative species and compared characteristics
across three strains with differing morphology and growth rate. Strains VGO657, CCAP 1119/48

and FE 126 were analyzed for cell size, total RNA yield cell^{-1} and transcript number cell^{-1} using

qRT-RPA (**Table 3, Figure S1**). Cell diameter and volume varied 1.8-fold and 3.4-fold,

respectively. The total RNA content, transcripts pg^{-1} RNA and transcripts cell^{-1}, all increased with

increasing cell size. Considering the qRT-RPA assay LoD of 0.01 ng reaction^{-1} *A. minutum*

VGO657 total RNA, this equates to the detection of 6 cells reaction^{-1}.

Table 3. Size, total RNA yield and *sxt*A4 transcript analysis based on qRT-RPA for three *A.*

minutum strains. Standard deviation is based on 3 replicates.

Strain	Cell diameter (µm)	Cell volume (µm^3)	Growth rate (day^{-1})	Total RNA (pg cell^{-1})	Transcripts pg^{-1}	Transcripts cell^{-1}
FE126	14.7 ± 1.7	1701 ± 557	0.19	5.1 ± 0.68	3.7 ± 2.2	19 ± 1
CCAP1119/48	16.8 ± 2.4	2625 ± 1025	0.16	9.1 ± 0.52	6.8 ± 1.2	62 ± 11
VGO657	22.0 ± 3.2	5718 ± 2226	0.15	15.9 ± 1.96	7.0 ± 0.7	112 ± 11

3.7 Assay performance with increasing sample complexity

We evaluated the influence of background RNA on *sxt*A4 RNA quantification using two types of

mock samples (**Figure 4**). In the first test, 1 ng RNA from three *A. minutum* strains was spiked

into 10 ng background RNA from a mixture of phytoplankton species (**Figure 4a**). These tests

showed that there was a degree of inhibition delaying the TTP by 1.7 minutes, on average for the

3 strains. Based on the equations of synthetic and total RNA standard curves (Figure 3b), this

would cause a quantitative underestimation of one order of magnitude (i.e. it would appear as 0.11

ng if the input was unknown). To further test this the effect of inhibition by more complex

background RNA, we used 10 ng total RNA from a natural sample, spiked with *A. minutum* total

RNA and over a wide range of concentrations (**Figure 4b**). The natural seawater sample contained

several non-target dinoflagellate species including *Peridiniales*, *Gymnodinium*, *Tripos spp.*,

Prorocentrales, *Protoperidinium* and *Gonyaulax,* as well as other phytoplankton and

microzooplankton. For the spike we used total RNA from *A. minutum* CCAP 1119/48, as this was

the representative strain with average size and *sxt*A4 transcript content. In this test, no inhibitory
effect was observed near the assays LoD at 0.1 ng, however, amplification was more variable
resulting in a higher standard deviation. The final LoD of the assay in complex samples was
therefore set as 0.1 ng reaction^{-1} of *A. minutum* CCAP 1119/48 total RNA which is equivalent to
11 *A. minutum* cells reaction^{-1}.

Figure 4. Assay performance in complex samples. 4a) qRT-RPA of total RNA from *three A.*
minutum strains (FE126, CCAP1119/48, VGO657) in the presence and absence of 10 ng
background total RNA 4b) qRT-RPA of total RNA from *A. minutum* CCAP 1119/48 in the
presence of 10 ng total RNA isolated from a natural seawater sample. In both cases, the
amplification curve from synthetic RNA is used for comparison (black).

4 Discussion

4.1 Methodological advance

This study demonstrates that qRT-RPA can robustly quantify *sxt*A4 transcripts across multiple
Alexandrium species, with comparable amplification kinetics over several orders of magnitude
despite interspecific sequence variation. This assay design ensures applicability across different
geographic locations, where the causative microorganism might vary or where multiple saxitoxin
producing species may coexist. Furthermore, as qRT-RPA is compatible with simple portable
devices, our assay will enable RNA-based testing for toxigenic *Alexandrium* species in the field.

By validating assay performance using isolated *sxt*A4 amplicons rather than genomic DNA, we

were able to separate sequence-dependent amplification behavior from confounding effects related

to genome size and variable gene copy number, which are known to complicate molecular

ecological analyses of dinoflagellates (Ruvindy et al., 2023). This approach revealed that probe–

template complementarity, rather than copy number variation, is the primary determinant of cross-

species performance in probe-based RPA, consistent with previous studies on mismatch sensitivity

in fluorescent RPA chemistries (Liu et al., 2019).

Targeting RNA enables preferential detection of metabolically active cells with the capacity for

toxin production, while also improving analytical sensitivity by exploiting the higher abundance

of transcripts relative to genomic copies (Geffroy et al., 2021; Murray et al., 2015). Furthermore,

the poor performance of genomic DNA in *sxt*A4 q-RPA reinforces RNA as the appropriate target

when DMSO-dependent amplification is incompatible with assay chemistry. Importantly, the

RNA-based assay achieved robust quantitative performance in less than 15 minutes, significantly

beyond conventional qPCR-based laboratory workflows. This demonstrates the suitability of

isothermal functional gene monitoring for early-warning HAB surveillance.

4.2 Performance characteristics

The qRT-RPA assay exhibited a quantitative range spanning four orders of magnitude, with limits

of detection of approximately 10^3 synthetic *sxt*A4 RNA copies reaction^{-1} and 0.1 ng total

Alexandrium minutum RNA reaction^{-1}, corresponding to approximately 10 cell equivalents

depending on strain. These sensitivities are comparable to those reported for qPCR-based saxitoxin

gene assays (Gao et al., 2015; Han-Sol et al., 2024; Murray et al., 2019; Yu et al., 2025) while

being achieved with substantially shorter runtimes. Cross-species testing using *sxt*A4 amplicons

revealed uniform amplification kinetics across most *Alexandrium* species, with a lower sensitivity

only seen for *A. tamarense* despite no mismatches with the probe and primer sequences indicating

a potentially strain-specific result. On the whole, our results indicate that the assay design

successfully minimizes sequence-dependent bias.

Inclusivity testing confirmed *sxt*A4 detection in all tested toxic *Alexandrium* strains and in *A.*
tamutum, a species previously reported to express *sxt*A transcripts despite the absence of
measurable saxitoxin production (Vingiani et al., 2020). This observation is consistent with earlier
work showing that *sxt*A4 expression is necessary but not sufficient for toxin accumulation and
reinforces the need to interpret transcript-based signals as indicators of toxigenic potential rather
than confirmed toxicity (Geffroy et al., 2021). In contrast, no amplification was observed for
Gymnodinium catenatum, despite its established toxigenicity and presence of *sxt*A4 (Muhammad
et al., 2025). This is consistent with the four mismatches within the probe-binding region that result
in amplification failure and make the assay specific to the genus *Alexandrium.*

Background non-target nucleic acids are known to have an inhibitory effect on RPA, most likely
due to non-specific primer binding at the low running temperature (Munawar, 2022; Rohrman and
Richards-Kortum, 2015). In our assay, we found moderate inhibition in the presence of complex
background RNA from cultured phytoplankton. This was expected as one of the primers is
degenerate, however, the probe design is considered more important for specificity (Liu et al.,
2019). The quantitative underestimation equated to less than one order of magnitude. However,
this effect did not persist when a more complex natural sample was used as the background RNA.
Perhaps a larger diversity of sequences causes a dilution of potentially interfering sequences. Our
results highlight that standard curves with natural background RNA are more representative for
future quantification of the target in unknown samples, and should be adopted as a standard
practice for qRT-RPA assays. Importantly, reliable detection at low target concentrations equating
to as little as 11 cell equivalents reaction^{-1} was retained under these conditions. To achieve
quantification of 100 *Alexandrium* cells L^{-1}, this would require that RNA from 1L of a water
sample be extracted and eluted into 50 μL water, of which 5 μL can be used for amplification. This
is in line with standard molecular biology protocols, making the assay highly suitable for HAB
monitoring programmes.

4.3 Comparison with qPCR- and isothermal-based approaches

Quantitative PCR assays for *Alexandrium* are well established and have demonstrated high sensitivity and strong correlations with toxin concentrations and microscopy, particularly when targeting functional genes of the saxitoxin biosynthesis pathway rather than ribosomal markers. Early LSU-based assays achieved very low detection limits but could not distinguish toxic from non-toxic strains (Garneau et al., 2011), motivating the development of qPCR assays targeting *sxt* genes. Universal and species-inclusive *sxt*A4 qPCR assays have since shown detection limits on the order of 30–50 cell equivalents L^{-1} and tighter correlations with toxin levels than rDNA-based assays (Murray et al., 2019; Savela et al., 2016; Stüken et al., 2013), while other toxin-gene targets such as *sxt*A1 and *sxt*B have similarly improved correspondence between molecular signal and toxicity at low cell abundances (Kim et al., 2024; Penna et al., 2015). Despite these strengths, qPCR and RT-qPCR workflows remain constrained to laboratory settings due to thermal cycling, multi-step processing and longer time-to-result, limiting their suitability for rapid, on-site decision making.

A growing body of work has therefore explored isothermal amplification strategies for HAB detection, primarily aiming at field deployability. Most reported approaches rely on DNA-based detection with qualitative or endpoint readouts, including RPA or LAMP coupled to lateral flow dipsticks or CRISPR–Cas signal amplification for species identification rather than functional risk assessment (e.g. haptophytes, *Karenia mikimotoi*, *Pseudo-nitzschia multiseries* (Wang et al., 2023; Yao et al., 2024; Yu et al., 2025). A notable exception is the use of RPA coupled to Nanopore sequencing, which enables portable, high-resolution taxonomic profiling but remains non-quantitative and DNA-based (Hatfield et al., 2020). Among isothermal methods, the only study to date targeting a toxin biosynthesis gene rather than a ribosomal marker is a LAMP assay targeting *dab*A in *Pseudo-nitzschia multistriata*, demonstrated on a custom-made field-deployable device; however, it is also DNA-based and lacks detailed quantitative and specificity characterization (Alrefaey et al., 2026). In contrast, the qRT-RPA assay presented here uniquely combines isothermal amplification, transcript-level detection and real-time quantification, making it well suited for early-warning applications where speed, functional relevance and portability are prioritized.

4.4 Conclusion

This study establishes the analytical performance of a universal qRT-RPA assay targeting the *sxt*A4 transcript, enabling rapid, cross-species detection of putatively toxigenic *Alexandrium* with functional relevance and short time-to-result. Validation under controlled conditions, including sequence diversity and increasing matrix complexity, provides a solid methodological foundation for RNA-based isothermal monitoring. As with any emerging molecular approach, targeted field testing with natural bloom samples and direct comparison to qPCR, microscopy and toxin measurements will be required to define operational performance and establish informed risk thresholds. With full compatibility to low-power portable fluorescence readers and sensor-integrated platforms, qRT-RPA is well positioned as a rapid early-warning tool in HAB monitoring programmes.

Acknowledgements

The authors would like to thanks Dr. Nuria Lluch, Dr. Rosa Isabel Figueroa and Dr. Esther Garces at Insitut de Ciencias del Mar, Spain, for contributing *Alexandrium minutum* and *Gymnodinium catenatum* cultures. This research was funded by the EU Horizon project AquaBioSens.

REFERENCES

Alrefaey, A.I., McQuillan, J.S., Schaap, A., Siracusa, F., Cardwell, C.L., Walk, J., Rogers, D., Forrester, R., Mowlem, M.C., Robidart, J.C., 2026. A lab-on-a-chip system integrating DNA purification and loop-mediated isothermal amplification for the quantification of the toxic diatom Pseudo-nitzschia multistriata. Sensors & Diagnostics.

Anderson, D.M., Alpermann, T.J., Cembella, A.D., Collos, Y., Masseret, E., Montresor, M., 2012. The globally distributed genus Alexandrium: Multifaceted roles in marine ecosystems and impacts on human health. Harmful Algae 14, 10-35.

Anderson, D.M., Fachon, E., Pickart, R.S., Lin B □, P., Fischer, A.D., Richlen, M.L., Uva, V.,

Brosnahan, M.L., McRaven, L., Bahr, F., Lefebvre, K., Grebmeier D □, J.M., Danielson, S.L.,

Lyu, Y., Fukai, Y., 2021a. Evidence for massive and recurrent toxic blooms of Alexandrium

catenella in the Alaskan Arctic. Proc Natl Acad Sci U S A 118(41):e2107387118.

Anderson, D.M., Fensin, E., Gobler, C.J., Hoeglund, A.E., Hubbard, K.A., Kulis, D.M.,

Landsberg, J.H., Lefebvre, K.A., Provoost, P., Richlen, M.L., Smith, J.L., Solow, A.R., Trainer,

V.L., 2021b. Marine harmful algal blooms (HABs) in the United States: History, current status

and future trends. Harmful Algae 102.

Commission., E., 2021. Commission Implementing Regulation (EU) 2021/1709 of 23 September

2021 amending Implementing Regulation (EU) 2019/627. (Official Journal L 339, p. 84.

Dai, L., Geng, H.X., Yu, R.C., Liu, Y., Zhao, J.Y., Wang, J.X., Zhang, Q.C., Kong, F.Z., Zhou,

M.J., 2020. Distribution of Alexandrium pacificum cysts in the area adjacent to the Changjiang

River estuary, China. Marine Pollution Bulletin 156.

Dai, Y., Yang, S., Zhao, D., Hu, C., Xu, W., Anderson, D.M., Li, Y., Song, X.P., Boyce, D.G.,

Gibson, L., Zheng, C., Feng, L., 2023. Coastal phytoplankton blooms expand and intensify in the

21st century. Nature 615(7951), 280-284.

Erdner, D.L., Percy, L., Keafer, B., Lewis, J., Anderson, D.M., 2010. A quantitative real-time PCR

assay for the identification and enumeration of Alexandrium cysts in marine sediments. Deep-Sea

Research Part II: Topical Studies in Oceanography 57(3-4), 279-287.

Gao, Y., Yu, R.C., Murray, S.A., Chen, J.H., Kang, Z.J., Zhang, Q.C., Kong, F.Z., Zhou, M.J.,

2015. High specificity of a quantitative PCR assay targeting a saxitoxin gene for monitoring toxic

algae associated with paralytic shellfish toxins in the Yellow Sea. Applied and Environmental

Microbiology 81(20), 6973-6981.

Garneau, M.È., Schnetzer, A., Countway, P.D., Jones, A.C., Seubert, E.L., Caron, D.A., 2011.

Examination of the seasonal dynamics of the toxic dinoflagellate Alexandrium catenella at

Redondo Beach, California, by quantitative PCR. Applied and Environmental Microbiology

77(21), 7669-7680.

Geffroy, S., Lechat, M.M., Le Gac, M., Rovillon, G.A., Marie, D., Bigeard, E., Malo, F., Amzil,
Z., Guillou, L., Caruana, A.M.N., 2021. From the sxtA4 Gene to Saxitoxin Production: What
Controls the Variability Among Alexandrium minutum and Alexandrium pacificum Strains?
Frontiers in Microbiology 12.

Gobler, C.J., Doherty, O.M., Hattenrath-Lehmann, T.K., Griffith, A.W., Kang, Y., Litaker, R.W.,
2017. Ocean warming since 1982 has expanded the niche of toxic algal blooms in the North
Atlantic and North Pacific oceans, Proceedings of the National Academy of Sciences of the United
States of America. National Academy of Sciences, pp. 4975-4980.

Gribble, M.O., Bennett, B.J., Liddie, J.M., Borchert, W., Pfluger, B.A., Segars, J.S., Keast, J.M.,
Hans, A., Kikkeri, N.S., Shin, C., Roland, H.B., Hoysala, S., Kohlhoff, J., Alahmad, B., Nagalli,
S., Rushton, W., Enevoldsen, H., Bell, M., Fortenberry, D., Harley, J.R., Schneider, A.L.C., 2025.
Global epidemiology of paralytic shellfish poisoning: a systematic search literature review. The
Lancet Planetary Health, 101271-101271.

Griffith, A.W., Doherty, O.M., Gobler, C.J., 2019. Ocean warming along temperate western
boundaries of the Northern Hemisphere promotes an expansion of Cochlodinium polykrikoides
blooms. Proceedings of the Royal Society B: Biological Sciences 286(1904).

Han-Sol, K., Taehee, K., Jaeyeon, P., Tae Gyu, P., Jang-Seu, K., 2024. Development of saxitoxin
biosynthesis gene sxtB-targeted qPCR assay for the quantification of toxic dinoflagellates
Alexandrium catenella (group I) and A. pacificum (group IV) occurring in the Korean coast.
Harmful Algae Volume 134,102603.

Hatfield, R.G., Batista, F.M., Bean, T.P., Fonseca, V.G., Santos, A., Turner, A.D., Lewis, A.,
Dean, K.J., Martinez-Urtaza, J., 2020. The Application of Nanopore Sequencing Technology to
the Study of Dinoflagellates: A Proof of Concept Study for Rapid Sequence-Based Discrimination
of Potentially Harmful Algae. Frontiers in Microbiology Volume 11 - 2020.

Hatfield, R.G., Bean, T., Turner, A.D., Lees, D.N., Lowther, J., Lewis, A., Baker-Austin, C., 2019.
Development of a TaqMan qPCR assay for detection of Alexandrium spp and application to
harmful algal bloom monitoring. Toxicon: X 2.

Hayashida, H., Matear, R.J., Strutton, P.G., 2020. Background nutrient concentration determines
phytoplankton bloom response to marine heatwaves. Global Change Biology 26(9), 4800-4811.

John, D.E., Wang, Z.A., Liu, X., Byrne, R.H., Corredor, J.E., López, J.M., Cabrera, A., Bronk,
D.A., Tabita, F.R., Paul, J.H., 2007. Phytoplankton carbon fixation gene (RuBisCO) transcripts
and air-sea CO"2 flux in the Mississippi River plume. ISME Journal 1(6), 517-531.

Kellmann, R., Mihali, T.K., Young, J.J., Pickford, R., Pomati, F., Neilan, B.A., 2008. Biosynthetic
intermediate analysis and functional homology reveal a saxitoxin gene cluster in cyanobacteria.
Applied and Environmental Microbiology 74(13), 4044-4053.

Kim, H.S., Kim, T., Park, J., Park, T.G., Ki, J.S., 2024. Development of saxitoxin biosynthesis
gene sxtB-targeted qPCR assay for the quantification of toxic dinoflagellates Alexandrium
catenella (group I) and A. pacificum (group IV) occurring in the Korean coast. Harmful Algae
134.

Kon, N.F., Teng, S.T., Hii, K.S., Yek, L.H., Mujahid, A., Lim, H.C., Lim, P.T., Leaw, C.P., 2015.
Spatial distribution of toxic Alexandrium tamiyavanichii (Dinophyceae) in the southeastern South
China Sea-Sulu Sea: A molecular-based assessment using real-time quantitative PCR (qPCR)
assay. Harmful Algae 50, 8-20.

Lewis, A.M., Coates, L.N., Turner, A.D., Percy, L., Lewis, J., 2018. A review of the global
distribution of Alexandrium minutum (Dinophyceae) and comments on ecology and associated
paralytic shellfish toxin profiles, with a focus on Northern Europe, Journal of Phycology.
Blackwell Publishing Inc., pp. 581-598.

Liu, H., Cao, T., Chen, H., Zhang, J., Li, W., Zhang, Y., Liu, H., 2023. Two-color lateral flow
nucleic acid assay combined with double-tailed recombinase polymerase amplification for
simultaneous detection of chicken and duck adulteration in mutton. Journal of Food Composition
and Analysis 118, 105209.

Liu, X., Yan, Q., Huang, J., Chen, J., Guo, Z., Liu, Z., Cai, L., Li, R., Wang, Y., Yang, G., Lan,
Q., 2019. Influence of design probe and sequence mismatches on the efficiency of fluorescent
RPA. World Journal of Microbiology and Biotechnology 35(6).

Muhammad, B.L., Kim, H.-S., Bui, Q.T.N., Ki, J.-S., 2025. Transcriptomic comparison unveils
saxitoxin biosynthesis genes in the marine dinoflagellate Gymnodinium catenatum. Harmful
Algae 147, 102872.

Munawar, M.A., 2022. Critical insight into recombinase polymerase amplification technology.
Expert Review of Molecular Diagnostics 22(7), 725-737.

Murray, S.A., Diwan, R., Orr, R.J.S., Kohli, G.S., John, U., 2015. Gene duplication, loss and
selection in the evolution of saxitoxin biosynthesis in alveolates. Molecular Phylogenetics and
Evolution 92, 165-180.

Murray, S.A., Ruvindy, R., Kohli, G.S., Anderson, D.M., Brosnahan, M.L., 2019. Evaluation of
sxtA and rDNA qPCR assays through monitoring of an inshore bloom of Alexandrium catenella
Group 1. Scientific Reports 9(1).

Murray, S.A., Wiese, M., Stüken, A., Brett, S., Kellmann, R., Hallegraeff, G., Neilan, B.A., 2011.
SxtA-based quantitative molecular assay to identify saxitoxin-producing harmful algal blooms in
marine waters. Applied and Environmental Microbiology 77(19), 7050-7057.

Parks, R., Bear, E., Coates, L., Maskrey, B., Algoet, M., 2020. Annual report on the results of the
Biotoxin and Phytoplankton Official Control Monitoring Programmes.

Penna, A., Perini, F., Dell'Aversano, C., Capellacci, S., Tartaglione, L., Giacobbe, M.G.,
Casabianca, S., Fraga, S., Ciminiello, P., Scardi, M., 2015. The sxt Gene and Paralytic Shellfish
Poisoning Toxins as Markers for the Monitoring of Toxic Alexandrium Species Blooms.
Environmental Science and Technology 49(24), 14230-14238.

Rabalais, N.N., Turner, R.E., Sen Gupta, B.K., Boesch, D.F., Chapman, P., Murrell, M.C., 2007.
Hypoxia in the Northern Gulf of Mexico: Does the Science Support the Plan to Reduce, Mitigate,
and Control Hypoxia?, pp. 753-772.

Rohrman, B., Richards-Kortum, R., 2015. Inhibition of Recombinase Polymerase Amplification
by Background DNA: A Lateral Flow-Based Method for Enriching Target DNA. Analytical
Chemistry 87(3), 1963-1967.

Ruvindy, R., Barua, A., Bolch, C.J.S., Sarowar, C., Savela, H., Murray, S.A., 2023. Genomic copy
number variability at the genus, species and population levels impacts in situ ecological analyses
of dinoflagellates and harmful algal blooms. ISME Communications 3(1).

Ruvindy, R., Bolch, C.J., MacKenzie, L., Smith, K.F., Murray, S.A., 2018. qPCR Assays for the
Detection and Quantification of Multiple Paralytic Shellfish Toxin-Producing Species of
Alexandrium. Frontiers in Microbiology 9.

Savela, H., Harju, K., Spoof, L., Lindehoff, E., Meriluoto, J., Vehniäinen, M., Kremp, A., 2016.
Quantity of the dinoflagellate sxtA4 gene and cell density correlates with paralytic shellfish toxin
production in Alexandrium ostenfeldii blooms. Harmful Algae 52, 1-10.

Senovilla-Herrero, D., Chen, Y., Liu, C., Suresh, S., Moore, H., McKinney, A., Service, M.,
Helyar, S., Mbadugha, L., Campbell, K., 2025. Environmental and Biological Factors of
Relevance to Shellfish Production in Northern Ireland: Insights From 20 Years of Regional
Monitoring Data. Reviews in Aquaculture 17(1), e12998.

Stüken, A., Dittami, S.M., Eikrem, W., McNamee, S., Campbell, K., Jakobsen, K.S., Edvardsen,
B., 2013. Novel hydrolysis-probe based qPCR assay to detect saxitoxin transcripts of
dinoflagellates in environmental samples. Harmful Algae 28, 108-117.

Stüken, A., Orr, R.J.S., Kellmann, R., Murray, S.A., Neilan, B.A., Jakobsen, K.S., 2011. Discovery
of nuclear-encoded genes for the neurotoxin saxitoxin in dinoflagellates. PLoS ONE 6(5).

Suleiman, M., Jelip, J., Rundi, C., Chua, T.H., 2017. Case report: Paralytic shellfish poisoning in
Sabah, Malaysia. American Journal of Tropical Medicine and Hygiene 97(6), 1731-1736.

Tan, M., Liao, C., Liang, L., Yi, X., Zhou, Z., Wei, G., 2022. Recent advances in recombinase
polymerase amplification: Principle, advantages, disadvantages and applications, Frontiers in
Cellular and Infection Microbiology. Frontiers Media S.A.

Toebe, K., Alpermann, T.J., Tillmann, U., Krock, B., Cembella, A., John, U., 2013. Molecular
discrimination of toxic and non-toxic Alexandrium species (Dinophyta) in natural phytoplankton
assemblages from the Scottish coast of the North Sea. European Journal of Phycology 48(1), 12-
26.

Turner, A.D., Dhanji-Rapkova, M., Baker, C., Algoet, M., 2014. Assessment of a semiquantitative
liquid chromatography-fluorescence detection method for the determination of paralytic shellfish
poisoning toxin levels in bivalve molluscs from Great Britain. Journal of AOAC International
97(2), 492-497.

Vandersea, M.W., Kibler, S.R., Van Sant, S.B., Tester, P.A., Sullivan, K., Eckert, G., Cammarata,
C., Reece, K., Scott, G., Place, A., Holderied, K., Hondolero, D., Litaker, R.W., 2017. QPCR
assays for Alexandrium fundyense and A. ostenfeldii (Dinophyceae) identified from Alaskan
waters and a review of species-specific Alexandrium molecular assays. Phycologia 56(3), 303-
320.

Vingiani, G.M., Stalberga, D., De Luca, P., Ianora, A., De Luca, D., Lauritano, C., 2020. De novo
transcriptome of the non-saxitoxin producing alexandrium tamutum reveals new insights on
harmful dinoflagellates. Marine Drugs 18(8).

Wang, L., Chen, X., Pan, F., Yao, G., Chen, J., 2023. Development of a rapid detection method
for Karenia mikimotoi by using CRISPR-Cas12a. Frontiers in Microbiology Volume 14 - 2023.

Yao, Y., Luo, N., Zong, Y., Jia, M., Rao, Y., Huang, H., Jiang, H., 2024. Recombinase Polymerase
Amplification Combined with Lateral Flow Dipstick Assay for the Rapid and Sensitive Detection
of Pseudo-nitzschia multiseries. International Journal of Molecular Sciences 25(2), 1350.

Yu, J., Shen, Y., Zhang, Q., Luo, X., Zong, Y., Zhou, C., Huang, H., Jiang, H., 2025. Development
of an RPA-CRISPR/LbaCas12a-Lateral Flow Assay for the Visual Detection of Chrysotila dentata
(Haptophyta). Microorganisms 13(9), 2203.

Yu, Z., Tang, Y., Gobler, C.J., 2023. Harmful algal blooms in China: History, recent expansion,
current status, and future prospects, Harmful Algae. Elsevier B.V.