OmniCell: Unified Foundation Modeling of Single-Cell and Spatial Transcriptomics for Cellular and Molecular Insights

Jiangshuan Pang[1,2†], Ping Qiu[4†], Youzhe He[5†], Baolong Li[1,2†], Yiting Deng[1,2†], Jun Wang[6] ,Adi Lin[2], Lei Cao[2], Fei Teng[4], Haoran Wang[2], Shuangsang Fang[2], Shengkang Li[4] , Ziqing Deng[2], Yong Zhang[4,*], Yuxiang Li[4,*], Shaoshuai Li[2,*], Xun Xu[3,*]

1. School of Artificial Intelligence, University of Chinese Academy of Sciences, Beijing 100049, China.

2. BGI Research, Beijing 102601, China.

3. BGI Research, Shenzhen 518083, China.

4. BGI Research, Wuhan 430074, China.

5. College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China.

6. Bioinformatics Centre, Department of Biology, University of Copenhagen, København Ø 2100, Denmark.

*Correspondence: xuxun@genomics.cn   (X.X.),   lishaoshuai@genomics.cn   (L.S.S.), zhangyong2@genomics.cn (Z.Y.),   liyuxiang@genomics.cn (L.Y.X.)

† These authors contributed equally as the first authors.

Abstract

Single-cell RNA sequencing (scRNA-seq) enables characterization of cellular heterogeneity but lacks spatial context, while Spatially Transcriptomics maps gene expression in tissues with limited single-cell resolution. Integrating the complementary strengths of these data into a unified framework remains challenging. Here, we present OmniCell, a foundation model for single-cell and spatial transcriptomics, pretrained on a large-scale corpus of 67 million single-cell and spatial transcriptomic profiles, enabling the unified multi-omics representation learning. As the first foundation model to jointly capture intra-cellular gene expression relationships and inter-cellular spatial dependencies within a unified framework, OmniCell explicitly represents tissue spatial topology by serializing spatially adjacent cells during input construction. Leveraging this unified modeling paradigm, OmniCell generates unified representations of genes, cells, and tissue spatial organization. In zero-shot evaluations, it reliably recovers cell-type structure and gene expression patterns, reconstructs co-expression relationships, and outperforms existing methods across all evaluated tasks, including cell-type deconvolution and spatial domain delineation. Applied to real spatial datasets, OmniCell resolves transitional zones at tumor margins and reveals associated inflammatory activation and immune-cell enrichment, demonstrating its capacity for high-resolution spatial profiling.

Introduction

In multicellular organisms, the diversity of cell types and the spatial architecture of tissues jointly shape physiological function and disease progression. The advent of scRNA-seq has enabled the dissection of cellular heterogeneity, regulatory networks, and state transitions at single-cell resolution, providing a detailed view of the molecular organization within tissues[1-3]. To achieve this, various scRNA-seq platforms have been developed, including droplet-based methods (10x Genomics[4], Drop-seq[5]), plate-based approaches (Smart-seq2[6], Smart-seq3[7]). However, because scRNA-seq requires physical dissociation of tissues prior to profiling, it inevitably disrupts the native spatial context, leading to the loss of critical information such as a cell's position within the tissue, its local neighborhood, and the microenvironment in which it resides[8, 9].

In recent years, spatial transcriptomics (ST) technologies have advanced rapidly, enabling researchers to measure gene expression directly within tissues while capturing spatial context. Imaging-based methods such as MERFISH[10], seqFISH[11], and STARmap[12] achieve subcellular resolution but typically target predefined gene panels, limiting genome-wide regulatory insights. In situ sequencing approaches, including BaristaSeq[13], enable direct RNA sequencing in tissue sections. Spatial barcoding technologies, such as Visium[14], Slide-seq[15], offer intermediate spatial resolution (10-100 μm) with transcriptome-wide coverage. More recently, genome-scale platforms like Stereo-seq[16] have produced high-resolution spatial expression maps approaching single-cell resolution. These advances enable the study of cell–cell interactions, tissue domains, and disease microenvironments. Nevertheless, current spatial techniques involve trade-offs between resolution and capture efficiency, and they often suffer from limited molecular coverage and substantial technical noise, particularly at near single-cell resolution, posing significant challenges for computational analysis.

As scRNA-seq and ST technologies mature, the amount of transcriptomic data available for analysis has expanded sharply. Large scRNA-seq collections and ST datasets, generated across diverse tissues, species, and experimental conditions, are now accumulating at an unprecedented pace, offering a broader view of gene regulation and cellular organization. This diversity raisean important question: whether such heterogeneous resources can be distilled into representations that capture biological structure shared across platforms and studies. Drawing on developments in language modeling, several groups have begun to explore large pretrained models for single-cell analysis, such as scGPT[17], scFoundation[18] and Geneformer[19]. They have demonstrated that gene expression can be treated as a molecular language, yielding rich cellular representations[20]. Building on this idea, several studies have extended pretrained

single-cell models to spatial data. For example, scGPT-Spatial[21] adapts scGPT via continual pretraining

on spot-level ST datasets. While effective for representation transfer, it is limited by spot-level aggregation,

which obscures cell-resolved heterogeneity, and separate modality pipelines for scRNA-seq and spatial

data, which hinder seamless cross-omics alignment. NicheFormer, despite being described as a unified

foundation model for single-cell and spatial data, does not explicitly capture intercellular spatial

dependencies [22]. Similarly, CellPLM is constrained by its reliance on cell-level scRNA-seq modeling,

which overlooks intracellular gene interactions, and by its modest spatial dataset of only 2 million spots,

which is insufficient to fully represent the diversity of spatial organization and cell–cell interactions [23].

These limitations underscore a key challenge: how to integrate complementary but heterogeneous data

within a single framework, enabling models to capture gene regulation, cellular states, and tissue spatial

organization simultaneously. To address this, we introduce OmniCell, a unified foundation model pretrained

on 67 million single-cell and spatial expression profiles to learn shared representations across gene, cellular,

and tissue spatial scales. For ST data, we construct spatially ordered sequences by concatenating gene

expression profiles from each cell and its neighboring cells, thereby enabling Transformer-style

self-attention[24] to capture tissue architecture and spatial context. In the gene space, OmniCell employs a

Mixture-of-Experts (MoE) expression embedding module, in which gene identity determines the routing of

expression values to gene-specific expert networks, while additional shared experts capture common

expression patterns across genes.The learnable Symmetric Bilinear Output Module is a central component of

OmniCell, integrating cell-level and gene-level representations through bilinear interactions. By jointly

capturing cellular expression patterns and spatial context, it produces stable embeddings that reflect gene

interactions and tissue architecture. This design enhances the model's generalization across platforms and

multi-omic datasets. Ablation experiments demonstrate that the module is essential for performance,

highlighting its role in generating coherent and biologically meaningful representations across modalities.

Building on this design, OmniCell achieves joint representation learning across genes, cells, and tissue

spatial organization within a unified framework. Even in zero-shot settings, it accurately recovers cellular

structures, captures cell-specific gene programs, and models cross-cell co-expression relationships.

Moreover, OmniCell consistently outperforms existing methods in tasks such as cell-type deconvolution and

spatial domain identification. In real tumor spatial datasets, it identifies transitional regions at tumor margins

and reveals key features such as inflammation pathway activation and immune cell enrichment,

demonstrating its remarkable capacity for spatially resolved biological insights.

Results

A unified foundation model for single-cell and spatial transcriptomics

OmniCell is a unified foundation model designed to jointly represent scRNA-seq and ST data. The model

encodes gene identity, expression level, and spatial coordinates within a shared, technology-agnostic

framework, allowing it to seamlessly integrate both dissociated and spatially resolved cells.   Building on

this architecture, OmniCell was pretrained on approximately 67 million cells drawn from scRNA-seq data

across 45 human tissues and ST data spanning 26 tissue types (**Figure 1A**). From this diverse training corpus,

the model learns coherent gene- and cell-level representations that capture both transcriptional programs and

spatial organization. This enables robust cross-modal reasoning and supports a wide range of downstream

analyses.

Model Overview. To jointly model scRNA-seq and ST data, OmniCell encodes cells as sequences of gene

tokens, enabling Transformer-style attention to capture intra-cellular gene interactions (**Figure 1B**).

Within this unified tokenization scheme, the two modalities are handled with tailored strategies. For

scRNA-seq, we represent each cell by a feature vector prioritizing its non-zero expressed genes. To augment

cellular representations in cases of limited non-zero signal, we specifically pad the vector with high-variance

zero-expression genes. For ST data, sequences incorporate a target cell alongside its spatial neighbors,

thereby capturing both intra- and inter-cellular dependencies.

A central challenge in modeling transcriptomic data lies in handling continuous gene expression values. We

address this by first applying a soft-rank transformation, which ranks genes within each cell and rescales

them to a fixed interval. This operation preserves the relative ordering of expression levels while mitigating

the influence of extreme values, effectively normalizing data for subsequent modeling. Beyond

normalization, we further introduce a gene-aware MoE value embedding that combines a unified gene

embedding matrix with a MoE-based value embedding module. This design adaptively encodes gene

expression in a context-sensitive manner, capturing subtle variations while maintaining biological fidelity

and cell-type specificity.

The architectural backbone of OmniCell comprises 10 Transformer layers with RMS normalization, 2D

rotary self-attention, and SwiGLU feedforward networks, totaling approximately 74 million parameters. At

the output stage, a learnable symmetric matrix integrates gene- and cell-level features: during training, it is

optimized under a soft-rank regression objective, while at inference, spectral decomposition projects cell

embeddings into a stable low-dimensional space. Together, this architecture produces robust, biologically

interpretable embeddings suitable for diverse downstream analyses, including cell-type annotation, spatial

domain delineation, and the construction of gene co-expression networks.

Downstream Tasks. We next evaluated OmniCell across both single-cell and spatial transcriptomic tasks,

(**Figure 1C**). At the gene level, OmniCell learns cell-type–specific gene representations that faithfully

capture transcriptional programs, enabling marker gene identification and co-expression pattern discovery.

At the cell level, the model generates embeddings that integrate transcriptional and spatial context,

supporting robust clustering, cell-type annotation, batch correction, spatial domain delineation, and spatial

deconvolution. Notably, across these diverse tasks, the model maintains stability under increased single-cell

noise and varying tissue contexts. Taken together, these results demonstrate that by jointly learning

high-quality gene- and cell-level embeddings, OmniCell effectively bridges dissociated and spatially

resolved data, yielding biologically interpretable representations that generalize across modalities.

Ablation Study. Beyond overall performance, we sought to dissect the contribution of key architectural

components through systematic ablation studies across clustering and cell-type annotation benchmarks

(**Figure 1D**). Removing the gene-aware MoE value embedder modestly reduced performance, reflecting its

role in capturing cell-type–specific expression patterns. Excluding spectral subspace projection impaired

embedding robustness, and eliminating the symmetric bilinear output module led to the largest performance

drop, confirming its central role in modeling cell–gene relationships. We also examined the impact of

training data composition: training on scRNA-seq data alone without spatial supervision degraded clustering

and annotation accuracy, highlighting the importance of integrating spatial transcriptomics to enrich data

diversity and provide structural constraints. Collectively, these ablation results demonstrate that each

component of OmniCell contributes synergistically to its robust, biologically interpretable performance.

Cell-Level and Dataset-Level Gene Embeddings Reveal Cell-Type–Specific Signatures and

Functional Modules in scRNA-seq and ST data.

We first evaluated the quality of cell-level gene embeddings by extracting contextualized representations

from the Transformer model and projecting them into a lower-dimensional space. For this analysis, we

employed Zhou's dataset, which enabled us to capture both global gene properties and cell-specific activities.

To assess how well these embeddings preserve cell-type information, we performed clustering on the gene

embeddings and computed the Adjusted Rand Index (ARI) against known cell-type labels. OmniCell

consistently achieved higher ARI scores compared to scGPT, scFoundation, and GeneFormer (**Figure 2A**),

indicating that its embeddings retain cell-type information more accurately. To further illustrate these

findings, we generated UMAP visualizations of the embeddings for several key genes, including *APOE*

(**Figure 2B**), *TREM2* (**Figure 2C**), and *LHFPL3* (**Supplementary Figure 2C**).

These visualizations reveal that OmniCell more clearly distinguishes between cell types, particularly in

resolving clusters of high and low expression within microglial cells—patterns with known associations to

Alzheimer's Disease that may reflect pathological alterations with diagnostic or therapeutic relevance. We

also examined how the model allocates attention across genes by aggregating attention weights into a

cell-by-gene attention matrix, which captures the relative importance of each gene across cell types.

Multiclass Receiver Operating Characteristic (ROC) analysis (**Figure 2D**) demonstrated that OmniCell

outperformed competing models, with all cell types achieving Area Under the Curve (AUC) values above

0.9, underscoring its robust capability in leveraging gene attention to differentiate cell populations.

Having demonstrated that OmniCell embeddings preserve cell-type information, we next investigated

whether they could be leveraged to identify marker genes for specific cell types. Using cell-level gene

embeddings generated by the foundation model (as described in the Methods), we ranked genes for each

annotated cell type in Zhou's scRNA-seq dataset based on their specificity in the embedding space and

filtered by transcriptional abundance to retain highly expressed, cell-type–specific markers. Encouragingly,

many of the embedding-derived markers corresponded to canonical genes reported in previous studies,

such as *SLC17A7* for excitatory neurons[25], *ERBB4* for inhibitory neurons[26], *GFAP* for astrocytes[27],

LRMDA for microglia[28], *PLP1* for oligodendrocytes[29], and *PTPRZ1* for oligodendrocyte precursor

cells (OPCs)[30] (**Figure 2E and Supplementary Figure 2B**). Notably, the majority of the

embedding-derived markers for most cell types were also present in the top marker gene lists identified by

conventional expression-based ranking methods (**Figure 2F**), supporting the biological validity of the

embeddings. Beyond known markers, OmniCell embeddings also identified additional cell-type–enriched

genes with strong discriminative power (AUC > 0.7), including *SYT1* for excitatory neurons[31] and

SPARCL1 for astrocytes[32] (**Figure 2G**). Together, these results revealed that OmniCell embeddings not

only recapitulate known cell-type–specific gene expression patterns but also enable the systematic

identification of novel markers directly from the representation space.

While cell-level embeddings capture gene properties within individual cells, understanding broader

co-expression relationships requires aggregating information across the entire dataset. To this end, we

generated dataset-level gene embeddings by integrating cell-level embeddings through a generalized

PageRank graph neural network [33] (GPR-GNN), which propagates each gene's embedding through a cell

similarity graph to reinforce co-expression relationships and context-dependent interactions

(**Supplementary Figure 2A**). Final embeddings were obtained by averaging across all cells, capturing both

intrinsic expression patterns and dataset-wide functional properties.   To validate these embeddings, we

constructed a gene co-embedding network using cell-type–specific gene sets identified previously. Louvain

community detection revealed six distinct modules (Modules 0–5) that exhibited highly cell-type–specific

activation patterns aligned with known transcriptional programs (**Figure 2H-I**): Module 0 for astrocytes;

Module 1 for oligodendrocytes; Module 2 for inhibitory neurons; Module 3 for microglia; Module 4 for

both excitatory and inhibitory neurons; and Module 5 for oligodendrocyte precursor cells. Gene Ontology

enrichment analysis confirmed functional coherence, with modules enriched for cell-type–appropriate

processes including neurotransmitter transport, myelination, synaptic signaling, immune function, and

oligodendrocyte differentiation (**Figure 2J**). We further assessed spatial coherence by performing joint

module identification on spatial transcriptomic data from Alzheimer's disease and control human

hippocampus samples. Five spatial modules emerged with region-specific expression patterns

corresponding to known anatomical boundaries: excitatory neuron–enriched modules showed peak activity

in the pyramidal cell layers of CA1–CA4 and the granule cell layer of the dentate gyrus, while

glial-enriched modules localized to the GC region, consistent with the spatial distribution of astrocytes and

oligodendrocyte-lineage cells (**Figure 2K-M and Supplementary Figure 2D, E**)

Collectively, these analyses demonstrate that dataset-level gene embeddings derived from OmniCell

capture biologically meaningful gene–gene relationships, organize genes into coherent functional modules

aligned with cell-type–specific transcriptional programs and anatomical structures, and provide a principled

framework for investigating coordinated gene expression patterns in both single-cell and spatial

transcriptomic contexts.

Omnicell Enables Cross-Species Generalization and Robust, Interpretable, Lineage-Aware Cell

Embeddings via a MoE Value Embedder.

To evaluate OmniCell's capacity for generalized cell representation learning, we benchmarked its

embeddings against those generated by existing pretrained models across human (kidney and blood) and

mouse (brain and muscle) datasets. As shown in **Figure 3A**, OmniCell consistently achieved the highest

clustering performance in both ARI and NMI, outperforming all baselines. Notably, although OmniCell was

pretrained exclusively on human data, it maintained clear cell-type separability in mouse datasets through

homology-based gene projection, demonstrating robust cross-species transferability. UMAP visualizations

further corroborated these quantitative results (**Figure 3B, 3C and Supplementary Figure 3A, 3B**),

showing that OmniCell embeddings form compact and biologically coherent clusters with superior

intra-cluster cohesion and inter-cluster separation compared to scGPT and GeneFormer.

Beyond clustering accuracy, a practical challenge in single-cell analysis is mitigating batch effects

while preserving biological variation. We therefore assessed OmniCell's performance on the Zhou dataset,

which comprises 11 distinct batches (**Figure 3D, 3E**). OmniCell achieved a favorable balance between

integration and biological preservation, effectively mixing technical batches while maintaining underlying

structure. Consistent results were observed in a multi-batch human pancreas dataset (**Supplementary**

Figure 3C, 3D), where diverse cell types remained well separated across experimental conditions,

highlighting OmniCell's robustness to technical variation. Given that real-world single-cell data often

suffer from dropout and missing values, we further evaluated embedding stability by applying progressive

random dropout to the Zhou dataset (**Figure 3F**). OmniCell retained substantially higher clustering

stability than existing models under increasing sparsity, demonstrating its ability to encode noise-resilient

transcriptional features. Together, these results establish OmniCell as a robust foundation model capable of

reliable generalization across species, platforms, and data quality conditions.

The strong performance observed above prompted us to investigate whether OmniCell's MoE value

embedder contributes to its representational capacity. We hypothesized that this modular routing

mechanism enables encoding of lineage- and function-specific transcriptional programs by dynamically

allocating computational resources. To test this, we computed mean routing weights and standardized

activations for each expert across annotated cell types in the human immune cell dataset. The resulting

heatmap (**Figure 3G**) revealed lineage-stratified activation patterns, with major populations—including

T/NK cells, B cells, myeloid cells, and erythroid progenitors—clearly delineated by distinct expert

preferences. Expert Preference Clustering ($Z > 1.0$; **Supplementary Figure 3E**) reinforced these findings,

highlighting clear lineage-specific associations across major immune lineages. These activation patterns

suggest that the MoE router autonomously partitions computation into biologically coherent modules

mirroring immune cell differentiation hierarchies, indicating the model encodes interpretable,

lineage-aware functional specialization. In summary, OmniCell excels at learning stable, biologically

meaningful cell embeddings, outperforming existing models in clustering, cross-species transfer, and

batch-effect mitigation. Its MoE architecture enables dynamic, lineage-specific routing that captures

cellular heterogeneity while maintaining interpretable functional organization.

Multi-Scale Spatial Transcriptomic Benchmarking of OmniCell Across Cellular and Tissue

Hierarchies.

To systematically evaluate OmniCell's capacity to capture spatially organized biological structure, we

conducted a multi-scale assessment spanning single-cell resolution to tissue-level architecture. At the

cellular scale, spatial clustering evaluates whether embeddings derived from individual cells faithfully

recover established cell-type identities. At the tissue scale, spatial domain delineation incorporates

multi-cell inputs, jointly encoding a central cell with its spatial neighbors to resolve mesoscale anatomical

organization. This hierarchical framework enables OmniCell to integrate molecular identity with spatial

context within a unified embedding space.

We first assessed spatial clustering using a comprehensive MERFISH-based mouse brain atlas

encompassing 31 spatial transcriptomic datasets that capture molecular and cellular dynamics across aging.

Single-cell embeddings were subjected to Leiden clustering across multiple resolutions (0.1–1.0), with

optimal clustering determined by concordance with reference annotations. OmniCell consistently achieved

superior performance relative to scGPT-spatial and Nicheformer, as quantified by both NMI and ARI

(**Figure 4A**). Representative visualization of slice 10_0 demonstrated that OmniCell embeddings yield

well-resolved clusters in UMAP space (**Figure 4B**) that accurately recapitulate anatomical organization

upon projection to spatial coordinates (**Figure 4C**), with comparable results for slice 10_2

(**Supplementary Figure 4A-B**). In contrast, baseline methods exhibited substantial intermixing of

spatially adjacent regions.

Moving from single-cell to tissue-scale analysis, we next evaluated spatial domain delineation on a

mouse cortex dataset with expert-curated regional annotations encompassing hippocampus, internal

capsule, cortical layers 2–6, pia Layer 1, ventricle, and white matter. In this multi-cell configuration, each

embedding integrates transcriptomic information from a central cell and its spatial neighborhood.

OmniCell achieved optimal reconstruction of cortical architecture (NMI = 0.6623, ARI = 0.6108),

significantly outperforming all competing methods (**Supplementary Figure 4C-D**). These results

demonstrate that explicit incorporation of spatial context enables OmniCell to resolve mesoscale tissue

organization beyond individual cell-type classification.

Having established OmniCell's performance on well-annotated datasets, we sought to challenge the

model with a more complex scenario: the Stereo-seq LC5-M hepatocellular carcinoma dataset, which

encompasses tumor core, peri-tumoral transition zone, and adjacent non-malignant parenchyma without

predefined annotations [34]. Comparative analysis across six methods revealed that only OmniCell

successfully delineated a discrete transitional domain (Cluster 3) at the tumor-parenchyma interface

(**Figure 4D**). Gene Ontology enrichment analysis of this transition zone revealed significant activation of

pathways including acute-phase response, inflammatory signaling, complement activation, coagulation

regulation, and copper ion detoxification (**Figure 4E**). Consistent with these inflammatory signatures,

spatial mapping of cell type distributions (**Figure 4F**) and quantitative composition analysis (**Figure 4G**)

demonstrated marked enrichment of immune populations—particularly macrophages and T/NK

cells—within the transition zone relative to tumor and paratumor tissues[34]. The transition zone also

exhibited pronounced activation of the copper detoxification pathway (Figure 4H) and elevated expression

of metallothionein family genes (*MT2A, MT1E, MT1X, MT1M, MT1F*)[35](**Figure 4I**). Given that

metallothioneins modulate immune cell function and oxidative stress responses within the tumor

microenvironment, their spatial co-localization with immune infiltrates reinforces the biological coherence

of this OmniCell-defined niche. Concordant spatial gradients observed for acute inflammatory response,

complement activation, and coagulation regulation pathways (**Supplementary Figure 4E**) further

delineate a spatially coordinated inflammatory-metabolic program characteristic of the invasive tumor

margin. These findings establish the transition zone as a molecularly and immunologically distinct

microenvironment uniquely resolved by OmniCell. More broadly, our multi-scale analyses demonstrate

that OmniCell embeddings effectively bridge molecular and spatial hierarchies—from accurate cell-type

recovery at single-cell resolution, through faithful anatomical domain reconstruction at tissue scale, to de

novo identification of functionally distinct niches within complex pathological tissues.

OmniCell achieves state-of-the-art performance in cell type annotation and spatial deconvolution.

OmniCell was designed to jointly leverage scRNA-seq and ST data, capturing spatially resolved contextual

information while learning fine-grained cellular features. By integrating these complementary modalities,

OmniCell enhances representation capacity, more faithfully characterizes cellular states, and resolves

cellular heterogeneity within complex biological environments. To quantify these advantages, we

systematically benchmarked OmniCell against leading foundation models for single-cell analysis, including

scGPT, scFoundation, and Geneformer. Across five datasets, OmniCell consistently achieved the highest

annotation performance as reflected by F1 score and accuracy (**Figure 5A**), demonstrating superior

generalizability across heterogeneous biological contexts. A particularly stringent test of annotation quality

is the ability to identify rare cell populations. On the Zheng68k dataset (**Figure 5B and 5C**), OmniCell

predicted CD34+ cells with 94% accuracy despite their comprising only 0.29% of the dataset—a

performance 13 percentage points higher than the next-best models, scGPT and Geneformer. For extremely

rare populations, OmniCell achieved 20% accuracy for CD4+/CD45RA+/CD25- Naive T cells (2.8% of

dataset) and 6% for CD4+ T Helper2 cells (0.14% of dataset), whereas all competing models failed entirely.

Summary metrics across all cell types (**Figure 5E**) and full confusion matrices (**Supplementary Figure 5A–**

B) further underscore OmniCell's capacity to capture biologically meaningful low-frequency states.

Robust annotation must also generalize across technical batches. Using the two-batch hPBMC dataset

with train-test splits across batches, UMAP projections showed that OmniCell maintained close alignment

between predicted and true labels (**Figure 5D**), outperformed competing models across all metrics (**Figure**

5E), and achieved the most accurate predictions in full-cell-type confusion matrices (**Supplementary**

Figure 5C). This demonstrates its ability to disentangle technical confounders for scalable cross-study

annotation. Consistent superiority was observed on the lung cancer dataset (**Figure 5E; Supplementary**

Figure 5D), confirming robust performance across complex, heterogeneous contexts. These advantages

extend naturally to spatial transcriptomics, where cell-type annotation must additionally respect tissue

architecture. Across 30 benchmark spatial datasets, OmniCell outperformed leading methods including

scGPT-spatial, Nicheformer, and cell2location (**Figure 5G**), with boxplots demonstrating the most stable F1

and accuracy distributions—indicative of stronger generalization. OmniCell further reconstructed spatial

maps that closely matched ground-truth cellular organization (**Figure 5F**), achieving higher fidelity than

competing methods (**Supplementary Figure 5E**). Confusion matrices from MERFISH-based mouse brain

atlas slices 9_0 and 11_2 (**Supplementary Figure 5F and 5G**) corroborated its superior accuracy in

recovering fine-grained tissue architecture.

In summary, OmniCell robustly integrates scRNA-seq and ST data, achieving superior performance in

cell-type annotation, rare cell detection, and spatial reconstruction. Its consistent results across diverse

datasets highlight both its robustness and utility for dissecting complex tissue architectures.

Discussion

In this study, we present OmniCell, a unified foundation model that integrates scRNA-seq and ST data

within a single architecture. By learning shared representations across genes, cells, and tissue spatial

organization, and by incorporating spatial neighborhood serialization, gene-specific MoE value embedder,

and a learnable symmetric bilinear output module, OmniCell achieves robust performance across diverse

tissues, platforms, and species. Importantly, it captures both cell-intrinsic gene programs and intercellular

spatial dependencies, enabling accurate zero-shot recovery of cellular structures, co-expression relationships,

and spatial domains.

We demonstrate OmniCell's capability across multiple downstream tasks at cellular, gene, and spatial

levels. At the cellular level, it enables integrative clustering, resolving heterogeneity across multi-slide and

multi-platform datasets while mitigating technical batch effects. At the gene level, OmniCell generates

cell-specific gene embeddings that capture expression programs of individual cell types, alongside dataset-level gene embeddings that reveal co-expression networks and regulatory relationships across cells. At the spatial level, the model produces spatially aware embeddings that enhance cell-type deconvolution and spatial domain delineation. In real tumor spatial datasets, OmniCell identifies transitional regions at tumor margins and highlights biologically relevant features such as immune cell enrichment and inflammation pathway activation, illustrating its capacity to provide meaningful insights across molecular, cellular, and tissue scales.

Despite its advances, OmniCell faces several limitations. Representation quality may decrease in extremely sparse spatial datasets, and rare cell types or lowly expressed genes may be less reliably captured. Training on tens of millions of cells and spots also requires substantial computational resources. Moreover, while OmniCell provides a unified multi-omic framework, modeling complex scenarios—such as dynamic spatial processes, additional omics layers, or cellular responses to perturbations—remains challenging, as models trained on control cells alone often fail to predict transcriptional shifts induced by chemical or genetic perturbations[36-38]. Future work could address these limitations by incorporating additional modalities, perturbation-aware objectives, and temporal or multi-modal spatial information, enhancing OmniCell′s applicability and biological insight.

Overall, OmniCell establishes a unified framework for multi-omic representation learning, bridging scRNA-seq and ST data. By capturing gene regulation, cellular states, and tissue spatial organization within a single architecture, it provides a versatile tool for exploring biological complexity across diverse contexts and lays the groundwork for future integrative studies in single-cell and spatial omics.

Methods

Data collection

Single-cell data for pretraining. scRNA-seq datasets were obtained from publicly available CELL×GENE repositories. For datasets originating from CELL×GENE, we used the CZ CELL×GENE Discover Census v.2023-07-15, retrieved from the project's publicly accessible Amazon S3 bucket. All datasets associated with this census release were downloaded directly from S3 to ensure full consistency with the published resource. The collected datasets span a broad range of human tissues, including brain, blood, lung, liver, digestive and immune systems, skin, and kidney, among others. The final corpus comprises 38 million human single cells across 45 tissues, covering multiple sequencing platforms, developmental stages, and disease contexts curated from the CZ CELL×GENE database. In total, single-cell profiles account for approximately 57.8% of all cells analyzed in this study.

Spatial transcriptomic data for pretraining. Spatially resolved transcriptomic data were generated

primarily using the STOmics Stereo-seq whole-transcriptome platform. These datasets originate from

large-scale internal resources produced at the BGI Research Institute. Stereo-seq data from multiple spatial

resolutions—including Bin20 and Cellbin—account for approximately 27.2% and 14.8% of the total cell

count, respectively. Portions of these datasets are publicly accessible through STOmics data portals,

whereas the remaining fraction is part of ongoing internal releases. All Stereo-seq data were processed

following the standard analysis pipeline, including spatial binning, UMI quantification, and spatial quality

assessment.

Datasets for downstream tasks and evaluations. To rigorously benchmark our framework across

diverse biological contexts, we curated a comprehensive collection of datasets spanning single-cell

analysis, batch integration, cell-type annotation, spatial domains, spatial deconvolution, and model

interpretability. This collection enables extensive evaluation across both single-cell and spatial

transcriptomic modalities.

For gene module analysis, we used the human single-nucleus transcriptomic dataset from Zhou[39]

and the human hippocampus dataset from both Alzheimer's disease (AD) and control samples profiled by

Stereo-seq[40]. Cell-level tasks, including clustering and annotation, were assessed using a diverse panel

of datasets primarily sourced from the CELL×GENE repository, covering multiple human and mouse

tissues (e.g., blood[41], kidney[42], brain[43], muscle[44]) and sequencing platforms to ensure

heterogeneity and cross-platform generalizability. Batch integration performance was further evaluated on

the hPBMC[45] and Zhou[39] datasets, while cell-type annotation benchmarks included five

CELL×GENE atlas-level datasets(e.g., Brain[46], Hippocampus[46], Cortex[47], Immune[48],

Great_Apes[49] in addition to the Zheng 68k[4] and hPBMC[45] datasets.

For spatial analyses—including clustering, domain delineation, and deconvolution—we employed

high-resolution spatial transcriptomic datasets, comprising the MERFISH mouse brain aging dataset (31

samples)[50] from the Allen Institute, the LC5-M Stereo-seq dataset[34] of invasive regions in human liver

cancer, and the osmFISH mouse cortex dataset[51] from the Linnarsson Lab. All datasets are publicly

accessible at https://modelscope.cn/datasets/PJSucas/OmniCell-test-data .

OmniCell architecture and pretraining

A unified foundation model was developed to generalize across scRNA-seq and ST data by encoding gene

identity, expression levels, and spatial information in an assay-agnostic format. Gene identities are

represented using Ensemble IDs (e.g., ENSG00000139618) and mapped to integer indices through a

standardized vocabulary to ensure consistency across platforms. Each cell is denoted as $i \in 1, 2, \ldots, N$,

where N is the total number of cells across all the datasets For each cell i, the model covers: (1) g_{ij}, the

integer-mapped identifier for the gene j; (2) v_{ij}, the raw expression value of the gene j in the cell i; and

(3) (x_i, y_i), the spatial coordinates of the cell i. While ST cells retain their native spatial positions,

scRNA-seq cells are assigned default coordinates (0, 0), preserving structural consistency while reflecting

the absence of spatial resolution in standard single-cell protocols.

Rank-Based Normalization of Gene Expression via Soft Rank Scaling

To harmonize expression values across sequencing technologies and mitigate platform-specific

variability[52], we implemented a rank-based normalization strategy. For each cell i, genes are ranked by

their expression levels in descending order, producing a rank ρ_{ij} for each gene j, where $\rho_{ij} \in \{1, 2, \ldots, G\}$,

and G represents the total number of unique expression values in cell i. To preserve expression ordering

while smoothing magnitude difference, we define a soft rank r_{ij} by linearly rescaling the ranks into a

predefined interval $[a, b]$ as follows:

$$r_{ij} = a + (b-a) \cdot \frac{G - \rho_{ij}}{G - 1} \tag{1}$$

In our implementation, we set $a = 0$ and $b = 5$, such that $r_{ij} \in [0, 5]$. This transformation retains the

relative expression order of genes within each cell while yielding a bounded, continuous representation

suitable for downstream modeling. The soft rank reduces the influence of extreme values and enables

effective integration across both scRNA-seq and ST datasets, without reliance on raw expression

magnitudes.

Construction of Input Sequences for Transformer Modeling

To integrate scRNA-seq and ST data into a unified Transformer-based modeling framework, we construct

gene-centric token sequences for each sample. The embedding of each token is derived from the sum of

two components: gene identity and expression level. This dual-component embedding design enables the

model to effectively combine discrete genetic signatures with context-dependent expression profiles,

facilitating a comprehensive analysis of transcriptional heterogeneity across diverse data modalities.

Unified Sequence Construction Across scRNA-seq and Spatial Transcriptomics. For scRNA-seq

data, each cell is treated as an independent sequence, with genes prioritized for inclusion based on

biological relevance: non-zero expression genes are selected first, followed by high-variance

zero-expression genes to maintain signal diversity. In the case of ST data, such as Stereo-seq, we extend

this approach by incorporating local tissue context. Each spatial token sequence is centered on a given cell

and includes gene-level profiles from its k nearest spatial neighbors, determined by Euclidean distance in

tissue coordinates. Using the same gene prioritization strategy as in scRNA-seq, n genes are selected from

each of the 1+k cells. The gene sequences from each cell are enclosed with <RNA_START> and

<RNA_END> tokens, resulting in a structured, multi-cellular input sequence that jointly encodes both

spatial relationships and transcriptional states. These special tokens serve as markers to denote the

boundaries of cellular contexts and guide position-aware encoding within the spatial neighborhood.

Gene Identity Embedding. Each gene is associated with a unique Ensemble identifier and mapped to

an integer index through a unified vocabulary. A shared embedding matrix $E_{\text{gene}} \in \mathbb{R}^{G \times d}$ then transforms

this index into a dense vector representation:

$$e_{ij} = E_{\text{gene}}(j) \tag{2}$$

where G is the total number of unique genes, d is the embedding dimension, and (i, j) refers to the

j-th gene in the i-th cell or spatial spot. This embedding captures global identity information and is

consistent across scRNA-seq and spatial transcriptomics.

Gene-Aware Value Embedder via MoE. Inspired by DeepSeek-V1[53], we introduce a gene-aware

MoE architecture to quantitatively embed each gene's expression level v_{ij}, capturing both the magnitude

and the biological context of transcriptional activity. This module is conditioned on the gene identity

embedding, enabling the model to adapt its encoding strategy based on gene-specific context. The

architecture integrates S shared experts that model general expression-magnitude patterns across all genes,

together with R routing experts that are dynamically selected through a learnable routing network,

allowing the expression magnitude of each gene to be encoded in a gene-sensitive and context-adaptive

manner.

The routing mechanism takes the gene embedding $\mathbf{e}_{ij}$ as input and produces a softmax distribution

over the routing experts:

$$p_{ij} = \text{softmax}(W_2 \cdot \text{ReLU}(W_1 \cdot e_{ij})) \in R^{r \times 1} \tag{3}$$

454   where r denotes the number of routing experts.

455   To enhance the representational capacity and generalization of the model, only the top k experts with

456   the highest probabilities (denoted as $\mathcal{T}_{ij}$) are activated for each gene expression embedding. A sparse

457   weight vector $\boldsymbol{\alpha}_{ij} \in \mathbb{R}^r$ is constructed, retaining the original routing probabilities for the selected experts

458   while assigning zero to the remainder:

459
$$\alpha_{ijk} = \begin{cases} p_{ijk}, & k \in \mathcal{T}_{ij} \\ 0, & \text{otherwise} \end{cases}. \tag{4}$$

460   Importantly, these sparse weights are not renormalized post-selection, preserving the relative magnitude of

461   expert contributions and maintaining the inherent sparsity.

462   Concretely, each expert (comprising both shared experts and routing experts) transforms the

463   expression value v_{ij} into an embedding vector. The shared experts, consisting of S linear

464   transformations, capture general patterns of expression magnitude:

465
$$\mathbf{h}_{ij}^{\text{shared}} = \sum_{s=1}^{S} \text{SharedExpert}_s(v_{ij}), \tag{5}$$

466   while the routing experts provide gene-specific transformations aggregated according to the sparse routing

467   weights:

468
$$\mathbf{h}_{ij}^{\text{routing}} = \sum_{k=1}^{R} \alpha_{ijk} \cdot \text{RoutingExpert}_k(v_{ij}). \tag{6}$$

469   The final embedding of the gene expression level is the sum of these two components:

470
$$h_{ij} = h_{ij}^{\text{shared}} + h_{ij}^{\text{routing}}. \tag{7}$$

This architecture enables the model to flexibly and selectively capture both global expression trends and

gene-specific nuances, with the sparse routing mechanism improving the model's expressive power and

generalization.

Load Balancing for Expert Utilization. To prevent expert collapse and encourage balanced expert

usage, we employ DeepSeek-V1's[53] load balancing loss. Let T be the total number of tokens in a batch

and K the number of active experts per token. The expected routing frequency for expert k is

approximated as:

478
$$f_k = \frac{1}{K \cdot T} \sum_{i,j} \mathbb{1}(k \in \mathcal{T}_{ij}) \cdot R, \tag{8}$$

479 and the average routing probability as:

$$P_k = \frac{1}{T} \sum_{i,j} p_{ijk}. \tag{9}$$

481 The load balancing loss is computed as:

$$\mathcal{L}_{\text{balance}} = \sum_{k=1}^{R} f_k \cdot P_k, \tag{10}$$

which encourages both the frequency and the confidence of expert routing to be uniformly distributed
across experts. This regularization promotes diversity among expert participation and improves
generalization.

Transformer with Two-Dimensional Rotary Positional Encoding

We developed a transformer-based encoder architecture incorporating two-dimensional rotary positional
encoding (2D-RoPE), inspired by the Qwen2-VL[54] model. The architecture follows the standard
Transformer encoder block structure, consisting of multi-head self-attention, feedforward networks, RMS
normalization, and residual connections. Spatial inductive bias is introduced via block-diagonal rotations
acting on the query and key vectors, parameterized by the 2D coordinates associated with each token. Let
the input to the Transformer be

$$\mathbf{E} \in \mathbb{R}^{B \times T \times d} \tag{11}$$

494 where B is the batch size, T is the sequence length and d is the model dimension. The model is
495 composed of L identical Transformer layers, each applying the following operations.

496   **Root Mean Square Normalization.** Each layer begins with RMS normalization applied to the input
497 $\mathbf{E}$. For a token embedding $\mathbf{e}_t \in \mathbb{R}^d$, the normalized vector is defined as:

$$\text{RMSNorm}(\mathbf{e}_t) = \frac{\mathbf{e}_t}{\sqrt{\frac{1}{d} \sum_{j=1}^{d} e_{t,j}^2 + \varepsilon}} \odot \boldsymbol{\gamma} \tag{12}$$

where ε is a small constant for numerical stability, and $\boldsymbol{\gamma} \in \mathbb{R}^d$ is a learned scaling vector.

Multi-Head Self-Attention with Two-Dimensional Rotary Encoding. We introduce a multi-head
self-attention mechanism enhanced with two-dimensional rotary encoding. Initially, token embeddings are

projected to form queries, keys, and values, which are then reshaped for multi-head attention. Each

attention head is divided into m four-dimensional subspaces, with each subspace further split into two 2D

planes corresponding to the spatial x- and y-dimensions. These planes are rotated independently using

log-spaced angular frequencies, allowing the model to capture positional relationships in both spatial

dimensions. Attention scores are computed for each head by performing dot-product interactions between

the rotated queries and keys, and the outputs from all heads are concatenated and projected to produce the

final attention output. This method ensures efficient integration of spatial information within the attention

mechanism.

Feedforward Network with SwiGLU. Following attention and residual connection, the output is

passed through a feedforward network:

512
$$\mathbf{E}_{res} = \mathbf{E} + \mathbf{E}_{attn}, \quad \mathbf{E}' = \mathbf{E}_{res} + \mathrm{FFN}(\mathrm{RMSNorm}(\mathbf{E}_{res})) \tag{13}$$

513 The feedforward network consists of a Swish-Gated Linear Unit:

514
$$\mathrm{FFN}(\mathbf{z}) = \mathbf{W}_3 \left[\mathrm{Swish}(\mathbf{W}_1\mathbf{z}) \odot (\mathbf{W}_2\mathbf{z}) \right], \quad \mathrm{Swish}(x) = x \cdot \sigma(x) \tag{14}$$

where $\sigma(x)$ denotes the sigmoid function，$\odot$: element-wise (Hadamard) product.

Layer Stacking. The above operations are repeated across L layers. Each layer processes the output

from the previous one, maintaining the dimensionality $\mathbf{E} \in \mathbb{R}^{B \times T \times d}$ throughout.

Symmetric Bilinear Output Module and Pretraining Objective

To more faithfully capture the subtle dynamics of expression levels, we reformulate the pretraining

objective as a soft-rank regression task. In contrast to conventional approaches that discretize expression

values into predefined bins for classification, our method preserves the intrinsic continuity and relative

ordering of expression by predicting their soft ranks within a local semantic context. This formulation

avoids the loss of relational information between expression levels introduced by discretization, enabling

finer-grained modeling of expression variation.

Let $\mathbf{E} \in \mathbb{R}^{B \times T \times d}$ denote the final Transformer representations, where B is the batch size, T the

sequence length, and d the hidden dimension. The sequence is partitioned into C semantic cells, each of

length $t + 2$ and comprising t content tokens flanked by two boundary markers. In our setup, we adopt

$C = 1$ for single-cell transcriptomes and $C = 10$ for spatial transcriptomics, where each cell

corresponds to a spatial region encompassing a focal cell and its neighbors.

For each cell, the t interior tokens are extracted and reshaped into $\mathbb{R}^{B \times C \times t \times d}$. A pooled cell-level

representation $\mathbf{h}_c \in \mathbb{R}^d$ is obtained via mean aggregation:

532
$$\mathbf{h}_c = \frac{1}{t} \sum_{i=1}^{t} \mathbf{e}_{c,i}. \tag{15}$$

533   We then apply a trainable symmetric matrix transformation:

534
$$\mathbf{W}_{\text{sym}} = \frac{1}{2}(\mathbf{W} + \mathbf{W}^\top), \tag{16}$$

535   and compute token-level scores via dot product:

536
$$s_{c,i} = \left(\mathbf{W}_{\text{sym}} \mathbf{h}_c\right)^\top \mathbf{e}_{c,i}. \tag{17}$$

537   These scores $s_{c,i} \in \mathbb{R}$ represent predicted soft ranks within each semantic cell. To supervise this

538   regression, ground-truth soft ranks $\hat{s}_{c,i}$ are derived from observed expression magnitudes. We adopt the

539   Huber loss (Robust estimation of a location parameter) as the training objective. The loss is computed over

540   masked positions as follows:

541
$$\mathcal{L}_{\text{MLM}} = \frac{1}{|\mathcal{M}|} \sum_{(c,i) \in \mathcal{M}} \mathcal{L}_\delta (s_{c,i} - \hat{s}_{c,i}), \tag{18}$$

542   where $\mathcal{M}$ denotes the set of masked positions, and $\mathcal{L}_\delta(\cdot)$ is the Huber loss function defined as:

543
$$\mathcal{L}_\delta(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } |x| \leq \delta, \\ \delta(|x| - \frac{1}{2}\delta) & \text{otherwise.} \end{cases} \tag{19}$$

544   We set the threshold parameter to $\delta = 1$ in all experiments. This formulation encourages stability

545   during optimization while preserving sensitivity to moderate errors.

546   To encourage uniform usage of semantic cells and avoid representational collapse, we incorporate a

547   load balancing regularization term $\mathcal{L}_{\text{balance}}$, previously introduced. The full pretraining loss is given by:

$$\mathcal{L}_{\text{total}} = \mathcal{L}_{\text{MLM}} + \alpha \cdot \mathcal{L}_{\text{balance}}, \quad \alpha = \begin{cases} \alpha_0 & \text{if } \mathcal{L}_{\text{MLM}} < 10, \\ 10 & \text{otherwise,} \end{cases} \tag{20}$$

549   where α_0 is a small constant (typically 0.05). This adaptive weighting ensures that cell utilization

550   remains balanced even during early or unstable phases of regression training.

Construction the Cell and Gene Embeddings

Cell Embedding. We compute cell embeddings in two steps. First, final-layer hidden states are

aggregated using a weighted averaging strategy in which each gene contributes based on whether it is

expressed (non-zero count), ensuring that only actively expressed genes influence the representation.

Second, an optional refinement step projects the aggregated embedding into a low-dimensional subspace

using a symmetric projection matrix learned during pretraining. This projection reduces technical

noise—including dropout artifacts and batch effects—while preserving biologically meaningful structure

in the resulting embedding space.

Specifically, the matrix $\mathbf{W}_{\text{sym}} \in \mathbb{R}^{d \times d}$ is trained directly during pretraining as a symmetric

transformation. We perform spectral decomposition:

$$\mathbf{W}_{\text{sym}} = \mathbf{Q} \Lambda \mathbf{Q}^{\top} \tag{21}$$

562   where Λ contains eigenvalues and $\mathbf{Q}$ contains orthonormal eigenvectors. We select the top-m

563   eigenvectors that collectively explain at least τ of the total spectral energy, computed using the absolute

564   values of eigenvalues. This forms an orthonormal basis $\mathbf{V} \in \mathbb{R}^{d \times m}$ used to denoise each cell embedding

565        $\mathbf{h}_c$:

$$\mathbf{h}_c = \mathbf{V}^{\top} \mathbf{h}_c \tag{22}$$

The projected vector $\mathbf{h}_c$ retains task-relevant variation while suppressing noise directions that dominate in

raw representations. In practice, this spectral projection improves the separation of cell types and reduces

technical confounding across spatial and single-cell modalities.

Gene Embedding. Gene embeddings were derived at two complementary levels, capturing both local

and global gene properties. At the cellular level, the final-layer hidden states from a Transformer encoder

were extracted, producing a contextualized embedding for every gene within each cell. To reduce technical

noise, these embeddings were projected onto a low-dimensional subspace as described previously, yielding

representations that reflect cell-state- or cell-type–specific gene characteristics.

Dataset-level gene embeddings were then obtained by integrating these cell-level gene embeddings

across all cells, with the explicit objective of bringing genes with similar expression patterns closer in the

embedding space. **Supplementary Figure 2A** illustrates this integration process, showing how cell-level

embeddings are combined to form dataset-level representations, which can then be utilized for further

analysis. A cell similarity graph was constructed based on the previously described cell embeddings,

connecting each cell to its k nearest neighbors, with edge weights determined using a decaying kernel[55,

56]. Each gene's cell-level embeddings were propagated through this graph using a generalized PageRank

graph neural network (GPR-GNN)[33], which aggregates information from neighboring cells while

optimizing reconstruction of observed gene expression. This procedure reinforces co-expression patterns

and context-dependent relationships, ensuring that genes exhibiting similar transcriptional behavior across

the dataset acquire similar dataset-level embeddings. The final embedding for each gene was obtained by

averaging its propagated embeddings across all cells, capturing both intrinsic expression patterns and

global, dataset-wide functional properties.

Pretraining of the OmniCell Model

All single-cell and spatial transcriptomic datasets were first converted from the AnnData (h5ad)

format into LMDB files to enable high-throughput sequential access and memory-efficient data streaming

during large-scale optimization. Pretraining was conducted in two sequential stages to hierarchically

integrate spatial context and single-cell precision within the same model.

In the first stage, spatial transcriptomic data were used to expose the model to tissue-level

organization and neighborhood-dependent gene co-variation, allowing it to capture local transcriptional

continuity across adjacent spots. Because spatial data are typically noisier and have lower molecular

capture efficiency, a moderate masking ratio was adopted to balance information removal and

reconstruction difficulty. Specifically, the non-zero expression values of central nodes were randomly

masked at a rate of 0.3, and those of neighboring nodes at 0.05, while zero entries were largely preserved.

This design guided the model to infer missing gene expression patterns from partially masked spatial

contexts, enhancing its ability to recover structured co-expression signals while maintaining training

stability.

In the second stage, the pretrained model was further optimized on scRNA-seq datasets, which

provide higher data quality and reduced technical noise. To fully exploit the intrinsic information

redundancy of transcriptomic data—analogous to pixel redundancy in natural images—a high masking

ratio was employed to strengthen contextual inference. The non-zero expression values of each cell were

masked at a rate of 0.3, while zero expression values were masked at a rate of 0.1, forcing the model to

predict large portions of the expression vector based on the remaining signals. This large-mask regime

encourages the model to learn robust gene–gene dependencies and to generalize effectively across diverse

cellular states.

Training was performed using a large batch size (3,200 cells per step) evenly distributed across 32

NVIDIA A100 GPUs (40 GB memory each), enabling efficient optimization at scale. Gradient clipping

(maximum norm = 1.0) was applied to stabilize optimization. The AdamW optimizer—a weight-decoupled

variant of Adam—was used with a learning rate of 1×10^{-6}, weight decay of 0.01, and momentum

coefficients $\beta_1 = 0.9$ and $\beta_2 = 0.99$. A cosine annealing scheduler with a minimum learning rate of 5×10^{-7}

ensured smooth and stable convergence.

Downstream tasks

Evaluation of Cell-Type Specificity of Gene Embeddings at the Cell Level

To assess the cell-type specificity captured by the learned gene embeddings, we analyzed the gene-level

representations derived from the final layer of each foundation model, including OmniCell, scGPT,

Geneformer, and scFoundation. This evaluation reflects how well each model learns the context-dependent

relationships for each gene, providing insight into how gene embeddings are organized across different cell

types. For every gene, we collected its embeddings across all cells and performed unsupervised clustering

in the corresponding embedding subspace. The resulting gene-specific clustering was then quantitatively

compared with the known cell-type annotations using the Adjusted Rand Index (ARI). This procedure

yields a cell-type specificity score for each gene, which reflects the degree to which its embedding

structure recapitulates known cellular identities. Higher ARI values indicate that the model captures more

distinct and biologically meaningful gene representations across diverse cell populations.**Gene–cell**

importance scores were obtained by pooling gene–gene attention weights across all layers of the

transformer. To further evaluate the performance of our model, we performed a multiclass ROC analysis

based on the gene–cell importance scores derived from the attention weights. For each gene, we calculated

a gene–cell importance matrix, which reflects how each gene's importance varies across different cells

based on the attention scores. This matrix was then used to assess the model's ability to discriminate

between cell types. Specifically, for each cell type, we computed a prototype importance profile by

averaging the importance scores of all cells in that type. The similarity between each cell's gene

importance vector and the prototype for each cell type was calculated using cosine similarity, yielding

predicted scores that reflect the likelihood of each cell belonging to a specific cell type. The resulting

multiclass ROC analysis, including AUC scores, allowed us to assess how well the model captures and

distinguishes the context-dependent gene relationships across diverse cell populations. This analysis was

conducted using the Zhou dataset, with 2000 highly variable genes selected for evaluation, providing a

comprehensive assessment of the model's ability to learn cell-type specificity in cell-level gene

embeddings.

Identification of Cell-Type–Specific Genes from OmniCell Cell-Level Gene Embeddings

To identify genes that characterize specific cell types or cellular states, we developed a method leveraging

cell-level gene embeddings generated by the foundation model. For each gene, its embedding distributions

across cells were compared between a target cell type and all other cells using the mean one-dimensional

Wasserstein distance across embedding dimensions, thereby quantifying how distinctively the gene is

represented in the target population. Genes were ranked by these distance scores, and the top candidates

were further filtered based on their transcriptional abundance in the corresponding single-cell data,

retaining only those with higher mean expression in the target cell type relative to others. This two-step

procedure—embedding-based discrepancy analysis followed by expression validation—yields

interpretable gene sets that reflect both representation-level and expression-level specificity, providing a

principled approach for discovering cell-type–specific or state-associated marker genes from foundation

model embeddings. To evaluate the consistency between marker genes identified from OmniCell

embeddings and those from conventional methods, we compared the embedding-derived markers with

cell-type markers generated by the Seurat[57] *FindAllMarkers* function. Results showed that the majority

of embedding-derived markers for most cell types overlapped with the top marker gene lists from

Seurat[57] *FindAllMarkers*, which strongly validated the biological reliability of the markers identified via

OmniCell embeddings. Beyond the overlapping known markers, OmniCell embeddings also identified

additional cell-type-enriched novel markers that were not detected by the conventional method. To assess

the discriminative power of these novel markers, we performed ROCanalysis and calculated the

AUCvalues. Further confirming the validity of both the overlapping and novel markers derived from

OmniCell embeddings.

Validation of Co-expression Structure in OmniCell Dataset-Level Gene Embeddings

To determine whether dataset-level gene embeddings learned by OmniCell capture biologically meaningful

co-expression structure, we constructed a gene–gene similarity network using cosine similarity or a

Gaussian-transformed Euclidean metric and applied soft thresholding to enhance high-confidence

relationships before identifying modules via Louvain community detection. We evaluated the biological

relevance of these embedding-derived modules using two independent datasets: the Zhou scRNA-seq

dataset and a Stereo-seq dataset comprising both Alzheimer's disease and control samples. Cell-type

marker genes identified from OmniCell cell-level embeddings and region-specific marker genes curated

from prior literature[40] served as biological priors for the scRNA-seq and ST datasets, respectively. We

then quantified whether these biological priors aligned with the embedding-derived network structure by

calculating expression score profiles for each module. Higher scores indicate that genes grouped together

in the embedding space also exhibit coordinated expression patterns at the transcriptomic level, a defining

property of biologically meaningful co-expression modules. Consistently higher scores in modules

enriched with known biological marker genes indicate that OmniCell's dataset-level gene embeddings

capture biologically coherent and context-dependent co-expression patterns rather than arbitrary latent

structure. These findings support that the learned embeddings provide a faithful representation of

regulatory organization across both scRNA-seq and ST datasets.

Single-Cell and Spatial Cell Clustering

To evaluate clustering performance, we assessed OmniCell alongside both single-cell and spatial

foundation models. The single-cell models included scGPT, Geneformer, and scFoundation, whereas

scGPT-spatial and Nicheformer represented spatial models. For models not pretrained on mouse datasets,

homologous gene mapping was applied to enable cross-species evaluation. To ensure fairness across
modalities, a unified gene selection strategy was applied: the top 2,000 highly variable genes were used for
all scRNA-seq datasets, including human kidney and blood, as well as mouse brain and muscle datasets,
while the full gene set was retained for the MERFISH mouse brain spatial transcriptomics dataset.
Embeddings generated by each model were clustered using the Leiden algorithm, with the resolution
systematically varied from 0.1 to 1.0; for each model, the resolution yielding the highest agreement with
reference annotations was selected as the final result. Clustering performance was quantified using
Adjusted Rand Index (ARI) and Normalized Mutual Information (NMI), providing complementary
measures of concordance between predicted clusters and biologically defined cell identities.

Batch correction evaluation

For batch correction tasks, model-derived cell embeddings were evaluated using the scIB[58]
benchmarking framework. Four baselines—PCA, scGPT, Geneformer, and scFoundation—were included
for comparison against OmniCell. Two human scRNA-seq datasets, hPBMC and the Zhou dataset, were
used to assess performance. To ensure comparability across models, the top 2,000 highly variable genes
were selected as input features for all evaluations. The scIB framework reports two composite metrics: a
biological conservation score, reflecting how well intrinsic cellular structure and annotations are preserved
across batches, and a batch correction score, measuring the effectiveness of batch integration based on
silhouette and graph connectivity–based metrics. Higher values in both scores indicate a more optimal
balance between maintaining biological fidelity and mitigating batch-driven technical variation.

Noise robustness evaluation

Noise robustness was assessed under increasing dropout perturbations using the Zhou dataset. Starting
from the original count matrix, random dropout noise was incrementally introduced from 0.0 to 1.0 in steps
of 0.05. To ensure fairness across methods, the top 2,000 highly variable genes were used as input features
for all models. Three foundation models—scGPT, Geneformer, and scFoundation—were evaluated
alongside OmniCell. For each noise level, cell embeddings were regenerated using the pretrained model
and subsequently subjected to Leiden clustering at a fixed resolution of 0.1, which was empirically
identified as optimal across models. Clustering consistency was quantified using Normalized Mutual
Information (NMI), selected as the primary metric due to its stability under varying cluster numbers and its
sensitivity to biological structure despite increasing data sparsity. All experiments were performed with a
fixed random seed to ensure reproducibility, and the resulting NMI–dropout curves capture each model's
robustness to transcriptional noise.

Spatial Domain Identification

To evaluate the model's ability to capture spatial tissue organization, we performed spatial domain

identification on two spatial transcriptomics datasets. For the mouse cortex slice with predefined

anatomical annotations, all 33 genes were used during inference; cell embeddings were generated in spatial

mode by integrating neighboring transcriptomes and averaging across the local neighborhood, followed by

construction of a k-nearest neighbor graph and Leiden clustering. The clustering resolution was optimized

from 0.1 to 1.0 (step size 0.1) by maximizing the combined Adjusted Rand Index (ARI) and Normalized

Mutual Information (NMI) relative to ground truth. For the LC5-M human lung cancer dataset, which

lacks predefined annotations but contains tumor, peri-tumoral transition, and non-malignant compartments,

we used the top 500 highly variable genes consistent with model training, while baselines used the

standard top 2,000 genes; embeddings were again generated in spatial mode, and a k-nearest neighbor

graph was used to accommodate broader tissue structure and tumor heterogeneity before applying Leiden

clustering at a fixed resolution of 0.1. Performance was compared against two foundation models

(scGPT-spatial and NicheFormer) and three expert spatial methods (SCAN-IT[44], SEDR[59], and

SpaGCN[60]). While ARI and NMI confirmed higher clustering accuracy on the mouse cortex dataset,

analysis of LC5-M revealed that only our model resolved a distinct peri-tumoral transitional domain. Gene

Ontology enrichment analysis identified significant activation of acute-phase response, inflammatory

signaling, complement activation, coagulation regulation, and copper ion detoxification pathways within

this region. Cell type composition analysis further revealed marked enrichment of immune populations.

Spatial gradient analysis using SLOPER[61] demonstrated elevated expression of metallothionein family

genes with directional streamlines delineating the spatial organization of these molecular programs.

Cell-Type Annotation

For quality estimation on cell embedding in supervised classification, we took the top 2,000 most variable

genes from each dataset and extracted cell embeddings from the corresponding models. Three pretrained

baseline models (scGPT, Geneformer, and scFoundation) were applied as baselines with respect to

OmniCell. ReLU activation and layer normalization were used to train the AdamW optimizer on a

three-layer multilayer perceptron (learning rate 1×10^{-4}, 20 epochs) in order to learn the cell types in the

embeddings. Assorted as 5-fold stratified cross-validation of five benchmark datasets (Brain, Hippocampus,

Cortex, Immune, Great_Apes), reciprocal cross-batch validation (training in first batch and testing on

second batch) and 80/20 stratified train–test split of large sparse datasets (e.g., zheng68k) were used. The

classification performance was determined by accuracy and macro-averaged F1-score with mean and

variance calculated across folds or batches. To maintain reproducibility, all experiments were performed on

fixed random seeds.

Spatial transcriptomic deconvolution

Cell embeddings derived from the foundation model were used to perform spatial cell-type deconvolution,

following a strategy conceptually similar to Tangram[62]. Single-cell reference profiles and spatial

transcriptomic data were aligned based on their shared genes. The single-cell embeddings served as the

reference space, and spatial embeddings were projected onto this space through an optimization procedure

to estimate the proportional contribution of each cell type to each spatial location. The resulting proportion

matrix defined the inferred cellular composition per spot, from which the dominant cell type was assigned

based on the highest contribution weight. Two spatial foundation models—scGPT-spatial and

Nicheformer—were included as baselines for comparison with OmniCell. Deconvolution performance was

evaluated using accuracy and macro-averaged F1 score.

Benchmarking against competing methods

Comparisons against foundation models

To thoroughly assess the OmniCell model, we performed benchmarking on various foundational

single-cell and spatial omics models (scGPT, Geneformer, scFoundation, scGPT-spatial, Nicheformer).

The extraction of embeddings for each foundational model as presented by the above-in-text

implementation described below: For scGPT we used the scGPT_human pre-trained weights from the

official repository and obtained cell embeddings for the model pre-trained on complete human data using

the pipeline described in the original paper. The Geneformer-V2-316M model (from Hugging Face's

official repository) has been utilized for Geneformer: to obtain embeddings from the penultimate layer and

to group gene-level embeddings into single-cell representations by mean pooling; for scFoundation we

obtain pre-trained weights (models.ckpt) from the official repository and to estimate embeddings using the

standard inference procedure presented by the authors; For the scGPT-spatial model we extracted the

scGPT_spatial_v1 model weights specifically pretrained on spatial transcriptomics data. For Nicheformer,

we applied pre-trained model parameters that were trained on the SpatialCorpus-110M dataset.

Embeddings provided by the base models were treated as task specific inputs.

Comparisons against spatial domain detection methods

We also compared OmniCell with specialized spatial domain detection methods (SpaGCN v1.2.7, SEDR,

and SCAN-IT) to evaluate its performance in spatial clustering tasks. For SpaGCN, we used the official

implementation version, utilizing the default hyperparameters proposed in the original paper. Training was

performed using a spatial graph built from the spatial coordinates of cells in an attempt to find spatially

coherent regions. The official implementation was applied for SEDR, adhering to their routine workflow

for producing low-dimensional representations necessary for spatial clustering by combining profiles of

gene expression with spatial locations. The official SCAN-IT implementation followed the recommended

spatial domain identification process to identify tissue domains and cellular microenvironments through

the combination of gene expression and spatial information.

Code availability

OmniCell is packaged and distributed as an open-source, publicly available repository at

BGIResearch/omnicell

Acknowledgement

Author contributions

Figure 1. Overview of the OmniCell Model, Pretraining Data, and Evaluation.

An overview of the OmniCell model, its pretraining data, architecture, downstream tasks, and ablation

study. **(A)** Composition of pretraining data. OmniCell was pretrained on 67 million cells comprising single-cell RNA sequencing (scRNA-seq; 57.8%) and Stereo-seq spatial transcriptomics (ST; 42.2%) data across diverse tissues and cell types. **(B)** Model architecture. OmniCell integrates scRNA-seq and ST data within a unified framework. For scRNA-seq, cells are encoded as ordered gene sequences based on expression. For ST, spatial context is incorporated through neighborhood graphs capturing local cellular relationships. Gene expression values are normalized via soft-rank transformation. A mixture-of-experts (MoE) gene-aware value embedding module adaptively encodes expression levels. The architecture comprises 10 Transformer layers, with a symmetric bilinear output module that jointly models cell–gene relationships to generate unified embeddings. **(C)** Downstream tasks. Model performance was assessed on ST-specific tasks (spatial clustering, domain identification, deconvolution, and gene module analysis) and scRNA-seq tasks (cell clustering, cell-type annotation, gene module analysis, batch correction, and marker gene identification). **(D)** Ablation analysis. Systematic removal of model components—including the MoE value embedder, spectral subspace projection, symmetric bilinear module, and ST pretraining data—quantifies their individual contributions to model performance.

Figure 2. Cell-level and dataset-level gene embeddings capture cell-type-specific signatures and functional modules.

(**A**) Benchmarking of cell-level gene embeddings across foundation models using Adjusted Rand Index (ARI) for clustering concordance with known cell types. (**B-C**) UMAP visualizations of cell-level embeddings for APOE and TREM2 expression (left) and corresponding cell-type annotations (right). (**D**) Assessment of cell-type separability via gene importance vectors. ROC curves depict the discriminative capacity of attention-derived gene importance score vectors across different cell types.(**E**) Dot plot of marker genes identified from cell-level embeddings across six major cell types (ASC, EX, INH, MG, ODC, OPC), with expression ratio (dot size) and level (color). (**F**) Proportion of embedding-derived markers that overlap with cell-type markers generated via the Seurat *FindAllMarkers* function. (**G**) Discriminative power (AUC) of novel embedding-derived marker genes not overlapping with those generated by the Seurat *FindAllMarkers* function. (**H**) UMAP visualization of the Zhou Alzheimer's disease scRNA-seq dataset was colored by cell type. (**I**) UMAP visualization of module expression scores for six embedding-derived gene modules (Modules 0-5). (**J**) Heatmap showing six modules of DEGs classified by OmniCell, with each correlated to distinct biological processes. (**K**) Anatomical parcellation of AD patients' hippocampus showing cornu ammonis subfields (CA1–CA4), dentate gyrus (DG), and granule cell layer (GC). (**L**) Spatial distribution of module expression scores in AD patients' hippocampus spatial transcriptomics data. (**M**) Dot plot of module expression across hippocampal regions, with expression ratio (dot size) and level (color).

Figure 3. Benchmarking OmniCell for cross-species generalization, batch integration, and expert modularity.

(A) OmniCell consistently achieved the highest clustering performance (ARI and NMI) across all datasets, surpassing existing models. Despite being pretrained solely on human data, OmniCell maintained clear cell-type separability in mouse datasets through homology-based gene projection, demonstrating strong cross-species transferability. **(B–C)** UMAP visualizations of representative human and mouse datasets show that OmniCell embeddings form compact and biologically coherent clusters with superior intra-cluster cohesion and inter-cluster separation compared with scGPT and GeneFormer.

(D) Scatterplot comparing batch-correction performance (x-axis) against biological conservation (y-axis) on the Zhou dataset. Each point represents a different method. **(E)** UMAP embeddings of the Zhou dataset (11 batches, 6 cell types) generated by five methods. Top row: cells colored by cell type (ASC, EX, INH, MG, ODC, OPC). Bottom row: same embeddings colored by batch (batches 11–21).

(F) Under progressive random dropout, OmniCell retained the highest clustering stability, indicating noise-resilient and biologically grounded representations. **(G)** Heatmap of mixture-of-experts (MoE) activation patterns across immune cell populations. Rows represent 16 cell types sorted by lineage class; columns denote individual experts (Expert_0 through Expert_9). Color scale indicates activation z-scores, with red reflecting positive and blue reflecting negative enrichment.

Figure 4. Multi-level spatial evaluation of OmniCell across MERFISH and Stereo-seq datasets.

(**A**) Boxplots comparing clustering performance (NMI and ARI) of OmniCell, scGPT-spatial, and Nicheformer across 31 MERFISH-based mouse brain atlas datasets. (**B**) UMAP visualization of cell embeddings from MERFISH slice 10_0 colored by cell type annotations. (**C**) Spatial coordinate maps comparing ground truth annotations with clustering results from OmniCell (NMI = 0.8975), scGPT-spatial (NMI = 0.8536), and Nicheformer (NMI = 0.3628). (**D**) Spatial segmentation of the Stereo-seq LC5-M liver cancer dataset by six methods (OmniCell, scGPT-Spatial, Nicheformer, SCAN-IT, SpaGCN, and SEDR). (**E**) Gene Ontology enrichment analysis of Cluster 3 (transition zone). (**F**) Spatial distribution of cell types in the Stereo-seq LC5-M liver cancer dataset. (**G**) Cell type composition across tumor, transition zone, and paratumor regions. (**H**) Spatial plot showing the activity of the detoxification of copper ion pathway, where white streamlines indicate the gradient direction. (**I**) Spatial expression patterns of the pathway genes of **H** (*MT2A, MT1E, MT1X, MT1M, MT1F*), with white streamlines depicting expression gradients.

Figure5. Benchmarking OmniCell for single-cell cell-type annotation and spatial transcriptomics deconvolution.

(**A**) Dot plots of cell-type annotation performance across five scRNA-seq datasets (Brain, Hippocampus, Cortex, Immune, Great Apes). F1 score (left) and accuracy (right) were evaluated using 5-fold cross-validation for OmniCell, scGPT, Geneformer, and scFoundation. (**B**) Confusion matrices illustrating OmniCell's accurate annotation of rare cell types in the Zheng68k dataset, compared with other models. (**C**) UMAP projections of the Zheng68k dataset showing original annotations (left) and predicted labels (right). Twelve cell types are displayed, including rare populations such as CD34+ progenitors and dendritic cells. (**D–E**) UMAP visualization of the hPBMC dataset comparing original annotations (left) with predicted labels (right) across eleven cell types. (**E**) Line plots of annotation metrics (F1, Accuracy, Precision, Recall) evaluated on three datasets (Zheng68k, hPBMC, LungCancer). Each line represents one method; error bars indicate variation across folds or batches. (**F**) Boxplots comparing spatial deconvolution performance across 30 spatial transcriptomics datasets. Macro F1 score (left) and accuracy (right) are shown for OmniCell, scGPT-spatial, Nicheformer, and cell2location. (**G**) Spatial reconstructions of mouse brain aging samples (MsBrainAgingSpatialDonor 9-0 and 11-2). Ground truth cell-type maps (left) are compared with OmniCell predictions (right).

Supplementary Figure 2. Extended analysis of gene embeddings and module characterization.

(**A**) Workflow for generating cell-level and dataset-level gene embeddings using OmniCell Transformer encoder, GPR-GNN graph propagation through cell-similarity networks, and linear aggregation across cells. (**B**) The radial heatmap demonstrates clear segregation of these markers, while the central Venn diagram highlights both unique and shared genes between excitatory and inhibitory lineages (42 EX-specific, 35 INH-specific, 34 shared). (**C**) UMAP visualizations of cell-level embeddings for *LHFPL3* expression (left) and corresponding cell-type annotations (right). (**D**) Anatomical parcellation of normal human hippocampus showing cornu ammonis subfields (CA1–CA4), dentate gyrus (DG), and granule cell layer (GC). (**E**) Spatial distribution of module expression scores in normal human hippocampus spatial transcriptomics data.

A

MouseBrain

OmniCell   scGPT   scFoundation   GeneFormer

Astro-1	ExN-L6-1	InN-Pvalb-1	Micro-2	Epen	InN-Calb2-2	InN-Vip	Olig-3	Micro-1
Astro-2	ExN-L6-2	InN-Pvalb-2	Micro-3	ExN-L2/3-1	InN-Chat	MSN-D1-1	Peri-1	ExN-L5-3
Endo-1	ExN-L6-3	InN-Pvalb-3	OPC	ExN-L2/3-2	InN-Lamp5	MSN-D1-2	Peri-2	InN-Olf-2
Endo-2	ExN-Olf	InN-Sst-1	Olig-1	ExN-L5-1	InN-Lhx6	MSN-D2	T cell	
Endo-3	InN-Calb2-1	InN-Sst-2	Olig-2	ExN-L5-2	InN-Olf-1	Macro	Vlmc	

B

HumanKidney

OmniCell   scGPT   scFoundation   GeneFormer

B cell	kidney collecting duct cell	monocyte	renal beta-intercalated cell
T cell	kidney distal convoluted tubule epithelial cell	natural killer cell	renal principal cell
columnar/cuboidal epithelial cell	kidney loop of Henle thick ascending limb epithelial cell	pericyte	vasa recta ascending limb cell
epithelial cell of proximal tubule	macrophage	podocyte	vasa recta descending limb cell
glomerular capillary endothelial cell	mesangial cell	renal alpha-intercalated cell	vascular associated smooth muscle cell

C

hPBMC

OmniCell   scGPT   scFoundation   GeneFormer   PCA

UMAP2
UMAP1

Cell Types   Batches

B cell	CD8 T cell	Hematopoietic stem cell	Megakaryocyte	Monocyte_CD14	0
CD4 T cell	Monocyte_FCGR3A	NK cell	Plasmacytoid dendritic cell		1

D

hPBMC

Avg Bio-conservation Score vs Avg Batch-correction Score

OmniCell, Geneformer, scGPT, PCA, scFoundation

E

Expert Preference Clustering (Z > 1.0)

Preference Score (0–1) vs Expert

Cell Group: Dendritic cells, Erythroid, Monocytes, Progenitors, T cells

Supplementary Figure 3. Extended evaluation of OmniCell generalization and expert modularity.

(**A**) UMAP projections of the MouseBrain dataset comparing embeddings from OmniCell, scGPT, scFoundation, and Geneformer. (**B**) UMAP projections of the HumanKidney dataset across the same four methods. (**C**) UMAP embeddings of the hPBMC dataset generated by five methods (OmniCell, scGPT, scFoundation, Geneformer, PCA). Upper row: cells colored by nine cell types . Lower row: same embeddings colored by batch identity. (**D**) Scatterplot of batch integration performance on the hPBMC dataset. Average batch-correction score (x-axis) is plotted against average bio-conservation score (y-axis) for each method. (**E**) Bar chart of expert preference clustering (z-score threshold > 1.0) across ten experts. Bars are colored by cell lineage group: dendritic cells, erythroid, monocytes, progenitors, and T cells.

Supplementary Figure 4. Extended spatial evaluation of OmniCell across MERFISH and Stereo-seq datasets.

(**A**) Spatial coordinate maps of MERFISH slice 10_2 comparing ground truth annotations with clustering results from OmniCell (NMI = 0.8792), scGPT-spatial (NMI = 0.8365), and Nicheformer (NMI = 0.3232). (**B**) UMAP visualization of cell embeddings from MERFISH slice 10_2 (MsBrainAgingSpatialDonor 10-2) colored by cell type annotations, comparing OmniCell, scGPT-spatial, and Nicheformer. (**C**) Spatial domain delineation on a mouse cortex dataset with gold-standard regional annotations. Ground truth shows anatomical regions including Hippocampus, Internal Capsule, cortical Layers 2 – 6, Pia Layer 1, Ventricle, and White matter. Comparison across methods demonstrates that OmniCell (NMI = 0.6623) best recapitulates cortical layer architecture, outperforming SCAN-IT (NMI = 0.6354), SpaGCN (NMI = 0.4319), SEDR (NMI = 0.2970), scGPT-spatial (NMI = 0.2778), and Nicheformer (NMI = 0.2249). (**D**) Quantitative comparison of ARI and NMI across all methods for spatial domain reconstruction. (**E**) Spatial plots showing the activity of enriched pathways in the transition zone, including acute inflammatory response, complement activation, and regulation of blood coagulation, where white streamlines indicate gradient directions.

Supplementary Figure 5. Extended benchmarking OmniCell for single-cell cell-type annotation and spatial transcriptomics deconvolution.

(**A**) Full confusion matrices for OmniCell, scGPT, scFoundation, and Geneformer on the Zheng68k dataset, showing OmniCell's superior accuracy and clearer diagonal dominance across all major and rare cell types. (**B**) Bar chart of cell-type frequency distribution in the Zheng68k dataset. (**C–D**) Bubble plots comparing predicted versus ground truth cell-type proportions on the hPBMC (**C**) and LungCancer (**D**) datasets. Dot size corresponds to cell-type proportion; color intensity reflects prediction accuracy. Each row represents a method; each column represents a cell type. (**E–G**) Spatial deconvolution evaluation on MERFISH mouse brain atlas slices. (**E**) Spatial reconstructions of slice 9_0 comparing ground truth annotations with predictions from OmniCell, scGPT-spatial, Nicheformer, and cell2location. (**F**) Corresponding confusion matrices for slice 9_0. (**G**) Spatial reconstructions and confusion matrices for slice 11_2. Cell types include neuronal, glial, vascular, and immune populations as annotated in the reference atlas.

Reference

1. Liao, J., et al., *Uncovering an Organ's Molecular Architecture at Single-Cell Resolution by Spatially Resolved Transcriptomics.* Trends Biotechnol, 2021. **39**(1): p. 43-58.

2. Lee, J., M. Yoo, and J. Choi, *Recent advances in spatially resolved transcriptomics: challenges and opportunities.* BMB Rep, 2022. **55**(3): p. 113-124.

3. Gulati, G.S., et al., *Profiling cell identity and tissue architecture with single-cell and spatial transcriptomics.* Nat Rev Mol Cell Biol, 2025. **26**(1): p. 11-31.

4. Zheng, G.X., et al., *Massively parallel digital transcriptional profiling of single cells.* Nat Commun, 2017. **8**: p. 14049.

5. Macosko, E.Z., et al., *Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets.* Cell, 2015. **161**(5): p. 1202-1214.

6. Picelli, S., et al., *Full-length RNA-seq from single cells using Smart-seq2.* Nat Protoc, 2014. **9**(1): p. 171-81.

7. Hagemann-Jensen, M., et al., *Single-cell RNA counting at allele and isoform resolution using Smart-seq3.* Nat Biotechnol, 2020. **38**(6): p. 708-714.

8. Hilts, K.E., et al., *Impact of Medicaid expansion on smoking prevalence and quit attempts among those newly eligible, 2011-2019.* Tob Prev Cessat, 2021. **7**: p. 16.

9. Moncada, R., et al., *Integrating microarray-based spatial transcriptomics and single-cell RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas.* Nat Biotechnol, 2020. **38**(3): p. 333-342.

10. Chen, K.H., et al., *RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells.* Science, 2015. **348**(6233): p. aaa6090.

11. Eng, C.L., et al., *Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH.* Nature, 2019. **568**(7751): p. 235-239.

12. Wang, X., et al., *Three-dimensional intact-tissue sequencing of single-cell transcriptional states.* Science, 2018. **361**(6400).

13. Chen, X., et al., *Efficient in situ barcode sequencing using padlock probe-based BaristaSeq.* Nucleic Acids Res, 2018. **46**(4): p. e22.

14. Stahl, P.L., et al., *Visualization and analysis of gene expression in tissue sections by spatial transcriptomics.* Science, 2016. **353**(6294): p. 78-82.

15. Rodriques, S.G., et al., *Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution.* Science, 2019. **363**(6434): p. 1463-1467.

16. Chen, A., et al., *Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays.* Cell, 2022. **185**(10): p. 1777-1792 e21.

17. Cui, H., et al., *scGPT: toward building a foundation model for single-cell multi-omics using generative AI.* Nat Methods, 2024. **21**(8): p. 1470-1480.

18. Hao, M., et al., *Large-scale foundation model on single-cell transcriptomics.* Nat Methods, 2024. **21**(8): p. 1481-1491.

19. Theodoris, C.V., et al., *Transfer learning enables predictions in network biology.* Nature, 2023. **618**(7965): p. 616-624.

20. Szalata, A., et al., *Transformers in single-cell omics: a review and new perspectives.* Nat Methods, 2024. **21**(8): p. 1430-1443.

21. Wang, C., et al., *scGPT-spatial: Continual Pretraining of Single-Cell Foundation Model for Spatial Transcriptomics.* bioRxiv, 2025.

22. Tejada-Lapuerta, A., et al., *Nicheformer: a foundation model for single-cell and spatial omics.* Nat
Methods, 2025.

23. Wen, H., et al., *CellPLM: Pre-training of Cell Language Model Beyond Single Cells.* 2023.

24. Vaswani, A., et al. *Attention Is All You Need.* 2017. arXiv:1706.03762 DOI:
10.48550/arXiv.1706.03762.

25. Zeng, H., et al., *Large-scale cellular-resolution gene profiling in human neocortex reveals*
species-specific molecular signatures. Cell, 2012. **149**(2): p. 483-96.

26. Gerfen, C.R. and D.J. Surmeier, *Modulation of striatal projection systems by dopamine.* Annu Rev
Neurosci, 2011. **34**: p. 441-66.

27. Eng, L.F., et al., *An acidic protein isolated from fibrous astrocytes.* Brain Res, 1971. **28**(2): p. 351-4.

28. Butovsky, O., et al., *Identification of a unique TGF-beta-dependent molecular and functional signature*
in microglia. Nat Neurosci, 2014. **17**(1): p. 131-43.

29. Nave, K.A. and H.B. Werner, *Myelination of the nervous system: mechanisms and functions.* Annu Rev
Cell Dev Biol, 2014. **30**: p. 503-33.

30. Spitzer, S.O., et al., *Oligodendrocyte Progenitor Cells Become Regionally Diverse and Heterogeneous*
with Age. Neuron, 2019. **101**(3): p. 459-471 e5.

31. Zeisel, A., et al., *Molecular Architecture of the Mouse Nervous System.* Cell, 2018. **174**(4): p. 999-1014
e22.

32. Jones, E.V., et al., *Astrocytes control glutamate receptor levels at developing synapses through*
SPARC-beta-integrin interactions. J Neurosci, 2011. **31**(11): p. 4154-65.

33. Chien, E., et al. *Adaptive Universal Generalized PageRank Graph Neural Network.* 2020.
arXiv:2006.07988 DOI: 10.48550/arXiv.2006.07988.

34. Wu, L., et al., *An invasive zone in human liver cancer identified by Stereo-seq promotes*
hepatocyte-tumor cell crosstalk, local immunosuppression and tumor progression. Cell Res, 2023.
33(8): p. 585-603.

35. Si, M. and J. Lang, *The roles of metallothioneins in carcinogenesis.* J Hematol Oncol, 2018. **11**(1): p.
107.

36. Wong, D.R., A.S. Hill, and R. Moccia, *Simple controls exceed best deep learning algorithms and reveal*
foundation model effectiveness for predicting genetic perturbations. Bioinformatics, 2025. **41**(6).

37. Ahlmann-Eltze, C., W. Huber, and S. Anders, *Deep-learning-based gene perturbation effect prediction*
does not yet outperform simple linear baselines. Nat Methods, 2025. **22**(8): p. 1657-1661.

38. Vinas Torne, R., et al., *Systema: a framework for evaluating genetic perturbation response prediction*
beyond systematic variation. Nat Biotechnol, 2025.

39. Zhou, Y., et al., *Human and mouse single-nucleus transcriptomics reveal TREM2-dependent and*
TREM2-independent cellular responses in Alzheimer's disease. Nat Med, 2020. **26**(1): p. 131-142.

40. Wang, P., et al., *Molecular pathways and diagnosis in spatially resolved Alzheimer's hippocampal atlas.*
Neuron, 2025. **113**(13): p. 2123-2140 e9.

41. De Simone, M., et al., *A comprehensive analysis framework for evaluating commercial single-cell RNA*
sequencing technologies. Nucleic Acids Res, 2025. **53**(2).

42. Zhang, Y., et al., *Single-cell analyses of renal cell cancers reveal insights into tumor microenvironment,*
cell of origin, and therapy response. Proc Natl Acad Sci U S A, 2021. **118**(24).

43. Becker, W.R., et al., *Single-cell analyses define a continuum of cell state and composition changes in*
the malignant transformation of polyps to colorectal cancer. Nat Genet, 2022. **54**(7): p. 985-995.

44. Cang, Z., et al., *SCAN-IT: Domain segmentation of spatial transcriptomics images by graph neural*

network. BMVC, 2021. **32**.

45.   Tran, H.T.N., et al., *A benchmark of batch-effect correction methods for single-cell RNA sequencing*
data. Genome Biol, 2020. **21**(1): p. 12.

46.   Siletti, K., et al., *Transcriptomic diversity of cell types across the adult human brain.* Science, 2023.
382(6667): p. eadd7046.

47.   Velmeshev, D., et al., *Single-cell analysis of prenatal and postnatal human cortical development.*
Science, 2023. **382**(6667): p. eadf0834.

48.   Ivanova, E.N., et al., *mRNA COVID-19 vaccine elicits potent adaptive immune response without the*
acute inflammation of SARS-CoV-2 infection. iScience, 2023. **26**(12): p. 108572.

49.   Jorstad, N.L., et al., *Comparative transcriptomics reveals human-specific cortical features.* Science,
2023. **382**(6667): p. eade9516.

50.   Allen, W.E., et al., *Molecular and spatial signatures of mouse brain aging at single-cell resolution.*
Cell, 2023. **186**(1): p. 194-208 e18.

51.   Codeluppi, S., et al., *Spatial organization of the somatosensory cortex revealed by osmFISH.* Nat
Methods, 2018. **15**(11): p. 932-935.

52.   Sarkar, A. and M. Stephens, *Separating measurement and expression models clarifies confusion in*
single-cell RNA sequencing analysis. Nat Genet, 2021. **53**(6): p. 770-777.

53.   Dai, D., et al. *DeepSeekMoE: Towards Ultimate Expert Specialization in Mixture-of-Experts Language*
Models. 2024. arXiv:2401.06066 DOI: 10.48550/arXiv.2401.06066.

54.   Wang, P., et al. *Qwen2-VL: Enhancing Vision-Language Model's Perception of the World at Any*
Resolution. 2024. arXiv:2409.12191 DOI: 10.48550/arXiv.2409.12191.

55.   Burkhardt, D.B., et al., *Quantifying the effect of experimental perturbations at single-cell resolution.*
Nat Biotechnol, 2021. **39**(5): p. 619-629.

56.   Moon, K.R., et al., *Visualizing structure and transitions in high-dimensional biological data.* Nat
Biotechnol, 2019. **37**(12): p. 1482-1492.

57.   Hao, Y., et al., *Integrated analysis of multimodal single-cell data.* Cell, 2021. **184**(13): p. 3573-3587
e29.

58.   Luecken, M.D., et al., *Benchmarking atlas-level data integration in single-cell genomics.* Nat Methods,
2022. **19**(1): p. 41-50.

59.   Xu, H., et al., *Unsupervised spatially embedded deep representation of spatial transcriptomics.*
Genome Med, 2024. **16**(1): p. 12.

60.   Hu, J., et al., *SpaGCN: Integrating gene expression, spatial location and histology to identify spatial*
domains and spatially variable genes by graph convolutional network. Nat Methods, 2021. **18**(11): p.
1342-1351.

61.   Wang, A., et al., *Mapping spatial gradients in spatial transcriptomics data with score matching.*
bioRxiv, 2025.

62.   Biancalani, T., et al., *Deep learning and alignment of spatially resolved single-cell transcriptomes with*
Tangram. Nat Methods, 2021. **18**(11): p. 1352-1362.

933