**che*Ck*OVER: An open framework and AI-ready global crayfish database**

**for next-generation biodiversity knowledge**

Running title: AI-interoperable biodiversity knowledge

Lucian PÂRVULESCU[1,2*], David LIVADARIU[1,3*], Victor I. BÂCU[4], Constantin I.

NANDRA[4], Teodor T. ȘTEFĂNUȚ[4] and World of Crayfish® Contributors[5]

[1] Crayfish Research Centre, Institute for Advanced Environmental Research, West University

of Timisoara, Oituz 4, 300086 Timisoara, Romania

[2] Department of Biology, Faculty of Chemistry, Biology, Geography, West University of

Timisoara, Pestalozzi 16A, 300115 Timisoara, Romania

[3] Department of Informatics, Faculty of Mathematics and Computer Science, West University

of Timisoara, Vasile Pârvan 4, 300223 Timisoara, Romania

[4] Artificial Intelligence Research Institute, Computer Science Department, Technical

University of Cluj-Napoca, Memorandumului 28, 400114 Cluj-Napoca, Romania

[5] World of Crayfish® Contributors (authors information provided in Table S1)

*Correspondence: lucian.parvulescu@e-uvt.ro, david.livadariu02@e-uvt.ro

## 20 Abstract

**21 Background**

Species occurrence records represent the backbone of biodiversity science, yet their utility is

often limited to spatial analyses, distribution maps, or presence-absence models. Current

biodiversity infrastructures rarely provide computational formats directly usable by modern

artificial intelligence (AI) systems, such as large language models (LLMs), which

increasingly mediate scientific communication and knowledge synthesis. Open frameworks

that convert biodiversity occurrences into structured, machine-accessible, provenance-rich

knowledge are therefore essential—particularly those enabling rapid integration of new

records, near real-time generation of spatial metrics, and production of both human-

interpretable reports and AI-consumable outputs. Such capabilities substantially reduce

latency between data acquisition and decision support, while ensuring biodiversity knowledge

remains traceable and verifiable in AI-mediated workflows.

**Results**

We introduce cheCkOVER, an open framework that converts raw species occurrence datasets

into standardized, API-ready, multi-layered outputs: biogeographic descriptors, dynamic

distribution maps, summary metrics, and structured JSON geo-narratives following a

canonical template. The framework stratifies processing by population origin (indigenous vs.

non-indigenous), enabling IUCN-aligned conservation metrics while simultaneously tracking

invasion dynamics. Each output embeds standardized citation metadata ensuring full

provenance traceability. We applied the pipeline to 111,729 validated crayfish (Astacidea)

occurrence records from 465 species, generating comprehensive species packages including

indigenous-range classifications (171 endemic, 287 regional, 5 cosmopolitan taxa) and non-

indigenous range tracking for 30 invasive species. This proof-of-concept demonstrates how

the framework transforms minimal datapoints—validated species occurrences—into

interoperable knowledge consumable by both humans and computational systems. The JSON

outputs are optimized for retrieval-augmented generation, enabling AI systems to dynamically

access and cite biodiversity knowledge with explicit source attribution.

**Conclusions**

che*CkOVER* is taxon-agnostic and establishes a reproducible pathway from biodiversity

occurrences to narrative-ready, AI-interoperable knowledge with immediate public utility via

the World of Crayfish® platform (https://world.crayfish.ro/), where each species page

integrates structured outputs. The open-source framework (GPL-3) combines a generalizable

processing pipeline with taxon-specific knowledge products, enabling flexible reuse across

conservation research, policy reporting, and AI-driven applications. This minimalist-to-

complex design extends the reach of biodiversity data beyond traditional analyses, positioning

occurrence repositories as active knowledge engines for next-generation biodiversity

informatics.

**Significance statement**

Biodiversity infrastructures remain underused by modern AI systems despite their central

role in science and society. che*CkOVER* embodies a minimalist-to-complex paradigm: from

the validated geographic occurrence of a species—a datapoint often perceived as trivial—it

derives structured, multi-layered outputs linking distribution, conservation status, and

standardized geographic indicators. These outputs are natively optimized for retrieval-

augmented generation and other machine-consumable workflows, enabling AI systems to

dynamically access and cite biodiversity knowledge with maintained provenance beyond their

pre-training corpora. Using a global crayfish dataset as proof-of-concept, we demonstrate how

raw occurrence records can scale into rich, interoperable biogeographic knowledge products

with immediate value for both human experts and computational systems. This positions

biodiversity databases as critical knowledge engines for next-generation science, policy, and

societal decision-making, providing standardized outputs directly incorporable into

conservation evaluation workflows where transparent, reproducible, and provenance-rich

occurrence-based metrics are essential.

**Keywords**: AI-ready knowledge; biodiversity informatics; geo-narrative; JSON database;

spatial analytics

## Introduction

Human societies have always relied on knowledge to guide decisions and shape progress [1–3]. From antiquity to the digital era, science has served as a trusted source of verified information [4], while new technologies have multiplied the ways knowledge flows into public and policy domains [5,6]. Today, artificial intelligence (AI), particularly large language models (LLMs), is emerging as a dominant mediator of communication, offering rapid access to information across disciplines and audiences [7–9].

Species distribution records are highly valuable for numerous purposes, from scientific research [10,11] and understanding species communities [12,13], to conservation planning and management [14], and ultimately to certain economic considerations [15]. In particular, the quality of stored and delivered biodiversity knowledge is essential [16]. Nature underpins human well-being [17,18], and its understanding depends on reliable data about both species and the habitats they occupy [19,20]. Biodiversity knowledge must not only be abundant, but also accurate, transparent and easy to integrate into decision-making [21]. In this domain, misinformation or poorly contextualized data can directly affect conservation outcomes, policy priorities, and public trust [22,23].

However the speed and scale of LLMs analyses come at a cost: LLMs generate fluent but often vague, erroneous, or unverifiable outputs [24,25]. Their training depends on heterogeneous and inconsistently curated data, leaving a fundamental gap between scientific knowledge production and the AI systems that now mediate it [26]. Simultaneously, scientific data are expanding at unprecedented rates [27]. Advances in modeling, genomics, and ecological monitoring—together with citizen science—have produced millions of biodiversity records and traits [28–30]. Dedicated infrastructures and portals have proliferated, enabling access to vast repositories of occurrence data, images, and observations [31]. Yet the value of these resources remains underexploited [32,33]. These resources typically generate outputs that AI systems cannot directly consume, leaving only fragments of biodiversity knowledge contextualized in their reporting [34,35].

While humans digest information through narratives and visualizations [36,37], AI requires structured inputs: machine-readable formats (e.g., JavaScript Object Notation – JSON) annotated with explicit metadata and provenance [38–40]. To enable the full integration of biodiversity science in the AI era [41,42], infrastructures must evolve from human-facing repositories into dual-purpose systems [43–45]. Here we present che*CkOVER*, the first generalizable framework designed to fill this gap. At its core, che*CkOVER*

deliberately returns to the roots of faunistics—validated raw occurrence records without prior

interpretation—and transforms them into standardized, multi-layered outputs that are both

human-readable and LLM-digestible. Using global crayfish (Astacidea) occurrence data [10]

as a proof of concept, we illustrate how the framework scales minimal datapoints into

interoperable, machine-readable knowledge packages that distinguish indigenous from non-

indigenous populations while maintaining unified outputs for species assessment. These

outputs are structured to support retrieval-augmented and other machine-consumable

workflows, positioning biodiversity infrastructures to interface effectively with emerging AI

technologies. In this way, che*CkOVER* reframes occurrence repositories as future-ready

knowledge engines, enabling broader reuse, clearer attribution, and improved integration

across scientific, conservation, and societal domains.

## Methods

### Framework description

che*CkOVER* is built as a modular architecture that transforms validated biodiversity

occurrence records into standardized, multi-layered knowledge packages. Each package is

generated at the species level and integrates visual, quantitative, and narrative components,

ensuring outputs are both human-readable and machine-interoperable. The name cheCkOVER

reflects the framework's systematic checking of occurrences over various boundaries,

signaling a step toward consolidated biodiversity knowledge infrastructure.

The workflow is organized around five conceptual components (Figure 1), internally

implemented through seven sequential processing phases that ensure methodological rigor

while accommodating complex distributional patterns. This hybrid design balances

conceptual clarity with implementation flexibility, enabling the framework to handle species

occurring exclusively as native populations, exclusively as introduced populations, or in both

contexts simultaneously.

*Phase 1: Data ingestion, standardization, and pre-enrichment.* Raw occurrence

records are loaded, cleaned, and standardized (species names, coordinates, status, temporal

data). Vernacular names are harvested from authoritative sources and embedded with ISO

639-3 language tags. Initial geospatial enrichment assigns continental context and

administrative boundaries (countries, subnational units via GADM v4), establishing the

foundational geographic framework for subsequent analyses.

*Phase 2: Population-origin stratification.* To maintain compatibility with IUCN Red

List methodology while simultaneously supporting invasion biology workflows, the

framework stratifies records by population origin (indigenous vs. non-indigenous). Species

are classified into three scenarios: (1) indigenous-only, (2) non-indigenous-only, or (3)

combined. This stratification guides all downstream metric calculations and ensures that

conservation-relevant indicators (EOO, AOO, fragmentation signals) derive strictly from

native-range data, while invasion dynamics are tracked separately.

*Phase 3: Indigenous-range analysis.* For native populations (including type locality),

the framework computes distributional categories (endemic, regional, cosmopolitan) based on

range extent and geographic scope. Occurrence records are overlaid with standardized

geospatial layers (ecoregions, protected areas, hydrographic basins) to generate species-level

summaries. For endemic and regional taxa, a conservative fragmentation signal is computed

via adaptive spatial clustering of native localities, providing early indications of potential

range disjunction without constituting a formal connectivity assessment.

*Phase 4: Non-indigenous-range analysis.* Alien, introduced, or cryptogenic

populations are processed through a parallel branch using the same geospatial layers but

without fragmentation analysis (as range structure reflects human-mediated dispersal rather

than natural processes). Non-indigenous metrics (EOO, AOO, basin occupancy) are computed

for informative purposes and support biosecurity prioritization, but do not contribute to

conservation status assessments.

*Phase 5: Scenario integration.* For species occurring in both native and introduced

contexts (Scenario 3), the framework merges data from both branches into unified outputs that

clearly distinguish indigenous from non-indigenous components. This ensures that species

pages present complete distributional footprints while preserving the methodological

separation required for IUCN-aligned assessments.

*Phase 6: Narrative and cartographic synthesis.* Species-level outputs are generated

following a standardized canonical geo-narrative template (Supplementary File S1), covering

taxonomic identity, indigenous and non-indigenous ranges, temporal patterns, data quality,

and provenance. Narratives are delivered in multiple formats: human-readable text

(Markdown, plain text), structured JSON optimized for LLM consumption, and machine-

readable metadata files. Geospatial products (KML/KMZ, GeoJSON) visualize indigenous

and non-indigenous ranges using distinct color schemes, with type localities highlighted when

available.

*Phase 7: Species package assembly.* Final outputs are bundled into self-contained,

publication-ready packages, each including maps, narratives, quantitative tables, citation files

(BibTeX, JSON, CITATION.cff), and a structured manifest. Privacy protection is ensured by

aggregating spatial information to administrative or ecological units, preventing exposure of

precise coordinates for sensitive species.

All outputs follow FAIR principles, include output-level versioning (package revision,

timestamp, and manifest), and are exposed through documented, API-ready endpoints to

support integration with external biodiversity infrastructures and AI-based systems [46]. The

framework implements incremental update capability: when new occurrence records are

added, only modified species are reprocessed, with results compared against previous versions

to track distributional changes over time. This version-aware design ensures computational

efficiency for large datasets while maintaining full reproducibility and traceability.

The che*CkOVER* pipeline was developed within a modern integrated development

environment combining Claude (Anthropic) AI-assisted code generation [47] with expert-

driven manual refinement. AI-generated code drafts were iteratively reviewed, validated,

optimized, and extended by the authors using real-world occurrence data subsets. This hybrid

development workflow ensured reproducibility, code transparency, and compliance with

biodiversity informatics best practices [48,49].

figure 1 here (one column)

**Figure 1. Conceptual workflow of the che*CkOVER* framework.** The system is organized around five

conceptual components: (1) integrated biodiversity input databases; (2) data ingestion, cleaning, and enrichment;

(3) analytical processing of distributional and spatial metrics; (4) automated report and narrative generation; and

(5) production of modular human- and machine-consumable outputs. Internally, these components are

implemented through seven sequential processing phases that stratify species by population origin (indigenous

vs. non-indigenous), enabling IUCN-aligned conservation metrics while simultaneously tracking invasion

dynamics. The workflow supports both immediate decision-ready reporting and continued context-aware

analysis through AI and LLM-based systems, with all outputs embedding provenance metadata for transparent

source attribution.

## Data inputs and preprocessing

che*CkOVER* relies on streamlined occurrence datasets that are easy to maintain and extend

over time. The framework was developed and validated using the World of Crayfish® (WoC®)

database [10], but accepts occurrence data from any taxonomic source following the same

core schema. Only core fields required for geo-narrative and quantitative metric generation

are used: the valid Latin binomial of the taxon (with synonymy resolved via external APIs),

population origin (indigenous vs. alien/introduced), year of observation, geographic

coordinates in standardized WGS84 decimal degrees, and a location-accuracy tag.

Bibliographic references for each record, including DOI/URL where available, are retained

for traceability and citation.

All occurrence records are processed through a series of validation and standardization

operations using the tidyverse suite. Species names are cleaned, coordinates coerced to

WGS84, population origin normalized (native/alien/introduced/cryptogenic), and accuracy

flagged (high/low). Duplicate records (species–year–coordinates–source) are removed,

ensuring consistency and reproducibility across datasets. Following preprocessing, records are

stratified by population origin, creating two parallel processing branches (indigenous and non-

indigenous) while maintaining a unified master dataset for scenario detection. Species are

then classified into three scenarios based on their population-origin profiles: Scenario 1

(indigenous-only), Scenario 2 (non-indigenous-only), or Scenario 3 (both population types

present). This classification guides downstream spatial resolution choices (e.g., finer

HydroBASINS levels for endemic taxa, coarser resolutions for cosmopolitan taxa) and

ensures internal consistency across quantitative metrics and visual outputs.

To support multilingual, real-world queries, vernacular (common) names are not

required as inputs; instead, they are harvested downstream in a dedicated step from

authoritative and community-curated services (e.g., ITIS; https://www.itis.gov/), or

supplemented with expert-reviewed multilingual names compiled in a static lookup table

(Table S2). This step is implemented using the *ritis* package for querying authoritative

vernacular-name services. All vernacular names are normalized with ISO 639-3 language

tags. The resulting language-aware name lists are embedded in species outputs to improve

discoverability without complicating the primary input schema.

**Geospatial enrichment**

To generate narrative-ready outputs, che*CkOVER* links validated occurrence records to a suite

of standardized geospatial layers through a two-stage enrichment process. Stage 1 (pre-

stratification) assigns continental context and administrative boundaries to all records prior to

population-origin splitting, establishing a unified geographic framework. Stage 2 (post-

stratification) performs branch-specific spatial overlays with ecoregions, protected areas, and

hydrographic basins, with resolution and layer selection adapted to each population type's

biogeographic context.

These layers provide named, interpretable geographic units—countries, administrative
regions, river basins, ecoregions, and protected areas—that allow species distributions to be
described using stable, widely recognized terminology. Ensuring correct names for these units
is essential, and the framework therefore integrates both programmatic attribution (via R
packages) and reproducible lookup tables for cases where names are absent from the original
polygons.
*Stage 1: Universal geographic context.* Country-level boundaries are sourced from
Natural Earth (via the *rnaturalearth* package), while sub-national administrative units are
retrieved through the geodata package, which provides standardized access to GADM v4.
Continental assignments are derived from Natural Earth medium-scale vector data. These
layers are applied uniformly to all occurrence records regardless of population origin,
ensuring that baseline geographic descriptors remain consistent across indigenous and non-
indigenous subsets.
*Stage 2: Branch-specific biogeographic enrichment.* Following population-origin
stratification, indigenous and non-indigenous records are processed through separate but
parallel spatial overlay workflows. This separation ensures that IUCN-aligned conservation
metrics derive exclusively from native-range data while enabling comprehensive tracking of
introduced populations for invasion biology applications.
Freshwater contexts are captured through HydroBASINS hierarchical watershed
boundaries (levels L06–L10) and HydroATLAS hydro-environmental descriptors, while
WWF Terrestrial Ecoregions of the World (TEOW) and Freshwater Ecoregions of the World
(FEOW) are assigned using the *ecoregions* and *feowR* packages. Because HydroBASINS
polygons do not include hydronym attributes, basin names were derived through a dedicated
offline geoprocessing step. HydroBASINS units (L06, L08, L10) were spatially intersected
with river-line datasets from the Global Runoff Data Centre (for L06–L08) and high-
resolution OpenStreetMap extracts (for L10). Each basin polygon was assigned the name of
the longest or most representative river crossing its area. This workflow yielded a static
hydronym lookup table linking HydroBASINS IDs to river names (Table S3), now integrated
into the che*Ck*OVER pipeline. When multiple HydroBASINS IDs share the same river name
(e.g., multiple segments of the Danube), records are aggregated by basin name and summed
for reporting, ensuring that outputs present deduplicated, human-readable geographic units
rather than technical polygon identifiers. In contrast, TEOW polygons include standardized
ecoregion names directly via the ecoregions package; therefore, no additional processing was
required. FEOW polygons, however, lack embedded name attributes in some distributions,

and a reproducible lookup table linking FEOW IDs to ecoregion names was compiled (Table

S4). Protected areas were intersected using the World Database on Protected Areas (WDPA,

Protected Planet), retrieved programmatically via the *wdpar* package to ensure license-

compliant access.

For non-indigenous populations fragmentation analysis is omitted, as introduced-range

structure reflects human-mediated dispersal patterns rather than natural biogeographic

processes. This design ensures that biosecurity-relevant spatial summaries (basin occupancy,

protected-area overlap) are generated without conflating anthropogenic range expansion with

evolutionarily meaningful fragmentation signals.

All geospatial operations—cropping, overlays, joins, reprojections, and geometry

handling—are executed using the *sf* package, which provides high-performance spatial data

processing and consistent handling of modern GIS geometries.

**Range metrics and distributional categorization**

For range metrics, the framework computes extent of occurrence (EOO) for native ranges as

minimum convex polygons encompassing native validated records, following IUCN

guidelines [50]. To avoid artificial inflation in species with disjunct ranges across continents

or islands, EOO is calculated as the sum of convex hulls per contiguous landmass (or

freshwater system), preventing polygons spanning oceans. Area of occupancy (AOO) is

calculated as the number of occupied grid cells (typically 2×2 km) multiplied by cell area.

Occurrences flagged as extinct are removed prior to spatial overlays and range calculations,

ensuring that EOO and AOO represent only the current distribution. Extinct localities remain

available for reporting but do not influence geospatial metrics. All outputs distinguish

between native and alien ranges to maintain ecological relevance and ensure compatibility

with IUCN conservation standards. To maintain full compatibility with IUCN Red List

methodology, alien/introduced occurrences are never included in metric calculations, but are

retained for spatial reporting and visualization. Accordingly, all geospatial display products

(GeoJSON, KML/KMZ) show both native and alien ranges when present, while the

corresponding EOO/AOO values included in reports and JSON summaries derive strictly

from the native subset of records.

To contextualize the breadth of native distributions prior to spatial enrichment,

che*Ck*O*VER* classifies taxa into three distributional categories using native-range occurrence

statistics only. (i) Endemic taxa are defined as species with a native extent of occurrence

(EOO) ≤5,000 km², restricted to ≤2 countries within a single continent. These thresholds are

consistent with restricted-range criteria commonly applied to freshwater invertebrates [51]

and reflect the naturally fragmented distributions of Astacidea across isolated drainage

systems [52,53]. (ii) Regional taxa are species whose native ranges exceed endemic

thresholds but remain confined to a single continent, regardless of any secondary

introductions outside their native area. (iii) Cosmopolitan taxa are defined as species whose

native ranges span more than one continent, indicating broad historical biogeographic

distributions rather than anthropogenic spread.

Records outside the native range (alien, introduced, or cryptogenic occurrences) are

explicitly retained and analyzed separately for descriptive statistics, protected-area

intersections, basin-level summaries, and cartographic outputs, but they do not influence

native-range classification. This separation ensures that biogeographic categorization reflects

evolutionary and historical range structure, while still capturing the full contemporary

distributional footprint of each species. This native-range classification guides downstream

spatial resolution choices (e.g., finer HydroBASINS levels for endemic taxa and coarser

resolutions for cosmopolitan taxa), ensuring internal consistency across quantitative metrics

and visual outputs.

**Fragmentation analysis (indigenous populations only)**

For endemic and regional taxa, che*CkOVER* additionally computes a simple distributional

fragmentation signal to summarize the spatial cohesion of known native localities. The

analysis is performed on straight-line (Euclidean) distances, calculated after transforming

occurrence coordinates to an equal-area projection (EPSG:6933) to ensure metric accuracy. A

full pairwise distance matrix is generated, and the mean inter-point distance is used as an

adaptive species-specific threshold for hierarchical clustering (complete linkage). This

approach allows fragmentation detection to scale naturally with each species' spatial

configuration. Clusters are derived using *hclust()* and *cutree()* at the adaptive height. Species

with fewer than five valid locality records, as well as cosmopolitan taxa whose ranges are

inherently multi-continental, are excluded from clustering. Non-indigenous populations are

excluded from fragmentation analysis by design, as introduced-range structure reflects

human-mediated dispersal rather than natural connectivity patterns.

**Computational optimization**

To minimize computational load during large-scale intersection operations, che*CkOVER*

implements an optimized, multi-stage spatial overlay procedure. For each species, the

framework first computes an early EOO polygon, buffered to account for spatial uncertainty.

All reference layers (countries, basins, protected areas) are dynamically cropped to this

bounding region prior to point-level intersections, reducing redundant geometry checks.

Spatial joins are executed in batches using indexed geometries (*sf* + *S2* engine), followed by

tabular aggregation per species and per geographic unit. When applicable, HydroBASINS

levels are dynamically selected (L10 for endemic, L8 for regional, L6 for cosmopolitan taxa),

and protected areas are pre-filtered to remove marine or micro-polygons. This adaptive

workflow reduces memory requirements and computation time while maintaining full spatial

accuracy.

**Narrative generation and outputs**

The core function of che*CkOVER* is to transform enriched species occurrence records into

structured geo-narratives that are both human-readable and machine-interoperable. Species-

level outputs follow a standardized canonical geo-narrative template (Supplementary File S1),

implemented using the glue package, ensuring consistency across taxa and enabling fully

reproducible reporting for both human users and automated systems.

*Canonical template structure.* The canonical template organizes biodiversity

knowledge into five standardized sections: (1) Taxonomic Identity, including scientific name,

higher taxonomy (Order > Superfamily > Family), vernacular names with full language

names (not ISO codes in output text), and type locality when available; (2) Indigenous Range

Overview, covering distribution category (endemic/regional/cosmopolitan), EOO/AOO

metrics, geographic distribution (countries, subnational units, hydrographic basins—all sorted

by record count descending), protected area coverage, biogeographic context (TEOW/FEOW

ecoregions), temporal context, auto-generated contextual statements, range dynamics (if

extinctions documented), and fragmentation assessment (endemic/regional only); (3) Non-

Indigenous Range Overview (informative section for species with introduced populations),

reporting EOO/AOO, countries, basins, and temporal patterns separately from native metrics;

(4) Data Quality, Traceability, and Provenance, documenting total records, indigenous/non-

indigenous breakdown, high-accuracy records (coordinate uncertainty <1 km), data quality

score (0–100 composite), bibliographic coverage (total references vs. DOI-linked references),

raw data provenance (WoC® citation), processing framework provenance (che*CkOVER*

version, processing date, data snapshot date), and interpretation note (disclaimer about non-

IUCN status); and (5) Formal Narrative Summary, a 300–500 word human-readable synthesis

structured in five paragraphs covering taxonomic identity, indigenous range

geography/biogeography, conservation context (protected areas, fragmentation, extinctions),

non-indigenous populations (if applicable), and data quality/provenance.

Validated data are organized into predefined narrative blocks following this template

structure. While the framework relies on a fixed narrative structure to guarantee comparability

and reproducibility, it remains extensible, allowing the addition of new sections or user-

defined analytical queries as the system evolves.

For Scenario 1 species (indigenous-only), outputs derive entirely from the indigenous

processing branch. For Scenario 2 species (non-indigenous-only), Section 2 (Indigenous

Range) is omitted, and Section 3 (Non-Indigenous Range) becomes the primary distributional

summary. For Scenario 3 species (both population types), the framework merges data from

both branches: Section 2 presents native-range metrics, Section 3 presents introduced-range

metrics, and both are clearly demarcated to prevent conflation of native vs. alien distributions

in downstream analyses.

Predefined query blocks are embedded within each narrative to support automated

responses to frequently asked questions, while the structured JSON outputs can be directly

consumed by large language models (LLMs) to address free-form prompts. When available,

type localities are highlighted as key descriptive attributes, explicitly linked to their

bibliographic sources, and incorporated into the indigenous range narrative. Vernacular names

harvested during earlier processing steps are automatically embedded in the JSON narratives

using ISO 639-3 language tags internally, but are rendered with full language names (e.g.,

"English: noble crayfish" not "eng: noble crayfish") in human-readable outputs, enabling

accurate querying with local or non-scientific terminology without compromising taxonomic

precision.

Each narrative includes a concise distributional summary, reported both in JSON and

tabular formats, presenting key metrics such as EOO, AOO, total numbers of validated

occurrences, and counts of records annotated as extinct. For endemic and regional taxa,

che*Ck*OVER additionally reports a conservative distributional fragmentation signal derived

exclusively from indigenous occurrence records. This signal contextualizes the spatial

cohesion of known localities by identifying geographically isolated clusters using adaptive,

species-specific clustering thresholds. Fragmentation outputs include (i) the number of

detected spatial clusters, (ii) their relative sizes expressed as occurrence counts and

percentages, and (iii) a categorical indicator of whether fragmentation was detected ("not

detected" for a single cohesive cluster; "detected" when multiple clusters are present). These

signals are intended as supportive descriptors aligned with IUCN criteria B and D [50],

without constituting a formal Red List assessment.

Final outputs are delivered in multiple interoperable formats: (i) canonical narratives

rendered via *rmarkdown* into Markdown files containing all five template sections; (ii) formal

narrative summaries (Section 5 only) exported as both plain text (TXT) and structured JSON

for streamlined LLM consumption; (iii) structured JSON files generated using *jsonlite*,

optimized for programmatic access and LLM-based workflows, with embedded metadata

blocks for each section; and (iv) geospatial artefacts (KML/KMZ and GeoJSON) supporting

visualization and spatial integration. Each JSON narrative embeds a persistent pointer to a de-

duplicated, species-level bibliography, ensuring full provenance, traceability, and

unambiguous citation even when accessed independently. A permanent "How to cite this

resource" entry is maintained in the che*CkOVER* documentation and will be updated

following publication with the definitive citation and DOI.

**Implementation**

The che*CkOVER* initial scripts were drafted and subsequently validated, optimized, and

extended by the authors, ensuring reproducibility and compliance. che*CkOVER* was

implemented in R v.4.3.3 [54] as a modular package, with all functionality exposed through

callable functions (Table 1). The workflow integrates sequential modules for ingestion,

enrichment, reporting, and packaging, yet each component can also be executed

independently. For transparency and reproducibility, all raw scripts are distributed within the

package (*inst*/*scripts_raw*/), enabling stepwise execution when required. The pipeline is

orchestrated by the *run_pipeline()* function, which reads a central configuration file (YAML),

sets paths and column mappings, filters taxa, and executes modules sequentially or selectively

with full logging and error handling.

**Table 1.** Core modules of the che*CkOVER* R package and their primary functions. A full inventory of supporting

R packages required by the pipeline, together with their roles and reference URLs, is provided in Table S5.

| Module | Core function |
| --- | --- |
| *ingest_clean()* | Ingests raw occurrence records (XLSX/TSV/DB), validates the input schema through configurable column mapping (default WoC®-compatible), and standardizes core fields. Species names are cleaned, coordinates coerced to WGS84, population origin normalized (native/type locality/alien/introduced/cryptogenic), and accuracy flagged. Duplicate records (species–year–coordinates–source) are removed; taxonomy may optionally be resolved via external APIs. Exports standardized tabular (CSV/TSV/RDS) and spatial (*sf*) objects. |
| *vernacular_names()* | Retrieves vernacular (common) names from prioritized sources (e.g., |

| Module | Core function |
|---|---|
| | ITIS), normalizes spelling, removes duplicates, applies ISO 639-3 tags internally, and supports manual curation. Outputs consolidated name tables (TSV) and injects them into JSON narratives with full language names (e.g., "English: noble crayfish") to enable queries using non-scientific names. |
| *split_by_population()* | Stratifies occurrence records by population origin (indigenous vs. non-indigenous), creating two parallel datasets while maintaining a unified master dataset. Enables IUCN-aligned metrics from native records while simultaneously tracking invasion dynamics. Exports separate TSV files for each population type. |
| *detect_species_scenarios()* | Classifies species into three scenarios based on population-origin profiles: Scenario 1 (indigenous-only), Scenario 2 (non-indigenous-only), or Scenario 3 (both population types present). This classification guides downstream processing branches and output merging strategies. Exports scenario classification table (TSV/JSON). |
| *enrich_with_continents()* | Assigns continental context to all occurrence records using Natural Earth medium-scale vector data. Applied uniformly before population-origin split to establish baseline geographic framework. |
| *enrich_with_gadm()* | Overlays occurrence records with country and subnational administrative boundaries (GADM v4.1) using the *geodata* package. Applied before population split to ensure consistent political geography across both branches. |
| *calculate_indigenous_metrics()* | Computes distributional categories (endemic/regional/cosmopolitan), EOO, AOO, and preliminary range statistics exclusively from native occurrence records. Implements IUCN-aligned thresholds (endemic: EOO $\leq$5,000 km², $\leq$2 countries, single continent). Guides downstream spatial resolution choices (HydroBASINS level selection). |
| *analyze_fragmentation()* | Detects distributional fragmentation patterns in endemic and regional taxa via adaptive Euclidean-distance clustering. Generates pairwise distance matrix, uses mean distance as species-specific threshold, and applies hierarchical clustering to identify spatially disjunct groups. Outputs cluster count, sizes (counts and percentages), and categorical fragmentation flag. Applied only to indigenous populations; cosmopolitan and non-indigenous taxa excluded by design. |
| *enrich_indigenous_spatial()* | Overlays native occurrence records with biogeographic layers: hydrographic basins (HydroBASINS L06/L08/L10, resolution adaptive to distributional category), ecoregions (FEOW/TEOW), and protected areas (WDPA). Implements optimized workflow: early EOO polygon crops reference layers, indexed geometry operations (*sf* + *S2* engine) batch-process intersections. Basin names assigned via offline hydronym lookup (Table S3); FEOW names via lookup table (Table S4). Aggregates records by basin name (deduplicating multiple polygon IDs for same river). Outputs enriched per-record spatial attributes and summary statistics. |
| *generate_indigenous_reports()* | Compiles quantitative species reports for native populations: record totals, temporal coverage, geographic tallies (countries, subnational units, basins, protected areas—all sorted by record count descending), density metrics, range statistics (EOO, AOO), fragmentation signals, and extinction summaries. Exports tidy tables (TSV), structured JSON summaries, and markdown reports. |
| *calculate_non_indigenous_metrics()* | Computes distributional categories (local/widespread), EOO, AOO, and range statistics exclusively from alien/introduced occurrence records. Metrics are informative (not IUCN-aligned) and support biosecurity prioritization. No fragmentation analysis applied. |
| *enrich_non_indigenous_spatial()* | Overlays alien/introduced records with the same geospatial layers as indigenous branch (basins, ecoregions, protected areas) but omits |

| Module | Core function |
|---|---|
| | fragmentation analysis. Ensures biosecurity-relevant spatial summaries without conflating human-mediated dispersal with natural biogeographic patterns. |
| *generate_non_indigenous_reports()* | Compiles quantitative reports for introduced populations: record totals, temporal coverage, geographic tallies (countries, basins, protected areas), range statistics. Statistics reported separately from native metrics to maintain IUCN compatibility. Exports TSV, JSON, and markdown. |
| *merge_scenario3_reports()* | For Scenario 3 species (both population types), merges data from indigenous and non-indigenous branches into unified outputs that clearly distinguish native from alien components. Ensures species pages present complete distributional footprints while preserving methodological separation required for IUCN assessments. |
| *generate_canonical_narratives()* | Generates structured geo-narratives following the canonical template (Supplementary File S1). Narratives cover five sections: (1) Taxonomic Identity (scientific name, higher taxonomy, vernacular names, type locality), (2) Indigenous Range Overview (distribution category, EOO/AOO, countries, subnational units, basins, protected areas, ecoregions, temporal context, fragmentation), (3) Non-Indigenous Range (if applicable), (4) Data Quality and Provenance (record counts, accuracy metrics, bibliographic coverage, processing metadata), (5) Formal Narrative Summary (300–500 word human-readable synthesis). Outputs canonical markdown (all sections), formal narrative TXT (Section 5), and formal narrative JSON (Section 5 structured). |
| *generate_maps()* | Produces standardized geospatial layers (KML/KMZ, GeoJSON) using the sf framework. Generates three map types per species: EOO convex hulls (yellow polygons), AOO grid cells (2×2 km, yellow), and HydroBASINS occupancy with population-type color coding (native: orange, introduced: purple, mixed: red-orange for Scenario 3). Type localities marked with red 2×2 km squares when available. Map resolution adapts to distributional category (L10 endemic, L8 regional, L6 cosmopolitan). Privacy-protected via spatial aggregation. |
| *generate_all_citations()* | Aggregates and de-duplicates bibliographic references from occurrence records, preserves DOIs/URLs, and generates multiple citation formats: APA-formatted lists, BibTeX files, structured JSON, and CITATION.cff metadata files. Embeds provenance links in all JSON outputs to enable source attribution in LLM-mediated workflows. |
| *export_species_packages()* | Bundles all species-level outputs into distributable packages: geospatial layers (KML/KMZ, GeoJSON), canonical narratives (Markdown, TXT, JSON), quantitative tables (CSV, TSV), citations (BibTeX, JSON, CFF), and metadata files. Generates manifests, checksums, and versioned outputs to support reproducibility and longitudinal tracking. Each package includes README documentation. |
| *run_pipeline()* | Orchestrates sequential or selective module execution across all seven processing phases. Supports YAML config files, CLI parameters, logging, resume mode (detects existing outputs, skips completed modules), and reproducibility reporting. Records version history and reports differences between current and earlier processed outputs. Implements optional parallel execution for indigenous/non-indigenous branches (Linux/macOS via multicore, Windows via multisession). |

**Deployment architecture and API accessibility**

che*CkOVER* outputs clear, well-structured, machine-consumable metadata that allow

seamless integration in complex processing workflows and support efficient integration with

existing LLM services through RESTful API endpoints (e.g., /api/llm/{species_name}). The

framework supports both local execution and server deployment configurations, ensuring

applicability across institutional scales from individual researchers to large biodiversity

platforms.

We comply with all the important aspects for providing a solution that is both API-

ready and LLM-ready: (1) machine consumable data and metadata, through standardized

formats (GeoJSON, KML/KMZ, MD, TSV etc.); (2) rich error semantics that improve API

discoverability and help AI agents and API integrators to understand data structure and

requirements, behavior and recover from errors; (3) speed and reliability ensured by a

containerization-layered technical stack, which provides seamless horizontal scalability at

each level: repository, processing, data conversions; (4) consistent REST end-points design

which, together with the standardized data and querying formats provides predictability; (5)

complete and well structured documentation available through OpenAPI specification.

Data storage combines species-level file hierarchies with a MariaDB relational

database that indexes taxonomic, geographic, and metadata attributes for efficient querying

and retrieval. For production environments, species packages can be exposed via documented

API endpoints that return structured JSON responses optimized for retrieval-augmented

generation workflows. The system is fully containerized (Docker/Singularity), supporting

horizontal scalability at each processing layer and ensuring portability across computing

environments.

**Reproducibility and privacy protection**

All code developed for che*CkOVER* is released as open source and publicly available in a

version-controlled *GitHub* repository. To ensure portability across systems, che*CkOVER* is

fully containerized and can be deployed both locally and within Docker or Singularity

environments. Stable releases are versioned and archived, ensuring long-term reproducibility.

This modular and container-ready design facilitates straightforward maintenance,

interoperability with other biodiversity infrastructures, and adaptability across taxonomic

groups.

Sensitive species records require protection against potential misuse, such as illegal

harvesting or habitat disturbance. che*CkOVER* ensures that raw geographic coordinates are

never released in its outputs. Instead, all products—including dynamic maps, quantitative

summaries, and biogeographic narratives—are based on aggregated or generalized spatial

information. Geographic descriptors are expressed in terms of administrative or ecological

units (e.g., countries, basins, ecoregions, protected areas), while statistical summaries report

counts, densities, and extents without exposing point-level data. To ensure full Unicode

compatibility across platforms (Windows/macOS/Linux), all tabular outputs are exported in

TSV format, which avoids encoding corruption of special characters and remains

interoperable with biodiversity infrastructures.

**Results**

**Implementation and dataset processing**

We implemented che*CkOVER* using global crayfish (Astacidea) occurrence records from the

WoC® database [10], comprising 115,486 occurrence records for 465 taxa spanning 1758–

2025, contributed by the World of Crayfish® consortium members (contributor details

provided in Table S1). While designed as a taxon-agnostic framework, the WoC® dataset

provided comprehensive validation due to its global scope and well-documented metadata.

The data ingestion and preprocessing module successfully processed 96.1% of input

records (111,729 from 116,274 raw records), with 3,801 records removed due to coordinate

validation failures and 744 duplicates eliminated. Taxonomy resolution through external APIs

standardized 465 species names, while vernacular name harvesting from ITIS and expert

inputs yielded 713 common names across 35 languages. Multiple language versions of

recognized common names were retained using special characters without forcing translation

into English. Type locality information was successfully identified and preserved for 120

species (25.8% of total), each linked to original taxonomic descriptions with full bibliographic

references.

Geospatial enrichment successfully assigned all validated records to standardized

geographic units spanning 97 countries, hierarchically nested hydrographic basins

(HydroBASINS levels L06–L10), and terrestrial and freshwater ecoregions (TEOW/FEOW),

providing comprehensive geographic coverage for demonstrating framework scalability.

Protected area coverage was calculated for all species distributions using WDPA-designated

boundaries, enabling assessment of conservation context and identification of protection gaps.

**Quantitative descriptors and metrics**

All quantitative metrics used for conservation assessment—including EOO, AOO,

distributional categories (endemic, regional, cosmopolitan), and fragmentation signals—were

calculated exclusively from native occurrence records, ensuring compatibility with IUCN Red

List methodology. Spatial statistics derived from geographic overlays (country counts, basin

tallies, ecoregion assignments, protected area intersections) are computed for both native and
alien records but reported separately in all outputs.

Although both the IUCN Red List of Threatened Species (RLTS) and the Red List of
Ecosystems (RLE) apply structured risk assessment frameworks, they address fundamentally
different conservation targets and are not interchangeable. RLTS evaluates extinction risk at
the species level, where evidence of population reduction is central, whereas RLE focuses on
ecosystem collapse inferred from changes in distribution, function, and integrity [50,55].
Consequently, inference rules and criteria developed for ecosystem-level assessments cannot
be directly transposed to species-level Red List evaluations. Within che*CkOVER*,
distributional metrics such as EOO and AOO are therefore treated strictly as descriptive
spatial evidence supporting range delineation, spatial structure, and temporal comparison, and
not as direct indicators of threat or decline under RLTS Criterion A, where pressures are only
relevant insofar as they can be linked to demonstrable population reduction.

che*CkOVER* framework computes a standardized set of quantitative distributional
descriptors from both indigenous (Table 2) and non-indigenous (Table 3) occurrence records,
while maintaining strict stratification for reporting and visualization purposes. For native-
range assessments, species were classified into three distributional categories based on
indigenous range extent and geographic scope: 171 endemic taxa (36.8%), 287 regional taxa
(61.7%), and 5 cosmopolitan taxa (1.1%). These classifications guide IUCN-aligned
conservation metrics and adaptive spatial resolution assignment.

For invasion biology applications, non-indigenous ranges were analyzed separately
across 30 species documented with alien populations (Table 3). Range extent varied
dramatically, with 8 species exhibiting localized distributions (median 4 alien records,
restricted to 1–2 countries) and 22 species displaying widespread invasions (median 212 alien
records, spanning multiple countries or continents). The global invasion footprint is
dominated by a small number of highly successful invaders: *Procambarus clarkii* (34,361
alien records across 50 countries on 5 continents), *Faxonius limosus* (7,484 records, 23
countries), and *Pacifastacus leniusculus* (6,408 records, 29 countries spanning 3 continents)
collectively account for 89.8% of all non-indigenous occurrence data. Despite representing
only 6.5% of all processed species (30 of 465), these invasive populations generated 53,729
alien occurrence records—approaching parity with the 58,000 indigenous records from 463
native-range species—reflecting a disproportionate occurrence indexing effort.

**Table 2. Distributional metrics by biogeographic category for crayfish species with native-range**

**occurrences.** Species classified as Endemic (n=171), Regional (n=287), or Cosmopolitan (n=5). All metrics

calculated from indigenous records only (58,000 total). Values shown as median (range). EOO = extent of

occurrence; AOO = area of occupancy.

| Metric | Endemic | Regional | Cosmopolitan |
|---|---|---|---|
| **Extent of occurrence (EOO, km²)** | | | |
| Median (range) | 839 (0.4–4,875) | 30,495 (5,098–5,635,595) | 723,779 (13,140–13,394,185) |
| Mean ± SD | 1,340 ± 1,238 | 250,791 ± 593,274 | 3,116,189 ± 5,308,643 |
| **Area of occupancy (AOO, km²)** | | | |
| Median (range) | 24 (4–292) | 128 (4–14,248) | 40 (16–8,376) |
| Mean ± SD | 42 ± 47 | 577 ± 1,475 | 2,611 ± 3,737 |
| **Occurrence records per species** | | | |
| Median (range) | 8 (3–94) | 34 (1–5,405) | 10 (4–2,735) |
| Mean ± SD | 13 ± 16 | 179 ± 454 | 848 ± 1,198 |
| Total records | 2,260 (3.9%) | 51,500 (88.8%) | 4,240 (7.3%) |
| **Geographic extent** | | | |
| Countries range | 0–1☐ | 0–30☐ | 2–23 |
| Continent | 1 | 1 | 2 |
| HydroBASINS resolution | L10 (100%) | L08 (100%) | L06 (100%) |

553 ☐ species with 0 countries are those with occurrence records in disputed or unassigned territories
554

555 **Table 3. Non-indigenous range metrics for invasive crayfish species.** Species classified as Local (n=8) or

556 Widespread (n=22) based on introduced-range extent. All metrics from non-indigenous records only (53,729

557 total). Values shown as median (range). These metrics are informative only and do not contribute to conservation

558 assessments. EOO = extent of occurrence; AOO = area of occupancy.

| Metric | Local | Widespread | All |
|---|---|---|---|
| **Extent of occurrence (EOO, km²)** | | | |
| Median (range) | 195 (0.4–3,438) | 6,066,323 (130,631–510,066,073) | 966,419 (0.4–510,066,073) |
| Mean ± SD | 598 ± 1,149 | 83,110,373 ± 149,259,184 | 59,777,773 ± 130,583,522 |
| **Area of occupancy (AOO, km²)** | | | |
| Median (range) | 12 (4–80) | 358 (4–51,124) | 20 (4–51,124) |
| Mean ± SD | 20 ± 24 | 4,949 ± 11,668 | 3,303 ± 9,682 |
| **Occurrence records per species** | | | |
| Median (range) | 4 (3–20) | 212 (1–34,361) | 6 (1–34,361) |
| Mean ± SD | 6 ± 6 | 2,440 ± 7,485 | 1,791 ± 6,280 |
| Total records | 50 (0.09%) | 53,679 (99.91%) | 53,729 (100%) |
| **Geographic extent** | | | |
| Countries range | 1–2 | 1–50 | 1–50 |
| Continent range | 1 | 1–6 | 1–6 |
| HydroBASINS resolution | L10 (100%) | L06 (100%) | Mixed |

The distributional analysis revealed significant variation in species' geographic

extents. Endemic species showed restricted EOO values (median: 839 km²), while

cosmopolitan taxa displayed extensive ranges spanning multiple continents. Alien range

stratification identified 30 species with documented non-indigenous records, representing

6.5% of the processed dataset. For endemic and regional taxa, che*CkOVER* assessed the

spatial cohesion of native occurrence records through a simple fragmentation analysis. Of 458

species in the endemic and regional categories, 362 (79.0%) had sufficient occurrence data

(≥5 validated indigenous localities) for reliable clustering analysis. Distributional

fragmentation was detected in all 362 analyzed species, with cluster counts ranging from 2 to

25 (median: 5 clusters), reflecting varying degrees of spatial disjunction across native ranges.

The remaining 96 species (21.0%) had insufficient data points for clustering. Cosmopolitan

species were excluded from this analysis by design.

### 573 **Output generation and validation**

The framework successfully generated complete species packages for all 465 processed taxa.

Each package contains standardized outputs across multiple formats, including visual

representations (displaying native and alien ranges using distinct colors, with type locality

when available), geo-narratives (TXT, MD, JSON), and bibliographic metadata provided both

as APA-formatted citations and machine-readable CITATION.cff. Geospatial outputs were

generated at resolutions scaled to distributional categories: endemic species were exported

using high-resolution basin units (HydroBASINS L10), regional species used intermediate

resolution (L08), and cosmopolitan species used broader basin levels (L06). Native and alien

ranges are visually distinguished using distinct colors in all map outputs. All exported layers

incorporate privacy protection by aggregating spatial information instead of exposing raw

coordinate data.

Geo-narrative generation produced structured JSON outputs optimized for LLM

consumption alongside human-readable text summaries. Each narrative included standardized

sections covering geographic distribution, ecological status, temporal patterns, and data

quality assessments. Pre-computed answers to common queries (e.g., "Is this species

invasive?", "What is its biogeographic and preliminary conservation status?") were embedded

to facilitate automated responses.

### 592 **Framework performance and scalability**

che*CkOVER* demonstrated robust performance across the complete crayfish dataset.

Processing 465 species with 111,729 validated occurrence records required approximately 41

hours on standard hardware, with linear scaling observed for larger datasets. Peak memory

requirements reached approximately 220 GB during intensive geospatial operations,

particularly during high-resolution GeoJSON/KML generation and protected-area

intersections for species with very large occurrence datasets. The modular architecture

enabled selective execution of pipeline components, supporting both complete workflow runs

and targeted updates. Individual modules executed independently without framework

dependencies, facilitating maintenance and customization for different taxonomic groups or

data sources. Output validation confirmed metadata completeness (100% for core

descriptors), comprehensive bibliographic traceability with documented provenance chains,

and API accessibility supporting programmatic integration with external platforms.

che*CkOVER* has modest baseline computational requirements for standard workflows,

but resource usage scales with dataset size and the complexity of spatial operations. For small

to moderate datasets (e.g., up to a few thousand occurrence records per species), the pipeline

typically requires ≤8 GB RAM. Intermediate steps involving spatial overlays (e.g., basin and

protected-area intersections) may require in the order of tens of gigabytes of memory. For

species with very large occurrence datasets (tens of thousands of records), memory-intensive

operations such as high-resolution GeoJSON/KML generation and range polygon rendering

(AOO/EOO) can require substantially higher RAM (≥128 GB), particularly when processed

at fine spatial resolutions. Disk requirements scale primarily with the number of generated

geospatial artifacts and cached intermediate products; several tens to hundreds of gigabytes

may be required for full multi-species runs. Users are therefore encouraged to adjust

computational resources according to dataset size and intended output resolution.

**Discussion**

The crayfish dataset serves as a proof-of-concept demonstration of che*CkOVER*'s technical

capabilities across diverse distributional scenarios. The quantitative patterns reported reflect

the characteristics of available occurrence data rather than definitive biogeographic or

ecological assessments. Occurrence datasets inherently reflect reporting effort, accessibility,

taxonomic expertise, and monitoring priorities rather than complete sampling of actual

distributions. che*CkOVER* outputs are designed as standardized, reproducible syntheses of

available data to support—but not replace—expert-driven assessments, field validation, and

integration with additional evidence sources. The framework's value lies in its ability to

transform heterogeneous occurrence records into consistent, interoperable knowledge

products regardless of data quality or taxonomic group, rather than in biological conclusions

derived from any single dataset. The following discussion focuses on the framework's

technical design, capabilities, and broader implications for biodiversity informatics.

**User interaction paradigms and accessibility**

che*CkOVER* is designed to serve diverse user communities through complementary modes of

interaction. Direct access via APIs and downloadable species packages supports traditional

workflows for researchers, conservation practitioners, and institutions, providing complete

datasets with full methodological transparency. In parallel, indirect access through LLM

intermediaries could create conversational interfaces, allowing users to query species

knowledge without prior taxonomic expertise or database literacy.

A distinctive feature is the integration of multilingual vernacular names, systematically

tagged by language. Vernacular names typically reflect local cultural perspectives rather than

direct translation of scientific nomenclature. Incorporating them expands accessibility for

global users, enabling queries phrased in national languages or traditional ecological

terminology to be accurately matched to validated species records. For Indigenous and local

communities, whose ecological knowledge is encoded in vernacular systems rather than

formal taxonomies, this creates new avenues for recognition and connection. By embedding

vernacular names while preserving scientific rigor, che*CkOVER* establishes a two-way bridge

between cultural knowledge and biodiversity infrastructures.

Beyond technical performance, che*CkOVER* aligns with broader principles in

interdisciplinary biodiversity research, which emphasize that effective integration requires

coordinated attention to people, process, and perspectives within collaborative teams [56]. By

combining computational infrastructure with expert validation, multilingual access pathways,

and culturally grounded vernacular systems, the framework operationalizes these principles in

practice. Its API-ready, AI-streamable JSON outputs and dynamic versioned database

facilitate dialogue between data scientists, taxonomists, conservation practitioners, and users

rooted in local ecological knowledge, reinforcing the view that durable biodiversity

infrastructures emerge not only from technical design, but from sustained mutual respect and

shared epistemic perspectives.

**Toward dynamic biodiversity knowledge systems**

By embedding API readiness and automated output versioning, che*CkOVER* transforms

biodiversity repositories from static archives into dynamic knowledge systems. Outputs can

be regenerated as databases expand, taxonomies change, or new spatial references become

available. This capacity ensures that users—and AI systems—access current information

while retaining archived output versions for temporal or methodological analyses.

The framework's handling of extinction events exemplifies this temporal dimension:

documented population losses are preserved as historical context while being excluded from

current range metrics, enabling transparent tracking of biodiversity change over time. Such

dynamism is particularly valuable in fast-moving contexts such as invasion biology, disease

ecology, and climate-change impact assessments [57,58]. The continuous-integration model

positions biodiversity infrastructures as living systems—responsive to scientific progress

while preserving methodological rigor.

**Transforming biodiversity informatics for the AI era**

Digital technologies are becoming indispensable for accelerating biodiversity knowledge and

conservation, while also highlighting practical constraints such as accessibility, skills,

collaboration capacity, and environmental costs [59,60]. In this context, che*CkOVER*

embodies a simple yet powerful principle: that complex, multi-layered insights can emerge

from the most minimal but accurate datapoint in biodiversity science—the validated

geographic occurrence. From these basic, traceable units, the framework constructs

interoperable outputs that expand into biogeographic, ecological, and conservation insights of

high complexity. This minimalist-to-complex design shows how what might appear as a

trivial piece of information—the location of a species at a given point in space and time—can

be transformed into a rich knowledge package while retaining fidelity to the original

observation.

This approach contrasts with much of traditional biodiversity informatics, where raw

occurrence data, spatial products, and analytical results exist in separate silos, often with

limited connectivity [61]. che*CkOVER* explicitly bridges these divides: species-level

narratives, georeferenced distribution layers, and conservation metrics are all produced as

linked outputs, maintaining lineage back to the primary evidence. Building on earlier efforts

to enrich biodiversity data and harmonize infrastructures [40,62], che*CkOVER* advances the

field by making outputs not only interoperable but also directly consumable by large language

models. Unlike prior initiatives that emphasized standards or database interoperability alone,

che*CkOVER* packages biodiversity information in formats designed to serve simultaneously

human researchers and AI-driven reasoning systems.

**Addressing the provenance crisis and pathways of AI uptake**

A defining challenge of AI-mediated science is the erosion of source attribution. LLMs often synthesize information from diverse sources without maintaining clear provenance chains, leading to responses that may be factually accurate but lack verifiable attribution [63–65]. che*Ck*OVER directly targets this challenge by embedding standardized citation metadata in every output, so that AI systems and other downstream tools can, when appropriately configured, provide not only accurate information but also direct pathways to primary sources. This approach has broader implications for scientific integrity in the AI era. As LLMs become primary interfaces for accessing scientific knowledge [66,67], the ability to trace information back to peer-reviewed publications becomes essential for maintaining trust in science-based decision-making [68,69]. che*Ck*OVER's structured outputs enable RAG approaches where AI systems can dynamically access and cite specific biodiversity knowledge packages rather than relying solely on pre-trained information that may become outdated or lack source attribution.

Equally important, however, is understanding how and when biodiversity resources become accessible to AI. Current LLMs are not continuously updated from the web; instead, they integrate new data only through periodic retraining by providers [70]. This means that che*Ck*OVER outputs will enter the permanent core of LLM knowledge only when included in future training snapshots. In the interim, near real-time interoperability can still be achieved through web indexation and RAG-enabled systems, as well as through direct API endpoints that allow models with web connectivity to retrieve updated outputs on demand. By structuring outputs as JSON packages with embedded provenance, che*Ck*OVER maximizes its visibility for both long-term model training and immediate dynamic retrieval.

**Version-aware and incremental updates**

Biodiversity occurrence datasets are inherently dynamic, with new records, corrections, and extinction annotations accumulating continuously over time. Recomputing full species-level analyses from scratch at each update becomes increasingly impractical for data-rich taxa, particularly when computationally expensive geospatial operations (e.g., basin overlays, AOO/EOO estimation, map generation) are involved. To address this, che*Ck*OVER is designed around a version-aware and incrementally updatable architecture.

At the core of this design is the separation between raw occurrence records and derived spatial knowledge units. While point-level occurrences constitute the primary input, che*Ck*OVER stores and exposes species-level outputs in terms of aggregated, semantically

meaningful spatial entities (e.g., hydrographic basins, countries, protected areas), together

with associated occurrence counts and temporal metadata. These aggregated units, rather than

individual points, represent the stable state of a species known distribution at a given

processing version. Each run produces a versioned snapshot that records: (i) the data snapshot

date, (ii) the che*CkOVER* software version, and (iii) the complete set of spatial units currently

occupied by the species, separately for indigenous and non-indigenous records. For each unit,

the snapshot retains occurrence counts and, where applicable, extinction annotations. This

enables downstream comparisons between successive runs without requiring access to the full

historical point dataset.

When new data becomes available, che*CkOVER* can operate in an incremental update

mode. Instead of reprocessing the entire occurrence dataset, the framework identifies records

added or modified since the previous snapshot and performs geospatial enrichment only on

this delta subset. Newly intersected spatial units are merged with those stored in the previous

snapshot, and occurrence counts are updated accordingly. Conversely, when extinction

annotations remove the last known occurrence from a spatial unit, that unit is flagged as

absent from the current distribution while remaining available for historical context.

Geospatial artefacts such as basin-level distribution maps are regenerated from the

updated set of occupied spatial units, rather than recomputed from all raw points. This

approach dramatically reduces memory requirements and processing time for large datasets,

while preserving full reproducibility and traceability. Historical and current ranges are

explicitly distinguished, enabling quantitative assessment of range contraction or expansion

across versions.

This incremental, version-aware strategy ensures that che*CkOVER* outputs remain

scalable as biodiversity datasets grow, supports longitudinal analyses of distributional change,

and aligns with FAIR principles by making both data provenance and processing history

explicit. Importantly, it allows computationally intensive species (e.g., globally invasive

crayfish with tens of thousands of records) to be re-evaluated efficiently as new data

accumulate, without compromising methodological consistency or interpretability.

**Scalability and generalizability across taxonomic domains**

While our proof of concept focuses on crayfish (Astacidea), che*CkOVER*'s modular

architecture is designed for broad applicability across taxa. The ingestion module accepts

heterogeneous occurrence tables by automatically standardizing commonly used coordinate

and date formats (e.g., decimal degrees, DMS, split-degree fields, multiple date conventions)

and by recognizing alternative column labels for latitude, longitude, and year. Once

harmonized, all datasets are processed through the same internal schema, ensuring that

standardized outputs remain consistent across species.

The framework is robust to complex distributional patterns—including multi-

continental ranges, invasive populations, and temporally stratified records—making it suitable

for marine taxa, terrestrial organisms, fossil occurrences, or trait-based datasets with spatial

components. Minor adjustments may be required—for example, alternative vernacular-name

sources or modified distributional thresholds—but the workflow and output architecture

remain unchanged. This design philosophy balances technical flexibility with standardized

outputs, enabling interoperability across biological domains while preserving taxon-specific

nuance [71,72].

**Scientific impact and institutional applications**

The framework's relevance for conservation and policy is immediate. IUCN Red List

assessments, for example, rely on metrics to evaluate extinction risk [50]. che*Ck*OVER

automates these calculations with explicit bibliographic grounding, offering more efficient

and consistent workflows across taxonomic groups. In addition to standard range metrics, the

framework also reports simple distributional fragmentation signals for endemic and regional

taxa, offering an early indication of whether known localities form cohesive or spatially

discontinuous units. While intentionally minimalistic, these signals provide useful first-pass

context for connectivity considerations and flag cases where more detailed ecological or

genetic assessments may be warranted. The framework's explicit separation of indigenous and

non-indigenous distributions supports multiple analytical workflows. For species assessment

under IUCN criteria, only native range metrics are biologically meaningful—che*Ck*OVER

ensures these are computed exclusively from indigenous records. For invasion biology

applications, comprehensive tracking of alien populations enables researchers to identify

biosecurity priorities and evaluate management interventions using the separately computed

non-indigenous metrics. This dual-output design allows the same occurrence dataset to serve

conservation assessment and invasion monitoring workflows without methodological

compromise. Similarly, protected area agencies could leverage dynamically updated species

packages to identify coverage gaps, monitor range shifts, or prioritize management

interventions based on up-to-date knowledge. This capacity is particularly critical for

freshwater taxa, where limited benefits were noted due to their specific requirements on

conservation goals [73].

Beyond institutional contexts, che*CkOVER* is designed to open biodiversity

knowledge to wider audiences. Once its outputs are exposed through user-facing interfaces or

AI-mediated tools, non-specialists (including educators, journalists, and citizens) could obtain

scientifically grounded answers to basic questions about species. Crucially, each output links

back to primary sources, providing any downstream system with explicit citation pathways

and countering the drift toward derivative summaries that accumulate errors across successive

retellings. In this way, che*CkOVER* has the potential to foster trust and engagement with

biodiversity science outside academic circles.

**Future developments and broader applications**

The modular architecture of che*CkOVER* enables several natural extensions that could

significantly expand its analytical scope. Integration with trait databases, molecular

repositories, and ecological interaction networks would support multidimensional species

profiles that combine distributional knowledge with functional, phylogenetic, and ecological

context. Multi-species analyses represent another promising direction, generating narratives

that capture community-level patterns such as co-distributions, predator-prey dynamics, or

pathogen-host associations within the same standardized framework.

User-customizable applications offer immediate practical value for research and

conservation communities. Researchers could upload occurrence datasets to benchmark

sampling effectiveness, identify geographic coverage gaps, or evaluate survey completeness

against known distributions. Conservation organizations could integrate monitoring data with

che*CkOVER* baselines to assess management outcomes, track population trends, or prioritize

intervention strategies based on updated distributional knowledge.

A particularly compelling direction builds on the integrated distributional status

summary, which consolidates standardized geographic indicators such as EOO, AOO, and the

number of known and extinct-population localities, while also incorporating temporal signals

when available. The per-species geospatial layers exported by che*CkOVER* are explicitly

designed as reusable inputs for downstream analytical tools capable of deriving more

advanced fragmentation or range-disjunction indicators. The preliminary fragmentation

signals produced for endemic and regional taxa further create a natural bridge toward such

advanced workflows [74], offering early pointers to potential spatial breaks that downstream

tools may refine.

For taxa of conservation concern, such implementations could incorporate custom,

time-restricted analyses which—together with other dedicated outputs such as ecological,

molecular, or threat overlays—can generate tailored range summaries that more precisely

contextualize distributional patterns. These refinements would strengthen the utility of

che*Ck*OVER outputs in conservation workflows by providing clearer evidence of spatial

patterns relevant to Red List criteria [50]. While not a substitute for formal, expert-driven

IUCN assessments, such standardized and provenance-rich summaries can support early-stage

prioritization, help flag species exhibiting spatial decline or emerging fragmentation, and

supply consistent, interoperable inputs that streamline subsequent evaluation efforts [75,76].

## Conclusion

che*Ck*OVER establishes a new paradigm for biodiversity informatics: transforming minimal

yet validated datapoints—species occurrence records—into comprehensive, AI-interoperable

knowledge products that serve both human experts and computational systems. By

systematically converting occurrence data into structured geo-narratives, quantitative metrics,

and machine-readable outputs, the framework repositions biodiversity databases from passive

archives into active knowledge engines capable of interfacing with emerging AI technologies.

Three technical innovations distinguish che*Ck*OVER from existing biodiversity

infrastructures. First, the framework implements strict stratification of indigenous and non-

indigenous occurrence records, enabling simultaneous support for IUCN-aligned conservation

assessments and invasion biology applications without methodological compromise. Second,

all outputs embed standardized citation metadata and provenance chains, directly addressing

the attribution crisis in AI-mediated science by ensuring that biodiversity knowledge remains

traceable to peer-reviewed sources even when accessed through LLM intermediaries. Third,

the version-aware architecture enables incremental updates and longitudinal tracking,

allowing outputs to evolve dynamically as datasets expand while maintaining full

reproducibility and historical context.

The framework's modular, taxon-agnostic design ensures broad applicability beyond

the crayfish proof-of-concept. Any taxonomic group with occurrence records can be

processed through the same standardized pipeline, generating consistent outputs regardless of

data density, geographic extent, or distributional complexity. By integrating multilingual

vernacular names and delivering outputs in multiple interoperable formats (JSON, Markdown,

KML/KMZ, GeoJSON), che*Ck*OVER expands access beyond traditional research

communities to encompass educators, policymakers, conservation practitioners, and local

communities whose ecological knowledge is encoded in vernacular terminology.

As biodiversity faces accelerating threats from global change, the need for rapid,

accurate, and traceable knowledge synthesis has never been more urgent. che*CkOVER*

demonstrates that occurrence repositories—often dismissed as simple data archives—can

function as sophisticated knowledge infrastructure when coupled with appropriate

computational frameworks. Rather than treating AI as external to biodiversity science, the

framework illustrates how specialized scientific communities can proactively shape the

integration of their knowledge domains into AI-mediated systems, ensuring that biodiversity

information remains authoritative, verifiable, and accessible across diverse user communities

and technological platforms.

The open-source implementation, containerized deployment architecture, and

comprehensive documentation position che*CkOVER* for adoption by biodiversity platforms,

research institutions, and conservation organizations worldwide. Because all outputs are fully

data-driven, they represent reproducible snapshots of available occurrence data rather than

static assessments, updating naturally as datasets grow and scientific understanding evolves.

This living-systems approach aligns with the dynamic nature of biodiversity knowledge itself,

ensuring that che*CkOVER* outputs remain relevant and actionable as global monitoring efforts

intensify and new occurrence records accumulate.

## Availability of source code and requirements

Project name: che*CkOVER*

Project homepage: https://world.crayfish.ro/

Source code repository: https://github.com/dadvilfed/checkcover/, together with a

demonstration dataset (RollData) containing sample crayfish occurrence data, canonical

species geo-narrative template (File S1), and the offline lookup tables required in the

workflow (provided in TSV format to ensure full Unicode compatibility and to support

machine-readable reuse): vernacular names expert supplement (Table S2), hydronyms (Table

S3), and freshwater ecoregions (Table S4)

Operating system(s): cross-platform (Windows 10/11, Linux Ubuntu 20.04+, macOS 11+)

Programming language: R (≥4.5.2)

Other requirements: pandoc (≥2.0), GDAL (≥3.0), PROJ (≥7.0) libraries for spatial operations

License: GPL-3

Any restrictions to use by non-academics: none

RRID: [not yet assigned, all persistent identifiers will be registered upon final release]

bio.tools ID: [not yet assigned]

Local installation requirements: For typical use cases (datasets with <10,000 records per

species): 16 GB RAM, 4 CPU cores, 100 GB disk space. For large-scale applications

(>20,000 records per species or processing hundreds of taxa): 256+ GB RAM, 16+ CPU

cores, 500+ GB disk space recommended.

Server deployment: RESTful API endpoints for species packages and search functionality,

MariaDB for metadata indexing, CDN integration for global accessibility, Docker

containerization supported

## Additional files

**Supplementary Table S1.** World of Crayfish® Contributors, including full names and

institutional affiliations, who provided validated occurrence data, metadata, or region-specific

expertise that supported data verification, contextual interpretation, and the completeness of

species-level information used in this study.

**Supplementary Table S2.** Expert-curated vernacular names lookup table, supplementing

ITIS-sourced names and providing multilingual coverage.

**Supplementary Table S3.** Hydronym lookup table, linking HydroBASINS unit IDs (levels 6,

8, and 10) to their assigned hydronyms, derived by spatial intersection of basin polygons with

named river-line datasets.

**Supplementary Table S4.** Lookup table for freshwater ecoregions, linking FEOW unit IDs to

their standardized ecoregion names.

**Supplementary Table S5.** Software dependencies used by the che*CkOVER* pipeline. All

packages are automatically installed during the standard workflow execution, except for the

auxiliary datasets listed in Supplementary Tables S2–S4.

**Supplementary File S1.** Canonical che*CkOVER* species geo-narrative template used for

standardized human- and LLM-readable outputs.

## Abbreviations

AI: artificial intelligence; AOO: area of occupancy; API: application programming interface;

CSV: comma-separated values; EOO: extent of occurrence; FAIR: Findable, Accessible,

Interoperable, Reusable; FEOW: Freshwater Ecoregions of the World; IUCN: International

Union for Conservation of Nature; JSON: JavaScript Object Notation; KML/KMZ: Keyhole

Markup Language / Keyhole Markup Language Zipped; LLM: large language model;

MRBW: Major River Basins of the World; RAG: retrieval-augmented generation; TEOW:

Terrestrial Ecoregions of the World; TSV: tab-separated values; WDPA: World Database on

Protected Areas; WoC: World of Crayfish® platform..

## Consent for publication

936 This study involves species-occurrence data sourced from community contributors, following

937 the anonymization protocol of the WoC® platform [10], ensuring that no precise coordinates

938 are released in any che*CkOVER* output. All authors, including WoC® Contributors, have

939 reviewed and approved the final manuscript and consent to its publication.

## Acknowledgments

942 We thank Marian Neagul (West University of Timişoara) for configuring and maintaining the

943 server environment hosting the dynamic data platform, and Alexandru E. Mizeranschi

944 (Research and Development Station for Bovine, Arad, Romania) for technical advice during

945 pipeline testing.

## Author contributions

L.P.: conceptualization, investigation, methodology, supervision, validation, writing—original

draft. D.L.: software, methodology, formal analysis, visualization, writing—original draft.

V.I.B.: methodology, visualization, writing—original draft. C.I.N.: methodology,

visualization, writing—original draft. T.T.□.: methodology, visualization, writing—original

draft. A.V.L.: methodology, visualization, writing—original draft. WoC® Contributors:

supervision, validation. All: writing—review & editing.

## Use of AI tools

956 ChatGPT 5.1 model [77] was used for language refinement of the manuscript. No scientific

957 interpretations were generated by AI. All content was reviewed and fully validated by the

958 authors, who take full responsibility for the final text.

## Funding

961 This work received no project-based or grant funding. The Article Processing Charge was

962 supported through doctoral publication funds provided by the West University of Timişoara.

## Data availability

The processed dataset used in this study has been deposited in GigaDB as a static, citable snapshot (DOI: …….. to be assigned upon GigaDB deposition, files will be made available to editors and reviewers through GigaDB's private access mechanism during peer review, and released publicly upon publication). All raw occurrence records originate from the WoC$^®$ platform (https://world.crayfish.ro), which functions as a dynamic, continuously updated repository. The platform provides interactive access to current and future versions of the dataset, as well as to all che*Ck*OVER-generated outputs across subsequent releases.

## Competing interests

The authors declare that they have no competing interests.

## Ethical approval

The study was approved by an institutional review board at the West University of Timisoara under the identifier UVT2025-086385/24.11.2025.

## References

980     1. Oreskes N. Models all the way down. *Metascience 2011 211*. Springer; 2011; doi: 10.1007/S11016-011-9558-9.

2. Gladwell M. The tipping point: How little things can make a big difference. Little, Brown and Company;

3. Swedlow B. Four States of Knowledge: A Cultural Theory of the Co-Production of Science, Social Order, and Scientific, Cultural, and Policy Change. *SSRN Electron J*. Elsevier BV; 2010; doi: 10.2139/SSRN.1673521.

4. Boivin N, Crowther A. Mobilizing the past to shape a better Anthropocene. *Nat Ecol Evol 2021 53*. Nature Publishing Group; 2021; doi: 10.1038/s41559-020-01361-4.

5. Marburger JH. Science, technology and innovation in a 21st century context. *Policy Sci*. Springer; 2011; doi: 10.1007/S11077-011-9137-3.

6. Cardoso AW, Hestir EL, Slingsby JA, Forbes CJ, Moncrieff GR, Turner W, et al.. BioSCape combines local knowledge and remote-sensing technology for inclusive biodiversity science. *Nat Rev Biodivers 2025 11*. Nature Publishing Group; 2025; doi: 10.1038/s44358-024-00007-8.

7. Moor M, Banerjee O, Abad ZSH, Krumholz HM, Leskovec J, Topol EJ, et al.. Foundation models for generalist medical artificial intelligence. *Nat 2023 6167956*. Nature Publishing Group; 2023; doi: 10.1038/s41586-023-05881-4.

8. Dwivedi YK, Kshetri N, Hughes L, Slade EL, Jeyaraj A, Kar AK, et al.. "So what if ChatGPT wrote it?" Multidisciplinary perspectives on opportunities, challenges and implications of generative conversational AI for research, practice and policy. *Int J Inf Manage*. Elsevier Limited; 2023; doi: 10.1016/J.IJINFOMGT.2023.102642.

9. Leslie D. Does the sun rise for ChatGPT? Scientific discovery in the age of generative AI. *AI Ethics 2023 54*. Springer; 2023; doi: 10.1007/S43681-023-00315-3.

10. Ion MC, Bloomer CC, Bărăscu TI, Oficialdegui FJ, Shoobs NF, Williams BW, et al.. World of Crayfish™: a web platform towards real-time global mapping of freshwater crayfish and their pathogens. *PeerJ*. PeerJ Inc.; 2024; doi: 10.7717/PEERJ.18229.

11. Cortés-Guzmán D, Sinclair J, Hof C, Kalusche JB, Haase P. Dispersal, glacial refugia and temperature shape biogeographical patterns in European freshwater biodiversity. *Glob Ecol Biogeogr*. John Wiley & Sons, Ltd; 2024; doi: 10.1111/GEB.13886.

12. Poelen JH, Simons JD, Mungall CJ. Global biotic interactions: An open infrastructure to share and analyze species-interaction datasets. *Ecol Inform*. Elsevier; 2014; doi:

10.1016/J.ECOINF.2014.08.005.

13. McClanahan TR, Friedlander AM, Chabanet P, Bruggemann JH, Wickel J, Azali MK.

Modeling the spatial distribution of numbers of coral reef fish species and community types in

the Western Indian Ocean faunal province. *Mar Ecol Prog Ser*. Inter-Research; 2024; doi:

10.3354/MEPS14538.

14. Rodrigues ASL, Pilgrim JD, Lamoreux JF, Hoffmann M, Brooks TM. The value of the

IUCN Red List for conservation. *Trends Ecol Evol*. Elsevier Ltd; 2006; doi:

10.1016/j.tree.2005.10.010.

15. Uysal Bayar F. Determination of natural populations to be included in breeding program

in St. John's wort species (Hypericum perforatum L.). *PeerJ*. PeerJ Inc.; 2024; doi:

10.7717/PEERJ.18336.

16. Mace GM. Whose conservation? *Science (80- )*. American Association for the

Advancement of Science; 2014; doi: 10.1126/SCIENCE.1254704.

17. Bowler DE, Bjorkman AD, Dornelas M, Myers-Smith IH, Navarro LM, Niamir A, et al..

Mapping human pressures on biodiversity across the planet uncovers anthropogenic threat

complexes. *People Nat*. Blackwell Publishing Ltd; 2020; doi: 10.1002/PAN3.10071.

18. Chaplin-Kramer R, Miller CR, Dee LE, Bennett NJ, Echeverri A, Gould RK, et al..

Wildlife's contributions to people. *Nat Rev Biodivers 2025 11*. Nature Publishing Group;

2025; doi: 10.1038/s44358-024-00006-9.

19. Pereira HM, Martins IS, Rosa IMD, Kim HJ, Leadley P, Popp A, et al.. Global trends and

scenarios for terrestrial biodiversity and ecosystem services from 1900 to 2050. *Science (80-*

*)*. American Association for the Advancement of Science; 2024; doi:

10.1126/science.adn3441.

20. Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, et al..

Biodiversity loss and its impact on humanity. *Nat 2012 4867401*. Nature Publishing Group;

2012; doi: 10.1038/nature11148.

21. Vinuesa R, Azizpour H, Leite I, Balaam M, Dignum V, Domisch S, et al.. The role of

artificial intelligence in achieving the Sustainable Development Goals. *Nat Commun 2020*

*111*. Nature Publishing Group; 2020; doi: 10.1038/s41467-019-14108-y.

22. Christie AP, Downey H, Frick WF, Grainger M, O'Brien D, Tinsley-Marshall P, et al.. A

practical conservation tool to combine diverse types of evidence for transparent evidence-

based decision-making. *Conserv Sci Pract*. John Wiley & Sons, Ltd; 2022; doi:

10.1111/CSP2.579.

23. Constantino-Panopio C, De | R M H, Pandithasekera S, Nyandire | N O, Rahman | S C,

Mckee | E, et al.. An evaluation of tools being promoted to help the private sector disclose its

impacts on biodiversity. *Conserv Sci Pract*. John Wiley & Sons, Ltd; 2025; doi:

10.1111/CSP2.70198.

24. Ayoub NF, Lee YJ, Grimm D, Divi V. Head-to-Head Comparison of ChatGPT Versus

Google Search for Medical Knowledge Acquisition. *Otolaryngol Neck Surg*. John Wiley &

Sons, Ltd; 2024; doi: 10.1002/OHN.465.

25. Culina A, Baglioni M, Crowther TW, Visser ME, Woutersen-Windhouwer S, Manghi P.

Navigating the unfolding open data landscape in ecology and evolution. *Nat Ecol Evol*.

Nature Publishing Group; 2018; doi: 10.1038/s41559-017-0458-2.

26. Kung TH, Cheatham M, Medenilla A, Sillos C, De Leon L, Elepaño C, et al.. Performance

of ChatGPT on USMLE: Potential for AI-assisted medical education using large language

models. *PLOS Digit Heal*. Public Library of Science; 2023; doi:

10.1371/JOURNAL.PDIG.0000198.

27. Affinito F, Williams JM, Campbell JE, Londono MC, Gonzalez A. Progress in developing

and operationalizing the Monitoring Framework of the Global Biodiversity Framework. *Nat

Ecol Evol 2024 812*. Nature Publishing Group; 2024; doi: 10.1038/s41559-024-02566-7.

28. Sullivan BL, Aycrigg JL, Barry JH, Bonney RE, Bruns N, Cooper CB, et al.. The eBird

enterprise: An integrated approach to development and application of citizen science. *Biol

Conserv*. Elsevier; 2014; doi: 10.1016/J.BIOCON.2013.11.003.

29. Branco V V., Capinha C, Rocha J, Correia L, Cardoso P. SPECTRE: Standardised Global

Spatial Data on Terrestrial SPecies and ECosystems ThREats. *Glob Ecol Biogeogr*. John

Wiley & Sons, Ltd; 2025; doi: 10.1111/GEB.13949.

30. Callaghan CT, Mesaglio T, Ascher JS, Brooks TM, Cabras AA, Chandler M, et al.. The

benefits of contributing to the citizen science platform iNaturalist as an identifier. *PLOS Biol*.

Public Library of Science; 2022; doi: 10.1371/JOURNAL.PBIO.3001843.

31. Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, et al.. Contribution of

citizen science towards international biodiversity monitoring. *Biol Conserv*. Elsevier; 2017;

doi: 10.1016/J.BIOCON.2016.09.004.

32. Costello MJ, Michener WK, Gahegan M, Zhang ZQ, Bourne PE. Biodiversity data should

be published, cited, and peer reviewed. *Trends Ecol Evol*. Elsevier; 2013; doi:

10.1016/j.tree.2013.05.002.

33. Amano T, Sutherland WJ. Four barriers to the global understanding of biodiversity

conservation: Wealth, language, geographical location and security. *Proc R Soc B Biol Sci*.

Royal Society; 2013; doi: 10.1098/RSPB.2012.2649.

34. Feng X, Smith AB, Boyle B, Chen X, Enquist BJ, Gallagher R, et al.. The next stage of
biodiversity informatics: community-driven synthesis and integration of biodiversity
databases. *Bioscience*. 2025; doi: 10.1093/BIOSCI/BIAF112.
35. Löffler F, Wesp V, König-Ries B, Klan F. Dataset search in biodiversity research: Do
metadata in data repositories reflect scholarly information needs? *PLoS One*. Public Library
of Science; 2021; doi: 10.1371/JOURNAL.PONE.0246099.
36. Stephens CR, González-Salazar C, Del M, Villalobos-Segura C, Marquet PA. Can
Ecological Interactions be Inferred from Spatial Data? *Biodivers Informatics*. The University
of Kansas; 2020; doi: 10.17161/BI.V15I1.9815.
37. Zyoud S, Zyoud AH. Mapping and visualizing global knowledge on planetary health in
the climate change context: a comprehensive exploration of insights, trends, and research
priorities. *Discov Sustain 2024 51*. Springer; 2024; doi: 10.1007/S43621-024-00499-6.
38. Zavalin V, Zavalina OL. Are we there yet? Evaluation of AI-generated metadata for
online information resources. *Inf Res an Int Electron J*. University of Boras; 2025; doi:
10.47989/IR30ICONF47215.
39. Yang X, Jia J, Zhou X, Wang S. The future of artificial intelligence: Time to embrace
more international collaboration. *Innov*. Elsevier; 2024; doi: 10.1016/J.XINN.2024.100703.
40. Vos RA, Biserkov JV, Balech B, Beard N, Blissett M, Brenninkmeijer C, et al.. Enriched
biodiversity data as a resource and service. *Biodivers Data J 2 e1125*. Pensoft Publishers;
2014; doi: 10.3897/BDJ.2.E1125.
41. Heberling JM, Miller JT, Noesgaard D, Weingart SB, Schigel D. Data integration enables
global biodiversity synthesis. *Proc Natl Acad Sci U S A*. National Academy of Sciences;
2021; doi: 10.1073/pnas.2018093118.
42. Pollock LJ, Kitzes J, Beery S, Gaynor KM, Jarzyna MA, Mac Aodha O, et al.. Harnessing
artificial intelligence to fill global shortfalls in biodiversity knowledge. *Nat Rev Biodivers*
*2025*. Nature Publishing Group; 2025; doi: 10.1038/s44358-025-00022-3.
43. Stenhouse A, Fisher N, Lepschi B, Schmidt-Lebuhn A, Rodriguez J, Turco F, et al.. A
vision of human–AI collaboration for enhanced biological collection curation and research.
*Bioscience*. 2025; doi: 10.1093/BIOSCI/BIAF021.
44. Jordão LSB, Morim MP, Baumgratz JFA, Simon MF, Eppinghaus ALC, Calfo VA.
TypeTaxonScript: sugarifying and enhancing data structures in biological systematics and
biodiversity research. *Biol Methods Protoc*. Oxford Academic; 2024; doi:
10.1093/BIOMETHODS/BPAE017.
45. Hegde H, Vendetti J, Goutte-Gattat D, Caufield JH, Graybeal JB, Harris NL, et al.. A

change language for ontologies and knowledge graphs. *Database*. Oxford Academic; 2025;

doi: 10.1093/DATABASE/BAAE133.

46. Wilkinson MD, Dumontier M, Aalbersberg IjJ, Appleton G, Axton M, Baak A, et al.. The

FAIR Guiding Principles for scientific data management and stewardship. *Sci Data 2016 31*.

Nature Publishing Group; 2016; doi: 10.1038/sdata.2016.18.

47. Anthropic: Essential grammar topics for high school graduates [Generative AI chat].

Claude Sonnet 4. https://claude.ai/share/329173b2-ec93-4663-ac68-4f65ea4f166d (2025).

Accessed 2025 Dec 19.

48. Canhos VP, Souza S de, Giovanni R De, Canhos DAL. Global Biodiversity Informatics:

setting the scene for a "new world" of ecological forecasting. *Biodivers Informatics*. The

University of Kansas; 2004; doi: 10.17161/BI.V1I0.3.

49. Guralnick R, Hill A. Biodiversity informatics: automated approaches for documenting

global biodiversity patterns and processes. *Bioinformatics*. Oxford Academic; 2009; doi:

10.1093/BIOINFORMATICS/BTN659.

50. IUCN Standards and Petitions Committee. Guidelines for Using the IUCN Red List

Categories and Criteria. 16th ed. IUCN.

51. Majer J. Saga of the short-range endemic. *Aust J Entomol*. John Wiley & Sons, Ltd; 2009;

doi: 10.1111/J.1440-6055.2009.00715.X.

52. Crandall KA, Buhay JE. Global diversity of crayfish (Astacidae, Cambaridae, and

Parastacidae—Decapoda) in freshwater. *Freshw Anim Divers Assess*. Springer, Dordrecht;

2007; doi: 10.1007/978-1-4020-8259-7_32.

53. Richman NI, Böhm M, Adams SB, Alvarez F, Bergey EA, Bunn JJS, et al.. Multiple

drivers of decline in the global status of freshwater crayfish (Decapoda: Astacidea). *Philos*

*Trans R Soc B Biol Sci*. Royal Society of London; 2015; doi: 10.1098/rstb.2014.0060.

54. R Core Team. R: A Language and Environment for Statistical Computing. R Core Team.

Viena, Austria;

55. IUCN. IUCN red list categories and criteria. Second edi. Switzerland and Cambridge, UK:

IUCN--The World Conservation Union;

56. Stenseke M, Harris FL, Crowe T. Advice for interdisciplinary biodiversity research.

*Conserv Sci Pract*. John Wiley & Sons, Ltd; 2025; doi: 10.1111/CSP2.70203.

57. Seebens H, Meyerson LA, Richardson DM, Lenzner B, Tricarico E, Courchamp F, et al..

Biological invasions: a global assessment of geographic distributions, long-term trends, and

data gaps. *Biol Rev*. John Wiley & Sons, Ltd; 2025; doi: 10.1111/BRV.70058.

58. Guareschi S, Cancellario | T, Oficialdegui | F J, Clavero | M. Insights from the past:

Invasion trajectory and niche trends of a global freshwater invader. *Glob Chang Biol*. John

Wiley & Sons, Ltd; 2024; doi: 10.1111/GCB.17059.

59. Freitas H, Gouveia AC. Biodiversity futures: digital approaches to knowledge and

conservation of biological diversity. *Web Ecol*. Copernicus Publications; 2025; doi:

10.5194/WE-25-29-2025.

60. Hasan Thung P, Damatashia P, Siti |, Kartikawati M, Muflihati |, Pritchard | Rose.

Enhancing data justice in community-led conservation: A case study from Indonesian Borneo.

*Conserv Sci Pract*. John Wiley & Sons, Ltd; 2026; doi: 10.1111/CSP2.70230.

61. Feng X, Enquist BJ, Park DS, Boyle B, Breshears DD, Gallagher R V., et al.. A review of

the heterogeneous landscape of biodiversity databases: Opportunities and challenges for a

synthesized biodiversity knowledge base. *Glob Ecol Biogeogr*. John Wiley & Sons, Ltd;

2022; doi: 10.1111/GEB.13497.

62. Bisby FA. The Quiet Revolution: Biodiversity Informatics and the Internet. *Science (80- )*.

American Association for the Advancement of Science; 2000; doi:

10.1126/SCIENCE.289.5488.2309.

63. Lim JS, Schneider E, Grover M, Zhang J, Peters D. Effects of AI versus human source

attribution on trust and forgiveness in the identical corporate apology statement for a data

breach scandal. *Public Relat Rev*. JAI; 2025; doi: 10.1016/J.PUBREV.2024.102520.

64. Sahebi S, Formosa P. The AI-mediated communication dilemma: epistemic trust, social

media, and the challenge of generative artificial intelligence. *Synthese*. Springer Science and

Business Media B.V.; 2025; doi: 10.1007/S11229-025-04963-2.

65. Hohenstein J, Jung M. AI as a moral crumple zone: The effects of AI-mediated

communication on attribution and trust. *Comput Human Behav*. Pergamon; 2020; doi:

10.1016/J.CHB.2019.106190.

66. Guo Q, Zhen J, Wu F, He Y, Qiao C. Can Students Make STEM Progress With the Large

Language Models (LLMs)? An Empirical Study of LLMs Integration Within Middle School

Science and Engineering Practice. *J Educ Comput Res*. SAGE Publications Inc.; 2025; doi:

10.1177/073563312413123.

67. Bewersdorff A, Hartmann C, Hornberger M, Seßler K, Bannert M, Kasneci E, et al..

Taking the next step with generative artificial intelligence: The transformative role of

multimodal large language models in science education. *Learn Individ Differ*. Elsevier Ltd;

2025; doi: 10.1016/j.lindif.2024.102601.

68. Suazo-Galdames IC, Saracostti M, Chaple-Gil AM. Scientific evidence and public policy:

a systematic review of barriers and enablers for evidence-informed decision-making. *Front*

*Commun*. Frontiers Media SA; 2025; doi: 10.3389/fcomm.2025.1632305.

69. Ulibarri N. Collaborative model development increases trust in and use of scientific

information in environmental decision-making. *Environ Sci Policy*. Elsevier; 2018; doi:

10.1016/J.ENVSCI.2018.01.022.

70. Kreuzberger D, Kuhl N, Hirschl S. Machine Learning Operations (MLOps): Overview,

Definition, and Architecture. *IEEE Access*. Institute of Electrical and Electronics Engineers

Inc.; 2023; doi: 10.1109/ACCESS.2023.3262138.

71. Otegui J, Guralnick RP. The geospatial data quality REST API for primary biodiversity

data. *Bioinformatics*. Oxford Academic; 2016; doi: 10.1093/BIOINFORMATICS/BTW057.

72. Cham H, Malek S, Milow P, Song C. Developing an ecological visualization system for

biodiversity data. *All Life*. Taylor & Francis; 2022; doi: 10.1080/26895293.2022.2066195.

73. Sinclair JS, Stubbington R, Welti EAR, Aroviita J, Baker NJ, Cañedo-Argüelles M, et al..

Current protected areas provide limited benefits for European river biodiversity. *Nat Commun

2025 161*. Nature Publishing Group; 2025; doi: 10.1038/s41467-025-67125-5.

74. Pârvulescu L, Iorgu E-I, Zaharia C, Ion MC, Satmari A, Krapal A-M, et al.. The future of

endangered crayfish in light of protected areas and habitat fragmentation. *Sci Reports 2020

101*. Nature Publishing Group; 2020; doi: 10.1038/s41598-020-71915-w.

75. Ion MC, Ács A-R, Laza AV, Lorincz I, Livadariu D, Lamoly AM, et al.. Conservation

status of the idle crayfish Austropotamobius bihariensis Pârvulescu, 2019. *Glob Ecol

Conserv*. Elsevier; 2024; doi: 10.1016/J.GECCO.2024.E02847.

76. Pârvulescu L. Community-Driven IUCN Red List Assessments for European Crayfish: A

Call to Action. *Freshw Crayfish*. International Association of Astacology; 2025; doi:

10.5869/FC.2025.V30-1.61.

77. OpenAI. High school grammar concepts [Generative AI chat]. ChatGPT.

1207

# che*Ck*OVER workflow