cheCkOVER: An open framework and AI-ready global crayfish database

for next-generation biodiversity knowledge

Running title: AI-interoperable biodiversity knowledge

Lucian PÂRVULESCU[1,2*], David LIVADARIU[1,3*], Victor I. BÂCU[4], Constantin I.

NANDRA[4], Teodor T. ȘTEFĂNUȚ[4] and World of Crayfish® Contributors[5]

[1] Crayfish Research Centre, Institute for Advanced Environmental Research, West University

of Timisoara, Oituz 4, 300086 Timisoara, Romania

[2] Department of Biology, Faculty of Chemistry, Biology, Geography, West University of

Timisoara, Pestalozzi 16A, 300115 Timisoara, Romania

[3] Department of Informatics, Faculty of Mathematics and Computer Science, West University

of Timisoara, Vasile Pârvan 4, 300223 Timisoara, Romania

[4] Artificial Intelligence Research Institute, Computer Science Department, Technical

University of Cluj-Napoca, Memorandumului 28, 400114 Cluj-Napoca, Romania

[5] World of Crayfish® Contributors (authors information provided in Table S1)

*Correspondence: lucian.parvulescu@e-uvt.ro, david.livadariu02@e-uvt.ro

Abstract

Background

Species occurrence records represent the backbone of biodiversity science, yet their utility is

often limited to spatial analyses, coarse distribution maps, or presence-absence models.

Current biodiversity infrastructures rarely provide computational formats directly usable by

modern artificial intelligence (AI) systems, such as large language models (LLMs), which

increasingly mediate scientific communication and knowledge synthesis. Open frameworks

that convert biodiversity occurrences into structured, machine-accessible knowledge are

therefore essential, particularly when they enable rapid integration of new records and near

real-time generation of spatial metrics and human-interpretable reports. Such capabilities

substantially reduce the latency between data acquisition and decision support, enabling rapid

production of updated range summaries and narrative outputs consumable by both domain

experts and AI-assisted decision-making systems.

Results

We introduce che*Ck*OVER, an open framework that converts raw species occurrence datasets into standardized, API-ready, multi-layered outputs: biogeographic descriptors, dynamic distribution maps, summary metrics, and structured JSON biogeographic-narratives, ready for rapid integration into conservation evaluation workflows. Each product embeds provenance metadata to ensure transparency, traceability, and citation. We applied the pipeline to 115,434 global crayfish (Astacidea) occurrence records, generating an AI-ready knowledge base of 458 species packages. This demonstrates how the framework transforms minimal datapoints, validated species occurrences, into interoperable knowledge consumable by both humans and computational systems. The JSON outputs are optimized for retrieval-augmented generation, enabling AI systems to dynamically access and cite biodiversity knowledge beyond their pre-training corpora.

Conclusions

che*Ck*OVER is taxon-generalizable and establishes a reproducible pathway from biodiversity occurrences to narrative-ready, AI-interoperable knowledge. The framework's dual output, a generalizable pipeline and taxon-specific knowledge databases, enables flexible reuse across conservation research, policy reporting, and AI-driven applications. This minimalist-to-complex design extends the reach of biodiversity data beyond traditional analyses, positioning occurrence repositories as pivotal engines for next-generation biodiversity knowledge systems.

Significance statement

Biodiversity infrastructures remain underused by modern AI systems despite their central role in science and society. che*Ck*OVER embodies a minimalist-to-complex paradigm: from the validated geographic occurrence of a species, a datapoint often perceived as trivial, it derives structured, multi-layered outputs linking distribution, status, and standardized geographic indicators. These outputs are natively optimized for retrieval-augmented generation and other machine-consumable workflows, enabling future AI systems to dynamically access and cite biodiversity knowledge beyond their pre-training corpora. Using a global crayfish dataset as proof of concept, we illustrate how raw occurrence records can scale into rich, interoperable biogeographic knowledge products with immediate value for both human synthesis and computational applications. This positions biodiversity databases as critical engines for next-

67 generation science, policy, and societal decision-making. The resulting standardized outputs

68 can be directly incorporated into conservation evaluation workflows, where transparent,

69 reproducible, and provenance-rich occurrence-based metrics are essential.

71 **Keywords**: AI-ready knowledge; biodiversity informatics; geo-narratives; JSON database;

72 spatial analytics

Introduction

Human societies have always relied on knowledge to guide decisions and shape progress [1–3]. From antiquity to the digital era, science has served as a trusted source of verified information [4], while new technologies have multiplied the ways knowledge flows into public and policy domains [5,6]. Today, artificial intelligence (AI), particularly large language models (LLMs), is emerging as a dominant mediator of communication, offering rapid access to information across disciplines and audiences [7–9].

Species distribution records are highly valuable for numerous purposes, from scientific research [10,11] and understanding species communities [12,13], to conservation planning and management [14], and ultimately to certain economic considerations [15]. In particular, the quality of stored and delivered biodiversity knowledge is essential [16]. Nature underpins human well-being [17,18], and its understanding depends on reliable data about both species and the habitats they occupy [19,20]. Biodiversity knowledge must not only be abundant, but also accurate, transparent and easy to integrate into decision-making [21]. In this domain, misinformation or poorly contextualized data can directly affect conservation outcomes, policy priorities, and public trust [22,23].

However the speed and scale of LLMs analyses come at a cost: LLMs generate fluent but often vague, erroneous, or unverifiable outputs [24,25]. Their training depends on heterogeneous and inconsistently curated data, leaving a fundamental gap between scientific knowledge production and the AI systems that now mediate it [26]. Simultaneously, scientific data are expanding at unprecedented rates [27]. Advances in modeling, genomics, and ecological monitoring—together with citizen science—have produced millions of biodiversity records and traits [28–30]. Dedicated infrastructures and portals have proliferated, enabling access to vast repositories of occurrence data, images, and observations [31]. Yet the value of these resources remains underexploited [32,33]. These resources typically generate outputs that AI systems cannot directly consume, leaving only fragments of biodiversity knowledge contextualized in their reporting [34,35].

While humans digest information through narratives and visualizations [36,37], AI requires structured inputs: machine-readable formats (e.g., JavaScript Object Notation – JSON) annotated with explicit metadata and provenance [38–40]. To enable the full integration of biodiversity science in the AI era [41,42], infrastructures must evolve from human-facing repositories into dual-purpose systems [43–45]. Here we present che*CkOVER*, the first generalizable framework designed to fill this gap. At its core, che*CkOVER*

deliberately returns to the roots of faunistics—validated raw occurrence records without prior

interpretation—and transforms them into standardized, multi-layered outputs that are both

human-readable and LLM-digestible. Using global crayfish (Astacidea) occurrence data [10]

as a proof of concept, we illustrate how the framework scales minimal datapoints into

interoperable, machine-readable knowledge packages enriched with provenance metadata.

These outputs are structured to support retrieval-augmented and other machine-consumable

workflows, positioning biodiversity infrastructures to interface effectively with emerging AI

technologies. In this way, che*CkOVER* reframes occurrence repositories as future-ready

knowledge engines, enabling broader reuse, clearer attribution, and improved integration

across scientific, conservation, and societal domains.

Methods

Framework description

che*CkOVER* is built as a modular architecture that transforms validated biodiversity

occurrence records into standardized, multi-layered knowledge packages. Each package is

generated at the species level and integrates visual, quantitative, and narrative components,

ensuring outputs are both human-readable and machine-interoperable. The name cheCkOVER

reflects the framework's systematic checking of occurrences over various boundaries,

signaling a step toward consolidated biodiversity knowledge infrastructure.

The workflow consists of five logical components implemented as a modular

processing pipeline (Figure 1). First, data ingestion and preprocessing involve loading

occurrence records, cleaning and standardizing fields (e.g., valid taxon name, geographic

coordinates, year of observation, native/alien status, and location accuracy), and optionally

resolving taxonomy through external services. At this stage, vernacular (common) names are

harvested from trusted sources, tagged by language and source, and embedded alongside the

Latin binomial for downstream narratives. Second, geospatial and quantitative enrichment

assigns an initial distributional category (endemic, regional, cosmopolitan) using interim

range statistics (e.g., EOO thresholds, country or continent counts). This early classification

guides downstream resolution choices (e.g., HydroBASINS level for overlays, map

generalization settings) and parameterization of range metrics. Occurrence and extinction

records are then overlaid with reference geospatial layers (countries, basins, ecoregions, and

protected areas), from which descriptive statistics are derived, including record counts,

temporal coverage, country-level distributions, density metrics, extent of occurrence (EOO),

and area of occupancy (AOO). Type locality information, when available, is treated as a

descriptive attribute and is explicitly linked to its bibliographic source. For endemic and

regional taxa, cheCkOVER additionally computes a simple distributional fragmentation signal

summarizing the spatial cohesion of known localities. Third, analytical results are transformed

into report-ready and machine-consumable outputs. Species-level artefact packages include

geospatial layers (KML/KMZ, GeoJSON), quantitative tables and reports, and LLM-ready

geo-narratives delivered in both machine and human-readable text. Each JSON narrative

embeds stable pointers to the full species-level bibliography, distributed as a separate, de-

duplicated artefact.

All outputs follow the FAIR principles, include output-level versioning (package

revision, timestamp, and manifest), and are exposed through documented, API-ready

endpoints to support integration with external biodiversity infrastructures and AI-based

systems [46].

Figure 1. Conceptual workflow of the che*Ck*O*VER* framework. The system can be read as five logical

components: (1) integrated biodiversity input databases; (2) data ingestion, cleaning, and enrichment; (3)

analytical processing of distributional and spatial metrics; (4) automated report and narrative generation; and (5)

production of modular human- and machine-consumable outputs. While internally implemented as independent

processing modules, the workflow is designed to support both immediate decision-ready reporting and continued

context-aware analysis through AI and LLM-based systems.

The che*CkOVER* pipeline was developed within a modern integrated development

environment combining Claude (Anthropic) AI-assisted code generation [47] with expert-

driven manual refinement. AI-generated code drafts were iteratively reviewed, validated,

optimized, and extended by the authors using real-world occurrence data subsets. This hybrid

development workflow ensured reproducibility, code transparency, and compliance with

biodiversity informatics best practices [48,49].

Data inputs and preprocessing

che*CkOVER* relies on streamlined occurrence datasets that are easy to maintain and extend

over time. The framework was developed and validated using the World of Crayfish® (WoC®)

database [10], but accepts occurrence data from any taxonomic source following the same

core schema. Only core fields required for geo-narrative and quantitative metric generation

are used: the valid Latin binomial of the taxon (with synonymy resolved via external APIs),

species status (indigenous vs. alien/introduced), year of observation, geographic coordinates

in standardized WGS84 decimal degrees, and a location-accuracy tag. Bibliographic

references for each record, including DOI/URL where available, are retained for traceability

and citation. All occurrence records are processed through a series of validation and

standardization operations using the *tidyverse* suite. Species names are cleaned, coordinates

coerced to WGS84, status normalized (native/alien/introduced), and accuracy flagged

(high/low). Duplicate records (species–year–coordinates–source) are removed, ensuring

consistency and reproducibility across datasets. Following preprocessing, species are

classified into distributional categories (endemic, regional, cosmopolitan), which then guide

the resolution of spatial overlays and downstream metrics. che*CkOVER* reads and

standardizes extinction tracking when present in input schemas (e.g., the Claim_extinction

field in WoC®). Records flagged as extinct are retained in the dataset for statistical reporting

but are excluded from all geospatial operations and quantitative metrics (dynamic maps,

ranges, EOO/AOO). Extinction events are therefore treated as documented historical

observations with ecological significance, preserving all attributes needed for transparent

reporting and future linkage across framework versions. This approach ensures temporal

completeness while maintaining accuracy of current distributional assessments, critical for

IUCN Red List evaluations and long-term monitoring [50].

To support multilingual, real-world queries, vernacular (common) names are not

required as inputs; instead, they are harvested downstream in a dedicated step from

authoritative and community-curated services (e.g., ITIS; https://www.itis.gov/), or

supplemented with expert-reviewed multilingual names compiled in a static lookup table

(Table S2). This step is implemented using the *ritis* package for querying authoritative

vernacular-name services. All vernacular names are normalized with ISO 639-3 language

tags. The resulting language-aware name lists are embedded in species outputs to improve

discoverability without complicating the primary input schema.

Geospatial enrichment

To generate narrative-ready outputs, che*CkOVER* links validated occurrence records to a suite

of standardized geospatial layers. These layers provide named, interpretable geographic

units—countries, administrative regions, river basins, ecoregions, and protected areas—that

allow species distributions to be described using stable, widely recognized terminology.

Ensuring correct names for these units is essential, and the framework therefore integrates

both programmatic attribution (via R packages) and reproducible lookup tables for cases

where names are absent from the original polygons. Country-level boundaries are sourced

from Natural Earth (via the *rnaturalearth* package), while sub-national administrative units

are retrieved through the *geodata* package, which provides standardized access to GADM v4.

Freshwater contexts are captured through HydroBASINS hierarchical watershed boundaries

(levels L06–L10) and HydroATLAS hydro-environmental descriptors, while WWF Terrestrial

Ecoregions of the World (TEOW) and Freshwater Ecoregions of the World (FEOW) are

assigned using the *ecoregions* and *feowR* packages. Because HydroBASINS polygons do not

include hydronym attributes, basin names were derived through a dedicated offline

geoprocessing step. HydroBASINS units (L06, L08, L10) were spatially intersected with

river-line datasets from the Global Runoff Data Centre (for L06–L08) and high-resolution

OpenStreetMap extracts (for L10). Each basin polygon was assigned the name of the longest

or most representative river crossing its area. This workflow yielded a static hydronym lookup

table linking HydroBASINS IDs to river names (Table S3), now integrated into the

che*CkOVER* pipeline. In contrast, TEOW polygons include standardized ecoregion names

directly via the *ecoregions* package; therefore, no additional processing was required. FEOW

polygons, however, lack embedded name attributes in some distributions, and a reproducible

lookup table linking FEOW IDs to ecoregion names was compiled (Table S4). Protected areas

were intersected using the World Database on Protected Areas (WDPA, Protected Planet),

retrieved programmatically via the *wdpar* package to ensure license-compliant access. All

geospatial operations—cropping, overlays, joins, reprojections, and geometry handling—are

executed using the *sf* package, which provides high-performance spatial data processing and
consistent handling of modern GIS geometries.

For range metrics, the framework computes extent of occurrence (EOO) for native
ranges as minimum convex polygons encompassing native validated records, following IUCN
guidelines [50]. To avoid artificial inflation in species with disjunct ranges across continents
or islands, EOO is calculated as the sum of convex hulls per contiguous landmass (or
freshwater system), preventing polygons spanning oceans. Area of occupancy (AOO) is
calculated as the number of occupied grid cells (typically 2×2 km) multiplied by cell area.
Occurrences flagged as extinct are removed prior to spatial overlays and range calculations,
ensuring that EOO and AOO represent only the current distribution. Extinct localities remain
available for reporting but do not influence geospatial metrics. All outputs distinguish
between native and alien ranges to maintain ecological relevance and ensure compatibility
with IUCN conservation standards. To maintain full compatibility with IUCN Red List
methodology, alien/introduced occurrences are never included in metric calculations, but are
retained for spatial reporting and visualization. Accordingly, all geospatial display products
(GeoJSON, KML/KMZ) show both native and alien ranges when present, while the
corresponding EOO/AOO values included in reports and JSON summaries derive strictly
from the native subset of records.

To contextualize the breadth of native distributions prior to spatial enrichment,
che*Ck*OVER classifies taxa into three distributional categories using native-range occurrence
statistics only. (i) Endemic taxa are defined as species with a native extent of occurrence
(EOO) ≤5,000 km², restricted to ≤2 countries within a single continent. These thresholds are
consistent with restricted-range criteria commonly applied to freshwater invertebrates [51]
and reflect the naturally fragmented distributions of Astacidea across isolated drainage
systems [52,53]. (ii) Regional taxa are species whose native ranges exceed endemic
thresholds but remain confined to a single continent, regardless of any secondary
introductions outside their native area. (iii) Cosmopolitan taxa are defined as species whose
native ranges span more than one continent, indicating broad historical biogeographic
distributions rather than anthropogenic spread. Records outside the native range (alien or
introduced occurrences) are explicitly retained and analyzed separately for descriptive
statistics, protected-area intersections, basin-level summaries, and cartographic outputs, but
they do not influence native-range classification. This separation ensures that biogeographic
categorization reflects evolutionary and historical range structure, while still capturing the full
contemporary distributional footprint of each species. This native-range classification guides

downstream spatial resolution choices (e.g., finer HydroBASINS levels for endemic taxa and

coarser resolutions for cosmopolitan taxa), ensuring internal consistency across quantitative

metrics and visual outputs.

For endemic and regional taxa, che*CkOVER* additionally computes a simple

distributional fragmentation signal to summarize the spatial cohesion of known native

localities. The analysis is performed on straight-line (Euclidean) distances, calculated after

transforming occurrence coordinates to an equal-area projection (EPSG:6933) to ensure

metric accuracy. A full pairwise distance matrix is generated, and the mean inter-point

distance is used as an adaptive species-specific threshold for hierarchical clustering (complete

linkage). This approach allows fragmentation detection to scale naturally with each species'

spatial configuration. Clusters are derived using hclust() and cutree() at the adaptive height.

Species with fewer than five valid locality records, as well as cosmopolitan taxa whose ranges

are inherently multi-continental, are excluded from clustering.

To minimize computational load during large-scale intersection operations,

che*CkOVER* implements an optimized, multi-stage spatial overlay procedure. For each

species, the framework first computes an early EOO polygon, buffered to account for spatial

uncertainty. All reference layers (countries, basins, protected areas) are dynamically cropped

to this bounding region prior to point-level intersections, reducing redundant geometry

checks. Spatial joins are executed in batches using indexed geometries (*sf* + S2 engine),

followed by tabular aggregation per species and per geographic unit. When applicable,

HydroBASINS levels are dynamically selected (L10 for endemic, L8 for regional, L6 for

cosmopolitan taxa), and protected areas are pre-filtered to remove marine or micro-polygons.

This adaptive workflow reduces memory requirements and computation time while

maintaining full spatial accuracy.

Narrative generation and outputs

The core function of che*CkOVER* is to transform enriched species occurrence records into

structured geo-narratives that are both human-readable and machine-interoperable. Species-

level outputs follow a standardized canonical geo-narrative template (Supplementary File S1),

implemented using the *glue* package, ensuring consistency across taxa and enabling fully

reproducible reporting for both human users and automated systems. Validated data are

organized into predefined narrative blocks covering taxonomic identity, indigenous and non-

indigenous range characteristics, occurrence origins, temporal coverage, data quality, and

biogeographic context (e.g., range limits, continental presence, overlap with protected areas).

While the framework relies on a fixed narrative structure to guarantee comparability and

reproducibility, it remains extensible, allowing the addition of new sections or user-defined

analytical queries as the system evolves.

Predefined query blocks are embedded within each narrative to support automated

responses to frequently asked questions, while the structured JSON outputs can be directly

consumed by large language models (LLMs) to address free-form prompts. When available,

type localities are highlighted as key descriptive attributes, explicitly linked to their

bibliographic sources, and incorporated into the indigenous range narrative. Vernacular names

harvested during earlier processing steps are automatically embedded in the JSON narratives

using ISO 639-3 language tags, enabling accurate querying with local or non-scientific

terminology without compromising taxonomic precision.

Each narrative includes a concise distributional summary, reported both in JSON and

tabular formats, presenting key metrics such as EOO, AOO, total numbers of validated

occurrences, and counts of records annotated as extinct. Extinction-related information is

reported transparently by aggregating annotations provided by primary data sources and

literature; che*CkOVER* does not infer extinction status, but summarizes the number of extinct

localities, their last recorded years, and the associated administrative or hydrological units.

For endemic and regional taxa, che*CkOVER* additionally reports a conservative

distributional fragmentation signal derived exclusively from indigenous occurrence records.

This signal contextualizes the spatial cohesion of known localities by identifying

geographically isolated clusters using adaptive, species-specific clustering thresholds.

Fragmentation outputs include (i) the number of detected spatial clusters, (ii) their relative

sizes expressed as occurrence counts and percentages, and (iii) a categorical indicator of

whether fragmentation was detected ("not detected" for a single cohesive cluster; "detected"

when multiple clusters are present). These signals are intended as supportive descriptors

aligned with IUCN criteria B and D [50], without constituting a formal Red List assessment.

Final outputs are delivered in multiple interoperable formats: (i) narrative reports for

human users, rendered via *rmarkdown* into TXT and MD; (ii) structured JSON files generated

using *jsonlite*, optimized for programmatic access and LLM-based workflows; and (iii)

geospatial artefacts (KML/KMZ and GeoJSON) supporting visualization and spatial

integration. Each JSON narrative embeds a persistent pointer to a de-duplicated, species-level

bibliography, ensuring full provenance, traceability, and unambiguous citation even when

accessed independently. A permanent "How to cite this resource" entry is maintained in the

che*CkOVER* documentation and will be updated following publication with the definitive

citation and DOI.

Implementation

The che*CkOVER* initial scripts were drafted and subsequently validated, optimized, and

extended by the authors, ensuring reproducibility and compliance. che*CkOVER* was

implemented in R v.4.3.3 [54] as a modular package, with all functionality exposed through

callable functions (Table 1). The workflow integrates sequential modules for ingestion,

enrichment, reporting, and packaging, yet each component can also be executed

independently. For transparency and reproducibility, all raw scripts are distributed within the

package (inst/scripts_raw/), enabling stepwise execution when required. The pipeline is

orchestrated by the run_pipeline() function, which reads a central configuration file (YAML),

sets paths and column mappings, filters taxa, and executes modules sequentially or selectively

with full logging and error handling.

Table 1. Core modules of the che*CkOVER* R package and their primary functions. A full inventory of supporting

R packages required by the pipeline, together with their roles and reference URLs, is provided in Table S5.

Module	Core function
ingest_clean()	Ingests raw occurrence records (XLSX/DB), validates the input schema through configurable column mapping (default WoC®-compatible), and standardizes core fields. Species names are cleaned, coordinates coerced to WGS84, status normalized (native/alien/introduced), and accuracy flagged. Duplicate records (species–year–coordinates–source) are removed; taxonomy may optionally be resolved via external APIs. Claim_extinction entries are tagged here and retained for reporting but excluded from geospatial and range calculations. Exports standardized tabular (CSV/TSV/RDS) and spatial (*sf*) objects.
vernacular_names()	Retrieves vernacular (common) names from prioritized sources (e.g., ITIS), normalizes spelling, removes duplicates, applies ISO 639-3 tags, and supports manual curation. Outputs consolidated name tables (TSV) and injects them into JSON narratives to enable queries using non-scientific names.
enrich_spatial()	Overlays validated occurrences with standardized geospatial layers: countries (Natural Earth), subnational units (GADM), hydrographic basins (HydroBASINS), ecoregions (FEOW/TEOW), major drainage systems (MRBW, FAO), and protected areas (WDPA). Implements an optimized workflow in which species' distributional categories (endemic, regional, cosmopolitan) guide basin resolution (L10, L8, L6). An early EOO polygon is used to dynamically crop reference layers before point-level intersections. Indexed geometry operations (*sf* + S2 engine) are batched and aggregated per species, balancing spatial precision and computational efficiency. Outputs enriched per-record spatial attributes and summary statistics for downstream modules.
generate_maps()	Produces standardized geospatial layers (KML/KMZ, GeoJSON) using the *sf* framework. Native and alien ranges are visually differentiated by color, with optional masking or aggregation for sensitive records. Map resolution adapts to the distributional category (L10 endemic, L8 regional, L6 cosmopolitan).

Module	Core function
	These layers are intended for downstream visualization, GIS workflows, and integration into external platforms.
build_reports()	Compiles quantitative species reports including record totals, temporal coverage, geographic tallies (countries, basins, protected areas), density metrics, and range statistics (EOO, AOO). For endemic and regional species, computes a simple distributional fragmentation signal based on native localities via Euclidean-distance clustering: a pairwise distance matrix is generated, the mean distance defines an adaptive threshold, and hierarchical clustering detects spatially cohesive groups. Cosmopolitan taxa are excluded from this analysis. Outputs include: number of clusters, cluster sizes (counts and percentages), and a categorical fragmentation flag ("none detected" vs. "detected"). Extinct localities are counted and summarized separately. Statistics for spatial overlays (countries, basins, ecoregions, protected areas) are computed for both native and alien records and reported distinctly. A unified distributional status summary consolidates these indicators. Exports tidy tables (TSV), JSON summaries, and human-readable reports (Markdown).
narrative_builder()	Generates structured geo-narratives using a standardized canonical template (Supplementary File S1), implemented using the *glue* package. Narratives cover distribution, indigenous vs. non-indigenous ranges, range limits, temporal patterns, and data-quality notes. Automatically annotates extinction events and incorporates type locality when available. Vernacular names are embedded with language tags. Outputs plain-text narratives, and LLM-ready JSON containing provenance metadata.
citation_manager()	Aggregates and de-duplicates references, preserves DOIs/URLs, and generates APA-formatted lists and a CITATION.cff metadata file. Embeds provenance links in all JSON outputs.
package_export()	Bundles all species-level outputs into distributable packages: geospatial layers (KML/KMZ, GeoJSON), quantitative tables (CSV, TSV), narratives (MD, JSON), and citation/metadata files. Generates manifests, checksums, and versioned outputs to support reproducibility and longitudinal tracking. Ensures provenance blocks are included in all JSON files.
run_pipeline()	Orchestrates sequential or selective module execution. Supports YAML config files, CLI parameters, logging, resume mode, and reproducibility reporting. Detects previous outputs in the working directory, records version history, and reports differences between current and earlier processed outputs.

cheCkOVER outputs clear, well-structured, machine-consumable metadata that allow

seamless integration in complex processing workflows and support efficient integration with

existing LLM services through RESTful API endpoints (e.g., /api/llm/{species_name}. We

comply with all the important aspects for providing a solution that is both API-ready and

LLM-ready: (1) machine consumable data and metadata, through standardized formats

(GeoJSON, KML/KMZ, MD, TSV etc.); (2) rich error semantics that improve API

discoverability and help AI agents and API integrators to understand data structure and

requirements, behavior and recover from errors; (3) speed and reliability ensured by a

containerization-layered technical stack, which provides seamless horizontal scalability at

each level: repository, processing, data conversions; (4) consistent REST end-points design

which, together with the standardized data and querying formats provides predictability; (5)

complete and well structured documentation available through OpenAPI specification. Data

storage combines species-level file hierarchies with a MariaDB relational database that

indexes taxonomic, geographic, and metadata attributes for efficient querying and retrieval.

Reproducibility and privacy protection

All code developed for che*CkOVER* is released as open source and publicly available in a

version-controlled GitHub repository. To ensure portability across systems, che*CkOVER* is

fully containerized and can be deployed both locally and within Docker or Singularity

environments. Stable releases are versioned and archived, ensuring long-term reproducibility.

This modular and container-ready design facilitates straightforward maintenance,

interoperability with other biodiversity infrastructures, and adaptability across taxonomic

groups.

Sensitive species records require protection against potential misuse, such as illegal

harvesting or habitat disturbance. che*CkOVER* ensures that raw geographic coordinates are

never released in its outputs. Instead, all products—including dynamic maps, quantitative

summaries, and biogeographic narratives—are based on aggregated or generalized spatial

information. Geographic descriptors are expressed in terms of administrative or ecological

units (e.g., countries, basins, ecoregions, protected areas), while statistical summaries report

counts, densities, and extents without exposing point-level data. To ensure full Unicode

compatibility across platforms (Windows/macOS/Linux), all tabular outputs are exported in

TSV format, which avoids encoding corruption of special characters and remains

interoperable with biodiversity infrastructures.

Results

Implementation and dataset processing

We implemented che*CkOVER* using global crayfish (Astacidea) occurrence records from the

WoC® database [10], comprising 115,434 occurrence records for 458 taxa spanning 1758-

2025. While designed as a taxon-agnostic framework, the WoC® dataset provided

comprehensive validation due to its global scope and well-documented metadata.

Taxonomy resolution through external APIs standardized 458 species names, while

vernacular name harvesting from ITIS and expert inputs yielded 713 common names across

35 languages. Multiple language versions of recognized common names were retained using

special characters without forcing translation into English. Type locality information was

successfully identified and preserved for 130 species (28.4% of total), each linked to original

taxonomic descriptions with full bibliographic references. Extinction tracking identified 135

documented population losses across 10 species. These extinction events span a temporal

range from 1971 to 2024, with 13.3% occurring post-2000. Extinct localities were retained in

statistical reporting but excluded from all current distributional analyses (maps, EOO/AOO

calculations), ensuring that range metrics reflect contemporary distributions while remaining

compatible with IUCN-aligned assessment workflows.

Quantitative descriptors and metrics

All quantitative metrics used for conservation assessment—including EOO, AOO,

distributional categories (endemic, regional, cosmopolitan), and fragmentation signals—was

calculated exclusively from native occurrence records, ensuring compatibility with IUCN Red

List methodology. Spatial statistics derived from geographic overlays (country counts, basin

tallies, ecoregion assignments, protected area intersections) are computed for both native and

alien records but reported separately in all outputs.

Although both the IUCN Red List of Threatened Species (RLTS) and the Red List of

Ecosystems (RLE) apply structured risk assessment frameworks, they address fundamentally

different conservation targets and are not interchangeable. RLTS evaluates extinction risk at

the species level, where evidence of population reduction is central, whereas RLE focuses on

ecosystem collapse inferred from changes in distribution, function, and integrity [50,55].

Consequently, inference rules and criteria developed for ecosystem-level assessments cannot

be directly transposed to species-level Red List evaluations. Within che*CkOVER*,

distributional metrics such as EOO and AOO are therefore treated strictly as descriptive

spatial evidence supporting range delineation, spatial structure, and temporal comparison, and

not as direct indicators of threat or decline under RLTS Criterion A, where pressures are only

relevant insofar as they can be linked to demonstrable population reduction.

che*CkOVER* framework computes a standardized set of quantitative distributional

descriptors EOO and AOO, calculated exclusively from native records (Table 2), while

indigenous and non-indigenous occurrences are stratified for reporting and visualization

purposes. Species are subsequently classified into endemic, regional, or cosmopolitan

distributional categories based on native-range geographic extent thresholds

Output generation and validation

The framework successfully generated complete species packages for all 458 processed taxa. Each package contains standardized outputs across multiple formats, including visual representations (display native and alien ranges using distinct colors, with type locality when available), geo-narratives (TXT, MD, JSON), and bibliographic metadata provided both as APA-formatted citations and machine-readable CITATION.cff. Geospatial outputs were generated at resolutions scaled to distributional categories: endemic species were exported using high-resolution basin units (HydroBASINS L10), regional species used intermediate resolution (L08), and cosmopolitan species used broader basin levels (L06). Native and alien ranges are visually distinguished using distinct colors in all map outputs. All exported layers incorporate privacy protection by aggregating spatial information instead of exposing raw coordinate data.

Geo-narrative generation produced structured JSON outputs optimized for LLM consumption alongside human-readable text summaries. Each narrative included standardized sections covering geographic distribution, ecological status, temporal patterns, and data quality assessments. Pre-computed answers to common queries (e.g., "Is this species invasive?", "What is its conservation preliminary status?") were embedded to facilitate automated responses.

Framework performance and scalability

che*Ck*OVER demonstrated robust performance across the complete crayfish dataset. Processing 458 species with 115,434 total records. The modular architecture enabled selective execution of pipeline components, supporting both complete workflow runs and targeted updates. Individual modules executed independently without framework dependencies, facilitating maintenance and customization for different taxonomic groups or data sources. Output validation confirmed metadata completeness (100% for core descriptors), bibliographic traceability, and API accessibility supporting programmatic integration with external platforms.

che*Ck*OVER has modest baseline computational requirements for standard workflows, but resource usage scales with dataset size and the complexity of spatial operations. For small to moderate datasets (e.g., up to a few thousand occurrence records per species), the pipeline typically requires ≤8 GB RAM. Intermediate steps involving spatial overlays (e.g., basin and protected-area intersections) may require in the order of tens of gigabytes of memory. For species with very large occurrence datasets (tens of thousands of records), memory-intensive operations such as high-resolution GeoJSON/KML generation and range polygon rendering

(AOO/EOO) can require substantially higher RAM (≥128 GB), particularly when processed

at fine spatial resolutions. Disk requirements scale primarily with the number of generated

geospatial artifacts and cached intermediate products; several tens to hundreds of gigabytes

may be required for full multi-species runs. Users are therefore encouraged to adjust

computational resources according to dataset size and intended output resolution.

Discussion

User interaction paradigms and accessibility

che*CkOVER* is designed to serve diverse user communities through complementary modes of

interaction. Direct access via APIs and downloadable species packages supports traditional

workflows for researchers, conservation practitioners, and institutions, providing complete

datasets with full methodological transparency. In parallel, indirect access through LLM

intermediaries could create conversational interfaces, allowing users to query species

knowledge without prior taxonomic expertise or database literacy.

A distinctive feature is the integration of multilingual vernacular names, systematically

tagged by language. Vernacular names typically reflect local cultural perspectives rather than

direct translation of scientific nomenclature. Incorporating them expands accessibility for

global users, enabling queries phrased in national languages or traditional ecological

terminology to be accurately matched to validated species records. For Indigenous and local

communities, whose ecological knowledge is encoded in vernacular systems rather than

formal taxonomies, this creates new avenues for recognition and connection. By embedding

vernacular names while preserving scientific rigor, che*CkOVER* establishes a two-way bridge

between cultural knowledge and biodiversity infrastructures.

Beyond technical performance, che*CkOVER* aligns with broader principles in

interdisciplinary biodiversity research, which emphasize that effective integration requires

coordinated attention to people, process, and perspectives within collaborative teams [56]. By

combining computational infrastructure with expert validation, multilingual access pathways,

and culturally grounded vernacular systems, the framework operationalizes these principles in

practice. Its API-ready, AI-streamable JSON outputs and dynamic versioned database

facilitate dialogue between data scientists, taxonomists, conservation practitioners, and users

rooted in local ecological knowledge, reinforcing the view that durable biodiversity

infrastructures emerge not only from technical design, but from sustained mutual respect and

shared epistemic perspectives.

Toward dynamic biodiversity knowledge systems

By embedding API readiness and automated output versioning, che*CkOVER* transforms biodiversity repositories from static archives into dynamic knowledge systems. Outputs can be regenerated as databases expand, taxonomies change, or new spatial references become available. This capacity ensures that users—and AI systems—access current information while retaining archived output versions for temporal or methodological analyses.

The framework's handling of extinction events exemplifies this temporal dimension: documented population losses are preserved as historical context while being excluded from current range metrics, enabling transparent tracking of biodiversity change over time. Such dynamism is particularly valuable in fast-moving contexts such as invasion biology, disease ecology, and climate-change impact assessments [57,58]. The continuous-integration model positions biodiversity infrastructures as living systems—responsive to scientific progress while preserving methodological rigor.

Transforming biodiversity informatics for the AI era

Digital technologies are becoming indispensable for accelerating biodiversity knowledge and conservation, while also highlighting practical constraints such as accessibility, skills, collaboration capacity, and environmental costs [59]. In this context, che*CkOVER* embodies a simple yet powerful principle: that complex, multi-layered insights can emerge from the most minimal but accurate datapoint in biodiversity science—the validated geographic occurrence. From these basic, traceable units, the framework constructs interoperable outputs that expand into biogeographic, ecological, and conservation insights of high complexity. This minimalist-to-complex design shows how what might appear as a trivial piece of information—the location of a species at a given point in space and time—can be transformed into a rich knowledge package while retaining fidelity to the original observation.

This approach contrasts with much of traditional biodiversity informatics, where raw occurrence data, spatial products, and analytical results exist in separate silos, often with limited connectivity [60]. che*CkOVER* explicitly bridges these divides: species-level narratives, georeferenced distribution layers, and conservation metrics are all produced as linked outputs, maintaining lineage back to the primary evidence. Building on earlier efforts to enrich biodiversity data and harmonize infrastructures [40,61], che*CkOVER* advances the field by making outputs not only interoperable but also directly consumable by large language

models. Unlike prior initiatives that emphasized standards or database interoperability alone,

che*CkOVER* packages biodiversity information in formats designed to serve simultaneously

human researchers and AI-driven reasoning systems.

Addressing the provenance crisis and pathways of AI uptake

A defining challenge of AI-mediated science is the erosion of source attribution. LLMs often

synthesize information from diverse sources without maintaining clear provenance chains,

leading to responses that may be factually accurate but lack verifiable attribution [62–64].

che*CkOVER* directly targets this challenge by embedding standardized citation metadata in

every output, so that AI systems and other downstream tools can, when appropriately

configured, provide not only accurate information but also direct pathways to primary

sources. This approach has broader implications for scientific integrity in the AI era. As LLMs

become primary interfaces for accessing scientific knowledge [65,66], the ability to trace

information back to peer-reviewed publications becomes essential for maintaining trust in

science-based decision-making [67,68]. che*CkOVER*'s structured outputs enable RAG

approaches where AI systems can dynamically access and cite specific biodiversity

knowledge packages rather than relying solely on pre-trained information that may become

outdated or lack source attribution.

Equally important, however, is understanding how and when biodiversity resources

become accessible to AI. Current LLMs are not continuously updated from the web; instead,

they integrate new data only through periodic retraining by providers [69]. This means that

che*CkOVER* outputs will enter the permanent core of LLM knowledge only when included in

future training snapshots. In the interim, near real-time interoperability can still be achieved

through web indexation and RAG-enabled systems, as well as through direct API endpoints

that allow models with web connectivity to retrieve updated outputs on demand. By

structuring outputs as JSON packages with embedded provenance, che*CkOVER* maximizes its

visibility for both long-term model training and immediate dynamic retrieval.

Version-aware and incremental updates

Biodiversity occurrence datasets are inherently dynamic, with new records, corrections, and

extinction annotations accumulating continuously over time. Recomputing full species-level

analyses from scratch at each update becomes increasingly impractical for data-rich taxa,

particularly when computationally expensive geospatial operations (e.g., basin overlays,

AOO/EOO estimation, map generation) are involved. To address this, che*CkOVER* is

designed around a version-aware and incrementally updatable architecture.

At the core of this design is the separation between raw occurrence records and

derived spatial knowledge units. While point-level occurrences constitute the primary input,

che*CkOVER* stores and exposes species-level outputs in terms of aggregated, semantically

meaningful spatial entities (e.g., hydrographic basins, countries, protected areas), together

with associated occurrence counts and temporal metadata. These aggregated units, rather than

individual points, represent the stable state of a species known distribution at a given

processing version. Each run produces a versioned snapshot that records: (i) the data snapshot

date, (ii) the che*CkOVER* software version, and (iii) the complete set of spatial units currently

occupied by the species, separately for indigenous and non-indigenous records. For each unit,

the snapshot retains occurrence counts and, where applicable, extinction annotations. This

enables downstream comparisons between successive runs without requiring access to the full

historical point dataset.

When new data becomes available, che*CkOVER* can operate in an incremental update

mode. Instead of reprocessing the entire occurrence dataset, the framework identifies records

added or modified since the previous snapshot and performs geospatial enrichment only on

this delta subset. Newly intersected spatial units are merged with those stored in the previous

snapshot, and occurrence counts are updated accordingly. Conversely, when extinction

annotations remove the last known occurrence from a spatial unit, that unit is flagged as

absent from the current distribution while remaining available for historical context.

Geospatial artefacts such as basin-level distribution maps are regenerated from the

updated set of occupied spatial units, rather than recomputed from all raw points. This

approach dramatically reduces memory requirements and processing time for large datasets,

while preserving full reproducibility and traceability. Historical and current ranges are

explicitly distinguished, enabling quantitative assessment of range contraction or expansion

across versions.

This incremental, version-aware strategy ensures that che*CkOVER* outputs remain

scalable as biodiversity datasets grow, supports longitudinal analyses of distributional change,

and aligns with FAIR principles by making both data provenance and processing history

explicit. Importantly, it allows computationally intensive species (e.g., globally invasive

crayfish with tens of thousands of records) to be re-evaluated efficiently as new data

accumulate, without compromising methodological consistency or interpretability.

Scalability and generalizability across taxonomic domains

While our proof of concept focuses on crayfish (Astacidea), che*Ck*OVER's modular

architecture is designed for broad applicability across taxa. The ingestion module accepts

heterogeneous occurrence tables by automatically standardizing commonly used coordinate

and date formats (e.g., decimal degrees, DMS, split-degree fields, multiple date conventions)

and by recognizing alternative column labels for latitude, longitude, and year. Once

harmonized, all datasets are processed through the same internal schema, ensuring that

standardized outputs remain consistent across species.

The framework is robust to complex distributional patterns—including multi-

continental ranges, invasive populations, and temporally stratified records—making it suitable

for marine taxa, terrestrial organisms, fossil occurrences, or trait-based datasets with spatial

components. Minor adjustments may be required—for example, alternative vernacular-name

sources or modified distributional thresholds—but the workflow and output architecture

remain unchanged. This design philosophy balances technical flexibility with standardized

outputs, enabling interoperability across biological domains while preserving taxon-specific

nuance [70,71].

Scientific impact and institutional applications

The framework's relevance for conservation and policy is immediate. IUCN Red List

assessments, for example, rely on metrics to evaluate extinction risk [50]. che*Ck*OVER

automates these calculations with explicit bibliographic grounding, offering more efficient

and consistent workflows across taxonomic groups. In addition to standard range metrics, the

framework also reports simple distributional fragmentation signals for endemic and regional

taxa, offering an early indication of whether known localities form cohesive or spatially

discontinuous units. While intentionally minimalistic, these signals provide useful first-pass

context for connectivity considerations and flag cases where more detailed ecological or

genetic assessments may be warranted. The framework's explicit separation of native and

alien distributions serves multiple conservation functions. For species assessment under IUCN

criteria, only native range metrics are biologically meaningful. However, comprehensive

tracking of alien populations remains essential for understanding invasion dynamics,

identifying biosecurity priorities, and evaluating management interventions. By computing

and displaying both ranges distinctly, che*Ck*OVER supports both conservation assessment

(native-focused) and invasion biology (requires both native and alien context). Similarly,

protected area agencies could leverage dynamically updated species packages to identify

coverage gaps, monitor range shifts, or prioritize management interventions based on up-to-

date knowledge. This capacity is particularly critical for freshwater taxa, where limited

benefits were noted due to their specific requirements on conservation goals [72].

Beyond institutional contexts, che*CkOVER* is designed to open biodiversity

knowledge to wider audiences. Once its outputs are exposed through user-facing interfaces or

AI-mediated tools, non-specialists (including educators, journalists, and citizens) could obtain

scientifically grounded answers to basic questions about species. Crucially, each output links

back to primary sources, providing any downstream system with explicit citation pathways

and countering the drift toward derivative summaries that accumulate errors across successive

retellings. In this way, che*CkOVER* has the potential to foster trust and engagement with

biodiversity science outside academic circles.

Future developments and broader applications

The modular architecture of che*CkOVER* enables several natural extensions that could

significantly expand its analytical scope. Integration with trait databases, molecular

repositories, and ecological interaction networks would support multidimensional species

profiles that combine distributional knowledge with functional, phylogenetic, and ecological

context. Multi-species analyses represent another promising direction, generating narratives

that capture community-level patterns such as co-distributions, predator-prey dynamics, or

pathogen-host associations within the same standardized framework.

User-customizable applications offer immediate practical value for research and

conservation communities. Researchers could upload occurrence datasets to benchmark

sampling effectiveness, identify geographic coverage gaps, or evaluate survey completeness

against known distributions. Conservation organizations could integrate monitoring data with

che*CkOVER* baselines to assess management outcomes, track population trends, or prioritize

intervention strategies based on updated distributional knowledge.

A particularly compelling direction builds on the integrated distributional status

summary, which consolidates standardized geographic indicators such as EOO, AOO, and the

number of known and extinct-population localities, while also incorporating temporal signals

when available. The per-species geospatial layers exported by che*CkOVER* are explicitly

designed as reusable inputs for downstream analytical tools capable of deriving more

advanced fragmentation or range-disjunction indicators. The preliminary fragmentation

signals produced for endemic and regional taxa further create a natural bridge toward such
advanced workflows, offering early pointers to potential spatial breaks that downstream tools
may refine.

For taxa of conservation concern, such implementations could incorporate custom,
time-restricted analyses which—together with other dedicated outputs such as ecological,
molecular, or threat overlays—can generate tailored range summaries that more precisely
contextualize distributional patterns. These refinements would strengthen the utility of
che*CkOVER* outputs in conservation workflows by providing clearer evidence of spatial
patterns relevant to Red List criteria [50]. While not a substitute for formal, expert-driven
IUCN assessments, such standardized and provenance-rich summaries can support early-stage
prioritization, help flag species exhibiting spatial decline or emerging fragmentation, and
supply consistent, interoperable inputs that streamline subsequent evaluation efforts [73].
668

Conclusion

che*CkOVER* is more than a technical solution to data formatting; it marks a shift toward
active integration between biodiversity science and AI-mediated knowledge systems. By
enabling occurrence repositories to be used as structured inputs for AI-assisted responses, the
framework positions biodiversity research as a foundational component for understanding
environmental challenges.

As global change accelerates, accurate and traceable knowledge becomes critical not
only for researchers and conservation practitioners but also for broader public discourse. AI
systems capable of delivering reliable answers on species distributions, conservation status,
and ecological relationships can strengthen the scientific basis of governance and policy. By
embedding multilingual vernacular names, che*CkOVER* further expands accessibility, making
biodiversity knowledge discoverable in national languages and cultural contexts, and opening
new pathways for educators, journalists, and local communities.

The framework outlines a concrete pathway by which biodiversity databases could
improve the accuracy and traceability of AI-assisted outputs while preserving scientific
integrity, offering a model that other data-intensive disciplines might adapt. By providing
standardized spatial summaries, including preliminary signals of potential range
fragmentation, che*CkOVER* also creates opportunities for more informed and nuanced AI-
mediated ecological reasoning. Rather than treating AI as a threat to traditional knowledge
dissemination, che*CkOVER* illustrates how specialized scientific communities can engage

constructively with emerging technologies to ensure authoritative, inclusive, and future-ready

knowledge systems.

691   Because che*CkOVER* is fully data-driven, all distributional outputs represent a

692   snapshot of the currently available dataset rather than a final assessment, and will update

693   periodically as new records are incorporated, ensuring that the various metrics provided

694   evolve dynamically and naturally with continued data acquisition.

Availability of source code and requirements

Project name: che*CkOVER*

Project homepage: https://world.crayfish.ro/

Source code repository: (link), together with a demonstration dataset (RollData) containing

sample crayfish occurrence data, canonical species geo-narrative template (File S1), and the

offline lookup tables required in the workflow (provided in TSV format to ensure full

Unicode compatibility and to support machine-readable reuse): vernacular names expert

supplement (Table S2), hydronyms (Table S3), and freshwater ecoregions (Table S4)

Operating system(s): cross-platform (Windows 10/11, Linux Ubuntu 20.04+, macOS 11+)

Programming language: R ($\geq$4.5.2)

Other requirements: pandoc ($\geq$2.0), GDAL ($\geq$3.0), PROJ ($\geq$7.0) libraries for spatial operations

License: GPL-3

Any restrictions to use by non-academics: none

RRID: [not yet assigned, all persistent identifiers will be registered upon final release]

bio.tools ID: [not yet assigned]

Local installation requirements: For typical use cases (datasets with <10,000 records per

species): 16 GB RAM, 4 CPU cores, 100 GB disk space. For large-scale applications

(>20,000 records per species or processing hundreds of taxa): 256+ GB RAM, 16+ CPU

cores, 500+ GB disk space recommended.

Server deployment: RESTful API endpoints for species packages and search functionality,

MariaDB for metadata indexing, CDN integration for global accessibility, Docker

containerization supported

Additional files

Supplementary Table S1. World of Crayfish® Contributors, including full names and

institutional affiliations, who provided validated occurrence data, metadata, or region-specific

expertise that supported data verification, contextual interpretation, and the completeness of

species-level information used in this study.

Supplementary Table S2. Expert-curated vernacular names lookup table, supplementing

ITIS-sourced names and providing multilingual coverage.

Supplementary Table S3. Hydronym lookup table, linking HydroBASINS unit IDs (levels 6,

8, and 10) to their assigned hydronyms, derived by spatial intersection of basin polygons with

named river-line datasets.

Supplementary Table S4. Lookup table for freshwater ecoregions, linking FEOW unit IDs to

their standardized ecoregion names.

Supplementary Table S5. Software dependencies used by the che*CkOVER* pipeline. All

packages are automatically installed during the standard workflow execution, except for the

auxiliary datasets listed in Supplementary Tables S2–S4.

Supplementary File S1. Canonical che*CkOVER* species geo-narrative template used for

standardized human- and LLM-readable outputs.

Abbreviations

AI: artificial intelligence; AOO: area of occupancy; API: application programming interface;

CSV: comma-separated values; EOO: extent of occurrence; FAIR: Findable, Accessible,

Interoperable, Reusable; FEOW: Freshwater Ecoregions of the World; IUCN: International

Union for Conservation of Nature; JSON: JavaScript Object Notation; KML/KMZ: Keyhole

Markup Language / Keyhole Markup Language Zipped; LLM: large language model;

MRBW: Major River Basins of the World; RAG: retrieval-augmented generation; TEOW:

Terrestrial Ecoregions of the World; TSV: tab-separated values; WDPA: World Database on

Protected Areas; WoC: World of Crayfish® platform..

Consent for publication

748 This study involves species-occurrence data sourced from community contributors, following

749 the anonymization protocol of the WoC® platform [10], ensuring that no precise coordinates

750 are released in any che*CkOVER* output. All authors, including WoC® Contributors, have

751 reviewed and approved the final manuscript and consent to its publication.

Acknowledgments

754 We thank Marian Neagul (West University of Timișoara) for configuring and maintaining the
755 server environment hosting the dynamic data platform, and Alexandru E. Mizeranschi
756 (Research and Development Station for Bovine, Arad, Romania) for technical advice during
757 pipeline testing.
758

Author contributions

L.P.: conceptualization, investigation, methodology, supervision, validation, writing—original
draft. D.L.: software, methodology, formal analysis, visualization, writing—original draft.
V.I.B.: methodology, visualization, writing—original draft. C.I.N.: methodology,
visualization, writing—original draft. T.T.Ș.: methodology, visualization, writing—original
draft. A.V.L.: methodology, visualization, writing—original draft. WoC® Contributors:
supervision, validation. All: writing—review & editing.
766

767 **Use of AI tools**

768 ChatGPT 5.1 model [74] was used for language refinement of the manuscript. No scientific
769 interpretations were generated by AI. All content was reviewed and fully validated by the
770 authors, who take full responsibility for the final text.
771

772 **Funding**

773 This work received no project-based or grant funding. The Article Processing Charge was
774 supported through doctoral publication funds provided by the West University of Timișoara.
775

Data availability

The processed dataset used in this study has been deposited in GigaDB as a static, citable
snapshot (DOI: to be assigned upon deposition). All raw occurrence records originate from
the WoC® platform (https://world.crayfish.ro), which functions as a dynamic, continuously
updated repository. The platform provides interactive access to current and future versions of
the dataset, as well as to all che*CkOVER*-generated outputs across subsequent releases.
782

783 **Competing interests**

784 The authors declare that they have no competing interests.
785

786 **Ethical approval**

References

1. Oreskes N. Models all the way down. *Metascience 2011 211*. Springer; 2011; doi: 10.1007/S11016-011-9558-9.

2. Gladwell M. The tipping point: How little things can make a big difference. Little, Brown and Company;

3. Swedlow B. Four States of Knowledge: A Cultural Theory of the Co-Production of Science, Social Order, and Scientific, Cultural, and Policy Change. *SSRN Electron J*. Elsevier BV; 2010; doi: 10.2139/SSRN.1673521.

4. Boivin N, Crowther A. Mobilizing the past to shape a better Anthropocene. *Nat Ecol Evol 2021 53*. Nature Publishing Group; 2021; doi: 10.1038/s41559-020-01361-4.

5. Marburger JH. Science, technology and innovation in a 21st century context. *Policy Sci*. Springer; 2011; doi: 10.1007/S11077-011-9137-3/METRICS.

6. Cardoso AW, Hestir EL, Slingsby JA, Forbes CJ, Moncrieff GR, Turner W, et al.. BioSCape combines local knowledge and remote-sensing technology for inclusive biodiversity science. *Nat Rev Biodivers 2025 11*. Nature Publishing Group; 2025; doi: 10.1038/s44358-024-00007-8.

7. Moor M, Banerjee O, Abad ZSH, Krumholz HM, Leskovec J, Topol EJ, et al.. Foundation models for generalist medical artificial intelligence. *Nat 2023 6167956*. Nature Publishing Group; 2023; doi: 10.1038/s41586-023-05881-4.

8. Dwivedi YK, Kshetri N, Hughes L, Slade EL, Jeyaraj A, Kar AK, et al.. "So what if ChatGPT wrote it?" Multidisciplinary perspectives on opportunities, challenges and implications of generative conversational AI for research, practice and policy. *Int J Inf Manage*. Elsevier Limited; 2023; doi: 10.1016/J.IJINFOMGT.2023.102642.

9. Leslie D. Does the sun rise for ChatGPT? Scientific discovery in the age of generative AI. *AI Ethics 2023 54*. Springer; 2023; doi: 10.1007/S43681-023-00315-3.

10. Ion MC, Bloomer CC, Bărăscu TI, Oficialdegui FJ, Shoobs NF, Williams BW, et al.. World of Crayfish^{TM}: a web platform towards real-time global mapping of freshwater crayfish and their pathogens. *PeerJ*. PeerJ Inc.; 2024; doi: 10.7717/PEERJ.18229.

11. Cortés-Guzmán D, Sinclair J, Hof C, Kalusche JB, Haase P. Dispersal, glacial refugia and temperature shape biogeographical patterns in European freshwater biodiversity. *Glob Ecol Biogeogr*. John Wiley & Sons, Ltd; 2024; doi: 10.1111/GEB.13886.

12. Poelen JH, Simons JD, Mungall CJ. Global biotic interactions: An open infrastructure to share and analyze species-interaction datasets. *Ecol Inform*. Elsevier; 2014; doi:

823 10.1016/J.ECOINF.2014.08.005.

824 13. McClanahan TR, Friedlander AM, Chabanet P, Bruggemann JH, Wickel J, Azali MK.

825 Modeling the spatial distribution of numbers of coral reef fish species and community types in

826 the Western Indian Ocean faunal province. *Mar Ecol Prog Ser*. Inter-Research; 2024; doi:

827 10.3354/MEPS14538.

828 14. Rodrigues ASL, Pilgrim JD, Lamoreux JF, Hoffmann M, Brooks TM. The value of the

829 IUCN Red List for conservation. *Trends Ecol Evol*. Elsevier Ltd; 2006; doi:

830 10.1016/j.tree.2005.10.010.

831 15. Uysal Bayar F. Determination of natural populations to be included in breeding program

832 in St. John's wort species (Hypericum perforatum L.). *PeerJ*. PeerJ Inc.; 2024; doi:

833 10.7717/PEERJ.18336/SUPP-1.

834 16. Mace GM. Whose conservation? *Science (80-)*. American Association for the

835 Advancement of Science; 2014; doi: 10.1126/SCIENCE.1254704.

836 17. Bowler DE, Bjorkman AD, Dornelas M, Myers-Smith IH, Navarro LM, Niamir A, et al..

837 Mapping human pressures on biodiversity across the planet uncovers anthropogenic threat

838 complexes. *People Nat*. Blackwell Publishing Ltd; 2020; doi:

839 10.1002/PAN3.10071/SUPPINFO.

840 18. Chaplin-Kramer R, Miller CR, Dee LE, Bennett NJ, Echeverri A, Gould RK, et al..

841 Wildlife's contributions to people. *Nat Rev Biodivers 2025 11*. Nature Publishing Group;

842 2025; doi: 10.1038/s44358-024-00006-9.

843 19. Pereira HM, Martins IS, Rosa IMD, Kim HJ, Leadley P, Popp A, et al.. Global trends and

844 scenarios for terrestrial biodiversity and ecosystem services from 1900 to 2050. *Science (80-*

845 *)*. American Association for the Advancement of Science; 2024; doi:

846 10.1126/SCIENCE.ADN3441/SUPPL_FILE/SCIENCE.ADN3441_MDAR_REPRODUCIBI

847 LITY_CHECKLIST.PDF.

848 20. Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, et al..

849 Biodiversity loss and its impact on humanity. *Nat 2012 4867401*. Nature Publishing Group;

850 2012; doi: 10.1038/nature11148.

851 21. Vinuesa R, Azizpour H, Leite I, Balaam M, Dignum V, Domisch S, et al.. The role of

852 artificial intelligence in achieving the Sustainable Development Goals. *Nat Commun 2020*

853 *111*. Nature Publishing Group; 2020; doi: 10.1038/s41467-019-14108-y.

854 22. Christie AP, Downey H, Frick WF, Grainger M, O'Brien D, Tinsley-Marshall P, et al.. A

855 practical conservation tool to combine diverse types of evidence for transparent evidence-

856 based decision-making. *Conserv Sci Pract*. John Wiley & Sons, Ltd; 2022; doi:

10.1111/CSP2.579.

23. Constantino-Panopio C, De | R M H, Pandithasekera S, Nyandire | N O, Rahman | S C,

Mckee | E, et al.. An evaluation of tools being promoted to help the private sector disclose its

impacts on biodiversity. *Conserv Sci Pract*. John Wiley & Sons, Ltd; 2025; doi:

10.1111/CSP2.70198.

24. Ayoub NF, Lee YJ, Grimm D, Divi V. Head-to-Head Comparison of ChatGPT Versus

Google Search for Medical Knowledge Acquisition. *Otolaryngol Neck Surg*. John Wiley &

Sons, Ltd; 2024; doi: 10.1002/OHN.465.

25. Culina A, Baglioni M, Crowther TW, Visser ME, Woutersen-Windhouwer S, Manghi P.

Navigating the unfolding open data landscape in ecology and evolution. *Nat Ecol Evol*.

Nature Publishing Group; 2018; doi: 10.1038/s41559-017-0458-2.

26. Kung TH, Cheatham M, Medenilla A, Sillos C, De Leon L, Elepaño C, et al.. Performance

of ChatGPT on USMLE: Potential for AI-assisted medical education using large language

models. *PLOS Digit Heal*. Public Library of Science; 2023; doi:

10.1371/JOURNAL.PDIG.0000198.

27. Affinito F, Williams JM, Campbell JE, Londono MC, Gonzalez A. Progress in developing

and operationalizing the Monitoring Framework of the Global Biodiversity Framework. *Nat*

Ecol Evol 2024 812. Nature Publishing Group; 2024; doi: 10.1038/s41559-024-02566-7.

28. Sullivan BL, Aycrigg JL, Barry JH, Bonney RE, Bruns N, Cooper CB, et al.. The eBird

enterprise: An integrated approach to development and application of citizen science. *Biol*

Conserv. Elsevier; 2014; doi: 10.1016/J.BIOCON.2013.11.003.

29. Branco V V., Capinha C, Rocha J, Correia L, Cardoso P. SPECTRE: Standardised Global

Spatial Data on Terrestrial SPecies and ECosystems ThREats. *Glob Ecol Biogeogr*. John

Wiley & Sons, Ltd; 2025; doi: 10.1111/GEB.13949.

30. Callaghan CT, Mesaglio T, Ascher JS, Brooks TM, Cabras AA, Chandler M, et al.. The

benefits of contributing to the citizen science platform iNaturalist as an identifier. *PLOS Biol*.

Public Library of Science; 2022; doi: 10.1371/JOURNAL.PBIO.3001843.

31. Chandler M, See L, Copas K, Bonde AMZ, López BC, Danielsen F, et al.. Contribution of

citizen science towards international biodiversity monitoring. *Biol Conserv*. Elsevier; 2017;

doi: 10.1016/J.BIOCON.2016.09.004.

32. Costello MJ, Michener WK, Gahegan M, Zhang ZQ, Bourne PE. Biodiversity data should

be published, cited, and peer reviewed. *Trends Ecol Evol*. Elsevier; 2013; doi:

10.1016/j.tree.2013.05.002.

33. Amano T, Sutherland WJ. Four barriers to the global understanding of biodiversity

conservation: Wealth, language, geographical location and security. *Proc R Soc B Biol Sci*.

Royal Society; 2013; doi: 10.1098/RSPB.2012.2649.

34. Feng X, Smith AB, Boyle B, Chen X, Enquist BJ, Gallagher R, et al.. The next stage of

biodiversity informatics: community-driven synthesis and integration of biodiversity

databases. *Bioscience*. 2025; doi: 10.1093/BIOSCI/BIAF112.

35. Löffler F, Wesp V, König-Ries B, Klan F. Dataset search in biodiversity research: Do

metadata in data repositories reflect scholarly information needs? *PLoS One*. Public Library

of Science; 2021; doi: 10.1371/JOURNAL.PONE.0246099.

36. Stephens CR, González-Salazar C, Del M, Villalobos-Segura C, Marquet PA. Can

Ecological Interactions be Inferred from Spatial Data? *Biodivers Informatics*. The University

of Kansas; 2020; doi: 10.17161/BI.V15I1.9815.

37. Zyoud S, Zyoud AH. Mapping and visualizing global knowledge on planetary health in

the climate change context: a comprehensive exploration of insights, trends, and research

priorities. *Discov Sustain 2024 51*. Springer; 2024; doi: 10.1007/S43621-024-00499-6.

38. Zavalin V, Zavalina OL. Are we there yet? Evaluation of AI-generated metadata for

online information resources. *Inf Res an Int Electron J*. University of Boras; 2025; doi:

10.47989/IR30ICONF47215.

39. Yang X, Jia J, Zhou X, Wang S. The future of artificial intelligence: Time to embrace

more international collaboration. *Innov*. Elsevier; 2024; doi: 10.1016/J.XINN.2024.100703.

40. Vos RA, Biserkov JV, Balech B, Beard N, Blissett M, Brenninkmeijer C, et al.. Enriched

biodiversity data as a resource and service. *Biodivers Data J 2 e1125*. Pensoft Publishers;

2014; doi: 10.3897/BDJ.2.E1125.

41. Heberling JM, Miller JT, Noesgaard D, Weingart SB, Schigel D. Data integration enables

global biodiversity synthesis. *Proc Natl Acad Sci U S A*. National Academy of Sciences;

2021; doi: 10.1073/PNAS.2018093118/SUPPL_FILE/PNAS.2018093118.SAPP.PDF.

42. Pollock LJ, Kitzes J, Beery S, Gaynor KM, Jarzyna MA, Mac Aodha O, et al.. Harnessing

artificial intelligence to fill global shortfalls in biodiversity knowledge. *Nat Rev Biodivers*

2025. Nature Publishing Group; 2025; doi: 10.1038/s44358-025-00022-3.

43. Stenhouse A, Fisher N, Lepschi B, Schmidt-Lebuhn A, Rodriguez J, Turco F, et al.. A

vision of human–AI collaboration for enhanced biological collection curation and research.

Bioscience. 2025; doi: 10.1093/BIOSCI/BIAF021.

44. Jordão LSB, Morim MP, Baumgratz JFA, Simon MF, Eppinghaus ALC, Calfo VA.

TypeTaxonScript: sugarifying and enhancing data structures in biological systematics and

biodiversity research. *Biol Methods Protoc*. Oxford Academic; 2024; doi:

10.1093/BIOMETHODS/BPAE017.

45. Hegde H, Vendetti J, Goutte-Gattat D, Caufield JH, Graybeal JB, Harris NL, et al.. A

change language for ontologies and knowledge graphs. *Database*. Oxford Academic; 2025;

doi: 10.1093/DATABASE/BAAE133.

46. Wilkinson MD, Dumontier M, Aalbersberg IjJ, Appleton G, Axton M, Baak A, et al.. The

FAIR Guiding Principles for scientific data management and stewardship. *Sci Data 2016 31*.

Nature Publishing Group; 2016; doi: 10.1038/sdata.2016.18.

47. Anthropic: Essential grammar topics for high school graduates [Generative AI chat].

Claude Sonnet 4. https://claude.ai/share/329173b2-ec93-4663-ac68-4f65ea4f166d (2025).

Accessed 2025 Dec 19.

48. Canhos VP, Souza S de, Giovanni R De, Canhos DAL. Global Biodiversity Informatics:

setting the scene for a "new world" of ecological forecasting. *Biodivers Informatics*. The

University of Kansas; 2004; doi: 10.17161/BI.V1I0.3.

49. Guralnick R, Hill A. Biodiversity informatics: automated approaches for documenting

global biodiversity patterns and processes. *Bioinformatics*. Oxford Academic; 2009; doi:

10.1093/BIOINFORMATICS/BTN659.

50. IUCN Standards and Petitions Committee. Guidelines for Using the IUCN Red List

Categories and Criteria. 16th ed. IUCN.

51. Majer J. Saga of the short-range endemic. *Aust J Entomol*. John Wiley & Sons, Ltd; 2009;

doi: 10.1111/J.1440-6055.2009.00715.X.

52. Crandall KA, Buhay JE. Global diversity of crayfish (Astacidae, Cambaridae, and

Parastacidae—Decapoda) in freshwater. *Freshw Anim Divers Assess*. Springer, Dordrecht;

2007; doi: 10.1007/978-1-4020-8259-7_32.

53. Richman NI, Böhm M, Adams SB, Alvarez F, Bergey EA, Bunn JJS, et al.. Multiple

drivers of decline in the global status of freshwater crayfish (Decapoda: Astacidea). *Philos*

Trans R Soc B Biol Sci. Royal Society of London; 2015; doi: 10.1098/rstb.2014.0060.

54. R Core Team. R: A Language and Environment for Statistical Computing. R Core Team.

Viena, Austria;

55. IUCN. IUCN red list categories and criteria. Second edi. Switzerland and Cambridge, UK:

IUCN--The World Conservation Union;

56. Stenseke M, Harris FL, Crowe T. Advice for interdisciplinary biodiversity research.

Conserv Sci Pract. John Wiley & Sons, Ltd; 2025; doi: 10.1111/CSP2.70203.

57. Seebens H, Meyerson LA, Richardson DM, Lenzner B, Tricarico E, Courchamp F, et al..

Biological invasions: a global assessment of geographic distributions, long-term trends, and

data gaps. *Biol Rev*. John Wiley & Sons, Ltd; 2025; doi: 10.1111/BRV.70058.

58. Guareschi S, Cancellario | T, Oficialdegui | F J, Clavero | M. Insights from the past:

Invasion trajectory and niche trends of a global freshwater invader. *Glob Chang Biol*. John

Wiley & Sons, Ltd; 2024; doi: 10.1111/GCB.17059.

59. Freitas H, Gouveia AC. Biodiversity futures: digital approaches to knowledge and

conservation of biological diversity. *Web Ecol*. Copernicus Publications; 2025; doi:

10.5194/WE-25-29-2025.

60. Feng X, Enquist BJ, Park DS, Boyle B, Breshears DD, Gallagher R V., et al.. A review of

the heterogeneous landscape of biodiversity databases: Opportunities and challenges for a

synthesized biodiversity knowledge base. *Glob Ecol Biogeogr*. John Wiley & Sons, Ltd;

2022; doi: 10.1111/GEB.13497.

61. Bisby FA. The Quiet Revolution: Biodiversity Informatics and the Internet. *Science (80-)*.

American Association for the Advancement of Science; 2000; doi:

10.1126/SCIENCE.289.5488.2309.

62. Lim JS, Schneider E, Grover M, Zhang J, Peters D. Effects of AI versus human source

attribution on trust and forgiveness in the identical corporate apology statement for a data

breach scandal. *Public Relat Rev*. JAI; 2025; doi: 10.1016/J.PUBREV.2024.102520.

63. Sahebi S, Formosa P. The AI-mediated communication dilemma: epistemic trust, social

media, and the challenge of generative artificial intelligence. *Synthese*. Springer Science and

Business Media B.V.; 2025; doi: 10.1007/S11229-025-04963-2/METRICS.

64. Hohenstein J, Jung M. AI as a moral crumple zone: The effects of AI-mediated

communication on attribution and trust. *Comput Human Behav*. Pergamon; 2020; doi:

10.1016/J.CHB.2019.106190.

65. Guo Q, Zhen J, Wu F, He Y, Qiao C. Can Students Make STEM Progress With the Large

Language Models (LLMs)? An Empirical Study of LLMs Integration Within Middle School

Science and Engineering Practice. *J Educ Comput Res*. SAGE Publications Inc.; 2025; doi:

10.1177/07356331241312365/SUPPL_FILE/SJ-PDF-1-JEC-

10.1177_07356331241312365.PDF.

66. Bewersdorff A, Hartmann C, Hornberger M, Seßler K, Bannert M, Kasneci E, et al..

Taking the next step with generative artificial intelligence: The transformative role of

multimodal large language models in science education. *Learn Individ Differ*. Elsevier Ltd;

2025; doi: 10.1016/j.lindif.2024.102601.

67. Suazo-Galdames IC, Saracostti M, Chaple-Gil AM. Scientific evidence and public policy:

a systematic review of barriers and enablers for evidence-informed decision-making. *Front*

Commun. Frontiers Media SA; 2025; doi: 10.3389/FCOMM.2025.1632305/XML.

68. Ulibarri N. Collaborative model development increases trust in and use of scientific

information in environmental decision-making. *Environ Sci Policy*. Elsevier; 2018; doi:

10.1016/J.ENVSCI.2018.01.022.

69. Kreuzberger D, Kuhl N, Hirschl S. Machine Learning Operations (MLOps): Overview,

Definition, and Architecture. *IEEE Access*. Institute of Electrical and Electronics Engineers

Inc.; 2023; doi: 10.1109/ACCESS.2023.3262138.

70. Otegui J, Guralnick RP. The geospatial data quality REST API for primary biodiversity

data. *Bioinformatics*. Oxford Academic; 2016; doi: 10.1093/BIOINFORMATICS/BTW057.

71. Cham H, Malek S, Milow P, Song C. Developing an ecological visualization system for

biodiversity data. *All Life*. Taylor & Francis; 2022; doi: 10.1080/26895293.2022.2066195.

72. Sinclair JS, Stubbington R, Welti EAR, Aroviita J, Baker NJ, Cañedo-Argüelles M, et al..

Current protected areas provide limited benefits for European river biodiversity. *Nat Commun*

2025 161. Nature Publishing Group; 2025; doi: 10.1038/s41467-025-67125-5.

73. Ion MC, Ács A-R, Laza AV, Lorincz I, Livadariu D, Lamoly AM, et al.. Conservation

status of the idle crayfish Austropotamobius bihariensis Pârvulescu, 2019. *Glob Ecol*

Conserv. Elsevier; 2024; doi: 10.1016/J.GECCO.2024.E02847.

74. OpenAI. High school grammar concepts [Generative AI chat]. ChatGPT.

1011