Friendship modulates physical effort and neural dynamics during group coordination in shared social networks

Qianliang Li[1*], Aliaksandr Dabranau[1], Kyveli Kompatsiari[1] and Ivana Konvalinka[1]

[1]Section for Cognitive Systems, DTU Compute, Technical University of Denmark, Kongens Lyngby, Denmark,

[*]Corresponding author. Email: glia@dtu.dk

ABSTRACT

Human social interactions are embedded within social networks, yet it remains unclear how existing interpersonal relationships influence the behavioural and neural dynamics of collective action. Here, we combined social network mapping with three-person EEG hyperscanning to investigate how friendships in shared networks modulate interpersonal coordination. We mapped social networks across seven undergraduate cohorts (N = 378) and recruited a subset of participants (N = 93) that consisted of triads of two mutually close friends and one non-friend from the same network. Triads performed a novel joint motor task in which participants produced forces towards a collective target. Non-friends consistently produced greater and more vigorous force than friends during joint action. This effect was strongest for non-friends embedded in smaller social networks. At the neural level, non-friends exhibited stronger alpha- and beta-band event-related desynchronization, higher inter-subject correlation, and increased inter-brain group phase synchrony. These neural signatures are consistent with increased action preparation, prediction, and attentional engagement during coordination. Taken together, these findings indicate that friendship and social network context are reflected in how individuals allocate physical effort and engage during collective action, and suggest that non-friends work harder and allocate more attention during joint tasks to compensate for the disparity in social standing. By linking real-world social relationships to behavioural and multi-brain neural dynamics, this work demonstrates that relational context is embodied in the mechanisms supporting human group coordination.

Keywords: Friendship, social networks, group coordination, joint action, EEG hyperscanning

1 Introduction

Friendship is one of the most powerful social ties humans form, shaping well-being, identity, and opportunities across the lifespan (Hartup and Stevens, 1997; van der Horst and Coffé, 2012; Dunbar, 2018; Alsarrani et al., 2022; Pezirkianidis et al., 2023). It forms the foundation of our social networks, which provide the broader societal context within which close relationships emerge and evolve. Yet despite its prominent role in social life, we know remarkably little about how friendship influences real-time moment-to-moment processes through which people interact and coordinate with one another.

Understanding this influence is crucial because coordination underlies many everyday behaviours - from moving furniture and cooking together, to group decision making at work and team sports. Such coordinated actions rely not only on cognitive mechanisms, such as the integration of sensory information from own and others' actions and the anticipation and adaptation to others' behaviour to allocate effort across shared tasks in real time (Sebanz et al., 2006; Knoblich et al., 2011; Vesper et al., 2017; Sebanz and Knoblich, 2021). To perform these actions, people also incorporate information about the relational structure of the group (Karau and Williams, 1993; Tomasello, 2014; Clark-Polner and Clark, 2014; Pan et al., 2017; Chierchia et al., 2020; Speer et al., 2024). Social identity theory illustrates this influence by positing that group membership becomes part of the self-concept and biases individuals toward in-group over out-group members (Tajfel and Turner, 1979). This raises the question of how high-level social relationships, such as friendship and group membership within shared so-

cial networks, modulate behavioural and neural coordination dynamics, particularly in interactions involving more than two people.

Competing predictions can be made regarding how friendship should influence effort allocation during collective action. On the one hand, close relationships and social familiarity may promote prosocial motivation, trust, and commitment to collective outcomes (Larzelere and Huston, 1980; Rempel et al., 1985; Karau and Williams, 1993; Van Lange et al., 1997; Wieselquist et al., 1999; Alarcon et al., 2016; Chierchia et al., 2020). From this perspective, friends may be expected to contribute more effort during coordination, reflecting greater concern for shared success or willingness to support one another. On the other hand, interactions with non-friends or out-group members may heighten arousal, vigilance, and sensitivity to evaluation (Cottrell and Neuberg, 2005; Holt-Lunstad et al., 2007), as well as motivation to signal competence or value to the group (Zajonc, 1965; Tetlock and Manstead, 1985; Leary and Kowalski, 1990; Williams and Karau, 1991; Baumeister and Leary, 1995; Karau and Wilhau, 2020). In this case, non-friends may contribute more effort during joint tasks, particularly when their role or standing within the group is uncertain.

Distinguishing between these possibilities requires methods that measure both how groups coordinate actions together and the neural mechanisms that are engaged during real-time interaction. Recent advances in social neuroscience and joint action have highlighted the need for interactive paradigms and multi-person neuroimaging for studying such processes (De Jaegher et al., 2010; Dumas, 2011; Konvalinka and Roepstorff, 2012; Schilbach et al., 2013; Dingemanse et al., 2023). Dyadic joint action studies show that successful coordination relies on real-time prediction of others' movements, the ability to generate predictable actions oneself - particularly when sensory access to partners is limited - and continuous mutual adaptation that allows individuals to maintain temporal alignment (Knoblich and Jordan, 2003; Keller et al., 2007; Konvalinka et al., 2010; Noy et al., 2011; Vesper et al., 2011; Pecenka and Keller, 2011; Keller et al., 2014; Vesper et al., 2017; Pezzulo et al., 2019; Konvalinka et al., 2023b). The neural mechanisms supporting these processes are more poorly understood, but interactive EEG studies demonstrate that sensorimotor coupling during coordination is accompa-

nied by enhanced mu-, alpha- and beta-band desynchronization (Tognoli et al., 2007; Perry et al., 2011; Dumas et al., 2012; Naeem et al., 2012; Konvalinka et al., 2014; Zamm et al., 2021; Zimmermann et al., 2022; Flösch et al., 2023), consistent with mechanisms of action preparation and execution, motor prediction, and increased attentional allocation to others' behaviour.

Multi-brain approaches have shown that people also synchronize neural rhythms with each other during action coordination (Dumas et al., 2010; Müller et al., 2013; Zamm et al., 2018; Fishburn et al., 2018; Gugnowska et al., 2022). Beyond oscillatory markers, inter-subject correlation (ISC) methods have shown that when people attend to the same visual or auditory stimulus, even in the absence of interaction, their neural signals become correlated as a result of common stimulus locking (Dmochowski et al., 2012; Cohen and Parra, 2016; Nastase et al., 2019; Madsen and Parra, 2022). Building on this principle, correlated components analysis (CorrCA) extracts shared neural components that are maximally aligned across individuals viewing or listening to a common stimulus (Parra et al., 2019). ISC has been shown to increase during periods of joint attention, and decrease with distraction (Golland et al., 2015; Pérez et al., 2021). It has thus been purposed to serve as a marker of joint attentional engagement, with higher correlation reflecting more consistent, time-locked processing of the task across individuals (Hasson et al., 2004; Dmochowski et al., 2012, 2014; Cohen et al., 2018; Nastase et al., 2019; Rai et al., 2025; He et al., 2025).

Despite this progress, relational context has rarely been incorporated into interactive neuroscience paradigms (Clark-Polner and Clark, 2014), as most joint action studies recruit strangers or pair participants with confederates (Clark-Polner and Clark, 2014), leaving open how naturally occurring social relationships influence coordination. Recent work in social network neuroscience has shown that people neurally encode properties of their social networks and that friends exhibit more similar neural responses during passive tasks (Schmälzle et al., 2017; Parkinson et al., 2018; Hyon et al., 2020a,b; D'Onofrio et al., 2022; Baek et al., 2023; Sijtsma et al., 2023; Speer et al., 2024; Shen et al., 2025). However, no studies have examined how friendship modulate interaction and brain dynamics during real-time interaction.

Here we address this gap by combining social network

mapping with three-person EEG hyperscanning to test how friendship dynamics within shared social networks influence physical effort allocation and neural engagement during group coordination. We mapped social networks across seven bachelor program study lines (Figure 1A), and recruited triads composed of two mutually close friends and one non-friend from the same study line networks. These triads engaged in a novel three-person coordination task (Figure 1B) - the "Force Game" - in which they were instructed to jointly coordinate their forces to reach specific target forces by applying pressure to individual pressure measuring devices, while receiving continuous or non-continuos visual feedback of joint - but not individual - contributions (Figure 1). This design allowed us to test competing hypotheses: whether friends contribute more effort due to prosocial motivation and relational commitment, or whether non-friends contribute more effort due to heightened arousal, evaluation sensitivity, or uncertainty about their role in the group. We show that non-friends consistently contributed greater physical force and produced it more vigorously than friends during joint tasks. This effect was strongest for non-friends from smaller class social networks, indicating that individuals within smaller networks may be more sensitive to how they are perceived in group settings with others from the same network and engage more effortfully during coordination. Non-friends also exhibited greater alpha and beta event-related desynchronization, higher neural inter-subject correlations, and stronger neural group phase synchrony, suggesting increased motivation, attention, and engagement. By combining relative friendship information from shared social networks, behavioural coordination, and multi-brain imaging, we provide evidence that relational context is embodied in the cognitive and motivational processes that support human group coordination.

2 Results

2.1 Social network mapping

We first mapped the social networks of seven bachelor program study lines, with class sizes ranging from 43 to 184 ($M = 83.14; SD = 44.26$). Students were asked to complete a questionnaire in which they checked off the individuals they considered to be friends or close friends (see Methods for definitions) from a roster of all of their classmates in the same study line. 378 students gave informed consent

to have their social networks mapped across the seven social networks (Figure 1A), ranging from 28 to 113 ($M = 54.0; SD = 27.09$) students per study line class.

A subset of the students ($N = 93$) were recruited as triads, consisting of two mutually reported close friends and one non-friend (defined as mutual lack of social ties to the friend-pair), to participate in the Force Game experiment (Figure 1B). These participants filled out an additional friendship questionnaire, where we asked participants to list their friends and close friends outside of the study line class social networks, to obtain a more complete estimate of the number of friends for each participant. Overall, the non-friends had significantly fewer friends (Supplementary Figure S1; FDR-corrected p-values for class close friends: $p < 0.01$; class friends $p < 0.05$; close friends and friends outside classroom $p < 0.05$).

2.2 Non-friends contribute greater and more vigorous forces compared to friends in group coordination

Participants were instructed to apply forces by pressing on individual pressure measuring devices in order to individually or jointly reach specific target forces visually shown on their individual screens at the start of each trial (Figure 1C), whilst three-person hyperscanning EEG was measured. Participants either performed the task alone (solo) or interactively. In interactive trials, they had to reach the target force together in triads (referred to as triadic trials), or in dyads where one person's device did not contribute to the total force (dyadic trials), a manipulation they were not aware of. Additionally, we manipulated whether the interaction recruited adaptive or predictive mechanisms, by either giving continuous visual feedback - displayed as a filled circle on the screen - or no feedback across the 4-second trial followed by discrete feedback of the reached force at the end of the trial. The feedback given in interactive trials was always of the joint force, hence individual forces had to be inferred. Each condition was repeated 30 times with varying target forces, represented by the radius of the target circle. An example of the trial averaged produced forces across the 4-second interaction from one triad can be seen in Figure 1D, depicting group interaction with continuous feedback.

To validate the Force Game paradigm, we estimated how well the participants performed the task. As expected, participants performed better with continuous visual feedback

Fig. 1. **Schematic of the Force Game setup.** A) Social networks of seven bachelor program study lines were mapped. B) Triads, consisting of two close friends and one non-friend from the same study line, performed the force game together. C) Timeline of the experimental paradigm. The blue arrows beneath the timeline indicate the repeated loops corresponding to different force levels and conditions. D) Example of the averaged forces across the 4-seconds triadic trials with continuous visual feedback from one triad. E-F) Behavioural performance, estimated with success rate, latency and stability around the target force, for trials E) without continuous visual feedback and F) with continuous feedback.

and performance deteriorated as more people contributed, i.e., with the best performance observed in solo trials, and worst in triadic ones (Figure 1E-F), likely reflecting greater perceptual demands placed on each participant when integrating the collective joint force.

To test whether different coordination mechanisms emerged between friends and non-friends, we first investigated how much force each participant contributed to the joint task. Here, a systematic asymmetry emerged: non-friends contributed greater physical force than friends (Figure 2), during both dyadic and triadic coordination, and both in interac-

tions with (FDR-corrected $p < 0.05$) and without continuous visual feedback (FDR-corrected $p < 0.01$).

We also performed a more temporally fine-grained analysis of the force dynamics, to better characterize the social timing during the task (Konvalinka et al., 2023a). The 1st derivative of the behavioural force time series, also commonly known as rate of force development (RFD) (Maffiuletti et al., 2016)) was estimated (Supplementary Figure S2). The peak RFD for each trial was quantified and averaged over trials (Figure 3), as a proxy of movement vigour (Marinovic et al., 2017). Non-friends had higher peak RFD in interactive

Fig. 2. **Non-friends produced greater physical force as compared to friends.** Non-friends contributed greater physical force to reach the targets in both the dyadic and triadic joint tasks, with and without continuous visual feedback. The striped line corresponds to equal force contribution from all participants during a trial. $^*p < 0.05$, $^{**}p < 0.01$. P-values were derived using non-parametric permutation tests with FDR multiple test correction.

dyadic and triadic trials with and without continuous visual feedback (FDR-corrected $p < 0.05$). No effect of friendship on peak RFD was observed in solo trials.

We did not find differences in performance between friends and non-friends. Friends and non-friends performed equally well in solo tasks (Supplementary Figure S3), and performance in dyadic friend – friend interactions were also on par with friend – non-friend interactions (Supplementary Figure S4).

2.3 Force asymmetry was primarily observed in smaller study line classes

We hypothesized that the force asymmetry between friends and non-friends may be explained by social disparity, and that non-friends from smaller classes or with fewer friends may feel more pressure to prove themselves, or be more sensitive to how they are perceived, hence contributing more force. Hence we investigated the effect of study line class size and

number of friends on the relative amount of force produced.

Strikingly, a clear correlation between the size of the study line classes and amount of relative force was observed, with greater force asymmetry observed in smaller classes. Non-friends from smaller classes contributed more force, which was negatively correlated with class size, such that the smaller the class size the non-friends came from the more force they contributed relative to the friends. This effect was significant for both dyadic and triadic interactions with continuous visual feedback (Figure 4; FDR corrected p-values for Dyadic: Friends $p = 0.003$, Non-friend $p = 0.002$; Triadic: Friends $p = 0.09$, Non-friend $p = 0.001$). While the same trend was observed for interactions without continuous feedback, it was not significant. Similar negative correlations in non-friends were observed when correlating RFD with class sizes, although the correlations did not survive multiple test correction (Supplementary Figure S5; lowest uncorrected p-values for non-friends with continuous visual feedback: $p = 0.038$;

Fig. 3. **Non-friends produced force more vigorously during interactive joint tasks compared to friends.** Non-friends produced force with more vigour, as evident by higher peak RFD compared to friends in interactive dyadic and triadic trials. $^*p < 0.05$. P-values were derived using non-parametric permutation tests with FDR multiple test correction. RFD: rate of force production.

no feedback: $p = 0.096$; lowest FDR corrected p-values with and without feedback: $p = 0.14$).

We hypothesized that non-friend participants with fewer friends overall may be more likely to contribute more force to the interaction relative to the two friends. We indeed observed a negative trend between the number of close friends outside the class social network and relative force in trials without continuous feedback for non-friends; however, this effect did not survive multiple test correction (Supplementary Figure S6; uncorrected p-values for non-friends in Dyadic interactions: $p = 0.02$, Triadic: $p = 0.01$; FDR corrected p-values for Dyadic and Triadic: $p = 0.17$).

2.4 Non-friends exhibit increased alpha and beta event-related desynchronization

To investigate the neural mechanisms underlying social coordination, we analysed evoked alpha and beta event related synchronization, EEG markers of sensorimotor prediction, preparation, execution, and attention. The Force Game was designed with 30 repeated trials for each condition, which

enabled us to increase the signal-to-noise ratio by averaging over trials to perform event related power analysis across classical EEG frequency bands (Supplementary Figure S7).

Across all conditions, we observed the expected sensorimotor alpha and beta event-related desynchronization (ERD) over left-hemispheric electrodes contralateral to the moving hand, which desynchronized shortly after the onset of the visual stimulus (the appearance of the target force) (Figure 5A). This pattern is consistent with classical findings linking alpha and beta desynchronization to motor preparation and execution during voluntary movement (Salmelin and Hari, 1994; Pfurtscheller and Lopes da Silva, 1999; Fujioka et al., 2012; Ross et al., 2017; Zamm et al., 2021).

We tested whether these neural responses differed as a function of friendship, i.e., whether non-friends exhibited larger desynchronization during interactive trials. When contrasting friends and non-friends, non-friends exhibited greater ERDs across all frequency bands, illustrated for alpha in Figure 5B. To account for multiple comparisons across time, space (electrodes), and frequency, we conducted

Fig. 4. **Greater force asymmetry was observed in smaller study line classes.** The effect of greater physical force produced by non-friends, and corresponding lower force by friends, was primarily observed for students recruited from smaller study line classes. The striped line corresponds to equal force contribution. Spearman correlation coefficients were computed and the p-values shown have been FDR multiple test corrected.

non-parametric cluster-based permutation tests (Maris and Oostenveld, 2007; Pernet et al., 2015; Sassenhagen and Draschkow, 2019). In trials with continuous visual feedback, non-friends had significantly greater ERDs across solo ($p = 0.006$), dyadic ($p = 0.033$) and triadic ($p = 0.01$) conditions (Supplementary Figure S8D-F). The clusters consistently spanned approximately 1.1s to 1.7s post-stimulus onset and extended across delta, theta, alpha and beta frequency ranges, across frontal, lateral-central and posterior regions. No significant differences in ERDs were observed between friends and non-friends in trials without continuous visual feedback (Supplementary Figure S8A-C).

Alpha and beta oscillatory activity have been commonly linked to top-down modulation of sensorimotor processing and are thought to play an important role in regulating attention, as they are often modulated by task demand and cognitive effort (Klimesch, 1999; Pfurtscheller and Lopes da Silva, 1999; Mathewson et al., 2012; Keehn et al., 2017; Stecher et al., 2025). Consistent with this interpretation, we observed

sustained alpha and beta ERD in trials with continuous visual feedback, whereas power rebounded toward the baseline approximately 2s after trial onset in trials without continuous feedback, where participants made an initial prediction of the required force but in the absence of feedback did not need to adjust this prediction across the trial (Figure 5A). The sustained ERD may thus reflect ongoing attentional demands required to continuously monitor and adapt to visual feedback, consistent with previous findings reporting greater alpha ERD when dyads have visual contact and hence feedback of each other's movements (Tognoli et al., 2007; Zimmermann et al., 2022). The increased sustained ERD during continuous visual feedback trials was highly significant across the whole brain in solo, dyadic and triadic conditions (Supplementary Figure S9; $p \leq 0.001$), with clusters consistently spanning approximately 0.7s to 3s in theta, alpha and beta frequency ranges.

Finally, we examined whether ERDs differed between interactive and non-interactive conditions. Alpha and beta ERD

were stronger during interactive trials compared to solo trials (Supplementary Figure S10; No Feedback Dyadic: $p = 0.016$, Triadic: $p = 0.51$; With Feedback Dyadic: $p = 0.002$, Triadic: $p = 0.017$), consistent with previous work reporting greater frontal and sensorimotor alpha suppression during social interaction (Konvalinka et al., 2014; Zimmermann et al., 2022). One potential explanation is that interaction increases attentional demands by requiring participants to predict and monitor others' actions in addition to their own (Demos and Palmer, 2023). Taken together, these results indicate that non-friends allocate more attention to the task, particularly when continuous feedback requires ongoing monitoring and adaptation.

2.5 Non-friends exhibit increased neural inter-subject correlations

To further validate whether non-friends were more engaged during the task, we analyzed inter-subject neural alignment using correlated components analysis (CorrCA, Parra et al. (2019)). CorrCA extracts neural components that are maximally similar across multiple individuals and enables computation of inter-subject correlation (ISC) during stimulus-locked processing, which has been proposed as a marker of attentional engagement (Dmochowski et al., 2012, 2014; Poulsen et al., 2017).

Applying CorrCA to event-related alpha and beta power revealed three components with ISC values greater than chance level in both frequency bands (Figure 6A for alpha power and Supplementary Figure S11A for beta). For the leading correlated component, i.e. the component with the greatest ISC across participants, non-friends exhibited greater ISC compared to friends in trials with continuous feedback across both alpha (Figure 6B; FDR corrected p-values for Solo: $p = 0.014$, Dyadic: $p = 0.015$, Triadic: $p = 0.001$) and beta frequency bands (Supplementary Figure S11B; FDR corrected p-values for Solo: $p = 0.001$, Dyadic: $p = 0.001$, Triadic: $p = 0.002$). The associated topographic distribution for the leading components showed global activation across all channels, with a slightly higher weights across lateral visual areas.

ISC was also higher for non-friends compared to friends in the second alpha correlated component in trials with continuous feedback (Supplementary Figure S12B; FDR corrected p-values for Solo: $p = 0.001$, Dyadic: $p = 0.003$,

Triadic: $p = 0.001$). This component exhibited a topographic distribution over sensorimotor areas (C2 in Figure 6A)), indicating enhanced alignment of sensorimotor-related neural activity across non-friends.

To exclude the possibility that higher ISC in non-friends was trivially driven by larger ERD magnitudes, we repeated the CorrCA analysis on individual-level standardized event related power. The friendship related ISC differences persisted under this normalization, primarily during trials with continuous visual feedback (Supplementary Figure S13), indicating that increased ISC reflects greater temporal alignment rather than differences in signal amplitude.

Finally, we assessed inter-subject neural alignment using a cluster phase-based inter-subject group synchrony metric (Richardson et al., 2012). This analysis revealed greater alpha and beta group phase synchrony in non-friends compared to friends (Supplementary Figures S14 and S15), indicating that non-friends had greater phase-locked neural dynamics.

Taken together, these analyses show increased neural ISC and group phase-synchrony in non-friends compared to friends, indicating greater stimulus-locked neural responses. Based on previous experimental and theoretical research (Hasson et al., 2004; Dmochowski et al., 2012, 2014; Cohen et al., 2018; Rai et al., 2025), this suggests greater attentional engagement in non-friends during the Force Game.

3 Discussion

Human interaction depends on the ability to coordinate actions while simultaneously inferring others' intentions, expectations, and levels of commitment (Sebanz et al., 2006; Michael et al., 2016; Vesper et al., 2017; Sebanz and Knoblich, 2021). Here we investigated how friendship versus non-friendship within real social networks modulate the behavioural and neural dynamics underlying group motor coordination. Across triads consisting of two close friends and one non-friend, non-friends consistently produced greater physical force and more vigorous force generation than friends. In addition, they exhibited greater alpha and beta ERD and higher neural ISC than friends - neural markers of attention and engagement. These effects were not explained by differences in maximal force capacity (Supplementary Figure S16) and baseline motor behaviour (i.e., RFD differences during solo conditions). Moreover, the greater alpha

Fig. 5. **Non-friends exhibit increased alpha/beta event related desynchronization.** A) Example plots of event related power from electrodes above left sensorimotor area in alpha and beta frequency bands, color-coded based on the different conditions of group sizes and feedback. B) Time averaged event related alpha power (0 to 4 seconds) from the left sensorimotor area, plotted separately for friends and non-friends. wFb: with feedback; nFb: no feedback, ERD: event-related desynchronization.

and beta ERD in non-friends could not be explained by the greater force produced, as ERD magnitude and amount of produced force were not correlated (Supplementary Figures S17 & S18). This indicates that the behavioural and neural differences between friends and non-friends are not related to basic motor or electrophysiological differences, but rather reflect higher level processes.

One potential explanation for the observed differences between friends and non-friends could be their difference in social identities, with the two friends likely perceiving themselves as an in-group, while the non-friends might perceive themselves as out-group members (Tajfel and Turner, 1979). To compensate for this social disparity, the non-friends might contribute more physical effort to the group task to increase social standing, rapport and affiliation (Krishna and Götz, 2024). An alternative explanation could be that the friends are more likely to distribute the work as they are part of the in-group, which could lead to social loafing (Karau and Williams, 1993), whereas the non-friend might have less trust and greater uncertainty about the group dynamics,

Fig. 6. **Non-friends exhibit increased inter-subject correlations.** A) Three correlated components fitted to alpha event related power had ISC above chance levels ($p < 0.05$ estimated with time-shifted/circular shuffled surrogate data). B) For the leading component C1, which has the highest ISC, non-friends exhibited greater ISC compared to friends. The topographic maps correspond to projections of the forward model weights, where the polarity is normalized so the Cz electrode weight is positive. ISC: inter-subject correlation.

which could result in an overcompensation of physical force (Williams and Karau, 1991). Being an out-group member could also induce feelings of ostracism, leading to increased arousal and attention, thus prompting the non-friends to over-contribute to the group task in order to increase their sense of belonging (Baumeister and Leary, 1995; Williams et al., 2000).

Interestingly, force asymmetry was strongest among individuals drawn from smaller class networks, whereas no differences were observed between friends and non-friends from larger classes. This pattern suggests that the effect is primarily driven by over-contribution from non-friends, rather than under-contribution from friends. One possible interpretation is that non-friends embedded in smaller, more tightly knit social networks may experience heightened sensitivity to how they are perceived by their peers from the same network, leading them to invest more effort during joint tasks (Zajonc, 1965; Tetlock and Manstead, 1985; Baumeister and Leary, 1995; Karau and Wilhau, 2020). In this context, increased force production may function as a behavioural signal of competence or willingness to affiliate. By contrast, in larger networks where interactions are more dispersed, non-

friends may experience less social pressure, or may be less familiar with the other two participants, resulting in reduced behavioural asymmetry relative to friends (Zajonc, 1965; Tetlock and Manstead, 1985; Karau and Williams, 1993; Baumeister and Leary, 1995; Karau and Wilhau, 2020). We additionally observed a trend indicating that non-friends with fewer close friends outside the class network tended to contribute greater relative force; however, this effect did not survive correction for multiple comparisons. While therefore interpreted with caution, this pattern is consistent with the broader account that over-contribution among non-friends may relate to increased motivation or arousal to demonstrate value in social contexts where belonging is less secure (Zajonc, 1965; Tetlock and Manstead, 1985; Baumeister and Leary, 1995; Krishna and Götz, 2024).

The increased motor vigour in non-friends further reinforces this interpretation. Movement vigour has previously been linked to reward, motivational states, and commitment (Desmurget and Turner, 2010; Thura and Cisek, 2017; Marinovic et al., 2017; Shadmehr et al., 2019; Marbaker et al., 2025). Increased vigour in the absence of external incentives therefore indicates a stronger internal motivation to succeed

in the joint task. This aligns with classical accounts of social facilitation, whereby the presence of others increases arousal and promotes enhanced performance on simple, well-learned motor tasks (Zajonc, 1965; Williams and Karau, 1991; Karau and Williams, 1993).

The neural results also suggest that the non-friends were more motivated and engaged. Non-friends showed stronger alpha and beta event-related desynchronization - a proposed marker of attention, motor preparation, execution, and prediction (Klimesch, 1999; Pfurtscheller and Lopes da Silva, 1999; Buschman and Miller, 2007; Klimesch et al., 2007; Klimesch, 2012; Kilavik et al., 2013; Stecher et al., 2025) - and these effects were most pronounced when participants continuously monitored the evolving group force. The alpha and beta ERD was also greater during interactive conditions compared to solo conditions, consistent with previous findings showing that social interaction recruits more motor and attentional resources (Konvalinka et al., 2014; Zimmermann et al., 2022). The non-friends also displayed higher stimulus-locked inter-subject correlation, which has been associated with engagement (Dmochowski et al., 2012, 2014; Cohen and Parra, 2016; Parra et al., 2019; Lambrechts et al., 2025), and greater stimulus-evoked group cluster phase synchrony (Richardson et al., 2012). The increased ISC and group phase synchrony among non-friends thus indicate more similar neural processing during the task, associated with higher attentional engagement with the feedback signal than observed among friends.

While the behavioural effects of friendship were observed for both feedback conditions, neural differences emerged exclusively in the continuous visual feedback condition. This suggests that relational context modulates neural processing primarily during periods that require fine-grained sensorimotor adaptation, as indicated by the ERD clusters contrasting friends and non-friends, which spanned 1-2 seconds corresponding to the period following the initial force ramp-up during which participants could actively fine-tune their force contributions around the target. This adaptive phase was absent in trials without continuous visual feedback, where participants had to rely on internal predictive models rather than ongoing correction in response to sensory feedback, hence the differences in task demands might mask the effect of friendship on the neural activity.

At first glance, our findings of greater neural similarity among non-friends may appear inconsistent with recent social network neuroscience findings showing that friends tend to exhibit more similar neural responses during passive video watching (Parkinson et al., 2018; Hyon et al., 2020b,a; D'Onofrio et al., 2022; Shen et al., 2025). However, the interactive nature of the current task differentiates it sharply from prior paradigms. Here, neural similarity is driven by task-evoked, stimulus-locked processing rather than intrinsic similarity in resting-state activity or perception during passive viewing. In this setting, stronger ISC in non-friends reflects greater alignment to the active task itself, rather than stable similarity in neural representations.

In general, it is important to bear in mind the experimental context in which the findings were generated, i.e. the cooperative force production task in this study. Although our paradigm offers a controlled means of quantifying joint motor coordination, it captures only one facet of social group behaviour, and the effects and dynamics of friendship may differ in tasks involving other social, cognitive, or motivational demands (Zajonc, 1965; Karau and Williams, 1993). Another limitation of the current study is that the social network analyses were limited to self-reported outdegree and class network sizes, due a high degree of incomplete network coverage. On average 66.4% of the students across all seven study lines reported their connections, hence reliable estimation of network parameters such as eigenvector centrality, density, clustering, and assortativity was precluded (Kossinets, 2006; Smith and Moody, 2013; Marks et al., 2013). The unique triad composition of two close friends and one non-friend from a shared class social network also posed recruitment challenges which limited our sample size. Nonetheless, combining network structure, real-world relational context, real-time behavioural coordination, and multi-brain EEG offers a powerful approach to understanding how social dynamics are embodied during interaction.

Overall, our findings demonstrate that differences in social relationship - friend versus non-friend - can systematically bias how individuals allocate physical effort, attentional resources, and neural engagement during group coordination. Non-friends appear to over-contribute, both behaviourally and cognitively, possibly reflecting increased evaluative concern, uncertainty, or a desire to signal value to familiar but

non-friend partners. Friends, by contrast, show behavioural mechanisms consistent with reduced vigilance in relation to non-friends, and potentially, social loafing. These results underscore the importance of incorporating relational context into joint action studies, and provide a neurobehavioural framework for understanding how social structure influences collective performance.

4 Methods

4.1 Social network mapping

378 students, across seven bachelor program study lines (with sizes ranging from 43 to 184 students, $M = 83.14; SD = 44.26$) at the Technical University of Denmark (DTU), gave informed consent to have their social networks in their study line programs mapped. Specifically, 28 to 113 ($M = 54.0; SD = 27.09$) students per study line filled our questionnaire, corresponding to a response rate ranging from 50.6% to 95.7% ($M = 66.4\%; SD = 13.05\%$) of the size of the study line classes.

In practice, we entered the classrooms, gave a brief 5 min presentation about the project and experiments, and asked students to fill out questionnaires in which they checked off the fellow students they considered a friend (defined as *a person whom you like to spend your free time, enjoy informal social activities, such as going for lunch, dinner, drinks, films, etc., and help each other out in a practical way*) (Burt, 1995; Spencer and Pahl, 2006; Parkinson et al., 2018), or close friend (defined as *a person whom you not only help each other and enjoy each other's company but also share personal information and provide emotional support*) (Spencer and Pahl, 2006), or leave it blank to indicate the fellow student was not a friend (Figure 1A). The social network surveys were administered during the students' first (2nd semester) or second (3rd or 4th semester) academic year in their bachelor program, as the courses taken by the students' in the early semesters of their bachelor programs are primarily mandatory courses, hence the students have a high overlap of common courses within each study program.

4.2 Experimental participants

From the social networks, 93 healthy right-handed participants (35 females) between the ages of 18 to 35 years (mean = 21.4, SD = 2.29) were recruited and gave informed consent to participate in the experimental task. Participants arriving in the lab were also asked to fill an additional questionnaire about their friends and close friends outside their study programs, divided into workplace, hobby-related, social media, studies and "other" categories to promote recall, prior to the experiment. Triads consisted of two reciprocal close friends and one non-friend from the same study program. The study was conducted according to the Declaration of Helsinki and was approved by the Scientific Ethics Committee for the Capital Region of Denmark (H-23013244). Participants in the experimental task were financially compensated for their participation.

4.3 The Force Game paradigm

The three participants sat facing away from each other, inside an electrically shielded room, in a triangular formation with 50 cm between each pair of chairs (Figure 1B). Each participant had a screen, placed 60 cm in front of the center of each chair, that displayed the experiment (Figure 1C). Each participant was equipped with a custom-made 3D printed pressure measuring (force) device, which we calibrated using a set of weights (295g, 500g, 1000g, 2000g and 5000g) to estimate the forces in Newtons (Supplementary Figure S19). The participants were instructed to hold the force devices with their right hand and press with their right thumbs to reach a target force, either individually or jointly. The screens would display the force (either solo or joint) as an expanding white circle (the bigger the force, the larger the circle) and the target force as a hollow blue circle. When the produced force was within ± 2N of the target force, the whole circle would turn blue indicating the participants successfully arrived at the target. The participants were explicitly instructed to keep the force at the target for the whole duration of the 4 second trial, following reaching the target. The radius of the displayed force and target force was based on the log-transformed forces in Newtons, to be visually sensitive to smaller forces. Practice trials were also performed prior to the experiment, to familiarize the participants with the devices, where the participants were instructed to freely press, press as hard as they could and reaching the target force individually or jointly with and without continuous visual feedback. Preliminary pilot runs estimated 50N as a reasonable max force and in order to enable all participants to be able to reach the target and taking fatigue into account, we set the target forces to vary as 8N, 10N, 12N, 14N or 16N (corresponding to around 15 to 30%

of max force production). The averaged max force during the practice trials from all 93 participants was 49.59N, confirming our pilot results.

The experiment consisted of 300 trials, across 10 conditions with varying group sizes (solo, duo and trio) with and without continuous visual feedback. There were three duo tasks, i.e. P1 and P2, P1 and P3 or P2 and P3 interacting while the last participant's force contribution was excluded from the joint force, a manipulation they were not aware of. In group trials, only the averaged collective joint force was shown, and the contribution from each participant was $1/2$ or $1/3$ the actual individually measured force in the duo and trio trials respectively, to take into account the increase in group size. Additionally, we manipulated whether the interaction recruited adaptive or predictive mechanisms, by either giving them continuous visual feedback, or no feedback across the 4-second trial, followed by discrete feedback of the reached force at the end of the trial. During the experiment, each condition was performed in blocks of five trials, corresponding to the five target forces (in randomized order), followed by a self-paced break. Prior to each 4 second trial, a fixation cross was presented on the screen (jittered 1 to 2 seconds duration, randomized for each trial to prevent EEG synchronization due to the predictability of stimulus onset timing). Prior to each block, a condition cue was presented with the text *Individual Task* or *Group Task* followed by a green emoji with a single person or an orange emoji with three people, and the text *Continuous Feedback* or *No Continuous Feedback*. No distinction in condition cue were given for duo or trio trials, and no explicit information about who was excluded in duo trials was provided. For trials with no continuous feedback, only the hollow blue circle corresponding to the target force was shown during the 4 second trial, followed by a static image of the joint force at the end of the trial, shown for 1 second. All 10 conditions were also administered in blocks (in randomized order), repeated six times (Figure 1C), to avoid the same conditions being consecutively performed (on top of the 5 repeated trials). Longer forced breaks of 1 min were also given once 100 or 200 trials had been completed to ensure the participants relaxed their thumbs. While performing the force game, simultaneous EEG was recorded. The experiment was designed using Psychopy (Peirce et al., 2019) and took around 45 min, and the whole procedure including EEG

setup and questionnaires took around 1 hour and 30 min.

4.4 Data acquisition

The custom-made force devices send a constant voltage to the Arduino, which converts the analogue voltage signal to a digital 8 bits signal and transfers it to the stimuli PC. The force devices have a pressure-sensitive button, which changes the resistance when pressed and thus the voltage will change accordingly. Based on the calibrations with constant weights, we observed measurement errors in the range of 0.1N. Each force device sends a continuous stream of numbers (more than 1000/sec), and to synchronize the measurements we stored the force measurements corresponding to when the screens updated the visual display. This resulted in the behavioural force data being acquired at 25 Hz. Despite having 60 Hz monitors, it took computational time to visually create the elements and display on the three screens simultaneously from one PC, hence the update frame rate and sampling rate was lower than 60 Hz. Nonetheless, visual inspection of the stimuli confirmed that 25 Hz was enough for the trials to visually seem continuous, and 25 Hz was enough to capture smooth force trajectories.

EEG was recorded using three synchronized, daisy-chained 64-channel Biosemi (Amsterdam, The Netherlands) ActiveTwo EEG set-ups in a 10–20 system, at a sampling frequency of 2048 Hz. EEG caps with 64 channels were positioned such that Cz was centered on the head, at the midpoint between the nasion and inion, and the left and right ear. Conductive gel (SignaGel Electrode Gel, Parker Laboritories, Farfield, NJ) was used to improve signal reception, and electrodes with offsets above $\pm$ 20 mV was adjusted in order to keep them as close to 0 as possible. Any visually bad electrodes at recording was noted down for exclusion during preprocessing.

The data from one participant was excluded from analysis, as it was clear from the behavioural force trajectories that the instructions were not followed (not pressing at all in many trials, resulting in failing to reach the target force in half of solo conditions with feedback, where the average for participants is 98.9%). This means the solo, duo and trio conditions involving the non-compliant participant was discarded, but the solo and duo conditions with the remaining two participants in the triad was kept.

4.5 Behavioural analysis

The behavioural force measurements were linearly interpolated to common timepoints, in fixed 40 ms intervals (25 Hz) starting from trial onset to end of trial at 4 seconds, in order to analyze across trials. A successful trial was defined as a trial where the force trajectory (solo or group) reached within $\pm 2N$ of the target force, and stayed at the target for 200 ms. The accuracy (success rate), latency to success and stability (duration the force stayed around the target) was computed. Relative force was computed as the relative force contribution in group tasks at time of success. The instantaneous rate of force development (RFD), defined as the first derivative of the force time series, was approximated by taking the difference in force between each sample, divided by the time difference: $RFD = \Delta F / \Delta t$. Coefficient of variability was computed 2 s after trial onset, to avoid the build-up of force when approaching the target, while focusing on the part where participants had to stay around the target. The 10 experimental conditions were collapsed into 6 conditions (3 x 2, corresponding to group size x visual feedback), by combining the two duo conditions where each participant were actively contributing to the joint force, and removing the duo condition that each participant was excluded from. The same 6 collapsed conditions were applied for EEG analysis.

4.6 EEG preprocessing

The EEG data were processed using MNE-Python 1.6.1 (Gramfort et al., 2013). First, the data were bandpass filtered at 1–40 Hz (Hamming), downsampled to 500 Hz, and segmented into epochs (-0.5 to 5.5 seconds relative to trial onset). Bad epochs and channels with gross artefacts were rejected by visual inspection. The bad channels were interpolated from adjacent channels using spherical spline interpolation (Perrin et al., 1989) followed by re-referencing to the common average. Ocular, heart and muscle related artefacts were removed using Picard independent component analysis (Ablin et al., 2018) with the number of components set to 32. The automatic detection of muscle-related components in MNE-python were employed to guide the manual determination of muscle artefacts. Autoreject 0.4.3 (Jas et al., 2017) was employed to catch any remaining artefacts by interpolating specific bad channels within each epoch, followed by a guided final manual visual inspection with highlighted bad epochs

determined by the algorithm. Using Autoreject this late in the preprocessing pipeline made it prone to false-positives if the signal was already clean, hence a manual check was performed and no epochs were dropped automatically. On average 3.45 (SD = 5.12) number of epochs were dropped, 1.1 (SD = 2.4) number of channels were interpolated and 15.66 (SD = 4.27) independent components were corrected. The EEG features were estimated in the five canonical frequency bands: delta (1–4 Hz), theta (4–8 Hz), alpha (8–13 Hz), and beta (13–30 Hz).

4.7 Event-related power

The time-frequency representation (TFR) of the epoched EEG data was computed using Morlet wavelets. We used a variable temporal window length (decreased temporal window with increased frequency), to obtain greater temporal resolution at higher frequencies. The power was computed for each frequency band and averaged across epochs for each of the 6 collapsed conditions. Percentage based baseline correction was applied on the averaged time-frequency representation using the pre-stimulus interval (-250 – 0 ms), i.e. on a single-subject level for each condition to avoid positive bias from performing percentage based baseline correction on a single-trial level (Hu et al., 2014).

4.8 Inter-subject correlation

ISC was estimated using correlated component analysis (Parra et al., 2019). Specifically, for each frequency and condition, we applied correlated component analysis on the TFRs, with participants on the first axis to compute stimulus-related power components that maximally correlates between subjects. To determine similar components across group size conditions (but separately for visual feedback), we concatenated the different group conditions along the first axis, in order to make ISC values comparable across group sizes. Importantly, to account for the positive bias induced by higher SNR in the TFRs from the collapsed duo condition (60 trials as opposed to the 30 trials), we performed CorrCA and calculated ISC separately for TFRs computed from each of the two uncollapsed included duo conditions for each participant, followed by averaging after ISC was computed, as a higher SNR condition can arbitrarily increase inter-brain synchrony estimates (Zimmermann et al., 2024). To determine an effect of friendship, ISC was separately computed among friends

and among non-friends, using all-to-all pairwise Pearson's correlations of the extracted components. To ensure the observed difference in ISC was not confounded by a difference in number of subjects for each group (2 friends for each non-friend), we repeated the ISC estimation on 30 randomly sampled friends as opposed to the full sample of friends, and observed no difference in mean ISC for friends. To rule out the increased ISC in non-friends were driven by increased ERD in non-friends, we also repeated correlated component analysis on subject-level standardized TFRs and observed the same results.

4.9 Group cluster phase synchrony

Group cluster phase synchrony was estimated following Richardson et al. (2012). Briefly, event-related potentials were computed by averaging across epochs for each condition, followed by filtering for each canonical frequency band. Hilbert transform was applied to estimate the analytic signal, and the phase was extracted. The group cluster-phase time series was estimated as the averaged phase time series across individuals, from which relative phase time series for each individual could be computed, followed by estimation of the degree of synchrony for each individual with respect to the group. Conceptually, the method is similar to the classical connectivity metric phase-locking value (Lachaux et al., 1999), except relative phase is not computed between individual electrodes or subjects, but between each individual to the group, for each electrode. Similar to ISC, we accounted for potential positive bias induced by higher SNR from collapsed duo conditions by performing the inter-subject analysis separately for each uncollapsed condition, before averaging the duo conditions after group synchrony was computed. To determine an effect of friendship, group cluster phase synchrony was separately computed among friends and among non-friends.

4.10 Statistical analysis

Results are shown as mean with standard error bars. Non-parametric permutation tests (10000 permutations) with FDR correction was used to test for group mean differences. The FDR correction was applied to each metric separately, but across conditions. QQ-plots were assessed to evaluate for normality, and Spearman's correlation was used to test for significant correlations.

Non-parametric cluster-based permutation tests (Maris and Oostenveld, 2007; Sassenhagen and Draschkow, 2019) were applied to handle the multiple comparison problems when taking time, space (electrodes) and frequency into account for estimating group differences in ERDs and group cluster phase synchrony. 1000 permutations were used for ERDs, due to high computational demand with the high dimensionality over time, space and frequency, while 10000 permutations were used for group cluster phase synchrony that only had space and frequency. A T-statistic with cluster forming threshold of 2 was employed. For all statistical tests, a p-value < 0.05 was considered significant for rejection of the null hypothesis.

Data and code availability

The EEG data and code that support the findings of this study are openly available on https://lab.compute.dtu.dk/glia/force-game-eeg-analysis.

Author contributions statement

Q.L. and I.K, conceived and designed the study and experiments. Q.L., A.D., K.K., and I.K. conducted the experiment and collected the data. Q.L. implemented the code. Q.L. and I.K. designed the analysis and generated the first draft of the manuscript. Q.L., A.D., K.K., and I.K. contributed to interpretation of the results and reviewed the manuscript.

Declaration of Competing Interests

The authors take full responsibility for the content of the publication. We declare we have no competing interests.

Acknowledgements

This work is supported by the Villum Young Investigator grant (project no. 37525) and the Carlsberg Semper Ardens: Accelerate grant (CF22-1251).

References

Ablin, P., Cardoso, J. F., and Gramfort, A. (2018). Faster independent component analysis by preconditioning with hessian approximations. *IEEE Transactions on Signal Processing*, 66(15):4040–4049. arXiv: 1706.08171.

Alarcon, G. M., Lyons, J. B., and Christensen, J. C. (2016). The effect of propensity to trust and familiarity on percep-

tions of trustworthiness over time. *Personality and Individual Differences*, 94:309–315.

Alsarrani, A., Hunter, R. F., Dunne, L., and Garcia, L. (2022). Association between friendship quality and subjective wellbeing among adolescents: a systematic review. *BMC Public Health*, 22(1):2420.

Baek, E. C., Hyon, R., López, K., Du, M., Porter, M. A., and Parkinson, C. (2023). Lonely Individuals Process the World in Idiosyncratic Ways. *Psychological Science*, 34(6):683–695. Publisher: SAGE Publications Inc.

Baumeister, R. F. and Leary, M. R. (1995). The need to belong: Desire for interpersonal attachments as a fundamental human motivation. *Psychological Bulletin*, 117(3):497–529. Place: US Publisher: American Psychological Association.

Burt, R. S. (1995). *Structural Holes: The Social Structure of Competition*. Harvard University Press.

Buschman, T. J. and Miller, E. K. (2007). Top-Down Versus Bottom-Up Control of Attention in the Prefrontal and Posterior Parietal Cortices. *Science*, 315(5820):1860–1862. Publisher: American Association for the Advancement of Science.

Chierchia, G., Tufano, F., and Coricelli, G. (2020). The differential impact of friendship on cooperative and competitive coordination. *Theory and Decision*, 89(4):423–452.

Clark-Polner, E. and Clark, M. S. (2014). Understanding and accounting for relational context is critical for social neuroscience. *Frontiers in Human Neuroscience*, 8. Publisher: Frontiers.

Cohen, S. S., Madsen, J., Touchan, G., Robles, D., Lima, S. F. A., Henin, S., and Parra, L. C. (2018). Neural engagement with online educational videos predicts learning performance for individual students. *Neurobiology of Learning and Memory*, 155:60–64.

Cohen, S. S. and Parra, L. C. (2016). Memorable Audiovisual Narratives Synchronize Sensory and Supramodal Neural Responses. *eNeuro*, 3(6). Publisher: Society for Neuroscience Section: New Research.

Cottrell, C. A. and Neuberg, S. L. (2005). Different Emotional Reactions to Different Groups: A Sociofunctional Threat-Based Approach to "Prejudice". *Journal of Personality and Social Psychology*, 88(5):770–789. Place: US Publisher: American Psychological Association.

De Jaegher, H., Di Paolo, E., and Gallagher, S. (2010). Can social interaction constitute social cognition? *Trends in Cognitive Sciences*, 14(10):441–447.

Demos, A. P. and Palmer, C. (2023). Social and nonlinear dynamics unite: musical group synchrony. *Trends in Cognitive Sciences*, 27(11):1008–1018. Publisher: Elsevier.

Desmurget, M. and Turner, R. S. (2010). Motor Sequences and the Basal Ganglia: Kinematics, Not Habits. *Journal of Neuroscience*, 30(22):7685–7690. Publisher: Society for Neuroscience Section: Brief Communications.

Dingemanse, M., Liesenfeld, A., Rasenberg, M., Albert, S., Ameka, F. K., Birhane, A., Bolis, D., Cassell, J., Clift, R., Cuffari, E., De Jaegher, H., Novaes, C. D., Enfield, N. J., Fusaroli, R., Gregoromichelaki, E., Hutchins, E., Konvalinka, I., Milton, D., Raczaszek-Leonardi, J., Reddy, V., Rossano, F., Schlangen, D., Seibt, J., Stokoe, E., Suchman, L., Vesper, C., Wheatley, T., and Wiltschko, M. (2023). Beyond Single-Mindedness: A Figure-Ground Reversal for the Cognitive Sciences. *Cognitive Science*, 47(1):e13230.

Dmochowski, J., Sajda, P., Dias, J., and Parra, L. (2012). Correlated Components of Ongoing EEG Point to Emotionally Laden Attention – A Possible Marker of Engagement? *Frontiers in Human Neuroscience*, 6.

Dmochowski, J. P., Bezdek, M. A., Abelson, B. P., Johnson, J. S., Schumacher, E. H., and Parra, L. C. (2014). Audience preferences are predicted by temporal reliability of neural processing. *Nature Communications*, 5(1):4567. Publisher: Nature Publishing Group.

Dumas, G. (2011). Towards a two-body neuroscience. *Communicative & Integrative Biology*, 4(3):349–352.

Dumas, G., Martinerie, J., Soussignan, R., and Nadel, J. (2012). Does the brain know who is at the origin of what in an imitative interaction? *Frontiers in Human Neuroscience*, 6.

Dumas, G., Nadel, J., Soussignan, R., Martinerie, J., and Garnero, L. (2010). Inter-Brain Synchronization during Social Interaction. *PLOS ONE*, 5(8):e12166. Publisher: Public Library of Science.

Dunbar, R. I. M. (2018). The Anatomy of Friendship. *Trends in Cognitive Sciences*, 22(1):32–51.

D'Onofrio, P., Norman, L. J., Sudre, G., White, T., and Shaw, P. (2022). The anatomy of friendship: neuroanatomic homophily of the social brain among classroom friends. *Cerebral Cortex*, 32(14):3031–3041.

Fishburn, F. A., Murty, V. P., Hlutkowsky, C. O., MacGillivray, C. E., Bemis, L. M., Murphy, M. E., Huppert, T. J., and Perlman, S. B. (2018). Putting our heads together: interpersonal neural synchronization as a biological mechanism for shared intentionality. *Social Cognitive and Affective Neuroscience*, 13(8):841–849.

Flösch, K.-P., Flaisch, T., Imhof, M. A., and Schupp, H. T. (2023). Alpha/beta oscillations reveal cognitive and affective brain states associated with role taking in a dyadic cooperative game. *Cerebral Cortex*, pages 1–16.

Fujioka, T., Trainor, L. J., Large, E. W., and Ross, B. (2012). Internalized Timing of Isochronous Sounds Is Represented in Neuromagnetic Beta Oscillations. *Journal of Neuroscience*, 32(5):1791–1802. Publisher: Society for Neuroscience Section: Articles.

Golland, Y., Arzouan, Y., and Levit-Binnun, N. (2015). The Mere Co-Presence: Synchronization of Autonomic Signals and Emotional Responses across Co-Present Individuals Not Engaged in Direct Interaction. *PLOS ONE*, 10(5):e0125804. Publisher: Public Library of Science.

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., and Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, 7:1–13.

Gugnowska, K., Novembre, G., Kohler, N., Villringer, A., Keller, P. E., and Sammler, D. (2022). Endogenous sources of interbrain synchrony in duetting pianists. *Cerebral Cortex*, 32(18):4110–4127.

Hartup, W. W. and Stevens, N. (1997). Friendships and adaptation in the life course. *Psychological Bulletin*, 121(3):355–370. Place: US Publisher: American Psychological Association.

Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., and Malach, R. (2004). Intersubject Synchronization of Cortical Activity During Natural Vision. *Science*, 303(5664):1634–1640. Publisher: American Association for the Advancement of Science.

He, H., Christensen, J. H., Soerensen, A. J. M., and Konvalinka, I. (2025). Physical proximity, social familiarity, and acoustic environment modulate heart rate synchrony in real-world social interactions. ISSN: 2692-8205 Pages: 2025.12.01.691366 Section: New Results.

Holt-Lunstad, J., Uchino, B. N., Smith, T. W., and Hicks, A. (2007). On the importance of relationship quality: The impact of ambivalence in friendships on cardiovascular functioning. *Annals of Behavioral Medicine*, 33(3):278–290.

Hu, L., Xiao, P., Zhang, Z. G., Mouraux, A., and Iannetti, G. D. (2014). Single-trial time–frequency analysis of electrocortical signals: Baseline correction and beyond. *NeuroImage*, 84:876–887.

Hyon, R., Kleinbaum, A. M., and Parkinson, C. (2020a). Social network proximity predicts similar trajectories of psychological states: Evidence from multi-voxel spatiotemporal dynamics. *NeuroImage*, 216:116492.

Hyon, R., Youm, Y., Kim, J., Chey, J., Kwak, S., and Parkinson, C. (2020b). Similarity in functional brain connectivity at rest predicts interpersonal closeness in the social network of an entire village. *Proceedings of the National Academy of Sciences*, 117(52):33149–33160. Publisher: Proceedings of the National Academy of Sciences.

Jas, M., Engemann, D. A., Bekhti, Y., Raimondo, F., and Gramfort, A. (2017). Autoreject: Automated artifact rejection for MEG and EEG data. *NeuroImage*, 159(December 2016):417–429. arXiv: 1612.08194 Publisher: Elsevier Ltd.

Karau, S. J. and Wilhau, A. J. (2020). Social Loafing and Motivation Gains in Groups: An Integrative Review. In *In-*

dividual Motivation within Groups, pages 3–51. Academic Press.

Karau, S. J. and Williams, K. D. (1993). Social Loafing: A Meta-Analytic Review and Theoretical Integration. *Journal of Personality and Social Psychology*, 65(4):681–706.

Keehn, B., Westerfield, M., Müller, R. A., and Townsend, J. (2017). Autism, Attention, and Alpha Oscillations: An Electrophysiological Study of Attentional Capture. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 2(6):528–536. Publisher: NIH Public Access.

Keller, P. E., Knoblich, G., and Repp, B. H. (2007). Pianists duet better when they play with themselves: On the possible role of action simulation in synchronization. *Consciousness and Cognition*, 16(1):102–111.

Keller, P. E., Novembre, G., and Hove, M. J. (2014). Rhythm in joint action: psychological and neurophysiological mechanisms for real-time interpersonal coordination. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369(1658):20130394.

Kilavik, B. E., Zaepffel, M., Brovelli, A., MacKay, W. A., and Riehle, A. (2013). The ups and downs of beta oscillations in sensorimotor cortex. *Experimental Neurology*, 245:15–26.

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Research Reviews*, 29(2-3):169–195. Publisher: Elsevier.

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored information. *Trends in Cognitive Sciences*, 16(12):606–617.

Klimesch, W., Sauseng, P., and Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews*, 53(1):63–88. Publisher: Elsevier.

Knoblich, G., Butterfill, S., and Sebanz, N. (2011). Psychological Research on Joint Action. Theory and Data. In *Psychology of Learning and Motivation - Advances in Research and Theory*, volume 54, pages 59–101. Elsevier Inc. ISSN: 00797421.

Knoblich, G. and Jordan, J. S. (2003). Action coordination in groups and individuals: Learning anticipatory control. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 29(5):1006–1016.

Konvalinka, I., Bauer, M., Stahlhut, C., Hansen, L. K., Roepstorff, A., and Frith, C. D. (2014). Frontal alpha oscillations distinguish leaders from followers: Multivariate decoding of mutually interacting brains. *NeuroImage*, 94:79–88.

Konvalinka, I., Kompatsiari, K., and Li, Q. (2023a). The fine-grained temporal dynamics of social timing: a window into sociality of embodied social agents. Comment on "The evolution of social timing" by L. Verga, S. A. Kotz, & A. Ravignani. *Physics of Life Reviews*, 47:95–98.

Konvalinka, I. and Roepstorff, A. (2012). The two-brain approach: how can mutually interacting brains teach us something about social interaction? *Frontiers in Human Neuroscience*, 6.

Konvalinka, I., Sebanz, N., and Knoblich, G. (2023b). The role of reciprocity in dynamic interpersonal coordination of physiological rhythms. *Cognition*, 230. Publisher: Elsevier B.V.

Konvalinka, I., Vuust, P., Roepstorff, A., and Frith, C. D. (2010). Follow you, Follow me: Continuous Mutual Prediction and Adaptation in Joint Tapping. *Quarterly Journal of Experimental Psychology*, 63(11):2220–2230. Publisher: SAGE Publications.

Kossinets, G. (2006). Effects of missing data in social networks. *Social Networks*, 28(3):247–268.

Krishna, A. and Götz, F. J. (2024). Motor coordination induces social identity—A novel paradigm for the investigation of the group performance-identity link. *British Journal of Social Psychology*, 63(4):1828–1843.

Lachaux, J. P., Rodriguez, E., Martinerie, J., and Varela, F. J. (1999). Measuring phase synchrony in brain signals. *Human Brain Mapping*, 8(4):194–208.

Lambrechts, L., Accou, B., Vanthornhout, J., Boets, B., and Francart, T. (2025). Interpersonal synchronization as an objective measure of listening engagement.

Larzelere, R. E. and Huston, T. L. (1980). The Dyadic Trust Scale: Toward Understanding Interpersonal Trust in Close Relationships. *Journal of Marriage and Family*, 42(3):595–604. Publisher: [Wiley, National Council on Family Relations].

Leary, M. R. and Kowalski, R. M. (1990). Impression Management: A Literature Review and Two-ComponentModel. *Psychological Bulletin*, 107(1):34–47.

Madsen, J. and Parra, L. C. (2022). Cognitive processing of a common stimulus synchronizes brains, hearts, and eyes. *PNAS Nexus*, 1(1):pgac020.

Maffiuletti, N. A., Aagaard, P., Blazevich, A. J., Folland, J., Tillin, N., and Duchateau, J. (2016). Rate of force development: physiological and methodological considerations. *European Journal of Applied Physiology*, 116:1091–1116.

Marbaker, R. M., Schmad, R. C., Al-Ghamdi, R. A., Sukumar, S., and Ahmed, A. A. (2025). Reward invigorates isometric gripping actions. *Journal of Neurophysiology*, 133(4):1282–1294. Publisher: American Physiological Society.

Marinovic, W., Poh, E., de Rugy, A., and Carroll, T. J. (2017). Action history influences subsequent movement via two distinct processes. *eLife*, 6:e26713. Publisher: eLife Sciences Publications, Ltd.

Maris, E. and Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, 164(1):177–190.

Marks, P. E. L., Babcock, B., Cillessen, A. H. N., and Crick, N. R. (2013). The Effects of Participation Rate on the Internal Reliability of Peer Nomination Measures. *Social Development*, 22(3):609–622.

Mathewson, K. J., Jetha, M. K., Drmic, I. E., Bryson, S. E., Goldberg, J. O., and Schmidt, L. A. (2012). Regional EEG alpha power, coherence, and behavioral symptomatology in autism spectrum disorder. *Clinical Neurophysiology*, 123(9):1798–1809. Publisher: Elsevier.

Michael, J., Sebanz, N., and Knoblich, G. (2016). The Sense of Commitment: A Minimal Approach. *Frontiers in Psychology*, 6. Publisher: Frontiers.

Müller, V., Sänger, J., and Lindenberger, U. (2013). Intra- and Inter-Brain Synchronization during Musical Improvisation on the Guitar. *PLOS ONE*, 8(9):e73852. Publisher: Public Library of Science.

Naeem, M., Prasad, G., Watson, D. R., and Kelso, J. A. S. (2012). Electrophysiological signatures of intentional social coordination in the 10–12 Hz range. *NeuroImage*, 59(2):1795–1803.

Nastase, S. A., Gazzola, V., Hasson, U., and Keysers, C. (2019). Measuring shared responses across subjects using intersubject correlation. *Social Cognitive and Affective Neuroscience*, 14(6):667–685.

Noy, L., Dekel, E., and Alon, U. (2011). The mirror game as a paradigm for studying the dynamics of two people improvising motion together. *Proceedings of the National Academy of Sciences of the United States of America*, 108(52):20947–20952. Publisher: National Academy of Sciences.

Pan, Y., Cheng, X., Zhang, Z., Li, X., and Hu, Y. (2017). Cooperation in lovers: An fNIRS-based hyperscanning study. *Human Brain Mapping*, 38(2):831–841. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hbm.23421.

Parkinson, C., Kleinbaum, A. M., and Wheatley, T. (2018). Similar neural responses predict friendship. *Nature Communications*, 9(1):332. Publisher: Nature Publishing Group.

Parra, L. C., Haufe, S., and Dmochowski, J. P. (2019). Correlated Components Analysis - Extracting Reliable Dimensions in Multivariate Data. *Neurons, Behavior, Data analysis, and Theory*, 2(1). arXiv:1801.08881 [cs, stat].

Pecenka, N. and Keller, P. E. (2011). The role of temporal prediction abilities in interpersonal sensorimotor synchronization. *Experimental Brain Research*, 211(3):505–515.

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., and Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, 51(1):195–203.

Pernet, C. R., Latinus, M., Nichols, T. E., and Rousselet, G. A. (2015). Cluster-based computational methods

for mass univariate analyses of event-related brain potentials/fields: A simulation study. *Journal of Neuroscience Methods*, 250:85–93. Publisher: Elsevier B.V.

Perrin, F., Pernier, J., and Bertrand, O. (1989). Spherical splines for scalp potential and current density mapping. *Electroencephalograp and clinical Neurophysiology*, pages 184–187.

Perry, A., Stein, L., and Bentin, S. (2011). Motor and attentional mechanisms involved in social interaction—Evidence from mu and alpha EEG suppression. *NeuroImage*, 58(3):895–904.

Pezirkianidis, C., Galanaki, E., Raftopoulou, G., Moraitou, D., and Stalikas, A. (2023). Adult friendship and wellbeing: A systematic review with practical implications. *Frontiers in Psychology*, 14. Publisher: Frontiers.

Pezzulo, G., Donnarumma, F., Dindo, H., D'Ausilio, A., Konvalinka, I., and Castelfranchi, C. (2019). The body talks: Sensorimotor communication and its brain and kinematic signatures. *Physics of Life Reviews*, 28:1–21.

Pfurtscheller, G. and Lopes da Silva, F. (1999). Event-related EEG/MEG synchronization and desynchronization: basic principles. *Clinical Neurophysiology*, 110(11):1842–1857. Publisher: Elsevier.

Poulsen, A. T., Kamronn, S., Dmochowski, J., Parra, L. C., and Hansen, L. K. (2017). EEG in the classroom: Synchronised neural recordings during video presentation. *Scientific Reports*, 7:1–9. arXiv: 1604.03019 Publisher: Nature Publishing Group.

Pérez, P., Madsen, J., Banellis, L., Türker, B., Raimondo, F., Perlbarg, V., Valente, M., Niérat, M. C., Puybasset, L., Naccache, L., Similowski, T., Cruse, D., Parra, L. C., and Sitt, J. D. (2021). Conscious processing of narrative stimuli synchronizes heart rate between individuals. *Cell Reports*, 36(11):109692. Publisher: Cell Press.

Rai, L. A., Lee, H., Becke, E., Trenado, C., Abad-Hernando, S., Sperling, M., Vidaurre, D., Wald-Fuhrmann, M., Richardson, D. C., Ward, J. A., and Orgs, G. (2025). Delta-band audience brain synchrony tracks engagement with live and recorded dance. *iScience*, 28(7):112922.

Rempel, J. K., Holmes, J. G., and Zanna, M. P. (1985). Trust in close relationships. *Journal of Personality and Social Psychology*, 49(1):95–112. Place: US Publisher: American Psychological Association.

Richardson, M., Garcia, R. L., Frank, T. D., Gregor, M., and Marsh, K. L. (2012). Measuring group synchrony: a cluster-phase method for analyzing multivariate movement time-series. *Frontiers in Physiology*, 3. Publisher: Frontiers.

Ross, B., Barat, M., and Fujioka, T. (2017). Sound-Making Actions Lead to Immediate Plastic Changes of Neuromagnetic Evoked Responses and Induced β-Band Oscillations during Perception. *Journal of Neuroscience*, 37(24):5948–5959. Publisher: Society for Neuroscience Section: Research Articles.

Salmelin, R. and Hari, R. (1994). Spatiotemporal characteristics of sensorimotor neuromagnetic rhythms related to thumb movement. *Neuroscience*, 60(2):537–550.

Sassenhagen, J. and Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology*, 56(6):1–8.

Schilbach, L., Timmermans, B., Reddy, V., Costall, A., Bente, G., Schlicht, T., and Vogeley, K. (2013). Toward a second-person neuroscience. *Behavioral and Brain Sciences*, 36(4):393–414.

Schmälzle, R., Brook O'Donnell, M., Garcia, J. O., Cascio, C. N., Bayer, J., Bassett, D. S., Vettel, J. M., and Falk, E. B. (2017). Brain connectivity dynamics during social interaction reflect social network structure. *Proceedings of the National Academy of Sciences*, 114(20):5153–5158.

Sebanz, N., Bekkering, H., and Knoblich, G. (2006). Joint action: Bodies and minds moving together. *Trends in Cognitive Sciences*, 10(2):70–76. Publisher: Elsevier Current Trends.

Sebanz, N. and Knoblich, G. (2021). Progress in Joint-Action Research. *Current Directions in Psychological Science*, 30(2):138–143.

Shadmehr, R., Reppert, T. R., Summerside, E. M., Yoon, T., and Ahmed, A. A. (2019). Movement Vigor as a Reflection of Subjective Economic Utility. *Trends in Neurosciences*, 42(5):323–336.

Shen, Y., Hyon, R., Wheatley, T., Kleinbaum, A. M., Welker, C. L., and Parkinson, C. (2025). Neural Similarity Predicts Whether Strangers Become Friends. *Nature Human Behaviour*.

Sijtsma, H., van Buuren, M., Hollarek, M., Walsh, R. J., Lee, N. C., Braams, B. R., and Krabbendam, L. (2023). Social network position, trust behavior, and neural activity in young adolescents. *NeuroImage*, 268:119882.

Smith, J. A. and Moody, J. (2013). Structural effects of network sampling coverage I: Nodes missing at random. *Social Networks*, 35(4):652–668.

Speer, S. P. H., Mwilambwe-Tshilobo, L., Tsoi, L., Burns, S. M., Falk, E. B., and Tamir, D. I. (2024). Hyperscanning shows friends explore and strangers converge in conversation. *Nature Communications*, 15(1):7781. Publisher: Nature Publishing Group.

Spencer, L. and Pahl, R. (2006). *Rethinking Friendship*. Princeton University Press.

Stecher, R., Cichy, R. M., and Kaiser, D. (2025). Decoding the rhythmic representation and communication of visual contents. *Trends in Neurosciences*, 48(3):178–188. Publisher: Elsevier.

Tajfel, H. and Turner, J. (1979). An Integrative Theory of Intergroup Conflict. In Hatch, M. J. and Schultz, M., editors, *The Social Psychology of Intergroup Relations*. Brooks/Cole Publishing Company.

Tetlock, P. E. and Manstead, A. S. (1985). Impression management versus intrapsychic explanations in social psychology: A useful dichotomy? *Psychological Review*, 92(1):59–77. Place: US Publisher: American Psychological Association.

Thura, D. and Cisek, P. (2017). The Basal Ganglia Do Not Select Reach Targets but Control the Urgency of Commitment. *Neuron*, 95(5):1160–1170.e5.

Tognoli, E., Lagarde, J., DeGuzman, G. C., and Kelso, J. A. S. (2007). The phi complex as a neuromarker of human social coordination. *Proceedings of the National Academy of Sciences*, 104(19):8190–8195. Publisher: Proceedings of the National Academy of Sciences.

Tomasello, M. (2014). The ultra-social animal. *European Journal of Social Psychology*, 44(3):187–194. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/ejsp.2015.

van der Horst, M. and Coffé, H. (2012). How Friendship Network Characteristics Influence Subjective Well-Being. *Social Indicators Research*, 107(3):509–529.

Van Lange, P. A. M., Rusbult, C. E., Drigotas, S. M., Arriaga, X. B., Witcher, B. S., and Cox, C. L. (1997). Willingness to sacrifice in close relationships. *Journal of Personality and Social Psychology*, 72(6):1373–1395. Place: US Publisher: American Psychological Association.

Vesper, C., Abramova, E., Bütepage, J., Ciardo, F., Crossey, B., Effenberg, A., Hristova, D., Karlinsky, A., McEllin, L., Nijssen, S. R., Schmitz, L., and Wahn, B. (2017). Joint action: Mental representations, shared information and general mechanisms for coordinating with others. *Frontiers in Psychology*, 7:2039. Publisher: Frontiers Media S.A.

Vesper, C., van der Wel, R. P. R. D., Knoblich, G., and Sebanz, N. (2011). Making oneself predictable: reduced temporal variability facilitates joint action coordination. *Experimental Brain Research*, 211(3):517–530.

Wieselquist, J., Rusbult, C. E., Foster, C. A., and Agnew, C. R. (1999). Commitment, pro-relationship behavior, and trust in close relationships. *Journal of Personality and Social Psychology*, 77(5):942–966. Place: US Publisher: American Psychological Association.

Williams, K. D., Cheung, C. K. T., and Choi, W. (2000). Cyberostracism: Effects of Being Ignored Over the Internet. *Journal of Personality and Social Psychology*, 79(5):748–762.

Williams, K. D. and Karau, S. J. (1991). Social Loafing and Social Compensation: The Effects of Expectations of Co-Worker Performance. *Journal of Personality and Social Psychology*, 61(4):570–581.

Zajonc, R. B. (1965). SOCIAL FACILITATION : A solution is suggested for an old unresolved social psychological problem. *Science*, 149(3681):269–274.

Zamm, A., Debener, S., Bauer, A.-K. R., Bleichner, M. G., Demos, A. P., and Palmer, C. (2018). Amplitude envelope correlations measure synchronous cortical oscillations in performing musicians. *Annals of the New York Academy of Sciences*, 1423(1):251–263.

Zamm, A., Debener, S., Konvalinka, I., Sebanz, N., and Knoblich, G. (2021). The sound of silence: an EEG study of how musicians time pauses in individual and joint music performance. *Social Cognitive and Affective Neuroscience*, 16(1-2):31–42.

Zimmermann, M., Lomoriello, A. S., and Konvalinka, I. (2022). Intra-individual behavioural and neural signatures of audience effects and interactions in a mirror-game paradigm. *Royal Society Open Science*, 9(2). Publisher: The Royal Society.

Zimmermann, M., Schultz-Nielsen, K., Dumas, G., and Konvalinka, I. (2024). Arbitrary methodological decisions skew inter-brain synchronization estimates in hyperscanning-EEG studies. *Imaging Neuroscience*, 2:1–19.