Identification and Regulation of a Hepatic Lipogenic Metabolon

Xiaotong Zhu[1, 2, 6], William B. McKean[3, 4], Tam Tran[1, 5], Xiaorong Fu[1, 2], Jay D. Horton[1, 2, 3, 7], Chai-Wan Kim[1, 2, 6]

Center for Human Nutrition, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046

Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046

Department of Molecular Genetics, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046

START Center for Cancer Research, Mountain Region, West Valley City, UT 84119

Center of Excellence for Leukemia Studies, Department of Pathology, St. Jude Children's Research Hospital, Memphis, TN 38105

These authors contributed equally

Corresponding author

SUMMARY

De novo lipogenesis (DNL) plays a key role in the excessive fat accumulation present in Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD). Most mechanistic studies and experimental strategies for improving hepatic steatosis in MASLD have focused on the transcriptional regulation of enzymes involved in DNL and triglyceride (TG) synthesis. Here, we provide evidence for a post-translational mechanism that enhances fatty acid (FA) and TG synthesis through the assembly of a multi-protein lipogenic metabolon in liver. Under anabolic conditions, acetyl-CoA carboxylase 1 (ACC1) interacts with additional key enzymes in the DNL and TG synthesis pathway. Immunofluorescence and electron microscopy reveal that this lipogenic metabolon localizes around lipid droplets (LDs) and in proximity to mitochondria and LD interfaces in the anabolic state. The formation of the lipogenic metabolon facilitates the efficient transfer of FA synthesis intermediates to enhance lipogenic flux. These findings uncover a new nutrient-responsive, post-translational regulatory mechanism for hepatic lipogenesis and highlight the lipogenic metabolon as a potential therapeutic target for metabolic liver diseases.

Keywords

Liver metabolism, Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD), lipogenic metabolon, *de novo* lipogenesis, fatty acid synthesis, triglyceride synthesis, ACC1, metabolic diseases, post-translational regulation

INTRODUCTION

De novo lipogenesis (DNL) is the process in which fatty acids (FAs) are synthesized in cells from acetyl-CoA, primarily derived from carbohydrate metabolism [1]. Key transcriptional regulators of DNL, including SREBP-1c and ChREBP, are tightly regulated by hormones such as insulin [2,3] and glucagon [2], and through post-translational modifications of the DNL enzymes including phosphorylation [2] and acetylation [4,5]. In a healthy liver, DNL contributes to less than 5% of total hepatic triglycerides (TGs) [6-8]. However, in Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD), this contribution rises to 25-30% [9-11], due to insulin resistance, which leads to an increase in hepatic FA and TG synthesis [2,12-14]. Excessive TG accumulation in liver is associated with progression to metabolic-associated steatohepatitis (MASH), cirrhosis, and hepatocellular carcinoma [15-17]. With over one-third of U.S. adults affected by MASLD [18], it is crucial to elucidate the mechanisms driving aberrant hepatic FA synthesis in insulin-resistant states to identify potential therapeutic targets for FA-driven metabolic disorders.

Our previous studies have shown that the excessive activation of SREBP-1c initiates the transcription of genes encoding the enzymes necessary for FA synthesis, as well as the enzyme catalyzing the first step in glycerolipid synthesis [12,13]. This transcriptional program is a major driver of elevated FA and TG synthesis in insulin-resistant states [2,12,13], subsequently leading to hepatic steatosis, increased VLDL secretion, and hypertriglyceridemia [13,19-24]. The requirement of SREBP activation for the development of hepatic steatosis and hypertriglyceridemia was demonstrated by studies in which SCAP, a protein required for SREBP activation, was deleted in obese, insulin resistant mice (*ob/ob* mice) that normally have very high levels of hepatic FA and TG synthesis. SCAP deletion completely normalized rates of FA synthesis, liver TGs, VLDL secretion, and plasma TG levels [24] in *ob/ob* mice and hamsters. These results demonstrate that enhanced DNL is a major driver of steatosis in insulin-resistant livers.

FA and TG synthesis requires sequential enzymatic reactions distributed across multiple subcellular compartments. ATP citrate lyase (ACL) produces acetyl-CoA from citrate, acetyl-CoA synthetase 2 (ACSS2) synthesizes acetyl-CoA from acetate, acetyl-CoA carboxylases (ACC) 1 and 2 convert acetyl-CoA to malonyl-CoA, and fatty acid synthase (FAS) uses malonyl-CoA and

acetyl-CoA to generate palmitate (C16:0). These enzymes (except for ACC2 that is predominantly
localized to the outer mitochondrial membrane) are predominantly localized in the cytosol [25].
Palmitate can then be further desaturated by stearoyl-CoA desaturase (SCD1) or elongated by
ELOVL fatty acid elongase 6 (ELOVL6) before further desaturation, but unlike the enzymes
involved in the synthesis of palmitate, these proteins reside in the endoplasmic reticulum (ER)
[26,27]. Additionally, the first enzyme in glycerolipid synthesis, glycerol-3-phosphate acyltransferase
1 (GPAM), which acylates glycerol-3-phosphate to initiate TG and phospholipid synthesis is
located in the mitochondria [28]. How the FA and glycerolipid synthesis intermediates are efficiently
transferred among these spatially distributed enzymes to sustain continuous and efficient lipid
synthesis remains unclear.

One conceptual framework that could address this challenge is the metabolon hypothesis. In
1970, A.M. Kuzin first introduced the concept of physical enzyme–enzyme complexes. This idea
gained momentum in 1972 when P.A. Srere applied it to enzymes of the citric acid cycle,
highlighting their potential functional interactions [29]. Srere further expanded on this idea by coining
the term "metabolon" during a 1985 lecture in Debrecen, Hungary. He defined it as "a
supramolecular complex of sequential metabolic enzymes and cellular structural elements" [30],
emphasizing the functional coordination within cellular metabolic pathways. This concept has
been applied to explain the spatial organization of metabolic pathways such as the citric acid cycle
[31,32] and glycolysis [33], prompting us to examine whether hepatic FA and TG synthesis might also
be coordinated through spatial organization of lipogenic enzymes.

Here, we show that under anabolic conditions—such as high-carbohydrate feeding and insulin
resistance—FA and TG synthesis enzymes assemble into a functional lipogenic metabolon. This
complex enhances lipogenic flux by coordinating the sequential enzymatic reactions required to
synthesize the end product. This metabolon, including accessory proteins MIG12 and SPOT14
(S14) [34], predominantly localizes around lipid droplets (LDs) in primary hepatocytes. Together,
these findings reveal a dynamic multi-enzyme complex that facilitates FA and TG synthesis and
provides new insights into the spatial organization and coordinated regulation of hepatic lipid
synthesis.

RESULTS

Identification of the lipogenic complex in liver

The first indication that a lipogenic complex might exist grew out of investigations focused on
determining the function of a poorly characterized protein, S14. S14 is highly expressed in

lipogenic tissues [35], and its gene transcription is regulated by SREBP-1c [36,37], carbohydrate-
responsive element-binding protein (ChREBP) [38], and thyroid hormone receptor (TR) [39], which
suggests that it may play a role in lipogenesis. Deletion of S14 in mice reduced lipid synthesis in
liver [34] and lactating mammary glands [40], although its exact function in lipogenesis has not been
elucidated. To further define the function of S14, we used an adenovirus that expressed S14 with
a Flag tag in mice and carried out immunoprecipitation of S14 from liver lysates to identify potential
interacting proteins. LC/MS analysis revealed that MIG12, ACC1, FAS, and β-tubulin all co-
immunoprecipitated with S14 (Figure 1A). MIG12, a protein we previously showed that promotes
the polymerization of ACC1 and ACC2 [41], shares 32% sequence identity with S14 [41] and can
form a heterodimer with S14 [34]. To further confirm that the association of these enzymes co-
immunoprecipitated with S14 reflects physical interaction in cells, we performed
immunofluorescence staining in primary hepatocytes isolated from rats refed a fat-free, high-
carbohydrate diet (FFD). ACC1, MIG12, and S14 prominently co-localized around LD–associated
ER regions. Quantitative analysis using Pearson's correlation (above Costes threshold)
demonstrated strong spatial overlap among the three enzyme pairs, with ACC1–S14 (R = 0.73),
ACC1–MIG12 (R = 0.52), and S14–MIG12 (R = 0.52) (Supplemental Table 1). These data
indicate that ACC1, S14, and MIG12 occupy shared subcellular domains under conditions of high
lipogenesis, supporting the existence of a coordinated lipogenic enzyme complex in hepatocytes
(Figure 1B). Representative ROIs and pixel-intensity histograms are shown in Figure S1, and
detailed colocalization metrics, including Manders' coefficients and Costes statistics, are provided
in Supplemental Table 1.

115   Co-localization of proteins involved in lipogenesis around LDs in hepatocytes

To investigate potential additional interactions among lipogenic proteins, we used antibodies
recognizing various lipogenic enzymes in primary hepatocytes to assess co-localization by
confocal immunofluorescence. As shown in Figure 1, co-localization was quantified using
Pearson's correlation coefficient (Pearson's R) in LD-enriched regions where the enzyme signals
were concentrated. Immunofluorescence imaging of primary hepatocytes isolated from rat livers
refed a FFD demonstrated that ACC1 co-localized with ACL (R ≈ 0.83), ACSS2 (R ≈ 0.46), and
FAS (R ≈ 0.55) around LDs, despite their known localization in the cytosol. GPAM, a mitochondrial
outer membrane protein involved in glycerolipid synthesis, also co-localized with ACC1 in LD-
associated regions (R ≈ 0.51). In contrast, the gluconeogenic enzyme PEPCK, which is unrelated
to lipogenesis, showed almost no overlap with ACC1 (R ≈ 0.11), likely because it remains confined
to the cytosol and does not redistribute to LD-proximal regions (Figure 2A, Supplemental Table

2). Comprehensive testing metrics, including Manders' coefficients and Costes significance
testing, are provided in Supplemental Table 2 and further support the robust spatial association
of ACC1 with multiple cytosolic lipogenic enzymes, whereas the non-lipogenic negative control,
PEPCK, showed only minimal overlap with ACC1. Adipose differentiation-related protein (ADRP),
an LD-resident protein, marked LDs where ACL, ACC1, and FAS also localized in primary
hepatocytes from refed rats (Figure 2B). Three-dimensional imaging further confirmed the
localization of ACC1 surrounding LDs in primary hepatocytes from refed rats (Figure 2C).
Additional imaging across various fasting and refeeding timepoints suggested that ACC1-
containing complexes assemble progressively during metabolic activation and accumulate
around LDs in rat primary hepatocytes (Figure S2).

Co-localization of proteins involved in lipogenesis at the interface of mitochondria
and LDs in hepatocytes

Electron microscopy (EM) with immunogold labeling using antibodies recognizing ACC1 or FAS
was used to determine their subcellular localization in primary hepatocytes from refed rats.
Consistent with the immunofluorescence images (Figure 2C; Figure S2), EM revealed that ACC1
localized predominantly around LDs (Figure 3A). This finding was further supported by
immunofluorescence imaging and EM performed on the same hepatocyte (Figure S3A). Double
immunogold labeling using different sizes of gold particles for ACC1 and FAS demonstrated their
co-localization around LDs (Figure 3B; Figure S3B–C). Although ACC1 and FAS were
occasionally observed near the ER, they were much more consistently enriched around LDs
(Figure S3D). This pattern aligned with staining of the ER marker protein disulfide isomerase
(PDI), which showed only limited overlap with ACC1, in contrast to the strong co-localization
observed between ACC1 and LDs (Figure S3E).

Notably, interactions between ACC1 and FAS were also observed at interfaces between
mitochondria and LDs (Figure 3C). Subcellular fractionation of liver lysates from FFD-refed rats
further demonstrated that ACL, ACC1, ACC2, FAS, SCD1, and GPAM were present in both LD
and mitochondrial fractions, regardless of their canonical localizations in the cytoplasm,
mitochondria, or ER (Figure 3D). EM frequently revealed mitochondria positioned adjacent to
LDs (Figure S3F), and GPAM localization was subsequently confirmed to be on the outer
mitochondrial membrane, including sites where mitochondria interface with LDs (Figure S3G).
Subcellular fractionation of liver lysates from FFD-refed rats further confirmed this finding,
whereas mitochondrial inner membrane proteins, such as prohibitin and cytochrome c, were
found only in the mitochondrial fraction (Figure S3H).

Combined, these findings demonstrate that lipogenic proteins co-localize around LDs and at
mitochondria–LD contact sites under high lipogenesis conditions.

Active dephosphorylated ACC promotes polymerization, enhances lipogenesis,
and increases LD localization

ACC activity is acutely regulated through phosphorylation and dephosphorylation. AMP-activated
protein kinase (AMPK) phosphorylates ACC at Ser-79 [42,43], inhibiting its enzymatic activity through
largely undefined mechanisms [44,45]. Fasting or glucagon treatment increases AMPK activity,
whereas insulin suppresses AMPK activity and stimulates the Mg^{2+}-dependent protein
phosphatase 2A (PP2A), which dephosphorylates ACC, thereby increasing its activity [46-50]. During
refeeding, ACC is converted from its phosphorylated inactive form to its dephosphorylated, active
form due to stimulation of protein phosphatase activity. This dephosphorylation activates ACC
and DNL in hepatocytes, allowing excess glucose to be converted into FAs and TGs for storage
[51,52].

We next determined whether the phosphorylated state of ACC affected the formation of the
lipogenic complex. For this study, we treated rat primary hepatocytes with 5-aminoimidazole-4-
carboxamide ribonucleoside (AICAR), an AMPK agonist, to induce ACC phosphorylation [53]. The
primary hepatocytes were incubated with 1 mM AICAR and cell lysates were prepared at 0, 1.5,
and 3 hours after treatment. SDS-PAGE immunoblot analysis of cell lysates showed an increase
in phosphorylated ACC levels in the AICAR-treated cells compared to the untreated cells;
however, the amount of total ACC protein was unchanged (Figure 4A). Blue native PAGE analysis
of cell lysates demonstrated that the polymerized, higher molecular weight form of ACC1 was
present only in untreated primary hepatocytes and was disrupted in AICAR-treated hepatocytes
(Figure 4A). Even with MIG12 overexpression, which robustly promotes ACC1 polymerization,
AICAR reduced the higher molecular weight form of ACC1 to baseline (Figure 4B). Consistent
with reduced polymerization, FA synthesis measured by ^{3}H-acetate incorporation into newly
synthesized FAs fell ~6-fold within 30 minutes of AICAR treatment (Figure 4C), while FA-synthesis
enzyme levels remained unchanged (Figure S4A). Collectively, these results imply that the rapid
decrease in DNL results from AMPK-mediated ACC phosphorylation, which disrupts ACC1
polymerization.

To further explore the role of dephosphorylated ACC in promoting DNL *in vitro*, we blocked
endogenous phosphatase activity in liver cytosol using a phosphatase inhibitor cocktail (Figure
4D). Immunoblot analysis confirmed that the inhibitor prevented ACC dephosphorylation,

whereas λ-phosphatase enhanced dephosphorylation (Figure S4B). Compared with the inhibitor-
treated group, untreated liver cytosol, which underwent dephosphorylation during incubation,
showed a 4-fold increase in palmitate synthesis rates in a citrate concentration-dependent manner
(Figure 4D). Furthermore, in primary hepatocytes from refed rats, ACC1 localized around LDs
under DMSO control, consistent with Figures 1–3, but this LD-associated pattern was significantly
reduced with AICAR treatment (Figure 4E).

Taken together, these findings demonstrate that the polymerization of ACC, mediated by
dephosphorylation, is critical for its localization with other lipogenic enzymes around LDs and for
regulating DNL in hepatocytes.

The interaction of ACC1 with lipid synthesis-associated proteins is enhanced when ACC is in the activated state

As demonstrated above, ACC1 polymerizes and co-localizes with other lipogenic proteins around
LDs under high DNL conditions. To determine how ACC1 interactions with other proteins change
upon activation, we performed ACC1 immunoprecipitation from mouse liver lysates incubated
with 5 mM $MgCl_2$ at 30°C for 30 minutes with or without a phosphatase inhibitor cocktail. With
endogenous phosphatases active, ACC1 co-immunoprecipitated ACC2, ACSS2, FAS, ACL,
GPAM, MIG12, S14, and SCD1 (Figure 5A). Notably, β-actin, a cytoskeletal protein, also showed
significant binding to ACC1, suggesting that cytoskeletal components might play a role in the
formation of the lipogenic complex. In contrast, these interactions were markedly reduced when
endogenous phosphatases were inhibited, suggesting that ACC1 interacts more strongly with
other proteins in a phosphatase-active state. Proteins unrelated to lipid synthesis, such as
PEPCK, prohibitin, cytochrome c, and PDI, did not bind ACC1 under either condition.

Notably, ACC2 maintained strong binding to ACC1 even in the presence of phosphatase
inhibitors, suggesting that ACC1 and ACC2 form a relatively stable heteromer. These
observations are supported by immunoprecipitation of liver lysates from hepatocyte-specific
ACC1 or ACC2 knockout mice, which revealed a clear reciprocal interaction between the two
enzymes (Figure S5A). Blue native PAGE analysis further supported this model: whereas ACC1
migrated similarly in wild-type and ACC2-deficient livers, ACC2 showed a markedly altered
pattern in ACC1-deficient livers. In normal liver, ACC2 predominantly appeared as a dimer or
tetramer, but in the absence of ACC1, ACC2 shifted toward higher–molecular-weight, more
polymerized species (Figure S5B). Considering that hepatic ACC2 expression is substantially

lower than ACC1, these findings suggest that, under physiological conditions, most endogenous ACC2 may exist as a heterodimer with ACC1.

Immunofluorescence analysis in cultured cells provided additional evidence for the ACC1–ACC2 heterodimer. ACC1 was detected predominantly in the cytosol when expressed alone (Figure S5C). However, when co-transfected with ACC2, a substantial fraction of ACC1 colocalized with ACC2 at the mitochondria (Figure S5D). In addition to this mitochondrial pool, ACC1 also displayed cytosolic localization that did not overlap with ACC2. These observations indicate that ACC1 is present in the cytosol but also at the mitochondrial membrane as a complex with ACC2.

To determine whether endogenous ACC2 influences ACC1-mediated protein interactions, we immunoprecipitated ACC1 from refed wild-type and ACC2 hepatocyte-specific knockout livers, using ACC1-deficient mice as a negative control. Co-precipitation of ACL, ACSS2, FAS, S14, SCD1, GPAM, β-actin, and β-tubulin was reduced to varying degrees in ACC2-deficient livers compared with wild-type (Figure 5B), suggesting that ACC2 contributes to the stability or assembly of ACC1-associated protein complexes. MIG12 binding to ACC1 was unaffected by phosphatase inhibition (Figure 5A) or by the absence of ACC2 (Figure 5B), consistent with its direct association with ACC1. As expected, PEPCK, prohibitin, cytochrome c, and calnexin were not co-immunoprecipitated with ACC1 in any group.

Given that ACL, ACSS2, ACC, FAS, SCD1, and the mitochondrial GPAM participate in ACC1-associated complexes, we next measured TG synthesis under phosphatase active versus phosphatase-inactive conditions. Glycerol 3-phosphate (G3P) was used as a substrate with excess oleoyl-CoA added to liver homogenates from refed rats to test whether TG synthesis via GPAM is enhanced when endogenous phosphatases are active, a condition that favors lipogenic complex formation (Figure S5E). The results demonstrated that TG synthesis was significantly higher under phosphatase-active conditions and increased further with rising $MgCl_2$ concentrations, paralleling $MgCl_2$-dependent ACC1 dephosphorylation (Figure S4B), whereas under phosphatase inactive conditions, TG synthesis remained unchanged with $MgCl_2$ (Figure S5E).

These results demonstrate that phosphorylation–dephosphorylation regulates both FA and TG synthesis in part by controlling ACC1-mediated lipogenic complex formation.

Lipogenic metabolon formation facilitates FA synthesis

To confirm the interactions between ACC1 and the other FA synthesis proteins, and to assess

the regulatory effects of phosphorylation and dephosphorylation on the activities of individual FA

synthesis enzymes *in vitro*, we conducted a comprehensive purification of recombinant ACSS2,

ACC1, FAS, MIG12, and S14 in mammalian cells using Ni-NTA resin and gel-filtration

chromatography (Figure 6A, S6A), and rPEPCK was purified as a negative control (Figure S6B).

The recombinant protein mixture was dephosphorylated with λ-phosphatase and

immunoprecipitation was performed with anti-ACC1 antibodies. Immunoblot analysis (Figure 6B)

revealed that all other recombinant FA synthesis proteins co-immunoprecipitated with rACC1,

whereas rPEPCK did not. These results confirm that recombinant ACC1 interacts with core

lipogenic factors *in vitro* using recombinant proteins (Figure 6B).

Next, we pre-incubated the individual recombinant proteins or their mixtures with λ-phosphatase

to simulate the DNL activated environment as described above. Parallel groups without λ-

phosphatase treatment were used as phosphorylated controls. Figure S6C shows that each

purified protein exhibited functional activity, and interestingly, all tested lipogenic proteins

displayed increased activity to varying degrees when dephosphorylated compared to their

phosphorylated forms. However, rACC1 exhibited the greatest increase in enzymatic activity upon

λ-phosphatase treatment compared with the other lipogenic enzymes. Next, we combined purified

lipogenic enzymes, rACL-rACC1-rFAS or rACSS2-rACC1-rFAS, and measured FA synthesis

from citrate or acetate, respectively. As predicted, rACL-rACC1-rFAS activity was nearly four

times higher with λ-phosphatase treatment compared with untreated conditions (Figure 6C).

Similarly, rACSS2-rACC1-rFAS activity increased by ~3.5-fold upon λ-phosphatase treatment

relative to untreated conditions (Figure 6C). Figure S6D shows the phosphorylation levels of

rACC1 alongside total rACC1 levels at various time points following λ-phosphatase treatment,

demonstrating that λ-phosphatase efficiently dephosphorylates proteins without altering their total

levels.

Inasmuch as each rACL, rACSS2, rACC1, and rFAS protein exhibited increased enzymatic

activity in the dephosphorylated state, these observations alone could not distinguish whether the

enhanced palmitate production resulted from cooperative complex formation or simply reflected

additive increases in the intrinsic activities of individual enzymes upon dephosphorylation. To

directly address this issue, we performed a FA-synthesis assay using (1) liver cytosol alone, (2)

recombinant ACL–ACC1–FAS alone, and (3) a mixture of liver cytosol and recombinant ACL–

ACC1–FAS, with protein amounts normalized across conditions. Each group was subjected to

either dephosphorylated or phosphorylated conditions under identical treatment settings. As

shown in Figure S6E, dephosphorylation was successfully achieved in all conditions. Under
dephosphorylated conditions, combining liver cytosol with recombinant ACL–ACC1–FAS
produced substantially greater palmitate than the expected additive activity calculated from
cytosol alone and recombinant enzymes assayed in parallel (Figure 6D). This enhancement
exceeded what would be predicted from simple additive effects of dephosphorylation on individual
enzymes and was supported by a significant interaction between phosphorylation state and
mixing condition (two-way ANOVA, $F_{(3,16)}$ = 133.4, P < 0.0001). Together, these data indicate
a strong synergistic increase in lipogenesis arising from cooperative interactions between
dephosphorylated cytosolic proteins and recombinant enzymes. In contrast, under
phosphorylated conditions, the palmitate produced by the combined reaction was
indistinguishable from the expected additive activity (Figure 6E), indicating that cooperative
assembly and synergistic flux enhancement are selectively enabled by dephosphorylation.

To directly assess complex formation under these conditions, ACC1 was immunoprecipitated
from the same reaction mixtures. Consistent with the results shown in Figure 5A,
dephosphorylation markedly enhanced the interactions of ACC1 with FAS and ACL, and β-tubulin
as determined by co-precipitation studies, suggesting that the lipogenic metabolon may be
functionally linked to cytoskeletal components (Figure S6F). No nonspecific binding was observed
with control rabbit IgG.

While it is not feasible to systematically test all cytosolic proteins that could potentially contribute
to metabolon assembly, Figure 5 identified structural proteins as representative non-lipogenic
factors that interact with ACC1 under dephosphorylated conditions. Consistent with this
observation, β-tubulin showed detectable co-localization with ACC1 around lipid droplets in
primary hepatocytes under lipogenic refeeding conditions (Pearson's R ≈ 0.32), whereas this
spatial association was markedly reduced upon fasting (Figure 6F). In parallel, pharmacological
perturbation of microtubule dynamics provided support for the role of cytoskeleton in lipogenesis.
Disrupting microtubule polymerization with nocodazole consistently decreased FA synthesis in
primary hepatocytes, whereas stabilizing microtubules with taxol, an agent with the opposite effect
of nocodazole, had minimal impact on lipogenesis (Figure 6G). These observations suggest that
intact cytoskeletal architecture facilitates lipogenesis by coordinating the nutrient-responsive
assembly and spatial organization of lipogenic enzyme complexes, rather than directly regulating
microtubule dynamics.

Collectively, these findings suggest that lipogenic activity is enhanced through post-translational
regulation that promotes the formation of a functional lipogenic metabolon under physiological

conditions favoring high DNL. This process likely involves dynamic and cooperative interactions
among DNL- and TG-related enzymes, aided by structural proteins and other accessory factors,
allowing for more integrated and efficient lipogenesis.

ACC1 anchors a multi-module lipogenic network integrating translation,
trafficking, and lipid metabolism

To further map the ACC1-centered protein network, we immunoprecipitated ACC1 from wild-type
and ACC-deficient liver lysates. In parallel, MIG12 and S14, two direct ACC1-binding partners
were immunoprecipitated, as was ACSS2, a representative lipogenic enzyme, to compare their
interaction profiles (Figure 7A). MIG12 and S14 pulldowns closely mirrored the ACC1
immunoprecipitation pattern: proteins enriched in wild-type lysates were markedly reduced in
ACC-deficient lysates, consistent with their direct association with ACC1 and the dependence of
their interactomes on ACC1 integrity. In contrast, ACSS2 immunoprecipitation showed a distinct
profile. Although several lipogenic enzymes were reduced in ACC-deficient lysates, ACSS2
retained clear interactions with multiple lipogenic proteins. This pattern suggests that ACC1
enhances or stabilizes the overall lipogenic assembly rather than acting as its sole organizing
component. Notably, β-actin binding to ACSS2 was unchanged regardless of ACC1 status,
implying that actin contributes to complex organization through an ACC1-independent
mechanism.

We next quantified proteins co-immunoprecipitated with ACC1 by mass spectrometry. Proteins
that were reduced by >20-fold in ACC1, MIG12, or S14 immunoprecipitations from ACC-deficient
lysates were classified as ACC1-dependent interactors. STRING network analysis of these
proteins resolved three highly connected clusters (Figure 7B, top). The largest cluster consisted
of ribosomal and translation-associated proteins, suggesting that a robust interface between
ACC1 and the translational machinery exists. A second cluster was enriched for vesicle/trafficking
components and cytoskeletal protein-binding factors, indicating that ACC1 is embedded within a
broader transport and structural scaffold. A third cluster consisted of enzymes involved in FA
synthesis and remodeling, consistent with ACC1's canonical metabolic function (Figure 7B,
bottom). DAVID functional annotations performed on the same dataset produced closely matching
categories (Figure S7 and Supplemental Table 3).

Together, these findings indicate that activated ACC1 functions as a central organizational hub
that coordinates core lipogenic enzymes—including FAS, ACL, ACSS2, SCD1, GPAM, and
ACC2—into a coherent metabolic assembly. Direct binding by MIG12 and S14 likely stabilizes

this structure, while cytoskeletal and trafficking components associate with the ACC1-centered

complex to support spatial organization and substrate delivery. The presence of ribosomal

proteins and transport-associated factors among ACC1-dependent interactors further suggests

that lipogenesis is integrated with protein synthesis and intracellular trafficking through multiple

points of connection.

DISCUSSION

Transcriptional control of FA and TG synthesis is well characterized [54], but much less is known

about how lipid synthesis is regulated post-transcriptionally. Here, we identified a previously

unrecognized post-translational mechanism in which hepatic lipogenic enzymes assemble into a

multi-protein metabolon under anabolic conditions. This metabolon provides a spatial basis for

efficient glycerolipid synthesis and introduces a new dimension to the regulation of DNL.

Using immunofluorescence and EM, we showed that enzymes involved in FA and TG synthesis,

along with accessory proteins like MIG12 and S14, co-localize around LDs and function as an

integrated multi-enzyme assembly rather than as isolated enzymes under physiological activation

states such as high carbohydrate feeding. These findings build on earlier LD proteomics reports

that identified ACC1 [55-59] and FAS [58] on LDs, extending them by providing direct structural

visualization of their co-localization in mammalian hepatocytes. Furthermore, the proximity

between cytosolic enzymes such as ACC1 and FAS and the mitochondrial enzyme GPAM at

contact sites suggests coordinated substrate transfer into the glycerolipid pathway. Subcellular

fractionation of liver lysates from high-carbohydrate-refed rats further supports these

observations.

Immunoprecipitation of ACC1 from liver lysates incubated under phosphatase-active conditions

revealed robust interactions between ACC1 and multiple lipogenic proteins, whereas the inclusion

of phosphatase inhibitors markedly reduced these associations (Figure 5A). This biochemical shift

was paralleled by *in vitro* FA and TG synthesis assays, in which phosphatase-active conditions

produced substantially higher lipid synthesis than phosphatase-inhibited ones. We initially

attributed this enhanced interaction and activity to ACC1 dephosphorylation, a hypothesis

supported by *in vivo* experiments using the AMPK agonist AICAR. In primary hepatocytes isolated

from high-carbohydrate–refed rats, AICAR treatment caused the activated, polymerized form of

ACC1 to dissociate into its inactive dimeric form, as shown by blue native PAGE. The extensive

polymerization of ACC1 driven by MIG12 overexpression was also significantly reduced by

AICAR-mediated ACC1 phosphorylation. Correspondingly, short-term AICAR treatment

significantly suppressed hepatic lipogenesis in primary hepatocytes without affecting the steady-
state levels of lipogenic enzymes and markedly reduced ACC1 localization around LDs. Notably,
ACC1 was not the only enzyme influenced by dephosphorylation, as phosphatase treatment also
increased the enzymatic activities of rACSS2 and rFAS, although to a lesser extent than ACC1.
These findings suggest that anabolic phosphatase-active states coordinate multiple components
of the lipogenic pathway, collectively enhancing metabolon assembly and catalytic output.

Another notable finding from the immunoprecipitation experiments was the strong interaction of
ACC1 with ACC2 and MIG12. Whereas most ACC1-associated proteins showed enhanced
binding under phosphatase-active conditions, ACC2 and MIG12 maintained stable interactions
with ACC1 irrespective of its phosphorylation state. This feature suggests that these two proteins
form a relatively stable core structural interaction with ACC1.

Additional insight into ACC2 was obtained from transfection studies in CHO-K1 cells. When ACC1
and ACC2 were co-expressed, the fraction of ACC1 that interacted with ACC2 and formed
heterodimers or higher-order heteromultimers and localized to mitochondria increased (Figure
S5D). ACC1 also retained a substantial cytosolic pool that did not overlap with ACC2, likely
reflecting cytosolic homomeric ACC1 assemblies. These observations indicate that ACC1 is
partitioned between heteromeric ACC1–ACC2 complexes at mitochondria and homomeric ACC1
assemblies in the cytosol and suggest that the subcellular localization and functional partitioning
of malonyl-CoA–producing acetyl-CoA carboxylase activity may be influenced by the relative
abundance of ACC2. These findings are consistent with the tissue-specific expression patterns
of ACC isoforms. In liver and adipose tissue, where malonyl-CoA is used primarily for lipogenesis,
ACC1 is the predominant isoform; whereas in muscle and heart, where malonyl-CoA mainly
inhibits mitochondrial fatty acid oxidation, ACC2 is the major form. Thus, cells may regulate the
relative abundance (stoichiometry) of ACC1 and ACC2 to redistribute acetyl-CoA carboxylase
activity between the cytosol and mitochondria according to metabolic demands. In this context,
ACC1–ACC2 interactions may serve as a regulatory node linking FA synthesis and oxidation.
Alterations in the stoichiometry of the two isoforms could therefore shift the balance between
lipogenesis and control of mitochondrial FA oxidation by modulating the subcellular distribution of
carboxylase activity.

Finally, although ACC2 has long been viewed primarily as a regulator of mitochondrial FA
oxidation, ACC2 deletion reduced ACC1 interactions with multiple lipogenic enzymes and
structural proteins (Figure 5B). This finding supports the view that ACC2 functions not only as a
metabolic enzyme that inhibits FA oxidation but also as a structural component essential for the

organization and maintenance of the lipogenic metabolon. Complementing these structural
insights, our biochemical assays demonstrated that assembling lipogenic enzymes within a
shared reaction environment generates catalytic outputs that exceed the sum of individual
activities. Under DNL-activated conditions, combining recombinant ACL–ACC1–FAS with liver
cytosol nearly doubled the FA synthesis compared with the expected additive activity, indicating
synergistic enhancement of pathway flux. This synergy suggests that enzyme co-assembly, rather
than simply increased intrinsic activity, facilitates more efficient substrate channeling during active
lipogenesis. These functional observations reinforce the multi-module organization described in
Figure 7, in which ACC1 immunocomplexes interact not only with lipogenic enzymes but also with
ER-proximal translation components, vesicle-movement proteins, and actin-linked scaffolds,
supporting the concept that lipogenic enzymes achieve maximal efficiency when spatially
integrated with cytoskeletal scaffolds and intracellular transport machinery, and when functionally
linked to translation-associated components.

Although metabolon-based regulation is less explored in lipogenesis, analogous mechanisms
have been established in the TCA cycle and glycolysis [29-31,60-63]. The substrate-responsive
assembly of PFK filaments in the presence of fructose-6-phosphate [64] provides a relevant
precedent and parallels ACC1 filament formation under anabolic conditions, suggesting that these
filaments may serve as structural scaffolds for metabolon assembly. Consistent with this concept,
a FA–synthesis metabolon has been described in the thermophile *Chaetomium thermophilum* [65]
and ACC–FAS co-localization has been visualized in *Saccharomyces cerevisiae* [66], where these
enzymes are reorganized or sequestered in response to glucose starvation. These observations
across lower eukaryotes support the idea that nutrient state governs the assembly and
disassembly of spatially organized lipogenic enzymes.

Recent work by Wijesinghe *et al.* [67] extends this concept to mammalian cells. In their study, ACC1
formed ring-like puncta and localized to the surface of lipid droplets (LDs), but this localization
was highly conditional: anabolic cues such as AMPK inhibition preserved ACC1–LD co-
localization, whereas large LDs generated by acute oleic-acid loading—which diminishes
anabolic drive—displayed reduced ACC1 staining. This pattern mirrors our observation that the
AMPK agonist AICAR suppresses ACC1 accumulation around LDs (Figure 4E), indicating that
ACC1 recruitment to LDs requires appropriate metabolic activation rather than merely reflecting
LD abundance. Together, these findings align with our model in which ACC1 polymerization and
its interactions with accessory factors generate a nutrient-responsive lipogenic complex in
hepatocytes. Collectively, the emerging evidence across species supports a broader framework

in which ACC1 integrates nutrient status with spatial enzyme organization to optimize lipogenesis

during anabolic transitions.

In conclusion, our study demonstrates that under anabolic conditions, hepatic lipogenic enzymes

assemble into a spatially organized, phosphorylation-responsive metabolon around LDs. This

multi-protein metabolon enhances catalytic efficiency facilitating improved transfer of

intermediates between sequential enzymes, thereby supporting efficient synthesis of FA and TGs.

These findings provide a mechanistic foundation for understanding hepatic lipid synthesis and

highlight therapeutic opportunities in metabolic disease by targeting the structure and regulation

of the lipogenic metabolon.

Limitations of the study

In this study, we present evidence for a novel post-translational regulatory mechanism that

enhances FA and TG synthesis in the liver via the assembly of a multi-protein metabolon. Ideally,

since the structure of ACC1 under different activation states has been reported [68,69], we would

have included structural analyses of this ACC1 mediated lipogenic metabolon. However,

detecting such metabolons presents greater challenges compared to stable protein assemblies.

The reason is that a metabolon is characterized as a transient structural-functional metabolon

formed by sequential enzymes within a metabolic pathway, exhibiting non-covalent interactions

and associating with various cellular structural elements, such as integral membrane proteins and

cytoskeletal components [30,70]. Consequently, the interactions among these proteins are more

transient and dynamic, enabling them to adapt to fluctuations in metabolic environments.

Additionally, the mRNA levels of the lipogenic proteins are all significantly downregulated during

fasting as a result of reduced SREBP-1c expression, which restricted our ability to compare the

co-localization of lipogenic proteins between fasting and refeeding conditions. Therefore, we

employed AICAR treatment and elevated phosphorylation levels to simulate catabolic states *in*

vivo and *in vitro*, respectively.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this manuscript, the authors used ChatGPT (OpenAI) solely for

language editing, including grammar correction and improvement of clarity. No content, data

interpretation, or scientific conclusions were generated by the tool. After using this assistance,

the authors thoroughly reviewed, revised, and approved all text, and they take full responsibility

for the integrity and accuracy of the final manuscript.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to the lead contact, Jay D. Horton (jay.horton@utsouthwestern.edu).

Materials availability

Custom antibodies generated in this study are available upon reasonable request and completion of an MTA. Cloning constructs are available from the Lead Contact. All other materials used in this study are specified in the key resources table.

Data and code availability

All data supporting the findings are available within the paper and Supplemental Information. No custom code was used.

ACKNOWLEDGEMENTS

We thank Norma Anderson, Judy Sanchez, Tuyet Dang, and Tessa Edwards for excellent technical assistance; Lisa Beaty for suspension cell culture and set up; Linda Donnelly for assistance with antibody generation; Jinhai Yu for providing TEV protease; and Chelsea Burroughs and Nancy Heard for help with graphics. The work was supported by grants NIH P01HL160487 and NIH P30DK127984.

AUTHOR CONTRIBUTIONS

X.Z., W.B.M., T.T., X.F., and C.-W.K. conducted the experiments. J.D.H., C.-W.K., and X.Z. designed the experiments. X.Z., C.-W.K., and J.D.H. wrote the manuscript. All authors reviewed and approved the article.

DECLARATION OF INTERESTS

J.D.H. is a consultant for AbbVie and insitro.

Figure Legends

Figure 1. Identification of the lipogenic complex in liver

(A) An adenovirus expressing S14-Flag was injected into mice, and liver lysates were prepared 3 days later. Immunoprecipitation was performed using anti-Flag antibodies. Eluates from the S14-Flag IP and an identically processed β-gal–injected control were separated on 8% and 15% SDS–PAGE gels, followed by silver staining. Protein bands that appeared specifically in the S14-Flag lanes were excised and analyzed by LC–MS/MS to identify potential S14-associated proteins.

(B) Primary hepatocytes were prepared from rats refed a FFD and plated on collagen-coated coverslips. Immunofluorescence confocal imaging was performed as described in Methods. Co-localization of ACC1 with MIG12 and S14 was confirmed by confocal microscopy using anti-mouse monoclonal ACC1, anti-MIG12 rabbit polyclonal antibodies, and anti-S14 rabbit polyclonal antibodies. Quantitative co-localization analysis was performed using Pearson's correlation (above Costes threshold), and full analytical metrics, including Manders' coefficients and Costes significance testing, are provided in Figure S1 and Supplemental Table 1.

Figure 2. Co-localization of proteins involved in lipogenesis around LDs in hepatocytes

(A) Co-localization of enzymes involved in lipogenesis in primary hepatocytes from rats refed a FFD using confocal microscopy as described in Methods. Quantitative co-localization analysis was performed using Pearson's correlation (above Costes threshold), and full analytical metrics are provided in Supplemental Table 2.

(B) Localization of enzymes involved in lipogenesis around LDs in primary hepatocytes from rats refed a FFD using confocal microscopy as described in Methods. Co-localization analysis was performed using Pearson's correlation as in panel A, with results summarized in Supplemental Table 2.

(C) A three-dimensional image stack was acquired using an OMX microscope (z-step size: 120 nm) to visualize the indicated proteins around lipid droplets, as described in Methods.

Figure 3. Co-localization of proteins involved in lipogenesis at the interface of mitochondria and LDs in hepatocytes

(A) Primary hepatocytes were prepared from rats refed a FFD and localization of ACC1 was determined using immunogold-EM. The 1.4 nm nanogold particles were enlarged by gold enhancement for clear visibility. Detailed protocol is described in Methods.

(B) Co-localization of ACC1 and FAS around LDs in primary hepatocytes from rats refed a FFD using immunogold-EM. Size of gold particles: ACC1 - 5 nm; FAS - 10 nm.

(C) Co-localization of ACC1 and FAS between mitochondria and LDs in primary hepatocytes from rats refed the FFD using immunogold-EM. Size of gold particles: ACC1 - 5 nm; FAS - 10 nm.

(D) Subcellular fractionation was performed with liver lysates from rats refed the FFD to isolate LDs as well as mitochondria, microsomes, and cytoplasm. Detailed protocols for the subcellular fractionation are described in Methods. Proteins from each fraction were subjected to SDS-PAGE and immunoblot analysis with the indicated antibodies. PLIN2, prohibitin, and α-tubulin were used as subcellular localization markers for lipid droplets, mitochondria, and cytosolic proteins, respectively.

Figure 4. Active dephosphorylated ACC promotes polymerization, enhances lipogenesis, and increases LD localization

(A) Primary hepatocytes from rats refed a FFD were treated with 1 mM AICAR for 0, 1.5, and 3 hours. Equal aliquots of cell lysate proteins were loaded and analyzed by BN-PAGE (above) or SDS-PAGE (bottom) for immunoblot analysis. An anti-ACC1 antibody was used to perform the immunoblot analysis of BN-PAGE. For SDS-PAGE, ACC immunoblots were performed using a fluorescently labeled streptavidin conjugate.

(B) Primary hepatocytes from rats refed a FFD were isolated. One set of primary hepatocytes were untreated, while the other two sets were infected with an adenovirus expressing MIG12. After 12 hours, 1 mM AICAR was added, and cells were harvested 3 hours later. Cell lysates were prepared and subjected to BN-PAGE and SDS-PAGE for immunoblot analysis. An anti-ACC1 antibody was used to perform the immunoblot analysis of BN-PAGE. For SDS-PAGE, a fluorescently labeled streptavidin conjugate to detect ACC and an anti-MIG12 antibody were used for immunoblot analysis.

(C) Primary rat hepatocytes prepared as in (B) were incubated with 1 mM AICAR for 30 minutes and FA synthesis rates were measured $ex\ vivo$ using ^{3}H-acetate as described in Methods. Individual data points are shown; bars indicate the mean (n = 2 technical replicates from one hepatocyte preparation).

(D) Liver cytosol from rats refed a FFD were pre-incubated with 1 mM $MnCl_2$ in the presence or

absence of a phosphatase inhibitor cocktail for 15 min at 30°C, then incubated with the indicated

concentrations of Na-citrate buffer (pH 7.1) for 15 min at 37°C. Palmitate synthesis (left) was

measured using ^{14}C-citrate. Palmitate synthesis rates (right) were calculated from the maximal

velocity (Vmax) of the enzymatic reaction. Data represent mean ± SEM from n = 3 technical

replicates. Statistics: multiple two-tailed unpaired Student's t-tests with Holm–Šidák correction,

comparing phosphatase inhibitor–treated (phosphorylated) vs untreated (dephosphorylated)

conditions at each citrate concentration (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$). The

detailed protocol for the enzymatic assay is described in Methods.

(E) Rat primary hepatocytes were isolated from rats fed a FFD for 2 days and incubated in medium

containing either (i) 25 mM glucose, 100 nM insulin, and 1% DMSO or (ii) 5 mM glucose and 1

mM AICAR. Following the indicated times, cells were stained with anti-ACC1 antibodies and

LipidTOX as described in Methods.

Figure 5. The interaction of ACC1 with lipid synthesis-associated proteins is

enhanced when ACC is in the activated state

(A) Liver lysates from refed mice were treated with or without a phosphatase inhibitor cocktail.

The ACC1 antibody was incubated with the lysate for 1 hour at RT and ACC1 was

immunoprecipitated. IgG was used as the control. The immunoprecipitation protocol is described

in Methods. Supernatant indicates the unbound fraction collected after bead pelleting; it was

mixed with SDS sample buffer and immunoblotted alongside the bead-bound material.

(B) ACC1 immunoprecipitation was carried out using lysates prepared from refed wild-type, ACC1

hepatocyte knockout, or ACC2 hepatocyte knockout mouse livers, respectively.

Figure 6. Lipogenic metabolon formation facilitates FA synthesis

(A) Human C-terminal His-Flag-tagged ACSS2, ACL, ACC1, FAS, MIG12, and S14 were

individually overexpressed in mammalian HEK 293S GnTI⁻ cells and purified using Ni-NTA and

gel filtration. Epitope tags were removed by TEV cleavage. Detailed protocols for purification were

described in Methods. The final purified recombinant proteins were subjected to SDS-PAGE for

Coomassie blue staining.

(B) Recombinant protein mixtures were incubated with λ-phosphatase and 1 mM $MnCl_2$ for 30

minutes at 30°C and then the protein mixture was incubated with 0.5 mM Na-citrate buffer for 20

minutes at 30°C. rACC1 was immunoprecipitated from the protein mixture using an anti-ACC1

antibody. Recombinant protein mixtures without rACC1 were used for control
immunoprecipitations. Proteins were subjected to SDS-PAGE for immunoblot analysis. ACC1
immunoblots were performed using a fluorescently labeled streptavidin conjugate.

(C) Enzymatic activity of reconstituted recombinant lipogenic enzyme modules was measured as
described in Methods. The left panel shows FA synthesis activity measured from citrate using
rACL–rACC1–rFAS, whereas the right panel shows FA synthesis activity measured from acetate
using rACSS2–rACC1–rFAS. Reactions were performed under phosphorylated or
dephosphorylated conditions as indicated. Data represent mean ± SEM from n = 3 technical
replicates. Relative synthesis rates were calculated based on Vmax values.

(D–E) Under dephosphorylated (D) or phosphorylated (E) conditions, ACL–ACC1–FAS activity
was measured in mixtures of liver cytosol (from FFD-refed rats) and recombinant proteins (rACL,
rACC1, and rFAS) and compared with the expected additive activity calculated from assays
performed in cytosol alone and recombinant proteins alone in parallel. Data represent mean ±
SEM from n = 3 technical replicates. Statistical analysis was performed using two-way ANOVA,
revealing a significant interaction between phosphorylation state and mixing condition. Post hoc
comparisons indicate that observed activity significantly exceeded the expected additive activity
only under dephosphorylated conditions. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

(F) Primary hepatocytes isolated from rats under refed or fasted conditions were fixed and stained
for ACC1 and β-tubulin, followed by confocal microscopy as described in Methods. Images show
representative localization patterns under each condition.

(G) Primary hepatocytes from FFD-fed rats were cultured for 8 hours and then treated with DMSO,
nocodazole, or taxol (20 μM) as indicated. Cells were subsequently incubated with [14]C-labeled
citrate for 2 or 4 hours, and citrate-derived FA synthesis was quantified as described in Methods.
Data points represent individual measurements from duplicate experiments.

Figure 7. ACC1 anchors a multi-module lipogenic network integrating translation,
trafficking, and lipid metabolism

(A) Liver lysates were prepared from *ob/ob* mouse livers and ACC1/ACC2–deleted *ob/ob* mouse
livers as described in Methods. To activate endogenous phosphatases, lysates were incubated
in 5 mM MgCl$_2$ for 30 minutes at 30°C. In parallel, anti-ACC1, anti-MIG12, anti-S14, or anti-
ACSS2 antibodies were pre-bound to protein G magnetic beads by incubating for 30 minutes at
room temperature. The activated lysates were then added directly to the antibody–bead

complexes and incubated for 1 hour at room temperature to perform immunoprecipitation. After
extensive washing, bound proteins were eluted by boiling and analyzed by SDS-PAGE and
immunoblot analysis was carried out with the indicated antibodies.

(B) Proteins co-immunoprecipitated with ACC1, MIG12, or S14 that were reduced by >20-fold in
ACC-deficient lysates were classified as ACC1-dependent interactors and subjected to STRING
network analysis. Three major functional clusters emerged—translation (red), protein
trafficking/transport and cytoskeletal protein binding (green), and FA metabolism (blue). These
clusters are displayed both as a network graph and summarized in the bar graph below using the
corresponding color scheme. The inset mini-graphs within each colored bar summarize
representative GO/KEGG terms for that cluster, with full enrichment analyses provided in Figure
S7.

Key resource table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-ACL	Abcam	Cat# ab40793
Rabbit anti-ACSS2	Abcam	Cat# ab66038
Rabbit anti-mouse ACC1	House antibodies[71]	508D or 510D
Mouse monoclonal anti-mouse ACC1	House antibodies	2B5
Rabbit anti-mouse ACC2	House antibodies	456D
Rabbit anti-human FAS	House antibodies	310D
Rabbit anti-mouse MIG12	House antibodies[72]	143D
Mouse monoclonal anti-mouse MIG12	House antibodies	6C4
Rabbit anti-mouse S14	House antibodies	39C
Rabbit anti-PEPCK	Abcam	Cat# ab70358
Mouse anti-β-Tubulin	ThermoFisher	Cat# MA5-16308
Rabbit anti-β-Actin	Cell Signaling	Cat# 4970S
Rabbit Anti-mouse GPAM	House antibodies	280D
Mouse monoclonal anti-rat GPAM	House antibodies	16F8
Anti-ADRP	Sigma	Cat# SAB4200452
Anti-SCD1	Cell Signaling	Cat# 2794S
Anti-PROHIBITIN	Santa Cruz	Cat# SC-56467

Anti-CYTOCHROME C	Millipore	Cat# MAC1800
Mouse anti-CPT1	Abcam	Cat# ab128568
Anti-Ser79-pACC1	Cell Signaling	Cat# 3661S
Anti-PDI	Cell Signaling	Cat# 2446S
Goat Anti-Rabbit IgG, F(ab')2	Jackson Immuno	Cat# 111-035-047
Goat Anti-Mouse IgG, F(ab')2	Jackson Immuno	Cat# 115-035-072
Li-COR IRDye 800 goat anti-Rabbit IgG	Fisher Scientific	Cat# NC9401842
Li-COR IRDye 680 goat anti-Mouse IgG	Fisher Scientific	Cat# NC0252290
Li-COR IRDye 680RD Streptavidin	Fisher Scientific	Cat# NC0337633
Bacterial and virus strains		
Stellar™ Competent Cells	Takara	Cat# 636764
TOP10 Competent Cells	ThermoFisher	Cat# C404003
Chemicals, peptides, and recombinant proteins		
LB broth	Research Products International	Cat# L24061
Freestyle 293 expression medium	Gibco	Cat# 12338018
Opti-MEM™ I Reduced Serum Medium	Gibco	Cat# 31985070
Fetal Calf Serum (FCS)	Sigma-Aldrich	Cat# F2442
Penicillin-Streptomycin solution (100x)	Corning	Cat# 30-002-CI
PEI 25K	Polysciences	Cat# 23966-1
Protease Inhibitor Cocktail Set III, EDTA-Free - Calbiochem	Millipore-Sigma	Cat# 539134-1ML
Protease Inhibitor Cocktail, for use in purification of Histidine-tagged proteins, DMSO solution	Sigma-Aldrich	Cat# P8849-5ML
Purified rTEV protease	Gift from Dr. Jinhai Yu, UT Southwestern Medical Center	N/A
Lambda Protein Phosphatase	New England Biolabs	Cat# P0753L
DL-Dithiothreitol	Sigma-Aldrich	Cat# 43819-5G
Adenosine 5'-triphosphate magnesium salt	Sigma-Aldrich	Cat# A9187-1G
Adenosine 5'-triphosphate disodium salt hydrate	Sigma-Aldrich	Cat# A2383-5G

Coenzyme A sodium salt hydrate	Sigma-Aldrich	Cat# C3144-25MG
Acetyl coenzyme A sodium salt	Sigma-Aldrich	Cat# A2056-25MG
Malonyl coenzyme A lithium salt	Sigma-Aldrich	Cat# M4263-10MG
NADPH, Tetrasodium Salt	Sigma-Aldrich	Cat# 481973-100MG
Sodium bicarbonate [14C]	American Radiolabeled Chemicals	Cat# ARC 0138C-50 µCi
Malonyl [2-14C] Coenzyme A	American Radiolabeled Chemicals	Cat# ARC 0528-25 µCi
Citric Acid, [1,5-14C]	Revvity	Cat# NEC160050UC
Acetic Acid, Sodium Salt, [1,2-14C]	Revvity	Cat# NEC553050UC
Lauryl Maltose Neopentyl Glycol	Anatrace	Cat# NG310 5 GM
3a70B Complete Counting Cocktail	Research Products International	Cat# 111154
In-Fusion® Snap Assembly Value Bundle	Takara	Cat# 638946
Platinum™ SuperFi II PCR Master Mix	ThermoFisher	Cat# 12368010
DreamTaq Green PCR Master Mix (2X)	ThermoFisher	Cat# K1081
Oleoyl coenzyme A dipotassium salt	ThermoFisher	Cat# CN10090405
sn-Glycerol 3-phosphate lithium salt	Sigma-Aldrich	Cat# 94124-10MG
Glycerol, L-[14C(U)] 3-phosphate disodium salt	American Radiolabeled Chemicals	Cat# ARC 0316-50 µCi
Sodium butyrate	Sigma-Aldrich	Cat# 303410-500G
HCS LipidTOX™ Deep Red Neutral Lipid Stain, for cellular imaging	Fisher Scientific	Cat# H34477
Clean-Blot™ IP Detection Reagent (HRP)	Thermo Scientific	Cat# 21230
AICAR	Abcam	Cat# ab120358
Critical commercial assays		
Pierce™ Coomassie (Bradford) Protein Assay Kit	ThermoFisher	Cat# 23200

Pierce BCA Protein Assay Kit	ThermoFisher	Cat# 23225
Silver staining Kit	ThermoFisher	Cat# 24612
Experimental models: Cell lines		
HEK-293S GnTI⁻	ATCC	Cat# CRL-3022; RRID: CVCL_A785
CHO-K1	ATCC	Cat# CCL-61; RRID: CVCL_0214
Experimental models: Organisms/strains		
C57BL/6J mouse	Jackson Laboratory	Stock# 000664
ob/ob mouse	Jackson Laboratory	Stock# 000632
Sprague Dawley	Charles River	SAS SD code 400
Phosphatase inhibitor cocktail		
Sodium Fluoride	Sigma-Aldrich	Cat# S7920
Imidazole	Sigma-Aldrich	Cat# I5513
Sodium molybdate	Sigma-Aldrich	Cat# M1003
Sodium tartrate dihydrate	Sigma-Aldrich	Cat# S4797
Sodium pyrophosphate	Sigma-Aldrich	Cat# P8135
β-glycerophosphate	Sigma-Aldrich	Cat# G9422
Recombinant DNA		
pEZT-BM	Addgene	Plasmid #74099
ATP citrate lyase (ACLY) (NM_001096) Human Tagged ORF Clone	Origene	Cat# RC200508
ACSS2 (NM_018677.4) cDNA ORF clone, Homo sapiens (human)	Genscript	Cat# OHu18043
ACL-Tev-Flag-His-XT3	IDT DNA	N/A
ACC1-Tev-Flag-His-XT3	IDT DNA	N/A
FAS-Tev-Flag-His-XT3	IDT DNA	N/A
MIG12-Tev-Flag-His_Human	IDT DNA	N/A
S14-Tev-Flag-His_Human	IDT DNA	N/A
PEPCK_Human-Tev-His	IDT DNA	N/A
pEZT-BM-ACL-3xFlag-10xHis	This paper	N/A
pEZT-BM-ACSS2-3xFlag-10xHis	This paper	N/A
pEZT-BM-ACC1-3xFlag-10xHis	This paper	N/A

pEZT-BM-FAS-3xFlag-10xHis	This paper	N/A
pEZT-BM-MIG12-3xFlag-10xHis	This paper	N/A
pEZT-BM-S14-3xFlag-10xHis	This paper	N/A
pEZT-BM-PEPCK-3xFlag-10xHis	This paper	N/A
ACC1-pcDNA3.1	This paper	N/A
ACC2-pcDNA3.1	This paper	N/A
Software and algorithms		
Image Studio™ Version 5.0	Li-COR	N/A
Prism 10	GraphPad	https://www.graphpad.com/
Adobe Illustrator 2023	Adobe	N/A
Other		
PD-10 desalting columns	Cytiva	Cat# 17085101
Superose 6 10/300 GL column	Cytiva	Cat# GE17-5172-01
Amicon Ultra centrifugal filter	Millipore	Cat# UFC903008
Ni-NTA Resin	Thermo Scientific	Cat# 88221
Slide-A-Lyzer dialysis cassette	Thermo Scientific	Cat# 66203
Dynabeads Protein A beads	Invitrogen	Cat# 10002D
Scintillation vials	Research Products International	Cat# 121040
Disposable Round-Bottom Rimless Glass Tubes	Corning	Cat# 14-962-15E
4–15% Criterion™ TGX™ Precast Midi Protein Gel, 26 well, 15 µl	BIO-RAD	Cat# 5671085
4–15% Mini-PROTEAN® TGX™ Precast Protein Gels, 15-well, 15 µl	BIO-RAD	Cat# 4561086
Dynabeads™ Protein G for Immunoprecipitation	Fisher Scientific	Cat# 10-004-D
Slide Warmer (110 V, for 23 slides)	Fisher Scientific	Cat# 50-930-986

Methods

Experimental model and study participant details

Animals and primary hepatocytes

Primary hepatocytes were isolated from 4-week-old male Sprague–Dawley rats fasted for 12
hours and refed for 12 hours with a fat-free diet, using collagenase perfusion as described by
Shimomura et al. [73] All rat studies were performed under an approved protocol from the UT
Southwestern Institutional Animal Care and Research Advisory Committee (IACUC #2015-
100838). Liver tissues were collected from wild-type male C57BL/6 mice, liver-specific ACC1
and/or ACC2 knockout male C57BL/6 mice, or ob/ob male C57BL/6 mice. All mice were
maintained under standard housing conditions with a 12-hour light/dark cycle (lights on at 6 AM
and off at 6 PM). Mouse studies were conducted under UT Southwestern IACUC approval (2015-
101135-G). Nocodazole and taxol were used at 20 µM to achieve robust disruption or stabilization
of microtubules, respectively, in primary hepatocytes.

Method Details

Antibody generation

For the generation of the mouse monoclonal antibody that recognizes mouse ACC1, a cDNA
encoding the N-terminal 95 amino acids of mouse ACC1 was cloned into a pGEX-4T-1 vector
and the recombinant protein was expressed in *Escherichia coli (E. coli)* and purified using
glutathione agarose. Monoclonal antibodies were generated by fusion of Sp2/mIL-6 (CRL-2016;
ATCC) mouse myeloma cells with splenic B lymphocytes derived from a female NZBWF1/J
mouse injected with purified mouse ACC1 protein [74,75].

For the generation of a rabbit polyclonal antibody against mouse ACC2, a cDNA encoding the N-
terminal 200 amino acids of mouse ACC2 was cloned into a pRSF-1b expression vector and
expressed in *E. coli*. The purified His-tagged protein was isolated using nickel affinity resin and
injected into rabbits as described [76].

For the generation of rabbit polyclonal antibody against human FAS, a cDNA encoding the C-
terminal 305 amino acids from human FAS was cloned into pGEX-4T-1 expression vector and
expressed in *E. coli.* The purified GST-fusion protein was injected into rabbits as described [19].

For the generation of a rabbit polyclonal antibody against mouse S14, recombinant (His)6-mouse
S14 was expressed in E. coli, purified by Ni-NTA affinity chromatography (Qiagen) according to
the manufacturer's protocol, and injected into rabbits as described [19].

For the generation of rabbit polyclonal antibody against mouse GPAM, a cDNA encoding the N-
terminal 150 amino acids from mouse GPAM was cloned into pRSF-1b expression and expressed
in *E. coli*. The purified His-tagged protein was injected into rabbits as described [19].

For the generation of mouse monoclonal antibody against rat GPAM, a cDNA encoding amino

acids 627 to 717 of rat GPAM was cloned into pRSF-1b expression vector and expressed in *E.*

coli. The purified His-tagged protein was injected into mice, and monoclonal antibodies were

produced as described above. For the generation of mouse monoclonal antibody against mouse

MIG12, full length mouse MIG12 cDNA was cloned into pRSF-1b and expressed in *E. coli* and

the purified protein (Ni-NTA agarose) was injected into mice. Monoclonal antibodies were

generated using the same hybridoma fusion strategy described for ACC1.

Immunofluorescence

Primary hepatocytes from refed rat livers were fixed in 4% paraformaldehyde (PFA) in PHEM

buffer (pH 7.4) for 30 minutes. After rinsing the cells with 0.1 M phosphate buffer, they were

permeabilized with pre-chilled (−20°C) methanol for 15 min at −20°C. Blocking was performed

using 1% BSA and 0.2% fish skin gelatin in 0.1 M phosphate buffer (pH 7.4) for 30 minutes at

room temperature. Primary antibodies were incubated overnight at 4°C in the blocking solution,

and Alexa Fluor 488- or 568-labeled secondary antibodies were used to visualize the localization

of each target protein. Imaging was conducted using a Zeiss LSM880 Airyscan microscope.

Colocalization analysis

Colocalization was quantified using the Coloc2 plugin in Fiji/ImageJ. To minimize background

noise and evaluate biologically relevant signal overlap, analysis was performed on lipid droplet–

enriched regions, which represent functional sites of lipogenic enzyme accumulation. CZI files

were imported directly into Fiji, and channels were split and processed separately to avoid

spectral bleed-through. Images used for Coloc2 analysis were converted to 8-bit format prior to

computation. Pearson's correlation coefficients were calculated using Costes automatic

thresholding, and above-threshold Pearson's R values were reported in the main text.

Thresholded Manders' coefficients (tM1 and tM2), Costes significance testing, and 2D intensity

histograms were generated by Coloc2 and are presented in Figure S1. All images were acquired

below saturation, identical ROIs were applied to paired channels for each analysis.

Pre-embedding immuno-electron microscopy

Primary hepatocytes from refed rat livers were fixed in 4% PFA/0.05% glutaraldehyde (GA) in

PHEM buffer (pH 7.4) for 30 minutes. After rinsing the cells with 0.1 M phosphate buffer, residual

aldehydes were quenched with 50 mM glycine in 0.1 M phosphate buffer for 15 minutes, followed

by permeabilization with 0.2 mg/mL saponin for 30 minutes at room temperature. Blocking was

performed using 5 µg/mL saponin, 1% BSA, 0.2% fish skin gelatin, and 5% goat serum in 0.1 M

phosphate buffer (pH 7.4) for 15 minutes at room temperature. The primary antibody was

incubated overnight at 4°C in the blocking solution. After incubation, cells were labeled with an

anti-rabbit or anti-mouse IgG secondary antibody conjugated to 1.4 nm gold. Following post-

fixation, the samples were subjected to gold enhancement, embedded in Embed-812 resin, and

cut into 60 nm sections. Electron micrographs were taken using the TEM JEOL 1400 Plus

microscope. For ACC1 and FAS double immunogold labeling, ACC1 and FAS primary antibodies

were bound to 5 nm and 10 nm colloidal gold-conjugated secondary antibodies, respectively. After

post-fixation, the cells were embedded in Embed-812 resin, sectioned into 60 nm slices, and

imaged using the JEOL 1400 Plus Transmission Electron Microscope. The pre-embedding

method followed the procedure described by Yamamoto *et al*. [77]

Liver subcellular fractionation

Fresh rat or mouse liver was homogenized in 50 mM Tris (pH 7.4) and 250 mM sucrose in the

presence of protease inhibitors. For lipid droplet isolation, half of the homogenate was transferred

to a new tube, and buffer A (20 mM HEPES, pH 7.4, 100 mM KCl, 2 mM $MgCl_2$) was gently added

to the top. The sample was centrifuged at 2,000 × g for 30 minutes, and the floating lipid layer

was transferred to a new tube. This lipid layer was mixed with an equal volume of buffer A-500S

(20 mM HEPES, pH 7.4, 100 mM KCl, 2 mM $MgCl_2$, and 500 mM sucrose) and gently overlaid

with buffer A. This step was repeated twice to isolate a pure lipid layer. A glass pipette was used

during lipid layer purification to minimize loss of lipid. Proteins from the lipid layer were extracted

using an organic solvent as described [78] and resuspended in SDS-lysis buffer (50 mM Tris, pH

6.8, 50 mM NaCl, 1% SDS). For mitochondria and microsome fractionations, the remaining half

of the liver homogenate was centrifuged at 1,000 × g for 10 minutes to remove the nuclear fraction.

The supernatant was centrifuged at 10,000 × g for 20 minutes to pellet the mitochondrial fraction.

The mitochondrial pellet was resuspended and washed twice more with homogenization buffer

by repeated centrifugation at 10,000 × g for 20 minutes. The final mitochondrial pellet was

resuspended in SDS-lysis buffer. The supernatant from the 10,000 × g centrifugation was further

centrifuged at 100,000 × g for 30 minutes to sediment the microsomal fraction, which contains the

endoplasmic reticulum. The microsomal pellet was rinsed with homogenization buffer and

resuspended in SDS-lysis buffer.

Recombinant protein expression and purification

Expression

The cDNAs encoding human ACSS2, ACL, ACC1, FAS, MIG12, S14, and PEPCK were cloned into the pEZT-BM vector containing a TEV protease cleavage site, an SSG linker, followed by a C-terminal 3 × Flag tag and 10 × His tag. All proteins were expressed in mammalian HEK 293S GnTI⁻ cells (ATCC). A total of 500 mL of HEK 293S GnTI⁻ cells were transfected at a density of 2–3 × 10⁶ cells/mL with 0.75 mg of plasmid DNA using PEI 25K (Polysciences). 16 hours post-transfection, 250 mL of Freestyle 293 expression medium (Gibco) supplemented with 2% fetal calf serum (FCS) and 1% penicillin-streptomycin solution (Corning) was added to the cultures. Sodium butyrate was added to a final concentration of 8 mM to enhance protein expression. Cells were harvested 72 hours post-transfection for subsequent protein purification.

Purification

The cell pellet from 400 mL culture was resuspended and homogenized in 100 mL buffer B (50 mM phosphate buffer, 500 mM NaCl, 1 mM DTT, 5 mM imidazole, 225 mM mannitol, 75 mM sucrose, protease inhibitor cocktail (Millipore-Sigma), protease inhibitor cocktail for use in purification of Histidine-tagged proteins (Sigma-Aldrich), pH 8.0). The homogenate was centrifuged at 135,000 × g for 45 minutes at 4 °C. The supernatant was collected and filtered through a 5 µm filter. The filtered supernatant was loaded on a 2 mL Ni-NTA resin (Thermo Scientific) that pre-equilibrated with buffer B and flowed through by gravity (~1 ml/min) at 4°C. The Ni-NTA was then washed sequentially with 10 CV buffer B and 25 CV buffer C (50 mM phosphate buffer, 500 mM NaCl, 1 mM DTT, 50 mM imidazole, 10% glycerol, pH 8.0) by gravity. Proteins were eluted in 5 CV buffer D (50 mM phosphate buffer, 500 mM NaCl, 1 mM DTT, 500 mM Imidazole, 10% glycerol, pH 8.0). The eluate was desalted using a PD-10 desalting column (Cytiva) pre-equilibrated with buffer E (10 mM HEPES, 50 mM NaCl, 100 mM KCl, 1 mM DTT, 5% glycerol, pH 7.4). Proteins were eluted in buffer E and incubated with TEV protease overnight at 4 °C.

Removal of TEV and tags

rACL, rACC1, and rFAS: the TEV-cleaved proteins were concentrated to 0.5 mL using a 30,000 MWCO Amicon Ultra centrifugal filter (Millipore) and then subjected to gel-filtration chromatography on a Superose 6 10/300 GL column (Cytiva) pre-equilibrated with buffer E. Purified proteins eluted as a single sharp peak in buffer E. Protein homogeneity was confirmed by Coomassie blue-stained SDS-PAGE.

rACSS2, rMIG12, rS14, and rPEPCK: the TEV-cleaved proteins were loaded on 2 mL Ni-NTA resin and flowed through by gravity (~1 ml/min) pre-equilibrated with buffer E. The Ni-NTA resin

769 was then washed sequentially with 5 CV buffer B, 5 CV buffer C and 5 CV buffer D by gravity
770 flow. The flow-through and eluted fractions without His tag signal were combined and
771 concentrated using an Amicon Ultra centrifugal filter (Millipore) (rACSS2 and rPEPCK: 30,000
772 MWCO, rMIG12: 10,000 MWCO, rS14: 3,000 MWCO) and then dialyzed using a 2,000 MWCO
773 Slide-A-Lyzer dialysis cassette (Thermo Scientific) in 2 L buffer E overnight at 4°C.

Co-immunoprecipitation of recombinant proteins

Recombinant ACC1, FAS, ACL, ACSS2, MIG12, S14, and PEPCK were mixed at 8: 64: 20: 40:
32: 96: 1 molar ratios, respectively. Groups lacking recombinant ACC1 served as controls. The
protein mixtures were incubated with 40 mM HEPES (pH 7.1), 1 mM $MnCl_2$, and λ-phosphatase
(New England Biolabs) for 30 minutes at 30°C. For phosphorylated groups, the mixtures were
incubated with a phosphatase inhibitor cocktail instead of lambda phosphatase, under the same
conditions. The reaction systems were then supplemented with 0.5 mM Na-citrate buffer (pH 7.1)
and incubated for 20 minutes at 30°C.

After pre-incubation, 5 µg of ACC1 polyclonal antibody was added to each reaction and incubated
for an additional 10 minutes at 30°C. Buffer F (50 mM NaCl, 100 mM KCl, 20 mM HEPES, 4 mM
$MgCl_2$, 1% LMNG, pH 7.1) was added, and the mixtures were incubated with rotation for 1 hour
at room temperature. Subsequently, 25 µL Dynabeads Protein A (Invitrogen) were added and
incubated with rotation for 20 minutes at room temperature. The beads were washed four times
with buffer F. Bound proteins were eluted by incubation with 1× SDS sample buffer for 10 minutes
at 95°C. Eluted proteins were analyzed by immunoblot analysis using the indicated antibodies.

Co-immunoprecipitation of liver lysates

Liver slices from fresh mouse (or rat) livers were homogenized in 10 mM Tris (pH 7.1), 250 mM
sucrose, and 1% LMNG (Lauryl maltose neopentyl glycol) in the presence of protease inhibitors.
The homogenate was centrifuged at 25,600 × g for 15 minutes at 20°C. The supernatant was
collected and filtered through a 0.45 µm filter. The clarified supernatant was divided into two tubes;
5 mM $MgCl_2$ was added to both, whereas phosphatase inhibitor cocktail was added to only one.
After incubation at 30°C for 30 minutes, NaCl and KCl were added to final concentrations of 50
mM and 100 mM, respectively. Mouse monoclonal ACC1 antibody or mouse IgG control antibody
was added to each tube, and the samples were incubated for 1 hour at room temperature.
Dynabeads Protein G (25 µL) were then added and incubated for a further 25 minutes at room
temperature. The beads were washed 5 times with 10 mM Tris (pH 7.1), 100 mM KCl, 50 mM

NaCl, 5 mM MgCl$_2$, and 1% LMNG. The beads were boiled in SDS-loading buffer for 5 minutes to elute proteins co-immunoprecipitated with ACC1.

Mass Spectrometry

Immunoprecipitates were resolved by SDS–PAGE, Coomassie-stained, and the excised gel regions were submitted to the UT Southwestern Proteomics Core Facility for LC–MS/MS analysis. Protein identification and label-free quantification were performed using the core's standard Complex Mixture ID workflow, and processed datasets were used for downstream analyses.

Radiometric FA synthesis in hepatocytes

Rat primary hepatocytes were cultured in 6-well dishes coated with collagen. Where indicated, after 8 hours of culture, cells were treated with DMSO, nocodazole, or taxol (20 µM). On the day of study, the culture medium was replaced with medium containing 1 mM acetate or citrate, supplemented with 1 µCi [^{14}C]-labeled acetate or citrate, respectively, and cells were harvested after the indicated incubation times by adding 1 mL of 0.1 N NaOH. A recovery carrier (^{3}H oleic acid), 1.5 mL of ethanol, and 0.75 mL of 75% KOH were added to the harvested cells, and the mixture was autoclaved for 45 minutes on slow exhaust for saponification. After saponification, 1.5 mL of ethanol and 1.5 mL of HCl were added sequentially with vigorous vortexing. After cooling, 1.5 mL of ethanol was added, and fatty acids were extracted three times with 3 mL of hexane. The solvent was evaporated under nitrogen, and the residue was re-dissolved in 30 µL of chloroform. Samples were spotted onto plastic-backed silica gel TLC plates and resolved in a solvent system consisting of heptanes:diethyl ether:acetic acid (90:30:1, v/v/v). After visualization with iodine vapor, each FA spot was excised and radioactivity was quantified by scintillation counting. FA synthesis was calculated based on the recovered radioactivity.

Recombinant protein enzyme assays

Dephosphorylation treatment

Different recombinant proteins were pre-incubated with 20 mM HEPES (pH 7.1), 1 mM MnCl$_2$, and lambda phosphatase for 30 minutes at 30°C in 96 well plates (Corning). The same groups without lambda phosphatase were used as phosphorylated groups.

ACSS2 and ACL (LC-MS/MS)

For the ACL activity assay, protein samples from dephosphorylation treatment were incubated at

37°C in a reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA, 4 mM

sodium citrate (pH 7.1), 5 mM Mg-ATP (pH 7.1), and 0.1 mM CoA. For the ACSS2 activity assay,

protein samples were incubated at 37°C with 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA,

4 mM sodium acetate (pH 7.1), 5 mM Mg-ATP (pH 7.1), and 0.1 mM CoA. At designated time

points (0, 4, 8, 16, and 32 minutes), reactions were terminated by the addition of 8 N HCl, followed

by thorough mixing. Each sample was then immediately spiked with a [1-^{13}C]acetyl-CoA internal

standard prior to further analysis.

Samples were centrifuged at 4°C for 10 minutes and 150 µL supernatant was loaded on an Oasis

HLB 1cc-30 mg solid-phase extraction column. The column was washed with water and methanol.

The eluent was lyophilized and dissolved in 50 µL of mobile phase of which 1 µL was subjected

to LC/MS/MS analyses. Analysis was performed on an API 3200 triple quadrupole LC/MS/MS

mass spectrometer (Applied Biosystems/Sciex Instruments) in positive electrospray ionization

mode. The mass spectrometer was equipped with a Shimadzu LC-20AD liquid chromatograph

(LC) and a SIL-20ACHT auto sampler. Chromatography was performed on a reverse-phase C18

column (Waters XBridge, 150 x 2.1 mm, 3 µm) with a mobile phase consisting of water/methanol

(95:5, v/v) and 4mM dibutylamine acetate (eluent A), and water/acetonitrile (25:75, v/v) (eluent

B). Multi-reaction monitoring (MRM) was carried out for the detection of the short-chain acyl-CoAs

in standard solutions and biological samples. Acetyl-CoA was quantitated by comparison of the

individual ion peak area with that of the internal standard. The analytical data were processed by

Analyst Software (version 1.6.2).

ACC1 (radiometric)

Following pre-incubation in dephosphorylation treatment step, rACC1 was incubated at 37°C in a

reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA, 5 mM NaHCO$_3$,

5 mM Mg-ATP (pH 7.1), 0.25 mM acetyl-CoA, and 0.225 µCi [^{14}C]NaHCO$_3$ (American

Radiolabeled Chemicals). At designated time points (0, 4, 8, 16, and 32 minutes), reactions were

terminated by the addition of 8 N HCl. The mixtures were transferred to scintillation vials

(Research Products International Corp.) and air-dried completely at 100°C. Dried samples were

then dissolved in 1 mL of methanol, and [^{14}C] radioactivity was quantified by liquid scintillation

counting (Beckman Coulter) following the addition of scintillation cocktail (Research Products

International Corp.).

FAS (radiometric)

Following pre-incubation in dephosphorylation treatment step, rFAS was incubated at 37°C in a reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA, 0.1 mM acetyl-CoA, 0.3 mM malonyl-CoA, 0.5 mM NADPH, and 0.02 µCi [^{14}C]malonyl-CoA (American Radiolabeled Chemicals). At designated time points (0, 4, 8, 16, and 32 minutes), reactions were terminated by adding HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning). Ethanol and hexane were added at a volume ratio of 1:1:2 (HCl/methanol:ethanol:hexane), and the mixtures were vortexed thoroughly. Samples were centrifuged at 2,000 × g for 5 minutes at room temperature. The upper organic phase was transferred to scintillation vials and air-dried completely at RT. Dried lipids were dissolved in 1 mL of methanol, and [^{14}C] radioactivity was quantified by liquid scintillation counting (Beckman Coulter) following the addition of scintillation cocktail (Research Products International Corp.).

ACL-ACC1-FAS (radiometric)

After pre-incubation in dephosphorylation treatment step, recombinant ACL (0.8 µg), ACC1 (0.4 µg), and FAS (3.2 µg) were incubated at 37 °C in a reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg ml^{-1} BSA, 4 mM sodium citrate (pH 7.1), 5 mM NaHCO$_3$, 5 mM Mg-ATP (pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.025 µCi [1,5-^{14}C]citric acid (Revvity). At designated time points (0, 4, 8, 16, and 32 minutes), reactions were terminated by adding HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning). Ethanol and hexane were added at a volume ratio of 1:1:2 (HCl/methanol:ethanol:hexane), and the mixtures were vortexed thoroughly. Samples were centrifuged at 2,000 × g for 5 minutes at room temperature. The upper organic phase was transferred to scintillation vials and air-dried completely at RT. Dried lipids were dissolved in 1 mL of methanol, and [^{14}C] radioactivity was quantified by liquid scintillation counting (Beckman Coulter) following the addition of scintillation cocktail (Research Products International Corp.).

ACSS2-ACC1-FAS (radiometric)

Following pre-incubation in dephosphorylation treatment step, recombinant ACSS2 (0.4 µg), ACC1 (0.4 µg), and FAS (3.2 µg) were incubated at 37 °C in a reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg ml^{-1} BSA, 4 mM sodium acetate (pH 7.1), 5 mM NaHCO$_3$, 5 mM Mg-ATP (pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.0225 µCi [1,2-^{14}C]acetic acid (Revvity). At designated time points (0, 4, 8, 16, and 32 minutes), reactions were terminated by adding HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning). Ethanol and hexane were added at a volume ratio of 1:1:2 (HCl/methanol:ethanol:hexane), and the mixtures were vortexed

thoroughly. Samples were centrifuged at 2,000 × g for 5 minutes at room temperature. The upper organic phase was transferred to scintillation vials and air-dried completely at RT. Dried lipids were dissolved in 1 mL of methanol, and [^{14}C] radioactivity was quantified by liquid scintillation counting (Beckman Coulter) following the addition of scintillation cocktail (Research Products International Corp.).

Cytosol-based citrate-to-palmitate assay

Mouse livers collected after fasting-refeeding treatment were homogenized in Buffer I (20 mM HEPES, 250 mM sucrose, protease inhibitor cocktail (Millipore-Sigma), pH 7.1) at a liver-to-buffer ratio of 1:5 (w/v). Homogenates were centrifuged at 6,000 × g for 10 minutes at 4 °C. The resulting supernatant was subjected to further centrifugation at 180,000 × g for 60 minutes at 4 °C. The supernatant was then loaded on PD-10 desalting column (Cytiva) that pre-equilibrated with buffer J (20 mM HEPES, 50 mM KCl, 1 mM DTT, and protease inhibitor cocktail (Millipore-Sigma), pH 7.1) and flowed through by gravity. Liver cytosols were eluted in buffer J.

For pre-incubation, 300 µg of liver cytosol was incubated with 20 mM HEPES (pH 7.1), 1 mM MnCl$_2$, and phosphatase inhibitor cocktail at 30 °C for 15 minutes. Parallel samples without phosphatase inhibitor cocktail served as the dephosphorylated groups.

Following pre-incubation, liver cytosol samples were incubated at 37 °C for 15 minutes in a reaction mixture containing 20 mM HEPES (pH 7.1), 5 mM MgCl$_2$, 1 mM DTT, 1 mg ml^{-1} BSA, varying concentrations of sodium citrate (pH 7.1), 5 mM NaHCO$_3$, 5 mM ATP (disodium salt, pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.05 µCi [1,5-^{14}C]citric acid (Revvity). Reactions were terminated by adding 1 N HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning), followed by vortexing with ethanol and hexane. The final solvent ratio of quenching solution:ethanol:hexane was 1:1:2. Samples were centrifuged at 2,000 × g for 5 minutes at room temperature (RT), and the upper organic phase was collected, transferred to scintillation vials, and air dried completely at RT. Dried lipids were dissolved in 1 mL methanol, and [^{14}C] radioactivity was quantified by liquid scintillation counting (Beckman Coulter) following the addition of scintillation cocktail (Research Products International Corp.).

TG synthesis in liver homogenates

The livers of fasted/refed rats were homogenized in buffer K (10 mM HEPES, 1 mM EDTA, 1 mM DTT, and protease inhibitor cocktail (Millipore-Sigma), pH 7.1) at a tissue-to-buffer ratio of 1:4

(w/v). The homogenate was filtered through gauze to remove unbroken tissue and then dialyzed
against dialysis buffer (10 mM HEPES, 50 mM KCl, 1 mM DTT, 1 mM PMSF, pH 7.1) for 2 hours.

The liver homogenates were pre-incubated with various concentrations of $MgCl_2$ (0, 2.5, 5, 10, 20
mM), 4 mM of Na-citrate buffer (pH 7.1), a phosphatase inhibitor cocktail, and 50 mM phosphate
buffer (pH 7.1) for 5 minutes at 37°C. Parallel groups without phosphatase inhibitor cocktail and
phosphate buffer served as dephosphorylated controls. After pre-incubation, samples were
incubated for 30 minutes at 37 °C in a reaction mixture containing 1 mM DTT, 1 mg/mL BSA,
10 mM oleoyl-CoA, 100 mM glycerol 3-phosphate, and 0.2 µCi [^{14}C]glycerol 3-phosphate
(American Radiolabeled Chemicals). Reactions were terminated by adding 1 N HCl/methanol
(6:4, v/v) in 16 × 100 mm glass tubes (Corning). The mixtures were then vortexed with phosphate-
buffered saline (PBS) and Folch reagent (chloroform:methanol = 2:1, v/v). The final solvent ratio
of HCl/methanol:PBS:Folch was 1:1:3. Samples were centrifuged at 2,000 × g for 5 minutes at
room temperature (RT), and the lower organic phases were transferred to new glass tubes and
air dried completely at RT. Dried lipids were reconstituted in 50 µL chloroform and vortexed.
Extracted lipids, along with a TG marker, were loaded onto thin-layer chromatography (TLC)
plates. Plates were developed in a TLC chamber saturated with a solvent system of
hexane:diethyl ether:acetic acid (80:20:1, v/v/v). TG dots were visualized using iodine vapor,
identified by comparison with the TG marker. TG dots were excised from the plates and
transferred to scintillation vials. Scintillation cocktail (Research Products International Corp.) was
added to each vial, followed by vortexing. [^{14}C] radioactivity was quantified by liquid scintillation
counting (Beckman Coulter).

Enzyme Synergy Assays (Recombinant ± Cytosol)

For dephosphorylation treatment, a mixture containing rACL (0.5 µg), rACC1 (0.25 µg), and rFAS
(2 µg) or an equal volume of protein purification buffer (Buffer E) (used for the cytosol only groups)
were incubated in a 25 µL total volume system with λ-phosphatase in the presence of 1 mM $MnCl_2$
at 30°C for 20 minutes. The resulting 22 µL reaction was combined with 20 µL of liver cytosol
(300 µg) or cytosol preparation buffer (Buffer J) (used for the recombinant protein only groups) in
a final volume of 50 µL and incubated at 30°C with 5 mM $MgCl_2$ for an additional 20 minutes.

To maintain proteins in a phosphorylated state, the same protein mixture or Buffer E was
incubated under identical conditions with phosphatase inhibitor cocktail instead of λ-phosphatase.
The resulting 22 µL reaction was then mixed with 20 µL of liver cytosol (300 µg) or Buffer J and
incubated in a 50 µL system containing 5 mM $MgCl_2$, phosphatase inhibitor cocktail, and 50 mM

phosphate buffer (pH 7.1) at 30°C for 20 minutes. Following the pre-incubation, samples were mixed with reaction buffer containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA, 4 mM sodium citrate (pH 7.1), 5 mM $NaHCO_3$, 5 mM Mg-ATP (pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.025 µCi [1,5-^{14}C]citric acid (Revvity) in a total volume of 75 µL. The reactions were incubated at 37°C for 15 minutes and terminated by adding 0.5 mL of cold methanol in 16 × 100 mm glass tubes (Corning). Next, 1.5 mL of methyl tert-butyl ether (MTBE) was added, and the samples were vortexed thoroughly and left to stand at RT for 5 minutes. Then, 0.5 mL of ddH_2O was added, followed by another round of vortexing and a 5-minute incubation at RT. Phase separation was achieved by centrifugation at 2,000 × g for 5 minutes at RT. The upper organic phase was transferred to scintillation vials and air-dried completely at RT. Dried lipids were dissolved in 1 mL of methanol, and [^{14}C] radioactivity was quantified by liquid scintillation counting (Beckman Coulter) after the addition of scintillation cocktail (Research Products International Corp.).

Phosphatase inhibitor cocktail preparation

The phosphatase inhibitor cocktail was prepared fresh using the following recipe: sodium fluoride (200 mM), imidazole (200 mM), sodium molybdate (115 mM), sodium orthovanadate (200 mM), sodium tartrate dihydrate (400 mM), sodium pyrophosphate (100 mM), and β-glycerophosphate (100 mM) in ddH_2O (final volume 20 mL). Sodium orthovanadate was prepared separately by dissolving 1.84 g of powder in 50 mL ddH_2O, adjusting the pH to 10, briefly boiling in a microwave (10 sec), cooling on ice to room temperature, and repeating the cycle three times. The solution was adjusted to 50 mL, aliquoted, and stored at –20°C until use.

Quantification and statistical analysis

Sample sizes (n) are indicated in the figure legends. Technical replicates are also noted in the figure legends when applicable. Unless otherwise stated, data are presented as mean ± SEM. Statistical analyses were performed using GraphPad Prism 10. The statistical test and model used for each experiment are specified in the corresponding figure legend. A significance threshold of $\alpha = 0.05$ was used.

For co-localization analyses, images were quantified using Fiji/ImageJ (Coloc2). Pearson's correlation coefficients were reported as values above the Costes threshold. Thresholded Manders' coefficients (tM1 and tM2), Costes significance testing, and 2D intensity histograms were generated using Coloc2 and are provided in the Supplemental Information.

Supplemental item titles

Figure S1. Representative regions of interest (ROIs) and pixel-intensity scatter plots used for colocalization analysis of ACC1, S14, and MIG12, related to Figure 1.

Figure S2. Sequential formation of the ACC1-containing lipogenic complex around LDs, related to Figure 2.

Figure S3. Electron microscope images, related to Figure 3.

Figure S4. Phosphorylated ACC levels in liver lysates from rats refed the FFD, related to Figure 4.

Figure S5. Physical association of ACC1 and ACC2 and regulation of TG synthesis by phosphorylation in an $MgCl_2$-dependent manner, related to Figure 5.

Figure S6. Lipogenic metabolon formation facilitates FA production, related to Figure 6.

Figure S7. DAVID functional annotation of ACC1-dependent interactors from ACC1, MIG12, and S14 immunoprecipitates, related to Figure 7.

Supplemental Table 1. Pearson's and Manders' colocalization coefficients together with Costes significance statistics for ACC1–MIG12, S14–MIG12, and S14–ACC1, related to Figure 1.

Supplemental Table 2. Quantitative co-localization analysis of ACC1 with cytosolic lipogenic enzymes in primary hepatocytes, related to Figure 2.

Supplemental Table 3. Original data of MS analysis of ACC1-dependent interactors from ACC1, MIG12, and S14 immunoprecipitates, related to Figure 7.

1006

References

1.   Sanders, F.W., and Griffin, J.L. (2016). De novo lipogenesis in the liver in health and disease: more than just a shunting yard for glucose. Biol Rev Camb Philos Soc *91*, 452-468. 10.1111/brv.12178.
2.   Horton, J.D., Goldstein, J.L., and Brown, M.S. (2002). SREBPs:activators of the complete program of cholesterol and fatty acid synthesis in the liver. J. Clin. Invest. *109*, 1125-1131.
3.   Samuel, V.T., and Shulman, G.I. (2016). The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. J. Clin. Invest. *126*, 12-22. .

4.   Miao, T., Kim, J., Kang, P., Fujiwara, H., Hsu, F.F., and Bai, H. (2022). Acetyl-CoA-mediated
autoacetylation of fatty acid synthase as a metabolic switch of de novo lipogenesis in
Drosophila. Proc Natl Acad Sci U S A *119*, e2212220119. 10.1073/pnas.2212220119.
5.   Lin, H.P., Cheng, Z.L., He, R.Y., Song, L., Tian, M.X., Zhou, L.S., Groh, B.S., Liu, W.R., Ji, M.B.,
Ding, C., et al. (2016). Destabilization of Fatty Acid Synthase by Acetylation Inhibits De Novo
Lipogenesis and Tumor Cell Growth. Cancer Res *76*, 6924-6936. 10.1158/0008-5472.CAN-
16-1597.
6.   Hellerstein, M.K. (1999). De novo lipogenesis in humans: metabolic and regulatory aspects.
Eur. J. Clin. Nutr. *53*, S53-65.
7.   Parks, E.J., Krauss, R.M., Christiansen, M.P., Neese, R.A., and Hellerstein, M.K. (1999).
Effects of a low-fat, high-carbohydrate diet on VLDL-triglyceride assembly, production, and
clearance. J Clin Invest *104*, 1087-1096. 10.1172/JCI6572.
8.   Schwarz, J.M., Neese, R.A., Turner, S., Dare, D., and Hellerstein, M.K. (1995). Short-term
alterations in carbohydrate energy intake in humans. Striking effects on hepatic glucose
production, de novo lipogenesis, lipolysis, and whole-body fuel selection. J Clin Invest *96*,
2735-2743. 10.1172/JCI118342.
9.   Donnelly, K.L., Smith, C.I., Schwarzenberg, S.J., Jessurun, J., Boldt, M.D., and Parks, E.J.
(2005). Sources of fatty acids stored in liver and secreted via lipoproteins in patients with
nonalcoholic fatty liver disease. J. Clin. Invest. *115*, 1343-1351.
10.   Ferre, P., and Foufelle, F. (2010). Hepatic steatosis: a role for de novo lipogenesis and the
transcription factor SREBP-1c. Diabetes Obes Metab *12 Suppl 2*, 83-92. 10.1111/j.1463-
1326.2010.01275.x.
11.   Lambert, J.E., Ramos-Roman, M.A., Browning, J.D., and Parks, E.J. (2014). Increased de novo
lipogenesis is a distinct characteristic of individuals with nonalcoholic fatty liver disease.
Gastroenterology *146*, 726-735. 10.1053/j.gastro.2013.11.049.
12.   Shimomura, I., Shimano, H., Korn, B.S., Bashmakov, Y., and Horton, J.D. (1998). Nuclear
sterol regulatory element-binding proteins activate genes responsible for the entire program
of unsaturated fatty acid biosynthesis in transgenic mouse liver. J. Biol. Chem. *273*, 35299-
35306.
13.   Shimomura, I., Bashmakov, Y., and Horton, J.D. (1999). Increased levels of nuclear SREBP-
1c associated with fatty livers in two mouse models of diabetes mellitus. J. Biol. Chem. *274*,
30028-30032.
14.   Hudgins, L.C., Hellerstein, M., Seidman, C., Neese, R., Diakun, J., and Hirsch, J. (1996).
Human fatty acid synthesis is stimulated by a eucaloric low fat, high carbohydrate diet. J Clin
Invest *97*, 2081-2091. 10.1172/JCI118645.
15.   Eslam, M., Newsome, P.N., Sarin, S.K., Anstee, Q.M., Targher, G., Romero-Gomez, M.,
Zelber-Sagi, S., Wai-Sun Wong, V., Dufour, J.F., Schattenberg, J.M., et al. (2020). A new
definition for metabolic dysfunction-associated fatty liver disease: An international expert
consensus statement. J Hepatol *73*, 202-209. 10.1016/j.jhep.2020.03.039.
16.   Cohen, J.C., Horton, J.D., and Hobbs, H.H. (2011). Human fatty liver disease: old questions
and new insights. Science *332*, 1519-1523. 10.1126/science.1204265.
17.   Eslam, M., Sanyal, A.J., and George, J. (2020). MAFLD: A Consensus-Driven Proposed
Nomenclature for Metabolic Associated Fatty Liver Disease. Gastroenterology *158*, 1999-
2014.e1991. 10.1053/j.gastro.2019.11.312.
18.   Browning, J.D., Szczepaniak, L.S., Dobbins, R., Nuremberg, P., Horton, J.D., Cohen, J.C.,
Grundy, S.M., and Hobbs, H.H. (2004). Prevalence of hepatic steatosis in an urban
population in the United States: Impact of ethnicity. Hepatology *40*, 1387-1395.

19.   Shimano, H., Horton, J.D., Hammer, R.E., Shimomura, I., Brown, M.S., and Goldstein, J.L.
1063   (1996). Overproduction of cholesterol and fatty acids causes massive liver enlargement in
1064   transgenic mice expressing truncated SREBP-1a. J. Clin. Invest. *98*, 1575-1584.

1065   20.   Horton, J.D., Bashmakov, Y., Shimomura, I., and Shimano, H. (1998). Regulation of sterol
1066   regulatory element binding proteins in livers of fasted and refed mice. Proc. Natl. Acad. Sci.
1067   U. S. A. *95*, 5987-5992.

1068   21.   Horton, J.D., Shimomura, I., Brown, M.S., Hammer, R.E., Goldstein, J.L., and Shimano, H.
1069   (1998). Activation of cholesterol synthesis in preference to fatty acid synthesis in liver and
1070   adipose tissue of transgenic mice overproducing sterol regulatory element-binding protein-
1071   2. J. Clin. Invest. *101*, 2331-2339.

1072   22.   Horton, J.D., Shimano, H., Hamilton, R.L., Brown, M.S., and Goldstein, J.L. (1999). Disruption
1073   of LDL receptor gene in transgenic SREBP-1a mice unmasks hyperlipidemia resulting from
1074   production of lipid-rich VLDL. J. Clin. Invest. *103*, 1067-1076.

1075   23.   Matsuda, M., Korn, B.S., Hammer, R.E., Moon, Y.A., Komuro, R., Horton, J.D., Goldstein, J.L.,
1076   Brown, M.S., and Shimomura, I. (2001). SREBP cleavage-activating protein (SCAP) is required
1077   for increased lipid synthesis in liver induced by cholesterol deprivation and insulin elevation.
1078   Genes Dev. *15*, 1206-1216.

1079   24.   Moon, Y.A., Liang, G., Xie, X., Frank-Kamenetsky, M., Fitzgerald, K., Koteliansky, V., Brown,
1080   M.S., Goldstein, J.L., and Horton, J.D. (2012). The Scap/SREBP pathway is essential for
1081   developing diabetic fatty liver and carbohydrate-induced hypertriglyceridemia in animals.
1082   Cell Metab. *15*, 240-246. 10.1016/j.cmet.2011.12.017.

1083   25.   Abu-Elheiga, L., Brinkley, W.R., Zhong, L., Chirala, S.S., Woldegiorgis, G., and Wakil, S.J.
1084   (2000). The subcellular localization of acetyl-CoA carboxylase 2. Proc. Natl. Acad. Sci. U. S.
1085   A. *97*, 1444-1449.

1086   26.   Man, W.C., Miyazaki, M., Chu, K., and Ntambi, J.M. (2006). Membrane topology of mouse
1087   stearoyl-CoA desaturase 1. J Biol Chem *281*, 1251-1260. 10.1074/jbc.M508733200.

1088   27.   Moon, Y.-A., Shah, N.A., Mohapatra, S., Warrington, J.A., and Horton, J.D. (2001).
1089   Identification of a mammalian long chain fatty acyl elongase regulated by sterol regulatory
1090   element-binding proteins. J. Biol. Chem. *276*, 45358-45366.

1091   28.   Gonzalez-Baró, M.R., Granger, D.A., and Coleman, R.A. (2001). Mitochondrial Glycerol
1092   Phosphate Acyltransferase Contains Two Transmembrane Domains with the Active Site in
1093   the N-terminal Domain Facing the Cytosol. Journal of Biological Chemistry *276*, 43182-
1094   43188. 10.1074/jbc.M107885200.

1095   29.   P.A., S. (1972). Is there an organization of Krebs cycle enzymes in the mitochondrial matrix?
1096   In: Mehlman M.A., Hanson R.W., editors. Energy Metabolism and the Regulation of Metabolic
1097   Processes in Mitochondria. Academic Press; New York, pp. 79–91.

1098   30.   P.A., S. (1985). The metabolon. Trends Biochem. Sci., 10:109–110.

1099   31.   Bulutoglu, B., Garcia, K.E., Wu, F., Minteer, S.D., and Banta, S. (2016). Direct Evidence for
1100   Metabolon Formation and Substrate Channeling in Recombinant TCA Cycle Enzymes. ACS
1101   Chem Biol *11*, 2847-2853. 10.1021/acschembio.6b00523.

1102   32.   Robinson, J.B., Jr., Inman, L., Sumegi, B., and Srere, P.A. (1987). Further characterization of
1103   the Krebs tricarboxylic acid cycle metabolon. J Biol Chem *262*, 1786-1790.

1104   33.   Kohnhorst, C.L., Kyoung, M., Jeon, M., Schmitt, D.L., Kennedy, E.L., Ramirez, J., Bracey, S.M.,
1105   Luu, B.T., Russell, S.J., and An, S. (2017). Identification of a multienzyme complex for glucose
1106   metabolism in living cells. J Biol Chem *292*, 9191-9203. 10.1074/jbc.M117.783050.

1107   34.   Colbert, C.L., Kim, C.W., Moon, Y.A., Henry, L., Palnitkar, M., McKean, W.B., Fitzgerald, K.,
1108   Deisenhofer, J., Horton, J.D., and Kwon, H.J. (2010). Crystal structure of Spot 14, a modulator
1109   of fatty acid synthesis. Proc Natl Acad Sci U S A *107*, 18820-18825. 1012736107 [pii]

10.1073/pnas.1012736107.

35.   Jump, D.B., and Oppenheimer, J.H. (1985). High basal expression and 3,5,3'-triiodothyronine
regulation of messenger ribonucleic acid S14 in lipogenic tissues. Endocrinology *117*, 2259-
2266. 10.1210/endo-117-6-2259.

36.   Horton, J.D., Shah, N.A., Warrington, J.A., Anderson, N.N., Park, S.W., Brown, M.S., and
Goldstein, J.L. (2003). Combined analysis of oligonucleotide microarray data from
transgenic and knockout mice identifies direct SREBP target genes. Proc. Natl. Acad. Sci. U.
S. A. *100*, 12027-12032.

37.   Mater, M.K., Thelen, A.P., Pan, D.A., and Jump, D.B. (1999). Sterol response element-binding
protein 1c (SREBP1c) is involved in the polyunsaturated fatty acid suppression of hepatic S14
gene transcription. J. Biol. Chem. *274*, 32725-32732.

38.   Ma, L., Tsatsos, N.G., and Towle, H.C. (2005). Direct role of ChREBP·Mlx in regulating hepatic
glucose-responsive genes. J. Biol. Chem. *280*, 12019-12027. 10.1074/jbc.M413063200.

39.   Seelig, S., Liaw, C., Towle, H.C., and Oppenheimer, J.H. (1981). Thyroid hormone attenuates
and augments hepatic gene expression at a pretranslational level. Proc. Natl. Acad. Sci. U.
S. A. *78*, 4733-4737.

40.   Zhu, Q., Anderson, G.W., Mucha, G.T., Parks, E.J., Metkowski, J.K., and Mariash, C.N. (2005).
The Spot 14 protein is required for de novo lipid synthesis in the lactating mammary gland.
Endocrinology *146*, 3343-3350.

41.   Kim, C.W., Moon, Y.A., Park, S.W., Cheng, D., Kwon, H.J., and Horton, J.D. (2010). Induced
polymerization of mammalian acetyl-CoA carboxylase by MIG12 provides a tertiary level of
regulation of fatty acid synthesis. Proc. Natl. Acad. Sci. U. S. A. *107*, 9626-9631.

42.   Ha, J., Daniel, S., Broyles, S.S., and Kim, K.H. (1994). Critical phosphorylation sites for acetyl-
CoA carboxylase activity. J Biol Chem *269*, 22162-22168.

43.   Locke, G.A., Cheng, D., Witmer, M.R., Tamura, J.K., Haque, T., Carney, R.F., Rendina, A.R.,
and Marcinkeviciene, J. (2008). Differential activation of recombinant human acetyl-CoA
carboxylases 1 and 2 by citrate. Arch. Biochem. Biophys. *475*, 72-79. S0003-9861(08)00196-
3 [pii]

10.1016/j.abb.2008.04.011 [doi].

44.   Kahn, B.B., Alquier, T., Carling, D., and Hardie, D.G. (2005). AMP-activated protein kinase:
Ancient energy gauge provides clues to modern understanding of metabolism. Cell
Metabolism *1*, 15-25. DOI: 10.1016/j.cmet.2004.12.003.

45.   Witters, L.A., Watts, T.D., Daniels, D.L., and Evans, J.L. (1988). Insulin stimulates the
dephosphorylation and activation of acetyl-CoA carboxylase. Proc. Natl. Acad. Sci. U. S. A.
85, 5473-5477.

46.   Berglund, E.D., Lustig, D.G., Baheza, R.A., Hasenour, C.M., Lee-Young, R.S., Donahue, E.P.,
Lynes, S.E., Swift, L.L., Charron, M.J., Damon, B.M., and Wasserman, D.H. (2011). Hepatic
glucagon action is essential for exercise-induced reversal of mouse fatty liver. Diabetes *60*,
2720-2729. 10.2337/db11-0455.

47.   Munday, M.R., Milic, M.R., Takhar, S., Holness, M.J., and Sugden, M.C. (1991). The short-term
regulation of hepatic acetyl-CoA carboxylase during starvation and re-feeding in the rat.
Biochem J *280 (Pt 3)*, 733-737. 10.1042/bj2800733.

48.   Berglund, E.D., Lee-Young, R.S., Lustig, D.G., Lynes, S.E., Donahue, E.P., Camacho, R.C.,
Meredith, M.E., Magnuson, M.A., Charron, M.J., and Wasserman, D.H. (2009). Hepatic energy
state is regulated by glucagon receptor signaling in mice. J Clin Invest *119*, 2412-2422.
10.1172/jci38650.

49. Kimball, S.R., Siegfried, B.A., and Jefferson, L.S. (2004). Glucagon represses signaling
through the mammalian target of rapamycin in rat liver by activating AMP-activated protein
kinase. J Biol Chem *279*, 54103-54109. 10.1074/jbc.M410755200.
50. Ragolia, L., and Begum, N. (1998). Protein phosphatase-1 and insulin action. Mol Cell
Biochem *182*, 49-58.
51. Kim, K.H. (1983). Regulation of acetyl-CoA carboxylase. Curr Top Cell Regul *22*, 143-176.
10.1016/b978-0-12-152822-5.50009-9.
52. Kim, K.H., Lopez-Casillas, F., Bai, D.H., Luo, X., and Pape, M.E. (1989). Role of reversible
phosphorylation of acetyl-CoA carboxylase in long-chain fatty acid synthesis. FASEB J *3*,
2250-2256. 10.1096/fasebj.3.11.2570725.
53. Fediuc, S., Gaidhu, M.P., and Ceddia, R.B. (2006). Regulation of AMP-activated protein
kinase and acetyl-CoA carboxylase phosphorylation by palmitate in skeletal muscle cells. J
Lipid Res *47*, 412-420. 10.1194/jlr.M500438-JLR200.
54. Wang, Y., Viscarra, J., Kim, S.-J., and Sul, H.S. (2015). Transcriptional regulation of hepatic
lipogenesis. Nature reviews. Molecular cell biology *16*, 678-689. 10.1038/nrm4074.
55. Umlauf, E., Csaszar, E., Moertelmaier, M., Schuetz, G.J., Parton, R.G., and Prohaska, R.
(2004). Association of stomatin with lipid bodies. J Biol Chem *279*, 23699-23709.
10.1074/jbc.M310546200.
56. Liu, P., Ying, Y., Zhao, Y., Mundy, D.I., Zhu, M., and Anderson, R.G. (2004). Chinese hamster
ovary K2 cell lipid droplets appear to be metabolic organelles involved in membrane traffic.
J Biol Chem *279*, 3787-3792. 10.1074/jbc.M311945200.
57. Bartz, R., Zehmer, J.K., Zhu, M., Chen, Y., Serrero, G., Zhao, Y., and Liu, P. (2007). Dynamic
activity of lipid droplets: protein phosphorylation and GTP-mediated protein translocation. J
Proteome Res *6*, 3256-3265. 10.1021/pr070158j.
58. Beller, M., Riedel, D., Jansch, L., Dieterich, G., Wehland, J., Jackle, H., and Kuhnlein, R.P.
(2006). Characterization of the Drosophila lipid droplet subproteome. Molecular & cellular
proteomics : MCP *5*, 1082-1094. 10.1074/mcp.M600011-MCP200.
59. Wan, H.C., Melo, R.C., Jin, Z., Dvorak, A.M., and Weller, P.F. (2007). Roles and origins of
leukocyte lipid bodies: proteomic and ultrastructural studies. FASEB J *21*, 167-178.
10.1096/fj.06-6711com.
60. Lyubarev, A.E., and Kurganov, B.I. (1989). Supramolecular organization of tricarboxylic acid
cycle enzymes. Biosystems *22*, 91-102. 10.1016/0303-2647(89)90038-5.
61. Kurganov B. I, L.A.E. (1988). Hypothetical structure of the complex of glycolytic enzymes
(glycolytic metabolon), formed on the membrane of erythrocytes. Molek. Biologia. *V.22,
No.6*, p. 1605–1613.
62. Graham, J.W., Williams, T.C., Morgan, M., Fernie, A.R., Ratcliffe, R.G., and Sweetlove, L.J.
(2007). Glycolytic enzymes associate dynamically with mitochondria in response to
respiratory demand and support substrate channeling. Plant Cell *19*, 3723-3738.
10.1105/tpc.107.053371.
63. Srere, P.A. (1974). Controls of citrate synthase activity. Life Sci *15*, 1695-1710.
10.1016/0024-3205(74)90172-6.
64. Webb, B.A., Dosey, A.M., Wittmann, T., Kollman, J.M., and Barber, D.L. (2017). The glycolytic
enzyme phosphofructokinase-1 assembles into filaments. J Cell Biol *216*, 2305-2313.
10.1083/jcb.201701084.
65. Kastritis, P.L., O'Reilly, F.J., Bock, T., Li, Y., Rogon, M.Z., Buczak, K., Romanov, N., Betts, M.J.,
Bui, K.H., Hagen, W.J., et al. (2017). Capturing protein communities by structural proteomics
in a thermophilic eukaryote. Mol Syst Biol *13*, 936. 10.15252/msb.20167412.

66. Suresh, H.G., da Silveira Dos Santos, A.X., Kukulski, W., Tyedmers, J., Riezman, H., Bukau,
B., and Mogk, A. (2015). Prolonged starvation drives reversible sequestration of lipid
biosynthetic enzymes and organelle reorganization in Saccharomyces cerevisiae. Mol Biol
Cell *26*, 1601-1615. 10.1091/mbc.E14-11-1559.
67. Wijesinghe, K.M., Kim, C.W., Schad, E.O., Li, S., Chen, S., Takeshima, E., Khandwala, C.B.,
Tillo, D., Lebensohn, A.M., Olzmann, J.A., et al. (2025). Plasma membrane accessible
cholesterol is regulated by ACC1 and lipid droplets. bioRxiv. 10.1101/2025.08.21.671640.
68. Zhou, F., Zhang, Y., Zhu, Y., Zhou, Q., Shi, Y., and Hu, Q. (2024). Filament structures unveil
the dynamic organization of human acetyl-CoA carboxylase. Sci Adv *10*, eado4880.
10.1126/sciadv.ado4880.
69. Hunkeler, M., Hagmann, A., Stuttfeld, E., Chami, M., Guri, Y., Stahlberg, H., and Maier, T.
(2018). Structural basis for regulation of human acetyl-CoA carboxylase. Nature *558*, 470-
474. 10.1038/s41586-018-0201-4.
70. Møller, B.L. (2010). Plant science. Dynamic metabolons. Science *330*, 1328-1329.
10.1126/science.1194971.
71. Kim, C.W., Addy, C., Kusunoki, J., Anderson, N.N., Deja, S., Fu, X., Burgess, S.C., Li, C.,
Ruddy, M., Chakravarthy, M., et al. (2017). Acetyl CoA Carboxylase Inhibition Reduces
Hepatic Steatosis but Elevates Plasma Triglycerides in Mice and Humans: A Bedside to
Bench Investigation. Cell Metab *26*, 394-406. e396. 10.1016/j.cmet.2017.07.009.
72. Kim, C.W., Moon, Y.A., Park, S.W., Cheng, D., Kwon, H.J., and Horton, J.D. (2010). Induced
polymerization of mammalian acetyl-CoA carboxylase by MIG12 provides a tertiary level of
regulation of fatty acid synthesis. Proc Natl Acad Sci U S A *107*, 9626-9631.
10.1073/pnas.1001292107.
73. Shimomura, I., Hammer, R.E., Ikemoto, S., Brown, M.S., and Goldstein, J.L. (1999). Leptin
reverses insulin resistance and diabetes mellitus in mice with congenital lipodystrophy.
Nature *401*, 73-76.
74. Lagace, T.A., Curtis, D.E., Garuti, R., McNutt, M.C., Park, S.W., Prather, H.B., Anderson, N.N.,
Ho, Y.K., Hammer, R.E., and Horton, J.D. (2006). Secreted PCSK9 decreases the number of
LDL receptors in hepatocytes and in livers of parabiotic mice. J Clin Invest *116*, 2995-3005.
10.1172/jci29383.
75. Herz, J., Kowal, R.C., Ho, Y.K., Brown, M.S., and Goldstein, J.L. (1990). Low density
lipoprotein receptor-related protein mediates endocytosis of monoclonal antibodies in
cultured cells and rabbit liver. J Biol Chem *265*, 21355-21362.
76. Engelking, L.J., Kuriyama, H., Hammer, R.E., Horton, J.D., Brown, M.S., Goldstein, J.L., and
Liang, G. (2004). Overexpression of Insig-1 in the livers of transgenic mice inhibits SREBP
processing and reduces insulin-stimulated lipogenesis. J. Clin. Invest. *113*, 1168-1175.
77. Yamamoto, A., and Masaki, R. (2010). Pre-embedding nanogold silver and gold
intensification. Methods Mol Biol *657*, 225-235. 10.1007/978-1-60761-783-9_18.
78. Ding, Y., Zhang, S., Yang, L., Na, H., Zhang, P., Zhang, H., Wang, Y., Chen, Y., Yu, J., Huo, C.,
et al. (2013). Isolating lipid droplets from multiple species. Nat Protoc *8*, 43-51.
10.1038/nprot.2012.142.

1244

1245

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7