Identification and Regulation of a Hepatic Lipogenic Metabolon

Xiaotong Zhu[1, 2, 6], William B. McKean[3, 4], Tam Tran[1, 5], Xiaorong Fu[1, 2], Jay D. Horton[1, 2, 3, 7], Chai-Wan Kim[1, 2, 6]

Center for Human Nutrition, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046

Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046

Department of Molecular Genetics, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046

START Center for Cancer Research, Mountain Region, West Valley City, UT 84119

Center of Excellence for Leukemia Studies, Department of Pathology, St. Jude Children's Research Hospital, Memphis, TN 38105

These authors contributed equally

Corresponding author

SUMMARY

De novo lipogenesis (DNL) plays a key role in the excessive fat accumulation present in Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD). Most mechanistic studies and experimental strategies for improving hepatic steatosis in MASLD have focused on the transcriptional regulation of enzymes involved in DNL and triglyceride (TG) synthesis. Here, we provide evidence for a post-translational mechanism that enhances fatty acid (FA) and TG synthesis through the assembly of a multi-protein lipogenic metabolon in liver. Under anabolic conditions, acetyl-CoA carboxylase 1 (ACC1) interacts with additional key enzymes in the DNL and TG synthesis pathway. Immunofluorescence and electron microscopy reveal that this lipogenic metabolon localizes around lipid droplets (LDs) and in proximity to mitochondria and LD interfaces in the anabolic state. The formation of the lipogenic metabolon facilitates the efficient transfer of FA synthesis intermediates to enhance lipogenic flux. These findings uncover a new nutrient-responsive, post-translational regulatory mechanism for hepatic

29 lipogenesis and highlight the lipogenic metabolon as a potential therapeutic target for metabolic
30 liver diseases.

Keywords

32 Liver metabolism, Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD), lipogenic
33 metabolon, *de novo* lipogenesis, fatty acid synthesis, triglyceride synthesis, ACC1, metabolic
34 diseases, post-translational regulation

INTRODUCTION

De novo lipogenesis (DNL) is the process in which fatty acids (FAs) are synthesized in cells
from acetyl-CoA, primarily derived from carbohydrate metabolism [1]. Key transcriptional
regulators of DNL, including SREBP-1c and ChREBP, are tightly regulated by hormones such
as insulin [2,3] and glucagon [2], and through post-translational modifications of the DNL enzymes
including phosphorylation [2] and acetylation [4,5]. In a healthy liver, DNL contributes to less than
5% of total hepatic triglycerides (TGs) [6-8]. However, in Metabolic Dysfunction-Associated
Steatotic Liver Disease (MASLD), this contribution rises to 25-30% [9-11], due to insulin resistance,
which leads to an increase in hepatic FA and TG synthesis [2,12-14]. Excessive TG accumulation in
liver is associated with progression to metabolic-associated steatohepatitis (MASH), cirrhosis,
and hepatocellular carcinoma [15-17]. With over one-third of U.S. adults affected by MASLD [18], it is
crucial to elucidate the mechanisms driving aberrant hepatic FA synthesis in insulin-resistant
states to identify potential therapeutic targets for FA-driven metabolic disorders.

Our previous studies have shown that the excessive activation of SREBP-1c initiates the
transcription of genes encoding the enzymes necessary for FA synthesis, as well as the enzyme
catalyzing the first step in glycerolipid synthesis [12,13]. This transcriptional program is a major
driver of elevated FA and TG synthesis in insulin-resistant states [2,12,13], subsequently leading to
hepatic steatosis, increased VLDL secretion, and hypertriglyceridemia [13,19-24]. The requirement
of SREBP activation for the development of hepatic steatosis and hypertriglyceridemia was
demonstrated by studies in which SCAP, a protein required for SREBP activation, was deleted
in obese, insulin resistant mice (*ob/ob* mice) that normally have very high levels of hepatic FA
and TG synthesis. SCAP deletion completely normalized rates of FA synthesis, liver TGs, VLDL
secretion, and plasma TG levels [24] in *ob/ob* mice and hamsters. These results demonstrate that
enhanced DNL is a major driver of steatosis in insulin-resistant livers.

FA and TG synthesis requires sequential enzymatic reactions distributed across multiple

subcellular compartments. ATP citrate lyase (ACL) produces acetyl-CoA from citrate, acetyl-

CoA synthetase 2 (ACSS2) synthesizes acetyl-CoA from acetate, acetyl-CoA carboxylases

(ACC) 1 and 2 convert acetyl-CoA to malonyl-CoA, and fatty acid synthase (FAS) uses malonyl-

CoA and acetyl-CoA to generate palmitate (C16:0). These enzymes (except for ACC2 that is

predominantly localized to the outer mitochondrial membrane) are predominantly localized in

the cytosol [25]. Palmitate can then be further desaturated by stearoyl-CoA desaturase (SCD1) or

elongated by ELOVL fatty acid elongase 6 (ELOVL6) before further desaturation, but unlike the

enzymes involved in the synthesis of palmitate, these proteins reside in the endoplasmic

reticulum (ER) [26,27]. Additionally, the first enzyme in glycerolipid synthesis, glycerol-3-phosphate

acyltransferase 1 (GPAM), which acylates glycerol-3-phosphate to initiate TG and phospholipid

synthesis is located in the mitochondria [28]. How the FA and glycerolipid synthesis intermediates

are efficiently transferred among these spatially distributed enzymes to sustain continuous and

efficient lipid synthesis remains unclear.

One conceptual framework that could address this challenge is the metabolon hypothesis. In

1970, A.M. Kuzin first introduced the concept of physical enzyme–enzyme complexes. This idea

gained momentum in 1972 when P.A. Srere applied it to enzymes of the citric acid cycle,

highlighting their potential functional interactions [29]. Srere further expanded on this idea by

coining the term "metabolon" during a 1985 lecture in Debrecen, Hungary. He defined it as "a

supramolecular complex of sequential metabolic enzymes and cellular structural elements" [30],

emphasizing the functional coordination within cellular metabolic pathways. This concept has

been applied to explain the spatial organization of metabolic pathways such as the citric acid

cycle [31,32] and glycolysis [33], prompting us to examine whether hepatic FA and TG synthesis

might also be coordinated through spatial organization of lipogenic enzymes.

Here, we show that under anabolic conditions—such as high-carbohydrate feeding and insulin

resistance—FA and TG synthesis enzymes assemble into a functional lipogenic metabolon.

This complex enhances lipogenic flux by coordinating the sequential enzymatic reactions

required to synthesize the end product. This metabolon, including accessory proteins MIG12

and SPOT14 (S14) [34], predominantly localizes around lipid droplets (LDs) in primary

hepatocytes. Together, these findings reveal a dynamic multi-enzyme complex that facilitates

FA and TG synthesis and provides new insights into the spatial organization and coordinated

regulation of hepatic lipid synthesis.

RESULTS

Identification of the lipogenic complex in liver

The first indication that a lipogenic complex might exist grew out of investigations focused on determining the function of a poorly characterized protein, S14. S14 is highly expressed in lipogenic tissues [35], and its gene transcription is regulated by SREBP-1c [36,37], carbohydrate-responsive element-binding protein (ChREBP) [38], and thyroid hormone receptor (TR) [39], which suggests that it may play a role in lipogenesis. Deletion of S14 in mice reduced lipid synthesis in liver [34] and lactating mammary glands [40], although its exact function in lipogenesis has not been elucidated. To further define the function of S14, we used an adenovirus that expressed S14 with a Flag tag in mice and carried out immunoprecipitation of S14 from liver lysates to identify potential interacting proteins. LC/MS analysis revealed that MIG12, ACC1, FAS, and β-tubulin all co-immunoprecipitated with S14 (Figure 1A). MIG12, a protein we previously showed that promotes the polymerization of ACC1 and ACC2 [41], shares 32% sequence identity with S14 [41] and can form a heterodimer with S14 [34]. To further confirm that the association of these enzymes co-immunoprecipitated with S14 reflects physical interaction in cells, we performed immunofluorescence staining in primary hepatocytes isolated from rats refed a fat-free, high-carbohydrate diet (FFD). ACC1, MIG12, and S14 prominently co-localized around LD–associated ER regions. Quantitative analysis using Pearson's correlation (above Costes threshold) demonstrated strong spatial overlap among the three enzyme pairs, with ACC1–S14 (R = 0.73), ACC1–MIG12 (R = 0.52), and S14–MIG12 (R = 0.52) (Supplemental Table 1). These data indicate that ACC1, S14, and MIG12 occupy shared subcellular domains under conditions of high lipogenesis, supporting the existence of a coordinated lipogenic enzyme complex in hepatocytes (Figure 1B). Representative ROIs and pixel-intensity histograms are shown in Figure S1, and detailed colocalization metrics, including Manders' coefficients and Costes statistics, are provided in Supplemental Table 1.

Co-localization of proteins involved in lipogenesis around LDs in hepatocytes

To investigate potential additional interactions among lipogenic proteins, we used antibodies recognizing various lipogenic enzymes in primary hepatocytes to assess co-localization by confocal immunofluorescence. As shown in Figure 1, co-localization was quantified using Pearson's correlation coefficient (Pearson's R) in LD-enriched regions where the enzyme signals were concentrated. Immunofluorescence imaging of primary hepatocytes isolated from rat livers refed a FFD demonstrated that ACC1 co-localized with ACL (R ≈ 0.83), ACSS2 (R ≈ 0.46), and FAS (R ≈ 0.55) around LDs, despite their known localization in the cytosol. GPAM, a mitochondrial outer membrane protein involved in glycerolipid synthesis, also co-localized with

ACC1 in LD-associated regions (R ≈ 0.51). In contrast, the gluconeogenic enzyme PEPCK,
which is unrelated to lipogenesis, showed almost no overlap with ACC1 (R ≈ 0.11), likely
because it remains confined to the cytosol and does not redistribute to LD-proximal regions
(Figure 2A, Supplemental Table 2). Comprehensive testing metrics, including Manders'
coefficients and Costes significance testing, are provided in Supplemental Table 2 and further
support the robust spatial association of ACC1 with multiple cytosolic lipogenic enzymes,
whereas the non-lipogenic negative control, PEPCK, showed only minimal overlap with ACC1.
Adipose differentiation-related protein (ADRP), an LD-resident protein, marked LDs where ACL,
ACC1, and FAS also localized in primary hepatocytes from refed rats (Figure 2B). Three-
dimensional imaging further confirmed the localization of ACC1 surrounding LDs in primary
hepatocytes from refed rats (Figure 2C). Additional imaging across various fasting and
refeeding timepoints suggested that ACC1-containing complexes assemble progressively during
metabolic activation and accumulate around LDs in rat primary hepatocytes (Figure S2).

Co-localization of proteins involved in lipogenesis at the interface of
mitochondria and LDs in hepatocytes

Electron microscopy (EM) with immunogold labeling using antibodies recognizing ACC1 or FAS
was used to determine their subcellular localization in primary hepatocytes from refed rats.
Consistent with the immunofluorescence images (Figure 2C; Figure S2), EM revealed that
ACC1 localized predominantly around LDs (Figure 3A). This finding was further supported by
immunofluorescence imaging and EM performed on the same hepatocyte (Figure S3A). Double
immunogold labeling using different sizes of gold particles for ACC1 and FAS demonstrated
their co-localization around LDs (Figure 3B; Figure S3B–C). Although ACC1 and FAS were
occasionally observed near the ER, they were much more consistently enriched around LDs
(Figure S3D). This pattern aligned with staining of the ER marker protein disulfide isomerase
(PDI), which showed only limited overlap with ACC1, in contrast to the strong co-localization
observed between ACC1 and LDs (Figure S3E).

Notably, interactions between ACC1 and FAS were also observed at interfaces between
mitochondria and LDs (Figure 3C). Subcellular fractionation of liver lysates from FFD-refed rats
further demonstrated that ACL, ACC1, ACC2, FAS, SCD1, and GPAM were present in both LD
and mitochondrial fractions, regardless of their canonical localizations in the cytoplasm,
mitochondria, or ER (Figure 3D). EM frequently revealed mitochondria positioned adjacent to
LDs (Figure S3F), and GPAM localization was subsequently confirmed to be on the outer
mitochondrial membrane, including sites where mitochondria interface with LDs (Figure S3G).

Subcellular fractionation of liver lysates from FFD-refed rats further confirmed this finding,
whereas mitochondrial inner membrane proteins, such as prohibitin and cytochrome c, were
found only in the mitochondrial fraction (Figure S3H).

Combined, these findings demonstrate that lipogenic proteins co-localize around LDs and at
mitochondria–LD contact sites under high lipogenesis conditions.

Active dephosphorylated ACC promotes polymerization, enhances lipogenesis,
and increases LD localization

ACC activity is acutely regulated through phosphorylation and dephosphorylation. AMP-
activated protein kinase (AMPK) phosphorylates ACC at Ser-79 [42,43], inhibiting its enzymatic
activity through largely undefined mechanisms [44,45]. Fasting or glucagon treatment increases
AMPK activity, whereas insulin suppresses AMPK activity and stimulates the Mg^{2+}-dependent
protein phosphatase 2A (PP2A), which dephosphorylates ACC, thereby increasing its activity [46-
50]. During refeeding, ACC is converted from its phosphorylated inactive form to its
dephosphorylated, active form due to stimulation of protein phosphatase activity. This
dephosphorylation activates ACC and DNL in hepatocytes, allowing excess glucose to be
converted into FAs and TGs for storage [51,52].

We next determined whether the phosphorylated state of ACC affected the formation of the
lipogenic complex. For this study, we treated rat primary hepatocytes with 5-aminoimidazole-4-
carboxamide ribonucleoside (AICAR), an AMPK agonist, to induce ACC phosphorylation [53]. The
primary hepatocytes were incubated with 1 mM AICAR and cell lysates were prepared at 0, 1.5,
and 3 hours after treatment. SDS-PAGE immunoblot analysis of cell lysates showed an
increase in phosphorylated ACC levels in the AICAR-treated cells compared to the untreated
cells; however, the amount of total ACC protein was unchanged (Figure 4A). Blue native PAGE
analysis of cell lysates demonstrated that the polymerized, higher molecular weight form of
ACC1 was present only in untreated primary hepatocytes and was disrupted in AICAR-treated
hepatocytes (Figure 4A). Even with MIG12 overexpression, which robustly promotes ACC1
polymerization, AICAR reduced the higher molecular weight form of ACC1 to baseline (Figure
4B). Consistent with reduced polymerization, FA synthesis measured by ³H-acetate
incorporation into newly synthesized FAs fell ~6-fold within 30 minutes of AICAR treatment
(Figure 4C), while FA-synthesis enzyme levels remained unchanged (Figure S4A). Collectively,
these results imply that the rapid decrease in DNL results from AMPK-mediated ACC
phosphorylation, which disrupts ACC1 polymerization.

To further explore the role of dephosphorylated ACC in promoting DNL *in vitro*, we blocked
endogenous phosphatase activity in liver cytosol using a phosphatase inhibitor cocktail (Figure
4D). Immunoblot analysis confirmed that the inhibitor prevented ACC dephosphorylation,
whereas λ-phosphatase enhanced dephosphorylation (Figure S4B). Compared with the
inhibitor-treated group, untreated liver cytosol, which underwent dephosphorylation during
incubation, showed a 4-fold increase in palmitate synthesis rates in a citrate concentration-
dependent manner (Figure 4D). Furthermore, in primary hepatocytes from refed rats, ACC1
localized around LDs under DMSO control, consistent with Figures 1–3, but this LD-associated
pattern was significantly reduced with AICAR treatment (Figure 4E).

Taken together, these findings demonstrate that the polymerization of ACC, mediated by
dephosphorylation, is critical for its localization with other lipogenic enzymes around LDs and for
regulating DNL in hepatocytes.

The interaction of ACC1 with lipid synthesis-associated proteins is enhanced when ACC is in the activated state

As demonstrated above, ACC1 polymerizes and co-localizes with other lipogenic proteins
around LDs under high DNL conditions. To determine how ACC1 interactions with other
proteins change upon activation, we performed ACC1 immunoprecipitation from mouse liver
lysates incubated with 5 mM $MgCl_2$ at 30°C for 30 minutes with or without a phosphatase
inhibitor cocktail. With endogenous phosphatases active, ACC1 co-immunoprecipitated ACC2,
ACSS2, FAS, ACL, GPAM, MIG12, S14, and SCD1 (Figure 5A). Notably, β-actin, a cytoskeletal
protein, also showed significant binding to ACC1, suggesting that cytoskeletal components
might play a role in the formation of the lipogenic complex. In contrast, these interactions were
markedly reduced when endogenous phosphatases were inhibited, suggesting that ACC1
interacts more strongly with other proteins in a phosphatase-active state. Proteins unrelated to
lipid synthesis, such as PEPCK, prohibitin, cytochrome c, and PDI, did not bind ACC1 under
either condition.

Notably, ACC2 maintained strong binding to ACC1 even in the presence of phosphatase
inhibitors, suggesting that ACC1 and ACC2 form a relatively stable heteromer. These
observations are supported by immunoprecipitation of liver lysates from hepatocyte-specific
ACC1 or ACC2 knockout mice, which revealed a clear reciprocal interaction between the two
enzymes (Figure S5A). Blue native PAGE analysis further supported this model: whereas ACC1
migrated similarly in wild-type and ACC2-deficient livers, ACC2 showed a markedly altered

pattern in ACC1-deficient livers. In normal liver, ACC2 predominantly appeared as a dimer or
tetramer, but in the absence of ACC1, ACC2 shifted toward higher–molecular-weight, more
polymerized species (Figure S5B). Considering that hepatic ACC2 expression is substantially
lower than ACC1, these findings suggest that, under physiological conditions, most endogenous
ACC2 may exist as a heterodimer with ACC1.

Immunofluorescence analysis in cultured cells provided additional evidence for the ACC1–
ACC2 heterodimer. ACC1 was detected predominantly in the cytosol when expressed alone
(Figure S5C). However, when co-transfected with ACC2, a substantial fraction of ACC1
colocalized with ACC2 at the mitochondria (Figure S5D). In addition to this mitochondrial pool,
ACC1 also displayed cytosolic localization that did not overlap with ACC2. These observations
indicate that ACC1 is present in the cytosol but also at the mitochondrial membrane as a
complex with ACC2.

To determine whether endogenous ACC2 influences ACC1-mediated protein interactions, we
immunoprecipitated ACC1 from refed wild-type and ACC2 hepatocyte-specific knockout livers,
using ACC1-deficient mice as a negative control. Co-precipitation of ACL, ACSS2, FAS, S14,
SCD1, GPAM, β-actin, and β-tubulin was reduced to varying degrees in ACC2-deficient livers
compared with wild-type (Figure 5B), suggesting that ACC2 contributes to the stability or
assembly of ACC1-associated protein complexes. MIG12 binding to ACC1 was unaffected by
phosphatase inhibition (Figure 5A) or by the absence of ACC2 (Figure 5B), consistent with its
direct association with ACC1. As expected, PEPCK, prohibitin, cytochrome c, and calnexin were
not co-immunoprecipitated with ACC1 in any group.

Given that ACL, ACSS2, ACC, FAS, SCD1, and the mitochondrial GPAM participate in ACC1-
associated complexes, we next measured TG synthesis under phosphatase active versus
phosphatase-inactive conditions. Glycerol 3-phosphate (G3P) was used as a substrate with
excess oleoyl-CoA added to liver homogenates from refed rats to test whether TG synthesis via
GPAM is enhanced when endogenous phosphatases are active, a condition that favors
lipogenic complex formation (Figure S5E). The results demonstrated that TG synthesis was
significantly higher under phosphatase-active conditions and increased further with rising $MgCl_2$
concentrations, paralleling $MgCl_2$-dependent ACC1 dephosphorylation (Figure S4B), whereas
under phosphatase inactive conditions, TG synthesis remained unchanged with $MgCl_2$ (Figure
S5E).

These results demonstrate that phosphorylation–dephosphorylation regulates both FA and TG
synthesis in part by controlling ACC1-mediated lipogenic complex formation.

Lipogenic metabolon formation facilitates FA synthesis

To confirm the interactions between ACC1 and the other FA synthesis proteins, and to assess
the regulatory effects of phosphorylation and dephosphorylation on the activities of individual FA
synthesis enzymes *in vitro*, we conducted a comprehensive purification of recombinant ACSS2,
ACC1, FAS, MIG12, and S14 in mammalian cells using Ni-NTA resin and gel-filtration
chromatography (Figure 6A, S6A), and rPEPCK was purified as a negative control (Figure S6B).
The recombinant protein mixture was dephosphorylated with λ-phosphatase and
immunoprecipitation was performed with anti-ACC1 antibodies. Immunoblot analysis (Figure 6B)
revealed that all other recombinant FA synthesis proteins co-immunoprecipitated with rACC1,
whereas rPEPCK did not. These results confirm that recombinant ACC1 interacts with core
lipogenic factors *in vitro* using recombinant proteins (Figure 6B).

Next, we pre-incubated the individual recombinant proteins or their mixtures with λ-phosphatase
to simulate the DNL activated environment as described above. Parallel groups without λ-
phosphatase treatment were used as phosphorylated controls. Figure S6C shows that each
purified protein exhibited functional activity, and interestingly, all tested lipogenic proteins
displayed increased activity to varying degrees when dephosphorylated compared to their
phosphorylated forms. However, rACC1 exhibited the greatest increase in enzymatic activity
upon λ-phosphatase treatment compared with the other lipogenic enzymes. Next, we combined
purified lipogenic enzymes, rACL-rACC1-rFAS or rACSS2-rACC1-rFAS, and measured FA
synthesis from citrate or acetate, respectively. As predicted, rACL-rACC1-rFAS activity was
nearly four times higher with λ-phosphatase treatment compared with untreated conditions
(Figure 6C). Similarly, rACSS2-rACC1-rFAS activity increased by ~3.5-fold upon λ-phosphatase
treatment relative to untreated conditions (Figure 6C). Figure S6D shows the phosphorylation
levels of rACC1 alongside total rACC1 levels at various time points following λ-phosphatase
treatment, demonstrating that λ-phosphatase efficiently dephosphorylates proteins without
altering their total levels.

Inasmuch as each rACL, rACSS2, rACC1, and rFAS protein exhibited increased enzymatic
activity in the dephosphorylated state, these observations alone could not distinguish whether
the enhanced palmitate production resulted from cooperative complex formation or simply
reflected additive increases in the intrinsic activities of individual enzymes upon

dephosphorylation. To directly address this issue, we performed a FA-synthesis assay using (1)
liver cytosol alone, (2) recombinant ACL–ACC1–FAS alone, and (3) a mixture of liver cytosol
and recombinant ACL–ACC1–FAS, with protein amounts normalized across conditions. Each
group was subjected to either dephosphorylated or phosphorylated conditions under identical
treatment settings. As shown in Figure S6E, dephosphorylation was successfully achieved in all
conditions. Under dephosphorylated conditions, combining liver cytosol with recombinant ACL–
ACC1–FAS produced substantially greater palmitate than the expected additive activity
calculated from cytosol alone and recombinant enzymes assayed in parallel (Figure 6D). This
enhancement exceeded what would be predicted from simple additive effects of
dephosphorylation on individual enzymes and was supported by a significant interaction
between phosphorylation state and mixing condition (two-way ANOVA, $F_{(3,16)}$ = 133.4, $P <$
0.0001). Together, these data indicate a strong synergistic increase in lipogenesis arising from
cooperative interactions between dephosphorylated cytosolic proteins and recombinant
enzymes. In contrast, under phosphorylated conditions, the palmitate produced by the
combined reaction was indistinguishable from the expected additive activity (Figure 6E),
indicating that cooperative assembly and synergistic flux enhancement are selectively enabled
by dephosphorylation.

To directly assess complex formation under these conditions, ACC1 was immunoprecipitated
from the same reaction mixtures. Consistent with the results shown in Figure 5A,
dephosphorylation markedly enhanced the interactions of ACC1 with FAS and ACL, and β-
tubulin as determined by co-precipitation studies, suggesting that the lipogenic metabolon may
be functionally linked to cytoskeletal components (Figure S6F). No nonspecific binding was
observed with control rabbit IgG.

While it is not feasible to systematically test all cytosolic proteins that could potentially contribute
to metabolon assembly, Figure 5 identified structural proteins as representative non-lipogenic
factors that interact with ACC1 under dephosphorylated conditions. Consistent with this
observation, β-tubulin showed detectable co-localization with ACC1 around lipid droplets in
primary hepatocytes under lipogenic refeeding conditions (Pearson's R ≈ 0.32), whereas this
spatial association was markedly reduced upon fasting (Figure 6F). In parallel, pharmacological
perturbation of microtubule dynamics provided support for the role of cytoskeleton in lipogenesis.
Disrupting microtubule polymerization with nocodazole consistently decreased FA synthesis in
primary hepatocytes, whereas stabilizing microtubules with taxol, an agent with the opposite
effect of nocodazole, had minimal impact on lipogenesis (Figure 6G). These observations

suggest that intact cytoskeletal architecture facilitates lipogenesis by coordinating the nutrient-responsive assembly and spatial organization of lipogenic enzyme complexes, rather than directly regulating microtubule dynamics.

Collectively, these findings suggest that lipogenic activity is enhanced through post-translational regulation that promotes the formation of a functional lipogenic metabolon under physiological conditions favoring high DNL. This process likely involves dynamic and cooperative interactions among DNL- and TG-related enzymes, aided by structural proteins and other accessory factors, allowing for more integrated and efficient lipogenesis.

ACC1 anchors a multi-module lipogenic network integrating translation, trafficking, and lipid metabolism

To further map the ACC1-centered protein network, we immunoprecipitated ACC1 from wild-type and ACC-deficient liver lysates. In parallel, MIG12 and S14, two direct ACC1-binding partners were immunoprecipitated, as was ACSS2, a representative lipogenic enzyme, to compare their interaction profiles (Figure 7A). MIG12 and S14 pulldowns closely mirrored the ACC1 immunoprecipitation pattern: proteins enriched in wild-type lysates were markedly reduced in ACC-deficient lysates, consistent with their direct association with ACC1 and the dependence of their interactomes on ACC1 integrity. In contrast, ACSS2 immunoprecipitation showed a distinct profile. Although several lipogenic enzymes were reduced in ACC-deficient lysates, ACSS2 retained clear interactions with multiple lipogenic proteins. This pattern suggests that ACC1 enhances or stabilizes the overall lipogenic assembly rather than acting as its sole organizing component. Notably, β-actin binding to ACSS2 was unchanged regardless of ACC1 status, implying that actin contributes to complex organization through an ACC1-independent mechanism.

We next quantified proteins co-immunoprecipitated with ACC1 by mass spectrometry. Proteins that were reduced by >20-fold in ACC1, MIG12, or S14 immunoprecipitations from ACC-deficient lysates were classified as ACC1-dependent interactors. STRING network analysis of these proteins resolved three highly connected clusters (Figure 7B, top). The largest cluster consisted of ribosomal and translation-associated proteins, suggesting that a robust interface between ACC1 and the translational machinery exists. A second cluster was enriched for vesicle/trafficking components and cytoskeletal protein-binding factors, indicating that ACC1 is embedded within a broader transport and structural scaffold. A third cluster consisted of enzymes involved in FA synthesis and remodeling, consistent with ACC1's canonical metabolic

function (Figure 7B, bottom). DAVID functional annotations performed on the same dataset
produced closely matching categories (Figure S7 and Supplemental Table 3).

Together, these findings indicate that activated ACC1 functions as a central organizational hub
that coordinates core lipogenic enzymes—including FAS, ACL, ACSS2, SCD1, GPAM, and
ACC2—into a coherent metabolic assembly. Direct binding by MIG12 and S14 likely stabilizes
this structure, while cytoskeletal and trafficking components associate with the ACC1-centered
complex to support spatial organization and substrate delivery. The presence of ribosomal
proteins and transport-associated factors among ACC1-dependent interactors further suggests
that lipogenesis is integrated with protein synthesis and intracellular trafficking through multiple
points of connection.

DISCUSSION

Transcriptional control of FA and TG synthesis is well characterized [54], but much less is known
about how lipid synthesis is regulated post-transcriptionally. Here, we identified a previously
unrecognized post-translational mechanism in which hepatic lipogenic enzymes assemble into a
multi-protein metabolon under anabolic conditions. This metabolon provides a spatial basis for
efficient glycerolipid synthesis and introduces a new dimension to the regulation of DNL.

Using immunofluorescence and EM, we showed that enzymes involved in FA and TG synthesis,
along with accessory proteins like MIG12 and S14, co-localize around LDs and function as an
integrated multi-enzyme assembly rather than as isolated enzymes under physiological
activation states such as high carbohydrate feeding. These findings build on earlier LD
proteomics reports that identified ACC1 [55-59] and FAS [58] on LDs, extending them by providing
direct structural visualization of their co-localization in mammalian hepatocytes. Furthermore,
the proximity between cytosolic enzymes such as ACC1 and FAS and the mitochondrial
enzyme GPAM at contact sites suggests coordinated substrate transfer into the glycerolipid
pathway. Subcellular fractionation of liver lysates from high-carbohydrate-refed rats further
supports these observations.

Immunoprecipitation of ACC1 from liver lysates incubated under phosphatase-active conditions
revealed robust interactions between ACC1 and multiple lipogenic proteins, whereas the
inclusion of phosphatase inhibitors markedly reduced these associations (Figure 5A). This
biochemical shift was paralleled by *in vitro* FA and TG synthesis assays, in which phosphatase-
active conditions produced substantially higher lipid synthesis than phosphatase-inhibited ones.
We initially attributed this enhanced interaction and activity to ACC1 dephosphorylation, a

hypothesis supported by *in vivo* experiments using the AMPK agonist AICAR. In primary
hepatocytes isolated from high-carbohydrate–refed rats, AICAR treatment caused the activated,
polymerized form of ACC1 to dissociate into its inactive dimeric form, as shown by blue native
PAGE. The extensive polymerization of ACC1 driven by MIG12 overexpression was also
significantly reduced by AICAR-mediated ACC1 phosphorylation. Correspondingly, short-term
AICAR treatment significantly suppressed hepatic lipogenesis in primary hepatocytes without
affecting the steady-state levels of lipogenic enzymes and markedly reduced ACC1 localization
around LDs. Notably, ACC1 was not the only enzyme influenced by dephosphorylation, as
phosphatase treatment also increased the enzymatic activities of rACSS2 and rFAS, although to
a lesser extent than ACC1. These findings suggest that anabolic phosphatase-active states
coordinate multiple components of the lipogenic pathway, collectively enhancing metabolon
assembly and catalytic output.

Another notable finding from the immunoprecipitation experiments was the strong interaction of
ACC1 with ACC2 and MIG12. Whereas most ACC1-associated proteins showed enhanced
binding under phosphatase-active conditions, ACC2 and MIG12 maintained stable interactions
with ACC1 irrespective of its phosphorylation state. This feature suggests that these two
proteins form a relatively stable core structural interaction with ACC1.

Additional insight into ACC2 was obtained from transfection studies in CHO-K1 cells. When
ACC1 and ACC2 were co-expressed, the fraction of ACC1 that interacted with ACC2 and
formed heterodimers or higher-order heteromultimers and localized to mitochondria increased
(Figure S5D). ACC1 also retained a substantial cytosolic pool that did not overlap with ACC2,
likely reflecting cytosolic homomeric ACC1 assemblies. These observations indicate that ACC1
is partitioned between heteromeric ACC1–ACC2 complexes at mitochondria and homomeric
ACC1 assemblies in the cytosol and suggest that the subcellular localization and functional
partitioning of malonyl-CoA–producing acetyl-CoA carboxylase activity may be influenced by the
relative abundance of ACC2. These findings are consistent with the tissue-specific expression
patterns of ACC isoforms. In liver and adipose tissue, where malonyl-CoA is used primarily for
lipogenesis, ACC1 is the predominant isoform; whereas in muscle and heart, where malonyl-
CoA mainly inhibits mitochondrial fatty acid oxidation, ACC2 is the major form. Thus, cells may
regulate the relative abundance (stoichiometry) of ACC1 and ACC2 to redistribute acetyl-CoA
carboxylase activity between the cytosol and mitochondria according to metabolic demands. In
this context, ACC1–ACC2 interactions may serve as a regulatory node linking FA synthesis and
oxidation. Alterations in the stoichiometry of the two isoforms could therefore shift the balance

between lipogenesis and control of mitochondrial FA oxidation by modulating the subcellular

distribution of carboxylase activity.

Finally, although ACC2 has long been viewed primarily as a regulator of mitochondrial FA

oxidation, ACC2 deletion reduced ACC1 interactions with multiple lipogenic enzymes and

structural proteins (Figure 5B). This finding supports the view that ACC2 functions not only as a

metabolic enzyme that inhibits FA oxidation but also as a structural component essential for the

organization and maintenance of the lipogenic metabolon. Complementing these structural

insights, our biochemical assays demonstrated that assembling lipogenic enzymes within a

shared reaction environment generates catalytic outputs that exceed the sum of individual

activities. Under DNL-activated conditions, combining recombinant ACL–ACC1–FAS with liver

cytosol nearly doubled the FA synthesis compared with the expected additive activity, indicating

synergistic enhancement of pathway flux. This synergy suggests that enzyme co-assembly,

rather than simply increased intrinsic activity, facilitates more efficient substrate channeling

during active lipogenesis. These functional observations reinforce the multi-module organization

described in Figure 7, in which ACC1 immunocomplexes interact not only with lipogenic

enzymes but also with ER-proximal translation components, vesicle-movement proteins, and

actin-linked scaffolds, supporting the concept that lipogenic enzymes achieve maximal

efficiency when spatially integrated with cytoskeletal scaffolds and intracellular transport

machinery, and when functionally linked to translation-associated components.

Although metabolon-based regulation is less explored in lipogenesis, analogous mechanisms

have been established in the TCA cycle and glycolysis [29-31,60-63]. The substrate-responsive

assembly of PFK filaments in the presence of fructose-6-phosphate [64] provides a relevant

precedent and parallels ACC1 filament formation under anabolic conditions, suggesting that

these filaments may serve as structural scaffolds for metabolon assembly. Consistent with this

concept, a FA–synthesis metabolon has been described in the thermophile *Chaetomium*

thermophilum [65] and ACC–FAS co-localization has been visualized in *Saccharomyces*

cerevisiae [66], where these enzymes are reorganized or sequestered in response to glucose

starvation. These observations across lower eukaryotes support the idea that nutrient state

governs the assembly and disassembly of spatially organized lipogenic enzymes.

Recent work by Wijesinghe *et al.* [67] extends this concept to mammalian cells. In their study,

ACC1 formed ring-like puncta and localized to the surface of lipid droplets (LDs), but this

localization was highly conditional: anabolic cues such as AMPK inhibition preserved ACC1–LD

co-localization, whereas large LDs generated by acute oleic-acid loading—which diminishes

anabolic drive—displayed reduced ACC1 staining. This pattern mirrors our observation that the AMPK agonist AICAR suppresses ACC1 accumulation around LDs (Figure 4E), indicating that ACC1 recruitment to LDs requires appropriate metabolic activation rather than merely reflecting LD abundance. Together, these findings align with our model in which ACC1 polymerization and its interactions with accessory factors generate a nutrient-responsive lipogenic complex in hepatocytes. Collectively, the emerging evidence across species supports a broader framework in which ACC1 integrates nutrient status with spatial enzyme organization to optimize lipogenesis during anabolic transitions.

In conclusion, our study demonstrates that under anabolic conditions, hepatic lipogenic enzymes assemble into a spatially organized, phosphorylation-responsive metabolon around LDs. This multi-protein metabolon enhances catalytic efficiency facilitating improved transfer of intermediates between sequential enzymes, thereby supporting efficient synthesis of FA and TGs. These findings provide a mechanistic foundation for understanding hepatic lipid synthesis and highlight therapeutic opportunities in metabolic disease by targeting the structure and regulation of the lipogenic metabolon.

Limitations of the study

In this study, we present evidence for a novel post-translational regulatory mechanism that enhances FA and TG synthesis in the liver via the assembly of a multi-protein metabolon. Ideally, since the structure of ACC1 under different activation states has been reported [68,69], we would have included structural analyses of this ACC1 mediated lipogenic metabolon. However, detecting such metabolons presents greater challenges compared to stable protein assemblies. The reason is that a metabolon is characterized as a transient structural-functional metabolon formed by sequential enzymes within a metabolic pathway, exhibiting non-covalent interactions and associating with various cellular structural elements, such as integral membrane proteins and cytoskeletal components [30,70]. Consequently, the interactions among these proteins are more transient and dynamic, enabling them to adapt to fluctuations in metabolic environments.

Additionally, the mRNA levels of the lipogenic proteins are all significantly downregulated during fasting as a result of reduced SREBP-1c expression, which restricted our ability to compare the co-localization of lipogenic proteins between fasting and refeeding conditions. Therefore, we employed AICAR treatment and elevated phosphorylation levels to simulate catabolic states *in vivo* and *in vitro*, respectively.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this manuscript, the authors used ChatGPT (OpenAI) solely for language editing, including grammar correction and improvement of clarity. No content, data interpretation, or scientific conclusions were generated by the tool. After using this assistance, the authors thoroughly reviewed, revised, and approved all text, and they take full responsibility for the integrity and accuracy of the final manuscript.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to the lead contact, Jay D. Horton (jay.horton@utsouthwestern.edu).

Materials availability

Custom antibodies generated in this study are available upon reasonable request and completion of an MTA. Cloning constructs are available from the Lead Contact. All other materials used in this study are specified in the key resources table.

Data and code availability

All data supporting the findings are available within the paper and Supplemental Information. No custom code was used.

ACKNOWLEDGEMENTS

We thank Norma Anderson, Judy Sanchez, Tuyet Dang, and Tessa Edwards for excellent technical assistance; Lisa Beaty for suspension cell culture and set up; Linda Donnelly for assistance with antibody generation; Jinhai Yu for providing TEV protease; and Chelsea Burroughs and Nancy Heard for help with graphics. The work was supported by grants NIH P01HL160487 and NIH P30DK127984.

AUTHOR CONTRIBUTIONS

X.Z., W.B.M., T.T., X.F., and C.-W.K. conducted the experiments. J.D.H., C.-W.K., and X.Z. designed the experiments. X.Z., C.-W.K., and J.D.H. wrote the manuscript. All authors reviewed and approved the article.

DECLARATION OF INTERESTS

J.D.H. is a consultant for AbbVie and insitro.

Figure Legends

Figure 1. Identification of the lipogenic complex in liver

(A) An adenovirus expressing S14-Flag was injected into mice, and liver lysates were prepared 3 days later. Immunoprecipitation was performed using anti-Flag antibodies. Eluates from the S14-Flag IP and an identically processed β-gal–injected control were separated on 8% and 15% SDS–PAGE gels, followed by silver staining. Protein bands that appeared specifically in the S14-Flag lanes were excised and analyzed by LC–MS/MS to identify potential S14-associated proteins.

(B) Primary hepatocytes were prepared from rats refed a FFD and plated on collagen-coated coverslips. Immunofluorescence confocal imaging was performed as described in Methods. Co-localization of ACC1 with MIG12 and S14 was confirmed by confocal microscopy using anti-mouse monoclonal ACC1, anti-MIG12 rabbit polyclonal antibodies, and anti-S14 rabbit polyclonal antibodies. Quantitative co-localization analysis was performed using Pearson's correlation (above Costes threshold), and full analytical metrics, including Manders' coefficients and Costes significance testing, are provided in Figure S1 and Supplemental Table 1.

Figure 2. Co-localization of proteins involved in lipogenesis around LDs in hepatocytes

(A) Co-localization of enzymes involved in lipogenesis in primary hepatocytes from rats refed a FFD using confocal microscopy as described in Methods. Quantitative co-localization analysis was performed using Pearson's correlation (above Costes threshold), and full analytical metrics are provided in Supplemental Table 2.

(B) Localization of enzymes involved in lipogenesis around LDs in primary hepatocytes from rats refed a FFD using confocal microscopy as described in Methods. Co-localization analysis was performed using Pearson's correlation as in panel A, with results summarized in Supplemental Table 2.

535   (C) A three-dimensional image stack was acquired using an OMX microscope (z-step size: 120

536   nm) to visualize the indicated proteins around lipid droplets, as described in Methods.

Figure 3. Co-localization of proteins involved in lipogenesis at the interface of mitochondria and LDs in hepatocytes

(A) Primary hepatocytes were prepared from rats refed a FFD and localization of ACC1 was

determined using immunogold-EM. The 1.4 nm nanogold particles were enlarged by gold

enhancement for clear visibility. Detailed protocol is described in Methods.

(B) Co-localization of ACC1 and FAS around LDs in primary hepatocytes from rats refed a FFD

using immunogold-EM. Size of gold particles: ACC1 - 5 nm; FAS - 10 nm.

(C) Co-localization of ACC1 and FAS between mitochondria and LDs in primary hepatocytes

from rats refed the FFD using immunogold-EM. Size of gold particles: ACC1 - 5 nm; FAS - 10

nm.

(D) Subcellular fractionation was performed with liver lysates from rats refed the FFD to isolate

LDs as well as mitochondria, microsomes, and cytoplasm. Detailed protocols for the subcellular

fractionation are described in Methods. Proteins from each fraction were subjected to SDS-

PAGE and immunoblot analysis with the indicated antibodies. PLIN2, prohibitin, and α-tubulin

were used as subcellular localization markers for lipid droplets, mitochondria, and cytosolic

proteins, respectively.

Figure 4. Active dephosphorylated ACC promotes polymerization, enhances lipogenesis, and increases LD localization

(A) Primary hepatocytes from rats refed a FFD were treated with 1 mM AICAR for 0, 1.5, and 3

hours. Equal aliquots of cell lysate proteins were loaded and analyzed by BN-PAGE (above) or

SDS-PAGE (bottom) for immunoblot analysis. An anti-ACC1 antibody was used to perform the

immunoblot analysis of BN-PAGE. For SDS-PAGE, ACC immunoblots were performed using a

fluorescently labeled streptavidin conjugate.

(B) Primary hepatocytes from rats refed a FFD were isolated. One set of primary hepatocytes

were untreated, while the other two sets were infected with an adenovirus expressing MIG12.

After 12 hours, 1 mM AICAR was added, and cells were harvested 3 hours later. Cell lysates

were prepared and subjected to BN-PAGE and SDS-PAGE for immunoblot analysis. An anti-

ACC1 antibody was used to perform the immunoblot analysis of BN-PAGE. For SDS-PAGE, a

fluorescently labeled streptavidin conjugate to detect ACC and an anti-MIG12 antibody were
used for immunoblot analysis.

(C) Primary rat hepatocytes prepared as in (B) were incubated with 1 mM AICAR for 30 minutes
and FA synthesis rates were measured *ex vivo* using ^{3}H-acetate as described in Methods.
Individual data points are shown; bars indicate the mean (n = 2 technical replicates from one
hepatocyte preparation).

(D) Liver cytosol from rats refed a FFD were pre-incubated with 1 mM $MnCl_2$ in the presence or
absence of a phosphatase inhibitor cocktail for 15 min at 30°C, then incubated with the
indicated concentrations of Na-citrate buffer (pH 7.1) for 15 min at 37°C. Palmitate synthesis
(left) was measured using ^{1}□C-citrate. Palmitate synthesis rates (right) were calculated from the
maximal velocity (Vmax) of the enzymatic reaction. Data represent mean ± SEM from n = 3
technical replicates. Statistics: multiple two-tailed unpaired Student's t-tests with Holm–Šidák
correction, comparing phosphatase inhibitor–treated (phosphorylated) vs untreated
(dephosphorylated) conditions at each citrate concentration (*P < 0.05, **P < 0.01, ***P < 0.001,
****P < 0.0001). The detailed protocol for the enzymatic assay is described in Methods.

(E) Rat primary hepatocytes were isolated from rats fed a FFD for 2 days and incubated in
medium containing either (i) 25 mM glucose, 100 nM insulin, and 1% DMSO or (ii) 5 mM
glucose and 1 mM AICAR. Following the indicated times, cells were stained with anti-ACC1
antibodies and LipidTOX as described in Methods.

Figure 5. The interaction of ACC1 with lipid synthesis-associated proteins is
enhanced when ACC is in the activated state

(A) Liver lysates from refed mice were treated with or without a phosphatase inhibitor cocktail.
The ACC1 antibody was incubated with the lysate for 1 hour at RT and ACC1 was
immunoprecipitated. IgG was used as the control. The immunoprecipitation protocol is
described in Methods. Supernatant indicates the unbound fraction collected after bead pelleting;
it was mixed with SDS sample buffer and immunoblotted alongside the bead-bound material.

(B) ACC1 immunoprecipitation was carried out using lysates prepared from refed wild-type,
ACC1 hepatocyte knockout, or ACC2 hepatocyte knockout mouse livers, respectively.

Figure 6. Lipogenic metabolon formation facilitates FA synthesis

(A) Human C-terminal His-Flag-tagged ACSS2, ACL, ACC1, FAS, MIG12, and S14 were
individually overexpressed in mammalian HEK 293S GnTI⁻ cells and purified using Ni-NTA and
gel filtration. Epitope tags were removed by TEV cleavage. Detailed protocols for purification
were described in Methods. The final purified recombinant proteins were subjected to SDS-
PAGE for Coomassie blue staining.

(B) Recombinant protein mixtures were incubated with λ-phosphatase and 1 mM $MnCl_2$ for 30
minutes at 30°C and then the protein mixture was incubated with 0.5 mM Na-citrate buffer for 20
minutes at 30°C. rACC1 was immunoprecipitated from the protein mixture using an anti-ACC1
antibody. Recombinant protein mixtures without rACC1 were used for control
immunoprecipitations. Proteins were subjected to SDS-PAGE for immunoblot analysis. ACC1
immunoblots were performed using a fluorescently labeled streptavidin conjugate.

(C) Enzymatic activity of reconstituted recombinant lipogenic enzyme modules was measured
as described in Methods. The left panel shows FA synthesis activity measured from citrate
using rACL–rACC1–rFAS, whereas the right panel shows FA synthesis activity measured from
acetate using rACSS2–rACC1–rFAS. Reactions were performed under phosphorylated or
dephosphorylated conditions as indicated. Data represent mean ± SEM from n = 3 technical
replicates. Relative synthesis rates were calculated based on Vmax values.

(D–E) Under dephosphorylated (D) or phosphorylated (E) conditions, ACL–ACC1–FAS activity
was measured in mixtures of liver cytosol (from FFD-refed rats) and recombinant proteins
(rACL, rACC1, and rFAS) and compared with the expected additive activity calculated from
assays performed in cytosol alone and recombinant proteins alone in parallel. Data represent
mean ± SEM from n = 3 technical replicates. Statistical analysis was performed using two-way
ANOVA, revealing a significant interaction between phosphorylation state and mixing condition.
Post hoc comparisons indicate that observed activity significantly exceeded the expected
additive activity only under dephosphorylated conditions. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$,
****$P < 0.0001$.

(F) Primary hepatocytes isolated from rats under refed or fasted conditions were fixed and
stained for ACC1 and β-tubulin, followed by confocal microscopy as described in Methods.
Images show representative localization patterns under each condition.

(G) Primary hepatocytes from FFD-fed rats were cultured for 8 hours and then treated with
DMSO, nocodazole, or taxol (20 µM) as indicated. Cells were subsequently incubated with ^{14}C-

labeled citrate for 2 or 4 hours, and citrate-derived FA synthesis was quantified as described in Methods. Data points represent individual measurements from duplicate experiments.

Figure 7. ACC1 anchors a multi-module lipogenic network integrating translation, trafficking, and lipid metabolism

(A) Liver lysates were prepared from *ob/ob* mouse livers and ACC1/ACC2–deleted *ob/ob* mouse livers as described in Methods. To activate endogenous phosphatases, lysates were incubated in 5 mM $MgCl_2$ for 30 minutes at 30°C. In parallel, anti-ACC1, anti-MIG12, anti-S14, or anti-ACSS2 antibodies were pre-bound to protein G magnetic beads by incubating for 30 minutes at room temperature. The activated lysates were then added directly to the antibody–bead complexes and incubated for 1 hour at room temperature to perform immunoprecipitation. After extensive washing, bound proteins were eluted by boiling and analyzed by SDS-PAGE and immunoblot analysis was carried out with the indicated antibodies.

(B) Proteins co-immunoprecipitated with ACC1, MIG12, or S14 that were reduced by >20-fold in ACC-deficient lysates were classified as ACC1-dependent interactors and subjected to STRING network analysis. Three major functional clusters emerged—translation (red), protein trafficking/transport and cytoskeletal protein binding (green), and FA metabolism (blue). These clusters are displayed both as a network graph and summarized in the bar graph below using the corresponding color scheme. The inset mini-graphs within each colored bar summarize representative GO/KEGG terms for that cluster, with full enrichment analyses provided in Figure S7.

Key resource table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-ACL	Abcam	Cat# ab40793
Rabbit anti-ACSS2	Abcam	Cat# ab66038
Rabbit anti-mouse ACC1	House antibodies[71]	508D or 510D
Mouse monoclonal anti-mouse ACC1	House antibodies	2B5
Rabbit anti-mouse ACC2	House antibodies	456D
Rabbit anti-human FAS	House antibodies	310D
Rabbit anti-mouse MIG12	House antibodies[72]	143D

Mouse monoclonal anti-mouse MIG12	House antibodies	6C4
Rabbit anti-mouse S14	House antibodies	39C
Rabbit anti-PEPCK	Abcam	Cat# ab70358
Mouse anti-β-Tubulin	ThermoFisher	Cat# MA5-16308
Rabbit anti-β-Actin	Cell Signaling	Cat# 4970S
Rabbit Anti-mouse GPAM	House antibodies	280D
Mouse monoclonal anti-rat GPAM	House antibodies	16F8
Anti-ADRP	Sigma	Cat# SAB4200452
Anti-SCD1	Cell Signaling	Cat# 2794S
Anti-PROHIBITIN	Santa Cruz	Cat# SC-56467
Anti-CYTOCHROME C	Millipore	Cat# MAC1800
Mouse anti-CPT1	Abcam	Cat# ab128568
Anti-Ser79-pACC1	Cell Signaling	Cat# 3661S
Anti-PDI	Cell Signaling	Cat# 2446S
Goat Anti-Rabbit IgG, F(ab')2	Jackson Immuno	Cat# 111-035-047
Goat Anti-Mouse IgG, F(ab')2	Jackson Immuno	Cat# 115-035-072
Li-COR IRDye 800 goat anti-Rabbit IgG	Fisher Scientific	Cat# NC9401842
Li-COR IRDye 680 goat anti-Mouse IgG	Fisher Scientific	Cat# NC0252290
Li-COR IRDye 680RD Streptavidin	Fisher Scientific	Cat# NC0337633
Bacterial and virus strains		
Stellar™ Competent Cells	Takara	Cat# 636764
TOP10 Competent Cells	ThermoFisher	Cat# C404003
Chemicals, peptides, and recombinant proteins		
LB broth	Research Products International	Cat# L24061
Freestyle 293 expression medium	Gibco	Cat# 12338018
Opti-MEM™ I Reduced Serum Medium	Gibco	Cat# 31985070
Fetal Calf Serum (FCS)	Sigma-Aldrich	Cat# F2442
Penicillin-Streptomycin solution (100x)	Corning	Cat# 30-002-CI
PEI 25K	Polysciences	Cat# 23966-1
Protease Inhibitor Cocktail Set III, EDTA-Free - Calbiochem	Millipore-Sigma	Cat# 539134-1ML
Protease Inhibitor Cocktail, for use in purification of	Sigma-Aldrich	Cat# P8849-5ML

Histidine-tagged proteins, DMSO solution		
Purified rTEV protease	Gift from Dr. Jinhai Yu, UT Southwestern Medical Center	N/A
Lambda Protein Phosphatase	New England Biolabs	Cat# P0753L
DL-Dithiothreitol	Sigma-Aldrich	Cat# 43819-5G
Adenosine 5'-triphosphate magnesium salt	Sigma-Aldrich	Cat# A9187-1G
Adenosine 5'-triphosphate disodium salt hydrate	Sigma-Aldrich	Cat# A2383-5G
Coenzyme A sodium salt hydrate	Sigma-Aldrich	Cat# C3144-25MG
Acetyl coenzyme A sodium salt	Sigma-Aldrich	Cat# A2056-25MG
Malonyl coenzyme A lithium salt	Sigma-Aldrich	Cat# M4263-10MG
NADPH, Tetrasodium Salt	Sigma-Aldrich	Cat# 481973-100MG
Sodium bicarbonate [14C]	American Radiolabeled Chemicals	Cat# ARC 0138C-50 µCi
Malonyl [2-14C] Coenzyme A	American Radiolabeled Chemicals	Cat# ARC 0528-25 µCi
Citric Acid, [1,5-14C]	Revvity	Cat# NEC160050UC
Acetic Acid, Sodium Salt, [1,2-14C]	Revvity	Cat# NEC553050UC
Lauryl Maltose Neopentyl Glycol	Anatrace	Cat# NG310 5 GM
3a70B Complete Counting Cocktail	Research Products International	Cat# 111154
In-Fusion® Snap Assembly Value Bundle	Takara	Cat# 638946
Platinum™ SuperFi II PCR Master Mix	ThermoFisher	Cat# 12368010
DreamTaq Green PCR Master Mix (2X)	ThermoFisher	Cat# K1081
Oleoyl coenzyme A dipotassium salt	ThermoFisher	Cat# CN10090405
sn-Glycerol 3-phosphate lithium salt	Sigma-Aldrich	Cat# 94124-10MG

Glycerol, L-[14C(U)] 3-phosphate disodium salt	American Radiolabeled Chemicals	Cat# ARC 0316-50 µCi
Sodium butyrate	Sigma-Aldrich	Cat# 303410-500G
HCS LipidTOX™ Deep Red Neutral Lipid Stain, for cellular imaging	Fisher Scientific	Cat# H34477
Clean-Blot™ IP Detection Reagent (HRP)	Thermo Scientific	Cat# 21230
AICAR	Abcam	Cat# ab120358
Critical commercial assays		
Pierce™ Coomassie (Bradford) Protein Assay Kit	ThermoFisher	Cat# 23200
Pierce BCA Protein Assay Kit	ThermoFisher	Cat# 23225
Silver staining Kit	ThermoFisher	Cat# 24612
Experimental models: Cell lines		
HEK-293S GnTI⁻	ATCC	Cat# CRL-3022; RRID: CVCL_A785
CHO-K1	ATCC	Cat# CCL-61; RRID: CVCL_0214
Experimental models: Organisms/strains		
C57BL/6J mouse	Jackson Laboratory	Stock# 000664
ob/ob mouse	Jackson Laboratory	Stock# 000632
Sprague Dawley	Charles River	SAS SD code 400
Phosphatase inhibitor cocktail		
Sodium Fluoride	Sigma-Aldrich	Cat# S7920
Imidazole	Sigma-Aldrich	Cat# I5513
Sodium molybdate	Sigma-Aldrich	Cat# M1003
Sodium tartrate dihydrate	Sigma-Aldrich	Cat# S4797
Sodium pyrophosphate	Sigma-Aldrich	Cat# P8135
β-glycerophosphate	Sigma-Aldrich	Cat# G9422
Recombinant DNA		
pEZT-BM	Addgene	Plasmid #74099
ATP citrate lyase (ACLY) (NM_001096) Human Tagged ORF Clone	Origene	Cat# RC200508

ACSS2 (NM_018677.4) cDNA ORF clone, Homo sapiens (human)	Genscript	Cat# OHu18043
ACL-Tev-Flag-His-XT3	IDT DNA	N/A
ACC1-Tev-Flag-His-XT3	IDT DNA	N/A
FAS-Tev-Flag-His-XT3	IDT DNA	N/A
MIG12-Tev-Flag-His_Human	IDT DNA	N/A
S14-Tev-Flag-His_Human	IDT DNA	N/A
PEPCK_Human-Tev-His	IDT DNA	N/A
pEZT-BM-ACL-3xFlag-10xHis	This paper	N/A
pEZT-BM-ACSS2-3xFlag-10xHis	This paper	N/A
pEZT-BM-ACC1-3xFlag-10xHis	This paper	N/A
pEZT-BM-FAS-3xFlag-10xHis	This paper	N/A
pEZT-BM-MIG12-3xFlag-10xHis	This paper	N/A
pEZT-BM-S14-3xFlag-10xHis	This paper	N/A
pEZT-BM-PEPCK-3xFlag-10xHis	This paper	N/A
ACC1-pcDNA3.1	This paper	N/A
ACC2-pcDNA3.1	This paper	N/A
Software and algorithms		
Image Studio™ Version 5.0	Li-COR	N/A
Prism 10	GraphPad	https://www.graphpad.com/
Adobe Illustrator 2023	Adobe	N/A
Other		
PD-10 desalting columns	Cytiva	Cat# 17085101
Superose 6 10/300 GL column	Cytiva	Cat# GE17-5172-01
Amicon Ultra centrifugal filter	Millipore	Cat# UFC903008
Ni-NTA Resin	Thermo Scientific	Cat# 88221
Slide-A-Lyzer dialysis cassette	Thermo Scientific	Cat# 66203
Dynabeads Protein A beads	Invitrogen	Cat# 10002D
Scintillation vials	Research Products International	Cat# 121040
Disposable Round-Bottom Rimless Glass Tubes	Corning	Cat# 14-962-15E

4–15% Criterion™ TGX™ Precast Midi Protein Gel, 26 well, 15 µl	BIO-RAD	Cat# 5671085
4–15% Mini-PROTEAN® TGX™ Precast Protein Gels, 15-well, 15 µl	BIO-RAD	Cat# 4561086
Dynabeads™ Protein G for Immunoprecipitation	Fisher Scientific	Cat# 10-004-D
Slide Warmer (110 V, for 23 slides)	Fisher Scientific	Cat# 50-930-986

Methods

Experimental model and study participant details

Animals and primary hepatocytes

Primary hepatocytes were isolated from 4-week-old male Sprague–Dawley rats fasted for 12 hours and refed for 12 hours with a fat-free diet, using collagenase perfusion as described by Shimomura et al. [73] All rat studies were performed under an approved protocol from the UT Southwestern Institutional Animal Care and Research Advisory Committee (IACUC #2015-100838). Liver tissues were collected from wild-type male C57BL/6 mice, liver-specific ACC1 and/or ACC2 knockout male C57BL/6 mice, or ob/ob male C57BL/6 mice. All mice were maintained under standard housing conditions with a 12-hour light/dark cycle (lights on at 6 AM and off at 6 PM). Mouse studies were conducted under UT Southwestern IACUC approval (2015-101135-G). Nocodazole and taxol were used at 20 µM to achieve robust disruption or stabilization of microtubules, respectively, in primary hepatocytes.

Method Details

Antibody generation

For the generation of the mouse monoclonal antibody that recognizes mouse ACC1, a cDNA encoding the N-terminal 95 amino acids of mouse ACC1 was cloned into a pGEX-4T-1 vector and the recombinant protein was expressed in *Escherichia coli (E. coli)* and purified using glutathione agarose. Monoclonal antibodies were generated by fusion of Sp2/mIL-6 (CRL-2016; ATCC) mouse myeloma cells with splenic B lymphocytes derived from a female NZBWF1/J mouse injected with purified mouse ACC1 protein [74,75].

For the generation of a rabbit polyclonal antibody against mouse ACC2, a cDNA encoding the N-terminal 200 amino acids of mouse ACC2 was cloned into a pRSF-1b expression vector and

expressed in *E. coli*. The purified His-tagged protein was isolated using nickel affinity resin and
injected into rabbits as described [76].

For the generation of rabbit polyclonal antibody against human FAS, a cDNA encoding the C-
terminal 305 amino acids from human FAS was cloned into pGEX-4T-1 expression vector and
expressed in *E. coli.* The purified GST-fusion protein was injected into rabbits as described [19].

For the generation of a rabbit polyclonal antibody against mouse S14, recombinant (His)6-
mouse S14 was expressed in E. coli, purified by Ni-NTA affinity chromatography (Qiagen)
according to the manufacturer's protocol, and injected into rabbits as described [19].

For the generation of rabbit polyclonal antibody against mouse GPAM, a cDNA encoding the N-
terminal 150 amino acids from mouse GPAM was cloned into pRSF-1b expression and
expressed in *E. coli*. The purified His-tagged protein was injected into rabbits as described [19].

For the generation of mouse monoclonal antibody against rat GPAM, a cDNA encoding amino
acids 627 to 717 of rat GPAM was cloned into pRSF-1b expression vector and expressed in *E.*
coli. The purified His-tagged protein was injected into mice, and monoclonal antibodies were
produced as described above. For the generation of mouse monoclonal antibody against mouse
MIG12, full length mouse MIG12 cDNA was cloned into pRSF-1b and expressed in *E. coli* and
the purified protein (Ni-NTA agarose) was injected into mice. Monoclonal antibodies were
generated using the same hybridoma fusion strategy described for ACC1.

Immunofluorescence

Primary hepatocytes from refed rat livers were fixed in 4% paraformaldehyde (PFA) in PHEM
buffer (pH 7.4) for 30 minutes. After rinsing the cells with 0.1 M phosphate buffer, they were
permeabilized with pre-chilled (–20°C) methanol for 15 min at –20°C. Blocking was performed
using 1% BSA and 0.2% fish skin gelatin in 0.1 M phosphate buffer (pH 7.4) for 30 minutes at
room temperature. Primary antibodies were incubated overnight at 4°C in the blocking solution,
and Alexa Fluor 488- or 568-labeled secondary antibodies were used to visualize the
localization of each target protein. Imaging was conducted using a Zeiss LSM880 Airyscan
microscope.

Colocalization analysis

Colocalization was quantified using the Coloc2 plugin in Fiji/ImageJ. To minimize background
noise and evaluate biologically relevant signal overlap, analysis was performed on lipid droplet–
enriched regions, which represent functional sites of lipogenic enzyme accumulation. CZI files

were imported directly into Fiji, and channels were split and processed separately to avoid
spectral bleed-through. Images used for Coloc2 analysis were converted to 8-bit format prior to
computation. Pearson's correlation coefficients were calculated using Costes automatic
thresholding, and above-threshold Pearson's R values were reported in the main text.
Thresholded Manders' coefficients (tM1 and tM2), Costes significance testing, and 2D intensity
histograms were generated by Coloc2 and are presented in Figure S1. All images were
acquired below saturation, identical ROIs were applied to paired channels for each analysis.

Pre-embedding immuno-electron microscopy

Primary hepatocytes from refed rat livers were fixed in 4% PFA/0.05% glutaraldehyde (GA) in
PHEM buffer (pH 7.4) for 30 minutes. After rinsing the cells with 0.1 M phosphate buffer,
residual aldehydes were quenched with 50 mM glycine in 0.1 M phosphate buffer for 15 minutes,
followed by permeabilization with 0.2 mg/mL saponin for 30 minutes at room temperature.
Blocking was performed using 5 µg/mL saponin, 1% BSA, 0.2% fish skin gelatin, and 5% goat
serum in 0.1 M phosphate buffer (pH 7.4) for 15 minutes at room temperature. The primary
antibody was incubated overnight at 4°C in the blocking solution. After incubation, cells were
labeled with an anti-rabbit or anti-mouse IgG secondary antibody conjugated to 1.4 nm gold.
Following post-fixation, the samples were subjected to gold enhancement, embedded in
Embed-812 resin, and cut into 60 nm sections. Electron micrographs were taken using the TEM
JEOL 1400 Plus microscope. For ACC1 and FAS double immunogold labeling, ACC1 and FAS
primary antibodies were bound to 5 nm and 10 nm colloidal gold-conjugated secondary
antibodies, respectively. After post-fixation, the cells were embedded in Embed-812 resin,
sectioned into 60 nm slices, and imaged using the JEOL 1400 Plus Transmission Electron
Microscope. The pre-embedding method followed the procedure described by Yamamoto *et al*.
[77]

Liver subcellular fractionation

Fresh rat or mouse liver was homogenized in 50 mM Tris (pH 7.4) and 250 mM sucrose in the
presence of protease inhibitors. For lipid droplet isolation, half of the homogenate was
transferred to a new tube, and buffer A (20 mM HEPES, pH 7.4, 100 mM KCl, 2 mM $MgCl_2$) was
gently added to the top. The sample was centrifuged at 2,000 × g for 30 minutes, and the
floating lipid layer was transferred to a new tube. This lipid layer was mixed with an equal
volume of buffer A-500S (20 mM HEPES, pH 7.4, 100 mM KCl, 2 mM $MgCl_2$, and 500 mM
sucrose) and gently overlaid with buffer A. This step was repeated twice to isolate a pure lipid

layer. A glass pipette was used during lipid layer purification to minimize loss of lipid. Proteins

from the lipid layer were extracted using an organic solvent as described [78] and resuspended in

SDS-lysis buffer (50 mM Tris, pH 6.8, 50 mM NaCl, 1% SDS). For mitochondria and microsome

fractionations, the remaining half of the liver homogenate was centrifuged at 1,000 × g for 10

minutes to remove the nuclear fraction. The supernatant was centrifuged at 10,000 × g for 20

minutes to pellet the mitochondrial fraction. The mitochondrial pellet was resuspended and

washed twice more with homogenization buffer by repeated centrifugation at 10,000 × g for 20

minutes. The final mitochondrial pellet was resuspended in SDS-lysis buffer. The supernatant

from the 10,000 × g centrifugation was further centrifuged at 100,000 × g for 30 minutes to

sediment the microsomal fraction, which contains the endoplasmic reticulum. The microsomal

pellet was rinsed with homogenization buffer and resuspended in SDS-lysis buffer.

745   Recombinant protein expression and purification

746   Expression

The cDNAs encoding human ACSS2, ACL, ACC1, FAS, MIG12, S14, and PEPCK were cloned

into the pEZT-BM vector containing a TEV protease cleavage site, an SSG linker, followed by a

C-terminal 3 × Flag tag and 10 × His tag. All proteins were expressed in mammalian HEK 293S

GnTI⁻ cells (ATCC). A total of 500 mL of HEK 293S GnTI⁻ cells were transfected at a density

of 2–3 × 10⁶ cells/mL with 0.75 mg of plasmid DNA using PEI 25K (Polysciences). 16

hours post-transfection, 250 mL of Freestyle 293 expression medium (Gibco) supplemented

with 2% fetal calf serum (FCS) and 1% penicillin-streptomycin solution (Corning) was added to

the cultures. Sodium butyrate was added to a final concentration of 8 mM to enhance protein

expression. Cells were harvested 72 hours post-transfection for subsequent protein

purification.

757   Purification

The cell pellet from 400 mL culture was resuspended and homogenized in 100 mL buffer B (50

mM phosphate buffer, 500 mM NaCl, 1 mM DTT, 5 mM imidazole, 225 mM mannitol, 75 mM

sucrose, protease inhibitor cocktail (Millipore-Sigma), protease inhibitor cocktail for use in

purification of Histidine-tagged proteins (Sigma-Aldrich), pH 8.0). The homogenate was

centrifuged at 135,000 × g for 45 minutes at 4 °C. The supernatant was collected and

filtered through a 5 µm filter. The filtered supernatant was loaded on a 2 mL Ni-NTA resin

(Thermo Scientific) that pre-equilibrated with buffer B and flowed through by gravity (~1 ml/min)

at 4°C. The Ni-NTA was then washed sequentially with 10 CV buffer B and 25 CV buffer C (50

mM phosphate buffer, 500 mM NaCl, 1 mM DTT, 50 mM imidazole, 10% glycerol, pH 8.0) by

gravity. Proteins were eluted in 5 CV buffer D (50 mM phosphate buffer, 500 mM NaCl, 1 mM

DTT, 500 mM Imidazole, 10% glycerol, pH 8.0). The eluate was desalted using a PD-10

desalting column (Cytiva) pre-equilibrated with buffer E (10 mM HEPES, 50 mM NaCl,

100 mM KCl, 1 mM DTT, 5% glycerol, pH 7.4). Proteins were eluted in buffer E and incubated

with TEV protease overnight at 4 °C.

Removal of TEV and tags

rACL, rACC1, and rFAS: the TEV-cleaved proteins were concentrated to 0.5 mL using a

30,000 MWCO Amicon Ultra centrifugal filter (Millipore) and then subjected to gel-filtration

chromatography on a Superose 6 10/300 GL column (Cytiva) pre-equilibrated with buffer E.

Purified proteins eluted as a single sharp peak in buffer E. Protein homogeneity was confirmed

by Coomassie blue-stained SDS-PAGE.

rACSS2, rMIG12, rS14, and rPEPCK: the TEV-cleaved proteins were loaded on 2 mL Ni-NTA

resin and flowed through by gravity (~1 ml/min) pre-equilibrated with buffer E. The Ni-NTA resin

was then washed sequentially with 5 CV buffer B, 5 CV buffer C and 5 CV buffer D by gravity

flow. The flow-through and eluted fractions without His tag signal were combined and

concentrated using an Amicon Ultra centrifugal filter (Millipore) (rACSS2 and rPEPCK: 30,000

MWCO, rMIG12: 10,000 MWCO, rS14: 3,000 MWCO) and then dialyzed using a 2,000 MWCO

Slide-A-Lyzer dialysis cassette (Thermo Scientific) in 2 L buffer E overnight at 4°C.

Co-immunoprecipitation of recombinant proteins

Recombinant ACC1, FAS, ACL, ACSS2, MIG12, S14, and PEPCK were mixed at 8: 64: 20: 40:

32: 96: 1 molar ratios, respectively. Groups lacking recombinant ACC1 served as controls. The

protein mixtures were incubated with 40 mM HEPES (pH 7.1), 1 mM $MnCl_2$, and λ-phosphatase

(New England Biolabs) for 30 minutes at 30°C. For phosphorylated groups, the mixtures were

incubated with a phosphatase inhibitor cocktail instead of lambda phosphatase, under the same

conditions. The reaction systems were then supplemented with 0.5 mM Na-citrate buffer (pH

7.1) and incubated for 20 minutes at 30°C.

After pre-incubation, 5 µg of ACC1 polyclonal antibody was added to each reaction and

incubated for an additional 10 minutes at 30°C. Buffer F (50 mM NaCl, 100 mM KCl, 20 mM

HEPES, 4 mM $MgCl_2$, 1% LMNG, pH 7.1) was added, and the mixtures were incubated with

rotation for 1 hour at room temperature. Subsequently, 25 µL Dynabeads Protein A (Invitrogen)

were added and incubated with rotation for 20 minutes at room temperature. The beads were
washed four times with buffer F. Bound proteins were eluted by incubation with 1× SDS sample
buffer for 10 minutes at 95°C. Eluted proteins were analyzed by immunoblot analysis using the
indicated antibodies.

Co-immunoprecipitation of liver lysates

Liver slices from fresh mouse (or rat) livers were homogenized in 10 mM Tris (pH 7.1), 250 mM
sucrose, and 1% LMNG (Lauryl maltose neopentyl glycol) in the presence of protease inhibitors.
The homogenate was centrifuged at 25,600 × g for 15 minutes at 20°C. The supernatant was
collected and filtered through a 0.45 μm filter. The clarified supernatant was divided into two
tubes; 5 mM $MgCl_2$ was added to both, whereas phosphatase inhibitor cocktail was added to
only one. After incubation at 30°C for 30 minutes, NaCl and KCl were added to final
concentrations of 50 mM and 100 mM, respectively. Mouse monoclonal ACC1 antibody or
mouse IgG control antibody was added to each tube, and the samples were incubated for 1
hour at room temperature. Dynabeads Protein G (25 μL) were then added and incubated for a
further 25 minutes at room temperature. The beads were washed 5 times with 10 mM Tris (pH
7.1), 100 mM KCl, 50 mM NaCl, 5 mM $MgCl_2$, and 1% LMNG. The beads were boiled in SDS-
loading buffer for 5 minutes to elute proteins co-immunoprecipitated with ACC1.

Mass Spectrometry

815 Immunoprecipitates were resolved by SDS–PAGE, Coomassie-stained, and the excised gel
816 regions were submitted to the UT Southwestern Proteomics Core Facility for LC–MS/MS
817 analysis. Protein identification and label-free quantification were performed using the core's
818 standard Complex Mixture ID workflow, and processed datasets were used for downstream
819 analyses.

Radiometric FA synthesis in hepatocytes

Rat primary hepatocytes were cultured in 6-well dishes coated with collagen. Where indicated,
after 8 hours of culture, cells were treated with DMSO, nocodazole, or taxol (20 μM). On the day
of study, the culture medium was replaced with medium containing 1 mM acetate or citrate,
supplemented with 1 μCi [^{14}C]-labeled acetate or citrate, respectively, and cells were harvested
after the indicated incubation times by adding 1 mL of 0.1 N NaOH. A recovery carrier (^{3}H oleic
acid), 1.5 mL of ethanol, and 0.75 mL of 75% KOH were added to the harvested cells, and the
mixture was autoclaved for 45 minutes on slow exhaust for saponification. After saponification,

1.5 mL of ethanol and 1.5 mL of HCl were added sequentially with vigorous vortexing. After

cooling, 1.5 mL of ethanol was added, and fatty acids were extracted three times with 3 mL of

hexane. The solvent was evaporated under nitrogen, and the residue was re-dissolved in 30 µL

of chloroform. Samples were spotted onto plastic-backed silica gel TLC plates and resolved in a

solvent system consisting of heptanes:diethyl ether:acetic acid (90:30:1, v/v/v). After

visualization with iodine vapor, each FA spot was excised and radioactivity was quantified by

scintillation counting. FA synthesis was calculated based on the recovered radioactivity.

Recombinant protein enzyme assays

Dephosphorylation treatment

Different recombinant proteins were pre-incubated with 20 mM HEPES (pH 7.1), 1 mM $MnCl_2$,

and lambda phosphatase for 30 minutes at 30°C in 96 well plates (Corning). The same groups

without lambda phosphatase were used as phosphorylated groups.

ACSS2 and ACL (LC-MS/MS)

For the ACL activity assay, protein samples from dephosphorylation treatment were incubated

at 37°C in a reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA,

4 mM sodium citrate (pH 7.1), 5 mM Mg-ATP (pH 7.1), and 0.1 mM CoA. For the ACSS2

activity assay, protein samples were incubated at 37°C with 20 mM HEPES (pH 7.1), 1 mM

DTT, 1 mg/mL BSA, 4 mM sodium acetate (pH 7.1), 5 mM Mg-ATP (pH 7.1), and 0.1 mM

CoA. At designated time points (0, 4, 8, 16, and 32 minutes), reactions were terminated by the

addition of 8 N HCl, followed by thorough mixing. Each sample was then immediately spiked

with a [1-^{13}C]acetyl-CoA internal standard prior to further analysis.

Samples were centrifuged at 4°C for 10 minutes and 150 µL supernatant was loaded on an

Oasis HLB 1cc-30 mg solid-phase extraction column. The column was washed with water and

methanol. The eluent was lyophilized and dissolved in 50 µL of mobile phase of which 1 µL was

subjected to LC/MS/MS analyses. Analysis was performed on an API 3200 triple quadrupole

LC/MS/MS mass spectrometer (Applied Biosystems/Sciex Instruments) in positive electrospray

ionization mode. The mass spectrometer was equipped with a Shimadzu LC-20AD liquid

chromatograph (LC) and a SIL-20ACHT auto sampler. Chromatography was performed on a

reverse-phase C18 column (Waters XBridge, 150 x 2.1 mm, 3 µm) with a mobile phase

consisting of water/methanol (95:5, v/v) and 4mM dibutylamine acetate (eluent A), and

water/acetonitrile (25:75, v/v) (eluent B). Multi-reaction monitoring (MRM) was carried out for the

detection of the short-chain acyl-CoAs in standard solutions and biological samples. Acetyl-CoA

was quantitated by comparison of the individual ion peak area with that of the internal standard.

The analytical data were processed by Analyst Software (version 1.6.2).

ACC1 (radiometric)

Following pre-incubation in dephosphorylation treatment step, rACC1 was incubated at 37°C in

a reaction mixture containing 20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg/mL BSA, 5 mM

NaHCO$_3$, 5 mM Mg-ATP (pH 7.1), 0.25 mM acetyl-CoA, and 0.225 µCi [^{1}□C]NaHCO$_3$

(American Radiolabeled Chemicals). At designated time points (0, 4, 8, 16, and 32 minutes),

reactions were terminated by the addition of 8 N HCl. The mixtures were transferred to

scintillation vials (Research Products International Corp.) and air-dried completely at 100°C.

Dried samples were then dissolved in 1 mL of methanol, and [^{1}□C] radioactivity was quantified

by liquid scintillation counting (Beckman Coulter) following the addition of scintillation cocktail

(Research Products International Corp.).

FAS (radiometric)

Following pre-incubation in dephosphorylation treatment step, rFAS was incubated at 37°C in a

reaction mixture containing 20□mM HEPES (pH□7.1), 1□mM DTT, 1□mg/mL BSA, 0.1□mM

acetyl-CoA, 0.3□mM malonyl-CoA, 0.5□mM NADPH, and 0.02 µCi [^{1}□C]malonyl-CoA

(American Radiolabeled Chemicals). At designated time points (0, 4, 8, 16, and 32 minutes),

reactions were terminated by adding HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes

(Corning). Ethanol and hexane were added at a volume ratio of 1:1:2

(HCl/methanol:ethanol:hexane), and the mixtures were vortexed thoroughly. Samples were

centrifuged at 2,000 × g for 5 minutes at room temperature. The upper organic phase was

transferred to scintillation vials and air-dried completely at RT. Dried lipids were dissolved in

1□mL of methanol, and [^{1}□C] radioactivity was quantified by liquid scintillation counting

(Beckman Coulter) following the addition of scintillation cocktail (Research Products

International Corp.).

ACL-ACC1-FAS (radiometric)

After pre-incubation in dephosphorylation treatment step, recombinant ACL (0.8 µg), ACC1 (0.4

µg), and FAS (3.2 µg) were incubated at 37 °C in a reaction mixture containing 20 mM HEPES

(pH 7.1), 1 mM DTT, 1 mg ml^{-1} BSA, 4 mM sodium citrate (pH 7.1), 5 mM NaHCO$_3$, 5 mM Mg-

ATP (pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.025 µCi [1,5-^{1}□C]citric acid (Revvity). At

designated time points (0, 4, 8, 16, and 32 minutes), reactions were terminated by adding
HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning). Ethanol and hexane were added
at a volume ratio of 1:1:2 (HCl/methanol:ethanol:hexane), and the mixtures were vortexed
thoroughly. Samples were centrifuged at 2,000 × g for 5 minutes at room temperature. The
upper organic phase was transferred to scintillation vials and air-dried completely at RT. Dried
lipids were dissolved in 1 mL of methanol, and [^{1}C] radioactivity was quantified by liquid
scintillation counting (Beckman Coulter) following the addition of scintillation cocktail (Research
Products International Corp.).

ACSS2-ACC1-FAS (radiometric)

Following pre-incubation in dephosphorylation treatment step, recombinant ACSS2 (0.4 μg),
ACC1 (0.4 μg), and FAS (3.2 μg) were incubated at 37 °C in a reaction mixture containing
20 mM HEPES (pH 7.1), 1 mM DTT, 1 mg ml^{-1} BSA, 4 mM sodium acetate (pH 7.1),
5 mM NaHCO$_3$, 5 mM Mg-ATP (pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.0225 μCi
[1,2-^{1}C]acetic acid (Revvity). At designated time points (0, 4, 8, 16, and 32 minutes), reactions
were terminated by adding HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning).
Ethanol and hexane were added at a volume ratio of 1:1:2 (HCl/methanol:ethanol:hexane), and
the mixtures were vortexed thoroughly. Samples were centrifuged at 2,000 × g for 5 minutes at
room temperature. The upper organic phase was transferred to scintillation vials and air-dried
completely at RT. Dried lipids were dissolved in 1 mL of methanol, and [^{1}C] radioactivity was
quantified by liquid scintillation counting (Beckman Coulter) following the addition of scintillation
cocktail (Research Products International Corp.).

Cytosol-based citrate-to-palmitate assay

Mouse livers collected after fasting-refeeding treatment were homogenized in Buffer I (20 mM
HEPES, 250 mM sucrose, protease inhibitor cocktail (Millipore-Sigma), pH 7.1) at a liver-to-
buffer ratio of 1:5 (w/v). Homogenates were centrifuged at 6,000 × g for 10 minutes at 4 °C.
The resulting supernatant was subjected to further centrifugation at 180,000 × g for
60 minutes at 4 °C. The supernatant was then loaded on PD-10 desalting column (Cytiva) that
pre-equilibrated with buffer J (20 mM HEPES, 50 mM KCl, 1 mM DTT, and protease inhibitor
cocktail (Millipore-Sigma), pH 7.1) and flowed through by gravity. Liver cytosols were eluted in
buffer J.

For pre-incubation, 300 µg of liver cytosol was incubated with 20 mM HEPES (pH 7.1),
1 mM $MnCl_2$, and phosphatase inhibitor cocktail at 30 °C for 15 minutes. Parallel samples
without phosphatase inhibitor cocktail served as the dephosphorylated groups.

Following pre-incubation, liver cytosol samples were incubated at 37 °C for 15 minutes in a
reaction mixture containing 20 mM HEPES (pH 7.1), 5 mM $MgCl_2$, 1 mM DTT, 1 mg ml^{-1}
BSA, varying concentrations of sodium citrate (pH 7.1), 5 mM $NaHCO_3$, 5 mM ATP
(disodium salt, pH 7.1), 0.1 mM CoA, 1 mM NADPH, and 0.05 µCi [1,5-^{14}C]citric acid
(Revvity). Reactions were terminated by adding 1 N HCl/methanol (6:4, v/v) in 16 × 100 mm
glass tubes (Corning), followed by vortexing with ethanol and hexane. The final solvent ratio of
quenching solution:ethanol:hexane was 1:1:2. Samples were centrifuged at 2,000 × g for
5 minutes at room temperature (RT), and the upper organic phase was collected, transferred to
scintillation vials, and air dried completely at RT. Dried lipids were dissolved in 1 mL methanol,
and [^{14}C] radioactivity was quantified by liquid scintillation counting (Beckman Coulter) following
the addition of scintillation cocktail (Research Products International Corp.).

TG synthesis in liver homogenates

The livers of fasted/refed rats were homogenized in buffer K (10 mM HEPES, 1 mM EDTA, 1
mM DTT, and protease inhibitor cocktail (Millipore-Sigma), pH 7.1) at a tissue-to-buffer ratio of
1:4 (w/v). The homogenate was filtered through gauze to remove unbroken tissue and then
dialyzed against dialysis buffer (10 mM HEPES, 50 mM KCl, 1 mM DTT, 1 mM PMSF,
pH 7.1) for 2 hours.

The liver homogenates were pre-incubated with various concentrations of $MgCl_2$ (0, 2.5, 5, 10,
20 mM), 4 mM of Na-citrate buffer (pH 7.1), a phosphatase inhibitor cocktail, and 50 mM
phosphate buffer (pH 7.1) for 5 minutes at 37°C. Parallel groups without phosphatase inhibitor
cocktail and phosphate buffer served as dephosphorylated controls. After pre-incubation,
samples were incubated for 30 minutes at 37 °C in a reaction mixture containing 1 mM DTT,
1 mg/mL BSA, 10 mM oleoyl-CoA, 100 mM glycerol 3-phosphate, and 0.2 µCi [^{14}C]glycerol
3-phosphate (American Radiolabeled Chemicals). Reactions were terminated by adding 1 N
HCl/methanol (6:4, v/v) in 16 × 100 mm glass tubes (Corning). The mixtures were then
vortexed with phosphate-buffered saline (PBS) and Folch reagent (chloroform:methanol = 2:1,
v/v). The final solvent ratio of HCl/methanol:PBS:Folch was 1:1:3. Samples were centrifuged at
2,000 × g for 5 minutes at room temperature (RT), and the lower organic phases were
transferred to new glass tubes and air dried completely at RT. Dried lipids were reconstituted in

50 μL chloroform and vortexed. Extracted lipids, along with a TG marker, were loaded onto

thin-layer chromatography (TLC) plates. Plates were developed in a TLC chamber saturated

with a solvent system of hexane:diethyl ether:acetic acid (80:20:1, v/v/v). TG dots were

visualized using iodine vapor, identified by comparison with the TG marker. TG dots were

excised from the plates and transferred to scintillation vials. Scintillation cocktail (Research

Products International Corp.) was added to each vial, followed by vortexing. [^{14}C] radioactivity

was quantified by liquid scintillation counting (Beckman Coulter).

Enzyme Synergy Assays (Recombinant ± Cytosol)

For dephosphorylation treatment, a mixture containing rACL (0.5 μg), rACC1 (0.25 μg), and

rFAS (2 μg) or an equal volume of protein purification buffer (Buffer E) (used for the cytosol only

groups) were incubated in a 25 μL total volume system with λ-phosphatase in the presence of 1

mM $MnCl_2$ at 30°C for 20 minutes. The resulting 22 μL reaction was combined with 20 μL of

liver cytosol (300 μg) or cytosol preparation buffer (Buffer J) (used for the recombinant protein

only groups) in a final volume of 50 μL and incubated at 30°C with 5 mM $MgCl_2$ for an additional

20 minutes.

To maintain proteins in a phosphorylated state, the same protein mixture or Buffer E was

incubated under identical conditions with phosphatase inhibitor cocktail instead of λ-

phosphatase. The resulting 22 μL reaction was then mixed with 20 μL of liver cytosol (300 μg)

or Buffer J and incubated in a 50 μL system containing 5 mM $MgCl_2$, phosphatase inhibitor

cocktail, and 50 mM phosphate buffer (pH 7.1) at 30°C for 20 minutes. Following the pre-

incubation, samples were mixed with reaction buffer containing 20 mM HEPES (pH 7.1), 1 mM

DTT, 1 mg/mL BSA, 4 mM sodium citrate (pH 7.1), 5 mM $NaHCO_3$, 5 mM Mg-ATP (pH 7.1), 0.1

mM CoA, 1 mM NADPH, and 0.025 μCi [1,5-^{14}C]citric acid (Revvity) in a total volume of 75 μL.

The reactions were incubated at 37°C for 15 minutes and terminated by adding 0.5 mL of cold

methanol in 16 × 100 mm glass tubes (Corning). Next, 1.5 mL of methyl tert-butyl ether (MTBE)

was added, and the samples were vortexed thoroughly and left to stand at RT for 5 minutes.

Then, 0.5 mL of ddH_2O was added, followed by another round of vortexing and a 5-minute

incubation at RT. Phase separation was achieved by centrifugation at 2,000 × g for 5 minutes at

RT. The upper organic phase was transferred to scintillation vials and air-dried completely at

RT. Dried lipids were dissolved in 1 mL of methanol, and [^{14}C] radioactivity was quantified by

liquid scintillation counting (Beckman Coulter) after the addition of scintillation cocktail

(Research Products International Corp.).

Phosphatase inhibitor cocktail preparation

The phosphatase inhibitor cocktail was prepared fresh using the following recipe: sodium fluoride (200 mM), imidazole (200 mM), sodium molybdate (115 mM), sodium orthovanadate (200 mM), sodium tartrate dihydrate (400 mM), sodium pyrophosphate (100 mM), and β-glycerophosphate (100 mM) in ddH$_2$O (final volume 20 mL). Sodium orthovanadate was prepared separately by dissolving 1.84 g of powder in 50 mL ddH$_2$O, adjusting the pH to 10, briefly boiling in a microwave (10 sec), cooling on ice to room temperature, and repeating the cycle three times. The solution was adjusted to 50 mL, aliquoted, and stored at –20°C until use.

Quantification and statistical analysis

Sample sizes (n) are indicated in the figure legends. Technical replicates are also noted in the figure legends when applicable. Unless otherwise stated, data are presented as mean ± SEM. Statistical analyses were performed using GraphPad Prism 10. The statistical test and model used for each experiment are specified in the corresponding figure legend. A significance threshold of $\alpha = 0.05$ was used.

For co-localization analyses, images were quantified using Fiji/ImageJ (Coloc2). Pearson's correlation coefficients were reported as values above the Costes threshold. Thresholded Manders' coefficients (tM1 and tM2), Costes significance testing, and 2D intensity histograms were generated using Coloc2 and are provided in the Supplemental Information.

Supplemental item titles

Figure S1. Representative regions of interest (ROIs) and pixel-intensity scatter plots used for colocalization analysis of ACC1, S14, and MIG12, related to Figure 1.

Figure S2. Sequential formation of the ACC1-containing lipogenic complex around LDs, related to Figure 2.

Figure S3. Electron microscope images, related to Figure 3.

Figure S4. Phosphorylated ACC levels in liver lysates from rats refed the FFD, related to Figure 4.

Figure S5. Physical association of ACC1 and ACC2 and regulation of TG synthesis by phosphorylation in an MgCl$_2$-dependent manner, related to Figure 5.

Figure S6. Lipogenic metabolon formation facilitates FA production, related to Figure 6.

Figure S7. DAVID functional annotation of ACC1-dependent interactors from ACC1, MIG12, and S14 immunoprecipitates, related to Figure 7.

Supplemental Table 1. Pearson's and Manders' colocalization coefficients together with Costes significance statistics for ACC1–MIG12, S14–MIG12, and S14–ACC1, related to Figure 1.

Supplemental Table 2. Quantitative co-localization analysis of ACC1 with cytosolic lipogenic enzymes in primary hepatocytes, related to Figure 2.

Supplemental Table 3. Original data of MS analysis of ACC1-dependent interactors from ACC1, MIG12, and S14 immunoprecipitates, related to Figure 7.

References

1. Sanders, F.W., and Griffin, J.L. (2016). De novo lipogenesis in the liver in health and disease: more than just a shunting yard for glucose. Biol Rev Camb Philos Soc *91*, 452-468. 10.1111/brv.12178.
2. Horton, J.D., Goldstein, J.L., and Brown, M.S. (2002). SREBPs:activators of the complete program of cholesterol and fatty acid synthesis in the liver. J. Clin. Invest. *109*, 1125-1131.
3. Samuel, V.T., and Shulman, G.I. (2016). The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. J. Clin. Invest. *126*, 12-22. .
4. Miao, T., Kim, J., Kang, P., Fujiwara, H., Hsu, F.F., and Bai, H. (2022). Acetyl-CoA-mediated autoacetylation of fatty acid synthase as a metabolic switch of de novo lipogenesis in Drosophila. Proc Natl Acad Sci U S A *119*, e2212220119. 10.1073/pnas.2212220119.
5. Lin, H.P., Cheng, Z.L., He, R.Y., Song, L., Tian, M.X., Zhou, L.S., Groh, B.S., Liu, W.R., Ji, M.B., Ding, C., et al. (2016). Destabilization of Fatty Acid Synthase by Acetylation Inhibits De Novo Lipogenesis and Tumor Cell Growth. Cancer Res *76*, 6924-6936. 10.1158/0008-5472.CAN-16-1597.
6. Hellerstein, M.K. (1999). De novo lipogenesis in humans: metabolic and regulatory aspects. Eur. J. Clin. Nutr. *53*, S53-65.
7. Parks, E.J., Krauss, R.M., Christiansen, M.P., Neese, R.A., and Hellerstein, M.K. (1999). Effects of a low-fat, high-carbohydrate diet on VLDL-triglyceride assembly, production, and clearance. J Clin Invest *104*, 1087-1096. 10.1172/JCI6572.
8. Schwarz, J.M., Neese, R.A., Turner, S., Dare, D., and Hellerstein, M.K. (1995). Short-term alterations in carbohydrate energy intake in humans. Striking effects on hepatic glucose production, de novo lipogenesis, lipolysis, and whole-body fuel selection. J Clin Invest *96*, 2735-2743. 10.1172/JCI118342.
9. Donnelly, K.L., Smith, C.I., Schwarzenberg, S.J., Jessurun, J., Boldt, M.D., and Parks, E.J. (2005). Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. J. Clin. Invest. *115*, 1343-1351.

10. Ferre, P., and Foufelle, F. (2010). Hepatic steatosis: a role for de novo lipogenesis and
the transcription factor SREBP-1c. Diabetes Obes Metab *12 Suppl 2*, 83-92.
10.1111/j.1463-1326.2010.01275.x.
11. Lambert, J.E., Ramos-Roman, M.A., Browning, J.D., and Parks, E.J. (2014). Increased
de novo lipogenesis is a distinct characteristic of individuals with nonalcoholic fatty liver
disease. Gastroenterology *146*, 726-735. 10.1053/j.gastro.2013.11.049.
12. Shimomura, I., Shimano, H., Korn, B.S., Bashmakov, Y., and Horton, J.D. (1998).
Nuclear sterol regulatory element-binding proteins activate genes responsible for the
entire program of unsaturated fatty acid biosynthesis in transgenic mouse liver. J. Biol.
Chem. *273*, 35299-35306.
13. Shimomura, I., Bashmakov, Y., and Horton, J.D. (1999). Increased levels of nuclear
SREBP-1c associated with fatty livers in two mouse models of diabetes mellitus. J. Biol.
Chem. *274*, 30028-30032.
14. Hudgins, L.C., Hellerstein, M., Seidman, C., Neese, R., Diakun, J., and Hirsch, J. (1996).
Human fatty acid synthesis is stimulated by a eucaloric low fat, high carbohydrate diet. J
Clin Invest *97*, 2081-2091. 10.1172/JCI118645.
15. Eslam, M., Newsome, P.N., Sarin, S.K., Anstee, Q.M., Targher, G., Romero-Gomez, M.,
Zelber-Sagi, S., Wai-Sun Wong, V., Dufour, J.F., Schattenberg, J.M., et al. (2020). A
new definition for metabolic dysfunction-associated fatty liver disease: An international
expert consensus statement. J Hepatol *73*, 202-209. 10.1016/j.jhep.2020.03.039.
16. Cohen, J.C., Horton, J.D., and Hobbs, H.H. (2011). Human fatty liver disease: old
questions and new insights. Science *332*, 1519-1523. 10.1126/science.1204265.
17. Eslam, M., Sanyal, A.J., and George, J. (2020). MAFLD: A Consensus-Driven Proposed
Nomenclature for Metabolic Associated Fatty Liver Disease. Gastroenterology *158*,
1999-2014.e1991. 10.1053/j.gastro.2019.11.312.
18. Browning, J.D., Szczepaniak, L.S., Dobbins, R., Nuremberg, P., Horton, J.D., Cohen,
J.C., Grundy, S.M., and Hobbs, H.H. (2004). Prevalence of hepatic steatosis in an urban
population in the United States: Impact of ethnicity. Hepatology *40*, 1387-1395.
19. Shimano, H., Horton, J.D., Hammer, R.E., Shimomura, I., Brown, M.S., and Goldstein,
J.L. (1996). Overproduction of cholesterol and fatty acids causes massive liver
enlargement in transgenic mice expressing truncated SREBP-1a. J. Clin. Invest. *98*,
1575-1584.
20. Horton, J.D., Bashmakov, Y., Shimomura, I., and Shimano, H. (1998). Regulation of
sterol regulatory element binding proteins in livers of fasted and refed mice. Proc. Natl.
Acad. Sci. U. S. A. *95*, 5987-5992.
21. Horton, J.D., Shimomura, I., Brown, M.S., Hammer, R.E., Goldstein, J.L., and Shimano,
H. (1998). Activation of cholesterol synthesis in preference to fatty acid synthesis in liver
and adipose tissue of transgenic mice overproducing sterol regulatory element-binding
protein-2. J. Clin. Invest. *101*, 2331-2339.
22. Horton, J.D., Shimano, H., Hamilton, R.L., Brown, M.S., and Goldstein, J.L. (1999).
Disruption of LDL receptor gene in transgenic SREBP-1a mice unmasks hyperlipidemia
resulting from production of lipid-rich VLDL. J. Clin. Invest. *103*, 1067-1076.
23. Matsuda, M., Korn, B.S., Hammer, R.E., Moon, Y.A., Komuro, R., Horton, J.D.,
Goldstein, J.L., Brown, M.S., and Shimomura, I. (2001). SREBP cleavage-activating
protein (SCAP) is required for increased lipid synthesis in liver induced by cholesterol
deprivation and insulin elevation. Genes Dev. *15*, 1206-1216.
24. Moon, Y.A., Liang, G., Xie, X., Frank-Kamenetsky, M., Fitzgerald, K., Koteliansky, V.,
Brown, M.S., Goldstein, J.L., and Horton, J.D. (2012). The Scap/SREBP pathway is
essential for developing diabetic fatty liver and carbohydrate-induced
hypertriglyceridemia in animals. Cell Metab. *15*, 240-246. 10.1016/j.cmet.2011.12.017.

25. Abu-Elheiga, L., Brinkley, W.R., Zhong, L., Chirala, S.S., Woldegiorgis, G., and Wakil,
S.J. (2000). The subcellular localization of acetyl-CoA carboxylase 2. Proc. Natl. Acad.
Sci. U. S. A. *97*, 1444-1449.

26. Man, W.C., Miyazaki, M., Chu, K., and Ntambi, J.M. (2006). Membrane topology of
mouse stearoyl-CoA desaturase 1. J Biol Chem *281*, 1251-1260.
10.1074/jbc.M508733200.

27. Moon, Y.-A., Shah, N.A., Mohapatra, S., Warrington, J.A., and Horton, J.D. (2001).
Identification of a mammalian long chain fatty acyl elongase regulated by sterol
regulatory element-binding proteins. J. Biol. Chem. *276*, 45358-45366.

28. Gonzalez-Baró, M.R., Granger, D.A., and Coleman, R.A. (2001). Mitochondrial Glycerol
Phosphate Acyltransferase Contains Two Transmembrane Domains with the Active Site
in the N-terminal Domain Facing the Cytosol. Journal of Biological Chemistry *276*,
43182-43188. 10.1074/jbc.M107885200.

29. P.A., S. (1972). Is there an organization of Krebs cycle enzymes in the mitochondrial
matrix? In: Mehlman M.A., Hanson R.W., editors. Energy Metabolism and the Regulation
of Metabolic Processes in Mitochondria. Academic Press; New York, pp. 79–91.

30. P.A., S. (1985). The metabolon. Trends Biochem. Sci., 10:109–110.

31. Bulutoglu, B., Garcia, K.E., Wu, F., Minteer, S.D., and Banta, S. (2016). Direct Evidence
for Metabolon Formation and Substrate Channeling in Recombinant TCA Cycle
Enzymes. ACS Chem Biol *11*, 2847-2853. 10.1021/acschembio.6b00523.

32. Robinson, J.B., Jr., Inman, L., Sumegi, B., and Srere, P.A. (1987). Further
characterization of the Krebs tricarboxylic acid cycle metabolon. J Biol Chem *262*, 1786-
1790.

33. Kohnhorst, C.L., Kyoung, M., Jeon, M., Schmitt, D.L., Kennedy, E.L., Ramirez, J.,
Bracey, S.M., Luu, B.T., Russell, S.J., and An, S. (2017). Identification of a multienzyme
complex for glucose metabolism in living cells. J Biol Chem *292*, 9191-9203.
10.1074/jbc.M117.783050.

34. Colbert, C.L., Kim, C.W., Moon, Y.A., Henry, L., Palnitkar, M., McKean, W.B., Fitzgerald,
K., Deisenhofer, J., Horton, J.D., and Kwon, H.J. (2010). Crystal structure of Spot 14, a
modulator of fatty acid synthesis. Proc Natl Acad Sci U S A *107*, 18820-18825.
1012736107 [pii]

10.1073/pnas.1012736107.

35. Jump, D.B., and Oppenheimer, J.H. (1985). High basal expression and 3,5,3'-
triiodothyronine regulation of messenger ribonucleic acid S14 in lipogenic tissues.
Endocrinology *117*, 2259-2266. 10.1210/endo-117-6-2259.

36. Horton, J.D., Shah, N.A., Warrington, J.A., Anderson, N.N., Park, S.W., Brown, M.S.,
and Goldstein, J.L. (2003). Combined analysis of oligonucleotide microarray data from
transgenic and knockout mice identifies direct SREBP target genes. Proc. Natl. Acad.
Sci. U. S. A. *100*, 12027-12032.

37. Mater, M.K., Thelen, A.P., Pan, D.A., and Jump, D.B. (1999). Sterol response element-
binding protein 1c (SREBP1c) is involved in the polyunsaturated fatty acid suppression
of hepatic S14 gene transcription. J. Biol. Chem. *274*, 32725-32732.

38. Ma, L., Tsatsos, N.G., and Towle, H.C. (2005). Direct role of ChREBP·Mlx in regulating
hepatic glucose-responsive genes. J. Biol. Chem. *280*, 12019-12027.
10.1074/jbc.M413063200.

39. Seelig, S., Liaw, C., Towle, H.C., and Oppenheimer, J.H. (1981). Thyroid hormone
attenuates and augments hepatic gene expression at a pretranslational level. Proc. Natl.
Acad. Sci. U. S. A. *78*, 4733-4737.

40. Zhu, Q., Anderson, G.W., Mucha, G.T., Parks, E.J., Metkowski, J.K., and Mariash, C.N. (2005). The Spot 14 protein is required for de novo lipid synthesis in the lactating mammary gland. Endocrinology *146*, 3343-3350.

41. Kim, C.W., Moon, Y.A., Park, S.W., Cheng, D., Kwon, H.J., and Horton, J.D. (2010). Induced polymerization of mammalian acetyl-CoA carboxylase by MIG12 provides a tertiary level of regulation of fatty acid synthesis. Proc. Natl. Acad. Sci. U. S. A. *107*, 9626-9631.

42. Ha, J., Daniel, S., Broyles, S.S., and Kim, K.H. (1994). Critical phosphorylation sites for acetyl-CoA carboxylase activity. J Biol Chem *269*, 22162-22168.

43. Locke, G.A., Cheng, D., Witmer, M.R., Tamura, J.K., Haque, T., Carney, R.F., Rendina, A.R., and Marcinkeviciene, J. (2008). Differential activation of recombinant human acetyl-CoA carboxylases 1 and 2 by citrate. Arch. Biochem. Biophys. *475*, 72-79. S0003-9861(08)00196-3 [pii]

10.1016/j.abb.2008.04.011 [doi].

44. Kahn, B.B., Alquier, T., Carling, D., and Hardie, D.G. (2005). AMP-activated protein kinase: Ancient energy gauge provides clues to modern understanding of metabolism. Cell Metabolism *1*, 15-25. DOI: 10.1016/j.cmet.2004.12.003.

45. Witters, L.A., Watts, T.D., Daniels, D.L., and Evans, J.L. (1988). Insulin stimulates the dephosphorylation and activation of acetyl-CoA carboxylase. Proc. Natl. Acad. Sci. U. S. A. *85*, 5473-5477.

46. Berglund, E.D., Lustig, D.G., Baheza, R.A., Hasenour, C.M., Lee-Young, R.S., Donahue, E.P., Lynes, S.E., Swift, L.L., Charron, M.J., Damon, B.M., and Wasserman, D.H. (2011). Hepatic glucagon action is essential for exercise-induced reversal of mouse fatty liver. Diabetes *60*, 2720-2729. 10.2337/db11-0455.

47. Munday, M.R., Milic, M.R., Takhar, S., Holness, M.J., and Sugden, M.C. (1991). The short-term regulation of hepatic acetyl-CoA carboxylase during starvation and re-feeding in the rat. Biochem J *280 (Pt 3)*, 733-737. 10.1042/bj2800733.

48. Berglund, E.D., Lee-Young, R.S., Lustig, D.G., Lynes, S.E., Donahue, E.P., Camacho, R.C., Meredith, M.E., Magnuson, M.A., Charron, M.J., and Wasserman, D.H. (2009). Hepatic energy state is regulated by glucagon receptor signaling in mice. J Clin Invest *119*, 2412-2422. 10.1172/jci38650.

49. Kimball, S.R., Siegfried, B.A., and Jefferson, L.S. (2004). Glucagon represses signaling through the mammalian target of rapamycin in rat liver by activating AMP-activated protein kinase. J Biol Chem *279*, 54103-54109. 10.1074/jbc.M410755200.

50. Ragolia, L., and Begum, N. (1998). Protein phosphatase-1 and insulin action. Mol Cell Biochem *182*, 49-58.

51. Kim, K.H. (1983). Regulation of acetyl-CoA carboxylase. Curr Top Cell Regul *22*, 143-176. 10.1016/b978-0-12-152822-5.50009-9.

52. Kim, K.H., Lopez-Casillas, F., Bai, D.H., Luo, X., and Pape, M.E. (1989). Role of reversible phosphorylation of acetyl-CoA carboxylase in long-chain fatty acid synthesis. FASEB J *3*, 2250-2256. 10.1096/fasebj.3.11.2570725.

53. Fediuc, S., Gaidhu, M.P., and Ceddia, R.B. (2006). Regulation of AMP-activated protein kinase and acetyl-CoA carboxylase phosphorylation by palmitate in skeletal muscle cells. J Lipid Res *47*, 412-420. 10.1194/jlr.M500438-JLR200.

54. Wang, Y., Viscarra, J., Kim, S.-J., and Sul, H.S. (2015). Transcriptional regulation of hepatic lipogenesis. Nature reviews. Molecular cell biology *16*, 678-689. 10.1038/nrm4074.

55. Umlauf, E., Csaszar, E., Moertelmaier, M., Schuetz, G.J., Parton, R.G., and Prohaska, R. (2004). Association of stomatin with lipid bodies. J Biol Chem *279*, 23699-23709. 10.1074/jbc.M310546200.

56. Liu, P., Ying, Y., Zhao, Y., Mundy, D.I., Zhu, M., and Anderson, R.G. (2004). Chinese hamster ovary K2 cell lipid droplets appear to be metabolic organelles involved in membrane traffic. J Biol Chem *279*, 3787-3792. 10.1074/jbc.M311945200.

57. Bartz, R., Zehmer, J.K., Zhu, M., Chen, Y., Serrero, G., Zhao, Y., and Liu, P. (2007). Dynamic activity of lipid droplets: protein phosphorylation and GTP-mediated protein translocation. J Proteome Res *6*, 3256-3265. 10.1021/pr070158j.

58. Beller, M., Riedel, D., Jansch, L., Dieterich, G., Wehland, J., Jackle, H., and Kuhnlein, R.P. (2006). Characterization of the Drosophila lipid droplet subproteome. Molecular & cellular proteomics : MCP *5*, 1082-1094. 10.1074/mcp.M600011-MCP200.

59. Wan, H.C., Melo, R.C., Jin, Z., Dvorak, A.M., and Weller, P.F. (2007). Roles and origins of leukocyte lipid bodies: proteomic and ultrastructural studies. FASEB J *21*, 167-178. 10.1096/fj.06-6711com.

60. Lyubarev, A.E., and Kurganov, B.I. (1989). Supramolecular organization of tricarboxylic acid cycle enzymes. Biosystems *22*, 91-102. 10.1016/0303-2647(89)90038-5.

61. Kurganov B. I, L.A.E. (1988). Hypothetical structure of the complex of glycolytic enzymes (glycolytic metabolon), formed on the membrane of erythrocytes. Molek. Biologia. *V.22, No.6*, p. 1605–1613.

62. Graham, J.W., Williams, T.C., Morgan, M., Fernie, A.R., Ratcliffe, R.G., and Sweetlove, L.J. (2007). Glycolytic enzymes associate dynamically with mitochondria in response to respiratory demand and support substrate channeling. Plant Cell *19*, 3723-3738. 10.1105/tpc.107.053371.

63. Srere, P.A. (1974). Controls of citrate synthase activity. Life Sci *15*, 1695-1710. 10.1016/0024-3205(74)90172-6.

64. Webb, B.A., Dosey, A.M., Wittmann, T., Kollman, J.M., and Barber, D.L. (2017). The glycolytic enzyme phosphofructokinase-1 assembles into filaments. J Cell Biol *216*, 2305-2313. 10.1083/jcb.201701084.

65. Kastritis, P.L., O'Reilly, F.J., Bock, T., Li, Y., Rogon, M.Z., Buczak, K., Romanov, N., Betts, M.J., Bui, K.H., Hagen, W.J., et al. (2017). Capturing protein communities by structural proteomics in a thermophilic eukaryote. Mol Syst Biol *13*, 936. 10.15252/msb.20167412.

66. Suresh, H.G., da Silveira Dos Santos, A.X., Kukulski, W., Tyedmers, J., Riezman, H., Bukau, B., and Mogk, A. (2015). Prolonged starvation drives reversible sequestration of lipid biosynthetic enzymes and organelle reorganization in Saccharomyces cerevisiae. Mol Biol Cell *26*, 1601-1615. 10.1091/mbc.E14-11-1559.

67. Wijesinghe, K.M., Kim, C.W., Schad, E.O., Li, S., Chen, S., Takeshima, E., Khandwala, C.B., Tillo, D., Lebensohn, A.M., Olzmann, J.A., et al. (2025). Plasma membrane accessible cholesterol is regulated by ACC1 and lipid droplets. bioRxiv. 10.1101/2025.08.21.671640.

68. Zhou, F., Zhang, Y., Zhu, Y., Zhou, Q., Shi, Y., and Hu, Q. (2024). Filament structures unveil the dynamic organization of human acetyl-CoA carboxylase. Sci Adv *10*, eado4880. 10.1126/sciadv.ado4880.

69. Hunkeler, M., Hagmann, A., Stuttfeld, E., Chami, M., Guri, Y., Stahlberg, H., and Maier, T. (2018). Structural basis for regulation of human acetyl-CoA carboxylase. Nature *558*, 470-474. 10.1038/s41586-018-0201-4.

70. Møller, B.L. (2010). Plant science. Dynamic metabolons. Science *330*, 1328-1329. 10.1126/science.1194971.

71. Kim, C.W., Addy, C., Kusunoki, J., Anderson, N.N., Deja, S., Fu, X., Burgess, S.C., Li, C., Ruddy, M., Chakravarthy, M., et al. (2017). Acetyl CoA Carboxylase Inhibition Reduces Hepatic Steatosis but Elevates Plasma Triglycerides in Mice and Humans: A Bedside to Bench Investigation. Cell Metab *26*, 394-406. e396. 10.1016/j.cmet.2017.07.009.

72. Kim, C.W., Moon, Y.A., Park, S.W., Cheng, D., Kwon, H.J., and Horton, J.D. (2010).
Induced polymerization of mammalian acetyl-CoA carboxylase by MIG12 provides a
tertiary level of regulation of fatty acid synthesis. Proc Natl Acad Sci U S A *107*, 9626-
9631. 10.1073/pnas.1001292107.
73. Shimomura, I., Hammer, R.E., Ikemoto, S., Brown, M.S., and Goldstein, J.L. (1999).
Leptin reverses insulin resistance and diabetes mellitus in mice with congenital
lipodystrophy. Nature *401*, 73-76.
74. Lagace, T.A., Curtis, D.E., Garuti, R., McNutt, M.C., Park, S.W., Prather, H.B.,
Anderson, N.N., Ho, Y.K., Hammer, R.E., and Horton, J.D. (2006). Secreted PCSK9
decreases the number of LDL receptors in hepatocytes and in livers of parabiotic mice. J
Clin Invest *116*, 2995-3005. 10.1172/jci29383.
75. Herz, J., Kowal, R.C., Ho, Y.K., Brown, M.S., and Goldstein, J.L. (1990). Low density
lipoprotein receptor-related protein mediates endocytosis of monoclonal antibodies in
cultured cells and rabbit liver. J Biol Chem *265*, 21355-21362.
76. Engelking, L.J., Kuriyama, H., Hammer, R.E., Horton, J.D., Brown, M.S., Goldstein, J.L.,
and Liang, G. (2004). Overexpression of Insig-1 in the livers of transgenic mice inhibits
SREBP processing and reduces insulin-stimulated lipogenesis. J. Clin. Invest. *113*,
1168-1175.
77. Yamamoto, A., and Masaki, R. (2010). Pre-embedding nanogold silver and gold
intensification. Methods Mol Biol *657*, 225-235. 10.1007/978-1-60761-783-9_18.
78. Ding, Y., Zhang, S., Yang, L., Na, H., Zhang, P., Zhang, H., Wang, Y., Chen, Y., Yu, J.,
Huo, C., et al. (2013). Isolating lipid droplets from multiple species. Nat Protoc *8*, 43-51.
10.1038/nprot.2012.142.

1275

1276

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7