Title: *Tbx1* Heterozygosity in the Oligodendrocyte Lineage Shifts Myelinated Axon
Composition in the Mouse Fimbria Without Behavioral Impairments

Authors:

Anne Marie Wells, PhD[1,2+], Takaki Tanifuji, MD, PhD[2+], Takeshi Takano, PhD[2,3+], Arumu Endo,
MD.[2], Gina Kang, B.Sc.[2], Marisa Esparza, B.Sc.[2], Qian Shi, PhD[4], Manzoor A. Bhat, PhD[4],
Noboru Hiroi, PhD[2, 4,5**]

AM Wells: wellsa1@livemail.uthscsa.edu
T Tanifuji: kjh.goodby@gmail.com
T. Takano: takanot@uthscsa.edu
A. Endo: endo@uthscsa.edu
G. Kang: kangg@uthscsa.edu
M. Esparza: esparzam1@uthscsa.edu
Q. Shi: shiQ@uthscsa.edu
M. Bhat: bhatm@uthscsa.edu
N. Hiroi: hiroi@uthscsa.edu
18

+ These authors equally contributed to the work.

[1]South Texas Medical Scientist Training Program (STX-MSTP), [2]Department of Pharmacology,
[4]Department of Cellular and Integrative Physiology, [5]Department of Cell Systems and Anatomy,
University of Texas Health Science Center, San Antonio, TX 78229, USA; [3]Tokyo Denki
University, Saitama 350-0394, Japan

ORCID: 0000-0002-6846-5969

**Corresponding Author
Noboru Hiroi, PhD
hiroi@uthscsa.edu
Department of Pharmacology
Department of Cellular and Integrative Physiology
Department of Cell Systems and Anatomy
University of Texas Health Science Center, San Antonio
Room 211B
7703 Floyd Curl Drive
San Antonio, TX 78229
+1 210 567 4169 (tel)
36

37 # of figures: 9
38 # of supplementary figures: 2
39 # of supplementary tables: 3
40
41

ABSTRACT

Constitutive heterozygosity of *Tbx1*, a T-box transcription factor gene located within the 22q11.2 deletion region, results in behavioral deficits and altered composition of myelinated axons in the fimbria, together with reduced levels of an oligodendrocyte precursor cell (OPC) marker, in mice. However, the cellular origins of these effects and the extent to which axonal changes causally contribute to behavioral impairments remain unclear. We hypothesized that *Tbx1* deficiency specifically within the oligodendrocyte lineage contributes to myelin and behavioral phenotypes. To test this hypothesis, we first demonstrated through *in vitro* siRNA knockdown that *Tbx1* regulates both OPCs and mature oligodendrocytes. Subsequently, we assessed the impact of *Tbx1* heterozygosity initiated in OPCs on behavioral and myelin phenotypes in male conditional PdgfrαCre;*Tbx1*$^{+/flox}$ mice. These mice exhibited Cre-mediated recombination in Pdgfrα-expressing brain regions and in the OPC progeny within the fimbria. At one month of age, the mutants displayed a higher rate of spontaneous alternation at the longest inter-trial interval in the T-maze compared to their wild-type littermates—an effect that was dissipated at two months. No significant phenotypic abnormalities were observed in conditional PdgfrαCre;*Tbx1*$^{+/flox}$ mice regarding neonatal ultrasonic vocalizations, social interaction, novel object approach, anxiety-like behavior (elevated plus maze), or open-field locomotion and thigmotaxis. Electron microscopic analysis revealed a compositional shift in myelinated axons within the fimbria of adult male mutants, characterized by an increased number of myelinated axons in the 300–800 nm diameter range and a decreased number in the ~1,200 nm and ~1,400 nm ranges, with myelin thickness remaining unchanged across diameters. These findings indicate that *Tbx1* heterozygosity in the oligodendrocyte lineage leads to a selective shift towards smaller myelinated axons in the fimbria and a transiently higher level of capacity for working memory and cognitive flexibility. However, it does not replicate the full spectrum of myelination abnormalities or the broader cognitive and social deficits observed in constitutive *Tbx1* heterozygotes, suggesting that *Tbx1* deficiency in non-oligodendrocyte lineage cells may lead to altered myelination and neurodevelopmental behavioral impairments.

Key Words

INTRODUCTION

Cognitive and social impairments are predictors of mental illness(1, 2). Infants later diagnosed with autism spectrum disorder (ASD) exhibit delays and deviations in various aspects of motor, social, and language development(3-11). Similarly, children who subsequently develop full-scale symptoms of schizophrenia display delayed development of social cognition, working memory, attention, and processing speed(12-16), with these deficits remaining integral to the disorder following its onset(17, 18).

These dimensional deficits in mental illness have genetic underpinnings. Copy number variations (CNVs)--which consist of deletions or duplications of up to several million base pairs at specific chromosomal loci encompassing numerous protein-coding genes—are associated with high odds ratios and penetrance rates for cognitive and social deficits(19, 20) as well as mental illnesses(21-23).

Individuals with hemizygous deletions of human chromosome 22q11.2, ranging from 1.5 Mb to 3.0 Mb, exhibit cognitive, social, and emotional impairments from childhood (24-26) and are diagnosed at elevated rates with anxiety disorders, attention-deficit hyperactivity disorder (ADHD), ASD, intellectual disability (ID), and schizophrenia(27, 28). Duplication of 22q11.2 is linked to epilepsy, ID, ADHD, and ASD at rates higher than those observed in non-carriers(29-45). However, the mechanisms through which over 40 protein-coding genes in 22q11.2 contribute to dimensional deficits and clinical diagnoses remain poorly understood in humans.

Rare inherited cases of variants in the human *TBX1* gene, a T-box transcription factor located within the 22q11.2 CNV, are associated with ASD, schizophrenia, and intellectual disability in the absence of 22q11.2 CNVs(46-51). However, due to their rarity, the statistical reliability of the association between *TBX1* variants and mental illness is limited. Moreover, these patients often carry variants in other single genes throughout the genome(49), complicating the establishment of causality from these associations.

The deletion or overexpression of small murine chromosomal regions orthologous to human chromosomes or single genes provides a complementary approach (1, 52-56). Constitutive overexpression of several hundred kilobase pair segments of the mouse ortholog of 22q11.2 recapitulates distinct social and cognitive deficits associated with 22q11.2 duplication(52, 57). Larger hemizygous deletions that encompass one of these segments result in deficits in prepulse inhibition (50). These studies in mouse models suggest the presence of driver genes for specific behavioral phenotypes within these segments.

The *Tbx1* gene is located within one of these small segments. The heterozygous deletion of *Tbx1* alone results in impairments in neonatal social communication(58-60), as well as post-pubertal and adult deficits in social interaction(58), social incentive learning(61), acoustic, but not non-acoustic, prepulse inhibition(50, 61), spatial memory acquisition in the Morris water maze, and the speed to complete simple discrimination and reversal phases of attentional set-shifting, without non-specific motor speed (62). The fimbria of post-pubertal *Tbx1* heterozygous mice displays a lack of large myelinated axons, enhanced myelination of medium-sized axons, a compositional shift of myelinated axons toward smaller sizes, and a gene expression profile indicative of defective oligodendrocyte precursor cells(62). Furthermore, the volume of the secondary motor cortex and amygdala, regions involved in the vocal network(63), is reduced in post-pubertal *Tbx1* heterozygous mice (61). Additionally, virally induced overexpression of *Tbx1* in the hippocampus negatively impacts the developmental maturation of working memory capacity(64).

The cellular and developmental origins of the effects of *Tbx1* deficiency on behavioral and
myelin phenotypes remain poorly understood. However, cells that express *Tbx1* are candidate
substrates. Recent single-cell transcriptomic analyses have revealed
that *Tbx1* is expressed in a more diverse array of cell clusters with signatures indicative
of distinct cell types in the mouse than previously indicated by *in situ* hybridization and
immunocytochemical studies. *Tbx1* is detectable in the neural tube, as well as endothelial cells,
radial glial cells, neural progenitor cells, oligodendrocyte progenitor cells, oligodendrocytes, and
various types of neurons and their progenitors (65, 66). During the embryonic period,
oligodendrocyte precursor cells emerge from embryonic neural progenitor cells in the medial
ganglionic eminence of the ventricular zones around E12.5, followed by a second wave from the
lateral ganglionic eminence at E15.5. Oligodendrocyte precursor cells continue to be generated
from neural progenitor cells in the subventricular zone during the neonatal period (67-71).
Given that single-cell transcriptomic analyses can detect rare or transitional cell types, these
studies suggest that *Tbx1* is present in radial glial cells and neural progenitor cells, as well as
their progeny, including oligodendrocyte and neuronal lineages.

Our previous work demonstrated that *Tbx1* heterozygosity initiated in neural progenitor
cells during the neonatal period recapitulates some behavioral deficits associated with
constitutive *Tbx1* heterozygosity(72). As neural progenitor cells during the neonatal period give
rise to oligodendrocyte precursor cells, oligodendrocytes, neural progenitor cells, and
neurons(67-71), we sought to determine whether the effects of *Tbx1* heterozygosity within the
oligodendrocyte lineage induces behavioral and anatomical phenotypes. Our findings indicate
that *Tbx1* heterozygosity in the oligodendrocyte lineage induces a compositional shift in
myelinated axons in the fimbria, while not altering myelin integrity or large myelinated axons, or
impairing behavior.

METHODS

Mice
All protocols for animal handling and use were approved by the Institutional Animal Care and
Use Committee (IACUC) of the University of Texas Health Science Center at San Antonio
(UTHSCSA) in accordance with National Institutes of Health (NIH) guidelines. Mice were
housed in a vivarium with a standard light phase from 7 am to 9 pm).
C57BL/6J mice: We used 1- and 2-month-old male C57BL6/J mice (Jax #000664, Jackson
Laboratory, Bar Harbor, ME) as stimulus mice for social interaction.

*PdgfrαCre;Tbx1$^{+/flox}$ m*ice: To conditionally induce *Tbx1* heterozygosity in oligodendrocyte
precursor cells, we chose a *Pdgfrα*Cre line that initiates Cre synthesis earlier than a *Cspg4*Cre
line to better induce recombination in the oligodendrocyte lineage(73). The *Pdgfrα*- promoter
drives Cre-based recombination most robustly in oligodendrocyte precursor cells (74), but also
in astrocytes, neurons, fibroblast-like cells, endothelial cells, vascular and leptomeningeal cells,
and pericytes in the mouse brain(74-77).

We crossed hemizygous *Pdgfrα*Cre mice with *Tbx1$^{+/flox}$* mice to generate *PdgfrαCre;Tbx1$^{+/flox}$*
mice. *PdgfraCre* breeder mice (C57BL/6-Tg(*Pdgfrα*-Cre)1Clc/J; Jax #013148, Jackson
Laboratory, Bar Harbor, ME) have a mixed C57BL/6J;C57BL/6N background. We backcrossed
non-congenic *Tbx1$^{+/flox}$* mice(78) to C57BL/6J for more than 10 generations and confirmed that
there was Cre-dependent recombination in a congenic *Tbx1$^{+/flox}$* mice(72). The genetic
background of *Pdgfrα*Cre and *Tbx1$^{+/flox}$* lines were mixed C57BL/6J;C57BL/6N and C57BL/6J,
respectively. These two C57BL/6 substrains have many behavioral differences(79). To avoid the
confounding effects of their unequal genetic backgrounds(80), we used only an F1 generation

derived from crosses of the *Pdgfrα*Cre and *Tbx1*$^{+/flox}$ lines. While both male and female mice
were born with all genotypes, only half as many females as males survived to complete
analyses up to 2 months of age. Because the sample sizes of females tested were not sufficient
for statistical reliability, our analyses were based on male mice only. No sex differences are
observed in social or cognitive dimensions among carriers of 22q11.2 deletion in humans (24).

*Pdgfrα*Cre*;ROSA-tdTomato* mice: We crossed hemizygous *Pdgfrα*Cre mice (C57BL/6-
Tg(*Pdgfrα*Cre)1Clc/J; Jax #013148, Jackson Laboratory, Bar Harbor, ME) with homozygous
congenic *ROSA-CAG-tdTomato* mice (B6.Cg-*Gt(ROSA)26Sor*$^{tm14(CAG-tdTomato)Hze}$/J, Jax # 007914,
Jackson Lab, Bar Harbor, ME) to generate *Pdgfrα*Cre*;ROSA-CAG-tdTomato* mice. This ROSA
line expresses tdTomato in the hippocampus without the action of Cre (
http://connectivity.brain-map.org/transgenic/experiment/81560256)

We determined the genotypes of the mice using our previously published protocol with primers
shown in **Table S1**. The sex of the mice was determined by inspection of their external genitalia.

***In vitro* cell culture of oligodendrocytes**
Lateral ventricle tissues, including the subventricular zone (SVZ), were obtained from postnatal
day 1-2 (P1-2) C57BL/6J pups. Cells were isolated in culture, with each culture derived from a
single mouse. The cells were cultured in a medium (DMEM/F12, HEPES [11320–032, Gibco,
Grand Island, NY, USA]) supplemented with N2 (17502048, Gibco, Grand Island, NY, USA), B27
(17504044, Gibco, Grand Island, NY, USA), epidermal growth factor (EGF) at a concentration of
20 ng/ml (AF100-15, Peprotech, Cranbury, NJ, USA). After two to four passages, the cells were
dissociated from the spheres using StemPro Accutase Cell Dissociation Reagent (A1110501,
Gibco, Grand Island, NY, USA). The cells were then divided into equal portions and plated on
each Poly-L-ornithine (P4957, Sigma, St. Louis, MO, USA) and Bovine Fibronectin (1030-FN,
R&D Systems, Minneapolis, MN, USA)-coated 6well plate (Corning, 351146, Corning, NY, USA,).
The cells were cultured for 144 hours in medium with 5% fetal bovine serum after withdrawing
EGF from the medium to induce and maintain differentiation. At the time of EGF withdrawal, we
applied *Tbx1* siRNA (10nM, cat#4390771, s74767, Invitrogen; Thermo Fisher Scientific, Waltham,
MA, USA) and control siRNA-A (10nM, cat#4390843, Invitrogen; Thermo Fisher Scientific,
Waltham, MA, USA) together with siLentFect™ Lipid Reagent for RNAi, 0.5 ml (BIO-RAD, cat#
1703360, Hercules, CA, USA).
197

Quantitative reverse transcription polymerase chain reaction (qRT-PCR)
In accordance with our published procedure(62), total RNA was extracted using an RNeasy
Plus Mini Kit (Cat#74134, Qiagen, Germantown, USA). Complementary DNA (cDNA) was
synthesized from total RNA using SuperScript IV VILO Master Mix (Cat#11766050,
Invitrogen, Carlsbad, USA). qRT-PCR reactions were performed in triplicate on a
QuantStudio 6 Flex Real-Time PCR System (Cat#4485694, Applied Biosystems, Waltham,
USA) using the TaqMan Fast Advanced Master Mix (Cat#4444963, Applied Biosystems,
Waltham, USA). We used TaqMan® Gene Expression Assays (cat# 4331182, Thermo
Fisher, Waltham, MA, USA). The Taqman probes are listed in the Supplementary Material
(**Table S2**). Data were analyzed using the ΔΔCt method and normalized to the reference
gene *Pgk1*; we had an identical pattern of results with another reference gene *18S*.
209

Whole brain tissue clearing and index matching

We used the SHIELD and SmartBatch+ (LifeCanvas Sciences, Cambridge, MA) standard pipelines for preserving and electrophoretic clearing(81) to visualize tdTomato in the brains of *Pdgfra*Cre;ROSA-CAG-tdTomato mice. We performed transcardial perfusion of 1-month old mice with ice-cold 0.9% NaCl in dH_2O, followed by 4% paraformaldehyde (PFA) in phosphate-buffered saline (PBS). Brains were preserved using SHIELD as follows: perfused brains were 1) removed from the cranium and fixed in 4% PFA in PBS overnight at 4°C with gentle shaking; 2) incubated in SHIELD OFF solution (dH_2O + SHIELD Buffer [cat#SH-Bf] + SHIELD Epoxy Buffer [cat#SH-Ex]) solution at 4°C with gentle shaking for 3 days; 3) incubated in pre-warmed SHIELD ON Buffer (cat#SH-ON) at 37°C for 1 day with gentle shaking; 4) incubated in the delipidation buffer (cat#DB) for 3 days at 45°C with gentle shaking. Final clearing by electrophoresis was performed with the SmartBatch+ device, with the delipidation buffer inside the clearing cup containing the samples and conduction buffer (cat#CB) for 30 h. Cleared brains were index-matched using a solution of 50% EasyIndex (RI = 1.52, cat#EI-500-1.52) + 50% dH_2O with shaking at 37°C for 1 day, then 100% EasyIndex for one additional day until brains were transparent(81).

Light sheet microscopy

We used a Zeiss Lightsheet 7 Microscope housed in the UTHSCSA Optical Imaging Core to perform volumetric light sheet imaging of cleared *Pdgfra*Cre;ROSA-CAG-tdTomato brains using a pair of 5×/0.1 focusing illumination objectives and a Fluar 2.5×/0.12 detection objective at identical settings. Image tiles were stitched together using Zeiss Zen Blue software (ver 3.4). 3D volume analysis was performed with Imaris (Oxford Instruments, ver 10.02). Peak intensity of the tdTomato signal was segmented, and volumetric measurements (mm^3) and peak intensities of the total volume (arbitrary units) were calculated for each volume in each mouse.

Immunofluorescence

*Pdgfra*Cre;ROSA-*tdTomato* mice were sacrificed at 1 month of age. Mice were deeply anesthetized using 4–5% isoflurane and perfused transcardially with 0.9% saline followed by 4% PFA. Brains were extracted and post-fixed in 4% PFA overnight at 4°C, followed by cryoprotection in glycerol overnight at 4°C. Coronal 60-µm thick sections were cut with a freezing microtome. For immunofluorescence labeling, sections were washed in 0.1 M PBS, blocked with 5% normal donkey serum in PBS, and incubated overnight at room temperature, as we described previously(62, 72). We used a well-validated primary rabbit antibody against myelin basic protein (1:200, MBP, ab40390, Abcam) and Donkey anti-rabbit IgG-Alexa Fluor 488 (1:200, A21206, Invitrogen), together with a nuclear marker (DAPI,1:25,000, D3571, Invitrogen). MBP is a marker of mature oligodendrocytes(82, 83). We captured images of sections that were mounted and cover-slipped with Vectashield® anti-fade mounting medium (H-1000) using a Keyence microscope (BZ-X800) microscope with BZX Hardware Module and Zeiss LSM 710 confocal microscope at the UT Health Optical Imaging Core.

Double staining of tdTomato with markers of oligodendrocytes

We captured a series of coronal brain sections from *Pdgfra*Cre*;ROSA-tdTomato* mice using a Keyence BZ-X800 microscope with BZX Hardware Module and a Zeiss LSM 710 confocal microscope at the UT Health Optical Imaging Core. We used the DAPI signal to identify the anatomical boundaries of landmark structures in each image, cross-matched with equivalent levels in the Allen Brain Atlas (Allen Reference Atlas – Mouse Brain [brain atlas]. Available from *atlas.brain-map.org*). We then used a color scale to represent the level of tdTomato intensity within the delineated anatomical boundaries.

264
Behavioral Analysis
265

Male neonatal mice were tested for vocalization resulting from maternal separation at P8 and
P12, followed by additional behavioral tests at 1 and 2 months of age, corresponding to
peripubertal and post-pubertal, adolescent periods(84), respectively; early signs of puberty
begin around 1 month of age(85). Because body weight can alter behavior in mice(86) and
Pdgfra regulates progenitor differentiation into adipocytes(87, 88), we measured body weight at
the beginning of behavioral tests. Our behavioral battery included reciprocal social interaction,
novel object approach, spontaneous alternation in a T-maze, elevated plus maze, and
locomotor activity and thigmotaxis in an inescapable open field(52, 57-59, 64, 89-91).
274

The order of the behavioral assays was based on the stress level; behaviors that occur in a
home cage-like setting was given first (i.e., social interaction and novel object approach). T-
maze and elevated plus maze tests permit choices and are thus considered less stressful than
an inescapable open field. At least a one-day interval was provided between behavioral tests to
reduce the likelihood of carryover effects(61, 62, 92), except for the T-maze, in which three
different delays were given on three consecutive days.
281

Mice were assigned randomly to experimental groups and tested during the light phase.
Experimenters were blinded to genotypes. Social interaction, novel object approach, and
behaviors in the elevated plus maze were recorded using a Basler GigE camera, with video
images stored in Ethovision v18 software (Noldus Information Technology, Leesburg, VA). Video
images of social interaction and novel object approach were manually scored according to our
established criteria. Various parameters of the elevated plus maze were automatically analyzed
by Ethovision. The time point of each arm entry was recorded for spontaneous alternation, and
the data were subsequently tallied manually. Locomotor activity and thigmotaxis in an
inescapable open field were analyzed using Med Associates Activity Monitor 7 Software
(Fairfax, VT).

Ultrasonic vocalization Pups were tested for vocalization induced by maternal separation at
postnatal days 8 and 12. A cage containing the mother and a litter was transferred to the test
room 30 min before testing. Ultrasonic vocalization was recorded for 5 min by
UltraSoundGate (Avisoft, Germany) connected to a computer equipped with Avisoft-
RECORDER software (Avisoft, Germany) in a test chamber (18 cm long × 18 cm wide × 30 cm
high). The sampling rate and lower cutoff frequency were set at 250 kHz (format, 16-bit) and 10
kHz, respectively. A frequency window from 15–150 kHz was used for analysis. Call detection
was provided by an automatic threshold-based algorithm and a hold-time mechanism (hold time
= 10 ms). Sonograms were inspected, and mechanical noises were eliminated from the
analysis. We used the VocalMat software, which has the lowest false positive and false negative
rates(93), to determine call types. This software typically detects only the most salient
components of the harmonic call type and classifies this call type differently. As the harmonic
call type is often affected in genetic mouse models of neuropsychiatric disorders(58, 59, 86, 94),
we manually inspected all call types and re-classified such cases as harmonic calls.

Reciprocal Social Interaction Test A test mouse and an age-matched unfamiliar C57BL/6J
mouse as a stimulus subject were simultaneously placed in a cage (28.5 cm long × 17.5 cm
wide × 12.5 cm high). Mouse behavior was recorded during two 5-min sessions with a 30-min
interval between sessions. Video images were scored manually for affiliative and aggressive
social interaction by a rater blinded to genotype. The following reciprocal social behaviors were

scored by the duration of interaction (minimum of 1 s): aggressive (tail rattle, bite/kicks, sideway
offense, boxing/wrestling), affiliative, non-aggressive (mount, pursuit, olfactory investigation,
allogrooming, escape/leap), or passive (side-by-side, submissive). We analyzed the sum of time
engaged in affiliative, active, and passive social interactions, but passive behavior was rarely
seen in our experimental set-up (72, 86, 91, 94).

<u>Novel Object Approach</u> A test mouse was placed in a cage (28.5 cm long × 17.5 cm wide × 12.5
cm high) with an empty, non-secured 50-mL Falcon tube (3 cm diameter × 8.5 cm long) lying on
its side. Videos of mouse interactions with the tube were used to manually score approach
behavior by two independent analysts blinded to genotype with an inter-rater reliability of >99%.
Approach behavior was scored for olfactory or non-olfactory investigations with a minimum
duration of 1 s. Time spent near the tube was also analyzed.

<u>Spontaneous Alternation in a T-Maze</u> A test mouse was placed in the start site of the long arm
of a black plexiglass T-maze (21.5 cm long × 10.5 cm wide × 20.5 cm high) and allowed to enter
the maze and explore either the left or right short arms (31 cm long × 10.25 cm wide × 20.5 cm
high) of the cage. We imposed 0 s, 15 s, or 30 s interval delays to maze reentry for 10 trials, and
the three interval trials were given on three consecutive days. We manually scored the
percentages of alternation and latency to reach the alternated and non-alternated arms.
329
<u>Elevated Plus Maze Test</u> Mice were placed in the center stage (5 × 5 cm) of the maze apparatus
with four arms (30 × 5 cm) and permitted to explore two closed arms and two open arms
extending from the center platform of the maze for 5 min. The maze was positioned 53 cm
above the floor. The time spent in visits and the frequency of visits to the two open and two
closed arms were analyzed.

<u>Open Field</u> Mice were placed in an open-field cage (27.3 cm × 27.3 cm × 20.2 cm; Med
Associates, Fairfax, VT) and permitted to explore the cage for 30 min. We measured the
distance and velocity of motor activity and duration of time spent in the center (19.05 cm ×
19.05 cm central square) versus the margin of an open field, using the activity monitor software
(Med Associates, Fairfax, VT).

Electron Microscopy (EM)
We prepared tissue for EM to characterize myelination in the fimbria of $PdgfraCre;Tbx1^{+/flox}$ mice
as we described previously(62). After behavioral assays were completed at 1 and 2 months of
age, 4–5-month-old male $PdgfraCre;Tbx1^{+/flox}$ (N=3) and five control mice, including
$PdgfraCre;Tbx1^{+/+}$ (N=2), wild-type (WT);$Tbx1^{+/flox}$ (N=2), and WT;$Tbx1^{+/+}$ (N=1), were
anesthetized with 4.5% isoflurane in a chamber, and anesthesia was maintained by a nose cone
with 2.0% isoflurane. The animals were transcardially perfused with 120 mL of 0.9% saline
followed by fixation with 120 mL of 0.1 M buffer (pH 7.4; cat#11653 Electron Microscopy Science,
Hatfield, PA) with 2.5% sodium glutaraldehyde (cat#16310, Electron Microscopy Sciences,
Hatfield, PA) and 2.5% paraformaldehyde (cat#19202, Electron Microscopy Sciences, Hatfield,
PA). Brains were extracted and post-fixed in the same solution at 4°C for 7 weeks. The fimbria
from the two hemispheres were obtained separately using a vibratome and placed in 0.1 M
sodium cacodylate buffer overnight. Tissues were rinsed with 0.1 M sodium cacodylate buffer to
remove aldehydes and placed in 2% osmium tetroxide solution (OsO$_4$; cat#19150, Electron
Microscopy Sciences, Hatfield, PA) in 0.1 M sodium cacodylate buffer for 1 h. Tissues were
dehydrated in a series of ethanol solutions and embedded in molds containing Polybed resin
(Poly/Red® 812 Embedding media, cat#08791-500 Polysciences, Inc., Warrington, PA). We cut
100-nm sections and collected them on square 150-mesh copper grids (cat#7551C,
Polysciences, Inc., Warrington, PA) and stained them with uranyl acetate (7 g in 100 mL deionized
water)/Reynold's lead citrate (1.33 g lead nitrate,1.76 g sodium citrate in 30 mL triple-distilled

water, 8 mL 1 N NaOH). Uranyl acetate stains membranous structures and structures containing nucleic acid; lead citrate binds to RNA-containing structures and the hydroxyl groups of carbohydrates.

Each fimbria section was viewed in the EM grid squares. Images were screened at 1,000× magnification. For each section, we used all grid images that contained fimbria tissue with round axons without wrinkles, folds, or tears in the sections. One image was captured at the center of each grid field at 20,000× magnification. Images were analyzed by MyelTracer(95). Only axons whose myelin was fully within the image were analyzed quantitatively.

We obtained multiple images for each animal from a section of the right and left hemispheres. The positions of sections sliced from the fimbria varied between the two hemispheres and did not necessarily match the two sides; in one case, only one hemisphere was available because of the accidental loss of a tissue slice. The number of images varied across tissue slices, resulting in unequal image numbers. Many axons from each grid square and many grid squares from each hemisphere were analyzed for each animal. As the number of images and positions of each image varied between the two hemispheres and did not match between the two sides, the hemisphere could not be used as a replicate. Thus, we pooled data for each animal as a technical replicate; data points within and across images were technical replicates. As there was considerable variability in the data, we could not average the values for each animal. The averages of the total axon diameter per genotype were also not appropriate for this data set, as axon numbers differed between genotypes at specific axon diameters (≥300 nm to <400 nm; ≥1,200 nm to <1300 nm) (genotype × axon diameter range, $F(27,5880) = 6.336$, $P < 1.0 \times 10^{-4}$). Thus, linear mixed models were used, as reported previously(62). Data were assessed for the number of myelinated axons, the thickness of myelination, and the g-ratio along a 100 nm unit axon diameter. Data from different images and hemispheres were embedded in random models.

Statistical Analysis
We used SPSS (v29.0.2.0 (20), IBM Corporation) to perform all statistical analyses. Among-group and between-group comparisons of the data were performed using analysis of variance (ANOVA) and Student's two-tailed t-test (α = 0.05). We determined normality and homogeneity of variance using the Shapiro-Wilk test and Levene's homogeneity of variance test, respectively If either assumption was violated, we analyzed the data using a linear mixed model, Kruskal Wallis tests, or Mann–Whitney U tests. The Greenhouse–Geisser correction was applied if sphericity was violated and the estimated epsilon was less than 0.75. If multiple tests were applied to a data set, the significance level was adjusted using the Benjamini–Hochberg correction, with a false discovery rate of 5%. We used GraphPad Prism software (v9; GraphPad Software, San Diego, CA) to generate all graphs. All statistical analyses are presented in Table S3.

Data Availability
All data that support the findings and conclusions are provided within the article. All raw data and additional information are available upon request.

List of abbreviations
CNVs, Copy number variations
ASD, Autism Spectrum disorder
scRNA-seq, single cell RNA-seq
Pdgfra, Platelet-derived growth factor receptor alpha

409 MAG, Myelin associated glycoprotein
410 MOG, Myelin oligodendrocyte glycoprotein,
411 MBP, myelin basic protein
412
413 **Declarations**

414 **Ethics approval and consent to participate**
415 All protocols for animal handling and use were approved by the Institutional Animal Care and
416 Use Committee (IACUC) of the University of Texas Health Science Center at San Antonio
417 (UTHSCSA) in accordance with National Institutes of Health (NIH) guidelines.
418
419 **Consent for publication**
420 Not applicable
421
422 **Availability of data and materials**
423 All data and materials are available upon request.
424
425 **Competing interests**
426 The authors declare no competing interests.
427

Funding
T. Takano, GK, ME, TH, and NH were supported by the National Institute of Health
(R01MH099660; R01DC015776). T. Tanifuji was supported by SENSHIN Medical Research
Foundation and Uehara Memorial Foundation. QS and MAB were supported by the National
Institute of Health (R01GM063074). AMW was funded by F30MH134482, UT Health San
Antonio CTSA T32 (T32R004545), UT Health San Antonio Neuroscience T32 (T32NS082145),
STX-MSTP (NIH T32GM113896/T32GM145432), and SfN NSP Fellowship (R25NS089462).
The Zeiss Lightsheet 7 microscope was funded by the NIH S10 grant 1S10OD030383. The
content is solely the responsibility of the authors and does not necessarily represent the official
views of the National Institutes of Health.

Authors' contributions
Anne Marie Wells: Conducted lightsheet brain analysis, perfused mice and applied MyelTracer
to images for electron microscopy, genotyping, social interaction assessments, and manuscript
writing.

Takaki Tanifuji: Performed qRT-PCR analysis, immunofluorescent staining, confocal imaging
and analyses, electron microscopy image capturing, and figure preparation.

Takeshi Takano: Conducted statistical analyses and prepared all figures except for
immunofluorescent images.

Arumu Endo: Performed qRT-PCR analysis and figure preparation.

Gina Kang: Conducted all behavioral testing.

Marisa Esparza: Managed breeder maintenance, ensured quality control of data input,
performed statistical analyses, conducted mouse perfusion, and carried out genotyping.

Qian Shi: Responsible for electron microscopy preparation and manuscript writing.

Manzoor A. Bhat: Engaged in electron microscopy preparation and manuscript writing.

Noboru Hiroi: Designed all experiments, supervised all personnel, prepared electron microscopy
samples, captured electron microscopy images, performed immunofluorescent staining and
prepared figures, and wrote the manuscript.

Acknowledgments
456   The LifeCanvas device was purchased with generous donations by 10 scientists at UT Health
457   San Antonio. We thank Dr. Bernice Morrow for providing $Tbx1^{+/flox}$ breeders, Ms. Monica D.
458   Alarcon for technical assistance with EM analysis, Dr. Lacey B. Sell for her help with MyelTracer
459   software, and Dr. Takeshi Hiramoto for valuable comments.

RESULTS

Effects of *Tbx1* Knockdown on Markers of Oligodendrocyte Lineage *In Vitro*

We aimed to identify the earliest stage in the oligodendrocyte lineage affected by *Tbx1* deficiency. To achieve this aim, we developed an *in vitro* screening assay utilizing neonatal neural progenitor cells derived from the subventricular zone of C57BL/6J pups sacrificed on postnatal day 2, which are capable of generating both neuronal precursor cells and oligodendrocyte precursor cells(71). We assessed the expression of *Cspg4*, a marker for oligodendrocyte precursor cells, as well as *Mag, Mbp, Mog*, and *Plp1*, which are indicative of maturing and mature myelinating oligodendrocytes (67), at 144 hours after EGF was withdrawn (i.e., differentiation) and *Tbx1* siRNA was applied.

The application of *Tbx1* siRNA resulted in a significant reduction in the mRNA levels of *Tbx1, Cspg4*, and all markers associated with mature oligodendrocytes, with *Pgk1* (**Figure 1**) and *18S* (**Table S3-Figure 1**) as reference genes. These *in vitro* findings align with our *in vivo* data , which demonstrate that *Cspg4* (also referred to as *Ng2*) was diminished in the fimbria of *Tbx1* heterozygous mice(62).

Given that our culture includes both neuronal (58, 72) and oligodendrocyte (see **Figure 1**) lineages, as well as residual neural progenitor cells, the mechanism by which how *Tbx1* knockdown alters the expression of myelin marker genes remains unclear. It is possible that *Tbx1* knockdown in oligodendrocyte precursor cells leads to a cell-autonomous reduction in myelin markers; alternatively, reduced *Tbx1* levels in neonatal neural progenitor cells or their neuronal progeny may indirectly influence the expression of these oligodendrocyte markers.

Localization of Recombination in *Pdgfrα-Cre*;ROSA-tdTomato Mice

Our data indicate that *Tbx1* deficiency begins to impact the stage of oligodendrocyte precursor cells (see **Figure 1**). Consequently, we planned to initiate *Tbx1* heterozygosity in oligodendrocyte precursor cells using a conditional heterozygous mouse model. This mouse employed Cre-based initiation of *Tbx1* heterozygosity, guided by the promoter of *Pdgfrα*, a gene expressed in oligodendrocyte precursor cells of both embryonic and post-embryonic origins.

To identify the brain regions and cell types where *Tbx1* recombination is anticipated, we utilized PdgfrαCre;ROSA-tdTomato mice as the best available proxy. Alternative methodologies are not applicable to *Tbx1*. Recent single-cell RNA sequencing analyses reveal that *Tbx1* is present in proliferating cells, including neural progenitor cells, lung AT2 cells, spermatogonia, spermatids, and embryonic mesenchymal cells(65, 66). However, although *Tbx1* mRNA and protein products are expressed in actively proliferating neural progenitor cells, their levels significantly decrease in differentiated cells (72). As a result, the levels of *Tbx1* mRNA and protein products in our conditional *Tbx1* heterozygous mice are expected to reflect reductions due to cell-cycle changes, recombination-induced heterozygosity, or a combination of both factors.

Detecting TBX1 protein in the mouse brain presents challenges, with notable exceptions in endothelial cells(96) and neonatal neural progenitor cells(58, 72) partly due to epitope masking caused by fixation(58). Additionally, approaches to localize TBX1 using reporters under the endogenous *Tbx1* gene or its promoter have not yielded detectable signals in cells other than endothelial cells within the mouse brain(50, 96, 97), making it difficult to assess its colocalization with Pdgfrα protein.

In theory, Pdgfra-positive and MBP-positive cells could be isolated via fluorescence-activated cell sorting (FACS). However, while *Tbx1* is present in oligodendrocyte precursor cells and immature oligodendrocytes, its expression is notably low in mice(65, 66). Evaluating further reduced *Tbx1* levels is technically challenging due to the floor effect in heterozygous mice.

We first assessed the extent of *PdgfraCre*-mediated recombination throughout the brain, employing a tissue clearing technique (SmartBatch+) and utilizing lightsheet microscopy to visualize the 3D volume of tdTomato signals in the brains of PdgfrαCre;ROSA-tdTomato mice (**Figure 2A**). The highest levels of tdTomato expression were observed in a pair of arch-shaped structures located along the lateral and third ventricles. Additionally, clusters of tdTomato signals were detected on the ventral surface of the posterior brain, while lower levels of signals were observed throughout other regions.

To identify brain regions exhibiting tdTomato signals in PdgfrαCre;ROSA-tdTomato mice, we analyzed a series of coronal sections from 1-month-old PdgfrαCre;ROSA-tdTomato mice. High levels of tdTomato signals were present in the striatum, superficial layers of the neocortex (**Figure 2BC**), choroid plexus within the lateral ventricle (**Figure 2ED**) and the fimbria (**Figure 2F**). This expression pattern, driven by the *Pdgfra* promoter, aligns with previously reported distributions of *Pdgfra* mRNA and protein, as well as recombination activity associated with the *Pdgfra* promoter in the mouse brain (98-100). Moreover, the intense staining of the choroid plexus is consistent with the prominently tdTomato+ structures resembling the shapes of the lateral and third ventricles (see **Figure 2A**).

Embryonic Pdgfra-positive cells give rise to distinct pre-oligodendrocyte precursor cells by the perinatal period; however, they exhibit similar profiles during the postnatal and adult periods(67, 101). PdgfraCre is expected to induce recombination in oligodendrocyte precursor cells(74). Once tdTomato is expressed through recombination in the PdgfraCre;ROSA-tdTomato mouse line, it remains expressed in their progeny, including mature oligodendrocytes and myelinated fibers. Given that myelination of the fimbria is selectively affected in constitutive *Tbx1* heterozygous mice(62), we examined whether tdTomato colocalized with MBP, a marker of mature oligodendrocytes and myelinated fibers, in the fimbria(102). MBP-positive fibers were present throughout the fimbria (**Figure 3AB**). TdTomato signals were more prominent laterally than medially (**Figure 3C**) and were colocalized with MBP-positive fibers (**Figure 3D**) in the fimbria. These findings confirm that tdTomato, activated by PdgfraCre in PdgfraCre;ROSA-tdTomato mice, is present in myelinated fibers in the fimbria.

Intense tdTomato signals were evident in the choroid plexus located laterally to the lateral surface of the fimbria (**Figure 2D**). The cell types in the mouse choroid plexus that express *Pdgfra* include fibroblasts, mural cells, and endothelial cells during embryonic development, as well as fibroblasts and macrophages during the post-embryonic period in the mouse brain. Although gene profiles indicative of oligodendrocyte precursor-like cells are detectable during the embryonic period, their precise location within the choroid plexus remains unclear (66, 103-105). The robust tdTomato signal observed in the choroid plexus at 1 month of age likely originates from some or all of these cell types and any developmental time point. It is important to note that tdTomato signals detected at one month of age reflects recombination that has occurred any developmental time prior to this age.

Effects of *Tbx1* Heterozygosity in Oligodendrocyte Precursor Cells on Behavior

The growth rate may influence behavior; PdgfrαCre;*Tbx1*$^{+/flox}$, referred to as o*Tbx1*$^{+/-}$ mice, may exhibit developmental delays, with their phenotypes potentially reflecting a lag in development

compared to their wild-type littermates. However, neonatal o*Tbx1*$^{+/+}$ and o*Tbx1*$^{+/-}$ pups
demonstrated comparable increases in body weight (**Figure S1**).

To investigate the impact of *Tbx1* heterozygosity in oligodendrocyte precursor cells and their
progeny on behavior, mice were assessed tested for social communication during the neonatal
period and subjected to a series of tests evaluating social, cognitive, anxiety-related, and motor
behaviors at one and two months of age. This timing aligns with the peal of myelination, which
occurs between the fourth and fifth postnatal weeks in rodents(106, 107) and the onset of
puberty around 1 month of age and adolescence around 2 months of age(84). Constitutive *Tbx1*
heterozygous mice exhibit impairments in 1) neonatal social communication as early as P7, 2)
social behaviors, 3) working memory/cognitive flexibility in a T-maze and attentional set shifting,
and 4) affect-related behaviors in an inescapable open field, although not in an elevated plus
maze during adolescence(58-62) .

We first assessed differences among the three control groups: PdgfraCre;*Tbx1*$^{+/+,}$ WT;*Tbx1*$^{+/flox}$,
and WT;*Tbx1*$^{+/+}$. No significant differences were observed among the controls for any behavioral
measure, expect at three time points regarding total margin time in the open field (see **Table
S3-Figure 8B**). When analyzing all four genotypes together, a significant genotype effect was
found without an interaction effect with time (Genotype, p=0.0256; Genotype x Time, p=0.665).
However, post-hoc comparisons did not reveal significant differences between WT;*Tbx1*$^{+/flox}$ and
WT;*Tbx1*$^{+/+}$ mice only at 5 min and 15 min (see **Table S3, Figure 8B**). Given the sample size
for these two control groups was n=6 each, we lacked confidence in these statistical values.
Consequently, combined all the control genotypes into a single control group, designated as
o*Tbx1*$^{+/+}$, for comparison with o*Tbx1*$^{+/-}$ (66).

Constitutive *Tbx1*$^{+/-}$ mice display various neonatal, peri-adolescent, and postnatal behavioral
phenotypes(58, 59). Thus, we evaluated mice for neonatal ultrasonic vocalization on P8 and
P12–P13 and conducted additional tests with peripubertal (1 month) and adolescent (2 months)
mice, allowing 1–2 day intervals between tasks to minimize potential carryover effects from prior
testing(92).

We found no differences among o*Tbx1*$^{+/-}$ pups regarding the number, percentage, and duration
of various neonatal vocal call types on P8 and P12 (**Figure 4A–C**). At 1 and 2 months of age,
mice were sequentially assessed for social interaction, novel object approach, spontaneous
alternation in a T-maze, anxiety-related behavior in an elevated plus maze, and locomotor
activity and thigmotaxis in an inescapable open field. o*Tbx1*$^{+/-}$ and o*Tbx1*$^{+/+}$ mice were
indistinguishable in social interaction (**Figure 5A**) and novel object approach (**Figure 5B**). In a
T-maze, o*Tbx1*$^{+/-}$ mice at 1 month of age exhibited higher spontaneous alternation rates at the
longest delay (**Figure 6A**); otherwise, o*Tbx1*$^{+/+}$ and o*Tbx1*$^{+/-}$ mice were indistinguishable in rates
of spontaneous alternation at 2 months of age and in latencies to correct choices at both ages
(**Figure 6B–D**).

The two genotypes were indistinguishable in the percentage of time spent in open arms (F**igure
7A**) and visits to open arms (**Figure 7B**) of the elevated plus maze.

We assessed motor activity and anxiety-related behavior in a high-stress open-field task(58,
62) where there is no option to escape to a closed space. In this task, o*Tbx1*$^{+/-}$ and o*Tbx1*$^{+/+}$
mice showed no differences in distance traveled at both 1 month and 2 months of age (**Figure
8A,B**) or in time spent in the margin zone (**Figure 8C,D**).

While conditional *Tbx1* heterozygosity in the oligodendrocyte cell lineage resulted in higher
spontaneous alternation scores at the longest inter-trial delay at one month of age, this

phenotype contrasts with the impaired spontaneous alternation observed in constitutive *Tbx1*
heterozygous mice (58, 62). It is noteworthy that *oTbx1$^{+/+}$* mice exhibited consistent decline in
alternation rates to chance levels at a 30 s delay, whereas *oTbx1$^{+/-}$* mice did show this decline
not from 15 s to 30 s delay (see **Figure 6A**, **1 month**), indicating a delayed decay of working
memory or cognitive flexibility. Furthermore, this conditional *Tbx1* heterozygosity did not
replicate the altered neonatal vocalizations, peri-adolescent or postnatal social interaction
deficits, or heightened responses to novel, non-social objects or thigmotaxis observed in
constitutive *Tbx1* heterozygous mice(58).

Effects of *Tbx1* Heterozygosity in Oligodendrocyte Precursor Cells on Myelinated Axons in the
Fimbria

Following behavioral testing, we examined myelinated axons in the fimbria utilizing electron
microscopy. Constitutive *Tbx1* heterozygosity distinctly results in an increased proportion of
myelinated axons with diameters ranging from 200 to 600 nm diameter, while concurrently
reducing the proportion of axons measuring ≥700 nm and <1,200 nm. Additionally, axons within
the 700–1,500 nm range demonstrated a thicker myelin sheath, with no larger myelinated axons
identified in constitutive *Tbx1* heterozygous mice (62). We evaluated whether the ultrastructural
alterations observed in the fimbria of constitutive *Tbx1$^{+/-}$* mice were mirrored in o*Tbx1$^{+/-}$* mice.

Densely packed axons were identified in the fimbria (**Figure 9A**). To evaluate relative myelin
thickness, we analyzed the ratio of inner axon diameter to outer fiber diameter (i.e., the g-ratio)
(**Figure 9B**). Similar to the findings in constitutive *Tbx1* mice, the g-ratio plateaued at
approximately 0.8, which is the optimal ratio for signal conduction in the brain (88). However, no
significant differences in g-ratios were noted between o*Tbx1$^{+/-}$* and o*Tbx1$^{+/+}$* mice across the full
range of axon diameters.

We also quantified the number and proportion of axons across different diameters. The fimbria
of o*Tbx1$^{+/-}$* mice exhibited an increased number of axons measuring ≥ 300 nm and <800 nm,
alongside decreased number of axons ≥1,200 nm and <1,500 nm diameters in comparison to
o*Tbx1$^{+/+}$* mice (**Figure 9C**). Consequently, the curve representing relative axon proportions (y-
axis) versus axon diameter (x-axis) shifted to the left in o*Tbx1$^{+/-}$* mice relative to o*Tbx1$^{+/+}$* mice
(**Figure 9D**). Consistent with g-ratio findings, myelin thickness showed no significant differences
between o*Tbx1$^{+/-}$* and *oTbx1$^{+/+}$* mice across all axon diameters (**Figure S2**). Thus, conditional
Tbx1 heterozygosity initiated in oligodendrocyte precursor cells selectively shifted the relative
proportions of small to medium myelinated axons in the fimbria without impacting myelin
thickness.

In summary, conditional *Tbx1* heterozygosity within the oligodendrocyte lineage resulted in a
shift toward smaller-sized myelinated axons while leaving myelin thickness unchanged in the
fimbria.

DISCUSSION

The objective of this study was to delineate the cellular origin of the diverse behavioral and myelin phenotypes observed in constitutive *Tbx1* heterozygous mice(58-62). Given that social and cognitive phenotypes are recapitulated by initiating *Tbx1* heterozygosity in neonatal progenitor cells (72), and that certain neonatal neural progenitor cells in the mouse brain are responsible for generating neurons and oligodendrocytes during the neonatal period (70, 71, 108), it is possible that *Tbx1* deficiency in the neuronal or oligodendrocyte lineage contributes to the observed behavioral phenotypes. This study aimed to investigate the impact of *Tbx1* heterozygosity in the oligodendrocyte lineage on the behavioral and ultrastructural phenotypes.

To achieve this aim, we crossed *Pdgfra*Cre mice with congenic *Tbx1*$^{+/flox}$ mice to generate *Pdgfra*Cre;*Tbx1*$^{+/flox}$ (o*Tbx1*$^{+/-}$) mice. We evaluated a range of social, cognitive, anxiety-related, and motor behaviors, as well as the ultrastructural composition of myelin and axons in the fimbria. The o*Tbx1*$^{+/-}$ mice selectively replicated the altered proportion of axon sizes in the fimbria observed in constitutive *Tbx1* heterozygous mice. In contrast, o*Tbx1*$^{+/-}$ exhibited higher levels of spontaneous alternation following a 30-second delay at one month of age—indicative of a delayed decay of working memory and cognitive flexibility-- contrary to the impaired spontaneous alternation observed in constitutive *Tbx1* heterozygous mice. This observation supports the notion that the final phenotype arises from the cumulative effects of various contributory and opposing factors (54, 80).

Other phenotypes associated with constitutive *Tbx1* heterozygosity were absent in o*Tbx1*$^{+/-}$ mice, including neonatal social communication, responses to social or non-social stimuli, and anxiety-like behaviors in an elevated plus maze and thigmotaxis. These findings indicate that *Tbx1* in the oligodendrocyte lineage does not significantly influence these phenotypes. Combined with the observation that *Tbx1* heterozygosity initiated in neonatal (P1–P5) neural progenitor cells recapitulates some social and cognitive deficits observed in constitutive *Tbx1* heterozygous mice (72), the current negative data suggest that *Tbx1* in neonatal neural progenitor cells or their neuronal lineage may play a principal role in determining the final phenotype (72).

We cannot dismiss the possibility that conditional *Tbx1* heterozygosity was incomplete. The *Pdgfra* promoter is expected to induce nearly complete recombination in oligodendrocyte precursor cells, with labels persisting in their mature progeny, oligodendrocytes (109, 110); however, variability in recombination may occur in certain cells (111). The apparent absence of behavioral impairments may be attributed to incomplete recombination in all or some individual oligodendrocyte precursor cells. Nonetheless, o*Tbx1*$^{+/-}$ mice showed an axonal phenotype at a presumed gene dosage reduction that did not significantly affect behavior, clearly indicating that behavioral phenotypes exhibit lower sensitivity to *Tbx1* heterozygosity in oligodendrocyte precursor cells compared to the axonal phenotype.

Constitutive *Tbx1* heterozygous mice exhibit increased myelin thickness of medium-sized axons and a lack of large (≥1,700 nm) myelinated axons in the fimbria(62); however, these characteristics were absent in o*Tbx1*$^{+/-}$ mice. On the other hand, the fimbria of o*Tbx1*$^{+/-}$ mice exhibited a greater number of axons with diameters ranging from ≥300 nm to <800 nm and a reduced number of axons with diameters from ≥1,200 nm to <1,500 nm, compared to o*Tbx1*$^{+/+}$ mice. This pattern partially aligns with the increase in axons measuring ≥200 nm and <400 nm, as well as the decrease in axons measuring ≥700 nm and <1,700 nm in constitutive *Tbx1* heterozygous mice. The slight differences in the ranges of diameters of altered axons might be

due to manual measurements in our previous study (62) and an automated measurements by
MyelTracer in the current study.

This shift in axonal composition may result from oligodendrocyte lineage effects in a non-cell-
autonomous manner, or from a non-specific, cell-autonomous effect of the *Pdgfra* promoter in
neurons. While *Pdgfra*Cre primarily induces recombination in oligodendrocyte precursor cells, it
also initiates recombination in neurons, astrocytes, pericytes, ependyma, perivascular
mesenchymal cells, and other cell types(74-77, 99, 110, 112).

The recombination initiated by *Pdgfra*Cre could have influenced the behavioral phenotype in
spontaneous alternation through interactions with other cell types and their progeny. In the
mouse brain, *Tbx1* is undetectable in astrocytes or microglia, yet it is expressed in endothelial
cells and neural progenitor cells. Furthermore, *Tbx1* is detected in immature neurons as well as
both excitatory and inhibitory neurons at various developmental stages, from embryonic
development to adulthood(65, 66, 113, 114). Additional research is required to clarify the
impacts of *Tbx1* deficiency across different cell types on phenotypic outcomes.

The general absence of phenotypic abnormalities may be attributed to the high mortality rate
observed in o*Tbx1*$^{+/-}$ litters, potentially skewing our sample toward mice that were less affected
by, or more resilient to, the mutation's effects. In our breeder colony, only 31% and 24% of
male pups survived to postnatal day (P7) and 2 months of age, respectively. Nevertheless, two
lines of evidence do not support this explanation. First, a greater number of mice were alive
during the neonatal period compared to 1 and 2 months of age, yet neonatal vocalizations on
P8 and P12 were normal. Second, our nestinCreERT;*Tbx1*$^{+/flox}$ mice also exhibited a high
mortality rate but exhibited more significant behavioral deficits (72).
698
Large-scale human brain imaging studies have identified alterations in white matter among
carriers of CNV(115, 116), as well as in idiopathic cases of ASD and schizophrenia (117, 118).
Moreover, individuals with hemizygous deletions of 22q11.2 exhibit changes in white matter
within the fornix/fimbria(119). Our present findings do not support the hypothesis that *Tbx1*
deficiency in the oligodendrocyte lineage contributes to the structural or behavioral
abnormalities related to 22q11.2 hemizygosity and protein-truncating variants of TBX1. Further
research is necessary to identify the specific cell types responsible for the behavioral and
structural phenotypes linked to *Tbx1* heterozygosity, 22q11.2 hemizygosity, and other CNVs, in
order to establish a generalizable mechanistic basis for dimensions associated with mental
illness.

Figure legends

Figure 1. The relative quantification (RQ) (mean ± SEM) of expression levels at each stage of oligodendrogenesis, as determined by quantitative reverse transcription polymerase chain reaction (qRT-PCR) using *Pgk1* as a reference gene. Three cell clones were derived from three different C57BL/6J pups at postnatal days 2 or 3, with data obtained from three culture wells for each clone, except for one *Tbx1* siRNA-treated clone where *Tbx1* mRNA was undetectable and thus excluded from the analysis. A linear mixed model showed that Tbx1 siRNA reduced expression of all genes tested equally (Treatment, $F_{(1,89)}=14.665$, $p=0.0002386$; Gene, $F_{(5,89)}=0.8440$, $p=0.5222$; Treatment x Gene, $F_{(5,89)}=0.8440$, $p=0.5222$). Mann-Whitney tests confirmed that *Tbx1* siRNA reduced expression of each gene. Statistically significant differences between control siRNA and *Tbx1* siRNA groups are indicated by *, **, and *** for $p <0.05$, $p<0.01$, and $p<0.005$, respectively; they remained significant following Benjamini-Hochberg correction at a 5% false discovery rate (FDR).

Figure 2. A. A representative lightsheet image of the brain from a 1-month-old male PdgfrαCre;ROSA-tdTomato mouse. Following SmartBatch+ tissue delipification, tdTomato signals were visualized using a Zeiss LSM 7 Lightsheet microscope with a 2.5x objective (N = 3). **B-D**. tdTomato expression in 1-month-old PdgfrαCre;ROSA-tdTomato mice is depicted together with blue DAPI signals at the levels of the anterior striatum (**B**), middle striatum (**C**), posterior striatum/fimbria (**D**), anterior lateral ventricle (**E**), middle lateral ventricle (**F**), and fimbria (**H**). Abbreviations: ChPx, choroid plexus; LV, lateral ventricle; Str, striatum; Sept, septum; CC, corpus callosum; CTX, cortex; HGCL, hippocampal granule cell layer; fmb, fimbria; SVZ, subventricular zone. N = 3.

Figure 3. A Representative Case of Confocal Images of the Fimbria of a 1-Month-Old Male PdgfrαCre;ROSA-tdTomato Mouse. Sections were stained for DAPI (**A**, blue) and MBP (**B**, green). **C**. Red tdTomato signals originated from the PdgfrαCre;ROSA-tdTomato genotype. **D**. A composite image of the three colors is presented; white arrows indicate an area where fibers are positive for both MBP and tdTomato. N = 4. ChPx, choroid plexus; fmb, fimbria.

Figure 4. The number (**A**), proportion (**B**), and duration (**C**) (mean ± standard error of the mean [SEM]) of 12 call types emitted on postnatal days (P) P8 and P12. The two genotypes exhibited differences in certain interaction effects: **A**. Number: Genotype, $F_{(1,912)}=0.1966$, $p = 0.6576$; Genotype x Postnatal day, $F_{(1, 912)}=4.835$, $p = 0.0281$; Genotype x Call type, $F_{(11, 912)}=0.7077$, $p = 0.7319$; Genotype x Postnatal day x Call types, $F_{(11, 912)}=0.3686$, $p = 0.9679$. **B**. Proportion: Genotype, $F_{(1,912)}=0.0107$, $p = 0.9176$; Genotype x Postnatal day, $F_{(1, 912)}=0.0107$, $p = 0.9176$; Genotype x Call type, $F_{(11, 912)}=1.3657$, $p = 0.1837$; Genotype x Postnatal day x Call types, $F_{(11, 912)}=0.7966$, $p = 0.6437$. **C**. Duration: Genotype, $F_{(1,560)}=0.0435$, $p = 0.8349$; Genotype x Postnatal day, $F_{(1, 560)}=0.10519$, $p = 0.7458$; Genotype x Call type, $F_{(11, 560)}=3.361$, $p = 0.00016$; Genotype x Postnatal day x Call types, $F_{(11, 560)}=1.0389$, $p = 0.4101$. However, Mann-Whitney U tests did not reveal any significant genotype effects at any postnatal day for any call type after applying Benjamini-Hochberg corrections at 5% false discovery rate (FDR). **Inset** in **A**: The total number of ultrasonic vocalizations (mean ± SEM) emitted at P8 and P12. The two genotypes did not differ on the two postnatal days (Genotype, $F_{(1,76)}=0.1559$, $p = 0.694$; Genotype x Postnatal day, $F_{(1, 76)}=1.103$, $p = 0.2969$). **Inset** in **C**: Duration: Genotype, $F_{(1,32)}=0.0002$, $p = 0.9878$; Genotype x Postnatal day, $F_{(1, 32)}=0.3617$, $p = 0.5518$). Ha, harmonic; Sd, step-down; Sh, short; Df, down frequency modulation; Uf, up frequency modulation; Ts, two steps; Fl, flat; Ch, chevron;

Co, complex; Su, step-up; Ms, multiple steps; Rc, reverse chevron. $oTbx1^{+/+}$ mice, N = 27;
$oTbx1^{+/-}$ mice, N = 11 at P8; $oTbx1^{+/+}$ mice, N = 30; $oTbx1^{+/-}$ mice, N = 12 at P12.

Figure 5. Interaction with Social and Non-Social Stimuli. **A**) Social Interaction. The $oTbx1^{+/-}$
(PdgfraCre;Tbx1$^{+/flox}$) mice and $oTbx1+/+$ mice did not differ in the amount of time (mean ±
SEM) spent in affiliative social interaction (1 month: genotype, F(1, 72) = 0.1118, p = 0.739;
genotype × session, F(1, 72) = 0.262, p = 0.6102; 2 months: genotype, F(1, 58) = 0.00547, p =
0.9413; genotype x session, F(1, 58) = 1.2175, p = 0.2744). The three genotypes of $oTbx1+/+$
(1 month, N = 8, PdgfraCre;$Tbx1^{+/+}$, N = 10, Wild-type;$Tbx1^{+/+}$, N = 9, Wild-type;$Tbx1^{+/flox}$; 2
months, N = 8, PdgfraCre;$Tbx1^{+/+}$, N = 6, Wild-type;$Tbx1^{+/+}$, N = 7, Wild-type;$Tbx1^{+/flox}$) did not
show significant differences (1 month: genotype, F(2, 48) = 1.686, p = 0.1960; genotype ×
session, F(2, 48) = 0.843, p = 0.4366; 2 months: genotype, F(2, 36) = 0.0392, p = 0.9616;
genotype x session, F(2, 36) = 0.2382, p = 0.7893), and were thus combined as $oTbx1+/+$.
$oTbx1^{+/+}$ mice , N = 27; $oTbx1^{+/-}$ mice, N = 11 at 1 month; $oTbx1^{+/+}$ mice, N = 21; $oTbx1^{+/-}$ mice,
N = 10 at 2 months. **B**) Approach to a novel, non-social object. The two genotypes did not differ
in the amount of time spent in affiliative social interaction (1 month: genotype, F(1, 70) =
2.2507E-05, p = 0.9962; genotype × session, F(1, 70) = 1.0305, p = 0.31354; 2 months:
genotype, F(1, 58) = 0.11634, p = 0.7342; genotype × session, F(1, 58) = 0.0821, p = 0.77545).
The three genotypes of $oTbx1^{+/+}$ (1 month: N = 8, PdgfraCre;$Tbx1^{+/+}$, N = 9, Wild-type;$Tbx1^{+/+}$, N
= 9, Wild-type;$Tbx1^{+/flox}$; 2 months: N = 8, PdgfraCre;$Tbx1^{+/+}$, N = 6, Wild-type;$Tbx1^{+/+}$, N = 7,
Wild-type;$Tbx1^{+/flox}$) did not differ (1 month: genotype, F(2, 46) = 0.1772, p = 0.8382; genotype ×
session, F(2, 46) = 0.9318, p = 0.4012; 2 months: genotype, F(2, 36) = 0.0580, p = 0.9437;
genotype × session, F(2, 36) = 0.6581, p = 0.5239) and were therefore combined as $oTbx1^{+/+}$.
$oTbx1^{+/+}$ mice, N = 26; $oTbx1^{+/-}$ mice, N = 11 at 1 month; $oTbx1^{+/+}$ mice, N = 21; $oTbx1^{+/-}$ mice,
N = 10 at 2 months.

Figure 6. Spontaneous alternation in a T-maze. Compared to male $oTbx1^{+/+}$ (N = 20) mice,
male $oTbx1^{+/-}$ (N = 10) mice performed at higher percentages of correct spontaneous
alternations (mean ± SEM) at the longest delay (30 s) in a T-maze at 1 month of age, but not at
2 months. Specifically, at 1 month: % alternation, genotype, F(1, 96) = 4.66, p = 0.0333;
genotype × delays, F(2,96) = 3.1509, p = 0.0473; at 2 months: genotype, F(1, 81) = 2.047, p =
0.1564; genotype × delays, F(2, 81) = 0.2439, p = 0.7841. The three genotypes of $oTbx1^{+/+}$ (1
month: N = 8, PdgfraCre;$Tbx1^{+/+}$, N = 7, Wild-type;$Tbx1^{+/+}$, N = 9, Wild-type;$Tbx1^{+/flox}$; 2 months:
N = 8, PdgfraCre;$Tbx1^{+/+}$, N = 6, Wild-type;$Tbx1^{+/+}$, N = 6, Wild-type;$Tbx1^{+/flox}$) did not differ (1
month: genotype, F(2, 63) = 0.1046, p = 0.9008; genotype × delays, F(4, 63) = 0.2648, p =
0.8995; 2 months: genotype, F(2, 51) = 0.1197, p = 0.8874; genotype × delays, F(4, 51) =
0.6401, p = 0.6363) and consequently were combined as $oTbx1^{+/+}$. $oTbx1^{+/-}$ mice exhibited
higher rates of correct alternations at the 30-second delay compared to $oTbx1+/+$ mice, as
determined by the Mann-Whitney test (p = 0.0027). **B**) The two groups did not exhibit significant
differences in the latency to correct choices in the T-maze at 1 month (genotype, F(1, 96) =
2.181, p = 0.1430; genotype × delays, F(2, 96) = 2.058, p = 0.1332) or at 2 months (genotype,
F(1, 81) = 1.1135, p = 0.2945; genotype × delays, F(2, 81) = 2.5825, p = 0.0818). The three
genotypes of $oTbx1^{+/+}$ (1 month, N=8), PdgfraCre;$Tbx1^{+/+}$ (N=7), Wild-type;$Tbx1^{+/+}$ (N=9), and
Wild-type;$Tbx1^{+/flox}$ (2 months, N=8), PdgfraCre;$Tbx1^{+/+}$ (N=6), Wild-type;$Tbx1^{+/+}$ (N=6), and
Wild-type;$Tbx1^{+/flox}$, demonstrated a significant interaction effect at 1 month of age (genotype,
F(2, 63) = 1.5787, p = 0.2143; genotype × session, F(4, 63) = 4.5625, p = 0.00267), but no
significant interaction at 2 months of age (genotype, F(2, 51) = 0.1631, p = 0.8500; genotype ×
session, F(4, 51) = 1.0611, p = 0.3853). However, Mann-Whitney U tests revealed that none of

the three genotypes significantly differed from one another at 1 month of age after applying the
Benjamini-Hochberg correction (p>0.05 for all pairs). The three genotypes were combined as
$oTbx1^{+/+}$. The sample sizes were as follows: $oTbx1^{+/+}$ mice, N = 24; $oTbx1^{+/-}$ mice, N = 10 at 1
month; $oTbx1^{+/+}$ mice, N = 20; $oTbx1^{+/-}$ mice, N = 9 at 2 months.

Figure 7. Anxiety-related behavior in an elevated T-maze.The $oTbx1^{+/-}$ and $oTbx1^{+/+}$ mice did
not differ significantly in the relative amount of time spent (mean +SEM) in the open arms of an
elevated plus maze (**A**,1 month,U = 100, p = 0.3440; 2 months, U = 98, p = 0.9483) or the
frequency of visits to the open arms of the elevated plus maze (**B**, 1 month,t(32) = 1.7412, p =
0.0912; 2 months, t(28) = 0.5925, p = 0.5583). The three genotypes of $oTbx1$+/+ (1 month, N=7,
PdgfraCre;$Tbx1^{+/+}$, N=7, Wild-type;$Tbx1^{+/+}$, N=6, Wild-type;$Tbx1$+/flox; 2 months, N=8,
PdgfraCre;$Tbx1^{+/+}$, N=6, Wild-type;$Tbx1^{+/+}$, N=6, Wild-type;$Tbx1^{+/flox}$) did not differ in the
percentage of time spent in open arms (1 month,genotype, F(2, 20) = 0.0745, p = 0.9285; 2
months,genotype, F(2, 17) = 2.126, p = 0.1500) or the percentage of visits to open arms (1
month,genotype, F(2, 20) = 0.1837, p = 0.8336; 2 months,genotype, F(2, 17) = 1.4821, p =
0.2551), and thus were combined as $oTbx1^{+/+}$. The sample sizes were: $oTbx1^{+/+}$ mice, N = 23;
$oTbx1^{+/-}$ mice, N = 11 at 1 month; $oTbx1^{+/+}$ mice, N = 20; $oTbx1^{+/-}$ mice, N = 10 at 2 months.

Figure 8. Motor Activity and Thigmotaxis. $oTbx1^{+/+}$ and $oTbx1^{+/-}$ mice exhibited indistinguishable
distances traveled at both 1 month and 2 months of age (**A**, 1 month, genotype, F(1, 198) =
0.6314, p = 0.4278; genotype × time, F(5, 198) = 0.9551, p = 0.4466; **B**, 2 months, genotype,
F(1, 28) = 0.0242, p = 0.87739; genotype × time, F(3.66, 102.485) = 0.5389, p = 0.69188). The
Greenhouse–Geisser correction was applied due to a violation of sphericity, with the estimated
epsilon being less than 0.75.The three genotypes of $oTbx1^{+/+}$ (1 month, N=8, PdgfraCre;$Tbx1^{+/+}$,
N=7, Wild-type;$Tbx1^{+/+}$, N=9, Wild-type;$Tbx1^{+/flox}$; 2 months, N=8, PdgfraCre;$Tbx1^{+/+}$, N=6, Wild-
type;$Tbx1^{+/+}$, N=6, Wild-type;$Tbx1^{+/flox}$) demonstrated differences at both ages (genotype, F(2,
21) = 5.475, p = 0.01220; genotype × time, F(7.10, 74.58) = 1.118, p = 0.3611; at 2
months,genotype, F(2, 17) = 9.533, p = 0.00167; genotype × time, F(10, 85) = 0.3955, p =
0.9453). Comparisons of all pairs using Student's t-tests indicated that no pair reached
significance after the Benjamini-Hochberg correction (p>0.05 for all pairs), resulting in their
combination as $oTbx1^{+/+}$.

$oTbx1^{+/+}$ and $oTbx1^{+/-}$ mice spent equal amounts of time in the margin zone (i.e., thigmotaxis) of
the inescapable open field at both 1 and 2 months of age (**C,** 1 month, genotype, F(1, 198) =
0.2009, p = 0.6545; genotype × time, F(5,198) = 2.1517, p = 0.0609; **D**, 2 months, genotype,
F(1, 168) = 0.08912, p = 0.7657; genotype × time, F(5, 168) = 0.9880, p = 0.4267). The three
genotypes of $oTbx1^{+/+}$ did not differ at 1 month of age (genotype, F(2, 21) = 0.3686, p = 0.6961;
genotype × time, F(5.485, 57.59) = 0.9884, p = 0.4375), but differences were observed at 2
months of age (genotype, F(2, 17) = 4.880, p = 0.0211; genotype × time, F(5.049, 42.917) =
0.7180, p = 0.6147). Comparisons of all pairs using Student's t-tests revealed that 2-month-old
Wild-type;$Tbx1^{+/flox}$ and Wild-type;$Tbx1^{+/+}$ differed at the first four time points, even after the
Benjamini-Hochberg correction (p<0.05). However, these groups had relatively small sample
sizes (N=6 for both), and PdgfraCre;$Tbx1^{+/+}$and Wild-type;$Tbx1^{+/flox}$ or PdgfraCre;$Tbx1^{+/+}$ and
Wild-type;$Tbx1^{+/flox}$ did not differ at any time point (p >0.05), thus they were combined as
$oTbx1^{+/+}$. Sample sizes included $oTbx1^{+/+}$ mice, N = 24; $oTbx1^{+/-}$ mice, N = 11 at 1 month;
$oTbx1^{+/+}$ mice, N = 20; $oTbx1^{+/-}$ mice, N = 10 at 2 months.

Figure 9. Electron Microscopy Analysis of Myelinated Axons. **A**. Representative electron
microscopy images of axons in the fimbria at 20,000× magnification. Genotypes: +/+, $oTbx1^{+/+}$;
+/-, $oTbx1^{+/-}$. Scale bar = 1,600 nm. **B**. G-ratios plotted against axon diameters for 4,423 and

2,317 axons from 5 *oTbx1$^{+/+}$* mice and 3 *oTbx1$^{+/-}$* mice, respectively. All axons with closed
myelin sheaths were analyzed using MyelTracer. The reciprocal function best fits the data
distributions. G-ratios increased as a function of axon diameter (*oTbx1+/+*, R = 0.6268, p <
0.001; *oTbx1+/−*, R = 0.6491, p < 0.001). There was no significant difference in the axon
diameter range for G-ratios between oTbx1+/+ and oTbx1+/− mice (genotype, p = 0.8131;
genotype × axon diameter, p = 0.221). **C**. Average number (± SEM) of axons per image field
versus axon diameter. Significant differences were observed between *oTbx1$^{+/+}$* and *oTbx1$^{+/-}$*
mice (χ^2 (27) = 109.7085, p = 6.22 × 10^{-12}), primarily due to a higher number of axons with
diameters ≥300 nm and <800 nm and a lower number of axons with diameters ≥1,200 nm and
<1,500 nm in *oTbx1$^{+/-}$* compared to *oTbx1$^{+/+}$* mice, as determined by Mann-Whitney U tests
applied to this diameter range (*, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001). The shift
in axon sizes toward smaller to medium diameters in *oTbx1$^{+/-}$* compared to *oTbx1$^{+/+}$* mice
resulted in a smaller average axon size for *oTbx1$^{+/-}$* mice (Mann-Whitney non-parametric tests,
p = 7.48 × 10^{-19}). **D**. Differences in the percentage of myelinated axons for *oTbx1$^{+/-}$* versus
oTbx1+/+ mice across axon diameters (Kolmogorov-Smirnov, p = 3.59 × 10^{-16}).

Table S1. Primers used for genotyping.

Mouse line	Forward (5' → 3')	Reverse (5' → 3')	Purpose
Pdgfra-Cre	oIMR1084 (GCGGTCTGGCAGTAAAAACTATC)	oIMR1085 (GTGAAACAGCATTGCTGTCACTT)	Generic cre - transgene
Pdgfra-Cre	oIMR7338 (CTAGGCCACAGAATTGAAAGATCT)	oIMR7339 (GTAGGTGGAAATTCTAGCATCATCC)	Generic cre – internal positive control
ROSA-CAG-tdTomato	oIMR9020 (AAGGGAGCTGCAGTGGAGTA)	oIMR9021 (CCGAAAATCTGTGGGAAGTC)	tdTomato wild-type control
ROSA-CAG-tdTomato	oIMR9103 (GGCATTAAAGCAGCGTATCC)	oIMR9105 (CTGTTCCTGTACGGCATGG)	tdTomato mutant
Tbx1$^{+/flox}$	2g 1F (TCTTCTTGGGGCTGTAGACT)	Tbx1 1R (TGACTGTGCTGAAGTGCATC)	LoxP site
Tbx1$^{+/−}$	KO1F (TTGGTGACGATCATCTCGGT)	KO1R (ATGATCTCCGCCGTGTCTAG) Mut2R (AGGTCCCTCGAAGAGGTTCA)	*Tbx1$^{+/+}$* *Tbx1$^{+/−}$*

866 **Table S2. Primers for qRT-PCR**

Gene	Assay ID
Tbx1	Mm00448949_m1
Cspg4	Mm00507257_m1
Mag	Mm00487538_m1
Mbp	Mm01266402_m1
Mog	Mm01279062_m1
Plp1	Mm01297210_m1

References

869
1.   Hiroi N, Yamauchi T. Modeling and Predicting Developmental Trajectories of
Neuropsychiatric Dimensions Associated With Copy Number Variations. Int J
Neuropsychopharmacol. 2019;22(8):488–500.
2.   Michelini G, Carlisi CO, Eaton NR, Elison JT, Haltigan JD, Kotov R, et al. Where do
neurodevelopmental conditions fit in transdiagnostic psychiatric frameworks? Incorporating a new
neurodevelopmental spectrum. World Psychiatry. 2024;23(3):333–57.
3.   Ozonoff S, Iosif AM, Baguio F, Cook IC, Hill MM, Hutman T, et al. A prospective study of the
emergence of early behavioral signs of autism. J Am Acad Child Adolesc Psychiatry.
2010;49(3):256–66.
4.   Ozonoff S, Young GS, Belding A, Hill M, Hill A, Hutman T, et al. The broader autism
phenotype in infancy: when does it emerge? J Am Acad Child Adolesc Psychiatry. 2014;53(4):398–
407.
5.   Esposito G, Hiroi N, Scattoni ML. Cry, baby, cry: Expression of Distress as a Biomarker and
Modulator in Autism Spectrum Disorder. Int J Neuropsychopharmacol. 2017;20(6):498–503.
6.   Sung S, Fenoglio A, Wolff JJ, Schultz RT, Botteron KN, Dager SR, et al. Examining the factor
structure and discriminative utility of the Infant Behavior Questionnaire-Revised in infant siblings of
autistic children. Child Dev. 2022;93(5):1398–413.
7.   Northrup JB, Leezenbaum NB, Campbell SB. Observed Social Emotional Behavior at 22
Months Predicts a Later ASD Diagnosis in High-Risk Siblings. J Autism Dev Disord. 2021;51(9):3187–
98.
8.   Miller M, Sun S, Iosif AM, Young GS, Belding A, Tubbs A, et al. Repetitive behavior with
objects in infants developing autism predicts diagnosis and later social behavior as early as 9
months. J Abnorm Psychol. 2021;130(6):665–75.
9.   Di Giorgio E, Rosa-Salva O, Frasnelli E, Calcagni A, Lunghi M, Scattoni ML, et al. Abnormal
visual attention to simple social stimuli in 4-month-old infants at high risk for Autism. Sci Rep.
2021;11(1):15785.
10.   Miller M, Iosif AM, Hill M, Young GS, Schwichtenberg AJ, Ozonoff S. Response to Name in
Infants Developing Autism Spectrum Disorder: A Prospective Study. J Pediatr. 2017;183:141–6 e1.
11.   Caravella KE, Roberts JE. Adaptive Skill Trajectories in Infants with Fragile X Syndrome
Contrasted to Typical Controls and Infants at High Risk for Autism. Res Autism Spectr Disord.
2017;40:1–12.
12.   Gur RC, Calkins ME, Satterthwaite TD, Ruparel K, Bilker WB, Moore TM, et al. Neurocognitive
growth charting in psychosis spectrum youths. JAMA Psychiatry. 2014;71(4):366–74.
13.   Seidman LJ, Shapiro DI, Stone WS, Woodberry KA, Ronzio A, Cornblatt BA, et al. Association
of Neurocognition With Transition to Psychosis: Baseline Functioning in the Second Phase of the
North American Prodrome Longitudinal Study. JAMA Psychiatry. 2016;73(12):1239–48.
14.   Meier MH, Caspi A, Reichenberg A, Keefe RS, Fisher HL, Harrington H, et al.
Neuropsychological decline in schizophrenia from the premorbid to the postonset period: evidence
from a population-representative longitudinal study. Am J Psychiatry. 2014;171(1):91–101.
15.   Bora E, Lin A, Wood SJ, Yung AR, McGorry PD, Pantelis C. Cognitive deficits in youth with
familial and clinical high risk to psychosis: a systematic review and meta-analysis. Acta Psychiatr
Scand. 2014;130(1):1–15.
16.   Sheffield JM, Karcher NR, Barch DM. Cognitive Deficits in Psychotic Disorders: A Lifespan
Perspective. Neuropsychol Rev. 2018;28(4):509–33.
17.   Tschentscher N, Woll CFJ, Tafelmaier JC, Kriesche D, Bucher JC, Engel RR, et al.
Neurocognitive Deficits in First-Episode and Chronic Psychotic Disorders: A Systematic Review
from 2009 to 2022. Brain Sci. 2023;13(2).

18.   Johansson M, Hjarthag F, Helldin L. Cognitive markers related to long-term remission status
in Schizophrenia Spectrum Disorders. Psychiatry Res. 2020;289:113035.

19.   Stefansson H, Meyer-Lindenberg A, Steinberg S, Magnusdottir B, Morgen K, Arnarsdottir S,
et al. CNVs conferring risk of autism or schizophrenia affect cognition in controls. Nature.
2014;505(7483):361–6.

20.   Kendall KM, Rees E, Escott-Price V, Einon M, Thomas R, Hewitt J, et al. Cognitive
Performance Among Carriers of Pathogenic Copy Number Variants: Analysis of 152,000 UK Biobank
Subjects. Biol Psychiatry. 2017;82(2):103–10.

21.   Malhotra D, Sebat J. CNVs: harbingers of a rare variant revolution in psychiatric genetics.
Cell. 2012;148(6):1223–41.

22.   Marshall CR, Howrigan DP, Merico D, Thiruvahindrapuram B, Wu W, Greer DS, et al.
Contribution of copy number variants to schizophrenia from a genome-wide study of 41,321
subjects. Nat Genet. 2017;49(1):27–35.

23.   Goh S, Thiyagarajan L, Dudding-Byth T, Mark P, Kirk EP. A systematic review and pooled
analysis of penetrance estimates of copy number variants associated with neurodevelopment.
Genet Med. 2024:101227.

24.   Gur RE, Yi JJ, Donald-McGinn DM, Tang SX, Calkins ME, Whinna D, et al. Neurocognitive
development in 22q11.2 deletion syndrome: comparison with youth having developmental delay
and medical comorbidities. Mol Psychiatry. 2014;19(11):1205–11.

25.   Morrison S, Chawner S, van Amelsvoort T, Swillen A, Vingerhoets C, Vergaelen E, et al.
Cognitive deficits in childhood, adolescence and adulthood in 22q11.2 deletion syndrome and
association with psychopathology. Transl Psychiatry. 2020;10(1):53.

26.   Chawner S, Owen MJ, Holmans P, Raymond FL, Skuse D, Hall J, et al. Genotype-phenotype
associations in children with copy number variants associated with high neuropsychiatric risk in the
UK (IMAGINE-ID): a case-control cohort study. Lancet Psychiatry. 2019;6(6):493–505.

27.   Zinkstok J, Boot E, Bassett AS, Hiroi N, Butcher NJ, Vingerhoets C, et al. The 22q11.2
deletion syndrome from a neurobiological perspective. Lancet Psychiatry. 2019;6(11):951–60.

28.   Schneider M, Debbane M, Bassett AS, Chow EW, Fung WL, van den Bree M, et al. Psychiatric
disorders from childhood to adulthood in 22q11.2 deletion syndrome: results from the International
Consortium on Brain and Behavior in 22q11.2 Deletion Syndrome. Am J Psychiatry.
2014;171(6):627–39.

29.   Ensenauer RE, Adeyinka A, Flynn HC, Michels VV, Lindor NM, Dawson DB, et al.
Microduplication 22q11.2, an emerging syndrome: clinical, cytogenetic, and molecular analysis of
thirteen patients. American Journal of Human Genetics. 2003;73(5):1027–40.

30.   Hassed SJ, Hopcus-Niccum D, Zhang L, Li S, Mulvihill JJ. A new genomic duplication
syndrome complementary to the velocardiofacial (22q11 deletion) syndrome. Clinical Genetics.
2004;65(5):400–4.

31.   Portnoi MF, Lebas F, Gruchy N, Ardalan A, Biran-Mucignat V, Malan V, et al. 22q11.2
duplication syndrome: two new familial cases with some overlapping features with
DiGeorge/velocardiofacial syndromes. American Journal of Medical Genetics A. 2005;137(1):47–51.

32.   Yobb TM, Somerville MJ, Willatt L, Firth HV, Harrison K, MacKenzie J, et al. Microduplication
and triplication of 22q11.2: a highly variable syndrome. American Journal of Human Genetics.
2005;76(5):865–76.

33.   de La Rochebrochard C, Joly-Helas G, Goldenberg A, Durand I, Laquerriere A, Ickowicz V, et
al. The intrafamilial variability of the 22q11.2 microduplication encompasses a spectrum from
minor cognitive deficits to severe congenital anomalies. Am J Med Genet A. 2006;140(14):1608–13.

34.   Alberti A, Romano C, Falco M, Cali F, Schinocca P, Galesi O, et al. 1.5 Mb de novo 22q11.21
microduplication in a patient with cognitive deficits and dysmorphic facial features. Clin Genet.
2007;71(2):177–82.

35. Engels H, Brockschmidt A, Hoischen A, Landwehr C, Bosse K, Walldorf C, et al. DNA
microarray analysis identifies candidate regions and genes in unexplained mental retardation.
Neurology. 2007;68(10):743–50.
36. Mukaddes NM, Herguner S. Autistic disorder and 22q11.2 duplication. World J Biol
Psychiatry. 2007;8(2):127–30.
37. Torres-Juan L, Rosell J, Morla M, Vidal-Pou C, Garcia-Algas F, de la Fuente MA, et al.
Mutations in TBX1 genocopy the 22q11.2 deletion and duplication syndromes: a new susceptibility
factor for mental retardation. Eur J Hum Genet. 2007;15(6):658–63.
38. Descartes M, Franklin J, de Stahl TD, Piotrowski A, Bruder CE, Dumanski JP, et al. Distal
22q11.2 microduplication encompassing the BCR gene. Am J Med Genet A. 2008;146A(23):3075–
81.
39. Ramelli GP, Silacci C, Ferrarini A, Cattaneo C, Visconti P, Pescia G. Microduplication
22q11.2 in a child with autism spectrum disorder: clinical and genetic study. Dev Med Child Neurol.
2008;50(12):953–5.
40. Wentzel C, Fernstrom M, Ohrner Y, Anneren G, Thuresson AC. Clinical variability of the
22q11.2 duplication syndrome. Eur J Med Genet. 2008;51(6):501–10.
41. Yu S, Cox K, Friend K, Smith S, Buchheim R, Bain S, et al. Familial 22q11.2 duplication: a
three-generation family with a 3-Mb duplication and a familial 1.5-Mb duplication. Clin Genet.
2008;73(2):160–4.
42. Lo-Castro A, Galasso C, Cerminara C, El-Malhany N, Benedetti S, Nardone AM, et al.
Association of syndromic mental retardation and autism with 22q11.2 duplication.
Neuropediatrics. 2009;40(3):137–40.
43. Portnoi MF. Microduplication 22q11.2: a new chromosomal syndrome. Eur J Med Genet.
2009;52(2-3):88–93.
44. Soysal Y, Vermeesch J, Davani NA, Sensoy N, Hekimler K, Imirzalioglu N. Molecular
characterization of microduplication 22q11.2 in a girl with hypernasal speech. Genet Mol Res.
2011;10(3):2148–54.
45. Wenger TL, Miller JS, DePolo LM, de Marchena AB, Clements CC, Emanuel BS, et al.
22q11.2 duplication syndrome: elevated rate of autism spectrum disorder and need for medical
screening. Mol Autism. 2016;7:27.
46. Singh T, Poterba T, Curtis D, Akil H, Al Eissa M, Barchas JD, et al. Rare coding variants in ten
genes confer substantial risk for schizophrenia. Nature. 2022;604(7906):509–16.
47. Alhazmi S, Alzahrani M, Farsi R, Alharbi M, Algothmi K, Alburae N, et al. Multiple Recurrent
Copy Number Variations (CNVs) in Chromosome 22 Including 22q11.2 Associated with Autism
Spectrum Disorder. Pharmgenomics Pers Med. 2022;15:705–20.
48. Alghamdi M, Al Khalifah R, Al Homyani DK, Alkhamis WH, Arold ST, Ekhzaimy A, et al. A
Novel TBX1 Variant Causing Hypoparathyroidism and Deafness. J Endocr Soc. 2020;4(2):bvz028.
49. Ogata T, Niihori T, Tanaka N, Kawai M, Nagashima T, Funayama R, et al. TBX1 mutation
identified by exome sequencing in a Japanese family with 22q11.2 deletion syndrome-like
craniofacial features and hypocalcemia. PLoS One. 2014;9(3):e91598.
50. Paylor R, Glaser B, Mupo A, Ataliotis P, Spencer C, Sobotka A, et al. Tbx1 haploinsufficiency
is linked to behavioral disorders in mice and humans: implications for 22q11 deletion syndrome.
Proc Natl Acad Sci U S A. 2006;103(20):7729–34.
51. Gong W, Gottlieb S, Collins J, Blescia A, Dietz H, Goldmuntz E, et al. Mutation analysis of
TBX1 in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects. J Med
Genet. 2001;38(12):E45.
52. Hiroi N, Zhu H, Lee M, Funke B, Arai M, Itokawa M, et al. A 200-kb region of human
chromosome 22q11.2 confers antipsychotic-responsive behavioral abnormalities in mice.

Proceedings of the National Academy of Sciences of the United States of America.
2005;102(52):19132–7.
53.   Hiroi N, Hiramoto T, Harper KM, Suzuki G, Boku S. Mouse models of 22q11.2-associated
autism spectrum disorder. Autism. 2012;S1(001):1–9.
54.   Hiroi N, Takahashi T, Hishimoto A, Izumi T, Boku S, Hiramoto T. Copy Number Variation at
22q11.2: from rare variants to common mechanisms of developmental neuropsychiatric disorders.
Mol Psychiatry. 2013;18:1153–65.
55.   Hiroi N, Nishi A. Dimensional deconstruction and reconstruction of CNV-associated
neuropsychiatric disorders. In: Pletnikov MV, Waddington JL, editors. Modeling the
Psychopathological Dimensions of Schizophrenia: From Molecules to Behavior. Handbook of
Behavioral Neuroscience. 23. London, UK2015. p. 285–302.
56.   Nishi A, Hiroi N. Genetic Mechanisms Emerging from Mouse Models of CNV-Associated
Neuropsychiatric Disorders. In: Abel T, Nickl-Jockschat T, editors. *The Neurobiology of*
Schizophrenia. New York: Academic Press/Elsevier; 2016. p. 397–417.
57.   Suzuki G, Harper KM, Hiramoto T, Funke B, Lee M, Kang G, et al. Over-expression of a human
chromosome 22q11.2 segment including TXNRD2, COMT and ARVCF developmentally affects
incentive learning and working memory in mice. Human Molecular Genetics. 2009;18(20):3914–25.
58.   Hiramoto T, Kang G, Suzuki G, Satoh Y, Kucherlapati R, Watanabe Y, et al. Tbx1:
identification of a 22q11.2 gene as a risk factor for autism spectrum disorder in a mouse model.
Hum Mol Genet. 2011;20(24):4775–85.
59.   Takahashi T, Okabe S, Broin PO, Nishi A, Ye K, Beckert MV, et al. Structure and function of
neonatal social communication in a genetic mouse model of autism. Mol Psychiatry.
2016;21(9):1208–14.
60.   Kato R, Machida A, Nomoto K, Kang G, Hiramoto T, Tanigaki K, et al. Maternal approach
behaviors toward neonatal calls are impaired by mother's experiences of raising pups with a risk
gene variant for autism. Dev Psychobiol. 2021;63(1):108–13.
61.   Hiramoto T, Sumiyoshi A, Kato R, Yamauchi T, Takano T, Kang G, et al. Highly demarcated
structural alterations in the brain and impaired social incentive learning in Tbx1 heterozygous mice.
Molecular Psychiatry. 2025;30(5):1876–86.
62.   Hiramoto T, Sumiyoshi A, Yamauchi T, Tanigaki K, Shi Q, Kang G, et al. Tbx1, a gene encoded
in 22q11.2 copy number variant, is a link between alterations in fimbria myelination and cognitive
speed in mice. Mol Psychiatry. 2022;27(2):929–38.
63.   Nieder A, Mooney R. The neurobiology of innate, volitional and learned vocalizations in
mammals and birds. Philos Trans R Soc Lond B Biol Sci. 2020;375(1789):20190054.
64.   Boku S, Izumi T, Abe S, Takahashi T, Nishi A, Nomaru H, et al. Copy number elevation of
22q11.2 genes arrests the developmental maturation of working memory capacity and adult
neurogenesis. Molecular Psychiatry. 2018;23(4):985–92.
65.   Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell
transcriptional landscape of mammalian organogenesis. Nature. 2019;566(7745):496–502.
66.   Wang R, Zhang P, Wang J, Ma L, E W, Suo S, et al. Construction of a cross-species cell
landscape at single-cell level. Nucleic Acids Res. 2023;51(2):501–16.
67.   Marques S, van Bruggen D, Vanichkina DP, Floriddia EM, Munguba H, Varemo L, et al.
Transcriptional Convergence of Oligodendrocyte Lineage Progenitors during Development. Dev
Cell. 2018;46(4):504–17 e7.
68.   Kessaris N, Fogarty M, Iannarelli P, Grist M, Wegner M, Richardson WD. Competing waves of
oligodendrocytes in the forebrain and postnatal elimination of an embryonic lineage. Nat Neurosci.
2006;9(2):173–9.

69.   Menn B, Garcia-Verdugo JM, Yaschine C, Gonzalez-Perez O, Rowitch D, Alvarez-Buylla A.
Origin of oligodendrocytes in the subventricular zone of the adult brain. J Neurosci.
2006;26(30):7907–18.
70.   Rushing G, Ihrie RA. Neural stem cell heterogeneity through time and space in the
ventricular-subventricular zone. Front Biol (Beijing). 2016;11(4):261–84.
71.   Fuentealba LC, Rompani SB, Parraguez JI, Obernier K, Romero R, Cepko CL, et al.
Embryonic Origin of Postnatal Neural Stem Cells. Cell. 2015;161(7):1644–55.
72.   Hiramoto T, Boku S, Kang G, Abe S, Nagashima M, Barbachan e Silva M, et al.
Transcriptional regulation of neonatal neural stem cells is a determinant of social behavior. BioRxiv.
2023;11(12):468452.
73.   Nishiyama A, Lin XH, Giese N, Heldin CH, Stallcup WB. Co-localization of NG2
proteoglycan and PDGF alpha-receptor on O2A progenitor cells in the developing rat brain. J
Neurosci Res. 1996;43(3):299–314.
74.   Rivers LE, Young KM, Rizzi M, Jamen F, Psachoulia K, Wade A, et al. PDGFRA/NG2 glia
generate myelinating oligodendrocytes and piriform projection neurons in adult mice. Nat
Neurosci. 2008;11(12):1392–401.
75.   Sil S, Periyasamy P, Thangaraj A, Chivero ET, Buch S. PDGF/PDGFR axis in the neural
systems. Mol Aspects Med. 2018;62:63–74.
76.   Brandebura AN, Morehead M, Heller DT, Holcomb P, Kolson DR, Jones G, et al. Glial Cell
Expansion Coincides with Neural Circuit Formation in the Developing Auditory Brainstem. Dev
Neurobiol. 2018;78(11):1097–116.
77.   Beiter RM, Rivet-Noor C, Merchak AR, Bai R, Johanson DM, Slogar E, et al. Evidence for
oligodendrocyte progenitor cell heterogeneity in the adult mouse brain. Sci Rep. 2022;12(1):12921.
78.   Arnold JS, Werling U, Braunstein EM, Liao J, Nowotschin S, Edelmann W, et al. Inactivation
of Tbx1 in the pharyngeal endoderm results in 22q11DS malformations. Development.
2006;133(5):977–87.
79.   Matsuo N, Takao K, Nakanishi K, Yamasaki N, Tanda K, Miyakawa T. Behavioral profiles of
three C57BL/6 substrains. Front Behav Neurosci. 2010;4:29.
80.   Hiroi N. Critical Reappraisal of Mechanistic Links of Copy Number Variants to Dimensional
Constructs of Neuropsychiatric Disorders in Mouse Models. Psychiatry and Clinical
Neurosciences. 2018;72(5):301–21.
81.   Yang D, Wang Y, Qi T, Zhang X, Shen L, Ma J, et al. Phosphorylation of pyruvate
dehydrogenase inversely associates with neuronal activity. Neuron. 2024;112(6):959–71 e8.
82.   Readhead C, Popko B, Takahashi N, Shine HD, Saavedra RA, Sidman RL, et al. Expression of
a myelin basic protein gene in transgenic shiverer mice: correction of the dysmyelinating
phenotype. Cell. 1987;48(4):703–12.
83.   Meschkat M, Steyer AM, Weil MT, Kusch K, Jahn O, Piepkorn L, et al. White matter integrity in
mice requires continuous myelin synthesis at the inner tongue. Nat Commun. 2022;13(1):1163.
84.   Arellano JI, Duque A, Rakic P. A coming-of-age story: adult neurogenesis or adolescent
neurogenesis in rodents? Front Neurosci. 2024;18:1383728.
85.   Bronson FH, Dagg CP, Snell GD. Reproduction. In: Green EL, editor. Biology of the
Laboratory Mouse. Second Edition ed. New York: Dover Publications, Inc.; 2007. p. Oneline
publication.
86.   Nakamura M, Ye K, MB ES, Yamauchi T, Hoeppner DJ, Fayyazuddin A, et al. Computational
identification of variables in neonatal vocalizations predictive for postpubertal social behaviors in a
mouse model of 16p11.2 deletion. Mol Psychiatry. 2021;26(11):6578–88.
87.   Gao Z, Daquinag AC, Su F, Snyder B, Kolonin MG. PDGFRalpha/PDGFRbeta signaling
balance modulates progenitor cell differentiation into white and beige adipocytes. Development.
2018;145(1).

88.   Marcelin G, Ferreira A, Liu Y, Atlan M, Aron-Wisnewsky J, Pelloux V, et al. A PDGFRalpha-
Mediated Switch toward CD9(high) Adipocyte Progenitors Controls Obesity-Induced Adipose Tissue
Fibrosis. Cell Metab. 2017;25(3):673–85.
89.   Harper KM, Hiramoto T, Tanigaki K, Kang G, Suzuki G, Trimble W, et al. Alterations of social
interaction through genetic and environmental manipulation of the 22q11.2 gene Sept5 in the
mouse brain. Human Molecular Genetics. 2012;21(15):3489–99.
90.   Suzuki G, Harper KM, Hiramoto T, Sawamura T, Lee M, Kang G, et al. Sept5 deficiency exerts
pleiotropic influence on affective behaviors and cognitive functions in mice. Human Molecular
Genetics. 2009;18(9):1652–60.
91.   Yamauchi T, Kang G, Hiroi N. Heterozygosity of murine Crkl does not recapitulate behavioral
dimensions of human 22q11.2 hemizygosity. Genes Brain and Behavior. 2020;20(5):e12719.
92.   Paylor R, Spencer CM, Yuva-Paylor LA, Pieke-Dahl S. The use of behavioral test batteries, II:
effect of test interval. Physiol Behav. 2006;87(1):95–102.
93.   Fonseca AH, Santana GM, Bosque Ortiz GM, Bampi S, Dietrich MO. Analysis of ultrasonic
vocalizations from mice using computer vision and machine learning. Elife. 2021;10.
94.   Sakamoto Y, Takano T, Shimoyama S, Hiramoto T, Hiroi N, Nakamura K. Prepartum
bumetanide treatment reverses altered neonatal social communication but non-specifically
reduces post-pubertal social behavior in a mouse model of fragile X syndrome. Genomic
Psychiatry. 2025;1(1):61–72.
95.   Kaiser T, Allen HM, Kwon O, Barak B, Wang J, He Z, et al. MyelTracer: A Semi-Automated
Software for Myelin g-Ratio Quantification. eNeuro. 2021;8(4).
96.   Cioffi S, Martucciello S, Fulcoli FG, Bilio M, Ferrentino R, Nusco E, et al. Tbx1 regulates brain
vascularization. Hum Mol Genet. 2014;23(1):78–89.
97.   Cioffi S, Flore G, Martucciello S, Bilio M, Turturo MG, Illingworth E. VEGFR3 modulates brain
microvessel branching in a mouse model of 22q11.2 deletion syndrome. Life Sci Alliance.
2022;5(12).
98.   Allen Brain Atlas. Mouse Brain. http://wwwbrain-maporg. 2017.
99.   Nait Oumesmar B, Vignais L, Baron-Van Evercooren A. Developmental expression of
platelet-derived growth factor alpha-receptor in neurons and glial cells of the mouse CNS. J
Neurosci. 1997;17(1):125–39.
100.   Zhu X, Bergles DE, Nishiyama A. NG2 cells generate both oligodendrocytes and gray matter
astrocytes. Development. 2008;135(1):145–57.
101.   Marques S, Zeisel A, Codeluppi S, van Bruggen D, Mendanha Falcao A, Xiao L, et al.
Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system. Science.
2016;352(6291):1326–9.
102.   Ozarkar SS, Patel RKR, Vulli T, Friar CA, Burette AC, Philpot BD. Regional analysis of myelin
basic protein across postnatal brain development of C57BL/6J mice. Front Neuroanat.
2025;19:1535745.
103.   Bastedo WE, Scott RW, Arostegui M, Underhill TM. Single-cell analysis of mesenchymal
cells in permeable neural vasculature reveals novel diverse subpopulations of fibroblasts. Fluids
Barriers CNS. 2024;21(1):31.
104.   Takeuchi A, Takahashi Y, Iida K, Hosokawa M, Irie K, Ito M, et al. Identification of Qk as a
Glial Precursor Cell Marker that Governs the Fate Specification of Neural Stem Cells to a Glial Cell
Lineage. Stem Cell Reports. 2020;15(4):883–97.
105.   Dani N, Herbst RH, McCabe C, Green GS, Kaiser K, Head JP, et al. A cellular and spatial map
of the choroid plexus across brain ventricles and ages. Cell. 2021;184(11):3056–74 e21.
106.   Downes N, Mullins P. The development of myelin in the brain of the juvenile rat. Toxicol
Pathol. 2014;42(5):913–22.

107. Thompson CL, Ng L, Menon V, Martinez S, Lee CK, Glattfelder K, et al. A high-resolution
spatiotemporal atlas of gene expression of the developing mouse brain. Neuron. 2014;83(2):309–
23.
108. Redmond SA, Figueres-Onate M, Obernier K, Nascimento MA, Parraguez JI, Lopez-
Mascaraque L, et al. Development of Ependymal and Postnatal Neural Stem Cells and Their Origin
from a Common Embryonic Progenitor. Cell Rep. 2019;27(2):429–41 e3.
109. O'Rourke M, Cullen CL, Auderset L, Pitman KA, Achatz D, Gasperini R, et al. Evaluating
Tissue-Specific Recombination in a Pdgfralpha-CreERT2 Transgenic Mouse Line. PLoS One.
2016;11(9):e0162858.
110. Kang SH, Fukaya M, Yang JK, Rothstein JD, Bergles DE. NG2+ CNS glial progenitors remain
committed to the oligodendrocyte lineage in postnatal life and following neurodegeneration.
Neuron. 2010;68(4):668–81.
111. Tripathi RB, Rivers LE, Young KM, Jamen F, Richardson WD. NG2 glia generate new
oligodendrocytes but few astrocytes in a murine experimental autoimmune encephalomyelitis
model of demyelinating disease. J Neurosci. 2010;30(48):16383–90.
112. Komitova M, Zhu X, Serwanski DR, Nishiyama A. NG2 cells are distinct from neurogenic
cells in the postnatal mouse subventricular zone. J Comp Neurol. 2009;512(5):702–16.
113. Rosenberg AB, Roco CM, Muscat RA, Kuchina A, Sample P, Yao Z, et al. Single-cell profiling
of the developing mouse brain and spinal cord with split-pool barcoding. Science.
2018;360(6385):176–82.
114. Yao Z, Liu H, Xie F, Fischer S, Adkins RS, Aldridge AI, et al. A transcriptomic and epigenomic
cell atlas of the mouse primary motor cortex. Nature. 2021;598(7879):103–10.
115. Thompson PM, Jahanshad N, Ching CRK, Salminen LE, Thomopoulos SI, Bright J, et al.
ENIGMA and global neuroscience: A decade of large-scale studies of the brain in health and
disease across more than 40 countries. Transl Psychiatry. 2020;10(1):100.
116. Kochunov P, Hong LE, Dennis EL, Morey RA, Tate DF, Wilde EA, et al. ENIGMA-DTI:
Translating reproducible white matter deficits into personalized vulnerability metrics in cross-
diagnostic psychiatric research. Hum Brain Mapp. 2020;43(1):194–206.
117. Koshiyama D, Fukunaga M, Okada N, Morita K, Nemoto K, Usui K, et al. White matter
microstructural alterations across four major psychiatric disorders: mega-analysis study in 2937
individuals. Mol Psychiatry. 2020;25(4):883–95.
118. Kelly S, Jahanshad N, Zalesky A, Kochunov P, Agartz I, Alloza C, et al. Widespread white
matter microstructural differences in schizophrenia across 4322 individuals: results from the
ENIGMA Schizophrenia DTI Working Group. Mol Psychiatry. 2018;23(5):1261–9.
119. Villalon-Reina JE, Martinez K, Qu X, Ching CRK, Nir TM, Kothapalli D, et al. Altered white
matter microstructure in 22q11.2 deletion syndrome: a multisite diffusion tensor imaging study.
Mol Psychiatry. 2020;25(11):2818–31.

1195

1196

1197

1198   **Figure 1**

1199

1200

1201

1202 **Figure 2**

1203
1204
1205

1206   **Figure 3**

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

 Figure 4

1233 **Figure 5**

1234

1235

1236 **Figure 6**

1237

1238 **Figure 7**

1239

1240

1241   **Figure 8**

1256 **Figure 9**

1257

1258

1259 **Supplementary Information**

1260

1261 **Figure S1.** Body weights o*Tbx1*$^{+/+}$ and o*Tbx1*$^{+/−}$ mice. o*Tbx1*$^{+/+}$ and o*Tbx1*$^{+/−}$ mice had
1262 indistinguishable body weight increases from P8 to 2 months of age (P8, U = 140, p = 0.800;
1263 P12, U = 153, p = 0.466; 1 month, U = 135.5, p = 0.7035; 2 months, U = 102, p = 0.917).
1264 o*Tbx1*$^{+/+}$ P8, N = 27, P12; N = 30; 1M, 1 month, N = 27; 2M, 2 months, N = 21: o*Tbx1*$^{+/−}$: P8, N
1265 = 11, P12, N = 12; 1M, N = 11; 2M, N = 10.

1266

1267
1268 **Supplementary Figure S2.** The myelin thickness of axons for o*Tbx1*$^{+/−}$ versus o*Tbx1*$^{+/+}$ mice
1269 versus axon diameter was similar along the axon diameter range (genotype, p = 0.931;
1270 genotype × axon diameter, p = 0.814).
1271

Figure	Control Gene	Gene	Treatment	# of Samples	Average	s.e.m	Shapiro-Wilk Test				Levene's Test				Linear Mixed Model					Mann Whitney U Test						
							Statistic	p_value	p_adj (FDR BH)	Reject (FDR5%)	Statistic	p_value	p_adj (FDR BH)	Reject (FDR5%)	Source	F	df1	df2	Sig.	Sample 1	Sample 2	Statistic	R (Z score)	p_value	p_adj (FDR BH)	Reject (FDR5%)
Figure 1	Pgk1	Tbx1	control siRNA	6	1	0.049186162	0.93572346	0.624966577	0.805579616	FALSE	1.758766194	0.21744683	0.297607562	FALSE	Corrected Model	2.006348255	11	89	0.03681448	control siRNA	Tbx1 siRNA	7	0.50847518	0.008658009	0.012987013	TRUE
			Tbx1 siRNA	5	0.464725499	0.116052639	0.932068851	0.610545167	0.805579616	FALSE					Gene	0.843979959	5	89	0.52216149							
		Cspg4	control siRNA	9	1	0.066746189	0.960660226	0.805205712	0.878406231	FALSE	1.522725608	0.235026707	0.297607562	FALSE	Treatment	14.66501979	1	89	0.00023857	control siRNA	Tbx1 siRNA	8	0.676423086	0.002756067	0.008268202	TRUE
			Tbx1 siRNA	9	0.685034037	0.037803054	0.980712867	0.967840714	0.967840714	FALSE					Gene * Treatment	0.843979957	5	89	0.522161492							
		MAG	control siRNA	9	1	0.041029197	0.908420558	0.304976014	0.731942433	FALSE	3.473378198	0.080821307	0.297607562	FALSE						control siRNA	Tbx1 siRNA	11	0.613984032	0.007774578	0.012987013	TRUE
			Tbx1 siRNA	9	0.670998004	0.157173093	0.903392871	0.272326671	0.731942433	FALSE																
		MBP	control siRNA	9	1	0.056959774	0.921773656	0.40712653	0.805579616	FALSE	0.679262555	0.421957432	0.421957432	FALSE						control siRNA	Tbx1 siRNA	6	0.718049122	0.00123406	0.00740436	TRUE
			Tbx1 siRNA	9	0.553891554	0.084308916	0.872112121	0.129519549	0.518078195	FALSE																
		MOG	control siRNA	9	1	0.064281655	0.948291674	0.671316347	0.805579616	FALSE	1.89729647	0.18734571	0.297607562	FALSE						control siRNA	Tbx1 siRNA	14	0.551544978	0.018757713	0.018757713	TRUE
			Tbx1 siRNA	9	0.60385684	0.126671201	0.85412067	0.082708168	0.496249006	FALSE																
		Plp1	control siRNA	9	1	0.06060843	0.939293908	0.574485759	0.805579616	FALSE	1.437463429	0.248006301	0.297607562	FALSE						control siRNA	Tbx1 siRNA	14	0.551544978	0.018757713	0.018757713	TRUE
			Tbx1 siRNA	9	0.700871358	0.116118855	0.84502056	0.065659769	0.496249006	FALSE																
	18S	Tbx1	control siRNA	6	1	0.078568382	0.92617953	0.550937859	0.734583812	FALSE	0.009347636	0.925096749	0.925096749	FALSE	Corrected Model	2.450555968	11	89	0.010091634	control siRNA	Tbx1 siRNA	6	0.554700196	0.004329004	0.007774578	TRUE
			Tbx1 siRNA	5	0.440976588	0.066877094	0.97650378	0.915107555	0.915107555	FALSE					Gene	1.101427491	5	89	0.365404553							
		Cspg4	control siRNA	9	1	0.044646855	0.916096471	0.360902781	0.662239178	FALSE	0.016880314	0.898245734	0.925096749	FALSE	Treatment	17.51187144	1	89	6.68502E-05	control siRNA	Tbx1 siRNA	4	0.759675158	0.000493624	0.002961744	TRUE
			Tbx1 siRNA	9	0.687687722	0.045939538	0.919279302	0.386306187	0.662239178	FALSE					Gene * Treatment	1.101427494	5	89	0.365404552							
		MAG	control siRNA	9	1	0.055152176	0.966495138	0.863205301	0.915107555	FALSE	1.450912436	0.24589768	0.643468053	FALSE						control siRNA	Tbx1 siRNA	11	0.613984032	0.007774578	0.007774578	TRUE
			Tbx1 siRNA	9	0.642342516	0.157925591	0.844966625	0.065569561	0.242978818	FALSE																
		MBP	control siRNA	9	1	0.053726606	0.931315789	0.493974756	0.734583812	FALSE	0.206705208	0.655469861	0.925096749	FALSE						control siRNA	Tbx1 siRNA	7	0.697236104	0.00185109	0.00555327	TRUE
			Tbx1 siRNA	9	0.544403139	0.082596116	0.85329064	0.080992939	0.242978818	FALSE																
		MOG	control siRNA	9	1	0.06650766	0.957538993	0.772242353	0.915107555	FALSE	1.178141059	0.293818671	0.643468053	FALSE						control siRNA	Tbx1 siRNA	11	0.613984032	0.007774578	0.007774578	TRUE
			Tbx1 siRNA	9	0.588913841	0.123648773	0.841639086	0.060228776	0.242978818	FALSE																
		Plp1	control siRNA	9	1	0.04046838	0.874188469	0.136290611	0.327097466	FALSE	1.045606567	0.321734027	0.643468053	FALSE						control siRNA	Tbx1 siRNA	11	0.613984032	0.007774578	0.007774578	TRUE
			Tbx1 siRNA	9	0.692323814	0.117369849	0.755073552	0.00620775	0.074492996	FALSE																

Figure 4A — Number

Figure	Measurement	Call type	PND	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MWU Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 4A	Number	Harmonics	P8	oTbx1+/+	27	25.44444444	6.455928465	0.734571278	1.2196E-05	2.54525E-05	TRUE	0.588767426	1	36	0.447899604	0.866000524	FALSE	Corrected Model	6.94053254	47	912	0	oTbx1+/+	oTbx1+/-	139	0.049603452	0.775387862	0.990742675	FALSE
				oTbx1+/-	11	18	5.412107639	0.869194567	0.075694919	0.0825276275	FALSE							Genotype	0.196622137	1	912	0.657566457							
			P12	oTbx1+/+	30	18.53333333	4.416201045	0.769232988	1.89257E-05	3.63374E-05	TRUE	0.080099472	1	40	0.778622211	0.866000524	FALSE	PND	6.147480911	1	912	0.013339656	oTbx1+/+	oTbx1+/-	180	0	1	1	FALSE
				oTbx1+/-	12	17.41666667	6.962516888	0.741135657	0.002162845	0.00288379	TRUE							Call type	19.63969375	11	912	0							
		step_down	P8	oTbx1+/+	27	1.518518519	0.424902772	0.641833842	6.62769E-07	2.65108E-06	TRUE	0.588478181	1	36	0.448010251	0.866000524	FALSE	Genotype * PND	4.835206967	1	912	0.028134731	oTbx1+/+	oTbx1+/-	140	0.044382036	0.799797921	0.990742675	FALSE
				oTbx1+/-	11	1	0.301511345	0.863411665	0.063748829	0.071161484	FALSE							PND * Call type	1.728524601	11	912	0.062712426							
			P12	oTbx1+/+	30	3.866666667	1.062131162	0.71998024	3.08876E-06	8.23669E-06	TRUE	0.194362482	1	40	0.661683974	0.866000524	FALSE	Call type * Genotype	0.707710242	11	912	0.731986987	oTbx1+/+	oTbx1+/-	174	0.025776963	0.879981095	0.990742675	FALSE
				oTbx1+/-	12	1.166666667	0.553370565	0.517821491	2.3982E-05	2.3982E-05	TRUE							Genotype * PND * Call type	0.368631844	11	912	0.967949606							
		short	P8	oTbx1+/+	27	19.92592593	3.665472193	0.834571779	0.000578805	0.000817137	TRUE	3.769928996	1	36	0.060039818	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	121	0.14358894	0.390409044	0.990742675	FALSE
				oTbx1+/-	11	12.27272727	2.157822594	0.916108012	0.287544787	0.287544787	FALSE																		
			P12	oTbx1+/+	30	27.53333333	5.654971027	0.831702948	0.000226076	0.000394014	TRUE	0.789588022	1	40	0.379537318	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	168.5	-0.049405846	0.773086018	0.990742675	FALSE
				oTbx1+/-	12	25.91666667	6.770544866	0.883340597	0.096735299	0.098793496	FALSE																		
		down_fm	P8	oTbx1+/+	27	17.40740741	3.629280755	0.831780851	0.000512651	0.00074674	TRUE	0.289267018	1	36	0.594000243	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	145.5	0.015664248	0.949461731	0.990742675	FALSE
				oTbx1+/-	11	15	4.872930796	0.815746427	0.01516132	0.017749838	TRUE																		
			P12	oTbx1+/+	30	14.46666667	3.822418243	0.713233709	2.44507E-06	6.90372E-06	TRUE	1.591281097	1	40	0.214450379	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	148.5	-0.135329056	0.400911456	0.990742675	FALSE
				oTbx1+/-	12	26.5	10.29599805	0.770105839	0.004369121	0.005242945	TRUE																		
		up_fm	P8	oTbx1+/+	27	17.7037037	4.965641153	0.629247069	4.62791E-07	2.46822E-06	TRUE	2.477639992	1	36	0.124223918	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	87	0.321117085	0.048527629	0.990742675	FALSE
				oTbx1+/-	11	4.909090909	1.889947308	0.722703338	0.00091824	0.001259304	TRUE																		
			P12	oTbx1+/+	30	27.83333333	7.521934719	0.702088237	1.6735E-06	5.35522E-06	TRUE	0.168045902	1	40	0.684041632	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	175	-0.022148083	0.901682953	0.990742675	FALSE
				oTbx1+/-	12	23.83333333	7.906492665	0.838153899	0.026294718	0.030051106	TRUE																		
		two_steps	P8	oTbx1+/+	27	1	0.301940542	0.694654405	3.26996E-06	8.26094E-06	TRUE	0.0764692	1	36	0.783722016	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	141.5	0.036549912	0.849168654	0.990742675	FALSE
				oTbx1+/-	11	1.181818182	0.572385609	0.571349978	1.12326E-05	2.45076E-05	TRUE																		
			P12	oTbx1+/+	30	0.7	0.215225089	0.653556228	3.5184E-07	2.12738E-06	TRUE	0.113222806	1	40	0.7382634	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	165	-0.064442408	0.690391077	0.990742675	FALSE
				oTbx1+/-	12	0.583333333	0.192996049	0.768340707	0.004181752	0.005146772	TRUE																		
		flat	P8	oTbx1+/+	27	17.7037037	4.250294045	0.742479801	1.6037E-05	3.20739E-05	TRUE	0.680374459	1	36	0.414890086	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	133	0.080931948	0.634139425	0.990742675	FALSE
				oTbx1+/-	11	12.45454545	3.866790157	0.875244677	0.090499207	0.094433956	FALSE																		
			P12	oTbx1+/+	30	14.86666667	3.159635267	0.817323029	0.000137908	0.000228261	TRUE	0.063139102	1	40	0.802887963	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	170.5	-0.040813525	0.815460496	0.990742675	FALSE
				oTbx1+/-	12	17.41666667	6.344239016	0.755882382	0.003080417	0.003891054	TRUE																		
		chevron	P8	oTbx1+/+	27	18.11111111	5.570407498	0.636691451	5.71795E-07	2.65108E-06	TRUE	0.0673775	1	36	0.796672834	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	110.5	0.198413808	0.237616458	0.990742675	FALSE
				oTbx1+/-	11	14.09090909	10.5833957	0.4472633	3.545636E-07	2.12738E-06	TRUE																		
			P12	oTbx1+/+	30	13.56666667	3.444363489	0.749773264	9.03E-06	2.064E-05	TRUE	0.345282084	1	40	0.560098838	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	160	0.08592321	0.591959808	0.990742675	FALSE
				oTbx1+/-	12	17.33333333	9.726045424	0.573818922	6.544E-05	0.000112183	TRUE																		
		complex	P8	oTbx1+/+	27	2.592592593	0.879100906	0.63989383	6.26769E-07	2.65108E-06	TRUE	0.506662225	1	36	0.481174865	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	112	0.190581684	0.250721005	0.990742675	FALSE
				oTbx1+/-	11	1.454545455	1.267521822	0.409144521	1.25664E-07	1.20637E-06	TRUE																		
			P12	oTbx1+/+	30	4.833333333	2.496841439	0.638445973	2.22498E-07	1.77998E-06	TRUE	0.031412363	1	40	0.860217983	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	161.5	-0.07947897	0.630588582	0.990742675	FALSE
				oTbx1+/-	12	4.583333333	2.496841439	0.560712337	5.1415E-05	9.14045E-05	TRUE																		
		step_up	P8	oTbx1+/+	27	3.703703704	0.802089179	0.814462734	0.000236307	0.000365895	TRUE	0.406675225	1	36	0.527697447	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	140	-0.044382036	0.799797921	0.990742675	FALSE
				oTbx1+/-	11	4.727272727	1.839421397	0.762873292	0.003061853	0.00389101054	TRUE																		
			P12	oTbx1+/+	30	3.2	0.848799016	0.710336447	2.21376E-06	6.64128E-06	TRUE	0.034791252	1	40	0.852976553	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	154.5	-0.109552093	0.50033814	0.990742675	FALSE
				oTbx1+/-	12	3.666666667	1.0396192	0.877785325	0.082109146	0.087583089	FALSE																		
		mult_steps	P8	oTbx1+/+	27	0.074074074	0.051361129	0.29382658	3.32643E-10	5.58344E-09	TRUE	0.028883184	1	36	0.866000524	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	146	-0.01305354	0.949461731	0.990742675	FALSE
				oTbx1+/-	11	0.090909091	0.090909091	0.344991028	2.2434E-08	2.69208E-07	TRUE																		
			P12	oTbx1+/+	30	0.466666667	0.046320556	0.275375307	4.40171E-11	2.11282E-09	TRUE	0.03421728	1	40	0.854180142	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	177	-0.012888482	0.945286072	0.990742675	FALSE
				oTbx1+/-	12	0.083333333	0.083333333	0.32692748	3.1129E-08	4.13925E-06	TRUE																		
		rev_chevron	P8	oTbx1+/+	27	1.592592593	0.475194545	0.708100617	5.03586E-06	1.20861E-05	TRUE	2.543697003	1	36	0.119478121	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	104.5	0.229742304	0.168812806	0.990742675	FALSE
				oTbx1+/-	11	0.363636364	0.278721995	0.47351861	7.28965E-07	2.69156E-06	TRUE																		
			P12	oTbx1+/+	30	2.133333333	0.97052342	0.474628329	2.91922E-09	4.67076E-08	TRUE	4.044048543	1	40	0.834828853	0.866000524	FALSE						oTbx1+/+	oTbx1+/-	123	-0.24488115	0.116641288	0.990742675	FALSE
				oTbx1+/-	12	2.833333333	1.407950011	0.621079922	0.000161712	0.000258739	TRUE																		

Figure 4A — Number (All)

Figure	Measurement	Call type	PND	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MWU Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 4A	Number	All	P8	oTbx1+/+	27	126.7777778	26.50132144	0.78554976	7.77043E-05	0.000157208	TRUE	1.121799982	1	36	0.296586227	0.593172454	FALSE	Corrected Model	0.636064898	3	76	0.594070046	oTbx1+/+	oTbx1+/-	126.5	0.114871152	0.504762746	0.73135923	FALSE
				oTbx1+/-	11	85.54545455	23.66407467	0.834964693	0.027126787	0.027126787	TRUE							Genotype	0.155970557	1	76	0.693999978							
			P12	oTbx1+/+	30	130.6	27.57400075	0.804300725	7.86042E-05	0.000157208	TRUE	0.000269074	1	40	0.98699405	0.98699405	FALSE	PND	1.419833311	1	76	0.237138636	oTbx1+/+	oTbx1+/-	166.5	-0.057998167	0.73135923	0.73135923	FALSE
				oTbx1+/-	12	143.3333333	45.18922058	0.79983592	0.009328736	0.01248315	TRUE							Genotype * PND	1.103292412	1	76	0.296873255							

Generalized Linear Mixed Model — Number (Harmonics etc.)

Measurement	Call type	PND	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene's Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	GLMM Source	F	df1	df2	Sig.	MWU Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Number	Harmonics	P8	PdgfraCre;Tbx1+/+	8	9.625	2.672194791	0.919298375	0.42419617	0.449148886	FALSE	1.65961678	2	24	0.211306457	0.559313096	FALSE	Corrected Model	4.565593657	71	612	0	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	28.5	0.091893381	0.798445998	0.974374778	FALSE
			WT;Tbx1+/flox	18.125	11.96786993	0.59934529	0.000156224	0.000749877	TRUE							Genotype	2.526849017	2	612	0.080743877	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	15.5	-0.539886352	0.024876	0.488370247	FALSE	
			WT;Tbx1+/+	11	42.27272727	11.79066727	0.863657084	0.064216294	0.087237229	FALSE							PND	0.381377323	1	612	0.537096747	WT;Tbx1+/flox	WT;Tbx1+/+	19	-0.473584519	0.04088275	0.644874038	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	11.22222222	7.025306988	0.600651882	9.73774E-05	0.000559321	TRUE	1.158509452	2	27	0.329073887	0.559313096	FALSE	Call type	18.61506233	11	612	0	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	40	-0.010406509	1	1	FALSE
			WT;Tbx1+/flox	11	11.66666667	5.525194818	0.765487459	0.008192575	0.016093411	TRUE							Genotype * PND * Call type	1.47748124	57	612	0.015566306	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	32.5	-0.291382252	0.258083093	0.743019943	FALSE	
			WT;Tbx1+/+	12	29.16666667	8.235946095	0.864345385	0.055443248	0.076753728	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	35	-0.294653333	0.193005137	0.644874038	FALSE	
	step_down	P8	PdgfraCre;Tbx1+/+	8	0.875	0.295048422	0.835213557	0.067235843	0.08964779	FALSE	1.361546454	2	24	0.275355808	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	31	0.026255252	0.95912979	1	FALSE
			WT;Tbx1+/flox	8	0.377964473	0.85994915	0.119934837	0.146341157	FALSE													PdgfraCre;Tbx1+/+	WT;Tbx1+/+	34.5	-0.179962117	0.49201291	0.906204631	FALSE	
			WT;Tbx1+/+	11	2.363636364	0.956059405	0.720871942	0.000869463	0.002504053	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	35.5	-0.161018736	0.544759334	0.906204631	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	0.888888889	0.611111111	0.570366044	4.29426E-05	0.000281079	TRUE	2.413977888	2	27	0.108515742	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	26.5	-0.291382252	0.258083093	0.743019943	FALSE
			WT;Tbx1+/flox	11	5.111111111	2.6793679	0.695406931	0.001254342	0.003473563	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	26.5	-0.42647193	0.058483312	0.743019943	FALSE	
			WT;Tbx1+/+	12	5.166666667	1.565892781	0.863856	0.054652247	0.076753728	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	47.5	-0.100802546	0.702085531	0.939137333	FALSE	
	short	P8	PdgfraCre;Tbx1+/+	8	15.125	4.834539053	0.867444039	0.142278889	0.165227097	FALSE	1.622768353	2	24	0.218268212	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	30.5	0.039382877	0.95912759	1	FALSE
			WT;Tbx1+/flox	8	13.25	4.087218754	0.849020808	0.093116392	0.117620706	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	27	-0.322037473	0.177370273	0.644874038	FALSE	
			WT;Tbx1+/+	11	28.27272727	7.311374587	0.875466193	0.091090759	0.11711669	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	27.5	-0.315261783	0.206001429	0.644874038	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	13	3.905124838	0.914834887	0.351194856	0.383121661	FALSE	2.151774119	2	27	0.135799339	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	33.5	-0.145691126	0.604812834	0.926521789	FALSE
			WT;Tbx1+/flox	9	33.33333333	13.89844116	0.807207052	0.024673583	0.039477739	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	34	-0.310161403	0.169421291	0.644874038	FALSE	
			WT;Tbx1+/+	12	34.08333333	8.745958663	0.909009298	0.207195733	0.22950912	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	46	-0.124064561	0.60159902	0.926521789	FALSE	
	down_fm	P8	PdgfraCre;Tbx1+/+	8	9.5	3.212698021	0.861074309	0.123069142	0.14768297	FALSE	1.921139644	2	24	0.168297426	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	32	0	1	1	FALSE
			WT;Tbx1+/flox	8	11.75	5.456810947	0.738429175	0.006087749	0.012504854	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	22	-0.416754377	0.075414781	0.644874038	FALSE	
			WT;Tbx1+/+	11	27.27272727	6.85842668	0.884384045	0.118227917	0.146361157	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	24.5	-0.369395925	0.1288402	0.644874038	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	12.88888889	9.075982415	0.490677123	4.96662E-06	5.55994E-05	TRUE	0.180736373	2	27	0.835656843	0.911625647	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	37.5	-0.062439054	0.863307281	1	FALSE
			WT;Tbx1+/flox	12	12.22222222	5.484264809	0.778607516	0.011607818	0.021429818	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	46.5	-0.116310526	0.651052972	0.937516279	FALSE	
			WT;Tbx1+/+	12	17.33333333	5.805205856	0.805205856	0.010743424	0.020355961	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	44	-0.155080702	0.507876025	0.906204631	FALSE	
	up_fm	P8	PdgfraCre;Tbx1+/+	8	10.375	4.788145704	0.799663368	0.028388995	0.044238038	TRUE	1.244964464	2	24	0.305890073	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	28	-0.105021006	0.720901321	0.939137333	FALSE
			WT;Tbx1+/flox	8	9.875	3.026298423	0.916331725	0.400811221	0.430722507	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	26	-0.340980854	0.151755709	0.644874038	FALSE	
			WT;Tbx1+/+	11	28.72727273	10.97773856	0.713897791	0.000706475	0.002248161	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	28	-0.303094092	0.206001429	0.644874038	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	10.11111111	7.091552965	0.529781177	1.43217E-05	0.000128896	TRUE	1.093047307	2	27	0.349570685	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	25	-0.32260178	0.190523596	0.81540225	FALSE
			WT;Tbx1+/flox	11	15.59479179	7.69584027	0.768342027	0.008270225	0.016093411	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	29	-0.387017514	0.081543225	0.644874038	FALSE	
			WT;Tbx1+/+	12	34.66666667	13.49485499	0.683792344	0.000590864	0.00202582	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	50.5	-0.054278246	0.8620692	1	FALSE	
	two_steps	P8	PdgfraCre;Tbx1+/+	8	0.125	0.125	0.418398447	1.04723E-06	1.88502E-05	TRUE	2.325091481	2	24	0.119400488	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	23	-0.29269264	0.382284382	0.834075016	FALSE
			WT;Tbx1+/flox	8	0.75	0.411877236	0.724167693	0.004213657	0.009480728	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	15.5	-0.539886352	0.02034876	0.488370247	FALSE	
			WT;Tbx1+/+	11	1.818181818	0.600275419	0.82936025	0.022898111	0.037469637	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	28	-0.303094092	0.206001429	0.644874038	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	0.333333333	0.333333333	0.389841298	3.21749E-07	1.1583E-05	TRUE	0.817105263	2	27	0.452335248	0.603113664	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	28.5	-0.249756216	0.340106952	0.78989045	FALSE
			WT;Tbx1+/flox	9	0.666666667	0.333333333	0.710163506	0.00186499	0.0049713	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	38.5	-0.24073088	0.310026197	0.769722013	FALSE	
			WT;Tbx1+/+	12	1	0.40824829	0.7517227	0.002785506	0.006907527	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	50.5	-0.054278246	1	1	FALSE	
	flat	P8	PdgfraCre;Tbx1+/+	8	11.875	3.656879525	0.882816897	0.200332641	0.225374222	FALSE	3.818027267	2	24	0.036336836	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	20.5	0.301935393	0.278623479	0.743019943	FALSE
			WT;Tbx1+/flox	8	6.375	2.744718955	0.801139132	0.029440249	0.044238038	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	28	-0.30304661	0.02034876	0.488370247	FALSE	
			WT;Tbx1+/+	11	30.18181818	8.790302696	0.855776158	0.050750623	0.073080898	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	16	-0.530414661	0.02034876	0.488370247	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	9.444444444	3.628914561	0.634351254	0.000241961	0.001024778	TRUE	0.955031557	2	27	0.39741914	0.59612871	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	36	-0.093658581	0.730440148	0.974374778	FALSE
			WT;Tbx1+/flox	9	12.44444444	4.972454991	0.834538516	0.05019806	0.073080898	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	32	-0.341177544	0.128534005	0.644874038	FALSE	
			WT;Tbx1+/+	12	20.75	5.615353682	0.900674282	0.16182923	0.184948255	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	36.5	-0.271391228	0.347766234	0.740772384	FALSE	
	chevron	P8	PdgfraCre;Tbx1+/+	8	11.375	5.990876483	0.735010786	0.005568815	0.011968073	TRUE	0.858879466	2	24	0.436252017	0.603113096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	31	0.026255252	0.95912979	1	FALSE
			WT;Tbx1+/flox	8	6.059231446	0.550872664	4.19244E-05	0.000281079	TRUE													PdgfraCre;Tbx1+/+	WT;Tbx1+/+	27.5	-0.312565783	0.206001429	0.644874038	FALSE	
			WT;Tbx1+/+	11	29.63636364	11.76610906	0.702313085	0.000500922	0.001800332	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	22	-0.416754377	0.075414781	0.743019943	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	11.22222222	7.827618088	0.560909917	3.32517E-05	0.000266014	TRUE	0.058536613	2	27	0.943263064	0.984274502	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	33	-0.156097635	0.545701357	0.906204631	FALSE
			WT;Tbx1+/flox	9	13.22222222	5.340562231	0.814027777	0.029492025	0.044238038	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	30	-0.248129123	0.277311729	0.743019943	FALSE	
			WT;Tbx1+/+	12	15.58333333	5.334931105	0.817775681	0.015028138	0.025762523	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	51.5	-0.038770175	0.917007451	1	FALSE	
	complex	P8	PdgfraCre;Tbx1+/+	8	1.5	0.80039053	0.604228412	0.000178175	0.000810733	TRUE	1.111310057	2	24	0.345479196	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	25	-0.131276258	0.645376845	0.937516279	FALSE
			WT;Tbx1+/flox	8	1.625	0.80039053	0.783670395	0.019092425	0.031968712	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	31.5	-0.20256732	0.510062422	0.906204631	FALSE	
			WT;Tbx1+/+	11	4.090909091	1.928044442	0.714449593	0.000718163	0.002248161	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	38.5	-0.104188594	0.71678442	0.939137333	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	3.888888889	1.795054936	0.451702288	1.72645E-06	2.48608E-05	TRUE	0.013210185	2	27	0.986883061	0.986883061	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	31	-0.177923671	0.436281366	0.937516279	FALSE
			WT;Tbx1+/flox	9	4	1.795054936	0.783337582	0.013157116	0.023682808	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	39	-0.232621053	0.310026197	0.769722013	FALSE	
			WT;Tbx1+/+	12	3.666666667	1.275250025	0.815437471	0.014110863	0.022730053	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	38.5	-0.104188594	0.71678442	0.939137333	FALSE	
	step_up	P8	PdgfraCre;Tbx1+/+	8	3.625	1.450831043	0.863678426	0.130614865	0.154168365	FALSE	0.399734807	2	24	0.674879426	0.809855311	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	30.5	0.039382877	0.95912759	1	FALSE
			WT;Tbx1+/flox	8	2.75	1.12994943	0.840049789	0.075410948	0.098719786	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	36.5	-0.14207356	0.599827278	0.926521789	FALSE	
			WT;Tbx1+/+	11	4.454545455	1.503714409	0.760819461	0.002878136	0.006907527	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	33	-0.17049062	0.49201291	0.906204631	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	2.555555556	1.415304356	0.6160407291	0.0001459	0.000749877	TRUE	1.307144496	2	27	0.287172685	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	36.5	0.083252072	0.796174414	0.974374778	FALSE
			WT;Tbx1+/flox	9	1.333333333	0.600925213	0.724908399	0.002770316	0.006907527	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	33	-0.325044476	0.147939986	0.644874038	FALSE	
			WT;Tbx1+/+	12	5.083333333	1.694278846	0.780364431	0.00565159	0.011968073	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	28.5	-0.395455789	0.081543225	0.644874038	FALSE	
	mult_steps	P8	PdgfraCre;Tbx1+/+	8	0	0	1	1	1	FALSE	1.580246914	2	24	0.226612028	0.559313096	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	32	0	1	1	FALSE
			WT;Tbx1+/flox	8	0.181818182	0.121967344	0.485720309	1.020866E-06	1.88502E-05	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	36	-0.151547046	0.544759334	0.906204631	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	0	0	1	1	1	FALSE	1.62	2	27	0.216549255	0.559313096	FALSE						WT;Tbx1+/flox	WT;Tbx1+/+	40.5	-0.139572632	0.553791719	0.906204631	FALSE	
			WT;Tbx1+/flox	9	0.166666667	0.11236664	0.464652258	9.81123E-06	0.000100916	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	45	-0.139572632	0.553791719	0.906204631	FALSE	
	rev_chevron	P8	PdgfraCre;Tbx1+/+	8	1.125	0.548943791	0.641202	0.00049057	0.00180332	TRUE	0.474061795	2	24	0.628174436	0.793483498	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	28	-0.105021006	0.720901321	0.939137333	FALSE
			WT;Tbx1+/flox	8	2.25	1.264205001	0.722041682	0.003988818	0.009264352	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	39	-0.094716904	0.777789468	0.974374778	FALSE	
			WT;Tbx1+/+	11	1.454545455	0.651761757	0.720728359	0.000865751	0.002504053	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	40	0.075773523	0.777789468	0.974374778	FALSE	
		P12	PdgfraCre;Tbx1+/+	9	2.77777778	2.777777778	0.389841298	3.21749E-07	1.1583E-05	TRUE	0.389841298	2	27	0.754642872	0.862448996	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	33	-0.156097635	0.545701357	0.906204631	FALSE
			WT;Tbx1+/flox	11	0.666666667	0.591160075	0.353161605	0.000539321	0.001800332	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	40	-0.217112982	0.345102575	0.78989045	FALSE	
			WT;Tbx1+/+	12	1.166666667	0.60233702	0.560702	0.000470016	0.00180332	TRUE												WT;Tbx1+/flox	WT;Tbx1+/+	47	-0.124064561	0.645376845	0.937516279	FALSE	

Repeated Measures 2 Way ANOVA — Number (All)

Measurement	Call type	PND	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene's Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	ANOVA Source	Correction	Type III Sum of Squares	df	Mean Square	F	Sig.	t-test Sample1	Sample2	Statistic	dof	Cohen's d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Number	All	P8	PdgfraCre;Tbx1+/+	8	75.125	22.42800983	0.919578326	0.426451218	0.426451218	FALSE	1.430162095	2	24	0.258940451	0.47061792	FALSE	Within-Subjects							PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-0.040423302	14	-0.020211651	0.968326472	0.968326472	FALSE
			WT;Tbx1+/flox	8	76.75	33.36370422	0.748178822	0.00779798	0.023193394	TRUE							PND	Sphericity Assumed	1107.334091	1	1107.334091	0.104222605	0.749615812	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-1.942536002	17	-0.902619145	0.068525485	0.25684836	FALSE
			WT;Tbx1+/+	11	200.7272727	52.31132172	0.852676341	0.046247915	0.069371873	FALSE							PND * Genotype	Sphericity Assumed	24001.49242	2	12000.74621	1.129513707	0.339774875	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-0.751221117	16	-0.341296056	0.46342642	0.58651612	FALSE
		P12	PdgfraCre;Tbx1+/+	9	78.33333333	46.90830298	0.573567292	4.68257E-05	0.000280954	TRUE	0.775145688	2	27	0.47061792	0.47061792	FALSE	Error(PND)	Sphericity Assumed	254992.8409	24	10624.7017			WT;Tbx1+/flox	WT;Tbx1+/+	-0.543641453	19	-0.341296056	0.695139663	0.361510052	FALSE
			WT;Tbx1+/flox	9	131.4444444	52.89667893	0.822522253	0.036792463	0.069371873	FALSE							Between-Subjects							WT;Tbx1+/flox	WT;Tbx1+/+	-1.389484067	19	-0.612704882	0.361510052	0.361510052	FALSE
			WT;Tbx1+/+	12	169.1666667	44.30897599	0.914009214	0.240072481	0.288086977	FALSE							Intercept		787339.8818	1	787339.8818	25.39936419	3.75383E-05	WT;Tbx1+/flox	WT;Tbx1+/+	-0.549424201	19	-0.2422733	0.589114473	0.706937367	FALSE
																	Genotype		112088.5152	2	56044.25758	1.807972061	0.18561632								
																	Error		743961.8182	24	30998.40909										

Figure 4B

Figure	Measurement	Call type	PND	Genotype	# of Samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MW Sample1	MW Sample2	MW Statistic	R (Z score)	MW p_value	MW p_adj(FDR BH)	MW Rejects(FDR 5%)
Figure 4B	Proportion	Harmonics	P8	oTbx1/+	27	17.03450018	2.203565508	0.951793015	0.236913264	0.27736187	FALSE	0.007209331	1	36	0.932805151	0.958177206	FALSE	Corrected Model	12.69567482	47	912	0	oTbx1/+	oTbx1/-	129	-0.101817612	0.546349031	0.974718567	FALSE
				oTbx1/-	11	19.61154304	3.599706317	0.960571766	0.788805196	0.79537552	FALSE							Genotype	0.010709424	1	912	0.917599736							
			P12	oTbx1/+	30	13.16752619	3.364700074	0.59169817	5.78851E-07	3.96926E-07	TRUE	0.873593574	1	40	0.355570832	0.948188885	FALSE	PND	1.295840311	1	912	0.255273381	oTbx1/+	oTbx1/-	157	0.098811692	0.536091044	0.974718567	FALSE
				oTbx1/-	12	8.505399165	1.955013986	0.873135149	0.07162638	0.090475427	FALSE							Call type	39.53182246	11	912	0							
		step_down	P8	oTbx1/+	27	1.336688279	0.313639637	0.787106931	8.25136E-05	0.000220036	TRUE	0.149229077	1	36	0.701546982	0.958177206	FALSE	Genotype * PND	0.010709424	1	912	0.917599736	oTbx1/+	oTbx1/-	143.5	-0.02610708	0.899119881	0.974718567	FALSE
				oTbx1/-	11	1.573785836	0.594457163	0.750182569	0.002090028	0.00480056	TRUE							PND * Call type	2.796512537	11	912	0.001369316							
			P12	oTbx1/+	30	1.789236696	0.427419086	0.770336628	1.97568E-05	5.92704E-05	TRUE	2.489159983	1	40	0.122509658	0.586275644	FALSE	Call type * Genotype	1.365715759	11	912	0.183698653	oTbx1/+	oTbx1/-	176.5	-0.015036562	0.945286072	0.974718567	FALSE
				oTbx1/-	12	9.940976217	8.252655216	0.398949146	4.43981E-09	1.28617E-05	TRUE							Genotype * PND * Call type	0.796636894	11	912	0.643654777							
		short	P8	oTbx1/+	27	23.4473742	4.012502925	0.666321874	1.3633E-06	5.45318E-06	TRUE	0.002788636	1	36	0.958177206	0.958177206	FALSE						oTbx1/+	oTbx1/-	142	0.033939204	0.849168654	0.974718567	FALSE
				oTbx1/-	11	21.62772654	4.72041667	0.830675006	0.023827234	0.035740851	TRUE																		
			P12	oTbx1/+	30	23.24444672	2.727284442	0.958858669	0.289563179	0.317734805	FALSE	0.18394001	1	40	0.670309944	0.958177206	FALSE						oTbx1/+	oTbx1/-	174	-0.025776963	0.879981095	0.974718567	FALSE
				oTbx1/-	12	22.86873026	4.926486399	0.944190502	0.55418241	0.578277298	FALSE																		
		down_fm	P8	oTbx1/+	27	12.75630401	1.76961371	0.920927048	0.041565027	0.058680038	FALSE	4.472020458	1	36	0.041444496	0.497333956	FALSE						oTbx1/+	oTbx1/-	105	-0.227131596	0.168812806	0.974718567	FALSE
				oTbx1/-	11	21.16328548	4.9646338	0.884678006	0.11924167	0.146758978	FALSE																		
			P12	oTbx1/+	30	10.37298456	1.542082257	0.928056836	0.043611676	0.059699769	FALSE	0.954744089	1	40	0.334387409	0.948188885	FALSE						oTbx1/+	oTbx1/-	156.5	-0.100959772	0.536091044	0.974718567	FALSE
				oTbx1/-	12	13.01185528	3.11935555	0.931758583	0.399125427	0.425733789	FALSE																		
		up_fm	P8	oTbx1/+	27	13.05872025	1.780737559	0.92226398	0.044774827	0.059699769	FALSE	5.242724489	1	36	0.028006319	0.497333956	FALSE						oTbx1/+	oTbx1/-	84	0.336781333	0.038051176	0.45661411	FALSE
				oTbx1/-	11	6.202009894	1.944137346	0.855210662	0.04989852	0.064733215	FALSE																		
			P12	oTbx1/+	30	17.31720233	2.688345341	0.892989337	0.005683312	0.010103665	TRUE	0.013930753	1	40	0.9066357	0.958177206	FALSE						oTbx1/+	oTbx1/-	165.5	0.062294328	0.710773696	0.974718567	FALSE
				oTbx1/-	12	14.90117688	3.627231491	0.915457904	0.250485063	0.286268643	FALSE																		
		two_steps	P8	oTbx1/+	27	1.22818806	0.583431602	0.439589798	4.31981E-09	6.9117E-08	TRUE	0.367415161	1	36	0.548219905	0.958177206	FALSE						oTbx1/+	oTbx1/-	137.5	0.057435576	0.751191124	0.974718567	FALSE
				oTbx1/-	11	0.657609323	0.30689217	0.709964395	0.000628547	0.001436678	TRUE																		
			P12	oTbx1/+	30	0.376106479	0.140892041	0.56127727	2.53783E-08	2.03026E-07	TRUE	0.031791176	1	40	0.859386723	0.958177206	FALSE						oTbx1/+	oTbx1/-	158	-0.094515531	0.5544243	0.974718567	FALSE
				oTbx1/-	12	0.420168462	0.189210931	0.706350029	0.000970717	0.002117929	TRUE																		
		flat	P8	oTbx1/+	27	13.18848553	1.439854436	0.955547333	0.291256905	0.317734805	FALSE	0.064044466	1	36	0.801653642	0.958177206	FALSE						oTbx1/+	oTbx1/-	146.5	-0.010442832	0.974718567	0.974718567	FALSE
				oTbx1/-	11	13.58957049	2.677180408	0.971073091	0.897150159	0.897150159	FALSE																		
			P12	oTbx1/+	30	10.75966512	1.248052332	0.916827679	0.02220858	0.03438791	TRUE	0.005399042	1	40	0.941791831	0.958177206	FALSE						oTbx1/+	oTbx1/-	164	0.068738568	0.670224551	0.974718567	FALSE
				oTbx1/-	12	9.627772862	1.980798048	0.90607655	0.189971969	0.227966362	FALSE																		
		chevron	P8	oTbx1/+	27	11.99601344	2.252797934	0.893206934	0.009383468	0.015531257	TRUE	0.562260857	1	36	0.458223333	0.958177206	FALSE						oTbx1/+	oTbx1/-	120	0.148810356	0.372960146	0.974718567	FALSE
				oTbx1/-	11	8.606354313	3.605578163	0.701831937	0.000493834	0.001185202	TRUE																		
			P12	oTbx1/+	30	7.944668927	1.340965584	0.883133352	0.00332987	0.006147453	TRUE	1.121536249	1	40	0.72920259	0.958177206	FALSE						oTbx1/+	oTbx1/-	149	0.133180976	0.400911456	0.974718567	FALSE
				oTbx1/-	12	5.704118406	2.467406	0.713398993	0.001137725	0.002374382	TRUE																		
		complex	P8	oTbx1/+	27	1.359067236	0.357237975	0.747017026	1.88041E-05	5.92704E-05	TRUE	1.462744932	1	36	0.234376731	0.803577363	FALSE						oTbx1/+	oTbx1/-	110.5	0.198413808	0.276164508	0.974718567	FALSE
				oTbx1/-	11	0.594390633	0.456151172	0.472920179	7.17045E-07	4.30227E-06	TRUE																		
			P12	oTbx1/+	30	1.444040234	0.348567383	0.776604891	2.52732E-05	7.13597E-05	TRUE	0.463312445	1	40	0.500001335	0.958177206	FALSE						oTbx1/+	oTbx1/-	153	-0.115996334	0.465855659	0.974718567	FALSE
				oTbx1/-	12	2.213403834	0.819843897	0.755306244	0.003037653	0.005832293	TRUE																		
		step_up	P8	oTbx1/+	27	3.459543469	0.754843072	0.828750014	0.000449811	0.001136364	TRUE	2.986552004	1	36	0.092528851	0.586275644	FALSE						oTbx1/+	oTbx1/-	128	-0.107039028	0.525351863	0.974718567	FALSE
				oTbx1/-	11	6.030970473	2.091188206	0.792426586	0.007478565	0.012820397	TRUE																		
			P12	oTbx1/+	30	2.930012431	0.802546816	0.690653443	1.14398E-06	5.26814E-06	TRUE	0.337019135	1	40	0.564811412	0.958177206	FALSE						oTbx1/+	oTbx1/-	174	-0.025776963	0.879981095	0.974718567	FALSE
				oTbx1/-	12	2.522648912	0.867273801	0.814436078	0.013736397	0.021978235	TRUE																		
		mult_steps	P8	oTbx1/+	27	0.053993663	0.038446941	0.300173998	2.61841E-10	6.28418E-09	TRUE	0.119494266	1	36	0.731595596	0.958177206	FALSE						oTbx1/+	oTbx1/-	147	-0.007832124	0.974718567	0.974718567	FALSE
				oTbx1/-	11	0.031897927	0.031897927	0.344991028	2.2434E-08	2.03026E-07	TRUE																		
			P12	oTbx1/+	30	0.016955782	0.011903467	0.282110274	5.00231E-11	2.40111E-09	TRUE	0.025753757	1	40	0.873310885	0.958177206	FALSE						oTbx1/+	oTbx1/-	177	-0.012888482	0.945286072	0.974718567	FALSE
				oTbx1/-	12	0.020627063	0.020627063	0.326927483	1.20728E-06	5.26814E-06	TRUE																		
		rev_chevron	P8	oTbx1/+	27	1.081121699	0.315727719	0.719082952	7.22694E-06	2.47781E-05	TRUE	2.201621745	1	36	0.146568911	0.586275644	FALSE						oTbx1/+	oTbx1/-	107	0.216688764	0.189914135	0.974718567	FALSE
				oTbx1/-	11	0.31085605	0.230553927	0.488995969	1.11768E-06	5.26814E-06	TRUE																		
			P12	oTbx1/+	30	0.647154535	0.267451787	0.507237554	6.39587E-09	7.67505E-08	TRUE	2.868746639	1	40	0.098089764	0.586275644	FALSE						oTbx1/+	oTbx1/-	99	-0.347989002	0.023533765	0.45661411	FALSE
				oTbx1/-	12	1.929789318	0.609606267	0.845055282	0.031914786	0.046421506	TRUE																		

(Second table)

Measurement	Call type	PND	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MW Sample1	MW Sample2	MW Statistic	R (Z score)	MW p_value	MW p_adj(FDR BH)	MW Rejects(FDR 5%)
Proportion	Harmonics	P8	PdgfraCre;Tbx1/+	8	14.33386512	3.180619627	0.920521609	0.434110293	0.529761713	FALSE	1.84713931	2	24	0.179413127	0.51429221	FALSE	Corrected Model	7.678337945	71	612	0	PdgfraCre;Tbx1/+	WT;Tbx1/flox	29.5	0.065638129	0.878477078	1	FALSE
			WT;Tbx1/flox	8	13.69827654	5.213100979	0.825142512	0.052831697	0.105663394	FALSE							Genotype	0.567660277	2	612	0.567148336	PdgfraCre;Tbx1/+	WT;Tbx1/+	25	-0.359924234	0.1288402	0.7730412	FALSE
			WT;Tbx1/+	11	21.42494286	2.955550922	0.956472392	0.727051807	0.805349694	FALSE							PND	1.529321287	1	612	0.216688691	WT;Tbx1/flox	WT;Tbx1/+	32	-0.227320569	0.351062422	0.901457961	FALSE
		P12	PdgfraCre;Tbx1/+	9	19.77849016	10.32058624	0.614034689	0.000190012	0.000915012	TRUE	1.199438418	2	27	0.316917223	0.51429221	FALSE	Call type	42.61123303	11	612	0	PdgfraCre;Tbx1/+	WT;Tbx1/flox	25	0.32260178	0.190250926	0.901457961	FALSE
			WT;Tbx1/flox	9	5.644679288	1.918519292	0.860758774	0.097722904	0.171610953	FALSE							Genotype * PND	0.567660277	2	612	0.567148336	PdgfraCre;Tbx1/+	WT;Tbx1/+	47	-0.108556491	0.651052992	0.901457961	FALSE
			WT;Tbx1/+	12	13.85143838	2.762225592	0.888610048	0.113077876	0.1893397	FALSE							PND * Call type	0.791648519	11	612	0.648626623	WT;Tbx1/flox	WT;Tbx1/+	26.5	-0.42647193	0.058483312	0.701799748	FALSE
	step_down	P8	PdgfraCre;Tbx1/+	8	1.794530722	0.846007146	0.772387788	0.014392817	0.041451314	TRUE	1.151437041	2	24	0.333040024	0.51429221	FALSE	Call type * Genotype	1.940248357	22	612	0.006372406	PdgfraCre;Tbx1/+	WT;Tbx1/flox	29.5	0.065638129	0.878477078	1	FALSE
			WT;Tbx1/flox	8	1.290974418	0.534434464	0.816551909	0.0429211	0.093646035	FALSE							Genotype * PND * Call type	1.033284006	22	612	0.419829765	PdgfraCre;Tbx1/+	WT;Tbx1/+	38.5	0.104188594	0.71678442	0.955712559	FALSE
			WT;Tbx1/+	11	1.0369584	0.305974108	0.896482842	0.167368171	0.262860212	FALSE												WT;Tbx1/flox	WT;Tbx1/+	42.5	0.028415071	0.967849488	1	FALSE
		P12	PdgfraCre;Tbx1/+	9	0.512440844	0.392071531	0.528195976	1.37202E-05	0.000142397	TRUE	3.768584833	2	27	0.036019673	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	24	-0.343414798	0.161497326	0.901457961	FALSE
			WT;Tbx1/flox	9	2.584253777	0.986144531	0.808525695	0.025540655	0.063411281	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	20	-0.418717895	0.084483312	0.701799748	FALSE
			WT;Tbx1/+	12	2.150570773	0.650897914	0.881745505	0.092280528	0.16782624	FALSE												WT;Tbx1/flox	WT;Tbx1/+	53	0.01550807	0.97722231	1	FALSE
	short	P8	PdgfraCre;Tbx1/+	8	27.79609334	7.835750345	0.802567038	0.030493177	0.073183624	TRUE	1.027435303	2	24	0.373136602	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	31	-0.026255252	0.959129759	1	FALSE
			WT;Tbx1/flox	8	30.52798165	10.4520263	0.674664094	0.001160659	0.004915731	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	27.5	-0.312565783	0.206001429	0.901457961	FALSE
			WT;Tbx1/+	11	15.13513667	1.98602887	0.910698292	0.248577402	0.357951459	FALSE												WT;Tbx1/flox	WT;Tbx1/+	24	0.378867615	0.108650208	0.74285928	FALSE
		P12	PdgfraCre;Tbx1/+	9	29.41714216	6.77204175	0.954397752	0.73826333	0.805378178	FALSE	2.18520834	2	27	0.131943177	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	32	0.176910653	0.489428219	0.901457961	FALSE
			WT;Tbx1/flox	9	21.11850489	4.315377521	0.818225622	0.032903072	0.076420038	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	41	0.201604912	0.3824448	0.901457961	FALSE
			WT;Tbx1/+	12	20.18438151	3.177084685	0.968703162	0.896753634	0.949503848	FALSE												WT;Tbx1/flox	WT;Tbx1/+	47	0.108556491	0.651052972	0.901457961	FALSE
	down_fm	P8	PdgfraCre;Tbx1/+	9	9.902998402	1.798693708	0.823580506	0.478929706	0.574715647	FALSE	1.301310466	2	24	0.290697828	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	29	-0.078765753	0.974274778	1	FALSE
			WT;Tbx1/flox	8	12.95875965	4.301199919	0.91573224	0.396200038	0.509400048	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	30	-0.265207331	0.272287052	0.901457961	FALSE
			WT;Tbx1/+	14	14.68419488	2.820054979	0.931509541	0.426325541	0.529231706	FALSE												WT;Tbx1/flox	WT;Tbx1/+	37	-0.132603665	0.599878278	0.901457961	FALSE
		P12	PdgfraCre;Tbx1/+	9	15.1261745	3.538818708	0.937379276	0.554598449	0.633826799	FALSE	1.089462096	2	27	0.350732163	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	21.5	0.395447343	0.11349239	0.74285928	FALSE
			WT;Tbx1/flox	9	6.768936551	2.39904953	0.877183695	0.1466369	0.239951291	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	37.5	0.255883158	0.277317729	0.901457961	FALSE
			WT;Tbx1/+	12	9.511128113	1.791519021	0.945887631	0.577853788	0.650085512	FALSE												WT;Tbx1/flox	WT;Tbx1/+	38.5	-0.240375088	0.310026197	0.901457961	FALSE
	up_fm	P8	PdgfraCre;Tbx1/+	8	10.52722533	3.589529281	0.898985786	0.282937423	0.399441068	FALSE	0.222556444	2	24	0.802103358	0.875021845	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	22	-0.262552516	0.328205128	0.901457961	FALSE
			WT;Tbx1/flox	8	15.15934247	3.299813459	0.929283224	0.509574132	0.594583706	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	36	-0.151547046	0.544759334	0.901457961	FALSE
			WT;Tbx1/+	13	13.11026403	2.688376967	0.960336023	0.775876897	0.833778158	FALSE												WT;Tbx1/flox	WT;Tbx1/+	38.5	0.104188594	0.955712559	FALSE
		P12	PdgfraCre;Tbx1/+	9	10.7373311	3.075778662	0.912907898	0.336760896	0.449014528	FALSE	1.438522228	2	27	0.254889982	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	22	-0.385040834	0.11349239	0.74285928	FALSE
			WT;Tbx1/flox	9	24.26204578	6.725552181	0.910146573	0.316901753	0.430508042	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	35	-0.294653333	0.193005137	0.901457961	FALSE
			WT;Tbx1/+	12	17.04347316	3.399092861	0.941065847	0.512002636	0.594583706	FALSE												WT;Tbx1/flox	WT;Tbx1/+	41.5	0.193850877	0.422107304	0.901457961	FALSE
	two_steps	P8	PdgfraCre;Tbx1/+	8	0.102459016	0.102459016	0.418398447	1.04723E-06	2.51336E-05	TRUE	1.00441335	2	24	0.381141088	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	22.5	-0.24942489	0.382284382	0.901457961	FALSE
			WT;Tbx1/flox	8	2.049948931	1.891510087	0.486985371	7.19813E-06	0.000103653	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	16.5	-0.520942971	0.0259321	0.701799748	FALSE
			WT;Tbx1/+	11	1.33652855	0.427577641	0.831222503	0.024225071	0.063411281	FALSE												WT;Tbx1/flox	WT;Tbx1/+	32.5	-0.217848879	0.391635641	0.901457961	FALSE
		P12	PdgfraCre;Tbx1/+	9	0.074906367	0.074906367	0.389841298	3.21749E-07	1.1583E-05	TRUE	1.013541908	2	27	0.376326518	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	26	-0.301788762	0.222418758	0.901457961	FALSE
			WT;Tbx1/flox	9	0.455038711	0.205119656	0.782599951	0.009527702	0.038708165	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	38	-0.248129123	0.277129729	0.901457961	FALSE
			WT;Tbx1/+	12	0.542807388	0.309332544	0.587077317	8.38703E-05	0.00070963	TRUE												WT;Tbx1/flox	WT;Tbx1/+	50.5	0.054278246	0.8620692	FALSE
	flat	P8	PdgfraCre;Tbx1/+	8	17.93017709	2.717057881	0.917100523	0.406781173	0.513828851	FALSE	0.524216516	2	24	0.598643331	0.75618105	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	13	-0.49884978	0.701799748	FALSE
			WT;Tbx1/flox	8	8.633543264	2.89229413	0.820293101	0.04699594	0.099520814	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	22.5	0.407282686	0.090841735	0.74285928	FALSE
			WT;Tbx1/+	13	13.05266786	1.437946293	0.927701172	0.388068488	0.50801693	FALSE												WT;Tbx1/flox	WT;Tbx1/+	23	-0.397810999	0.090841735	0.74285928	FALSE
		P12	PdgfraCre;Tbx1/+	9	10.49163426	2.3312487	0.893212814	0.215226195	0.322839292	FALSE	0.040346651	2	27	0.960514231	0.960514231	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	29	0.239349707	0.340106952	0.901457961	FALSE
			WT;Tbx1/flox	9	7.38091094	2.106347882	0.89566068	0.22790882	0.33488643	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	39	-0.232621053	0.310204358	0.901457961	FALSE
			WT;Tbx1/+	12	13.4947539	1.915033856	0.923022175	0.311949399	0.430508042	FALSE												WT;Tbx1/flox	WT;Tbx1/+	26.5	-0.42647193	0.058483312	0.701799748	FALSE
	chevron	P8	PdgfraCre;Tbx1/+	8	11.11468622	3.975984082	0.839208898	0.073925068	0.143854186	FALSE	0.991480484	2	24	0.385719158	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	29	0.078765753	0.798445998	0.974374778	FALSE
			WT;Tbx1/flox	8	8.28992559	2.59542819	0.854717441	0.106307995	0.182242278	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	36	-0.151547046	0.544759334	0.901457961	FALSE
			WT;Tbx1/+	11	15.33231531	4.329267764	0.896602764	0.167938469	0.262860212	FALSE												WT;Tbx1/flox	WT;Tbx1/+	32	-0.227320569	0.351062422	0.901457961	FALSE
		P12	PdgfraCre;Tbx1/+	9	8.44801746	3.058254192	0.834102105	0.049637303	0.102111023	FALSE	1.856606996	2	27	0.175598067	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	37	0.072845563	0.796147414	0.974374778	FALSE
			WT;Tbx1/flox	9	6.795574321	1.76969935	0.807793859	0.025055817	0.063411281	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	52	-0.03101614	0.917007451	1	FALSE
			WT;Tbx1/+	12	8.398978482	2.216508759	0.903793823	0.177537843	0.271972866	FALSE												WT;Tbx1/flox	WT;Tbx1/+	48.5	-0.085294386	0.754370088	0.974374778	FALSE
	complex	P8	PdgfraCre;Tbx1/+	9	0.976069158	0.63900965	0.568475515	6.77437E-05	0.000609693	TRUE	0.3998853	2	24	0.674781143	0.771178449	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	26	-0.157531509	0.573737374	FALSE
			WT;Tbx1/flox	8	1.68881032	0.777065407	0.807628446	0.034526318	0.077684215	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	35	-0.170490427	0.49202191	0.901457961	FALSE
			WT;Tbx1/+	11	1.397798141	0.529558045	0.635967329	0.00450508	0.017071912	TRUE												WT;Tbx1/flox	WT;Tbx1/+	44	0	1	FALSE
		P12	PdgfraCre;Tbx1/+	9	1.35985243	0.834393015	0.639968592	0.000281573	0.00153755	TRUE	0.065201783	2	27	0.937025452	0.960514231	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	34	-0.135284617	0.604812834	0.901457961	FALSE
			WT;Tbx1/flox	9	1.639512949	0.546565381	0.736084074	0.003737285	0.014949138	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	34	-0.155080702	0.507876025	0.901457961	FALSE
			WT;Tbx1/+	12	1.360576551	0.50435214	0.777657114	0.005278143	0.019001316	TRUE												WT;Tbx1/flox	WT;Tbx1/+	46	0.124064561	0.60159902	0.901457961	FALSE
	step_up	P8	PdgfraCre;Tbx1/+	8	4.58948929	1.934601202	0.843751974	0.082294537	0.155926491	FALSE	2.38559384	2	24	0.113511907	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	31	-0.026255252	0.959129759	FALSE
			WT;Tbx1/flox	8	3.489325269	1.336233123	0.849076133	0.093236803	0.167826246	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	43.5	-0.00947169	0.557773523	0.777989468	0.974374778	FALSE
			WT;Tbx1/+	11	2.6161052	0.811090133	0.802232956	0.01006582	0.031589754	TRUE												WT;Tbx1/flox	WT;Tbx1/+	40	0.05773523	0.777989468	0.974374778	FALSE
		P12	PdgfraCre;Tbx1/+	9	3.389790988	1.520322048	0.773329537	0.010091172	0.031589754	TRUE	1.256794019	2	27	0.300686319	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	28	-0.260162729	0.297326203	0.901457961	FALSE
			WT;Tbx1/flox	9	3.84752633	0.532565223	0.603414342	0.000104925	0.00075546	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	46	-0.124064561	0.60159902	0.901457961	FALSE
			WT;Tbx1/+	12	4.156623362	1.54243506	0.656505111	0.000331387	0.001590656	TRUE												WT;Tbx1/flox	WT;Tbx1/+	46	-0.465242105	0.033865206	0.701799748	FALSE
	mult_steps	P8	PdgfraCre;Tbx1/+	8			1			FALSE	1.488233273	2	24	0.245874864	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	32				FALSE
			WT;Tbx1/flox	8			1			FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	36	-0.151547046	0.544759334	0.901457961	FALSE
			WT;Tbx1/+	11	0.132529899	0.091610889	0.511703575	2.09968E-06	3.77942E-05	TRUE												WT;Tbx1/flox	WT;Tbx1/+	36	-0.151547046	0.544759334	0.901457961	FALSE
		P12	PdgfraCre;Tbx1/+	9			1			FALSE	1.582954373	2	27	0.223840796	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	40.5				FALSE
			WT;Tbx1/flox	9			1			FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	45	-0.139572632	0.553791719	0.901457961	FALSE
			WT;Tbx1/+	12	0.042389454	0.028911445	0.485515454	1.38441E-05	0.000142397	TRUE												WT;Tbx1/flox	WT;Tbx1/+	45	-0.139572632	0.553791719	0.901457961	FALSE
	rev_chevron	P8	PdgfraCre;Tbx1/+	8	0.932406314	0.485217797	0.736969104	0.005855571	0.020076244	TRUE	0.998042961	2	24	0.383388724	0.51429221	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	27	-0.131276258	0.645376845	0.901457961	FALSE
			WT;Tbx1/flox	8	1.698111905	0.810075328	0.792567662	0.02382121	0.063411281	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	44	0	1	FALSE
			WT;Tbx1/+	11	0.740558193	0.375948876	0.684843395	0.000298968	0.00153755	TRUE												WT;Tbx1/flox	WT;Tbx1/+	37	-0.132603665	0.599878278	0.901457961	FALSE
		P12	PdgfraCre;Tbx1/+	9	0.624219725	0.624219725	0.389841298	3.21749E-07	1.1583E-05	TRUE	0.46841942	2	27	0.630983165	0.757179799	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	33	-0.156079635	0.564376845	0.901457961	FALSE
			WT;Tbx1/flox	9	0.29356793	0.173122527	0.653259574	0.000403018	0.00181358	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	40	-0.217112982	0.345102575	0.901457961	FALSE
			WT;Tbx1/+	12	0.929545596	0.476189549	0.644326098	0.000257894	0.00153755	TRUE												WT;Tbx1/flox	WT;Tbx1/+	44	-0.108556491	0.651052972	0.901457961	FALSE

Figure 4C — Duration (by call type)

Figure	Measurement	Call type	PND	Genotype	# Samples	Average	s.e.m.	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MW Sample1	MW Sample2	MW Statistic	R (Z score)	MW p_value	MW p_adj(FDR BH)	MW Rejects(FDR 5%)
Figure 4C	Duration	Harmonics	P8	oTbx1/+	23	47.92295829	2.699256034	0.979313314	0.8941468	0.989708006	FALSE	0.000929168	1	31	0.975877848	0.985623162	FALSE	Corrected Model	20.26123774	47	560	0	oTbx1/+	oTbx1/-	129	-0.101817612	0.602683656	0.911355311	FALSE
				oTbx1/-	10	44.69960544	4.401566176	0.974668384	0.930436969	0.989708006	FALSE							Genotype	0.043473793	1	560	0.834911507							
			P12	oTbx1/+	22	54.74005891	3.852574293	0.918354988	0.070288621	0.311429934	FALSE	1.537197377	1	29	0.224974682	0.490853852	FALSE	PND	0.165222066	1	560	0.684548787	oTbx1/+	oTbx1/-	97	-0.356581323	0.948931589	0.990189484	FALSE
				oTbx1/-	9	52.60944711	3.167958994	0.916101158	0.360939831	0.770815088	FALSE							Call type	60.43094494	11	560								
		step_down	P8	oTbx1/+	18	27.09201646	3.053456892	0.865561366	0.510047655	0.989708006	FALSE	0.2059045	1	23	0.654249229	0.962256678	FALSE	Genotype * PND	0.105188211	1	560	0.745811343	oTbx1/+	oTbx1/-	86	-0.529037945	0.710031204	0.911355311	FALSE
				oTbx1/-	7	20.10412698	3.302056133	0.904883921	0.361587048	0.770815088	FALSE							PND * Call type	1.398862882	11	560	0.169112377							
			P12	oTbx1/+	15	23.33660106	1.7963783	0.982755959	0.98487395	0.989708006	FALSE	2.068498506	1	20	0.165834425	0.429742989	FALSE	Call type * Genotype	3.360831125	11	560	0.000163775	oTbx1/+	oTbx1/-	68	-0.481169978	0.298245614	0.911355311	FALSE
				oTbx1/-	7	28.09701587	8.430810214	0.715816438	0.005502878	0.077620809	FALSE							Genotype * PND * Call type	1.038937083	11	560	0.410076829							
		short	P8	oTbx1/+	27	7.74224882	0.312538479	0.945187688	0.163532123	0.528334551	FALSE	0.171090279	1	36	0.68159848	0.962256678	FALSE						oTbx1/+	oTbx1/-	132	0.086153364	0.611651952	0.911355311	FALSE
				oTbx1/-	11	7.774337147	0.846332788	0.846332788	0.038221501	0.229332896	FALSE																		
			P12	oTbx1/+	26	7.072236643	0.334744164	0.87922591	0.005544343	0.077620809	FALSE	0.00038636	1	34	0.984432657	0.985623162	FALSE						oTbx1/+	oTbx1/-	133	-0.201919544	0.930816808	0.990189484	FALSE
				oTbx1/-	10	7.017617959	0.485436046	0.96597141	0.851189494	0.989708006	FALSE																		
		down_fm	P8	oTbx1/+	24	30.66279906	1.957850728	0.966806293	0.589142263	0.907498758	FALSE	3.431740525	1	33	0.072916956	0.35000139	FALSE						oTbx1/+	oTbx1/-	137	-0.060046284	0.874932414	0.990189484	FALSE
				oTbx1/-	11	30.69520938	1.801882739	0.955081701	0.709225118	0.930857968	FALSE																		
			P12	oTbx1/+	25	35.67096042	2.959919864	0.969138086	0.62329185	0.907498758	FALSE	2.160735587	1	32	0.151341379	0.429742989	FALSE						oTbx1/+	oTbx1/-	101	-0.339396681	0.673085541	0.911355311	FALSE
				oTbx1/-	9	37.35198732	3.365400362	0.94418776	0.626606286	0.907498758	FALSE																		
		up_fm	P8	oTbx1/+	23	19.62332101	0.925405003	0.955981197	0.387208449	0.770815088	FALSE	5.35177982	1	30	0.027773942	0.221903537	FALSE						oTbx1/+	oTbx1/-	91	-0.063333672	0.621111142	0.911355311	FALSE
				oTbx1/-	9	23.60520165	3.653717179	0.921376705	0.403760284	0.770815088	FALSE																		
			P12	oTbx1/+	24	19.72581213	1.368821557	0.947432995	0.238222212	0.667022192	FALSE	0.058377287	1	32	0.810620958	0.985623162	FALSE						oTbx1/+	oTbx1/-	165	-0.064442408	0.092918255	0.557509532	FALSE
				oTbx1/-	10	15.11884466	1.853801309	0.986200121	0.987008006	0.989708006	FALSE																		
		two_steps	P8	oTbx1/+	12	56.71407407	8.656626098	0.946003079	0.579481781	0.907498758	FALSE	2.000905298	1	14	0.179059579	0.429742989	FALSE						oTbx1/+	oTbx1/-	28	-0.629180629	0.683516484	0.911355311	FALSE
				oTbx1/-	4	50.34666667	5.410990188	0.792466283	0.088181523	0.341274908	FALSE																		
			P12	oTbx1/+	10	58.62755556	4.194056899	0.951295912	0.683848679	0.926504662	FALSE	0.000336509	1	14	0.985623162	0.985623162	FALSE						oTbx1/+	oTbx1/-	53	-0.545612386	0.010989011	0.263736264	FALSE
				oTbx1/-	6	34.70222222	5.58446381	0.918257356	0.492899448	0.840512896	FALSE																		
		flat	P8	oTbx1/+	26	19.78944829	0.81484035	0.984566271	0.953157961	0.989708006	FALSE	1.174282732	1	34	0.286148033	0.572296066	FALSE						oTbx1/+	oTbx1/-	113	-0.185360268	0.565958548	0.911355311	FALSE
				oTbx1/-	10	21.47446293	2.726545128	0.82753104	0.031245172	0.218716204	FALSE																		
			P12	oTbx1/+	24	22.07510842	0.90491284	0.950715065	0.280761451	0.736998808	FALSE	2.759012943	1	31	0.106790325	0.4271613	FALSE						oTbx1/+	oTbx1/-	129	-0.219104186	0.414164975	0.911355311	FALSE
				oTbx1/-	9	21.11847067	0.935953023	0.891232073	0.205424145	0.616272435	FALSE																		
		chevron	P8	oTbx1/+	21	36.64371988	2.271811058	0.919187725	0.0835924	0.341274908	FALSE	0.003024766	1	27	0.956545111	0.985623162	FALSE						oTbx1/+	oTbx1/-	81	-0.352445581	0.904925626	0.990189484	FALSE
				oTbx1/-	8	37.73476406	3.772893681	0.954048574	0.751886845	0.956946893	FALSE																		
			P12	oTbx1/+	20	41.88753169	2.667991417	0.975014746	0.855094075	0.989708006	FALSE	0.204884413	1	25	0.654710508	0.962256678	FALSE						oTbx1/+	oTbx1/-	87	-0.395542928	0.369771292	0.911355311	FALSE
				oTbx1/-	7	37.00915927	3.607004024	0.965664983	0.865602672	0.989708006	FALSE																		
		complex	P8	oTbx1/+	12	53.79841981	3.593523806	0.928657532	0.366100401	0.770815088	FALSE	0.004785745	1	12	0.945986497	0.985623162	FALSE						oTbx1/+	oTbx1/-	16	-0.691837621	0.549450549	0.911355311	FALSE
				oTbx1/-	2	47.81714286	8.92952381																						
			P12	oTbx1/+	15	62.12476912	6.394717172	0.939049602	0.370569885	0.770815088	FALSE	0.257959301	1	21	0.616881976	0.962256678	FALSE						oTbx1/+	oTbx1/-	68	-0.481169978	0.635523359	0.911355311	FALSE
				oTbx1/-	8	53.32536201	7.106570063	0.86379683	0.130967751	0.458387129	FALSE																		
		step_up	P8	oTbx1/+	21	39.52578272	2.436556474	0.955946565	0.43852523	0.800785202	FALSE	2.443787387	1	28	0.129224079	0.429742989	FALSE						oTbx1/+	oTbx1/-	52	-0.503866645	0.056277595	0.557509532	FALSE
				oTbx1/-	9	50.31356252	6.763535213	0.971306562	0.905666947	0.989708006	FALSE																		
			P12	oTbx1/+	19	36.25782364	3.631192924	0.956213474	0.500305295	0.840512896	FALSE	0.000525987	1	25	0.981884673	0.985623162	FALSE						oTbx1/+	oTbx1/-	42	-0.592870152	0.074893866	0.557509532	FALSE
				oTbx1/-	8	54.54926984	8.836116188	0.641596675	0.000484111	0.020332659	TRUE																		
		mult_steps	P8	oTbx1/+	2	45.0133333	37.3333333					1.84044E+31	1	1	1.48395E-16	1.78074E-15	TRUE						oTbx1/+	oTbx1/-	1	-0.770158862	1	1	FALSE
				oTbx1/-	1	48.21333333																							
			P12	oTbx1/+	2	13.01333333	5.76					2.80384E+31	1	1	1.20227E-16	1.78074E-15	TRUE						oTbx1/+	oTbx1/-	0	-0.773308893	0.666666667	0.911355311	FALSE
				oTbx1/-	1	52.90666667																							
		rev_chevron	P8	oTbx1/+	12	42.56651852	2.153044286	0.841997564	0.029282464	0.218716204	FALSE	4.749155614	1	12	0.049960362	0.299762171	FALSE						oTbx1/+	oTbx1/-	13	-0.707501869	0.923076923	0.990189484	FALSE
				oTbx1/-	2	38.4	14.50666667																						
			P12	oTbx1/+	9	47.50971016	5.892589066	0.94786489	0.666441116	0.926504662	FALSE	0.537574475	1	15	0.474743246	0.87644907	FALSE						oTbx1/+	oTbx1/-	52	-0.549908546	0.138790621	0.666194981	FALSE
				oTbx1/-	8	37.21680672	5.066803973	0.837518036	0.0710207	0.311429934	FALSE																		

Figure 4C — Duration (All)

Figure	Measurement	Call type	PND	Genotype	# of Samples	Average	s.e.m.	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	2-way ANOVA Source	Correction	Type III SS	df	Mean Square	F	Sig.	t-test Sample1	t-test Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 4C	Duration	All	P8	oTbx1/+	27	260.2677275	23.10002451	0.966685951	0.517069697	0.689426263	FALSE	0.024977278	1	36	0.875306878	0.875306878	FALSE	Within subjects							oTbx1/+	oTbx1/-	0.529251423	36	0.189310895	0.599879665	0.770649486	FALSE
				oTbx1/-	11	237.7105327	35.22954873	0.985315174	0.988519132	0.988519132	FALSE							PND	Sphericity Assumed	11487.37597	1	11487.37597	0.746726852	0.393039589							
			P12	oTbx1/+	27	266.3561112	29.14567445	0.965607464	0.490966678	0.689426263	FALSE	0.933710604	1	34	0.34304749	0.68069498	FALSE	PND * Genotype	Sphericity Assumed	5543.314971	1	5543.314971	0.361737769	0.551780088		oTbx1/+	oTbx1/-	-0.293732063	36	-0.105066661	0.770649486	0.770649486	FALSE
				oTbx1/-	11	281.6978642	39.45653524	0.85749799	0.053434357	0.213737428	FALSE							Error(PND)	Sphericity Assumed	490372.0159	32	15324.1255									
																		Between Subjects													
																		Intercept		3819990.943	1	3819990.943	166.4686802	3.16998E-14							
																		Genotype		5.440297392	1	5.440297392	0.000237079	0.987810766							
																		Error		734310.5625	32	22947.20508									

Duration — PdgfraCre;Tbx1 (by call type)

Measurement	Call type	PND	Genotype	# of samples	Average	s.e.m.	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	GLMM Source	F	df1	df2	Sig.	t-test Sample1	t-test Sample2	Statistic	dof	Cohen's d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Duration	Harmonics	P8	PdgfraCre;Tbx1/+	7	40.90886772	4.558427742	0.938468978	0.624970931	0.989653678	FALSE	0.002206996	2	20	0.997795681	0.997795681	FALSE	Corrected Model	11.02882805	67	364	0	PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.282121004	10	-0.165193145	0.783603353	0.968298081	FALSE
			WT;Tbx1/flox	5	42.93581128	5.625100406	0.880300822	0.310676402	0.861420933	FALSE							Genotype	5.347190073	2	364	0.706904756	PdgfraCre;Tbx1/+	WT;Tbx1/+	-2.489154851	16	-1.203490884	0.024196155	0.968298081	FALSE
			WT;Tbx1/+	11	54.65335546	3.322269435	0.906698369	0.2227774	0.745742576	FALSE							PND	1.596530724	1	364	0.207204201	WT;Tbx1/flox	WT;Tbx1/+	-1.891444267	14	-1.020169172	0.079436704	0.968298081	FALSE
		P12	PdgfraCre;Tbx1/+	6	47.11113757	5.674386478	0.953771049	0.763849974	0.989653678	FALSE	2.463344678	2	19	0.111885654	0.830248368	FALSE	Call type	52.12777172	11	364	0	PdgfraCre;Tbx1/+	WT;Tbx1/flox	-1.457113075	11	-0.810662878	0.173032593	0.968298081	FALSE
			WT;Tbx1/flox	6	64.71990508	11.31009545	0.964217505	0.851580079	0.989653678	FALSE							PND * Genotype	5.022025307	2	364	0.007055228	PdgfraCre;Tbx1/+	WT;Tbx1/+	-1.144156066	14	-0.576600459	0.271744981	0.968298081	FALSE
			WT;Tbx1/+	9	54.02024841	3.022231296	0.965456458	0.853254002	0.989653678	FALSE							PND * Call type	1.821393368	11	364	0.048996774	WT;Tbx1/flox	WT;Tbx1/+	0.179193122	13	0.57539704	0.294768343	0.968298081	FALSE
		step_down	P8	PdgfraCre;Tbx1/+	5	26.92266667	3.372016607	0.877170705	0.296686073	0.861420933	FALSE	0.430873811	2	15	0.657736097	0.996273433	FALSE	Call type * Genotype	0.536675492	20	364	0.950317556	PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.570697045	8	0.360946503	0.583874953	0.968298081	FALSE
				WT;Tbx1/flox	5	23.83644444	4.227757401	0.930576706	0.60030857	0.989653678	FALSE							PND * Genotype * Call type	0.998274051	20	364	0.463368195	PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.273985489	11	-0.156195976	0.789172445	0.968298081	FALSE
				WT;Tbx1/+	8	29.23259259	6.236899259	0.834931578	0.066786083	0.493875425	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.623418251	11	-0.353450384	0.545719109	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	2	27.97511111	1.379555556	0.93123362	0.604806917	0.989653678	FALSE	1.403826013	2	12	0.283250233	0.890215017	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.118223257	5	0.098912673	0.910493384	0.968298081	FALSE
				WT;Tbx1/flox	5	27.04863492	4.662940802	0.942444012	0.635276389	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.142292666	8	0.11249223	0.89036177	0.968298081	FALSE
				WT;Tbx1/+	8	27.35695268	2.060533643	0.943233087	0.560789013	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	0.069161795	11	0.039420029	0.94610209	0.968298081	FALSE
		short	P8	PdgfraCre;Tbx1/+	8	7.91781593	0.530604626	0.831355391	0.061324236	0.493875425	FALSE	0.02965329	2	24	0.970817565	0.997795681	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.11190297	14	0.055951485	0.912488936	0.968298081	FALSE
				WT;Tbx1/flox	8	7.81333203	0.768281296	0.655720075	0.00070145	0.042953152	TRUE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.533322905	17	0.247813921	0.600713923	0.968298081	FALSE
				WT;Tbx1/+	11	7.56286763	0.416805441	0.985154377	0.980045907	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	0.307715843	17	0.142983302	0.762036619	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	7	7.224727273	0.302944896	0.892752792	0.248184669	0.756963239	FALSE	0.62103357	2	23	0.54615906	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.681216815	13	0.352563197	0.507688111	0.968298081	FALSE
				WT;Tbx1/flox	7	6.733424679	0.691876485	0.83817731	0.095540452	0.5827976	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.059016841	17	0.027422777	0.953626898	0.968298081	FALSE
				WT;Tbx1/+	11	7.176941981	0.650036053	0.89527921	0.161741928	0.684425775	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.449666572	16	-0.21741099	0.658982876	0.968298081	FALSE
		down_fm	P8	PdgfraCre;Tbx1/+	7	31.22903484	2.618699941	0.966327518	0.870816941	0.989653678	FALSE	2.420919834	2	21	0.113215687	0.830248368	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	1.329497727	11	0.739664253	0.210596869	0.968298081	FALSE
				WT;Tbx1/flox	5	24.00876305	5.03714095	0.858242658	0.183218167	0.698519264	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.718770319	16	-0.347085833	0.483188151	0.968298081	FALSE
				WT;Tbx1/+	11	33.93194139	2.490502409	0.977074081	0.947711079	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-1.993865557	15	-1.011925509	0.064675183	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	8	35.27615686	5.281399559	0.962996667	0.838114522	0.989653678	FALSE	0.086686621	2	22	0.917276034	0.997795681	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.526963161	12	-0.284592634	0.607823392	0.968298081	FALSE
				WT;Tbx1/flox	6	40.01672804	7.675950561	0.936991462	0.63696165	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.25976169	17	0.120700916	0.79816736	0.968298081	FALSE
				WT;Tbx1/+	11	33.38767159	4.005392441	0.958638001	0.754588316	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	0.823481277	15	0.418948552	0.422203461	0.968298081	FALSE
		up_fm	P8	PdgfraCre;Tbx1/+	6	22.08635809	1.767895755	0.979031784	0.94664488	0.989653678	FALSE	0.803986224	2	20	0.46148733	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	2.090998344	11	1.163324085	0.060536036	0.968298081	FALSE
				WT;Tbx1/flox	7	18.68497536	1.231492248	0.778875725	0.025161975	0.383720116	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	1.06895429	14	0.55200622	0.303175501	0.968298081	FALSE
				WT;Tbx1/+	10	19.50234072	1.538468802	0.88695579	0.156663332	0.684425775	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.858608983	15	-0.423127126	0.404068462	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	7	18.25207174	1.882487619	0.811423279	0.0325187	0.405419699	FALSE	0.393935778	2	21	0.679277341	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.362025824	12	-0.195310942	0.723627738	0.968298081	FALSE
				WT;Tbx1/flox	7	19.37050535	2.449587654	0.952865475	0.755690423	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.790464576	15	-0.389545196	0.441575344	0.968298081	FALSE
				WT;Tbx1/+	10	21.00614516	2.58398818	0.900295495	0.220738485	0.745742576	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.440145385	15	-0.216906014	0.666107981	0.968298081	FALSE
		two_steps	P8	PdgfraCre;Tbx1/+	1	56.74666667						0.756333907	2	9	0.497030518	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.596519781	2		0.6114663	0.968298081	FALSE
				WT;Tbx1/flox	3	39.37185185	14.56837035	0.765693459	0.034986854	0.426839615	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.18779369	7		0.856366123	0.968298081	FALSE
				WT;Tbx1/+	8	63.21333333	11.47831855	0.944422712	0.655048847	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-1.135840723	9	-0.768967804	0.285363816	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	1	59.59111111						1.718803649	2	7	0.247024372	0.890215017	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.365685768	3		0.738863479	0.968298081	FALSE
				WT;Tbx1/flox	4	69.89333333	5.261383104	0.948534927	0.706977093	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.303654689	4		0.776530465	0.968298081	FALSE
				WT;Tbx1/+	5	54.22222222	7.218198127	0.955559829	0.776776713	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	1.028937284	7	0.690232114	0.337744317	0.968298081	FALSE
		flat	P8	PdgfraCre;Tbx1/+	9	18.38285418	2.161523708	0.938127232	0.552725901	0.989653678	FALSE	1.88909965	2	23	0.173933952	0.890215017	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.259433573	13	-0.12420963	0.799363748	0.968298081	FALSE
				WT;Tbx1/flox	7	20.03418318	1.026599508	0.918007443	0.45407321	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.247055629	17	-0.114796915	0.807840052	0.968298081	FALSE
				WT;Tbx1/+	11	19.07941271	1.060503268	0.930035527	0.411213037	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	0.066877881	16	0.032335019	0.947507498	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	7	22.06600313	2.099649318	0.855242162	0.137169031	0.684425775	FALSE	0.661867473	2	21	0.526320871	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.466876518	11	-0.259746116	0.649699228	0.968298081	FALSE
				WT;Tbx1/flox	6	23.32211358	1.547504437	0.833187707	0.114362185	0.6341903	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.288188199	16	0.139337201	0.776902466	0.968298081	FALSE
				WT;Tbx1/+	11	21.40033624	1.281598243	0.987590136	0.987590136	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	0.922891212	15	0.468385027	0.370671424	0.968298081	FALSE
		chevron	P8	PdgfraCre;Tbx1/+	6	36.00005604	4.652989959	0.795458709	0.053475915	0.493875425	FALSE	0.059069292	2	18	0.942823397	0.997795681	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.549602398	10	0.317313092	0.594652508	0.968298081	FALSE
				WT;Tbx1/flox	6	32.45211329	4.474684324	0.990198852	0.989653678	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.715913679	13	-0.377319639	0.486706	0.968298081	FALSE
				WT;Tbx1/+	9	39.86723351	3.153606108	0.913333661	0.33990915	0.901498181	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-1.397947944	13	-0.736783259	0.185255454	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	5	35.15140741	5.45098299	0.794380663	0.072866866	0.493875425	FALSE	0.724169595	2	17	0.499090145	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-1.458624868	9	-0.88324122	0.178666611	0.968298081	FALSE
				WT;Tbx1/flox	6	43.83518788	3.031502235	0.981537083	0.958938137	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-1.236549181	12	-0.639009947	0.239909947	0.968298081	FALSE
				WT;Tbx1/+	9	44.33138549	4.612113275	0.930351025	0.484695498	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.079848715	13	-0.042083968	0.937573691	0.968298081	FALSE
		complex	P8	PdgfraCre;Tbx1/+	2	52.06552381	5.47352381	0.983042133	0.919621758	0.989653678	FALSE	4.243778405	2	9	0.050328058	0.830248368	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.289710054	4	0.250896786	0.786434528	0.968298081	FALSE
				WT;Tbx1/flox	4	50.53333333	2.727266981	0.934042153	0.614480606	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.346516593	6	-0.281819178	0.741754886	0.968298081	FALSE
				WT;Tbx1/+	6	56.55277613	6.97908285	0.944235993	0.693480606	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.669843422	8	-0.432382007	0.52181452	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	5	56.86289128	15.15889488	0.98798584	0.790240544	0.989653678	FALSE	0.007045782	2	12	0.992983086	0.997795681	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.587160054	6	-0.428801075	0.578253405	0.968298081	FALSE
				WT;Tbx1/flox	4	68.77497258	12.75070208	0.964095962	0.836176977	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.160572581	8	-0.110805608	0.876410673	0.968298081	FALSE
				WT;Tbx1/+	5	59.62971429	9.270122584	0.944156119	0.676391711	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	0.349144991	10	0.564242133	0.739847339	0.968298081	FALSE
		step_up	P8	PdgfraCre;Tbx1/+	5	40.9686811	3.783911882	0.84138729	0.168793223	0.684425775	FALSE	1.646688998	2	18	0.220420241	0.890215017	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.440276927	9	-0.266600919	0.670120851	0.968298081	FALSE
				WT;Tbx1/flox	6	43.3817284	3.877075903	0.690721619	0.00492818	0.150209498	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.685188585	13	0.347288934	0.505259557	0.968298081	FALSE
				WT;Tbx1/+	10	36.49076612	4.16921906	0.943592325	0.593632891	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	1.112199041	14	0.574337115	0.28478682	0.968298081	FALSE
			P12	PdgfraCre;Tbx1/+	5	32.81066667	8.360663706	0.944022431	0.694498426	0.989653678	FALSE	0.168774818	2	16	0.8461834	0.997795681	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.323406166	7	-0.21708165	0.755689769	0.968298081	FALSE
				WT;Tbx1/flox	4	28.91555556	8.444243272	0.92068423	0.54709467	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.949195702	13	-0.593958838	0.359836407	0.968298081	FALSE
				WT;Tbx1/+	10	40.91830937	4.426603923	0.969189671	0.883231963	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-1.373626201	12	-0.812432792	0.194785619	0.968298081	FALSE
		mult_steps	P8	PdgfraCre;Tbx1/+	0																								
				WT;Tbx1/flox	0																								
				WT;Tbx1/+	2	45.01333333	37.33333333																						
			P12	PdgfraCre;Tbx1/+	0																								
				WT;Tbx1/flox	0																								
				WT;Tbx1/+	2	13.01333333	5.76																						
		rev_chevron	P8	PdgfraCre;Tbx1/+	3	43.94666667	3.205526093	0.998031496	0.91523587	0.989653678	FALSE	0.132365162	2	9	0.877695297	0.997795681	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	0.501057483	5	0.382688974	0.637605545	0.968298081	FALSE
				WT;Tbx1/flox	4	41.984	2.412836781	0.991194337	0.963506274	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.252803413	6	0.184621509	0.808858345	0.968298081	FALSE
				WT;Tbx1/+	5	42.20444444	4.861369009	0.690556819	0.007608686	0.154709958	FALSE												WT;Tbx1/flox	WT;Tbx1/+	-0.037327957	7	-0.025040355	0.971265737	0.971265737	FALSE
			P12	PdgfraCre;Tbx1/+	1	43.6224						0.468689127	2	6	0.646945397	0.996273433	FALSE						PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.618955639	3		0.599052199	0.968298081	FALSE
				WT;Tbx1/flox	3	58.64296296	12.13379604	0.982868766	0.749306217	0.989653678	FALSE												PdgfraCre;Tbx1/+	WT;Tbx1/+	0.113246424	4		0.915291352	0.968298081	FALSE
				WT;Tbx1/+	5	41.60722051	7.264631898	0.944978233	0.701332904	0.989653678	FALSE												WT;Tbx1/flox	WT;Tbx1/+	1.297665641	6	0.947680991	0.242835829	0.968298081	FALSE

Duration — PdgfraCre;Tbx1 (All)

Measurement	Call type	PND	Genotype	# of samples	Average	s.e.m.	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	RM 2 Way ANOVA Source	Correction	Type III SS	df	Mean Square	F	Sig.	t-test Sample1	t-test Sample2	Statistic	dof	Cohen's d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Duration	All	P8	PdgfraCre;Tbx1/+	8	213.0079521	37.53124163	0.926702066	0.486557768	0.862529525	FALSE	0.211873328	2	24	0.810564284	0.810564284	FALSE	Within Subjects							PdgfraCre;Tbx1/+	WT;Tbx1/flox	-0.093039895	14	-0.046519947	0.92719021	0.92719021	FALSE
			WT;Tbx1/flox	8	218.4553202	44.93729257	0.767937109	0.039833943	0.862529525	FALSE							PND	Sphericity Assumed	3902.027524	1	3902.027524	0.312510909	0.582056653	PdgfraCre;Tbx1/+	WT;Tbx1/+	-0.329883921	17	-0.178493026	0.745673888	0.862529525	FALSE
			WT;Tbx1/+	11	325.0474966	30.83990383	0.924511428	0.308039398	0.862529525	FALSE							PND * Genotype	Sphericity Assumed	40899.39978	2	20449.69989	1.637803489	0.218281319	PdgfraCre;Tbx1/+	WT;Tbx1/+	-2.027150892	14	-0.941936916	0.062036353	0.862529525	FALSE
		P12	PdgfraCre;Tbx1/+	9	180.1755783	41.40608696	0.939344896	0.575019684	0.862529525	FALSE	0.390880507	2	21	0.680689304	0.810564284	FALSE	Error(PND)	Sphericity Assumed	262207.0967	21	12486.05222			PdgfraCre;Tbx1/+	WT;Tbx1/flox	-1.831107564	14	-0.922791474	0.088446699	0.862529525	FALSE
			WT;Tbx1/flox	7	320.0533716	68.70128939	0.980379221	0.961358291	0.961358291	FALSE							Between Subjects							PdgfraCre;Tbx1/+	WT;Tbx1/+	-2.123475695	18	-0.954564363	0.047806477	0.862529525	FALSE
			WT;Tbx1/+	11	302.6964725	39.66885402	0.929910487	0.409948701	0.862529525	FALSE							Intercept		3049614.801	1	3049614.801	129.5812445	1.91413E-10	WT;Tbx1/flox	WT;Tbx1/+	-0.235645859	16	0.113933307	0.816697222	0.92719021	FALSE
																		Genotype		120624.2088	2	60312.1044	2.562722855	0.100956354							
																		Error		494222.0692	21	23534.38425									

Figure 5A

Figure	Measurement	Month	Session	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 5A	Social Interaction (s)	1	1	oTbx1+/+	27	16.40740741	3.092707745	0.822716916	0.000347817	0.000695635	TRUE	0.0722396	1	36	0.789637762	0.789637762	FALSE	Corrected Model	0.497244222	3	72	0.685351082	oTbx1+/-	oTbx1+/+	138.5	-0.05221416	0.775387862	0.849168654	FALSE
				oTbx1+/-	11	16.09090909	6.185693972	0.732690896	0.001237266	0.001649687	TRUE							Genotype	0.111765542	1	72	0.739114848	oTbx1+/-	oTbx1+/+	142	-0.033939204	0.849168654	0.849168654	FALSE
			2	oTbx1+/+	27	14.40740741	3.858708012	0.644196044	7.09586E-07	2.83834E-06	TRUE	0.344949992	1	36	0.56065175	0.789637762	FALSE	Session	1.379597591	1	72	0.244039836	oTbx1+/-	oTbx1+/+	142	-0.033939204	0.849168654	0.849168654	FALSE
				oTbx1+/-	11	11	3.311138228	0.855904878	0.050946602	0.050946602	FALSE							Genotype * Session	0.262132613	1	72	0.610225677							
		2	1	oTbx1+/+	21	24.23809524	7.075411737	0.65323431	7.59541E-06	1.51908E-05	TRUE	0.093406833	1	29	0.762073072	0.762073072	FALSE	Corrected Model	1.489687533	3	58	0.226836385	oTbx1+/-	oTbx1+/+	93.5	0.087281656	0.662541772	0.723978187	FALSE
				oTbx1+/-	10	30.3	16.01808006	0.521146224	6.44244E-06	1.51908E-05	TRUE							Genotype	0.005469838	1	58	0.941298006							
			2	oTbx1+/+	21	19.61904762	5.080061956	0.764962029	0.000200312	0.000267082	TRUE	0.24965746	1	29	0.621086413	0.762073072	FALSE	Session	4.309691377	1	58	0.042338451	oTbx1+/-	oTbx1+/+	95.5	-0.072102238	0.723978187	0.723978187	FALSE
				oTbx1+/-	10	15.2	5.133333333	0.841080498	0.045458931	0.045458931	TRUE							Genotype * Session	1.217519133	1	58	0.274405205							

Measurement	Month	Session	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Social Interaction (s)	1	1	PdgfraCre;Tbx1+/+	8	19	6.450359901	0.852384539	0.100707919	0.151061878	FALSE	1.333074967	2	24	0.282495218	0.282495218	FALSE	Corrected Model	1.081487416	5	48	0.382584793	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	25.5	0.245049015	0.370382559	0.745047522	FALSE
			WT;Tbx1+/+	10	20	6.151783842	0.853382418	0.063721582	0.151061878	FALSE							Genotype	1.686430667	2	48	0.195965476	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	37	0.062828086	0.828557064	0.828557064	FALSE
			WT;Tbx1+/flox	9	10.11111111	2.468942894	0.870970279	0.125930494	0.151116593	FALSE							Session	0.495285736	1	48	0.484979032	WT;Tbx1+/+	WT;Tbx1+/flox	35.5	-0.177951304	0.496698348	0.745047522	FALSE
		2	PdgfraCre;Tbx1+/+	8	11.625	4.37907973	0.852137342	0.100130923	0.151061878	FALSE	1.941945267	2	24	0.165308228	0.282495218	FALSE	Genotype*Session	0.843209591	2	48	0.436601196	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	31	0.116690007	0.672974085	0.807568902	FALSE
			WT;Tbx1+/+	10	23.3	9.316711389	0.746606164	0.003244757	0.01946854	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	29.5	-0.219898302	0.408245349	0.745047522	FALSE
			WT;Tbx1+/flox	9	7	1.748014747	0.925306341	0.437971545	0.437971545	FALSE												WT;Tbx1+/+	WT;Tbx1+/flox	29	-0.29970746	0.21102427	0.745047522	FALSE
	2	1	PdgfraCre;Tbx11+/+	8	25.375	10.82480204	0.646705991	0.000554496	0.001663489	TRUE	0.105027705	2	18	0.900847215	0.972131	FALSE	Corrected Model	0.499425251	5	36	0.774566604	PdgfraCre;Tbx11+/+	WT;Tbx1+/flox	27	0.029880715	0.955089355	0.955089355	FALSE
			WT;Tbx1+/+	6	26.16666667	19.80642432	0.554416353	0.000124267	0.000745601	TRUE							Genotype	0.03924997	2	36	0.961551418	PdgfraCre;Tbx11+/+	WT;Tbx1+/+	13.5	0.362284419	0.228438228	0.89044289	FALSE
			WT;Tbx1/flox	7	21.28571429	7.870119155	0.815067345	0.057543741	0.067157791	FALSE							Session	1.75456665	1	36	0.193651322	WT;Tbx1+/+	WT;Tbx1+/flox	14.5	0.257539377	0.445221445	0.89044289	FALSE
		2	PdgfraCre;Tbx11+/+	8	17.875	6.418270517	0.835164751	0.067157791	0.067157791	FALSE	0.028309139	2	18	0.972131	0.972131	FALSE	Genotype*Session	0.238208993	2	36	0.789270203	PdgfraCre;Tbx11+/+	WT;Tbx1+/flox	26	0.05976143	0.866511267	0.955089355	FALSE
			WT;Tbx1+/+	6	22.83333333	12.78910126	0.642868731	0.00146069	0.002921381	TRUE												PdgfraCre;Tbx11+/+	WT;Tbx1+/+	20.5	-0.120761473	0.754578755	0.955089355	FALSE
			WT;Tbx1+/flox	7	18.85714286	9.174053919	0.752239945	0.013389431	0.020084147	TRUE												WT;Tbx1+/+	WT;Tbx1+/flox	15	-0.237728656	0.445221445	0.89044289	FALSE

Table 1

Figure	Measurement	Month	Session	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 5B	Sniffing novel object (s)	1	1	oTbx1+/+	26	14.57692308	2.634781017	0.846840904	0.001236135	0.00247227	TRUE	0.08242971	1	35	0.775724316	0.775724316	FALSE	Corrected Model	2.521664315	3	70	0.064800787	oTbx1+/-	oTbx1+/+	129	-0.076479138	0.65844134	0.934817788	FALSE
				oTbx1+/-	11	12.36363636	3.739889953	0.818474025	0.016467715	0.016467715	TRUE							Genotype	2.25065E-05	1	70	0.996228266							
			2	oTbx1+/+	26	8.115384615	1.778786901	0.802681024	0.000195641	0.000782565	TRUE	1.057934255	1	35	0.310742865	0.621485731	FALSE	Session	3.706544227	1	70	0.05826308	oTbx1+/-	oTbx1+/+	140	-0.016388387	0.934817788	0.934817788	FALSE
				oTbx1+/-	11	10.36363636	4.101602186	0.789282155	0.006799356	0.009065808	TRUE							Genotype * Session	1.03048056	1	70	0.31354175							
		2	1	oTbx1+/+	21	8	1.657594356	0.868273317	0.008915895	0.01783179	TRUE	2.542177104	1	29	0.12168399	0.243367981	FALSE	Corrected Model	0.495546864	3	58	0.686790912	oTbx1+/-	oTbx1+/+	104	0.007589709	0.983438982	0.983438982	FALSE
				oTbx1+/-	10	6.8	1.41264134	0.911900064	0.294299164	0.294299164	FALSE							Genotype	0.116338641	1	58	0.734272506							
			2	oTbx1+/+	21	6.333333333	1.731134136	0.786296484	0.000408019	0.001632078	TRUE	0.028910762	1	29	0.86616589	0.86616589	FALSE	Session	0.920506319	1	58	0.341324669	oTbx1+/-	oTbx1+/+	99	0.045538256	0.81932528	0.983438982	FALSE
				oTbx1+/-	10	5.9	2.008039398	0.823579501	0.027996164	0.037328218	TRUE							Genotype * Session	0.082129844	1	58	0.775452844							

Table 2

Measurement	Month	Session	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Generalized Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Sniffing novel object (s)	1	1	PdgfraCre;Tbx1+/+	8	12	3.9551052	0.861423719	0.124057733	0.148869279	FALSE	0.476633313	2	23	0.626867139	0.952913718	FALSE	Corrected Model	3.093385391	5	46	0.017260974	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	30.5	-0.128395008	0.672974085	0.931427396	FALSE
			WT;Tbx1+/+	9	14	3.989569735	0.818691778	0.033304831	0.066609661	FALSE							Genotype	0.177160277	2	46	0.838214318	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	30	-0.140028008	0.605841218	0.931427396	FALSE
			WT;Tbx1+/flox	9	17.44444444	5.725457554	0.864838833	0.10819449	0.148869279	FALSE							Session	12.98794535	1	46	0.000767292	WT;Tbx1+/flox	WT;Tbx1+/+	37	0.072845563	0.796174414	0.931427396	FALSE
		2	PdgfraCre;Tbx1+/+	8	7	3.5152728	0.759035249	0.010275376	0.030826129	TRUE	0.048332199	2	23	0.952913718	0.952913718	FALSE	Genotype*Session	0.931759073	2	46	0.401165818	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	28	-0.186704011	0.480707528	0.931427396	FALSE
			WT;Tbx1+/+	9	9.555555556	3.667087518	0.721553597	0.00253196	0.01519176	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	26	-0.233380014	0.370382559	0.931427396	FALSE
			WT;Tbx1+/flox	9	7.666666667	2.254624876	0.919315229	0.386600395	0.386600395	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	39	-0.031219527	0.931427396	0.931427396	FALSE
	2	1	PdgfraCre;Tbx1+/+	8	7.5	2.692582404	0.907626652	0.337653424	0.37778911	FALSE	0.627763682	2	18	0.545071894	0.545071894	FALSE	Corrected Model	0.477274668	5	36	0.790748408	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	27	-0.029880715	0.955089355	1	FALSE
			WT;Tbx1+/+	6	8	2.695675549	0.87228892	0.23548219	0.353223285	FALSE							Genotype	0.057987693	2	36	0.943749519	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	23.5	-0.017251639	1	1	FALSE
			WT;Tbx1+/flox	7	8.571428571	3.537938214	0.813275039	0.055283937	0.110567873	FALSE							Session	1.061158579	1	36	0.309822637	WT;Tbx1+/flox	WT;Tbx1+/+	19	-0.079242885	0.835664336	1	FALSE
		2	PdgfraCre;Tbx1+/+	8	8	4.110092109	0.764055086	0.011666808	0.070000849	FALSE	1.298312466	2	18	0.297363734	0.545071894	FALSE	Genotype*Session	0.658142803	2	36	0.523930478	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	27	-0.029880715	0.955089355	1	FALSE
			WT;Tbx1+/+	6	6.333333333	1.452966315	0.90065047	0.37778911	0.37778911	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	19.5	-0.155264751	0.662004662	1	FALSE
			WT;Tbx1+/flox	7	4.428571429	2.191511143	0.796592563	0.037898712	0.110567873	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	14	-0.277350098	0.365967366	1	FALSE

Figure 6A

Figure	Measurement	Month	Delay (s)	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 6A	% alternation	1	0	oTbx1+/+	24	87.96296296	2.209774585	0.817882433	0.000586058	0.001758173	TRUE	1.26349718	1	32	0.269351926	0.808055777	FALSE	Corrected Model	66.06011981	5	96	0	oTbx1+/-	oTbx1+/+	113	0.045374261	0.809114032	0.809114032	FALSE
				oTbx1+/-	10	88.88888889	3.3126933	0.84073699	0.04503041	0.054036492	FALSE																		
			15	oTbx1+/+	24	51.85185185	2.184325276	0.813035203	0.000484908	0.001758173	TRUE	0.003101076	1	32	0.955937164	0.955937164	FALSE	Genotype	4.660889678	1	96	0.033349839	oTbx1+/-	oTbx1+/+	113	0.045374261	0.809114032	0.809114032	FALSE
				oTbx1+/-	10	53.33333333	3.228814032	0.884986306	0.14879822	0.14879822	FALSE																		
			30	oTbx1+/+	24	44.90740741	2.059150213	0.905249105	0.027853786	0.041780679	TRUE	0.011740247	1	32	0.914392707	0.955937164	FALSE	Delay	122.9828701	2	96	0	oTbx1+/-	oTbx1+/+	43	0.499116867	0.002700336	0.008101007	TRUE
				oTbx1+/-	10	57.77777778	2.771598064	0.819704545	0.025134415	0.041780679	TRUE							Genotype * Delay	3.150937861	2	96	0.04727228							
		2	0	oTbx1+/+	20	88.88888889	2.418254167	0.832136365	0.002714676	0.008144029	TRUE	0.005608641	1	27	0.940853676	0.940853676	FALSE	Corrected Model	38.5469499	5	81	0	oTbx1+/-	oTbx1+/+	80	-0.087537622	0.660118692	0.660118692	FALSE
				oTbx1+/-	9	85.18518519	4.899539465	0.790885567	0.016062009	0.032124018	TRUE							Genotype	2.046863951	1	81	0.156366686							
			15	oTbx1+/+	20	54.44444444	2.265657383	0.898171934	0.038114561	0.057171842	FALSE	2.474961012	1	27	0.127318695	0.381956086	FALSE	Delay	81.56087251	2	81	0	oTbx1+/-	oTbx1+/+	72.5	-0.153190838	0.443859589	0.660118692	FALSE
				oTbx1+/-	9	48.14814815	5.237828009	0.911520399	0.326661423	0.337937343	FALSE							Genotype * Delay	0.243946056	2	81	0.784103458							
			30	oTbx1+/+	20	51.11111111	2.47140896	0.800994201	0.000893986	0.005363913	TRUE	0.039618489	1	27	0.843719703	0.940853676	FALSE						oTbx1+/-	oTbx1+/+	75	-0.131306433	0.501389665	0.660118692	FALSE
				oTbx1+/-	9	49.38271605	3.266359643	0.913067364	0.337937343	0.337937343	FALSE																		

Measurement	Month	Delay (s)	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Generalized Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
% alternation	1	0	PdgfraCre;Tbx1+/+	8	86.11111111	2.777777778	0.827205306	0.05551859	0.124916827	FALSE	0.400924437	2	21	0.674718601	0.961718489	FALSE	Corrected Model	31.01103484	8	63	0	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	26.5	-0.221711013	0.423447141	1	FALSE
			WT;Tbx1+/+	7	87.3015873	4.489566865	0.664435629	0.001497441	0.006738486	TRUE							Genotype	0.104620037	2	63	0.900822824	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	27.5	-0.014940358	1	1	FALSE
			WT;Tbx1+/flox	9	90.12345679	4.320987654	0.775541227	0.01070126	0.032103779	TRUE							Delay	121.2150635	2	63	0	WT;Tbx1+/+	WT;Tbx1+/flox	27.5	0.105851225	0.757692308	1	FALSE
		15	PdgfraCre;Tbx1+/+	8	50	4.199605256	0.567903737	6.66981E-05	0.000600283	TRUE	0.122464503	2	21	0.885364514	0.961718489	FALSE	Genotype*Delay	0.26479949	4	63	0.899485819	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	35	-0.023338001	0.962566845	1	FALSE
			WT;Tbx1+/+	7	53.96825397	4.489566865	0.893581377	0.293870591	0.293870591	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	22.5	-0.164343934	0.612587413	1	FALSE
			WT;Tbx1+/flox	9	51.85185185	3.207501495	0.872821722	0.131797654	0.175446148	FALSE												WT;Tbx1+/+	WT;Tbx1+/flox	27	-0.119082628	0.680594406	1	FALSE
		30	PdgfraCre;Tbx1+/+	8	45.83333333	3.893138268	0.871515193	0.155952131	0.175446148	FALSE	0.039106145	2	21	0.961718489	0.961718489	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	35.5	-0.011669001	1	1	FALSE
			WT;Tbx1+/+	7	44.44444444	3.428963332	0.8577129	0.144397137	0.175446148	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	27	0.029880715	0.955089355	1	FALSE
			WT;Tbx1+/flox	9	44.44444444	3.703703704	0.853289618	0.080990849	0.145783528	FALSE												WT;Tbx1+/+	WT;Tbx1+/flox	30	0.039694209	0.918181818	1	FALSE
	2	0	PdgfraCre;Tbx1+/+	8	87.5	3.893138268	0.735991362	0.005710635	0.019119856	TRUE	0.205732484	2	17	0.816047931	0.925960593	FALSE	Corrected Model	22.23991673	8	51	3.94129E-14	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	22	-0.069006556	0.851814852	0.94971695	FALSE
			WT;Tbx1+/+	6	90.74074074	5.30289669	0.772718964	0.032937266	0.074108848	TRUE							Genotype	0.119696984	2	51	0.887437723	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	19.5	-0.155264751	0.662004662	0.94971695	FALSE
			WT;Tbx1+/flox	6	88.88888889	4.05720413	0.853191607	0.167002833	0.214717928	FALSE							Delay	87.4546486	2	51	0	WT;Tbx1+/+	WT;Tbx1+/flox	15	-0.138675049	0.699134199	0.94971695	FALSE
		15	PdgfraCre;Tbx1+/+	8	55.55555556	2.969569355	0.848910992	0.092877818	0.139316727	FALSE	0.476	2	17	0.629298498	0.925960593	FALSE	Genotype*Delay	0.640087093	4	51	0.636335206	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	17	0.241522946	0.413586414	0.94971695	FALSE
			WT;Tbx1+/+	6	57.40740741	5.30289669	0.907578534	0.420658529	0.473240845	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	23	-0.034503278	0.94971695	0.94971695	FALSE
			WT;Tbx1+/flox	6	50	3.79516695	0.701259348	0.000673285	0.019119856	TRUE												WT;Tbx1+/+	WT;Tbx1+/flox	13	-0.231125082	0.484848485	0.94971695	FALSE
		30	PdgfraCre;Tbx1+/+	8	52.77777778	3.482128725	0.826041799	0.053987435	0.097177383	FALSE	0.077272727	2	17	0.925960593	0.925960593	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	23	0.034503278	0.94971695	0.94971695	FALSE
			WT;Tbx1+/+	6	48.14814815	5.493480361	0.665788341	0.002641354	0.019119856	TRUE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	19.5	-0.155264751	0.662004662	0.94971695	FALSE
			WT;Tbx1+/flox	6	51.85185185	4.684855793	0.915458969	0.473270921	0.473270921	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	15.5	0.115562541	0.818181818	0.94971695	FALSE

Figure 5B

Figure	Measurement	Month	Delay (s)	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 5B	Latency to correct arm (s)	1	0	oTbx1+/+	24	7.060929233	0.514093404	0.800175638	0.00029653	0.00177918	TRUE	0.277615077	1	32	0.601902435	0.601902435	FALSE	Corrected Model	16.83605715	5	96	6.44695E-12	oTbx1+/-	oTbx1+/+	110	0.064820372	0.723944044	0.896531195	FALSE
				oTbx1+/-	10	7.045238095	0.530928951	0.926337453	0.412848339	0.476660672	FALSE							Genotype	2.181306251	1	96	0.142968861							
			15	oTbx1+/+	24	23.71111111	3.024286701	0.822986774	0.000717253	0.00215176	TRUE	0.326055759	1	32	0.571979595	0.601902435	FALSE	Delay	27.42527721	2	96	3.79112E-10	oTbx1+/-	oTbx1+/+	116	-0.025928149	0.896531195	0.896531195	FALSE
				oTbx1+/-	10	21.89666667	3.777162368	0.850065677	0.058193748	0.11638749 6	FALSE							Genotype * Delay	2.058495873	2	96	0.13324324							
			30	oTbx1+/+	24	30.76111111	2.763625603	0.951467639	0.29143235	0.437148525	FALSE	1.123976658	1	32	0.297000607	0.601902435	FALSE						oTbx1+/-	oTbx1+/+	67	-0.343547973	0.045949085	0.137847254	FALSE
				oTbx1+/-	10	20.47833333	3.235213323	0.932865391	0.476660672	0.476660672	FALSE																		
		2	0	oTbx1+/+	20	8.249603175	1.084890689	0.650063636	1.01826E-05	6.10959E-05	TRUE	1.416496538	1	27	0.244340874	0.366511312	FALSE	Corrected Model	9.617022113	5	81	3.11418E-07	oTbx1+/-	oTbx1+/+	79	0.096291384	0.626843222	0.626843222	FALSE
				oTbx1+/-	10	10.25432099	2.448449218	0.825388972	0.039632942	0.118898826	FALSE							Genotype	1.113528338	1	81	0.294453611							
			15	oTbx1+/+	20	21.45690476	2.262232887	0.949405569	0.358157001	0.358157001	FALSE	1.772805112	1	27	0.194168243	0.366511312	FALSE	Delay	18.44615477	2	81	2.50312E-07	oTbx1+/-	oTbx1+/+	58	0.28012039	0.13969359	0.419080769	FALSE
				oTbx1+/-	9	32.12037037	6.192265298	0.86589138	0.11106244	0.145532979	FALSE							Genotype * Delay	2.582518592	2	81	0.081796465							
			30	oTbx1+/+	20	27.41833333	3.194822863	0.924555501	0.121277482	0.145532979	FALSE	0.387060947	1	27	0.539067153	0.539067153	FALSE						oTbx1+/-	oTbx1+/+	71	-0.166321482	0.390153575	0.585230362	FALSE
				oTbx1+/-	9	23.54814815	4.343311557	0.852295522	0.078981247	0.145532979	FALSE																		

Measurement	Month	Delay (s)	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Generalized Linear Mixed Model Source	F	df1	df2	Sig.	Mann-Whitney U test Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Latency to correct arm (s)	1	0	PdgfraCre;Tbx1+/+	8	6.936507937	1.245302549	0.49801402	9.77951E-06	8.80156E-05	TRUE	0.118489741	2	21	0.888850552	0.888850552	FALSE	Corrected Model	12.80465967	8	63	9.75433E-11	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	26	-0.233380014	0.370382559	0.666688606	FALSE
			WT;Tbx1+/+	7	7.238095238	0.859979629	0.889504742	0.272090107	0.61220274	FALSE							Genotype	1.578749645	2	63	0.214282282	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	27	-0.029880715	0.955089355	1	FALSE
			WT;Tbx1+/flox	7	7.033730159	0.619516622	0.953498416	0.728441817	0.964816546	FALSE							Delay	41.38453208	2	63	3.34022E-12	WT;Tbx1+/flox	WT;Tbx1+/+	31.5	0	1	1	FALSE
		15	PdgfraCre;Tbx1+/+	8	32.51875	7.148675811	0.867888501	0.143716676	0.61220274	FALSE	1.806979696	2	21	0.188754388	0.283131582	FALSE	Genotype*Delay	4.562533663	4	63	0.0026726	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	30	0.140028008	0.605841218	0.778938708	FALSE
			WT;Tbx1+/+	7	13.89285714	1.511419745	0.881281719	0.232198499	0.61220274	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	8	0.597614305	0.020512821	0.140034965	FALSE
			WT;Tbx1+/flox	7	23.51851852	3.323532402	0.936177907	0.542290058	0.964816546	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	14	0.463099109	0.071153846	0.213461538	FALSE
		30	PdgfraCre;Tbx1+/+	8	33.4	5.751512051	0.954254508	0.753939808	0.964816546	FALSE	2.91193178	2	21	0.076529454	0.229588362	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	20	0.373408022	0.138790621	0.312278898	FALSE
			WT;Tbx1+/+	7	37.25952381	4.89605426	0.981124342	0.964816546	0.964816546	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	22	-0.179284291	0.535819736	0.778938708	FALSE
			WT;Tbx1+/flox	9	23.36111111	2.668082668	0.967019227	0.868103653	0.964816546	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	11	-0.542487527	0.031118881	0.140034965	FALSE
	2	0	PdgfraCre;Tbx1+/+	8	7.634672619	0.856014268	0.893506903	0.252187548	0.567421983	FALSE	2.318063852	2	17	0.128762601	0.193143916	FALSE	Corrected Model	5.716167127	8	51	3.39007E-05	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	20	-0.138013112	0.662004662	0.754578755	FALSE
			WT;Tbx1+/+	6	6.136574074	0.419288306	0.935332466	0.621863247	0.799538461	FALSE							Genotype	0.163060535	2	51	0.849980901	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	14.5	0.327781141	0.282384282	0.754578755	FALSE
			WT;Tbx1+/flox	6	11.18253968	3.263533536	0.781374891	0.039698931	0.357290375	FALSE							Delay	20.11273606	2	51	3.63144E-07	WT;Tbx1+/flox	WT;Tbx1+/+	12.5	0.25423759	0.484848485	0.754578755	FALSE
		15	PdgfraCre;Tbx1+/+	6	18.46458333	3.639106975	0.925920356	0.479712667	0.799538461	FALSE	3.251300612	2	17	0.063728018	0.191184054	FALSE	Genotype*Delay	1.061101788	4	51	0.385349838	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	16	-0.276026224	0.344988345	0.754578755	FALSE
			WT;Tbx1+/+	6	20.77301587	2.072925902	0.928919732	0.571785689	0.799538461	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	18	-0.207019668	0.490842491	0.754578755	FALSE
			WT;Tbx1+/flox	6	26.13055556	5.381154386	0.960050918	0.820132956	0.820132956	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	15	0.138675049	0.699134199	0.754578755	FALSE
		30	PdgfraCre;Tbx1+/+	6	28.00625	6.083776841	0.851422944	0.0984807	0.44316315	FALSE	0.906736253	2	17	0.422502853	0.422502853	FALSE						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	21	0.103509834	0.754578755	0.754578755	FALSE
			WT;Tbx1+/+	6	30.63333333	6.500957194	0.873464834	0.240368777	0.567421983	FALSE												PdgfraCre;Tbx1+/+	WT;Tbx1+/+	21	0.103509834	0.754578755	0.754578755	FALSE
			WT;Tbx1+/flox	6	23.41944444	3.546480824	0.954883247	0.779566398	0.820132956	FALSE												WT;Tbx1+/flox	WT;Tbx1+/+	13	-0.231125082	0.484848485	0.754578755	FALSE

Table 1

Figure	Measurement	Month	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test				Levene test						Mann-Whitney U test						
							Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 7A	% open arms time	1	oTbx1+/+	23	13.92453346	2.578767506	0.871431687	0.00682799	0.01365598	TRUE	0.03061532	1	32	0.862203787	0.862203787	FALSE	oTbx1+/-	oTbx1+/+	100	0.167302625	0.343976779	0.343976779	FALSE
			oTbx1+/-	11	16.98607089	3.279767612	0.906204785	0.219762803	0.219762803	FALSE													
		2	oTbx1+/+	20	5.617823791	1.293061822	0.865730567	0.009893527	0.019787054	TRUE	0.129376135	1	28	0.721778502	0.721778502	FALSE	oTbx1+/-	oTbx1+/+	98	0.016064387	0.948316451	0.948316451	FALSE
			oTbx1+/-	10	6.108311403	1.697112495	0.861318188	0.079087694	0.079087694	FALSE													

Table 2

Measurement	Month	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test				Levene test						One way ANOVA						Student t test							
						Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Sample1	Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
% open arms time	1	PdgfraCre;Tbx1+/+	7	12.90725421	4.103497891	0.886610333	0.257444394	0.386166591	FALSE	0.107045624	2	20	0.898995957	0.898995957	FALSE	Intercept	4421.166501	1	4421.166501	26.47361987	4.92643E-05	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-0.090770003	14	-0.045743782	0.928961264	0.928961264	FALSE
		WT;Tbx1+/+	7	15.44729701	4.180055074	0.973508997	0.922513345	0.922513345	FALSE							Genotype	24.86679522	2	12.43339761	0.074450271	0.928509658	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-0.43363151	12	-0.231785792	0.672249124	0.928961264	FALSE
		WT;Tbx1+/flox	9	13.53137901	5.135617804	0.770175921	0.009280317	0.02784095	TRUE							Error	3340.054381	20	167.0027191			WT;Tbx1+/flox	WT;Tbx1+/+	-0.277229778	14	-0.139710676	0.785655127	0.928961264	FALSE
	2	PdgfraCre;Tbx1+/+	8	7.962568057	2.541302393	0.9223745	0.449425688	0.674138532	FALSE	1.779086372	2	17	0.19881424	0.19881424	FALSE	Intercept	563.5759562	1	563.5759562	18.85031974	0.000443371	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.934686976	12	1.044850385	0.076951997	0.126596757	FALSE
		WT;Tbx1+/+	6	6.17210563	1.907847722	0.972621635	0.90955982	0.90955982	FALSE							Genotype	127.1072061	2	63.55360307	2.125721876	0.14997335	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.529258771	12	0.285832405	0.606278344	0.606278344	FALSE
		WT;Tbx1+/flox	6	1.937216264	1.116534507	0.751259955	0.02049345	0.061480349	FALSE							Error	508.2561668	17	29.89742157			WT;Tbx1+/flox	WT;Tbx1+/+	-1.915762576	10	-1.106060039	0.084397838	0.126596757	FALSE

Figure 7B

Figure	Measurement	Month	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Student t test Sample1	Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 7B	% open arm visits	1	oTbx1+/+	23	38.76689665	3.457186322	0.959194931	0.447303171	0.472928936	FALSE	0.034575631	1	32	0.853661423	0.853661423	FALSE	oTbx1+/-	oTbx1+/+	1.741236711	32	0.638317885	0.091249849	0.091249849	FALSE
			oTbx1+/-	11	48.92821047	4.328945934	0.935843176	0.472928936	0.472928936	FALSE														
		2	oTbx1+/+	20	16.20798761	2.580004845	0.933143261	0.177461607	0.339087396	FALSE	0.734877731	1	28	0.398584802	0.398584802	FALSE	oTbx1+/-	oTbx1+/+	0.59246246	28	0.229459724	0.558295668	0.558295668	FALSE
			oTbx1+/-	10	19.18868564	4.938885044	0.91781454	0.339087396	0.339087396	FALSE														

Measurement	Month	Genotype	# of samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Levene test Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	One way ANOVA Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Student t test Sample1	Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
% open arm visits	1	PdgfraCre;Tbx1+/+	7	35.6769162	4.664325787	0.928344033	0.536905413	0.536905413	FALSE	0.54245905	2	20	0.589631629	0.589631629	FALSE	Intercept	33992.03738	1	33992.03738	114.4766364	1.00324E-09	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-0.38767913	14	-0.195371917	0.704083822	0.851686321	FALSE
		WT;Tbx1+/+	7	41.14160143	5.028883164	0.831111044	0.081981393	0.219437109	FALSE							Genotype	109.0958965	2	54.54794824	0.183703776	0.833570659	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-0.796719857	12	-0.425864677	0.441092294	0.851686321	FALSE
		WT;Tbx1+/flox	9	39.32322217	7.418792132	0.877086869	0.146291406	0.219437109	FALSE							Error	5938.685561	20	296.934278			WT;Tbx1+/+	WT;Tbx1+/flox	-0.190455488	14	-0.095980544	0.851686321	0.851686321	FALSE
	2	PdgfraCre;Tbx1+/+	8	20.47904805	3.951856074	0.921801353	0.444650175	0.666975262	FALSE	0.062288917	2	17	0.939824807	0.939824807	FALSE	Intercept	4860.798667	1	4860.798667	38.36512241	9.80937E-06	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.794257938	12	0.969010037	0.097985827	0.293957482	FALSE
		WT;Tbx1+/+	6	16.68154542	5.036940952	0.970955927	0.898793324	0.898793324	FALSE							Genotype	375.5691249	2	187.7845625	1.482138681	0.255070525	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.602275965	12	0.325266196	0.558199622	0.558199622	FALSE
		WT;Tbx1+/flox	6	10.03968254	4.156071184	0.906492643	0.413726943	0.666975262	FALSE							Error	2153.872376	17	126.698375			WT;Tbx1+/+	WT;Tbx1+/flox	-1.017097195	10	-0.537221339	0.33308331	0.499624965	FALSE

Figure 8A — Total Distance (cm)

Figure	Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	Levene df1	Levene df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MWU Sample1	MWU Sample2	MWU Statistic	R (Z score)	MWU p_value	MWU p_adj(FDR BH)	MWU Rejects(FDR 5%)
Figure 8A	Total Distance (cm)	1	5	oTbx1+/+	24	523.3911589	30.20140117	0.975883028	0.809668571	0.809668571	FALSE	4.130581659	1	33	0.050226926	0.301361554	FALSE	Corrected Model	14.18810262	11	198	0	oTbx1+/-	oTbx1+/+	125	0.042043748	0.82012655	0.902572012	FALSE
				oTbx1+/-	11	575.0133513	81.09884483	0.924914551	0.361718736	0.569470319	FALSE							Genotype	0.631374662	1	198	0.4278029							
			10	oTbx1+/+	24	428.858175	25.3219597	0.947354113	0.237276704	0.569470319	FALSE	1.663209327	1	33	0.206136149	0.412272297	FALSE	Time	27.84739692	5	198	0	oTbx1+/-	oTbx1+/+	126	0.036037499	0.84744042	0.902572012	FALSE
				oTbx1+/-	11	463.1100966	54.68330524	0.916262967	0.288732783	0.569470319	FALSE							Genotype * Time	0.955127959	5	198	0.446565974							
			15	oTbx1+/+	24	353.2098723	29.23769273	0.966833314	0.589787776	0.707745332	FALSE	0.140317736	1	33	0.710361842	0.896645927	FALSE						oTbx1+/-	oTbx1+/+	128	0.024024999	0.902572012	0.902572012	FALSE
				oTbx1+/-	11	411.012008	55.7960521	0.755590857	0.002459007	0.029508078	TRUE																		
			20	oTbx1+/+	24	323.8313083	27.24334579	0.975754462	0.806709631	0.809668571	FALSE	0.017135892	1	33	0.896645927	0.896645927	FALSE						oTbx1+/-	oTbx1+/+	120	0.072074997	0.687299671	0.902572012	FALSE
				oTbx1+/-	11	349.0746296	38.67701388	0.902850991	0.200221378	0.569470319	FALSE																		
			25	oTbx1+/+	24	264.9099236	32.94277932	0.903535486	0.02559625	0.153577498	FALSE	0.040522424	1	33	0.841698265	0.896645927	FALSE						oTbx1+/-	oTbx1+/+	93	0.23424374	0.173954599	0.902572012	FALSE
				oTbx1+/-	11	339.7488583	47.44333904	0.940451418	0.525819716	0.701092954	FALSE																		
			30	oTbx1+/+	24	246.8655593	33.32994361	0.947492768	0.238941106	0.569470319	FALSE	2.249666174	1	33	0.143150583	0.412272297	FALSE						oTbx1+/-	oTbx1+/+	111	-0.126131245	0.472220258	0.902572012	FALSE
				oTbx1+/-	11	201.5505166	37.02036109	0.926826438	0.37964688	0.569470319	FALSE																		

Total Distance (cm) — Repeated Measures ANOVA / Student t test

Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	Levene df1	Levene df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	RM ANOVA Source	Sphericity	Type III Sum of Squares	df	Mean Square	F	Sig.	t Sample1	t Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Total Distance (cm)	1	5	PdgfraCre;Tbx1+/+	8	620.0439824	61.18380332	0.896772958	0.270160506	0.686126804	FALSE	0.78676576	2	21	0.468282323	0.702423485	FALSE	**Within subjects**							PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.782324381	15	0.866053977	0.094944335	0.18988867	FALSE
			WT;Tbx1+/+	7	441.6039626	47.42794277	0.93556056	0.599115546	0.898673318	FALSE							Time	Greenhouse-Geisser	1232066.871	3.551512016	346913.3331	18.43930669	6.42253E-10	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.254227774	13	1.166673712	0.042077619	0.182512156	FALSE
			WT;Tbx1+/flox	9	501.0898018	31.91778323	0.918522745	0.380149455	0.760298911	FALSE							Time * Genotype	Greenhouse-Geisser	149400.5872	7.103024031	21033.37769	1.11797635	0.361061454	WT;Tbx1+/+	WT;Tbx1+/flox	1.0781018	14	0.543312238	0.299214264	0.414296673	FALSE
		10	PdgfraCre;Tbx1+/+	8	528.1841483	47.57087669	0.949171774	0.702877111	0.973214461	FALSE	1.114125751	2	21	0.346839968	0.693679937	FALSE	Error(Time)	Greenhouse-Geisser	1403165.787	74.58175233	18813.79484			PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.69065136	15	1.307421554	0.016770132	0.182512156	FALSE
			WT;Tbx1+/+	7	384.9748501	26.01161759	0.970333	0.900838417	0.980892361	FALSE							**Between subjects**							PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.532927569	13	1.31091456	0.024986314	0.182512156	FALSE
			WT;Tbx1+/flox	9	374.699896	33.33231151	0.977856565	0.952350694	0.980892361	FALSE							Intercept		18280351.96	1	18280351.96	417.5782475	2.43882E-15	WT;Tbx1+/+	WT;Tbx1+/flox	-0.231693882	14	-0.116762741	0.820128066	0.820128066	FALSE
		15	PdgfraCre;Tbx1+/+	8	435.548186	56.01819145	0.984272937	0.980892361	0.980892361	FALSE	2.264993593	2	21	0.12860899	0.678212575	FALSE	Genotype		479372.5079	2	239686.254	5.475155296	0.012198718	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.870796877	15	0.909043883	0.081011856	0.182512156	FALSE
			WT;Tbx1+/+	7	331.29903	29.99137947	0.789411464	0.03213434	0.431236082	FALSE							Error		919318.4597	21	43777.06951			PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.571756012	13	0.813461019	0.140019642	0.22912305	FALSE
			WT;Tbx1+/flox	9	297.0620262	48.93223185	0.905661169	0.286676525	0.686126804	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-0.554475257	14	-0.279429264	0.588002519	0.661502834	FALSE
		20	PdgfraCre;Tbx1+/+	8	411.8585995	57.63072458	0.973883362	0.926616722	0.980892361	FALSE	1.597337999	2	21	0.226070858	0.678212575	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.92146405	15	0.933663704	0.073882743	0.182512156	FALSE
			WT;Tbx1+/+	7	260.7565684	42.47375038	0.889114132	0.270074414	0.686126804	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.054788232	13	1.064851326	0.060276832	0.182512156	FALSE
			WT;Tbx1+/flox	9	294.6429583	26.46424959	0.856295547	0.087368002	0.524208011	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.70823831	14	0.35691856	0.490422016	0.630542591	FALSE
		25	PdgfraCre;Tbx1+/+	8	337.0979663	67.94142154	0.902611055	0.304945246	0.686126804	FALSE	0.479360415	2	21	0.62579012	0.750948144	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.955850532	15	0.467124166	0.647099748	0.761902345	FALSE
			WT;Tbx1+/+	7	280.8983661	57.58848895	0.918866143	0.460635174	0.763439422	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.620765095	13	0.321276459	0.545484347	0.654581216	FALSE
			WT;Tbx1+/flox	9	188.3073192	38.18211176	0.832731267	0.04791512	0.431236082	FALSE														PdgfraCre;Tbx1+/flox	WT;Tbx1+/+	-1.390976009	13	-0.700986064	0.188559711	0.264888417	FALSE
		30	PdgfraCre;Tbx1+/+	8	285.4238129	54.60920401	0.924397037	0.466546314	0.763439422	FALSE	0.105755822	2	21	0.900120365	0.900120365	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.663575002	15	0.808352151	0.116943757	0.210498763	FALSE
			WT;Tbx1+/+	7	310.8749546	62.46942788	0.966178335	0.869648503	0.980892361	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-0.308231337	13	-0.159524872	0.762789645	0.807659625	FALSE
			WT;Tbx1+/flox7	9	162.8064708	49.74066212	0.878647793	0.15193339	0.683790257	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-1.879743998	14	-0.881116514	0.081166536	0.182512156	FALSE

Figure 8B

Figure	Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	RM ANOVA	Method	Type III Sum of Squares	df	Mean Square	F	Sig.	Sample1	Sample2	t Statistic	dof	cohens_d	t p_value	t p_adj(FDR BH)	t Rejects(FDR 5%)
Figure 8B	Total Distance (cm)	2	5	oTbx1+/+	20	582.5324674	39.81366145	0.961805113	0.580510261	0.964595553	FALSE	2.027577552	1	28	0.165516111	0.496548332	FALSE	Within Subjects							oTbx1+/+	oTbx1+/-	-0.200159227	28	-0.077521335	0.842802741	0.9640554	FALSE
				oTbx1+/-	10	565.9321735	87.02480436	0.952194518	0.694501499	0.964595553	FALSE							Time	Greenhouse-Geisser	1995853.076	3.660164031	545290.6098	26.86051187	1.71481E-14								
			10	oTbx1+/+	20	465.0152343	28.92161572	0.976249885	0.877119898	0.964595553	FALSE	3.110488944	1	28	0.088700281	0.496548332	FALSE	Time * Genotype	Greenhouse-Geisser	40039.66219	3.660164031	10939.30814	0.538860217	0.691877512	oTbx1+/+	oTbx1+/-	0.243505064	28	0.094309106	0.809389409	0.9640554	FALSE
				oTbx1+/-	10	478.4165818	52.4441209	0.848827274	0.056251555	0.675018657	FALSE							Error(Time)	Greenhouse-Geisser	2080522.01	102.4845929	20300.82719										
			15	oTbx1+/+	20	361.7175911	28.74548525	0.979861932	0.932306056	0.964595553	FALSE	0.864912718	1	28	0.360316817	0.530231581	FALSE	Between Subjects							oTbx1+/+	oTbx1+/-	0.892772198	28	0.345769186	0.379588252	0.9640554	FALSE
				oTbx1+/-	10	409.8879946	50.49069356	0.979893494	0.964595553	0.964595553	FALSE							Intercept		23339048.01	1	23339048.01	306.4591211	1.2929E-16								
			20	oTbx1+/+	20	348.215865	27.97211407	0.978671917	0.915818525	0.964595553	FALSE	0.608602693	1	28	0.441859651	0.530231581	FALSE	Genotype		1846.055051	1	1846.055051	0.024240081	0.87739263	oTbx1+/+	oTbx1+/-	-0.897736865	28	-0.347691993	0.376980718	0.9640554	FALSE
				oTbx1+/-	10	300.6010342	50.30866599	0.93172014	0.465023493	0.964595553	FALSE							Error		2132399.721	28	76157.13288										
			25	oTbx1+/+	20	299.1055367	24.94704284	0.974019572	0.836412834	0.964595553	FALSE	0.876820379	1	28	0.357080786	0.530231581	FALSE								oTbx1+/+	oTbx1+/-	-0.703264717	28	-0.272373254	0.487697769	0.9640554	FALSE
				oTbx1+/-	10	264.0208722	50.34200893	0.938023496	0.531237334	0.964595553	FALSE																					
			30	oTbx1+/+	20	255.362162	39.42744784	0.949277247	0.356252313	0.964595553	FALSE	0.036907756	1	28	0.849039501	0.849039501	FALSE								oTbx1+/+	oTbx1+/-	-0.045470044	28	-0.017610472	0.9640554	0.9640554	FALSE
				oTbx1+/-	10	252.3293176	51.6114029	0.940717398	0.56102342	0.964595553	FALSE																					

Second table

Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	RM ANOVA	Method	Type III Sum of Squares	df	Mean Square	F	Sig.	Sample1	Sample2	t Statistic	dof	cohens_d	t p_value	t p_adj(FDR BH)	t Rejects(FDR 5%)
Total Distance (cm)	2	5	PdgfraCre;Tbx1+/+	8	675.3377481	67.21069573	0.960970017	0.819288618	0.90097253	FALSE	0.513944781	2	17	0.60713361	0.845882843	FALSE	Within subjects							PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.318789314	12	1.25228374	0.038853127	0.145180864	FALSE
			WT;Tbx1+/+	6	576.9360505	60.27675931	0.947046048	0.716310322	0.90097253	FALSE							Time	Sphericity Assumed	1366149.157	5	273229.8314	19.2608065	9.05321E-13	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	1.049097419	12	0.566577362	0.314803416	0.404747249	FALSE
			WT;Tbx1+/flox	6	464.38851	54.01971119	0.932193779	0.597146871	0.90097253	FALSE							Time * Genotype	Sphericity Assumed	56104.56039	10	5610.456039	0.395498206	0.94529822	WT;Tbx1+/+	WT;Tbx1+/flox	-1.390491666	10	-0.802800737	0.194547247	0.291820871	FALSE
		10	PdgfraCre;Tbx1+/+	8	563.3077705	39.36525666	0.91827703	0.414039464	0.90097253	FALSE	0.142545825	2	17	0.86817344	0.86817344	FALSE	Error(Time)	Sphericity Assumed	1205792.482	85	14185.7939			PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	3.212491801	12	1.734943863	0.007457649	0.134237677	FALSE
			WT;Tbx1+/+	6	423.8291928	36.86712279	0.939986406	0.659089302	0.90097253	FALSE							Between subjects							PdgfraCre;Tbx1+/+	WT;Tbx1+/+	2.504995321	12	1.352852094	0.02766074	0.145180864	FALSE
			WT;Tbx1+/flox	6	375.1445608	42.57179879	0.896122527	0.351518024	0.90097253	FALSE							Intercept		16804988	1	16804988	552.5642347	2.10342E-14	WT;Tbx1+/+	WT;Tbx1+/flox	-0.864483291	10	-0.499109661	0.407581081	0.489097297	FALSE
		15	PdgfraCre;Tbx1+/+	8	412.9964819	53.1536322	0.970729632	0.903671601	0.903671601	FALSE	1.538099909	2	17	0.243233282	0.518147214	FALSE	Genotype		579823.9342	2	289911.9671	9.532554405	0.001673302	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.951155687	12	1.053744506	0.074776993	0.154898089	FALSE
			WT;Tbx1+/+	6	384.9906002	24.8709437	0.960076551	0.820330406	0.90097253	FALSE							Error		517018.1288	17	30412.83111			PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.427254885	12	0.23074401	0.676759782	0.716569181	FALSE
			WT;Tbx1+/flox	6	270.0727277	45.65896481	0.885730035	0.296415795	0.90097253	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-2.210243188	10	-1.2760845	0.051539445	0.154618335	FALSE
		20	PdgfraCre;Tbx1+/+	8	383.3626534	47.27633929	0.962488762	0.833447547	0.90097253	FALSE	0.356989008	2	17	0.704902369	0.845882843	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.870093843	12	1.009966107	0.086054494	0.154898089	FALSE
			WT;Tbx1+/+	6	392.8098267	35.00674422	0.93743225	0.638585233	0.90097253	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-0.150587648	12	-0.081326625	0.882802665	0.882802665	FALSE
			WT;Tbx1+/flox	6	256.7595188	45.97104537	0.889647485	0.316344886	0.90097253	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-2.354527304	10	-1.359386973	0.040328018	0.145180864	FALSE
		25	PdgfraCre;Tbx1+/+	8	364.6078814	41.60013821	0.951382549	0.72516182	0.90097253	FALSE	1.850493244	2	17	0.187452819	0.518147214	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.664033886	12	1.438742736	0.02063703	0.145180864	FALSE
			WT;Tbx1+/+	6	296.9929018	29.92382991	0.964127822	0.8509185	0.90097253	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/+	1.232745856	12	0.665758853	0.241276061	0.334074546	FALSE
			WT;Tbx1+/flox	6	213.881712	34.05386391	0.910326291	0.438546471	0.90097253	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-1.833054905	10	-1.058479037	0.096646438	0.154898089	FALSE
		30	PdgfraCre;Tbx1+/+	8	325.6745739	75.49961398	0.953377125	0.745179396	0.90097253	FALSE	1.463867976	2	17	0.259073607	0.518147214	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	1.942251174	12	1.048935519	0.075945879	0.154898089	FALSE
			WT;Tbx1+/+	6	277.1861365	55.16889694	0.929847805	0.578927051	0.90097253	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.485421275	12	0.262157451	0.636119338	0.715634256	FALSE
			WT;Tbx1+/flox	6	139.7883048	43.83595818	0.905323563	0.40635214	0.90097253	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-1.949896567	10	-1.125773308	0.079761998	0.154898089	FALSE

Figure 8C — Total Margin Time (s)

Figure	Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MWU Sample1	MWU Sample2	MWU Statistic	R (Z score)	MWU p_value	MWU p_adj(FDR BH)	MWU Rejects(FDR 5%)
Figure 8C	Total Margin Time (s)	1	5	oTbx1+/+	24	187.9541667	9.818642574	0.989732318	0.995574805	0.995574805	FALSE	0.044942279	1	33	0.833414161	0.88610226	FALSE	Corrected Model	6.693346578	11	198	1.67687E-09	oTbx1+/-	oTbx1+/+	79	0.318331237	0.061404045	0.200151386	FALSE
				oTbx1+/-	11	224.8954545	14.69387698	0.929218242	0.402999507	0.592165394	FALSE							Genotype	0.200852805	1	198	0.654523373							
			10	oTbx1+/+	24	177.46875	8.621859233	0.960282408	0.444124046	0.592165394	FALSE	1.749233955	1	33	0.195066221	0.88610226	FALSE	Time	8.251156703	5	198	4.22922E-07	oTbx1+/-	oTbx1+/+	80	0.312324987	0.066717129	0.200151386	FALSE
				oTbx1+/-	11	202.8863636	9.029917675	0.968324319	0.869129208	0.948140954	FALSE							Genotype * Time	2.151693449	5	198	0.06094955							
			15	oTbx1+/+	24	199.6520833	11.65783522	0.959487016	0.428203171	0.592165394	FALSE	0.97523234	1	33	0.330564543	0.88610226	FALSE						oTbx1+/-	oTbx1+/+	118	0.084087497	0.63644326	0.63644326	FALSE
				oTbx1+/-	11	211.4636364	13.35909988	0.948946345	0.630518333	0.756622	FALSE																		
			20	oTbx1+/+	24	224.575	13.89428188	0.864461767	0.004105455	0.012316364	TRUE	0.073888102	1	33	0.787451039	0.88610226	FALSE						oTbx1+/-	oTbx1+/+	105	-0.162168743	0.351863535	0.540916088	FALSE
				oTbx1+/-	11	212.5727273	18.03323745	0.905116019	0.213240511	0.426481022	FALSE																		
			25	oTbx1+/+	24	238.9916667	11.76924598	0.78965836	0.000200552	0.001244162	TRUE	0.459512459	1	33	0.502577518	0.88610226	FALSE						oTbx1+/-	oTbx1+/+	109.5	-0.135140619	0.450763407	0.540916088	FALSE
				oTbx1+/-	11	221.3590909	16.82044727	0.889299635	0.136293398	0.327104154	FALSE																		
			30	oTbx1+/+	24	255.1479167	10.74757408	0.790566413	0.00020736	0.001244162	TRUE	0.020836853	1	33	0.88610226	0.88610226	FALSE						oTbx1+/-	oTbx1+/+	109	-0.138143744	0.429851852	0.540916088	FALSE
				oTbx1+/-	11	247.8272727	16.96605483	0.755966079	0.002486921	0.009947684	TRUE																		

Total Margin Time (s)

Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	RM ANOVA Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Sample1	Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Total Margin Time (s)	1	5	PdgfraCre;Tbx1+/+	8	192.4	19.42002142	0.940119442	0.612255906	0.795951402	FALSE	0.155922533	2	21	0.856606813	0.96116231	FALSE	Within subjects						PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.203680516	15	0.098970941	0.841340647	0.991826617	FALSE
			WT;Tbx1+/+	7	183.6142857	18.45076045	0.951934955	0.747275189	0.836773791	FALSE							Time	Greenhouse-Geisser	113778.7452	2.742258347	41490.89212	12.81272325	3.10589E-06	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.325226959	13	0.168320942	0.750184167	0.991826617	FALSE
			WT;Tbx1+/flox	9	187.3777778	15.59257549	0.878358778	0.150890078	0.388003057	FALSE							Time * Genotype	Greenhouse-Geisser	17553.4018	5.484516693	3200.537582	0.988351906	0.437454084	WT;Tbx1+/+	WT;Tbx1+/flox	0.156685574	14	0.078962107	0.877729762	0.991826617	FALSE
		10	PdgfraCre;Tbx1+/+	8	176.525	17.77269173	0.852568912	0.10114029	0.303420871	FALSE	0.275926248	2	21	0.761577722	0.96116231	FALSE	Error(Time)	Greenhouse-Geisser	186482.8892	57.58742528	3238.257107			PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-0.043066059	15	-0.020926343	0.96621693	0.991826617	FALSE
			WT;Tbx1+/+	7	178.4642857	12.14288305	0.937809188	0.619073312	0.795951402	FALSE							Between subjects						PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-0.045246236	13	-0.087424034	0.931666773	0.991826617	FALSE
			WT;Tbx1+/flox	9	177.5333333	15.42819389	0.958946118	0.787226042	0.836773791	FALSE							Intercept		6567827.009	1	6567827.009	673.365646	1.94254E-17	WT;Tbx1+/+	WT;Tbx1+/flox	-0.022804731	14	-0.045251736	0.964545789	0.991826617	FALSE
		15	PdgfraCre;Tbx1+/+	8	190.74375	21.81628945	0.976457289	0.943303265	0.943303265	FALSE	0.039686801	2	21	0.96116231	0.96116231	FALSE	Genotype		7190.956262	2	3595.478131	0.368625948	0.696071568	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-0.187656445	15	-0.386193777	0.704775853	0.991826617	FALSE
			WT;Tbx1+/+	7	207.3857143	23.9748646	0.793291374	0.035136728	0.128541595	FALSE							Error		204828.3396	21	9753.730455			PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-0.266236486	13	-0.514417762	0.615593079	0.991826617	FALSE
			WT;Tbx1+/flox	9	201.5555556	17.9268991	0.939932071	0.581183632	0.795951402	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-0.100323913	14	-0.199074094	0.845066999	0.991826617	FALSE
		20	PdgfraCre;Tbx1+/+	8	217.70625	22.27686468	0.930651593	0.522025005	0.795951402	FALSE	1.011932031	2	21	0.380567208	0.96116231	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.464184705	15	0.225553224	0.649180802	0.991826617	FALSE
			WT;Tbx1+/+	7	262.8928571	13.23386002	0.882178317	0.236294206	0.531661963	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-1.679155527	13	-0.869045415	0.116981237	0.991826617	FALSE
			WT;Tbx1+/flox	9	200.8777778	27.81605266	0.821368373	0.035705999	0.128541595	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-1.833493431	14	-0.923993838	0.088073727	0.991826617	FALSE
		25	PdgfraCre;Tbx1+/+	8	241.09375	16.03556994	0.937214563	0.541105179	0.795951402	FALSE	1.824164579	2	21	0.186009019	0.96116231	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.620368342	15	0.30144483	0.544326164	0.991826617	FALSE
			WT;Tbx1+/+	7	260.0642857	6.701019061	0.956735323	0.790286358	0.836773791	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-1.035608398	13	-0.535978266	0.319265725	0.991826617	FALSE
			WT;Tbx1+/flox	9	220.7333333	27.39387604	0.785658445	0.013990261	0.106253968	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-1.234919708	14	-0.622341035	0.237302673	0.991826617	FALSE
		30	PdgfraCre;Tbx1+/+	8	266.31875	13.4645697	0.780658686	0.017089995	0.106253968	FALSE	0.45938567	2	21	0.637870373	0.96116231	FALSE								PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.608023955	15	0.295446536	0.552267228	0.991826617	FALSE
			WT;Tbx1+/+	7	249.7428571	17.73063138	0.907420109	0.378299157	0.756598315	FALSE														PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.755605871	13	0.391063191	0.463357322	0.991826617	FALSE
			WT;Tbx1+/flox	9	249.4222222	23.24614531	0.741618208	0.004333717	0.078006906	FALSE														WT;Tbx1+/+	WT;Tbx1+/flox	-0.01042843	14	-0.005255435	0.991826617	0.991826617	FALSE

Figure 8D — Table 1

Figure	Measurement	Month	Time (min)	Genotype	# of samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects(FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects(FDR 5%)	LMM Source	F	df1	df2	Sig.	MWU Sample1	MWU Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure 8D	Total Margin Time (s)	2	5	oTbx1+/+	20	206.8325	12.8543966	0.930050585	0.154763372	0.371432094	FALSE	0.018345627	1	28	0.893228286	0.893228286	FALSE	Corrected Model	6.617692766	11	168	3.86918E-09	oTbx1+/-	oTbx1+/+	95	0.040160966	0.845778642	0.845778642	FALSE
				oTbx1+/-	10	214.465	18.6721823	0.924244799	0.393721373	0.472465648	FALSE							Genotype	0.089121278	1	168	0.765666179							
			10	oTbx1+/+	20	189.3575	12.22481583	0.94401407	0.285201157	0.427801736	FALSE	0.528505896	1	28	0.473270499	0.893228286	FALSE	Time	11.39367746	5	168	1.76945E-09	oTbx1+/-	oTbx1+/+	91	0.07228974	0.713163831	0.845778642	FALSE
				oTbx1+/-	10	198.145	15.69639635	0.863931639	0.08488746	0.25466238	FALSE							Genotype * Time	0.987985386	5	168	0.426725453							
			15	oTbx1+/+	20	219.36	12.95180672	0.943204905	0.275461326	0.427801736	FALSE	0.293845397	1	28	0.59205509	0.893228286	FALSE						oTbx1+/-	oTbx1+/+	64	-0.289158958	0.11962883	0.717772982	FALSE
				oTbx1+/-	10	192.6	16.68604041	0.922021218	0.374121624	0.472465648	FALSE																		
			20	oTbx1+/+	20	225.285	15.17396884	0.870559134	0.012018056	0.048072225	TRUE	0.875553509	1	28	0.357423105	0.893228286	FALSE						oTbx1+/-	oTbx1+/+	87	-0.104418513	0.588437383	0.845778642	FALSE
				oTbx1+/-	10	219.055	16.35034395	0.973597897	0.922052465	0.922052465	FALSE																		
			25	oTbx1+/+	20	243.54	12.86533725	0.847368866	0.004817268	0.028903611	TRUE	0.13461714	1	28	0.716450392	0.893228286	FALSE						oTbx1+/-	oTbx1+/+	85	-0.120482899	0.529969714	0.845778642	FALSE
				oTbx1+/-	10	240.37	14.0183693	0.908087267	0.268109319	0.427801736	FALSE																		
			30	oTbx1+/+	20	258.69	10.77157297	0.763194126	0.000256532	0.003078381	TRUE	0.021812291	1	28	0.883645943	0.893228286	FALSE						oTbx1+/-	oTbx1+/+	77	-0.184740446	0.328175706	0.845778642	FALSE
				oTbx1+/-	10	246.795	13.5509685	0.941920819	0.574578388	0.626182787	FALSE																		

Table 2 — Total Margin Time (s), 3 genotypes

Descriptive / Shapiro-Wilk / Levene:

Time (min)	Genotype	# samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj	SW Rejects	Levene Statistic	df1	df2	Levene p_value	Levene p_adj	Levene Rejects
5	PdgfraCre;Tbx1+/+	8	215.05	20.7141213	0.914114559	0.383949912	0.589193044	FALSE	2.353529985	2	17	0.125229683	0.324892303	FALSE
	WT;Tbx1+/+	6	155.8916667	13.65221238	0.920982548	0.512465635	0.658884388	FALSE						
	WT;Tbx1+/flox	6	246.8166667	15.77535349	0.902472395	0.388750333	0.589193044	FALSE						
10	PdgfraCre;Tbx1+/+	8	192.06875	20.35318275	0.869602244	0.149385243	0.537786875	FALSE	0.703121363	2	17	0.508876155	0.610651385	FALSE
	WT;Tbx1+/+	6	149.775	19.35028747	0.957394761	0.799454028	0.899385782	FALSE						
	WT;Tbx1+/flox	6	225.325	13.08559482	0.903135512	0.392795363	0.589193044	FALSE						
15	PdgfraCre;Tbx1+/+	8	210.59375	24.92307171	0.904635296	0.317826456	0.589193044	FALSE	2.176534062	2	17	0.144013566	0.324892303	FALSE
	WT;Tbx1+/+	6	186.4166667	12.48941997	0.975341879	0.926200355	0.96371726	FALSE						
	WT;Tbx1+/flox	6	263.99166667	13.16809539	0.926026806	0.549786816	0.659744179	FALSE						
20	PdgfraCre;Tbx1+/+	8	214.0375	28.49870415	0.834033939	0.065373269	0.392239615	FALSE	1.521800567	2	17	0.246616393	0.36992459	FALSE
	WT;Tbx1+/+	6	189.425	21.96758123	0.982597585	0.96371726	0.96371726	FALSE						
	WT;Tbx1+/flox	6	276.1416667	10.47147249	0.752640586	0.021139106	0.190251956	FALSE						
25	PdgfraCre;Tbx1+/+	8	231.325	19.64722127	0.925406735	0.47524764	0.658035195	FALSE	2.026321741	2	17	0.162446151	0.324892303	FALSE
	WT;Tbx1+/+	6	218.1083333	28.53045863	0.887468781	0.305136927	0.589193044	FALSE						
	WT;Tbx1+/flox	6	285.2583333	6.691492318	0.829589393	0.106631799	0.479843093	FALSE						
30	PdgfraCre;Tbx1+/+	8	258.3625	18.83312467	0.716307697	0.00343956	0.061912072	FALSE	0.476100124	2	17	0.629238835	0.629238835	FALSE
	WT;Tbx1+/+	6	232.4333333	21.96380229	0.884449517	0.290118872	0.589193044	FALSE						
	WT;Tbx1+/flox	6	285.3833333	6.690623621	0.871648175	0.232854122	0.589193044	FALSE						

Repeated Measures ANOVA:

Source		Type III Sum of Squares	df	Mean Square	F	Sig.
Within subjects						
Time	Greenhouse-Geisser	63034.55554	2.524553611	24968.59455	12.07218449	2.00728E-05
Time * Genotype	Greenhouse-Geisser	7497.715639	5.049107223	1484.958688	0.717969736	0.61474568
Error(Time)	Greenhouse-Geisser	88764.99903	42.9174114	2068.274766		
Between subjects						
Intercept		5924566.762	1	5924566.762	563.1055124	1.79933E-14
Genotype		102679.8172	2	51339.90862	4.879645051	0.021148604
Error		178861.0353	17	10521.23737		

Student t test:

Time	Sample1	Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
5	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-1.14816594	12	-0.620080478	0.273274002	0.351352288	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	2.204819261	12	1.190738493	0.047722279	0.107375127	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	4.358294506	10	2.516262506	0.00142495	0.014632215	TRUE
10	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-1.267237854	12	-0.684386661	0.229111121	0.343666681	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	1.46200151	12	0.789571057	0.169430463	0.277249849	FALSE
	WT;Tbx1+/+	WT;Tbx1+/+	3.234229662	10	1.867283366	0.008956536	0.040304412	TRUE
15	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	-1.712823843	12	-0.925030599	0.1124384	0.20238912	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.780634653	12	0.421590897	0.450142746	0.506410589	FALSE
	WT;Tbx1+/flox		4.274352212	10	2.4677984	0.001625802	0.014632215	TRUE
20	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-1.803787117	12	-0.974156382	0.09640862	0.192817241	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	0.644726307	12	0.348192001	0.531236864	0.562486091	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	3.563351857	10	2.057302154	0.005152424	0.030914546	TRUE
25	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-1.232996056	12	0.041448022	0.107375127		FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	0.395110604	12	0.213384114	0.699694601	0.699694601	FALSE
	WT;Tbx1+/+	WT;Tbx1+/+	2.291444377	10	1.322966028	0.044903228	0.143166836	FALSE
30	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-1.190208127	12	-0.642785854	0.256978702	0.351352288	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	0.897571615	12	0.484744074	0.387067972	0.107375127	FALSE
	WT;Tbx1+/+	WT;Tbx1+/+	2.306159612	10	1.331461873	0.043792782	0.107375127	FALSE

Table 3 — Total Margin Time (s), 4 genotypes

Descriptive / Shapiro-Wilk / Levene:

Time (min)	Genotype	# samples	Average	s.e.m	SW Statistic	SW p_value	SW p_adj	SW Rejects	Levene Statistic	df1	df2	Levene p_value	Levene p_adj	Levene Rejects
5	PdgfraCre;Tbx1+/+	8	215.05	20.7141213	0.914114559	0.383949912	0.59058206	FALSE	2.021947392	3	26	0.13547732	0.485523172	FALSE
	PdgfraCre;Tbx1+/flox	10	214.465	18.6721823	0.924244799	0.393721373	0.59058206	FALSE						
	WT;Tbx1+/+	6	155.8916667	13.65221238	0.920982548	0.512465635	0.683287514	FALSE						
	WT;Tbx1+/flox	6	246.8166667	15.77535349	0.902472395	0.388750333	0.59058206	FALSE						
10	PdgfraCre;Tbx1+/+	8	192.06875	20.35318275	0.869602244	0.149385243	0.59058206	FALSE	0.46733694	3	26	0.707587378	0.707587378	FALSE
	PdgfraCre;Tbx1+/flox	10	198.145	15.69639635	0.863931639	0.08488746	0.50932476	FALSE						
	WT;Tbx1+/+	6	149.775	19.35028747	0.957394761	0.799454028	0.913661747	FALSE						
	WT;Tbx1+/flox	6	225.325	13.08559482	0.903135512	0.392795363	0.59058206	FALSE						
15	PdgfraCre;Tbx1+/+	8	210.59375	24.92307171	0.904635296	0.317826456	0.59058206	FALSE	1.366979197	3	26	0.274814022	0.500050269	FALSE
	PdgfraCre;Tbx1+/flox	10	192.6	16.68604041	0.922021218	0.374121624	0.59058206	FALSE						
	WT;Tbx1+/+	6	186.4166667	12.48941997	0.975341879	0.926200355	0.96371726	FALSE						
	WT;Tbx1+/flox	6	263.99166667	13.16809539	0.926026806	0.549786816	0.689494065	FALSE						
20	PdgfraCre;Tbx1+/+	8	214.0375	28.49870415	0.834033939	0.065373269	0.50932476	FALSE	1.188722003	3	26	0.333366846	0.500050269	FALSE
	PdgfraCre;Tbx1+/flox	10	219.055	16.35034395	0.973597897	0.922052465	0.96371726	FALSE						
	WT;Tbx1+/+	6	189.425	21.96758123	0.982597585	0.96371726	0.96371726	FALSE						
	WT;Tbx1+/flox	6	276.1416667	10.47147249	0.752640586	0.021139106	0.253669274	FALSE						
25	PdgfraCre;Tbx1+/+	8	231.325	19.64722127	0.925406735	0.47524764	0.670937845	FALSE	1.856257447	3	26	0.161841057	0.485523172	FALSE
	PdgfraCre;Tbx1+/flox	10	240.37	14.0183693	0.908087267	0.268109319	0.59058206	FALSE						
	WT;Tbx1+/+	6	218.1083333	28.53045863	0.887468781	0.305136927	0.59058206	FALSE						
	WT;Tbx1+/flox	6	285.2583333	6.691492318	0.829589393	0.106631799	0.511832633	FALSE						
30	PdgfraCre;Tbx1+/+	8	258.3625	18.83312467	0.716307697	0.00343956	0.082549429	FALSE	0.505235134	3	26	0.682057595	0.707587378	FALSE
	PdgfraCre;Tbx1+/flox	10	246.795	13.5509685	0.941920819	0.574578388	0.689494065	FALSE						
	WT;Tbx1+/+	6	232.4333333	21.96380229	0.884449517	0.290118872	0.59058206	FALSE						
	WT;Tbx1+/flox	6	285.3833333	6.690623621	0.871648175	0.232854122	0.59058206	FALSE						

Repeated Measures ANOVA:

Source		Type III Sum of Squares	df	Mean Square	F	Sig.
Within subject						
Time	Greenhouse-Geisser	80940.86107	2.72028392	29754.56366	13.33360251	1.23598E-06
Time * Genotype	Greenhouse-Geisser	13331.91631	8.160851761	1633.642749	0.7320673	0.665388825
Error(Time)	Greenhouse-Geisser	157831.4927	70.72738193	2231.547222		
Between Subject						
Intercept		8537113.515	1	8537113.515	899.8162227	1.06132E-21
Genotype		103791.7875	3	34597.2625	3.646569534	0.025572586
Error		246678.0947	26	9487.619026		

Student t test:

Time	Sample1	Sample2	Statistic	dof	cohens_d	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
5	PdgfraCre;Tbx1+/+	PdgfraCre;Tbx1+/flox	-0.020957687	16	-0.009941104	0.983538546	0.983538546	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-1.189356816	14	-0.614181219	0.254080479	0.440534918	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	2.207266313	14	1.139827423	0.044489921	0.143166836	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	-1.14816594	12	-0.620080478	0.273274002	0.447175639	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/+	2.204819261	12	1.190738493	0.047722279	0.143166836	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	4.358294506	10	2.516262506	0.00142495	0.02926443	TRUE
10	PdgfraCre;Tbx1+/+	PdgfraCre;Tbx1+/flox	0.240552709	16	0.114104169	0.81295694	0.860777937	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-1.19168395	14	-0.615382946	0.253195562	0.440534918	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	1.917509151	14	0.990197551	0.075806824	0.194931833	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	-1.267237854	12	-0.684386661	0.229111121	0.440534918	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/+	1.46200151	12	0.789571057	0.169430463	0.358793922	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	3.234229662	10	1.867283366	0.008956536	0.072454791	FALSE
15	PdgfraCre;Tbx1+/+	PdgfraCre;Tbx1+/flox	-0.620273775	16	-0.294221685	0.543811849	0.699186664	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-2.973663831	14	-1.535593399	0.010063165	0.074454791	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.259805567	14	0.134163018	0.798795484	0.860777937	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	-1.712823843	12	-0.925030599	0.1124384	0.252986399	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/+	0.780634653	12	0.421590897	0.450142746	0.623274571	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	4.274352212	10	2.4677984	0.001625802	0.02926443	TRUE
20	PdgfraCre;Tbx1+/+	PdgfraCre;Tbx1+/flox	0.160447456	16	0.076106911	0.874536855	0.899523622	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-2.501145356	14	-1.291585908	0.025410373	0.143166836	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	1.09363096	14	0.5647486	0.292577527	0.457947434	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	-1.803787117	12	-0.974156382	0.09640862	0.231380689	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/+	0.644726307	12	0.348192001	0.531236864	0.699186664	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	3.563351857	10	2.057302154	0.005152424	0.061829092	FALSE
25	PdgfraCre;Tbx1+/+	PdgfraCre;Tbx1+/flox	0.384737119	16	0.182496839	0.705498097	0.79368536	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-2.35775279	14	-1.217538305	0.033465057	0.143166836	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.786078068	14	0.405928969	0.444923628	0.623274571	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	-1.232996056	12	0.041448022	0.699694601	0.79368536	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/+	0.395110604	12	0.213384114	0.699694601	0.79368536	FALSE
	WT;Tbx1+/+	WT;Tbx1+/flox	2.291444377	10	1.322966028	0.044903228	0.143166836	FALSE
30	PdgfraCre;Tbx1+/+	PdgfraCre;Tbx1+/flox	-0.51135554	16	-0.24255723	0.616086234	0.739303481	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	-2.091602839	14	-1.080099061	0.055183309	0.152815317	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/+	0.591033005	14	0.305208131	0.563919383	0.700037854	FALSE
	PdgfraCre;Tbx1+/flox	WT;Tbx1+/flox	-1.190208127	12	-0.642785854	0.256978702	0.440534918	FALSE
	PdgfraCre;Tbx1+/+	WT;Tbx1+/flox	0.897571615	12	0.484744074	0.387067972	0.580601957	FALSE
	WT;Tbx1+/flox	WT;Tbx1+/+	2.306159612	10	1.331461873	0.043792782	0.143166836	FALSE

Figure	Measurement	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk test		Levene test				Linear Mixed Model					Measurement	Fisher's r-to-Z transformation					
						Statistic	p_value	Statistic	df1	df2	p_value	factor	F	df1	df2	p_value		N	Pearson's r	p_value (correlation)	Z transformed correlation	Z statistic	p_value (Z statistic)
Figure 9B	g-ratio	oTbx1+/+	4423	0.742132168	0.001178885	0.983206272	1.62E-22	1.670711318	1	6738	0.196207608	Genotype	0.055858571	1	6736	0.813172794	g-ratio vs Axon Diameter (oTbx1+/+)	4423	0.626809	0.00E+00	0.736142587	-1.466330096	0.14255838
		oTbx1+/-	2317	0.736721572	0.001594201	0.98563242	1.56E-14					Axon Diameter	4117.938128	1	6736	0	g-ratio vs Axon Diameter (oTbx1+/-)	2317	0.649115	2.53E-277	0.773767499		
												Genotype:Axon Diameter	1.497334759	1	6736	0.22112471							

Figure 9C — Axon Number

Figure	Measurement	Axon Diameter	Genotype	# of samples	Average	s.e.m	SW statistic	SW p_value	SW p_adj(FDR BH)	SW Rejects (FDR 5%)	Levene Statistic	df1	df2	Levene p_value	Levene p_adj(FDR BH)	Levene Rejects (FDR 5%)	LMM (session random) Source	F	df1	df2	Sig.	LMM (mouse+session random) Source	F	df1	df2	Sig.	MW Statistic	MW R (Z score)	MW p_value	MW p_adj(FDR BH)	MW Rejects (FDR 5%)	MW (300-1400nm) Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects (FDR 5%)
Figure 9C	Axon Number	200	oTbx1+/+	145	0.131034483	0.035632184	0.341078043	8.80283E-23	5.47732E-22	TRUE	0.036618062	1	210	0.848429647	0.961107197	FALSE	Corrected Model	133.9074722	55	5880	0	Corrected Model	133.7784885	55	5880	0	4856	-0.002248086	0.996365794	0.996365794	FALSE					
			oTbx1+/-	67	0.119402985	0.045226142	0.358457863	1.42881E-15	3.63697E-15	TRUE							genotype	7.655858881	1	5880	0.005676512	genotype	0.561757486	1	5880	0.453582974										
		300	oTbx1+/+	145	0.662068966	0.111254037	0.56666255	6.46568E-19	2.78522E-18	TRUE	8.05053552	1	210	0.004994777	0.139853743	FALSE	binned_Axon_Diameter	246.5223833	27	5880	0	binned_Axon_Diameter	246.5223833	27	5880	0	3574.5	-0.212196465	0.00037454	0.00524356	TRUE	3574.5	-0.212196465	0.00037454	0.00224724	TRUE
			oTbx1+/-	67	1.313432836	0.207512627	0.771918416	7.85361E-09	1.25658E-08	TRUE							genotype * binned_Axon_Diameter	6.336275377	27	5880	0	genotype * binned_Axon_Diameter	6.336275377	27	5880	0										
		400	oTbx1+/+	145	1.8	0.180622593	0.749555611	5.68057E-13	1.1361E-12	TRUE	0.146243949	1	210	0.70253768	0.961107197	FALSE											2713	-0.354680685	1.48147E-07	4.14811E-06	TRUE	2713	-0.354680685	1.48147E-07	1.77776E-06	TRUE
			oTbx1+/-	67	3.268656716	0.268656716	0.922051668	0.000442592	0.000495703	TRUE																										
		500	oTbx1+/+	145	3.04137931	0.201497236	0.927050591	9.02246E-07	1.29553E-06	TRUE	0.171565279	1	210	0.679147179	0.961107197	FALSE											3448.5	-0.233035713	0.00062617	0.00584425	TRUE	3448.5	-0.233035713	0.00062617	0.002504679	TRUE
			oTbx1+/-	67	4.253731343	0.309171785	0.915763259	0.000238829	0.000278634	TRUE																										
		600	oTbx1+/+	145	4.020689655	0.241349494	0.9389707745	6.20047E-06	8.26729E-06	TRUE	1.592773869	1	210	0.20833013	0.729155453	FALSE											4021	-0.138349449	0.042188689	0.147660412	FALSE	4021	-0.138349449	0.042188689	0.072323467	FALSE
			oTbx1+/-	67	5.059701493	0.405123562	0.891388595	2.6243E-05	3.26579E-05	TRUE																										
		700	oTbx1+/+	145	3.806896552	0.19605705	0.943566203	1.3749E-05	1.79057E-05	TRUE	0.742076712	1	210	0.389978983	0.880618111	FALSE											3861.5	-0.16472929	0.015562111	0.087147822	FALSE	3861.5	-0.16472929	0.015562111	0.037349066	TRUE
			oTbx1+/-	67	4.865671642	0.359591407	0.867092669	3.69424E-06	5.04579E-06	TRUE																										
		800	oTbx1+/+	145	3.593103448	0.176540081	0.946822121	1.67654E-05	2.13378E-05	TRUE	0.310912228	1	210	0.577714827	0.951530303	FALSE											4841	0.002728949	0.968922578	0.996365794	FALSE	4841	0.002728949	0.968922578	0.968922578	FALSE
			oTbx1+/-	67	3.671641791	0.28484093	0.922504842	0.000463096	0.000508498	TRUE																										
		900	oTbx1+/+	145	2.620689655	0.160658656	0.954623818	0.000107914	0.000128578	TRUE	0.7278997	1	210	0.394536889	0.880618111	FALSE											4439	0.069216072	0.308357041	0.411142721	FALSE	4439	0.069216072	0.308357041	0.411142721	FALSE
			oTbx1+/-	67	3.014925373	0.261758774	0.920220673	0.000369011	0.000421727	TRUE																										
		1000	oTbx1+/+	145	2.668965517	0.155380483	0.9350098397	3.20007E-06	4.4801E-06	TRUE	0.684857725	1	210	0.408858409	0.880618111	FALSE											4552.5	0.05044421	0.456622851	0.741419141	FALSE	4552.5	0.05044421	0.456622851	0.547947421	FALSE
			oTbx1+/-	67	2.432835821	0.207528892	0.930607498	0.001061447	0.001143097	TRUE																										
		1100	oTbx1+/+	145	1.972413793	0.103086853	0.925580502	7.20232E-07	1.09008E-06	TRUE	6.478227093	1	210	0.011638138	0.162933933	FALSE											4436	0.069712245	0.298569389	0.626862409	FALSE	4436	0.069712245	0.298569389	0.411142721	FALSE
			oTbx1+/-	67	1.880597015	0.19667336	0.894105256	3.31308E-05	4.03331E-05	TRUE																										
		1200	oTbx1+/+	145	1.448275862	0.097345533	0.886540353	3.87729E-09	6.38613E-09	TRUE	3.69443342	1	210	0.055947631	0.391633419	FALSE											3707	0.190282177	0.003824339	0.026770374	TRUE	3707	0.190282177	0.003824339	0.0114730017	TRUE
			oTbx1+/-	67	0.970149254	0.114467183	0.829086244	2.4814E-07	3.85995E-07	TRUE																										
		1300	oTbx1+/+	145	1.144827586	0.093351451	0.832913399	1.46897E-11	2.65362E-11	TRUE	0.321086438	1	210	0.571560581	0.951530303	FALSE											4732.5	0.020673857	0.752577581	0.996365794	FALSE	4732.5	0.020673857	0.752577581	0.820993725	FALSE
			oTbx1+/-	67	1.029850746	0.112474228	0.847671866	8.85128E-07	1.29553E-06	TRUE																										
		1400	oTbx1+/+	145	0.910344828	0.080656574	0.778914571	1.6278E-13	3.37617E-13	TRUE	0.011174349	1	210	0.915914309	0.961107197	FALSE											3984	0.144468911	0.023473363	0.093893452	FALSE	3984	0.144468911	0.023473363	0.046946726	TRUE
			oTbx1+/-	67	0.656716418	0.113638558	0.720096111	5.37756E-10	9.12555E-10	TRUE																										
		1500	oTbx1+/+	145	0.537931034	0.067820962	0.664089799	9.36116E-17	2.66433E-16	TRUE	0.241876622	1	210	0.623367675	0.961107197	FALSE											4530.5	0.054082809	0.358207091	0.626862409	FALSE					
			oTbx1+/-	67	0.47761194	0.104961083	0.599906087	3.1492E-12	6.08121E-12	TRUE																										
		1600	oTbx1+/+	145	0.372413793	0.050802777	0.625598192	1.17036E-17	3.8677E-17	TRUE	6.61207E-05	1	210	0.993519823	0.993519823	FALSE											4780.5	-0.012735096	0.820278888	0.996365794	FALSE					
			oTbx1+/-	67	0.373134328	0.070006205	0.64398396	1.80164E-11	3.15287E-11	TRUE																										
		1700	oTbx1+/+	145	0.317241379	0.042326328	0.607726872	4.70168E-18	1.7553E-17	TRUE	0.107856086	1	210	0.742925516	0.961107197	FALSE											4852	0.00090965	0.987880394	0.996365794	FALSE					
			oTbx1+/-	67	0.343283582	0.072283423	0.60768044	4.24081E-12	7.91618E-12	TRUE																										
		1800	oTbx1+/+	145	0.262068966	0.045926697	0.525665522	1.0219E-19	4.76887E-19	TRUE	1.66949102	1	210	0.197746792	0.729155453	FALSE											4529	0.054330895	0.249807126	0.626862409	FALSE					
			oTbx1+/-	67	0.164179104	0.050313423	0.439795375	1.44797E-14	3.24346E-14	TRUE																										
		1900	oTbx1+/+	145	0.124137931	0.029169144	0.382323027	3.6823E-22	2.06209E-21	TRUE	0.981731842	1	210	0.322911885	0.880618111	FALSE											4626.5	-0.038205287	0.34420146	0.626862409	FALSE					
			oTbx1+/-	67	0.119104478	0.051176825	0.462751687	2.90876E-14	6.26502E-14	TRUE																										
		2000	oTbx1+/+	145	0.151724138	0.032944622	0.41921103	1.4113E-21	7.1848E-21	TRUE	0.095627397	1	210	0.757448003	0.961107197	FALSE											4831	0.004382858	0.916278431	0.996365794	FALSE					
			oTbx1+/-	67	0.134328358	0.041974785	0.403025091	4.94005E-15	1.15268E-14	TRUE																										
		2100	oTbx1+/+	145	0.096551724	0.026486709	0.326505065	5.39777E-23	3.77844E-22	TRUE	0.561931824	1	210	0.454321862	0.908643725	FALSE											4710.5	-0.024312455	0.495563538	0.741419141	FALSE					
			oTbx1+/-	67	0.134328358	0.047054932	0.38790673	3.21939E-15	7.83851E-15	TRUE																										
		2200	oTbx1+/+	145	0.082758621	0.024958679	0.294410527	1.89092E-23	1.51274E-22	TRUE	0.036086921	1	210	0.8495196	0.961107197	FALSE											4849	0.001405822	0.966419652	0.996365794	FALSE					
			oTbx1+/-	67	0.074626866	0.032347005	0.289320052	2.35055E-16	6.26812E-16	TRUE																										
		2300	oTbx1+/+	145	0.027586207	0.013648669	0.151343524	2.64901E-25	2.96689E-24	TRUE	1.882778001	1	210	0.171482662	0.729155453	FALSE											4723.5	0.022162374	0.17251315	0.483036819	FALSE					
			oTbx1+/-	67				1	1	FALSE																										
		2400	oTbx1+/+	145	0.055172414	0.021395989	0.218779445	1.83508E-24	1.71274E-23	TRUE	0.012885239	1	210	0.90973217	0.961107197	FALSE											4842.5	0.002480863	0.924256179	0.996365794	FALSE					
			oTbx1+/-	67	0.059701493	0.036094986	0.207609415	3.26996E-17	1.01732E-16	TRUE																										
		2500	oTbx1+/+	145	0.027586207	0.013648669	0.151343524	2.64901E-25	2.96689E-24	TRUE	5.644026015	1	210	0.018414322	0.171867004	FALSE											4846.5	-0.001819299	0.929852964	0.996365794	FALSE					
			oTbx1+/-	67	0.029850746	0.020947148	0.162513554	1.17412E-17	3.8677E-17	TRUE																										
		2600	oTbx1+/+	145	0.006896552	0.006896552	0.056594372	2.12105E-26	5.93893E-25	TRUE	2.176243568	1	210	0.14165388	0.729155453	FALSE											4601	-0.042422754	0.019005936	0.088694366	FALSE					
			oTbx1+/-	67	0.059701493	0.02916443	0.252678573	9.51548E-17	2.66433E-16	TRUE																										
		2700	oTbx1+/+	145			1	1		FALSE	0.928222721	1	210	0.336433275	0.880618111	FALSE											4785	-0.011990837	0.144023867	0.448074253	FALSE					
			oTbx1+/-	67	0.014925373	0.014925373	0.10137707	3.12269E-18	1.24907E-17	TRUE																										
		2800	oTbx1+/+	145	0.013793103	0.009719273	0.093628466	5.5507E-26	1.03613E-24	TRUE																		4790.5	0.011081187	0.338862118	0.626862409	FALSE					
			oTbx1+/-	67						FALSE																										
		2900	oTbx1+/+	145	0.006896552	0.006896552	0.056594372	2.12105E-26	5.93893E-25	TRUE	0.460888365	1	210	0.497956586	0.929518961	FALSE											4824	0.005540594	0.503105846	0.741419141	FALSE					
			oTbx1+/-	67						FALSE																										

Measurement	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk Statistic	p_value	Levene Statistic	df1	df2	p_value	Mann-Whitney U Statistic	p_value	Alternative (H1)
Axon Diameter (nm)	oTbx1+/+	4423	913.5795028	5.511747597	0.929378211	1.60238E-41	2.629408218	1	6738	0.104947594	5790975.5	7.48066E-19	greater
	oTbx1+/-	2317	845.8037375	7.58800961	0.897995234	9.79998E-37							(oTbx1+/+ > oTbx1+/-)

Figure	# of samples	Chi squared test			Kolmogorov-Smirnov test	
		Statistic	dof	p_value	Distance	p_value
Figure 9D	6740	109.7085231	27	6.22E-12	0.108947998	3.59E-16

Figure	Measurement	PND	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk test				Levene test						Mann-Witney U test						
							Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Figure S1	Weight	P8	oTbx1+/+	27	4.6	0.125064086	0.922948078	0.046515575	0.186062298	FALSE	0.639894198	1	36	0.428996273	0.428996273	FALSE	oTbx1+/+	oTbx1+/-	140	0.044382036	0.799797921	0.91732983	FALSE
			oTbx1+/-	11	4.454545455	0.237931861	0.980904629	0.970981671	0.970981671	FALSE													
		P12	oTbx1+/+	30	6.54	0.155482194	0.986169448	0.955380174	0.970981671	FALSE	1.973777187	1	40	0.167766267	0.428996273	FALSE	oTbx1+/+	oTbx1+/-	153	0.115996334	0.465855659	0.91732983	FALSE
			oTbx1+/-	12	6.225	0.351861931	0.915390828	0.249993706	0.499987413	FALSE													
		1M	oTbx1+/+	27	16.15925926	0.252241194	0.863150358	0.002121269	0.016970151	TRUE	0.856989714	1	36	0.360747027	0.428996273	FALSE	oTbx1+/+	oTbx1+/-	135.5	-0.067878408	0.703523286	0.91732983	FALSE
			oTbx1+/-	11	16.37272727	0.447232075	0.942333972	0.548313562	0.80806296	FALSE													
		2M	oTbx1+/+	21	23.78571429	0.385837723	0.943028092	0.24996769	0.499987413	FALSE	1.523319488	1	29	0.227026429	0.428996273	FALSE	oTbx1+/+	oTbx1+/-	102	0.022769128	0.91732983	0.91732983	FALSE
			oTbx1+/-	10	23.53	0.736515822	0.94467026	0.60604722	0.80806296	FALSE													

Measurement	PND	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk test				Levene test						Mann-Witney U test						
						Statistic	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Statistic	df1	df2	p_value	p_adj(FDR BH)	Rejects(FDR 5%)	Sample1	Sample2	Statistic	R (Z score)	p_value	p_adj(FDR BH)	Rejects(FDR 5%)
Weight	P8	PdgfraCre;WT	8	4.35	0.205287255	0.866045239	0.137839188	0.413517564	FALSE	0.127476314	2	24	0.880906332	0.880906332	FALSE	PdgfraCre;WT	WT;Tbx1+/flox	18	-0.367573522	0.160528361	0.872562731	FALSE
		WT;Tbx1+/flox	8	4.7375	0.203485257	0.753919531	0.009024632	0.054147793	FALSE							PdgfraCre;WT	WT;WT	29.5	-0.274679021	0.272287052	0.872562731	FALSE
		WT;WT	11	4.681818182	0.225153758	0.91199385	0.257472583	0.617934199	FALSE							WT;Tbx1+/flox	WT;WT	40.5	-0.066301833	0.840385277	1	FALSE
	P12	PdgfraCre;WT	9	6.477777778	0.227777778	0.921953577	0.408659184	0.700558602	FALSE	1.743649374	2	27	0.194001478	0.776005912	FALSE	PdgfraCre;WT	WT;Tbx1+/flox	30.5	-0.20813018	0.436281366	0.872562731	FALSE
		WT;Tbx1+/flox	9	6.788888889	0.217590623	0.947010301	0.657291765	0.876389021	FALSE							PdgfraCre;WT	WT;WT	52.5	0.023262105	0.97227231	1	FALSE
		WT;WT	12	6.4	0.31454633	0.994973684	0.999999306	0.999999306	FALSE							WT;Tbx1+/flox	WT;WT	42	0.186096842	0.422107304	0.872562731	FALSE
	1M	PdgfraCre;WT	8	16.3625	0.287189273	0.934767327	0.560437988	0.840656982	FALSE	0.176982173	2	24	0.838878203	0.880906332	FALSE	PdgfraCre;WT	WT;Tbx1+/flox	31	0.116690007	0.672974085	1	FALSE
		WT;Tbx1+/flox	9	15.78888889	0.590537525	0.763341553	0.007737799	0.054147793	FALSE							PdgfraCre;WT	WT;WT	40	0	1	1	FALSE
		WT;WT	10	16.33	0.381240665	0.925384317	0.404055022	0.700558602	FALSE							WT;Tbx1+/flox	WT;WT	42	-0.056195149	0.842105263	1	FALSE
	2M	PdgfraCre;WT	8	24.3875	0.578309575	0.794950867	0.025269633	0.101078534	FALSE	0.889390233	2	18	0.428210148	0.856420296	FALSE	PdgfraCre;WT	WT;Tbx1+/flox	26	0.05976143	0.866511267	1	FALSE
		WT;Tbx1+/flox	7	23.91428571	0.525279323	0.958737507	0.807832827	0.969399392	FALSE							PdgfraCre;WT	WT;WT	14.5	0.327781141	0.382384282	0.872562731	FALSE
		WT;WT	6	22.83333333	0.889818958	0.974279891	0.919850787	0.999999306	FALSE							WT;Tbx1+/flox	WT;WT	14	0.277350098	0.365967366	0.872562731	FALSE

*including P11 and P13

**At the beginning of the behavioral battery (RSI) at each age

Figure	Measurement	Genotype	# of Samples	Average	s.e.m	Shapiro-Wilk test Statistic	p_value	Levene test Statistic	df1	df2	p_value	Linear Mixed Model factor	F	df1	df2	p_value	Measurement	N	Pearson's r	p_value (correlation)	Z transformed correlation	Z statistic	p_value (Z statistic)
Figure S2	myelin thickness (nm)	oTbx1+/+	4423	298.4267488	1.598178651	0.883548319	0.00E+00	18.9486008	1	6738	1.36273E-05	Genotype	0.007407354	1	6736	0.931416522	myelin thickness vs Axon Diameter (oTbx1+/+)	4423	0.309679	6.35E-99	0.320189803	-2.190862775	0.028461724
		oTbx1+/-	2317	281.3171899	1.943145919	0.89264971	1.85E-37					Axon Diameter	755.8280666	1	6736	0	myelin thickness vs Axon Diameter (oTbx1+/-)	2317	0.359582	1.15E-71	0.376405671		
												Genotype:Axon Diameter	0.05543768	1	6736	0.813865028							