A comprehensive analysis of flagellar mutations in *Vibrio cholerae*: Trade-off mechanisms between motility loss and host adaptability

Guozhong Chen [a,b,1], Zixin Qin [a,1], Fenxia Fan [c,1], Mei Luo [a,1], Hongou Wang [d], Baoshuai Xue [e], Shucheng Li [b], Shiyong Chen [f], Xiaoman Yang [a], Xiang Mao [g], Liwen Yi [a], Chunrong Yi [a], Wei Li [a, *], Xiaoyun Liu [d,h,**], Biao Kan [c, ***], Zhi Liu [a, *]

[a]Key Laboratory of Molecular Biophysics of the Ministry of Education/Department of Biotechnology, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China.
[b]State Key Laboratory of Microbial Technology, Shandong University, Qingdao, Shandong 266237, China.
[c]National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China.
[d]Department of Microbiology and Infectious Disease Center, NHC Key Laboratory of Medical Immunology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China.
[e]Key Laboratory of Zoonosis, Ministry of Education, College of Veterinary Medicine, Jilin University, Changchun, Jilin 130015, China.
[f] School of Marine Science and Engineering, Qingdao Agricultural University, Qingdao, Shandong 266109, China.
[g]Institute of Hygiene Toxicology, Wuhan Centre for Disease Prevention and Control, Wuhan, Hubei 430024, China.
[h]Department of Infectious Diseases, Peking University Third Hospital, Beijing 100191, China.

*Corresponding author: Key Laboratory of Molecular Biophysics of the Ministry of Education/Department of Biotechnology, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China.

**Department of Microbiology and Infectious Disease Center, NHC Key Laboratory of Medical Immunology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China.

Department of Infectious Diseases, Peking University Third Hospital, Beijing 100191, China.

***National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China.

E-mail addresses: hulwei_009@163.com (W. Li), xiaoyun.liu@bjmu.edu.cn (X. Liu), kanbiao@icdc.cn (B. Kan), zhiliu@hust.edu.cn (Z. Liu)

[1] These four authors contributed equally to this work.

Abstract

Cholera, caused by *Vibrio cholerae*, is characterized by severe watery diarrhea and poses a significant threat to global public health and the aquaculture industry. The flagellum of *V. cholerae* serves not only as a motility organelle for host seeking and colonization but also as a key environmental sensor and signal transducer, directly regulating virulence and biofilm formation. Despite its central role, the trade-off mechanisms between motility loss and enhanced host adaptability remain poorly understood. To address this, we analyzed 3,135 cholera-related samples and found that flagellar mutations in *V. cholerae* cluster primarily in genes such as *flhB*, *fliA*, *fliF*, *fliD*, and *fliM*. Using a flagellar mutant library, we observed that colonization ability was inversely correlated with FlgM secretion. Mechanistically, high intracellular FlgM inactivates the σ^{28} factor FliA, leading to transcriptional derepression of the quorum-sensing master regulator *hapR*. HapR in turn upregulates methionine metabolism and transport, thereby enhancing *V. cholerae* colonization in vivo. Through further validation with a uniquely constructed 6N-labeled mutant pool in an adult mouse model, we demonstrated that motility loss promotes environmental adaptability via the FlgM-FliA-HapR-methionine axis. Our study demonstrates that the secretion level of the flagellar regulator FlgM drives an adaptive shift in *V. cholerae* from motility to colonization within the host. This shift is mediated through quorum sensing and methionine metabolic reprogramming, revealing a novel mechanism that provides new insights into bacterial evolution and pathogenicity.

Keywords: *Vibrio cholerae*, flagellum, FlgM, quorum sensing, methionine metabolism

1. Introduction

Cholera, caused by *Vibrio cholerae*, is characterized by severe watery diarrhea that can be lethal. Infection occurs through the consumption of contaminated food or water. Since 1817, there have been seven global pandemics that resulted in massive human deaths, substantial economic losses and considerable healthcare burdens [1]. The WHO reports a large number of cholera cases or outbreaks, from 44 countries in 2022 to 45 countries in 2023, which poses a serious challenge of cholera prevention and control [2].

The life cycle of *V. cholerae* is closely linked to its pathogenic mechanism. Initially, the bacteria enter the host via contaminated water, survive gastric acid, and colonize the small intestine. Subsequently, their flagella facilitate penetration of the mucus layer, allowing access to the intestinal epithelial surface. As infection progresses, *V. cholerae* forms biofilms and employs quorum-sensing regulation. Upon reaching high cell density, the bacteria detach from the epithelial surface, enter the intestinal lumen, and are ultimately excreted in host feces. They then return to aquatic environments, thereby completing the transmission cycle [3, 4].

V. cholerae is a Gram-negative bacterium with a single polar flagellum, by which it
swims rapidly and this motility contributes to quick transmission among different
habitats [5,6]. As the propeller of *V. cholerae* movement, the flagellum is a
sophisticated device composed of many different structural and functional components.
Its biosynthesis involves expression of proteins involved in a series of fine and accurate
regulations, acting on transcriptional, translational, and post-translational levels [7].
Transcriptional regulation of *V. cholerae* genes has been well-studied, and is built upon
a four-tiered transcriptional hierarchy [3]. The flagellum of *V. cholerae* is also involved
in many other biological activities, such as promoting adherence [8], influencing the
infectivity [9], triggering c-di-GMP production [10], and sensing mucus layer
mechanical   attraction   [4].   Interestingly,   a   high   number   of
single nucleotide polymorphisms (SNPs) were detected in the Haiti and Nepal isolates
of the 7th cholera pandemic, with 7 out of 54 high quality SNPs belonging to flagellum-
associated genes [11]. Similarly, a large number of mutants with impaired motility have
been isolated from the laboratory [12-17], the natural environment [18-20], and clinical
samples [21]. However, the relationship between motility loss and host adaptability in
V. cholerae remains unclear.
Here, we analyzed 3,135 cholera-related samples and identified a recurrent pattern
of flagellar mutations in *V. cholerae*, with notable clustering in genes
including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM*. Screening a flagellar mutant library revealed
an inverse correlation between bacterial colonization capacity and FlgM secretion.
Mechanistically, elevated intracellular FlgM inhibits the σ^{28} factor FliA, which
derepresses transcription of the quorum-sensing master regulator *hapR*. Subsequent
upregulation of methionine metabolism and transport by HapR enhances *V.*
cholerae colonization in vivo. Using a uniquely constructed 6N-labeled mutant pool in
an adult mouse model, we further confirmed that loss of motility promotes host
adaptability through the FlgM-FliA-HapR-methionine regulatory axis. Together, our
data will shed light on trade-off mechanisms between motility loss and host adaptability.

2.   Materials and methods

2.1 In silico analysis flagellar gene mutations in V. cholerae

A total of 3,135 samples in Table S1 were retrieved from the NCBI SRA database by
searching for "cholera" or "Vibrio cholerae" in combination with metagenomics-
related keywords. These valid sequencing runs were selected for subsequent analysis
of flagellar gene mutations. For each dataset, raw reads were quality-filtered using fastp
v0.23.2 [22] and assembled with metaWRAP v1.3.2 [23]. Resulting contigs were
binned and taxonomically classified using GTDB-Tk v2.4.0 [24]. Bins identified as *V.*
cholerae were retained for protein prediction with Prodigal v1.2.6 [25]. A custom
reference library of 51 flagellar proteins in Table S2 was constructed based on the *V.*
cholerae N16961 genome. Flagellar mutations across 1258 assembled *V. cholerae* bins
were then identified in Table S3 and quantified using DIAMOND BLASTP v0.9.30-3
[26]. As controls, the *hapR* gene (which carries a frameshift mutation in N16961)

served as a positive control, while the housekeeping
genes *asd* (VC2107), *mdh* (VC0432), and *dnaE* (VC2245) were used as negative
controls, respectively [27,28].

2.2 Bacterial strains, plasmids, and growth conditions

All *V. cholerae* mutants were derived from strain E1 Tor C6706 and cultured in Luria-
Bertani (LB) medium, M9 minimal medium containing 2 mM $MgSO_4$, 0.1 mM $CaCl_2$
and 0.2% casein acid hydrolysate, or defined medium containing appropriate antibiotics
(streptomycin 100 μg mL^{-1}, kanamycin 50 μg mL^{-1}, ampicillin 100 μg mL^{-1},
chloramphenicol 25 μg mL^{-1} for *E. coli* and 1.0 μg mL^{-1} for *V. cholerae*) at 37 °C unless
otherwise noted. This defined medium was prepared with the omission of Met and
supplementation with 0.5% glucose [29]. In-frame deletion mutants were constructed
by cloning the regions flanking a target gene into suicide vector pWM91 that contains
a *sacB* counter-selectable marker [30]. Complementation of *flgM* was achieved by
inserting P_{lac}-*flgM* into chromosomal *lacZ* using the suicide plasmid pJL1 [31].
Overexpression of *flgM* was obtained by cloning *flgM-his$_6$* downstream of the *tac*
promoter of pMal-c2x. Plasmid pBAD24 containing a pBAD-inducible promoter was
used to overexpress *flgM-his$_6$*, *flhB*, *flaA-his$_6$*, *fliA-flag*, and *hapR-flag*. The fragment
of pBAD-*fliA-flag* was cloned into pYM24 derived from pACYC117 [32], achieving
coexistence with the plasmids pMal-*flgM-his$_6$* and pBBR-*hapR-luxCDABE*. All
constructed mutants and plasmids were confirmed by sequencing. All bacterial strains,
primers and plasmids used in this study are listed in Table S4&5.

2.3 Motility assay

V. cholerae strains were incubated overnight at 37 °C for 12 h. Single colonies were
inoculated with sterile toothpicks into LB plates containing 0.3% agar, and these soft
agar plates were incubated at 37 °C for 8 h. Photographs were taken and analyzed for
colony width as a measure of motility using a gel imaging instrument (GenoSens 1860).

2.4 H_2O_2 killing experiments in vitro

Overnight cultures of flagellar mutants were 1:1,000 sub-cultured into M9 minimal
medium containing 2 mM $MgSO_4$, 0.1 mM $CaCl_2$ and 0.2% casein acid hydrolysate
and grown at 37°C, 200 rpm until mid-log phase. H_2O_2 was directly added into the
cultures at appropriate concentrations, followed by stationary incubation at 37°C for 30
min. Serial dilutions of the samples were plated on LB agar for enumeration of colony-
forming units (CFU) and the survival rate in presence of H_2O_2 was calculated by
comparing CFUs of H_2O_2-challenged and unchallenged samples [32].

2.5 Biofilm formation

Colonies grown on LB agar plates were resuspended in LB broth and grown to an
optical density at 600 nm (OD$_{600}$) of ~0.6. A 1:100 dilution was inoculated into 10 ×
75 mm borosilicate glass tubes containing 1 mL M9 minimal medium with 2 mM
MgSO$_4$, 0.1 mM CaCl$_2$ and 0.2% casein acid hydrolysate. Biofilms were allowed to
form by static incubation at 37 °C for 96 h. The biofilms were stained with crystal violet
and quantified with a microplate reader (Spark, Tecan) at 570 nm.

2.6 In vivo colonization assays

All animal experiments were carried out in strict accordance with the animal protocols
that were approved by the Ethical Committee of Huazhong University of Science and
Technology (Permit Number: SYXK (E) 2022-0094).
The streptomycin-treated adult mouse model was used to assess the competition ability
of mutants or complement strains during co-colonization with C6706 WT. ROS
resistance in the adult mice assay was performed as previously described [33]. For this,
five-week-old CD-1 mice were provided with drinking water with or without 10 g L^{-1}
of the antioxidant N-acetylcysteine for one week. Afterwards, 5 g L^{-1} streptomycin and
0.05 g L^{-1} aspartame were added to the drinking water. One day after streptomycin
treatment, the mice were administrated with 100 μL of 10% (w/v) NaHCO$_3$ followed
by 100 μL of a 1:1 mixture of WT and mutant (approximately 10^9 CFU for each strain
per mouse). Fecal pellets were collected from each mouse at 1, 3, and 5 days after
gavage, homogenized, serially diluted in LB and plated on LB agar containing
streptomycin and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal) to
quantify the bacterial loads. The competition index CI was calculated as the ratio of
mutant to WT colonies normalized with the input ratio.
Five-day-old CD-1 mice were transferred to an incubator and left to adapt for 2 h at
30 °C before inoculation. The mice were then administrated with a 50 μL of a 1:1
mixture of WT and mutant *V. cholerae* (approximately 10^6 CFU for each strain per
mouse) and returned to the 30 °C incubator. The animals were sacrificed 18 h after
gavage. Samples of the small intestine were removed and homogenized in 1.5 ml of LB,
serially diluted, and plated on LB agar containing streptomycin and X-Gal for
quantification of bacterial load. The CI was calculated as the ratio of mutant to WT
colonies normalized with the input ratio.

2.7 Growth curve determination and luminescence detection

Overnight LB cultures of targeted strains (with or without plasmid) were inoculated in
fresh medium at a ratio of 1:100 and grown with shaking to mid-log phase at 37 °C.
The bacteria were harvested by centrifugation, washed three times and resuspended to
an OD$_{600}$ of 1.0 with fresh pre-warmed medium (LB or M9), and then diluted 1:1,000
in 96-well microplates that were incubated at 37 °C under shaking. The growth kinetics
were determined by measuring OD$_{600}$ or OD$_{655}$ using a microplate reader (Spark, Tecan)
combined (where applicable) with luminescence detection. If necessary, 0.1 mM

isopropyl-β-D-thiogalactoside (IPTG) or 0.2% arabinose was added. Readouts were
performed every 30 min for 20 h. The luminescence assay was standardized at OD_{600}.
All experiments were performed in triplicate.

2.8 Transmission electron microscopy (TEM) imaging

Single colonies of target strains were inoculated into 50 μL of saline and gently mixed
with a clipped, sterile pipette tip. Five microliters of sample were added to a copper
supporting mesh. Excess liquid was removed after standing in a ventilated and clean
environment for 5 min. The mesh was stained with freshly prepared 1%
phosphotungstic acid for 2 min. Excessive dye was removed and the mesh was dried
for approximately 1 h prior to transmission electron microscopy (HT7700, HITACHI).

2.9 Secretion of FlgM

To investigate the secretion of FlgM, culture supernatant of the *V. cholerae* mutants
containing pBAD24-*flgM-his₆* was collected and the protein was detected according to
a previous method with slight modification [4]. All mutant sediment samples were
standardized with CFU (10^9). Proteins in the 400mL supernatant were concentrated
with 20% trichloroacetic acid (TCA). These, together with the proteins in the precipitate
and supernatant were separated by SDS-PAGE, transferred to a polyvinylidene fluoride
(PVDF) membrane. The proteins on the Western blot were detected with primary 6×
His-tagged rabbit antibodies at a dilution of 1:5,000, and as secondary antibody
horseradish peroxidase [HRP]-conjugated goat anti-rabbit immunoglobulin G was used
at a dilution of 1:10,000. Immunoblotting was performed using the Pierce ECL Western
Blotting Substrate Kit (Thermo Scientific).

2.10 Proteomic analysis

Overnight LB cultures were inoculated 1:100 in fresh LB medium and incubated until
OD_{600} reached 0.9 (approximately 10^9 CFU per sample). Then the culture was
centrifuged. The pellets were for bacterial proteome samples. For the analysis of the
bacterial secretome, the 400 mL of culture supernatant was filtered through a 0.22 μm
microporous membrane to remove intact bacteria. The proteins in the filtrate were
precipitated overnight at 4 °C with 20% TCA containing 0.5 mM sodium deoxycholate.
After centrifugation at 10,000 g for 1 h at 4 °C, the protein pellets were washed three
times with cold acetone for LC-MS/MS experiments, that were carried out as previously
described [32]. The secretome experiments were performed in duplicates, and bacterial
proteome samples were analyzed in triplicates. Proteins with Fold changes >1.5 and *P*<
0.05 were considered as significant differences.

2.11 Gene expression analysis via RT-PCR and qPCR

227 Overnight LB cultures were inoculated 1:100 in fresh LB medium containing
228 appropriate antibiotics or, if needed, 0.2% arabinose, and incubated at 37 °C. RNA for

real-time PCR (RT-PCR) was extracted using an RNA extraction kit. The RT-PCR reaction was performed using SYBR green fluorescent dye and CFX Connect Real-time Detection System (Bio-Rad). The genes *recA* or *secA* or 16S rRNA were used for normalization. Relative expression levels were calculated using the $2^{-\Delta\Delta CT}$ method. All experiments were performed in triplicate. For reverse transcription, approximately 400 ng of RNA was extracted using the HiScript II 1st Strand cDNA Synthesis Kit. The RT-qPCR primers used for detection of mRNA of specific genes are listed in Table S5. All primers were tested for PCR specificity and efficiency.

2.12 Protein purification

The proteins HapR-His$_6$ were expressed in *E. coli* BL21 (DE3) and purified using the pET system and BeyoGold™ His-tag Purification Resin (P2218, Beyotime).

2.13 EMSA

The purified recombinant proteins HapR containing 6×His-tags were used for EMSA. The promoters of their target genes were amplified by PCR and purified. These amplicons (50 ng) were incubated with 6× His-tagged protein in 20 μL containing 4μL of 5× binding buffer (50 mM Tris-HCl, 0.25 M KCl, 2.5 mM DTT, 5 mM MgCl$_2$, 0.25 mg mL^{-1} BSA, 2.5 mM EDTA, 1% glycerol, pH 8.3) at 25 °C for 30 min. DNA-protein complexes were analyzed by electrophoresis on a 5% native polyacrylamide gel at 4 °C and 80 V cm^{-1} for 1.5 h. After staining the gel in ethidium bromide for 10 min, a gel imaging system (GenoSens 1860) was used to produce images. The EMSA primers used in this study are listed in Table S5.

2.14 Synthesis of fluorescent monomers

FITC-D-Glucose was synthesized as previously described [34]. Briefly, D-glucosamine hydrochloride (77.6 mg, 0.36 mmol, 1.2 equiv) and Et3N (91.1 mg, 0.9 mmol, 3.0 equiv) were mixed in DMF (10 mL). After 1 h of stirring, fluorescein (116.8 mg, 0.3 mmol, 1.0 equiv) was added. The reaction mixture was stirred for 16 h at RT in the dark, after which the solid product was obtained by filtering and concentrated *in vacuo*.

2.15 Bacterial metabolic activity assay in vitro

Bacterial cultures in LB were harvested at OD$_{600}$ ~0.4 by centrifugation and the cell pellet was washed 3 times with Met-free MM medium. The cells were resuspended and incubated at 37°C for 30 minutes. A solution of 10 mL Met-free MM medium and 1 μL of FITC-D-Glu was added and 500 μL of this was added to transwell wells and 200 μL of this was added to the transwell chambers. To each well, 100 μL of bacterial suspension was added and the plates were incubated for 30 min at 37 °C in the dark. The supernatant was washed off with PBS buffer and then assayed by Flow Cytometry.

2.16 Bacterial metabolic activity assay in vivo

Bacterial metabolic activity was assayed in vivo as previously described with following
modifications [35]. The CD-1 mice were administered with WT and mutants for
competition colonization experiments as described above. At 48 h post-inoculation, the
mice were administered with 100 μL FITC-D-Glu (0.1 mM) in PBS. Two hours after
the gavage, the animals were sacrificed and the small intestine was dissected and rinsed
with 2 mL PBS, cut longitudinally and the intestinal content was scraped off using a
cell scraper. The intestinal content was suspended into a 1.5 mL PBS buffer with 2 steel
beads and shaken to release bacteria. Centrifugation at 3000 rpm for 1 min removed
large tissue fragments and impurities, and the supernatant was filtered through 40 μm
cell strainers to remove smaller cell debris. The filtrate was centrifuged (8,000 rpm, 5
min), the bacterial pellets were washed twice with 1.5 mL PBS and resuspended in PBS
for flow cytometry analysis. In the FITC channel, cells with strong signals of
approximately 10% were selected for sorting, with 10^5 events obtained for each sample,
and the collected samples were diluted and plated on LB agar containing streptomycin
and X-Gal for quantification of the bacterial index.

2.17 Detection of amino acid profiles in ΔflhB and ΔflaA by HPLC and MS

Overnight LB cultures of ΔflaA and ΔflhB were sub-cultured with an inoculum of 1:100
in MM medium and shaken at 37 °C until OD_{600}=0.5. The bacteria were collected by
centrifugation (>200mg) and samples of 60 mg were suspended in 50 μL water,
vortexed for 60 s, and mixed with 400 μL methanol-acetonitrile solution (1:1, v/v) and
50 μL of a mixture of 16 isotopic internal standards (50 μM). The mixture was vortexed
for 60 s and twice subjected to low-temperature ultrasound for 30 min. Proteins were
precipitated at -20 °C for 1 h and after centrifugation at 14,000 g and 4 °C for 20 min
and the supernatant was collected, freeze-dried, and stored at -80 °C prior to high-
performance liquid chromatography (HPLC). HPLC was performed using an Agilent
1290 Infinity LC ultra-high-performance liquid chromatography system. The mobile
phase consisted of 25 mM ammonium formate + 0.08% formic acid (solution A) and
0.1% formic acid acetonitrile (solution B). The samples were loaded into an automatic
sampler at 4 °C, and the column temperature was set at 40 °C. The flow rate was 250
μL min^{-1}, and the injection volume was 1 μL. The gradient elution was as follows: 0-
12 min, linear change from 90% B to 70% B; 12-18 min, linear change from 70% B to
50% B; 18-25 min, linear change from 50% B to 40% B; 30-30.1 min, linear change
from 40% B to 90% B; 30.1-37 min, maintained at 90% B. At regular intervals between
experimental samples a quality control (QC) sample was inserted to evaluate the
stability and reproducibility of the system.
Mass spectrometry (MS) analysis was performed using a 5500 QTRAP mass
spectrometer (SCIEX) in positive ion mode. The ESI source conditions for the 5500
QTRAP were as follows: source temperature 500 °C, ion Source Gas 1 (Gas 1): 40, Ion
Source Gas 2 (Gas 2): 40, Curtain gas (CUR): 30, IonSapary Voltage Floating (ISVF)
5,500 V. The analytes were detected using the MRM mode.

 2.18 In vivo colonization assay of 6N-labeled mutants

Briefly, using C6706 (*vc1807*:: CmR) as a template, amplify the CmR fragment and
insert a 6N site as a marker in the middle of one end primer (6N-R) [36], then clone it
into the pWM91 vector. Sequence tags of 6N were inserted into the *lacZ* position of the
26 mutants and the WT strain with the 6N-labeled plasmids pWM91-CmR-6N by
conjugation experiments. After labeling the bacterial strains, confirmed by sequencing,
they were sub-cultured and adjusted to an OD_{600} of 1.0. After harvesting by
centrifugation, the bacterial pellets were stored at -20°C. Using the methionine-
supplemented and water-restricted mouse model, the mice were gavaged with 100 μL
of the mixed bacterial samples per mouse. On day 5 after gavage, mouse feces were
collected and total DNA was extracted, along with the input samples. Amplicons
produced by PCR with primers targeting both ends of *LacZ* were sequenced. The
sequenced raw data was analyzed by Vsearch for merging paired-end reads and Python
to calculate the proportion of reads from the 6N-labeled mutants among the total reads.

2.19 Statistical analysis

Results are presented as mean ± standard deviation. When two groups were compared,
t-tests were performed. One-way analysis of variance (ANOVA) was used for two or
more treatment groups. Statistical analysis was performed using the software GraphPad
Prism v.9 for Windows. Differences with a p value less than 0.05 were considered to
be statistically significant.

3.   Results

3.1 High-frequency mutations in the flagellar genes of V. cholerae

The life cycle of *V. cholerae* primarily alternates between aquatic environments and the
human host [3]. To investigate its adaptation across these niches, we categorized
cholera-related metagenomic sequencing samples from the SRA database in Table S1
into five groups based on their origin: human, clinical, aquatic (including water and
sewage), fish, and amoeba-derived. These samples were collected from 19 countries
(Fig. 1A). A total of 1,258 metagenome-assembled genomes (MAGs) of *V. cholerae* in
Table S3 were reconstructed from these samples. Their flagellar protein mutations were
then systematically analyzed using the 51 flagellar proteins of *V. cholerae* N16961 in
Table S2 as reference amino acid sequences. For quality control, the mutation-
prone *hapR* gene, which carries a frameshift mutation in the N16961 genome, served
as   a   positive   control,   while   the   conserved   housekeeping
genes *asd* (VC2107), *mdh* (VC0432), and *dnaE* (VC2245) were used as negative
controls, as established in prior studies [27,28].
Our mutation frequency analysis revealed that across all sampled habitats, flagellar
gene   mutations   in *V.   cholerae* consistently   clustered   in   key   genes,
including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM* (Fig. 1B). Notably, mutations were also

detected in the primary flagellar regulator gene *flrA*, a finding consistent with previous
reports [16,37]. Functionally, these mutated genes are involved in critical flagellar
structures and regulatory hierarchies, including Class I and IV transcriptional regulation,
the type III secretion system (T3SS) apparatus, the filament-capping complex, and the
C-ring switch structure (Figure 1C, D). This indicates that these loci have key
evolutionary and pathological significance in regulating the host adaptability balance.

3.2 Phenotypic screening of the flagellar mutant library

To elucidate the environmental adaptability conferred by the flagellar gene mutations
identified through the above big data-driven analysis, a systematic mutant library
targeting 26 flagellar-related regulons of *V. cholerae* listed in Table S4 was constructed.
Subsequently, this library was subjected to comprehensive phenotypic screening,
assessing motility, biofilm formation, growth rates in both rich and minimal media,
colonization efficiency in infant and adult mice, as well as resistance to H_2O_2. The H_2O_2
assay is particularly relevant as it reflects bacterial survival under host-derived reactive
oxygen species (ROS) stress. Of the investigated mutants, 22 mutants had completely
lost their motility. Deletion of chemotaxis-related genes *cheV-3*, *flgO* and *flgP* resulted
in significantly impaired motility (Fig. 2A, Supplementary Fig. S1), which is consistent
with previous reports [6, 38]. The colonization ability in infant mice was seriously
damaged (Supplementary Fig. S2) and biofilm formation of the 26 mutants were mostly
enhanced, which were consistent with earlier findings (Supplementary Fig. S3) [5,6,10].
Surprisingly, we observed completely opposite results in the adult mice, as some of the
mutants were able to outcompete the WT strain (Fig. 2B), although their growth rates
were consistently lower than that of WT in minimum or rich medium (Supplementary
Fig. S4 and S5), and others were attenuated. In total, five mutants (Δ*flgBCDE*, Δ*fliK*,
Δ*fliKLMNOPQRflhB*, Δ*fliE,* and Δ*flgP*) produced no significance difference in CI (blue
in Fig. 1D), nine mutants (Δ*flrA*, Δ*rpoN*, Δ*flrC*, Δ*fliEFG*, Δ*fliHIJ,* Δ*fliA*, Δ*flgFGHIJ,*
Δ*fliPQR*, and Δ*flhB*) displayed enhanced CI (red in Fig. 2B), and twelve mutants (Δ*flhA*,
Δ*flhA* operon, Δ*flgB*, Δ*motY*, Δ*flaA*, Δ*flgK*, Δ*flgO*, Δ*flgT*, Δ*cheV-2*, Δ*vc1384*, Δ*flgM*,
and Δ*cheV-3*) were attenuated for CI (green in Fig. 2B). Of these, the Δ*flgFGHIJ* mutant
produced the highest CI with a 1000-fold increased CI compared to WT, while the
Δ*flgO* mutant had the lowest CI (1/1000-fold of WT). Of the other mutants, Δ*flhB*,
impaired in production of the flagellar type III secretion system (fT3SS) protein FlhB,
was enhanced in adaptability, and Δ*flaA*, missing the major flagellin filament protein,
was attenuated. We furthermore analyzed the various phenotypes of all 26 mutants in more
detail, and found that only the colonization potential in the adult mouse model differed
among the mutants. All other phenotypes (e.g. motility, biofilm development,
proliferation in LB/M9, H_2O_2 resistance, infant mice colonization) exhibited a unipolar
distribution. Correlation analysis indicated that adult mice colonization was not
associated with other phenotypes (Supplementary Fig. S6).

3.3 FlgM secretion negatively regulating adaptability

Considering FlaA and FlhB are structural flagellar component without a regulatory
function, they were confirmed through transmission electron microscopy that the 2
mutants lacked flagella (Supplementary Fig. S7A), and that overexpression of the
relevant genes restored their motility capability (Supplementary Fig. S7B). So, we
choose Δ*flhB*, Δ*flaA* and WT normalized with CFU (Supplementary Fig. S7C) were to
explore the underlying mechanism of these opposite colonization properties by
sequenced proteins in the supernatant (Fig. 2C). These proteins from Δ*flhB* and Δ*flaA*
were compared to a WT normalized control, in an attempt to seeking differentially
expressed proteins. This detected 36 proteins that were differently secreted between the
Δ*flhB* and Δ*flaA* mutants (Table S6, Supplementary Fig. S7D), including 15 flagellar
assembly proteins, 9 cellular processes related proteins, 2 secreted hemolysin proteins,
2 cell membrane associated proteins, 2 reactive oxygen species (ROS)-resistance
proteins, and 6 other proteins.

Hemolysins HlyA and Tlh are secreted toxin proteins that play a role in the virulence
of *V. cholerae* [6, 39]. However, their deletions in Δ*flhB* to give Δ*flhB*Δ*hlyA* and
Δ*flhB*Δ*tlh*, Δ*flhB*Δ*katG* and didn't affect their CI in the adult mice model
(Supplementary Fig. S8A, B). Colonization in this model can also be enhanced by ROS-
resistance, for which KatG and catalase are held responsible [32]. Experiments were
performed to investigate this, for which N-acetylcysteine (NAC) was employed to
remove ROS from the murine intestine (Supplementary Fig. S8C), but this revealed no
difference in the CI of Δ*flhB* or Δ*flaA* compared to that in absence of NAC
(Supplementary Fig. S8D, E). As not all mutants but the majority are resistant to ROS,
this did not seem to explain the enhanced colonization of particular flagellar mutants
(Supplementary Fig. S9).

Interestingly, the secretion of FlgM in Δ*flhB* was reduced about 9-fold compared
with that of Δ*flaA* (Fig. 2C). FlgM is an anti-σ^{28} factor, and functions as a chaperone
protein of σ^{28} factor FliA [40]. Our previous research had demonstrated that FlgM can
be secreted by Δ*flgD* through broken flagella, which activates FliA activity, to then
repress the expression of *hapR* that subsequently upregulates the expression of
virulence gene *tcpA* and enhances *V. cholerae* colonization in infant mice [4].
Introduction of a double-mutation where *flgM* was knocked out in a Δ*flhB* background
dramatically reversed the Δ*flhB* colonization pattern, from a CI of 176.3293 for Δ*flhB*
to 0.0098 for Δ*flhB*Δ*flgM* (*P*= 0.0005) (Fig. 2D). In addition, the overexpression of
FlgM in Δ*flaA* (*lacZ*::P*lac*-*flgM*) also significantly reversed the phenotype of Δ*flaA*,
from CI=0.0130 for Δ*flaA* to 0.2285 for Δ*flaA* (*P*= 0.0146) (Fig. 2E). Based on these
observations, we analyzed FlgM secretion in all 26 mutants and found that this
negatively correlated (R^2=-0.4417, *P*=0.0239) with their CI in adult mice (Fig. 2E, F).
Together, these data suggest that changes in intracellular FlgM levels in the various
flagellar mutants may contribute to their diverse colonization properties in adult mice.

3.4 Intracellular FlgM drives hapR expression

V. cholerae employs cell-to-cell communication, so-called quorum sensing, of which
HapR is the master regulator, which is in charge of many important regulation activities
in response to different environmental signals [41, 42]. In the Δ*flhB* mutant, a high
concentration of intracellular anti-σ^{28} factor FlgM is available to bind to σ^{28} factor FliA,
thereby inhibiting its function; this leads to an increase in *hapR* transcription (Fig. 3A,
left) [4]. We sequentially knocked out *flgM*, *fliA*, and *hapR* in a Δ*flhB* background.
Whereas the double mutant Δ*flhB*Δ*flgM* had a reduced CI (0.0010) compared to Δ*flhB*
(CI=210.5072, *P*=0.002), this phenotype was more than reversed in the triple mutant
Δ*flhB*Δ*flgM*Δ*fliA*   (CI=616.9768),   whereas   the   quadruple   mutant
Δ*flhB*Δ*flgM*Δ*fliA*Δ*hapR* had a reduced CI (1.408) again (Fig. 3A, right). The order of
sequential mutation was then reserved by inactivating (in that order) *flgM*, *fliA*, and
hapR in a Δ*flaA* background, which resulted in similar switches in colonization ability
(Fig. 3B). This suggests that *hapR* expression may be responsible for the different
colonization abilities of these flagellar mutants in the applied adult mouse model.

This hypothesis was confirmed by determining that the transcription of *hapR* was
significantly upregulated in those mutants with enhanced colonization, and
significantly downregulated or unchanged in the other flagellar mutants (Fig. 3C). And
In addition, the knockout of *hapR* significantly decreased the CI (Fig. 3D), suggesting
that a FlgM-FliA-*hapR* pathway contributes to the colonization potential.

We found that overexpression of FliA (by means of pBAD-*fliA*) in Δ*flgM*Δ*fliA*
significantly repressed *hapR* expression (*P*=0.020), while overexpression of FlgM in
Δ*flgM*Δ*fliA* derepressed *hapR* transcription (*P*=0.003) (Fig. 3E). Similar results were
obtained with *E. coli* DH5α (Supplementary Fig. S10), suggesting that *hapR* expression
may be regulated by FliA directly.

Previous studies had reported that the VarS/VarA-CsrA/BCD pathway, three
autoinducer systems (CAI-1, AI-2, DPO), and VqmA can regulate *hapR* expression
(Fig. 3F) [43-45]. We determined the expression of each key gene within the quorum
sensing network in the Δ*fliA* mutant, which suggested that FlgM-FliA may indirectly
inhibit HapR degradation through the VarS/VarA-CsrA/BCD pathway (Fig. 3G). To
compare the priority of direct and indirect regulation of *hapR* by FlgM-FliA, the *varA*
and *hapR* genes were individually knocked out in a Δ*flhB* background. Adult mice
colonization results revealed that this drastically reduced colonization compared to
Δ*flhB* (*P*=0.0026 for *varA* and *P*=0.0023 for *hapR*); no significantly difference was
observed between Δ*flhB*Δ*varA* and Δ*flhB*Δ*hapR* (Fig. 3H). The double mutation of
varA and *flhB* consistently decreased the expression level of *hapR* (Supplementary Fig.
S11). Taken together, it is concluded that interaction between intracellular FlgM and
FliA derepresses *hapR* transcription, resulting in an enhanced colonization ability of *V.*
cholerae in adult mice.

3.5 HapR monitors methionine metabolism

We next compared precipitated protein profiles of Δ*flhB* and Δ*flaA*, using WT as a
normalization control in Table S7. The cellular protein levels of flagellin proteins FlaD,

methionine sulfoxide reductase MsrC, serine protease VesC, and hypothetical proteins VC1075 were upregulated (Fig. 4A). However, no significantly difference was observed among Δ*flhB* and Δ*flhB*Δ*flaD,* Δ*flhB*Δ*vesC,* Δ*flhB*Δ*vc1075* in the adult mice (Supplementary Fig. S12). The MsrC was the highest enriched cellular protein in Δ*flhB* (Log$_2$Ratio=3.000, *P*=0.002), relative to Δ*flaA* (Fig. 4A). Therefore, we now hypothesize that cellular methionine contributes to the different colonization patterns of the 26 flagellar mutants in adult mice.

In *V. cholerae*, cellular methionine metabolism is finely regulated via biosynthesis, transformation and transportation systems [Fig. 4B] [46]. As expected, the addition of methionine significantly reduced Δ*flhB* and it increased Δ*flaA* colonization in adult mice (Fig. 4C), whereas Δ*msrC* drastically reduced the colonization advantage (Fig. 4D, CI=0.015). However, knockout of the methionine biosynthesis genes *msrC* or/and *metR* in a Δ*flhB* background didn't reduce colonization advantage, unless the methionine transportation genes *metI* and *metT* were also inactivated (Δ*flhB*Δ*msrC*Δ*metR*Δ*metI*Δ*metT vs.* WT compared to Δ*flhB vs.* WT, FC=6.341, *P*=0.001) (Fig. 4E). This suggests that the cellular methionine yield is precisely regulated by a range of metabolic and transportation genes in *V. cholerae*, which is not surprising, since methionine is an essential amino acid for bacterial metabolism and proliferation. Subsequently, the amino acid content in the precipitates of Δ*flaA*, Δ*flhB*, and WT were quantified. We found lower serine concentration (*P*=0.040) and higher concentrations of cystine (a disulfide of two oxidized cysteine molecules, *P*=0.018) in Δ*flhB* compared with Δ*flaA*, but no variation in methionine concentrations could be demonstrated by intracellular amino acid assays (Fig. 4F). Beyond bacterial biosynthesis and uptake, intracellular methionine can also be increased via other amino acid pathways. The KEGG pathway shows that L-cystathionine, a downstream product of serine and methionine, can be converted to cysteine, from which cystine is formed (Fig. 4G) [47], so that a higher active methionine metabolism in Δ*flhB* may relate to lower serine concentrations and higher cystine concentrations. Interestingly, we also found that *hapR* is involved in the expression of enzymes related to other amino acid metabolism by analyzing the transcriptome from our Mr.VC database [48, 49], including serine and cysteine metabolism, suggesting that *hapR* may be involved in the regulation of methionine metabolic flow.

Tsou et al. have described that HapR is a global regulator in *V. cholerae,* playing an important role in virulence regulation, biofilm development, natural transformation, and stress response [42]. By means of sequence alignments we identified sequences characteristic of the HapR-binding motif 1 in front of *metT*, *msrC*, *metI*, and a HapR-binding motif 2-like sequence in front of *metR* (Fig. 4H). Indeed, HapR directly promotes the transcription of these four genes, as demonstrated by RT-qPCR (Fig. 4I), and binding of HapR to promoter DNA was demonstrated by EMSA experiments (Fig. 4J, Supplementary Fig. S13). We observed that it rescued the growth defect of Δ*flaA* in methionine-limited medium (MM) (Fig. 4K) and adding methionine to the medium did not revive the growth of Δ*flhB*Δ*metR*Δ*msrC*Δ*metI*Δ*metT* (Fig. 4L).

3.6 Methionine metabolism affects V. cholerae colonization

To examine the metabolic activity of $\Delta flhB$, $\Delta flaA$ and WT in vitro, we used FITC-
labeled glucose method (Fig. 5A). The metabolic activity is often labeled by marking
bacterial cell wall precursors [50, 51], such as GlcNAc-1-phosphate, which is originally
derived from glucose [52]. Fluorescein Isothiocyanate (FITC) is a commonly used
fluorescent probe molecule that can form a covalent complex with glucose molecules.
After FITC-labeled glucose is utilized, the detection of fluorescence signals can
indicate bacterial metabolic activity. A transwell co-culture system with WT showed
that the population of $\Delta flhB$ was highly metabolically active and 1.7-fold higher than
WT (P=0.012), while $\Delta flaA$ was 0.6-fold lower (P=0.003) than WT (Fig. 5B). We also
investigated bacterial metabolism in vivo by administering FITC-labeled glucose to
mice that were colonized with either a $\Delta flaA$ and WT mixture or a $\Delta flhB$ and WT
mixture, after which the bacterial content was isolated from their intestine and subjected
to flow cytometry to sort those bacterial cells with high metabolic activity. These were
enumerated on X-Gal plates, which revealed that the population of $\Delta flhB$ was 7-fold
higher (P=0.017), while the population of $\Delta flaA$ was 0.2-fold lower compared to WT
(P=0.001) (Fig. 5C). Thus, both in vitro co-culture and in vivo in adult mice co-
colonization experiments provided strong evidence that bacterial metabolic activity
contributed benefited the bacteria. In combination with the other results, this suggest
that differences in methionine-related metabolic activity may contribute to the different
colonization patterns of the flagellar mutants.

3.7 Ecological simulation of flagellar mutants reveals trade-off mechanisms between
motility loss and host adaptability.

To assess the host adaptability of the flagellar mutants, the previously characterized
mutants were labeled using a novelty developed 6N-labeling method and subsequently
administered to adult mice via oral gavage. Colonization competition was then assessed
by high-throughput sequencing (Fig. 5D, Table S8). In the absence of methionine
supplementation, the competition indices (CIs) of all 26 mutants showed a scattered
and diverse distribution, consistent with the head-to-head colonization outcomes
presented earlier (Fig. 2B). Upon methionine addition, the distribution of CIs for all
flagellar mutants became markedly more constrained, as detailed in Supplementary
Dataset 6. Further pairwise analysis revealed that methionine supplementation could
reverse the colonization pattern of individual flagellar mutants (Fig. 5E). Collectively,
these findings demonstrate that ecological simulation of flagellar mutants reveals trade-
off mechanisms between motility loss and host adaptability.

4. Discussion

By employing big data analysis on *V. cholerae* life cycle samples, this study
elucidates the trade-off mechanism whereby flagellar mutations result in motility loss

yet enhance host adaptability through the FlgM-FliA-HapR-methionine axis, a driver
of metabolic reprogramming (Fig. 6).

Although flagellar mutations are prevalent across the life cycle environments of *V.*
cholerae, their evolutionary paths across distinct host niches remain unclear. Previous
work indicates that the cyclin gene *dps*, rather than DNA repair or ROS/RNS
scavenging systems, significantly increases flagellar mutation frequency in an adult
mouse model [15]. In amoeba hosts, mutations in the transcriptional
regulator *flrA* provide a competitive edge, while intense microbial competition in the
gut may further shape *V. cholerae* evolution [16]. In addition, the *flrA* in *V.*
cholerae senses intracellular levels of the second messenger cyclic diguanylate
monophosphate (c-di-GMP) in response to dietary signals. Under a high c-di-GMP
environment induced by casein, FlrA represses the expression of the type VI secretion
system (T6SS), thereby impairing the ability of *V. cholerae* to compete with intestinal
commensal bacteria via T6SS and ultimately leading to a colonization defect in casein-
fed mice [53]. Furthermore, intense microbial competition in the gut may serve as a
major driving force for the evolution of *V. cholerae* within the intestinal environment
[54]. Integrating comparative genomics with experimental evolution assays may
unravel whether these mutations represent adaptations to host niches or collateral
effects of pathogen transmission dynamics.

Methionine is an essential amino acid utilized by bacteria in various physiological
and metabolic processes, including polyamine synthesis, DNA synthesis, redox balance,
and DNA and histone methylation [55]. Also, it is important to note that methionine
plays an essential role in the immune system through its metabolites. For example, in
response to antigens, T cells activate the methionine cycle to support proliferation and
differentiation in cancer, indicating the importance of the methionine cycle to T cell
immunity [56]. Methionine supplementation or restriction can intervene in the natural
antioxidant capacity of an organism by leading to the production of endogenous
enzymes that reduce oxidative stress and DNA damage [57]. These data imply that *V.*
cholerae may employ the methionine pathway as a means of immune evasion. To
successfully propagate and cause disease, *V. cholerae* also evades the immune system
by repressing the mannose-sensitive hemagglutinin pilus via the transcription factor
ToxT, preventing secretory IgA binding and enabling intestinal colonization to bypass
host defenses [58].

Methionine metabolism is a central node within the interconnected amino acid
network of *V. cholerae* [59]. Beyond its metabolic functions, methionine serves as a
precursor for S-adenosylmethionine (SAM), a cofactor involved in critical processes
including the synthesis of the quorum-sensing autoinducer CAI-1 [60,61]. Similarly,
threonine, another methionine-derived metabolite, may contribute to autoinducer DPO
synthesis, and in *Pseudomonas aeruginosa*, SAM participates in producing AI-1, AI-2,
and AHLs [62]. Analyzing methionine flux could therefore help identify key

metabolites that enhance colonization in flagellar mutants, and future work should
examine whether intermediate methionine metabolites or other amino acids influence *V.*
cholerae colonization through bacterial or host metabolic remodeling.

This study found that HapR binds to promoters of methionine metabolism-related
genes (*metR*, *msrC*, *metI*, *metT*), a result not reported in prior systematic screens of
HapR targets. The discrepancy may stem from limitations in bioinformatic prediction
accuracy and variability in HapR-binding sequences [42]. Beyond methionine
regulation, HapR suppresses toxin-coregulated pilus (TCP) and cholera toxin (CTX)
genes by repressing *aphA*, while activating extracellular proteases and modulating
T6SS genes [63-65]. However, in adult and infant mouse models, this virulence factors
do not substantially affect colonization [39,66]. HapR also inhibits biofilm formation
via cyclic di-GMP modulation and *vpsT* repression [67-69], although flagellar mutants
show enhanced biofilm capacity, they still exhibit colonization defects, indicating
biofilm is not a major driver of colonization in these strains. Additionally, HapR and
RpoS jointly facilitate stress adaptation, luminal dispersal, and host exit. Although the
flagellar mutants of *V. cholerae* exhibit enhanced biofilm formation capacity, they still
display a colonization defect (Supplementary Fig. S3, Fig. 2B), suggesting that biofilm
formation is also not a major factor contributing to enhanced colonization in these
mutants. In addition, HapR and RpoS collectively facilitates stress adaptation to host
factors, dispersal within the intestinal lumen, and eventual exit of *V. cholerae* from the
host [70]. In the precipitated proteome, we observed that MsrC was the most highly
enriched cellular protein in Δ*flhB* (Log2Ratio = 3.000, P = 0.002) relative to Δ*flaA* (Fig.
4A). Through experimental validation, we concluded that methionine contributes to the
differential colonization patterns among the 26 flagellar mutants in adult mice.
The flagellum renders bacteria motility, but it is also a major antigen recognized by
the host [71, 72]. Extensive studies have shown that flagellar mutants of *V. cholerae*
are frequently present in clinical and environmental samples, and specifically the *V.*
cholerae Haiti strains contain multiple mutational hotspots in flagellum gene regions
according to SNP analysis [11]. Besides mutations, *V. cholerae* also actively loses
flagellar filaments during environmental changes such as mucus penetration [4],
biofilm development [10], and host Lypd8 promotes the segregation antibody capture
[72]. This is also demonstrated for other pathogenic bacteria. The freshwater bacterium
Caulobacter crescentus loses its flagella through ejection trigged by ClpA [73]. The
loss of the polar flagellum in *V. fischeri* and in plant-associated *Methylobacteria* has
also been reported [74, 75]. Bacterial flagella also detach during starvation [76].
We demonstrated that the trade-off mechanism between motility loss and host
adaptability operates via the sophisticated FlgM-FliA-HapR-methionine metabolic
reprogramming axis. However, this study has the following limitations: (1) The
mechanistic focus on the FlgM-FliA-HapR-methionine axis, though insightful, may not
capture the full complexity of the flagellar regulatory network, and deeper metabolic
validation is needed; (2) It does not directly simulate or verify how this trade-off
process dynamically occurs and becomes fixed within a population under

environmental pressure, lacking direct experimental evolution evidence in human; (3) The adaptive efficacy and stability of the revealed mechanism in such highly complex real-world environments require further evaluation; (4) Time-course experiments were not provided to track the changes in FlgM secretion, HapR expression, and methionine metabolism over the course of infection.

4. Conclusion

Based on extensive sample analysis, this study reveals that flagellar mutations in *V. cholerae* cluster in key structural and regulatory genes, underscoring their central role in host adaptation. Specifically, the secretion level of the flagellar regulator FlgM drives a phenotypic shift from motility to host adaptability. This transition is mediated through the FlgM-FliA-HapR-methionine axis, which balances the loss of motility with enhanced host adaptability. Ultimately, impaired motility promotes environmental fitness by activating quorum sensing and reprogramming methionine metabolism, elucidating a novel evolutionary trade-off strategy between movement and adaptability at the molecular level.

Environmental implications

V. cholerae can persist in both aquatic environments and the human intestinal tract, posing substantial impacts on environmental and human health. Genetic mutations enable the bacterium to survive under complex environmental stresses, thereby increasing the risk of human infection. This study demonstrates that *V. cholerae* can acquire flagellar gene mutations across different environmental niches within its life cycle. Notably, mutations in key genes enhance its adaptability within the host intestine, a process mediated by the FlgM-FliA-HapR-methionine metabolic axis. Our findings elucidate the trade-off mechanism between motility loss and host adaptability, offering novel perspectives on bacterial evolution and pathogenicity.

CRediT authorship contribution statement

Guozhong Chen: Conceptualization, Methodology, Investigation, Project administration, Funding acquisition, Visualization, Writing-original draft, Writing-review & editing. **Zixin Qin:** Methodology, Investigation, Visualization. **Fenxia Fan:** Investigation, Writing-review & editing, Funding acquisition. **Mei Luo:** Writing-review & editing. **Hongou Wang:** Methodology. **Baoshuai Xue:** Methodology. **Shucheng Li:** Methodology. **Shiyong Chen:** Supervision. **Xiaoman Yang:** Investigation. **Xiang Mao:** Investigation. **Liwen Yi:** Investigation. **Chunrong Yi:** Investigation. **Wei Li:** Investigation, Supervision. **Xiaoyun Liu:** Project administration, Funding acquisition. **Biao Kan:** Supervision, Project administration. **Zhi Liu:** Conceptualization, Methodology, Visualization, Project administration, Funding acquisition, Supervision, Writing-original draft, Writing-review & editing.

Declaration of Competing Interest

The authors declare no financial/commercial conflicts of interests.

Acknowledgments

We are grateful to all the members of our team. We thank Core Facilities for Life
Science, Huazhong University of Science and Technology for the technical support.
We also thank Core Facilities for Life and Environmental Sciences, State Key
laboratory of Microbial Technology of Shandong University for technical guidance.
This study was supported by National Natural Science Foundation of China under the
grant numbers 31770132 (to Z. L), (22174003 and 21974002 to X. Y. L) and National
Key Research and Development Program of China (2022YFA1304500 to X. Y. L),
National Key Research and Development Program of China (2021YFC2300304 to F.
X. F), Shandong Province Science Foundation for Youths (ZR2024QC312 to G. Z. C).

Supplementary Material

Supplementary material is available online.

Data Availability

The proteomic data that support the findings of this study are openly available in the
iProX database at https://www.iprox.cn//page/SCV017.html?query=IPX0005998000,
accession number IPX0005998000. The raw data of 6N-Label is openly available in
the NCBI database at https://www.ncbi.nlm.nih.gov/sra/?term=PRJNA1099378,
reference number PRJNA1099378. The codes for 6N-Label relative abundance analysis,
bins acquisition and species annotation from cohort-based studies, and analyzing high-
frequency mutations in the flagellar genes of *V. cholerae* of this study are openly
available in Zenodo at https://doi.org/10.5281/zenodo.14920964, reference number
14920964.

References

1. Kanungo S, Azman AS, Ramamurthy T, Deen J, Dutta S. Cholera. *Lancet* 2022;
399: 1429-1440. https://doi.org/10.1016/s0140-6736(22)00330-0.
2. World Health Organization. Cholera. https://www.who.int/news-room/fact-
sheets/detail/cholera (10 February 2025, date last accessed).
3. D. Chac, C. N. Dunmire, J. Singh, A. A. Weil. Update on environmental and
hostfactors impacting the risk of *Vibrio cholerae* infection. ACS Infectious
Diseases,2021, 7(5): 1010-1019. https://doi.org/10.1021/acsinfecdis.0c00914.
4. Liu Z, Miyashiro T, Tsou A, Hsiao A, Goulian M, Zhu J. Mucosal penetration
primes *Vibrio cholerae* for host colonization by repressing quorum sensing. *Proc
Natl Acad Sci U S A* 2008; **105**: 9769-9774.
https://doi.org/10.1073/pnas.0802241105.

5.   Echazarreta MA, Klose KE. *Vibrio* flagellar synthesis. Front Cell Infect Microbiol
2019; 9: 131. https://doi.org/10.3389/fcimb.2019.00131.

6.   Syed KA, Beyhan S, Correa N, Queen J, Liu J, Peng F, et al. The *Vibrio cholerae*
flagellar regulatory hierarchy controls expression of virulence factors. *J Bacteriol*
2009; **191**: 6555–6570. https://doi.org/10.1128/jb.00949-09.

7.   Sowa Y, Berry RM. Bacterial flagellar motor. *Q Rev Biophys* 2008; **41**: 103-132.
https://doi.org/10.1017/s0033583508004691.

8.   Attridge SR, Rowley D. The role of the flagellum in the adherence of *Vibrio*
cholerae. *J Infect Dis* 1983; **147**: 864-872.
https://doi.org/10.1093/infdis/147.5.864.

9.   Butler SM, Camilli A. Both chemotaxis and net motility greatly influence the
infectivity of *Vibrio cholerae*. *Proc Natl Acad Sci U S A* 2004; **101**: 5018-5023.
https://doi.org/10.1073/pnas.0308052101.

10. Wu DC, Zamorano-Sánchez D, Pagliai FA, Park JH, Floyd KA, Lee CK, et al.
Reciprocal c-di-GMP signaling: Incomplete flagellum biogenesis triggers c-di-
GMP signaling pathways that promote biofilm formation. *PLoS Genet* 2020; **16**:
e1008703. https://doi.org/10.1371/journal.pgen.1008703.

11. Katz LS, Petkau A, Beaulaurier J, Tyler S, Antonova ES, Turnsek MA, et al.
Evolutionary dynamics of *Vibrio cholerae* O1 following a single-source
introduction to Haiti. *mBio* 2013; **4**: 1-10. https://doi.org/10.1128/mbio.00398-13.

12. Mostow P, Richardson K. High-frequency spontaneous mutation of classical
Vibrio cholerae to a nonmotile phenotype. *Infect Immun* 1990; **58**: 3633-3639.
https://doi.org/10.1128/iai.58.11.3633-3639.1990.

13. Jubair M, Atanasova KR, Rahman M, Klose KE, Yasmin M, Yilmaz O, et al.
Vibrio cholerae persisted in microcosm for 700 days inhibits motility but promotes
biofilm formation in nutrient-poor lake water microcosms. *PLoS One* 2014; **9**:
e92883. https://doi.org/10.1371/journal.pone.0092883.

14. Giacomucci S, Mathieu-Denoncourt A, Vincent AT, Jannadi H, Duperthuy M.
Experimental evolution of *Vibrio cholerae* identifies hypervesiculation as a way to
increase motility in the presence of polymyxin B. *Front Microbiol* 2022; **13**: 1-18.
https://doi.org/10.3389/fmicb.2022.932165.

15. Hoque MM, Noorian P, Espinoza-Vergara G, Manuneedhi Cholan P, Kim M,
Rahman MH, et al. Adaptation to an amoeba host drives selection of virulence-
associated traits in *Vibrio cholerae*. *ISME J* 2022; **16**: 856-867.
https://doi.org/10.1038/s41396-021-01134-2.

16. Luo M, Chen G, Yi C, Xue B, Yang X, Ma Y, et al. Dps-dependent in vivo
mutation enhances long-term host adaptation in *Vibrio cholerae*. *PLoS Pathog*
2023; **19**: e1011250. https://doi.org/10.1371/journal.ppat.1011250.

17. Klose KE, Mekalanos JJ. Distinct roles of an alternative sigma factor during both
free-swimming and colonizing phases of the *Vibrio cholerae* pathogenic cycle.
Mol Microbiol 1998; **28**: 501-520. https://doi.org/10.1046/j.1365-
2958.1998.00809.x.

18. Azarian T, Ali A, Johnson JA, Mohr D, Prosperi M, Veras NM, et al.
Phylodynamic analysis of clinical and environmental *Vibrio cholerae* isolates from
Haiti reveals diversification driven by positive selection. *mBio* 2014; **5**: e01824-
14. https://doi.org/10.1128/mbio.01824-14.

19. Daboul J, Weghorst L, DeAngelis C, Plecha SC, Saul-McBeth J, Matson JS.
Characterization of *Vibrio cholerae* isolates from freshwater sources in northwest
Ohio.   *PLoS   One   2020;   **15**:   e0238438.
https://doi.org/10.1371/journal.pone.0238438.

20. Mavian C, Paisie TK, Alam MT, Browne C, de Rochars VMB, Nembrini S, et al.
Toxigenic *Vibrio cholerae* evolution and establishment of reservoirs in aquatic
ecosystems. *Proc Natl Acad Sci U S A* 2020; **117**: 7897-7904.
https://doi.org/10.1073/pnas.1918763117.

21. Mwaba J, Debes AK, Murt KN, Shea P, Simuyandi M, Laban N, et al. Three
transmission events of *Vibrio cholerae* O1 into Lusaka, Zambia. *BMC Infect Dis*
2021; **21**: 570. https://doi.org/10.1186/s12879-021-06259-5.

22. Chen, Shifu. Ultrafast one-pass FASTQ data preprocessing, quality control, and
deduplication using fastp. *iMeta* 2023; **2**: 1-5. https://doi.org/10.1002/imt2.107.

23. Uritskiy, Gherman V., DiRuggiero Jocelyne, Taylor James. MetaWRAP—a
flexible pipeline for genome-resolved metagenomic data analysis. *Microbiome*
2018; **6**: 158. https://doi.org/10.1186/s40168-018-0541-1.

24. Chaumeil, Pierre-Alain, Mussig Aaron J, Hugenholtz Philip, Parks Donovan H.
GTDB-Tk v2: memory friendly classification with the genome taxonomy database.
Bioinformatics   2022;   **38**:   5315-5316.
https://doi.org/10.1093/bioinformatics/btac672.

25. Hyatt, Doug, Chen Gwo-liang, LoCascio Philip F, Land Miriam L, Larimer Frank
W, Hauser Loren J. Prodigal: Prokaryotic gene recognition and translation
initiation   site   identification.   *BMC   Bioinformatics*   2010;   **11**:119.
https://doi.org/10.1186/1471-2105-11-119.

26. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. Nature Methods 2015;**12**: 59-60. https://doi.org/10.1038/nmeth.3176

27. Farhadi T, Nezafat N, Ghasemi Y. In silico phylogenetic analysis of *Vibrio
cholerae* isolates based on three housekeeping genes. Int J Comput Biol Drug Des.
2015;8(1):62-74. https://doi.org/10.1504/ijcbdd.2015.068789

28. Rui YY, Kan B, Gao SY, Liu YQ, Qi GM. Sequence analysis of housekeeping
genes *recA*, *dnaE*, and *mdh* in different serogroups or biotypes of *Vibrio cholerae*
isolated from China. Nan Fang Yi Ke Da Xue Xue Bao. 2006;26(12):1720-1723.
https://www.j-smu.com/CN/Y2006/V26/I12/1720

29. Houot L, Watnick PI. A novel role for enzyme I of the *Vibrio cholerae*
phosphoenolpyruvate phosphotransferase system in regulation of growth in a
biofilm. *J Bacteriol* 2008; **190**: 311-320. https://doi.org/10.1128/jb.01410-07.

30. Metcalf WW, Jiang W, Daniels LL, Kim SK, Haldimann A, Wanner BL.
Conditionally replicative and conjugative plasmids carrying lacZα for cloning,

mutagenesis, and allele replacement in bacteria. *Plasmid* 1996; **35**: 1-13. https://doi.org/10.1006/plas.1996.0001.

31. Xia A, Qian M, Wang C, Huang Y, Liu Z, Ni L, et al. Optogenetic modification of *Pseudomonas aeruginosa* enables controllable twitching motility and host infection. *ACS Synth Biol* 2021; **10**: 531-541. https://doi.org/10.1021/acssynbio.0c00559.

32. Ma Y, Yang X, Wang H, Qin Z, Yi C, Shi C, et al. CBS-derived H_2S facilitates host colonization of *Vibrio cholerae* by promoting the iron-dependent catalase activity of KatB. *PLoS Pathog* 2021; **17**: e1009763. https://doi.org/10.1371/journal.ppat.1009763.

33. Liu Z, Wang H, Zhou Z, Sheng Y, Naseer N, Kan B, et al. Thiol-based switch mechanism of virulence regulator AphB modulates oxidative stress response in *Vibrio cholerae. Mol Microbiol* 2016; **102**: 939-949. https://doi.org/10.1111/mmi.13524.

34. Hong S, Zheng DW, Zhang QL, Deng WW, Song WF, Cheng SX, et al. An RGB-emitting molecular cocktail for the detection of bacterial fingerprints. *Chem Sci* 2020; **11**: 4403-4409. https://doi.org/10.1039/d0sc01704c.

35. Lin L, Song J, Du Y, Wu Q, Gao J, Song Y, et al. Quantification of bacterial metabolic activities in the gut by d-Amino acid-based in vivo labeling. *Angew Chemie - Int Ed* 2020; **59**: 11923-11926. https://doi.org/10.1002/anie.202004703.

36. Dalia AB, McDonough EK, Camilli A. Multiplex genome editing by natural transformation. *Proc Natl Acad Sci U S A* 2014; **111**: 8937-8942. https://doi.org/10.1073/pnas.1406478111.

37. Hoque MM, Noorian P, Espinoza-Vergara G, Manuneedhi Cholan P, Kim M, Rahman MH, et al. Adaptation to an amoeba host drives selection of virulence-associated traits in *Vibrio cholerae. ISME J* 2022; 16: 856-867. https://doi.org/10.1038/s41396-021-01134-2.

38. Martinez RM, Dharmasena MN, Kirn TJ, Taylor RK. Characterization of two outer membrane proteins, FlgO and FlgP, that influence *Vibrio cholerae* motility. *J Bacteriol* 2009; **191**: 5669-5679. https://doi.org/10.1128/jb.00632-09.

39. Olivier V, Salzman NH, Satchell KJF. Prolonged Colonization of Mice by *Vibrio cholerae* El Tor O1 Depends on Accessory Toxins. *Infect Immun* 2007; **75**: 5043-5051. https://doi.org/10.1128/iai.00508-07.

40. Correa NE, Barker JR, Klose KE. The *Vibrio cholerae* FlgM homologue is an anti-σ^{28} factor that is secreted through the sheathed polar flagellum. *J Bacteriol* 2004; **186**: 4613-4619. https://doi.org/10.1128/jb.186.14.4613-4619.2004.

41. Qin Z, Yang X, Chen G, Park C, Liu Z. Crosstalks between gut microbiota and *Vibrio cholerae. Front Cell Infect Microbiol* 2020; **10**: 582554. https://doi.org/10.3389/fcimb.2020.582554.

42. Tsou AM, Cai T, Liu Z, Zhu J, Kulkarni R V. Regulatory targets of quorum sensing in *Vibrio cholerae*: Evidence for two distinct HapR-binding motifs. *Nucleic Acids Res* 2009; **37**: 2747-2756. https://doi.org/10.1093/nar/gkp121.

43. Tsou AM, Liu Z, Cai T, Zhu J. The VarS/VarA two-component system modulates the activity of the *Vibrio cholerae* quorum-sensing transcriptional regulator HapR. *Microbiology* 2011; **157**: 1620-1628. https://doi.org/10.1099/mic.0.046235-0.

44. Herzog R, Peschek N, Fröhlich KS, Schumacher K, Papenfort K. Three autoinducer molecules act in concert to control virulence gene expression in *Vibrio cholerae*. *Nucleic Acids Res* 2019; **47**: 3171-3183. https://doi.org/10.1093/nar/gky1320.

45. Liu Z, Hsiao A, Joelsson A, Zhu J. The transcriptional regulator VqmA increases expression of the quorum-sensing activator HapR in *Vibrio cholerae*. *J Bacteriol* 2006; **188**: 2446-2453. https://doi.org/10.1128/jb.188.7.2446-2453.2006

46. Vanhove AS, Jugder BE, Barraza D, Watnick PI. Methionine availability in the arthropod intestine is elucidated through identification of *Vibrio cholerae* methionine acquisition systems. *Appl Environ Microbiol* 2020; **86**: e00371-20. https://doi.org/10.1128/aem.00371-20.

47. Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 2000; **28**: 27-30. https://doi.org/10.1093/nar/28.1.27.

48. Zhang Z, Chen G, Hu J, Hussain W, Fan F, Yang Y, et al. Mr.Vc: A database of microarray and RNA-seq of *Vibrio cholerae*. *Database* 2019; **2019**: 1-8. https://doi.org/10.1093/database/baz069.

49. Zhang Z, Chen G, Hussain W, Qin Z, Liu J, Su Y, et al. Mr.Vc v2: An updated version of database with increased data of transcriptome and experimental validated interactions. *Front Microbiol* 2022; **13**: 1047259. https://doi.org/10.3389/fmicb.2022.1047259.

50. Calabretta PJ, Hodges HL, Kraft MB, Marando VM, Kiessling LL. Bacterial cell wall modification with a glycolipid substrate. *J Am Chem Soc* 2019; **141**: 9262-9272. https://doi.org/10.1021/jacs.9b02290.

51. Xu Y, Hernández-Rocamora VM, Lorent JH, Cox R, Wang X, Bao X, et al. Metabolic labeling of the bacterial peptidoglycan by functionalized glucosamine. *iScience* 2022; **25**: 104753. https://doi.org/10.1016/j.isci.2022.104753.

52. Sadamoto R, Matsubayashi T, Shimizu M, Ueda T, Koshida S, Koda T, et al. Bacterial surface engineering utilizing glucosamine phosphate derivatives as cell wall precursor surrogates. *Chem - A Eur J* 2008; **14**: 10192-10195. https://doi.org/10.1002/chem.200801734.

53. Liu, R. et al. Diet modulates *Vibrio cholerae* colonization and competitive outcomes with the gut microbiota. *Cell Host and Microbe* 2025; **33**, 2085-2099.e5. https://doi.org/10.1016/j.chom.2025.11.004.

54. Jiang S, Zhang C, Han Z, Ma W, Wang S, Huo D, Cui W, Zhai Q, Huang S, Zhang J. Native microbiome dominates over host factors in shaping the probiotic genetic evolution in the gut. *NPJ Biofilms Microbiomes*. 2023; 1:80. https://doi.org/10.1038/s41522-023-00447-8.

55. Martínez Y, Li X, Liu G, Bin P, Yan W, Más D, et al. The role of methionine on
metabolism, oxidative stress, and diseases. *Amino Acids* 2017; **49**: 2091-2098.
https://doi.org/10.1007/s00726-017-2494-2.

56. Zhao T, Lum JJ. Methionine cycle-dependent regulation of T cells in cancer
immunity. *Front Oncol* 2022; **12**: 1-9. https://doi.org/10.3389/fonc.2022.969563.

57. Martínez Y, Li X, Liu G, Bin P, Yan W, Más D, et al. The role of methionine on
metabolism, oxidative stress, and diseases. *Amino Acids* 2017; **49**: 2091-2098.
https://doi.org/10.1007/s00726-017-2494-2.

58. Hsiao A, Liu Z, Joelsson A, Zhu J. *Vibrio cholerae* virulence regulator-coordinated
evasion of host immunity. *Proc Natl Acad Sci U S A* 2006; **103**: 14542-14547.
https://doi.org/10.1073/pnas.0604650103.

59. Bogard RW, Davies BW, Mekalanos JJ. MetR-regulated *Vibrio cholerae*
metabolism is required for virulence. *mBio* 2012; **3**.e00236-12.
https://doi.org/10.1128/mbio.00236-12.

60. Hondorp ER, Matthews RG. Methionine. *EcoSal Plus* 2006; **2**: 185-185.
https://doi.org/10.1128/ecosalplus.3.6.1.7.

61. Winans SC. A new family of quorum sensing pheromones synthesized using S-
adenosylmethionine and Acyl-CoAs. *Mol Microbiol* 2011; **79**: 1403-1406.
https://doi.org/10.1111/j.1365-2958.2011.07551.x.

62. Heurlier, K., Dénervaud, V. & Haas, D. Impact of quorum sensing on fitness of
Pseudomonas aeruginosa. *International Journal of Medical Microbiology* 2006,
296, 93-102. https://doi.org/10.1016/j.ijmm.2006.01.043.

63. Cho JY, Liu R, Macbeth JC, Hsiao A. The interface of *Vibrio cholerae* and the gut
microbiome.   *Gut   Microbes*.   2021,   13:1937015.
https://doi.org/10.1080/19490976.2021.1937015.

64. 2. Hsiao A, Zhu J. Pathogenicity and virulence regulation of *Vibrio cholerae* at the
interface of host-gut microbiome interactions. *Virulence*. 2020; **11**:1582-1599.
https://doi.org/10.1080/21505594.2020.1845039.

65. 3. Qin Z, Yang X, Chen G, Park C, Liu Z. Crosstalks between gut microbiota and
Vibrio cholerae. *Front Cell Infect Microbiol* 2020; **10**: 582554.
https://doi.org/10.3389/fcimb.2020.582554.

66. Silva AJ, Leitch GJ, Camilli A, Benitez JA. Contribution of
hemagglutinin/protease and motility to the pathogenesis of El Tor biotype cholera.
Infect Immun. 2006;**74**:2072-9. https://doi.org/10.1128/iai.74.4.2072-2079.2006.

67. Waters CM, Lu W, Rabinowitz JD, Bassler BL. Quorum sensing controls biofilm
formation in *Vibrio cholerae* through modulation of cyclic di-GMP levels and
repression of *vpsT*. *J Bacteriol*. 2008; **190**:2527-36.
https://doi.org/10.1128/jb.01756-07

68. Srivastava D, Harris RC, Waters CM. Integration of cyclic di-GMP and quorum
sensing in the control of *vpsT* and *aphA* in *Vibrio cholerae*. *J Bacteriol*. 2011
Nov;193(22):6331-41. https://doi.org/10.1128/jb.05167-11

69. Johnson GE, Fei C, Wingreen NS, Bassler BL. Analysis of gene expression within
individual cells reveals spatiotemporal patterns underlying *Vibrio cholerae* biofilm
development. *PLoS Biol*. 2025; **23**: e3003187.
https://doi.org/10.1371/journal.pbio.3003187

70. Joelsson A, Kan B, Zhu J. Quorum sensing enhances the stress response in *Vibrio*
cholerae. *Appl Environ Microbiol*. 2007; **73**:3742-6.
https://doi.org/10.1128/aem.02804-06.

71. Khan F, Tabassum N, Anand R, Kim YM. Motility of *Vibrio* spp.: regulation and
controlling strategies. *Appl Microbiol Biotechnol* 2020; **104**: 8187-8208.
https://doi.org/10.1007/s00253-020-10794-7.

72. Okumura R, Kurakawa T, Nakano T, Kayama H, Kinoshita M, Motooka D, et al.
Lypd8 promotes the segregation of flagellated microbiota and colonic epithelia.
Nature 2016; **532**: 117-121. https://doi.org/10.1038/nature17406.

73. Grünenfelder B, Tawfilis S, Gehrig S, Østerås M, Eglin D, Jenal U. Identification
of the protease and the turnover signal responsible for cell cycle-dependent
degradation of the Caulobacter FliF motor protein. *J Bacteriol* 2004; **186**: 4960-
4971. https://doi.org/10.1128/jb.186.15.4960-4971.2004.

74. Ruby EG, Asato LM. Growth and flagellation of *Vibrio fischeri* during initiation
of the sepiolid squid light organ symbiosis. *Arch Microbiol* 1993; **159**: 160-167.
https://doi.org/10.1007/bf00250277.

75. Doerges L, Kutschera U. Assembly and loss of the polar flagellum in plant-
associated methylobacteria. *Naturwissenschaften* 2014; **101**: 339-346.
https://doi.org/10.1007/s00114-014-1162-6.

76. Zhu S, Gao B. Bacterial flagella loss under starvation. *Trends Microbiol* 2020; **28**:
785-788. https://doi.org/10.1016/j.tim.2020.05.002.

Figures and Tables

Fig. 1. High-frequency mutations in the flagellar genes of *V. cholerae* from the life
cycle. (**A**), World distribution map of the collected samples, with areas in pink
indicating no sample collection. (**B**) Spectrum and count statistics of flagellar gene
mutations. The *hapR* and housekeeping genes (*asd*, *dnaE*, *mdh*) were used as controls.
(**C**) Schematic diagram of the *V. cholerae* flagellum assembly. (**D**) *V. cholerae* flagellar
transcription hierarchy.

Fig. 2. FlgM secretion regulates colonization of *V. cholerae* flagellar mutants. (**A**), Motility assay. (**B**), Competition assay in adult mice. The horizontal bar indicates the average of CI values. The CI values at 5 days post inoculation were shown. The grey dashed lines from top to bottom represent 10-fold enhanced colonization and 10-fold reduced colonization, respectively. (**C**), Volcano plot of the distribution of altered proteins. Red dots represent up-regulated expression proteins, and green dots represent down-regulated expression proteins and grey dots represent no significant expression difference (threshold > 1.5, p < 0.05). (**D** and **E**), Effect of FlgM on the colonization of the Δ*flhB* or Δ*flaA* mutant. The CI values at 5 days post inoculation were shown. (**F**), Western blot (WB) assay for different secretion of FlgM in the *V. cholerae* flagellar mutants. sup: supernatant, p: pellets. (**G**), Correlation analysis of FlgM secretion and CI values in adult mice model (ImageJ and Pearson method). The colors of red, green dots represent a higher, lower of colonization than WT, respectively. Significance was determined by *t*-test; *P*-values: ns, not significant, *, < 0.05, **, < 0.01, ***, < 0.001.

Fig. 3. Intracellular FlgM deactivated FliA to promote *hapR* expression on transcriptionally and post-translationally. (**A** and **B**), Schematic diagram of FlgM secretion in Δ*flhB* and Δ*flaA*, and the corresponding role of FlgM-FliA-*hapR* pathway in colonization, respectively. (**C**), RT-PCR results of *hapR* expression in flagellar mutants with enhanced colonization. The *recA* was used for normalization and Δ*luxO* was used as a positive control. (**D**), *In vivo* competition experiments of different *hapR* mutants in adult mice. The CI values at 5 days post inoculation were shown. (**E**), Effect of FlgM and FliA overexpression on *hapR* expression. (**F**), Quorum sensing (QS) regulatory network of *V. cholerae*. (**G**), Gene expression in the QS regulatory pathway. Gene expression was normalized by *16S rRNA*, except for small RNAs called *csrBCD* which were normalized by *secA*. (**H**), Role of the VarS/VarA-CsrA/BCD system in adult mice colonization. The CI values at 5 days post inoculation were shown. The colors of red, green, blue dots represent a higher, lower of colonization than WT or same level, respectively. Significance was determined by *t*-test and one-way ANOVA analysis; *P*-values: *, < 0.05, **, < 0.01, ***, < 0.001, ****, < 0.0001.

Fig. 4. HapR regulates *V. cholerae* colonization through methionine metabolism and transportation. (**A**), Volcano map of the precipitated proteomic of Δ*flhB*/Δ*flaA*. Red dots represent up-regulated expression proteins, and green dots represent down-regulated expression proteins and grey dots represent no significant expression difference (threshold > 1.5, *p* < 0.05). (**B**), Schematic diagram of methionine (Met) metabolism. (**C**), Effect of Met addition (in drinking water) *in vivo* on the colonization of Δ*flhB* and Δ*flaA*. (**D**), Effect of *msrC* in precipitated proteomic on the colonization of Δ*flhB* mutant. (**E**), Effect of indicated mutants on the colonization of Δ*flhB*, Δ*flhB*Δ*msrC*, Δ*flhB*Δ*metR*, Δ*flhB*Δ*metR*Δ*msrC*, Δ*flhB*Δ*metR*Δ*msrC*Δ*metI*Δ*metT*. (**F**), Amino acids and its derivatives were quantified by UPHLC. The column diagram represents the contents of amino acid and its derivatives in Δ*flhB* and Δ*flaA*. The line chart represents the ratio of amino acid and its derivative content of Δ*flhB*/Δ*flaA*. (**G**), Metabolic pathway of methionine converted to other amino acids. Methionine can be converted to homocysteine by demethylation, which subsequently forms cysteine together with serine. Two molecules of cysteine synthesize cystine, while serine itself can be converted to cysteine. The metabolic pathway was annotated by Kyoto Encyclopedia of Genes and Genomes. (**H**), Promoter sequence alignment of *metR*, *msrC*, *metI*, and *metT* with conserved two HapR binding Motifs. Yellow shading represents the conserved binding sequence of Motif 1 and aqua shading represents the conserved binding sequence of Motif 2. (**I**), Effect of *hapR* overexpression on the expression of *metR*, *msrC*, *metI*, and *metT*. The bars represent the corresponding standard deviations. (**J**), EMSA for HapR binding to the promoters of *metR*, *msrC*, *metI*, and *metT*. (**K**), Growth curve of Δ*flhB* and Δ*flaA* in the MM medium with or without 0.01mM L-Met. (**L**), Growth curve of WT, Δ*flhB*Δ*metR*Δ*msrC*Δ*metI*Δ*metT* in the MM medium with or without 0.01mM L-Met. The colors of red, green, blue dots represent a higher, lower of colonization than WT or same level, respectively. The CI values at 5 days post inoculation were shown. The horizontal bars indicate the averages of CI. Significance was determined by *t*-test and one-way ANOVA methods; *P*-values: ns, not significant, *, < 0.05, **, < 0.01, ***, < 0.001.

1016

Fig. 5. Methionine-related metabolic activity affects *V. cholerae* colonization. (**A**),
The workflow for metabolic activity assay with FITC-D-Glu-labeled method in vitro
and in vivo. The figure was created by BioRender. (**B**), Flow cytometry analysis of
wild-type co-with Δ*flhB* and wild-type co-with Δ*flaA* FITC-D-Glu-labeled in transwell
chamber at 30 min. (**C**), The bacterial metabolic activity of wild-type co-with Δ*flhB*
and wild-type co-with Δ*flaA in vivo* at 48h post-inoculation. (**D**), The workflow of
colonization assay of 6N-labeled mutants *in vivo*. (**E**), Colonization assay of 6N-labeled
mutants *in vivo*. The raw data was analyzed by Vsearch and Python. The adjusted CIs
of all flagellar mutants were analyzed by was calculated as the reads of mutant to WT
colonies normalized with the input reads. The colors of red, green, blue dots represent
a higher, lower of colonization than WT or same level, respectively. Significance was
determined by *t*-test and one-way ANOVA methods; *P*-values: ns, not significant, *, <
0.05, **, < 0.01, ***, < 0.001.

1030

1031

1032

1033

Fig. 6. Trade-off mechanisms between motility loss and host adaptability Mechanism. *Vibrio cholerae* predominantly cycles between aquatic reservoirs and the human host. Humans become infected by ingesting water or food contaminated with *V. cholerae*. The bacteria then colonize the small intestine, where they adapt by modulating their gene expression profile. Ultimately, they are shed back into the aquatic environment via feces, completing the transmission cycle and enabling potential new rounds of infection. A systematic analysis of cholera-associated samples collected from diverse environmental niches revealed a recurrent pattern of flagellar gene mutations in *V. cholerae* populations. Follow-up experimental investigations elucidated a fundamental trade-off mechanism, whereby the loss of bacterial motility is counterbalanced by enhanced adaptation to the host. In details, we analyzed 3,135 cholera-related samples and identified a recurrent pattern of flagellar mutations in *V. cholerae*, with notable clustering in genes including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM*. A high-throughput screen of a comprehensive flagellar mutant library established a clear inverse correlation between a strain's capacity to colonize the host and its level of FlgM secretion. Specifically, elevated intracellular concentrations of FlgM inhibit the alternative sigma factor σ^{28} (FliA). This inhibition, in turn, relieves the repression on the transcription of *hapR*, which encodes the master regulator of quorum sensing. The subsequent upregulation of HapR activates genes involved in methionine metabolism and transport. This metabolic reprogramming ultimately enhances the colonization fitness of *V. cholerae* within the host. Employing a specially constructed, competitively matched pool of 6N-labeled mutant strains in an adult mouse model, we provided direct *in vivo* validation. These experiments confirmed that the loss of motility confers a selective advantage for environmental persistence, a phenotype mediated through the defined FlgM-FliA-HapR-methionine regulatory axis.