Anti-σ^{28} factor FlgM secretion regulates *Vibrio cholerae* adaptability in adult mice through quorum sensing and methionine metabolism

Running title: Mechanism of *Vibrio cholerae* fitness in the host gut by FlgM secretion

Guozhong Chen [a,b,1], Zixin Qin [a,1], Fenxia Fan [c,1], Mei Luo [a,1], Hongou Wang [d], Baoshuai Xue [e], Shucheng Li [b], Shiyong Chen [f], Xiaoman Yang [a], Xiang Mao [g], Liwen Yi [a], Chunrong Yi [a], Wei Li [a,*], Xiaoyun Liu [d,h,*], Biao Kan [c,*], Zhi Liu [a,*]

[a]Key Laboratory of Molecular Biophysics of the Ministry of Education/Department of Biotechnology, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China.
[b]State Key Laboratory of Microbial Technology, Shandong University, Qingdao, Shandong 266237, China.
[c]National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China.
[d]Department of Microbiology and Infectious Disease Center, NHC Key Laboratory of Medical Immunology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China.
[e]Key Laboratory of Zoonosis, Ministry of Education, College of Veterinary Medicine, Jilin University, Changchun, Jilin 130015, China.
[f] School of Marine Science and Engineering, Qingdao Agricultural University, Qingdao, Shandong 266109, China.
[g]Institute of Hygiene Toxicology, Wuhan Centre for Disease Prevention and Control, Wuhan, Hubei 430024, China.
[h]Department of Infectious Diseases, Peking University Third Hospital, Beijing 100191, China.

*Corresponding author: Department of Microbiology and Infectious Disease Center, NHC Key Laboratory of Medical Immunology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China. Department of Infectious Diseases, Peking University Third Hospital, Beijing 100191, China. *E-mail addresses*: xiaoyun.liu@bjmu.edu.cn (X. Liu).
National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China. *E-mail addresses*: kanbiao@icdc.cn (B. Kan).
Key Laboratory of Molecular Biophysics of the Ministry of Education/Department of Biotechnology, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China. *E-mail addresses*: hulwei_009@163.com (W. Li), zhiliu@hust.edu.cn (Z. Liu, Tel/Fax: 0086-27-87792072).
[1] These four authors contributed equally to this work.

Abstract

Introduction: Cholera, caused by *Vibrio cholerae*, is a severe diarrheal disease
threatening global health. The flagellum of *V. cholerae* functions not only for motility
but also as an environmental sensor regulating virulence. The trade-off mechanism
between motility loss and enhanced host adaptation remains unclear.

Objective: This study aimed to elucidate how flagellar mutations, which lead to a loss
of motility, affect host adaptability in *V. cholerae* and to uncover the underlying
molecular basis.

Methods: We analyzed 3,135 cholera-related samples to identify mutation hotspots in
flagellar genes. A relevant flagellar mutant library was constructed and assessed for
host adaptability. Multiple approaches, including molecular, genetic, transcriptomic,
and proteomic analyses, were applied to dissect the FlgM-mediated pathway, with key
findings revalidated in an adult mouse model using a specifically constructed
6N-labeled mutant pool.

Results: Big data analysis reveals those flagellar mutations in *V. cholerae* cluster in
key structural and regulatory genes, including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM*. Flagellar
mutations led to a scenario where the secretion level of the regulator FlgM was
negatively correlated with host adaptability. Intracellular FlgM inactivates either the
σ^{28} factor FliA directly or acts through the VarS/VarA-CsrA/BCD system, ultimately
leading to the derepression of the quorum-sensing master regulator named *hapR*.
HapR can directly bind to the promoters of genes involving in methionine
transportation to regulate adaptability of *V. cholerae*. The 6N-labeled mutants pool
experiment reconfirmed that motility loss promotes host adaptability via the
FlgM-FliA-HapR-methionine axis.

Conclusion: Our findings demonstrate that secretion levels of the flagellar regulator
FlgM drive an adaptive shift in *V. cholerae* from motility to host adaptability,
mediated through quorum sensing and methionine metabolic reprogramming. This
reveals a novel mechanism underlying bacterial evolution and pathogenicity.

Keywords: *Vibrio cholerae*, flagellum, FlgM, quorum sensing, methionine
metabolism

Introduction
Cholera, caused by *Vibrio cholerae*, is characterized by severe watery diarrhea
that can be lethal. Infection occurs through the consumption of contaminated food or
water. Since 1817, there have been seven global pandemics that resulted in massive
human deaths, substantial economic losses and considerable healthcare burdens [1].
The WHO reports a large number of cholera cases or outbreaks, from 44 countries in
2022 to 45 countries in 2023, which poses a serious challenge of cholera prevention
and control [2].

The life cycle of *V. cholerae* is closely linked to its pathogenic mechanism. Initially,
the bacteria enter the host via contaminated water, survive gastric acid, and colonize
the small intestine. Subsequently, their flagella facilitate penetration of the mucus
layer, allowing access to the intestinal epithelial surface. As infection progresses, *V.*
cholerae forms biofilms and employs quorum-sensing regulation. Upon reaching high
cell density, the bacteria detach from the epithelial surface, enter the intestinal lumen,
and are ultimately excreted in host feces. They then return to aquatic environments,
thereby completing the transmission cycle [3, 4].
V. cholerae is a Gram-negative bacterium with a single polar flagellum, by which it
swims rapidly and this motility contributes to quick transmission among different
habitats [5,6]. As the propeller of *V. cholerae* movement, the flagellum is a
sophisticated device composed of many different structural and functional
components. Its biosynthesis involves expression of proteins involved in a series of
fine and accurate regulations, acting on transcriptional, translational, and
post-translational levels [7]. Transcriptional regulation of *V. cholerae* genes has been
well-studied, and is built upon a four-tiered transcriptional hierarchy [3]. The
flagellum of *V. cholerae* is also involved in many other biological activities, such as
promoting adherence [8], influencing the infectivity [9], triggering c-di-GMP
production [10], and sensing mucus layer mechanical attraction [4]. Interestingly, a
high number of single nucleotide polymorphisms (SNPs) were detected in the Haiti
and Nepal isolates of the 7[th] cholera pandemic, with 7 out of 54 high quality SNPs
belonging to flagellum-associated genes [11]. Similarly, a large number of mutants
with impaired motility have been isolated from the laboratory [12-17], the natural
environment [18-20], and clinical samples [21]. However, the relationship between
motility loss and host adaptability in *V. cholerae* remains unclear.
Here, we analyzed 3,135 cholera-related samples and identified a recurrent pattern
of flagellar mutations in *V. cholerae*, with notable clustering in genes
including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM*. Screening a flagellar mutant library revealed
an inverse correlation between bacterial colonization capacity and FlgM secretion.
Mechanistically, elevated intracellular FlgM inhibits the σ^{28} factor FliA, which
derepresses transcription of the quorum-sensing master regulator *hapR*. Subsequent
upregulation of methionine metabolism and transport by HapR enhances *V.*
cholerae colonization in vivo. Using a uniquely constructed 6N-labeled mutant pool
in an adult mouse model, we further confirmed that loss of motility promotes host
adaptability through the FlgM-FliA-HapR-methionine regulatory axis. Together, our
data will shed light on trade-off mechanisms between motility loss and host
adaptability.

Materials and methods

In silico analysis flagellar gene mutations in V. cholerae

A total of 3,135 samples in Table S1 were retrieved from the NCBI SRA database by
searching for "cholera" or "Vibrio cholerae" in combination with

metagenomics-related keywords. These valid sequencing runs were selected for
subsequent analysis of flagellar gene mutations. For each dataset, raw reads were
quality-filtered using fastp v0.23.2 [22] and assembled with metaWRAP v1.3.2 [23].
Resulting contigs were binned and taxonomically classified using GTDB-Tk v2.4.0
[24]. Bins identified as *V. cholerae* were retained for protein prediction with Prodigal
v1.2.6 [25]. A custom reference library of 51 flagellar proteins in Table S2 was
constructed based on the *V. cholerae* N16961 genome. Flagellar mutations across
1258 assembled *V. cholerae* bins were then identified in Table S3 and quantified
using DIAMOND BLASTP v0.9.30-3 [26]. As controls, the *hapR* gene (which carries
a frameshift mutation in N16961) served as a positive control, while the housekeeping
genes *asd* (VC2107), *mdh* (VC0432), and *dnaE* (VC2245) were used as negative
controls, respectively [27,28]. The codes for 6N-Label relative abundance analysis,
bins acquisition and species annotation from cohort-based studies, and analyzing
high-frequency mutations in the flagellar genes of *V. cholerae* of this study are openly
available in Zenodo at https://doi.org/10.5281/zenodo.14920964, reference number
14920964.

Bacterial strains, plasmids, and growth conditions

All *V. cholerae* mutants were derived from strain E1 Tor C6706 and cultured in
Luria-Bertani (LB) medium (Qingdao Hope Bio-technology, China), M9 minimal
medium containing 2 mM $MgSO_4$, 0.1 mM $CaCl_2$ and 0.2% casein acid hydrolysate
(Sangon Biotech (Shanghai) Co., Ltd., China), or defined medium containing
appropriate antibiotics (streptomycin 100 μg mL^{-1}, kanamycin 50 μg mL^{-1}, ampicillin
100 μg mL^{-1}, chloramphenicol 25 μg mL^{-1} for *E. coli* and 1.0 μg mL^{-1} for *V. cholerae*)
or 1mM isopropyl β-D-1-thiogalactopyranoside (IPTG) at 37 °C unless otherwise
noted. Antibiotic and IPTG powders were purchased from Sangon Biotech (Shanghai)
Co., Ltd.. This defined medium was prepared with the omission of Met and
supplementation with 0.5% glucose (Sangon Biotech (Shanghai) Co., Ltd., China)
[29]. In-frame deletion mutants were constructed by cloning the regions flanking a
target gene into suicide vector pWM91 that contains a *sacB* counter-selectable marker
[30]. Complementation of *flgM* was achieved by inserting P_{lac}-*flgM* into chromosomal
lacZ using the suicide plasmid pJL1 [31]. Overexpression of *flgM* was obtained by
cloning *flgM-his₆* downstream of the *tac* promoter of pMal-c2x. Plasmid pBAD24
containing a pBAD-inducible promoter was used to overexpress *flgM-his₆*, *flhB*,
flaA-his₆, *fliA-flag*, and *hapR-flag*. The fragment of pBAD-*fliA-flag* was cloned into
pYM24 derived from pACYC117 [32], achieving coexistence with the plasmids
pMal-*flgM-his₆* and pBBR-*hapR-luxCDABE*. All constructed mutants and plasmids
were confirmed by sequencing. All bacterial strains, primers and plasmids used in this
study are listed in Table S4&5.

Motility assay

V. cholerae strains were incubated overnight at 37 °C for 12 h. Single colonies were
inoculated with sterile toothpicks into LB plates containing 0.3% select agar
(Invitrogen, USA), and these soft agar plates were incubated at 37 °C for 8 h.
Photographs were taken and analyzed for colony width as a measure of motility using
a gel imaging instrument (GenoSens 1860, China).

H_2O_2 killing experiments in vitro

Overnight cultures of flagellar mutants were 1:1,000 sub-cultured into M9 minimal
medium containing 2 mM $MgSO_4$, 0.1 mM $CaCl_2$ and 0.2% casein acid hydrolysate
and grown at 37°C, 200 rpm until mid-log phase. H_2O_2 (Sinopharm Chemical
Reagent Co., Ltd., China) was directly added into the cultures at appropriate
concentrations, followed by stationary incubation at 37°C for 30 min. Serial dilutions
of the samples were plated on LB agar for enumeration of colony-forming units
(CFU) and the survival rate in presence of H_2O_2 was calculated by comparing CFUs
of H_2O_2-challenged and unchallenged samples [32].

Biofilm formation

Colonies grown on LB agar plates were resuspended in LB broth and grown to an
optical density at 600 nm (OD_{600}) of ~0.6. A 1:100 dilution was inoculated into 10 ×
75 mm borosilicate glass tubes containing 1 mL M9 minimal medium with 2 mM
$MgSO_4$, 0.1 mM $CaCl_2$ and 0.2% casein acid hydrolysate. Biofilms were allowed to
form by static incubation at 37 °C for 96 h. The biofilms were stained with crystal
violet (Solarbio Life Sciences Co., Ltd., China) and resolved with DMSO (Sinopharm
Chemical Reagent Co., Ltd., China), quantifying with a microplate reader (Spark,
Tecan, Switzerland) at 570 nm.

In vivo colonization assays

The streptomycin-treated adult mouse model was used to assess the competition
ability of mutants or complement strains during co-colonization with C6706 WT.
ROS resistance in the adult mice assay was performed as previously described [33].
For this, five-week-old CD-1 mice (Jiangsu GemPharmatech Co., Ltd., China) were
provided with drinking water with or without 10 g L^{-1} of the antioxidant
N-acetylcysteine (MERCK, Germany) for one week. Afterwards, 5 g L^{-1} streptomycin
and 0.05 g L^{-1} aspartame (Macklin Inc., China) were added to the drinking water. One
day after streptomycin treatment, the mice were administrated with 100 μL of 10%
(w/v) $NaHCO_3$ (Sangon Biotech (Shanghai) Co., Ltd., China) followed by 100 μL of
a 1:1 mixture of WT and mutant (approximately 10^9 CFU for each strain per mouse).
Fecal pellets were collected from each mouse at 1, 3, and 5 days after gavage,
homogenized, serially diluted in LB and plated on LB agar containing streptomycin
and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal) (Sangon Biotech

(Shanghai) Co., Ltd., China) to quantify the bacterial loads. The competition index CI
was calculated as the ratio of mutant to WT colonies normalized with the input ratio.
Five-day-old CD-1 mice were transferred to an incubator and left to adapt for 2 h at
30 °C before inoculation. The mice were then administrated with a 50 μL of a 1:1
mixture of WT and mutant *V. cholerae* (approximately 10^6 CFU for each strain per
mouse) and returned to the 30 °C incubator. The animals were sacrificed 18 h after
gavage. Samples of the small intestine were removed and homogenized in 1.5 ml of
LB, serially diluted, and plated on LB agar containing streptomycin and X-Gal for
quantification of bacterial load. The CI was calculated as the ratio of mutant to WT
colonies normalized with the input ratio.

Growth curve determination and luminescence detection

Overnight LB cultures of targeted strains (with or without plasmid) were inoculated in
fresh medium at a ratio of 1:100 and grown with shaking to mid-log phase at 37 °C.
The bacteria were harvested by centrifugation, washed three times and resuspended to
an OD_{600} of 1.0 with fresh pre-warmed medium (LB or M9), and then diluted 1:1,000
in 96-well microplates that were incubated at 37 °C under shaking. The growth
kinetics were determined by measuring OD_{600} or OD_{655} using a microplate reader
(Spark, Tecan, Switzerland) combined (where applicable) with luminescence
detection. If necessary, 0.1 mM isopropyl-β-D-thiogalactoside (IPTG) or 0.2%
arabinose was added. Readouts were performed every 30 min for 20 h. The
luminescence assay was standardized at OD_{600}. All experiments were performed in
triplicate.

Transmission electron microscopy (TEM) imaging

Single colonies of target strains were inoculated into 50 μL of saline and gently mixed
with a clipped, sterile pipette tip. Five microliters of sample were added to a copper
supporting mesh. Excess liquid was removed after standing in a ventilated and clean
environment for 5 min. The mesh was stained with freshly prepared 1%
phosphotungstic acid (Sangon Biotech (Shanghai) Co., Ltd., China) for 2 min.
Excessive dye was removed and the mesh was dried for approximately 1 h prior to
transmission electron microscopy (HT7700, HITACHI, Japan).

Secretion of FlgM

To investigate the secretion of FlgM, culture supernatant of the *V. cholerae* mutants
containing pBAD24-*flgM-his₆* was collected and the protein was detected according
to a previous method with slight modification [4]. All mutant sediment samples were
standardized with CFU (10^9). Proteins in the 400mL supernatant were concentrated
with 20% trichloroacetic acid (TCA) (Sinopharm Chemical Reagent Co., Ltd., China).
These, together with the proteins in the precipitate and supernatant were separated by
SDS-PAGE, transferred to a polyvinylidene fluoride (PVDF) membrane (Millipore,

USA). The proteins on the Western blot were detected with primary $6\times$ His-tagged rabbit antibodies (Sangon Biotech (Shanghai) Co., Ltd., China) at a dilution of 1:5,000, and as secondary antibody horseradish peroxidase [HRP]-conjugated goat anti-rabbit immunoglobulin G (Sangon Biotech (Shanghai) Co., Ltd., China) was used at a dilution of 1:10,000. Immunoblotting was performed using the Pierce ECL Western Blotting Substrate Kit (Thermo Scientific, USA).

Proteomic analysis

Overnight LB cultures were inoculated 1:100 in fresh LB medium and incubated until OD_{600} reached 0.9 (approximately 10^9 CFU per sample). Then the culture was centrifuged. The pellets were for bacterial proteome samples. For the analysis of the bacterial secretome, the 400 mL of culture supernatant was filtered through a 0.22 μm microporous membrane to remove intact bacteria. The proteins in the filtrate were precipitated overnight at 4 °C with 20% TCA containing 0.5 mM sodium deoxycholate. After centrifugation at 10,000 g for 1 h at 4 °C, the protein pellets were washed three times with cold acetone for LC-MS/MS experiments, that were carried out as previously described [32]. The secretome experiments were performed in duplicates, and bacterial proteome samples were analyzed in triplicates. Proteins with Fold changes >1.5 and *P*< 0.05 were considered as significant differences.

Gene expression analysis via RT-qPCR

Overnight LB cultures were inoculated 1:100 in fresh LB medium containing appropriate antibiotics or, if needed, 0.2% arabinose, and incubated at 37 °C. RNA for real-time PCR (RT-PCR) was extracted using an RNA extraction kit (Vazyme Biotech Co., Ltd. China). The RT-PCR reaction was performed using SYBR green fluorescent dye (Yeasen Biotechnology (Shanghai) Co., Ltd., China) and CFX Connect Real-time Detection System (Bio-Rad, USA). The genes *recA* or *secA* or 16S rRNA were used for normalization. Relative expression levels were calculated using the $2^{-\Delta\Delta CT}$ method. All experiments were performed in triplicate. For reverse transcription, approximately 400 ng of RNA was extracted using the HiScript II 1st Strand cDNA Synthesis Kit (Yeasen Biotechnology (Shanghai) Co., Ltd., China). The RT-qPCR primers used for detection of mRNA of specific genes are listed in Table S5. All primers were tested for PCR specificity and efficiency.

Protein purification

The proteins HapR-His$_6$ were expressed in *E. coli* BL21 (DE3) and purified using the pET system and BeyoGold™ His-tag Purification Resin (Beyotime, China).

EMSA

The purified recombinant proteins HapR containing 6×His-tags were used for EMSA. The promoters of their target genes were amplified by PCR and purified. These

amplicons (50 ng) were incubated with 6× His-tagged protein in 20 μL containing
4μL of 5× binding buffer (50 mM Tris-HCl, 0.25 M KCl, 2.5 mM DTT, 5 mM $MgCl_2$,
0.25 mg mL^{-1} BSA, 2.5 mM EDTA, 1% glycerol, pH 8.3) (Sangon Biotech
(Shanghai) Co., Ltd., China) at 25 °C for 30 min. DNA-protein complexes were
analyzed by electrophoresis on a 5% native polyacrylamide gel at 4 °C and 80 V cm^{-1}
for 1.5 h. After staining the gel in ethidium bromide for 10 min, a gel imaging system
(GenoSens 1860, China) was used to produce images. The EMSA primers used in this
study are listed in Table S5.

Synthesis of fluorescent monomers

FITC-D-Glucose was synthesized as previously described [34]. Briefly,
D-glucosamine hydrochloride (77.6 mg, 0.36 mmol, 1.2 equiv) and Et3N (91.1 mg,
0.9 mmol, 3.0 equiv) were mixed in DMF (10 mL) (Sinopharm Chemical Reagent Co.,
Ltd., China). After 1 h of stirring, fluorescein (116.8 mg, 0.3 mmol, 1.0 equiv) was
added. The reaction mixture was stirred for 16 h at RT in the dark, after which the
solid product was obtained by filtering and concentrated *in vacuo*.

Bacterial metabolic activity assay in vitro

Bacterial cultures in LB were harvested at OD_{600} ~0.4 by centrifugation and the cell
pellet was washed 3 times with Met-free MM medium. The cells were resuspended
and incubated at 37□ for 30 minutes. A solution of 10 mL Met-free MM medium and
1 μL of FITC-D-Glu was added and 500 μL of this was added to transwell wells and
200 μL of this was added to the transwell chambers. To each well, 100 μL of bacterial
suspension was added and the plates were incubated for 30 min at 37 °C in the dark.
The supernatant was washed off with PBS buffer and then assayed by Flow
Cytometry (Agilent Technologies Inc., USA).

Bacterial metabolic activity assay in vivo

Bacterial metabolic activity was assayed in vivo as previously described with
following modifications [35]. Briefly, the CD-1 mice were administered with WT and
mutants for competition colonization experiments as described above. At 48 h
post-inoculation, the mice were administered with 100 μL FITC-D-Glu (0.1 mM) in
PBS (Sangon Biotech (Shanghai) Co., Ltd., China). Two hours after the gavage, the
animals were sacrificed and the small intestine was dissected and rinsed with 2 mL
PBS, cut longitudinally and the intestinal content was scraped off using a cell scraper.
The intestinal content was suspended into a 1.5 mL PBS buffer with 2 steel beads and
shaken to release bacteria. Centrifugation at 3000 rpm for 1 min removed large tissue
fragments and impurities, and the supernatant was filtered through 40 μm cell
strainers to remove smaller cell debris. The filtrate was centrifuged (8,000 rpm, 5
min), the bacterial pellets were washed twice with 1.5 mL PBS and resuspended in
PBS for flow cytometry analysis. In the FITC channel, cells with strong signals of

approximately 10% were selected for sorting, with 10^5 events obtained for each
sample, and the collected samples were diluted and plated on LB agar containing
streptomycin and X-Gal for quantification of the bacterial index.

Detection of amino acid profiles in ΔflhB and ΔflaA by HPLC and MS

Overnight LB cultures of ΔflaA and ΔflhB were sub-cultured with an inoculum of
1:100 in MM medium and shaken at 37 °C until OD_{600}=0.5. The bacteria were
collected by centrifugation (>200mg) and samples of 60 mg were suspended in 50 μL
water, vortexed for 60 s, and mixed with 400 μL methanol-acetonitrile solution (1:1,
v/v) and 50 μL of a mixture of 16 isotopic internal standards (50 μM). The mixture
was vortexed for 60 s and twice subjected to low-temperature ultrasound for 30 min.
Proteins were precipitated at -20 °C for 1 h and after centrifugation at 14,000 g and
4 °C for 20 min and the supernatant was collected, freeze-dried, and stored at -80 °C
prior to high-performance liquid chromatography (HPLC). HPLC was performed
using an Agilent 1290 Infinity LC ultra-high-performance liquid chromatography
system (USA). The mobile phase consisted of 25 mM ammonium formate + 0.08%
formic acid (solution A) and 0.1% formic acid acetonitrile (solution B). The samples
were loaded into an automatic sampler at 4 °C, and the column temperature was set at
40 °C. The flow rate was 250 μL min^{-1}, and the injection volume was 1 μL. The
gradient elution was as follows: 0-12 min, linear change from 90% B to 70% B; 12-18
min, linear change from 70% B to 50% B; 18-25 min, linear change from 50% B to
40% B; 30-30.1 min, linear change from 40% B to 90% B; 30.1-37 min, maintained at
90% B. At regular intervals between experimental samples a quality control (QC)
sample was inserted to evaluate the stability and reproducibility of the system.
Mass spectrometry (MS) analysis was performed using a 5500 QTRAP mass
spectrometer (SCIEX) in positive ion mode. The ESI source conditions for the 5500
QTRAP were as follows: source temperature 500 °C, ion Source Gas 1 (Gas 1): 40,
Ion Source Gas 2 (Gas 2): 40, Curtain gas (CUR): 30, IonSapary Voltage Floating
(ISVF) 5,500 V. The analytes were detected using the MRM mode.

In vivo colonization assay of 6N-labeled mutants

Briefly, using C6706 (*vc1807*:: CmR) as a template, amplify the CmR fragment and
insert a 6N site as a marker in the middle of one end primer (6N-R) [36], then clone it
into the pWM91 vector. Sequence tags of 6N were inserted into the *lacZ* position of
the 26 mutants and the WT strain with the 6N-labeled plasmids pWM91-CmR-6N by
conjugation experiments. After labeling the bacterial strains, confirmed by sequencing,
they were sub-cultured and adjusted to an OD_{600} of 1.0. After harvesting by
centrifugation, the bacterial pellets were stored at -20°C. Using the
methionine-supplemented and water-restricted mouse model, the mice were gavaged
with 100 μL of the mixed bacterial samples per mouse. On day 5 after gavage, mouse
feces were collected and total DNA was extracted, along with the input samples.

Amplicons produced by PCR with primers targeting both ends of *LacZ* were
sequenced. The sequenced raw data was analyzed by Vsearch for merging paired-end
reads and Python to calculate the proportion of reads from the 6N-labeled mutants
among the total reads.

Statistical analysis

Results are presented as mean ± standard deviation. When two groups were compared,
t-tests were performed. One-way analysis of variance (ANOVA) was used for two or
more treatment groups. Statistical analysis was performed using the software
GraphPad Prism v.9 for Windows. Differences with a p value less than 0.05 were
considered to be statistically significant.

Results

High-frequency mutations in the flagellar genes of V. cholerae

The life cycle of *V. cholerae* primarily alternates between aquatic environments and
the human host [3]. To investigate its adaptation across these niches, we categorized
cholera-related metagenomic sequencing samples from the SRA database in Table S1
into five groups based on their origin: human, clinical, aquatic (including water and
sewage), fish, and amoeba-derived. These samples were collected from 19 countries
(Fig. 1A). A total of 1,258 metagenome-assembled genomes (MAGs) of *V.*
cholerae in Table S3 were reconstructed from these samples. Their flagellar protein
mutations were then systematically analyzed using the 51 flagellar proteins of *V.*
cholerae N16961 in Table S2 as reference amino acid sequences. For quality control,
the mutation-prone *hapR* gene, which carries a frameshift mutation in the N16961
genome, served as a positive control, while the conserved housekeeping
genes *asd* (VC2107), *mdh* (VC0432), and *dnaE* (VC2245) were used as negative
controls, as established in prior studies [27,28].
Our mutation frequency analysis revealed that across all sampled habitats, flagellar
gene mutations in *V. cholerae* consistently clustered in key genes,
including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM* (Fig. 1B). Notably, mutations were also
detected in the primary flagellar regulator gene *flrA*, a finding consistent with
previous reports [16,37]. Functionally, these mutated genes are involved in critical
flagellar structures and regulatory hierarchies, including Class I and IV transcriptional
regulation, the type III secretion system (T3SS) apparatus, the filament-capping
complex, and the C-ring switch structure (Figure 1C and D). This indicates that these
loci have key evolutionary and pathological significance in regulating the host
adaptability balance.

Phenotypic screening of the flagellar mutant library

To elucidate the environmental adaptability conferred by the flagellar gene mutations
identified through the above big data-driven analysis, a systematic mutant library
targeting 26 flagellar-related regulons of *V. cholerae* listed in Table S4 was
constructed using the primes in Table S5. Subsequently, this library was subjected to
comprehensive phenotypic screening, assessing motility, biofilm formation, growth
rates in both rich and minimal media, colonization efficiency in infant and adult mice,
as well as resistance to H_2O_2. The H_2O_2 assay is particularly relevant as it reflects
bacterial survival under host-derived reactive oxygen species (ROS) stress. Of the
investigated mutants, 22 mutants had completely lost their motility. Deletion of
chemotaxis-related genes *cheV-3*, *flgO* and *flgP* resulted in significantly impaired
motility (Fig. 2A, Fig. S1), which is consistent with previous reports [6, 38]. The
colonization ability in infant mice was seriously damaged (Fig. S2) and biofilm
formation of the 26 mutants were mostly enhanced, which were consistent with earlier
findings (Fig. S3) [5,6,10]. Surprisingly, we observed completely opposite results in
the adult mice, as some of the mutants were able to outcompete the WT strain (Fig.
2B), although their growth rates were consistently lower than that of WT in minimum
or rich medium (Fig. S4 and 5), and others were attenuated. In total, five mutants
(Δ*flgBCDE*, Δ*fliK*, Δ*fliKLMNOPQRflhB*, Δ*fliE,* and Δ*flgP*) produced no significance
difference in CI (blue in Fig. 1D), nine mutants (Δ*flrA*, Δ*rpoN*, Δ*flrC*, Δ*fliEFG*,
Δ*fliHIJ,* Δ*fliA*, Δ*flgFGHIJ,* Δ*fliPQR*, and Δ*flhB*) displayed enhanced CI (red in Fig.
2B), and twelve mutants (Δ*flhA*, Δ*flhA* operon, Δ*flgB*, Δ*motY*, Δ*flaA*, Δ*flgK*, Δ*flgO*,
Δ*flgT*, Δ*cheV-2*, Δ*vc1384*, Δ*flgM*, and Δ*cheV-3*) were attenuated for CI (green in Fig.
2B). Of these, the Δ*flgFGHIJ* mutant produced the highest CI with a 1000-fold
increased CI compared to WT, while the Δ*flgO* mutant had the lowest CI (1/1000-fold
of WT). Of the other mutants, Δ*flhB*, impaired in production of the flagellar type III
secretion system (fT3SS) protein FlhB, was enhanced in adaptability, and Δ*flaA*,
missing the major flagellin filament protein, was attenuated. We furtherly analyzed
the various phenotypes of all 26 mutants in more detail, and found that only the
colonization potential in the adult mouse model differed among the mutants. All other
phenotypes (e.g. motility, biofilm development, proliferation in LB/M9, H_2O_2
resistance, infant mice colonization) exhibited a unipolar distribution. Correlation
analysis indicated that adult mice colonization was not associated with other
phenotypes (Fig. S6).

FlgM secretion negatively regulating adaptability

Considering FlaA and FlhB are structural flagellar component without a regulatory
function, they were confirmed through transmission electron microscopy that the 2
mutants lacked flagella (Fig. S7A), and that overexpression of the relevant genes
restored their motility capability (Fig. S7B). So, we choose Δ*flhB*, Δ*flaA* and WT
normalized with CFU (Fig. S7C) were to explore the underlying mechanism of these
opposite colonization properties by sequenced proteins in the supernatant (Fig. 2C).
These proteins from Δ*flhB* and Δ*flaA* were compared to a WT normalized control, in

an attempt to seeking differentially expressed proteins. This detected 36 proteins that
were differently secreted between the Δ*flhB* and Δ*flaA* mutants (Table S6, Fig. S7D),
including 15 flagellar assembly proteins, 9 cellular processes related proteins, 2
secreted hemolysin proteins, 2 cell membrane associated proteins, 2 reactive oxygen
species (ROS)-resistance proteins, and 6 other proteins.

Hemolysins HlyA and Tlh are secreted toxin proteins that play a role in the
virulence of *V. cholerae* [6, 39]. However, their deletions in Δ*flhB* to give Δ*flhB*Δ*hlyA*
and Δ*flhB*Δ*tlh*, Δ*flhB*Δ*katG* and didn't affect their CI in the adult mice model (Fig.
S8A and B). Colonization in this model can also be enhanced by ROS-resistance, for
which KatG and catalase are held responsible [32]. Experiments were performed to
investigate this, for which N-acetylcysteine (NAC) was employed to remove ROS
from the murine intestine (Fig. S8C), but this revealed no difference in the CI of
Δ*flhB* or Δ*flaA* compared to that in absence of NAC (Fig. S8D and E). As not all
mutants but the majority are resistant to ROS, this did not seem to explain the
enhanced colonization of particular flagellar mutants (Fig. S9).

Interestingly, the secretion of FlgM in Δ*flhB* was reduced about 9-fold compared
with that of Δ*flaA* (Fig. 2C). FlgM is an anti-σ^{28} factor, and functions as a chaperone
protein of σ^{28} factor FliA [40]. Our previous research had demonstrated that FlgM can
be secreted by Δ*flgD* through broken flagella, which activates FliA activity, to then
repress the expression of *hapR* that subsequently upregulates the expression of
virulence gene *tcpA* and enhances *V. cholerae* colonization in infant mice [4].
Introduction of a double-mutation where *flgM* was knocked out in a Δ*flhB* background
dramatically reversed the Δ*flhB* colonization pattern, from a CI of 176.3293 for Δ*flhB*
to 0.0098 for Δ*flhB*Δ*flgM* (*P*= 0.0005) (Fig. 2D). In addition, the overexpression of
FlgM in Δ*flaA* (*lacZ*::P$_{lac}$-*flgM*) also significantly reversed the phenotype of Δ*flaA*,
from CI=0.0130 for Δ*flaA* to 0.2285 for Δ*flaA* (*P*= 0.0146) (Fig. 2E). Based on these
observations, we analyzed FlgM secretion in all 26 mutants and found that this
negatively correlated (R^2=-0.4322, *P*=0.0275) with their CI in adult mice (Fig. 2E and
F). Together, these data suggest that changes in intracellular FlgM levels in the
various flagellar mutants may contribute to their diverse colonization properties in
adult mice.

Intracellular FlgM drives hapR expression

V. cholerae employs cell-to-cell communication, so-called quorum sensing, of which
HapR is the master regulator, which is in charge of many important regulation
activities in response to different environmental signals [41, 42]. In the Δ*flhB* mutant,
a high concentration of intracellular anti-σ^{28} factor FlgM is available to bind to σ^{28}
factor FliA, thereby inhibiting its function; this leads to an increase in *hapR*
transcription (Fig. 3A, left) [4]. We sequentially knocked out *flgM*, *fliA*, and *hapR* in a
Δ*flhB* background. Whereas the double mutant Δ*flhB*Δ*flgM* had a reduced CI (0.0010)
compared to Δ*flhB* (CI=210.5072, *P*=0.002), this phenotype was more than reversed
in the triple mutant Δ*flhB*Δ*flgM*Δ*fliA* (CI=616.9768), whereas the quadruple mutant

Δ*flhB*Δ*flgM*Δ*fliA*Δ*hapR* had a reduced CI (1.408) again (Fig. 3A, right). The order of
sequential mutation was then reserved by inactivating (in that order) *flgM*, *fliA*, and
hapR in a Δ*flaA* background, which resulted in similar switches in colonization ability
(Fig. 3B). This suggests that *hapR* expression may be responsible for the different
colonization abilities of these flagellar mutants in the applied adult mouse model.

This hypothesis was confirmed by determining that the transcription of *hapR* was
significantly upregulated in those mutants with enhanced colonization, and
significantly downregulated or unchanged in the other flagellar mutants (Fig. 3C).
And In addition, the knockout of *hapR* significantly decreased the CI (Fig. 3D),
suggesting that a FlgM-FliA-*hapR* pathway contributes to the colonization potential.

We found that overexpression of FliA (by means of pBAD-*fliA*) in Δ*flgM*Δ*fliA*
significantly repressed *hapR* expression (*P*=0.020), while overexpression of FlgM in
Δ*flgM*Δ*fliA* derepressed *hapR* transcription (*P*=0.003) (Fig. 3E). Similar results were
obtained with *E. coli* DH5α (Fig. S10), suggesting that *hapR* expression may be
regulated by FliA directly.

Previous studies had reported that the VarS/VarA-CsrA/BCD pathway, three
autoinducer systems (CAI-1, AI-2, DPO), and VqmA can regulate *hapR* expression
(Fig. 3F) [43-45]. We determined the expression of each key gene within the quorum
sensing network in the Δ*fliA* mutant, which suggested that FlgM-FliA may indirectly
inhibit HapR degradation through the VarS/VarA-CsrA/BCD pathway (Fig. 3G). To
compare the priority of direct and indirect regulation of *hapR* by FlgM-FliA, the *varA*
and *hapR* genes were individually knocked out in a Δ*flhB* background. Adult mice
colonization results revealed that this drastically reduced colonization compared to
Δ*flhB* (*P*=0.0026 for *varA* and *P*=0.0023 for *hapR*); no significantly difference was
observed between Δ*flhB*Δ*varA* and Δ*flhB*Δ*hapR* (Fig. 3H). The double mutation of
varA and *flhB* consistently decreased the expression level of *hapR* (Fig. S11). Taken
together, it is concluded that interaction between intracellular FlgM and FliA
derepresses *hapR* transcription, resulting in an enhanced colonization ability of *V.*
cholerae in adult mice.

HapR monitors methionine metabolism

We next compared precipitated protein profiles of Δ*flhB* and Δ*flaA*, using WT as a
normalization control in Table S7. The cellular protein levels of flagellin proteins
FlaD, methionine sulfoxide reductase MsrC, serine protease VesC, and hypothetical
proteins VC1075 were upregulated (Fig. 4A). However, no significantly difference
was observed among Δ*flhB* and Δ*flhB*Δ*flaD,* Δ*flhB*Δ*vesC,* Δ*flhB*Δ*vc1075* in the adult
mice (Fig. S12). The MsrC was the highest enriched cellular protein in Δ*flhB*
(Log$_2$Ratio=3.000, *P*=0.002), relative to Δ*flaA* (Fig. 4A). Therefore, we now
hypothesize that cellular methionine contributes to the different colonization patterns
of the 26 flagellar mutants in adult mice.

In *V. cholerae*, cellular methionine metabolism is finely regulated via biosynthesis,
transformation and transportation systems [Fig. 4B] [46]. As expected, the addition of

methionine significantly reduced Δ*flhB* and it increased Δ*flaA* colonization in adult
mice (Fig. 4C), whereas Δ*msrC* drastically reduced the colonization advantage (Fig.
4D, CI=0.015). However, knockout of the methionine biosynthesis genes *msrC* or/and
metR in a Δ*flhB* background didn't reduce colonization advantage, unless the
methionine transportation genes *metI* and *metT* were also inactivated
(Δ*flhB*Δ*msrC*Δ*metR*Δ*metI*Δ*metT vs.* WT compared to Δ*flhB vs.* WT, FC=6.341,
P=0.001) (Fig. 4E). This suggests that the cellular methionine yield is precisely
regulated by a range of metabolic and transportation genes in *V. cholerae*, which is
not surprising, since methionine is an essential amino acid for bacterial metabolism
and proliferation. Subsequently, the amino acid content in the precipitates of Δ*flaA*,
Δ*flhB*, and WT were quantified. We found lower serine concentration (*P*=0.040) and
higher concentrations of cystine (a disulfide of two oxidized cysteine molecules,
P=0.018) in Δ*flhB* compared with Δ*flaA*, but no variation in methionine
concentrations could be demonstrated by intracellular amino acid assays (Fig. 4F).
Beyond bacterial biosynthesis and uptake, intracellular methionine can also be
increased via other amino acid pathways. The KEGG pathway shows that
L-cystathionine, a downstream product of serine and methionine, can be converted to
cysteine, from which cystine is formed (Fig. 4G) [47], so that a higher active
methionine metabolism in Δ*flhB* may relate to lower serine concentrations and higher
cystine concentrations. Interestingly, we also found that *hapR* is involved in the
expression of enzymes related to other amino acid metabolism by analyzing the
transcriptome from our Mr.VC database [48, 49], including serine and cysteine
metabolism, suggesting that *hapR* may be involved in the regulation of methionine
metabolic flow.

Tsou et al. have described that HapR is a global regulator in *V. cholerae,* playing an
important role in virulence regulation, biofilm development, natural transformation,
and stress response [42]. By means of sequence alignments we identified sequences
characteristic of the HapR-binding motif 1 in front of *metT*, *msrC*, *metI*, and a
HapR-binding motif 2-like sequence in front of *metR* (Fig. 4H). Indeed, HapR directly
promotes the transcription of these four genes, as demonstrated by RT-qPCR (Fig. 4I),
and binding of HapR to promoter DNA was demonstrated by EMSA experiments (Fig.
4J, Fig. S13, Table S5). We observed that it rescued the growth defect of Δ*flaA* in
methionine-limited medium (MM) (Fig. 4K) and adding methionine to the medium
did not revive the growth of Δ*flhB*Δ*metR*Δ*msrC*Δ*metI*Δ*metT* (Fig. 4L).

Methionine metabolism affects V. cholerae colonization

To examine the metabolic activity of Δ*flhB*, Δ*flaA* and WT in vitro, we used
FITC-labeled glucose method (Fig. 5A). The metabolic activity is often labeled by
marking bacterial cell wall precursors [50, 51], such as GlcNAc-1-phosphate, which is
originally derived from glucose [52]. Fluorescein Isothiocyanate (FITC) is a
commonly used fluorescent probe molecule that can form a covalent complex with
glucose molecules. After FITC-labeled glucose is utilized, the detection of

fluorescence signals can indicate bacterial metabolic activity. A transwell co-culture
system with WT showed that the population of Δ*flhB* was highly metabolically active
and 1.7-fold higher than WT (P=0.012), while Δ*flaA* was 0.6-fold lower (P=0.003)
than WT (Fig. 5B). We also investigated bacterial metabolism in vivo by
administering FITC-labeled glucose to mice that were colonized with either a Δ*flaA*
and WT mixture or a Δ*flhB* and WT mixture, after which the bacterial content was
isolated from their intestine and subjected to flow cytometry to sort those bacterial
cells with high metabolic activity. These were enumerated on X-Gal plates, which
revealed that the population of Δ*flhB* was 7-fold higher (P=0.017), while the
population of Δ*flaA* was 0.2-fold lower compared to WT (P=0.001) (Fig. 5C). Thus,
both in vitro co-culture and in vivo in adult mice co-colonization experiments
provided strong evidence that bacterial metabolic activity contributed benefited the
bacteria. In combination with the other results, this suggest that differences in
methionine-related metabolic activity may contribute to the different colonization
patterns of the flagellar mutants.

Ecological simulation of flagellar mutants reveals trade-off mechanisms between
motility loss and host adaptability

To assess the host adaptability of the flagellar mutants, the previously characterized
mutants were labeled using a novelty developed 6N-labeling method (using the
primes in Table S5) and subsequently administered to adult mice via oral gavage.
Colonization competition was then assessed by high-throughput sequencing (Fig. 5D,
Table S8). In the absence of methionine supplementation, the competition indices
(CIs) of all 26 mutants showed a scattered and diverse distribution, consistent with the
head-to-head colonization outcomes presented earlier (Fig. 2B). Upon methionine
addition, the distribution of CIs for all flagellar mutants became markedly more
constrained, as detailed in Supplementary Dataset 6. Further pairwise analysis
revealed that methionine supplementation could reverse the colonization pattern of
individual flagellar mutants (Fig. 5E). Collectively, these findings demonstrate that
ecological simulation of flagellar mutants reveals trade-off mechanisms between
motility loss and host adaptability.

Discussion

By employing big data analysis on *V. cholerae* life cycle samples, this study
elucidates the trade-off mechanism whereby flagellar mutations result in motility loss
yet enhance host adaptability through the FlgM-FliA-HapR-methionine axis, a driver
of metabolic reprogramming (Fig. 6).

Although flagellar mutations are prevalent across the life cycle environments of *V.*
cholerae, their evolutionary paths across distinct host niches remain unclear. Previous
work indicates that the cyclin gene *dps*, rather than DNA repair or ROS/RNS
scavenging systems, significantly increases flagellar mutation frequency in an adult

mouse model [15]. In amoeba hosts, mutations in the transcriptional
regulator *flrA* provide a competitive edge, while intense microbial competition in the
gut may further shape *V. cholerae* evolution [16]. In addition, the *flrA* in *V.*
cholerae senses intracellular levels of the second messenger cyclic diguanylate
monophosphate (c-di-GMP) in response to dietary signals. Under a high c-di-GMP
environment induced by casein, FlrA represses the expression of the type VI secretion
system (T6SS), thereby impairing the ability of *V. cholerae* to compete with intestinal
commensal bacteria via T6SS and ultimately leading to a colonization defect in
casein-fed mice [53]. Furthermore, intense microbial competition in the gut may serve
as a major driving force for the evolution of *V. cholerae* within the intestinal
environment [54]. Integrating comparative genomics with experimental evolution
assays may unravel whether these mutations represent adaptations to host niches or
collateral effects of pathogen transmission dynamics.

Methionine is an essential amino acid utilized by bacteria in various physiological
and metabolic processes, including polyamine synthesis, DNA synthesis, redox
balance, and DNA and histone methylation [55]. Also, it is important to note that
methionine plays an essential role in the immune system through its metabolites. For
example, in response to antigens, T cells activate the methionine cycle to support
proliferation and differentiation in cancer, indicating the importance of the methionine
cycle to T cell immunity [56]. Methionine supplementation or restriction can intervene
in the natural antioxidant capacity of an organism by leading to the production of
endogenous enzymes that reduce oxidative stress and DNA damage [57]. These data
imply that *V. cholerae* may employ the methionine pathway as a means of immune
evasion. To successfully propagate and cause disease, *V. cholerae* also evades the
immune system by repressing the mannose-sensitive hemagglutinin pilus via the
transcription factor ToxT, preventing secretory IgA binding and enabling intestinal
colonization to bypass host defenses [58].

Methionine metabolism is a central node within the interconnected amino acid
network of *V. cholerae* [59]. Beyond its metabolic functions, methionine serves as a
precursor for S-adenosylmethionine (SAM), a cofactor involved in critical processes
including the synthesis of the quorum-sensing autoinducer CAI-1 [60,61]. Similarly,
threonine, another methionine-derived metabolite, may contribute to autoinducer DPO
synthesis, and in *Pseudomonas aeruginosa*, SAM participates in producing AI-1, AI-2,
and AHLs [62]. Analyzing methionine flux could therefore help identify key
metabolites that enhance colonization in flagellar mutants, and future work should
examine whether intermediate methionine metabolites or other amino acids
influence *V. cholerae* colonization through bacterial or host metabolic remodeling.

This study found that HapR binds to promoters of methionine metabolism-related
genes (*metR*, *msrC*, *metI*, *metT*), a result not reported in prior systematic screens of
HapR targets. The discrepancy may stem from limitations in bioinformatic prediction

accuracy and variability in HapR-binding sequences [42]. Beyond methionine regulation, HapR suppresses toxin-coregulated pilus (TCP) and cholera toxin (CTX) genes by repressing *aphA*, while activating extracellular proteases and modulating T6SS genes [63-65]. However, in adult and infant mouse models, this virulence factors do not substantially affect colonization [39,66]. HapR also inhibits biofilm formation via cyclic di-GMP modulation and *vpsT* repression [67-69], although flagellar mutants show enhanced biofilm capacity, they still exhibit colonization defects, indicating biofilm is not a major driver of colonization in these strains. Additionally, HapR and RpoS jointly facilitate stress adaptation, luminal dispersal, and host exit. Although the flagellar mutants of *V. cholerae* exhibit enhanced biofilm formation capacity, they still display a colonization defect (Fig. S3, Fig. 2B), suggesting that biofilm formation is also not a major factor contributing to enhanced colonization in these mutants. In addition, HapR and RpoS collectively facilitates stress adaptation to host factors, dispersal within the intestinal lumen, and eventual exit of *V. cholerae* from the host [70]. In the precipitated proteome, we observed that MsrC was the most highly enriched cellular protein in $\Delta flhB$ (Log2Ratio = 3.000, P = 0.002) relative to $\Delta flaA$ (Fig. 4A). Through experimental validation, we concluded that methionine contributes to the differential colonization patterns among the 26 flagellar mutants in adult mice.

The flagellum renders bacteria motility, but it is also a major antigen recognized by the host [71, 72]. Extensive studies have shown that flagellar mutants of *V. cholerae* are frequently present in clinical and environmental samples, and specifically the *V. cholerae* Haiti strains contain multiple mutational hotspots in flagellum gene regions according to SNP analysis [11]. Besides mutations, *V. cholerae* also actively loses flagellar filaments during environmental changes such as mucus penetration [4], biofilm development [10], and host Lypd8 promotes the segregation antibody capture [72]. This is also demonstrated for other pathogenic bacteria. The freshwater bacterium *Caulobacter crescentus* loses its flagella through ejection trigged by ClpA [73]. The loss of the polar flagellum in *V. fischeri* and in plant-associated *Methylobacteria* has also been reported [74, 75]. Bacterial flagella also detach during starvation [76].

We demonstrated that the trade-off mechanism between motility loss and host adaptability operates via the sophisticated FlgM-FliA-HapR-methionine metabolic reprogramming axis. However, this study has the following limitations: (1) The mechanistic focus on the FlgM-FliA-HapR-methionine axis, though insightful, may not capture the full complexity of the flagellar regulatory network, and deeper metabolic validation is needed; (2) It does not directly simulate or verify how this trade-off process dynamically occurs and becomes fixed within a population under environmental pressure, lacking direct experimental evolution evidence in human; (3) The adaptive efficacy and stability of the revealed mechanism in such highly complex real-world environments require further evaluation; (4) Time-course experiments

were not provided to track the changes in FlgM secretion, HapR expression, and
methionine metabolism over the course of infection.

5. Conclusion

Based on extensive sample analysis, this study reveals that flagellar mutations in *V.*
cholerae cluster in key structural and regulatory genes, underscoring their central role
in host adaptation. Specifically, the secretion level of the flagellar regulator FlgM
drives a phenotypic shift from motility to host adaptability. This transition is mediated
through the FlgM-FliA-HapR-methionine axis, which balances the loss of motility
with enhanced host adaptability. Ultimately, impaired motility promotes
environmental fitness by activating quorum sensing and reprogramming methionine
metabolism, elucidating a novel evolutionary trade-off strategy between movement
and adaptability at the molecular level.

Credit Authors Statement

Guozhong Chen: Conceptualization, Methodology, Investigation, Project
administration, Funding acquisition, Visualization, Writing-original draft,
Writing-review & editing. **Zixin Qin:** Methodology, Investigation, Visualization.
Fenxia Fan: Investigation, Writing-review & editing, Funding acquisition. **Mei Luo:**
Writing-review & editing. **Hongou Wang:** Methodology. **Baoshuai Xue:**
Methodology. **Shucheng Li:** Methodology. **Shiyong Chen:** Supervision. **Xiaoman**
Yang: Investigation. **Xiang Mao:** Investigation. **Liwen Yi:** Investigation. **Chunrong**
Yi: Investigation. **Wei Li:** Investigation, Supervision. **Xiaoyun Liu:** Project
administration, Funding acquisition. **Biao Kan:** Supervision, Project administration.
Zhi Liu: Conceptualization, Methodology, Visualization, Project administration,
Funding acquisition, Supervision, Writing-original draft, Writing-review & editing.

Acknowledgments

We are grateful to all the members of our team. We thank Core Facilities for Life
Science, Huazhong University of Science and Technology for the technical support.

Funding

We also thank Core Facilities for Life and Environmental Sciences, State Key
laboratory of Microbial Technology of Shandong University for technical guidance.
This study was supported by National Natural Science Foundation of China under the
grant numbers 31770132 (to Z. L), (22174003 and 21974002 to X. Y. L) and National
Key Research and Development Program of China (2022YFA1304500 to X. Y. L),
National Key Research and Development Program of China (2021YFC2300304 to F.
X. F), Shandong Province Science Foundation for Youths (ZR2024QC312 to G. Z.
C).

Ethics statement

Only publicly available cholera data from NCBI database were used in this study. Informed consent was obtained from all subjects in the original studies. For those datasets, ethical review and approval can be accessed in the original studies.

All animal experiments were carried out in strict accordance with the animal protocols that were approved by the Ethical Committee of Huazhong University of Science and Technology (Approval no. 2017-S560).

Declaration of Competing Interest

The authors declare no financial/commercial conflicts of interests.

Supplementary Material

Supplementary material is available online.

Data Availability

The proteomic data that support the findings of this study are openly available in the iProX database at https://www.iprox.cn//page/SCV017.html?query=IPX0005998000, accession number IPX0005998000. The raw data of 6N-Label is openly available in the NCBI database at https://www.ncbi.nlm.nih.gov/sra/?term=PRJNA1099378, reference number PRJNA1099378.

References

1. Kanungo S, Azman AS, Ramamurthy T, Deen J, Dutta S. Cholera. Lancet 2022; 399: 1429-1440.
2. World Health Organization [https://www.who.int/]. The organization; Cholera kills more people for second consecutive year, while prevention and treatment available. Available from: https://www.who.int/news/item/12-09-2025-cholera-kills-more-people-for-secon d-consecutive-year-while-prevention-and-treatment-available.
3. D. Chac, C. N. Dunmire, J. Singh, A. A. Weil. Update on environmental and hostfactors impacting the risk of *Vibrio cholerae* infection. ACS Infectious Diseases,2021, 5: 1010-1019.
4. Liu Z, Miyashiro T, Tsou A, Hsiao A, Goulian M, Zhu J. Mucosal penetration primes *Vibrio cholerae* for host colonization by repressing quorum sensing. Proc Natl Acad Sci U S A 2008; 105: 9769-9774.
5. Echazarreta MA, Klose KE. *Vibrio* flagellar synthesis. Front Cell Infect Microbiol 2019; 9: 131.

6.   Syed KA, Beyhan S, Correa N, Queen J, Liu J, Peng F, et al. The *Vibrio cholerae*
flagellar regulatory hierarchy controls expression of virulence factors. J Bacteriol
2009; 191: 6555-6570.

7.   Sowa Y, Berry RM. Bacterial flagellar motor. Q Rev Biophys 2008; 41: 103-132.

8.   Attridge SR, Rowley D. The role of the flagellum in the adherence of *Vibrio*
cholerae. J Infect Dis 1983; 147: 864-872.

9.   Butler SM, Camilli A. Both chemotaxis and net motility greatly influence the
infectivity of *Vibrio cholerae*. Proc Natl Acad Sci U S A 2004; 101: 5018-5023.

10. Wu DC, Zamorano-Sánchez D, Pagliai FA, Park JH, Floyd KA, Lee CK, et al.
Reciprocal c-di-GMP signaling: Incomplete flagellum biogenesis triggers
c-di-GMP signaling pathways that promote biofilm formation. PLoS Genet 2020;
16: e1008703.

11. Katz LS, Petkau A, Beaulaurier J, Tyler S, Antonova ES, Turnsek MA, et al.
Evolutionary dynamics of *Vibrio cholerae* O1 following a single-source
introduction to Haiti. mBio 2013; 4: 1-10.

12. Mostow P, Richardson K. High-frequency spontaneous mutation of classical
Vibrio cholerae to a nonmotile phenotype. Infect Immun 1990; 58: 3633-3639.

13. Jubair M, Atanasova KR, Rahman M, Klose KE, Yasmin M, Yilmaz O, et al.
Vibrio cholerae persisted in microcosm for 700 days inhibits motility but promotes
biofilm formation in nutrient-poor lake water microcosms. PLoS One 2014; 9:
e92883.

14. Giacomucci S, Mathieu-Denoncourt A, Vincent AT, Jannadi H, Duperthuy M.
Experimental evolution of *Vibrio cholerae* identifies hypervesiculation as a way to
increase motility in the presence of polymyxin B. Front Microbiol 2022; 13: 1-18.

15. Hoque MM, Noorian P, Espinoza-Vergara G, Manuneedhi Cholan P, Kim M,
Rahman MH, et al. Adaptation to an amoeba host drives selection of
virulence-associated traits in *Vibrio cholerae*. ISME J 2022; 16: 856-867.

16. Luo M, Chen G, Yi C, Xue B, Yang X, Ma Y, et al. Dps-dependent in vivo
mutation enhances long-term host adaptation in *Vibrio cholerae*. PLoS Pathog
2023; 19: e1011250.

17. Klose KE, Mekalanos JJ. Distinct roles of an alternative sigma factor during both
free-swimming and colonizing phases of the *Vibrio cholerae* pathogenic cycle.
Mol Microbiol 1998; 28: 501-520.

18. Azarian T, Ali A, Johnson JA, Mohr D, Prosperi M, Veras NM, et al.
Phylodynamic analysis of clinical and environmental *Vibrio cholerae* isolates
from Haiti reveals diversification driven by positive selection. mBio 2014; 5:
e01824-14.

19. Daboul J, Weghorst L, DeAngelis C, Plecha SC, Saul-McBeth J, Matson JS.
Characterization of *Vibrio cholerae* isolates from freshwater sources in northwest
Ohio. PLoS One 2020; 15: e0238438.

20. Mavian C, Paisie TK, Alam MT, Browne C, de Rochars VMB, Nembrini S, et al.
Toxigenic *Vibrio cholerae* evolution and establishment of reservoirs in aquatic
ecosystems. Proc Natl Acad Sci U S A 2020; 117: 7897-7904.
21. Mwaba J, Debes AK, Murt KN, Shea P, Simuyandi M, Laban N, et al. Three
transmission events of *Vibrio cholerae* O1 into Lusaka, Zambia. BMC Infect Dis
2021; 21: 570.
22. Chen, Shifu. Ultrafast one-pass FASTQ data preprocessing, quality control, and
deduplication using fastp. iMeta 2023; 2: e107.
23. Uritskiy, Gherman V., DiRuggiero Jocelyne, Taylor James. MetaWRAP-a flexible
pipeline for genome-resolved metagenomic data analysis. Microbiome 2018; 6:
158.
24. Chaumeil, Pierre-Alain, Mussig Aaron J, Hugenholtz Philip, Parks Donovan H.
GTDB-Tk v2: memory friendly classification with the genome taxonomy
database. Bioinformatics 2022; 38: 5315-5316.
25. Hyatt, Doug, Chen Gwo-liang, LoCascio Philip F, Land Miriam L, Larimer Frank
W, Hauser Loren J. Prodigal: Prokaryotic gene recognition and translation
initiation site identification. BMC Bioinformatics 2010; 11:119.

26. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. Nature Methods 2015;12: 59-60.
27. Farhadi T, Nezafat N, Ghasemi Y. In silico phylogenetic analysis of *Vibrio*
cholerae isolates based on three housekeeping genes. Int J Comput Biol Drug
Des. 2015; 8:62-74.
28. Rui YY, Kan B, Gao SY, Liu YQ, Qi GM. Sequence analysis of housekeeping
genes *recA*, *dnaE*, and *mdh* in different serogroups or biotypes of *Vibrio cholerae*
isolated from China. Nan Fang Yi Ke Da Xue Xue Bao. 2006; 26:1720-1723.
29. Houot L, Watnick PI. A novel role for enzyme I of the *Vibrio cholerae*
phosphoenolpyruvate phosphotransferase system in regulation of growth in a
biofilm. J Bacteriol 2008; 190: 311-320.
30. Metcalf WW, Jiang W, Daniels LL, Kim SK, Haldimann A, Wanner BL.
Conditionally replicative and conjugative plasmids carrying lacZα for cloning,
mutagenesis, and allele replacement in bacteria. Plasmid 1996; 35: 1-13.
31. Xia A, Qian M, Wang C, Huang Y, Liu Z, Ni L, et al. Optogenetic modification of
Pseudomonas aeruginosa enables controllable twitching motility and host
infection. ACS Synth Biol 2021; 10: 531-541.
32. Ma Y, Yang X, Wang H, Qin Z, Yi C, Shi C, et al. CBS-derived H_2S facilitates
host colonization of *Vibrio cholerae* by promoting the iron-dependent catalase
activity of KatB. PLoS Pathog 2021; 17: e1009763.
33. Liu Z, Wang H, Zhou Z, Sheng Y, Naseer N, Kan B, et al. Thiol-based switch
mechanism of virulence regulator AphB modulates oxidative stress response in
Vibrio cholerae. Mol Microbiol 2016; 102: 939-949.

34. Hong S, Zheng DW, Zhang QL, Deng WW, Song WF, Cheng SX, et al. An
RGB-emitting molecular cocktail for the detection of bacterial fingerprints. Chem
Sci 2020; 11: 4403-4409.

35. Lin L, Song J, Du Y, Wu Q, Gao J, Song Y, et al. Quantification of bacterial
metabolic activities in the gut by d-Amino acid-based in vivo labeling. Angew
Chemie - Int Ed 2020; 59: 11923-11926.

36. Dalia AB, McDonough EK, Camilli A. Multiplex genome editing by natural
transformation. Proc Natl Acad Sci U S A 2014; 111: 8937-8942.

37. Hoque MM, Noorian P, Espinoza-Vergara G, Manuneedhi Cholan P, Kim M,
Rahman MH, et al. Adaptation to an amoeba host drives selection of
virulence-associated traits in *Vibrio cholerae*. ISME J 2022; 16: 856-867.

38. Martinez RM, Dharmasena MN, Kirn TJ, Taylor RK. Characterization of two
outer membrane proteins, FlgO and FlgP, that influence *Vibrio cholerae* motility. J
Bacteriol 2009; 191: 5669-5679.

39. Olivier V, Salzman NH, Satchell KJF. Prolonged Colonization of Mice by *Vibrio
cholerae* El Tor O1 Depends on Accessory Toxins. Infect Immun 2007; 75:
5043-5051.

40. Correa NE, Barker JR, Klose KE. The *Vibrio cholerae* FlgM homologue is an
anti-σ^{28} factor that is secreted through the sheathed polar flagellum. J Bacteriol
2004; 186: 4613-4619.

41. Qin Z, Yang X, Chen G, Park C, Liu Z. Crosstalks between gut microbiota and
Vibrio cholerae. Front Cell Infect Microbiol 2020; 10: 582554.

42. Tsou AM, Cai T, Liu Z, Zhu J, Kulkarni R V. Regulatory targets of quorum
sensing in *Vibrio cholerae*: Evidence for two distinct HapR-binding motifs.
Nucleic Acids Res 2009; 37: 2747-2756.

43. Tsou AM, Liu Z, Cai T, Zhu J. The VarS/VarA two-component system modulates
the activity of the *Vibrio cholerae* quorum-sensing transcriptional regulator HapR.
Microbiology 2011; 157: 1620-1628.

44. Herzog R, Peschek N, Fröhlich KS, Schumacher K, Papenfort K. Three
autoinducer molecules act in concert to control virulence gene expression in *Vibrio
cholerae*. Nucleic Acids Res 2019; 47: 3171-3183.

45. Liu Z, Hsiao A, Joelsson A, Zhu J. The transcriptional regulator VqmA increases
expression of the quorum-sensing activator HapR in *Vibrio cholerae*. J Bacteriol
2006; 188: 2446-2453.

46. Vanhove AS, Jugder BE, Barraza D, Watnick PI. Methionine availability in the
arthropod intestine is elucidated through identification of *Vibrio cholerae*
methionine acquisition systems. Appl Environ Microbiol 2020; 86: e00371-20.

47. Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
Res 2000; 28: 27-30.

48. Zhang Z, Chen G, Hu J, Hussain W, Fan F, Yang Y, et al. Mr.Vc: A database of
microarray and RNA-seq of *Vibrio cholerae*. Database 2019; 2019: 1-8.

49. Zhang Z, Chen G, Hussain W, Qin Z, Liu J, Su Y, et al. Mr.Vc v2: An updated
version of database with increased data of transcriptome and experimental
validated interactions. Front Microbiol 2022; 13: 1047259.

50. Calabretta PJ, Hodges HL, Kraft MB, Marando VM, Kiessling LL. Bacterial cell
wall modification with a glycolipid substrate. J Am Chem Soc 2019; 141:
9262-9272.

51. Xu Y, Hernández-Rocamora VM, Lorent JH, Cox R, Wang X, Bao X, et al.
Metabolic labeling of the bacterial peptidoglycan by functionalized glucosamine.
iScience 2022; 25: 104753.

52. Sadamoto R, Matsubayashi T, Shimizu M, Ueda T, Koshida S, Koda T, et al.
Bacterial surface engineering utilizing glucosamine phosphate derivatives as cell
wall precursor surrogates. Chem - A Eur J 2008; 14: 10192-10195.

53. Liu, R. et al. Diet modulates *Vibrio cholerae* colonization and competitive
outcomes with the gut microbiota. Cell Host and Microbe 2025; 33, 2085-2099.e5.

54. Jiang S, Zhang C, Han Z, Ma W, Wang S, Huo D, Cui W, Zhai Q, Huang S, Zhang
J. Native microbiome dominates over host factors in shaping the probiotic genetic
evolution in the gut. NPJ Biofilms Microbiomes. 2023; 1:80.

55. Martínez Y, Li X, Liu G, Bin P, Yan W, Más D, et al. The role of methionine on
metabolism, oxidative stress, and diseases. Amino Acids 2017; 49: 2091-2098.

56. Zhao T, Lum JJ. Methionine cycle-dependent regulation of T cells in cancer
immunity. *Front Oncol* 2022; 12: 1-9.

57. Martínez Y, Li X, Liu G, Bin P, Yan W, Más D, et al. The role of methionine on
metabolism, oxidative stress, and diseases. *Amino Acids* 2017; 49: 2091-2098.

58. Hsiao A, Liu Z, Joelsson A, Zhu J. *Vibrio cholerae* virulence
regulator-coordinated evasion of host immunity. Proc Natl Acad Sci U S A.
2006;103 :14542-14547

59. Bogard RW, Davies BW, Mekalanos JJ. MetR-regulated *Vibrio cholerae*
metabolism is required for virulence. mBio 2012; 3.e00236-12.

60. Hondorp ER, Matthews RG. Methionine. *EcoSal Plus* 2006; 2: 1-36.

61. Winans SC. A new family of quorum sensing pheromones synthesized using
S☐adenosylmethionine and Acyl☐CoAs. Mol Microbiol 2011; 79: 1403-1406.

62. Heurlier, K., Dénervaud, V. & Haas, D. Impact of quorum sensing on fitness of
Pseudomonas aeruginosa. International Journal of Medical Microbiology 2006,
296, 93-102.

63. Cho JY, Liu R, Macbeth JC, Hsiao A. The interface of *Vibrio cholerae* and the
gut microbiome. Gut Microbes. 2021, 13:1937015.

64. 2. Hsiao A, Zhu J. Pathogenicity and virulence regulation of *Vibrio cholerae* at
the interface of host-gut microbiome interactions. Virulence. 2020;
11:1582-1599.

65. 3. Qin Z, Yang X, Chen G, Park C, Liu Z. Crosstalks between gut microbiota and
Vibrio cholerae. Front Cell Infect Microbiol 2020; 10: 582554.

66. Silva AJ, Leitch GJ, Camilli A, Benitez JA. Contribution of
hemagglutinin/protease and motility to the pathogenesis of El Tor biotype cholera.
Infect Immun. 2006;74:2072-9.

67. Waters CM, Lu W, Rabinowitz JD, Bassler BL. Quorum sensing controls biofilm
formation in *Vibrio cholerae* through modulation of cyclic di-GMP levels and
repression of *vpsT*. J Bacteriol. 2008; 190:2527-36.

68. Srivastava D, Harris RC, Waters CM. Integration of cyclic di-GMP and quorum
sensing in the control of *vpsT* and *aphA* in *Vibrio cholerae*. J Bacteriol. 2011
Nov;193(22):6331-41.

69. Johnson GE, Fei C, Wingreen NS, Bassler BL. Analysis of gene expression
within individual cells reveals spatiotemporal patterns underlying *Vibrio cholerae*
biofilm development. PLoS Biol. 2025; 23: e3003187.

70. Joelsson A, Kan B, Zhu J. Quorum sensing enhances the stress response in *Vibrio
cholerae*. Appl Environ Microbiol. 2007; 73:3742-6.

71. Khan F, Tabassum N, Anand R, Kim YM. Motility of *Vibrio* spp.: regulation and
controlling strategies. Appl Microbiol Biotechnol 2020; 104: 8187-8208.

72. Okumura R, Kurakawa T, Nakano T, Kayama H, Kinoshita M, Motooka D, et al.
Lypd8 promotes the segregation of flagellated microbiota and colonic epithelia.
Nature 2016; 532: 117-121.

73. Grünenfelder B, Tawfilis S, Gehrig S, Østerås M, Eglin D, Jenal U. Identification
of the protease and the turnover signal responsible for cell cycle-dependent
degradation of the Caulobacter FliF motor protein. *J Bacteriol* 2004; 186:
4960-4971.

74. Ruby EG, Asato LM. Growth and flagellation of *Vibrio fischeri* during initiation of
the sepiolid squid light organ symbiosis. Arch Microbiol 1993; 159: 160-167.

75. Doerges L, Kutschera U. Assembly and loss of the polar flagellum in
plant-associated methylobacteria. Naturwissenschaften 2014; 101: 339-346.

76. Zhu S, Gao B. Bacterial flagella loss under starvation. Trends Microbiol 2020; 28:
785-788.

Figures

Fig. 1. High-frequency mutations in the flagellar genes of *V. cholerae* from the life cycle. (**A**), World distribution map of the collected samples, with areas in pink indicating no sample collection. (**B**) Spectrum and count statistics of flagellar gene mutations. The *hapR* and housekeeping genes (*asd*, *dnaE*, *mdh*) were used as controls. (**C**) Schematic diagram of the *V. cholerae* flagellum assembly. (**D**) *V. cholerae* flagellar transcription hierarchy.

Fig. 2. FlgM secretion regulates colonization of *V. cholerae* flagellar mutants. (**A**), Motility assay. (**B**), Competition assay in adult mice. The horizontal bar indicates the average of CI values. The CI values at 5 days post inoculation were shown. The grey dashed lines from top to bottom represent 10-fold enhanced colonization and 10-fold reduced colonization, respectively. (**C**), Volcano plot of the distribution of altered proteins. Red dots represent up-regulated expression proteins, and green dots represent down-regulated expression proteins and grey dots represent no significant expression difference (threshold > 1.5, *p* < 0.05). (**D** and **E**), Effect of FlgM on the colonization of the Δ*flhB* or Δ*flaA* mutant. The CI values at 5 days post inoculation were shown. (**F**), Western blot (WB) assay for different secretion of FlgM in the *V. cholerae* flagellar mutants. sup: supernatant, p: pellets. (**G**), Correlation analysis of FlgM secretion and CI values in adult mice model (ImageJ and Pearson method). The colors of red, green dots represent a higher, lower of colonization than WT, respectively. Significance was determined by *t*-test; *P*-values: ns, not significant, *, < 0.05, **, < 0.01, ***, < 0.001.

Fig. 3. Intracellular FlgM deactivated FliA to promote *hapR* expression on
transcriptionally and post-translationally. (**A** and **B**), Schematic diagram of FlgM
secretion in Δ*flhB* and Δ*flaA*, and the corresponding role of FlgM-FliA-*hapR* pathway
in colonization, respectively. (**C**), RT-PCR results of *hapR* expression in flagellar
mutants with enhanced colonization. The *recA* was used for normalization and Δ*luxO*
was used as a positive control. (**D**), *In vivo* competition experiments of different *hapR*
mutants in adult mice. The CI values at 5 days post inoculation were shown. (**E**),
Effect of FlgM and FliA overexpression on *hapR* expression. (**F**), Quorum sensing
(QS) regulatory network of *V. cholerae*. (**G**), Gene expression in the QS regulatory
pathway. Gene expression was normalized by *16S rRNA*, except for small RNAs
called *csrBCD* which were normalized by *secA*. (**H**), Role of the
VarS/VarA-CsrA/BCD system in adult mice colonization. The CI values at 5 days
post inoculation were shown. The colors of red, green, blue dots represent a higher,
lower of colonization than WT or same level, respectively. Significance was

determined by *t*-test and one-way ANOVA analysis; *P*-values: *, < 0.05, **, < 0.01,
, < 0.001, *, < 0.0001.

Fig. 4. HapR regulates *V. cholerae* colonization through methionine metabolism
and transportation. (**A**), Volcano map of the precipitated proteomic of Δ*flhB*/Δ*flaA*.
Red dots represent up-regulated expression proteins, and green dots represent
down-regulated expression proteins and grey dots represent no significant expression
difference (threshold > 1.5, *p* < 0.05). (**B**), Schematic diagram of methionine (Met)
metabolism. (**C**), Effect of Met addition (in drinking water) *in vivo* on the
colonization of Δ*flhB* and Δ*flaA*. (**D**), Effect of *msrC* in precipitated proteomic on the
colonization of Δ*flhB* mutant. (**E**), Effect of indicated mutants on the colonization of
Δ*flhB*,   Δ*flhB*Δ*msrC*,   Δ*flhB*Δ*metR*,   Δ*flhB*Δ*metR*Δ*msrC*,
Δ*flhB*Δ*metR*Δ*msrC*Δ*metI*Δ*metT*. (**F**), Amino acids and its derivatives were quantified
by UPHLC. The column diagram represents the contents of amino acid and its
derivatives in Δ*flhB* and Δ*flaA*. The line chart represents the ratio of amino acid and
its derivative content of Δ*flhB*/Δ*flaA*. (**G**), Metabolic pathway of methionine
converted to other amino acids. Methionine can be converted to homocysteine by
demethylation, which subsequently forms cysteine together with serine. Two
molecules of cysteine synthesize cystine, while serine itself can be converted to
cysteine. The metabolic pathway was annotated by Kyoto Encyclopedia of Genes and
Genomes. (**H**), Promoter sequence alignment of *metR*, *msrC*, *metI*, and *metT* with
conserved two HapR binding Motifs. Yellow shading represents the conserved
binding sequence of Motif 1 and aqua shading represents the conserved binding
sequence of Motif 2. (**I**), Effect of *hapR* overexpression on the expression of *metR*,
msrC, *metI*, and *metT*. The bars represent the corresponding standard deviations. (**J**),
EMSA for HapR binding to the promoters of *metR*, *msrC*, *metI*, and *metT*. (**K**),
Growth curve of Δ*flhB* and Δ*flaA* in the MM medium with or without 0.01mM L-Met.
(**L**), Growth curve of WT, Δ*flhB*Δ*metR*Δ*msrC*Δ*metI*Δ*metT* in the MM medium with
or without 0.01mM L-Met. The colors of red, green, blue dots represent a higher,

lower of colonization than WT or same level, respectively. The CI values at 5 days
post inoculation were shown. The horizontal bars indicate the averages of CI.
Significance was determined by *t*-test and one-way ANOVA methods; *P*-values: ns,
not significant, *, < 0.05, **, < 0.01, ***, < 0.001.

1001

Fig. 5. Methionine-related metabolic activity affects *V. cholerae* colonization. (**A**),
The workflow for metabolic activity assay with FITC-D-Glu-labeled method in vitro
and in vivo. The figure was created by BioRender. (**B**), Flow cytometry analysis of
wild-type co-with Δ*flhB* and wild-type co-with Δ*flaA* FITC-D-Glu-labeled in
transwell chamber at 30 min. (**C**), The bacterial metabolic activity of wild-type
co-with Δ*flhB* and wild-type co-with Δ*flaA in vivo* at 48h post-inoculation. (**D**),
workflow of colonization assay of 6N-labeled mutants *in vivo*. (**E**), Colonization
assay of 6N-labeled mutants *in vivo*. The raw data was analyzed by Vsearch and
Python. The adjusted CIs of all flagellar mutants were analyzed by was calculated as
the reads of mutant to WT colonies normalized with the input reads. The colors of red,
green, blue dots represent a higher, lower of colonization than WT or same level,
respectively. Significance was determined by *t*-test and one-way ANOVA methods;
P-values: ns, not significant, *, < 0.05, **, < 0.01, ***, < 0.001.

1015

Fig. 6. Anti-σ²⁸ factor FlgM secretion regulates *Vibrio cholerae* adaptability in adult mice through quorum sensing and methionine metabolism. *Vibrio cholerae* predominantly cycles between aquatic reservoirs and the human host. Humans become infected by ingesting water or food contaminated with *V. cholerae*. The bacteria then colonize the small intestine, where they adapt by modulating their gene expression profile. Ultimately, they are shed back into the aquatic environment via feces, completing the transmission cycle and enabling potential new rounds of infection. During the life cycle of *Vibrio cholerae*, flagellar gene mutations are highly prevalent, clustering in genes including *flhB*, *fliA*, *fliF*, *fliD*, and *fliM*. Follow-up experimental investigations elucidated a fundamental trade-off mechanism of *Vibrio cholerae* fitness in the host gut by FlgM secretion. The level of FlgM secretion is primarily regulated by the rod complex and the flagellar type III secretion system (fT3SS). Specifically, elevated intracellular concentrations of FlgM inhibit the alternative sigma factor σ²⁸ (FliA). FliA itself via the VarS/VarA-CsrA/BCD system affects the expression of quorum sensing master regulator called *hapR*. Moreover, HapR can directly bind to the promoters of genes involving in Met transportation to regulate methionine availability and metabolic activity by labeled FITC-D-Glucose to regulates *V. cholerae* colonization in the gut.

Table legends

Table S1. Sample collection. Samples were collected from 19 countries. Based on their sources, the cholera-related metagenomic sequencing samples obtained from the SRA database were classified into five categories: human-derived, clinical, aquatic (including water and sewage), fish-derived, and amoeba-derived.

Table S2. Structure and function of *Vibrio cholerae* flagella. The flagella of *V. cholerae* are divided into many flagellar components which play a different role in flagellar biosynthesis. Their functions were summarized by literature search, and literature sources were added.

1044

1045 **Table S3. In silico analysis of flagellar gene mutations in *V. cholerae*.** Mutations in
1046 flagellar genes were analyzed using the Diamond software.

1047

1048 **Table S4. Strains and public Plasmids used in this study.** The strains or public
1049 plasmids were descripted. The public plasmids were obtained from anywhere.

1050

1051 **Table S5. The primers used in this study.** These primers involve plasmid
1052 construction, Reverse Transcription Polymerase Chain Reaction, Electrophoretic
1053 Mobility Shift Assay, *lacZ* sequencing.

1054

1055 **Table S6. Altered proteins in secretory proteomics data.** The up-regulated and
1056 down-regulated expression proteins of Δ*flhB*/Δ*flaA* were list (Fold change > 1.5, *p* <
1057 0.05, *t-test*). The involved functions of these altered proteins were explained.

1058

1059 **Table S7. Differentially expressed proteins in precipitated proteomics data.**
1060 Bacterial proteome samples of Δ*flhB* or Δ*flaA* were analyzed in triplicates. Proteins
1061 with Fold changes >1.5 and *p*< 0.05 (*t-test*) were considered as significant differences.

1062

1063 **Table S8. Relative abundance analysis of 6N-labelled LacZ sequence of *V.***
1064 ***cholerae* wild-type and flagellar regulon mutants' libraries.** The sequenced raw
1065 data was analyzed by Vsearch for merging paired-end reads and Python to calculate
1066 the proportion of reads from the 6N-labeled mutants among the total reads.