Repurposed small molecule toxin inhibitors neutralise a diversity of venoms

from the Neotropical viperid snake genus *Bothrops*

Authors: Rachel H. Clare[1,2*], Adam Westhorpe[2], Emma Stars[2], T.D. Kazandjian[2], Laura-

Oana Albulescu[2], Stefanie K. Menzies[3,2] and Nicholas R. Casewell[2]

[1]Biology Department, Edge Hill University, St Helens Road, Ormskirk, L39 4QP, UK; [2]Centre

for Snakebite Research and Interventions, Liverpool School of Tropical Medicine, Liverpool,

L3 5QA, UK; [3]Biomedical & Life Sciences, Lancaster University, Bailrigg, Lancaster, UK, LA1

4YW

*For correspondence: rachel.clare@edgehill.ac.uk

Abstract

Abstract

Snakebite globally claims more than 100,000 lives per year and results in morbidity for

400,000 survivors. Current treatment uses antibody-based antivenoms which are

constrained by their efficacy, safety and cost. In this study we evaluated the efficacy of

previously described repurposed drugs against viperid snakes of the medically important

Bothrops genus. Despite variable toxin representation and bioactivity across this central and

south American genus, we found that the lead inhibitors targeting metalloproteinases

(marimastat and DMPS) and phospholipases (varespladib), demonstrated pan-species

neutralisation in enzymatic assays, whilst nafamostat (serine protease inhibitor) had variable

activity. The metalloproteinase inhibitors protected against the procoagulant and

haemorrhagic effects of several venoms in phenotypic assays. Collectively these findings

demonstrate that repurposed drugs may be of great value as early interventions for the

treatment of bothropic envenoming in the Neotropics and thus provides a strong rationale for

their progression into future preclinical and clinical evaluation for snakebite indication.

Introduction

Introduction

Snakebite envenoming is an acute, potentially lethal event that affects several million people

each year, primarily those working and living in rural areas of the world's tropical and sub-

tropical regions. Snakebite is classified by the World Health Organization as a priority

neglected tropical disease, and is responsible for over 100,000 deaths annually, while

several times that number of survivors suffer from long-term disabling or debilitating health

conditions caused by venomous snakebites [1]. Current treatment for snakebite consists of

intravenously delivered antivenoms, which are serotherapies consisting of polyclonal

antibodies generated from venom-immunised animals (e.g. equines, ovines, camelids) [2,3].

Multiple venoms are often used in the immunising mixture, resulting in a broad range of

antibodies that target the diversity of toxins present. Nonetheless, antivenom therapies are

generally constrained in their efficacy to particular geographical regions or snake species

due to venom toxin variation, which is highly variable across medically important venomous

snakes [4].

The cocktail of toxins produced in the venom gland varies not only across snake species, but

can also vary between populations and individuals of the same species, and has been

shown to be affected by the age, sex and ecological factors experienced by the producing

animal [5-8]. The application of 'venomics' approaches over the past two decades has

enhanced the identification and characterisation of snake venom toxin diversity at the

proteomic level [9]. This advancement, alongside functional characterisation, has supported

the consensus that certain venom toxin families are often the dominant drivers of pathology

following snakebite, due to both their high abundance and toxicity; namely, the snake venom

metalloproteinases (SVMP), phospholipases A2 (PLA_2), snake venom serine proteases

(SVSP) and three finger toxins (3FTx) [10]. This knowledge has paved the way for the

discovery and development of alternative therapeutic approaches to antivenom, focusing on

the development of rational data-informed therapeutic molecules which inhibit specific

venom toxin families or sub-families, unlike conventional antivenom. Recent advances [11]

include the identification of toxin-specific monoclonal antibodies [12] or nanobodies [13],

computationally designed binding proteins [13,14] and synthetic inhibitors [2,10,15]. Repurposed

drugs (synthetic inhibitors), defined by the new therapeutic use for existing molecules

previously developed for other indications, are showing considerable promise for snakebite

envenoming. This is partly because three of the four priority venom toxin families to

neutralise are enzymes, lending themselves to potential generic toxin family inhibition (i.e.

across diverse isoforms) by small molecule drugs via binding to the active site. Several such

drugs have been shown to rescue the lethal and pathological effects of snake venoms in

small animal models, including SVMP-induced coagulopathy and haemorrhage, PLA_2-

mediated neurotoxicity, and localised tissue damage via cytotoxicity (caused by PLA_2, SVMP

and/or cytotoxic 3FTx) [15-19].

Several of these repurposed drugs have now entered or are entering clinical trials as orally

bioavailable therapeutics amenable for clinical evaluation as front-line snakebite treatments.

For example, the PLA_2 inhibitor varespladib recently completed a Phase II trial for snakebite

in the USA and India [20,21], with a second Phase II study ongoing at the time of writing.

Similarly, the SVMP inhibiting metal chelator DMPS has undergone Phase I studies [22] to

dose optimise for snakebite indication, and is planned to enter Phase II evaluations in 2026,

potentially alongside another SVMP inhibitor, marimastat, which has a distinct mode of

action. Potential key benefits of such small molecules over traditional polyclonal antibody-

based antivenoms include clearly established and acceptable safety profiles, broad-

spectrum toxin inhibition, low-cost manufacture, improved tissue penetration and oral

bioavailability [15]. The latter provides a potential paradigm shift for treatment, from: i) the

requirement for intravenous delivery (in tertiary healthcare settings) of antivenom which

leads to treatment delays, ii) the reliance on cold chain storage and iii) the risk of serious

adverse events from animal-derived antibodies, resulting in a vision of community based,

rapid delivery of low cost, well tolerated, temperature stable, oral medications. Substantially

reducing the time frame between bite and treatment, which in many settings such as the

remote areas of the Amazon has been reported to be more than five hours, holds much

promise for improving patient outcomes for this time sensitive, acute, life-threatening

condition [23,24].

In this study we focused on assessing the potential utility of small molecule drugs in the

context of the Neotropics. Unlike many other locations at risk of snakebite where a range of

genera are indicated in the majority of severe envenoming cases, medically important

snakebites in Central and South America are often dominated by snakes of the genus

Bothrops [25-28]. This group of diverse pit vipers (family Viperidae) includes species that range

geographically from Mexico to Argentina, as well as certain Caribbean islands (Figure 1A).

Conservative estimates of snakebite incidence in Latin America were previously reported as

50.37/100,000 people per year (80,329 snakebites per year), resulting in 540 deaths per

year (0.5781/100,000 per year) [29]. A more recent publication focusing on South America

reported the continent as experiencing the third highest incidence of snakebite after Asia and

Africa at 21.7/100,000 population per year, resulting in the fourth highest mortality rate at

0.03/100,000 population per year, after Asia, Africa and North America [30]. The fact that that

Costa Rica and Nicaragua are classified within the North American dataset further

exacerbates the snakebite incidence in the wider biogeographical realm of the Neotropics.

There are several commercial antivenom options to treat bothropic envenoming in the

region, including the polyvalent antibotrópico antivenom from Instituto Butantan (Brazil) and

PoliVal-ICP from Instituto Clodomiro Picado (Costa Rica). However, each antivenom is

designed to cover only certain species, and delays between bite and accessing antivenom

treatment remain, with a clinical epidemiological study of *Bothrops* bites in Brazil [24]

estimating an average of three hours between accident and initial medical care for 65% of

cases, but increasing to up to six hours for most cases in northern and northeastern regions

of the country [24]. Indeed, antivenom distribution centres are disproportionately spatially

distributed compared to snakebite incidents, with the northern regions that experience higher

incidences of *Bothrops* envenoming having fewer centres [24]. These findings were mirrored in

a systematic review of antivenom use in the Americas, which identified an average of 5.7

hours between bite and antivenom administration, with the treatment time ranging from 1.5

to 19 hours [28].

The typical clinical presentation of 'bothropic syndrome' following *Bothrops* envenoming

includes local tissue damage (bruising, blistering, dermonecrosis, myonecrosis, and

oedema), pain, incoagulable state, haemorrhage, circulatory shock and acute kidney injury.

The latter three cause the most serious systemic and life-threatening effects [24-26,31-33]. The

reported pathologies linked to *Bothrops* envenoming are attributed to the dominant venom

protein families, namely the PLA_2, SVMP and SVSP enzymes (Figure 1B). To investigate the

potential utility and therapeutic value of small molecule drugs for bothropic envenoming, in

this study we tested the ability of previously described repurposed drugs to neutralise a wide

range of *Bothrops* venoms in a panel of toxin-specific enzymatic assays, followed by

phenotypic assays measuring venom-induced coagulopathy and general haemotoxicity. Our

findings demonstrate that the lead repurposed snakebite drugs currently under study

(varespladib, marimastat and DMPS) provide broad, albeit often variable, inhibitory potency

against the diverse enzymatic activities of *Bothrops* venoms. Furthermore, the SVMP

inhibitors show much promise for inhibiting the coagulopathic and haemotoxic effects caused

by these venoms, providing a strong rationale for their future preclinical and clinical

evaluation.

Results

***In vitro* profiling of *Bothrops* venom activity**

Variation in venom composition

Figure 1B summarises the published reports of the venom toxin compositions of the species

tested in this manuscript. These prior studies demonstrate clear interspecies variation in

venom toxin composition, but also intraspecies variation for several cases (Supplemental

Table 1). In this manuscript we did not seek to formally characterise venom composition, but

instead used SDS-PAGE to provide an overview of the venom protein compositions of the

species used in this study (Figure 2A). In line with the literature, notable differences were

observed between the species in terms of the proportion and abundance of proteins of all

molecular weights. The seven venoms could be broadly assigned into two groups. The first

group, containing *B. alternatus*, *B. jararaca* and *B. lanceolatus* venoms, exhibited abundant

protein bands at ~55-60 kDa (likely PIII SVMPs) alongside less abundant lower molecular

mass proteins at ~20-35 kDa (likely SVSP, cysteine-rich secretory proteins and/or PI SVMP)

and ~12-16 kDa (likely PLA$_2$ and C-type lectin-like proteins) [34]. The sample of *B. jararaca*

used in this study had a dominant protein band ~55-60 kDa, which if assumed to be PIII

SVMPs, was in line with previous proteomic studies (Figure 1B, Supplementary Table 1) [35].

The second group, containing *B. moojeni*, *B. neuwiedi*, *B. asper* and *B. atrox* venoms, had a

much higher abundance of lower molecular weight toxins (<25 kDa), including prominent

bands at ~12-16 kDa, and lower abundance of higher molecular weight toxins.

Comparisons of SVMP activity

Quantification of venom SVMP activity used a consistent venom and substrate concentration

and was measured kinetically. The area under the curve (AUC) of each kinetic reaction was

calculated, and this value was used to compare SVMP activity between the venoms (Figure

2B). All venoms demonstrated some degree of SVMP activity; *B. jararaca* and *B. lanceolatus*

displayed the most activity (AUC = 4.24×10^7 and 4.18×10^7 respectively) followed by *B.*

alternatus (3.82×10^7) and *B. atrox* (3.55×10^7), then a moderate drop in SVMP activity of

1.4 to 2.3-fold was observed with the final three venoms (*B. asper*, *B. neuwiedi* and *B.*

moojeni; 2.62×10^7, 2.31×10^7 and 1.86×10^7, respectively). Only the four most active

venoms had converted all the substrate present in the reaction by the end of the experiment,

causing a plateau in fluorescent signal. A general trend was seen between the total intensity

of the higher molecular weight SDS-PAGE bands that corresponded with PIII SVMP proteins

(i.e. ~55-60 kDa in Figure 2A) and SVMP assay activity, with the exception of *B. atrox*,

however the potential contribution of other SVMP subclasses to this activity remains unclear.

Comparison of PLA$_2$ activity

Venom activity in the PLA$_2$ assay also showed considerable variation. As with the SVMP

assay, the AUC of the curve generated from the kinetic read of the assay was used to

compare PLA$_2$ specific substrate conversion rate, and therefore PLA$_2$ activity, of the venoms

tested at a matched concentration of 100 ng per reaction (Figure 2C). *B. lanceolatus* and *B.*

atrox venom demonstrated very similar and potent PLA$_2$ activities (AUC = 1832.0 and

1703.0, respectively), with activity 1.2 to 1.5-fold higher than the activity seen to the next

grouping of venoms, which all displayed very similar activities; *B. asper* (1449.8), *B.*

neuwiedi (1408.5), *B. moojeni* (1269.5), and *B. jararaca* (1214.5). *B. alternatus* displayed no

PLA$_2$ activity at this dose (105.2 compared to the 99.56 assay background) and was

distinctly lower than any of the other tested venoms, but this correlates with the previously

described venom composition of *B. alternatus* (2% PLA$_2$; Supplementary Table 1) [35].

Correlations with the SDS-PAGE in the absence of proteomic data are challenging, as all

venoms showed protein bands that corresponded with the predicted molecular weight of
PLA$_2$ toxins (~12-14 kDa; though these also overlap with other venom toxins), including *B.*
alternatus, though the strongest bands were observed in *B. atrox* and *B. asper* venom which
were the second and third most active in this enzymatic activity assay. While *B. lanceolatus*
showed bands of weaker intensity at this molecular weight, its venom profile showed three
distinct bands in this molecular region, which may cumulatively account for the highest
functional activity seen across all of the venoms tested.
191
Comparison of SVSP activity
Quantification of the kinetic profiles of SVSP activity revealed more extensive variation
compared with the findings from the PLA$_2$ and SVMP activity assays. The venoms of *B.*
atrox and *B. alternatus* presented with low levels of enzymatic SVSP activity (mean AUCs:
2213 and 2504), compared to the moderate activity of *B. asper*, *B. neuwiedi* and *B. jararaca*
(mean AUCs: 3916, 4073 and 5214) (Figure 2D). The highest venom activity was observed
for *B. moojeni* and *B. lanceolatus* (mean AUCs: 6552 and 7143)*,* which exhibited greater
than 2.5-fold higher activity than *B. alternatus and B. atrox* venoms and between 1.2 and
1.8-fold than that of the moderately active venoms of *B. asper*, *B. neuwiedi* and *B. jararaca*.
201
Comparison of coagulation profiles
The plasma coagulation assay was conducted in a similar manner to the previous assays, in
which a fixed venom dose (100 ng per well) was used to compare the coagulopathic
activities of each venom. This assay utilises a biological product, citrated bovine plasma, and
as such is more representative of the phenotypic effects of the venom as a whole, in
comparison to the SVMP, SVSP, and PLA$_2$ assays which are specific to one toxin family. All
venoms demonstrated an overall procoagulant activity, as indicated by each of the curves
initiating clotting earlier (within 4 minutes) and plateauing earlier (by 17 minutes) than the no-
venom clotting control (initiated after 7 minutes and plateauing at ~21 minutes) (Figure 3A).
As with the other assays, there was variance between each venom with regards to the
potency of the observed procoagulant effect; *B. moojeni*, *B. neuwiedi* and *B. atrox* were the
most procoagulant and had caused complete clotting of the plasma by the second or third
read timepoint (< 7 minutes, latter two venoms overlapped in kinetic reads). The venoms of
B. asper, *B. jararaca* and *B. lanceolatus* exhibited comparable procoagulant profiles,
demonstrating complete clotting by the fourth timepoint (~ 10 minutes), while *B. alternatus*
venom demonstrated the least potent procoagulant profile, with clotting plateauing around
timepoint 6 (~ 17 minutes), which was only marginally earlier than the no-venom control.
219
220   ***In vitro* drug inhibition**

Inhibition of SVMP activity by MMPi and metal chelators

Following on from the SVMP assay in which variable activities were demonstrated across all venoms, a panel of 9 known matrix metalloproteinase (MMP) inhibitors were screened as dose response curves against a fixed dose of each venom with subsequent EC_{50} calculations to allow for potency comparisons (Figure 4A). Marimastat, a matrix metalloproteinase inhibitor (MMPi) that has demonstrated single digit nanomolar EC_{50}s against a broad range of different snake venoms [16,18,36,37], was included as a gold standard. The top dose of marimastat, 10 μM, inhibited the SVMP activity of all venoms down to baseline (Figure 4B) and demonstrated low nanomolar inhibition in the SVMP assay against most of the venoms tested (EC_{50} range 1.8 – 10.6 nM). DMPS, a heavy metal chelator that has recently completed a Phase I safety trial for snakebite indication [22], demonstrated sub-micromolar EC_{50}s against all but two of the venoms (*B. asper* = 2019.0 nM, *B. atrox* = 2753.0 nM, with EC_{50}s for the remainder ranging from 196.1 – 579.6 nM). These findings are in line with previously published data demonstrating the lower *in vitro* potency of DMPS when compared to MMPis due to their different mechanisms of action [37,38]. Dimercaprol, another heavy metal chelator, demonstrated a pattern of inhibition in line with DMPS, albeit at slightly lower potency (EC_{50} range 228.4 – 4490 nM). The remaining 6 MMPis demonstrated comparable activity to previous studies [38-40]. Prinomastat, batimastat, XL-784 and CTS-1027 demonstrated low nanomolar pan-species inhibition (EC_{50} = 1.5 – 15.4 nM across all four compounds) similar to marimastat, with the exception of CTS-1027 against *B. jararaca* (61.6 nM), and XL-784 against *B. lanceolatus*, *B. atrox,* and *B. jararaca* (EC_{50}s of 34.0 nM for *B. lanceolatus*, 36.3 nM for *B. atrox*, and 41.1 nM for *B. jararaca*). *B. jararaca* and *B. lanceolatus* were not inhibited at any dose of tanomastat (EC_{50} > 10 μM), with the remaining venoms being inhibited at single digit micromolar (EC_{50} range = 1.1 – 5.1 μM). Doxycycline, a tetracycline antibiotic that has been shown to also target matrix metalloproteinases [41], and specifically SVMPs [42], was included as an unrelated compound family [38], though it displayed minimal inhibitory capacity against any of the venoms tested (all EC_{50}s > 10 μM).

Inhibition of PLA$_2$ activity by varespladib

Varespladib has previously been shown to potently inhibit PLA$_2$ activity in a wide variety of venoms, with quoted EC_{50}s in the sub-nanomolar range [21,43], as well as having been employed in a Phase II trial to determine efficacy against snakebite when given in combination with antivenom [44]. In this study, varespladib mirrored the results seen in the literature, with pan-species inhibition at the top dose of 10 μM completely reducing PLA$_2$ activity to baseline in all venoms, and dose response EC_{50}s in the sub- to single-digit nanomolar range (0.2 – 1.6 nM) (Figure 4A and C). Against most venoms, varespladib

displayed similar EC_{50}s of between 0.2 and 0.3 nM, while, perhaps surprisingly, *B. alternatus*

demonstrated the highest EC_{50} of 1.6 nM despite having the lowest PLA_2 activity of the

tested venoms (Figure 2C). However, in an attempt to standardise the relative activities of

the different venoms, we conducted these inhibition experiments using variable venom

concentrations, so the presence of a higher concentration of venom (250 ng for *B. alternatus*

compared with 25-50 ng for the other venoms) seems likely to explain the modest increase

in EC_{50} observed for this species.

Inhibition of SVSP activity by nafamostat

The SVSP activity across the seven venoms was highly variable (Figure 2) and to a greater

extent than the SVMP and PLA_2 activities. Nafamostat is a serine protease inhibitor which

has previously been shown to inhibit the SVSP activities of certain snake venoms at high

doses (10 µM fully inhibited 1 µg venom activities [37]). However, in this study using *Bothrops*

venoms, the 10 µM dose of nafamostat inhibited less than 80% of the SVSP activity of all

venoms at the same 1 µg of venom (Figure 4D). This weak inhibitory activity justified no

further assessment of EC_{50} testing, particularly given that higher dose testing at 80 µM

resulted in only three venoms (*B. asper*, *B. atrox* and *B. jararaca*) being inhibited at greater

than 80%.

Inhibition of coagulopathic venom toxins in bovine plasma

Based on the findings from the *in vitro* assays above, three representative SVMP inhibitors

(marimastat, prinomastat and DMPS), the PLA_2 inhibitor varespladib, and the SVSP inhibitor

nafamostat were tested in the bovine plasma coagulation assay to determine whether any

drug could restore normal coagulation in the presence of the various venoms. First, we

assessed whether any of the compounds affected coagulation in the absence of venom

(Figure 3B). While most had no effect, the serine protease inhibitor nafamostat was shown to

be inherently anticoagulant at 5 µM and higher (Figure 3B) and so was excluded from

downstream dose-response inhibition experiments. For the dose-response experiments,

venom doses were adjusted to provide similar coagulation profiles for each venom to

normalise the window between venom-induced clotting and no-venom clotting across all

venoms (50-250 ng used). Marimastat and prinomastat displayed moderate to potent

inhibition of all venoms at 5 µM (Figure 3C–3I, marimastat 61.8 to 129.1%, prinomastat 67.0

to 121.9%), with a fairly broad EC_{50} range seen in both compounds (marimastat 16.2 to

1357.1 nM, prinomastat 50.9 to 824.4 nM; Table 3J). Though marimastat exhibited EC_{50}s

below 100 nM against more of the venoms (three vs one with prinomastat), it also

demonstrated a high EC_{50} of 1357 nM against *B. lanceolatus*, a venom that prinomastat

inhibited at a comparably lower EC_{50} of 485.5 nM. DMPS required a substantially higher

dose for EC_{50} inhibition than any of the MMPis, which correlates with the lower potency of

this compound in the SVMP assay. At a matched dose of 5 µM, DMPS was only weakly

inhibitory against *B. alternatus, B. asper, B. atrox, B. jararaca* and *B. neuwiedi* venoms

(30.7%, 11.9%, 16.8%, 17.0%, and 26.1% inhibition, respectively, Figure 3, panels C-F, I).

Perhaps surprisingly, for *B. lanceolatus*, a venom that required a higher venom dose to

match the procoagulant profiles of the other venoms (250 ng vs 50-100 ng), 5 µM of DMPS

inhibited the venom by 75% (Figure 3G). In *B. moojeni*, 5 µM of DMPS inhibited the venom

by 53% (Figure 3H). Interestingly, despite all venoms displaying procoagulant profiles, which

could be assumed to be primarily driven by SVMPs, the PLA_2 inhibitor varespladib

demonstrated moderate inhibitory capacity of the procoagulant venom profiles at higher

doses and was capable of generating EC_{50}s for three of the venoms (*B. alternatus, B. asper,*

and *B. atrox,* EC_{50}s all > 50 µM, Figure 3J). However, at the matched 5 µM doses used for

comparative purposes, varespladib had no effect on venom coagulopathy, suggesting this

effect is predominately driven by SVMP toxins (Figure 3, panels C-I).

As marimastat displayed mediocre potency against *B. jararaca* venom in single-inhibitor

experiments (79.2% inhibition at 5 µM and an EC_{50} of 267.0 nM in the coagulation assay,

Figure 3), we postulated that the remaining procoagulant effect of the venom following

inhibition with 5 µM marimastat could have been due to SVSP activity, as described for other

species [45,46]. To explore this, we used nafamostat in combination with marimastat and also

included *B. atrox* venom for comparison (96.7 % inhibition at 5 µM and an EC_{50} of 24.7 nM),

alongside drug only controls to further assess the inherently anticoagulant effect of

nafamostat at lower drug doses. As expected, the procoagulant effect of *B. jararaca* venom

was not fully inhibited by 5 µM marimastat (64.9% inhibition), while 5 µM nafamostat only

modestly inhibited procoagulant effects (13.0% inhibition) (Figure 5A and B). When

combining the two inhibitors, an increase in venom inhibition was observed, with 5 µM doses

of both drugs resulting in a percentage inhibition greater than the effect of marimastat alone

(82.5% vs 64.9%, respectively). This increase in inhibition is roughly equivalent to an

additive effect of both compounds alone. Experiments with *B. atrox* venom showed that the

procoagulant activity of this venom was near fully inhibited by 5 µM marimastat alone

(96.4%), with similarly high inhibition at 2.5 µM (89.4%) and 1.25 µM (83.5%) (Figure 5C and

D), whilst nafamostat displayed no effect on the procoagulant venom profile, even at the 5

µM drug dose shown to be anticoagulant (153.9% drug only inhibition, Figure 5E and F),

indicating that the anticoagulant effect of this compound cannot alter the procoagulant

effects of the SVMPs present in the venom. When combining 5 µM marimastat and

nafamostat with *B. atrox* venom, the resulting coagulation profile shifted to being

anticoagulant (i.e. >100% inhibition, 136.6%), likely the result of the anticoagulant effects of

nafamostat being realised once procoagulant SVMPs were inhibited by marimastat.

Inhibition of coagulopathic venom toxins by human whole blood thromboelastography

To further investigate the plasma coagulopathy results in a more clinically relevant system

we spiked *B. atrox* and *B. jararaca* venom into human whole blood and evaluated the

resulting coagulation profiles using thromboelastography [26]. The methodology utilised in this

study involved the addition of venom with or without marimastat, added to calcium chloride

and whole blood immediately before reading on a ROTEM instrument. Pilot experiments with

nafamostat only revealed direct anticoagulant effects (clotting time [CT]: 3600 at 5 µM and

1368 seconds at 1.25 µM, compared to 682.7 for 5 µM marimastat only and 644.3 seconds

for vehicle control), and thus was excluded from further study. As in the bovine plasma

assay, both *B. atrox* and *B. jararaca* venom presented with a rapid clotting time (mean CT:

B. atrox, 133.3s; *B. jararaca*, 184.3s; vs no venom control, 664.3s) and similar clotting

strength (mean maximum clot firmness [MCF]: *B. atrox,* 68.3; *B. jararaca,* 66.7; vs no venom

control, 58.0) over a similar time frame (clot formation time [CFT]: *B. atrox*, 67.3; *B. jararaca*,

71.0; vs no venom control, 207.7) (Figure 6). The reduced CT time caused by both venoms

was partially inhibited by marimastat, resulting in restoration of 67% and 62% of the longer

CT of the no venom control against *B. atrox* and *B. jararaca* venom, respectively. Despite

this similar inhibition of the CT, there was a notable difference in the ability of marimastat to

reduce the increased clotting strength induced by the two venoms, with *B. jararaca* venom

inhibited by 74% compared with 29% with *B. atrox*. Due to the previously described

interfering effect of nafamostat in this assay, we were unable to explore whether parallel

inhibition of SVSPs might further restore the clotting strength induced by *B. atrox* venom

closer to baseline, though data from the previously described bovine plasma assay (Figure

5) suggested little effect by nafamostat for this venom, hinting that perhaps other toxins

families might be responsible for this phenomenon.

Inhibition of venom lethality in an *in vivo* chicken egg model of envenoming

To further evaluate the potential protective effects of marimastat seen in the coagulation

assay above, we investigated *in vivo* protection in an insensate chicken egg model of

snakebite envenoming. Using *B. atrox* venom as a model, we first demonstrated that topical

application of venom to the vitelline vein of embryos at a dose of 20 µg resulted in clear

observable pathology, characterised by rapid destruction of the vasculature within 1 hour

(Supplementary Figure 1), and lethality by the end of the experimental time course of 6

hours (80% lethality, n=20). Topical treatment with marimastat (0.5 µg, 1.0 µg and 5.0 µg,

n=5 per group) immediately after venom dosing resulted in dose-dependent increases in

efficacy against vasculature damage and venom-induced lethality, with the highest dose

providing complete protection against lethal venom effects (0.5 µg, 60% survival; 1.0 µg,

80% survival; 5.0 µg, 100% survival) (Figure 7 and Supplementary Figure 1). To

contextualise these findings gained with the MMPi marimastat, we repeated these

experiments using the lead SVMP-inhibiting metal chelator, DMPS, at the same therapeutic

doses. The lowest dose of DMPS (0.5 µg) resulted in observable venom pathology and a

survival curve near identical to the venom control (20% survival), indicative of no protection

(Figure 7 and Supplementary Figure 1). However, higher doses of DMPS (1.0 and 5.0 µg)

showed clear evidence of venom neutralisation and provided complete protection against

venom-induced lethality (100% survival) (Figure 7 and Supplementary Figure 1). These

findings further emphasise the promising preclinical potency of this drug [37], despite its

reduced *in vitro* SVMP inhibitory potency in comparison with MMPis.

Finally, to further investigate whether inhibition of SVSP toxins might also be beneficial for *in*

vivo protection against *Bothrops* envenoming, we repeated these *B. atrox* chicken embryo

experiments using the SVSP inhibitor nafamostat. As outlined above, due to its potent off

target anticoagulant effects, we were unable to robustly evaluate the value of inhibiting

SVSP toxins with nafamostat in the coagulation assays, but control doses of nafamostat

alone (5.0 µg) had no observable effect on the embryos (100% survival, Supplementary

Figure 1), facilitating use in these haemotoxicity and lethality focused experiments. Perhaps

unsurprisingly, embryos dosed with nafamostat showed evidence of pathology at all tested

drug doses (0.5, 1.0, 5.0 µg), although at the end of experiment six hours post-dosing, some

evidence of modest efficacy was observed, with 40-60% survival achieved across the three

treatment groups, which compared somewhat favourably with 20% survival reported in the

venom only control group. These findings suggest that inhibition of SVSPs *in vivo* is likely to

be beneficial in the case of *B. atrox* envenoming, though likely of secondary importance

when compared with SVMP inhibition, which can convey complete protection from venom-

induced lethality.

Discussion

Using a wide range of *in vitro* approaches, including venom composition profiling by SDS-

PAGE and toxin specific activity assays, we show that *Bothrops* species have highly variable

SVMP, SVSP and PLA$_2$ activities (Figure 2 and 3), agreeing with the widely reported

literature describing their variable venom composition (Figure 1). For example, *B. alternatus*

and *B. atrox* had the lowest SVSP activities, comparably high SVMP activities, yet

completely distinct PLA$_2$ activities (lowest and second highest). Conversely, *B. moojeni* and

B. neuwiedi presented with the lowest SVMP activity, were in the lower range for PLA_2

activity, but had variable SVSP activity. Crude comparisons between the literature reports of

proteomic venom composition (Figure 1) to the enzymatic activity of the venoms in this study

show similar rankings for *B. alternatus* and *B. atrox* (*B. alternatus*: low PLA_2, low SVSP and

high SVMP. *B. atrox*: high PLA_2, low SVSP and high SVMP), but there was limited

correlation for the remaining venoms. This lack of correlation is not surprising as the

literature clearly describes evidence of variation in venom composition within individual

Bothrops species caused by various factors, such as venom sampling location inclusive of

habitat variation, gender or ontogeny differences and wild caught versus captive bred

(Supplementary Table 1) [5,7,8,47]. It should further be noted that the data we show for each

species in Figure 1 are not uniform, with some species represented by the venom

composition based on a single publication sampling a small pool of individuals, while other

species have multiple publications compromising analysis of multiple pools of venom from

diverse individuals (Supplementary Table 1). Future proteomic characterisations of the

specific venom samples used in this study, which were all sourced from a historical

collection (except for *B. lanceolatus*), would be informative in this regard.

Despite the variation in toxin activity across the *Bothrops* species selected in this study, we

observed pan-species nanomolar neutralisation at the enzymatic level for two small

molecule drugs entering snakebite clinical trials, and which neutralise SVMP (marimastat)

and PLA_2 (varespladib) toxins (Figure 4). The second lead SVMP inhibiting drug, the metal

chelator DMPS, has reduced *in vitro* neutralising potency across the venoms tested (EC_{50}

~200 nM – 3 µM). However, the serine proteinase inhibitor nafamostat required greater than

10 µM to achieve full neutralisation against any of the venoms. SVSP toxins are present in

the venoms of various *Bothrops* species, can be of comparable abundance to SVMP toxins

in certain species [7,48], and had high enzymatic activity in the *B. jararaca*, *B. lanceolatus*, *B.*

moojeni and *B. neuwiedi* venoms tested here (Figure 2). Despite the potentially important

coagulopathic effect of SVSPs on the haemostatic system, small molecule inhibitors of

SVSPs have received limited attention, in contrast to the recent drug discovery efforts

activity against PLA_2 and SVMP toxins. Here we used the previously described, repurposed,

serine protease inhibitor nafamostat, which was previously used in small molecule

combination therapies to investigate its value in preventing venom lethality preclinically [37].

However, nafamostat showed limited additive value compared to a combination of SVMP

and PLA_2 inhibitors in that study, while the low *in vitro* inhibitory potency of the drug

described here (>10 µM EC_{50}) coupled with its innate anticoagulant nature via interactions

with cognate serine proteases such as thrombin [49], severely hampered our assessment of

the role SVSPs play in the coagulation disturbances caused by *Bothrops* venoms in the

plasma coagulation and whole blood TEG assays. Nonetheless, evidence of modest

protection conferred by nafamostat against the venom of *B. atrox* in the *in vivo* chicken egg

embryo assay further suggests that SVSPs are likely important pathological components of

Bothrops venoms, as previously described [50]. There therefore remains a strong rationale for

future drug discovery activities to identify novel broad-spectrum inhibitors against SVSP

toxins, with a focus on optimising their potency and selectively to avoid off target effects

against mammalian serine proteases involved in the coagulation cascade [51].

Viperid PLA_2 toxins induce coagulopathy primarily through their anticoagulant activity.

Although we identified clear enzymatic activity of PLA_2 across many of our sampled species,

we observed no anticoagulant activity in either our plasma or whole blood testing, even

when procoagulant SVMP toxins were inhibited. Despite this lack of coagulopathic effect, the

potent pan-species inhibitory activity of varespladib observed in the PLA_2 enzymatic assay

(EC_{50}s ranging from 0.14 to 1.45 nM) is likely still to be beneficial at inhibiting non-coagulant

PLA_2-associated pathology, such as tissue damage, inflammation and myotoxicity [43,52]. The

potential value of a PLA_2 inhibitor against *Bothrops* venoms is further highlighted by prior

reports of poor antivenom performance against certain PLA_2 isoforms due to either their lack

of inclusion in the immunisation mixture or due to poor tissue penetration of the antivenom

[25,31,35].

Challenges associated with the neutralising efficacy of antivenoms, such as Instituto

Butantan's polyvalent antivenom antibotrópico, have also been reported against certain

SVMP isoforms [31,35]. The pan-species activity of MMPis like marimastat is widely reported to

be facilitated by broad spectrum inhibitory effects against varying isoforms of SVMPs [38,53]. In

our studies, marimastat demonstrated pan-species inhibitory activity across diverse venoms

with nanomolar EC_{50}s in the enzymatic assay (1.8 to 10.6 nM) and plasma coagulation

assay (16.2 to 592.9 nM, except for *B. lanceolatus*, 1357.0 nM). These results were further

supported via the use of TEG assessment of coagulopathy in human blood, where the

distinct venoms of two WHO category one (highest medical importance) species, *B. atrox*

and *B. jararaca*, were used [54]. In Brazil, *B. atrox* is implicated in 80-90% of snakebites [24], but

this species also extends into the Amazonian regions of many other countries (Figure 1A),

while *B. jararaca* is found in the southern states of Brazil, northeastern Paraguay and

northern Argentina, where it is a leading cause of snakebite, especially in the densely

populated areas of southeastern Brazil [7]. Despite their differing *in vitro* profiles, in the TEG

human whole blood assay, marimastat rescued the procoagulant activity of *B. atrox* and *B.*

jararaca venom by inhibiting the induced rapid clotting time and increased clotting strength,

reverting the clotting profile towards normal parameters seen in the absence of venom.

Further, marimastat provided protection in an *in vivo* model of haemotoxicity, with chicken

egg embryos dosed with *B. atrox* venom fully protected against lethality and the vascular

destructive effects caused by this venom at the 5 µg therapeutic dose tested (vs 20 µg

venom). These findings provide further confidence in the potential therapeutic value of

marimastat, which remains a lead candidate repurposed drug currently approaching clinical

development. Future evaluation of its safety and efficacy in a Phase II clinical trial against *B.*

atrox envenomings will be revealing.

As seen in previous work, there was clear superiority in terms of the *in vitro* SVMP inhibitory

potency of matrix metalloproteinase inhibitors over metal chelators (Figure 4A) [38].

Marimastat and prinomastat exhibited EC_{50}s in the low nanomolar range (from 0.7 nM for

prinomastat vs *B. asper* to 10.6 nM for marimastat vs *B. lanceolatus*) in the SVMP assay,

resulting in at least a 20-fold increase in potency, and often >100-fold potency, over the lead

metal chelator DMPS (EC_{50} range of 196.1 to 2019 nM). A similar trend was observed in the

coagulation assay (EC_{50} range of 16.2-1357.0 nM for marimastat and 50.9-824.4 nM for

prinomastat vs 1.2 µM to >320 µM for DMPS), although the anti-coagulopathic potency of

DMPS against *B. lanceolatus* venom was dramatically lower than all other venoms tested

(1242 nM) and highly comparable to the EC_{50} of marimastat against this venom (1357 nM).

Given that envenomings by *B. lanceolatus* are known to often be somewhat distinct to bites

by other *Bothrops* species by often presenting with severe thrombotic complications [55],

further evaluation of the potential protective effects of the already licensed drug DMPS in a

preclinical setting would be of great interest. Finally, despite the considerable differences in

in vitro potency mentioned above, DMPS provided highly comparable protective effects to

marimastat against the *in vivo* pathology caused in chicken egg embryos by *B. atrox* venom.

In this pilot study measuring *in vivo* haemotoxicity, treatment with 1 µg and 5 µg of

marimastat resulted in 80% and 100% protection against lethal venom effects, while DMPS

was fully protective at both these doses. These findings further highlight the apparent *in*

vitro/*in vivo* potency disconnect previously described for this metal chelator [56] and highlight

that caution should be applied when triaging SVMP inhibitors with different mechanisms of

action to MMPis based solely on *in vitro* enzymatic inhibitory potency.

This manuscript reinforces prior work demonstrating that, although *Bothrops* species

dominate the snakebite incidence across the Neotropics resulting in bothropic syndrome, the

diversity of venom composition found across this genus of snakes is substantial. Here we

evidence this variation in a comparative manner, highlighting both similarities and

differences in the enzymatic activity (SVMP, SVSP and PLA_2) and coagulopathic (*in vitro*

bovine plasma coagulation and TEG human whole blood) effects of various *Bothrops*

venoms. Despite such variation, we show that three of the leading small molecule treatments
currently progressing into clinical trials (marimastat, DMPS and varespladib) can collectively
neutralise all seven venoms in terms of their enzymatic PLA$_2$ and SVMP activity, while the
SVMP inhibitors were also effective against the action of coagulotoxins and provided *in vivo*
protection against lethal haemotoxicity in a chicken embryo model. Future priorities in this
space include: i) evaluating the diversity of toxin isoforms that are inhibited by these specific
drugs, including those described to have limited neutralisation by available antivenoms [9,25,35],
ii) undertaking drug discovery activities to identify novel SVSP inhibitors specific to venom
toxins, and iii) progressing lead repurposed drugs and drug combinations into conventional
murine preclinical models [11] to evaluate their efficacy against systemic and local envenoming
in vivo. Overall, this study provides convincing evidence of the potential value of small
molecule-based toxin inhibitors for the treatment of snakebite in the Neotropics, adding
further weight to the recent paradigm shift towards early therapeutic interventions via oral
dosing in community settings [15,22,44]. Indeed, certain settings within the Neotropics provide
exciting potential for the robust, future evaluation of the efficacy of small molecule drugs
against snakebite - in particular the use of SVMP inhibitors against *B. atrox* - and the
outcomes of future clinical trials have the potential to provide valuable proof of concept for
the future translation of safe and effective oral snakebite drugs.
534
535

536   **Author contribution**
537   R.H.C., N.R.C., A.W. and S.K.M. conceptualised the project and wrote the manuscript, with
538   funding acquired by N.R.C. The *in vitro* bioassays were performed by A.W., R.H.C. and
539   S.K.M. The *in vivo* embryo assays were performed by E.S., L-O.A. and T.D.K. All authors
540   analysed the results and reviewed the manuscript.
541

Acknowledgements
We gratefully acknowledge MicroPharm Limited for provision of *Bothrops lanceolatus*
venom. We also thank Camille Abada and Iara Cardoso for venepuncture and anonymised
donors for their provision of blood samples. Our thanks are extended to Dr. Charlotte
Dawson for the establishment of the chicken egg model and training the authors on its use.
This work was funded by Wellcome (#221712/Z/20/Z to N.R.C.).
548

Methods
Venoms

Representative venoms were selected to cover the diversity of the *Bothrops* genus. All,

except for *Bothrops lanceolatus* (which was gifted by MicroPharm Limited), were historical

samples sourced from the herpetarium facility at the Centre for Snakebite Research and

Interventions (CSRI) at the Liverpool School of Tropical Medicine (LSTM). The species were:

Bothrops alternatus (Brazil), *B. asper* (Costa Rica – Atlantic), *B. atrox* (Colombia), *B.*

jararaca (Brazil), *B. lanceolatus* (Martinique), *B. moojeni* (Brazil) and *B. neuwiedi* (Brazil).

Crude venoms were stored lyophilised at 2–8 °C before reconstitution to 10 mg/mL in sterile

Phosphate Buffered Saline (PBS, pH 7.4) (Gibco, Cat.no. 10010023) prior to use.

Drugs

The small molecule drugs used in this study were selected based on their previously

reported inhibitory activity against snake venom SVMP, PLA_2 or SVSP toxins [37,38,43,53]. The

SVMP-inhibiting matrix metalloproteinase (MMP) inhibitors were sourced from

MedChemExpress - prinomastat hydrochloride (Cat. no. HY-12170A), XL-784 (HY-19485,

98.25%) and CTS-1027 (HY-10398, 99.24%); Sigma-Aldrich - marimastat (Cat. no. M2699),

batimastat (Cat. no. SML0041) and doxycycline (Cat. no. D9891); and Cayman chemicals -

tanomastat (Cat. no. 9258). The SVMP inhibiting metal chelators were dimercaprol (Cat. no.

64046, Sigma-Alrich) and DMPS (Cat. no. H56578, Alfa Aesar). The PLA_2 inhibitor was

varespladib (Cat no: SML1100, Sigma) and the SVSP inhibitor was nafamostat mesylate

(Cat. no. ab141432, Abcam). All drugs were resuspended in dimethyl sulfoxide (DMSO)

(Cat. no. D2650-100ML, Sigma-Aldrich).

SDS-PAGE gel electrophoresis

Five micrograms of each venom were mixed in an equal volume of 2 X sample loading buffer

(62.5 mM Tris-Cl pH 6.8, 25% v/v glycerol, 2% SDS, 0.75% bromophenol blue) with 100 mM

dithiothreitol, incubated at 100°C for 5 minutes, before loading on a 4–20% Mini-Protean

TGX gel (BioRad, Cat. no. 456-1096) with the addition of a molecular weight marker (5 µL,

PageRuler Prestained Protein Ladder, Thermo, Cat. no. 26616) to one lane of the gel. After

electrophoresis at 200 V for 30 minutes, the gel was stained with Coomassie blue (50%

methanol, 40% deionized water, 10% glacial acetic acid and 0.1% Coomassie Brilliant Blue)

for 1 hour at room temperature with gentle shaking, then de-stained for 2 hours at room

temperature in 50% methanol, 40% deionized water and 10% glacial acetic acid. Gels were

rinsed in deionised water and imaged on a GelDoc (Bio-Rad) under white light.

SVMP *in vitro* assay

The SVMP assay utilises a quenched fluorescent substrate for MMPs previously utilised to

assess the activity of SVMPs [36-38,53]. One microgram per well of venom was added to a flat-

bottomed 384-well plate (Greiner, Cat. no. 781101) in 15 µl PBS, before incubation at 37°C

for 25 minutes (to keep conditions identical to the drug inhibition assays described below).

Plates were then allowed to acclimatise to room temperature for 5 minutes, before 75 µl of

fluorescent substrate (Bio Techne, Cat. no. ES010) at a final 7.5 µM reaction concentration

in SVMP assay buffer (50mM Tris HCl pH 7.5, 150mM NaCl) was added to each well.

Immediately following addition of substrate, the fluorescence (excitation 320 nm, emission

420 nm) was read kinetically on a CLARIOstar Plus (BMG labtech). Venom inhibition assays

were conducted in the same manner, with an additional preliminary step in which various

concentrations of the drugs (dose range of 10 µM – 0.17 nM) were created in DMSO. These

were then stamped as 0.91 µL droplets at 100x the desired final concentration (final well

volume being 91 µL) onto flat-bottomed 384-well plate wells prior to venom addition, allowing

drug-venom interaction prior to substrate addition. For both protocols the AUC over 108

minutes was calculated for every condition to quantify activity. This time frame was selected

based on prior knowledge of active venoms having capacity to fully convert the substrate [36-

38,53]. For the inhibitor testing the AUC was converted to a percentage of venom inhibition by

normalising to the negative and positive controls. These values were then plotted as a dose-

response curve, and EC_{50} values calculated.

PLA$_2$ *in vitro* assay

The PLA$_2$ assay uses a commercially available colorimetric assay kit (Abcam, Cat. no.

ab133089) and relies on the cleavage of dithiol groups from a substrate (PLA2 Diheptanoyl

Thio-PC) by PLA$_2$s. A third component, DTNB, is added to the reaction, and binds to the

freely available thiol groups, producing a colorimetric change. This kit has been adapted for

venom PLA$_2$ activity assessment [37,43]. For venom activity assays, 100 ng of each venom in a

10 µL volume were added to appropriate wells of a flat-bottomed 384-well plate (Greiner,

Cat. no. 781101) then incubated for 25 minutes at 37°C for consistency with later dose-

response experiments. Following incubation, the plates were allowed to acclimatise to room

temperature prior to addition of 5 µL DTNB (resuspended in H_2O) and 30 µL of PLA$_2$

substrate (diluted in supplied assay buffer), and the plates read kinetically using an

absorbance protocol (405 nm) on a CLARIOstar Plus. Thereafter, for each venom, the

highest venom dose that gave a linear increase in absorbance over the 15-minute read time

was selected for drug inhibition studies, in which dose-response curves of varespladib (final

concentration range of 11.1 µM – 1.11 pM) were created in DMSO and first stamped onto a

flat-bottomed 384-well plate as 0.5 µL droplets at 90x the desired final concentration (45uL

final well volume) to allow for drug-venom interaction prior to substrate addition. The

calculated AUCs over 12 minutes (based on the Abam guidelines) were normalised to a

percentage inhibition (compared to the positive and negative controls) and plotted as dose

response curves, allowing for EC_{50} to be calculated.

SVSP *in vitro* assay

The serine protease activity of the venoms was tested using a commercial chromogenic

broad-spectrum peptide substrate (S-2288, Quadratech Diagnostics Ltd) to quantify the

cleavage of the substrate by serine protease via absorbance, as previously described [37]. The

substrate was diluted in water to make a 6 mM stock solution. The reaction was performed

with the addition of venom, buffer and substrate at a volume ratio of 1:1:1 with a final volume

of 45 µL per well of a 384-well plate (Greiner, Cat. no. 781101). For each of the venoms, 1

µg was added per well (15 µL of 0.07 µg/µL diluted in PBS from a 10 mg/mL stock) prior to

the addition of 15 µL of buffer (100 mM Tris pH 8.5, 100 mM NaCl) and incubated for 30

minutes at 37 ºC. Following this incubation, 15 µL of the substrate stock solution was added

per well (2 mM final concentration) and absorbance immediately measured kinetically at 405

nm on a CLARIOstar Plus plate reader at 37 ºC. Drug activity was investigated identically to

above, with an additional preliminary step in which dose-response curves were created in

DMSO, via a 12-point curve ranging from 80 µM to 39 nM in 1:2 dilution steps. From each

dose 0.5 µL was added to the venom in each relevant well and preincubated for 30 minutes

at 37 ºC before the buffer and substrate was added. For analysis, the AUC was calculated

over a 35-minute time interval and all data normalised by subtracting the no venom control

(PBS). This time frame was selected based on prior knowledge of active venoms having

capacity to fully convert the substrate. The inhibitor data was normalised to a percentage

inhibition value due to the variation in venom activity.

Bovine plasma coagulation assay

The plasma assay utilises citrated bovine plasma, which is incoagulable until the addition of

calcium. Factors involved in the coagulation cascade are targets for specific venom toxins,

and so addition of venom and calcium to the citrated bovine plasma can cause the plasma to

clot faster than normal (procoagulant toxins) or to not clot at all (anticoagulant toxins). In

venom activity experiments, 10 µL of venom at different doses (1 µg – 5 ng) was added to

384-well plates (Greiner, Cat. no. 781101) to determine the most appropriate venom dose

for downstream drug inhibition studies. Following a 25-minute incubation at 37°C for

consistency with future dose response studies, the plates were acclimatised to room

temperature for 5 minutes, before the addition of 20 µL of 20mM $CaCl_2$, followed

immediately by 20 µL of citrated bovine plasma (Biowest, Cat. no. S0260), which had been

centrifuged for 5 minutes at 3000 RCF to pellet any particulate matter. Immediately following

plasma addition, the plates were read kinetically on a CLARIOstar Plus at 595 nm

absorbance. To provide comparable procoagulant clotting profiles, doses were selected that

induced complete clotting before the negative control (no venom) wells had initiated clotting.

Preliminary drug only experiments were conducted to identify any inherently pro- or anti-

coagulant effects, with only compounds that had no inherent effect on coagulation

progressed to dose-response venom inhibition experiments. In these experiments, dose-

response curves of inhibitory compounds (as identified in the previous *in vitro* assays) were

created in DMSO and 0.5 µL stamped per well of a 384-well flat-bottomed plate at 100x the

desired final concentration (final doses of 80 µM – 0.8 nM for all compounds except DMPS

tested at 320 µM – 1.3 nM due to reduced potency). Thereafter the experimental workflow

was as described above, with the addition of venom, incubation of venom and drug, followed

by $CaCl_2$ and plasma. As with other assays, AUCs were calculated and normalised to the

positive and negative controls to determine percentage inhibition, which was used to plotted

as dose response curves, and generate EC_{50} values.

Human blood thromboelastography

The clotting profiles of two representative venoms (*B. atrox* and *B. jararaca*) and the effects

of the SVMP and SVSP inhibitors marimastat and nafamostat were tested using

thromboelastography [26,39,53,57-59]. Blood from healthy consenting donors were collected

according to ethically approved protocols (LSTM research tissue bank, REC ref.

11/H1002/9) and used up to 4 hours post-sampling with three independent replicates per

experimental condition. The blood was collected into BD Vacutainer tubes with ACD-A

anticoagulant solution (sodium citrate: 22.0 g/L, dextrose: 24.5 g/L, citric acid: 8.0 g/L and

antimycotic [potassium sorbate] reagent: 0.15 g/L (Fisher Scientific, Cat. no. BD 366645)).

For each experiment 1.2 mL of whole blood was pre-heated at 37 °C for 5 minutes on a heat

block. During this time all reagents were added into the pre-heated sample cup at the

following volumes: 12 µL venom sample or PBS control, 15 µL drug sample or PBS control

and 20 µL $CaCl_2$ (Star-tem, Cat. no. 503-01), before the final addition of 253 µL of whole

blood. Viscoelasticity data were then recorded immediately at 37 °C for 60 min using a

ROTEM Delta™ (Werfen). Venom concentrations were at 0.6 µg/reaction (12 µL of 50

µg/mL) diluted in PBS from the 10 mg/mL venom stock. The drug concentrations were

diluted in PBS from the 10 mM stock to a 5 µM reaction concentration (15 µL of 100 µM

working solution). The negative control (corresponding to spontaneous coagulation of whole

blood following recalcification) consisted of no venom or drugs but included $CaCl_2$. The

positive control consisted of venom and $CaCl_2$ without drug treatment. To ensure the $CaCl_2$

did not interfere with the drug treatments, each drug was run with $CaCl_2$ but no venom. The

parameters assessed by ROTEM include coagulation time (CT) and maximum clot firmness

(MCF), visualised graphically in an XY plot to demonstrate a clotting profile using Prism v10

software (GraphPad). Additional measures of clot formation time (CFT), alpha angle, and amplitude in 5 minutes (A5) were also reported. These parameters are defined as follows: CT (seconds), time from the start of the measurement until the initiation of clotting classified as 2 mm amplitude; MCF (mm), maximum amplitude of clot firmness reached during the run time, used as a proxy for 'clotting strength'; CFT (seconds), time interval between the initiation of clotting (2 mm amplitude) until a clot firmness of 20 mm is achieved; alpha angle (∘), the angle between the baseline and tangent to the clotting curve through the 2 mm point. Each parameter is dependent upon different elements of the clotting process. CT represents the coagulation activation via the enzymatic activity of coagulation factors. CFT and alpha angle are dependent on thrombin generation, platelet count/function as well as fibrinogen levels and fibrin polymerization. MCF and A5 are dependent on platelet count/function, fibrin concentration/formation and factor XIII [26,60].

Chicken egg *in vivo* assay of haemotoxicity

Egg embryos have previously been used to assay venom pathology (e.g., haemorrhage, coagulation, inflammation and lethality) in a vascularised environment and as an efficacy read out for snakebite treatments [61-63]. Fertilised chicken eggs (Medeggs Ltd, UK) on day 1 post-fertilisation were placed horizontally in an incubator at 37°C until day 5, at which point they were sprayed with 70% ethanol and candled to mark embryo position. Damaged and infertile eggs were discarded. A sterile windowing procedure was then conducted to provide visibility of embryos; 6-8 mL albumin was removed using a 23G needle and 10 mL syringe, and the shell was reinforced with clear tape before the marked area was removed using sharp dissection scissors. Eggs were then covered with parafilm and returned to the incubator until day 6. Venom neutralisation studies for marimastat, DMPS or nafamostat were tested at a drug dose of 0.5 µg, 1 µg or 5 µg per egg, resuspended in 5% DMSO, with control experiments demonstrating no effect of 5% DMSO or drug only activity on embryo pathology. On day 6 post-fertilisation, eggs were randomly assigned to dose groups. The group size was 5 for all drug treatment groups, with control groups having increased sample sizes via incorporation across multiple independent experiments (PBS negative control, n=12; venom only control, n=20). Egg embryos were dosed with 10 µL PBS, or 20 µg *B. atrox* venom in 2 µL PBS (5% DMSO) followed by either 9 µL PBS (5% DMSO) or 9 µL treatment (5% DMSO). Doses were pipetted directly onto the vitelline vein on the ventral side of the embryo, with second doses applied immediately after the first at the same location. Embryo survival was determined thereafter using observation of the embryo's heartbeat at multiple timepoints, up to 6 hours, and plotted as Kaplan-Meier survival curves. Pathology was also monitored prior to dosing and throughout the time course using a microscope (Motic SMZ-171) and a mounted smartphone used to capture representative

images of selected embryos at a consistent magnification and orientation. At the end of the

time course, any surviving embryos were culled by cervical dislocation.

Statistics

All calculations (including EC_{50} and AUC) and figures (including dose response curves and

Kaplan-Meier survival curves) were generated using Prism v10 software (GraphPad).

Standard deviation (SD) for all kinetic reads and standard error for AUC is reported based on

Prism v10. The outputs from the thromboelastography profiles are calculated by the

integrated software on the ROTEM sigma (Werfen). For combination drug coagulation

assays, ordinary one-way ANOVA with Šídák's multiple comparisons test was used to

calculate significant differences ($p = <0.05$) between matched conditions. The sample sizes

for HTS assays including the SVMP, PLA_2 and coagulation experiment are the average of

the means from independent assays ($n >2$ within each independent assay). For the serine

protease assay, thromboelastography and chicken egg model the sample sizes are

individual values due to lower throughput and venom availability.

References

References

1 Gutierrez, J. M. *et al.* Snakebite envenoming. *Nat Rev Dis Primers* **3**, 17079 (2017).
https://doi.org/10.1038/nrdp.2017.79
2 Menzies, S. K., Patel, R. N. & Ainsworth, S. Practical progress towards the
development of recombinant antivenoms for snakebite envenoming. *Expert Opin
Drug Discov* **20**, 799–819 (2025). https://doi.org/10.1080/17460441.2025.2495943
3 Pucca, M. B. *et al.* History of Envenoming Therapy and Current Perspectives. *Front
Immunol* **10**, 1598 (2019). https://doi.org/10.3389/fimmu.2019.01598
4 Casewell, N. R., Jackson, T. N. W., Laustsen, A. H. & Sunagar, K. Causes and
Consequences of Snake Venom Variation. *Trends Pharmacol Sci* **41**, 570–581
(2020). https://doi.org/10.1016/j.tips.2020.05.006
5 Alape-Giron, A. *et al.* Studies on the venom proteome of Bothrops asper:
perspectives and applications. *Toxicon* **54**, 938–948 (2009).
https://doi.org/10.1016/j.toxicon.2009.06.011
6 Menezes, M. C., Furtado, M. F., Travaglia-Cardoso, S. R., Camargo, A. C. &
Serrano, S. M. Sex-based individual variation of snake venom proteome among
eighteen Bothrops jararaca siblings. *Toxicon* **47**, 304–312 (2006).
https://doi.org/10.1016/j.toxicon.2005.11.007
7 Goncalves-Machado, L. *et al.* Combined venomics, venom gland transcriptomics,
bioactivities, and antivenomics of two Bothrops jararaca populations from geographic
isolated regions within the Brazilian Atlantic rainforest. *J Proteomics* **135**, 73–89
(2016). https://doi.org/10.1016/j.jprot.2015.04.029
8 Amorim, F. G. *et al.* Proteopeptidomic, Functional and Immunoreactivity
Characterization of Bothrops moojeni Snake Venom: Influence of Snake Gender on
Venom Composition. *Toxins (Basel)* **10** (2018).
https://doi.org/10.3390/toxins10050177

9   Calvete, J. J., Juarez, P. & Sanz, L. Snake venomics. Strategy and applications. *J*
Mass Spectrom **42**, 1405–1414 (2007). https://doi.org/10.1002/jms.1242
10   Gutierrez, J. M. *et al.* The Search for Natural and Synthetic Inhibitors That Would
Complement Antivenoms as Therapeutics for Snakebite Envenoming. *Toxins (Basel)*
13 (2021). https://doi.org/10.3390/toxins13070451
11   Gutierrez, J. M., N, R. C. & Laustsen, A. H. Progress and Challenges in the Field of
Snakebite Envenoming Therapeutics. *Annu Rev Pharmacol Toxicol* **65**, 465–485
(2025). https://doi.org/10.1146/annurev-pharmtox-022024-033544
12   Ledsgaard, L. *et al.* Discovery and optimization of a broadly-neutralizing human
monoclonal antibody against long-chain alpha-neurotoxins from snakes. *Nat*
Commun **14**, 682 (2023). https://doi.org/10.1038/s41467-023-36393-4
13   Ahmadi, S. *et al.* Nanobody-based recombinant antivenom for cobra, mamba and
rinkhals bites. *Nature* (2025). https://doi.org/10.1038/s41586-025-09661-0
14   Vazquez Torres, S. *et al.* De novo designed proteins neutralize lethal snake venom
toxins. *Nature* **639**, 225–231 (2025). https://doi.org/10.1038/s41586-024-08393-x
15   Clare, R. H., Hall, S. R., Patel, R. N. & Casewell, N. R. Small Molecule Drug
Discovery for Neglected Tropical Snakebite. *Trends Pharmacol Sci* **42**, 340–353
(2021). https://doi.org/10.1016/j.tips.2021.02.005
16   Rudresha, G. V., Khochare, S., Casewell, N. R. & Sunagar, K. Preclinical evaluation
of small molecule inhibitors as early intervention therapeutics against Russell's viper
envenoming in India. *Commun Med (Lond)* **5**, 226 (2025).
https://doi.org/10.1038/s43856-025-00900-z
17   Dawson, C. A. *et al.* (2024). https://doi.org/10.1101/2024.03.13.584772
18   Hall, S. R. *et al.* Repurposed drugs and their combinations prevent morbidity-
inducing dermonecrosis caused by diverse cytotoxic snake venoms. *Nat Commun*
14, 7812 (2023). https://doi.org/10.1038/s41467-023-43510-w
19   Gutierrez, J. M., Lewin, M. R., Williams, D. J. & Lomonte, B. Varespladib (LY315920)
and Methyl Varespladib (LY333013) Abrogate or Delay Lethality Induced by
Presynaptically Acting Neurotoxic Snake Venoms. *Toxins (Basel)* **12** (2020).
https://doi.org/10.3390/toxins12020131
20   Gerardo, C. J. *et al.* Oral varespladib for the treatment of snakebite envenoming in
India and the USA (BRAVO): a phase II randomised clinical trial. *BMJ Glob Health* **9**
(2024). https://doi.org/10.1136/bmjgh-2024-015985
21   Lewin, M. R. *et al.* Varespladib in the Treatment of Snakebite Envenoming:
Development History and Preclinical Evidence Supporting Advancement to Clinical
Trials in Patients Bitten by Venomous Snakes. *Toxins (Basel)* **14** (2022).
https://doi.org/10.3390/toxins14110783
22   Abouyannis, M. *et al.* Development of an oral regimen of unithiol for the treatment of
snakebite envenoming: a phase 1 open-label dose-escalation safety trial and
pharmacokinetic analysis in healthy Kenyan adults. *EBioMedicine* **113**, 105600
(2025). https://doi.org/10.1016/j.ebiom.2025.105600
23   Mise, Y. F., Lira-da-Silva, R. M. & Carvalho, F. M. Time to treatment and severity of
snake envenoming in Brazil. *Rev Panam Salud Publica* **42**, e52 (2018).
https://doi.org/10.26633/RPSP.2018.52
24   da Silva, W. *et al.* Who are the most affected by Bothrops snakebite envenoming in
Brazil? A Clinical-epidemiological profile study among the regions of the country.
PLoS Negl Trop Dis **17**, e0011708 (2023).
https://doi.org/10.1371/journal.pntd.0011708
25   Sousa, L. F. *et al.* Diversity of Phospholipases A(2) from Bothrops atrox Snake
Venom: Adaptive Advantages for Snakes Compromising Treatments for Snakebite
Patients. *Toxins (Basel)* **14** (2022). https://doi.org/10.3390/toxins14080543
26   Larreche, S. *et al.* Bothrops atrox and Bothrops lanceolatus Venoms In Vitro
Investigation: Composition, Procoagulant Effects, Co-Factor Dependency, and
Correction Using Antivenoms. *Toxins (Basel)* **15** (2023).
https://doi.org/10.3390/toxins15100614

Hui Wen, F. *et al.* Snakebites and scorpion stings in the Brazilian Amazon: identifying research priorities for a largely neglected problem. *PLoS Negl Trop Dis* **9**, e0003701 (2015). https://doi.org/10.1371/journal.pntd.0003701

Guerra-Duarte, C., de Sousa, V. P., de Oliveira-Sousa, G. & Gomes Mol, M. P. Use of snake antivenom in the Region of the Americas: a systematic review. *Bull World Health Organ* **103**, 785–798 (2025). https://doi.org/10.2471/BLT.24.291941

Kasturiratne, A. *et al.* The global burden of snakebite: a literature analysis and modelling based on regional estimates of envenoming and deaths. *PLoS Med* **5**, e218 (2008). https://doi.org/10.1371/journal.pmed.0050218

Afroz, A., Siddiquea, B. N., Chowdhury, H. A., Jackson, T. N. & Watt, A. D. Snakebite envenoming: A systematic review and meta-analysis of global morbidity and mortality. *PLoS Negl Trop Dis* **18**, e0012080 (2024). https://doi.org/10.1371/journal.pntd.0012080

Calvete, J. J. *et al.* Snake population venomics and antivenomics of Bothrops atrox: Paedomorphism along its transamazonian dispersal and implications of geographic venom variability on snakebite management. *J Proteomics* **74**, 510–527 (2011). https://doi.org/10.1016/j.jprot.2011.01.003

Albuquerque Barbosa, F. B. *et al.* Dermatopathological findings of Bothrops atrox snakebites: A case series in the Brazilian Amazon. *PLoS Negl Trop Dis* **18**, e0012704 (2024). https://doi.org/10.1371/journal.pntd.0012704

Larreche, S. *et al.* Bleeding and Thrombosis: Insights into Pathophysiology of Bothrops Venom-Related Hemostasis Disorders. *Int J Mol Sci* **22** (2021). https://doi.org/10.3390/ijms22179643

Xiao, H., Pan, H., Liao, K., Yang, M. & Huang, C. Snake Venom PLA(2), a Promising Target for Broad-Spectrum Antivenom Drug Development. *Biomed Res Int* **2017**, 6592820 (2017). https://doi.org/10.1155/2017/6592820

Sousa, L. F. *et al.* Comparison of phylogeny, venom composition and neutralization by antivenom in diverse species of bothrops complex. *PLoS Negl Trop Dis* **7**, e2442 (2013). https://doi.org/10.1371/journal.pntd.0002442

Menzies, S. K. *et al.* In vitro and in vivo preclinical venom inhibition assays identify metalloproteinase inhibiting drugs as potential future treatments for snakebite envenoming by Dispholidus typus. *Toxicon X* **14**, 100118 (2022). https://doi.org/10.1016/j.toxcx.2022.100118

Albulescu, L. O. *et al.* A therapeutic combination of two small molecule toxin inhibitors provides broad preclinical efficacy against viper snakebite. *Nat Commun* **11**, 6094 (2020). https://doi.org/10.1038/s41467-020-19981-6

Clare, R. H. *et al.* Snakebite drug discovery: high-throughput screening to identify novel snake venom metalloproteinase toxin inhibitors. *Front Pharmacol* **14**, 1328950 (2023). https://doi.org/10.3389/fphar.2023.1328950

Rucavado, A. *et al.* A murine experimental model of the pulmonary thrombotic effect induced by the venom of the snake Bothrops lanceolatus. *PLoS Negl Trop Dis* **18**, e0012335 (2024). https://doi.org/10.1371/journal.pntd.0012335

Youngman, N. J., Lewin, M. R., Carter, R., Naude, A. & Fry, B. G. Efficacy and Limitations of Chemically Diverse Small-Molecule Enzyme-Inhibitors against the Synergistic Coagulotoxic Activities of Bitis Viper Venoms. *Molecules* **27** (2022). https://doi.org/10.3390/molecules27051733

Stechmiller, J., Cowan, L. & Schultz, G. The role of doxycycline as a matrix metalloproteinase inhibitor for the treatment of chronic wounds. *Biol Res Nurs* **11**, 336–344 (2010). https://doi.org/10.1177/1099800409346333

Arens, D. K. *et al.* Doxycycline-Mediated Inhibition of Snake Venom Phospholipase and Metalloproteinase. *Mil Med* **189**, e2430–e2438 (2024). https://doi.org/10.1093/milmed/usae184

Albulescu, L. O. *et al.* Optimizing drug discovery for snakebite envenoming via a high-throughput phospholipase A2 screening platform. *Front Pharmacol* **14**, 1331224 (2023). https://doi.org/10.3389/fphar.2023.1331224

44 Gerardo, C. J. *et al.* Oral varespladib for the treatment of snakebite envenoming in
India and the USA (BRAVO): a phase II randomised clinical trial. *BMJ Glob Health* **9**
(2024). https://doi.org/10.1136/bmjgh-2024-015985
45 Latinovic, Z. *et al.* The Procoagulant Snake Venom Serine Protease Potentially
Having a Dual, Blood Coagulation Factor V and X-Activating Activity. *Toxins (Basel)*
12 (2020). https://doi.org/10.3390/toxins12060358
46 Torres-Huaco, F. D. *et al.* Rapid purification and procoagulant and platelet
aggregating activities of Rhombeobin: a thrombin-like/gyroxin-like enzyme from
Lachesis muta rhombeata snake venom. *Biomed Res Int* **2013**, 903292 (2013).
https://doi.org/10.1155/2013/903292
47 Nunez, V. *et al.* Snake venomics and antivenomics of Bothrops atrox venoms from
Colombia and the Amazon regions of Brazil, Peru and Ecuador suggest the
occurrence of geographic variation of venom phenotype by a trend towards
paedomorphism. *J Proteomics* **73**, 57–78 (2009).
https://doi.org/10.1016/j.jprot.2009.07.013
48 Cavecci-Mendonca, B. *et al.* Preliminary Insights of Brazilian Snake Venom
Metalloproteomics. *Toxins (Basel)* **15** (2023). https://doi.org/10.3390/toxins15110648
49 Zhao, C. *et al.* Delayed administration of nafamostat mesylate inhibits thrombin-
mediated blood-spinal cord barrier breakdown during acute spinal cord injury in rats.
J Neuroinflammation **19**, 189 (2022). https://doi.org/10.1186/s12974-022-02531-w
50 Serrano, S. M. The long road of research on snake venom serine proteinases.
Toxicon **62**, 19–26 (2013). https://doi.org/10.1016/j.toxicon.2012.09.003
51 Romanazzi, M. *et al.* The Versatility of Serine Proteases from Brazilian Bothrops
Venom: Their Roles in Snakebites and Drug Discovery. *Biomolecules* **15** (2025).
https://doi.org/10.3390/biom15020154
52 Vargas-Valerio, S. *et al.* Localization of Myotoxin I and Myotoxin II from the venom of
Bothrops asper in a murine model. *Toxicon* **197**, 48–54 (2021).
https://doi.org/10.1016/j.toxicon.2021.04.006
53 Chong, D. J. W. *et al.* Discovery and Development of DC-174 as a Novel Oral
Snakebite Treatment. *bioRxiv* (2025). https://doi.org/10.1101/2025.05.23.655830
54 World Health Organisation, W. *Snakebite information and data platform*,
<https://www.who.int/teams/control-of-neglected-tropical-diseases/snakebite-
envenoming/snakebite-information-and-data-platform> (2023).
55 Rapon, C. *et al.* Thromboinflammatory complications of Bothrops snakebite
envenoming: the case of B. lanceolatus endemic to the Caribbean Island of
Martinique. *Frontiers in Immunology* **16** (2025).
https://doi.org/10.3389/fimmu.2025.1625165
56 Albulescu, L. O. *et al.* Preclinical validation of a repurposed metal chelator as an
early-intervention therapeutic for hemotoxic snakebite. *Sci Transl Med* **12** (2020).
https://doi.org/10.1126/scitranslmed.aay8314
57 Roszko, P. J., Kavanaugh, M. J., Boese, M. L., Longwell, J. J. & Earley, A. S.
Rotational thromboelastometry (ROTEM) guided treatment of an Afghanistan viper
envenomation at a NATO military hospital. *Clin Toxicol (Phila)* **55**, 229–230 (2017).
https://doi.org/10.1080/15563650.2016.1263857
58 Dang, X. T., Nguyen, T. X., Nguyen, H. T. T. & Ha, H. T. Correlations between
rotational thromboelastometry (ROTEM) and standard coagulation tests following
viper snakebites. *J Int Med Res* **50**, 3000605211067321 (2022).
https://doi.org/10.1177/03000605211067321
59 Bernardoni, J. L. *et al.* Functional variability of snake venom metalloproteinases:
adaptive advantages in targeting different prey and implications for human
envenomation. *PLoS One* **9**, e109651 (2014).
https://doi.org/10.1371/journal.pone.0109651
60 Gorlinger, K., Bhardwaj, V. & Kapoor, P. M. Simulation in coagulation testing using
rotational thromboelastometry: A fast emerging, reliable point of care technique. *Ann*
Card Anaesth **19**, 516–520 (2016). https://doi.org/10.4103/0971-9784.185546

61   Sells, P. G., Ioannou, P. & Theakston, R. D. A humane alternative to the measurement of the lethal effects (LD50) of non-neurotoxic venoms using hens' eggs. *Toxicon* **36**, 985–991 (1998). https://doi.org/10.1016/s0041-0101(98)00004-x

62   Kazandjian, T. D. *et al.* Convergent evolution of pain-inducing defensive venom components in spitting cobras. *Science* **371**, 386–390 (2021). https://doi.org/10.1126/science.abb9303

63   Verity, E. E., Stewart, K., Vandenberg, K., Ong, C. & Rockman, S. Potency Testing of Venoms and Antivenoms in Embryonated Eggs: An Ethical Alternative to Animal Testing. *Toxins (Basel)* **13** (2021). https://doi.org/10.3390/toxins13040233

973   **Figures**

974
Figure 1. Variation in the distribution and venom composition of *Bothrops* species. A)
Geographical distribution of the *Bothrops* species relevant to this manuscript. The distribution is
based on the ICUN Red List accessed in April 2025 presented using QGIS 3.4. The images of *B.*
asper, *B. atrox*, *B. jararaca, B. moojeni* and *B. neuwiedi* are held under all rights reserved copyright
and have been published with permission by Wolfgang Wüster, while the image of *B. lanceolatus* is
published with permission by Jonathan Florentin. The image of *B. alternatus* was accessed via
Wikimedia commons and is held respectively under Creative Commons Attribution-Share Alike 2.0
Generic licence by Cláudio Timm. B) A summary of the published percentage venom compositions for
the three key toxins of relevance to this manuscript; snake venom metalloproteinases (SVMP) in
beige, snake venom serine proteases (SVSP) in orange, and snake venom phospholipase A2 (PLA$_2$)
in green. The extracted data (Supplementary Table 1) vary from an individual to large pools of
specimens including captive and wild caught. The data is summarised from the following number of
publications per species; *B. alternatus* (n=2), *B. asper* (n=1), *B. atrox* (n=6), *B. jararaca* (n=3), *B.*
lanceolatus (n=2)*, B. moojeni* (n=2) *and B. neuwiedi* (n=1). The data is displayed as a box and violin
plot (interleaved low-high with line at the mean) using Prism software version 10 (GraphPad).

990
991
Figure 2. *In vitro* profiling of venom samples from seven different *Bothrops* species. A) Protein
profiles of the venom protein components per species. Whole venom (5 µg per lane) was prepared
under denaturing and reducing conditions, then separated by SDS-PAGE using a 4-20% gel. The gel
was then stained with Coomassie blue and destained to show all proteinaceous components. M =
molecular weight marker, and approximate masses of the markers are given in kDa. B-D) enzymatic
activity of the seven *Bothrops* species in toxin specific assays. The results were analysed using area
under the curve (AUC) in seconds and plotted using Prism version 10 software (GraphPad). A) Snake
venom metalloproteinase (SVMP) activity using fluorogenic substrate ES010 read at an excitation
wavelength of 320 nm and an emission wavelength of 405 nm over 108 minutes (n=3, ±SE), with 1 µg
of venom per reaction. C) phospholipase A_2 (PLA$_2$) activity using commercially available secretory
PLA$_2$ kit (Abcam) read at an absorbance wavelength of 405 nm over 12 minutes (n= ≥2, ±SE), with
100 ng of venom per reaction. D) snake venom serine protease (SVSP) activity using commercially
available chromogenic substrate S-2288 read at an absorbance of 405 nm over 35 minutes (n=4 ±SE)
with 1 µg of venom per reaction.

1006

1007
1008
**Figure 3. *In vitro* coagulation profile of seven different *Bothrops* species and inhibition of
procoagulant activity by small molecule drugs.** A) The coagulopathic profile of the seven *Bothrops*
species at a comparable 100 ng dose in bovine plasma over 25 minutes at an absorbance of 595 nm
(n=2, ±SD). B) Coagulation profile in the absence of venom for SVMP inhibitors (DMPS, a metal
chelator, and the MMP inhibitors marimastat and prinomastat), the PLA$_2$ inhibitor varespladib and the
serine protease inhibitor nafamostat. The profiles indicate no direct coagulopathy for any of the
inhibitors except for nafamostat which has a strong anticoagulant profile at this dose (n=2, ±SD). C-I)
The percentage inhibition in the coagulation assay of the small molecules (excluding nafamostat due
to the inherent anticoagulant activity) in each of the seven *Bothrops* venoms, tested at 5 µM (n=2,
±SE). The adjusted dose for each venom indicated in the graph title was selected to provide
comparable profiles for all seven venoms. J) Dose response testing was performed for the four
inhibitors in the coagulation assays and EC$_{50}$s (nM) calculated using Prism v10 software (GraphPad)
as presented this table (n=2, ±95% CI).

1022

1023

A)

Venom	SVMPi EC$_{50}$ (nM)									PLA$_2$i EC$_{50}$ (nM)
	DMPS	dimercaprol	marimastat	prinomastat	batimastat	tanomastat	doxycycline	XL-784	CTS-1027	varespladib
B. alternatus	216.1 (154.7 - 291.6)	228.4 (163.4 - 310.6)	1.8 (1.5 - 2.1)	2.0 (1.7 - 2.3)	1.5 (1.3- 1.8)	5126.0 (2872.0 - 16814.0)	> 10000 (N/A)	6.3 (4.6 - 8.5)	8.5 (7.1 - 10.2)	1.6 (1.2 - 2.3)
B. asper	2019.0 (849.4 - 8877)	3561.0 (1050 - 59680)	2.8 (2.3 - 3.5)	0.7 (0.6 - 0.8)	3.9 (2.6 - 5.6)	1687.0 (746.0 - 5836.0)	> 10000 (N/A)	13.3 (9.0 - 19.2)	1.5 (1.2 - 1.9)	0.2 (0.2 - 0.2)
B. atrox	2753.0 (1574 - 4490)	4490.0 (2601 - 6189)	8.2 (6.7 - 10.1)	2.4 (2.0 - 2.8)	9.8 (7.9 - 12.0)	3352.0 (2184.0 - 5788.0)	> 10000 (N/A)	36.3 (26.6 - 49.1)	8.5 (6.6 - 10.9)	0.2 (0.1 - 0.3)
B. jararaca	387.8 (321.6 - 472.3)	432.3 (341.5 - 554.4)	7.7 (6.6 - 8.8)	7.5 (7.2 - 7.9)	5.0 (4.7 - 5.4)	> 10000 (N/A)	> 10000 (N/A)	41.1 (35.9 - 47.1)	61.6 (54.7 - 69.3)	0.2 (0.1 - 0.4)
B. lanceolatus	266.5 (245.2 - 289.2)	300.4 (268.3 - 336.9)	10.6 (9.4 - 11.9)	8.6 (8.1 - 9.1)	6.2 (5.8 - 6.7)	> 10000 (N/A)	> 10000 (N/A)	34.0 (30.2 - 38.3)	15.4 (13.7 - 17.2)	0.2 (0.2 - 0.3)
B. moojeni	579.6 (358.3 - 891.0)	657.7 (411.8 - 1024)	3.2 (2.4 - 4.1)	3.7 (2.9 - 4.7)	6.6 (4.9 - 8.7)	1109.0 (633.7 - 2099.0)	> 10000 (N/A)	10.5 (7.3 - 14.8)	3.5 (2.6 - 4.7)	0.2 (0.1 - 0.4)
B. neuwiedi	196.1 (103.4 - 347.4)	255.2 (153.9 - 405.6)	2.8 (2.3 - 3.3)	3.3 (2.4 - 4.5)	2.8 (2.2 - 3.5)	> 10000 (N/A)	> 10000 (N/A)	8.8 (6.1 - 12.5)	3.9 (2.9 - 5.2)	0.3 (0.2 - 0.5)

1024
1025

Figure 4. *In vitro* inhibition of seven different *Bothrops* species by small molecule drugs. The enzymatic activity of the seven *Bothrops* species with relevant inhibitors in the three toxin specific assays are presented, with the results analysed using area under the curve (AUC) and plotted using Prism version 10 software (GraphPad). A) Dose response testing was performed in the SVMP and PLA$_2$ assays using varespladib for the latter, but a wider panel of MMPis and metal chelators with prior evidence of SVMP inhibition in the SVMP assay. This allowed for the calculation of EC$_{50}$s (with 95% CI shown) using Prism v10 software (GraphPad) which are displayed in this table (n=2). B) Snake venom metalloproteinase (SVMP) activity and inhibition by the MMP inhibitor, marimastat (AUC over 178 minutes, n=4, ±SE). C) phospholipase A$_2$ (PLA$_2$) activity and inhibition by the PLA$_2$ inhibitor, varespladib (AUC over 13 minutes, n= ≥3, ±SE). D) snake venom serine protease (SVSP) activity and inhibition by the SP inhibitor, nafamostat (AUC over 35 minutes, n=4, ±SE).

1037
1038
1039
1040

1041

1042
**Figure 5. *In vitro* coagulation profile of two coagulopathic *Bothrops* species and inhibition by
combination therapy of two small molecule drugs.** The effect of *Bothrops jararaca* (A-B, purple)
and *Bothrops atrox* (C-D, green) with and without small molecule inhibitors is presented alongside the
direct effect of these inhibitors (E and F) in the coagulation assay. (A and C) The kinetic curves
demonstrate the procoagulant effect of both venoms, compared to normal clotting control (black
circles). The SVSP inhibitor nafamostat (orange triangles) fails to rescue the procoagulant effect,
whilst marimastat (5 μM, white triangles) fully inhibits the procoagulant effect of *B. atrox* and partial
rescue of *B. jararaca* (n=6, ± SD). (B and D) Percentage inhibition of coagulopathy by *Bothrops*
species in single treatment by small molecule inhibitors (n=6, ± SE) matched that of the kinetic curves
for the top 5 μM dose of marimastat with subsequent lower doses slightly reducing the percentage
inhibition, nafamostat resulted in minor inhibition at any doses. Combination testing of marimastat and
nafamostat, for both venoms appears to reflect an additive effect, however in the no-venom control
(E, kinetic curve; F, % inhibition), nafamostat shows a dose-dependent increase beyond 100% in the
absence of venom, demonstrating that it is directly anticoagulant at higher doses. Marimastat has no
effect on plasma clotting, indicating that the effects seen in graphs (A to D) are through inhibition of
venom.
1059

1060

Treatment	CT	MCF	CFT	Alpha angle	A5
No venom (normal clotting)	664.3 (±49)	58.0 (±4)	207.7 (±41)	52.7 (±5)	27.7 (±4)
marimastat only 5 µM	682.7 (±28)	59.3 (±3)	187.3 (±67)	56.0 (±10)	30.3 (±8)
B. atrox only	133.3 (±15)	68.3 (±1)	67.3 (±8)	76.3 (±1)	51.3 (±2)
B. atrox + marimastat 5 µM	490.3 (±15)	65.3 (±1)	163.0 (±9)	62.7 (±1)	40.3 (±2)
B. jararaca only	184.3 (±21)	66.7 (±2)	71.0 (±11)	75.3 (±2)	48.3 (±3)
B. jararaca + marimastat	479.7 (±11)	60.3 (±2)	133.0 (±10)	64.3 (±2)	37.7 (±3)

1061
1062
**Figure 6. Thromboelastography (TEG) profiles of two representative *Bothrops* species and
inhibition by the SVMP inhibitor marimastat.** A and B) Thromboelastography profile of respectively
B. atrox and *B. jararaca* using an XY plot of clotting time vs clotting strength. Both venoms (0.6 µg per
300 µL reaction) have strong procoagulant activity (circles) compared to the no venom controls (black
diamond) or drug only control (grey diamond). Marimastat inhibits the procoagulant activity of both
venoms (squares) (n >3). C) The parameters reported by the ROTEM instrument are displayed
including the clotting time (CT; seconds), maximum clot firmness (MCF; mm), clot formation time
(CFT; seconds), alpha angle (º) and amplitude at 5 minutes (A5; mm), each is the mean of the
triplicate results ± the related range.

1072
Figure 7. Survival curves of chicken embryos dosed with *B. atrox* venom with and without
representative small molecule drugs. Groups of 6-day post-fertilisation chicken egg embryos were
dosed on to the vitelline vein with either PBS (negative control) or *B. atrox* venom (20 µg) with or
without a subsequent dose of 0.5 µg, 1 µg and 5 µg of A) marimastat, B) DMPS and C) nafamostat.
All treatment group sizes were n=5, the PBS control contained an n=12, while the venom only control
contained an n=20. D) Representative pathology images of chicken embryos 1 hour post-dosing with
B. atrox venom with and without 5 µg of the inhibitory drugs, showing varying degrees of protection
against the vascular damage shown in the venom only control (see also Supplementary Figure 1).

1081

1082

1083