Systemic dissemination, reproductive tissue tropism, and gestational stage-dependent placental responses during Mayaro virus infection in a murine model

Arévalo A.P.[1], Perbolianachis P.[2,3], Pórfido J.L.[1], Pereira-Gómez M.[2,4], Greif G.[5], Hurtado J.[2,3], Fajardo A.[2,3], Varela B.[6], Verdes J.M.[6], Moratorio G.[2,3*], Crispo M[1*].

[1]Laboratory Animal Biotechnology Unit, Institut Pasteur de Montevideo, Uruguay

[2]Experimental Virus Evolution Laboratory, Institut Pasteur de Montevideo, Uruguay

[3]Laboratory of Molecular Virology, Nuclear Research Center, Faculty of Sciences, Udelar

[4]Microbiology Unit, Department of Pathobiology, Faculty of Veterinary Medicine, Udelar

[5]Host-Pathogen Interaction Laboratory, Institut Pasteur de Montevideo, Uruguay

[6]Pathology Unit, Department of Pathobiology, Faculty of Veterinary Medicine, Udelar

* Corresponding authors: crispo@pasteur.edu.uy; moratorio@pasteur.edu.uy

ABSTRACT

Mayaro virus (MAYV) is an emerging mosquito-borne alphavirus associated with acute febrile illness and persistent arthralgia, with increasing reports in Latin America and potential for geographic expansion. However, its dissemination dynamics, tissue tropism, and transmission routes remain incompletely defined. Here, we developed a murine model to characterize systemic viral dissemination, tissue tropism, and the impact of gestational stage on maternal–fetal infection. A nanoluciferase-expressing MAYV reporter enabled non-invasive *in vivo* imaging, revealing rapid and widespread systemic dissemination under controlled conditions. Complementary infection with wild-type (WT) virus confirmed broad tropism across lymphoid, musculoskeletal, and reproductive tissues. At 24 hours post-infection (hpi), viral distribution was relatively homogeneous, whereas at 48 hpi, tissue-specific accumulation was evident, including increased viral loads in male reproductive tissues. Hematological and biochemical analyses indicated early systemic alterations consistent with inflammatory and immune cell redistribution. Infectious viral particles were detected in reproductive tissues of both sexes, supporting the plausibility of non-vector transmission. Gestational stage significantly influenced infection outcomes: early gestation (infected at 7.5–8.5 dpc; analyzed at 9.5–10.5 dpc) was associated with placental permissiveness and detection of infective virus in fetal tissues, whereas mid-gestation (infected at 13.5–14.5 dpc; analyzed at 15.5–16.5 dpc) showed no detectable infectious virus in fetuses despite sustained viral presence in maternal and placental compartments. Together, these findings establish a robust experimental framework for investigating MAYV pathogenesis and identify tissue tropism, sex-related differences, and gestational stage as key determinants of viral dissemination and maternal–fetal infection.

KEYWORDS: Mayaro virus; viral pathogenesis; Emerging arbovirus; Vertical transmission; Systemic dissemination; Placental infection; Murine model; *in vivo* imaging

AUTHOR SUMMARY

Mayaro virus is an emerging mosquito-borne virus that causes fever and long-lasting joint pain, with increasing reports in Latin America and potential for wider geographic spread. Despite its growing importance, several key aspects of MAYV pathogenesis remain to be fully elucidated, including the mechanisms driving systemic dissemination and tissue tropism, as well as its potential to impact pregnancy or be transmitted through non-vector routes. We used a mouse model with a luminescent reporter virus to track Mayaro virus dissemination in vivo, revealing rapid and widespread spread across multiple tissues. In studies using wild-type virus, detection occurred not only in muscle and immune-related organs but also in reproductive tissues of both males and females. Over time, viral distribution shifted from an initially uniform pattern to a more tissue-specific accumulation, including increased viral load in reproductive tissues. Importantly, gestational stage influenced viral distribution. When infection occurred early in gestation, infectious viral particles were detected in fetal tissues. In contrast, at later stages of pregnancy, infectious virus was no longer detectable in fetuses, even though it remained present in maternal and placental tissues. Altogether, these findings improve our understanding of Mayaro virus infection and highlight gestational timing as a key factor determining the risk of fetal exposure. They also support the potential for alternative transmission routes and provide a framework for future studies on this emerging virus.

INTRODUCTION

Mayaro virus (MAYV), an emerging arbovirus of the *Alphavirus* genus and *Togaviridae* family, was first identified in Trinidad and Tobago in 1954 and has since been responsible for sporadic outbreaks across Central and South America. Imported cases have also been reported in Europe and North America, raising concerns at the international public health level as reflected in the epidemiological alert issued by the Pan American Health Organization / World Health Organization in May 2019, which emphasized the need to strengthen epidemiological surveillance and laboratory diagnostic capacity (1). The virus has already been detected in Brazil, Peru, Venezuela, Suriname, Colombia, Ecuador, Bolivia, and Panama, with serological evidence also reported in Costa Rica, Guatemala, Mexico, and Haiti (2–5). Multiple anthropogenic factors—including climate change, urbanization, human mobility, and the expanding distribution of mosquito vectors such as *Aedes aegypti* and *Aedes albopictus*—contribute to the increasing risk of MAYV geographic expansion, even in countries without active transmission, such as Uruguay (6,7). A recent study estimated that 58.9 million people in Latin America reside in ecologically suitable areas for MAYV transmission (6–8). From a genomic perspective, MAYV possesses a positive-sense single-stranded RNA genome of approximately 11.5 kb, which encodes both non-structural (nsP1–nsP4) and structural proteins (CP, E3, E2, TF, 6k, and E1) through two open reading frames (ORFs). The virion is enveloped in a lipid membrane containing E1 and E2 glycoproteins and exhibits high genetic plasticity, which facilitates its adaptation to a broad range of hosts. The viral replication cycle begins with the fusion of the viral envelope to the host cell membrane, followed by translation of viral proteins and the assembly of new virions. MAYV is classified into three genotypes: D (dispersed), L (limited), and N (new), the latter identified in Peru (9–14). In addition to its sylvatic cycle, MAYV is proposed to establish urban transmission

through vectors such as *Aedes spp.*, *Culex spp.*, and *Sabethes spp.*, with potential vertebrate reservoirs including marsupials, rodents, birds and alligators (15–17).

Clinically, Mayaro disease is generally a self-limiting illness characterized by fever, rash, headache, photophobia, myalgia and persistent polyarthralgia, which may last for months or even years, significantly impacting patients' quality of life. Hemorrhagic and neurological complications, as well as myocarditis, have also been reported (18). The clinical similarity to other arboviral diseases, such as Dengue and Chikungunya, combined with limited diagnostic capacity in endemic regions, hinders accurate differential diagnosis and contributes to underestimation of the actual disease burden (18,19). Beyond its typical acute presentation, increasing evidence from other arboviral infections, including Zika and Chikungunya viruses (ZIKV; CHIKV), indicates that clinical outcomes and transmission dynamics may extend beyond the classical vector-borne paradigm, raising questions about the potential involvement of vulnerable populations such as pregnant women and neonates (20–23). Although no specific antiviral treatments or licensed vaccines are currently available, several vaccine candidates are under development, with sustained epidemiological surveillance remaining essential to anticipate and mitigate future outbreaks (3,9,24).

Animal models, particularly murine species, have been instrumental in elucidating the pathogenesis of MAYV and in the preclinical evaluation of therapeutic and vaccine candidates. These models faithfully reproduce hallmark clinical manifestations—including ruffled fur, altered gait, and ocular irritation—as well as key histopathological features such as arthritis, myositis and tenosynovitis (25,26). Disease outcomes vary depending on host age and immune status (25,27–29). While the musculoskeletal system represents the site of the most evident clinical damage, the liver and spleen act as major systemic sites of viral replication, functioning as central hubs for Mayaro virus

amplification and dissemination (10,15,28,30). The development of experimental

platforms that enable robust and reproducible antiviral testing is essential for advancing

MAYV research. In this context, the integration of *in vivo* imaging technologies

constitutes a major methodological advance, as it allows real-time, non-invasive

monitoring of viral dissemination dynamics using tagged viruses (27,31) .

Here, we establish and refine a murine model as a preclinical platform for the future

evaluation of antiviral drugs and therapeutic interventions against MAYV. To validate

early systemic dissemination under controlled conditions, we employed a nanoluciferase-

expressing MAYV reporter for longitudinal *in vivo* imaging. Complementary

experiments using wild-type MAYV were performed to characterize tissue tropism, viral

load, hematological and biochemical alterations, and placental transcriptional responses

following controlled infection. We further assessed gestational stage–dependent placental

permissiveness and explored the biological plausibility of vertical and sexual

transmission. We hypothesize that gestational timing modulates placental susceptibility

to MAYV, defining critical windows of maternal–fetal vulnerability. Together, this

integrated framework enables mechanistic investigation of systemic dissemination,

reproductive involvement, and maternal–fetal risk during MAYV infection.

MATERIALS AND METHODS

Ethics statement

All procedures and experiments were conducted in accordance with the Ethics Committee

on Animal Use of the Institut Pasteur de Montevideo (CEUAIP #010-22, in compliance

with National Law 18.611 "Procedures for the Use of Animals in Experimentation,

Teaching, and Scientific Research Activities" and strictly followed ARRIVE guidelines.

MAYV infective clones

This study employed wild-type MAYV and nano Luc-expressing MAYV (MAYV-nLuc)

infectious clones based on the TRVL 4675 strain, a genotype D virus originally isolated

from a human case in Trinidad in the 1950s and kindly provided by Dr. James Weger-

Lucarelli (31–33).

Cell culture

Vero cells were maintained in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% fetal bovine serum (FBS), penicillin (100 U/mL), and

streptomycin (100 μg/mL) and maintained at 37 °C, with 5% CO_2.

Virus generation

Infectious virus was rescued from both WT and MAYV-nLuc infectious clones by

transfection of Vero cells using Lipofectamine™ 3000 (Invitrogen), according to the

manufacturer's instructions. Viral supernatants were collected when transfected cells

exhibited ≈80% cytopathic effect (CPE) and stored at -80°C as passage P0.

Working viral stocks were generated by sequential infection of Vero cells to produce P1

and P2 passages. The P2 viral stocks were titrated by plaque assay, verified by sequencing

and stored at -80°C until further use.

Mice

BALB/cJ mice (#000651) were bred and maintained at the Laboratory Animal

Biotechnology Unit's animal facility of Institut Pasteur de Montevideo. Mice were housed

in IVC Isolator Biocontainment unit (ISOCAGE-N, Tecniplast, Milan, Italy) and

acclimatized for one week prior to the experimental procedures. The cages were prepared

with wood chip bedding (Toplit 6, SAFE, France), plastic tunnels as enrichment and paper

napkins as nesting material. Environmental conditions were maintained at $20 \pm 1°C$, with

30–70% relative humidity, and a 14/10-hour light/dark cycle. The mice were provided

with *ad libitum* autoclaved standard diet (D113, SAFE) and filtered, autoclaved water.

Mice were specific pathogen-free (SPF) certified by IDEXX (USA) quarterly, in

accordance with FELASA standards.

Animal experimental design

Intraperitoneal (i.p.) inoculation was selected to enable controlled and synchronized

assessment of early systemic dissemination of Mayaro virus under standardized

experimental conditions. Although this route does not reflect natural mosquito-borne

transmission, it enables precise dose administration and reproducible systemic exposure,

thereby facilitating analysis of early viral dissemination dynamics. A nanoluciferase-

expressing MAYV reporter (MAYV-nLuc) was employed for non-invasive *in vivo*

imaging. This reporter-based approach was used specifically to define spatial and

temporal dissemination patterns. Complementary experiments using wild-type MAYV

were subsequently performed to quantify infectious viral loads, characterize

hematological and biochemical alterations, and assess placental transcriptional responses

following controlled infection. Viral doses were optimized according to the specific

objectives of reporter-based imaging and pathogenesis studies. Sex-specific analyses

were incorporated where biologically relevant, including assessment of reproductive

tissue tropism and transmission experiments. This integrated experimental framework

enabled systematic evaluation of viral dissemination, reproductive involvement, and

gestational stage–dependent placental permissiveness (Fig 1).

Figure 1. Experimental design for longitudinal assessment of early systemic dissemination of Mayaro virus in mice. Mice were i.p. inoculated with a bioluminescent Mayaro virus to enable longitudinal, non-invasive monitoring of early viral dissemination by *in vivo* imaging. Parallel groups were infected with wild-type Mayaro virus to evaluate tissue tropism and transmission-related outcomes. Blood and lymphoid, visceral, musculoskeletal, and reproductive tissues were collected for downstream analysis.

- *In vivo* imaging evaluation with MAYV-nLuc infection

In vivo bioluminescence imaging frequency was performed over a 10-day observation window to capture the peak of viral replication and the onset of immune responses associated with recovery from acute infection, based on previous reports (25,26,30). Age groups were selected according to their susceptibility to MAYV infection, with younger animals exhibiting higher susceptibility (34). For this study, four- and eight-week-old BALB/cJ female mice were i.p. infected with 1 x 10^4 PFU of MAYV-nLuc resuspended in 100 µL PBS (n=4 per age group). Mock-infected animals received PBS only (n = 2; one animal per age group). The viral dose was selected based on pilot optimization experiments to ensure robust reporter detection while maintaining animal welfare. To perform *in vivo* luminescence imaging, the animals were i.p. injected with 20 µg of Furimazine (TargetMol®, MA, USA) in 100 µL of PBS, anesthetized with isoflurane (3–4% for induction and 1-2 % for maintenance) and imaged at defined time points using an In-Vivo Xtreme II™ system (Bruker Daltonik GmbH, Billerica, MA, USA). Image

acquisition and analyses were performed using MolecularImaging software V.7.5.3.22464 (Bruker Daltonik Gmbh, Billerica, MA, USA). Thresholds and exposure times were kept constant across time points for longitudinal comparability. Bioluminescent signal reflects reporter expression and may not directly correlate with infectious viral titers. Imaging was performed at 1-, 2-, 6- and 10- day post infection (dpi). Animals were necropsied at 6 dpi (n=2 per age group) and at 10 dpi (remaining animals). Organs collected for confirmation of infectious viral particles included the liver, kidneys, spleen, and *quadriceps femoris* muscle. Peripheral tissues exhibiting luminescent signal—tail, ear, forelimb or hindlimb—were also sampled.

- **Viral tropism of Mayaro WT and health status**

Twelve BALB/cJ female and twelve BALB/cJ male mice (4 weeks old) were i.p. infected with 1×10^5 PFU of MAYV WT resuspended in 100 μL of PBS (n=24). Mock-infected mice received the PBS diluent alone (n=4). The viral dose was selected based on prior pilot experiments as the lowest dose that consistently allowed detection of infectious viral particles across tissues during acute infection. Blood samples were collected pre and post-infection for hematological and biochemical parameter analyses at 0-, 24- and 48-hour post infection (hpi). At 24 hpi, six animals of each sex were euthanized under deep anesthesia induced by ketamine-xylazine mixture administered via the i.p. route (ketamine 110 mg/kg; Pharmaservice, Ripoll Vet, Montevideo, Uruguay; xylazine 13 mg/kg; Seton 2%; Calier, Montevideo, Uruguay), followed by cervical dislocation. Organs collected for quantification of infectious viral particles included the liver, kidneys, spleen, *quadriceps femoris* muscle, ovaries, uterus, seminal vesicles and testes. The same procedures were performed on the remaining animals at 48 hpi.

- **MAYV WT vertical transmission**

Seventeen BALB/cJ female mice (8-weeks old) were synchronized in their estrous cycle

by exposing them to male dirty bedding for 72 h prior to mating (35). Mating was then

performed by placing one proven male in a cage with two females for 48 h, and the

presence of a vaginal plug was checked daily from the following morning as a positive

sign of mating to be considered day 0.5 of pregnancy. At 7.5-8.5-day post-coitum (dpc),

six females were i.p. infected with 1×10^5 PFU of MAYV WT resuspended in 100 µL of

PBS, while two mock-infected received only PBS diluent (n = 6 infected; n = 2 mock).

The females were euthanized and necropsied at 48 hpi (9.5-10.5 dpc). Blood samples

were collected prior to euthanasia for post-infection hematological and biochemical

analyses. Organs collected for quantification of infectious viral particles and histological

analysis included the liver, kidneys, *quadriceps femoris* muscle, ovaries, uterus, placenta

and fetus. The same procedures were performed on the remaining pregnant females

infected at 13.5–14.5 dpc (n = 7 infected; n = 2 mock), which were necropsied at 48 hpi

(15.5-16.5 dpc).

- **MAYV WT sexual transmission**

To evaluate sexual transmission in both directions, BALB/cJ mice were mated in a 1:1

ratio under controlled conditions. For female-to-male transmission, seven 8-week-old

females were synchronized in their estrous cycles by exposure to male dirty bedding for

72 h prior to mating, as described above, and were infected i.p. with 1×10^5 PFU of

MAYV in 100 µL of PBS 12 h before pairing with uninfected males. For male-to-female

transmission, six synchronized 8-week-old females were paired with males infected i.p.

with the same viral dose and schedule. Infected animals were necropsied at 48 hpi,

whereas their uninfected partners at 48 h after mating. The liver, spleen, *quadriceps*

femoris muscle, ovaries, uterus, seminal vesicles, and testes were collected for infectious

viral particle detection by plaque assay and for RT-qPCR analysis in organs from exposed

animals. Sperm from the bilateral cauda epididymides was collected for plaque assay and

RT-qPCR analysis. For analytical purposes, an exposed animal was defined as an

uninfected individual mated with an infected partner.

Hematology and biochemistry profile

For hematological analysis, 20 µL of blood per animal were collected and stored in 0.5

mL microtubes containing potassium EDTA salts (W anticoagulant, Wiener Lab,

Rosario, Argentina) at a 1:10 ratio (EDTA: blood). Samples were analyzed using an Auto

Hematology Analyzer BC-5000 Vet (Mindray, Shenzhen, China). The parameters

evaluated included white blood cells (WBC), neutrophils (Neu), lymphocytes (Lym),

monocytes (Mon), eosinophils (Eos), basophils (Bas), red blood cells (RBC), hemoglobin

(HGB), hematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular

hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) and platelets

(PLT).

For biochemical blood profile analysis, 100 µL of whole blood per animal was collected

and mixed with 0.5 µL of sodium heparin (Heparin Na, 20 U/mL). Using the Chemistry

Analyzer Pointcare® V3 (Tianjin MNCHIP Technologies Co. Ltd, Tianjin, China), the

following parameters were evaluated: total protein (TP), albumin (ALB), globulin (GLO),

total bilirubin (Tbil), alanine aminotransferase (ALT), aspartate aminotransferase (AST),

gamma-glutamyl transferase (GGT), blood urea nitrogen (BUN), and creatinine (CRE).

Tissue collection and processing

Samples collected from each organ of interest were weighed, rapidly frozen in liquid

nitrogen under a biosafety cabinet and stored at -70°C until processing. Samples were

homogenized using zirconium beads (0.5 mm/tube) and 800 µL of sterile 1× DPBS in a

Precellys® 24 homogenizer (Bertin Technologies, France), (2 min, power 10).

Homogenates were clarified by centrifugation (6,000 × g, 5 min), and supernatants were

stored at −70 °C for viral titration or RT-qPCR analysis.

Viral titration

For viral titration, confluent Vero cell monolayers in 12-well plates (2.6×10^6 cells per

plate) were inoculated with tissue homogenate–derived supernatants and their 10-fold

serial dilutions (200 µL per well) and incubated at 37 °C with 5% CO_2 for 1 h, with gentle

rocking every 15 min to ensure uniform inoculum distribution. Mock-infected wells

served as negative controls. After adsorption, cells were overlaid with DMEM containing

0.8% agarose, 1.6% FBS, penicillin, and streptomycin and incubated for 48 h.

Monolayers were then fixed with 4% formaldehyde, stained with crystal violet, and

plaques were counted to determine viral titers, expressed as plaque-forming units per

gram of tissue (PFU/g).

Mouse sperm collection

The bilateral cauda epididymides were collected in a 60 mm plastic Petri dish and incised

with several cuts using fine scissors, releasing the sperm into a 50 µL drop of Dulbecco's

Phosphate Buffered Saline 1x (DPBS, Sigma-Aldrich), and stored at −80°C until further

processing. Later, sperm samples were thawed and resuspended in 150µL of PBS.

Zirconium beads were added to each sample and disruption was performed using a Bullet

Blender® STORM 24 (Next Advance Inc., New York, USA) for one minute at speed 8.

After centrifugation, 100 µL of supernatant was divided into two aliquots. To assess the

presence of MAYV, one aliquot was subjected to cell culture amplification followed by

plaque assay, whereas the other was analyzed directly for viral RNA by RT-qPCR.

Viral RNA extraction and detection

Viral RNA from tissues was extracted from clarified tissue homogenate supernatants, as

described above, using the QIAamp® Viral RNA Mini Kit (QIAGEN), according to the

manufacturer's instructions. RNA from sperm samples was extracted from TRIzol-treated

homogenates using the Direct-zol RNA Miniprep Kit (Zymo Research), following the

manufacturer's guidelines. RNA samples were eluted in 25µL of nuclease-free water and

stored at -80°C until further use (36). Detection was performed using a one-step real-time

reverse transcription PCR (RT-qPCR) assay, employing the primers and probe listed in

Table 1, which target an 86-bp region of the MAYV nsP1 gene. Reactions were carried

out using TaqMan™ Fast Virus 1-Step Master Mix (Thermo Fisher Scientific) on a

QuantStudio™ 7 Pro Real-Time PCR System (Applied Biosystems). Due to limited

sample availability, sperm-derived RNA samples were analyzed in technical duplicates.

The assay was used for qualitative detection (presence/absence) of viral RNA.

Table 1: primers and probe used for MAYV detection by qPCR.

MAY-qPCR-Fw	5'-AGGACCCAGAGGAACACTAATA-3'
MAY-qPCR-Rv	5'-CGATACTCTTTCGCCCACTTAC-3'
MAY-qPCR-Probe	5'-/56-FAM/TACCTGCTT/ZEN/CCAGTGGTGTCACAA/3IABkFQ/-3'

Cell culture–based isolation of virus from sperm samples

Due to the limited volume of sperm samples, processed samples were subjected to

passage in Vero cells to allow potential biological amplification prior to titration. Vero

cells were seeded in 96-well plates at 5,000 cells per well in 100 µL of DMEM

supplemented with 5% FBS, penicillin (100 U/mL), and streptomycin (100 µg/mL). The

following day, cells were inoculated with 100 µL of processed sperm samples. Mock-

infected wells receiving culture DMEM alone served as negative controls. After 72 h of

incubation at 37 ºC, supernatants were harvested and the presence of infectious viral

particles was assessed by plaque assay. Cytopathic effects were monitored using inverted

optic microscope Zeiss Primovert, Carl Zeiss Suzhou Co., LTD, China.

Transcriptomics in placenta

For the RNA-seq experiment, three placentas per condition (control uninfected; infected

at 7.5-8.5 dpc and infected at 13.5-14.5 dpc) were selected and processed for total RNA

extraction. Placentas stored at −80 °C were disrupted in TRI Reagent using a Bullet

Blender® STORM 24 (Next Advance Inc., New York, USA) with 0.5 mm zirconium

oxide beads (ZROB05, Next Advance Inc.) for 10 min at maximum speed. Following

homogenization, the samples were centrifuged to remove beads and debris. The upper

aqueous phase was collected and purified using the Direct-zol RNA kit (Zymo Research,

USA) according to the manufacturer's instructions. RNA was quantified with RNA HS

Qubit (Invitrogen, USA). RNA quality assessment, library construction and sequencing

were performed by Macrogen (Seoul, South Korea). Libraries were prepared using

TruSeq stranded total RNA with Ribo-Zero Gold kit (Illumina, USA). Sequencing was

performed on an Illumina NovaSeq X platform. Paired-end reads of 100 base pairs were

obtained.

Raw paired-end RNA-seq reads were aligned to the *Mus musculus* reference genome

(GRCm39, Ensembl release 114) using STAR v2.7.8a. Aligned reads were quantified at

the gene level using featureCounts v2.0.2 (Subread package). The Ensembl GTF

annotation file (Mus_musculus.GRCm39.114.gtf) was used.

Downstream analysis was conducted in R v4.3.0 using the DESeq2 package. The raw

count matrix was filtered to retain genes with a minimum of 10 total reads across all

samples. A DESeq2 dataset was constructed with experimental conditions as the main design factor. Variance stabilizing transformation (VST) was applied to normalize data prior to visualization and clustering. Differential expression analyses were performed for multiple contrasts of interest, and log2 fold changes were shrinkage-corrected using the "normal" method of lfcShrink. Genes with an adjusted p-value <0.05 were considered significantly differentially expressed.

Volcano plots were generated with ggplot2, where significant genes were highlighted (adjusted p <0.05). Heatmaps of differentially expressed genes were created with pheatmap R package, using variance-stabilized expression values scaled by gene (row-wise z- scores). Color schemes were defined with the RColorBrewer package. Additional hierarchical clustering of differentially expressed genes was performed to assess global expression patterns across experimental groups. Gene ontology analysis was performed using Metascape (37).

Immunohistochemistry

Immediately after the necropsy, placenta and fetus from mid-stage of pregnancy were fixed in 10% neutral buffered formalin (pH 7.4) for further processing. The samples were then decalcified by immersion and agitation in a 3% aqueous nitric acid solution for 24 h. Once decalcification was complete, histological processing for paraffin embedding began. Once the paraffin blocks were obtained from each sample, successive 4μm-thick sections were made with a manual rotary microtome (Slee, Germany), and the successive sections were mounted on slides and prepared for immunohistochemistry against MAYV immunoreaction, using a rabbit polyclonal primary antibody (dilution 1:100) according to a previous immunofluorescence protocol developed for immunocytochemistry in cell cultures (38,39). Histological and immunohistochemical blinded examinations were

370 performed using transmitted light microscope (Nikon Eclipse E200®, Nikon Corporation,

371 China) by two different veterinary pathologists.

372 Data Analysis

Statistical analyses were performed using GraphPad Prism version 10 (GraphPad

Software, San Diego, CA, USA). Viral load data (PFU/g) were $\log_{10}$-transformed to

approximate a normal distribution. Values below the limit of detection (LOD) were

imputed as LOD/2 (630 PFU/g) prior to transformation to enable statistical analysis

without introducing bias due to non-detectable values. For viral load analyses, mixed-

effects models (REML) were applied with time point, tissue, and/or experimental group

as fixed effects, and individual animal as a random effect to account for repeated measures

and intra-subject variability. Post hoc comparisons were performed using Tukey's

multiple comparisons test. Hematological and biochemical parameters were analyzed

using mixed-effects models for repeated measures when longitudinal data were available,

with time and/or condition as fixed effects and individual animal as a random effect.

When model assumptions were not met, non-parametric tests were applied, including

Mann–Whitney U test for two-group comparisons and Kruskal–Wallis test followed by

Dunn's multiple comparisons test for multiple groups. Normality of data distributions

was assessed using the Shapiro–Wilk test and visual inspection of residuals. RNA-seq

differential expression analysis was performed using DESeq2. Genes with fewer than 10

total counts across all samples were filtered out. Variance stabilizing transformation

(VST) was applied for data normalization prior to visualization and clustering.

Differentially expressed genes were identified using adjusted p-values (Benjamini–

Hochberg correction), with a significance threshold of adjusted $p < 0.05$. Data are

presented as individual values with median lines unless otherwise specified. All statistical

tests were two-tailed, and a p value < 0.05 was considered statistically significant.

RESULTS

Bioluminescence reveals rapid dissemination and longitudinal persistence of MAYV-nLuc signal. Luminescence was first detected at 24 hpi (1 dpi) and remained detectable through the predetermined endpoint at 10 dpi. Signals were identified in the forelimbs, hindlimbs, tail, ears, and abdominal region. The abdominal signal decreased after 6 dpi, whereas residual luminescence persisted in peripheral sites, remaining detectable mainly in the limbs at 10 dpi (Fig 2).

To assess the presence of infectious virus, plaque assays were performed on tissues collected at 6 and 10 dpi. Infectious viral particles were selectively recovered from the peripheral tissues of 4-week-old mice —including forelimbs, hindlimbs, tail and ears— while visceral organs (liver, spleen, muscle, and kidneys) remained negative. Viral loads in these peripheral compartments spanned nearly two orders of magnitude, ranging from 3.17 to 5.47 $\log_{10}$ PFU/g (1.49×10^3 to 2.98×10^5 PFU/g), with the highest titers detected in tail and limb samples. In contrast, tissues from 8-week-old animals that exhibited luminescent signals *in vivo* did not yield detectable infectious virus. This apparent discrepancy may be attributable to differences in analytical sensitivity between reporter-based imaging and plaque assays, the latter of which requires a minimum threshold of viable, replication-competent virions.

Figure 2. *In vivo* **bioluminescence imaging of MAYV-nLuc infection in females at 4 and 8 weeks-old**

following i.p. inoculation (representative image). Luminescent signals (yellow arrows) were detected in

the limbs, tail, ears, and abdominal region at 1 (24 hpi), 2 (48 hpi), 6 and 10 dpi. Abdominal signals declined

after 6 dpi, whereas residual luminescence persisted mainly in the limbs at later time points. Images from

all animals in each age group are shown. All images were acquired and displayed using identical exposure

settings and a fixed color scale to enable direct comparisons over time. C, mock-infected control.

Temporal dynamics and tissue tropism of MAYV infection in 4-week-old mice. Viral

titers exhibited time- and tissue-dependent differences during acute infection (Fig 3A).

Mixed-effects analysis with Tukey's multiple comparisons identified a significant

increase in seminal vesicles at 48 hpi compared with 24 hpi (Mean difference = 0.9452,

SE of difference = 0.3397, p = 0.0284). Model-based estimates are reported as mean

differences ($\pm$ SE), while graphical representations show individual values with median

lines due to non-normal data distribution. At 24 hpi, no significant inter-tissue differences

were detected, consistent with a relatively homogeneous early dissemination pattern. By

48 hpi, tissue-specific differences emerged, with spleen titers exceeding those in liver

(Mean difference = 0.8391, SE of difference = 0.2150, p = 0.0352) and muscle (Mean

difference = 1.257, SE of difference = 0.2840, p = 0.0156). This substantial difference

suggests the spleen as a primary site of viral accumulation compared to muscular tissue

Sex-based analyses revealed transient differences at 24 hpi (Fig 3B). Specifically,

significant female–male differences were observed in the spleen (mean difference =

1.593, SE of difference = 0.5278, p = 0.0109) and kidneys (mean difference = 2.038, SE

of difference = 0.4765, p = 0.0159). In both tissues, males exhibited substantially higher

viral titers than females at this time point. Organ-specific viral distribution also varied

within each sex at 24 hpi. Among females, viral loads in the kidneys were significantly

higher than in the muscle (mean difference = 1.078, SE of difference = 0.2911, p =

0.0493). Conversely, in males, the spleen showed a significantly greater viral burden

compared to the liver (mean difference = 0.5601, SE of difference = 0.1205, p = 0.0205).

By 48 hpi, no significant differences in viral loads were observed either between sexes or

among tissues within each sex (Fig 3C). This lack of statistical significance (p > 0.05 for

all comparisons) indicates that the early sex-dependent effects and tissue-specific

disparities observed at 24 hpi are not sustained as the infection progresses toward later

stages.

Figure 3. **Early systemic dissemination with time-dependent tissue tropism and transient sex effects.**

Viral loads (log$_{10}$ PFU/g) across tissues at 24 and 48 hours post-infection (hpi) and sex-based comparisons

at each time point. (A) Viral loads across tissues at 24 and 48 hpi. (B) Sex-based comparison of viral loads

across tissues at 24 hpi. (C) Sex-based comparison at 48 hpi. Viral loads were analyzed using mixed-effects

models (REML) with time point and tissue as fixed effects and individual animal as a random effect,

followed by Tukey's multiple comparisons test. Values below the limit of detection were assigned a value

of LOD/2 (630 PFU/g) prior to transformation. Each dot represents an individual animal; horizontal lines

indicate median values. Only statistically significant comparisons are indicated. Statistical significance was

defined as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (***).

Early hematological alterations during acute MAYV infection. Hematological

analysis revealed alterations in leukocyte profiles during acute infection. Total white

blood cell counts (WBC) decreased at 24 and 48 hpi, accompanied by increased

neutrophil (NEU) and decreased lymphocyte (LYM) percentages (Fig 4A–C). Platelet

counts (PLT) also decreased at both time points (Fig 4D). Eosinophils (EOS) decreased,

basophils (BAS) increased at 48 hpi, and monocytes (MON) remained stable in infected

animals but decreased in mock controls (Fig S1A–C). Red blood cell counts (RBC)

increased at 48 hpi, while hemoglobin (HGB) and hematocrit (HCT) increased at both

time points (Fig S1D–F). Mean corpuscular volume (MCV) increased at 24 and 48 hpi,

mean corpuscular hemoglobin (MCH) increased at 24 hpi, and mean corpuscular

hemoglobin concentration (MCHC) remained unchanged (Fig S3A–C). Red cell

distribution width (RDW-CV and RDW-SD) decreased at both time points. Mean platelet

volume (MPV) increased, while platelet distribution width (PDW) decreased at 48 hpi.

Procalcitonin (PCT) decreased in infected animals without changes in mock controls (Fig

S3D–H). Statistically significant differences are indicated in the corresponding figures.

Biochemical and metabolic parameters. Aspartate transaminase (AST) at both 24 and

48 hpi and increased alanine transaminase (ALT) at 48 hpi (Fig 4E–F). Creatinine (CRE)

remained stable, whereas blood urea nitrogen (BUN) decreased at 24 hpi and showed a

decreasing trend at 48 hpi, including in mock animals (Fig S1G–H). Total protein (TP)

decreased at 48 hpi, while albumin (ALB), globulin (GLO), and gamma-glutamyl

transferase (GGT) remained unchanged (Fig S2A–C, E). Total bilirubin (TBIL) decreased

at 24 hpi (Fig S2D). Glucose (GLU) increased at both time points, and body weight (BW)

decreased at 24 and 48 hpi (Fig S2G–H). Statistical significance is indicated in the

corresponding figures.

Figure 4. Early hematological and hepatic changes following acute MAYV infection.

Hematological and hepatic parameters during acute MAYV infection at 24 and 48 hours post-infection

(hpi) compared with mock controls. (A) Total white blood cell (WBC) count, (B) neutrophil percentage

(NEU), (C) lymphocyte percentage (LYM), (D) platelet count (PLT), (E) alanine transaminase (ALT), and

(F) aspartate transaminase (AST). Each dot represents an individual animal; horizontal lines indicate

median values. Statistical analyses were performed using mixed-effects models (REML) for repeated

measures with time and condition as fixed effects and individual animal as a random effect, or non-

parametric tests when assumptions were not met. Differences were considered statistically significant at p

< 0.05 (*), p < 0.01 (**) and p < 0.001 (***).

Gestational stage–dependent differences in MAYV distribution during pregnancy.

At 48 hpi, viral titers exhibited significant variations based on gestational stage across

multiple tissues (Fig 5). Mixed-effects modeling followed by Tukey's multiple comparison test revealed significantly higher infectious viral loads at 48 hpi in tissues analyzed at 15.5–16.5 dpc (pregnant mice infected at 13.5-14.5 dpc) compared with those analyzed at 9.5–10.5 dpc (pregnant mice infected at 7.5-8.5 dpc). This stage-dependent increase was observed in the spleen (mean difference = 0.5751, SE of difference = 0.1624, p = 0.0213), liver (mean difference = 1.924, SE of difference = 0.1552, p = 0.0011), kidneys (mean difference = 1.691, SE of difference = 0.3097, p = 0.0055), uterus (mean difference = 1.660, SE of difference = 0.6606, p = 0.0403) and placenta (mean difference = 0.7328, SE of difference = 0.2277, p = 0.0061). However, infectious virus remained undetectable in fetal tissues at this same stage (15.5–16.5 dpc), with titers falling below the limit of detection. This suggests that even when the placenta is significantly infected, the virus remained undetectable at the fetal compartment in an infectious form. Within early gestation (9.5–10.5 dpc), tissue-specific differences were observed, with the spleen showing higher titers than the liver and kidneys. At mid-gestation, infectious viral particles remained detectable in maternal and placental compartments, supporting ongoing systemic dissemination despite restricted recovery of infectious virus from fetuses.

MAYV WT distribution in maternal and fetal tissue

Figure 5. Gestational stage shapes maternal–fetal distribution and restricts recovery of infectious virus from mid-gestation fetuses. Infectious viral loads ($\log_{10}$ PFU/g) in maternal and fetal tissues at 48 hours post-infection (hpi) according to gestational stage. Females were infected at early (7.5–8.5 dpc; analyzed at 9.5–10.5 dpc) or mid-gestation (13.5–14.5 dpc; analyzed at 15.5–16.5 dpc). Each dot represents an individual animal; horizontal lines indicate median values. Viral loads were analyzed using mixed-effects models (REML) with gestational stage and tissue as fixed effects and individual animal as a random effect, followed by Tukey's multiple comparisons test. Values below the limit of detection were assigned a value of LOD/2 (630 PFU/g) prior to transformation. Only statistically significant comparisons are indicated. Only statistically significant comparisons are indicated. Statistical significance was defined as $p < 0.05$ (*), $p < 0.01$ (**) and $p < 0.001$ (***).

Acute MAYV infection induces leukocyte redistribution without overt hematological cytopenias at 48 hpi. Infected females at both gestational stages exhibited a shift in leukocyte distribution, characterized by increased neutrophil and decreased lymphocyte percentages compared with mock controls (Fig S4A–B). Total white blood cell counts (WBC) did not differ between infected and mock animals (Fig S4C), indicating leukocyte redistribution rather than overall leukocytosis. Red blood cell counts, platelet numbers, and procalcitonin levels remained comparable between groups (Fig

S4D–F). Given the limited mock sample size, these findings should be interpreted with

caution.

Immunohistochemistry confirms placental and fetal localization of MAYV at mid

stage. Immunohistochemical analysis revealed clear differences in MAYV antigen

distribution between infected and non-infected tissue at fetus and placenta from an

infected female at mid-gestation. In placentas from infected females, moderate granular

cytoplasmic staining was observed, marking a substantial proportion of the giant

trophoblast cells within the maternal–fetal junction zone (basal plate) indicating focal

placental infection. In contrast, placentas from non-infected pregnant females showed

only nonspecific erythrocyte staining and overall negative to mild background labeling.

In fetus from infected female, intense immunostaining was detected across multiple

organs, particularly in the visceral epithelium of the intestine, the developing bone

(growth cartilage), and other collagen-rich tissues including cartilage and dermis.

Calcified bone exhibited no staining or only minimal labeling in both intensity and extent.

Fetuses from non-infected females were largely negative, with nonspecific erythrocyte

staining and minimal background labeling in collagen-rich tissues (Fig 6). This discrepancy

may reflect detection limits of plaque assays or persistence of viral antigens following partial viral

clearance.

Figure 6. IHC against MAYV. A) Placenta of an uninfected mouse (control): Negative to mild (-/+)

immunostaining of isolated trophoblasts (foam giant cells) from the basal plate of the maternal-fetal

junction is observed. Scale bar = 50µm, 40×. B) Placenta of an experimentally infected mouse: Mild to

moderate, granular intracytoplasmic staining of foci of trophoblasts (foam giant cells) from the basal plate

of the maternal-fetal junction is observed (asterisks). Scale bar = 50µm, 40×. C) Fetus of the same

uninfected mouse of Figure A: No immunostaining against MAYV is observed (-). Scale bar = 500µm, 4×.

D) Fetus of the same experimentally infected mouse of Figure B: There is intense immunostaining in several

organs, particularly in uncalcified developing bone and growth cartilage (asterisks), as well as other

collagen-rich tissues (cartilage and dermis) (arrowheads) and the perimysium of skeletal muscles (arrows).

Already calcified bone shows no staining or minimal staining in intensity and extent. Scale bar = 500µm,

4×.

Placental transcriptional responses to MAYV are gestational stage-dependent. To

investigate placental transcriptional responses to MAYV infection, differential gene

expression was performed. The profiles revealed gene expression patterns influenced by

viral infection at early and mid-stages: 208 genes were upregulated and 60 downregulated

at the early stage, and 139 genes were upregulated and 1 downregulated at the mid stage

(Fig 7A and Table S1). Among the upregulated genes, 50 were shared between both

stages, leaving 158 stage-specific upregulated genes at the early stage and 89 at the mid

stage. These data suggest a robust transcriptional response to infection during the early

stage of pregnancy, while the response in mid pregnancy, although sharing common pathways, is comparatively attenuated.

Gene Ontology (GO) enrichment analysis of the 50 shared upregulated genes revealed significant association with antiviral defense pathways, including responses to viral infection, interferon-stimulated genes (ISGs) activity and cytokine-mediated signaling (Fig 7B).

The 158 genes uniquely upregulated in the early stage were enriched not only for the immune pathways but also for distinct biological processes, including heart development, myeloid cell development, regulation of plasminogen activation, connective tissue development, and adipogenesis (Fig 7C). In contrast, the 89 late-stage-specific upregulated genes were again predominantly associated with immune functions, specifically type II interferon signaling, negative regulation of innate immune response, and innate antiviral immune response (Fig 7D).

Finally, GO analysis of the 60 genes downregulated specifically at the early stage identified enrichment for processes such as multi-organism reproductive process, antigen processing and presentation of peptide antigen via MHC class I, lymphocyte activation, signaling by NTRKs, insulin secretion, thermogenesis, extracellular matrix organization, and adrenergic signaling in cardiomyocytes (Fig 7E).

Figure 7. Transcriptomic profiles and Gene Ontology enrichment of placental response to MAYV

infection at early and mid-gestational stages. A) Comparison of placentas infected at 7.5–8.5 dpc (early)

and 13.5–14.5 dpc (mid) versus controls revealed 208 up-/60 down-regulated genes at early stage and 139

up-/1 down-regulated at mid stage, with 50 genes shared between stages. B) Shared genes were enriched

for antiviral and interferon-mediated responses. C) Early-stage-specific genes were associated with

immune, developmental, and metabolic pathways. D) Mid-stage-specific genes were mainly related to type

II interferon and innate immune modulation. E) Early-stage down-regulated genes were enriched for

reproductive, metabolic, and signaling processes.

MAYV detection in reproductive tissues and sperm. Following mating, the liver,

spleen, *quadriceps femoris* muscle, ovaries, uterus, seminal vesicles and testes were

analyzed for the presence/absence of MAYV by plaque assay in all animals. RT-qPCR

was performed on the spleen of infected animals and on the corresponding organs from

exposed animals. To detect the presence/absence of MAYV in sperm, RT-qPCR was

performed and, following cell culture amplification and plaque assay.

Infectious viral particles were detected in the spleen (7/7), ovaries (3/7), and muscle (1/7)

of infected females mated with uninfected males. In contrast, all organs analyzed from

exposed males were negative by both plaque assay and RT-qPCR. However, after cell

culture amplification of sperm samples followed by plaque assay, infectious virus was

detected in one exposed male (1/7), which had been paired with the female positive for

infectious virus in both the spleen and muscle.

In infected males mated with uninfected females, infectious viral particles were

consistently detected in the testes (6/6) and in the spleen (3/6). Plaque assays performed

after cell culture amplification of sperm revealed infectious virus in four males (4/6),

consistent with the cytopathic effect observed (Fig 8). Viral RNA was also detected by

RT-qPCR in direct unamplified sperm samples from the same four animals. No infectious

virus was detected in the seminal vesicles. Among exposed females, infectious viral

particles tested positive in two animals: the uterus and ovary of one female and the spleen

of another, although the RT-qPCR resulted below the limit of detection.

Overall, viral detection in exposed animals was limited and inconsistent across analytical

approaches.

Figure 8. Cytopathic effect of MAYV in sperm cell culture–based amplification (Representative image)**.**

(A) Sperm extracted from the bilateral cauda epididymides. (B´D) Representative images of Vero cells

inoculated with processed sperm samples showing a MAYV-positive sample (B) a MAYV-negative sample

(C) and Mock-infected Vero cells (D). Cell morphology was examined by a Zeiss Primovert inverted light

microscope (40×).

DISCUSSION

Elucidating the tissue tropism and viral dynamics of MAYV is essential given the limited

evidence available for this emerging arbovirus and the relevance of murine models for

characterizing infection *in vivo*. Here, we provide evidence of MAYV dissemination to

fetal tissues in mice, identifying anatomical targets not previously described. Together,

these findings support a model in which gestational timing critically determines placental

permissiveness and fetal exposure to MAYV, identifying early gestation as a window of

heightened vulnerability during acute infection. Although direct temporal equivalence

between murine and human gestation is not possible, early murine gestation (7.5-8.5 dpc)

represents a developmental window of active placentation and early organogenesis that

broadly parallels the first trimester of human pregnancy, whereas mid-gestation (13.5-14.5 dpc) reflects a more mature placental interface (21,40). Considering that related alphaviruses may cross the placental barrier, our results underscore the importance of including pregnancy models when assessing MAYV transmission routes (41).

To examine early dissemination under controlled conditions, we employed a nanoluciferase-expressing MAYV reporter to evaluate intraperitoneal delivery as a strategy to achieve systemic exposure. *In vivo* imaging showed rapid and sustained viral spread, with peripheral luminescent patterns consistent with the musculoskeletal involvement previously described in experimental models and human disease (18,25,29).

The incomplete overlap between bioluminescent signal and recovery of infectious virus, particularly in older animals, likely reflects differences in analytical sensitivity between reporter detection and plaque assays or the persistence of viral material following clearance of replicative particles. In contrast, the consistent recovery of infectious virus in younger mice supports age-dependent variation in susceptibility and viral maintenance within target tissues.

This imaging approach enabled longitudinal, non-invasive monitoring of viral dynamics over a 10-day period, encompassing both acute infection and early recovery phases (25,26). By providing temporal and anatomical resolution of dissemination events that are difficult to capture in natural transmission settings, the model links viral spread to measurable tissue outcomes.

Together, these observations expand the experimental landscape for MAYV research, supporting mechanistic investigations of tissue tropism, host responses, and transmission potential under standardized conditions, and offering a framework for future studies addressing reproductive involvement and non-vector routes of spread.

MAYV WT exhibited rapid and systemic dissemination at 24 and 48 hpi, with infectious

virus detected across multiple organs, including detection in reproductive tissues of

pregnant females. These results are consistent with other murine alphavirus models,

including IFNAR−/− mice, in which widespread viral detection at early time points

reflects acute viremia (28). This pattern reflects an initial homogeneous dissemination

phase at 24 hpi followed by the emergence of tissue-specific viral accumulation at 48 hpi,

consistent with previous murine MAYV models showing that viral burden is influenced

by host age, immune status, and route of infection, with predominant viral accumulation

in skeletal muscle, limbs, and lymphoid organs such as the spleen (10,30). Infectious

particles were also detected in muscle, supporting the myopathy and myositis previously

described during MAYV infection (10,29). The increase in viral load in seminal vesicles

at 48 hpi further supports progressive involvement of reproductive tissues during acute

infection, consistent with the detection of infectious virus in male reproductive organs.

The early hematological alterations observed in our study partially align with previous

experimental and clinical reports of MAYV infection. In IFNAR−/− mice model, early

leukopenia predominantly driven by lymphopenia was reported during acute infection

(29), consistent with the leukopenia and lymphocyte reduction detected in our model. In

contrast, a BALB/c intraplantar infection model described increases in monocytes,

neutrophils, and lymphocytes during the acute phase (25), indicating that hematological

profiles may vary depending on experimental conditions. In human cases, leukopenia,

thrombocytopenia, and elevated ALT and AST have been documented among reported

laboratory abnormalities (18), which parallels the transient cytopenias and

aminotransferase elevations observed here. Together, these comparisons suggest that

early systemic hematological and biochemical changes are reproducible features of

MAYV infection, although their magnitude and direction may differ across models and

clinical settings. Despite the limited mock sample size in pregnant cohorts, the consistent leukocyte redistribution observed at 48 hpi across gestational stages supports the interpretation of an early systemic inflammatory response rather than global hematological impairment.

MAYV belongs to the Alphavirus genus (18,30), which includes viruses capable of affecting the nervous system and embryonic development. The detection of infectious particles in reproductive organs (uterus, ovary, testes, seminal vesicles) at both time points highlights a tissue tropism previously described in immunodeficient mice (15) and raises the possibility of sexual or vertical transmission. Transient and early sex-based differences were observed at 24 hpi. These differences were no longer evident at 48 hpi. While most studies focused on age and nutritional status as determinants of disease outcome (25,31,42), these observations suggest a potential sex-dependent component of MAYV infection that warrants further investigation, particularly given similar findings reported for CHIKV (43).

A gestational stage–dependent susceptibility was evident in our model. At the early stage of pregnancy (7.5–8.5 dpc infection; analyzed at 9.5–10.5 dpc), fetuses contained infectious particles at 48 hpi, indicating that MAYV can cross the maternal–fetal barrier during acute infection. In contrast, our data suggest a more restrictive placental environment during mid-gestation (13.5–14.5 dpc; analyzed at 15.5–16.5 dpc), as no infectious particles were recovered from fetal tissues despite significant placental viral loads. It is critical to interpret this 'restriction' specifically as the absence of infectious viral particles detectable by plaque assay. While this confirms a lack of replication-competent virions in the fetus at this stage, it does not preclude the potential presence of viral RNA or non-infectious antigens that might be detectable through more sensitive

molecular methods. This is supported by our immunohistochemistry results, where viral antigen was detected in fetal tissues at mid-gestation despite the absence of recoverable infectious virus, suggesting either focal infection below the limit of detection or the persistence of viral antigens following partial viral clearance. This timing-dependent susceptibility is consistent with observations from ZIKV infection, where placental infection and transplacental transmission are strongly associated with maternal infection occurring during the first trimester of human pregnancy, whereas infections acquired at later gestational stages are less frequently linked to fetal exposure (20,44). This window—analogous to early human embryogenesis (41)— displayed a broad transcriptional response (208 upregulated, 60 downregulated genes), including alterations in developmental and tissue-remodeling programs, as well as downregulation of hormonal signaling, MHC I presentation, and ECM organization. These disruptions are compatible with teratogenesis described for alphaviruses such as Semliki Forest virus (42) and ZIKV (21).

In contrast, at mid-gestation infection (13.5–14.5 dpc), infectious viral particles were not detected in fetuses, and transcriptional activation was more restricted (139 upregulated, 1 downregulated), dominated by interferon-mediated pathways. This profile aligns with increased placental maturity and antiviral competence (21). The presence of infectious viruses in placenta and uterus, but not in fetal tissues, supports restricted fetal infection at this stage (40,41). Together, these findings support the concept of a conserved gestational window of vulnerability to vertical transmission among arboviruses, in which early placentation is associated with increased permissiveness at the maternal–fetal interface. The 50 shared upregulated genes across gestational stages—mostly ISGs—reflect a conserved antiviral signature, but timing-dependent transcriptional divergence underscores critical periods of vulnerability, including disruption of angiogenic and

developmental pathways (45) and inflammatory responses characteristic of MAYV

(25,26). Viral antigen was detected by immunohistochemistry in placental and fetal

tissues at mid-gestation, despite the absence of recoverable infectious virus, suggesting

focal or low-level infection below plaque assay detection limits or persistence of viral

antigens following partial viral clearance.

While direct extrapolation from murine models to humans is limited, these findings

provide a biologically plausible framework to explore potential maternal–fetal risks with

MAYV infection in human during pregnancy.

Limitations should be considered when interpreting these findings. Intraperitoneal

inoculation does not fully recapitulate natural mosquito-borne transmission and may

influence early viral kinetics. However, this route is widely used in murine models of

systemic arbovirus infection and was deliberately selected to enable controlled

assessment of early systemic dissemination and tissue tropism while minimizing

procedure-related distress (46,47) and local inflammatory confounders. Although

intraplantar models better reproduce articular manifestations (25,26,28), intraperitoneal

delivery provides a standardized framework to evaluate systemic viral dynamics.

Previous murine studies have reported systemic MAYV dissemination and detection of

viral RNA in gonadal tissues, suggesting broader tissue distribution (48), although

evidence supporting a functional role of reproductive infection in transmission has

remained indirect. In the present study, infectious virus was detected in reproductive

organs and sperm, supporting the biological plausibility of sexual transmission, though

not establishing it as an efficient route and therefore requiring dedicated follow-up

studies. Detection among exposed animals was infrequent, consistent with minimal or

transient exposure rather than sustained transmission. If sexual transmission occurs, it is

likely to involve low-level infection and restricted replication within gonadal tissues,

potentially limiting detectability and contributing to the scarcity of documented cases, a

challenge also described for other arboviruses in semen (41).

Taken together, these findings broaden the understanding of MAYV pathogenesis by

integrating virological, transcriptomic, histological, and imaging approaches within a

unified murine model. We provide evidence of expanded tissue tropism, including

reproductive and fetal involvement, and identify gestational timing as a determinant of

placental permissiveness and fetal exposure. Detection of infectious virus in reproductive

tissues supports the biological plausibility of non-vector transmission, although its

epidemiological relevance remains uncertain. While extrapolation to humans requires

caution, these results establish a preclinical framework for investigating MAYV systemic

dissemination, maternal–fetal infection, and potential transmission routes.

CONCLUSION

This study establishes a murine model of MAYV infection that reveals rapid systemic

dissemination followed by tissue-specific viral distribution, broad tissue tropism, and

early involvement of reproductive organs. Transient sex-dependent differences were

observed during the early phase of infection, suggesting a potential but limited role of sex

in modulating viral dynamics. Gestational stage emerged as a key determinant of

outcomes: specifically, infection initiated during early pregnancy (7.5–8.5 dpc) and

analyzed at 9.5–10.5 dpc identified a window of increased susceptibility associated with

the detection of infectious viral loads in fetal tissue. In contrast, infection occurring during

mid-gestation (13.5–14.5 dpc) and analyzed at 15.5–16.5 dpc was associated with

restricted fetal infection despite sustained viral presence in maternal and placental tissues. The detection of infectious viral particles in reproductive tissues supports the biological plausibility of alternative transmission routes, although their epidemiological relevance remains uncertain. These findings provide a comprehensive experimental framework to advance the understanding of MAYV pathogenesis, systemic dissemination, and maternal–fetal and non-vector transmission dynamics.

REFERENCES

1. Organización Panamericana de la Salud, Organización Mundial de la Salud. Alerta epidemiológica: fiebre de Mayaro. Washington (DC): OPS/OMS; 2019. Available from: https://www.paho.org

2. Blohm G, Elbadry MA, Mavian C, Stephenson C, Loeb J, White S, et al. Mayaro as a Caribbean traveler: evidence for multiple introductions and transmission of the virus into Haiti. Int J Infect Dis. 2019;87:151–153. doi:10.1016/j.ijid.2019.07.031

3. Ganjian N, Riviere-Cinnamond A. Mayaro virus in Latin America and the Caribbean. Rev Panam Salud Publica. 2020;44:e14. doi:10.26633/RPSP.2020.14

4. Valencia-Marín BS, Gandica ID, Aguirre-Obando OA. The Mayaro virus and its potential epidemiological consequences in Colombia: an exploratory biomathematics analysis. Parasit Vectors. 2020;13(1):519. doi:10.1186/s13071-020-04354-1

5. Brustolin M, Bartholomeeusen K, Rezende T, Ariën KK, Müller R. Mayaro virus, a potential threat for Europe: vector competence of autochthonous

vector species. Parasit Vectors. 2024;17(1):93. doi:10.1186/s13071-024-06293-7

6. Celone M, Beeman S, Han BA, Potter AM, Pecor DB, Okech B, et al. Understanding transmission risk and predicting environmental suitability for Mayaro virus in Central and South America. PLoS Negl Trop Dis. 2024;18(1):e0011859. doi:10.1371/journal.pntd.0011859

7. García-Romero C, Carrillo Bilbao GA, Navarro JC, Martin-Solano S, Saegerman C. Arboviruses in mammals in the Neotropics: a systematic review. Viruses. 2023;15(2):417. doi:10.3390/v15020417

8. Fundación iO. Aumento significativo de casos de fiebre Mayaro en Acre, Brasil. 2025. Available from: https://fundacionio.com/aumento-significativo-de-casos-de-fiebre-mayaro-en-acre-brasil/

9. Andreolla AP, Borges AA, Bordignon J, Duarte dos Santos CN. Mayaro virus: the state of the art for antiviral drug development. Viruses. 2022;14(8):1787. doi:10.3390/v14081787

10. Diagne CT, Bengue M, Choumet V, Hamel R, Pompon J, Missé D. Mayaro virus pathogenesis and transmission mechanisms. Pathogens. 2020;9(9):738. doi:10.3390/pathogens9090738

11. Marinho MS, Ferreira GM, Grosche VR, Nicolau-Junior N, Campos TL, Santos IA, et al. Evolutionary profile of Mayaro virus in the Americas: an update into genome variability. Viruses. 2024;16(5):809. doi:10.3390/v16050809

12.   Cereghino C, Roesch F, Carrau L, Hardy A, Ribeiro-Filho HV, Henrion-

Lacritick A, et al. The E2 glycoprotein holds key residues for Mayaro virus

adaptation to the urban Aedes aegypti mosquito. PLoS Pathog.

2023;19(4):e1010491. doi:10.1371/journal.ppat.1010491

13.   Mendonça DC, Reis EVS, Arias NEC, Valencia HJ, Bonjardim CA. A study

of the MAYV replication cycle: correlation between the kinetics of viral

multiplication and viral morphogenesis. Virus Res. 2022;323:199002.

doi:10.1016/j.virusres.2022.199002

14.   Ribeiro-Filho HV, Coimbra LD, Cassago A, Rocha RPF, Guerra JVS, de

Felicio R, et al. Cryo-EM structure of the mature and infective Mayaro virus

at 4.4 Å resolution reveals features of arthritogenic alphaviruses. Nat

Commun. 2021;12(1):3038. doi:10.1038/s41467-021-23400-9

15.   Krokovsky L, Lins CRB, Guedes DRD, Wallau GL, Ayres CFJ, Paiva

MHS. Dynamics of Mayaro virus transmission in Aedes aegypti, Culex

quinquefasciatus mosquitoes, and a mouse model. Viruses. 2023;15(3):799.

doi:10.3390/v15030799

16.   Pereira TN, Virginio F, Souza JI, Moreira LA. Emergent arboviruses: a

review about Mayaro virus and Oropouche orthobunyavirus. Front Trop

Dis. 2021;2:737436. doi:10.3389/fitd.2021.737436

17.   de Thoisy B, Gardon J, Alba Salas R, Morvan J, Kazanji M. Mayaro virus

in wild mammals, French Guiana. Emerg Infect Dis. 2003;9(10):1326–

1329. doi:10.3201/eid0910.030161

18.   Silva-Ramos CR, Mejorano-Fonseca JA, Hidalgo M, Rodríguez-Morales

AJ, Faccini-Martínez ÁA. Clinical, epidemiological, and laboratory features

of Mayaro virus infection: a systematic review. Curr Trop Med Rep.

2023;10(4):309–319. doi:10.1007/s40475-023-00308-6

19.   de Curcio JS, Salem-Izacc SM, Pereira Neto LM, Nunes EB, Anunciação

CE, Silveira-Lacerda EP. Detection of Mayaro virus in Aedes aegypti

mosquitoes   circulating   in   Goiânia,   Brazil.   Microbes   Infect.

2022;24(4):104948. doi:10.1016/j.micinf.2022.104948

20.   Brasil P, Pereira JP Jr, Moreira ME, Ribeiro Nogueira RM, Damasceno L,

Wakimoto M, et al. Zika virus infection in pregnant women in Rio de

Janeiro.   N   Engl   J   Med.   2016;375(24):2321–2334.

doi:10.1056/NEJMoa1602412

21.   Vermillion MS, Lei J, Shabi Y, Baxter VK, Crilly NP, McLane M, et al.

Intrauterine Zika virus infection of pregnant immunocompetent mice

models transplacental transmission and adverse perinatal outcomes. Nat

Commun. 2017;8:14575. doi:10.1038/ncomms14575

22.   Ramful D, Carbonnier M, Pasquet M, Bouhmani B, Ghazouani J,

Noormahomed T, et al. Mother-to-child transmission of chikungunya virus

infection.   Pediatr   Infect   Dis   J.   2007;26(9):811–815.

doi:10.1097/INF.0b013e3180616d4f

23.   Gérardin P, Barau G, Michault A, Bintner M, Randrianaivo H, Choker G, et

al. Multidisciplinary prospective study of mother-to-child chikungunya

virus infections on the island of La Réunion. PLoS Med. 2008;5(3):e60.

doi:10.1371/journal.pmed.0050060

24.   Pancetti FGM, Honório NA, Urbinatti PR, Lima-Camara TN. Twenty-eight

years of Aedes albopictus in Brazil: a rationale to maintain active

entomological and epidemiological surveillance. Rev Soc Bras Med Trop. 2015;48(1):87–89. doi:10.1590/0037-8682-0155-2014

25. Mota MT, Costa VV, Sugimoto MA, Guimarães GF, Queiroz-Junior CM, Moreira TP, et al. In-depth characterization of a novel live-attenuated Mayaro virus vaccine candidate using an immunocompetent mouse model. Sci Rep. 2020;10:5306. doi:10.1038/s41598-020-62084-x

26. Santos FM, Dias RS, de Oliveira MD, Costa ICTA, Fernandes LS, Pessoa CR, et al. Animal model of arthritis and myositis induced by the Mayaro virus. PLoS Negl Trop Dis. 2019;13(5):e0007375. doi:10.1371/journal.pntd.0007375

27. da Silva Caetano CC, Camini FC, Almeida LT, Ferraz AC, da Silva TF, Lima RLS, et al. Mayaro virus induction of oxidative stress is associated with liver pathology in a non-lethal mouse model. Sci Rep. 2019;9:12347. doi:10.1038/s41598-019-51713-9

28. de Carvalho AC, Dias CSB, Coimbra LD, Rocha RPF, Borin A, Fontoura MA, et al. Characterization of systemic disease development and paw inflammation in a susceptible mouse model of Mayaro virus infection. Int J Mol Sci. 2023;24(5):4799. doi:10.3390/ijms24054799

29. Rosa RB, de Castro EF, de Oliveira Santos D, da Silva MV, Pena LJ. Mouse models of Mayaro virus. Viruses. 2023;15(9):1803. doi:10.3390/v15091803

30. Acosta-Ampudia Y, Monsalve DM, Rodríguez Y, Pacheco Y, Anaya JM, Ramírez-Santana C. Mayaro: an emerging viral threat? Emerg Microbes Infect. 2018;7(1):163. doi:10.1038/s41426-018-0163-5

31. Weger-Lucarelli J, Carrau L, Levi LI, Rezelj V, Vallet T, Blanc H, et al.
Host nutritional status affects alphavirus virulence, transmission, and
evolution. PLoS Pathog. 2019;15(11):e1008089.
doi:10.1371/journal.ppat.1008089

32. Powers AM, Aguilar PV, Chandler LJ, Brault AC, Meakins TA, Watts D,
et al. Genetic relationships among Mayaro and Una viruses suggest distinct
patterns of transmission. Am J Trop Med Hyg. 2006;75(3):461–469.
doi:10.4269/ajtmh.2006.75.461

33. Chuong C, Bates TA, Weger-Lucarelli J. Infectious cDNA clones of two
strains of Mayaro virus for studies on viral pathogenesis and vaccine
development. Virology. 2019;535:227–231.
doi:10.1016/j.virol.2019.07.013.

34. Figueiredo CM, Neris RLS, Gavino-Leopoldino D, da Silva MOL, Almeida
JS, dos Santos JS, et al. Mayaro virus replication restriction and induction
of muscular inflammation in mice are dependent on age, type I interferon
response, and adaptive immunity. Front Microbiol. 2019;10:2246.
doi:10.3389/fmicb.2019.02246

35. Whitten WK. Modification of the oestrous cycle of the mouse by external
stimuli associated with the male. J Endocrinol. 1956;13(4):399–404.
doi:10.1677/joe.0.0130399

36. Roszkowski M, Mansuy IM. High-efficiency RNA extraction from sperm
cells using guanidinium thiocyanate supplemented with tris(2-
carboxyethyl)phosphine. Front Cell Dev Biol. 2021;9:648274.
doi:10.3389/fcell.2021.648274

37. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun. 2019;10(1):1523. doi:10.1038/s41467-019-09234-6

38. Llamas-González YY, Campos D, Pascale JM, Arbiza J, González-Santamaría J. A functional ubiquitin-proteasome system is required for efficient replication of New World Mayaro and Una alphaviruses. Viruses. 2019;11(4):370. doi:10.3390/v11040370

39. Valdés-Torres P, Campos D, Bhakta M, Galán-Jurado PE, Durant-Archibold AA, González-Santamaría J. Honokiol and alpha-mangostin inhibit Mayaro virus replication through different mechanisms. Molecules. 2022;27(21):7362. doi:10.3390/molecules27217362

40. Megli CJ, Coyne CB. Infections at the maternal–fetal interface: an overview of pathogenesis and defence. Nat Rev Microbiol. 2022;20(2):67–82. doi:10.1038/s41579-021-00610-y

41. Hcini N, Lambert V, Picone O, Carod JF, Carles G, Pomar L, et al. Arboviruses and pregnancy: are the threats visible or hidden? Trop Dis Travel Med Vaccines. 2024;10(1):1–12. doi:10.1186/s40794-023-00213-w

42. Atkins GJ, Carter J, Sheahan BJ. Effect of alphavirus infection on mouse embryos. Infect Immun. 1982;35(2):546–553.

43. Taylor M, Rayner JO. Immune response to chikungunya virus: sex as a biological variable and implications for natural delivery via the mosquito. Viruses. 2023;15(9):1869. doi:10.3390/v15091869

44. Hoen B, Schaub B, Funk AL, Ardillon V, Boullard M, Cabié A, et al. Pregnancy outcomes after ZIKV infection in French territories in the Americas. N Engl J Med. 2018;378(11):985–994. doi:10.1056/NEJMoa1709481

45. Wu Y, Su K, Zhang Y, Liang L, Wang F, Chen S, et al. A spatiotemporal transcriptomic atlas of mouse placentation. Cell Discov. 2024;10(1):74. doi:10.1038/s41421-024-00740-6

46. European Parliament, Council of the European Union. Directive 2010/63/EU on the protection of animals used for scientific purposes. 2010. Available from: https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32010L0063

47. National Research Council (US). Recognition and alleviation of pain in laboratory animals. Washington (DC): National Academies Press; 2009

48. Rosa RB, De Castro EF, De Oliveira Santos D, Dantas WM, Tanaka CA, Mendes RPG, et al. A comprehensive murine clinical model for development of countermeasures and studying Mayaro virus infection. PLoS Negl Trop Dis. 2025;19(7):e0013333. doi:10.1371/journal.pntd.0013333

ACKNOWLEDGMENTS

We thank José González Santamaría from the Department of Genomics and Proteomics Research, Gorgas Memorial Institute for Health Studies, Panama, for providing the primary antibody used in this study.

AUTHOR CONTRIBUTIONS

Conceptualization: APA, GM and MC. Methodology: APA, PP, JLP, MPG, BV, JMV, GM and MC. Investigation: APA, PP, JLP, MPG, GG, BV, JMV, GM and MC. Visualization: APA, PP, JLP, MPG, GG, BV, JMV, GM and MC. Statistical Analysis: APA. Supervision: GM and MC. Writing—original draft: APA, PP, JLP, GG, JMV, GM and MC. Writing—review & editing: APA, PP, JLP, MPG, GG, JH, AF, BV, JMV, GM and MC. Project administration: GM and MC. Funding acquisition: GM and MC. All authors agreed to submit the manuscript and have read and approved its final version. Authors take full responsibility for its content.

FUNDING

Institut Pasteur de Montevideo and FOCEM (COF 03/11), G4 program Institut Pasteur de Montevideo.

COMPETING INTERESTS

The authors declare no competing interests.