Genome-wide identification of heat shock proteins (HSP70) in the Cuban painted snail *Polymita picta*: evolutionary constraints face climate change challenges

ABSTRACT

Climate change poses a critical threat for biodiversity. Among ectothermal animals, terrestrial molluscs are considered highly vulnerable to temperature increase and drought due to their low vagility and specialized ecological requirements. In response to thermal stress, heat shock proteins (HSP70) play a crucial role in the response syndrome and phenotypic plasticity and mediate the adaptation to fluctuating environments. Here, we performed the first genome-wide identification and characterization of the HSP70 family in a Neotropical terrestrial gastropod of high conservation concern: the Cuban painted snail *Polymita picta*. Using comprehensive bioinformatic approaches, we identified HSP70 paralogs, analysed their structural features and assessed sequence variation, highlighting their potential relevance for thermoresistance. Our findings reveal an HSP70 repertoire of ten diversified sequences with signatures of functional specialization and possible adaptive variation linked to environmental temperature regimes. Using novel molecular resources, this pioneering analysis establishes a baseline for early warning systems, integrating stress-related biomarkers into conservation management and climate change adaptation strategies for threatened terrestrial molluscs.

INTRODUCTION

Ectothermic organisms, such as terrestrial molluscs, are extremely susceptible to fluctuations in temperature and humidity due to lacking acclimation capacity and a limited increase in physiological variance (Morley et al., 2019; Miyahira et al., 2022; Staikou et al., 2024; Noble et al., 2025). Their low vagility, or dispersal capability, poses a challenge for movement and colonization of new habitats with favourable environmental conditions (Kramarenko, 2014). Gastropods are currently exposed to environmental pressures primarily related to human activities, which force them to adapt or acclimate to rapidly changing conditions (Nicolai and Ansart, 2017). However, acclimation does not necessarily result in complete compensation in response to environmental changes (Huey et al., 1999). Often, increases in physiological rates are only partially compensated for (Havird et al., 2020). Disruption of cellular or organismal homeostasis caused by internal or external factors is defined as physiological or primary biological stress (Kültz, 2020). This latter encompasses three major dimensions: intrinsic developmental stress, environmental stress, and aging. The environmental stress arises when external conditions exceed critical thresholds, posing a threat to homeostatic balance (Hayflick, 2007). Within this framework, external factors are termed "stressors," while the internal reaction they provoke is referred to as "stress" or the "stress response syndrome" (Van Straalen, 2003).

In response to stress, physiological acclimation is mediated by endocrine and epigenetic processes, and relies on a highly evolutionarily conserved stress proteasome that modifies underlying cellular mechanisms to maintain physiological performance (Taff & Vitousek, 2016; Kültz, 2020). It is that populations experiencing high but predictable environmental variability will evolve acclimation, particularly when the fitness costs of plasticity are low (Chevin & Hoffmann, 2017). This capacity has also been associated with various bioecological traits such as habitat type, latitude, body size, and life-history stage (Rohr et al., 2018; Morley et al., 2019).

In general, habitat loss, meteorological phenomena, drought and high temperatures constitute the main threats for terrestrial gastropods affecting species survival, fertility and longevity. Gastropods have evolved several adaptations guaranteeing survival and reproduction in dry and hot environments (Astor et al., 2017; Nicolai and Ansart, 2017). These adaptations involve thermodynamic, physiological, cellular and molecular processes (Schweizer et al., 2019). Aestivation, the induction of chaperones, changes in tissue and

cell structure, as well as modifications in mucus production and pigmentation, demonstrate the interrelationship of functions across different levels of biological organization in response to thermal stress and dehydration (Scheil et al., 2012; Bhagat et al., 2016; Schweizer et al., 2019; Tull et al., 2020). Contrasting with the numerous studies on heat effects on macro-physiology and morphology in terrestrial snails, research focusing on tissue, cells or molecular aspects are scarce (Schweizer et al., 2019).

One of the most studied molecular systems directly related to thermal stress response syndrome is the heat shock protein (HSP) family HSP70. These cellular chaperones are ubiquitous, highly conserved proteins distributed in all organisms, from bacteria to humans (Feder et al., 1996; Wheeler and Wong, 2012). According to its variable molecular weight, HSPs are organized in various subfamilies: small HSPs, HSP40, HSP70, HSP90 and HSP110 (Lindquist and Craig 1988). The HSPs with a molecular weight, MW ~ 70 kDa comprise a set of functionally related molecules known as HSP68, HSP70, HSC70 (cognate heat shock protein), HSP75 (mortalin or mitochondrial heat shock protein) and HSP78 (BiP: binding immunoglobulin or endoplasmic reticulum heat shock protein); all characterized by a highly conserved N-terminal ATPase domain (Wang et al., 2021). In general, HSP70 functions in buffering against intrinsic and extrinsic stress (Murphy, 2013). The roles of HSP70 are variable and specific functions are related to specific stressors. The HSP70 family promotes protein folding and proteastasis, participates in the degradation of misfolded sequences and intervenes in translocation, and prevents deleterious aggregation (Mayer and Bukau, 2005; Li et al., 2025). Additionally, essential functions in response to environmental stress such as heat, hypoxia, pathogen infection, exposure to organochlorines and heavy metals, physiological stress and immune responses have also been observed (Moseley, 1997; Mosser et al., 1998; Tsan and Gao, 2009; Deane and Woo, 2011; Chen et al., 2018; Shi et al., 2020). In molluscs, HSP70 proteins are localized in cytoplasm, nuclei, endoplasmic reticulum and mitochondrion and each subcellular location is determined by specific motifs (Rosenzweig et al., 2019).

The HSP70 proteins are classified into two distinct groups based on their function and regulation patterns: constitutive expressed heat shock proteins (HSCs), which are expressed in the cells under normal conditions, and inducible heat shock proteins (HSPs), which are expressed under specific stress conditions (Boone and Vijayan, 2002). In general, HSP70 contains three main conserved domains: N-terminal ATPase domain (~ 40 kDa), also called the nucleotide-binding domain (NBD); it controls the interaction of the chaperone with the client protein. A second substrate-binding domain (SBD) (~ 18 kDa),

binds hydrophobic regions of client proteins during early folding. And lastly, a third C-terminal domain with variable length (~10 kDa) (Nikolaidis and Nei, 2004; Daugaard et al., 2007; Melikov et al., 2024). Until now, HSP70 in terrestrial molluscs has been predominately studied via Western blot in Mediterranean species (e.g. Di Lellis et al., 2012, 2014; Dieterich et al., 2013, 2015; Staikou et al., 2024). The development of genomic technologies and the recent increasing availability of whole genome sequences of mollusc species, enables us to assess the HSP70 gene family within an evolutionary context and its role in the adaptation to different environments (e.g. Hu et al., 2022; Lu et al., 2024).

In the Neotropical Cuban archipelago, the genus *Polymita* Beck, 1837, includes six endemic tree snail species: *P. muscarum*, *P. versicolor*, *P. venusta*, *P. sulphurosa*, *P. brocheri* and *P. picta* (González-Guillén, 2021) all inhabiting East Cuba to northeast of Sabana-Camagüey (de la Torre, 1950). The "polimitas" are admired worldwide because of their remarkable shell colour polymorphism (Alfonso y Berovides, 1993), arousing the interest of evolutionary biologists, conservationists and collectors. They were recently awarded as Mollusc of the Year 2022 (Mayer-Bömoser et al., 2022). All *Polymita* species are considered threatened (Maceira-Filgueira, 2016) and listed in Appendix I by the Convention on International Trade in Endangered Species (CITES, 2022).

Since its description as *Helix* (*Polymita*) *picta* (Born, 1778), this charismatic species has received growing research interest from different disciplines: taxonomy, behaviour, reproductive anatomy, shell polymorphism and ecology (e.g. Andrews, 1932; de la Torre, 1950; Moreno, 1950; Berovides et al., 1986; Alfonso and Berovides, 1987, 1989a, 1989b; Berovides and Alfonso, 1992; Berovides et al., 1986). Recently, its shell's luminance was quantified and its potential link to thermoresistance was discussed (Gordillo-Pérez et al., 2025). Simultaneously, the full sequencing of the mitochondrial and nuclear genes, provided new insights into its phylogeny (Lewis et al., 2025; Reyes-Tur et al., 2025).

In a recent study on six Cuban species of land snails found that ecological niche models predicted a loss of favourable habitats for all species by 2050 (Martínez y Hernández 2017). Besides, Mancina et al. (2017), conducted similar research on the six *Polymita* species, reporting habitat decrease in all cases and the vulnerability status of the genus in three hypothetical climate change scenarios for 2050. Despite the effort made in research and conservation of Cuban land snails, there are still no molecular studies investigating the resilience of these animals to climatic variability.

The aim of this paper is to conduct a genome-wide identification of the 70-kDa heat shock

protein (HSP70) family in *Polymita picta*, considering its role in thermal stress response and phenotypic plasticity as key mechanisms of climate-change adaptation. As far as we know, this research explores for the first time an HSP family at genome level in a Neotropical terrestrial gastropod.

MATERIAL AND METHODS

Genome-wide identification of HSP70 family members in the *Polymita picta* genome

The identification of HSP70 proteins in the *P. picta* genome was based on sequence homology and conserved domain analysis following methodologies carried out in similar studies (Hu et al., 2022; Lu et al., 2024). The genome of *Polymita picta* was obtained from NCBI (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1250545), and a genome-wide TBLASTN search was performed using all available mollusc HSP sequences from NCBI (1,064 sequences). The resulting TBLASTN hits (78,994) were filtered in R (R Core Team, 2022) to retain sequences with an E-value ≤ e-5.

To verify the presence of conserved protein domains in the candidate sequences, a hidden Markov model (HMM) profile for HSP70 (PF00012) was downloaded from the Pfam protein family database (https://pfam.xfam.org/), and processed using the software HMMER 3.0. The presence of conserved HSP70 domains was further confirmed using the NCBI Conserved Domains Database (http://www.ncbi.nlm.nih.gov/cdd/), with an E-value of <0.0001 and all other parameters set to their default values.

Full-length sequence extraction and translation from genomic data

The 56 HSP70 protein fragments identified through HMM were re-aligned to the *P. picta* genome assembly (version v4665_2) in NCBI using BLAST+ v2.13.0 (TBLASTN) (Camacho et al., 2009). The corresponding BLAST hits were provided in BED format, containing genomic coordinates for each identified region. The BED file containing the genomic coordinates of the HSP70 hits was used to extract the corresponding DNA sequences from the genome. Each hit provided the start and end positions for sequence extraction. If the sequence was identified on the negative strand of the genome, it was reverse complemented to ensure proper reading frame translation. To identify the best open reading frame (ORF), we tested three possible reading frames for each extracted

sequence. This was done by translating the sequence from three different starting positions corresponding to the three potential frames. The number of stop codons ("*") was counted in each translated sequence. The reading frame with the fewest stop codons was selected as the optimal frame for translation. Sequences with internal stop codons were discarded, and the longest continuous sequence without stop codons was retained for further analysis. After selecting the optimal reading frame for each sequence, the corresponding protein sequence was translated (Supplementary Data 1: https://doi.org/10.5281/zenodo.16744586). These procedures were carried out in R software (R Core Team, 2022). We manually checked the R extraction using Integrative Genomic Viewer (IGV) ver. 2.19.3. (Thorvaldsdóttir et al., 2013). Each translated sequence was then annotated with its genomic coordinates (scaffold, start, end, and strand direction).

Physicochemical profiling of *Polymita picta* HSP70s

The physicochemical properties, including molecular weight (MW), number of amino acids, theoretical isoelectric point (pI), grand average of hydropathy (GRAVY), and instability index (II), were analysed using the ExPASy ProtParam tool. The signal peptide prediction in *P. picta* HSP70s (*Pp*HSP70) was carried out through the online tool Signal P 5.0 (http://services.healthtech.dtu.dk/service.php?SignalP-5.0. The conserved motifs in *Pp*HSP70 were analysed using the Multiple EM for Motif Elicitation online tool (http://meme-suite.org/meme/tools/meme).

Motifs analysis in *Polymita picta* HSP70s and comparative sequences

A total of 34 protein sequences were analysed for conserved motifs, including the 10 HSP70 paralogs from *P. picta* and additionally HSP70 sequences from the molluscan species *Biomphalaria glabrata*, *Ostrea edulis*, *Magallana gigas*, *Mya arenaria*, *Aplysia californica*, *Dreissena polymorpha* and *Crassostrea virginica* and one HSP70 from *Drosophila melanogaster*. For helping identifying subfamilies motifs, sequences annotated as HSP70 cytosolic, HSC70, HSP75 and HSP78 were included (Supplementary Data 2: https://doi.org/10.5281/zenodo.16744775). We selected the aforementioned species to enable orthologous-based comparisons of HSP genes with those of *Polymita*, prioritizing taxa with available reference sequences and leveraging the high evolutionary conservation

of HSPs (Wheeler and Wong, 2012).

Homology and gene duplication

Two criteria were considered to identify gene duplications: (i) sequence similarity greater than 70% and (ii) the shorter aligned sequence covering at least 70% of the longer one (Gu et al., 2002; Yang et al., 2008; Lu et al., 2024). The sequences were aligned using MAFFT v7 (https://mafft.cbrc.jp/alignment/server/index.html) and loaded into R using the Biostrings package. Metadata including sequence names, scaffolds, and strand orientation were parsed from the FASTA headers using regular expressions and stored in a structured data frame. A pairwise comparison between all sequences was conducted to compute percentage identity. A custom R function compared aligned amino acids for each pair and calculated identity as the proportion of matching positions over the total alignment length (R Core Team, 2022). In all pairs selection pressure rates for duplicated genes were estimated using Ka/Ks_Calculator 3.0.

Phylogenetic analysis

For the phylogenetic analysis, the sequences of *P. picta* HSP70 that had been identified were re-aligned to the same set of orthologous used for motif identification, ensuring consistency across comparative analyses. The major heat shock 70 kDa protein from the model species *Drosophila melanogaster* was used as an out-group. The LG+G4 amino acid substitution model was selected for phylogenetics through the ModelFinder function (-m MFP) (Kalyaanamoorthy et al., 2017) followed by tree inference using IQ-TREE v2.3.6 (Nguyen et al., 2015; Minh et al., 2020). Branch support was estimated through ultrafast bootstraps (Hoang et al., 2018) and Shimodaira-Hasegawa like approximate likelihood ratio test (SH~aLRT) (Guindon et al., 2010). In both algorithms 10 000 replications were performed. Nodes with an ultrafast bootstrap value ≥ 95 and a likelihood ratio ≥80 were considered as well as supported (Cruz-Laufer et al., 2025). Finally, phylogenetic trees were visualized using FigTree v1.4.4.

RESULTS

Genome-wide identification and physicochemical profiling of heat shock proteins 70 kDa (HSP70) in *Polymita picta*

In this study, 56 partial sequences of HSP70 were identified across the genome of *P. picta* using a TBLASTN search with 1,064 mollusc sequences from NCBI. After screening using HMM and CDD-NCBI, 22 sequences (39.2%) showed significant similarity to a query containing the ATPase-like domain of the ASKHA superfamily (Acetate and Sugar Kinases/Hsc70/Actin), while the remaining sequences (60.8%) were associated with the PTZ00009 domain, which is also annotated for heat shock 70 kDa proteins.

Heat shock proteins were identified at 10 unique positions in the genome and were named consecutively *Pp*HSP70_1 to *Pp*HSP70_10. Information on the identifier is followed by the scaffold where it is located, the start and end coordinates, and the strand, e.g., *Pp*HSP70_1_ptg000811l_1: 167165-168904 (-) (Supplementary Data 3: https://doi.org/10.5281/zenodo.16744894).

The ten HSP70 protein sequences identified were distributed across five scaffolds (Fig. 1). *Pp*HSP70_1 to *Pp*HSP70_6 were closely located on both strands of ptg000811l_1 (only *Pp*HSP70_5 and *P*pHSP70_6 on the positive strand), while the remaining four sequences were distributed across the positive strand of scaffolds ptg004791l_1, ptg010661l_1, ptg018870l_1, and ptg018978l_1.

Figure 1. Scaffold distribution of *Polymita picta* heat shock proteins 70 kDa (HSP70).

The 10 different identified HSP70 proteins consist of amino acid sequences ranging from 635 to 672, with an average of approximately 641 residues (Table 1). The molecular weights varied between 69,305.92 Da and 73,549.12 Da, with an average of 69,725.06 Da. The isoelectric points (pI) were consistently low, ranging from 5.58 to 5.80 (mean pI = 5.67), indicating a generally acidic nature. Additionally, all proteins exhibited negative GRAVY values (–0.47 to –0.38; mean = –0.44), suggesting overall hydrophilicity. The instability indices for all sequences were below 40, ranging from 28.23 to 30.69 (mean = 29.24), classifying them as stable proteins. The aliphatic index values ranged from 79.43 to 84.32 (mean = 81.15), supporting moderate to high predicted thermostability (Table 1).

Table 1. Physico-chemical parameters of *Polymita picta* heat shock protein 70 kDa.

Sequence	Number of Amino Acids	Molecular Weight	Isoelectric Point	GRAVY	Instability Index	Aliphatic Index
*Pp*HSP70_1	637	69536.36	5.66	-0.44	28.81	81.46
*Pp*HSP70_2	637	69550.26	5.58	-0.46	30.69	81
*Pp*HSP70_3	645	70355.17	5.59	-0.45	29.52	80.76
*Pp*HSP70_4	637	69607.35	5.66	-0.47	29.84	80.55
*Pp*HSP70_5	637	69580.28	5.58	-.0.46	30.2	80.55
*Pp*HSP70_6	637	69607.35	5.66	-0.47	29.84	80.55
*Pp*HSP70_7	672	73549.12	5.8	-0.38	29.44	84.32
*Pp*HSP70_8	635	69305.92	5.76	-0.47	28.23	79.43
*Pp*HSP70_9	641	69925.64	5.69	-0.44	28.44	80.34
*Pp*HSP70_10	635	69317.22	5.77	-0.42	30.34	83.56

All *P. picta* HSP70 proteins appear to be primarily localized in the cytoplasm, with prediction scores ranging from 0.47 to 0.78 (mean = 0.73) (Supplementary Data 4: https://doi.org/10.5281/zenodo.16745227). However, two proteins, *Pp*HSP70_3 and *Pp*HSP70_8, showed nuclear localization signals (NLS), implying potential dual localization in both the cytoplasm and nucleus. Additionally, *Pp*HSP70_7 displayed a peroxisomal targeting signal, with a relatively lower cytoplasmic probability (0.47), but higher probabilities for other compartments such as the endoplasmic reticulum (0.52), Golgi apparatus (0.47), and lysosome/vacuole (0.46). Probabilities for mitochondrial or cell membrane localization were generally low across all proteins (mean < 0.34).

Signal peptide prediction using SignalP 5.0 indicated that none of the *Pp*HSP70 proteins possess a classical N-terminal signal peptide for secretion via the Sec/SPI pathway (General secretion pathway / Signal Peptidase I). All sequences were classified as "OTHER" with high probability (average OTHER probability = 0.9975 ± 0.0003) (Supplementary Data 5: https://doi.org/10.5281/zenodo.16745405). The SP (Sec/SPI)

scores were consistently low across all sequences (average SP probability = 0.0025 ±
0.0003), suggesting that these proteins are not secreted through the classical secretory
pathway.

Motif Analysis of HSP70 Family in *Polymita picta* and Comparative Sequences

A total of 21 motifs were identified across the dataset, revealing distinct differences in motif
architecture that correlate with subcellular localization and functional HSP70 subfamilies.
Notably, mitochondrial HSP70s (HSP75) shared a unique motif at the N-terminal region
(motif 21), which was absent from all other sequences. Both HSP75 and HSP78
(endoplasmic reticulum) displayed an additional motif in the IIA Actin-like ATPase
subdomain of NBD, which was not found in cytosolic HSP70s, including those identified in
P. picta. The three HSP70 cognate proteins (HSC70) lacked a conserved motif (motif 15)
in the SBD, which was consistently present in all inducible HSP70s, including all *P. picta*
paralogs and the *D. melanogaster* inducible HSP70.

Minor variations were detected in the height and sequence of two short motifs (~8 amino
acids), located near the IA Actin-like ATPase subdomain of NBD and in the central region
of the SBD. These changes correspond to single amino acid polymorphisms (SAPs).
Sequence alignment (Supplementary Data 6: https://doi.org/10.5281/zenodo.16746341),
confirmed that these SAPs are scattered and mostly conservative. For example, in the C-
terminal region, just before the EEVD sequence (see motif 16 in Fig. 2), an insertion of
four or five residues was detected between positions 638-642 in PpHSP70_7 (SGSSR),
*Pp*HSP70_8 (AHSQ), *Pp*HSP70_9 (AGSQ), and *Pp*HSP70_10 (TGGHS). Additionally, a
residue deletion at position 635 is observed in *Pp*HSP70_10, and a second deletion at
position 627 appears in *Pp*HSP70_7, *Pp*HSP70_8, and *Pp*HSP70_10. The N-terminal
region displayed slight variability in the residues following the initiating methionine:
*Pp*HSP70_1 starts with MA, *Pp*HSP70_2 with MR, *Pp*HSP70_5 with MT, and *Pp*HSP70_6
with MS. Notably, *Pp*HSP70_10 lacks the first five amino acids.

Figure 2. Distribution of conserved motifs of different subfamilies of heat shock proteins 70 kDa (HSP70) in *Polymita picta* and comparative sequences.

Sequence Identity and Evolutionary Constraints

We computed pairwise amino acid identity percentages to assess the degree of similarity among the identified HSP70 paralogous proteins in *P. picta*. As shown (Fig. 3), proteins located within the same scaffold (e.g., ptg000811l_1) generally exhibit very high identity (>97%). In contrast, lower identity values were observed for sequences located in different scaffolds. The minimum identity, 83.26 %, was found between *Pp*HSP70_7 and *Pp*HSP70_10.

Figure 3. Matrix of identical positions (pID) percentages between *Polymita picta* heat shock protein 70 kDa (HSP70) sequences.

The Ka/Ks ratios for the various HSP70 sequence pairs were calculated to assess selective pressure and potential evolutionary divergence. The results, presented in (Supplementary Data 7: https://doi.org/10.5281/zenodo.16751419), include the comparison of 45 pairwise combinations of HSP70 sequences in *P. picta*. As a general trend, all calculated Ka/Ks ratios are below 1. For example, the sequence pair *Pp*HSP70_1 vs *Pp*HSP70_10 exhibited the lowest ratio of 0.0127. In contrast, the pairs *Pp*HSP70_3 vs *Pp*HSP70_4 and *Pp*HSP70_3 vs *Pp*HSP70_6 both exhibited the highest Ka/Ks ratio of

0.1642. Notably, the pair *Pp*HSP70_4 vs *Pp*HSP70_6 returned a NaN value due to the sequences being identical but located at different loci.

Phylogenetic analysis of *Polymita picta* HSP70 and comparative sequences

The phylogenetic analysis revealed the presence of two well defined major clades (Fig. 4). The first clade grouped all *P. picta* sequences together with four cytosolic HSP70 sequences from the pulmonate *B. glabrata*, with strong statistical support (SH-aLRT/UFBOOT = 100/100). Within this clade, two main branches are identified: one branch contains exclusively the four *B. glabrata* sequences, supported by very high values (SH-aLRT/UFBOOT = 100/100 and 98.8/100) and the other branch grouping the ten *Pp*HSP70 sequences. The internal relationships among the *P. picta* sequences display heterogeneous support. In particular, the placement of PpHSP70_10 relative to the other paralogs exhibited low resolution. Similarly, the lowest bootstrap values were observed at the nodes grouping *Pp*HSP70_1, *Pp*HSP70_3, *Pp*HSP70_4, and *Pp*HSP70_6. By contrast, the remaining nodes within the paralogs clade showed better resolved relationships. The second major clade comprised sequences from all other molluscan species corresponding to HSC, HSP75, and HSP78 also with high support (SH-aLRT/UFBOOT = 99.4/100). Within this clade, a secondary branch grouped other cytosolic HSP70 sequences, although their internal relationships remained unresolved.

Figure 4. Phylogenetic analysis of *Polymita picta* heat shock protein 70 kDa (HSP70) amino acid sequences and orthologous. The node grouping *P. picta* sequences is highlighted in yellow.

DISCUSSION

To ultimately provide policy makers with sound advice on the protection of vulnerable fauna in a changing climate, more specifically in the sub-tropics, we focused on the HSP70 protein family, given its promising roles in coping with the physical challenges caused by climate change, of the Cuban painted snail *Polymita picta*. Therefore, we performed a genome-wide identification of the chaperones and analysed its molecular characteristics, variation and evolution.

The genome-wide identification of HSP70s in *Polymita picta* and its potential relationship with thermoresistance

The heat shock proteins of 70 kDa (HSP70) is a well characterized protein family playing a

crucial role in the cells' housekeeping and stress response, from bacteria to mammals (Karlin and Brocchieri, 1998; Clerico et al., 2019; Rosenzweig et al., 2019). However, until now related studies in molluscs mainly focused on species of economic interest being a food source or a threat as an invasive species (Zhang et al., 2012; Cheng et al., 2016; Hu et al., 2019, 2022; Lu et al., 2024). Unfortunately, studies addressing the HSP70s in land snails from a genomic angle are totally lacking. This evidently complicates understanding the evolution and diversity of these essential molecules.

Besides many new research opportunities, the recent *P. picta* full genome sequencing (Reyes-Tur et al., 2025), facilitated this first HSP70 screening. The genome assembly contained 1.85 Gbp with 22628 contigs and a contig N50 of 124.2 Kbp. Moreover, the genome completeness corresponds to 88.4 % and the high heterozygosity, 2.41 % are as expected (Vendrami et al., 2020; Bean et al., 2022; Gundappa et al., 2022; Reyes-Tur et al., 2025). Therefore, we interpreted our results at scaffold level and conservatively (to avoid identifying alleles as duplications). We identified 10 different genes coding for full length heat shock proteins The number of HSP70 variants retrieved, is comparable with those found in several fish species; or even the sea cucumber *Apostichopus japonicus* (Xu et al., 2018; Ma et al., 2020; Gao et al., 2018, 2022; Zheng et al., 2023). Strikingly, in bivalve species the number of observed HSP70s proteins, including several tandem duplicated sequences, is manyfold e.g. 65 genes in the clam *Telligarga granosa* (Kim et al., 2024) or 113 variants in the pacific oyster *M. gigas* (Lu et al., 2024).

The 10 HSP70 identified in this study were distributed in five different scaffolds. Four sequences, *Pp*HSP70_7 to *Pp*HSP70_10 were located in four of the five different scaffolds, which suggest they represent putative dispersed duplications, being majorly a product of transposition (Wang, 2011). The other sequences, from *Pp*HSP70_1 to *Pp*HSP70_6 are located in the same scaffold in each other's vicinity (< 10 bp apart); four of these in the negative strand and two in the positive one. Most likely, this study included tandem duplications (Reams et al., 2015), a common phenomenon well described in mollusc genomes (Zheng et al., 2022; Lu et al., 2024). Consequently, providing them with functional variation, relevant for adapting to different and/or changing environments, coping with diseases and potentially useful in conservation management (Hawkins et al., 2024; Sutherland et al., 2024, 2025).

Overall, *P. picta* HSP70 proteins show high conservation in size, with an average of 641 residues, low isoelectric points indicating their acidic nature and negative GRAVY values, which suggest their hydrophilic character. Therefore, all molecules can be classified as

stable proteins. Additionally, due to their aliphatic indexes we consider them to have a moderately to high thermostability. These physico-chemical characteristics are typical of the HSP70 family (Dayrit et al., 2023; Singh et al., 2025).

The 10 identified sequences are most likely located in the cytoplasm, however, two proteins, *Pp*HSP70_3 and *Pp*HSP70_8 also showed signals corresponding to a nuclear position suggesting a potential dual function both in cytoplasm and nucleus. Moreover, one sequence, *Pp*HSP70_7 may have a broader subcellular distribution, showing target signals for peroxisome, endoplasmic reticulum, Golgi apparatus and lysosome/vacuole. This possibly reflects a specialized or a distinct stress-related function from the other members (Daugaard et al., 2007). *Polymita picta*'s HSP70 proteins therefore can be considered as predominantly cytoplasmic chaperones, with potential minor roles in other organelles for a subset of sequences (Peyretaillade et al., 1998; Kaneko, 2022; Hasnain et al., 2022; Yadav et al., 2019). Importantly, our results confirm that none of the ten sequences is secreted through the classical secretory pathway (Sec/SPI), supporting our conclusion that all *Pp*HSP70 in *P. picta* are primarily cytoplasmic chaperones.

Motifs analysis of HSP70 family in *Polymita picta* and comparative sequences

The identified motifs correspond to canonical HSP70 domains, supporting functional conservation across the family despite gene duplications. The unique motifs in organellar subfamilies and the absence of specific ones for HSC70s highlight the structural divergence between cytosolic and non-cytosolic members of the HSP70 family. For the case of this chaperone family, major differences are observed at C-terminal. The three HSP70 cognate proteins (HSC70) included in our analysis lacked a conserved motif (motif 15) in the SBD, consistently present in all inducible HSP70s, including all *P. picta* paralogs and the *D. melanogaster* inducible HSP70, what suggests functional divergence between constitutive and inducible variants. Some identified changes among *P. picta* paralogous are based on single amino acid polymorphisms (SAPs) that do not disrupt motif presence but may indicate allelic or functional divergence. With the exception of *Pp*HSP70_8, which lacks the motif 16 and exhibits some differences, potentially related to the exon-intron structure of the C-terminal region (see distance between motifs 8 and 15 in *Pp*HSP70_3), the motif patterns are conserved. This suggests that the identified chaperons in *P. picta* have unique biological functions that vary discretely (Xu et al., 2018; Zhang et al., 2013; Huang et al., 2018). Notably, *Pp*HSP70_10 lacks the first five amino acids, potentially

indicating an alternative start codon but the sequence possesses the full first identified motif on N-terminal (motif 4).

Despite these minor substitutions, no major insertions or deletions were detected, and all sequences retained the expected domain architecture and length consistent with functional HSP70 proteins. The preservation of key domains such as the nucleotide-binding domain, substrate-binding domain, and the C-terminal EEVD motif suggests that all sequences maintain their chaperone function. Taken together, the high degree of sequence and motif conservation, combined with modest SAPs, supports the hypothesis that the HSP70 sequences in *P. picta* may represent recent gene duplications or allelic variants within a smaller number of loci in comparison to marine molluscs species. Terrestrial ectotherms are generally less able to acclimatize and exhibit smaller increases in physiological rates than aquatic ones (Noble et al., 2025). In any case, further genomic, transcriptomics, and expression analyses are necessary to elucidate the functional relevance of the observed polymorphisms. Motif-based comparisons between *Polymita* HSPs sequences and orthologs provide preliminary context but should be interpreted with caution. Our analyses were necessarily constrained by the availability and quality of reference sequences and functional inferences from motif presence/absence remain indirect. Accordingly, we use the existing experimental literature to contextualize putative roles and to guide hypotheses about protein function. More functionally informative comparisons - specially with orthologs from molluscan lineages most closely related to *Polymita* (e.g. terrestrial gastropods) - will depend on the continued expansion of high quality genomic and transcriptomics resources for these taxa.

Sequence Identity and Evolutionary Constraint

The amino acid sequences of *Pp*HSP70_1 to *Pp*HSP70_6 found in the same scaffold demonstrate high levels of homology (>97%). This suggests either recent gene duplications, as previously proposed, or strong functional constraints. Despite the possible presence of allelic variants among these *P. picta* sequences, tandem duplication of HSP70 genes has been reported in several molluscs species. For example HSP70 genes are distributed across all 19 chromosomes of *M. mercenaria*, and the HSPa12 gene appears to be duplicated 39 times on chromosome 7 (Hu et al., 2022). A similar high-density collinear gene block has been observed in the chromosome 14 of *C. sinensis* (Wei et al., 2020), as well as in other bivalves belonging to the subclass Pteriomorpha (marine

species) and the order Veneroida (marine and fresh water species) (Zhang et al., 2012; Cheng et al., 2016; Powell et al., 2018; Peng et al., 2020; Gao et al., 2022). Lower identity values were found in the sequences from *Pp*HSP70_7 to *Pp*HSP70_10, but still presenting more than 83 % of homology, suggesting older duplication events followed by varying degrees of sequence divergence. It is understood that duplications can exhibit homology values ranging from 70 to 100 % in the case of HSP70 proteins (Lu et al., 2024).

All Ka/Ks ratios between *Pp*HSP70 pairs are below 0.16, indicating strong purifying selection. The duplicated genes typically remain under purifying selection during the early stages of evolution (Kondrashov et al., 2002), to ensure normal cellular function, while those that respond properly to stress tend to be amplified (Evgen'ev et al., 2014; Metzger et al., 2016). It is well known, gene duplication provides material for evolutionary novelty and tandem duplications is recognized as an important driving force in this process (Roth et al., 2007; Chen et al., 2013; Long et al., 2013; Singh et al., 2025). Generally, variability in HSP70 gene copy number is assumed to reflect regulatory differences, particularly in relation to the factor temperature. Furthermore, genomic variability resulting from gene duplications has been reported to directly influence physiological responses to environmental fluctuations (Zhang et al., 2012). Based on the observed homology among *P. picta* HSP70 sequences and the selective pressure ratios, it appears that the gene duplication events in this species are recent and still subject to purifying selection

Our preliminary genomic evidence suggests that *P. picta* may harbor an expanded HSP70 gene family relative to molluscs that routinely experience broad thermal regimes (e.g. intertidal oysters), yet *Polymita* inhabits comparatively thermally stable terrestrial microhabitats. To elucidate how the architecture of the *Pp*HSP70 family relates to the species's life history, it is necessary to consider both, the evolutionary trajectory of *Polymita* in prehistoric Eastern Cuba and the evolution of its habitats. Historically, the archipelago has experienced profound paleoclimatic fluctuations since early Miocene, when the origin of its modern fauna is estimated (MacPhee & Iturralde-Vinent, 2005). Episodes of tectonic uplift, Pleistocene sea-level oscillations and recurrent drought-humid cycles recorded in speleothem and lacustrin archives (Fensterer et al., 2013; Peros et al., 2015, 2017) likely shaped population connectivity, niche distributions and thermotolerance in ancestral *Polymita* lineages. It is therefore expected that the basal architecture of the HSP70 family was modeled under these long term selective regimes. Moreover, the evolution of thermal stress genes has also been linked to ecological drivers beyond temperature per se (Staikou et al., 2024). Future functional studies that profile differential

expression of individual paralogs across environmental gradients will be crucial to disentangle the relative contribution of ancient versus ongoing selective pressures to *Polymita*'s HSP70 complement.

Conservation perspectives

In terms of conservation, over the past centuries, Cuba has undergone dramatic habitat transformation, including the loss of ~ 85 % of native forests within seven decades (del Risco, 1995) and persistent deforestation-driven fragmentation of habitats. Although recent reports indicate a slight recovery of forest cover, ~ 32.2 % (ONEI, 2025), climatic stressors are intensifying: annual mean temperatures are rising, heat extremes are breaking historical records and rainy season shows increased thermal variability (Centella et al., 1999; Febles, 2009; Fonseca-Rivero et al., 2024; Harbott et al., 2023; Campbell, 2023). Moreover, Cuba's protected-area system covers only 15.94% of the national territory (ONEI, 2025), in a region that forms part of a biodiversity hotspot. Protected areas do not always encompass the full extent of species' distributions (Mancina and Cruz, 2017), and interactions with surrounding ecosystems remain poorly investigated. In addition to this, the country has experienced deep political and economic crises in recent decades, resulting in a shortage of vital resources and hunger, particularly in rural areas. In response, the use of firewood for cooking and charcoal production as surviving strategies, lack of and deficiencies in management plans have become a direct threat to the forest ecosystems (Maal-Bared, 2006; Alonso and Clark, 2015).

In an adaptive context, evidence of an expanded HSP70 chaperone family in *Polymita* does not, by itself, imply resilience to climate change or heat stress. Gene duplication is common but only rarely yields clear adaptive gains and increase in copy number can be neutral or even deleterious (Qian and Zhang, 2014). Classic experiments with *Drosophila* further show that adding HSP70 copies can improve survival under severe heat stress yet reduce performance under benign or moderately stressful conditions -underscoring that "more copies" does not straightforwardly translate into greater thermal tolerance- (Feder et al., 1996; Roberts et al., 2000). More broadly, thermal tolerance itself limited plasticity across ectotherms, cautioning against equating gene-family size with climate resilience (Gunderson & Stillman, 2015).

The finds presented in this research regarding *P. picta* HSP70, are the result of a genome-wide identification in one individual used for the genome sequencing (Reyes-Tur at al.,

2025); so, intra and interspecific variations in the HSP70 system on the genus and the study of other stress related molecules should be taken into account for a better understanding of the thermoresistance molecular mechanisms. We understand that more in-depth experimental research is needed to improve our understanding of *Polymita*'s resilience to global warming in the Anthropocene, and more specifically in the context of Cuba. The early detection of "more thermoresistant" haplotypes could constitute a strategy for genetic conservation, population reinforcement and re-introduction.

REFERENCES

AJAYI, O. O., PETERS, S. O., DE DONATO, M., SOWANDE, S. O., MUJIBI, F. D., MORENIKEJI, O. B., THOMAS, B. N., ADELEKE, M. A. & IMUMORIN, I. G. 2018. Computational genome-wide identification of heat shock protein genes in the bovine genome. F1000Research, 7, 1504.

ALFONSO, M. & BEROVIDES, B. 1987. Genética ecológica de *Polymita picta roseolimbata* (Mollusca: Pulmonata) en un agroecosistema de la región de Maisí. Ciencias Biológicas, 18, 33-44.

ALFONSO, M. & BEROVIDES, V. 1989. Fenotipos de los morfos de color y bandas de la concha en *Polymita picta roseolimbata* (Mollusca: Pulmonata) de Maisí (Cuba). Revista Biología, 3, 213-222.

ALFONSO, M. & BEROVIDES, V. 1989. Variación de los patrones de color y bandas de la concha en el género *Polymita* (Mollusca: Pulmonata). Ciencias Biológicas, 21-22.

ALFONSO, M. & BEROVIDES, V. 1993. Conservation problems of land snails in Cuba. Tentacle, 3, 20-23.

ALONSO, G. & CLARK, I. 2015. Cuba confronts climate change. Medicc review, 17, 10-13.

AMARO-VALDÉS, S. & ÁLVAREZ, V. B. 2012. Lista roja de la fauna cubana, Editorial AMA.

ANDREWS, E. 1932. Habitat of a Cuban snail, *Polymita picta*, Born. The Nautilus, 46, 22-25.

ARENAL, I. M., MAYO, A. H., INFANTE, J. C., PIT, L. L., FRAGUELA, E. G., RIVERA, C. F., HERNÁNDEZ, A. G., ZALDÍVAR, B. V., DE LA MAZA SANTANA, D. G. & ÁLVAREZ, M. M. 2024. Estado del Clima en Cuba 2023. Resumen ampliado. Revista cubana de meteorología, 30, 1-16.

ASTOR, T., VON PROSCHWITZ, T., STRENGBOM, J., BERG, M. P. & BENGTSSON, J. 2017. Importance of environmental and spatial components for species and trait composition in terrestrial snail communities. Journal of Biogeography, 44, 1362-1372.

BEAN, T. P., TANGUY, A., PEÑALOZA, C., GUNDAPPA, M. K., BOUTET, I., HOUSTON, R. D., MACQUEEN, D. J. & BOUDRY, P. 2022. Two parallel chromosome-level reference genomes to support restoration and aquaculture of European flat oyster *Ostrea edulis*. Evolutionary applications, 15, 1709-1712.

BECK, C. H. H. 1837. *Index molluscorum praesentis aevi musei principis augustissimi* Christiani Frederici. (No Title).

BEROVIDES, A. V. 1986. Polimorfismo genético de *Polymita picta roseolimbata* Torre, 1950 en la region de Maisí. Cuba Carib J Sci, 22, 179-189.

BEROVIDES, V. & ALFONSO, M. 1992. Evidencias experimentales de la selección climática en *Polymita picta roseolimbata* (Gastropoda: Fruticicollidae) de Maisí, Cuba. Ciencias Biológicas, 25, 1-8.

BHAGAT, J., INGOLE, B. & SINGH, N. 2016. Glutathione S-transferase, catalase, superoxide dismutase, glutathione peroxidase, and lipid peroxidation as biomarkers of oxidative stress in snails: A review. Invertebrate Survival Journal, 13, 336-349.

BOONE, A. N. & VIJAYAN, M. M. 2002. Constitutive heat shock protein 70 (HSC70) expression in rainbow trout hepatocytes: effect of heat shock and heavy metal exposure. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology, 132, 223-233.

BORN, I. 1778. *Index rerum naturalium Musei Caesarei Vindobonensis*. Verzeichniss der Natürlichen Seltenheiten des KK Naturalien Kabinets zu Wien. Erster Theil, Schalthiere. Museum of Comparative Zoology--Biodiversity Heritage Library digitization project.

BORO, R. C. & JAYAKUMAR, S. 2020. Studies on enzymes Superoxide dismutase and Catalase in midgut of *Samia ricini* under thermal stress condition. Journal of Entomological Research, 44, 87-91.

CAMACHO, C., COULOURIS, G., AVAGYAN, V., MA, N., PAPADOPOULOS, J., BEALER, K. & MADDEN, T. L. 2009. BLAST+: architecture and applications. BMC bioinformatics, 10, 421.

CAMPBELL-LENDRUM, D., NEVILLE, T., SCHWEIZER, C. & NEIRA, M. 2023. Climate change and health: three grand challenges. Nature medicine, 29, 1631-1638.

CENTELLA, A., GUTIÉRREZ, T., LIMIA, M. & JASPE, R. R. 1999. Climate change scenarios for impact assessment in Cuba. Climate Research, 12, 223-230.

CHEN, B., FEDER, M. E. & KANG, L. 2018. Evolution of heat-shock protein expression underlying adaptive responses to environmental stress. Molecular ecology, 27, 3040-3054.

CHEN, J., HUANG, Q., GAO, D., WANG, J., LANG, Y., LIU, T., LI, B., BAI, Z., LUIS GOICOECHEA, J. & LIANG, C. 2013. Whole-genome sequencing of *Oryza brachyantha* reveals mechanisms underlying *Oryza* genome evolution. Nature Communications, 4, 1595.

CHENG, J., XUN, X., KONG, Y., WANG, S., YANG, Z., LI, Y., KONG, D., WANG, S., ZHANG, L. & HU, X. 2016. Hsp70 gene expansions in the scallop *Patinopecten yessoensis* and their expression regulation after exposure to the toxic dinoflagellate *Alexandrium catenella*. Fish & shellfish immunology, 58, 266-273.

CHEVIN, L.-M. & HOFFMANN, A. A. 2017. Evolution of phenotypic plasticity in extreme environments. Philosophical Transactions of the Royal Society B: Biological Sciences, 372, 20160138.

CLERICO, E. M., MENG, W., POZHIDAEVA, A., BHASNE, K., PETRIDIS, C. & GIERASCH, L. M. 2019. Hsp70 molecular chaperones: multifunctional allosteric holding and unfolding machines. Biochemical Journal, 476, 1653-1677.

CNAP 2013. Plan del Sistema Nacional de Áreas Protegidas 2014–2020. Ministerio de Ciencias Tecnología y Medio Ambiente, La Habana, Cuba.

CRUZ-LAUFER, A. J., VANHOVE, M. P., BACHMANN, L., BARSON, M., BASSIROU, H., BITJA NYOM, A. R., GERAERTS, M., HAHN, C., HUYSE, T. & KASEMBELE, G. K. 2025. Adaptive evolution of stress response genes in parasites aligns with host niche diversity. BMC biology, 23, 10.

DAUGAARD, M., KIRKEGAARD-SØRENSEN, T., OSTENFELD, M. S., AABOE, M., HØYER-HANSEN, M., ØRNTOFT, T. F., ROHDE, M. & JAATTELA, M. 2007. Lens epithelium-derived growth factor is an Hsp70-2 regulated guardian of lysosomal stability in human cancer. Cancer research, 67, 2559-2567.

DAUGAARD, M., ROHDE, M. & JÄÄTTELÄ, M. 2007. The heat shock protein 70 family: Highly homologous proteins with overlapping and distinct functions. FEBS letters, 581, 3702-3710.

DAYRIT, G. B., BURIGSAY, N. P. F., VERA CRUZ, E. M. & SANTOS, M. D. 2023. In silico characterization, homology modeling and structure-based functional annotation of Nile Tilapia (*Oreochromis niloticus*) Hsp70 and Hsc70 proteins. bioRxiv, 2023.12. 27.573401.

DE LA TORRE, C. 1950. El Género *Polymita*. Memorias de la Sociedad Cubana de Historia Natural "Felipe Poey", 20, 1-20.

DEANE, E. E. & WOO, N. Y. 2011. Advances and perspectives on the regulation and expression of piscine heat shock proteins. Reviews in Fish Biology and Fisheries, 21, 153-185.

DI LELLIS, M. A., SEIFAN, M., TROSCHINSKI, S., MAZZIA, C., CAPOWIEZ, Y., TRIEBSKORN, R. & KÖHLER, H.-R. 2012. Solar radiation stress in climbing snails: behavioural and intrinsic features define the Hsp70 level in natural populations of *Xeropicta derbentina* (Pulmonata). Cell Stress and Chaperones, 17, 717-727.

DI LELLIS, M. A., SEREDA, S., GEISSLE, A., PICOT, A., ARNOLD, P., LANG, S., TROSCHINSKI, S., DIETERICH, A., HAUFFE, T. & CAPOWIEZ, Y. 2014. Phenotypic diversity, population structure and stress protein-based capacitoring in populations of *Xeropicta derbentina*, a heat-tolerant land snail species. Cell Stress and Chaperones, 19, 791-800.

DIETERICH, A., FISCHBACH, U., LUDWIG, M., DI LELLIS, M., TROSCHINSKI, S., GÄRTNER, U., TRIEBSKORN, R. & KÖHLER, H.-R. 2013. Daily and seasonal changes in heat exposure and the Hsp70 level of individuals from a field population of *Xeropicta derbentina* (Krynicki 1836)(Pulmonata, Hygromiidae) in Southern France. Cell Stress and Chaperones, 18, 405-414.

DIETERICH, A., TROSCHINSKI, S., SCHWARZ, S., DI LELLIS, M., HENNEBERG, A., FISCHBACH, U., LUDWIG, M., GÄRTNER, U., TRIEBSKORN, R. & KÖHLER, H.-R. 2015. Hsp70 and lipid peroxide levels following heat stress in *Xeropicta derbentina* (Krynicki 1836)(Gastropoda, Pulmonata) with regard to different colour morphs. Cell Stress and Chaperones, 20, 159-168.

ENRIQUE, D. R. 1995. Los bosques de Cuba: su historia y características, La Habana, Cuba, Editorial Científico-Técnica.

EVGEN'EV, M. B., GARBUZ, D. G. & ZATSEPINA, O. G. 2014. Heat shock proteins and whole body adaptation to extreme environments, Springer.

FEBLES, G. 2009. The global climatic change and its impacts in Cuba. Actions for the future. Cuban Journal of Agricultural Science, 43.

FEDER, M. E., CARTAÑO, N. V., MILOS, L., KREBS, R. A. & LINDQUIST, S. L. 1996. Effect of engineering Hsp70 copy number on Hsp70 expression and tolerance of ecologically relevant heat shock in larvae and pupae of *Drosophila melanogaster*. Journal of experimental biology, 199, 1837-1844.

FENSTERER, C., SCHOLZ, D., HOFFMANN, D. L., SPÖTL, C., SCHRÖDER-RITZRAU, A., HORN, C., PAJON, J. M. & MANGINI, A. 2013. Millennial-scale climate variability

during the last 12.5 ka recorded in a Caribbean speleothem. Earth and Planetary Science Letters, 361, 143-151.

FODEN, W., MACE, G. M., VIÉ, J.-C., ANGULO, A., BUTCHART, S. H., DEVANTIER, L., DUBLIN, H. T., GUTSCHE, A., STUART, S. & TURAK, E. 2008. Species susceptibility to climate change impacts. Wildlife in a changing world–an analysis of the 2008 IUCN Red List of threatened species, 77.

FONSECA-RIVERO, C., ALPIZAR-TIRZO, M., CUTIÉ-CANCINO, V., GONZÁLEZ-GARCÍA, I. T., HERNÁNDEZ-SOSA, M., VALDERÁ-FIGUEREDO, N., HERNÁNDEZ-GONZÁLEZ, D. & CABRERA-MEDINA, A. 2018. Estado del clima en Cuba 2017. Resumen ampliado. Revista cubana de meteorología, 24, 226-237.

GAO, L., YUAN, Z., LI, Y. & MA, Z. 2022. Genome-wide comparative analysis of DNAJ genes and their co-expression patterns with HSP70s in aestivation of the sea cucumber *Apostichopus japonicus*. Functional & Integrative Genomics, 22, 317-330.

GAO, L., YUAN, Z., YU, S., YANG, Y., LI, Y. & HE, C. 2018. Genome-wide identification of HSP70/110 genes in sea cucumber *Apostichopus japonicus* and comparative analysis of their involvement in aestivation. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 28, 162-171.

GAO, Y., LI, J.-N., PU, J.-J., TAO, K.-X., ZHAO, X.-X. & YANG, Q.-Q. 2022. Genome-wide identification and characterization of the HSP gene superfamily in apple snails (Gastropoda: Ampullariidae) and expression analysis under temperature stress. International Journal of Biological Macromolecules, 222, 2545-2555.

GONZÁLEZ-GUILLÉN, A. 2021. *Polymita* uncoiled, Collierville, Memphis: Fundcraft Publishing.

GORDILLO-PÉREZ, M. J., BEENAERTS, N., SÁNCHEZ, D. L., SMEETS, K., ARIAS-SOSA, Y. C. & REYES-TUR, B. 2025. Shell colour luminance of Cuban painted snails, *Polymita picta* and *Polymita muscarum* (Gastropoda: Cepolidae). PloS one, 20, e0314008.

GU, Z., CAVALCANTI, A., CHEN, F.-C., BOUMAN, P. & LI, W.-H. 2002. Extent of gene duplication in the genomes of *Drosophila*, nematode, and yeast. Molecular biology and evolution, 19, 256-262.

GUINDON, S., DUFAYARD, J.-F., LEFORT, V., ANISIMOVA, M., HORDIJK, W. & GASCUEL, O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Systematic biology, 59, 307-321.

GUNDAPPA, M. K., PEÑALOZA, C., REGAN, T., BOUTET, I., TANGUY, A., HOUSTON, R. D., BEAN, T. P. & MACQUEEN, D. J. 2022. Chromosome-level reference genome for

European flat oyster (*Ostrea edulis* L.). Evolutionary Applications, 15, 1713-1729.

GUNDERSON, A. R. & STILLMAN, J. H. 2015. Plasticity in thermal tolerance has limited potential to buffer ectotherms from global warming. Proceedings of the Royal Society B: Biological Sciences, 282, 20150401.

HARBOTT, M., WU, H., KUHNERT, H., JIMENEZ, C., GONZÁLEZ-DÍAZ, P. & RIXEN, T. 2023. A warming southern Gulf of Mexico: Reconstruction of anthropogenic environmental changes from a *Siderastrea siderea* coral on the northern coast of Cuba. Paleoceanography and Paleoclimatology, 38, e2023PA004717.

HASNAIN, P. & KANEKO, G. 2022. Phylogenetic annotation of *Caenorhabditis elegans* heat shock protein 70 genes. Micropublication Biology, 2022, 10.17912/micropub. biology. 000633.

HAVIRD, J. C., NEUWALD, J. L., SHAH, A. A., MAURO, A., MARSHALL, C. A. & GHALAMBOR, C. K. 2020. Distinguishing between active plasticity due to thermal acclimation and passive plasticity due to Q10 effects: Why methodology matters. Functional Ecology, 34, 1015-1028.

HAWKINS, S. J., MIESZKOWSKA, N., MROWICKI, R., LAB, M. B. A. G. A., OF LIFE, W. S. I. T. & CONSORTIUM, D. T. O. L. 2024. The genome sequence of the black-footed limpet, *Patella depressa* (Pennant, 1777). Wellcome Open Research, 9, 47.

HAYFLICK, L. 2007. Biological aging is no longer an unsolved problem. Annals of the New York academy of Sciences, 1100, 1-13.

HOANG, D. T., CHERNOMOR, O., VON HAESELER, A., MINH, B. Q. & VINH, L. S. 2018. UFBoot2: improving the ultrafast bootstrap approximation. Molecular biology and evolution, 35, 518-522.

HU, B., LI, M., YU, X., XUN, X., LU, W., LI, X., LI, Y., LOU, J., WANG, S. & ZHANG, L. 2019. Diverse expression regulation of Hsp70 genes in scallops after exposure to toxic *Alexandrium dinoflagellates*. Chemosphere, 234, 62-69.

HU, Z., SONG, H., FENG, J., ZHOU, C., YANG, M.-J., SHI, P., YU, Z.-L., LI, Y.-R., GUO, Y.-J. & LI, H.-Z. 2022. Massive heat shock protein 70 genes expansion and transcriptional signatures uncover hard clam adaptations to heat and hypoxia. Frontiers in Marine Science, 9, 898669.

HUANG, X., LI, S., GAO, Y. & ZHAN, A. 2018. Genome-wide identification, characterization and expression analyses of heat shock protein-related genes in a highly invasive ascidian *Ciona savignyi*. Frontiers in Physiology, 9, 1043.

HUEY, R. B., BERRIGAN, D., GILCHRIST, G. W. & HERRON, J. C. 1999. Testing the

adaptive significance of acclimation: a strong inference approach. American Zoologist, 39, 323-336.

ONEI. 2025. Anuario Estadístico de Cuba 2024. La Habana, Cuba: ONEI.

KALYAANAMOORTHY, S., MINH, B. Q., WONG, T. K. F., VON HAESELER, A. & JERMIIN, L. S. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods, 14, 587-589.

KAMPINGA, H. H. & CRAIG, E. A. 2010. The HSP70 chaperone machinery: J proteins as drivers of functional specificity. Nature reviews Molecular cell biology, 11, 579-592.

KANEKO, G. 2022. Phylogenetic annotation of *Drosophila melanogaster* heat shock protein 70 genes. microPublication Biology, 2022, 10.17912/micropub. biology. 000558.

KARLIN, S. & BROCCHIERI, L. 1998. Heat Shock Protein 70 Family: Multiple Sequence Comparisons, Function, and Evolution. Journal of Molecular Evolution, 47, 565-577.

KIM, J., KIM, H. J., CHOI, E., CHO, M., CHOI, S., JEON, M. A., LEE, J. S. & PARK, H. 2024. Expansion of the HSP70 gene family in *Tegillarca granosa* and expression profiles in response to zinc toxicity. Cell Stress and Chaperones, 29, 97-112.

KONDRASHOV, F. A., ROGOZIN, I. B., WOLF, Y. I. & KOONIN, E. V. 2002. Selection in the evolution of gene duplications. Genome biology, 3, research0008. 1.

KRAMARENKO, S. 2014. Active and passive dispersal of terrestrial mollusks: a review. Ruthenica, Russian Malacological Journal, 24.

KÜLTZ, D. 2003. Evolution of the cellular stress proteome: from monophyletic origin to ubiquitous function. Journal of Experimental Biology, 206, 3119-3124.

KÜLTZ, D. 2020. Defining biological stress and stress responses based on principles of physics. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 333, 350-358.

LEWIS, A. M., REYES-TUR, B. & DAVISON, A. 2025. The mitochondrial genome and phylogenetic placement of the world's most beautiful snail, *Polymita picta*. Journal of Molluscan Studies, 91, eyaf006.

LI, S., YANG, H., DUAN, Y., WU, L., HU, C., YU, B. & ZHAO, Y. 2025. Role of heat shock proteins in response to temperature stress and their effect on apoptosis in *Drosophila melanogaster*. International Journal of Biological Macromolecules, 306, 141320.

LINDQUIST, S. & CRAIG, E. A. 1988. The heat-shock proteins. Annual review of genetics, 22, 631-677.

LIU, J., WU, J., SILAIYIMAN, S., OUYANG, L., CAO, Z. & SHEN, C. 2025. Comparative Genomics Reveals Gene Duplication and Evolution in 26 Aurantioideae Species.

Horticulturae, 11, 209.

LONG, M., VANKUREN, N. W., CHEN, S. & VIBRANOVSKI, M. D. 2013. New gene evolution: little did we know. Annual review of genetics, 47, 307-333.

LU, H., LIU, C., YANG, C., HE, Z., WANG, L. & SONG, L. 2024. Genome-wide identification of the HSP70 genes in Pacific oyster *Magallana gigas* and their response to heat stress. Cell Stress and Chaperones, 29, 589-602.

MA, F. & LUO, L. 2020. Genome-wide identification of Hsp70/110 genes in rainbow trout and their regulated expression in response to heat stress. PeerJ, 8, e10022.

MAAL-BARED, R. 2006. Comparing environmental issues in Cuba before and after the Special Period: Balancing sustainable development and survival. Environment international, 32, 349-358.

MACEIRA-FILGUEIRA, D. 2016. *Polymita*. In: HIDALGO-GATO M, E. J., RODRÍGUEZ-LEÓN R. (ed.) Libro rojo de invertebrados terrestres de Cuba. La Habana.

MACEIRA-FILGUEIRA, D. 2016. *Polymita muscarum*. Libro rojo de invertebrados terrestres de Cuba, 115-116.

MACPHEE, R. D. & VINENT, M. A. I. The interpretation of Caribbean paleogeography: reply to Hedges. Proceedings of the International Symposium" Insular Vertebrate Evolution: the Palaeontological Approach": September, 16-19 Mallorca, 2005. Societat d'Història Natural de Balears, 175-184.

MANCINA, C. A. & FLORES, D. D. C. 2017. Diversidad biológica de Cuba: métodos de inventario, monitoreo y colecciones biológicas, Editorial AMA.

MANCINA, C. A., HERNÁNDEZ, M., MARTÍNEZ, D. & ESTRADA, R. 2017. An assessment of the potential effects of climate change on the distribution of painted land snail species, genus *Polymita* (Gastropoda: Cepolidae). Tentacle, 25, 17-19.

MARTÍNEZ, D. & HERNÁNDEZ, M. 2017. Posible efecto del cambio climático sobre la distribución actual y futura de moluscos terrestres endémicos cubanos.

MAYER, M. & BUKAU, B. 2005. Hsp70 chaperones: cellular functions and molecular mechanism. Cellular and molecular life sciences, 62, 670.

MAYER-BÖMOSER, S., SIGWART, J. & GREVE, C. 2022. The Cuban painted snail is "mollusc of the ear 2022". Senckenberg Institute press release [Cited March 18, 2022].

MELIKOV, A. & NOVAK, P. 2024. Heat Shock Protein Network: the Mode of Action, the Role in Protein Folding and Human Pathologies. Folia Biologica (00155500), 70.

METZGER, D. C., HEMMER-HANSEN, J. & SCHULTE, P. M. 2016. Conserved structure and expression of hsp70 paralogs in teleost fishes. Comparative Biochemistry and

Physiology Part D: Genomics and Proteomics, 18, 10-20.

MINH, B. Q., SCHMIDT, H. A., CHERNOMOR, O., SCHREMPF, D., WOODHAMS, M. D., VON HAESELER, A. & LANFEAR, R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. Molecular biology and evolution, 37, 1530-1534.

MIYAHIRA, I. C., CLAVIJO, C., CALLIL, C. T., CUEZZO, M. G., DARRIGRAN, G., GOMES, S. R., LASSO, C. A., MANSUR, M. C. D., PENA, M. S. & RAMÍREZ, R. 2022. The conservation of non-marine molluscs in South America: where we are and how to move forward. Biodiversity and Conservation, 31, 2543-2574.

MORENO, A. 1950. Estudio anatómico del género *Polymita* Beck. Memorias de la Sociedad Cubana de Historia Natural "Felipe Poey, 20, 21-35.

MORLEY, S., PECK, L., SUNDAY, J., HEISER, S. & BATES, A. 2019. Physiological acclimation and persistence of ectothermic species under extreme heat events. Global Ecology and Biogeography, 28, 1018-1037.

MOSELEY, P. L. 1997. Heat shock proteins and heat adaptation of the whole organism. Journal of applied physiology, 83, 1413-1417.

MOSSER, D. D. & BOLS, N. C. 1988. Relationship between heat-shock protein synthesis and thermotolerance in rainbow trout fibroblasts. Journal of Comparative Physiology B, 158, 457-467.

MROWICKI, R., UHL, R., LAB, M. B. A. G. A., OF LIFE, W. S. I. T. & CONSORTIUM, D. T. O. L. 2025. The genome sequence of the Pacific oyster, *Magallana gigas* (Thunberg, 1793). Wellcome Open Research, 9, 284.

MURPHY, M. E. 2013. The HSP70 family and cancer. Carcinogenesis, 34, 1181-1188.

NGUYEN, L.-T., SCHMIDT, H. A., VON HAESELER, A. & MINH, B. Q. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular biology and evolution, 32, 268-274.

NICOLAI, A. & ANSART, A. 2017. Conservation at a slow pace: terrestrial gastropods facing fast-changing climate. Conservation Physiology, 5, cox007.

NIKOLAIDIS, N. & NEI, M. 2004. Concerted and nonconcerted evolution of the Hsp70 gene superfamily in two sibling species of nematodes. Molecular biology and evolution, 21, 498-505.

NOBLE, D. W., KAR, F., BUSH, A., SEEBACHER, F. & NAKAGAWA, S. 2025. Limited plasticity but increased variance in physiological rates across ectotherm populations under climate change. Functional Ecology, 39, 1176-1193.

PENG, J., LI, Q., XU, L., WEI, P., HE, P., ZHANG, X., ZHANG, L., GUAN, J., ZHANG, X. & LIN, Y. 2020. Chromosome-level analysis of the *Crassostrea hongkongensis* genome reveals extensive duplication of immune-related genes in bivalves. Molecular ecology resources, 20, 980-994.

PEROS, M., COLLINS, S., G'MEINER, A. A., REINHARDT, E. & PUPO, F. M. 2017. Multistage 8.2 kyr event revealed through high-resolution XRF core scanning of Cuban sinkhole sediments. Geophysical Research Letters, 44, 7374-7381.

PEROS, M., GREGORY, B., MATOS, F., REINHARDT, E. & DESLOGES, J. 2015. Late-Holocene record of lagoon evolution, climate change, and hurricane activity from southeastern Cuba. The Holocene, 25, 1483-1497.

PEYRETAILLADE, E., BROUSSOLLE, V., PEYRET, P., MÉTÉNIER, G., GOUY, M. & VIVARES, C. P. 1998. *Microsporidia*, amitochondrial protists, possess a 70-kDa heat shock protein gene of mitochondrial evolutionary origin. Molecular biology and evolution, 15, 683-689.

POWELL, D., SUBRAMANIAN, S., SUWANSA-ARD, S., ZHAO, M., O'CONNOR, W., RAFTOS, D. & ELIZUR, A. 2018. The genome of the oyster *Saccostrea* offers insight into the environmental resilience of bivalves. DNA Research, 25, 655-665.

QIAN, W. & ZHANG, J. 2014. Genomic evidence for adaptation by gene duplication. Genome research, 24, 1356-1362.

REAMS, A. B. & ROTH, J. R. 2015. Mechanisms of gene duplication and amplification. Cold Spring Harbor perspectives in biology, 7, a016592.

REYES-TUR, B., CHEN, Z., GORDILLO-PÉREZ, M. J., HAMADOU, A. B., GERHEIM, C., GREVE, C. & SIGWART, J. D. 2025. Draft genome of the Cuban painted land snail *Polymita picta*, International Mollusc of the Year 2022.

ROBERTS, S. & FEDER, M. 2000. Changing fitness consequences of hsp70 copy number in transgenic Drosophila larvae undergoing natural thermal stress. Functional Ecology, 14, 353-357.

RODRÍGUEZ, L. C., COLLINS, L., FERNÁNDEZ, M., HARMON, S., HIRTLE, J., KLEPAK, H. P., MARTIN, G. M., RIDEL, G. M., MILLER, V. & VÁZQUEZ, L. L. 2021. Disaster preparedness and climate change in Cuba: management and adaptation, Bloomsbury Publishing PLC.

ROHR, J. R., CIVITELLO, D. J., COHEN, J. M., ROZNIK, E. A., SINERVO, B. & DELL, A. I. 2018. The complex drivers of thermal acclimation and breadth in ectotherms. Ecology letters, 21, 1425-1439.

ROSENZWEIG, R., NILLEGODA, N. B., MAYER, M. P. & BUKAU, B. 2019. The Hsp70 chaperone network. Nature reviews molecular cell biology, 20, 665-680.

ROTH, C., RASTOGI, S., ARVESTAD, L., DITTMAR, K., LIGHT, S., EKMAN, D. & LIBERLES, D. A. 2007. Evolution after gene duplication: models, mechanisms, sequences, systems, and organisms. Journal of Experimental Zoology Part B: Molecular and Developmental Evolution, 308, 58-73.

SCHEIL, A. E., GÄRTNER, U. & KÖHLER, H.-R. 2012. Colour polymorphism and thermal capacities in *Theba pisana* (OF Müller 1774). Journal of Thermal Biology, 37, 462-467.

SCHWEIZER, M., TRIEBSKORN, R. & KÖHLER, H. R. 2019. Snails in the sun: Strategies of terrestrial gastropods to cope with hot and dry conditions. Ecology and Evolution, 9, 12940-12960.

SHI, L., XU, Y., MAO, C., WANG, Z., GUO, S., JIN, X., YAN, S. & SHI, B. 2020. Effects of heat stress on antioxidant status and immune function and expression of related genes in lambs. International Journal of Biometeorology, 64, 2093-2104.

SINGH, M. K., HAN, S., JU, S., RANBHISE, J. S., HA, J., YEO, S. G., KIM, S. S. & KANG, I. 2025. Hsp70: A Multifunctional Chaperone in Maintaining Proteostasis and Its Implications in Human Disease. Cells, 14, 509.

SOMERO, G. N. 2020. The cellular stress response and temperature: Function, regulation, and evolution. Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 333, 379-397.

STAIKOU, A., SAGONAS, K., SPANOUDI, O., SAVVIDOU, K., NAZLI, Z., FEIDANTSIS, K. & MICHAELIDIS, B. 2024. Activities of antioxidant enzymes and Hsp levels in response to elevated temperature in land snail species with varied latitudinal distribution. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 269, 110908.

STRAALEN, N. M. V. 2003. Peer reviewed: ecotoxicology becomes stress ecology. Environmental science & technology, 37, 324A-330A.

SUTHERLAND, B. J., DIVILOV, K., THOMPSON, N. F., DELOMAS, T. A., LUNDA, S. L., LANGDON, C. J. & GREEN, T. J. 2025. Pedigree-based genome-wide imputation using a low-density amplicon panel for the highly polymorphic Pacific oyster *Crassostrea* (*Magallana*) *gigas*. bioRxiv, 2025.02. 11.637717.

SUTHERLAND, B. J. G., THOMPSON, N. F., SURRY, L. B., GUJJULA, K. R., CARRASCO, C. D., CHADARAM, S., LUNDA, S. L., LANGDON, C. J., CHAN, A. M., SUTTLE, C. A. & GREEN, T. J. 2024. An amplicon panel for high-throughput and low-cost

genotyping of Pacific oyster. G3 Genes|Genomes|Genetics, 14.

TAFF, C. C. & VITOUSEK, M. N. 2016. Endocrine flexibility: optimizing phenotypes in a dynamic world? Trends in ecology & evolution, 31, 476-488.

TEAM, R. C. 2022. R: a language and environment for statistical computing and graphics. Foundation for Statistical Computing.

THORVALDSDÓTTIR, H., ROBINSON, J. T. & MESIROV, J. P. 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Briefings in bioinformatics, 14, 178-192.

TSAN, M.-F. & GAO, B. 2009. Heat shock proteins and immune system. Journal of Leucocyte Biology, 85, 905-910.

TULL, T., SCHMIDT, D. & KÖHLER, H.-R. 2020. Role of ambient wavelength, shell size and pigmentation intensity in the heating of Mediterranean *Theba pisana* (Eupulmonata: Helicidae). Journal of Molluscan Studies, 86, 249-253.

VENDRAMI, D. L., DE NOIA, M., TELESCA, L., BRODTE, E. M. & HOFFMAN, J. I. 2020. Genome-wide insights into introgression and its consequences for genome-wide heterozygosity in the *Mytilus* species complex across Europe. Evolutionary Applications, 13, 2130-2142.

WANG, W., LIU, Q., LIU, Q. & HENDRICKSON, W. A. 2021. Conformational equilibria in allosteric control of Hsp70 chaperones. Molecular cell, 81, 3919-3933. e7.

WANG, Y., WANG, X., TANG, H., TAN, X., FICKLIN, S. P., FELTUS, F. A. & PATERSON, A. H. 2011. Modes of gene duplication contribute differently to genetic novelty and redundancy, but show parallels across divergent angiosperms. PloS one, 6, e28150.

WEI, M., GE, H., SHAO, C., YAN, X., NIE, H., DUAN, H., LIAO, X., ZHANG, M., CHEN, Y. & ZHANG, D. 2020. Chromosome-level clam genome helps elucidate the molecular basis of adaptation to a buried lifestyle. Science, 23.

WHEELER, D. S. & WONG, H. R. 2012. Heat shock proteins. Encyclopedia of Intensive Care Medicine. Springer.

XU, K., XU, H. & HAN, Z. 2018. Genome-wide identification of Hsp70 genes in the large yellow croaker (*Larimichthys crocea*) and their regulated expression under cold and heat stress. Genes, 9, 590.

YADAV, K., YADAV, A., VASHISTHA, P., PANDEY, V. P. & DWIVEDI, U. N. 2019. Protein misfolding diseases and therapeutic approaches. Current Protein and Peptide Science, 20, 1226-1245.

YANG, S., ZHANG, X., YUE, J.-X., TIAN, D. & CHEN, J.-Q. 2008. Recent duplications

dominate NBS-encoding gene expansion in two woody species. Molecular Genetics and Genomics, 280, 187-198.

ZHANG, J., LI, J., LIU, B., ZHANG, L., CHEN, J. & LU, M. 2013. Genome-wide analysis of the *Populus* Hsp90 gene family reveals differential expression patterns, localization, and heat stress responses. BMC genomics, 14, 1-14.

ZHANG, Z. & ZHANG, Q. 2012. Molecular cloning, characterization and expression of heat shock protein 70 gene from the oyster *Crassostrea hongkongensis* responding to thermal stress and exposure of Cu2+ and malachite green. Gene, 497, 172-180.

ZHENG, W., XU, X., CHEN, Y., WANG, J., ZHANG, T., E, Z., CHEN, S. & LIU, Y. 2023. Genome-wide identification, molecular characterization, and involvement in response to Abiotic and Biotic stresses of the HSP70 Gene Family in Turbot (*Scophthalmus maximus*). International Journal of Molecular Sciences, 24, 6025.