Graft-Induced Reprogramming Under Arterial Hemodynamic Stress is Associated with

a Shared Proteomic Phenotype in Arterial and Venous Conduits Following Coronary

Artery Bypass Grafting

Eun Na Kim, MD.PhD.[1†], Suk Ho Sohn, MD.PhD.[2†], Jiyoung Yu, PhD.[3†], Joon Seo Lim,

PhD.[4], Jiwon Koh, MD.PhD.[1], Jaemoon Koh, MD.PhD.[1], Changha Hwang, PhD.[5,6],

Kyunggon Kim, PhD.[7‡,] Ho Young Hwang, MD.PhD.[2‡], Se Jin Oh, MD.PhD.[8‡]

[1]Department of Pathology, Seoul National University Hospital, Seoul National University

College of Medicine, Seoul, Republic of Korea

[2]Department of Thoracic and Cardiovascular Surgery, Seoul National University Hospital,

Seoul National University College of Medicine, Seoul, Republic of Korea

[3]Clinical Proteomics Core Lab, Convergence Research Center, Asan Medical Center, Seoul,

Republic of Korea

[4]Clinical Research Center, Asan Medical Center, University of Ulsan College of Medicine,

Seoul, Republic of Korea

[5]Department of Applied Statistics, Dankook University, Gyeonggi-do, Republic of Korea

[6]Arontier, Seoul, Republic of Korea

[7]Department of Digital Medicine, BK21 Project, Asan Medical Center, University of Ulsan

College of Medicine, Seoul, Republic of Korea

[8]Department of Thoracic and Cardiovascular Surgery, SMG-SNU Boramae Medical Center,

Seoul National University College of Medicine, Seoul, Republic of Korea

SHORT TITLE: Proteomic Convergence of Arterial/Venous Grafts

†These authors have contributed equally to this work and share the first authorship

‡These authors have contributed equally to this work and share the corresponding authorship

CORRESPONDING AUTHORS

Kyunggon Kim, PhD

Department of Digital Medicine, BK21 Project

Asan Medical Center, University of Ulsan College of Medicine

88, Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Republic of Korea

Email: kimkyunggon@gmail.com

Ho Young Hwang, MD, PhD

Department of Thoracic and Cardiovascular Surgery

Seoul National University Hospital, Seoul National University College of Medicine

101 Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea

Email: scalpel@snu.ac.kr

Se Jin Oh, MD, PhD (Lead contact)

Department of Thoracic and Cardiovascular Surgery

SMG-SNU Boramae Medical Center, Seoul National University College of Medicine

20, Boramae-ro 5-gil, Dongjak-gu, Seoul 07061, Republic of Korea

Email: wpwnn@snu.ac.kr

THE TOTAL WORD COUNT OF THE MANUSCRIPT: 8128

ABSTRACT

Background: Autologous venous conduits remain the mainstay of coronary artery bypass

grafting (CABG); however, their long-term patency is limited by adverse vascular

remodeling. While arterial and venous grafts undergo structural and molecular adaptation to

arterial hemodynamics, the mechanisms underlying successful adaptation remain poorly

characterized owing to the scarcity of functioning human graft tissues.

Methods: We analyzed bypass grafts explanted en bloc during orthotopic heart

transplantation, comprising patent vein grafts (VGP, n=19), occluded vein grafts (VGO,

n=9), and patent arterial grafts (AGP, n=13), alongside freshly harvested controls (vein,

n=23; artery, n=11). Quantitative histomorphometry, label-free LC–MS/MS proteomics,

pathway analysis, and immunohistochemistry were performed.

Results: While patent arterial and venous grafts exhibited medial elastin reinforcement,

adventitial neovascularization, and organized intimal remodeling, occluded vein grafts

demonstrated elastic layer disruption, medial degeneration, and collagen-dense fibrosis.

Proteomic analysis revealed a convergent adaptive signature in patent grafts enriched for

RNA metabolism, protein synthesis, cytoskeletal dynamics, and extracellular matrix

organization, with shared activation of NR4A3, STAT1, and EGFR signaling. Vein-specific

adaptation involved SRC–PTGES activation with concomitant STAT3 suppression, whereas

arterial grafts engaged IGF1–RUNX2-associated proliferative programs. Failed venous grafts

exhibited angiotensin-driven fibroinflammatory remodeling. Endothelial nitric oxide synthase

expression was markedly upregulated in the luminal endothelium and adventitial neovessels

of patent grafts.

Conclusions: Despite distinct vascular origins, arterial and venous grafts converged toward a

shared adaptive phenotype characterized by structural maturation and vasoprotective

signaling. Targeting common adaptive pathways—including NR4A3, STAT1, EGFR, and

adventitial eNOS—may offer therapeutic strategies to enhance long-term graft durability.

WHAT ARE THE CLINICAL IMPLICATIONS?

This study provides human tissue–based evidence that patent coronary bypass grafts share a

common adaptive remodeling program, irrespective of whether the conduit is arterial or

venous. Rather than simple venous "arterialization," successful grafts converged on a

phenotype characterized by coordinated biosynthetic activity, structural maturation, and

vasoprotective signaling, whereas occluded vein grafts exhibited a distinct angiotensin-driven

fibroinflammatory remodeling program. Importantly, patent grafts demonstrated conserved

upregulation of NR4A3, STAT1, EGFR, and endothelial nitric oxide synthase (eNOS) in

both luminal and adventitial endothelial compartments, identifying molecular features that

distinguish adaptive from maladaptive remodeling in humans.

These findings are clinically relevant because they highlight shared, potentially targetable

pathways that may support graft durability across conduit types and provide mechanistic

support for surgical strategies that preserve perivascular adipose tissue during vein

harvesting. Moreover, the association of graft failure with angiotensin-centered extracellular

matrix remodeling suggests a modifiable biological axis that warrants further investigation.

Together, this work establishes a human molecular framework for graft adaptation that may

inform future biomarker development, therapeutic targeting, and precision strategies aimed at

improving outcomes after coronary artery bypass grafting.

NON-STANDARD ABBREVIATIONS AND ACRONYMS

AC	Artery control
AGP	Artery graft, patent
CABG	Coronary artery bypass grafting
DAB	3,3′-Diaminobenzidine
ECM	Extracellular matrix
eNOS	Endothelial nitric oxide synthase
FFPE	Formalin-fixed, paraffin-embedded
IGF1	Insulin-like growth factor 1
IPA	Ingenuity Pathway Analysis
LC–MS/MS	Liquid chromatography–tandem mass spectrometry
PCA	Principal component analysis
PGC-1α	Peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PPARGC1A)
PPI	Protein–protein interaction
PVAT	Perivascular adipose tissue
RUNX2	Runt-related transcription factor 2
SRC	Src proto-oncogene, non-receptor tyrosine kinase
SRP	Signal recognition particle
STAT	Signal transducer and activator of transcription
VC	Vein control
VEGF	Vascular endothelial growth factor
VGO	Vein graft, occluded
VGP	Vein graft, patent

INTRODUCTION

Coronary artery bypass grafting (CABG) remains a primary revascularization strategy for multivessel coronary artery disease; however, patient outcomes continue to heavily depend on long-term graft performance[1]. Arterial grafts, characterized by their elastic architecture and intrinsic compatibility with the high-pressure coronary circulation, exhibit superior long-term patency[2]. In contrast, venous grafts are abruptly exposed to arterial hemodynamics that differ fundamentally from their low-pressure venous origin and must undergo extensive adaptive remodeling to maintain patency[3]. Although arterial conduits—including the internal mammary artery, radial artery, and other elastic or muscular arteries—provide the most durable options[4], their limited availability and anatomical constraints often necessitate the use of venous grafts in patients with multivessel disease[5]. Consequently, mitigating venous graft failure has become a key determinant of long-term outcomes following CABG.

Failure of this adaptive process—particularly in venous grafts—precipitates early thrombosis, excessive intimal hyperplasia, and accelerated atherosclerosis, events that collectively contribute to their significantly lower patency compared to arterial conduits[6,7]. Because graft failure can result in recurrent ischemia, repeat revascularization, and even sudden cardiac death, elucidating the biological mechanisms that preserve graft integrity is essential for improving CABG outcomes.

Although the mechanisms contributing to venous graft failure have been extensively described, the biological processes that enable successful graft adaptation remain poorly understood. Previous human studies have primarily examined conduits at the time of harvesting[8] or on grafts removed following clinical failure[9], offering limited insight into the structural and molecular changes that occur in vivo during graft function. As a result, the histomorphological features, molecular pathways, and vasoprotective mechanisms that support adaptation to coronary hemodynamic stress remain incompletely characterized.

Recent attention has focused on strategies such as no-touch harvesting and the

preservation of perivascular adipose tissue (PVAT), both of which have been associated with

improved graft performance[10,11]. Proposed mechanisms include enhanced endothelial

integrity and increased endothelial nitric oxide synthase (eNOS) expression in freshly

harvested no-touch conduits[12,13]. However, these observations are limited to pre-implant

tissue, and direct evidence linking PVAT-associated or endothelial eNOS to graft patency in

humans remains lacking.

A further conceptual gap concerns the traditional assumption that venous conduits

undergo "arterialization" after implantation[3]. Although this term is widely used to describe

the adaptive transition of veins exposed to arterial pressure, the molecular basis, temporal

progression, and histomorphological correlates of this process in clinically patent grafts have

never been systematically defined. It remains uncertain whether venous conduits undergo a

unidirectional transition toward an arterial phenotype while arterial grafts largely preserve

their native architecture, or whether more complex adaptive programs are involved.

In this study, we leveraged a rare clinical circumstance that offers uniquely valuable

biological insights by enabling the direct examination of both arterial and venous bypass

grafts explanted en bloc at the time of orthotopic heart transplantation. Using an integrated

approach that combines quantitative histomorphology, high-resolution proteomics, and

targeted immunohistochemistry, we examined not only occluded venous grafts but also

successfully functioning venous and arterial grafts. By contrasting these specimens, we aimed

to define the adaptive remodeling pathways, signaling trajectories, and conduit-specific

mechanisms that support graft patency in humans.

METHODS

Study Approval and Patient Samples

This study was approved by the institutional review boards of Seoul National University

Hospital (J-2306-002-1435) and SMG-SNU Boramae Medical Center (10-2023-25) on June

2, 2023. Written informed consent was obtained from all participants.

A total of eight patients who had previously undergone CABG and subsequently

received orthotopic heart transplantation for end-stage heart failure (Figure 1A) were

enrolled. Patient demographics are summarized in Table S1. All vein grafts were saphenous

vein conduits, and all artery grafts were internal mammary arteries. Accordingly, the terms

"vein graft" and "artery graft" were used to refer to these respective conduit types.

Seven patients underwent both procedures at Seoul National University Hospital and

SMG-SNU Boramae Medical Center and received vein grafts harvested using the no-touch

technique, configured in Y- or I-graft arrangements, and anastomosed to the in-situ arterial

graft. One patient, who underwent CABG at an external institution, received conventionally

harvested vein grafts with aortocoronary anastomosis. Graft patency prior to transplantation

was assessed using coronary angiography or multi-detector computed tomography. The

median interval from CABG to transplantation was 74 months (range, 0.2–226 months).

Histomorphological Analysis

For histological assessment, we analyzed 41 formalin-fixed, paraffin-embedded (FFPE) graft

blocks, including 13 artery graft patent (AGP) blocks from 5 patients, 19 vein graft patent

(VGP) blocks from 6 patients, and 9 vein graft obstructed (VGO) blocks from 2 patients. As

controls, freshly harvested graft tissues were obtained from 28 patients undergoing CABG,

yielding 11 artery control (AC) blocks from 11 patients and 23 vein control (VC) blocks from

23 patients.

FFPE tissue blocks were sectioned into 4-µm-thick slides and stained with hematoxylin

and eosin, elastin, and Masson's trichrome (MT) to evaluate histological features, including

elastic fiber content, the formation of internal and external elastic laminae, and intimal

fibrosis. Slides were digitized using a whole-slide scanner (Aperio GT450 DX, Leica

Biosystems). Manual annotations were performed using QuPath (https://qupath.github.io/),

followed by quantification of intima, media, and adventitia thickness and area, as well as

luminal, medial, and adventitial circumferences. Additionally, the number and depth of clefts

within the hyperplastic intima were quantified (Figure S1).

Details on the methodologies regarding proteomic analysis, LC-MS/MS, database search,

canonical pathway analysis, immunohistochemistry, and statistical analysis are provided in

Supplemental Methods.

RESULTS

Histomorphological Characteristics of Patent Arterial and Venous Grafts

We analyzed 41 FFPE vessel and graft specimens to characterize baseline vascular

architecture and graft-associated morphological remodeling (Figure 1B-D). After exposure

to arterial hemodynamics, VGP underwent pronounced structural remodeling,

characterized by substantial intimal and medial thickening, the development of a well-

defined internal elastic lamina, and an increased presence of elastic fibers interwoven

within the medial layer. In addition, a newly formed, wavy elastic layer between the media

and adventitia was consistently observed, representing a structural reorganization that

renders the venous wall similar to the layered organization typical of arterial conduits. In

contrast, VGO showed aneurysmal dilatation, loss of organized elastic layers, medial

degeneration, and collapse of the normal intima–media–adventitia boundaries, consistent

with structural failure rather than adaptive remodeling. Detailed results of quantitative

morphometric analysis and intimal measurements are provided in **Supplemental Results**.

Immunohistochemical Analysis of Patent and Occluded Vein Grafts

Histochemical and immunohistochemical staining (Figure 1E) revealed distinct differences in

intimal organization between patent and occluded vein grafts. In VGP, elastic staining

demonstrated a well-organized intima and media. Masson's trichrome staining indicated

fibrosis; however, the collagen matrix maintained a structured pattern with interspersed

αSMA-positive myofibroblasts, suggesting an orderly and adaptive remodeling process. In

contrast, VGO displayed a fundamentally different morphology. Elastic staining revealed

significant disruption of the elastic lamina structures. Masson's Trichrome staining

demonstrated dense, disorganized collagen consistent with a non-cellular fibrotic matrix, and

αSMA staining confirmed the absence of smooth muscle–type cells within the thickened

intima.

Proteomics

Global Proteomic Profiling Reveals Condition-Specific Molecular Signatures

To characterize the molecular landscape of vascular remodeling during graft adaptation

and failure, we performed quantitative proteomic analysis using LC-MS/MS on five

experimental groups: VC, AC, VGP, AGP, and VGO. A total of 4,836 proteins were

identified across all vascular samples (Figure 2A). Unsupervised two-dimensional PCA

revealed a clear separation between AC and VC samples. In contrast, AGP and VGP samples

clustered closely together. VGO samples formed a distinct, isolated cluster, separate from

both native vessels and patent grafts (Figure 2B).

Volcano Plot Analysis of Graft-Associated Proteomic Changes

225 Volcano plot analysis (Figure 2C) revealed extensive proteomic remodeling across all graft

226 types. Only a very small subset of proteins differed between AGP and VGP, underscoring the

227 proteomic similarity between the two conduits after grafting. In contrast, the comparison

228 between VGP and VGO showed marked divergence, with 475 proteins enriched in VGP and

229 220 enriched in VGO.

Canonical Pathway Enrichment Analysis of AGP and VGP

232 Canonical pathway enrichment analysis (Figure 2D) demonstrated that AGP (vs. AC) and

233 VGP (vs. VC) share a broad set of commonly enriched pathways. Both comparisons showed

234 significant activation of translation-associated programs, indicating similarities in

235 biosynthetic and proteostasis-related pathways in both graft types.

Hierarchical Clustering Revealed Group-Specific Proteomic Signatures

Hierarchical clustering of the 2,492 differentially expressed proteins revealed distinct group-

specific expression patterns across vascular samples (Figure 2E)—VC in Cluster 7, AC in

Cluster 6, VGO in Cluster 5, and both AGP and VGP in Cluster 4. To further characterize the

biological processes represented by each cluster, we employed a three-tiered analytical

workflow consisting of: (1) pathway enrichment analysis using Ingenuity Pathway Analysis,

which offers a function-centered interpretation of overrepresented biological programs; (2)

protein–protein interaction (PPI) network analysis using the STRING database[14], which

delineates the structural and functional connectivity among proteins; and (3) integrative

functional annotation using the Metascape[15] enrichment bubble plot (GO/KEGG/Reactome).

These analyses were conducted on cluster-specific protein subsets derived from the

hierarchical clustering: AC (Cluster 6, n=171 proteins), VC (Cluster 7, n=340 proteins), VGP

+ AGP (Cluster 4, n=741 proteins), and VGO (Cluster 5, n=210 proteins).

Integrated Functional Analysis Revealed Fundamentally Distinct Remodeling Programs

Represented by Clusters 4 and 5

Cluster 4 (VGP + AGP) captured the shared proteomic signature of patent arterial and venous

grafts. Pathway enrichment analysis (Figure 3A) demonstrated coordinated activation of

RNA metabolism, mRNA processing, translation initiation and elongation, ribonucleoprotein

biogenesis, and SRP-dependent cotranslational protein targeting to membranes, along with

pathways supporting extracellular matrix (ECM) organization and cytoskeletal adaptation.

PPI network analysis (Figure 3B) revealed densely interconnected ribosomal and translation

hubs, as well as organized ECM-remodeling subnetworks, consistent with an active,

reparative remodeling phenotype. In contrast, Cluster 5 (VGO) represented the molecular

landscape of failed vein grafts. Pathway enrichment analysis revealed predominant activation

of complement and coagulation cascades, acute inflammatory signaling, wound healing, and

disorganized ECM remodeling. PPI networks (Figure S4) formed extensive thrombo-

inflammatory modules and collagen-rich ECM clusters, indicating a shift toward a

maladaptive, pro-inflammatory, and pro-thrombotic phenotype.

Metascape-based functional annotation (Figure 3C) further distinguished the adaptive

and maladaptive programs of these clusters. Cluster 4 showed enrichment in RNA

metabolism, VEGFA–VEGFR2 signaling, receptor tyrosine kinase pathways, protein

maturation, intracellular transport, lysosomal activity, and immune-regulatory processes,

reflecting a biosynthetically vigorous, structurally adaptive, and immunologically balanced

program characteristic of successful grafts. In contrast, Cluster 5 was enriched in complement

and coagulation cascades, NABA matrisome pathways, extracellular matrix organization,

platelet activation, plasma lipoprotein assembly, and IGFBP-mediated IGF transport

regulation. This enrichment is consistent with a matrix-disruptive, thrombosis-prone, and

growth factor–dysregulated environment associated with graft obstruction.

Canonical pathway analysis further revealed distinct activation patterns across graft

conditions, with biosynthetic and vascular remodeling pathways predominating in functional

grafts, while inflammatory and coagulative pathways characterized occluded grafts.

Activation of Proliferative and Biosynthetic Programs and Suppression of Contractile

Regulators in Patent Grafts

Upstream regulator analysis of patent grafts compared to native vessels revealed coordinated

activation of proliferative and biosynthetic signaling networks. In AGP, key activated

upstream nodes included IGF1, RUNX2, NR4A3, PGC-1α, EGFR, and MYC (Figure 4A). In

VGP, STAT1, SRC, PTGES, resistin (RETN), and G6PD were identified as activated nodes,

forming a coherent regulatory network (Figure 4A,B). In contrast, regulators involved in

vascular contractility and cytoskeletal integrity, including serum response factor (SRF) and

myocardin-related transcription factor A (MRTFA), were predicted to be inhibited in VGP.

This transcriptional profile indicates phenotypic modulation toward a synthetic phenotype,

characterized by enhanced extracellular matrix production, increased biosynthetic capacity,

and elevated proliferative activity—hallmarks of adaptive vascular remodeling during vein

graft transformation in the arterial circulation.

Immunohistochemical Validation of Key Regulatory Nodes

To validate pathway-level predictions, immunohistochemical analysis was performed on key

upstream regulators identified through IPA analysis (Figure 4C). Shared adaptive regulators

across both graft types—NR4A3, EGFR, and STAT1—demonstrated significantly increased

expression in both arterial grafts (AGP vs. AC, P<0.01) and venous grafts (VGP vs. VC,

P<0.01), establishing these as conserved adaptive regulators during adaptation to arterial

hemodynamics. Notably, the expressions of NR4A3 and EGFR were markedly reduced in

VGO, consistent with regression of proliferative signaling during graft failure. Graft-Type–

Specific Adaptive Signatures AGP exhibited significant increases in IGF1 and RUNX2

expression compared to native AC, validating pathway predictions of growth factor–driven

proliferative and osteogenic signaling in arterial graft adaptation. Conversely, VGP showed

elevated expression of RETN (resistin), SRC, and PTGES compared to native VC,

confirming graft-type–specific activation of metabolic stress and biosynthetic mediator

pathways predicted by IPA analysis.

PGC-1α—a factor predicted by the proteomic model to vary by graft type—showed an

upward trend in both arterial and venous grafts (AGP > AC; VGP > VC), although the

differences did not reach statistical significance (AC vs. AGP, P=0.177; VC vs. VGP,

P=0.104). Similarly, c-Myc expression tended to be higher in AGP compared with AC, but

this difference was not statistically significant (P=0.114) (Figure S14).

Collectively, these immunohistochemical findings validate the pathway analysis

predictions and establish both conserved adaptive programs (NR4A3, EGFR, STAT1, PGC-

1α) and graft-type–specific regulatory signatures (IGF1/RUNX2 in arterial grafts;

RETN/SRC/PTGES in venous grafts) that operate during vascular graft adaptation to arterial

circulation.

EGFR–STAT3 Signaling Differences Between VGP and VGO

Upstream regulator analysis predicted preferential activation of EGFR in VGP, identifying

EGFR as a key regulatory node with predicted inhibitory effects on STAT3 (Figure 5A). This

prediction aligns with prior immunohistochemical findings (Figure 4C), which demonstrate

elevated EGFR expression in VGP and AGP, whereas VGO exhibits marked EGFR

downregulation.

To evaluate the corresponding STAT3 response, immunohistochemical staining was

performed (Figure 5B). VGP displayed minimal STAT3 nuclear positivity, whereas VGO

demonstrated pronounced STAT3 upregulation within the fibrotic neointima. Quantitative

analysis revealed a trend toward higher STAT3 expression in VGO (P=0.08), supporting

reciprocal signaling dynamics characterized by EGFR-dominant/STAT3-suppressed activity

in patent grafts versus STAT3-dominant activation in occluded grafts.

Angiotensin Activation and ECM Remodeling in VGO

Upstream regulator analysis comparing VGP and VGO identified angiotensinogen as one of

the most strongly activated regulatory nodes associated with graft failure (Figure 5C). This

activation was accompanied by the coordinated upregulation of multiple extracellular matrix–

related proteins in VGO. The angiotensin-centered regulatory network (Figure 5D) revealed

elevated expression of fibrillar collagens (COL5A1, COL5A2), small leucine-rich

proteoglycans (decorin, biglycan, lumican), and thrombospondin-2, which collectively

promote collagen fibrillogenesis and matrix stabilization. Regulator-effects modeling

predicted enhanced matrix tensile strength as a downstream consequence in VGO, consistent

with maladaptive fibrotic remodeling and reduced vascular compliance.

To validate these proteomic predictions, immunohistochemical staining for

angiotensinogen was performed on matched VGP and VGO sections (Figure 5E). VGP

exhibited minimal angiotensinogen immunoreactivity, showing only faint background

staining. In contrast, VGO demonstrated marked angiotensinogen accumulation throughout

the media and adventitia, with intense deposition in regions of dense extracellular matrix

expansion and structural distortion. Quantitative analysis confirmed significantly elevated

angiotensinogen expression in VGO compared with VGP (P<0.01), supporting the

association between angiotensin pathway activation and fibroinflammatory remodeling in

failed grafts.

Enhanced eNOS Expression in Luminal and Adventitial Neovascular Endothelial Cells

Following Arterial Grafting

Expression of eNOS was evaluated by immunohistochemical staining across all vessel groups

(Figure 6A). AC and VC demonstrated weak to moderate eNOS immunoreactivity in the

luminal endothelium. Both AGP and VGP showed markedly elevated eNOS expression,

whereas VGO showed minimal signal, consistent with near-complete endothelial denudation.

Single-cell optical density analysis demonstrated enhanced eNOS expression in patent

grafts. Histogram distributions showed rightward shifts in AGP and VGP compared with

their respective controls (Figure 6B). Using a threshold of mean DAB optical density > 0.2,

the proportion of eNOS-positive cells was significantly higher in AGP versus AC (P<0.001)

and in VGP versus VC (P<0.01) (Figure 6C), distinguishing successfully adapted conduits

from failed grafts.

Adventitial Neovascularization and Associated Upregulation of eNOS

To evaluate adaptive remodeling of the vessel wall, we quantified CD31-positive adventitial

neovascularization (Figure S1E–H). Total microvascular content, measured as the cumulative

CD31-positive vessel perimeter, was significantly higher in AGP compared with AC

(P<0.001). Venous grafts showed a trend toward increased vasa vasorum relative to VC,

although this did not reach statistical significance. VGO demonstrated markedly reduced

adventitial microvascular density (Figure 6D).

Notably, robust eNOS immunoreactivity was observed in the endothelial cells lining

adventitial neovessels of patent grafts (Figure 6E, arrows). Quantification of eNOS-positive

microvessels—defined by endothelial DAB staining with nuclear co-localization—revealed

significant increases in AGP vs. AC and in VGP vs. VC (Figure 6F). VGO exhibited a

marked reduction in eNOS-positive neovessels, suggesting that eNOS expression in both

luminal and adventitial endothelium is associated with graft patency.

DISCUSSION

By leveraging a unique opportunity to analyze patent arterial and venous grafts explanted en

bloc during cardiac transplantation, we provide the first comprehensive molecular

characterization of successful graft adaptation in humans. Our findings challenge the

traditional concept of venous "arterialization" and reveal that both arterial and venous grafts

converge toward a shared adaptive phenotype characterized by coordinated activation of

biosynthetic, vasoprotective, and stress-responsive pathways. We also identified specific

molecular signatures—including NR4A3, STAT1, EGFR, and eNOS upregulation—that

distinguish patent from occluded grafts, and demonstrate that angiotensinogen-driven

fibroinflammatory remodeling is associated with graft failure. These findings provide a

mechanistic framework for developing targeted therapies to enhance graft durability and

improve long-term outcomes following CABG.

Histological Signatures of Graft Adaptation

While previous studies have predominantly focused on pathological processes leading to

graft failure[16], our histological analysis of functionally patent grafts reveals adaptive

remodeling signatures that have been largely overlooked in the literature. Traditional

conceptualizations of vein graft adaptation have emphasized "arterialization" as a

unidirectional transformation toward an arterial phenotype[3]. However, our findings

demonstrate that both arterial and venous grafts undergo substantial structural reorganization

that does not simply recapitulate native arterial architecture.

We found that arterial grafts, rather than preserving their native elastic artery

characteristics, transitioned toward a muscular artery-like configuration characterized by

marked medial thickening and an augmented internal elastic lamina. In contrast, venous

grafts developed features absent in native veins—including a prominent internal elastic

lamina, extensive medial muscularization accompanied by elastic fiber deposition, and the

formation of a novel external elastic layer. Most strikingly, despite their distinct

embryological origins and baseline morphologies, both graft types converged toward similar

caliber and structural organization while retaining unique architectural features not found in

either native arteries or veins. This morphological convergence, coupled with the

preservation of graft-specific characteristics, challenges the classical paradigm of simple

arterialization and instead supports the emergence of a distinct adaptive phenotype uniquely

tailored to the extraordinary hemodynamic demands of the coronary circulation. In this

setting, grafts are subjected to intense pulsatile stress as the first major branches arising

directly from the aortic root, receiving the full force of ventricular ejection without the

dampening effect experienced by more distal vascular territories. In contrast, occluded grafts

failed to achieve this adaptive remodeling, exhibiting severe aneurysmal dilatation, loss of

elastic integrity, and medial necrosis — indicating that successful adaptation requires

coordinated structural reorganization rather than passive exposure to arterial flow.

Adaptive Versus Pathologic Intimal Hyperplasia in Vein Grafts

Intimal hyperplasia has traditionally been regarded as a pathological hallmark of vein graft

failure in CABG, contributing to progressive luminal narrowing and eventual occlusion[17].

However, our findings fundamentally challenge this binary perspective by demonstrating that

substantial intimal thickening occurs in both functionally patent and occluded grafts, albeit

with markedly distinct structural and cellular phenotypes. Patent grafts exhibited well-

organized intimal architecture with preserved luminal patency, suggesting that intimal

hyperplasia may represent an adaptive, actively regulated biological response essential for

successful graft remodeling rather than an inherently pathological process.

This reconceptualization is strongly supported by previous studies. Jiang and colleagues,

using a bilateral jugular vein graft model that exposed grafts to varying hemodynamic

conditions, demonstrated that intimal hyperplasia develops as an adaptive response to

mechanical stress; their findings showed that grafts maintain patency through coordinated

outward remodeling[18]. Similarly, a recent animal study has reported that patent grafts exhibit

concurrent intimal hyperplasia alongside positive remodeling—characterized by lumen

preservation despite wall thickening—indicating that intimal hyperplasia is a component of

successful environmental adaptation rather than an inevitable pathological process[19].

Histologically, patent grafts in our cohort exhibited a structurally organized intima with well-

preserved endothelial integrity and a clearly demarcated internal elastic lamina, as

demonstrated by elastic tissue staining. Notably, α-SMA-positive myofibroblasts were

consistently identified within the intima of patent grafts, indicating controlled fibromuscular

remodeling mediated by tightly regulated cellular proliferation and differentiation. The

mechanistic role of myofibroblasts in adaptive intimal remodeling has been elegantly

elucidated by Vasudevan and colleagues through ultrastructural analyses of rat vein grafts[20].

They documented that adventitial fibroblasts undergo coordinated migration into the intima

and subsequent differentiation into myofibroblasts, which facilitate intimal thickening

through organized matrix synthesis. Importantly, their ultrastructural findings revealed that

adaptive intimal thickening comprises well-organized, multilayered myofibroblast assemblies

rather than disorganized collagenous fibrosis—observations that closely align with our

immunohistochemical characterization of human patent grafts. In contrast, occluded grafts

exhibited profoundly disorganized intimal architecture, fragmented elastic laminae, and

excessive extracellular matrix deposition without coordinated cellular organization. This

distinction underscores that the qualitative characteristics of intimal hyperplasia—rather than

merely its presence or extent—determine the long-term fate of the graft.

Embryonic Arterialization Signaling

The convergent adaptive remodeling observed in both arterial and venous grafts may reflect the reactivation of embryonic arterialization signaling pathways. During vascular development, arterial identity is established through the coordinated activation of VEGF-Notch-ERK signaling cascades, which promote arterial-specific gene expression and structural maturation. This developmental program is particularly well-characterized in coronary artery embryogenesis, where VEGFR2-ERK-eNOS and NOTCH4 signaling axes are distinctly upregulated and activated in endothelial cells during vascular morphogenesis[21–23]. Specifically, VEGFR2 signaling activates the downstream ERK pathway, which is critical for endothelial proliferation, branching morphogenesis, and lumen formation in both coronary and general vascular embryology. eNOS is also upregulated and activated downstream of VEGFR2 in response to VEGF stimulation in the endothelium, mediating vasodilation and angiogenic responses required for vessel remodeling. Regarding NOTCH4, this receptor (along with NOTCH1) is specifically expressed in embryonic vascular endothelium and becomes markedly upregulated and activated during arterial specification and network patterning. Activation of NOTCH4 in endothelial cells is essential for correct branching morphogenesis and vascular patterning in the developing embryo, including the coronary vasculature, with increased NOTCH4 expression further supporting arterial specification from progenitor cells[22,23]

Our proteomic analysis revealed that patent grafts exhibited significant enrichment of key components of the developmental arterialization program: VEGFR2 signaling (Figure 3A and Figure S9B), ERK1/2 MAPK pathways (FDR $< 5 \times 10^{-4}$; Supplementary Data), and NOTCH4 signaling across all comparison groups, with consistently positive directionality (AGP vs. AC: z-score=3.317; VGP vs. VC: z-score=2.673; VGP vs. VGO: z-score=4.472). This pathway enrichment was accompanied by enhanced eNOS expression in both the

luminal endothelium and adventitial neovessels (Figure 6). The concurrent activation of

VEGFR2-ERK-eNOS and NOTCH4 pathways—mirroring the signaling architecture that

drives arterial specification during coronary embryogenesis—strongly supports the

hypothesis that successful graft adaptation involves partial recapitulation of developmental

arterialization programs. This coordinated pathway reactivation may represent an

evolutionarily conserved adaptive mechanism whereby adult vessels redeploy embryonic

programs to achieve structural and functional optimization under hemodynamic stress.

Graft Failure is Characterized by Angiotensin-Driven Fibroinflammatory Remodeling

and Loss of Biosynthetic Resilience

Our upstream regulator and regulator effects analysis identified angiotensin as a key activated

node in occluded vein grafts. The angiotensin-centered network predicted a coordinated

upregulation of extracellular matrix proteins (COL5A1, COL5A2, DCN, BGN, LUM,

THBS2), consistent with fibrotic remodeling and reduced vascular compliance.

Immunohistochemistry confirmed increased angiotensin expression in the medial and

adventitial regions of occluded vein grafts, supporting the hypothesis of an angiotensin-

driven fibroinflammatory remodeling mechanism underlying graft occlusion in our cohort.

This observation aligns with mechanistic insights from the renin-angiotensin system

literature: angiotensin II mediates vascular remodeling through vascular smooth muscle cell

proliferation, migration, extracellular matrix production, inflammation, and oxidative stress,

and is implicated in the pathogenesis of intimal hyperplasia and graft degeneration[24].

Moreover, polymorphism studies of the angiotensin-converting enzyme (ACE) gene in

saphenous vein graft atherosclerosis further support a link between angiotensin signaling and

graft pathology[25]. Given that we observed significantly greater luminal deposition of

angiotensin in occluded grafts compared to patent grafts, our data strengthen the hypothesis

that angiotensin signaling is not merely a bystander but may function as an upstream driver of

graft failure.

From a therapeutic perspective, although no large randomized trials have specifically

evaluated angiotensin blockade for improving saphenous vein graft patency, preclinical and

indirect clinical evidence suggest that targeting the renin-angiotensin axis may be beneficial.

For example, ACE inhibitors and angiotensin receptor blockers (ARBs) reduce vascular

remodeling and intimal hyperplasia in experimental models[24]. Observational analyses in post-

CABG populations have examined ACE inhibitor use, with one cohort study reporting that

post-CABG ACE/ARB therapy was associated with a lower hazard of major adverse

cardiovascular events at 1 and 5 years[26]. Although this does not directly measure graft-

specific patency, it is biologically plausible that the beneficial effect partly reflects improved

graft adaptation.

Thus, in light of our finding that angiotensin accumulation is specifically associated with

occluded grafts—and that this pattern corresponds to a predicted upstream regulatory

network driving extracellular matrix remodeling—we propose that angiotensin signaling

represents a novel, modifiable pathway in graft failure.

Potential Role of eNOS in Perivascular Adipose Tissue as a Key Vasoprotective

Modulator

eNOS is a key regulator of vascular homeostasis, mediating anti-thrombotic, anti-

inflammatory, and vasodilatory effects that maintain conduit integrity. Previous clinical and

experimental studies have demonstrated that increased eNOS activity within the luminal

endothelium correlates with improved graft patency and reduced vascular complications in

saphenous vein grafts used for coronary bypass surgery[27,28]. However, these investigations

have primarily focused on endothelial eNOS localized to the main vessel lumen, without

exploring its potential roles in the perivascular compartment.

In our study, we identified a distinct spatial pattern of eNOS upregulation extending

beyond the endothelium to the adventitia and PVAT in patent saphenous vein grafts. Notably,

intense eNOS expression was observed in the vasa vasorum and surrounding supportive

structures—an expression pattern absent in pre-implantation segments. This finding suggests

that the PVAT-adventitial domain may actively participate in graft adaptation, serving as a

secondary vasoprotective interface that complements luminal endothelial function.

PVAT is a metabolically active layer that release paracrine mediators—such as

adipokines and hydrogen sulfide—which enhance eNOS activity and exert anti-inflammatory

and anti-fibrotic effects on the vascular wall. Our histological evidence supports the concept

that preserving PVAT during no-touch vein harvesting not only maintains endothelial

integrity but also preserves an eNOS-enriched microenvironment within the perivascular

compartment. This dual-source nitric oxide signaling likely contributes to the superior long-

term patency observed with the no-touch technique.

Limitations and Future Study

This study has several limitations that warrant acknowledgment. First, the number of

available graft specimens was limited, reflecting the rarity and clinical challenges of

obtaining well-preserved human graft tissues explanted during cardiac transplantation.

Despite the small cohort size, each sample provided valuable biological insights into long-

term graft adaptation—a phenomenon rarely accessible in human studies. Second, surgical

techniques were not fully standardized across cases; variations in harvesting methods and

anastomotic configurations, including Y-graft constructs, may have influenced hemodynamic

and remodeling responses. Third, all molecular analyses were performed on bulk tissue,

precluding cell type- and layer-specific resolution. Although immunohistochemistry provided

partial spatial context for key markers such as eNOS, it could not fully delineate molecular

signals originating from distinct vascular compartments, including the intima, media, and

adventitia.

To address these limitations, future studies should incorporate spatially resolved

molecular profiling techniques—such as spatial transcriptomics and proteomics—to

characterize cell type- and layer-specific remodeling signatures within arterial and venous

grafts. Particular emphasis should be placed on the intimal compartment, where

hemodynamic and metabolic cues converge to influence the trajectory of intimal remodeling.

Spatial omics approaches will be essential for identifying the molecular regulators that

determine whether intimal proliferation stabilizes adaptively or progresses toward

pathological occlusion. Integrating these spatial insights with functional validation in

experimental graft models will enhance mechanistic understanding and guide the

development of targeted therapeutic strategies to improve long-term graft patency.

Conclusion

In this study, we found that arterial and venous grafts with long-term patency undergo a

convergent remodeling process and establish a distinct, graft-specific adaptive phenotype

characterized by structural maturation, reconstitution of elastic architecture, and a shared

biosynthetic-proteomic program uniquely shaped by the extraordinary hemodynamic

environment of the coronary circulation. Adventitial eNOS-positive neovascularization

emerged as a hallmark of adaptive remodeling, suggesting a vasoprotective mechanism that

sustains graft homeostasis. Collectively, these findings identify a biosynthetic-vasoprotective

remodeling axis as a central determinant of graft durability and propose that therapeutic

modulation of shared adaptive pathways may enhance long-term graft patency following

coronary artery bypass grafting.

SOURCES OF FUNDING

This study was supported by grant No 0420254100 [2025–3000] from the SNUH Research

Fund; and a focused clinical research grant-in-aid from the Seoul Metropolitan Government

Seoul National University (SMG-SNU) Boramae Medical Center (04-2023-0005).

DISCLOSURES

None.
587

588 **SUPPLEMENTAL MATERIAL**

589 SUPPLEMENTAL METHODS

590 SUPPLEMENTAL RESULTS

591 Tables S1 and S2

592 Figures S1–S14

REFERENCES

1. Gharibeh L, Ferrari G, Ouimet M, Grau JB. Conduits' Biology Regulates the Outcomes of Coronary Artery Bypass Grafting. *JACC: Basic Transl Sci*. 2021;6(4):388-396. doi:10.1016/j.jacbts.2020.11.015

2. Taggart DP. Current status of arterial grafts for coronary artery bypass grafting. *Ann Cardiothorac Surg*. 2013;2(4):427-430. doi:10.3978/j.issn.2225-319x.2013.07.21

3. Muto A, Model L, Ziegler K, Eghbalieh SDD, Dardik A. Mechanisms of Vein Graft Adaptation to the Arterial Circulation. *Circ J*. 2010;74(8):1501. doi:10.1253/circj.cj-10-0495

4. Gaudino M, Benedetto U, Fremes S, Ballman K, Biondi-Zoccai G, Sedrakyan A, Nasso G, Raman J, Buxton B, Hayward PA, Moat N, Collins P, Webb C, Peric M, Petrovic I, Yoo KJ, Hameed I, Franco AD, Moscarelli M, Speziale G, Puskas JD, Girardi LN, Hare DL, Taggart DP, Investigators R. Association of Radial Artery Graft vs Saphenous Vein Graft With Long-term Cardiovascular Outcomes Among Patients Undergoing Coronary Artery Bypass Grafting. *JAMA*. 2020;324(2):179-187. doi:10.1001/jama.2020.8228

5. Cuminetti G, Gelsomino S, Curello S, Lorusso R, Maessen JG, Hoorntje JCA. Contemporary use of arterial and venous conduits in coronary artery bypass grafting: anatomical, functional and clinical aspects. *Neth Hear J*. 2017;25(1):4-13. doi:10.1007/s12471-016-0919-2

6. Gaudino M, Antoniades C, Benedetto U, Deb S, Franco AD, Giammarco GD, Fremes S, Glineur D, Grau J, He GW, Marinelli D, Ohmes LB, Patrono C, Puskas J, Tranbaugh R, Girardi LN, Taggart DP, Ruel M, Bakaeen FG. Mechanisms, Consequences, and Prevention

of Coronary Graft Failure. *Circulation*. 2017;136(18):1749-1764.

doi:10.1161/circulationaha.117.027597

7. Vries MR de, Simons KH, Jukema JW, Braun J, Quax PHA. Vein graft failure: from

pathophysiology to clinical outcomes. *Nat Rev Cardiol*. 2016;13(8):451-470.

doi:10.1038/nrcardio.2016.76

8. Dashwood MR, Savage K, Dooley A, Shi-Wen X, Abraham DJ, Souza DSR. Effect of

Vein Graft Harvesting on Endothelial Nitric Oxide Synthase and Nitric Oxide Production.

Ann Thorac Surg. 2005;80(3):939-944. doi:10.1016/j.athoracsur.2005.03.042

9. Bikdeli B, Hassantash SA, Pourabdollah M, Kalantarian S, Sadeghian M, Afshar H, Sabeti

S, Marzban M, Ahmadi H, Mohammadi F. Histopathologic Insight into Saphenous Vein

Bypass Graft Disease. *Cardiology*. 2012;123(4):208-215. doi:10.1159/000343269

10. Tian M, Wang X, Feng W, Wang H, Liu S, Liu Z, Chen Y, Miao Q, Su P, Li X, Wang Y,

Lu B, Chen K, Zhang C, Hu S. No-touch versus conventional vein in coronary artery bypass

grafting: three year follow-up of multicentre randomised PATENCY trial. *BMJ*.

2025;389:e082883. doi:10.1136/bmj-2024-082883

11. Mikami T, Dashwood MR, Kawaharada N, Furuhashi M. An Obligatory Role of

Perivascular Adipose Tissue in Improved Saphenous Vein Graft Patency in Coronary Artery

Bypass Grafting. *Circ J*. 2024;88(6):845-852. doi:10.1253/circj.cj-23-0581

12. Saito T, Kurazumi H, Suzuki R, Matsunaga K, Tsubone S, Lv B, Kobayashi S, Nagase T,

Mizoguchi T, Samura M, Suehiro K, Harada T, Morikage N, Mikamo A, Hamano K.

Perivascular Adipose Tissue Is a Major Source of Nitric Oxide in Saphenous Vein Grafts

Harvested via the No-Touch Technique. *J Am Hear Assoc: Cardiovasc Cerebrovasc Dis*.

2020;11(3):e020637. doi:10.1161/jaha.120.020637

13. Dashwood MR, Dooley A, Shi-Wen X, Abraham DJ, Souza DSR. Does Periadventitial

Fat-Derived Nitric Oxide Play a Role in Improved Saphenous Vein Graft Patency in Patients

Undergoing Coronary Artery Bypass Surgery? *J Vasc Res*. 2007;44(3):175-181.

doi:10.1159/000099833

14. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT,

Legeay M, Fang T, Bork P, Jensen LJ, Mering C von. The STRING database in 2021:

customizable protein–protein networks, and functional characterization of user-uploaded

gene/measurement sets. *Nucleic Acids Res*. 2021;49(D1):D605-D612.

doi:10.1093/nar/gkaa1074

15. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C,

Chanda SK. Metascape provides a biologist-oriented resource for the analysis of systems-

level datasets. *Nat Commun*. 2019;10(1):1523. doi:10.1038/s41467-019-09234-6

16. Owens CD, Gasper WJ, Rahman AS, Conte MS. Vein graft failure. *J Vasc Surg*.

2015;61(1):203-216. doi:10.1016/j.jvs.2013.08.019

17. Goldstein DJ, Chang HL, Mack MJ, Voisine P, Gammie JS, Marks ME, Iribarne A,

Vengrenyuk Y, Raymond S, Taylor BS, Dagenais F, Ailawadi G, Chu MWA, DiMaio JM,

Narula J, Moquete EG, O'Sullivan K, Williams JB, Crestanello JA, Scavo V, Puskas JD,

Acker MA, Gillinov M, Gelijns AC, O'Gara PT, Moskowitz AJ, Alexander JH, Bagiella E.

Intimal hyperplasia, saphenous vein graft disease, and clinical outcomes: Insights from the

CTSN VEST randomized trial. *J Thorac Cardiovasc Surg*. 2024;167(5):1782-1792.e5.

doi:10.1016/j.jtcvs.2022.10.034

18. Jiang Z, Wu L, Miller BL, Goldman DR, Fernandez CM, Abouhamze ZS, Ozaki CK,

Berceli SA. A novel vein graft model: adaptation to differential flow environments. *Am J*

Physiol-Hear Circ Physiol. 2004;286(1):H240-H245. doi:10.1152/ajpheart.00760.2003

19. Yao G, Li H, Zuo X, Wang C, Xiao Y, Zhao Y, Wang X. Oscillatory shear stress

promotes vein graft intimal hyperplasia via NADPH oxidase-related pathways. *Front Surg*.

2023;10:1073557. doi:10.3389/fsurg.2023.1073557

20. Stark VK, Warner TF, Hoch JR. An ultrastructural study of progressive intimal

hyperplasia in rat vein grafts. *J Vasc Surg*. 1997;26(1):94-103. doi:10.1016/s0741-

5214(97)70152-6

21. Kroll J, Waltenberger J. VEGF-A Induces Expression of eNOS and iNOS in Endothelial

Cells via VEGF Receptor-2 (KDR). *Biochem Biophys Res Commun*. 1998;252(3):743-746.

doi:10.1006/bbrc.1998.9719

22. Luo W, Garcia-Gonzalez I, Fernández-Chacón M, Casquero-Garcia V, Sanchez-Muñoz

MS, Mühleder S, Garcia-Ortega L, Andrade J, Potente M, Benedito R. Arterialization

requires the timely suppression of cell growth. *Nature*. 2021;589(7842):437-441.

doi:10.1038/s41586-020-3018-x

23. Wythe JD, Dang LTH, Devine WP, Boudreau E, Artap ST, He D, Schachterle W, Stainier

DYR, Oettgen P, Black BL, Bruneau BG, Fish JE. ETS Factors Regulate Vegf-Dependent

Arterial Specification. *Dev Cell*. 2013;26(1):45-58. doi:10.1016/j.devcel.2013.06.007

24. Osgood MJ, Harrison DG, Sexton KW, Hocking KM, Voskresensky IV, Komalavilas P,

Cheung-Flynn J, Guzman RJ, Brophy CM. Role of the Renin–Angiotensin System in the

Pathogenesis of Intimal Hyperplasia: Therapeutic Potential for Prevention of Vein Graft

Failure? *Ann Vasc Surg*. 2012;26(8):1130-1144. doi:10.1016/j.avsg.2011.12.001

25. Zeinali N, Hashemi M, Mirmohammadsadeghi M, Mirmohammadsadeghi H, Eskandari

N, Sabzghabaee AM. Association of Angiotensin-Converting Enzyme Genotype,

Insertion/Deletion Polymorphism and Saphenous Vein Graft Atherosclerosis in Iranian

Patients. *Rev Bras Cir Cardiovasc*. 2015;30(5):557-561. doi:10.5935/1678-9741.20150069

26. Kalavrouziotis D, Buth KJ, Cox JL, Baskett RJ. Should all patients be treated with an

angiotensin-converting enzyme inhibitor after coronary artery bypass graft surgery? The

impact of angiotensin-converting enzyme inhibitors, statins, and β-blockers after coronary

artery bypass graft surgery. *Am Hear J*. 2011;162(5):836-843. doi:10.1016/j.ahj.2011.07.004

27. Otsuka F, Yahagi K, Sakakura K, Virmani R. Why is the mammary artery so special and

what protects it from atherosclerosis? *Ann Cardiothorac Surg*. 2013;2(4):519-526.

doi:10.3978/j.issn.2225-319x.2013.07.06

28. Collins MJ, Li X, Lv W, Yang C, Protack CD, Muto A, Jadlowiec CC, Shu C, Dardik A.

Therapeutic strategies to combat neointimal hyperplasia in vascular grafts. *Expert Rev*

Cardiovasc Ther. 2012;10(5):635-647. doi:10.1586/erc.12.33

Figure Legends

Figure 1.

(A) Explanted heart with attached saphenous vein grafts harvested using the no-touch

technique. Gross photograph of an explanted heart from a patient with prior coronary artery

bypass grafting (yellow arrow). The saphenous vein graft conduits remain attached to the

heart, with perivascular supportive tissue preserved as a result of the no-touch harvesting

technique.

(B) Histomorphologic remodeling of arterial and venous conduits following grafting.

Blue arrowheads indicate the internal elastic lamina, marking the boundary between the

intima (I) and media (M). Green arrowheads indicate the external elastic lamina, marking the

boundary between the media (M) and adventitia. Yellow labels denote the intima (I) and

media (M) within each graft.

(C) Quantitative morphometric analysis of arterial and venous graft remodeling.

Boxplots show the lumen area (upper left), lumen circumference (upper right), intimal area

(lower left), and medial thickness (lower right) across native arterial conduits (AC), arterial

grafts (AGP), native venous conduits (VC), patent venous grafts (VGP), and occluded venous

grafts (VGO). Statistical comparisons are indicated above each plot (*$P<0.05$; **$P<0.01$;

***$P<0.001$; ****$P<0.0001$; ns, not significant).

(D) Intimal and medial morphology across arterial and venous graft states.

Elastic staining of native arteries (AC) and veins (VC), their patent graft counterparts (AGP,

VGP), and occluded vein grafts (VGO).

(E) Histochemical and immunohistochemical characteristics of patent versus occluded vein

grafts.

Elastic staining showed a well-organized intima–media structure in patent vein grafts (VGP),

whereas occluded grafts (VGO) exhibited disrupted elastic layers. Masson Trichrome

staining demonstrated structured fibrosis with preserved layering in VGP, but dense,

disorganized collagen in VGO. αSMA staining revealed abundant, organized smooth muscle–

type cells in the VGP intima (blue arrow) and near-complete absence of αSMA-positive cells

in the fibrotic VGO intima (red arrow).

Figure 2

(A) Number of identified proteins

Venn diagram showing overlap of identified proteins across five vascular tissue groups.

A total of 4,836 proteins were identified across all samples using mass spectrometry–based

proteomic analysis. The five groups included vein control (VC, n=4,490 proteins), artery

control (AC, n=4,260), patent vein grafts (VGP, n=4,591), patent artery grafts (AGP,

n=4,605), and occluded vein grafts (VGO, n=4,298). Among these, 4,012 proteins were

commonly detected across all five groups, while subsets of proteins were uniquely identified

in specific conditions. The number of shared and group-specific proteins is indicated within

each section of the Venn diagram.

(B) Unsupervised 2-dimensional PCA showing distinct clustering patterns across vascular

groups. AC and VC samples separated widely along PC1, reflecting their differing baseline

proteomic profiles. AGP and VGP samples clustered closely, indicating similar post-grafting

proteomic changes. VGO samples formed a cluster that was distinct from both native vessels

and patent grafts.

(C) Differential protein expression among native vessels and graft types.

Volcano plot comparing VC and AC showing 702 proteins enriched in VC and 164 enriched

in AC, indicating clear baseline proteomic differences between veins and arteries.

Comparison of AC and AGP identified 134 proteins enriched in AC and 472 enriched in

AGP, consistent with extensive arterial graft remodeling. Comparison of VC and VGP

revealed 225 proteins enriched in VC and 380 enriched in VGP, reflecting substantial

remodeling following venous grafting. Direct comparison of AGP and VGP showed only a

minimal number of differentially expressed proteins, underscoring the post-grafting similarity

between arterial and venous conduits. Comparison of VGP and VGO identified 475 proteins

enriched in VGP and 220 enriched in VGO, highlighting the molecular divergence between

functional and occluded vein grafts.

(D) Canonical pathway enrichment analysis of arterial and venous grafts.

(Left) Canonical pathway enrichment analysis for AGP versus AC showed significant

activation of translation-related pathways (e.g., initiation, elongation, termination, and SRP-

dependent targeting), EIF2/GCN2 signaling, nonsense-mediated decay, and proteostasis-

associated programs, including microautophagy and BAG2 signaling.

(Right) Canonical pathway enrichment analysis for VGP versus VC showed similar

enrichment in translation-associated pathways, including EIF2/GCN2, and proteostasis

pathways, along with additional processes related to extracellular matrix organization,

collagen biosynthesis, integrin signaling, and neutrophil degranulation.

(E) Hierarchical clustering reveals group-specific proteomic signatures across vascular

samples. Clustering of the 2,492 most variable proteins generated distinct groups

corresponding to major vascular phenotypes. Cluster 7 consisted predominantly of VC

samples, while Cluster 6 was largely composed of AC samples, reflecting baseline proteomic

differences between native veins and arteries. Cluster 5 was enriched with VGO samples and

displayed a molecular pattern distinct from all other groups. Cluster 4 contained proteins

consistently upregulated in both AGP and VGP, highlighting the close similarity of

proteomic profiles in functional arterial and venous grafts.

Figure 3. Integrated functional analysis differentiates adaptive remodeling in patent grafts from maladaptive processes in occluded grafts.

(A) Pathway enrichment analysis of Cluster 4 (VGP + AGP) and Cluster 5 (VGO).

Cluster 4 showed coordinated enrichment in RNA metabolism, mRNA processing, translation

initiation/elongation, ribonucleoprotein biogenesis, and pathways supporting extracellular

matrix (ECM) organization and cytoskeletal adaptation. In contrast, Cluster 5 exhibited

predominant enrichment of complement and coagulation cascades, acute inflammatory

signaling, wound healing, and disorganized ECM remodeling.

(B) Protein–protein interaction (PPI) networks for Cluster 4

Cluster 4 networks revealed densely interconnected ribosomal/translation hubs and organized

ECM-remodeling subnetworks, consistent with an active, reparative remodeling phenotype.

(C) Metascape pathway enrichment analysis of functional and obstructed grafts (Cluster 4

and Cluster 5). Pathway enrichment analysis was conducted using Metascape

(www.metascape.org), which integrates multiple curated ontologies, including GO, KEGG,

and Reactome. (A) Pathways enriched in functional grafts, including vein graft patent (VGP)

and artery graft patent (AGP) groups. (B) Pathways enriched in the vein graft obstructed

(VGO) group. Each bubble represents a biological pathway, where the x-axis shows

enrichment significance ($-\log_{10}$(q-value)), the y-axis shows pathway names, bubble size

reflects the percentage of overlapping genes (Overlap%), and bubble color reflects

significance strength, where a more intense pink indicates higher significance. Cluster 4

(VGP + AGP) demonstrated enrichment of pathways related to RNA metabolism, VEGFA–

VEGFR2 signaling, receptor tyrosine kinase activity, protein maturation, intracellular

transport, lysosomal processes, and immune-regulatory programs—reflecting a

biosynthetically active, structurally adaptive, and immune-balanced phenotype characteristic

of functional grafts. In contrast, Cluster 5 (VGO) was enriched for complement and

coagulation pathways, NABA matrisome and ECM-associated programs, platelet activation,

plasma lipoprotein assembly, and IGFBP-mediated IGF transport regulation—defining a

matrix-disruptive, thrombosis-prone, and growth factor–dysregulated environment associated

with graft obstruction.

Figure 4. Upstream regulatory programs and immunohistochemical validation in arterial and

venous grafts

(A) Arterial grafts (AGP vs. AC). Upstream regulator analysis revealed coordinated

activation of metabolic and proliferative regulators, including IGF1, RUNX2, NR4A3, PGC-

1α, EGFR, and MYC.

(B) Venous grafts (VGP vs. VC). Patent vein grafts showed activation of proliferative–

biosynthetic regulators, including STAT1, SRC, PTGES, RETN, and G6PD, accompanied by

predicted inhibition of vascular contractility regulators such as SRF and MRTFA—indicating

a shift toward a synthetic, remodeling-competent phenotype.

(C) Immunohistochemical validation. IHC confirmed increased expression of NR4A3,

EGFR, and STAT1 in both AGP and VGP ($p<0.01$), identifying these as conserved adaptive

regulators across graft types. Arterial grafts showed specific upregulation of IGF1 and

RUNX2, while venous grafts exhibited higher expression of RETN, SRC, and PTGES

compared with their respective controls, validating graft-type–specific adaptive signaling

predicted by upstream analyses.

Figure 5. Distinct upstream signaling patterns and angiotensin-associated extracellular matrix

remodeling in patent and occluded venous grafts

(A) Upstream regulator analysis comparing VGP and VGO using Ingenuity Pathway

Analysis. Orange nodes indicate predicted activated regulators, while blue nodes indicate

inhibited regulators. In VGP relative to VGO, STAT3 downregulation, coupled with

reciprocal EGFR upregulation, emerged as a key regulatory axis distinguishing successful

graft adaptation.

(B) STAT3 expression assessed by immunohistochemistry and quantitative analysis. Upper

panels: Representative STAT3 staining in VGP (left) and VGO (right) demonstrating

minimal STAT3 expression in patent grafts versus increased nuclear positivity in occluded

grafts. Scale bars=100 μm. Lower panel: Quantification of STAT3-positive cells showing a

trend toward elevated expression in VGO (P=0.08). n=8–10 per group.

(C) Upstream regulator analysis identified angiotensin as a central activated node in VGO.

The angiotensin-centered regulatory network illustrates predicted downstream targets related

to the extracellular matrix. Node size reflects network connectivity, while connecting edges

represent predicted regulatory or protein–protein interactions.

(D) Expanded angiotensin regulatory network highlighting extracellular matrix structural

proteins (orange), matrix-associated signaling molecules (blue), and downstream predicted

biological functions (pink). Color intensity reflects the predicted activation state. Upregulated

matrix components include fibrillar collagens (COL5A1, COL5A2), small leucine-rich

proteoglycans (decorin, biglycan, lumican), and thrombospondin-2, which collectively

contribute to enhanced matrix tensile strength in occluded grafts.

(E) Immunohistochemical validation of the expression of angiotensinogen, the precursor of

angiotensin. Left column: Two representative cases of patent venous grafts (VGP), each from

a different patient, demonstrating minimal angiotensinogen staining throughout the vessel

wall. Right column: Two representative cases of occluded venous grafts (VGO), each from a

different patient, showing prominent angiotensinogen deposition in the media and adventitia,

with dense accumulation within fibrotic regions. All images are at the same magnification.

Scale bars=200 μm. Right panel: Quantitative H-score analysis revealed significantly

elevated angiotensinogen expression in VGO versus VGP (**$P<0.01$). Box plots display the

median and interquartile range (n=8–12 per group).

Figure 6. Upregulation of eNOS in luminal and adventitial endothelial cells in arterial grafts

(A-C) eNOS expression in the luminal endothelium.

(A) Representative immunohistochemistry showing weak eNOS expression in control vessels

(AC, VC) and marked upregulation in patent grafts (AGP, VGP). Occluded grafts (VGO)

exhibit endothelial denudation with minimal eNOS signal. Scale bars=50 µm.

(B) Histogram analysis of single-cell optical density revealed a rightward shift in the graft

endothelium, indicating enhanced eNOS expression.

(C) Proportion of eNOS-positive cells (optical density >0.2) was significantly higher in AGP

versus AC (***$P<0.001$) and in VGP versus VC (**$P<0.01$).

(D, E) eNOS expression in the adventitial vasa vasorum.

(D) Representative images show enhanced neovascularization with eNOS-positive

endothelium in AGP (green arrow) and VGP (cyan arrow) compared with controls. VGO

(magenta arrow) demonstrated sparse vasculature lacking eNOS expression. Scale bars=100

μm.

(E) Left: Total microvascular perimeter was significantly larger in AGP versus AC

(***$P<0.001$). Right: eNOS-positive adventitial vessel count was significantly higher in AGP

versus AC (***$P<0.001$) and in VGP versus VC (**$P<0.01$). Box plots display median and

interquartile range. Statistical analysis was performed using one-way ANOVA with Tukey's

post hoc test.

880