Neural Integration of Affective Prosodic and Semantic Cues in Non-literal Forms of Speech Understanding

Adrien Wittmann,[1*] Leonardo Ceravolo,[2] Audrey Mayr,[2] Didier Grandjean[2]

[1]Neuroscience of Emotions and Affective Dynamics lab, Swiss Center for Affective Sciences,

Department of Psychology and Educational Sciences, University of Geneva,

Biotech campus, Chemin des Mines 9, 1202 Geneva, Switzerland

[2]Neuroscience of Emotions and Affective Dynamics lab, Swiss Center for Affective Sciences,

Department of Psychology and Educational Sciences, University of Geneva,

Unimail, Chemin des Mines 9, 1202 Geneva, Switzerland

*Correspondence: adrien.wittmann@unige.ch

March 26, 2026

Keywords: integration, irony, prosody, sarcasm, semantics, ToM

Abstract: Emotions in speech are conveyed through both semantics and prosody—voice 'melody', yet how listeners integrate these cues in the brain remains unclear. We investigated non-literal forms of speech, such as irony and sarcasm, where understanding beyond literal meaning relies on the dynamic interplay between affective prosodic and semantic cues, alongside theory of mind (ToM). We used functional magnetic resonance imaging while participants listened to short dialogues between two characters, which varied in prosody and semantics to convey either literal or non-literal meanings. Behaviorally, semantics and prosody interacted in shaping participants' evaluations, although the data suggest a prosody dominance effect. Non-literal speech engaged a distributed network including the bilateral inferior frontal gyrus, temporal speech regions, and ToM areas, with ROI analyses revealing heterogeneous prosody-semantics integration profiles across regions and tasks. Taken together, our data clarify the behavioral and neural underpinnings of the integration of prosody and semantics

in non-literal speech and open new venues in this field.

Author note. The authors made the following contributions. Adrien Wittmann: Conceptualization, Methodology, Data curation, Formal analysis, Software, Investigation, Validation, Visualisation, Writing – Original Draft Preparation, Writing – Review & Editing; Leonardo Ceravolo: Methodology, Software, Writing – Review & Editing; Audrey Mayr: Conceptualization, Software, Methodology, Investigation, Validation; Didier Grandjean: Funding acquisition, Supervision, Conceptualization, Writing – Review & Editing.

Correspondence concerning this article should be addressed to Adrien Wittmann, Neuroscience of Emotions and Affective Dynamics lab, Swiss Center for Affective Sciences, Department of Psychology and Educational Sciences, University of Geneva, Biotech campus, Chemin des Mines 9, 1202 Geneva, Switzerland. E-mail: adrien.wittmann@unige.ch

1   Introduction

Human communication employs various modalities, yet language stands out for its capacity to convey precise, abstract, and elaborated expressions. Language may serve many purposes, among them the communication of emotions or attitudes. These can be conveyed in different ways in speech—either through prosody, the intonation and rhythm of the voice, or through semantics, using words and their meanings. These two types of emotional information are differentially communicated (Castelluccio et al., 2016), interactive (Astésano et al., 2004), and activate common brain regions (Buchanan et al., 2000; Meyer et al., 2003; R. L. C. Mitchell, 2007). However, the way these different cues are integrated at the behavioral and brain levels is still little understood.

A particularly revealing context for studying this integration is non-literal speech, such as irony and sarcasm. Irony and sarcasm are forms of communication that express ideas opposite to their literal meaning (Bosco et al., 2017; Matsui et al., 2016; Searle, 1979). Irony involves incongruity between explicit and intended meaning, often creating humor or satire—it can express praise through seemingly negative language or, in sarcastic irony, convey derision through superficially positive language. Understanding these non-literal forms of speech hinges on the incongruence between the lexico-semantic content of the utterance and the context or expectations about the speaker's attitude, while paralinguistic cues—such as tone of voice—play an important role in facilitating their recognition (Mauchand et al., 2020; Nakamura et al., 2022). To better understand prosodic and semantic integration in this context, researchers have often used an "incongruity" paradigm (Kotz et al., 2015; Lin et al., 2020; R. L. C. Mitchell, 2006; Pell

et al., 2011; Wittfoth et al., 2009). When emotional prosodic and semantic cues are incongruent, as in irony and sarcasm, listeners must integrate these cues to effectively decipher the intended emotional message (Matsui et al., 2016). Beyond integration, listeners need to make inferences about others' feelings, attitudes, and mental states—concepts referred to as theory of mind (ToM; Zhu and Wang 2020). The primary goal of our research was thus to shed light on the brain network involved in the integration of semantic and prosodic affective cues during the processing of non-literal forms of speech using functional Magnetic Resonance Imaging (fMRI), across different tasks mirroring a hierarchy of processing demands, from lower-order perceptual decoding of prosody and semantics, through pragmatic inference of ironic and sarcastic meaning, to higher-order mentalizing processes.

To date, many studies have examined the neural networks involved in perceiving emotions in speech (Castelluccio et al., 2016; Frühholz & Belin, 2018; Seydell-Greenwald et al., 2020). One key channel is prosody, expressed through suprasegmental features such as tone modulation (Lin et al., 2020), which involves acoustic changes in pitch (F0), loudness, rhythm, articulation rate, and timbre (Brück et al., 2011; Scherer, 2003; Wildgruber et al., 2006). Although lesion studies have long suggested right-hemisphere dominance in emotional prosody processing (Borod et al., 2002; Ross & Monnot, 2008; Witteman et al., 2011), this strict lateralization has been challenged by evidence of bilateral STC involvement (Brück et al., 2011; Kotz et al., 2006), and more recent models describe it as a dynamic and multi-stage process encompassing sensory analysis, emotional inference, and explicit evaluation (Brück et al., 2011; Frühholz & Grandjean, 2012; Kotz & Paulmann, 2011; Schirmer & Kotz, 2006; Wildgruber et al., 2009). Basic acoustic features are first extracted in the bilateral auditory cortices, particularly in the superior temporal cortex (STC) or temporal voice-sensitive areas (TVAs; Belin et al. 2000; Grandjean 2021), which are consistently activated regardless of attention or task demands (Ethofer et al., 2006; Grandjean et al., 2005; Witteman et al., 2011), before being integrated into an emotional percept in higher temporal regions and subsequently evaluated explicitly. During this final stage, the right IFG and orbitofrontal cortex (OFC) are primarily recruited (Brück et al., 2011; Kotz & Paulmann, 2011), whereas the left IFG appears to integrate emotional prosody with other aspects of speech such as semantics, with greater activation observed when prosodic and semantic cues are incongruent (Kotz et al., 2015; R. L. C. Mitchell, 2006; Schirmer et al., 2004; Wiethoff et al., 2008). However, the precise neural mechanisms supporting this integration—particularly in irony and sarcasm, which feature incongruence between contextual cues, prosody, and semantics—remain poorly understood.

Emotion in speech is also conveyed through words and utterances whose meanings, associations, and contexts may evoke emotional significance. Neuroimaging and lesion studies have identified key regions for semantic processing, primarily in the left hemisphere. The anterior temporal lobe (ATL), including the anterior STS and gyrus, is crucial for sentence comprehension (Friederici, 2011; Matchin & Hickok,

2020), and has been shown to support compositional semantic processing (Humphries et al., 2006; Vandenberghe et al., 2002; Westerlund & Pylkkänen, 2014) or syntactic processing, or both (Rogalsky & Hickok, 2009). The middle temporal gyrus (MTG) integrates lexical information in context (Binder et al., 2009; Hoffman et al., 2015), and the IFG responds more strongly to abstract and emotional words during semantic processing (Hoffman et al., 2015; Pauligk et al., 2019). Processing emotional sentences increases activation in the left IFG, the posterior left STS, and the medial prefrontal cortex (mPFC)—part of the mentalizing network (Beaucousin et al., 2006). Finally, the angular gyrus (AG) acts as a supramodal integration hub for high-level semantic tasks such as concept retrieval and combination (Binder et al., 2009).

Research on the neural processing of irony and sarcasm has mainly focused on reading, identifying key regions such as the IFG and the mPFC (Bosco et al., 2017; Filik et al., 2019; Shibata et al., 2010; Uchiyama et al., 2006). However, only a few studies have examined the auditory processing of irony. Matsui et al. (2016) investigated how contextual and prosodic cues interact during sarcasm processing, finding significant activations in the rostroventral left IFG, proposed to integrate discourse context, utterance meaning, and affective prosody. Nakamura et al. (2022) examined context-prosody interactions during sarcasm comprehension, reporting activations for context-content incongruity in the anterior rostral mPFC, temporal pole, and bilateral cerebellum, and for content-prosody incongruity in the bilateral amygdala, with interaction effects in the salience network and left IFG. Finally, Obert et al. (2016) found that irony processing activated the bilateral IFG, posterior left STG, mPFC, and caudate. Notably, Spotorno et al. (2012) showed that irony comprehension increases activity in the ToM network— including the mPFC, bilateral temporoparietal junction (TPJ), and precuneus (PCun)—as well as the bilateral IFG, with enhanced mPFC–IFG coupling during irony comprehension. Taken together, these findings indicate that irony comprehension engages brain regions supporting both semantic processing and ToM. Nevertheless, how these networks interact to integrate semantic, prosodic, and contextual information—and how such integration relates to higher-order mentalizing—remains unclear and warrants further investigation.

ToM—the ability to infer others' intentions, beliefs, perspectives, and emotions, and to recognize that others have mental states distinct from one's own (R. L. C. Mitchell & Phillips, 2015)—is central to irony and sarcasm comprehension, as understanding a speaker's intended meaning requires inferring their mental state. ToM and pragmatic abilities are developmentally linked (Bosco & Gabbatore, 2017), and ToM deficits are associated with impaired irony comprehension in autism spectrum disorder (Happé, 1993; Saban-Bezalel et al., 2019; Song et al., 2024). The ToM network encompasses the bilateral TPJ, associated with perspective-taking and representing others' goals and intentions; the mPFC, which engages in reflective reasoning and predicts others' actions and emotions; and the PCun, involved in

mental imagery (Frith & Frith, 2006; J. P. Mitchell, 2009; Saxe & Baron-Cohen, 2006; Schurz et al., 2014; Tamir et al., 2016; Van Overwalle, 2009; Van Overwalle & Baetens, 2009).

According to the predictive processing framework (Friston, 2005, 2010), irony comprehension has been proposed to rely on precision-modulated prediction error minimization across pragmatic, social, and emotional components that jointly help infer the speaker's intended, non-literal meaning (Fabry, 2021). In our view, irony comprehension may hinge on identifying and resolving prediction errors that stem from incongruencies between the context and the semantic content, as well as between the semantic content and prosody within an utterance. To minimize these prediction errors, a dynamic integration between contextual, semantic, and prosodic cues is thus needed. To process this integration, the brain engages in evidence accumulation across these different cues in a precision-modulated and bottom-up fashion. These integrated representations are then relayed to higher-order regions that also engage in prediction error minimization through refinement of predictions regarding others' intentions, emotions, and the use of irony given sensory inputs and social contexts. These prediction errors are then back-propagated to lower-order regions to gain precision by assigning greater weight to relevant cues, enhancing future interpretative frameworks for irony comprehension. The left IFG may be a key integration hub that combines prosodic and semantic cues in a bottom-up manner and transmits this information to higher-order and social cognition regions (Tettamanti et al., 2017), and subsequently receives cognitive evaluations fed back to refine perceptual integration through top-down modulation of lower-order processing in temporal regions (Cope et al., 2023; He et al., 2024). This view is supported by evidence of its functional connectivity with superior and middle temporal regions involved in speech comprehension and emotional prosody processing (Ethofer et al., 2012; Frühholz & Grandjean, 2012; Huang et al., 2024; Leitman et al., 2010; Turken & Dronkers, 2011; Yue et al., 2013), as well as with the mentalizing network (Mori & Haruno, 2022; Spotorno et al., 2012; Tettamanti et al., 2017). The present study thus examines this processing hierarchy empirically—from lower-order perceptual decoding in speech-related regions, through integrative mechanisms in the IFG, to higher-order mentalizing in the ToM network.

To this end, we designed scenarios featuring short conversations between two characters: the first character's sentence provides the context, while the second character's response serves as the target statement. By manipulating the semantics of the context, as well as the semantics and prosody of the target statement, we created distinct conditions that differed in their potential to be perceived as ironic or sarcastic. In the present fMRI study, we used two literal conditions—positive sincerity (positive context, positive statement, positive prosody) and negative sincerity (negative context, negative statement, negative prosody)—and two non-literal conditions—sarcastic irony (negative context, positive statement, negative prosody) and praise irony (positive context, negative statement, positive prosody). It should be noted that because context valence and prosody valence co-vary across these conditions, our analyses speak

to the joint influence of these cues as they naturally co-occur in non-literal speech, rather than to their fully independent contributions. We then developed five tasks in which participants were required to attend to and evaluate specific information, with each task focusing on different components involved in understanding non-literal speech, including emotional prosody and semantic decoding, irony and sarcasm inferences, as well as ToM processes. In the prosody and semantic evaluation tasks, participants were specifically required to focus solely on the modality corresponding to the task, whereas in the irony, sarcasm, and ToM tasks, they were instructed to use and integrate all available cues.

In Hypothesis 1 (behavioral data), we predicted an interaction between prosodic and semantic cues across all tasks, with a reduced effect size in tasks where participants were instructed to focus exclusively on a single modality (i.e., prosody or semantics) compared to tasks requiring integration of both cues (i.e., irony, sarcasm, and ToM tasks). Indeed, we still expected that prosodic information would continue to influence semantic judgments, and conversely, even in tasks with unimodal instructions.

For fMRI data, in Hypothesis 2 we predicted that the integration of prosodic and semantic cues in non-literal speech processing would trigger enhanced activity in key brain networks implicated in language comprehension, multimodal integration, and socio-cognitive inference. Specifically, we hypothesized heightened activations for the contrast (Non-literal > Literal) across our different tasks in: i) the left IFG as a marker of non-literal speech processing (Matsui et al., 2016; Rapp et al., 2012) and highlighting its potential role as a key hub for integrating prosodic, semantic, and contextual cues (Schirmer & Kotz, 2006); ii) the speech processing network (Friederici, 2011, 2012), including the bilateral MTG, STG, STS, and ATL, driven by greater demands for prosodic and semantic decoding and integration; iii) the ToM network (Schurz et al., 2014), including the mPFC, the TPJ, and the PCun supporting increased demands for perspective taking and inference on others' intents.

Building on our initial analyses, we extracted beta values from ROIs defined by peak activations within the predicted brain regions of Hypothesis 2 that reached significance, allowing us to examine how prosodic and semantic cues interact within the neural circuits supporting non-literal speech processing. If Hypothesis 2i is supported, we predicted in Hypothesis 3 a significant prosody-by-semantics interaction across all tasks within the left IFG, consistent with its proposed role as a key integration hub bridging lower-order perceptual processing and higher-order pragmatic and ToM regions. Mirroring the behavioral patterns anticipated in Hypothesis 1, we expected this interaction to be stronger in integrative tasks (irony, sarcasm, and ToM), which require the combination of prosodic, semantic, and contextual cues.

We further anticipated that the prosody-by-semantics interaction would extend beyond the left IFG to other regions of the identified network in a task-specific manner. This reflects the idea that the network supports integration at multiple levels—from lower-order perceptual decoding in temporal regions to

higher-order social inference in ToM regions—and that the interaction may therefore emerge selectively depending on each task's processing demands. In Hypotheses 4 and 5, we generated predictions guided by clear theoretical expectations but treated them as exploratory, given the limited prior evidence on task- and region-specific patterns. Specifically, we reasonably predicted in Hypothesis 4 that the interaction would be detectable in temporal regions identified in Hypothesis 2ii during at least the prosody and semantics tasks, where attention is directed to individual linguistic channels. In Hypothesis 5, we predicted that prosody and semantics would interact within ToM network regions identified in Hypothesis 2iii at least during the ToM task, where participants integrate multiple cues to infer communicative intent.

2 Materials and Methods

The study was approved by the Commission cantonale d'éthique de la recherche (CCER; approval number 2021-01219) and conducted in accordance with the Declaration of Helsinki and local regulations. All participants provided written informed consent and received monetary compensation for their participation.

2.1 Participants

In our fMRI study, 45 right-handed participants (23 males and 22 females, $M_{\text{age}} = 23.04$, $SD_{\text{age}} = 4.65$) were recruited at the University of Geneva, achieving a statistical power of 0.95 to detect an effect size of Cohen's $f^2 = 0.3$ at an alpha level of 0.05 with two predictors. Eligibility was determined via self-report and standard MRI safety screening. Specifically, right-handedness, native French proficiency, absence of hearing or speech impairments, absence of psychiatric history, and absence of claustrophobia were all confirmed by self-declaration. Participants were aged between 18 and 45 years, and pregnancy was also assessed by self-report. MRI compatibility was verified through our institution's standard safety questionnaire, which was completed by all participants prior to scanning.

2.2 Materials and Tasks

Our stimuli consisted of recorded sentences in French featuring short dialogues between two characters. Specifically, the first character delivered an utterance, referred to as the context ($M_{duration} = 2.10$, $SD_{duration} = 0.33$), and the second character responded with the target statement (i.e., the one that could be ironic or not; $M_{duration} = 1.06$, $SD_{duration} = 0.21$). We created 16 scenarios in which the

context's semantics, and the statement's semantics and prosody were manipulated. For a given scenario, the context could be stated either positively or negatively (e.g., "my husband won a lot of money in the lottery" vs. "this player lost a lot of money in the lottery") with a monotone prosody, while the target statement could be stated either positively or negatively (e.g., "he is lucky" vs. "he is unlucky") and delivered with a positive or negative prosody. A first pilot study examined whether our prosodic manipulations produced the intended effects on perceived emotional valence and intensity—specifically, whether positive prosody was rated as more positive and more intense than negative prosody. Systematically combining these factors—context valence, statement valence, and statement prosody—yielded eight distinct experimental conditions, each varying in their potential to be perceived as ironic or sarcastic. For the main study, we narrowed our focus to a subset of these conditions to create a clear contrast between literal and non-literal speech. This selection was guided by theoretical relevance and by the results of a second pilot study, in which we piloted our experimental tasks and identified the conditions most clearly perceived as sarcastic and ironic. We first selected the fully literal conditions—those in which semantic valence and prosody were fully congruent—resulting in the *negative sincerity* condition (negative context, negative statement, negative prosody) and the *positive sincerity* condition (positive context, positive statement, positive prosody). To represent non-literal speech, we then selected the most relevant ironic conditions based on both theoretical considerations and the estimated marginal mean results from Pilot Study 2 (see Table S4). Specifically, we included the *sarcastic irony* condition (negative context, positive statement, negative prosody), which was perceived as the most sarcastic in the sarcasm task of Pilot Study 2. We also included its positive-context counterpart, *praise irony* (positive context, negative statement, positive prosody), which was the condition with a positive context rated as the most ironic in the irony task of Pilot Study 2. These choices ensured relevance by providing non-literal forms of speech characterized by incongruence between the context and the target utterance and by featuring incongruence between prosody and semantic cues within the target statement, which was critical to capture their integration. Importantly, they ensured symmetry between sarcastic and praise forms of irony, enabling balanced analyses.

Participants then had to evaluate these different conditions in 5 different tasks. In the first task, participants were asked to evaluate the prosodic emotional valence of the target statement on a 5-point Likert scale from *very negative* (1) to *very positive* (5). In a second task, they were asked to evaluate the semantic emotional valence of the target statement from *very negative* (1) to *very positive* (5). In a third task, they were required to first evaluate the degree of perceived irony from *not ironic at all* (1) to *very ironic* (5). In a fourth task, they were asked to evaluate the degree of perceived sarcasm from *not sarcastic at all* (1) to *very sarcastic* (5). Finally, we wanted participants to focus on ToM processes. To achieve this, we drew inspiration from the social faux-pas task (Baron-Cohen et al., 1999).

More specifically, a social faux-pas is a task used to measure ToM abilities in which participants are presented with short scenarios and are asked to evaluate social mistakes, specifically whether someone did or said something that could have hurt or made another character feel uncomfortable, without the direct intention to do so. In the context of our experiment, we merely asked participants whether the second character (i.e., the one making the target statement) could have made the first character feel *not uncomfortable at all* (1) to *very uncomfortable* (5). By doing so, we aimed to target "meta-meta emotional representations" since participants needed to first infer the intent of the second character and then the emotional response of the first character regarding this intent. In the prosodic and semantic tasks, they were explicitly instructed to attend only to the channel relevant to the task while ignoring the other channel. For the irony, sarcasm, and the ToM tasks, they were asked to integrate both channels along with contextual and emotional information available.

2.3   fMRI Task Procedure

Before the fMRI session, participants were provided with detailed instructions for the different tasks. Particular attention was devoted to ensuring that participants fully understood the task instructions, especially the distinction between irony and sarcasm. For instance, we clarified that sarcasm represents a specific pejorative subtype of irony—meaning that sarcastic statements should always be perceived as ironic—while irony can also take other forms, such as praise irony. Auditory stimuli were presented via MR-compatible headphones, and the task was displayed on an MR-compatible screen that participants could see through a tilted mirror attached to the head coil. Participants performed the task using a response box in their right hand. Before starting the tasks, a training phase consisting of one random example of each task was conducted within the MRI scanner. This allowed us to adjust the headphone volume according to the MRI noise level. Participants subsequently completed the experiment in five sessions, each corresponding to a different task, with each session lasting approximately 10 minutes. In each task, participants were presented with all scenarios of all conditions, resulting in 64 trials per task and 320 trials in total. Each trial began with a 1-second inter-trial interval during which a fixation cross was presented at the center of the screen. For each trial, the context and the statement were separated by a fixation cross on a black screen, with the duration varying between 0.5 and 1.5 seconds to minimize habituation and expectation effects. Participants then evaluated the target statement depending on the task they were performing. The evaluation could last up to a maximum of 5 seconds, but participants could directly proceed to the next stimulus after validating their response. For each participant and trial, the voices of the context and the statement were assigned randomly. In addition to ensuring that the voice of the context differed from that of the statement, we also implemented the constraint that participants

could not encounter the same set of voices for the same scenario of a given condition in another task. This minimized habituation and promoted generalizability. Finally, the task order and the stimuli within each task were randomized. A typical session lasted approximately 1.5 hours in total, including setup, instructions, a training phase, and five scanning runs of approximately 10 minutes each, with short breaks provided between runs. To assess potential time-on-task effects, mean reaction times and the proportion of missing or invalid responses were examined across runs (see Supplementary Figure S4). Neither measure showed a systematic increase consistent with progressive fatigue.

Figure 1: Experimental plan summary.

2.4   Behavioral Analyses

Behavioral data were analyzed using RStudio software (R Core Team, 2021). Mixed-model linear regressions were conducted using the function `lmer` from the `lme4` package (Bates et al., 2015), *p*-values

were derived using the `Anova` function from the `car` package with the Type II sum of squares method (Fox & Weisberg, 2019), the marginal and conditional R^2 values were computed using the `performance` package (Lüdecke et al., 2021), and estimated marginal means were computed using the `emmeans` package (Lenth, 2024). Post-hoc power analysis was conducted using G*Power (Faul et al., 2007). In all our models, we added random intercepts for the participants and for the stimuli of the contexts and the statements. We also calculated the estimated marginal means for the interaction between prosody and semantics to gain a clearer understanding of how each condition was evaluated across tasks. The full set of mixed-model formulas are displayed in Table S6.

2.5   Image Acquisition

Imaging was performed at the Brain and Behaviour Laboratory (BBL) of the University of Geneva on a 3T Siemens Trio System (Siemens, Erlangen, Germany) using a T2*-weighted gradient echo planar imaging sequence with 440 volumes per run (EPI; $3.0 \times 3.0 \times 3.0$ mm voxels, slice thickness = 3 mm, gap = 1 mm, 36 slices, TR = 650 ms, TE = 30 ms, flip angle = 64°, matrix = 64×64, field of view = 192 mm). The number of volumes per run varied according to the length of the randomly assigned stimuli and the response time of the participants ($M_{\text{scans}} = 832.112$, $\min_{\text{scans}} = 738$, $\max_{\text{scans}} = 1024$). Additionally, a T1-weighted, magnetization-prepared, rapid-acquisition, gradient echo anatomical scan ($0.8 \times 0.8 \times 0.8$ mm voxels, slice thickness = 0.8 mm, 192 slices, RT = 2000 ms, TE = 2.49 ms, flip angle = 9°, matrix = 288×288, FOV = 230 mm) was acquired.

2.6   Image Analysis

Analysis of functional images was performed using Statistical Parametric Mapping software version 12 (Ashburner et al., 2014). Preprocessing steps included realignment to the first volume of the time series to correct for subject motion, coregistration, normalization into the Montreal Neurological Institute space (Collins et al., 1994), and spatial smoothing with an isotropic Gaussian filter of 6 mm full width at half maximum. Anatomical locations were identified based on a standardized coordinate database using the Automated Anatomical Labeling (AAL) atlas, integrated within the xjView toolbox (http://www.alivelearn.net/xjview). Finally, the brain plots were created using the visualization tool of the CONN toolbox (Whitfield-Gabrieli & Nieto-Castanon, 2012). For the analyses, we used a general linear model to perform the first-level analysis, treating each task as a separate session. Each trial was convolved with the hemodynamic response function, time-locked to the onset of the statement and modeled with a duration extending to the end of the evaluation. This approach was chosen to capture the full processing cascade

of non-literal speech comprehension—from initial perceptual decoding through pragmatic inference to evaluative judgment—within a single regressor. Crucially, the epoch is bounded by the participant's reaction time, ensuring that the modeled window corresponds precisely to the moment at which the full comprehension process has been completed and the meaning resolved. Sensitivity analyses separating the statement and evaluation periods into distinct regressors are reported in the Supplementary Material. Then, separate regressors were created for each condition (positive sincerity, negative sincerity, praise irony, and sarcastic irony) of each task (prosody, semantic, sarcasm, irony, and ToM), resulting in a total of 20 regressors of interest (design matrix: "prosody.positive_sincerity," "prosody.negative_sincerity," ... "ToM.praise_irony," "ToM.sarcastic_irony"). Additionally, for each run, six motion parameters were included as regressors of no interest to account for movement in our data. This resulted in a design matrix of 50 columns plus the 4 constants. To examine our main effects and interaction, we used our regressors of interest to compute 10 simple contrasts for each participant, corresponding to the simple effect of literal and non-literal conditions for each task. We then used these contrasts in a flexible factorial second-level analysis with three factors: the participants' factor (data independence set to "true", variance set to "unequal"), the non-literality factor (independence set to "false", variance set to "unequal"), and the task factor (independence set to "false", variance set to "unequal"). All results displayed in the figures were thresholded using a voxel-wise correction for multiple comparisons using a corrected False Discovery Rate (FDR) $p < .05$ and an arbitrary cluster size of $k > 10$ voxels to remove very small clusters.

fMRI ROI Analyses

Beta weights were extracted from regions of interest defined by the MNI peak coordinates of regions hypothesized in H2 and found to be significant in the flexible factorial analyses. Beta estimates were then averaged across a $3 \times 3 \times 3$ voxel neighbourhood (27 voxels) centered on that location, yielding one mean beta weight per condition, per task, per ROI, and per participant. Beta weights were then used as dependent variables in a series of 2×2 repeated-measures ANOVAs, with prosody valence and semantic valence as within-subject factors. One ANOVA was conducted for each ROI $\times$ task combination. All analyses were conducted in R using the `rstatix` package.

3   Results

Prior to the main study, two preliminary studies were conducted to validate the stimuli and pilot the experimental tasks. Full methods and results are reported in the Supplementary Material. In the main study presented in this section, participants listened to short dialogues between two characters, where

the first character's sentence served as the context and the second character's sentence as the target statement. The semantic valence of the context, the semantic valence of the target statement, and its prosody were manipulated to create literal (positive and negative sincerity) and non-literal (sarcastic and praise irony) conditions. Across five tasks, participants evaluated different aspects of the target statement: prosody, semantics, irony, sarcasm, and ToM. While the prosody and semantic tasks required focus on a single modality, the irony, sarcasm, and ToM tasks required integration of prosodic, semantic, and contextual cues. If this manipulation successfully modulated the degree of cross-channel integration, we would expect a gradient of prosody-by-semantics interaction strength across tasks—from absent or attenuated in attention-constrained tasks to strongest in explicitly integrative tasks. Importantly, however, selective attention instructions modulate rather than eliminate cross-channel interference, and we therefore predicted an interaction across all tasks, with attenuation rather than absence in the prosody and semantic conditions. Building on this, we investigated the interaction between target statement semantics and prosody across our different tasks to better understand their interplay when focusing on different emotional and pragmatic aspects. Also, participants' brain responses were recorded using fMRI and the data were analyzed to examine our fMRI hypotheses. Analyses focused on the time window from the onset of the target statement to the end of the evaluation phase, allowing us to capture both early, lower-level integration mechanisms and higher-order integration involved in social-pragmatic interpretation.

3.1   Behavioral Results

To test our first hypothesis, according to which an interaction between prosody and semantics existed across our tasks—with a reduced effect size in tasks where participants were instructed to focus exclusively on a single modality—we computed a linear mixed model including the interaction between prosody and semantics on participants' evaluations. For behavioral results, statistical modeling power is reported for fixed effects (R^2c) and for the combined fixed and random effects (R^2m). We found a significant interaction between semantics and prosody in the semantic task $F(1, 329.51) = 6.54$, $p = .011$, $R^2m = 0.76$, $R^2c = 0.85$; in the irony task $F(1, 403.43) = 4299.90$, $p < .001$, $R^2m = 0.65$, $R^2c = 0.72$; in the sarcasm task $F(1, 428.00) = 2501.43$, $p < .001$, $R^2m = 0.64$, $R^2c =$ not computable due to singularity; in the ToM task $F(1, 419.56) = 600.64$, $p < .001$, $R^2m = 0.26$, $R^2c = 0.50$; but not in the prosody task. These findings are consistent with study piloting results (see Supplementary Material). Full mixed model linear regression results are reported in Table S6, estimated marginal means from the interaction models in Table S7, plots of the predicted values in Fig. 2, and plots of the actual values in Fig. S5.

Figure 2: Plots of the predicted values of the interaction model between semantics and prosody in our A) prosody, B) semantic, C) irony, D) sarcasm, and E) ToM tasks. Each boxplot displays the distribution of predicted evaluation scores as a function of prosody (x-axis) and semantic valence (fill color). Green (positive) and red (negative) correspond to the two semantic conditions. Each point represents an individual data value. The boxplots show the median (horizontal line), interquartile range (box), and data spread (whiskers). Panels are faceted by the context (positive or negative).

Whole-brain fMRI Results

To examine our second hypothesis, we conducted a flexible factorial second-level analysis and examined the main effect contrast Non-literal $>$ Literal. Whole-brain analysis revealed the highest significant clusters of activation in the left IFG (Fig. 3A,E)—with peak activation in the pars orbitalis (IFGorb), extending into the pars triangularis (IFGtri) and pars opercularis (IFGop). Further activations were found in the bilateral medial superior frontal gyrus (mSFG; Fig. 3B,D,F)—including the mPFC, in the bilateral supplementary motor area, in the right IFG (Fig. 3C,E)—predominantly in the IFGtri and IFGop with lesser involvement of the IFGorb, in the bilateral anterior insula (Fig. 3A,C)—with left hemisphere dominance, in the bilateral TPJ (Fig. 3A,C)—including the left AG and inferior parietal lobule, as well as the right AG, in the left PCun—extending to the posterior cingulate cortex (PCC; Fig. 3D,F), in the bilateral MTG (Fig. 3A,C)—extending to the STS in the left hemisphere, in the left middle cingulate cortex (MCC; Fig. 3D), and in the bilateral middle frontal gyrus (Fig. 3A,B,C). Full peak MNI coordinates for this contrast are reported in Table S8, corresponding brain activations are displayed in Fig. 3, and brain activations for the contrast Literal $>$ Non-literal are shown in Fig. S6.

ROI Analyses Results

Based on the brain regions that were significant in the Non-literal $>$ Literal contrast and predicted in Hypothesis 2, we extracted beta values at peak coordinates for each task and condition. We thus selected peak activity betas in the left IFGorb $[-44, 22, -8]$, right IFGtri $[56, 24, 18]$, left MTG $[-62, -34, -6]$, right MTG $[62, -30, -6]$, left TPJ $[-44, -56, 40]$, right TPJ $[50, -56, 30]$, left mPFC $[-4, 42, 28]$, and right mPFC $[6, 42, 36]$, and left precuneus $[-4, -66, 32]$; a visual representation of the anatomical localization of these ROIs is provided in Figure 4, panel C. All Prosody $\times$ Semantics interaction effects were corrected for multiple comparisons using the False Discovery Rate method (Benjamini-Hochberg, 1995; 45 comparisons; FDR-corrected $p < .05$).

Hypothesis 3 predicted a significant prosody-by-semantics interaction across all tasks within the left IFG, consistent with its proposed role as a key integration hub bridging lower-order perceptual processing and higher-order pragmatic inference. This hypothesis was supported, with the left IFG showing a significant prosody-by-semantics interaction across all tasks. However, the prediction that this interaction would be strongest in integrative tasks was only partially confirmed: while the sarcasm and ToM tasks showed stronger interactions than the prosody and semantic tasks as expected, the irony task showed a weaker interaction than the prosody task, departing from the predicted pattern. Exploratory analyses further revealed that the right IFG showed significant prosody-semantics interactions across all tasks, though

Figure 3: Whole-brain results of the contrast Non-literal > Literal across our different tasks in a sagittal A and C), medial D and F), superior B), and inferior E) view. All activations are thresholded at a voxelwise $p < 0.05$ FDR. The colorbar represents T statistics. IFG: inferior frontal gyrus; tri: pars triangularis; op: pars opercularis; orb: pars orbitalis; aINS: anterior insula; STS: superior temporal sulcus; MTG: mid temporal gyrus; TPJ: temporo-parietal junction; IPL: inferior parietal lobule; PCun: precuneus; PCC: posterior cingulate cortex; MCC: middle cingulate cortex; ACC: anterior cingulate cortex; SMA: supplementary motor area; SFG: superior frontal gyrus; mPFC: medial prefrontal cortex; pOFC: posterior orbitofrontal cortex.

with stronger effects in non-integrative than integrative tasks. The prosody-by-semantics interaction beta values for the left IFG are displayed in Figure 4, panel A.

Hypothesis 4 predicted that the prosody-by-semantics interaction would extend to temporal regions at least during the prosody and semantics tasks, where attention is directed toward individual linguistic channels. This prediction was confirmed. Moreover, these exploratory analyses revealed that the interaction between prosody and semantics in both the left and right MTG extended beyond the predicted tasks, reaching significance during the ToM task as well.

Finally, Hypothesis 5 predicted that prosody and semantics would interact within ToM network regions—including the TPJ, precuneus, and mPFC—at least during the ToM task. This prediction was confirmed across all mentalizing network ROIs except the right TPJ. Exploratory analyses revealed that the left mPFC showed significant interactions across four tasks—semantics, irony, sarcasm, and ToM—with the exception of the prosody task. The right mPFC similarly showed significant interactions across four tasks—prosody, semantics, sarcasm, and ToM—with the exception of the irony task. The left TPJ showed significant interactions during the prosody, semantics, sarcasm, and ToM tasks, but not during irony. The right TPJ showed significant interactions during the prosody and semantics tasks but did not reach significance for any of the integrative tasks. The left precuneus showed a more restricted pattern, with significant interactions during the prosody, sarcasm, and ToM tasks only. F-values and significance of the prosody-by-semantics interaction for each task and each ROI are displayed in Figure 4, panel B. Full interaction results for each ROI are reported in Supplementary Table S9, and plots of the beta values for all ROIs are provided in Supplementary Figure S9.

4   General Discussion

The primary objective of our research was to investigate how prosody and semantics interact in non-literal forms of speech and to elucidate the brain mechanisms supporting their integration—both at the level of combining linguistic channels and in relation to higher-level cognitive processes, such as theory of mind. At the behavioral level, we observed significant interactions between prosodic and semantic factors across all tasks except the prosody task. As predicted, interactions were attenuated in non-integrative tasks (prosodic, semantic) compared to integrative tasks (irony, sarcasm, ToM). However, this prediction was only partially confirmed: rather than being merely reduced, the interaction was entirely absent in the prosody task. This asymmetry suggests that while prosody can influence semantic judgments, semantics may not equivalently penetrate prosodic judgments. We then conducted whole-brain imaging analyses focusing on the main effect contrast Non-literal > Literal speech as a first step to identify the

Figure 4: **Prosody $\times$ Semantics interaction across ROIs and tasks.** Panel A displays the distribution of individual beta values (signal change %) extracted from the left IFG $[-44, 22, -8]$ as a function of prosody valence (x-axis: Positive, Negative) and semantic valence (fill color: blue = Positive, orange = Negative), across all five tasks (Prosody, Semantic, Irony, Sarcasm, ToM). Each dot represents an individual subject's beta value. Bars display the mean signal change (%) and error bars reflect the standard error of the mean ($\pm$ SE). Diamond shapes indicate the group mean per condition. Significance brackets indicate the Prosody $\times$ Semantics interaction effect (*** $p < .001$, ** $p < .01$, * $p < .05$; FDR-corrected). Panel B displays a summary heatmap of the Prosody $\times$ Semantics interaction F-values derived from 2×2 repeated-measures ANOVAs, across all nine ROIs (rows) and five tasks (columns). Coloured cells indicate significant interactions, with darker blue reflecting higher F-values; grey cells indicate non-significant effects (ns). Stars indicate FDR-corrected significance levels (* $p < .05$, ** $p < .01$, *** $p < .001$; Benjamini-Hochberg, 45 comparisons). The left IFG row is highlighted with a blue border. Panel C displays the anatomical localisation of the regions of interest identified as significant in Hypothesis 2, color-coded by region: IFG (blue): inferior frontal gyrus; MTG (orange): middle temporal gyrus; TPJ (green): temporoparietal junction; mPFC (pink): medial prefrontal cortex; and PCun (red): precuneus.

network involved in non-literal speech processing, followed by a targeted ROI analysis directly examining the prosody-by-semantics interaction within these regions across tasks. Our hypothesis was partially confirmed for the Non-literal > Literal speech contrast, with significant activity in the bilateral IFG— showing a clear left dominance, the left MTG and STS, the right MTG, as well as the bilateral TPJ, mPFC, and left PCun. Within these regions, the ROI analyses revealed that the prosody-by-semantics interaction was detectable across the network, though with markedly different profiles: bilateral IFG showed sensitivity across all tasks, temporal regions were most strongly engaged during lower-order tasks but extended to ToM, the left and right mPFC both showed sensitivity across four tasks, while TPJ and precuneus showed more selective patterns. Overall, these findings suggest that prosody-semantics integration in non-literal speech is distributed across a fronto-temporal and mentalizing network whose components engage selectively depending on the level of processing demanded.

In the current study, we investigated the interaction between prosody and semantics on participants' evaluations across multiple tasks. Our findings revealed significant interactions in tasks involving irony, sarcasm, and ToM, with a weaker interaction observed in the semantic task. No interaction was found in the prosody task. This result further replicates findings from Pilot Study 2 (see Supplementary Material), which demonstrated an identical pattern. The observation that participants instructed to attend semantic content only while ignoring prosodic cues still had their semantic evaluation modulated by perceived prosody valence, without the reverse effect, supports the prosody dominance hypothesis. For instance, studies using a cross-channel Stroop task with participants asked to attend only one channel (i.e., prosody or semantics) and evaluate word emotional valence showed that prosody led to faster and more accurate responses than semantics (Filippi et al., 2017; Lin et al., 2020; Schirmer & Kotz, 2003). This effect persisted even in incongruent conditions but also showed less interference of semantics on prosody, highlighting the salience and relevance of prosodic cues (Schirmer & Kotz, 2003). Similarly, Ben-David et al. (2016) developed a new assessment tool that requires participants to attend only one channel to categorize discrete emotions, providing measures of both the dominance of the attended channel and the ability to selectively ignore the other one. Overall, their results demonstrate prosody dominance, also with prosodic cues proving more difficult to selectively ignore, which is in line with the present findings.

We hypothesized that the left IFG would serve as a key integration hub for prosodic and semantic content during non-literal speech processing. The left IFGorb emerged as the peak activation in the Non-literal > Literal contrast and showed a significant prosody-by-semantics interaction across all five tasks, displaying both the strongest task-specific interaction effect and the highest cumulative interaction strength across tasks, supporting its proposed role as an integration hub operating across multiple stages of the processing hierarchy. This is consistent with a large body of evidence showing the convergence

of semantic and emotional cue processing—including prosody—within the IFGorb (Belyk et al., 2017), which has also been specifically implicated in integrating discourse context and utterance-level information with prosodic cues during sarcasm comprehension (Matsui et al., 2016). Interestingly, the pattern of prosody–semantics interaction in the left IFGorb did not mirror the behavioral asymmetry between tasks. Although participants successfully inhibited semantic information in the prosody task behaviorally, the left IFGorb still showed a robust interaction in this condition, even stronger than in the irony task. This dissociation suggests that the conjunction of prosodic and semantic cues may be computed relatively automatically in the left IFGorb, prior to top-down control processes that enable behavioral suppression of semantic information. It is important to acknowledge, however, that activity in the left IFG may also reflect effortful processing more broadly, including hesitation or conflict monitoring during tasks involving ambiguous acoustic signals, competing emotional cues, or even non-human vocalizations (Ceravolo et al., 2023; Frühholz & Grandjean, 2013; Moss et al., 2005), a demand that was likely present across all five of our tasks given the inherent competition between prosodic and semantic cues in our stimuli. Moreover, the right IFGtri also showed sensitivity to the conjunction of prosodic and semantic cues across all tasks, suggesting that it may contribute to this integration process independently of task demands as well. Notably, its sensitivity was strongest in the two non-integrative tasks. This pattern may reflect a role in the active inhibition of the task-irrelevant channel rather than meaning-oriented integration, consistent with accounts proposing an inhibitory function of the right IFG during speech processing along a posterior-to-anterior gradient from more automatic inhibitory mechanisms to higher-order social–cognitive processes (Aron et al., 2014; Hartwigsen et al., 2019; Neef et al., 2016).

Within the temporal speech network, activity during non-literal speech processing was observed primarily in the bilateral MTG, extending into the STS. The MTG showed sensitivity to the prosody–semantics interaction mainly during non-integrative tasks—consistent with its role in lower-level perceptual decoding of individual linguistic channels—but not during integrative tasks, with the notable exception of the ToM task. This pattern suggests that the MTG contributes to prosody–semantics integration when attention is directed toward individual linguistic channels, and may re-engage during ToM processing to ground higher-order social inferences in the perceptual properties of the utterance. The left IFG may constitute a key interface linking perceptual speech processing in temporal regions with higher-order social-cognitive networks. In this view, the IFG integrates prosodic and semantic information through its connectivity with superior and middle temporal regions involved in speech comprehension and emotional prosody processing (Ethofer et al., 2012; Frühholz & Grandjean, 2012; Huang et al., 2024; Leitman et al., 2010; Turken & Dronkers, 2011; Yue et al., 2013), and monitors their conjunction before relaying this representation to the theory-of-mind network, including the mPFC, TPJ, and precuneus (Tettamanti et al., 2017). Consistent with this account, reciprocal IFGorb–MTG interactions have been proposed to support the selection

and maintenance of lexical–semantic representations within broader contextual frameworks (Turken & Dronkers, 2011), enabling their integration into the higher-order social and pragmatic inferences that characterize non-literal speech comprehension.

In line with this proposed fronto–temporal interface, we also observed robust engagement of the mentalizing network during non-literal speech processing. This finding aligns with previous evidence of co-activation between speech and ToM networks during non-literal comprehension (Hauptman et al., 2023), suggesting that non-literal interpretation engages both core language mechanisms and social inference processes. Moreover, all ToM regions except the right TPJ displayed a significant prosody–semantics interaction at least during the ToM task, indicating that the integration of prosodic and semantic cues is supported by a broader neural architecture in which social-cognitive regions actively contribute to resolving the meaning of non-literal utterances. Within this network, the bilateral mPFC is consistently reported in non-literal speech processing (Hauptman et al., 2023; Rapp et al., 2012) and has been implicated in predicting and integrating others' beliefs, intentions, and emotions to support perspective-taking and social understanding (Frith & Frith, 2006; Van Overwalle, 2009). In the ROI analysis, the left mPFC showed the strongest sensitivity across tasks—particularly in the ToM task—consistent with its proposed role in orchestrating top-down social inference (Tettamanti et al., 2017). We also observed activation in the bilateral TPJ, primarily located in the angular gyrus and associated with the mentalizing network, where it is thought to support the representation of others' intentions and goals based on integrated socially and emotionally relevant cues (Frith & Frith, 2006; J. P. Mitchell, 2009; Saxe & Baron-Cohen, 2006; Schurz et al., 2014; Tamir et al., 2016; Van Overwalle, 2009; Van Overwalle & Baetens, 2009). However, the two hemispheres showed distinct interaction profiles in the ROI analysis: the left TPJ showed sensitivity to the prosody–semantics conjunction across all tasks except irony, whereas the right TPJ was sensitive only during non-integrative tasks. This hemispheric asymmetry may reflect distinct functional profiles of these regions: the right TPJ's engagement in non-integrative tasks is consistent with its proposed role in attentional reorienting (Chang et al., 2013; Kubit & Jack, 2013), particularly when participants must suppress one channel while attending to another, whereas the broader engagement of the left TPJ aligns with its more language-anchored role in contextual and semantic integration (Branzi & Lambon Ralph, 2023; Branzi et al., 2021), supporting prosody–semantics conjunction across a wider range of processing demands. Finally, we observed activity in the left precuneus in the Non-literal > Literal contrast, previously linked to irony processing and theory of mind (Spotorno et al., 2012). Despite its reported involvement in the processing of emotional prosody and affective semantics (Castelluccio et al., 2016), it showed relatively selective sensitivity to the prosody-semantics conjunction in the ROI analysis, appearing less responsive to semantic content, which is more in line with its proposed role in mental imagery and perspective-taking (Cavanna & Trimble, 2006; Schurz et al., 2014).

Even though we did our best to control for potentially confounding factors and variables, several limitations should be acknowledged. In our behavioral and fMRI analyses, the context and prosody factors were partially nested, as their levels co-varied across the four experimental conditions, thereby limiting the ability to disentangle their unique and independent contributions to the interaction between prosody and semantics. Our analysis of interactions thus reflects the combined effect of these cues as they naturally appear in non-literal speech, rather than their completely independent contributions. That said, the prosody dominance effect we report suggests that prosody carries meaningful and non-redundant information that listeners integrate even when instructed to ignore it, and is therefore unlikely to be fully reducible to context. Furthermore, we designed five distinct tasks to target the different mechanisms involved in non-literal speech perception (i.e., prosodic and semantic decoding, irony and sarcasm inference, and theory-of-mind processes). Because participants listened to the same dialogues across all five tasks, we cannot fully rule out the possibility that participants were making pragmatic inferences spontaneously even in tasks requiring attention to a single channel, or that responses in later tasks were influenced by judgments formed in earlier sessions. Although task order was randomized and systematic task-dependent modulation was observed at both behavioral and neural levels, with RTs consistent with greater pragmatic inference demands in integrative tasks, this remains a structural limitation of our design. More broadly, the use of five distinct tasks—while necessary to capture the full hierarchy of processing demands from lower-order perceptual decoding to higher-order mentalizing—introduced inherent trade-offs: it limited the number of trials per task and consequently statistical power, as we sought to balance experimental time constraints and minimize habituation and fatigue, and may have increased the risk of carry-over effects across sessions. Although random intercepts for stimuli in our behavioral models partially absorb item-level variability, no explicit matching was performed across items with respect to social or emotional salience or personal relevance, and no analogous control was implemented at the fMRI inference level. Similarly, although voices were randomly assigned to trials to promote generalizability, variability in actor performance may still have introduced some noise across participants' evaluations and brain responses. Lastly, our stimuli consisted of short sentences, which may pose challenges to the ecological validity of the study, as they do not accurately reflect the natural use of irony in everyday conversational contexts. This may have further limited the recruitment of ToM processes and associated brain networks (Spotorno et al., 2012).

5   Conclusion

In this study, we investigated the neural mechanisms underlying the integration of prosodic and semantic cues in non-literal forms of speech, such as irony and sarcasm. To this end, we designed five tasks

targeting distinct cognitive processes, including prosody and semantics evaluation, irony and sarcasm inference, and theory of mind. At the behavioral level, participants' responses revealed systematic interactions between semantic and prosodic factors across all tasks except the prosody-only condition, suggesting a prosody dominance effect. Our fMRI analyses first identified the network recruited by non-literal speech comprehension through a whole-brain Non-literal > Literal contrast, revealing activation in the bilateral inferior frontal gyri, the broader speech network, and ToM regions. Building directly on this, a targeted ROI analysis examined the prosody-by-semantics interaction within these regions across all five tasks. This revealed that the network supporting non-literal speech comprehension is functionally heterogeneous: the left IFGorb—which also showed the strongest activation in the Non-literal > Literal contrast—showed the prosody-by-semantics interaction across all tasks, consistent with its proposed role as a key integration hub, while other regions engaged more selectively as a function of task demand. These findings provide preliminary support for a distributed and hierarchically organized account of prosody-semantics integration, though further research is needed to clarify the dynamic interplay between these systems—particularly through functional connectivity or electrophysiological approaches—to better understand how they cooperate during the comprehension of non-literal language.

Data and Code Availability

Our behavioral and fMRI data, along with the behavioral analysis scripts, are available at https://doi.org/10.26037/ya

Author Contributions

W.A. conceived and conducted the second and third experiments, analyzed the data of all experiments, and wrote the manuscript. C.L. contributed to the design of the experiments and data analysis, and reviewed the manuscript. M.A. created the stimuli and conceived and conducted the first pilot experiment. D.G. acquired the funding, contributed to the design of the experiments, and reviewed the manuscript.

Funding

This work was supported by funds from the NCCR Evolving Language, Swiss National Science Foundation Agreement #51NF40_180888.

Declaration of Competing Interests

The authors declare no conflicts of interest.

References

Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: One decade on. *Trends in Cognitive Sciences*, *18*(4), 177–185. https://doi.org/10.1016/j.tics.2013.12.003

Ashburner, J., Barnes, G., Chen, C., Daunizeau, J., Flandin, G., Friston, K., Kiebel, S., Kilner, J., Litvak, V., Moran, R., et al. (2014). SPM12 manual. *Wellcome Trust Centre for Neuroimaging, London, UK*, *2464*(4). https://doi.org/10.1109/wincom55661.2022.9966463

Astésano, C., Besson, M., & Alter, K. (2004). Brain potentials during semantic and prosodic processing in french. *Cognitive Brain Research*, *18*(2), 172–184. https://doi.org/10.1016/j.cogbrainres.2003.10.002

Baron-Cohen, S., O'Riordan, M., Stone, V., Jones, R., & Plaisted, K. (1999). Recognition of faux pas by normally developing children and children with Asperger syndrome or high-functioning autism. *Journal of Autism and Developmental Disorders*, *29*(5), 407–418. https://doi.org/10.1023/a:1023035012436

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Beaucousin, V., Lacheret, A., Turbelin, M., Morel, M., Mazoyer, B., & Tzourio-Mazoyer, N. (2006). fMRI study of emotional speech comprehension. *Cerebral Cortex*, *17*(2), 339–352. https://doi.org/10.1093/cercor/bhj151

Belin, P., Zatorre, R. J., Lafaille, P., Ahad, P., & Pike, B. (2000). Voice-selective areas in human auditory cortex. *Nature*, *403*(6767), 309–312. https://doi.org/10.1038/35002078

Belyk, M., Brown, S., Lim, J., & Kotz, S. A. (2017). Convergence of semantics and emotional expression within the IFG pars orbitalis. *NeuroImage*, *156*, 240–248. https://doi.org/10.1016/j.neuroimage.2017.04.020

Ben-David, B. M., Multani, N., Shakuf, V., Rudzicz, F., & Van Lieshout, P. H. H. M. (2016). Prosody and semantics are separate but not separable channels in the perception of emotional speech: Test for rating of emotions in speech. *Journal of Speech, Language, and Hearing Research*, *59*(1), 72–89. https://doi.org/10.1044/2015_jslhr-h-14-0323

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex*, *19*(12), 2767–2796. https://doi.org/10.1093/cercor/bhp055

Borod, J. C., Bloom, R. L., Brickman, A. M., Nakhutina, L., & Curko, E. A. (2002). Emotional processing deficits in individuals with unilateral brain damage. *Applied Neuropsychology*, *9*(1), 23–36. https://doi.org/10.1207/s15324826an0901_4

Bosco, F. M., & Gabbatore, I. (2017). Sincere, deceitful, and ironic communicative acts and the role of the theory of mind in childhood. *Frontiers in Psychology*, *8*. https://doi.org/10.3389/fpsyg.2017.00021

Bosco, F. M., Parola, A., Valentini, M. C., & Morese, R. (2017). Neural correlates underlying the comprehension of deceitful and ironic communicative intentions. *Cortex*, *94*, 73–86. https://doi.org/10.1016/j.cortex.2017.06.010

Branzi, F. M., & Lambon Ralph, M. A. (2023). Semantic-specific and domain-general mechanisms for integration and update of contextual information. *Human Brain Mapping*, *44*(17), 5547–5566. https://doi.org/10.1002/hbm.26454

Branzi, F. M., Pobric, G., Jung, J., & Lambon Ralph, M. A. (2021). The left angular gyrus is causally involved in context-dependent integration and associative encoding during narrative reading. *Journal of Cognitive Neuroscience*, *33*(6), 1082–1095. https://doi.org/10.1162/jocn_a_01698

Brück, C., Kreifelts, B., & Wildgruber, D. (2011). Emotional voices in context: A neurobiological model of multimodal affective information processing. *Physics of Life Reviews*, *8*(4), 383–403. https://doi.org/10.1016/j.plrev.2011.10.002

Buchanan, T. W., Lutz, K., Mirzazade, S., Specht, K., Shah, N. J., Zilles, K., & Jäncke, L. (2000). Recognition of emotional prosody and verbal components of spoken language: An fMRI study. *Cognitive Brain Research*, *9*(3), 227–238. https://doi.org/10.1016/s0926-6410(99)00060-9

Castelluccio, B. C., Myers, E. B., Schuh, J. M., & Eigsti, I. (2016). Neural substrates of processing anger in language: Contributions of prosody and semantics. *Journal of Psycholinguistic Research*, *45*(6), 1359–1367. https://doi.org/10.1007/s10936-015-9405-z

Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: A review of its functional anatomy and behavioural correlates. *Brain*, *129*(3), 564–583. https://doi.org/10.1093/brain/awl004

Ceravolo, L., Debracque, C., Pool, E., Gruber, T., & Grandjean, D. (2023). Frontal mechanisms underlying primate calls recognition by humans. *Cerebral Cortex Communications*, *4*(4). https://doi.org/10.1093/texcom/tgad019

Chang, C.-F., Hsu, T.-Y., Tseng, P., Liang, W.-K., Tzeng, O. J. L., Hung, D. L., & Juan, C.-H. (2013). Right temporoparietal junction and attentional reorienting. *Human Brain Mapping*, *34*(4), 869–877. https://doi.org/10.1002/hbm.21476

Collins, D. L., Neelin, P., Peters, T. M., & Evans, A. C. (1994). Automatic 3D intersubject registration of MR volumetric data in standardized Talairach space. *Journal of Computer Assisted Tomography*, *18*(2), 192–205. https://doi.org/10.1097/00004728-199403000-00005

Cope, T. E., Sohoglu, E., Peterson, K. A., Jones, P. S., Rua, C., Passamonti, L., Sedley, W., Post, B., Coebergh, J., Butler, C. R., Garrard, P., Abdel-Aziz, K., Husain, M., Griffiths, T. D., Patterson, K., Davis, M. H., & Rowe, J. B. (2023). Temporal lobe perceptual predictions for speech are instantiated in motor cortex and reconciled by inferior frontal cortex. *Cell Reports*, *42*(5), 112422. https://doi.org/10.1016/j.celrep.2023.112422

Ethofer, T., Anders, S., Erb, M., Herbert, C., Wiethoff, S., Kissler, J., Grodd, W., & Wildgruber, D. (2006). Cerebral pathways in processing of affective prosody: A dynamic causal modeling study. *NeuroImage*, *30*(2), 580–587. https://doi.org/10.1016/j.neuroimage.2005.09.059

Ethofer, T., Bretscher, J., Gschwind, M., Kreifelts, B., Wildgruber, D., & Vuilleumier, P. (2012). Emotional voice areas: Anatomic location, functional properties, and structural connections revealed by combined fMRI/DTI. *Cerebral Cortex*, *22*(1), 191–200. https://doi.org/10.1093/cercor/bhr113

Fabry, R. E. (2021). Getting it: A predictive processing approach to irony comprehension. *Synthese*, *198*(7), 6455–6489. https://doi.org/10.1007/s11229-019-02470-9

Faul, F., Erdfelder, E., Lang, A., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, *39*(2), 175–191. https://doi.org/10.3758/BF03193146

Filik, R., Țurcan, A., Ralph-Nearman, C., & Pitiot, A. (2019). What is the difference between irony and sarcasm? An fMRI study. *Cortex*, *115*, 112–122. https://doi.org/10.1016/j.cortex.2019.01.025

Filippi, P., Ocklenburg, S., Bowling, D. L., Heege, L., Güntürkün, O., Newen, A., & De Boer, B. (2017). More than words (and faces): Evidence for a Stroop effect of prosody in emotion word processing. *Cognition and Emotion*, *31*(5), 879–891. https://doi.org/10.1080/02699931.2016.1177489

Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (Third). Sage. https://doi.org/10.32614/cran.package.car

Friederici, A. D. (2011). The brain basis of language processing: From structure to function. *Physiological Reviews*, *91*(4), 1357–1392. https://doi.org/10.1152/physrev.00006.2011

Friederici, A. D. (2012). The cortical language circuit: From auditory perception to sentence comprehension. *Trends in Cognitive Sciences*, *16*(5), 262–268. https://doi.org/10.1016/j.tics.2012.04.001

Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *360*(1456), 815–836. https://doi.org/10.1098/rstb.2005.1622

Friston, K. (2010). The free-energy principle: A unified brain theory? *Nature Reviews Neuroscience*, *11*(2), 127–138. https://doi.org/10.1038/nrn2787

Frith, C. D., & Frith, U. (2006). The neural basis of mentalizing. *Neuron*, *50*(4), 531–534. https://doi.org/10.1016/j.neuron.2006.05.001

Frühholz, S., & Belin, P. (2018). *The Oxford handbook of voice perception*. Oxford University Press. https://doi.org/10.1093/oxfordhb/9780199982295.001.0001

Frühholz, S., & Grandjean, D. (2012). Towards a fronto-temporal neural network for the decoding of angry vocal expressions. *NeuroImage*, *62*(3), 1658–1666. https://doi.org/10.1016/j.neuroimage.2012.06.015

Frühholz, S., & Grandjean, D. (2013). Processing of emotional vocalizations in bilateral inferior frontal cortex. *Neuroscience & Biobehavioral Reviews*, *37*(10), 2847–2855. https://doi.org/10.1016/j.neubiorev.2013.10.007

Grandjean, D. (2021). Brain networks of emotional prosody processing. *Emotion Review*, *13*(1), 34–43. https://doi.org/10.1177/1754073919898522

Grandjean, D., Sander, D., Pourtois, G., Schwartz, S., Seghier, M. L., Scherer, K. R., & Vuilleumier, P. (2005). The voices of wrath: Brain responses to angry prosody in meaningless speech. *Nature Neuroscience*, *8*(2), 145–146. https://doi.org/10.1038/nn1392

Happé, F. G. E. (1993). Communicative competence and theory of mind in autism: A test of relevance theory. *Cognition*, *48*(2), 101–119. https://doi.org/10.1016/0010-0277(93)90026-r

Hartwigsen, G., Neef, N. E., Camilleri, J. A., Margulies, D. S., & Eickhoff, S. B. (2019). Functional Segregation of the Right Inferior Frontal Gyrus: Evidence From Coactivation-Based Parcellation. *Cerebral Cortex*, *29*(4), 1532–1546. https://doi.org/10.1093/cercor/bhy049

Hauptman, M., Blank, I., & Fedorenko, E. (2023). Non-literal language processing is jointly supported by the language and theory of mind networks: Evidence from a novel meta-analytic fMRI approach. *Cortex*, *162*, 96–114. https://doi.org/10.1016/j.cortex.2023.01.013

He, Y., Shao, X., Liu, C., Fan, C., Jefferies, E., Zhang, M., & Li, X. (2024). Diverse frontoparietal connectivity supports semantic prediction and integration in sentence comprehension. *The Journal of Neuroscience*, e1404242024. https://doi.org/10.1523/jneurosci.1404-24.2024

Hoffman, P., Binney, R. J., & Lambon Ralph, M. A. (2015). Differing contributions of inferior prefrontal and anterior temporal cortex to concrete and abstract conceptual knowledge. *Cortex*, *63*, 250–266. https://doi.org/10.1016/j.cortex.2014.09.001

Huang, C., Li, A., Pang, Y., Yang, J., Zhang, J., Wu, X., & Mei, L. (2024). How the intrinsic functional connectivity patterns of the semantic network support semantic processing. *Brain Imaging and Behavior*, *18*(3), 539–554. https://doi.org/10.1007/s11682-024-00849-y

Humphries, C., Binder, J. R., Medler, D. A., & Liebenthal, E. (2006). Syntactic and semantic modulation of neural activity during auditory sentence comprehension. *Journal of Cognitive Neuroscience*, *18*(4), 665–679. https://doi.org/10.1162/jocn.2006.18.4.665

Kotz, S. A., Dengler, R., & Wittfoth, M. (2015). Valence-specific conflict moderation in the dorso-medial PFC and the caudate head in emotional speech. *Social Cognitive and Affective Neuroscience*, *10*(2), 165–171. https://doi.org/10.1093/scan/nsu021

Kotz, S. A., Meyer, M., & Paulmann, S. (2006). Lateralization of emotional prosody in the brain: An overview and synopsis on the impact of study design. In *Progress in brain research* (pp. 285–294). Elsevier. https://doi.org/10.1016/s0079-6123(06)56015-7

Kotz, S. A., & Paulmann, S. (2011). Emotion, language, and the brain. *Language and Linguistics Compass*, *5*(3), 108–125. https://doi.org/10.1111/j.1749-818x.2010.00267.x

Kubit, B., & Jack, A. I. (2013). Rethinking the role of the rTPJ in attention and social cognition in light of the opposing domains hypothesis: Findings from an ALE-based meta-analysis and resting-state functional connectivity. *Frontiers in Human Neuroscience*, *7*. https://doi.org/10.3389/fnhum.2013.00323

Leitman, D. I., Wolf, D. H., Ragland, J. D., Laukka, P., Loughead, J., Valdez, J. N., Javitt, D. C., Turetsky, B., & Gur, R. (2010). "It's not what you say, but how you say it": A reciprocal temporo-frontal network for affective prosody. *Frontiers in Human Neuroscience*, *4*, 19. https://doi.org/10.3389/fnhum.2010.00019

Lenth, R. V. (2024). *Emmeans: Estimated marginal means, aka least-squares means* [R package version 1.10.6]. https://doi.org/10.1080/00031305.1980.10483031

Lin, Y., Ding, H., & Zhang, Y. (2020). Prosody dominates over semantics in emotion word processing: Evidence from cross-channel and cross-modal Stroop effects. *Journal of Speech, Language, and Hearing Research*, *63*(3), 896–912. https://doi.org/10.1044/2020_jslhr-19-00258

Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance: An R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software*, *6*(60), 3139. https://doi.org/10.21105/joss.03139

Matchin, W., & Hickok, G. (2020). The cortical organization of syntax. *Cerebral Cortex*, *30*(3), 1481–1498. https://doi.org/10.1093/cercor/bhz180

Matsui, T., Nakamura, T., Utsumi, A., Sasaki, A. T., Koike, T., Yoshida, Y., Harada, T., Tanabe, H. C., & Sadato, N. (2016). The role of prosody and context in sarcasm comprehension: Behavioral and fMRI evidence. *Neuropsychologia*, *87*, 74–84. https://doi.org/10.1016/j.neuropsychologia.2016.04.031

Mauchand, M., Vergis, N., & Pell, M. D. (2020). Irony, prosody, and social impressions of affective stance. *Discourse Processes*, *57*(2), 141–157. https://doi.org/10.1080/0163853X.2019.1581588

Meyer, M., Alter, K., & Friederici, A. (2003). Functional MR imaging exposes differential brain responses to syntax and prosody during auditory sentence comprehension. *Journal of Neurolinguistics*, *16*(4), 277–300. https://doi.org/10.1016/S0911-6044(03)00026-5

Mitchell, J. P. (2009). Inferences about mental states. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1521), 1309–1316. https://doi.org/10.1098/rstb.2008.0318

Mitchell, R. L. C. (2006). How does the brain mediate interpretation of incongruent auditory emotions? The neural response to prosody in the presence of conflicting lexico-semantic cues. *European Journal of Neuroscience*, *24*(12), 3611–3618. https://doi.org/10.1111/j.1460-9568.2006.05231.x

Mitchell, R. L. C. (2007). fMRI delineation of working memory for emotional prosody in the brain: Commonalities with the lexico-semantic emotion network. *NeuroImage*, *36*(3), 1015–1025. https://doi.org/10.1016/j.neuroimage.2007.03.016

Mitchell, R. L. C., & Phillips, L. H. (2015). The overlapping relationship between emotion perception and theory of mind. *Neuropsychologia*, *70*, 1–10. https://doi.org/10.1016/j.neuropsychologia.2015.02.018

Mori, K., & Haruno, M. (2022). Resting functional connectivity of the left inferior frontal gyrus with the dorsomedial prefrontal cortex and temporoparietal junction reflects the social network size for active interactions. *Human Brain Mapping*, *43*(9), 2869–2879. https://doi.org/10.1002/hbm.25822

Moss, H. E., Abdallah, S., Fletcher, P., Bright, P., Pilgrim, L., Acres, K., & Tyler, L. K. (2005). Selecting among competing alternatives: Selection and retrieval in the left inferior frontal gyrus. *Cerebral Cortex*, *15*(11), 1723–1735. https://doi.org/10.1093/cercor/bhi049

Nakamura, T., Matsui, T., Utsumi, A., Sumiya, M., Nakagawa, E., & Sadato, N. (2022). Context-prosody interaction in sarcasm comprehension: A functional magnetic resonance imaging study. *Neuropsychologia*, *170*, 108213. https://doi.org/10.1016/j.neuropsychologia.2022.108213

Neef, N. E., Bütfering, C., Anwander, A., Friederici, A. D., Paulus, W., & Sommer, M. (2016). Left posterior-dorsal area 44 couples with parietal areas to promote speech fluency, while right area 44 activity promotes the stopping of motor responses. *NeuroImage*, *142*, 628–644. https://doi.org/10.1016/j.neuroimage.2016.08.030

Obert, A., Gierski, F., Calmus, A., Flucher, A., Portefaix, C., Pierot, L., Kaladjian, A., & Caillies, S. (2016). Neural correlates of contrast and humor: Processing common features of verbal irony. *PLOS ONE*, *11*(11), e0166704. https://doi.org/10.1371/journal.pone.0166704

Pauligk, S., Kotz, S. A., & Kanske, P. (2019). Differential impact of emotion on semantic processing of abstract and concrete words: ERP and fMRI evidence. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-50755-3

Pell, M. D., Jaywant, A., Monetta, L., & Kotz, S. A. (2011). Emotional speech processing: Disentangling the effects of prosody and semantic cues. *Cognition & Emotion*, *25*(5), 834–853. https://doi.org/10.1080/02699931.2010.516915

R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria. https://www.R-project.org/

Rapp, A. M., Mutschler, D. E., & Erb, M. (2012). Where in the brain is nonliteral language? A coordinate-based meta-analysis of functional magnetic resonance imaging studies. *NeuroImage*, *63*(1), 600–610. https://doi.org/10.1016/j.neuroimage.2012.06.022

Rogalsky, C., & Hickok, G. (2009). Selective attention to semantic and syntactic features modulates sentence processing networks in anterior temporal cortex. *Cerebral Cortex*, *19*(4), 786–796. https://doi.org/10.1093/cercor/bhn126

Ross, E. D., & Monnot, M. (2008). Neurology of affective prosody and its functional–anatomic organization in right hemisphere. *Brain and Language*, *104*(1), 51–74. https://doi.org/10.1016/j.bandl.2007.04.007

Saban-Bezalel, R., Dolfin, D., Laor, N., & Mashal, N. (2019). Irony comprehension and mentalizing ability in children with and without autism spectrum disorder. *Research in Autism Spectrum Disorders*, *58*, 30–38. https://doi.org/10.1016/j.rasd.2018.11.006

Saxe, R., & Baron-Cohen, S. (2006). Editorial: The neuroscience of theory of mind. *Social Neuroscience*, *1*(3–4), 1–9. https://doi.org/10.1080/17470910601117463

Scherer, K. (2003). Vocal communication of emotion: A review of research paradigms. *Speech Communication*, *40*(1–2), 227–256. https://doi.org/10.1016/s0167-6393(02)00084-5

Schirmer, A., & Kotz, S. A. (2003). ERP evidence for a sex-specific Stroop effect in emotional speech. *Journal of Cognitive Neuroscience*, *15*(8), 1135–1148. https://doi.org/10.1162/089892903322598102

Schirmer, A., & Kotz, S. A. (2006). Beyond the right hemisphere: Brain mechanisms mediating vocal emotional processing. *Trends in Cognitive Sciences*, *10*(1), 24–30. https://doi.org/10.1016/j.tics.2005.11.009

Schirmer, A., Zysset, S., Kotz, S. A., & Yves Von Cramon, D. (2004). Gender differences in the activation of inferior frontal cortex during emotional speech perception. *NeuroImage*, *21*(3), 1114–1123. https://doi.org/10.1016/j.neuroimage.2003.10.048

Schurz, M., Radua, J., Aichhorn, M., Richlan, F., & Perner, J. (2014). Fractionating theory of mind: A meta-analysis of functional brain imaging studies. *Neuroscience & Biobehavioral Reviews*, *42*, 9–34. https://doi.org/10.1016/j.neubiorev.2014.01.009

Searle, J. R. (1979). *Expression and meaning*. Cambridge University Press. https://doi.org/10.1017/cbo9780511609213

Seydell-Greenwald, A., Chambers, C. E., Ferrara, K., & Newport, E. L. (2020). What you say versus how you say it: Comparing sentence comprehension and emotional prosody processing using fMRI. *NeuroImage*, *209*, 116509. https://doi.org/10.1016/j.neuroimage.2019.116509

Shibata, M., Toyomura, A., Itoh, H., & Abe, J. (2010). Neural substrates of irony comprehension: A functional MRI study. *Brain Research*, *1308*, 114–123. https://doi.org/10.1016/j.brainres.2009.10.030

Song, Y., Nie, Z., & Shan, J. (2024). Comprehension of irony in autistic children: The role of theory of mind and executive function. *Autism Research*, *17*(1), 109–124. https://doi.org/10.1002/aur.3051

Spotorno, N., Koun, E., Prado, J., Van Der Henst, J., & Noveck, I. A. (2012). Neural evidence that utterance-processing entails mentalizing: The case of irony. *NeuroImage*, *63*(1), 25–39. https://doi.org/10.1016/j.neuroimage.2012.06.046

Tamir, D. I., Thornton, M. A., Contreras, J. M., & Mitchell, J. P. (2016). Neural evidence that three dimensions organize mental state representation: Rationality, social impact, and valence. *Proceedings of the National Academy of Sciences*, *113*(1), 194–199. https://doi.org/10.1073/pnas.1511905112

Tettamanti, M., Vaghi, M. M., Bara, B. G., Cappa, S. F., Enrici, I., & Adenzato, M. (2017). Effective connectivity gateways to the theory of mind network in processing communicative intention. *NeuroImage*, *155*, 169–176. https://doi.org/10.1016/j.neuroimage.2017.04.050

Turken, U., & Dronkers, N. F. (2011). The neural architecture of the language comprehension network: Converging evidence from lesion and connectivity analyses. *Frontiers in Systems Neuroscience*, *5*. https://doi.org/10.3389/fnsys.2011.00001

Uchiyama, H., Seki, A., Kageyama, H., Saito, D. N., Koeda, T., Ohno, K., & Sadato, N. (2006). Neural substrates of sarcasm: A functional magnetic-resonance imaging study. *Brain Research*, *1124*(1), 100–110. https://doi.org/10.1016/j.brainres.2006.09.088

Van Overwalle, F. (2009). Social cognition and the brain: A meta-analysis. *Human Brain Mapping*, *30*(3), 829–858. https://doi.org/10.1002/hbm.20547

Van Overwalle, F., & Baetens, K. (2009). Understanding others' actions and goals by mirror and mentalizing systems: A meta-analysis. *NeuroImage*, *48*(3), 564–584. https://doi.org/10.1016/j.neuroimage.2009.06.009

Vandenberghe, R., Nobre, A. C., & Price, C. J. (2002). The response of left temporal cortex to sentences. *Journal of Cognitive Neuroscience*, *14*(4), 550–560. https://doi.org/10.1162/08989290260045800

Westerlund, M., & Pylkkänen, L. (2014). The role of the left anterior temporal lobe in semantic composition vs. semantic memory. *Neuropsychologia*, *57*, 59–70. https://doi.org/10.1016/j.neuropsychologia.2014.03.001

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). CONN: A functional connectivity toolbox for corre-
lated and anticorrelated brain networks. *Brain Connectivity*, *2*(3), 125–141. https://doi.org/10.
1089/brain.2012.0073

Wiethoff, S., Wildgruber, D., Kreifelts, B., Becker, H., Herbert, C., Grodd, W., & Ethofer, T. (2008).
Cerebral processing of emotional prosody—influence of acoustic parameters and arousal. *Neu-
roImage*, *39*(2), 885–893. https://doi.org/10.1016/j.neuroimage.2007.09.028

Wildgruber, D., Ackermann, H., Kreifelts, B., & Ethofer, T. (2006). Cerebral processing of linguistic
and emotional prosody: fMRI studies. In S. Anders, G. Ende, M. Junghofer, J. Kissler, & D.
Wildgruber (Eds.), *Progress in brain research* (pp. 249–268, Vol. 156). Elsevier. https://doi.org/
10.1016/S0079-6123(06)56013-3

Wildgruber, D., Ethofer, T., Grandjean, D., & Kreifelts, B. (2009). A cerebral network model of speech
prosody comprehension. *International Journal of Speech-Language Pathology*, *11*(4), 277–281.
https://doi.org/10.1080/17549500902943043

Witteman, J., van Ijzendoorn, M. H., van de Velde, D., van Heuven, V. J., & Schiller, N. O. (2011). The
nature of hemispheric specialization for linguistic and emotional prosodic perception: A meta-
analysis of the lesion literature. *Neuropsychologia*, *49*(13), 3722–3738. https://doi.org/10.1016/
j.neuropsychologia.2011.09.028

Wittfoth, M., Schröder, C., Schardt, D. M., Dengler, R., Heinze, H., & Kotz, S. A. (2009). On emo-
tional conflict: Interference resolution of happy and angry prosody reveals valence-specific effects.
Cerebral Cortex, *20*(2), 383–392. https://doi.org/10.1093/cercor/bhp106

Yue, Q., Zhang, L., Xu, G., Shu, H., & Li, P. (2013). Task-modulated activation and functional connec-
tivity of the temporal and frontal areas during speech comprehension. *Neuroscience*, *237*, 87–95.
https://doi.org/10.1016/j.neuroscience.2012.12.067

Zhu, N., & Wang, Z. (2020). The paradox of sarcasm: Theory of mind and sarcasm use in adults.
Personality and Individual Differences, *163*, 110035. https://doi.org/10.1016/j.paid.2020.110035