Title:

-    Structural basis of CO_2 valence coding in *Drosophila*

Authors:

Example: Javorski Dominik[1], Bergkirchner Beate[1], Ensinger Gernot[1], Lingl Alexander[1], Navolic
Jelena[2], Batawi Ashwaq[3], Hummel Thomas[1]

Affiliations:

[1]Department of Neurosciences and Development, University of Vienna, Djerassiplatz 1, 1030
Vienna, Austria

[2]Center for Molecular Neurobiology Hamburg, University Medical Center Hamburg-
Eppendorf, Falkenried 94, 20251 Hamburg, Germany

[3]King Abdulaziz University, 21589 Jeddah, Saudi Arabia

*Corresponding author: Email: thomas.hummel@univie.ac.at

Abstract

In the olfactory system, glomerular sensory channels of single receptor identity support

reliable odor recognition for appropriate approach or avoidance behaviors. For many olfactory

stimuli, the assigned sensory value is innate but modulated by the internal state and previous

experiences. How context-dependent modulation of innate valence coding supports distinct

behavioral responses is poorly understood. Here we show that CO_2 sensory information in

Drosophila, intrinsically aversive but modified by attractive food signals, diverges from the

canonical glomerular channel already in the antennal lobe and is relayed via the polarized local

interneuron LN23. LN23 relays sensory input via an extraglomerular CO_2 pathway and

manipulation of LN23 activity revealed a dominant role in CO_2-induced avoidance behavior.

The extraglomerular CO_2 pathway projects to the posterior lateral protocerebrum (PLP)

adjacent to the canonical Lateral Horn (LH) olfactory processing center and segregates into

anatomically distinct valence channels. Connectome data together with functional

characterization showed the convergence of parallel CO_2 channels onto two interconnected

third-order neurons. These neurons integrate additional sensory modalities via distinct

mechanisms: while the glomerular CO_2 pathway converges with food relay neurons onto

separated dendritic domains of the PD5 interneuron in the LH, the extraglomerular pathways

integrating CO_2 information with antennal humidity and temperature modalities establish

antagonistic inputs onto the PLP interneuron PV9. This early anatomical divergence of a

defined olfactory channel followed by separated multi-modal integration provides a structural

basis for context-dependent valence coding and appropriate behavioral responses.

Introduction

The ability to recognize and react to environmental stimuli is essential for the survival of animals. Insect olfactory systems allow the sensing and integration of volatile signals to identify food resources and avoid potential threats (Wilson, 2013). Although CO_2 is a component of the atmosphere, local changes in ambient CO_2 levels can induce context-dependent behavioral responses (Faucher et al. 2006; Hansson & Stensmyr, 2011). Walking *Drosophila* show robust CO_2-induced avoidance (Suh et al. 2004; Kwon et al. 2007; Faucher et al. 2006), while in more aroused states such as flight or foraging, CO_2 becomes an attractive olfactory cue (van Breugel et al. 2018) which is linked to dopamine levels affecting mushroom body output neurons (Lewis et al. 2015; Sachse & Beshel 2016). At the circuit level, prolonged CO_2 exposure leads to olfactory habituation, mediated by plasticity of GABAergic interneurons in the antennal lobe (Das et al. 2011; Olsen & Wilson, 2008; Nagel & Wilson, 2016; Sachse et al. 2007; Semelidou et al. 2018). Interestingly, *Drosophila* avoidance behavior directly correlates with the activity of CO_2-specific olfactory receptor neurons (MacWilliam et al. 2018), raising the question of how CO_2-induced sensory activity is transformed into central valence representations and modulated to support context-dependent behavior (Frechter et al. 2019).

The peripheral CO_2 circuit follows the canonical organization of olfactory sensory channels. A single population of ORNs (ab1C) co-express the gustatory receptors Gr21a and Gr63a and projects ipsilaterally to the V glomerulus in the ventral antennal lobe (Suh et al. 2004; Kwon et al. 2007; Vosshall & Stocker 2007). CO_2 sensation is transmitted to higher brain centers through two glomerular PNv types: a bilateral cholinergic PN (PNv^{bi}) projecting to both the mushroom Body (MB) and lateral Horn (LH), and a LH specific relay via a unilateral PN (PNv^{uni}) (Grabe et al. 2016; Lin et al. 2013; Sachse et al. 2007; Tanaka et al. 2012). Although distinct spatial LH domains encoding opposite hedonic values and odor intensity have been identified (Frechter et al. 2019; Das Chakraborty et al. 2022), glomerular PNs provide synaptic input to a diverse set of LH neurons making the identification of central brain representation and context-dependent modulation of a given sensory channel difficult (Dolan et al. 2019; Scheffer et al. 2020).

Here we describe the identification of a novel CO_2 relay channel, morphologically and
functionally separate from the canonical glomerular circuit via a polarized olfactory
interneuron, LN23. The LN23 pathway mediates a robust CO_2-induced aversion largely
unaffected by glomerular circuit plasticity but allows early multi-modal integration.
Furthermore, the parallel pathways of glomerular and extraglomerular PNs converge onto a
small group of LH interneurons as the core domain for CO_2 valence coding in the central brain.

Material and Methods
92

93 ***Drosophila* Stocks**
94

VT031497 (PNv^{bi} Gal4 and the alternative line: R53A05; BDSC#38859; BDSC: Bloomington
Drosophila Stock Center), NP7273 (PNv^{uni} Gal4), R44A02 (LN23 Gal4; BDSC#50196) were used
to target neurons PNv^{bi}, PNv^{uni} and LN23 respectively. LN23-lexA (BDSC#53644) and PNv^{bi}-lexA
(VT031497 lexA) were used in sybGRASP (BDSC#93197 / reverse direction BDSC#64315) and
tGRASP (BDSC#79040/reverse direction BDSC#79039) experiments to express the
postsynaptic GFP fragment. Gr21-Gal4 (BDSC#57600) was used to target ORNs that project
specifically to the V-glomerulus. To target LNv1 and LNv2, we used NP1227P from the Max
Planck Institute and NP2426-P from the Kyoto Stock Center. The Gal4 driver line BDSC #40444
was used for l2LN19. UAS-*shibire*^{ts} (Gift from Barry Dickson) was used to inhibit neurons at
restrictive 31°C. UAS-ReaCh (BDSC#53741) was used in optogenetic experiments to artificially
activate specific neuronal subsets which are labelled by the Gal4 lines. CantonS and w1118
wild-type flies were used for control experiments. The Gr63a^1 (BDSC#9941) mutant stock was
used in behavior experiments as a control for flies which are unable to detect CO_2. In addition
to our main LN23 line (BDSC#50196) an additional line also labeling LN23 (BDSC#48408) was
used for optogenetics. UAS-CaLexA (BDSC#66542) was used in combination with the Gal4 lines
to label active neurons. To further activate or silence specifically active neurons, a CaLexA
stock without GFP was used (BDSC#66543) in combination with lexAop-ReaCh or lexAop-shi^{ts}
respectively. UAS-*DenMark::cherry* was used to label dendrites of neurons while UAS-*syb::GFP*
and UAS-*syt::GFP* specifically labeled presynaptic terminals. To confirm the location of active
T-bars in LN23 and PNv^{bi} neurons, we also applied UAS-*brp::GFP* (BDSC#36291 and
BDSC#36292).

Immunohistochemistry

Flies were dissected in PBS (phosphate buffer solution) and brains were fixed in 2% PFA (Paraformaldehyde). After the fixation brains were washed 3 times with 0.3% PBT (phosphate buffer with TritonX) and then blocked with goat serum. After 1 hour of blocking the first AB (Antibody) was applied and the brains were kept in a 4°C fridge overnight. The next day brains were washed again 3 times before applying the second AB. After the staining protocol was complete, brains were mounted and subsequently scanned with a confocal microscope. Anti-Ncad rat (1:10, DHSB) was used to label all neuropiles in the fly brain in all experiments. For CaLexA experiments anti GFP rabbit (1:5000, Invitrogen) was used additionally to enhance the signal strength, according to Masuyama et al. (2012). GRASP experiments, DenMark and labeling of presynaptic terminals (*syt::GFP* and *syb::GFP*) didn't require any antibodies besides anti-Ncad since the endogenous signal was sufficient and changes after CO_2 incubation should not be distorted. For MCFO, first Antibodies were Anti-Ncad rat (1:10) or Anti-ollas rat (1:2000), Anti-flag mouse (1:2000) and Anti-HA rabbit (1:2000). As second antibodies anti-rat 647 (1:500), anti-mouse (highly cross-absorbed, 1:300) and anti-rabbit (1:500) were applied.

MCFO (Multicolor flip-out)

To label single cells in a usually rather broad expressing Gal4-line, we took advantage of a method established by Nern et al. 2015, called MCFO. They recombined four constructs, consisting of an UAS-site, followed by a stop-cassette to suppress transcription and one of four different transmembrane proteins, which can, if expressed, be labeled via antibodies. These four independent constructs can also be expressed together in the same cell, which leads to a mixture of colors and increased probability of one cell getting labeled in one distinct color. The stochastic excision of the stop cassette is achieved by a heat sensitive enzyme, whose activation is triggered by incubating the flies at 37°C in a water bath.

GRASP

To visualize synaptic contacts two different GRASP tools were used. SybGRASP is activity dependent because one fragment of GFP is located on vesicle membranes. As vesicles fuse with the presynaptic membrane, which is dependent on incoming action potentials, the large GFP fragment is exposed in the synaptic cleft and can reconstitute with the small GFP fragment

located on the postsynaptic membrane (Macpherson et al. 2015). tGRASP, or targeted GRASP

(Shearin et al. 2018), is activity independent and the two GFP fragments are bound to pre and

postsynaptic specific proteins (Cacophony and Telencephalin respectively). A comparison of

these two GRASP techniques allowed us to make assumptions about changes of synaptic

activity and changes to synaptic density. No antibodies against GFP were used since the

endogenous expression was sufficient. GFP intensities were quantified with FIJI, which is

measured as the mean gray value of a defined region of interest (ROI) above a threshold. To

subtract the mean gray value of the background noise, intensity was measured in another

glomerulus (same size of ROI) which was not showing any signal. Each datapoint in the

diagrams represents one quantified glomerulus. The same applies to intensity measurements

for the following two sections.

DenMark::cherry / Syt:GFP, syb::GFP

UAS-*syt::GFP* and UAS-*syb::GFP* were used to label presynaptic terminals of target neurons.

Synaptotagmin (*syt*) and synaptobrevin (*syb*) are proteins of the vesicle membrane and

therefore a Gal4 driven expression of the GFP tagged proteins shows strong signal in the

presynaptic terminals. UAS-*DenMark* (Dendritic Marker) is a somatodendritic marker (*cherry*)

to label dendrites. *GFP/Cherry* intensities were quantified with FIJI.

CaLexA

CaLexA (Masuyama et al. 2012) was used to label active cells. It is based on the NFAT

transcription factor, which is calcium responsive. As Ca^+ enters a cell, a chimeric lexA-VP16-

NFAT, enters the nucleus where expressed lexA can bind to lexAop inducing the expression of

GFP. The stronger and longer the excitation lasts, higher GFP intensities are expected. GFP

levels were quantified in FIJI as it was described in the section about GRASP. There was no

selection for gender and all flies tested were 2 days old at the start of the experiments. In the

experimental condition where flies were kept under the elevated level of 5% CO_2 the duration

of the exposure was 3 days. We also used a CaLexA construct where expressed lexA is not

expressing GFP but instead driving the expression of a temperature sensitive shibire protein

to silence neurons. Further we also used this stock in combination with lexA driven expression

of ReaCh to specifically further activate physiologically active neurons. 99% spermidine was

used as an inhibitor to GR21a receptors. 5µl were put on a small piece of cotton which was

then transferred to a vial containing the flies which were used in the experiments after 3 days

of exposure.

Behavioral experiments

For behavioral experiments, flies were kept in total darkness, in 25°C incubators in vials

containing fly food. Flies which were used in optogenetics were raised on fly food containing

all-trans-retinal to enhance their response to light exposure (Vilinsky et al. 2018). All flies were

age matched and tested when they were 3-4 days old. Since there was no observed difference

between male and female flies, both sexes were used in all behavioral experiments. As we did

not observe any behavioral artifacts caused by either green or red LED light we opted for the

green LEDs, since they elicited a stronger response in the flies (compared to Inagaki et al.

2014).

Flies were immediately transferred to empty vials which can be connected to the CO_2 or

optogenetic setup after they were taken out of the incubators. The whole procedure was also

done in darkness (red light was used to be able to handle the flies, which is barely visible for

the flies) to minimize visual distraction affecting the fly's behavior. The whole apparatus was

shifted sideways, and the flies were gently moved to the elevator segment of the setup. The

elevator was lowered to a position short above the decision point and the flies were given 60

seconds rest. After the resting period either a green LED light or CO_2 flow was activated and

the elevator was lowered to the decision point.

The animals were given 20 seconds to move inside the behavioral setup and make a choice.

Afterwards the elevator was pulled up again, so that the flies remained in either the test tube,

control tube or the central elevator area. Each of these populations was counted and a

preference index (PI) calculated, defined as the percentage of flies located in the control arm

(air-filled /dark) of the setup, minus the percentage of flies located in the test arm (CO_2

filled/light). Since in some of the tested populations, a few animals remained in the central

elevator part of the arena, their decision was defined as neutral. Their number was divided by

two and added to both the test and control population for the PI. Each datapoint in the

diagrams represents the PI of a group of flies (n=15-40).

Hemibrain analyses

Connectivity data were obtained from the neuPrint interface (neuPrint, Janelia Research Campus; https://neuprint.janelia.org/) built on the Hemibrain connectome (Scheffer et al. 2020). It was used to study and model the neurons involved in the V-circuit. The main goal was to identify synaptic connectivity between the different layers of information processing within the circuit. Finally, the putative circuit was reconstructed in 3D and PN/LN output zones were identified. Neuronbridge.janelia.org was used to gather 3D images from l2_PNm17 and sm_PNm1. FlyWire.ai was a tool to quickly get information about the predicted neurotransmitter identities of neurons.

Statistics

All diagrams and statistical analyses were done by using GraphPad Prism 9 software. Normally distributed data were compared with a student's t-test, whereas datasets that did not meet normality assumptions were analyzed with a Mann–Whitney U test. Behavioral data spanning multiple days were tested with a one-way ANOVA. Statistical significance is indicated in the figures by asterisks, with $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), and $p < 0.0001$ (****).

Results

Glomerular circuit dynamics of CO_2 perception

The V glomerulus at the ventral AL surface is innervated by a single population of about 40 CO_2-sensitive olfactory receptor neurons (ORNv) expressing the two gustatory receptors GR21a and GR63a (Kwon et al. 2007; Jones et al. 2007) (Fig.1A-B). Each ORNv axon forms 3-5 terminal branches tightly packed in a non-overlapping fashion with other ORN terminals (Fig. 1, C-C´´), providing sensory input to glomerular inter- and projection neurons (Fig1D-G´) (Grabe et al. 2016).

To identify relay neurons of the V glomerulus with different physiological properties we screened a collection of Gal4 lines using the CaLexA activity reporter to compare intracellular

Ca^{2+} levels under ambient CO_2 conditions (Koyama et al. 2021; Masuyama et al. 2012) (Fig. 1H-

O). Gal4 lines for ORNv show strong CaLexA reporter activity (Fig.1H), in line with a high default

activity of CO_2-receptive sensory neurons due to the absence of $GABA_B$-mediated presynaptic

inhibition (MacWilliam et al. 2018; Olsen & Wilson, 2008; Root et al. 2008). In contrast to

elevated Ca^{2+} levels in ORNv, the two uniglomerular PN classes of the V glomerulus, unilateral

V_l2PNs (PNv[uni]) and bilateral V_ilPNs (PNv[bi], Fig. 1D) do not show any significant level of

CaLexA activation under ambient air conditions (Fig. 1I, J). Two main classes of local

interneurons, LNv1 and LNv2 display moderate levels of CaLexA activity at atmospheric CO_2

levels (Fig. 1K, L). A strong CaLexA-GFP signal, comparable to ORNv can be observed for l2LN23

(LN23), a less characterized olfactory class of sparsely innervating local interneurons (Fig.1F,

F', M and N) (Liou et al. 2018). Fluorescence signal of Ca^{2+}-induced GFP expression quickly

decreases in LN23 following sensory deprivation in adult flies via removal of both antennae

(Fig.1O). Furthermore, an extended exposure to the ORNv inhibitor spermidine (MacWilliam

et al. 2018) results in a reduced CaLexA activity not only in ORNv, but also LN23, indicating a

preferred physiological coupling of sensory neurons to LN23 interneurons under ambient CO_2

conditions (Fig. 1O).

A predominant relay of CO_2-induced sensory activity by LN23 is supported by the modified

activity-dependent GRASP technique (Macpherson et al. 2015). Expression of the presynaptic

syb::GFP[1-10] in ORNv and the postsynaptic *GFP[11]* in AL neurons revealed a significantly stronger

GRASP signal for ORNv→LN23 connectivity compared to the ORNv→PNv[bi] synaptic activity

(Fig.2F and G). Similarly, the activity-independent tGRASP (Shearin et al. 2018) is most

prominent for ORNv→LN23 (Fig.2I and J), which is in line with connectome data

(neuPrintExplorer, hemibrain; Scheffer et al. 2020; Schlegel et al. 2021). In summary, ORNv

activity at ambient CO_2 levels seems to be mainly represented by the LN23 class of local

interneurons and less by glomerular PN relay neurons, thereby identifying a new circuit

component for the ORNv-induced behavioral aversion.

Cell type plasticity of glomerular relay neurons following chronic CO_2 stimulation

We next determined how the overall activity of V glomerulus relay- and interneurons changes following long-lasting sensory stimulation. Adult *Drosophila melanogaster* avoids ambient levels of CO_2 in the T-maze choice assay (Faucher et al. 2006; Suh et al. 2004) (Fig. 2A) but shows behavioral habituation following chronic CO_2 exposure (Das et al. 2011, Fig. 2B, C). The reduction of CO_2-induced aversive responses is associated with an increase in the synaptic volume of the V glomerulus (Fig.2D; Sachse et al. 2007 and Das et al. 2011). At the ORN level, mosaic analysis revealed an extension in the surface area and size of individual synaptic boutons of ORNv axons (Fig.2E). Compared to ambient CO_2 levels, prolonged elevated CO_2 (5%) sensation results in neuron-type specific changes of CaLexA activity (Fig. 1 H'-N') (Masuyama et al. 2012). While PNv^{bi} neurons showed a strong increase in Ca^{2+} levels following CO_2 habituation (Fig.1I'), whereas no CaLexA-dependent GFP expression was observed in PNv^{uni} (Fig.1J'), indicating different physiological properties of the two main glomerular CO_2 relay channels (Lin et al. 2013). In addition to chronic CO_2 stimulation, short pulses of elevated CO_2 also did not induce Ca^{2+}-mediated-GFP expression in PNv^{uni} (Fig.1O). No significant changes in CaLexA-GFP levels could be detected for LN23 neurons following prolonged CO_2 exposure compared to the already high expression at ambient CO_2 conditions (Fig1M', N', O).

As GABAergic local interneurons are associated with olfactory habituation (Das et al. 2011; Sadanandappa et al. 2013) we compared changes in CaLexA activation in LNv1 and LNv2 (Das et al. 2011, Sachse et al. 2007). While an increase in the Ca^{2+} reporter signal could be observed for both LN types following chronic CO_2 exposure, the activity of LNv1 is significantly higher compared to LNv2 (Fig.1K' and L'). In line with the cell-type specific CaLexA expression, previous connectome analysis showed significant differences of LN input for CO_2 relay neurons: (neuPrintExplorer, hemibrain): While both PNv^{uni} and LN23 do not receive major LN-type input, PNv^{bi} gains GABAergic modulation by lLN1 neurons ("bc" subtype, 16% of input) and a group of multiple lLN2 neuron types (X, F, P, 19%) (Schlegel et al. 2021; Bates et al. 2020). These results indicate cell type specific physiological changes of glomerular neurons following CO_2-induced behavioral habituation, in which PNv^{bi} shows dynamic changes in glomerular connectivity while both PNv^{uni} and LN23 neurons maintained their characteristic activity seen in ambient CO_2 conditions.

We next tested how the cell-type specific changes in neuronal activity during habituation

impact their synaptic connectivity within the V glomerulus. A strong increase of activity-

dependent sybGRASP levels for ORNv input onto PNv[bi] could be detected following chronic

CO_2 stimulation (Fig.2F'). In contrast, no significant changes in sybGRASP levels were observed

for ORNv input onto LN23 (Fig.2G′). Of note, compared to ambient CO_2 conditions, the

standard deviation for ORNv→LN23 sybGRASP levels was reduced in flies chronically exposed

to CO_2, indicating that sensory input onto LN23 also increases in the context of habituation

(Fig.2H). In addition, activity-independent tGRASP expression showed that ORNs increase

their synaptic connections to the two main relay neurons, PNv[bi] and LN23 (Fig.2I', J'), which is

in line with morphological changes of ORN axon terminals described above (Fig.2D, E).

Following prolonged CO_2 exposure, the PNv[bi] class displayed increased sybGRASP levels on

PNv[uni] and LN23 (Fig.2L-M'''), while no significant changes could be observed for LN23 on PNv[bi]

(Fig.2N-N'''). Taken together, CO_2-induced ORNv activity seems to be relayed via two distinct

glomerular output channels dependent on the stimulus strength: LN23 neurons indicate

default ambient CO_2 levels while PNv[bi] signals at conditions of extended elevated CO_2.

Identity, innervation patterns and polarity of second-order CO_2 sensitive neurons

To test if the difference in CaLexA activity might be related to the organization of presynaptic

sites within CO_2 relay neurons we visualized the active zones using *syt::GFP/brp::GFP*

transgenes together with the dendrite marker *DenMark::mCherry* (Kittel et al. 2006, Nicolai et

al. 2010; Wagh et al. 2006). PNv[bi], PNv[uni] and LN23 differ significantly in their expression and

localization of presynaptic proteins within the dendritic compartment (Fig.3A-C). While PNv[bi]

shows prominent dendritic *syt::GFP* localization throughout the glomerulus (Fig.3C), no

syt::GFP signal can be detected within PNv[uni] dendrites (Fig.3A), indicating different cellular

polarity of the two main glomerular PNv classes. Despite the significant expression level,

syt::GFP appeared more scattered within LN23 dendrites and less restricted to the glomerulus

neuropil compared to PNv[bi] (Fig.3B). In fact, single optical sections allow the localization of the

active zone marker *Brp::GFP* in LN23 dendrites outside the glomerular neuropile (Fig.3F) but

in PNv[bi] confined to the V glomerulus (Fig.3G). In line with these results hemibrain EM data

lists enriched presynaptic site in LN23 dendrites outside the V glomerulus in contrast to the

restriction PNv[bi] presynaptic sites to the glomerular neuropil (Fig.3I)

(https://neuprint.janelia.org/; Schlegel et al. 2021; Bates et al. 2020).

Following chronic sensory stimulation, PNv[bi] showed a significant increase in the density of

presynaptic sites while no significant change in the amount presynaptic *syt::GFP* labeling of

LN23 dendrites can be observed (Fig.3D). Like the subcellular organization at ambient CO_2

conditions, presynaptic sites of PNv[bi] dendrites are uniformly distributed throughout the V-

glomerulus (Fig.3D, last image), while most presynaptic sites of LN23 dendrites are found

outside of the glomerular boundaries (Fig.3D, middle image). In contrast to CO_2 relay neurons

PNv[bi] and LN23, no dendritic localization of presynaptic marker transgenes could be detected

for PNv[uni] (Fig.3D, first image), neither under ambient nor chronic CO_2 condition, further

supporting a differential synaptic profile of the two PN classes. All *syt::GFP* and

DenMark::cherry intensities were quantified using ImageJ (Fig.3H). An additional presynaptic

marker *syb::GFP* revealed a similar expression pattern: while PNv[bi] showed a strong signal

within the glomerular boundaries, which increases following CO_2 incubation, LN23 dendrites

show a robust GFP signal adjacent to the neuropil border of the V glomerulus (Fig.3E). As both,

PNv[bi] and PNv[uni] are described as being excitatory, cholinergic neurons (Bates et al. 2020; Lin

et al. 2013; Schlegel et al. 2021; Tanaka et al. 2012) we investigated the neurotransmitter

identity of LN23. Anti-GABA immunostaining does not label LN23 cell bodies (Supplemental

Fig.3A), excluding an inhibitory function similar to canonical interglomerular LNs. These results

revealed not only major differences in neuronal organization among the 3 classes of CO_2 relay

neurons, but also a distinct response profile in intraglomerular plasticity of dendro-dendritic

connections. CO_2 input diverges into two more robust feed-forward channels (PNv[uni] towards

LH and LN23 within the AL) while the dynamic PNv[bi] pattern indicates intraglomerular

processing.

Developmental profile of the inter-glomerular CO_2 relay neuron

Having established the polarity of LN23, our next objective was to investigate the

development of this neuron, specifically focusing on how interglomerular relay neurons are

integrated into stereotyped antennal lobe circuits. While the glomerulus-specific assembly of

olfactory receptor neuron (ORN) axons with projection neuron (PN) dendrites has been

extensively documented (Hong & Luo 2014), less is known about the mechanisms underlying

the incorporation of relay neurons into these circuits. At the beginning of pupal development,

PN dendrites innervate the early precursor of the adult AL and occupy distinct spatial domains

before ORN axons enter the AL by around 20h APF (Jefferis et al. 2001; Jefferis et al. 2004).

LN23 neurons can be traced back to the late larval stage with a cellular morphology related to

SEZ (subesophageal zone) interneurons (Truman & Bate, 1988; Winding et al. 2023, Fig. 4).

During early pupal stages, larval LN23 precursor neurons retract from the SEZ to accumulate

at the ventral region of the AL precursor neuropile (Fig4A-C'). Here LN23 processes target a

region which is devoid of dendrites of uniglomerular PNs (Fig4F, F'). Before ORN axon arrival,

LN23 neurons establish a separate dendritic branch, which extends dorsally before bending

back to the posterior AL and merges with the main dendritic domain (Fig. 4D-E').

Loss of the secreted molecule *Wnt5*, which affects PN dendrite patterning (Hing et al. 2020;

Wu et al. 2014), leads to a continued growth of the dorsal LN23 branch along PN axon tracts

which exit the AL towards higher olfactory brain centers (Fig.4I-K). These results indicate that

LN23 neurons display an intrinsic polarity similar to uniglomerular PNs, but instead of relaying

sensory information from AL glomeruli to the LH/MB, a *Wnt5*-dependent mechanism seems

to prevent AL exit and target the presynaptic process to the extraglomerular region in the

ventral AL. In wild-type, ORN axons innervate the developing V glomerulus only at the

ipsilateral AL, in contrast to the majority of bilateral ORN classes (Fig.4F-H) (Couto et al. 2005).

To gain insight into the mechanisms which support glomerulus-specific ORNv input on LN23,

we analyzed candidate genes which are expressed on growing ORNs. Mutations in the

protocadherin-type cell adhesion molecule Flamingo (*fmi*), which is highly enriched on ORN

axons in the developing V glomerulus, affect their synaptic targeting. Here, *fmi* mutant ORNv

axons are no longer restricted to the glomerular boundaries but extend into the posterior AL

next to the output domain of LN23 (Suppl.Fig.4A, B), indicating cellular interactions to restrict

class-specific axon-dendrite recognition within the glomerular domain.

In summary, despite being morphologically classified as a local interneuron, the LN23 class

shows cellular and developmental features similar to those of olfactory projection neurons: a

highly polarized neuronal morphology in the adult olfactory system is established by a

glomerulus-specific dendrite targeting for singular ORN input and a default axonal extension

along olfactory PN tracts rerouted by external signals (Komiyama & Luo, 2006).

LN23 mediates CO_2-induced aversion

To test the role of the different CO_2 relay neurons on *Drosophila* behavior, we expressed the *shibire*[ts] transgene to block synaptic transmission in a cell type specific manner (Chen et al. 1991; Kitamoto 2001). Neuronal silencing of each of the two glomerular PNs led to a reduction of innate CO_2 avoidance, with stronger effects observed for PNv[uni] compared to PNv[bi] (Fig. 5A). While CO_2 avoidance can still be observed following the neuronal silencing of each PN class, *shibire*[ts] induced silencing of LN23 fully abolished aversion behavior towards CO_2 (Fig. 5A). To test for cell type specificity of the behavioral phenotype following LN23 silencing, we used the Ca^{2+} dependent LexA activation of the CaLexA construct for shi[ts] expression and observed a reduction of the innate CO_2 aversion (Fig. 5B) (Masuyama et al. 2012). To increase lexA levels we chronically exposed flies to high levels of CO_2. Interestingly, increased CaLexA mediated shi[ts] expression via chronic exposure not only suppressed behavioral aversion but led to a significant level of attraction towards CO_2 (Fig.5B). These results indicate a prominent function of LN23 in CO_2 valence coding: Blocking LN23 activity is sufficient to fully suppress CO_2-induced behavioral aversion and together with the GABAergic suppression of PNv[bi] during habituation an attraction towards CO_2 is observed, which phenocopies the silencing of ORNv (Jones et al. 2007; Kwon et al. 2007, MacWilliam et al. 2018; Suh et al. 2004).

To further determine if these relay neurons are sufficient for behavioral aversion, we optogenetically activated each neuron type and measured the behavioral response (Inagaki et al. 2014; Klapoetke et al. 2014; Pulver et al. 2009) (see M&M). Light-induced activation of ORNv (*GR21a-Gal4*) expressing UAS-ReaCh led to strong aversion (Fig. 5C). Similarly, optogenetic activation of LN23 is sufficient to trigger behavioral aversion via two independent Gal4 driver lines (LN23 Gal4 and the alternative line we used: VT48408-Gal4; Fig.5C) or by $Ca2^+$ induced LexA activation (Fig.5D). In contrast, no change in light-induced activation for both PNv[bi] (two independent driver lines, see Methods) nor PNv[uni] could be observed (Fig.5C). These results support a predominant role of LN23 in CO_2-induced avoidance behavior. Compared to the more dynamic activity of PNv[bi] following acute and chronic sensory stimulation, LN23 shows little changes in intrinsic activity, representing ORNv activity independent of changes in CO_2 levels or temporal pattern. Based on these findings we propose a ground state of ORNv/LN23 mediated aversion induced by ambient CO_2 default state, which is modified either by the parallel PN channels or downstream of LN23.

Third-order neurons postsynaptic to LN23 have a putative role in valence coding

To gain further insights into the unique organization of the LN23 pathway compared to the glomerular relay channel, we analyzed the postsynaptic candidate neurons within the posterior AL. Two PN types, l2_PNm17 and sm_PNm1 are receiving a large proportion of their synaptic input (27% and 18%, respectively) from LN23 (Fig.6A, neuPrint, Janelia Research Campus; https://neuprint.janelia.org/; Schlegel et al. 2021). Like the two PN classes of the V glomerulus, PNm1 and PNm17 show distinct relay properties: While input into PNm1 is mostly restricted to the LN23-CO_2 channel, dendrites of PNm17 extend into adjacent VP1 and VP3-5 glomeruli to receive direct input from thermo- and humidity- receptor neurons (neuPrint, Janelia Research Campus; https://neuprint.janelia.org/; Frank et al. 2015; Marin et al. 2020). Both types of non-olfactory relay neurons target multiple distinct brain areas adjacent to the canonical olfactory LH, the posterior lateral protocerebrum (PLP), superior clamp (SCL) and the superior lateral protocerebrum (SLP) (Fig6 A, B, C) (Ito et al. 2014). To visualize the trajectories of the PNs, we used MCFO (Aso et al. 2014) to create single cell clones, which show that PNm1 and PNm17 axons turn close before entering the LH to target adjacent brain regions (Fig. 6B, C). Of note, compared to glomerular PNv[uni] and PNv[bi], PNm1 and PNm17 receive only minor input from GABAergic LNs (neuPrint, Janelia Research Campus; https://neuprint.janelia.org/), indicating reduced gain control properties of the extra-glomerular ORN→LN23→PNm pathway (Olsen & Wilson, 2008).

Optogenetic activation of each of the two PNm class neurons downstream of LN23 induce opposite behavioral responses: While activity of the "CO_2-only" class PNm1 results in strong aversion, the "multi-modal" PNm17 initiates behavioral attraction (Fig.6D). Furthermore, CO_2-induced aversive behavior could be attenuated following shi[ts]-mediated silencing of PNm1 but was not changed by silencing PNm17 (Fig.6 E, F). Interestingly, FlyWire.ai (Dorkenwald et al. 2024; Eckstein et al. 2024; Schlegel et al. 2024; Zheng et al. 2018) revealed distinct neurotransmitter identities for the two PNm classes, GABA expression for PNm1, while PNm17 shows a cholinergic synaptic profile. Together with the behavioral responses following optogenetic activation this suggests a circuit motif in which the default activity of the ORNv→LN23→PNm1 channel by ambient CO_2 sensation blocks a central brain region important for the generation of behavioral attraction (Dolan et al. 2019).

Connectivity in higher brain centers and lateral communication in the antennal lobe

As the glomerular PNs and the extra-glomerular LN23→PNm channels target adjacent central brain regions we asked whether the two parallel CO_2 pathways converge onto common downstream neurons. The hemibrain dataset (https://neuprint.janelia.org/; Plaza et al. 2022; Scheffer et al. 2020) identified three main types of glutamatergic LH neurons (FlyWire.ai; Dorkenwald et al. 2024; Eckstein et al. 2024; Schlegel et al. 2024; Zheng et al. 2018) as PNv or PNm targets: LH^{PD5c1} as the primary target of PNv^{bi} (Fig.6 G), LH^{PV9b1} downstream of PNm1 (Fig.6 H) and LH^{CENT4} as the major postsynaptic partner of PNv^{uni} (Fig.6 I). All 3 LH target neurons of the parallel CO_2 pathways are directly interconnected (Fig. 8): While LH^{PV9b1} not only receives input from PNm1 but also from PNv^{bi}, LH^{PD5c1}, the main downstream target of PNv^{bi}, is also targeted by LH^{CENT4}, the primary postsynaptic partner of PNv^{uni}. This core motif of CO_2 perception receives additional input from other sensory modalities. LH^{PD5c1} gains input from "food-sensing" glomeruli (DM4, DC4/DP1, and VM7) via class-specific cholinergic PNs, indicating a neuronal integrator of CO_2 and food-related cues (Couto et al. 2005; Dumenil et al. 2025; Min et al. 2013). In addition, LH^{CENT4} is also a main target for $l2PN^{VP4+VL1}$, mediating the perception of dryness and pyrrolidine-type odorants, thereby adding another layer of multi-modal sampling to the LN23 → PNm1 pathway (Fig.8) (Li et al. 2022; Choi et al. 2022; Ni 2021). Interestingly, the convergence of the glomerular CO_2 and food channels is not only observed for LH neurons downstream of AL relay pathways but seems to occur already at the level of AL glomeruli. Hemibrain describes with l2LN19 another class of polarized local interneuron (Fig.7A, B) with enriched postsynaptic sites at "food-sensing" glomeruli DM4, DC4/DP1, and VM7 (Fig.7A) as well as targeted output onto PNv^{bi} and PNv^{uni} in the V-glomerulus (Fig.7B), but merely with LN23 (Fig.7C) (https://neuprint.janelia.org/; Schlegel et al. 2021; Bates et al.2020). Accordingly, optogenetic activation of LN19 led to an aversive behavioral response (Fig.7D), while inhibition had no effect (Fig.7E).

Based on these distinct connectivity patterns and behavioral response profiles of individual relay neurons we are proposing the following circuit mechanism for CO_2 valence coding (Fig.8): The activity of CO_2-responsive sensory neurons (ORNv, black) is relayed by parallel PN channels and converge onto two central brain target neurons of opposite intrinsic valence identity. The glomerular PNs target the ventral domain of LH^{PD5} to trigger behavioral aversion and extraglomerular PNs activated via the lateral LN23 relay converge onto LH^{PV9} which

mediates attraction. While PNm17 supports LH^{PV9} activity, GABAergic PNm1 blocks LH^{PV9}

through which the negative valence state of LH^{PD5} triggers avoidance. This core aversion

pathway (PNv^{bi}/PNv^{uni}/PNm1), without prominent GABAergic modulation, directly represents

the activity of CO_2 sensory neurons, is modified either via the activation of the dorsal LH^{PD5}

domain by fruit odor PNs ($PN^{DM/VM/DC}$) or by an increase of PNm17 through non-olfactory

sensory neurons to support a context-dependent behavioral response repertoire.

Discussion

In *Drosophila*, the activation of a single class of CO_2 receptive sensory neurons induces

avoidance behavior, which is modified by multiple external and internal stimuli (Bräcker et al.

2013; Faucher et al. 2006; Dissegna et al. 2021; Suh et al. 2004; Turner and Ray, 2009). How

does the *Drosophila* brain transform ORNv input into flexible behavioral responses towards

CO_2? Here we identified a novel CO_2 relay channel with unique morphological and

physiological properties compared to the canonical glomerular PN pathway. First, instead of

direct transmission of sensory input by glomerular PNs in the anterior AL towards higher brain

centers, the ORNv induced activation of the local interneuron LN23 relays peripheral CO_2

sensation to an extraglomerular neuropil of the posterior AL generating two parallel output

channels with different target areas. Second, in contrast to the high level of neuronal plasticity

observed for PNs within glomerular circuits, LN23 analysis revealed very little dynamic

changes under different physiological conditions. This might be due to distinct intrinsic

features of PNs and LN23 such as the expression of ion channels or receptor molecules but

could also be explained by the absence of glomerular GABAergic neurons at the

extraglomerular CO_2 pathway. Third, following the manipulation of neuronal activity, only

LN23, but no individual PNs are able to induce an aversive behavioral response similar to

ORNv. Finally, the extraglomerular CO_2 pathway includes a structural segregation of opposite

valence-encoding elements, in which the "CO_2-only" neuron induces aversion while activation

of the parallel "multi-modal" channel determines behavioral attraction. These findings are

consistent with a designated circuit motif encoding a default behavioral state of avoidance

even under atmospheric CO_2 levels. As the "CO_2-only" relay neuron PNm1 has a predicted

GABAergic identity (FlyWire.ai; Dorkenwald et al. 2024; Eckstein et al. 2024; Schlegel et al.

2024; Zheng et al. 2018), the previously reported ORNv induced avoidance (MacWilliam et al.

2018; Das et al. 2011) could be explained with the inhibition of a premotor area by the LN23

channel.

The interplay between the extraglomerular default channel and the glomerular dynamic

pathway in encoding behavioral avoidance was demonstrated in the context of habituation.

While the increased GABAergic inhibition of glomerular PNs following chronic CO_2 sensation

reduces innate aversion, the simultaneous silencing of LN23 not only eliminates all aversive

response features but reverses CO_2 valence to trigger attraction. Interestingly, the reduced

aversion through inhibition of PN activity in the context of habituation seems partially

compensated by increasing PNv^{bi} synaptic plasticity. We observed a strong gain of presynaptic

sites of glomerular PNv^{bi} dendrites onto LN23, which might maintain a balanced level of

aversion activity for local CO_2 levels following habituation (Das et al. 2011; Sachse et al. 2007).

Although both PNv^{uni} and PNv^{bi} are excitatory cholinergic projection neurons (Tanaka et al.

2012; Lin et al. 2013; Bates et al. 2020; Schlegel et al. 2021), they differ in their cellular

organization and behavioral function. In contrast to PNv^{bi}, PNv^{uni} dendrites lack presynaptic

sites, therefore specifies a feed-forward channel similar to LN23. Based on these results we

are proposing complementary function of the three relay neurons in mediating flexible CO_2

behavioral response patterns: While LN23 encodes default avoidance under ambient

conditions, supported by PNv^{uni} as a sensor for elevated local levels of CO_2, PNv^{bi} defines the

main component of experience-dependent modulation by LNs and neurons of the MB calyx

(CA) (Bräcker et al. 2013).

Interestingly, developmental analysis revealed a polarized growth pattern of LN23 that

resembles projection neurons more than typical local interneurons (Jefferis et al. 2004;

Komiyama & Luo 2006). LN23 originates from a larval interneuron located in the

suboesophageal zone (SEZ), which undergoes remodeling into an adult olfactory interneuron

during early pupal stages (Truman & Bate, 1988). We hypothesize that the secreted *Wnt5*

protein, a critical factor for initial projection neuron dendrite patterning, supports the

targeting of LN23 processes to the posterior antennal lobe (AL) region. Like the genetically

defined recognition mechanisms that guide glomerulus-specific ORN-PN connectivity, growing

ORNv axons appear to target LN23 dendrites as synaptic partners. Since LN23 dendrites are

organized into a defined glomerular input and extraglomerular output zone, we propose

Flamingo-dependent expression on ingrowing ORN axons to prevent sensory input to LN23
outside the glomerular domain.

The divergence of CO_2-induced ORN activation into parallel relay channels allows for early
convergence of additional sensory information. This includes not only the integration of non-
olfactory cues by the positive valence branch PNm17 downstream of LN23 (Frank et al. 2015;
Li et al. 2022), but also the sampling of food-odor-sensitive glomeruli by the polarized
interneuron LN19 for V-glomerulus input (Couto et al. 2005; Dumenil et al. 2025; Min et al.
2013). The convergence of different parallel CO_2 channels onto a common set of lateral horn
(LH) interneurons suggests a potential hub for motor action control (Dolan et al. 2019). The
results presented in this study support a conceptual model in which the default state of
behavioral aversion, induced through inhibition of an LH motor hub by the "CO_2-only"
ORNv→LN23→PNm1 pathway, is modified by the inclusion of additional glomerular and
extraglomerular projection neuron classes to allow approach behavior. Further studies on the
physiological architecture of the CO_2 core circuit within and downstream of the LH may reveal
deeper insights into the multi-modal processing underlying flexible, context-dependent
behaviors.

Additionally, despite the conserved molecular receptor identity of CO_2 sensory neurons
among insects, species-specific behavioral responses to local CO_2 concentrations are evident
(Turner & Ray 2009). In many species, such as mosquitoes, elevated CO_2 levels signal potential
food sources and elicit an attractive response (Tauxe et al. 2013; Siju 2009; Ghaninia 2007).
The identification of a dedicated CO_2 aversion channel in *D. melanogaster* opens new avenues
for comparative research into circuit motifs and provides insight into the neural basis of
behavioral flexibility across insect evolution (Oram & Card 2022).

References

Aso, Y., Hattori, D., Yu, Y., Johnston, R. M., Iyer, N. A., Ngo, T. T., Dionne, H., Abbott, L. F.,
Axel, R., Tanimoto, H., & Rubin, G. M. (2014). The neuronal architecture of the mushroom
body provides a logic for associative learning. *eLife*, *3*, e04577.
https://doi.org/10.7554/eLife.04577

Bates, A. S., Schlegel, P., Roberts, R. J. V., Drummond, N., Tamimi, I. F. M., Turnbull, R., Zhao,
X., Marin, E. C., Popovici, P. D., Dhawan, S., Jamasb, A., Javier, A., Serratosa Capdevila, L., Li,
F., Rubin, G. M., Waddell, S., Bock, D. D., Costa, M., & Jefferis, G. S. X. E. (2020). Complete
Connectomic Reconstruction of Olfactory Projection Neurons in the Fly Brain. *Current biology*
: CB, *30*(16), 3183–3199.e6. https://doi.org/10.1016/j.cub.2020.06.042

Bräcker, L. B., Siju, K. P., Varela, N., Aso, Y., Zhang, M., Hein, I., Vasconcelos, M. L., &
Grunwald Kadow, I. C. (2013). Essential role of the mushroom body in context-dependent CO_2
avoidance in Drosophila. *Current biology : CB*, *23*(13), 1228–1234.
https://doi.org/10.1016/j.cub.2013.05.029

Das Chakraborty, S., Chang, H., Hansson, B. S., & Sachse, S. (2022). Higher-order olfactory
neurons in the lateral horn support odor valence and odor identity coding
in *Drosophila*. *eLife*, *11*, e74637. https://doi.org/10.7554/eLife.74637

Chen, M. S., Obar, R. A., Schroeder, C. C., Austin, T. W., Poodry, C. A., Wadsworth, S. C., &
Vallee, R. B. (1991). Multiple forms of dynamin are encoded by shibire, a Drosophila gene
involved in endocytosis. *Nature*, *351*(6327), 583–586. https://doi.org/10.1038/351583a0

Choi, K., Kim, W. K., & Hyeon, C. (2022). Olfactory responses of *Drosophila* are encoded in the
organization of projection neurons. *eLife*, *11*, e77748. https://doi.org/10.7554/eLife.77748

Couto, A., Alenius, M., & Dickson, B. J. (2005). Molecular, anatomical, and functional
organization of the Drosophila olfactory system. *Current biology : CB*, *15*(17), 1535–1547.
https://doi.org/10.1016/j.cub.2005.07.034

Das, S., Sadanandappa, M. K., Dervan, A., Larkin, A., Lee, J. A., Sudhakaran, I. P., Priya, R.,

Heidari, R., Holohan, E. E., Pimentel, A., Gandhi, A., Ito, K., Sanyal, S., Wang, J. W., Rodrigues,

V., & Ramaswami, M. (2011). Plasticity of local GABAergic interneurons drives olfactory

habituation. *Proceedings of the National Academy of Sciences of the United States of*

America, *108*(36), E646–E654. https://doi.org/10.1073/pnas.1106411108

Dissegna, A., Turatto, M., & Chiandetti, C. (2021). Context-Specific Habituation: A

Review. *Animals : an open access journal from MDPI*, *11*(6), 1767.

https://doi.org/10.3390/ani11061767

Dolan, M. J., Frechter, S., Bates, A. S., Dan, C., Huoviala, P., Roberts, R. J. V., Schlegel, P.,

Dhawan, S., Tabano, R., Dionne, H., Christoforou, C., Close, K., Sutcliffe, B., Giuliani, B., Feng,

L., Costa, M., Ihrke, G., Meissner, G. W., Bock, D. D., Aso, Y., Rubin, G. M., & Jefferis, G. S. X.

E. (2019). Neurogenetic dissection of the Drosophila lateral horn reveals major outputs,

diverse behavioural functions, and interactions with the mushroom body. *eLife, 8*, e43079.

https://doi.org/10.7554/eLife.43079

Dorkenwald, S., Matsliah, A., Sterling, A. R., Schlegel, P., Yu, S. C., McKellar, C. E., Lin, A.,

Costa, M., Eichler, K., Yin, Y., Silversmith, W., Schneider-Mizell, C., Jordan, C. S., Brittain, D.,

Halageri, A., Kuehner, K., Ogedengbe, O., Morey, R., Gager, J., Kruk, K., … FlyWire

Consortium (2024). Neuronal wiring diagram of an adult brain. *Nature*, *634*(8032), 124–138.

https://doi.org/10.1038/s41586-024-07558-y

Dumenil, C., Yildirim, G., & Haase, A. (2025). Differential Coding of Fruit, Leaf, and Microbial

Odours in the Brains of *Drosophila suzukii* and *Drosophila melanogaster*. *Insects*, *16*(1), 84.

https://doi.org/10.3390/insects16010084

Eckstein, N., Bates, A. S., Champion, A., Du, M., Yin, Y., Schlegel, P., Lu, A. K., Rymer, T.,

Finley-May, S., Paterson, T., Parekh, R., Dorkenwald, S., Matsliah, A., Yu, S. C., McKellar, C.,

Sterling, A., Eichler, K., Costa, M., Seung, S., Murthy, M., … Funke, J. (2024). Neurotransmitter

classification from electron microscopy images at synaptic sites in Drosophila

melanogaster. *Cell*, *187*(10), 2574–2594.e23. https://doi.org/10.1016/j.cell.2024.03.016

Faucher, C., Forstreuter, M., Hilker, M., & de Bruyne, M. (2006). Behavioral responses of
Drosophila to biogenic levels of carbon dioxide depend on life-stage, sex and olfactory
context. *The Journal of experimental biology*, *209*(Pt 14), 2739–2748.
https://doi.org/10.1242/jeb.02297

Frank, D. D., Jouandet, G. C., Kearney, P. J., Macpherson, L. J., & Gallio, M. (2015).
Temperature representation in the Drosophila brain. *Nature*, *519*(7543), 358–361.
https://doi.org/10.1038/nature14284

Frechter, S., Bates, A. S., Tootoonian, S., Dolan, M. J., Manton, J., Jamasb, A. R., Kohl, J.,
Bock, D. D., & Jefferis, G. S. X. E. (2019). Functional and anatomical specificity in a higher
olfactory centre. *eLife*, 8, e44590. https://doi.org/10.7554/eLife.44590

Ghaninia, M. (2007). *Olfaction in Mosquitoes.* Doctoral Thesis, Swedish University of
Agricultural Sciences. https://pub.epsilon.slu.se/1560/.

Grabe, V., Baschwitz, A., Dweck, H. K. M., Lavista-Llanos, S., Hansson, B. S., & Sachse, S.
(2016). Elucidating the neuronal architecture of olfactory glomeruli in the Drosophila antennal
lobe. *Cell Reports*, 16(12), 3401–3413. https://doi.org/10.1016/j.celrep.2016.08.063

Hansson, B. S., & Stensmyr, M. C. (2011). Evolution of insect olfaction. *Neuron, 72*(5), 698–
711. https://doi.org/10.1016/j.neuron.2011.11.003

Hing, H., Reger, N., Snyder, J., & Fradkin, L. G. (2020). Interplay between axonal Wnt5-Vang
and dendritic Wnt5-Drl/Ryk signaling controls glomerular patterning in the Drosophila
antennal lobe. *PLoS genetics*, *16*(5), e1008767.
https://doi.org/10.1371/journal.pgen.1008767

Hong, W., & Luo, L. (2014). Genetic control of wiring specificity in the fly olfactory system.
Genetics, 196(1), 17–29. https://doi.org/10.1534/genetics.113.154336

Inagaki, H. K., Jung, Y., Hoopfer, E. D., Wong, A. M., Mishra, N., Lin, J. Y., Tsien, R. Y., &
Anderson,D.J.(2014). Optogenetic control of *Drosophila* using a red-shifted channelrhodopsin
(ReaChR). *Nature Methods, 11*(3), 325–332. https://doi.org/10.1038/nmeth.2765

Ito, K., Shinomiya, K., Ito, M., Armstrong, J. D., Boyan, G., Hartenstein, V., Harzsch, S.,

Heisenberg, M., Homberg, U., Jenett, A., Keshishian, H., Restifo, L. L., Rössler, W., Simpson,

J. H., Strausfeld, N. J., Strauss, R., Vosshall, L. B., & Insect Brain Name Working Group (2014).

A systematic nomenclature for the insect brain. *Neuron*, *81*(4), 755–765.

https://doi.org/10.1016/j.neuron.2013.12.017

Jefferis, G. S., Marin, E. C., Stocker, R. F., & Luo, L. (2001). Target neuron prespecification in

the olfactory map of Drosophila. *Nature*, *414*(6860), 204–208.

https://doi.org/10.1038/35102574

Jefferis, G. S., Vyas, R. M., Berdnik, D., Ramaekers, A., Stocker, R. F., Tanaka, N. K., Ito, K., &

Luo, L. (2004). Developmental origin of wiring specificity in the olfactory system of

Drosophila. *Development (Cambridge, England)*, *131*(1), 117–130.

https://doi.org/10.1242/dev.00896

Jones, W. D., Cayirlioglu, P., Kadow, I. G., & Vosshall, L. B. (2007). Two chemosensory

receptors together mediate carbon dioxide detection in *Drosophila*. *Nature*, 445(7123), 86–

90. https://doi.org/10.1038/nature05466

Kitamoto T. (2001). Conditional modification of behavior in Drosophila by targeted expression

of a temperature-sensitive shibire allele in defined neurons. *Journal of neurobiology*, *47*(2),

81–92. https://doi.org/10.1002/neu.1018

Kittel, R. J., Wichmann, C., Rasse, T. M., Fouquet, W., Schmidt, M., Schmid, A., Wagh, D. A.,

Pawlu, C., Kellner, R. R., Willig, K. I., Hell, S. W., Buchner, E., Heckmann, M., & Sigrist, S. J.

(2006). Bruchpilot promotes active zone assembly, Ca2+ channel clustering, and vesicle

release. *Science (New York, N.Y.)*, *312*(5776), 1051–1054.

https://doi.org/10.1126/science.1126308

Klapoetke, N. C., Murata, Y., Kim, S. S., Pulver, S. R., Birdsey-Benson, A., Cho, Y. K.,

Morimoto, T. K., Chuong, A. S., Carpenter, E. J., Tian, Z., Wang, J., Xie, Y., Yan, Z., Zhang, Y.,

Chow, B. Y., Surek, B., Melkonian, M., Jayaraman, V., Constantine-Paton, M., Wong, G. K., …

Boyden, E. S. (2014). Independent optical excitation of distinct neural populations. *Nature*

methods, *11*(3), 338–346. https://doi.org/10.1038/nmeth.2836

Komiyama, T., & Luo, L. (2006). Development of wiring specificity in the olfactory
system. *Current opinion in neurobiology*, *16*(1), 67–73.
https://doi.org/10.1016/j.conb.2005.12.002

Koyama, T., Rodrigues, M. A., Athanasiadis, A., Shingleton, A. W., & Mirth, C. K. (2021).
A nutrient-responsive hormonal circuit mediates an inter-organ growth coordination in
Drosophila.*Nature Communications, 12*, 7168.
https://doi.org/10.1038/s41467-021-25445-2

Kwon, J. Y., A. Dahanukar, L. A. Weiss, and J. R. Carlson. (2007). The Molecular Basis of CO_2
Reception in Drosophila. *Proceedings of the National Academy of Sciences USA* 104 (9): 3574–
78. https://doi.org/10.1073/pnas.0700079104.

**Lewis, L. P. C., Siju, K. P., Aso, Y., Friedrich, A. B., Bulteel, A. J. B., Rubin, G. M., & Grunwald
Kadow, I. C. (2015).** A higher brain circuit for immediate integration of conflicting sensory
information in *Drosophila*. *Current Biology, 25*(17), 2203–2214.
https://doi.org/10.1016/j.cub.2015.07.015

Li, S., Li, B., Gao, L., Wang, J., & Yan, Z. (2022). Humidity response in Drosophila olfactory
sensory neurons requires the mechanosensitive channel TMEM63. *Nature*
communications, *13*(1), 3814. https://doi.org/10.1038/s41467-022-31253-z

Lin, H. H., Chu, L. A., Fu, T. F., Dickson, B. J., & Chiang, A. S. (2013). Parallel neural pathways
mediate CO_2 avoidance responses in Drosophila. *Science (New York, N.Y.), 340*(6138), 1338–
1341. https://doi.org/10.1126/science.1236693

**Liou, N. F., Lin, S. H., Chen, Y. J., Tsai, K. T., Yang, C. J., Lin, T. Y., Wu, T. H., Lin, H. J., Chen, Y.
T., Gohl, D. M., Silies, M., & Chou, Y. H. (2018).** Diverse populations of local interneurons
integrate into the Drosophila adult olfactory circuit. *Nature communications*, *9*(1), 2232.
https://doi.org/10.1038/s41467-018-04675-x

**Macpherson, L. J., Zaharieva, E. E., Kearney, P. J., Alpert, M. H., Lin, T. Y., Turan, Z., Lee, C.
H., & Gallio, M. (2015).** Dynamic labelling of neural connections in multiple colours by trans-
synaptic fluorescence complementation. *Nature communications*, *6*, 10024.
https://doi.org/10.1038/ncomms10024

MacWilliam, D., J. Kowalewski, A. Kumar, C. Pontrello, and A. Ray. (2018). Signaling Mode of

the Broad-Spectrum Conserved CO_2 Receptor Is One of the Important Determinants of Odor

Valence in Drosophila. *Neuron* 97 (5): 1153–67.e4.

https://doi.org/10.1016/j.neuron.2018.01.028.

Marin, E. C., Büld, L., Theiss, M., Sarkissian, T., Roberts, R. J. V., Turnbull, R., Tamimi, I. F. M.,

Pleijzier, M. W., Laursen, W. J., Drummond, N., Schlegel, P., Bates, A. S., Li, F., Landgraf, M.,

Costa, M., Bock, D. D., Garrity, P. A., & Jefferis, G. S. X. E. (2020). Connectomics Analysis

Reveals First-, Second-, and Third-Order Thermosensory and Hygrosensory Neurons in the

Adult Drosophila Brain. *Current biology : CB*, *30*(16), 3167–3182.e4.

https://doi.org/10.1016/j.cub.2020.06.028

Masuyama, K., Zhang, Y., Rao, Y., & Wang, J. W. (2012). Mapping neural circuits with activity-

dependent nuclear import of a transcription factor. *Journal of neurogenetics*, *26*(1), 89–102.

https://doi.org/10.3109/01677063.2011.642910

Min, S., Ai, M., Shin, S. A., & Suh, G. S. (2013). Dedicated olfactory neurons mediating

attraction behavior to ammonia and amines in Drosophila. *Proceedings of the National*

Academy of Sciences of the United States of America, *110*(14), E1321–E1329.

https://doi.org/10.1073/pnas.1215680110

Nagel, K. I., & Wilson, R. I. (2016). Mechanisms underlying population response dynamics in

inhibitory interneurons of the *Drosophila* antennal lobe. *Journal of Neuroscience, 36*(14),

4325–4338. https://doi.org/10.1523/JNEUROSCI.3887-15.2016

Nern, A., Pfeiffer, B. D., & Rubin, G. M. (2015). Optimized tools for multicolor stochastic

labeling reveal diverse stereotyped cell arrangements in the fly visual system. *PNAS*, 112(22),

E2967–E2976. https://doi.org/10.1073/pnas.1506763112

Ni L. (2021). The Structure and Function of Ionotropic Receptors in *Drosophila*. *Frontiers in*

molecular neuroscience, *13*, 638839. https://doi.org/10.3389/fnmol.2020.638839

Nicolaï, L. J., Ramaekers, A., Raemaekers, T., Drozdzecki, A., Mauss, A. S., Yan, J., Landgraf,

M., Annaert, W., & Hassan, B. A. (2010). Genetically encoded dendritic marker sheds light on

neuronal connectivity in Drosophila. *Proceedings of the National Academy of Sciences of the*

United States of America, 107(47), 20553–20558. https://doi.org/10.1073/pnas.1010198107

Olsen, S. R., & Wilson, R. I. (2008). Lateral presynaptic inhibition mediates gain control in an

olfactory circuit. *Nature*, *452*(7190), 956–960. https://doi.org/10.1038/nature06864

Oram, T. B., & Card, G. M. (2022). Context-dependent control of behavior in

Drosophila. *Current opinion in neurobiology*, *73*, 102523.

https://doi.org/10.1016/j.conb.2022.02.003

Plaza, S. M., Clements, J., Dolafi, T., Umayam, L., Neubarth, N. N., Scheffer, L. K., & Berg, S.

(2022). *neu*Print: An open access tool for EM connectomics. *Frontiers in neuroinformatics*, *16*,

896292. https://doi.org/10.3389/fninf.2022.896292

Pulver, S. R., Pashkovski, S. L., Hornstein, N. J., Garrity, P. A., & Griffith, L. C. (2009).

Temporal dynamics of neuronal activation by Channelrhodopsin-2 and TRPA1 determine

behavioral output in *Drosophila* larvae. *Journal of Neurophysiology, 101*(6), 3075–3088.

https://doi.org/10.1152/jn.00071.2009

Root, C. M., Masuyama, K., Green, D. S., Enell, L. E., Nässel, D. R., Lee, C. H., & Wang, J. W.

(2008). A presynaptic gain control mechanism fine-tunes olfactory behavior. *Neuron*, *59*(2),

311–321. https://doi.org/10.1016/j.neuron.2008.07.003

Sachse, S., & Beshel, J. (2016). The good, the bad, and the hungry: how the central brain codes

odor valence to facilitate food approach in Drosophila. *Current opinion in neurobiology*, *40*,

53–58. https://doi.org/10.1016/j.conb.2016.06.012

Sachse, S., Rueckert, E., Keller, A., Okada, R., Tanaka, N. K., Ito, K., & Vosshall, L. B. (2007).

Activity-dependent plasticity in an olfactory circuit. *Neuron*, *56*(5), 838–850.

https://doi.org/10.1016/j.neuron.2007.10.035

Sadanandappa, M. K., Redondo, B. B., Michels, B., Rodrigues, V., Gerber, B., VijayRaghavan,

K., Buchner, E., & Ramaswami, M. (2013). Synapsin function in GABA-ergic interneurons is

required for short-term olfactory habituation. *Journal of Neuroscience, 33*(42), 16576–16585.

https://doi.org/10.1523/JNEUROSCI.3142-13.2013

Scheffer, L. K., Xu, C. S., Januszewski, M., Lu, Z., Takemura, S. Y., Hayworth, K. J., Huang, G.
B., Shinomiya, K., Maitlin-Shepard, J., Berg, S., Clements, J., Hubbard, P. M., Katz, W. T.,
Umayam, L., Zhao, T., Ackerman, D., Blakely, T., Bogovic, J., … Meinertzhagen, I. A. (2020).
A connectome and analysis of the adult Drosophila central brain. *eLife, 9*, e57443.
https://doi.org/10.7554/eLife.57443

Schlegel, P., Bates, A. S., Stürner, T., Jagannathan, S. R., Drummond, N., Hsu, J., Serratosa
Capdevila, L., Javier, A., Marin, E. C., Barth-Maron, A., Tamimi, I. F., Li, F., Rubin, G. M., Plaza,
S. M., Costa, M., & Jefferis, G. S. X. E. (2021). Information flow, cell types and stereotypy in a
full olfactory connectome. *eLife*, *10*, e66018. https://doi.org/10.7554/eLife.66018

Schlegel, P., Yin, Y., Bates, A. S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D. S., Gkantia,
M., Dos Santos, M., Munnelly, E. J., Badalamente, G., Serratosa Capdevila, L., Sane, V. A.,
Fragniere, A. M. C., Kiassat, L., Pleijzier, M. W., Stürner, T., Tamimi, I. F. M., Dunne, C. R.,
Salgarella, I., … Jefferis, G. S. X. E. (2024). Whole-brain annotation and multi-connectome cell
typing of Drosophila. *Nature*, *634*(8032), 139–152. https://doi.org/10.1038/s41586-024-
07686-5

Semelidou, O., Acevedo, S. F., & Skoulakis, E. M. (2018). Temporally specific engagement of
distinct neuronal circuits regulating olfactory habituation in *Drosophila*. *eLife*, *7*, e39569.
https://doi.org/10.7554/eLife.39569

Shearin, H. K., Quinn, C. D., Mackin, R. D., Macdonald, I. S., & Stowers, R. S. (2018). t-GRASP,
a targeted GRASP for assessing neuronal connectivity. *Journal of neuroscience methods*, *306*,
94–102. https://doi.org/10.1016/j.jneumeth.2018.05.014

Siju, K. P. (2009). *Neuromodulation in the Chemosensory System of Mosquitoes:*
Neuroanatomy and Physiology. Doctoral Thesis, Swedish University of Agricultural Sciences.
https://pub.epsilon.slu.se/1956/.

Suh, G. S., Wong, A. M., Hergarden, A. C., Wang, J. W., Simon, A. F., Benzer, S., Axel, R., &
Anderson, D. J. (2004). A single population of olfactory sensory neurons mediates an innate
avoidance behaviour in Drosophila. *Nature*, *431*(7010), 854–859.
https://doi.org/10.1038/nature02980

Tanaka, N. K., Endo, K., & Ito, K. (2012). Organization of antennal lobe-associated neurons in
adult Drosophila melanogaster brain. *The Journal of comparative neurology*, *520*(18), 4067–
4130. https://doi.org/10.1002/cne.23142

Tauxe, G. M., MacWilliam, D., Boyle, S. M., Guda, T., & Ray, A. (2013). Targeting a dual
detector of skin and CO2 to modify mosquito host seeking. *Cell*, *155*(6), 1365–1379.
https://doi.org/10.1016/j.cell.2013.11.013

Truman, James W., and Michael Bate. (1988). Spatial and Temporal Patterns of Neurogenesis
in the Central Nervous System of *Drosophila melanogaster*. *Developmental Biology* 125 (1):
145–57.

Turner, S. L., & Ray, A. (2009). Modification of CO_2 avoidance behaviour in Drosophila by
inhibitory odorants. *Nature*, *461*(7261), 277–281. https://doi.org/10.1038/nature08295

van Breugel, F., Huda, A., & Dickinson, M. H. (2018). Distinct activity-gated pathways mediate
attraction and aversion to CO_2 in Drosophila. *Nature*, *564*(7736), 420–424.
https://doi.org/10.1038/s41586-018-0732-8

Vilinsky, I., Hibbard, K. L., Johnson, B. R., & Deitcher, D. L. (2018). Probing Synaptic
Transmission and Behavior in *Drosophila* with Optogenetics: A Laboratory Exercise. *Journal of*
undergraduate neuroscience education : JUNE : a publication of FUN, Faculty for
Undergraduate Neuroscience, *16*(3), A289–A295.

Vosshall, L. B., & Stocker, R. F. (2007). Molecular architecture of smell and taste in
Drosophila. *Annual review of neuroscience*, *30*, 505–533.
https://doi.org/10.1146/annurev.neuro.30.051606.094306

Wagh, D. A., Rasse, T. M., Asan, E., Hofbauer, A., Schwenkert, I., Dürrbeck, H., Buchner, S.,
Dabauvalle, M. C., Schmidt, M., Qin, G., Wichmann, C., Kittel, R., Sigrist, S. J., & Buchner, E.
(2006). Bruchpilot, a protein with homology to ELKS/CAST, is required for structural integrity
and function of synaptic active zones in Drosophila. *Neuron*, *49*(6), 833–844.
https://doi.org/10.1016/j.neuron.2006.02.008

Wilson, R. I. (2013). Early olfactory processing in Drosophila: Mechanisms and principles.
Annual Review of Neuroscience, 36, 217–241. https://doi.org/10.1146/annurev-neuro-
062111-150533

Winding, M., Pedigo, B. D., Barnes, C. L., Patsolic, H. G., Park, Y., Kazimiers, T., Fushiki, A.,

Andrade, I. V., Khandelwal, A., Valdes-Aleman, J., Li, F., Randel, N., Barsotti, E., Correia, A.,

Fetter, R. D., Hartenstein, V., Priebe, C. E., Vogelstein, J. T., Cardona, A., & Zlatic, M. (2023).

The connectome of an insect brain. *Science (New York, N.Y.)*, *379*(6636), eadd9330.

https://doi.org/10.1126/science.add9330

Wu, Y., Helt, J. C., Wexler, E., Petrova, I. M., Noordermeer, J. N., Fradkin, L. G., & Hing, H.

(2014). Wnt5 and drl/ryk gradients pattern the Drosophila olfactory dendritic map. *The*

Journal of neuroscience : the official journal of the Society for Neuroscience, *34*(45), 14961–

14972. https://doi.org/10.1523/JNEUROSCI.2676-14.2014

Zheng, Z., Lauritzen, J. S., Perlman, E., Robinson, C. G., Nichols, M., Milkie, D., Torrens, O.,

Price, J., Fisher, C. B., Sharifi, N., Calle-Schuler, S. A., Kmecova, L., Ali, I. J., Karsh, B.,

Trautman, E. T., Bogovic, J. A., Hanslovsky, P., Jefferis, G. S. X. E., Kazhdan, M., Khairy, K., …

Bock, D. D. (2018). A Complete Electron Microscopy Volume of the Brain of Adult Drosophila

melanogaster. *Cell*, *174*(3), 730–743.e22. https://doi.org/10.1016/j.cell.2018.06.019

Figure 1. Principal neurons of the V glomerulus show distinct response properties for CO_2. (A–C) About 40 olfactory sensory neurons (ORNv) innervate the ipsilateral V glomerulus at the ventral AL surface (A). Single cell clones revealed the terminal branch pattern of ORNv axons (B). Each ORNv axon forms 3–5 non-overlapping terminal branches (C). **(D–G)** Morphology of central neurons of the V glomerulus, including 2 types of projection neurons (PNv^{uni} and PNv^{bi}, D), 2 classes of broad local interneurons (LNv1 and LNv2, E/E') and 2 classes of sparse LNs (LN23 and LN19, F, G). Gal4 expression lines for sparse interneurons LN23 and LN19 (F', G'). **(H–N)** CaLexA reporter expression in different neuron classes under ambient air (H–N) and elevated CO_2 levels (H'–N'). CaLexA activity could be observed in ORNv, LNv1/2 and LN19, but not in glomerular PNs under ambient air conditions. **(H'-N')** Following chronic CO_2 stimulation, ORNv and LN23 show high activity, comparable to the GFP expression levels at ambient air. Based on the CaLexA-GFP intensity, previous activity was higher for PNv^{bi} following stimulation as was the case for LNv1 and LNv2. PNv^{uni} showed no activity as indicated by the GFP levels. **(O)** Quantification of GFP signal intensity in ORNv, LN23, PNv^{bi}, PNv^{uni}, LNv1, and LNv2. The experimental conditions were, that the flies were either kept in ambient air or in 5% CO_2 for 3 consecutive days. For ORNs, LN23 and PNv^{bi} we also exposed the flies to spermidine, which is a strong GR21a (ORNv) inhibitor. The effect was so strong for LN23, so that we also wanted to compare it to a complete bilateral ablation of both antennae. Further, since continuous exposure to CO_2 did not produce any GFP signal for PNv^{uni}, we tried a pulsed presentation of the CO_2 stimulus, but that also showed no effect. The white bars at the lower right corners indicate the scale of 20 ym. This also applies to the following figures.

A

B

Supplementary Figure 1. (A) Schematic of the canonical olfactory pathway (left) and the identified AL relay pathway (right). **(B)** CaLexA::GFP expression in vORN cell bodies on the 3[rd] antennal segment.

Figure 2. Circuit dynamics of the V glomerulus during CO_2-induced habituation. (A) T-maze setup to determine CO_2 valence identity of adult flies. **(B)** Performance index (PI) of naïve flies compared to flies, which have experienced either chronic CO_2 exposure, removal of both antennae or a mutation of the CO_2 receptor GR63a, each of it leading to a loss of innate aversion response towards CO_2. A value of 100 for the PI represents a 100% attraction towards ambient air for a tested group of flies, while a negative value of -100 would indicate a 100% attraction towards 5% CO_2. **(C)** Temporal pattern of CO_2-induced behavioral habituation. It became evident, that even after 1 day of exposure, that aversion towards 5% CO_2 already decreased drastically. **(D, E)** Structural changes of the V glomerulus volume (D) and axonal arborization of ORNv (E) following 5 days of chronic CO_2 stimulation. **(F–Q)** Synaptic connectivity among neurons of the V glomerulus analyzed by transgenic GRASP. Activity-dependent sybGRASP indicates enhanced ORNv→PNv[bi] connectivity (F′, H), but no significant changes for ORNv→LN23 (G′, H) during behavioral habituation. Activity-independent tGRASP expression demonstrates increased synaptic connectivity of ORNv axons with PNv[bi] and LN23, consistent with morphological changes observed in ORN terminals during habituation (I′, J′, K). ORN input is restricted to the glomerulus but not seen in the extraglomerular region (arrows, J).Differences in SybGRASP levels among central neurons of the V glomerulus PNv[bi] neurons showed higher GFP intensity after prolonged exposure to elevated CO_2 levels (O, P compared to Q). The GFP levels for LN23 remained similar to the ambient air control (L-Q)

Figure 3. Neuronal polarity of CO_2 glomerular circuit components. (A-C) confocal sections illustrate *syt::GFP* and *DenMark::Cherry* expression in LN23 Quantification of GFP/cherry levels *syt::GFP* and *DenMark::Cherry* revealed distinct localization of presynaptic sites in PNv^{bi}, PNv^{uni}, and LN23. PNv^{bi} displayed prominent presynaptic labeling throughout the glomerular neuropil, while PNv^{uni} showed no detectable presynaptic signal, despite clear dendritic arborizations. LN23 exhibited scattered presynaptic puncta largely outside the glomerular boundaries. **(D)** PNv^{uni} LN23 and PNv^{bi} show different responses following habituation. While LN23 did not show increased *syt::GFP* or *DenMark::cherry* signal, both were increased for PNv^{bi} while for PNv^{uni} only *DenMark::cherry* was amplified . **(E)** *Syb::GFP* labeling further confirmed that presynaptic sites in PNv^{bi} were restricted to the glomerular neuropil and increased after chronic CO_2 exposure, whereas LN23 signals remained scattered along the glomerular margin. **(F-G)** Higher-resolution optical sections confirm the presence of *brp::GFP* puncta in LN23 dendrites outside the V glomerulus **(F, arrow)** but confined to the V glomerulus in PNv^{bi}. **(H)** Quantification of *syt::GFP* and *DenMark::cherry*. For LN23 we differentiated between intensity measurements in the center of the glomerulus and measurements near the glomerular boundary. **(I)** Hemibrain reconstructions show the enrichment of LN23 presynaptic sites (colored spheres) in extraglomerular domains compared to the intraglomerular distribution in PNv^{bi}

A

LN23::GFP

Supplementary Figure 3. (A) CaLexA labeling of LN23 combined with anti-GABA staining revealed no colocalization, indicating that LN23 is not GABAergic.

1000

1001

1002

1003

1004

1005

Figure 4. Early LN23 development. (A-E') At the beginning of pupal development (0h APF), larval LN23 neurons remodel their ventral processes (arrow in A) and start extending into the developing adult AL (arrow in B). While a broad arborization can be observed in the ventral AL (arrows in C and D), a small process extends in the dorsal AL to target posterior to the V glomerulus (arrowheads in C and D). At 40h APF, the main LN23 dendrites are restricted to the glomerular boundaries (arrow in E). **(F-G'')** In the reorganization from the larval to adult AL, LN23 processes (green) synchronize with uni-glomerular PN dendrites (red). Compared to wild-type **(H)**, loss of *Wnt5* **(I-K)** leads to a larger separation of ventral and dorsal LN23 branches (arrows in I and J), which later extend along axons towards higher brain centers (arrow in K).

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

Supplementary Figure 4. Loss of the cell adhesion molecule Flamingo in ORNv. In contrast to the (A) wild-type axonal convergence of ORNv within the glomerular boundaries, (B) *fmi*-mutant ORNv axons overshoot the ventral glomerular domain and extend into the posterior AL (arrow).

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

Figure 5. Activity of LN23 controls CO$_2$ aversion behavior. (A) Cell type–specific silencing with UAS-*shibire*^{ts} revealed that neuronal activity of all three V-glomerulus relay neurons contributes to innate CO$_2$ avoidance. Silencing of PNv^{uni} or PNv^{bi} reduced avoidance behavior, with a stronger effect in PNv^{uni}. In contrast, LN23 silencing abolished CO$_2$ aversion entirely. Control experiments performed at room temperature (rt; exp stands for experimental condition at restrictive temperatures), where *shibire*^{ts} should remain permissive, showed partial reduction of avoidance in PNv^{uni} and LN23. Since experiments with PNv^{bi} Gal4 (VT031497) showed inconclusive results, we used the alternative PNv^{bi} (alt. PNv^{bi}) Gal4 driver line (R53A05; BDSC#38859). **(B)** Cell type–specific expression of *shibire*^{ts} via CaLexA confirmed the essential role of LN23 in CO$_2$ behavior. LN23 CaLexA-*shibire*^{ts} flies showed reduced avoidance, and chronic CO$_2$ exposure, which should boost CaLexA activity, shifted behavior from avoidance to attraction. **(C)** Optogenetic activation using UAS-ReaChR demonstrated sufficiency of LN23 for driving aversive behavior. Light activation of ORNv (GR21a-Gal4) and LN23 (independent drivers LN23-Gal4 and alternative line 48408-Gal4) triggered robust avoidance, while activation of PNv^{bi} or PNv^{uni} failed to elicit significant changes. **(D)** CaLexA-driven ReaChR expression in LN23 also induced light-mediated aversion, confirming cell type specificity.

Figure 6. Two postsynaptic neurons of LN23 convey opposing odor valence. (A) Connectomic analysis (neuPrint, Janelia Research Campus) revealed two major postsynaptic partners of LN23, PNm1 and PNm17. **(B, C)** Single-cell MCFO clones of sm_PNm1 and l2_PNm17 show axonal projections bypassing the canonical olfactory lateral horn (LH) and targeting adjacent regions including the posterior lateral protocerebrum (PLP), superior clamp (SCL), and superior lateral protocerebrum (SLP). **(D)** Optogenetic activation of PNm1 induced strong aversion, whereas PNm17 activation elicited attraction, demonstrating opposing behavioral valence. **(E, F)** Silencing experiments with *shibire*[ts] confirmed that CO_2 aversion requires PNm1 activity but is unaffected by PNm17 silencing. **(G–I)** Hemibrain connectome data identified distinct downstream LH partners for the two CO_2 relay pathways: LHPD5c1 as the primary postsynaptic target of PNv[bi], LHPV9b1 downstream of PNm1, and LHCENT4 downstream of PNv[uni]. The abbreviation "rt" stands for the room temperature control of *shibire*[ts] while "exp" is the experimental condition at the restrictive temperature. The midline in the brain is indicated by the dashed line and "ml.".

A LN19 input

B LN19 output

C

Postsynaptic partner	# of connections		Presynaptic partners	# of connections
PNv^{bi}	223		ORNv	292
ORNv	201		ORN_VM7v	128
vLN24	158		VM7d_adPN	91
VM7d_adPN	140		ORN_DC4	72
PNv^{uni}	134		DM4_adPN	43

D

E

1142 **Figure 7. The "food" odor associated l2LN19.** **(A)** LN19 gets input from DM4, DC4/DP1 and VM7 while **(B)** the output is within the V
1143 glomerulus. **(C)** Table illustrating the post,- and presynaptic partners of LN19 as they can be found on https://neuprint.janelia.org/ **(D)**
Optogenetic activation of LN19 using a specific driver line induced strong behavioral aversion, indicating that LN19 activity can bias olfactory
1144 responses towards avoidance. **(E)** In contrast, silencing LN19 via shibire^{ts} did not alter innate CO_2 avoidance, suggesting that while LN19 is
sufficient to promote aversive responses, it is not essential for baseline CO_2-driven behavior. The abbreviation "rt" stands for the room
1145 temperature control of shibire^{ts} while "exp" is the experimental condition at the restrictive temperature.

Figure 8. Convergence of parallel channels for CO_2 valence coding. The activity of CO_2-responsive sensory neurons (ORNv, black) is relayed by 4 parallel PN channels to converge onto 2 central brain target neurons of opposite intrinsic valence identity. First, the glomerular PNv[bi] (red) and PNv[uni] (pink) target the ventral domain of LH[PD5] (directly or via LN[cent4], respectively) to trigger behavioral aversion (red). Second, the extraglomerular PNm1 (yellow) and PNm17 (turquoise), activated via the lateral LN23 relay (gray), converge onto LH[PV9] (directly or via CRE[108], respectively) which mediates attraction (green). While PNm17 supports LH[PV9] activity, GABAergic PNm1 blocks LH[PV9] thereby resulting in an aversive response. This core aversion domain (PNv[bi]/PNv[uni]/PNm1) can be modified either via the activation of the dorsal LH[PD5] domain by fruit odor PNs (PN[DM/VM/DC], green) or an increase of PNm17 by non-olfactory sensory neurons (turquoise). PNv[bi] is providing balancing activity between channels, for example through direct input onto LH[PV9] or via receiving LN19-mediated inhibition by food PNs.