Exploring DNA methylation profiles in the pathogenesis of human osteoporosis via whole-genome bisulfite sequencing

Yinyin Zhang[1], Guoying Wu[2*], Jialu Hou[1*], Yeling Zhong[1], Yukai Zhang[1], Shishuo Xiong[1], Zehua guo[1],Ying Li[1+]

1. The Third Clinical Medical School, Guangzhou University of Chinese Medicine, Guangzhou 510405, China
2. Guangdong Provincial Hospital of Integrated Traditional Chinese and Western Medicine, Guangzhou 528000, China

*These authors contributed equally: Author Yinyin Zhang;
+Corresponding author: Ying Li, Email: Y.Li66@outlook.com;

Abstract:

Background: Osteoporosis is a prevalent bone metabolic disorder characterized by reduced bone mass, disruption of bone microarchitecture, and increased bone fragility, leading to a heightened risk of fracture. This condition significantly impairs patients' quality of life and increases mortality risk. Emerging evidence suggests that DNA methylation may play a crucial role in regulating the expression of genes related to bone metabolism, thereby influencing the development of osteoporosis. However, the precise relationship between DNA methylation and osteoporosis remains unclear and warrants further investigation.

Results: Our study revealed significant differences in both the quantity and ratio of DNA methylation between individuals with osteoporosis and non-osteoporosis controls, with differences predominantly occurring in CpG islands. GO/KEGG enrichment analyses highlighted distinct osteoporosis-related gene pathways. Notably, we identified six genes, MSX1, HOXD4, AXIN2, WNT5A, TGFB1, and STAT3, respectively, that are potentially involved in the pathogenesis of osteoporosis and are broadly involved in various diseases and biological processes.

Conclusions: These findings indicate distinct methylation patterns between osteoporosis patients and healthy individuals, with differential methylation levels in genes associated with osteoporosis. This research offers new insights into the epigenetic mechanisms underlying osteoporosis.

Keywords: DNA methylation, Genes, Human, Osteoporosis, WGBS

Introductions

Osteoporosis is a prevalent metabolic bone disease characterized by diminished bone mass and microarchitectural deterioration, leading to enhanced bone fragility and increased fracture risk. It currently affects an estimated 200 million individuals globally, with approximately one-third of women and one-fifth of men over the age of 50 experiencing osteoporotic fractures. Driven by aging populations and lifestyle factors, its incidence continues to rise, posing a substantial public health burden due to high morbidity, mortality, and associated healthcare costs [1]. Hip fractures represent a particularly severe outcome, associated with a one-year mortality rate of 20–24% and a significant loss of independence; about one-third of survivors become fully dependent or require institutional

care within a year post-fracture [2].

The pathogenesis of osteoporosis is multifaceted, involving a complex interplay of genetic,

hormonal, and environmental factors. In recent years, epigenetic mechanisms, particularly DNA

methylation, have emerged as critical regulators in bone homeostasis and the development of

osteoporosis[3]. DNA methylation, the addition of a methyl group to the fifth carbon of cytosine

within CpG dinucleotides, is a key heritable epigenetic modification that modulates gene expression

without altering the underlying DNA sequence. This process is essential for governing fundamental

biological programs such as embryonic development, genomic imprinting, and cellular

differentiation [4].

Aberrant DNA methylation patterns are implicated in a wide array of pathologies, including cancer,

neurological disorders, and metabolic diseases like osteoporosis [5, 6]. In bone biology, altered

methylation can dysregulate the expression of genes pivotal for bone formation and resorption. For

instance, promoter hypermethylation of osteogenic genes may suppress their expression, impairing

osteoblast function, while hypomethylation of genes promoting osteoclastogenesis can exacerbate

bone loss[7-9]. Thus, DNA methylation serves as a vital layer of regulation for bone stability, and

its dysregulation appears integral to the pathogenesis of osteoporosis.

Furthermore, DNA methylation profiles hold promise as potential biomarkers for early risk

stratification, diagnosis, and prognosis. Epigenome-wide association studies (EWAS) have

identified specific differentially methylated regions (DMRs) correlated with bone mineral density

(BMD) and fracture risk [10]. Such epigenetic signatures could facilitate the identification of high-

risk individuals and inform personalized therapeutic strategies. Additionally, the reversible nature

of DNA methylation opens novel therapeutic avenues. Epigenetic drugs, such as DNA

methyltransferase inhibitors, which aim to restore normative gene expression patterns, have shown

potential in modulating bone metabolism [11].

To comprehensively map the DNA methylome, whole-genome bisulfite sequencing (WGBS) has

become a powerful high-throughput technology. WGBS combines bisulfite conversion of DNA

with next-generation sequencing to achieve single-base-pair resolution of methylation states across

the entire genome [12]. During bisulfite treatment, unmethylated cytosines are deaminated to uracil,

while methylated cytosines remain unchanged. This allows for the precise quantification of

methylation levels at CpG sites as well as in non-CpG contexts [13].

This study employed WGBS to perform a genome-wide analysis of DNA methylation in individuals

with and without osteoporosis. By systematically comparing differential methylation patterns, we

aimed to identify genes and regulatory pathways closely associated with osteoporosis, thereby

elucidating the role of epigenetic dysregulation in its etiology.

Methods

Ethical Statement

This study was conducted in accordance with the Declaration of Helsinki and was approved by the

Ethics Committee of The Third Affiliated Hospital of Guangzhou University of Chinese Medicine

(Approval No. 20240531-009, Date: June 1, 2024). The ethical approval is valid from June 1, 2024

to June 1, 2025. All participants, or their legal guardians, provided written informed consent prior

to enrollment in the study.

Study Subjects and Sample Collection

This case-control study enrolled 20 participants aged 50–75 years, including 10 patients diagnosed with osteoporosis and 10 non-osteoporotic controls. All participants provided written informed consent prior to enrollment, and the study protocol was approved by the appropriate institutional review board. Participants were categorized into two groups: the osteoporosis group (O group, n=10) and the non-osteoporosis control group (N group, n=10). Bone mineral density (BMD) was measured for all subjects using dual-energy X-ray absorptiometry (DXA; Hologic Discovery, USA). Age and body mass index (BMI) were matched between groups to ensure comparability. Bone tissue specimens were obtained during elective lumbar spinal surgery. After dissection to remove adjacent soft tissues, samples were immediately snap-frozen in liquid nitrogen and stored at –80°C until DNA extraction **(Figure 1)**.

Figure 1 Whole-genome bisulfite sequencing analysis flowchart

DNA Extraction from Bone Tissue

Bone samples were rinsed with deionized water and a mild detergent to remove surface contaminants. A sterile surgical blade was used to remove a thin outer layer, exposing the inner bone matrix. Tissues were flash-frozen in liquid nitrogen and mechanically pulverized into a fine powder. Bone powder was decalcified in 0.5 M EDTA (pH 8.0) at 4°C for 72 hours with gentle agitation. Decalcified tissue was subsequently digested in lysis buffer, and genomic DNA was isolated using a commercial DNA extraction kit according to the manufacturer's instructions. DNA

100 purity and concentration were assessed using a NanoDrop spectrophotometer and Qubit fluorometer,
101 respectively.

Library Preparation and Whole-Genome Bisulfite Sequencing

Genomic DNA was fragmented by sonication, followed by end repair, 3′ adenylation, and ligation
of methylated adapters. Adapter-ligated DNA underwent bisulfite conversion using the EZ DNA
Methylation-Gold Kit. Converted libraries were amplified by PCR, purified, and assessed for quality
using an Agilent 2100 Bioanalyzer. Quantification was performed with a Qubit® 2.0 Fluorometer.
Pooled libraries were loaded onto an Illumina NovaSeq 6000 platform for 150 bp paired-end
sequencing, following standard Illumina protocols. Sequencing metrics, including cluster density
and phasing rates, were monitored in real time **(Table 1)**.

Tabel 1 Reagents for library construction and quality inspection

Product/Reagent	Manufacturer	Catalog No.
SureSelectXT Methyl Reagent kit, HSQ,16	Agilent	G9651A
EZ-DNA Methylation-Gold Kit	Zymo	D5006
SureSelectXT Human Methyl-Seq Capture Library	Agilent	5190-4661
ExKubit dsDNA HS Analysis Kit	Excell	NGS00-3012
Agilent High Sensitivity DNA Kit	Agilent	5067-4626

Sequencing Data Quality Control

112 Raw sequencing reads were evaluated for quality using FastQC. Base calling quality was expressed
113 as Q-scores, where $Q = -10 \log_{10}(E)$ and E represents the estimated error probability. Due to
114 bisulfite conversion of unmethylated cytosines to thymines, a reduction in C-content and a
115 corresponding increase in T-content were expected and confirmed. Per-sample sequencing output
116 targeted ~20 Gb, with >90% of bases achieving $Q \geq 20$ (error rate < 1%).

Read Preprocessing and Alignment

Adapter sequences and low-quality bases were trimmed using Trim Galore (v0.4.1) with the
following criteria: removal of reads with >50% of bases having Phred score < 20, trimming of 3′-
end bases with Q < 20, and discarding reads shorter than 70 bp after trimming. Cleaned reads were
aligned to the human reference genome (GRCh37/hg19) using Bismark (v0.15.0) in conjunction
with Bowtie 2 (v2.2.9)[14]. PCR duplicates were removed using the deduplicate_bismark tool.
Mapping efficiency, coverage depth, and bisulfite conversion rates were calculated for each sample.

Methylation Calling and Differential Analysis

Cytosine methylation levels were extracted using bismark_methylation extractor[15]. Methylation ratios (mC/C) were calculated for each CpG site covered by at least 5 reads. Genome-wide methylation profiles were summarized and visualized using the R package methylKit (v0.9.5)[16, 17]. Differentially methylated regions (DMRs) were identified with the DSS package using a smoothing-based approach. Regions with $|\Delta\beta| > 0.1$ (10% methylation difference) and an adjusted p-value < 0.05 (FDR correction) were considered significant. CpG islands were annotated using the annotatr package in R based on UCSC definitions.

Functional Enrichment Analysis

Genes associated with significant DMRs were subjected to functional annotation. Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analyses were performed using the clusterProfiler package in R. A hypergeometric test was applied, with Bonferroni-corrected p-values ≤ 0.05 considered statistically significant.

Protein-Protein Interaction Network Analysis

To investigate the functional associations among the genes associated with differentially methylated regions (DMRs), a protein-protein interaction (PPI) network was constructed using the STRING database (version 12.0; https://string-db.org). The list of DMR-associated genes was submitted to STRING, and interactions were retrieved with a minimum required interaction score set to 0.400 (medium confidence). The resulting network was visualized within the STRING platform, and hub genes were identified based on their connectivity degree. This analysis aimed to reveal potential core regulatory modules and functional clusters among the epigenetically dysregulated genes in osteoporosis.

Statistical Analysis

Clinical and demographic variables were analyzed using R 4.5.2. Continuous variables conforming to normal distribution are presented as mean ± standard deviation and compared via independent two-sample t-tests. Categorical variables are expressed as counts and compared using chi-square tests. Non-normally distributed data were analyzed with nonparametric tests. Spearman's rank correlation was used to assess associations between methylation levels and clinical parameters. A two-tailed p-value < 0.05 was considered statistically significant.

Results

Baseline Characteristics of the Study Cohort

A total of 20 participants were enrolled in this case-control study and categorized into two groups: the osteoporosis group (O group, n=10) and the non-osteoporosis control group (N group, n=10).

The two groups were well-matched at baseline. The mean age was 65.20 (±4.44) years in the O

group and 63.60 (±9.37) years in the N group (P=0.85). There were no statistically significant

differences in height (157.80 ± 8.66 cm vs. 161.20 ± 10.81 cm, P=0.448), weight (60.10 ± 8.48 kg

vs. 62.30 ± 10.89 kg, P=0.62), or body mass index (23.69 ± 3.22 kg/m² vs. 23.94 ± 3.19 kg/m²,

P=0.86) between the O and N groups. The cohort was predominantly female (85%), with a

comparable sex distribution between groups (80% female in O group vs. 90% in N group,

P>0.99)**(Table 2)**. These results confirm that the two groups were comparable in terms of major

baseline characteristics, minimizing potential confounding effects in the subsequent epigenetic

analysis.

Table 2 Demographic and clinical characteristics at baseline

Variable	Overall, n = 20[1]	N group n = 10 (25%)[1]	O group N = 10 (75%)[1]	P-value[2]
Age(years)	64.40 ±7.18	63.60 ±(9.37)	65.20 ±(4.44)	0.85
Height(cm)	159.50± (9.69)	161.20 ±(10.81)	157.80 ±(8.66)	0.448
Weight(kg)	61.20± (9.57)	62.30 ±(10.89)	60.10 ±(8.48)	0.62
BMI(kg/m²)	23.81± (3.12)	23.94± (3.19)	23.69 ±(3.22)	0.86
Sex				
Female ,No. (%)	17 (85.00%)	9 (90.00%)	8 (80.00%)	>0.99
Male,No. (%)	3 (15.00%)	1 (10.00%)	2 (20.00%)	

[1]n (%); Mean±SD; Median [IQR], [2]Fisher's exact test; Two Sample t-test; Wilcoxon rank sum test

Genome-wide mapping of DNA methylation and genome coverage statistics

To comprehensively profile genome-wide DNA methylation differences in bone tissue between

osteoporotic patients and healthy controls, we conducted whole-genome bisulfite sequencing on a

cohort of 20 clinical specimens. Following stringent quality control and filtering, high-quality

sequencing data were obtained for all samples in both the osteoporosis group (O group, n=10) and

the normal control group (N group, n=10). Each sample yielded approximately 20 Gb of clean data

with 150-base pair paired-end reads. Subsequent alignment and quality assessment demonstrated

that all samples achieved a mapping efficiency exceeding 80% and a genome coverage breadth of

over 99%. The average sequencing depth across the genome was consistently maintained at ≥150-

fold for all samples, ensuring robust detection of methylation states **(Table 3)**

Tabel 3 Whole genome methylation sequencing data

Group	Clean base	Clean reads	Mapped (%)	Base coverage rate (%)	Mean depth
O	28.35±7.55	189126106±50348380	84.81±2.01	99.48±0.02	201.55±60.72
N	21.04±1.90	139598858±13666689	81.71±0.52	99.53±0.02	153.36±15.80

180 O: osteoporosis; N: non-osteoporosis; Clean base: Clean data after sequencing; Clean reads: number of reads that

181 have been filtered; Mapped (%): ratio of reads to the Clean reads on the only match; Base coverage rate (%): the

182 percentage of bases in a genome or target region that are covered by at least one sequencing read, indicating the

183 extent of the genome that has been successfully sequenced; Mean depth: the average sequencing depth of each site
184 in the captured region.

Methylation level and distribution characteristics

To compare the genome-wide distribution of DNA methylation contexts between osteoporotic and
healthy bone tissues, we analyzed the proportion of methylated cytosines in CG, CHG, and CHH
sequence contexts. In the osteoporosis group, methylation occurred predominantly in the CG
context (91.35%), with minor fractions in CHG (4.49%) and CHH (4.16%). Similarly, the normal
control group showed 91.75% CG, 4.25% CHG, and 4.00% CHH methylation **(Figure 2A)**,
confirming CG methylation as the dominant form in both groups.
We next quantified methylation levels in each sequence context using stringent bisulfite conversion
efficiency thresholds. The osteoporosis group exhibited methylation levels of $48.99 \pm 2.46\%$ (CG),
$2.41 \pm 0.08\%$ (CHG), and $2.23 \pm 0.06\%$ (CHH), while the normal group showed $46.84 \pm 1.01\%$
(CG), $2.17 \pm 0.04\%$ (CHG), and $2.04 \pm 0.06\%$ (CHH) **(Table 3)**. A statistically significant difference
between groups was observed specifically in CG methylation ($P < 0.05$).
To further characterize CG methylation patterns, we generated dual-parameter frequency
histograms assessing both methylation probability and coverage depth. Methylation probability
distribution revealed a bimodal pattern in both groups, with peaks at <10% and >90% methylation
frequencies and few sites at intermediate levels **(Figure 2B)**. Coverage analysis indicated a
progressive decrease in the proportion of CpG sites with increasing sequencing depth **(Figure 2C)**.
Comparative assessment demonstrated that the osteoporosis group consistently displayed higher
methylation probabilities and greater per-site coverage depths than the normal controls across
individually mapped cytosine positions.

Figure 2 The average ratio and distribution characteristics of group O and Group N DNA methylation types. **(A)** Methylation ratio in different backgrounds. O group, osteoporosis group. N group, normal group. The blue, orange and gray colors represent mCG, mCHG and mCHH, respectively. **(B)** Average percentage of single base methylation probability per group. The horizontal coordinate indicates the probability of methylation, and the vertical coordinate indicates the number of CpG sites for this degree of methylation. The probability of methylation is determined by the amount of C measured at each site/the amount of (C+T) measured at that. **(C)** The coverage depth of methylation sites was detected in each CpG background. The abscissa represents the coverage of each base, and the ordinate represents the proportion of CpG under that coverage.

Tabel 4 Average methylation levels in different backgrounds

Group	MeanCG(%)	MeanCHG(%)	MeanCHH(%)
O	48.99 ± 2.46*	2.41 ± 0.08	2.23 ± 0.06
N	46.84 ± 1.01	2.17 ± 0.04	2.04 ± 0.06

Data represent means ± SD (n = 10 samples replicates for O and N group), *refers as significant difference ($P < 0.05$).MeanCG(%): average methylation levels in CG regions, mCG counts/ (CG counts on the genome)*100%; MeanCHG(%): average methylation levels in CHG region, mCHG counts/ (CHG counts on the genome)*100%; MeanCHH(%): average methylation levels in CHH region, mCHH counts/ (CHH counts on the genome)

Differential DNA methylation regions distribution and characteristics

Volcano plot analysis of DMRs revealed 996 regions with significant methylation alterations (795
hypermethylated and 201 hypomethylated) out of 2,484 analyzed regions **(Figure 3A)**,
corresponding to a hyper- to hypomethylation ratio of 3.96:1, demonstrating a genome-wide shift
toward hypermethylation in osteoporotic bone.
Genomic annotation of DMRs indicated their predominant enrichment within CpG islands (CGIs)
(1,145 regions, 46.4%), with decreasing density in CGI shores (496 regions, 20.1%) and shelves
(407 regions, 16.4%) **(Figure 3B)**. Functional compartment analysis further showed that DMRs
were significantly enriched in intronic regions (33.7%) and promoter/enhancer elements (24.9%),
while being minimally represented in 5′ UTR (2.1%) and 3′ UTR regions (1.7%) **(Figure 3C)**.
Distance-frequency distribution analysis relative to CGIs confirmed an inverse relationship
between DMR abundance and genomic distance from CGIs **(Figure 3D)**. To characterize DMR
architecture, we analyzed the distribution of region lengths. Length-frequency histograms revealed
a strong negative correlation between DMR length and abundance, with over 90% of DMRs
spanning less than 1,000 bp **(Figure 3E)**.

Figure 3 Differential DNA methylation regions distribution and characteristics. **(A)** Distribution of differential

methylation regions in CpG islands and different components of CPG islands. Island indicates that DMRs overlaps CGI; Shore indicates that the distance between DMRs and CGI is between 1-2kb. Shelf indicates a distance between 2 and 4kb. OpenSea indicates that the distance between the two is greater than 4kb. **(B)** Distribution of differential methylation regions on different gene elements. **(C)** The frequency of the difference between methyl region segments and CGI relative positions. The horizontal coordinate refers to the upstream and downstream position of DMRs relative to CGI, and the vertical coordinate refers to the frequency of differential methylation segments at specific locations. **(D)** Length distribution of differential methylation regions. The horizontal coordinate refers to the length of DMRs, and the total left side refers to the frequency of DMRs for that length.

GO and KEGG enrichment

To elucidate the biological functions of the DMRs identified between osteoporotic patients and
healthy individuals, pathway enrichment analyses were conducted based on the GO and KEGG
databases. In addition, PPI network analysis was performed using the STRING database. A total of
996 significant DMRs were analyzed, and the top 10 regions showing the most pronounced
differences, along with osteoporosis-related GO terms and KEGG pathways, are presented.
In the GO enrichment analysis, DMRs were significantly enriched in biological processes such as
gland development, epithelial cell proliferation, embryonic organ development, and embryonic
organ morphogenesis **(Figure 4A)**. KEGG pathway analysis revealed significant enrichment in
pathways including non-small cell lung cancer, chronic myeloid leukemia, and parathyroid hormone
synthesis **(Figure 4B)**.
GO entries containing keywords associated with bone biology—such as "bone," "osteoclast,"
"osteoblast," "ossification," "mineral," "cartilage," "chondrocyte," "skeletal," "calcium," "Wnt,"
"BMP," "TGF," "collagen," "extracellular matrix," and "ECM"—were screened. This search
identified five GO terms related to osteoporosis: "embryonic skeletal system
development","embryonic skeletal system morphogenesis," "regulation of extracellular matrix
organization," "positive regulation of BMP signaling pathway," and "positive regulation of
extracellular matrix organization" **(Figure 4C)**. Notably, no KEGG pathways directly associated
with osteoporosis were identified.

B

C

268
269

Figure 4. Functional enrichment analysis of DMRs between osteoporotic patients and healthy controls. **(A)** GO enrichment analysis of biological processes. The bubble chart displays significantly enriched GO terms, with the bubble size representing the number of genes or DMRs mapped, and the color intensity indicating the statistical significance. Representative terms include gland development and epithelial cell proliferation. **(B)** KEGG pathway enrichment analysis. The bubble chart illustrates significantly enriched KEGG pathways, where the size and color of the bubbles denote the gene count and the level of enrichment significance, respectively. Key pathways identified include non-small cell lung cancer and chronic myeloid leukemia. **(C)** Screening of osteoporosis-related GO terms via keyword filtering. This bubble chart presents the five GO terms identified as relevant to bone metabolism and extracellular matrix organization, with the bubble size and color similarly reflecting gene counts and statistical significance.

Differential gene screening and validation in public databases

Through analysis of osteoporosis-related pathways, we identified 19 genes and constructed a PPI network for these genes. Genes with node degrees ≥3 were preliminarily screened, resulting in 11 candidate genes: MSX1, HOXD4, AXIN2, WNT5A, TGFB1, STAT3, RUNX2, NOTCH1, HOXD9, MSX2, BMP7, and HOXD3**(Figure 5)**.

Subsequently, we selected the dataset GSE230664 from the GEO database, which is also related to osteoporosis research, to validate the expression of these 11 genes. The results showed that six genes—MSX1, HOXD4, AXIN2, WNT5A, TGFB1, and STAT3—exhibited expression trends consistent with the methylation data. However, none of the genes showed statistically significant differences in expression between the two groups**(Table 5)**.

Figure 5 Construction of differential methylation gene networks related to bone development. Analysis of the interaction between DMRs related to muscle development using STRING software according to the interplay index (confidence > 0.4). The interplay index between genes was represented by edge width and transparency. Dark and wide edges indicated high confidence.

Table 5. Validation of differentially expressed genes in the GEO database.

Gene	Methylation Change (Δβ)	Expression Change (logFC)	Expression adj.P.Val	Methylation-Expression Concordance
MSX1	+0.155	-0.750	0.348	Concordant (Hypermethylation/Downregulation)
HOXD4	+0.095	-0.804	0.080	Concordant (Hypermethylation/Downregulation)
AXIN2	+0.112	-0.282	0.663	Concordant (Hypermethylation/Downregulation)
WNT5A	+0.099	-0.121	0.973	Concordant (Hypermethylation/Downregulation)
TGFB1	-0.129	+1.109	0.330	Concordant (Hypomethylation/Upregulation)
STAT3	-0.112	+1.316	0.436	Concordant (Hypomethylation/Upregulation)
RUNX2	+0.114	+0.860	0.402	Discordant
NOTCH1	+0.116	+0.607	0.424	Discordant
HOXD9	+0.129	+0.581	0.653	Discordant
MSX2	+0.101	+0.867	0.227	Discordant
BMP7	+0.112	+0.048	0.963	Discordant
HOXD3	+0.101	+0.299	0.666	Discordant

Δβ: Average methylation difference (O group-N group), positive value indicates hypermethylation. logFC: log2 fold

change (O group/N group), negative value indicates downregulation.

Discussion

The concept of epigenetics, introduced by the British biologist Conrad Hal Waddington in 1942[18],
describes how interactions between genes and their products influence cellular development without
altering the DNA sequence itself. Epigenetic modifications affect gene activity by altering DNA
and chromatin chemical modifications, primarily through DNA methylation, histone modification,
noncoding RNA regulation, and chromatin remodeling. As research has advanced, epigenetics has
been implicated in the onset and progression of various diseases, including cancer, neurological
disorders, and metabolic diseases.
DNA methylation is a key epigenetic modification in which a methyl group is added to the 5th
carbon atom of cytosine within DNA, forming 5-methylcytosine. This process is catalyzed by DNA
methyltransferases (DNMT), primarily at CpG dinucleotides[19]. DNMT1 maintains methylation
patterns during DNA replication, transferring methylation marks from the parent strand to the
daughter strand. DNMT3A and DNMT3B establish new methylation patterns during development
and differentiation, whereas DNMT3L facilitates DNMT3A and DNMT3B but lacks catalytic
activity itself[20]. DNA methylation is generally associated with gene silencing, particularly when
promoter regions are highly methylated[21], which suppresses transcription. During development,
specific methylation patterns change in response to gene expression needs, and methylation helps
suppress transposons and repetitive sequences, maintaining genomic stability.
Osteoporosis development is characterized by an imbalance in bone remodeling, with decreased
bone formation and increased bone resorption. Factors such as genetics, the environment, hormonal
changes, nutrition, and physical activity influence this balance[22]. DNA methylation also plays a
critical role in osteoporosis by regulating the expression of genes related to bone metabolism. The
methylation status of specific genes can alter osteoblast and osteoclast function, impacting
osteoporosis development. For instance[23], hypermethylation of the RUNX2 gene, which is

associated with osteoblast differentiation, can inhibit osteoblast function and reduce bone formation.
Conversely, hypomethylation of the RANKL gene, which is linked to osteoclast differentiation, can
increase osteoclast activity and increase bone resorption.
Our results suggest that DNA methylation is involved in various biological processes, cellular
functions, and disease progression. Although this study cannot directly establish a causal
relationship between DNA methylation and osteoporosis, evidence points to a significant
association between the two. We observed differences in the methylation levels and expression of
methylation-related genes between the osteoporosis and non-osteoporosis groups. Enrichment
analysis further revealed differential methylation of several genes associated with osteoporosis,
indicating that altered methylation levels of certain genes may be associated with this disease.
Through the integrative analysis of whole-genome methylation profiles from bone tissue and an
independent transcriptomic dataset (GSE230664), this study identified several potential
epigenetically regulated targets, such as MSX1, AXIN2, and TGFB1, which exhibited consistent
directions of change in both methylation and expression. However, we also noted that some genes
displaying significant differential methylation in our cohort did not show statistically significant
expression differences in the public dataset. This incomplete correspondence between epigenetic
alterations and transcriptional output is a common phenomenon in complex biological systems and
may be attributed to the following interrelated factors.
First, the function of DNA methylation, particularly at regions outside promoter CpG islands—such
as enhancers, gene bodies, or silencers—is highly dependent on cell type, differentiation stage, and
specific microenvironmental signals[24]. The bone tissue samples used in this study contain
heterogeneous cell populations, including osteoblast progenitors, mature osteoblasts, and osteocytes.
A key methylation alteration in a specific progenitor subpopulation may have its corresponding
expression signal diluted in bulk RNA analysis. For instance, RUNX2, the master regulator of
osteogenic differentiation, exhibits stringent stage-specific expression. The methylation changes
observed in its promoter region may significantly affect its transcriptional activity only during
specific differentiation windows or within particular cell subsets, which is difficult to capture in
bulk sequencing data.
Second, gene expression is co-regulated by a multi-layered network, including transcription factors,
histone modifications, and non-coding RNAs. DNA methylation represents one layer within this
network, and its effects may be compensated for or buffered by other, stronger regulatory layers.
Methylation in a gene's promoter region could be counteracted by a potent enhancer or specific
activating histone modifications, thereby maintaining expression stability. Conversely, even if the
methylation status remains unchanged, alterations in other regulatory layers may drive expression
changes[25]. This network robustness explains why a single methylation change is sometimes
insufficient to produce a detectable expression phenotype.
MSX1 and HOXD4 both encode homeobox transcription factors that are crucial for skeletal
patterning and osteoblast commitment[26]. Their hypermethylation and reduced expression suggest
an early impairment in osteogenic lineage determination. This epigenetic silencing may lead to a
diminished osteoprogenitor cell pool or reduced differentiation capacity, a hallmark of age-related
bone loss.
Concurrently, the epigenetic repression of AXIN2 and WNT5A points to a coordinated disruption
of Wnt signaling[27, 28]. Silencing of AXIN2 could theoretically result in sustained activation of
canonical Wnt/β-catenin signaling; however, its concurrent downregulation with WNT5A may

reflect a more complex, tissue-specific rewiring of the Wnt network that favors bone resorption over
formation. This apparent paradox warrants further dissection in future functional studies.
In contrast, the observed hypomethylation and increased expression of TGFB1 align with its
complex, stage-dependent role in bone metabolism. Although TGFB1 is essential for early bone
formation, its sustained overexpression in the bone microenvironment is a well-recognized driver
of abnormal bone remodeling. It promotes the recruitment of osteoclast precursors and stimulates
bone resorption, while ultimately leading to fibrous tissue deposition and impaired bone quality[29].
Thus, its epigenetic activation may be a key mechanism driving the high-turnover imbalance
observed in some osteoporotic patients.
Similarly, the epigenetic upregulation of STAT3 links epigenetic alteration to inflammatory bone
loss. Persistent STAT3 activation can promote osteoclast differentiation and survival while
inhibiting osteoblast function, creating a vicious cycle that exacerbates net bone loss[30]. This
positions STAT3 as a potential epigenetic mediator connecting chronic low-grade inflammation—
often associated with aging—to the pathogenesis of osteoporosis.
Importantly, these genes do not function in isolation but act as nodes within interconnected
pathways. The silencing of osteogenic transcription factors and specific Wnt components, coupled
with the activation of catabolic signals, paints a coherent picture of a biased epigenetic program that
disrupts bone homeostasis[31]. This network-level dysregulation suggests that targeting a single
gene may be insufficient. Instead, the identified epigenetic marks themselves—particularly the
reversible promoter hypermethylation of genes such as MSX1 and AXIN2—hold potential as novel
therapeutic targets for demethylating agents or as diagnostic biomarkers for patient stratification[32].
While this integrative study provides novel insights into the epigenetic landscape of osteoporosis,
several limitations must be acknowledged. First, the sample size of our discovery cohort, although
utilizing precious bone biopsy specimens, remains modest, which may constrain the statistical
power to detect more subtle methylation alterations. Second, the correlative nature of our bulk tissue
analysis precludes the establishment of direct causality between the observed DNA methylation
changes and downstream gene expression shifts or the osteoporotic phenotype. Third, the
observational nature of our study, without direct experimental perturbation, limits causal inference.
The functional impact of the prioritized epigenetic alterations on osteoblast or osteoclast activity
warrants future validation using targeted in vitro or in vivo models. Finally, the clinical
translatability of our findings necessitates validation in larger, independent, and well-characterized
longitudinal cohorts to rigorously assess the prognostic value and generalizability of the identified
epigenetic signatures.
Future research should extend these findings along several critical directions. For instance, applying
single-cell multi-omics technologies to bone samples would enable the direct mapping of
methylation and expression changes to specific cell types, thereby clarifying the precise cellular
origins and impacts of the observed epigenetic dysregulation[33]. Subsequently, direct functional
testing of our prioritized candidate genes, particularly MSX1 and AXIN2, is essential. Employing
CRISPR-dCas9-based epigenetic editing tools to specifically reverse promoter hypermethylation at
these loci in relevant bone cell models would be crucial for conclusively establishing their causal
role in driving osteogenic dysfunction[34].
Nevertheless, our work delineates a coherent set of interconnected, epigenetically dysregulated
genes and pathways. Further investigation along the proposed directions will not only solidify the
pathophysiological understanding of epigenetic contributions to osteoporosis but will also facilitate

the development of novel epigenetic diagnostics and targeted therapeutic strategies.

Conclusion

This study employed whole-genome bisulfite sequencing to profile genome-wide DNA methylation
differences between patients with osteoporosis and non-osteoporosis controls. The osteoporotic
group exhibited a higher global methylation burden and a greater number of differentially
methylated genes compared to the control group. Through integrated enrichment analysis and
validation using public databases, we prioritized six genes—MSX1, HOXD4, AXIN2, WNT5A,
TGFB1, and STAT3—as potential epigenetic drivers of osteoporosis, likely acting through distinct
biological pathways. We acknowledge that the limited sample size and the multiple-gene screening
approach may introduce bias in identifying robust differential methylation signatures; thus, these
findings require further validation. Functional experiments are also necessary to test the mechanistic
hypotheses generated from these methylome data. Nonetheless, this work contributes to the growing
understanding of epigenetic dysregulation in osteoporosis and may inform the development of novel
diagnostic and therapeutic strategies.

Abbreviations

EWAS: Epigenome-wide association studies; DMRs: Differentially methylated regions; WGBS:
Whole-genome bisulfite sequencing; BMD: Bone mineral density; BMI: Body mass index; CGI:
CpG island; GO: Kyoto Encyclopedia of Genes and Genomes; KEGG: Kyoto Encyclopedia of
Genes and Genomes; PPI: Protein-Protein Interaction ; FDR: False discovery rate; DNMT: DNA
Methyltransferase; RUNX2: Runt-related transcription factor 2; MSX1: Msh Homeobox 1; HOXD4:
Homeobox D4; AXIN2: Axin 2; WNT5A: Wnt Family Member 5A; TGFB1: Transforming Growth
Factor Beta 1; STAT3: Signal Transducer And Activator Of Transcription 3.

Data availability

436 Sequence data that support the findings of this study have been deposited in the NCBI with the
437 primary accession code PRJNA1154659

Acknowledgements

439 We sincerely thank all members of Ying Li's group for their contributions to sample collection and
440 technical assistance. All images were created by the author team.

Authors' contributions

442 YYZ and GYW conducted the experiments and data analysis. JLH designed and supervised the
443 experiment and assisted with manuscript revisions. YLZ provided technical support. YKZ and SSX
444 coordinated the research. ZHG and YL performed the surgeries and provided the samples. YYZ
445 drafted the manuscript. All authors read and approved the final version of the manuscript.

Funding

447 This work was financially supported by the National Natural Science Foundation of China (NSFC-

82374482). The funding bodies had no role in the design of the study, the collection, analysis, and interpretation of data, or in writing the manuscript.

Ethics approval

All experimental procedures, including human care and tissue sample collection, were approved by and conducted in accordance with the guidelines of the Medical Ethics Committee of the Third Affiliated Hospital of Guangzhou University of Chinese Medicine, China (Approval ID: PJ-XS-20240531-009).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

1. Johnston CB, Dagar M: **Osteoporosis in Older Adults**. *The Medical clinics of North America* 2020, **104**(5):873-884.

2. Ebeling PR, Nguyen HH, Aleksova J, Vincent AJ, Wong P, Milat F: **Secondary Osteoporosis**. *Endocr Rev* 2022, **43**(2):240-313.

3. Xu Z, Yu Z, Chen M, Zhang M, Chen R, Yu H, Lin Y, Wang D, Li S, Huang L *et al*: **Mechanisms of estrogen deficiency-induced osteoporosis based on transcriptome and DNA methylation**. *Frontiers in cell and developmental biology* 2022, **10**:1011725.

4. Moore LD, Le T, Fan G: **DNA methylation and its basic function**. *Neuropsychopharmacology* 2013, **38**(1):23-38.

5. Hannon E, Dempster EL, Mansell G, Burrage J, Bass N, Bohlken MM, Corvin A, Curtis CJ, Dempster D, Di Forti M *et al*: **DNA methylation meta-analysis reveals cellular alterations in psychosis and markers of treatment-resistant schizophrenia**. *eLife* 2021, **10**.

6. Liang X, Aouizerat BE, So-Armah K, Cohen MH, Marconi VC, Xu K, Justice AC: **DNA methylation-based telomere length is associated with HIV infection, physical frailty, cancer, and all-cause mortality**. *Aging cell* 2024, **23**(7):e14174.

7. Visconti VV, Cariati I, Fittipaldi S, Iundusi R, Gasbarra E, Tarantino U, Botta A: **DNA Methylation Signatures of Bone Metabolism in Osteoporosis and Osteoarthritis Aging-Related Diseases: An Updated Review**. *International journal of molecular sciences* 2021, **22**(8).

8. Xu F, Li W, Yang X, Na L, Chen L, Liu G: **The Roles of Epigenetics Regulation in Bone Metabolism and Osteoporosis**. *Frontiers in cell and developmental biology* 2020, **8**:619301.

9. Yang S, Duan X: **Epigenetics, Bone Remodeling and Osteoporosis**. *Curr Stem Cell Res Ther* 2016.

10. Wen B, Zhang Y, He J, Tan L, Xiao G, Wang Z, Cui W, Wu B, Wang X, He L *et al*: **Causal impact of DNA methylation on refracture in elderly individuals with osteoporosis - a prospective cohort study**. *BMC Musculoskelet Disord* 2024, **25**(1):432.

11.   Li B, Zhao J, Ma JX, Li GM, Zhang Y, Xing GS, Liu J, Ma XL: **Overexpression of DNMT1**
leads to hypermethylation of H19 promoter and inhibition of Erk signaling pathway in
disuse osteoporosis. *Bone* 2018, **111**:82-91.

12.   Cilia C, Friggieri D, Vassallo J, Xuereb-Anastasi A, Formosa MM: **Whole Genome**
Sequencing Unravels New Genetic Determinants of Early-Onset Familial Osteoporosis
and Low BMD in Malta. *Genes (Basel)* 2022, **13**(2).

13.   Bagger FO, Borgwardt L, Jespersen AS, Hansen AR, Bertelsen B, Kodama M, Nielsen FC:
Whole genome sequencing in clinical practice. *BMC Med Genomics* 2024, **17**(1):39.

14.   Krueger F, Andrews SR: **Bismark: a flexible aligner and methylation caller for Bisulfite-**
Seq applications. *Bioinformatics* 2011, **27**(11):1571-1572.

15.   Ryan DP, Ehninger D: **Bison: bisulfite alignment on nodes of a cluster**. *BMC Bioinformatics*
2014, **15**(1):337.

16.   Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, Mason CE:
methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation
profiles. *Genome Biol* 2012, **13**(10):R87.

17.   Park Y, Wu H: **Differential methylation analysis for BS-seq data under general**
experimental design. *Bioinformatics* 2016, **32**(10):1446-1453.

18.   Mattei AL, Bailly N, Meissner A: **DNA methylation: a historical perspective**. *Trends Genet*
2022, **38**(7):676-707.

19.   Jones PA: **Functions of DNA methylation: islands, start sites, gene bodies and beyond**.
Nature reviews Genetics 2012, **13**(7):484-492.

20.   Prasad R, Yen TJ, Bellacosa A: **Active DNA demethylation-The epigenetic gatekeeper of**
development, immunity, and cancer. *Advanced genetics (Hoboken, NJ)* 2021, **2**(1):e10033.

21.   Wang D, Wu W, Callen E, Pavani R, Zolnerowich N, Kodali S, Zong D, Wong N, Noriega S,
Nathan WJ *et al*: **Active DNA demethylation promotes cell fate specification and the DNA**
damage response. *Science (New York, NY)* 2022, **378**(6623):983-989.

22.   Wang P, Cao Y, Zhan D, Wang D, Wang B, Liu Y, Li G, He W, Wang H, Xu L: **Influence of**
DNA methylation on the expression of OPG/RANKL in primary osteoporosis.
International journal of medical sciences 2018, **15**(13):1480-1485.

23.   Galbraith K, Snuderl M: **DNA methylation as a diagnostic tool**. *Acta neuropathologica*
communications 2022, **10**(1):71.

24.   Greenberg MVC, Bourc'his D: **The diverse roles of DNA methylation in mammalian**
development and disease. *Nat Rev Mol Cell Biol* 2019, **20**(10):590-607.

25.   Dhar GA, Saha S, Mitra P, Nag Chaudhuri R: **DNA methylation and regulation of gene**
expression: Guardian of our health. *Nucleus (Calcutta)* 2021, **64**(3):259-270.

26.   Ishii M, Han J, Yen HY, Sucov HM, Chai Y, Maxson RE, Jr.: **Combined deficiencies of Msx1**
and Msx2 cause impaired patterning and survival of the cranial neural crest. *Development*
2005, **132**(22):4937-4950.

27.   Suthon S, Perkins RS, Bryja V, Miranda-Carboni GA, Krum SA: **WNT5B in Physiology and**
Disease. *Frontiers in cell and developmental biology* 2021, **9**:667581.

28.   Velasco J, Zarrabeitia MT, Prieto JR, Perez-Castrillon JL, Perez-Aguilar MD, Perez-Nuñez MI,
Sañudo C, Hernandez-Elena J, Calvo I, Ortiz F *et al*: **Wnt pathway genes in osteoporosis and**
osteoarthritis: differential expression and genetic association study. *Osteoporos Int* 2010,
21(1):109-118.

29.   Langdahl BL, Carstens M, Stenkjaer L, Eriksen EF: **Polymorphisms in the transforming**
growth factor beta 1 gene and osteoporosis. *Bone* 2003, **32**(3):297-310.
30.   Hou X, Tian F: **STAT3-mediated osteogenesis and osteoclastogenesis in osteoporosis**. *Cell*
Commun Signal 2022, **20**(1):112.
31.   Gao Y, Chen N, Fu Z, Zhang Q: **Progress of Wnt Signaling Pathway in Osteoporosis**.
Biomolecules 2023, **13**(3).
32.   Mues G, Kapadia H, Wang Y, D'Souza RN: **Genetics and human malformations**. *J Craniofac*
Surg 2009, **20 Suppl 2**(Suppl 2):1652-1654.
33.   Feng S, Li J, Tian J, Lu S, Zhao Y: **Application of Single-Cell and Spatial Omics in**
Musculoskeletal Disorder Research. *International journal of molecular sciences* 2023, **24**(3).
34.   Cai R, Lv R, Shi X, Yang G, Jin J: **CRISPR/dCas9 Tools: Epigenetic Mechanism and**
Application in Gene Transcriptional Regulation. *International journal of molecular sciences*
2023, **24**(19).
544

Top 10 KEGG Enrichment Pathways

Osteoporosis-Related GO Enrichment Terms (n = 5)

DMRs Genetic Annotation

DMRs CGI Annotation

Top 10 GO Enrichment Terms