**Title:**

Enhanced tRNA array method version 2 for simultaneous in vitro synthesis of 21 tRNAs

**Authors and affiliations**

Ryota Miyachi[a], Anna Irie[a], and Norikazu Ichihashi[a,b, c]*

[a] Department of Life Science, Graduate School of Arts and Science, The University of Tokyo, 3-8-

1 Komaba, Meguro-ku, Tokyo 153-8902, Japan

[b] Komaba Institute for Science, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-

8902, Japan

[c] Research Center for Complex Systems Biology, Universal Biology Institute, The University of

Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan

**Correspondence:**

Norikazu Ichihashi

Department of Life Science, Graduate School of Arts and Science, The University of Tokyo, 3-8-1

Komaba, Meguro-ku, Tokyo 153-8902, Japan

Tel: 81-3-5465-7307; Fax: 81-3-5465-7307; Email: ichihashi@bio.c.tokyo-u.ac.jp

**Keywords**

tRNA; cell-free synthetic biology; in vitro transcription; PURE system

**Abstract**

Transfer RNAs (tRNAs) play an essential role in translation, and their simultaneous *in vitro* synthesis remains a key challenge in bottom-up synthetic biology. We previously developed the tRNA array method that enables the simultaneous *in vitro* synthesis of 21 tRNAs from a single DNA template; however, the translational activity was substantially lower than that achieved using individually prepared 21 tRNAs for some proteins. Here, we identify the tRNA groups (PIEN group) that limit translation in the tRNA array method and improve the translational activities through sequence modification and incorporation of a leader sequence into the array construct. The resulting tRNA array method version 2 produces a tRNA set that allows translation at levels similar to those achieved with individually prepared tRNAs for multiple reporter proteins under both translation-coupled and uncoupled conditions. This tRNA synthesis scheme provides an improved platform for constructing self-reproducible gene expression systems.

**Introduction**

Living systems possess the ability to reproduce the molecular components that constitute themselves. In vitro reconstitution of a minimal molecular system that exhibits this property is one of the central challenges in bottom-up synthetic biology(1–9). In particular, the construction of a self-reproductive central dogma system is an essential requirement for sustaining life-like systems. Although many components, including translation proteins(10–18), ribosomal components(19–21), transfer RNAs (tRNA)(22–24), and DNA replication machinery(25–31), have been successfully synthesized in an *E. coli*-based reconstituted gene expression system (PURE system)(32) for self-regeneration, the efficiency of these syntheses remains a major challenge. Among these components, tRNA poses a large obstacle(33), as its synthesis requires multiple processing steps that have not been fully reconstituted *in vitro*.

In recent years, the use of *in vitro*-synthesized tRNAs in reconstituted gene expression systems has advanced substantially, enabling a wide range of applications, including genetic code expansion or engineering(33–36). In addition, two research groups, including ours, have demonstrated that functional tRNAs can be expressed within the PURE system and coupled directly with translation reactions(22–24). Li *et al.* achieved sustained translation under a microfluidic chemostat system by expressing 21 tRNAs using nicked DNA templates(24). We also succeeded in the simultaneous expression of 21 tRNAs from a single polycistronic DNA template in the PURE system by combining the self-cleaving HDV ribozyme (HDVR)(37) and RNase P digestion, which we named the tRNA array method(23). While this method provided a useful means for both genetic codon engineering and constructing a self-reproducible translation system, there remains a shortcoming: the translation level with tRNA prepared by this method was significantly lower than that with individually prepared tRNAs for some proteins.

In this study, we aimed to improve the tRNA array method to achieve translation efficiencies comparable to those obtained with individually prepared tRNAs for multiple proteins. To this end,

we first identified the tRNA groups that limited translation using the original tRNA array method. We then improved the activities and expression levels of the insufficient tRNAs by introducing mutations to stabilize the cloverleaf structure and introducing leader sequences to enhance processing efficiency. By integrating these approaches, we succeeded in enhancing the expression levels of at least three proteins to levels comparable to those obtained with individually prepared tRNA mixtures under both translation-uncoupled and -coupled conditions.

**Materials and Methods**

**DNA preparation**

DNA templates for preparation of tRNA variants were generated by PCR using forward oligonucleotides containing a T7 promoter and the corresponding tRNA sequences, together with 2'-O-methyl−modified reverse primers (Table S1). The combinations of tRNA variant sequences and primers used in this study are listed in Table S2. DNA fragment encoding L_PIEN_v2 was obtained using Twist Bioscience's artificial gene synthesis service. Replacement of the original PIEN region in the pTwist_21tRNAs plasmid with the improved PIEN sequences was performed by amplifying the corresponding fragments using specific primers (primers 20 and 21 for the insert fragment and primers 22 and 23 for the vector fragment) and assembling them using In-Fusion cloning kit (Takara). The resulting DNA sequences are listed in Table S3. A linear DNA template encoding all 21 tRNAs (a single 21 tRNA template) was amplified by PCR using primers 24 and 21 for use in expression-coupled translation experiments.

The luciferase template DNA was prepared by PCR amplification using primers 25 and 26 (Table S1) and a plasmid (pUC-T7p-Fluc_21tRNA) constructed in our previous study(22), followed by column purification. DNA fragments encoding sfGFP and GUS were obtained using an artificial gene synthesis service (Twist Bioscience), as described previously(23), and used for DNA templates used for translation experiments in the PURE system after PCR amplification using the same primer pair (primers 25 and 26) and column purification. The sequences of all reporter gene templates are listed in Table S3. All DNA concentrations were quantified based on the $A_{260}$.

**Individual tRNA and M1 RNA preparation**

The sequences of the 21 tRNAs used in this study were based on our previous work(22,23) and are listed in Table S4. Individually-prepared IVS tRNAs were synthesized essentially as described previously. Briefly, among the 21 tRNAs, 15 species initiating with guanosine at the 5′ end were individually prepared by *in vitro* transcription whereas the remaining six non-G-start tRNAs were chemically synthesized (Invitrogen), as in our earlier studies. The DNA template for M1 RNA was

prepared by PCR using a plasmid encoding M1 RNA as described previously and primers 27 and 28
(Table S3).

*In vitro* transcription was performed in a mixture containing 40 mM Tris-HCl pH 8.0, 10 mM
magnesium chloride, 2 mM spermidine, 5 mM DTT, 1 U/µl T7 RNA polymerase (Takara), 2 mM
each NTPs(ATP, GTP, CTP, and UTP), 3 mM GMP, 5 µg/µl template DNA, 2 U/µl inorganic
pyrophosphatase (New England Biolab), 0.8 U/µl RNasin (Promega). The mixture was incubated at
37°C for 12 h, and the RNA product was individually purified using the PureLink RNA Mini Kit
(Invitrogen). Both tRNAs and M1 RNA were dissolved in water and stored at -80 °C until use. RNA
concentrations were determined based on the A260.

**Simultaneous 21 tRNAs preparation using the tRNA array method**

Simultaneous synthesis of 21 tRNAs was performed using the tRNA array method with a single
linear DNA template encoding all 21 tRNAs (25 nM). *In vitro* transcription was carried out with the
single 21 tRNAs template (25 nM) in the presence of Tris-HCl pH 8.0 (40 mM), magnesium chloride
(10 mM), spermidine (2 mM), T7 RNA polymerase (1.0 U/µL, Takara) and T4 PNK (0.1 U/µL, New
England Biolab), NTPs (2 mM each: ATP, GTP, CTP, and UTP), GMP (3 mM), 0.2 U/µl inorganic
pyrophosphatase (New England Biolab), and 0.8 U/µl RNasin (Promega) at 37°C  for 12 h. The
resulting array-derived pre-tRNA transcripts were purified using the PureLink RNA Mini Kit
(Invitrogen). The purified premature tRNA transcripts were subsequently processed by RNase P
digestion in buffer R (see RNase P digestion in buffer R section) with RNase P (4 µM M1 RNA and
6 µM C5 protein) at 37 °C for 12 h for digestion. Following processing, the resulting tRNA products
were purified again using the PureLink RNA Mini Kit and used for subsequent translation assays. C5
protein was purified as described previously(22,23)

**tfPURE system preparation**

All components of the laboratory-prepared PURE system were expressed and purified essentially
as described previously, using affinity chromatography with a histidine tag, followed by gel-filtration
chromatography (38). EF-Tu was further purified by two rounds of affinity chromatography in a
stringent buffer to remove residual tRNA, as previously reported(22). Ribosomes were further
purified by ultrafiltration to remove residual tRNAs, following the same procedure used in our
previous work. The detailed composition of the tfPURE system used in this study is summarized in
Table S5. The composition A is the same composition we used in our previous study(22). The
compositions B and C are the compositions optimized for four and 21 tRNA expressions, respectively,
in our previous study (22,23).

**tRNA design**

Secondary structure prediction of each tRNA variant was performed using the RNAfold Web Server from the ViennaRNA package(39). For each sequence, both the minimum free energy (MFE) structure and the centroid structure were obtained, and formation of the canonical cloverleaf structure was evaluated. A cloverleaf structure was defined by the presence of a well-formed acceptor stem, D arm, anticodon arm, and T arm in the predicted secondary structure. In addition, ensemble diversity and the frequency of the MFE structure within the structural ensemble were calculated to assess structural stability and heterogeneity. These parameters were used to compare predicted secondary structures and to select tRNA variants for experimental validation.

**RNase P digestion in buffer R**

RNase P digestion of pre-tRNA transcripts was performed in buffer R, consisting of 50 mM Tris-HCl (pH 7.6), 60 mM $NH_4Cl$, 10 mM $Mg(OAc)_2$, and 5 mM spermidine. Pre-tRNA substrates (250 nM) were incubated with M1 RNA (500 nM) and C5 protein (750 nM) in buffer R at 37 °C for 12 h.

**Urea-PAGE**

Denaturing polyacrylamide gel electrophoresis was performed using 8% polyacrylamide gels (acrylamide: bisacrylamide = 19:1) containing 8 M urea, 0.1% ammonium persulfate, and 0.1% N, N, N', N'-tetramethylethylene-diamine in Tris-borate-EDTA buffer. RNA samples were mixed with stripping buffer composed of 50 mM EDTA, 90% formamide, and 0.025% bromophenol blue prior to electrophoresis. Following electrophoresis, RNA was visualized by staining with SYBR Green II (Takara). An RNA size marker (Dyna Marker RNA Low II Easy Load, Bio Dynamics Laboratory Inc.) was loaded alongside samples in all urea-PAGE analyses.

**RT-qPCR for tRNA quantification**

For quantification of individual tRNA species, 21-tRNA mixtures prepared by the tRNA array method were purified by size selection using the RNA Clean & Concentrator system (Zymo Research) to enrich the tRNA fraction. The purified 21-tRNA mixtures were diluted to 500 pM as the total 21 tRNAs and analyzed by RT-qPCR using the One Step TB Green PrimeScript PLUS RT-PCR Kit (Takara) with tRNA-specific primer sets (Supplementary Table S1). The following primer pairs were used: Gly, primers 29 and 30; Pro_v1, primers 31 and 32; Pro_v2, primers 33 and 32; Ile_v1, primers 34 and 35; Ile_v2, primers 36 and 35; Glu, primers 37 and 38; Asn_v1, primers 39 and 40; and Asn_v2, primers 41 and 40. Reactions were performed on a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific) under the following conditions: reverse transcription at 42 °C for 30 min; initial denaturation at 95 °C for 10 sec; followed by 50 cycles of 95 °C for 5 sec and 60 °C for 30 sec, with fluorescence measured at the end of each cycle.

For quantification, standard curves were generated using serial dilutions of individually prepared
tRNAs with known concentrations, and the concentrations of the corresponding tRNAs in the 21-
tRNA mixtures were determined from the measured Ct values. Primer specificity was evaluated by
comparing amplification of the target tRNA with that of the corresponding non-target 20-tRNAs
mixture.

**Translation with purified 21 tRNAs**

Translation reactions were carried out in the laboratory-prepared tRNA-free PURE system (tfPURE,
composition A, Table S6) supplemented with a prepared 21-tRNA mixture, a reporter gene encoding
DNA template (1 nM). The reaction mixture was incubated at 30 °C for 16 h. The total concentration
of the individually prepared 21 tRNAs was 240 ng/μL (40 ng/μL each for PIEN and 4.7 ng/μL each
for the others). When 21 tRNAs prepared by the tRNA array method were used, the total tRNA
concentration was also set to 240 ng/μL. For assays involving tRNA variants, PURE systems lacking
the respective tRNA were prepared.

After incubation, reporter protein production was quantified using three independent assays. For
luciferase, a 1 μL aliquot of the translation reaction was mixed with 30 μL of Luciferase Assay
Reagent (Promega), and luminescence was measured using a GloMax luminometer (Promega). For
sfGFP, fluorescence was monitored for 24 h using an Mx3005P (Agilent Technologies). For GUS, a
2 μL aliquot of the translation reaction was added to 8 μL of GUS reaction buffer (final concentrations:
50 mM HEPES-KOH, pH 7.4; 0.01% Triton X-100; 5 μM TokyoGREEN-βGlcU(Na); Sekisui
Medical, Japan)(40). Fluorescence was recorded every 1 min for 60 min using an Mx3005P with
FAM detection settings. GUS activity was calculated from the slope of the linear portion of the
fluorescence time course.

**Translation coupled with tRNA expression**

tRNA-expression coupled translation reactions were performed in the tfPURE system (composition
B or C, Table S6) by addition of reporter gene encoding DNA templates (1 nM) and tRNA template
DNA. The concentrations of tRNA template DNA, T7 RNA polymerase (Takara), T4 PNK (New
England Biolab), and RNase P were adjusted depending on the experimental design and are specified
in the corresponding figure legends. The reaction mixture was incubated at 30°C for 24 h. After
incubation, translation outputs were quantified using reporter-specific assays for luciferase, sfGFP, or
GUS, as described above.

## Results

### Identification of limiting tRNAs

In our previous work, we constructed a tRNA array method that encodes all 21 tRNA species
required for translation, which are transcribed from five polycistronic gene groups, in which each
tRNA gene in the groups is directly linked in an end-to-end manner (Fig. 1A). After transcription, the
five transcripts are processed through self-cleavage of HDVR at the 3'-end and digestion with RNase
P to generate mature-length tRNA products. The translation level of luciferase using the 21 tRNAs
synthesized by this method was at a similar level to that using individually prepared *in vitro*-
synthesized (IVS) tRNA mixtures of all 21 species, but was significantly lower when using sfGFP
and beta-glucuronidase (GUS) as reporter genes(23).

To examine whether the reduced translation level was caused by insufficiency of some tRNAs, we
performed a supplementation assay (Fig. 1B), in which we added five individually-prepared tRNA
mixtures to the 21 tRNA mixtures prepared by the tRNA array method and compared the translation
efficiency of three reporter proteins (luciferase, sfGFP, and GUS) in a tRNA-free PURE system
(tfPURE). The compositions of the five added tRNA mixtures were based on the five array groups
(GARCQ, DfMHKV, SmMFT, LWY, and PIEN; see Fig. 1A). The total concentration of the 21
tRNAs was set to 240 ng/μL, under which, translation output increased linearly with tRNA
concentration (Fig. S1). The concentration of each supplemented tRNA was 40 ng/μL for PIEN
tRNAs (tRNA$^{Pro, Ile, Glu, Asn}$) and 4.7 ng/μL for the others, which was the same relative composition as
in the reconstituted 21 tRNAs system reported previously(22,33), but uniformly scaled down to 40%
of the original concentrations.

Under these conditions, supplementation with the PIEN tRNA mixture increased translation by
approximately 3- to 11-fold for all three reporter proteins: firefly luciferase, sfGFP, and GUS. In
contrast, supplementation with the GARCQ tRNAs enhanced translation of luciferase (approximately
three-fold) and GUS (approximately 20-fold), but did not increase sfGFP translation, indicating that
the optimal tRNA composition varies depending on the protein to some extent. Because the addition
of PIEN tRNAs consistently enhanced translation across all three proteins, we focused subsequent
optimization efforts on this group.

### Sequence modification of tRNA$^{Ile}$, tRNA$^{Asn}$, and tRNA$^{Pro}$

We attempted to improve the translational activity of tRNAs in the PIEN group. Previous studies
have shown that stabilization of the canonical cloverleaf structure can enhance tRNA
functionality(41,42). Secondary structure prediction revealed that the tRNA$^{Ile}$ and tRNA$^{Asn}$ used here,
which are the same as *E. coli* native sequences, do not form standard cloverleaf structures as their
minimum free energy (MFE) structures (Fig. S2). In addition, while the MFE structure of tRNA$^{Pro}$ is
predicted to form a cloverleaf, the anticodon stem region exhibits reduced structural stability (Fig.

S2). These non-standard structures may impair their translational activities. We therefore introduced
mutations to tRNA$^{Ile}$, tRNA$^{Asn}$, and tRNA$^{Pro}$ to adopt stable cloverleaf structures.

To make stable cloverleaf structures, we first introduced point mutations primarily into loop
regions while avoiding identity elements(43) and generated various tRNA variants. We then predicted
the secondary structures of the resulting variants using the ViennaRNA package(39) and evaluated
the frequency at which the MFE structure formed the canonical cloverleaf conformation and the
ensemble diversity, which quantifies the structural heterogeneity of the RNA ensemble. Variants
exhibiting a high MFE frequency and low ensemble diversity were selected for the following
translation assay. In the translation assay, tfPURE was supplemented with either a tRNA variant or
the original tRNA, together with a mixture of the remaining 20 tRNAs and DNA encoding luciferase.
The transcription and translation reactions were carried out by incubating the mixture at 30 °C for 16
h, and the luciferase activity was measured.

For tRNA$^{Ile}$ and tRNA$^{Asn}$, we selected five point mutants that exhibit increased frequencies of the
cloverleaf structure for translation assay (Figs. 2A and 2C). The percentage of MFE structures that
adopt a cloverleaf conformation are shown below in blue. For tRNA$^{Ile}$, three out of five tested variants
exhibited similar translation activity to that of the original tRNA (Fig. 2B). For tRNA$^{Asn}$, three of the
five variants showed similar activities to that of the original tRNA (Fig. 2D). In contrast, some tRNA
variants (C32G and U33A in tRNA$^{Ile}$, and C32G and A38G in tRNA$^{Asn}$), which contain mutations in
the anticodon loop, significantly reduced translation activity, indicating that these regions are
essential for translation activity, although they have not been reported as identity elements.

For tRNA$^{Pro}$, we selected seven sequences that exhibit higher frequencies of cloverleaf
conformation (Fig. 2E). In these variants, we additionally introduced a mutation (G1C) at the 5'-
terminus. The original nucleotide at this position in wild-type tRNA$^{Pro}$ is C, but was mutated to G in
the previous array method to allow transcription to start immediately downstream of the T7
promoter(23). To examine the possibility that this mutation could impair translation activity, we
selected four tRNA$^{Pro}$ variants that reverted this position to C (G1C). Among these variants, the
G1C_U40A_C16A variant exhibited the highest translation activity, reaching approximately 2.5-fold
higher activity than the original tRNA (Fig. 2F).

**Evaluation of PIEN array arrangement**

We next examined the arrangement of the PIEN group on the DNA template. In the tRNA array
method, transcription generates premature transcripts that are first processed by HDVR self-cleavage
at the 3'-end. In our previous study, we observed that the PIEN group exhibited relatively low
efficiency at this self-cleavage step(23). Because this processing step may depend on the order of
tRNA genes within the array, we constructed four PIEN group arrays with different tRNA orders and
synthesized the corresponding premature transcripts in the absence of RNase P to directly compare

HDVR self-cleavage efficiencies (Fig. 3A). The self-cleavage efficiency varied among these four
constructs, and the original PIEN arrangement showed the highest self-cleavage efficiency (Fig. 3B,
blue arrowheads). In addition, the NIPE arrangement exhibited a larger proportion of uncleaved
products of larger size. This is likely due to predominant transcription termination at the second stem
loop of the tandem T7 terminator(44) for unknown reasons, resulting in longer RNA species.

Next, we examined whether the gene arrangement affected the yield of RNase P processing. To
enable direct comparison after self-cleavage, pre-tRNAs lacking the HDV ribozyme sequence were
prepared by *in vitro* transcription. These pre-tRNAs were incubated with RNase P in buffer R (see
Methods) at 37 °C for 12 h, and the cleavage products were analyzed by urea-PAGE (Fig. 3C). The
EIPN and NIPE arrangements exhibited the largest products, while the PIEN arrangement exhibited
slightly smaller products (Fig. 3D, the bands indicated as "mature-sized tRNAs"). It should be noted
that in our previous study (23), we reported translation with these constructs but did not analyze each
HDVR and RNase P cleavages. The present analysis was therefore performed to clarify these
processing steps.

**PIEN array version 2**

Based on these results, we decided to use the original PIEN tRNA order because it exhibited the
highest HDVR self-cleavage (Fig. 3B) and modest RNase P digestion (Fig. 3D). We then proceeded
to replace the original tRNA sequences in the PIEN array with the cloverleaf-stabilized variants that
showed higher translation activities (tRNA$^{Pro}$_G1C_U40A_C16A, tRNA$^{Ile}$_G16A, and
tRNA$^{Asn}$_C17G, respectively). Because the T7 RNA polymerase we used starts RNA synthesis from
G nucleotides, to adopt the tRNA$^{Pro}$, which has a non-G at the 5'-terminus, we introduced a 27-
nucleotide leader sequence containing 5'-G upstream of the tRNA$^{Pro}$_G1C_U40A_C16A gene (Fig.
4A). The first eight nucleotides of the leader sequence were designed for efficient transcription
initiation(45) and the last six nucleotides were designed to enhance RNase P cleavage efficiency(46).
The intervening sequence was adopted from a previously reported leader sequence(33). The resulting
PIEN construct was designated as L_PIEN_v2.

We first compared the efficiency of HDVR self-cleavage of L_PIEN_v2 with that of the previous
construct (PIEN_v1) by performing *in vitro* transcription followed by urea-PAGE analysis (Fig. 4B).
Quantification of the ratio of cleaved (blue arrowheads) to uncleaved (yellow arrowheads) products
revealed that the original PIEN construct yielded approximately 25% cleaved products, whereas
L_PIEN_v2 demonstrated an efficiency of approximately 53%, corresponding to a 2.1-fold
improvement. When only the tRNA sequences were replaced without the leader sequence (PIEN_v2),
the self-cleavage efficiency was intermediate between that of PIEN_v1 and L_PIEN_v2 (Fig. S3).

We next evaluated the translation activity of the expressed tRNA set by a translation-coupling
reaction (Fig. 4C). In this assay, the template DNA encoding the PIEN arrays was used at a

concentration of 25 nM, corresponding to half the DNA concentration used in our previous study(23),
to evaluate translation activity under an unsaturated condition. Under this condition, the original PIEN
construct exhibited a luciferase activity of $8.8×10^4$, whereas L_PIEN_v2 exhibited a value of $4.6×10^6$,
representing an approximately 52-fold increase in translation activity. This large improvement may
reflect the combined effects of enhanced HDVR processing, improved tRNA function, and the non-
proportional increase in translation with increasing PIEN tRNA concentration in this range (Fig. S4).

**Construction and evaluation of tRNA array version 2**

L_PIEN_v2 was incorporated into the polycistronic DNA construct encoding all 21 tRNAs used in
our previous study as the tRNA array version 2. Using this DNA, we synthesized 21 tRNAs under a
translation-uncoupled condition, purified them, and evaluated the translation activity for three
reporter genes (luciferase, sfGFP, and GUS) at a total tRNA concentration of 240 ng/µL (Fig. 5A).
For luciferase translation, the original tRNA array (v1) yielded a luminescence signal of
approximately $4.1×10^5$, whereas tRNA array version 2 (v2) produced a signal of approximately
$4.3×10^6$, corresponding to an approximately 11-fold increase (Fig. 5B). Similarly, sfGFP translation
was enhanced by approximately 4.1-fold (Fig. 5C) and GUS translation increased by approximately
5.1-fold with tRNA array version 2 (Fig. 5D). Furthermore, for all three reporter genes, tRNA array
version 2 supported translation activities comparable to, or exceeding, those achieved with
individually prepared 21 IVS tRNA mixtures (IVS) at the same concentration, the composition of
which is based on a previous study(33). RT-qPCR analysis of purified 21-tRNA mixtures showed that
the abundances of the redesigned PIEN-group tRNAs were increased in version 2 relative to version
1, whereas tRNA$^{Gly}$, the construct of which was unchanged, exhibits a slight decrease (Fig. S5).

We compared the performance of tRNA array version 2 in the translation-coupled method. In this
experiment, all 21 tRNAs were synthesized in situ within tfPURE and directly utilized for translation
(Fig. 5E). First, we measured tRNA synthesis in tfPURE. Quantification of total tRNA production
revealed that tRNA array version 2 yielded 780 ng/µl, approximately 1.3-fold higher total tRNA levels
than version 1 (Fig. S6). When measuring translated luciferase activity, tRNA array version 2
supported an approximately three-fold higher translation than that with version 1, and an
approximately 2.4-fold increase relative to individually prepared 21 IVS tRNA mixtures at the same
concentration (Fig. 5F). Similarly, translation of sfGFP was enhanced by approximately six-fold
compared with version 1 and reached a level comparable to that obtained with individually prepared
21 IVS tRNAs (Fig. 5G). GUS translation increased by approximately 7.8-fold relative to version 1,
reaching approximately 50% of the level achieved with individually prepared 21 IVS tRNAs (Fig.
5H). The absolute GUS activity values differed more markedly between the translation-uncoupled
and translation-coupled reactions than those of luciferase or sfGFP, possibly because the two reactions
differ substantially in reaction composition and time-dependent tRNA supply. The oligomeric GUS
reaction may be more sensitive to these changes.

**Discussion**

In this study, we identified the tRNA groups that limited translation activity in the tRNA array method established in our previous work(23), and constructed and developed tRNA array version 2 by improving both tRNA activity and pre-tRNA processing efficiency. Specifically, we identified the PIEN group as a bottleneck for translation of all measured proteins, enhanced the activities of tRNAs (especially tRNA$^{Pro}$) through stabilization of their cloverleaf structures, and redesigned the PIEN array by incorporating a leader sequence. As a result, the translation activity of 21 tRNAs prepared using tRNA array version 2 was improved to levels similar to those achieved with individually prepared tRNAs across multiple reporter proteins, enabling stable expression of all 21 tRNAs in translation-coupled reaction in the PURE system. Our results highlight that quantitative tuning of tRNA composition is critical for efficient translation and represents a key requirement for constructing self-reproducible gene expression systems(47).

Our results demonstrated that translation activity strongly depended on the relative composition of the 21 tRNAs. Translation activity with the previous tRNA array method was increased by supplementation with the PIEN tRNA group for all tested reporter genes (Fig. 1), indicating that reduced translation output from array-derived tRNAs was primarily attributable to insufficient levels of these tRNA species. Consistently, the PIEN tRNA set was used at higher concentrations in a previous study that used an *in vitro*-transcribed tRNA set (33). Considering that the relative expression level of each group was similar, as shown in our previous study(23), this result suggests that efficient translation does not require uniform concentrations of all 21 tRNAs, but requires adjusted concentrations that contain more of the highly-demanded tRNA species, such as PIEN tRNAs.

This study demonstrated that tRNA$^{Pro}$, tRNA$^{Ile}$, and tRNA$^{Asn}$ were predicted to either fail to stably adopt the canonical cloverleaf structure or to exhibit high structural heterogeneity, and some of their variants with more frequent cloverleaf structures exhibited enhanced translation activity. These results suggest that the canonical cloverleaf structure plays a certain role in translation, probably through interactions with translation factors(48,49). Considering that their original sequences are derived from *E. coli* native tRNAs, this result further suggests that native tRNA sequences are not always fully suited to in vitro translation using unmodified tRNAs. However, some mutants (e.g., G18U and G18A in Fig. 2) that stabilized the cloverleaf structures did not enhance translation, indicating that other unknown factors are also important for tRNA function.

The introduction of leader sequences was another key factor contributing to the improved translation performance of tRNA array version 2. Although the leader sequence was designed to allow transcription of tRNA$^{Pro}$ variants that start with 5'-C, an unexpected improvement was observed in HDVR self-cleavage efficiency. This finding suggests that incorporation of leader sequences and modification of tRNA sequences altered the overall secondary structure of the pre-tRNA, potentially generating conformations that are more favorable for HDVR cleavage. In addition, insertion of the

leader sequence is expected to mitigate heterogeneity at the 5'-end of tRNA transcripts, which can arise from imprecise transcription initiation by T7 RNA polymerase(50,51). Further investigation will be required to determine how such leader sequences behave in other sequence contexts and whether additional optimization is possible.

In this study, we observed differences in processing efficiency among four IPEN arrangements (IPEN, PIEN, EIPN, and NIPE) (Figs. 3B and 3D), as well as among PIEN_v1, PIEN_v2, and L_PIEN_v2 (Fig. S3). To investigate whether these differences were associated with structural changes, we analyzed the predicted secondary structures using ViennaRNA (Fig. S7). The precursor RNAs of the EIPN and NIPE constructs, both of which showed lower HDV ribozyme (HDVR) self-cleavage efficiency, adopted relatively similar predicted structures both with (Fig. S7A) and without (Fig. S7B) the HDVR and terminator sequences. However, we have no direct evidence that this structural tendency accounts for the reduced cleavage efficiency. Likewise, no clear structural tendencies were identified among the predicted structures of PIEN_v1, PIEN_v2, and L_PIEN_v2 (Fig. S7C). Taken together, these results suggest that the mechanistic basis of cleavage efficiency remains poorly understood, and elucidating this mechanism represents an important direction for future research. Although tRNA array version 2 constructed in this study exhibited high translation performance as a simultaneous expression system based on unmodified tRNAs, there remains considerable room for further improvement. For example, the same optimization strategy as conducted for the PIEN group in this study can be applied to the other tRNA groups. In addition, because both HDVR and RNase P processing efficiencies are sequence dependent, systematic design of tRNA order within the array and of leader sequences may enable more flexible control of tRNA composition tailored to specific translation demands. In the longer term, customization of the PURE system using translation factors better suited for unmodified tRNAs(52) could further advance this platform toward a self-reproducible translation system with translation performance approaching that of native tRNAs.

**Acknowledgement**

The authors declare no conflict of interest associated with this manuscript. This work was supported by JST, CREST Grant Number JPMJCR20S1, Japan, and Kakenhi Grant Numbers 22H05402, 24H01111, and 23KJ0815.

Ryota Miyachi: Conceptualization, Methodology, Data curation, Investigation, Writing - Original Draft. Anna Irie: Methodology, Investigation. Norikazu Ichihashi: Conceptualization, Supervision, Writing- Reviewing and Editing, Funding acquisition.

Supporting Information includes supplemental figures (Figures S1–S7) and tables (Tables S1–S6).

**References**

1.　Rothschild, L. J., Averesch, N. J. H., Strychalski, E. A., Moser, F., Glass, J. I., Cruz Perez,
R., Yekinni, I. O., Rothschild-Mancinelli, B., Roberts Kingman, G. A., Wu, F. and other 7
authors.: Building Synthetic Cells─From the Technology Infrastructure to Cellular Entities.
ACS Synth. Biol., **13,** 974–997 (2024).

2.　Buddingh', B. C. & Van Hest, J. C. M.: Artificial Cells: Synthetic Compartments with Life-
like Functionality and Adaptivity. Acc. Chem. Res., **50,** 769–777 (2017).

3.　Tror, S., Jeon, S. M., Nguyen, H. T., Huh, E. & Shin, K.: A Self-Regenerating Artificial Cell,
that is One Step Closer to Living Cells: Challenges and Perspectives. Small Methods, **7,**
2300182 (2023).

4.　Noireaux, V., Maeda, Y. T. & Libchaber, A.: Development of an artificial cell, from self-
organization to computation and  self-reproduction. Proc. Natl. Acad. Sci. U. S. A., **108,**
3473–3480 (2011).

5.　Amy Yewdall, N., Mason, A. F. & Van Hest, J. C. M.: The hallmarks of living systems:
towards creating artificial cells. Interface Focus, **8,** 20180023 (2018).

6.　Duim, H. & Otto, S.: Towards open-ended evolution in self-replicating molecular systems.
Beilstein Journal of Organic Chemistry, **13,** 1189 (2017).

7.　De Capitani, J. & Mutschler, H.: The Long Road to a Synthetic Self-Replicating Central
Dogma. Biochemistry, **62,** 1221–1232 (2023).

8.　Kazuta, Y., Matsuura, T., Ichihashi, N. & Yomo, T.: Synthesis of milligram quantities of
proteins using a reconstituted in vitro  protein synthesis system. J. Biosci. Bioeng., **118,** 554–
557 (2014).

9.　Ivanov, I., Castellanos, S. L., Balasbas, S., Otrin, L., Maruscaroniccaron, N.,
Vidakovicacute-Koch, T. & Sundmacher, K.: Bottom-Up Synthesis of Artificial Cells:
Recent Highlights and Future Challenges. Annu. Rev. Chem. Biomol. Eng., **12,** 287–308
(2021).

10.　Hagino, K. & Ichihashi, N.: In Vitro Transcription/Translation-Coupled DNA Replication
through Partial Regeneration of 20 Aminoacyl-tRNA Synthetases. ACS Synth. Biol., **12,**
1252–1263 (2023).

11. Hagino, K., Masuda, K., Shimizu, Y. & Ichihashi, N.: Sustainable regeneration of 20
aminoacyl-tRNA synthetases in a reconstituted system toward self-synthesizing artificial
systems. Science Advances , **11,** 6269 (2025).

12. Lavickova, B., Laohakunakorn, N. & Maerkl, S. J.: A partially self-regenerating synthetic
cell. Nat. Commun., **11,** 6340 (2020).

13. Doerr, A., Foschepoth, D., Forster, A. C. & Danelon, C.: In vitro synthesis of 32 translation-
factor proteins from a single template reveals impaired ribosomal processivity. Sci. Rep., **11,**
1898 (2021).

14. Libicher, K. & Mutschler, H.: Probing self-regeneration of essential protein factors required
for In vitro translation activity by serial transfer. Chemical Communications, **56,** 15426–
15429 (2020).

15. Libicher, K., Hornberger, R., Heymann, M. & Mutschler, H.: In vitro self-replication and
multicistronic expression of large synthetic genomes. Nat. Commun., **11,** 904 (2020).

16. Wei, E. & Endy, D.: Experimental tests of functional molecular regeneration via a standard
framework for coordinating synthetic cell building. bioRxiv, 2021.03.03.433818 (2021).
doi:10.1101/2021.03.03.433818

17. Mottaghi, S. S. & Maerkl, S. J.: PURE makes PURE: reconstitution of the PURE cell-free
system from self-synthesized non-ribosomal proteins. bioRxiv, 2025.12.17.694911 (2025).
doi:10.64898/2025.12.17.694911

18. Schwarz-Schilling, M., Cohen, I., Dupin, A., Avidan, N., Barak, Y., Shimizu, Y., Daube, S. S.
& Bar-Ziv, R. H.: Autonomous biogenesis of all thirty proteins of the Escherichia coli
translation machinery. Nature Communications 2026 17:1, **17,** 1028 (2026).

19. Shimojo, M., Amikura, K., Masuda, K., Kanamori, T., Ueda, T. & Shimizu, Y.: In vitro
reconstitution of functional small ribosomal subunit assembly for  comprehensive analysis of
ribosomal elements in E. coli. Commun. Biol., **3,** 142 (2020).

20. Levy, M., Falkovich, R., Daube, S. S. & Bar-Ziv, R. H.: Autonomous synthesis and assembly
of a ribosomal subunit on a chip. Sci. Adv., **6,** eaaz6020 (2020).

21. Aoyama, R., Masuda, K., Shimojo, M., Kanamori, T., Ueda, T. & Shimizu, Y.: In vitro
reconstitution of the Escherichia coli 70S ribosome with a full set of  recombinant ribosomal
proteins. J. Biochem., **171,** 227–237 (2022).

22. Miyachi, R., Shimizu, Y. & Ichihashi, N.: Transfer RNA Synthesis-Coupled Translation and
DNA Replication in a Reconstituted  Transcription/Translation System. ACS Synth. Biol.,
**11,** 2791–2799 (2022).

23.    Miyachi, R., Masuda, K., Shimizu, Y. & Ichihashi, N.: Simultaneous in vitro expression of
490            minimal 21 transfer RNAs by tRNA array method. Nature Communications 2025, **16,** 7418
491            (2025).

492    24.    Li, F., Baranwal, A. K. & Maerkl, S. J.: Continuous in situ synthesis of a complete set of
493            tRNAs sustains steady-state translation in a recombinant cell-free system. Nature
494            Communications 2025, **16,** 6212 (2025).

495    25.    Sakatani, Y., Ichihashi, N., Kazuta, Y. & Yomo, T.: A transcription and translation-coupled
496            DNA replication system using rolling-circle replication. Sci. Rep., **5,** 10404 (2015).

497    26.    Sakatani, Y., Yomo, T. & Ichihashi, N.: Self-replication of circular DNA by a self-encoded
498            DNA polymerase through rolling-circle replication and recombination. Sci. Rep., **8,** 13089
499            (2018).

500    27.    Okauchi, H. & Ichihashi, N.: Continuous Cell-Free Replication and Evolution of Artificial
501            Genomic DNA in a Compartmentalized Gene Expression System. ACS Synth. Biol., **10,**
502            3507–3517 (2021).

503    28.    Fujiwara, K., Katayama, T. & Nomura, S. I. M.: Cooperative working of bacterial
504            chromosome replication proteins generated by a reconstituted protein expression system.
505            Nucleic Acids Res., **41,** 7176–7183 (2013).

506    29.    van Nies, P., Westerlaken, I., Blanken, D., Salas, M., Mencía, M. & Danelon, C.: Self-
507            replication of DNA by its encoded proteins in liposome-based synthetic  cells. Nat.
508            Commun., **9,** 1583 (2018).

509    30.    Abil, Z., Restrepo Sierra, A. M., Stan, A. R., Châne, A., del Prado, A., de Vega, M.,
510            Rondelez, Y. & Danelon, C.: Darwinian Evolution of Self-Replicating DNA in a Synthetic
511            Protocell. Nature Communications 2024, **15,** 9091 (2024).

512    31.    María, A., Sierra, R., Ramirez Gomez, F., Van Tongeren, M., Sierra Heras, L. & Danelon, C.:
513            A synthetic cell with integrated DNA self-replication and lipid biosynthesis. Nature
514            Communications 2026, **17,** 2727 (2026).

515    32.    Shimizu, Y., Inoue, A., Tomari, Y., Suzuki, T., Yokogawa, T., Nishikawa, K. & Ueda, T.:
516            Cell-free translation reconstituted with purified components. Nat. Biotechnol., **19,** 751–755
517            (2001).

518    33.    Hibi, K., Amikura, K., Sugiura, N., Masuda, K., Ohno, S., Yokogawa, T., Ueda, T. &
519            Shimizu, Y.: Reconstituted cell-free protein synthesis using in vitro transcribed tRNAs.
520            Commun. Biol., **3,** 350 (2020).

521    34.    Iwane, Y., Hitomi, A., Murakami, H., Katoh, T., Goto, Y. & Suga, H.: Expanding the amino
522            acid repertoire of ribosomal polypeptide synthesis via the  artificial division of codon boxes.
523            Nat. Chem., **8,** 317–325 (2016).

35. Fujino, T., Tozaki, M. & Murakami, H.: An Amino Acid-Swapped Genetic Code. ACS
Synth. Biol., **9,** 2703–2713 (2020).

36. Fujino, T., Sonoda, R., Higashinagata, T., Mishiro-Sato, E., Kano, K. & Murakami, H.:
Ser/Leu-swapped cell-free translation system constructed with natural/in vitro transcribed-
hybrid tRNA set. Nature Communications 2024, **15,** 4143 (2024).

37. Schürer, H., Lang, K., Schuster, J. & Mörl, M.: A universal method to produce in vitro
transcripts with homogeneous 3' ends. Nucleic Acids Res., **30,** e56 (2002).

38. Fujii, S., Matsuura, T., Sunami, T., Nishikawa, T., Kazuta, Y. & Yomo, T.: Liposome display
for in vitro selection and evolution of membrane proteins. Nature Protocols 2014 9:7, **9,**
1578–1591 (2014).

39. Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R. & Hofacker, I. L.: The Vienna RNA
Websuite. Nucleic Acids Res., **36,** W70–W74 (2008).

40. Ahmad, S., Hughes, M. A., Lane, K. T., Redinbo, M. R., Yeh, L. A. & Scott, J. E.: A High
Throughput Assay for Discovery of Bacterial β-Glucuronidase Inhibitors. Curr. Chem.
Genomics, **5,** 13 (2011).

41. Helmling, C., Chan, A. I. & Cunningham, C. N.: Engineered initiator tRNAs can effectively
start translation at non-AUG start codons and diversify N-terminal amino acids for mRNA
Display. Nucleic Acids Res., **53,** gkaf003 (2025).

42. Spinck, M., Guppy, A. & Chin, J. W.: Automated orthogonal tRNA generation. Nature
Chemical Biology 2025, **21,** 657–667 (2025).

43. Cervettini, D., Tang, S., Fried, S. D., Willis, J. C. W., Funke, L. F. H., Colwell, L. J. & Chin,
J. W.: Rapid discovery and evolution of orthogonal aminoacyl-tRNA synthetase–tRNA pairs.
Nature Biotechnology 2020, **38,** 989–999 (2020).

44. Calvopina-Chavez, D. G., Gardner, M. A. & Griffitts, J. S.: Engineering efficient termination
of bacteriophage T7 RNA polymerase transcription. G3: Genes|Genomes|Genetics, **12,**
jkac070 (2022).

45. Conrad, T., Plumbom, I., Alcobendas, M., Vidal, R. & Sauer, S.: Maximizing transcription of
nucleic acids with efficient T7 promoters. Commun. Biol., **3,** 439 (2020).

46. Niland, C. N., Anderson, D. R., Jankowsky, E. & Harris, M. E.: The contribution of the C5
protein subunit of Escherichia coli ribonuclease P to specificity for precursor tRNA is
modulated by proximal 5′ leader sequences. RNA, **23,** 1502–1511 (2017).

47. Nishizawa, C., Aburaya, S., Kosaka, Y., Sugase, K. & Aoki, W.: Optimizing in vitro
expression balance of central dogma-related genes using parallel reaction monitoring. J.
Biosci. Bioeng., **138,** 97–104 (2024).

48. Schultz, S. K., Katanski, C. D., Halucha, M., Peña, N., Fahlman, R. P., Pan, T. & Kothe, U.:
Modifications in the T arm of tRNA globally determine tRNA maturation, function, and
cellular fitness. Proceedings of the National Academy of Sciences, **121,** e2401154121
(2024).

49. Eargle, J., Black, A. A., Sethi, A., Trabuco, L. G. & Luthey-Schulten, Z.: Dynamics of
Recognition between tRNA and Elongation Factor Tu. J. Mol. Biol., **377,** 1382 (2008).

50. Takatsuki, H., Miyachi, R., Seo, K., Hagino, K. & Ichihashi, N.: Single-subunit RNA
polymerases, KpnP, Ro45Iw, and CD23823, with precise terminal synthesis. Journal of
Biological Chemistry, **301,** 110359 (2025).

51. Pleiss, J. A., Derrick, M. L. & Uhlenbeck, O. C.: T7 RNA polymerase produces 5' end
heterogeneity during in vitro transcription from certain templates. RNA, **4,** 1313 (1998).

52. Clifton, B. E., Fariz, M. A., Uechi, G. I. & Laurino, P.: Evolutionary repair reveals an
unexpected role of the tRNA modification m1G37 in aminoacylation. Nucleic Acids Res.,
**49,** 12467 (2021).

**Figures**

**Figure 1. Identification of limiting tRNA groups.**

(A) Schematic of 21 tRNAs synthesis by the tRNA array method. The 21 tRNA genes are divided
into five groups and encoded in a single DNA template. After transcription, five transcripts are
processed into mature-sized tRNAs through HDVR self-cleavage and RNase P digestion. (B)
Schematic of the supplementation assay. The purified 21 tRNAs (240 ng/μL) synthesized by the
tRNA array method and an individually-synthesized additional tRNA mixture (PIEN: 40 ng/μL
each; others: 4.7 ng/μL each), corresponding to one of the five groups, were used for translation of
three reporter proteins (firefly luciferase, sfGFP, and GUS) in a tRNA-free PURE system (tfPURE;
composition A, see Tables S5 and S6) containing reporter-encoding DNA template (1 nM) and T7
RNA polymerase (0.42 U/μL) at 30 °C for 16 h. (C) Luciferase activities. (D) sfGFP fluorescence.
(E) GUS activities. GUS activity was quantified by an enzymatic assay after the reaction, and the
slope of fluorescence increase over time was used as a measure of activity. Error bars indicate
standard deviations (N = 3).

**Figure 2. Sequence modification of tRNAs to stabilize cloverleaf structures.**

(A, C, E) Minimum free energy (MFE) secondary structures of tRNA$^{Ile}$(A), tRNA$^{Asn}$(C), and tRNA$^{Pro}$ (E) variants predicted using ViennaRNA(39). The percentages indicate the frequency with which the corresponding minimum free energy (MFE) structure was predicted to adopt the canonical cloverleaf conformation among the ensemble. The number of mutational site follows the universal conventional tRNA positions (i.e., anticodons are 34–36 for all tRNAs). (B, D, F) Translation activity assays of tRNA$^{Ile}$ (B), tRNA$^{Asn}$ (D), and  tRNA$^{Pro}$ (F) variants. Translation reactions were performed in the  tfPURE system (composition A) containing a 20-tRNAs mixture

(500 ng/μL) lacking the target tRNA, one of the target tRNA variants (tRNA$^{Ile}$ or tRNA$^{Asn}$ at 13
ng/μL, tRNA$^{Pro}$ at 25 ng/μL), a DNA template encoding luciferase (1 nM), and T7 RNA polymerase
(0.42 U/μL). Reactions were incubated at 30 °C for 16 h and luciferase activity was measured. Error
bars indicate standard deviations (N = 3).

**Figure 3.   Evaluation of PIEN group arrangement.**

(A) Schematic of the HDVR self-cleaving assay of PIEN group constructs. Transcription was
performed using templates with different tRNA orders in the absence of RNase P, and the resulting
transcripts were analyzed by urea-PAGE. (B) Urea-PAGE analysis of transcripts from PIEN group
constructs, followed by SYBR Green II staining. Predicted bands corresponding to transcripts
containing HDVR and those lacking HDVR are indicated by blue and yellow arrowheads,
respectively. (C) Schematic of the experimental workflow for the RNase P digestion assay. Prepared
pre-tRNAs (250 nM) lacking HDVR were used as substrates and incubated with RNase P (500 nM
M1 RNA and 750 nM C5 protein) in buffer R (see Methods). (D) Urea-PAGE analysis of the
digested products, followed by SYBR green II staining. Individually prepared PIEN tRNA mixtures
were included as size controls. Undigested pre-tRNA species are indicated by red arrows.

**Figure 4.  Cleavage of and translation with PIEN array version 2.**

(A) Design of the improved PIEN construct (L_PIEN_v2), in which tRNA$^{Pro}$, tRNA$^{Ile}$, and tRNA$^{Asn}$ were replaced with the higher-activity variants (tRNA$^{Pro}$_G1C_U40A_C16A, tRNA$^{Ile}$_G16A, and tRNA$^{Asn}$_C17G, respectively) and a leader sequence was inserted upstream of tRNA$^{Pro}$. (B) Urea-PAGE analysis of transcripts generated from the previous construct (PIEN_v1) and L_PIEN_v2 templates, followed by SYBR green II staining. Predicted bands corresponding to the uncleaved transcript containing HDVR and the transcript after self-cleavage of HDVR are indicated by yellow and blue arrowheads, respectively. (C) Schematic of translation assay using PIEN tRNAs. Reactions were performed in tfPURE (composition B) supplemented with the tRNA$^{PIEN}$ array template DNA (25 nM), luciferase template DNA (1 nM), RNase P (1 μM M1 RNA and 1.5 μM C5 protein), T4 polynucleotide kinase (0.094 U/μL), T7 RNA polymerase (1.7 U/μL), and 17 IVS tRNAs that lack PIEN tRNAs. After incubation at 30 °C for 24 h, luciferase activity was measured. (D) Measured luciferase activities. Error bars indicate standard deviations (N = 3).

**Figure 5. Translation with 21 tRNAs prepared by using tRNA array ver. 2.**

(A) Schematic of translation assays using 21 tRNAs prepared by the translation-uncoupled method. Twenty-one tRNAs were simultaneously expressed using the tRNA array method from a 21-tRNA array version 2, in which the PIEN group was replaced with version 2 variants. The resulting 21-tRNA mixture (240 ng/μL) was purified and added to tfPURE (composition A) together with a reporter gene (luciferase, sfGFP, or GUS)–encoding DNA template (1 nM) and T7 RNA polymerase (0.42 U/μL), followed by incubation at 30 °C for 16 h before activity measurements. (B) Luciferase activities. (C) sfGFP fluorescence. (D) GUS activity. (E) Schematic of translation-coupled *in situ* synthesis of all 21 tRNAs using tRNA array version 2. Reactions were performed in tfPURE (composition C) supplemented with the 21 tRNA template (50 nM), a reporter gene (luciferase, sfGFP, or GUS)–encoding DNA(1 nM), T7 RNA polymerase (3.4 U/μL), T4 polynucleotide kinase (0.094 U/μL), and RNase P (4 μM M1 RNA and 6 μM C5 protein). When using individually prepared 21 IVS tRNAs (IVS), the concentration was set to 780 ng/μL. After incubation at 30 °C for 24 h, translation activities were measured. (F) Luciferase activities. (G) sfGFP fluorescence.  (H) GUS activity. Error bars indicate standard deviations (N = 3).