Arousal modulates functional connectivity through structured and

hemispherically asymmetric community architecture during wakefulness

Xiangyu Kong[1], Siyu Li[1], Gaolang Gong[1,2,3*]

[1] State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, Beijing, China

[2] Beijing Key Laboratory of Brain Imaging and Connectomics, Beijing Normal

University, Beijing 100875, China

[3] Chinese Institute for Brain Research, Beijing 102206, China

[*] **Corresponding author**
Dr. Gaolang Gong
National Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern
Institute for Brain Research, Beijing Normal University,
#19 Xinjiekouwai Street, Beijing 100875, China
Tel: +86 10 58804678; Fax: +86 10 58804678

E-mail: gaolang.gong@bnu.edu.cn

Abstract

Arousal fluctuates continuously during wakefulness, yet how these moment-to-moment variations shape large-scale functional connectivity (FC) remains unclear. Here, we combined 7T fMRI with concurrent pupillometry to quantify, for every functional connection, how time-varying FC covaries with spontaneous arousal in the awake human brain. Rather than exerting a uniform influence across the connectome, arousal organized FC into a low-dimensional set of seven connectivity communities, each defined by characteristic network compositions. These communities exhibited systematic hemispheric asymmetries, specifically identifying a "left-hemisphere centripetal architecture" where the left hemisphere serves as a structural sink for the asymmetric convergence of arousal-modulated signals. Importantly, hemispheric asymmetry did not arise from global shifts in connectivity strength, but instead reflected structured spatial heterogeneity embedded within community architecture. This modular and asymmetric organization was highly preserved during naturalistic movie watching, indicating that arousal-related modulation of FC reflects intrinsic principles that generalize across awake cognitive contexts. Together, these findings demonstrate that moment-to-moment arousal fluctuations shape large-scale FC through structured, hemispherically asymmetric network organization during wakefulness.

Introduction

Arousal is a core dimension of brain state that shapes perception, attention, and the coordination of large-scale neural systems. Across major transitions—such as wakefulness to sleep, anesthesia, or disorders of consciousness—substantial reorganization of functional connectivity (FC) has been consistently observed (Chow et al., 2013; Tagliazucchi et al., 2016; Demertzi et al., 2019; Banks et al., 2020; Damaraju et al., 2020; Huang et al., 2020; Jang et al., 2024). These state-based findings demonstrate that arousal fundamentally influences whole-brain communication patterns. However, such approaches typically contrast discrete arousal states and therefore provide limited insight into how moment-to-moment fluctuations in arousal within the awake brain influence the organization of functional connectivity.

Increasing evidence shows that arousal varies continuously even during stable wakefulness, including during resting-state fMRI and ongoing cognitive engagement. These spontaneous fluctuations, often indexed by pupil diameter (Reimer et al., 2014; McGinley et al., 2015; Joshi & Gold, 2020), modulate neural gain, sensory responses, and behavioral performance. Yet despite their ubiquity and behavioral relevance, it remains largely unknown how fine-grained arousal variations are expressed across the functional connectome. Prior work has primarily examined regional signal amplitude or isolated networks (Yellin et al., 2015; Schneider et al., 2016; Breeden et al., 2017; Podvalny et al., 2021; Sobczak et al., 2021; Lloyd et al., 2023), thereby leaving unresolved the fundamental question of whether the awake brain's FC is uniformly sensitive to arousal or whether arousal instead imprints structured spatial patterns across functional networks.

A parallel question is whether arousal-modulated connectivity patterns show hemispheric asymmetry, given longstanding evidence for lateralized arousal, vigilance,

and alerting mechanisms. Studies of unihemispheric sleep and lateralized arousal

dynamics in animals (Rattenborg et al., 2000; Lyamin et al., 2016; Mascetti, Gian

Gastone, 2016; Reicher et al., 2021; Fenk et al., 2023; Libourel et al., 2023),

asymmetries in human EEG-based vigilance (Tamaki et al., 2016), and

right-lateralized alerting functions in attention (Heilman & Abell, 1980; Sturm &

Willmes, 2001; Shulman et al., 2010; Corbetta & Shulman, 2011) all suggest that

arousal may modulate left and right hemispheric systems differently. Nevertheless,

these observations have yet to be linked to the organization of whole-brain functional

interactions, and it remains unknown whether such asymmetries manifest at the level

of large-scale FC, and whether they reflect organized connectivity patterns rather than

non-specific global effects.

To address these questions, we combined high-field fMRI with concurrent

pupillometry to quantify, for every functional connection, how its connectivity

covaries with spontaneous arousal fluctuations during wakefulness. This edgewise

measure of arousal–time varying functional connectivity (tvFC) coupling provides a

comprehensive map of where in the connectome arousal leaves its strongest imprint,

without imposing predefined states or regional assumptions. Using this framework,

we first test whether arousal sensitivity is spatially homogeneous or segregates into

distinct sets of connections with similar coupling profiles. We next assess whether

these spatially organized arousal-modulated patterns show systematic hemispheric

asymmetry, with particular emphasis on attentional systems that show known

lateralization. Finally, we evaluate the cross-context stability of this organizational

structure by comparing resting state and naturalistic movie watching in the same

participants. Together, these analyses delineate how moment-to-moment arousal

fluctuations shape large-scale functional architecture in the awake human brain.

Results

The processing procedure of estimating arousal–tvFC coupling from fMRI and
pupillometry was illustrated in Figure 1. Here, we use the term arousal–tvFC coupling
to refer to the regression-based estimate of how spontaneous arousal fluctuations
modulate each functional connection over time.

Figure 1. Pipeline for estimating arousal–tvFC coupling from fMRI and pupillometry. **(A)**
Concurrent 7T fMRI and eye tracking were collected during resting state and naturalistic movie
watching. **(B)** tvFC was computed over sliding windows for each pair of brain regions. **(C)** Pupil
diameter was preprocessed to obtain a continuous arousal time series. Arousal–tvFC coupling for
each connection was then defined as the Pearson correlation between its windowed FC time
course and the corresponding arousal fluctuations. **(D)** These edgewise coupling values yielded a
dense arousal–tvFC coupling matrix per run, which served as the input for analyses of community
structure, hemispheric asymmetry, and cross-paradigm consistency. **(E)** The connections were
categorized based on the hemispheres of the connected regions: Left-Left (LL) and Right-Right
(RR) represent intra-hemispheric connections; Left-Right (LR) and Right-Left (RL) represent
inter-hemispheric connections. This classification enabled hemisphere-specific analyses of
arousal-tvFC coupling.

While our primary inferential analyses were conducted at the run level to leverage the
high-density sampling of the HCP 7T dataset, we further validated the robustness of
these findings using participant-level statistical summaries and resampling to account

for within-participant dependencies (see Figure. S1-S2 in Supplementary Material).

Arousal–tvFC coupling reveals seven distinct connectivity communities

For each functional connection, we quantified how strongly its time-varying
connectivity covaried with moment-to-moment arousal, producing an edgewise
arousal–tvFC coupling matrix per run. Across all participants and runs, these coupling
profiles showed clear structure: connections did not exhibit uniform arousal
sensitivity but instead formed separable groups.

Unsupervised clustering of edgewise coupling patterns identified seven stable
connectivity communities (Fig. 2A). This solution was consistently favored across a
broad range of cluster numbers, indicating that arousal–tvFC coupling is inherently
low-dimensional. The seven communities captured distinct patterns of arousal
sensitivity across the connectome. Projecting each community into canonical network
pairs space (Thomas Yeo et al., 2011) showed that these communities were not
random mixtures of connections. Instead, each displayed a characteristic distribution
across network pairs (Fig. 2B). Some communities were enriched in network pairs
linking heteromodal and unimodal systems (Mesulam, 1998), while others were
dominated by heteromodal–heteromodal (H-H) or unimodal–unimodal (U-U) network
pairs. Community participation entropy further highlighted this structure:
unimodal–unimodal connections showed low entropy, indicating a restricted
participation in a few communities, whereas H-H and heteromodal–unimodal (H-U)
connections showed significantly higher entropy, indicating more diverse engagement
across communities ($F(2,88)=12.24$, $p = 4.38\times10^{-6}$).

Together, these results demonstrate that arousal does not uniformly modulate the
connectome but instead engages a small number of organized connectivity

communities, each with distinct network-level compositions.

The robustness of the seven-community architecture was cross-validated using both

split-half and participant-level resampling strategies. As shown in **Figure S1**, both

approaches yielded high alignment accuracy and consistently high Dice coefficients

across all communities, confirming that the identified clusters are not artifacts of

specific data partitioning.

Figure 2. **Arousal–tvFC coupling partitions the connectome into seven distinct connectivity**
communities. (A) Unsupervised clustering of edgewise arousal–tvFC coupling identified seven
stable communities, demonstrating that arousal-linked modulation is organized into
low-dimensional structure rather than uniformly distributed across connections. **(B)** Mapping
communities onto network-pair space revealed distinct and reproducible composition profiles
(upper panel), with some communities dominated by heteromodal interactions and others enriched
in H–U or U–U network pairs (bottom panel). **(C)** Community participation entropy varied
systematically across connection modal types: U–U network pairs showed lower entropy,
indicating concentrated engagement in a small subset of communities, whereas H-H and H–U
network pairs exhibited significantly higher entropy (F(2,88)=12.24, p = 4.38×10^{-6}), reflecting

broader distribution across communities. **(D)** Dominant-community assignments confirmed this
organization, showing that heteromodal interactions load onto multiple arousal-sensitive
communities, whereas U–U network pairs show more restricted community involvement.

Arousal-modulated community architecture exhibit systematic hemispheric asymmetry

We next asked whether these arousal-modulated communities express hemispheric

biases. Using integration and segregation indices derived from LL, RR, LR, and RL

edge categories, we quantified, for each community, whether arousal preferentially

modulated intra- versus inter-hemispheric connectivity and whether these effects

favored one hemisphere.

At the network-pair level, several pairs showed significant lateralization compared

with a spatial permutation null model (Fig. 3A–B). Importantly, lateralization was not

global: only specific network pairs within communities exhibited robust hemispheric

biases, while others remained symmetric. Some communities showed rightward

integration, indicating stronger arousal-related modulation of network pairs within the

right hemisphere or between the right hemisphere and the rest of the brain, whereas

others showed leftward segregation, reflecting preferential influence on

within-left-hemisphere interactions.

Figure 3. Arousal-modulated communities architecture exhibit community-specific hemispheric asymmetry. (A) Integration indices for each network pair revealed significant leftward or rightward deviations from a spatial permutation null, indicating that arousal differentially modulates between- and within-hemisphere interactions for specific network pairs. **(B)** Segregation indices identified network pairs showing hemisphere-specific strengthening of within-hemisphere connectivity, further demonstrating that lateralization is localized rather than global. **(C–D)** Community-averaged integration and segregation values showed that hemispheric biases vary across communities and are not determined solely by connection modal type, underscoring the community-specific nature of the asymmetry. **(E)** Aggregating all significantly lateralized network pairs, each community exhibited a distinct integration–segregation profile, revealing unique hemispheric signatures across communities. **(F)** Low similarity among communities' integration and segregation patterns confirmed that arousal imposes multiple, community-specific forms of hemispheric asymmetry, rather than a single unified left- or right-dominant pattern.

Averaging indices across connection modal types reinforced this community-specific patterning (Fig. 3C–D). Notably, identical connection modal types (e.g., H-H) could show leftward bias in one community but rightward bias in another, demonstrating that lateralization is tied to the community architecture rather than to the canonical network pair class. Aggregating all significantly lateralized network pairs, each

community expressed a unique lateralization signature (Fig. 3E), and there was no

clear cluster of lateralization patterns among communities (Fig. 3F), indicating

heterogeneous, rather than unified, hemispheric influences of arousal on FC. Instead,

arousal imprints distinct left–right biases across different communities.

The robustness of these hemispheric lateralization of community architecture was

further supported by the aforementioned resampling validations (Fig. S2). For the

mean integration and segregation indices across network pairs, the empirical values

consistently fell within the center of the resampling distributions. Specifically, for the

network-pair specific lateralization signatures, the directional biases observed in each

community remained stable during resampling, with empirical values largely aligned

with the center of their respective distributions. This high degree of consistency

confirms that the reported community-specific asymmetry is a stable and

representative feature of arousal-modulated organization, ensuring that our findings

are not skewed by specific sample compositions or outliers.

211 Gradients of community affiliation entropy and the centripetal
212 lateralization architecture of community affiliation

Having established that arousal-modulated functional connections can be organized

into modular communities with distinct hemispheric lateralization at the network level,

we next explored the nodal-level mapping of these communities and the substantial

variability in how regions participate in arousal-modulated communities. For each

region, we computed its community affiliation for each brain region, defined as the

proportion of edges connected to that region that were assigned to each of the seven

communities, separately for LL, LR, RL, and RR edges (Fig. 4 A–B).

Next, we investigated the nodal-level community participation flexibility by

calculating the community affiliation entropy for each region. Nodal-level affiliation

entropy showed a clear network gradient (Fig. 4C–D). Default modal network (DMN),

frontoparietal network (FPN), and limbic (LIMB) regions exhibited lower entropy,

indicating focused participation in fewer arousal-modulated communities. In contrast,

dorsal attention network (DAN), somatomotor network (SMN), and visual (VIS)

regions showed broader participation across communities (t(798) = -23.81, p =

4.24×10^{-95}).

To evaluate lateralization at the nodal level, we derived metrics of segregation and

integration based on these affiliation profiles. We observed a common organizational

principle across several key communities: while no significant hemispheric bias in

integration was detected in any community (Fig. 4E), Communities 3, 5, 6, and 7

exhibited robust leftward segregation biases (Fig. 4F). This dissociation between

segregation and integration reveals a pronounced centripetal lateralization architecture.

A detailed examination of the regional affiliation profiles indicated a consistent

directional bias in signal flow across these communities: whereas left-hemisphere

projections were predominantly intra-hemispheric (LL > LR), right-hemisphere inputs

were biased contralaterally toward the left side (RL > RR). These findings suggest

that these communities collectively represent a "left-hemisphere centripetal

architecture", where the left hemisphere serves as a preferential convergence of

arousal-modulated signals, preferentially aggregating both ipsilateral and contralateral

inputs.

The robustness of these observations was further supported by the aforementioned

resampling validations (Fig. S3). Regarding the hemispheric lateralization of

affiliation profiles, the directional biases observed across regions and communities

remained highly stable during resampling, with empirical values consistently aligned

with the center of their respective distributions. This high degree of consistency

confirms that the reported centripetal affiliation architecture is a stable and

representative feature of arousal-modulated lateralization, ensuring that our findings

are not skewed by specific sample compositions or outliers.

Figure 4. Characterizing the spatial distribution, entropy, and hemispheric divergence of
regional community affiliation. **(A–B)** Nodal-level community affiliation matrices, computed
separately for LL, LR, RL, and RR edges, showed substantial heterogeneity in how nodes
distribute their arousal–tvFC coupling across the seven communities, with distinct patterns
emerging across canonical networks. For visualization purposes, Figure A is restricted to
displaying values where the proportion exceeds 0.2. **(C–D)** Region-level community affiliation
entropy revealed a systematic network gradient, in which heteromodal systems displayed more
selective participation, whereas unimodal network showed broader, more distributed engagement
across communities (t(798) = -23.81, p = 4.24×10^{-95}) **(E)** No integration bias was detected in any
community. **(F)** Significant leftward segregation biases were identified within specific

communities (communities 3, 5, 6, and 7). These asymmetries were primarily localized in regions
belonging to the DMN, FPN, LIMB, and VAN. The color of each box corresponds to the
community identity. Statistical significance: * p_FDR < 0.01; ** p_FDR < 0.001.

Arousal-tvFC coupling lateralization arises from spatial heterogeneity rather than mean shifts

In the preceding analyses, we mainly focused on the lateralization of the
organizational patterns of the decomposed communities at the network-pairs and
nodal levels. However, how the strength of the arousal–tvFC coupling is spatially
distributed and whether lateralization exists in this distribution remains elusive. We
therefore next examined the spatial distribution of arousal–tvFC coupling strength and
its lateralization properties.

To test whether hemispheric biases reflected simple shifts in mean arousal–tvFC
coupling strength, we compared mean integration and segregation across communities
and whole connectome. Although the inter-community difference for segregation was
statistically significant, the mean difference between communities was very small.
This indicates limited hemispheric imbalance.

Although the mean arousal modulation on FC showed no significant lateralization,
prior results suggested that its spatial pattern is highly community specific. We
therefore hypothesized that the key information lies in the spatial heterogeneity
(distribution gradient) of the modulation strength, not the overall strength. We
therefore quantified spatial heterogeneity by ranking network pairs along a
"lateralization axis" (Yang et al., 2025) and computing the slope of integration or
segregation values along this axis. All communities showed slopes significantly
steeper than the whole-brain baseline (all p_FDR < 0.001; Fig. 5B), demonstrating
that lateralization arises from spatial heterogeneity—not uniform shifts.

Figure 5. Spatial heterogeneity—not mean shifts—drives arousal-modulated hemispheric asymmetry. (A) Mean integration and segregation values showed no consistent hemispheric bias across individual communities or at the whole-brain (WB) level, indicating minimal imbalance in overall modulation strength. **(B)** Spatial heterogeneity, quantified as the slope of integration or segregation values ranked along a lateralization axis, was significantly steeper in every community compared with the whole-brain baseline (all p_FDR < 0.0001). These effects demonstrate that hemispheric asymmetry arises from spatially patterned variation, not from uniform shifts in mean modulation. **(C)** Leave-one-out analyses revealed that this heterogeneity reflects distributed contributions from many network pairs, rather than being driven by a few extreme edges. **(D)** Contribution patterns for integration and segregation were strongly correlated (rho ≈ 0.72, p = 2.35×10⁻²⁶), indicating coordinated spatial organization across metrics. **(E–F)** Similarity in contribution patterns between communities was positively associated with similarity in communities' intrinsic structure (integration: rho ≈ 0.79, p = 1.35×10⁻¹¹; segregation: rho ≈ 0.64, p = 5.82×10⁻⁷), showing that spatial heterogeneity is constrained by each community's underlying connectivity architecture. Statistical note: * p_FDR < 0.01; ** p_FDR < 0.001; *** p_FDR < 0.0001.

Leave-one-out analyses showed that heterogeneity does not depend on a small set of

extreme network pairs. Instead, most pairs contributed modestly, producing broad,

community-specific gradients (Fig. 5C). Contribution patterns for integration and

segregation were highly correlated (rho $\approx$ 0.72, p = 2.35×10^{-26}). Furthermore,

similarity in contribution patterns tracked similarity in intrinsic community structure

(integration: rho $\approx$ 0.79, p = 1.35×10^{-11}; segregation: rho $\approx$ 0.64, p = 5.82×10^{-7}),

indicating that spatial heterogeneity is shaped by underlying connectivity architecture

(Fig. 5E–F).

Thus, arousal-driven lateralization is best understood as structured spatial

heterogeneity within communities, rather than gross hemispheric dominance.

The robustness of these observations was further supported by the aforementioned

resampling validations (Fig. S4). For the mean integration and segregation indices, the

empirical values consistently fell within the center of the resampling distributions,

reinforcing the absence of a uniform hemispheric mean shift across the population.

Conversely, for the spatial heterogeneity metrics, the slopes observed in each

community remained stable during resampling, with the empirical values largely

aligned with the center of their respective distributions. This high degree of

consistency confirms that the reported spatial heterogeneity is a stable and

representative feature of arousal-modulated lateralization, ensuring that our findings

are not skewed by specific sample compositions or outliers.

Community structure and hemispheric asymmetry are preserved during movie watching

325   To assess whether these organizational principles generalize beyond the resting state,

326   we applied the same analytic pipeline to naturalistic movie-watching data from the

327   same participants.

The seven-community structure was highly preserved across paradigms. After aligning communities using Hungarian algorithm (Kuhn, 1955), we found that the overall community structure showed consistent correspondence (average dice ≈ 0.46; Fig. 6A–B). One heteromodal-dominated community (community 6) showed near-perfect correspondence (rho ≈ 0.94), suggesting that its arousal sensitivity is largely independent of external stimulation. Other communities also demonstrated moderate cross-paradigm similarity, with only two showing low similarity—likely reflecting context-dependent modulation under rich sensory input.

Figure 6. Community structure and hemispheric asymmetry of arousal–tvFC coupling are preserved across resting state and movie watching. (A) Communities derived from rest and movie data were aligned using Hungarian algorithm, revealing robust correspondence between paradigms (average dice ≈ 0.46). **(B)** Network-pair composition profiles for each community were strongly correlated across paradigms (mean rho ≈ 0.58), indicating that the modular organization of arousal–tvFC coupling is stable across cognitive contexts.

These findings indicate that the modular, asymmetric organization of arousal–tvFC coupling is not specific to rest but reflects intrinsic principles that persist across cognitive contexts.

Discussion

In this study, we investigated how moment-to-moment fluctuations in arousal shape large-scale functional connectivity in the awake human brain. By combining high-field fMRI with concurrent pupillometry, we quantified arousal–tvFC coupling at the level of individual edges and showed that arousal does not exert a uniform or diffuse influence across the connectome. Instead, it modulates connectivity through a set of seven distinct communities, each defined by characteristic network compositions and hemispheric patterns. These findings demonstrate that fluctuations in arousal, even within stable wakefulness, impose a structured and asymmetric organization on whole-brain functional interactions.

Although arousal is often conceptualized as a global modulatory state, our results show that its impact on tvFC is highly organized. Specifically, we show the presence of low-dimensional, reproducible communities suggests that arousal modulates the connectome through structured spatial patterns rather than homogeneous gain modulation. We hypothesize that this structured macroscopic architecture reflects the differentiated projection patterns of subcortical neuromodulatory systems, such as the locus coeruleus–noradrenergic pathway (Aston-Jones & Cohen, 2005; Jordan, 2024) and thalamus (Magnin et al., 2010; Lewis et al., 2015; Liu et al., 2018). This organized

pattern of modulation further supports the view advocated in recent years that arousal

states should be viewed as possessing spatiotemporal dynamics and regional

complexity (Siclari & Tononi, 2017; Nir & De Lecea, 2023), and is strongly supported

by prior work showing that spontaneous arousal fluctuations influence distributed

cortical responses in selective ways (Reimer et al., 2014; McGinley et al., 2015). The

hierarchical community pattern—where unimodal interactions cluster into fewer

patterns while heteromodal systems participate broadly—is compatible with theories

that place heteromodal cortex at the apex of large-scale integrative gradients

(Mesulam, 1998; Margulies et al., 2016). Together, these observations suggest that the

arousal-sensitive connectome reflects not merely regional susceptibility but an intrinsic

network-level architecture, in which state-dependent modulation is aligned with

established large-scale organizational principles.

A major goal of this work was to determine whether arousal imposes systematic

hemispheric asymmetry on FC. We found clear evidence that it does, but in a

community-specific rather than global manner. Certain communities showed

rightward integration, others leftward segregation, and others no significant

bias—indicating that hemispheric asymmetry emerges from the organization of

arousal-sensitive connectivity motifs rather than a single overarching hemispheric

dominance. This distributed asymmetry resonates with converging evidence across

species: unihemispheric vigilance in birds and marine mammals (Rattenborg et al.,

2000; Lyamin et al., 2016; Mascetti, Gian Gastone, 2016; Reicher et al., 2021; Fenk et

al., 2023; Libourel et al., 2023), asymmetric EEG signatures related to vigilance in

humans (Tamaki et al., 2016), and classic findings of right-lateralized alerting and

reorienting functions (Heilman & Abell, 1980; Sturm & Willmes, 2001; Shulman et al.,

2010; Corbetta & Shulman, 2011). This suggests that hemispheric specialization may

arise from distinct modes of arousal-modulated reconfiguration rather than fixed

structural asymmetries.

Despite the robustness of these hemispheric differences, mean integration and

segregation of arousal-tvFC coupling strength across the entire community or whole

brain showed minimal global lateralization. Instead, arousal-driven asymmetry

manifested as spatially heterogeneous gradients within communities. Arousal does not

uniformly shift connectivity toward one hemisphere; rather, it selectively amplifies

lateralization in specific connectivity motifs. This distributed, gradient-like pattern

complements recent work highlighting macroscale cortical gradients and manifold

structure as fundamental organizational principles (Margulies et al., 2016; Huntenburg

et al., 2018). We further found that communities with similar intrinsic topology

exhibited similar heterogeneity signatures, suggesting that baseline connectivity

architecture constrains how arousal shapes hemispheric interactions. This provides a

mechanistic explanation for why traditional hemisphere-level metrics often obscure

arousal-modulated asymmetries—these effects are expressed not as global biases but

as structured, topology-dependent gradients.

Another important observation is the stability of arousal–tvFC organization across

cognitive contexts. While the overall spatial layout of the seven-community

architecture showed moderate reorganization between resting-state and naturalistic

movie-watching, specific motifs—most notably communities 6 and 7—demonstrated

near-perfect correspondence across paradigms. This robustness aligns with prior

evidence that intrinsic connectivity organization persists across tasks (Cole et al., 2014;

Krienen et al., 2014) and that spontaneous fluctuations in arousal modulate cortical

dynamics even during rich sensory stimulation (Tanner et al., 2023). By demonstrating

that arousal-driven network modulation generalizes across both internally and

externally oriented states, our findings indicate that arousal acts as a stable organizing

axis of large-scale brain communication, rather than merely a background

physiological fluctuation.

Despite the important findings of this study, several limitations should be noted. First,

to ensure a mathematically rigorous assessment of hemispheric asymmetry, our

analysis was restricted to a symmetric cortical parcellation. Consequently, while we

demonstrate that arousal-modulated connectivity follows a structured macroscopic

architecture, we did not explicitly analyze the subcortical nuclei hypothesized to drive

these patterns. We hypothesize that the presence of these low-dimensional cortical

communities reflects coordinated motifs rather than a homogeneous gain modulation,

potentially   mirroring   the   differentiated   projection   patterns   of   subcortical

neuromodulatory systems. For instance, the locus coeruleus–noradrenergic pathway

(Chandler et al., 2014; Schwarz & Luo, 2015) and thalamus (Hwang et al., 2017;

Shine, 2019; Müller et al., 2020; Shine et al., 2023) possess extensive yet non-uniform

projections that may anchor the community-specific and hemispherically asymmetric

patterns observed here. In the absence of direct subcortical-cortical integration in our

current framework, this link remains hypothetical. Future investigations incorporating

high-resolution subcortical data will be essential to empirically bridge the gap

between these large-scale cortical communities and the underlying physiological

scaffolding of the Ascending Reticular Activating System (ARAS). Second, while

pupillometry is a well-established and accessible proxy for central arousal (Joshi &

Gold, 2020), it remains an indirect peripheral marker. Integrating these findings with

concurrent EEG will be essential to provide a more granular, multi-modal

characterization of arousal states. Third, we utilized sliding-window FC to track

time-varying connectivity. Although this is a robust and widely validated technique,

employing alternative dynamic approaches with higher temporal resolution—such as

edge FC (Faskowitz et al., 2020) , multiplication of temporal derivatives (Shine et al.,

2015) or dynamic conditional correlation (Lindquist et al., 2014)—may offer

complementary perspectives on the transient dynamics of arousal-modulated states.

Fourth, the generalizability of our approach to external cohorts warrants caution

regarding pupillary data integrity. In contexts where high-fidelity eye-tracking is

technically demanding—such as in clinical settings involving patients with restricted

compliance or in naturalistic fMRI studies—the prevalence of blink artifacts and

signal dropouts may bias the estimation of arousal-modulated states. Excessive

reliance on data interpolation in such cases could artificially smooth temporal

fluctuations, leading to an overestimation of community stability. Future applications

should therefore prioritize high-frequency sampling and potentially incorporate

multi-modal physiological features (e.g., respiratory or cardiac signals) to

cross-validate arousal dynamics when pupillary data is suboptimal (Meissner et al.,

2023; Bolt et al., 2025; Weijs et al., 2025). Fifth, the findings reported here were

derived exclusively from ultra-high-field (7T) imaging data. The superior BOLD

sensitivity of 7T fMRI was instrumental in resolving the fine-scale community

architecture of arousal–tvFC coupling, which involves subtle signals that may be

challenging to detect at lower field strengths. Given that 3T remains the most

common parameter for neuroimaging research and clinical applications, future

investigations are needed to determine the extent to which these organizational

principles generalize to standard field strength data. Validating these communities in

large-scale 3T datasets will be essential to establish their broader applicability across

different imaging environments. Sixth, our findings were derived using a single

high-resolution cortical parcellation. While the specific choice of atlas can influence

fine-grained regional connectivity, it is important to note that our primary

conclusions—such as hemispheric asymmetries and community-level

preferences—were identified and interpreted at the macroscopic network and system

level. By aggregating signals across broad functional systems, this approach likely

mitigates the dependency on precise regional boundary definitions. Nevertheless,

future studies employing alternative parcellation schemes would be valuable to further

confirm that these organizational principles are not specific to the current atlas but

represent a generalizable feature of the arousal-modulated connectome.

In summary, moment-to-moment fluctuations in arousal modulate functional

connectivity through a small number of structured connectivity communities, each

with distinct hemispheric characteristics. These asymmetries arise not from global

shifts but from spatially heterogeneous gradients embedded within community

structure. The reproducibility of these communities across resting state and

naturalistic stimulation suggests that they reflect stable and intrinsic principles by

which arousal dynamically shapes large-scale brain interactions during wakefulness.

Methods and Materials

Participants and datasets

We used data from the Human Connectome Project (HCP) 7T dataset, which includes

resting-state fMRI and naturalistic movie-watching runs with simultaneous eye

tracking (Van Essen et al., 2013). Across participants, up to four resting runs and four

movie runs were available, each approximately 15–16 minutes in length. All

procedures were approved by the Washington University Institutional Review Board,

and written informed consent was obtained from all participants.

MRI acquisition and preprocessing

All imaging was collected on a Siemens 7T scanner (TR = 1 s, TE = 22.2 ms, voxel

size = 1.6 mm isotropic, multiband factor = 5, 900 volumes/run). We used the HCP

minimally preprocessed data (Glasser et al., 2013), followed by additional denoising:

linear detrending; regression of 24 head-motion parameters; regression of white

matter and CSF signals; band-pass filtering (0.01–0.1 Hz); scrubbing of frames with

FD > 0.2 mm; and interpolation across removed frames.

ROI parcellation

Analyses were performed using a 400-ROI symmetric parcellation with explicitly matched left–right homologues (Yan et al., 2023). Vertexwise BOLD signals were averaged within each ROI to obtain regional time series. All subsequent hemispheric analyses relied on this explicit homotopic structure.

Eye tracking preprocessing

Pupil diameter was extracted from the raw eye-tracking stream and cleaned following established procedures (Gonzalez-Castillo et al., 2022): Removal of samples outside MRI acquisition; detection of blinks and short missing segments (<1 s); linear interpolation across missing segments; removal of brief physiologically implausible excursions (<1 ms within long closures); smoothing with a 200 ms Hanning window; down-sampling to 1 Hz to match the temporal resolution of the fMRI data. The resulting time series served as a continuous arousal index for each run.

Quality control

The final analyzed sample for the resting-state consisted of N = 139 healthy participants (mean age = 29.1±3.5 years, 77 female). Runs were excluded if (a) more than 20% of frames exceeded motion thresholds, (b) eye tracking did not cover the full fMRI time series, or (c) more than 90% of samples were classified as eye closure. After applying these criteria, 485 of the initial 723 scans were retained for analysis. The same quality-control pipeline was applied to the movie-watching dataset, yielding 513 usable scans out of the original 725. After rigorous quality control, 139 participants (485 runs) were retained for the final analysis. Detailed information on data retention and run distribution per participant is summarized in Figure S9.

Time-varying functional connectivity

518   Time-varying FC between each pair of ROIs was estimated using sliding-window

519   correlations: window length: 30 s; step size: 5 s. Within each window, Pearson

520   correlation coefficients were computed and Fisher-z transformed. This procedure

521   yielded a tvFC time series for each edge in each run.

Arousal estimation from pupil size

523   Pupil data were processed using the same sliding-window parameters. The mean pupil

524   size within each window was taken as an index of moment-to-moment arousal level.

Estimating arousal–tvFC coupling

For each functional connection, arousal–tvFC coupling was defined as the Pearson

correlation between its time-varying FC and the pupil-derived arousal fluctuations

across windows. Thus, each run produced a 400×400 symmetric matrix of coupling

values, later vectorized into edgewise features.

These matrices were concatenated across runs to form the dataset used for community

detection and all subsequent analyses.

Community detection on arousal–tvFC coupling

To identify brain regions sharing similar arousal-related modulation profiles, we

performed community detection on the edgewise coupling values. Prior to clustering,

coupling values were z-scored across runs to ensure comparability. In this analytical

framework, brain edges were treated as observations, while the individual runs served

as feature dimensions, effectively representing each edge by its unique across-run

coupling motif.

We employed the k-means clustering algorithm (Euclidean distance) to explore a

range of cluster solutions from K = 2 to 15. To ensure the stability of the results and

avoid local optima, each K was repeated 250 times with random initializations. The

optimal number of clusters was determined by evaluating clustering quality and

reproducibility (e.g., maximizing silhouette stability). It is important to clarify that

"communities" in this context refer to clusters of edges that exhibit similar

arousal-modulation motifs within a high-dimensional feature space, rather than

topological modules typically derived from graph-theoretic algorithms like modularity

maximization (Blondel et al., 2008). This procedure consistently identified seven

distinct communities, each representing an arousal-sensitive connectivity motif.

Mapping communities to networks and computing entropy

Each edge was assigned to one of seven communities. Edges were then mapped to

Yeo's 7 canonical networks (DMN, FPN, LIMB, VAN, DAN, SMN and VIS)

(Thomas Yeo et al., 2011).

For each network pair (i, j), we computed the proportion $p_{i,j,k}$ of its edges

belonging to each community k. Shannon entropy $H_{i,j}$ quantified how broadly a

network pair participated across communities, calculated as:

$$H_{i,j} = -\sum_{k=1}^{7} p_{i,j,k} \log_2\left(p_{i,j,k}\right)$$

556   A higher $H_{i,j}$ indicates a broader, more uniform participation across the seven

557   communities, while a lower $H_{i,j}$ indicates that the edges are primarily concentrated

558   in a few communities.

559   Additionally, the dominant community for each network pair is defined as

$$\mathrm{Dom}(i, j) = \arg\max_{k} p_{i,j,k}$$

Entropy and dominant-community assignments jointly characterized both the diversity and primary affiliation of each network pair within the community structure.

Nodal-level community affiliation entropy A_r quantified how broadly each region r participated across communities. The entropy was calculated using the proportion $q_{r,k}$ of all incident edges of the region r belonging to the community k:

$$A_r = -\sum_{k=1}^{7} q_{r,k} \log_2\left(q_{r,k}\right)$$

Quantifying hemispheric asymmetry: integration and segregation indices

To evaluate hemispheric biases in arousal–tvFC coupling, we categorized all functional edges into four types based on their nodal locations: LL (within left hemisphere), RR (within right hemisphere), LR (left-to-right), and RL (right-to-left). Following established frameworks for lateralization (Gotts et al., 2013), we calculated two complementary indices to capture the nature of this asymmetry.

The integration index provides a measure of the overall hemispheric dominance of arousal-modulated connections. A positive value indicates that arousal-modulated edges are preferentially concentrated in the left hemisphere (encompassing both its intra-hemispheric and commissural connections) relative to the right. It is defined as:

$$Integartion = (LL + LR) - (RR + RL)$$

The segregation index assesses whether arousal preferentially modulates local, intra-hemispheric communication versus long-range, inter-hemispheric communication. A positive value reflects a "segregated" left-hemisphere bias, where arousal strengthens connections within the left hemisphere more than it strengthens its

580 communication with the contralateral hemisphere. It is defined as:

$$Segregation = (LL - LR) - (RR - RL)$$

581 Indices were computed for each network pair, each community (weighted by edge
582 count), and all significantly lateralized subsets. Statistical significance was assessed
583 relative to a null model (see below).

Spatial heterogeneity of lateralization

To quantify the spatial distribution characteristics of the arousal-tvFC coupling
strength features (e.g., integration, segregation) within each community k, we first
projected the edgewise coupling matrix for each participant onto the network-pair
level, following the same procedure described in the previous section. The average
coupling value of each network pair (i,j) within each community k is $C_{i,j,k}$.
Network pairs were then sorted by their lateralization values to define the
"lateralization axis" (Yang et al., 2025).

Spatial heterogeneity was quantified by performing a linear regression of the
network-pair coupling values against their rank along the axis:

$$\bar{C}_{i,j,k} = \beta_k \cdot axis + \varepsilon_{i,j,k} \,,$$

the regression slope β_k indexed the spatial heterogeneity, where a larger $|\beta_k|$
indicates a greater difference in lateralization value within the community.

The influence of each individual network pair (i,j) on the overall spatial
heterogeneity β_k was assessed using a leave-one-out method. The contribution $D_{i,j,k}$
was defined as the difference between the new slope $\bar{\beta}_{i,j,k}$ (after removing the pair)
and the original slope β_k:

$$D_{i,j,k} = \bar{\beta}_{(i,j,k)} - \beta^{(k)}$$

600 The sign of $D_{i,j,k}$ indicates its modulatory role: positive values enhance spatial
601 heterogeneity, whereas negative values reduce it.

Null model for hemispheric asymmetry

To determine whether observed lateralization exceeded chance, we construct a null
model for hemispheric asymmetry.

For each run, ROI indices were permuted identically for rows and columns,
preserving matrix symmetry and degree distribution while disrupting hemispheric
structure. 10,000 permutations were performed. For each iteration, clustering and
asymmetry indices were recomputed. p-values were FDR-corrected across
comparisons.

Cross-paradigm validation using movie watching

To assess the context-independence of arousal–tvFC organization, we applied all
analyses to movie-watching runs. Community structures from rest and movie data
were matched using Hungarian assignment (Kuhn, 1955). The overall similarity
between the community architectures derived from the two paradigms was quantified
by the mean Dice coefficient. Community-level correspondence was quantified by
Spearman correlation between network-pair community profiles across paradigms.

Robustness and validation

To ensure the stability and generalizability of our findings, we performed extensive
robustness analyses across multiple biological scales. We first employed two
complementary resampling strategies—500-iteration split-half reliability tests and

participant-level resampling—with the entire analytical pipeline re-executed for each

iteration. The stability of community partitions was quantified using Dice coefficients,

while the reliability of hemispheric asymmetry indices was assessed by their average

deviation from the full-dataset estimates. Crucially, we further confirmed that these

organizational patterns were not driven by non-neural confounds, as the identified

community architecture and lateralization remained highly stable even after explicitly

controlling for head motion and the global signal in the arousal–tvFC coupling model.

A series of sensitivity analyses regarding sliding-window parameters, temporal lags,

and alternative pupillometry preprocessing pipelines further supported the robustness

of our results. Detailed procedures and supporting results for these validations are

provided in Supplementary Methods S1–S6 and Figures S1–S9.

632 Data availability

633 Raw and preprocessed HCP data can be accessed at https://db.humanconnectome.org/.

634 Source data to replicate the results of the study are openly available at

635 https://github.com/kongxy6478/Arousal-modulates-functional-connectivity.

636 Code availability

637 All analyses were implemented in Python (NumPy, SciPy, scikit-learn) using custom

638 scripts. Visualization was performed with Matplotlib, Seaborn, and Surfplot.

639 Computer codes used to calculate the communities, analyse results and reproduce the

640 figures of the study are openly available at

641 https://github.com/kongxy6478/Arousal-modulates-functional-connectivity.

643 Acknowledgment

This work is supported by the National Natural Science Foundation of China (Nos. T2325006, 82021004), STI 2030-Major Projects (Nos.2021ZD0201701, 2021ZD0200500); the Fundamental Research Funds for the Central Universities (No. 2233200020).

Reference

Aston-Jones, G., & Cohen, J. D. (2005). AN INTEGRATIVE THEORY OF LOCUS COERULEUS-NOREPINEPHRINE FUNCTION: Adaptive Gain and Optimal Performance. In *Annual Review of Neuroscience* (Vol. 28, Issue Volume 28, 2005, pp. 403–450). Annual Reviews. https://doi.org/10.1146/annurev.neuro.28.061604.135709

Banks, M. I., Krause, B. M., Endemann, C. M., Campbell, D. I., Kovach, C. K., Dyken, M. E., Kawasaki, H., & Nourski, K. V. (2020). Cortical functional connectivity indexes arousal state during sleep and anesthesia. *NeuroImage*, *211*, 116627. https://doi.org/10.1016/j.neuroimage.2020.116627

Blondel, V. D., Guillaume, J.-L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment*, *2008*(10), P10008. https://doi.org/10.1088/1742-5468/2008/10/P10008

Bolt, T., Wang, S., Nomi, J. S., Setton, R., Gold, B. P., deB.Frederick, B., Yeo, B. T. T.,

Chen, J. J., Picchioni, D., Duyn, J. H., Spreng, R. N., Keilholz, S. D., Uddin, L.

Q., & Chang, C. (2025). Autonomic physiological coupling of the global fMRI

signal.   *Nature   Neuroscience*,   *28*(6),   1327–1335.

https://doi.org/10.1038/s41593-025-01945-y

Breeden, A. L., Siegle, G. J., Norr, M. E., Gordon, E. M., & Vaidya, C. J. (2017).

Coupling between spontaneous pupillary fluctuations and brain activity relates

to inattentiveness. *European Journal of Neuroscience*, *45*(2), 260–266.

https://doi.org/10.1111/ejn.13424

Chandler, D. J., Gao, W.-J., & Waterhouse, B. D. (2014). Heterogeneous organization

of the locus coeruleus projections to prefrontal and motor cortices. *Proceedings*

of   the   National   Academy   of   Sciences,   *111*(18),   6816–6821.

https://doi.org/10.1073/pnas.1320827111

Chow, H. M., Horovitz, S. G., Carr, W. S., Picchioni, D., Coddington, N., Fukunaga, M.,

Xu, Y., Balkin, T. J., Duyn, J. H., & Braun, A. R. (2013). Rhythmic alternating

patterns of brain activity distinguish rapid eye movement sleep from other states

of consciousness. *Proceedings of the National Academy of Sciences*, *110*(25),

10300–10305. https://doi.org/10.1073/pnas.1217691110

Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014).

Intrinsic and Task-Evoked Network Architectures of the Human Brain. *Neuron*,

83(1), 238–251. https://doi.org/10.1016/j.neuron.2014.05.014

Corbetta, M., & Shulman, G. L. (2011). Spatial Neglect and Attention Networks. In

Annual Review of Neuroscience (Vol. 34, Issue Volume 34, 2011, pp. 569–599).

Annual Reviews. https://doi.org/10.1146/annurev-neuro-061010-113731

Damaraju, E., Tagliazucchi, E., Laufs, H., & Calhoun, V. D. (2020). Connectivity

dynamics from wakefulness to sleep. *NeuroImage*, *220*, 117047.

https://doi.org/10.1016/j.neuroimage.2020.117047

Demertzi, A., Tagliazucchi, E., Dehaene, S., Deco, G., Barttfeld, P., Raimondo, F.,

Martial, C., Fernández-Espejo, D., Rohaut, B., Voss, H. U., Schiff, N. D., Owen,

A. M., Laureys, S., Naccache, L., & Sitt, J. D. (2019). Human consciousness is

supported by dynamic complex patterns of brain signal coordination. *Science*

Advances, *5*(2), eaat7603. https://doi.org/10.1126/sciadv.aat7603

Faskowitz, J., Esfahlani, F. Z., Jo, Y., Sporns, O., & Betzel, R. F. (2020). Edge-centric

functional network representations of human cerebral cortex reveal overlapping

system-level architecture. *Nature Neuroscience*, *23*(12), 1644–1654.

https://doi.org/10.1038/s41593-020-00719-y

Fenk, L. A., Riquelme, J. L., & Laurent, G. (2023). Interhemispheric competition

during sleep. *Nature*, *616*(7956), 312–318.

https://doi.org/10.1038/s41586-023-05827-w

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson,

J. L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., &

Jenkinson, M. (2013). The minimal preprocessing pipelines for the Human

Connectome Project. *NeuroImage*, *80*, 105–124.

https://doi.org/10.1016/j.neuroimage.2013.04.127

Gonzalez-Castillo, J., Fernandez, I. S., Handwerker, D. A., & Bandettini, P. A. (2022).

Ultra-slow fMRI fluctuations in the fourth ventricle as a marker of drowsiness.

NeuroImage, *259*, 119424. https://doi.org/10.1016/j.neuroimage.2022.119424

Gotts, S. J., Jo, H. J., Wallace, G. L., Saad, Z. S., Cox, R. W., & Martin, A. (2013). Two

distinct forms of functional lateralization in the human brain. *Proceedings of the*

National Academy of Sciences, *110*(36), E3435–E3444.

https://doi.org/10.1073/pnas.1302581110

Heilman, K. M., & Abell, T. V. D. (1980). Right hemisphere dominance for attention:

The mechanism underlying hemispheric asymmetries of inattention (neglect).

Neurology, *30*(3), 327–327. https://doi.org/10.1212/WNL.30.3.327

Huang, Z., Zhang, J., Wu, J., Mashour, G. A., & Hudetz, A. G. (2020). Temporal circuit

of macroscale dynamic brain activity supports human consciousness. *Science*

Advances, *6*(11), eaaz0087. https://doi.org/10.1126/sciadv.aaz0087

Huntenburg, J. M., Bazin, P. L., & Margulies, D. S. (2018). Large-Scale Gradients in

Human Cortical Organization. *Trends Cogn Sci*, *22*(1), 21–31.

https://doi.org/10/gcr9h4

Hwang, K., Bertolero, M. A., Liu, W. B., & D'Esposito, M. (2017). The Human

Thalamus Is an Integrative Hub for Functional Brain Networks. *The Journal of*

Neuroscience, *37*(23), 5594–5607.

https://doi.org/10.1523/JNEUROSCI.0067-17.2017

Jang, H., Mashour, G. A., Hudetz, A. G., & Huang, Z. (2024). Measuring the dynamic

balance of integration and segregation underlying consciousness, anesthesia,

and sleep in humans. *Nature Communications*, *15*(1), 9164.

https://doi.org/10.1038/s41467-024-53299-x

Jordan, R. (2024). The locus coeruleus as a global model failure system. *Trends in*

Neurosciences, *47*(2), 92–105. https://doi.org/10.1016/j.tins.2023.11.006

Joshi, S., & Gold, J. I. (2020). Pupil Size as a Window on Neural Substrates of

Cognition. *Trends in Cognitive Sciences*, *24*(6), 466–480.

https://doi.org/10.1016/j.tics.2020.03.005

Krienen, F. M., Yeo, B. T. T., & Buckner, R. L. (2014). Reconfigurable task-dependent

functional coupling modes cluster around a core functional architecture. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *369*(1653), 20130526. https://doi.org/10.1098/rstb.2013.0526

Kuhn, H. W. (1955). The Hungarian method for the assignment problem. *Naval Research Logistics Quarterly*, *2*(1–2), 83–97. https://doi.org/10.1002/nav.3800020109

Lewis, L. D., Voigts, J., Flores, F. J., Schmitt, L. I., Wilson, M. A., Halassa, M. M., & Brown, E. N. (2015). Thalamic reticular nucleus induces fast and local modulation of arousal state. *eLife*, *4*, e08760. https://doi.org/10.7554/eLife.08760

Libourel, P.-A., Lee, W. Y., Achin, I., Chung, H., Kim, J., Massot, B., & Rattenborg, N. C. (2023). Nesting chinstrap penguins accrue large quantities of sleep through seconds-long microsleeps. *Science*, *382*(6674), 1026–1031. https://doi.org/10.1126/science.adh0771

Lindquist, M. A., Xu, Y., Nebel, M. B., & Caffo, B. S. (2014). Evaluating dynamic bivariate correlations in resting-state fMRI: A comparison study and a new approach. *NeuroImage*, *101*, 531–546. https://doi.org/10.1016/j.neuroimage.2014.06.052

Liu, X., De Zwart, J. A., Schölvinck, M. L., Chang, C., Ye, F. Q., Leopold, D. A., & Duyn, J. H. (2018). Subcortical evidence for a contribution of arousal to fMRI studies of brain activity. *Nature Communications*, *9*(1), 395. https://doi.org/10.1038/s41467-017-02815-3

Lloyd, B., De Voogd, L. D., Mäki-Marttunen, V., & Nieuwenhuis, S. (2023). Pupil size reflects activation of subcortical ascending arousal system nuclei during rest. *eLife*, *12*, e84822. https://doi.org/10.7554/eLife.84822

Lyamin, O. I., Lapierre, J. L., Kosenko, P. O., Kodama, T., Bhagwandin, A., Korneva, S. M., Peever, J. H., Mukhametov, L. M., & Siegel, J. M. (2016). Monoamine Release during Unihemispheric Sleep and Unihemispheric Waking in the Fur Seal. *Sleep*, *39*(3), 625–636. https://doi.org/10.5665/sleep.5540

Magnin, M., Rey, M., Bastuji, H., Guillemant, P., Mauguière, F., & Garcia-Larrea, L. (2010). Thalamic deactivation at sleep onset precedes that of the cerebral cortex in humans. *Proceedings of the National Academy of Sciences*, *107*(8), 3829–3833. https://doi.org/10.1073/pnas.0909710107

Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., Bezgin, G., Eickhoff, S. B., Castellanos, F. X., Petrides, M., Jefferies, E., & Smallwood, J. (2016). Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proc Natl Acad Sci U S A,*

20191114, *113*(44), 12574–12579. https://doi.org/10/f89vsk

Mascetti, Gian Gastone, G. G. (2016). Unihemispheric sleep and asymmetrical sleep:

Behavioral, neurophysiological, and functional perspectives. *Nature and*

Science of Sleep, *Volume 8*, 221–238. https://doi.org/10.2147/NSS.S71970

McGinley, M. J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C. R.,

Tolias, A. S., Cardin, J. A., & McCormick, D. A. (2015). Waking State: Rapid

Variations Modulate Neural and Behavioral Responses. *Neuron*, *87*(6),

1143–1161. https://doi.org/10.1016/j.neuron.2015.09.012

Meissner, S. N., Bächinger, M., Kikkert, S., Imhof, J., Missura, S., Carro Dominguez,

M., & Wenderoth, N. (2023). Self-regulating arousal via pupil-based

biofeedback. *Nature Human Behaviour*, *8*(1), 43–62.

https://doi.org/10.1038/s41562-023-01729-z

Mesulam, M. (1998). From sensation to cognition. *Brain*, *121*(6), 1013–1052.

https://doi.org/10.1093/brain/121.6.1013

Müller, E. J., Munn, B., Hearne, L. J., Smith, J. B., Fulcher, B., Arnatkevičiūtė, A.,

Lurie, D. J., Cocchi, L., & Shine, J. M. (2020). Core and matrix thalamic

sub-populations relate to spatio-temporal cortical connectivity gradients.

NeuroImage, *222*, 117224. https://doi.org/10.1016/j.neuroimage.2020.117224

Nir, Y., & De Lecea, L. (2023). Sleep and vigilance states: Embracing spatiotemporal

dynamics. *Neuron*, *111*(13), 1998–2011.

https://doi.org/10.1016/j.neuron.2023.04.012

Podvalny, E., King, L. E., & He, B. J. (2021). Spectral signature and behavioral

consequence of spontaneous shifts of pupil-linked arousal in human. *eLife*, *10*,

e68265. https://doi.org/10.7554/eLife.68265

Rattenborg, N. C., Amlaner, C. J., & Lima, S. L. (2000). Behavioral,

neurophysiological and evolutionary perspectives on unihemispheric sleep.

Neuroscience & Biobehavioral Reviews, *24*(8), 817–842.

https://doi.org/10.1016/S0149-7634(00)00039-7

Reicher, V., Kis, A., Simor, P., Bódizs, R., & Gácsi, M. (2021). Interhemispheric

asymmetry during NREM sleep in the dog. *Scientific Reports*, *11*(1), 18817.

https://doi.org/10.1038/s41598-021-98178-3

Reimer, J., Froudarakis, E., Cadwell, C. R., Yatsenko, D., Denfield, G. H., & Tolias, A.

S. (2014). Pupil Fluctuations Track Fast Switching of Cortical States during

Quiet Wakefulness. *Neuron*, *84*(2), 355–362.

https://doi.org/10.1016/j.neuron.2014.09.033

Schneider, M., Hathway, P., Leuchs, L., Sämann, P. G., Czisch, M., & Spoormaker, V. I.

(2016). Spontaneous pupil dilations during the resting state are associated with

activation of the salience network. *NeuroImage*, *139*, 189–201.

https://doi.org/10.1016/j.neuroimage.2016.06.011

Schwarz, L. A., & Luo, L. (2015). Organization of the Locus

Coeruleus-Norepinephrine System. *Current Biology*, *25*(21), R1051–R1056.

https://doi.org/10.1016/j.cub.2015.09.039

Shine, J. M. (2019). Neuromodulatory Influences on Integration and Segregation in the

Brain. *Trends in Cognitive Sciences*, *23*(7), 572–583.

https://doi.org/10.1016/j.tics.2019.04.002

Shine, J. M., Koyejo, O., Bell, P. T., Gorgolewski, K. J., Gilat, M., & Poldrack, R. A.

(2015). Estimation of dynamic functional connectivity using Multiplication of

Temporal Derivatives. *NeuroImage*, *122*, 399–407.

https://doi.org/10.1016/j.neuroimage.2015.07.064

Shine, J. M., Lewis, L. D., Garrett, D. D., & Hwang, K. (2023). The impact of the

human thalamus on brain-wide information processing. *Nature Reviews*

Neuroscience, *24*(7), 416–430. https://doi.org/10.1038/s41583-023-00701-0

Shulman, G. L., Pope, D. L. W., Astafiev, S. V., McAvoy, M. P., Snyder, A. Z., &

Corbetta, M. (2010). Right Hemisphere Dominance during Spatial Selective

Attention and Target Detection Occurs Outside the Dorsal Frontoparietal

Network. *The Journal of Neuroscience*, *30*(10), 3640–3651.

https://doi.org/10.1523/JNEUROSCI.4085-09.2010

Siclari, F., & Tononi, G. (2017). Local aspects of sleep and wakefulness. *Current*

Opinion in Neurobiology, *44*, 222–227.

https://doi.org/10.1016/j.conb.2017.05.008

Sobczak, F., Pais-Roldán, P., Takahashi, K., & Yu, X. (2021). Decoding the brain

state-dependent relationship between pupil dynamics and resting state fMRI

signal fluctuation. *eLife*, *10*, e68980. https://doi.org/10.7554/eLife.68980

Sturm, W., & Willmes, K. (2001). On the Functional Neuroanatomy of Intrinsic and

Phasic Alertness. *NeuroImage*, *14*(1), S76–S84.

https://doi.org/10.1006/nimg.2001.0839

Tagliazucchi, E., Roseman, L., Kaelen, M., Orban, C., Muthukumaraswamy, S. D.,

Murphy, K., Laufs, H., Leech, R., McGonigle, J., Crossley, N., Bullmore, E.,

Williams, T., Bolstridge, M., Feilding, A., Nutt, D. J., & Carhart-Harris, R.

(2016). Increased Global Functional Connectivity Correlates with

LSD-Induced Ego Dissolution. *Current Biology*, *26*(8), 1043–1050.

https://doi.org/10.1016/j.cub.2016.02.010

Tamaki, M., Bang, J. W., Watanabe, T., & Sasaki, Y. (2016). Night Watch in One Brain

Hemisphere during Sleep Associated with the First-Night Effect in Humans.

Current Biology, *26*(9), 1190–1194. https://doi.org/10.1016/j.cub.2016.02.063

Tanner, J. C., Faskowitz, J., Byrge, L., Kennedy, D. P., Sporns, O., & Betzel, R. F.

(2023). Synchronous high-amplitude co-fluctuations of functional brain

networks during movie-watching. *Imaging Neuroscience*, *1*, imag–1–00026.

https://doi.org/10.1162/imag_a_00026

Thomas Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D.,

Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R.,

Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human

cerebral cortex estimated by intrinsic functional connectivity. *Journal of*

Neurophysiology, *106*(3), 1125–1165. (21653723).

https://doi.org/10.1152/jn.00338.2011

Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., Ugurbil, K.,

& W. U-Minn HCP Consortium. (2013). The WU-Minn Human Connectome

Project: An overview. *NeuroImage*, *80*, 62–79.

https://doi.org/10.1016/j.neuroimage.2013.05.041

Weijs, M. L., Missura, S., Potok-Szybińska, W., Bächinger, M., Badii, B.,

Carro-Domínguez, M., Wenderoth, N., & Meissner, S. N. (2025). Modulating

cortical excitability and cortical arousal by pupil self-regulation. *Nature*

Communications, *16*(1), 4552. https://doi.org/10.1038/s41467-025-59837-5

Yang, H., Wu, G., Li, Y., Xu, X., Cong, J., Xu, H., Ma, Y., Li, Y., Chen, R., Pines, A.,

Xu, T., Sydnor, V. J., Satterthwaite, T. D., & Cui, Z. (2025). Connectional axis

of individual functional variability: Patterns, structural correlates, and relevance

for development and cognition. *Proceedings of the National Academy of*

Sciences, *122*(12), e2420228122. https://doi.org/10.1073/pnas.2420228122

Yellin, D., Berkovich-Ohana, A., & Malach, R. (2015). Coupling between pupil

fluctuations and resting-state fMRI uncovers a slow build-up of antagonistic

responses in the human cortex. *NeuroImage*, *106*, 414–427.

https://doi.org/10.1016/j.neuroimage.2014.11.034