De novo design of metalloproteases for targeted amyloid-β cleavage

Yannan Qu [1, 2, †], Chentong Wang [1, 2, †], Hongli Zhu[1,2], Yanjun Wang[1,2], Longxing Cao[1,2*]

1. State Key Laboratory of Gene Expression, School of Life Sciences, Westlake University, Hangzhou 310024, China.

2. Artificial Intelligence Drug Design Core Laboratory, Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, Zhejiang, 310024, China.

*Corresponding author. E-mail: caolongxing@westlake.edu.cn

† These authors are co-first authors.

Abstract

De novo protein design has not yet achieved the creation of proteases capable of selectively cleaving any desired peptide bond within a native protein with high precision. Here, we report the use of the flow-based generative model Proteus2 to design metalloproteases by generating enzyme–substrate complexes conditioned on a target peptide sequence and a predefined catalytic motif. Our approach employs a two-step encapsulation strategy to create clamp-like metalloproteases that bind the target peptide in a manner that maximizes substrate sequence specificity. The generative process simultaneously optimizes the precise positioning of the target peptide bond in a catalytically competent configuration and accurately scaffolds the transition state catalytic residues—both essential for specific and efficient catalysis. Using this strategy, we designed zinc metalloproteases targeting three distinct cleavage sites within the aggregation-prone regions of amyloid-β (Aβ), a key pathogenic factor in Alzheimer's disease. Experimental characterization validated five enzymes, each capable of precise cleavage at the intended sites with high specificity and minimal or undetectable activity on non-cognate substrates. On average, these enzymes accelerated peptide bond hydrolysis by more than 10^7-fold relative to the uncatalyzed reaction , and enabled efficient digestion of the Aβ peptide into smaller segments when enzymes targeting different sites were combined. Cryo-EM structures of three designed enzymes in complex with Aβ peptide, each targeting a distinct cleavage site, revealed close agreement with the design models. Together, these results demonstrate the potential of sequence-guided, generative approaches for developing programmable, sequence-specific proteolysis and lay the groundwork for future applications in basic research and therapeutic development.

Introduction

Proteases are among biology's most powerful regulators[1,2]. By cutting specific peptide bonds they activate hormones[3], terminate signalling cascades[4], remodel extracellular matrices[5], and control protein homostasis[1]. Their catalytic activity also underpins a wide spectrum of industrial and medical applications—from laundry detergents and food processing to wound debridement and thrombolytic therapy[6,7]. Yet the utility of natural proteases is circumscribed by their hard-wired substrate preferences[8,9]. An enduring objective is therefore to create "molecular scalpels" whose cleavage site can be programmed at will, enabling precise dissection of cellular pathways and highly selective therapeutic intervention[10,11]. Despite decades of effort, very few platform delivers custom proteases that combine robust catalytic efficiency with single-bond specificity[12-14].

Metalloproteases are prime candidates for such re-engineering, and their catalytic chemistry has been well studied[15-17]. In a typical zinc metalloprotease, the zinc ion is coordinated by three amino-acid ligands and positioned adjacent to a general base that activates a water molecule for nucleophilic attack on the peptide bond, while a neighbouring residue stabilises the tetrahedral oxyanion intermediate[17] (**Fig. 1a,b**). This minimalist engine is embedded in diverse scaffolds—including matrix metalloproteinases[18], ADAMs[19], astacins[20], thermolysins[21], and bacterial virulence factors[22]—that collectively drive processes from extracellular-matrix turnover to cytokine maturation. Despite their natural versatility, repurposing these enzymes for novel targets remains challenging. While efforts to reprogram specificity through directed evolution or rational mutagenesis have yielded promising advances[12,13,23], certain structural and functional limitations persist. As a result, repurposed natural metalloproteases have achieved only modest improvements in selectivity and therapeutic potential[24].

De novo protein design holds the promise of sidestepping the intrinsic limitations of natural

enzymes by constructing entirely new scaffolds around a chosen chemical task[25-29]. Recent

breakthroughs in deep-learning based structure generation have indeed delivered artificial

metallohydrolases whose catalytic efficiencies approach those of native enzymes[26,28,30]; however,

these successes have been confined almost exclusively to model reactions such as the hydrolysis

of activated ester bonds in small, hydrophobic substrates—reactions that present relatively low

activation barriers and low specificity demands. Extending this progress to true proteolysis is

vastly more challenging. Amide bonds are orders of magnitude more inert than esters[31,32], and a

protease must not only accelerate their cleavage but also capture a highly flexible, polar peptide

and orient one particular bond in a near-attack conformation. Moreover, practical applications

require a high level of sequence selectivity, such that the designed substrate binding groove

preferentially accommodates the target sequence while minimizing activity toward closely related

peptides. Natural proteases achieve this balance with folds that have been evolutionarily honed for

cognate substrates; even slight structural perturbations can disrupt the sub-angstrom precision of

the catalytic residues[9,33]. Consequently, conventional sequence redesign has limited success in

retargets these enzymes while maintaining native-like catalytic efficiency[33-36]. The central

challenge, therefore, is to design new structures that simultaneously preserve the strict metal-

dependent catalytic geometry, lock the scissile bond into position, and rigorously discriminate

against off-target sequences—an objective that has so far remained out of reach.

Computational strategy for designing sequence specific metalloprotease

We set out to develop a general strategy for the de novo design of sequence-specific

metalloproteases. To establish catalytic motifs for design, we focused on two representative zinc-

dependent endopeptidases: PPEP-1 (PDB: 6r4w) [37] and Fungalysin (PDB: 4k90) [38]. Both enzymes

are secreted by microorganisms and play critical roles in survival or pathogenicity by hydrolyzing

extracellular host or cell-surface proteins. Despite sharing a common catalytic mechanism, these

two enzymes exhibit distinct active site architectures. Notably, their oxyanion hole compositions

differ: PPEP-1 employs a tyrosine, whereas Fungalysin utilizes a histidine residue. Structurally,

PPEP-1 features a broad, clamp-like groove formed by two flexible loops that close over the

substrate, conferring high sequence specificity for cleavage at Pro-Pro amide bonds. To reprogram

this specificity, we developed a strategy that removes the peptide-binding loops from PPEP-1,

while retaining the core protein structure (**Fig. 1c**). The excised regions were then rebuilt, enabling

the design of new peptide-binding interfaces. Proteases designed through this approach are denoted

"PP" (peptide-binding pocket designed protease) followed by a numerical identifier. In contrast,

Fungalysin lacks an extended, pre-formed substrate-binding pocket and instead features an open

substrate binding groove, resulting in a broader substrate specificity for peptide bonds flanked by

hydrophobic side chains. For this scaffold, we extracted only the minimalist catalytic motif—

comprising the zinc-coordinating residues, the general base glutamate, and the oxyanion hole

histidine (**Fig. 1c**). This minimalist approach necessitates de novo generation of the entire

surrounding protein scaffold to accurately position the catalytic motif and design a novel peptide-

binding interface for sequence-specific recognition. Proteases generated using this minimalist

catalytic motif are designated "DP" (de novo protease).

To enable precise co-design of enzyme functionality and substrate specificity, we employed

Proteus2, our generative flow-based model significantly advanced over its predecessor[39]. To

explicitly endow the structural generative process with a deep understanding of substrate sequence

constraints and specificity, we integrated a customized protein language model pre-trained on

UniRef50[40] with an auxiliary structural decoding objective (see Methods). Proteus2 was trained

on experimental structures from the Protein Data Bank[41] and high-confidence AlphaFold2[42]

predictions using a dual-modality motif scaffolding task. In this framework, sequence and structure

conditions are treated as independent masking tracks, leveraging geometry-aware sequence

semantics to control structure generation. This decoupled conditioning strategy empowers the

model to process hybrid inputs comprising structurally defined catalytic motifs and coordinate-

free substrate sequences, thereby generating metalloproteases with stabilized catalytic sites and

precise substrate specificity.

We leveraged Proteus2 to co-generate metalloprotease–substrate complex structures using the

extracted active motif configurations as templates. To determine the catalytically competent

conformation of the substrate relative to the catalytic residues, we used AlphaFold3[43] to model the

substrate peptide configuration within the template protease. The predictions revealed that the

substrate adopted a β-strand conformation in both enzymes (**Extended Data Fig. 1**). From these

predicted structures, we extracted the P2–P1′ backbone atoms of the Fungalysin substrate peptide

(where P1 is the residue immediately preceding, and P1′ immediately following, the cleavage site),

and the P3–P1 backbone atoms of the PPEP-1 substrate. In both cases, these segments form strand

pairing interactions with the enzyme's strand adjacent to the active site, thereby enabling the

construction of a complete target active site. To maximize structural complementarity and

specificity for the substrate peptide, we developed a 'two-step encapsulation' strategy based on

Proteus2 (**Fig. 1c,d**). Unlike conventional one-step generative approaches—which frequently yield

suboptimal packing and limited control over overall enzyme topology—our method proceeds

sequentially. The first step generates the protein backbone surrounding the catalytic glutamate,

stabilizing the general base and achieving partial pre-wrapping of the substrate. In the second step,

the scaffold is constructed around the oxyanion hole to complete substrate peptide encapsulation.

This stepwise process minimizes steric clashes, substantially enhances the packing density at the

enzyme–substrate interface, and consistently generates clamp-like enzymes that precisely grasp

the substrate peptide—thereby improving sequence-specific recognition. Following backbone

generation, protein sequences were optimized using LigandMPNN[44], and the resulting protease–

substrate–zinc complex structures were validated with AlphaFold3 to ensure convergence to the

intended catalytic geometry.

De novo proteases facilitate the direct hydrolysis of Aβ peptide

We selected the amyloid-β (Aβ) peptide as the substrate for design, given its central role in

Alzheimer's disease (AD) pathology, where the aggregation of neurotoxic Aβ plaques is a hallmark

event[45]. The pathological self-assembly of Aβ is primarily driven by intermolecular interactions

within two hydrophobic regions: the central hydrophobic core (residues 17–21) and the C-terminal

segment (residues 30–42) (**Fig. 2a**). These domains act as critical nucleation sites for β-sheet

stacking, facilitating the formation of stable, neurotoxic fibrils[46,47]. Consequently, these regions

represent key therapeutic targets for catalytic interventions aimed at destabilizing amyloid

architecture and mitigating AD progression[48]. To demonstrate the generalizability of our

computational methodology, we selected three distinct cleavage sites within Aβ: one within the

central hydrophobic core and two within the C-terminal hydrophogic region. We employed both

the peptide-binding pocket (PP) and de novo protease (DP) strategies to design metalloproteases

targeting each site. On average, 26 designs were selected from each strategy–substrate pair for

experimental characterization (**Extended Data Table 2**). These designed enzymes were expressed

in *Escherichia coli*, purified using Ni-NTA magnetic beads, with the His-tag subsequently removed

via HRV3C protease. Final purification was performed by size exclusion chromatography. To

assess catalytic activity, we engineered fusion substrates in which the targeted Aβ fragment was

inserted between the mannose-binding protein (MBP) and a small ubiquitin-like modifier (SUMO)

tag. Successful proteolytic cleavage results in two lower-molecular-weight fragments, readily

detectable by SDS-PAGE (**Fig. 2b**). Upon overnight incubation of the designed proteases with

their respective substrates in the presence of zinc, we observed that the DP strategy yielded active

enzymes for all three targeted sites: DP720-S1, DP622-S2, and DP221-S3. In contrast, the PP

strategy generated active designs only for the first site, resulting in PP507-S1 and PP532-S1 (**Fig.**

2c). We attribute the limited success of the PP designs at the remaining sites to steric constraints

imposed by excessive preservation of the template backbone, which may be incompatible with the

sequence and geometry of these cleavage sites. In contrast, the DP strategy allows broader

sampling of the conformational space, facilitating the discovery of active variants. Proteolytic

specificity was confirmed by mass spectrometry analysis of the cleavage products, with observed

molecular weights matching the theoretical predictions and validating site-specific cleavage

(**Extended Data Fig. 2**). Notably, for DP221-S3, in addition to the major mass spectrometric peak

corresponding to the intended cleavage products, a very minor peak—representing approximately

10% of the main product's abundance—was also observed, indicating an alternative cleavage site

shifted by one residue toward the C-terminus (**Extended Data Fig. 2e**). We next characterized

their catalytic efficiency by monitoring the time course of substrate cleavage (**Extended Data Fig.**

3). At a 1:2 enzyme-to-substrate molar ratio, DP622-S2 exhibited the highest activity, achieving

complete substrate degradation within 4 hours (**Extended Data Fig. 3b**). DP720-S1 and PP507-

S1 achieved full digestion in 8 hours (**Extended Data Fig. 3a,c**), while PP532-S1 required nearly

12 hours for complete cleavage (**Extended Data Fig. 3e**). In contrast, DP221-S3 displayed slower

kinetics, cleaving more than half of the substrate after 12 hours (**Extended Data Fig. 3d**).

Collectively, these results indicate our design strategies enables the de novo generation of

sequence-specific metalloproteases capable of targeting disease-relevant sites within Aβ.

Validation of catalytic mechanism

To elucidate the catalytic mechanism and confirm the functional roles of key active site residues,

we conducted a series of site-directed mutagenesis experiments (**Fig. 3a**). For designs generated

by both the DP and PP strategies, the catalytic glutamate and oxyanion hole residues (histidine in

the DP designs and tyrosine in the PP designs) were mutated to alanine (**Fig. 3b**). Additionally, in

the DP designs, we mutated the newly generated second-sphere residues that form hydrogen bonds

with the zinc-coordinating residues, to evaluate their contribution to catalytic activity. Functional

assays revealed that substitution of the catalytic glutamate with alanine universally abolished

proteolytic activity across both DP and PP designs, highlighting its indispensable role as the

general base in catalysis (**Fig. 3b**). Interestingly, in the DP designs, mutation of the oxyanion hole

histidine to alanine had minimal or even positive effects on protease function; notably, the H191A

variant of DP221-S3 exhibited enhanced substrate degradation. By contrast, in the PP designs,

alanine substitution of the oxyanion hole tyrosine resulted in partial reduction of activity. These

findings suggest that the oxyanion stabilization function may be partially compensated by solvent

molecules in the absence of the native residue. Furthermore, mutagenesis of the second

coordination sphere residues in the DP designs underscored their importance in tuning catalytic

efficiency. The D128A mutation in DP720-S1 and the N111A mutation in DP221-S3 each resulted

in a marked reduction in activity. In DP622-S2, single mutations (D126A or Y91F) had minimal

impact on catalytic activity, whereas the double mutant (D126A/Y91F) led to a dramatic decrease

in activity (**Fig. 3b**). These results suggest that second-shell interactions play context-dependent

roles in stabilizing the primary coordination environment and modulating the electrostatic

landscape of the active site. Collectively, these findings support the conclusion that the observed

catalytic activity arises from the designed active sites and highlight the importance of precise active

site architecture for efficient catalysis.

Characterization of the catalytic kinetics

To quantitatively assess the kinetics of our designed proteases, we developed a specialized

fluorescence-based assay. The intrinsic hydrophobicity of the Aβ target sequences posed

significant challenges for synthesizing peptides compatible with standard fluorophore-quencher

systems, due to marked solubility and aggregation issues. As an alternative, we engineered a

FRET-based biosensor by inserting the targeted Aβ sequences directly between the fluorescent

protein pair mTurquoise2 (donor) and mVenus (acceptor) (**Extended Data Table 3**). In this system,

enzymatic cleavage of the substrate separates the fluorescent proteins, resulting in a measurable

decrease in FRET efficiency. Kinetic parameters were determined by fitting initial reaction

velocities to the Michaelis-Menten model (**Fig. 3c**). Analysis revealed that the turnover numbers

(k_{cat}) for our designs were generally on the order of 10^{-4} s^{-1}, with Michaelis constants (K_m) in the

micromolar range. These k_{cat} values represent an average $> 10^7$-fold rate enhancement relative to

the uncatalyzed hydrolysis of peptide bonds[31]. Among the designed proteases, DP622-S2

demonstrated the highest catalytic efficiency, achieving a k_{cat}/K_m of 160.8 $M^{-1}s^{-1}$, driven by a

favorable combination of a high k_{cat} (0.0009 s^{-1}) and a low K_m (5.36 μM). The PP507-S1 and

PP532-S1 exhibited moderate efficiencies of 28.4 and 24.3 $M^{-1}s^{-1}$, respectively, while DP221-S3

and DP720-S1 displayed the lowest efficiencies at 4.2 and 3.8 $M^{-1}s^{-1}$. With the exception of DP720-

S1, these kinetic findings were in close concordance with our time-course degradation assays. The

discrepancy observed for DP720-S1—which showed effective substrate degradation comparable

to PP507-S1 in previous assays but exhibited low catalytic efficiency in the FRET system—may

be attributable to steric hindrance, as the bulky fluorescent proteins could restrict substrate access

to the active site. Notably, the K_m value of our lead design, DP622-S2, closely parallels those of

naturally occurring Aβ-degrading proteases such as Insulin-Degrading Enzyme, which also exhibit

K_m values in the low micromolar range[49]. Although the turnover rates of our designs are lower than

those of natural enzymes, achieving micromolar substrate affinity demonstrates that our

computational strategy can successfully encode robust recognition for disease-relevant targets.

To further elucidate the functional roles of the oxyanion hole and second coordination sphere

residues, we characterized the catalytic kinetics of the corresponding mutants, including single

mutants of DP622-S2 (Y91F, D126A, or H172A), DP221-S3 (H191A), as well as the DP622-S2

double mutant (Y91F/D126A) (**Extended Data Fig. 4**). Intriguingly, all tested single mutations in

the DP622-S2 series (Y91F, D126A, and H172A) (**Extended Data Fig. 4a-c**) showed enhanced

catalytic efficiency (k_{cat}/K_m) compared to the parental design, whereas the double mutant

Y91F/D126A decreased catalytic efficiency (**Extended Data Fig. 4d**), consistent with the SDS-

PAGE results. The observed improvements were predominantly driven by increased turnover

numbers (k_{cat}); notably, the DP622-S2 D126A variant exhibited a threefold increase in k_{cat}

(**Extended Data Fig. 4b**). This suggests that optimizing the catalytic microenvironment can

significantly modulate the activation energy barrier of the rate-limiting step. Most mutations led

to a modest decrease in substrate affinity, as reflected by an increase in K_m; however, the DP622-

S2 Y91F variant maintained a K_m nearly identical to the original, resulting in a twofold

enhancement in overall catalytic efficiency **(Extended Data Fig. 4a)**. The DP221-S3 H191A

mutation had a synergistic effect, increasing k_{cat} while simultaneously decreasing K_m, resulting in

a 2.7-fold improvement in catalytic efficiency **(Extended Data Fig. 4e)**. Together, these findings

indicate that second-shell coordination and oxyanion hole residues have a substantial impact on

catalytic efficiency, largely by modulating k_{cat}—a parameter that remains challenging to accurately

model in current design protocols.

Characterization the substrate specificity

We next assessed the substrate specificity of the designed metalloproteases by evaluating their

cleavage activity against three distinct MBP–Aβ_segment–SUMO fusion constructs (Substrates 1,

2, and 3) **(Fig. 4)**. SDS-PAGE analysis of the cleavaged products revealed that PP507-S1 exhibited

the highest degree of selectivity, catalyzing cleavage exclusively of Substrate 1, with no detectable

activity against the other two substrates **(Fig. 4b)**. In contrast, PP532-S1 displayed broader

substrate tolerance, efficiently processing all three substrates **(Fig. 4c)**, while PP720-S1 showed

measurable, though reduced, activity against Substrate 2 **(Fig. 4a)**. DP622-S2 and DP221-S3

exhibited partial cross-reactivity with each other's substrates, albeit at considerably lower

efficiencies compared to their cognate targets **(Fig. 4d,e)**. We speculate that the observed cross-

reactivity for DP622-S2 and DP221-S3 may be attributed to extensive sequence overlap between

the target sequences and the use of glycine-serine (GS) linkers flanking the inserted Aβ segments,

which may facilitate recognition of similar sequence motifs across different constructs **(Extended**

Data Table 1). Notably, the flanking sequence of the S1 cleavage site contains a high proportion

of bulky aromatic and large hydrophilic residues, whereas the regions flanking the S2 and S3 sites

are primarily composed of small aliphatic amino acids and glycines. This composition may

underlie the high specificity of PP507 and likely prevents the S2/S3 enzymes from cleaving the

S1 substrate due to steric occlusion.

We next evaluated the substrate specificity of the active site mutants (**Extended Data Fig. 5**). The

second-shell and oxyanion hole mutants in DP622 and DP720 exhibited specificity profiles similar

to those of their respective parent designs (**Extended Data Fig. 5a,b**). Notably, the DP221-S3

H191A variant showed a remarkable increase in selectivity, virtually abolishing cleavage activity

toward non-target substrates (**Extended Data Fig. 5c**). However, mass spectrometry analysis of

the cleavage products revealed that the scissile bond shifted almost entirely to the original minor

cleavage site—one residue closer to the C-terminus—compared to the parent design (**Extended**

Data Fig. 6b). In contrast, the DP622-S2 Y91F mutant retained the intended cleavage site,

consistent with the parental enzyme (**Extended Data Fig. 6a**). Nonetheless, the stringent

specificity demonstrated by PP507-S1 underscores the potential of our approach to generate

custom, sequence-specific proteases. We anticipate that future iterations incorporating explicit

negative design strategies during the design and filtering stages will further enhance substrate

specificity.

To further evaluate the cleavage activity of our most effective designs and select variants—

including PP507, DP622 Y91F, and DP221 H191A, we incubated all three enzymes together with

a full-length synthetic Aβ42 peptide. Mass spectrometry analysis revealed a diverse mixture of

peptide species. Notably, alongside with several partially cleaved intermediate fragments, we

successfully detected the expected fully cleaved products (DAEFRHDSGYEVHHQKL,

VFFAEDVGSNKGAIIGL, and VVIA) (**Extended Data Fig. 7**). These results confirm that our de

novo metalloproteases are capable of catalyzing the site-specific hydrolysis of the full-length Aβ42

peptide as designed.

Structural characterization of the protease-substrate complex

To assess the structural accuracy of our computational models, we determined the cryo-EM

structures of PP507-S1, DP622-S2, and DP221-S3 in complex with the Aβ42 peptide (**Fig. 5a**). To

overcome the resolution limitations inherent to the small molecular weight of the designed

proteases, each protease was rigidly fused with a large helical bundle protein generated by Proteus2

(**Extended Data Fig. 8**). Minimal interfacial interactions were computationally introduced to

rigidify the fusion construct, ensuring that neither the overall fold nor the active site architecture

of the protease was perturbed. To capture the pre-catalytic Aβ42-bound state, the catalytic

glutamate in each protease was mutated to glutamine (E to Q). The resulting fusion proteins, with

inactivated active sites, were incubated with the Aβ42 peptide and subjected to cryo-EM structure

determination (**Extended Data Fig. 9**). Focused refinement of the protease region within the

fusion protein yielded structures at resolutions ranging from 2.98 to 3.44 Å (**Extended Data Fig.**

10). The reconstructions revealed clear, continuous density for the Aβ42 substrate, allowing

confident assignment of the peptide backbone and local side chains within the binding cleft (**Fig.**

5b). The resolved peptide sequence confirms that all three proteases specifically recognize the

designed Aβ segment, consistent with their high substrate specificity.

Superimposing the experimentally determined cryo-EM structures onto their respective

computational models revealed a high degree of global structural agreement, with Cα root-mean-

square deviation (RMSD) values ranging from 1.13 to 1.88 Å. This high structural fidelity

underscores the robustness of our flow-based generative strategy in accurately constructing the

overall protein structures (**Fig. 5c**). Furthermore, detailed inspection of the superimposed active

sites confirmed that the designed catalytic geometries were accurately recapitulated in the

experimental structures (**Fig. 5d**). Critical spatial arrangements were evaluated by measuring the

distances between the substrate P1 backbone carbonyl oxygen, the coordinating zinc ion, and the

designed oxyanion hole residue. In PP507-S1 and DP221-S3, we observed local structural

relaxation within the binding pocket. For PP507-S1, the P1 oxygen–zinc distance expanded to 5.1

Å in the cryo-EM structure compared to the designed 2.1 Å, and its distance to the tyrosine

oxyanion hole increased from 2.7 Å to 4.4 Å. Similarly, DP221-S3 exhibited an enlarged P1

oxygen–zinc distance of 3.2 Å (designed 2.3 Å) and a relaxed distance of 4.4 Å to the histidine

oxyanion hole (designed 2.1 Å). In contrast, the active site of DP622-S2 demonstrated remarkable

structural fidelity, maintaining a tightly coordinated P1 oxygen–zinc distance of 2.3 Å (closely

matching the designed 2.1 Å) and faithfully preserving the designed 2.8 Å distance to the histidine

oxyanion hole. The precise geometric pre-organization of the DP622-S2 active site is consistent

with our kinetic observations (**Fig. 3c**), which demonstrate both higher substrate affinity and a

moderately improved turnover rate for DP622-S2 compared to PP507-S1 and DP221-S3.

Collectively, these structural insights, together with biochemical data, underscore that achieving

high precision in active site geometry remains critical for enhancing the efficiency of de novo

designed enzymes.

Discussion

By explicitly integrating target sequence information into the flow-based structure generation process, Proteus2 enables the direct generation of sequence-specific metalloproteases. Our two-step encapsulation strategy facilitates the creation of clamp-like proteases that envelop the substrate peptide, thereby maximizing both specificity and discrimination. The successful design of metalloproteases capable of cleaving three distinct native Aβ peptide sequences extends de novo protein design beyond stable binding, enabling functional proteolysis of physiological targets. The atomic-level agreement between the design models and experimentally determined protease-substrate complex structures further demonstrates that our computational approach can achieve both high structure accuracy and precise substrate discrimination. Our specificity characterization also indicates that, for future enzyme design, careful selection of the target site is critical; regions enriched in distinctive side chains, such as bulky aromatic or highly charged residues, may facilitate the development of enzymes with higher specificity. While a certain degree of nonspecific cleavage was seen in some designes, the clamp-like architecture provides a robust starting point for further sequence redesign or structure-guided refinement to enhance specificity.

Our successful de novo design of metalloproteases, starting from minimalist catalytic motifs, represents a major advance in the de novo enzyme design field by bridging the gap between model reactions and catalysis relevant for therapeutic intervention. On average, our designs accelerate peptide bond cleavage by more than 10^7-fold compared to the uncatalyzed reaction in aqueous solution[31]. Importantly, these metalloproteases exhibit substrate affinities (K_m) comparable to those of natural enzymes, indicating that our design strategies can accommodate the structural and chemical complexity of physiological polypeptides[49]. However, our mutational analysis also

indicates that key microenvironmental factors affecting turnover rates (k_{cat}) remain challenging to

accurately model with current approach. While our structural characterization indicates that

improving the modeling precision of the active site might represent a promising direction for

enhancing catalytic efficiency, further advancements will necessitate careful consideration of

solvent effects, second-shell coordination, and the dynamic nature of the catalytic process to

achieve optimal catalytic proficiency.

The aggregation of Aβ into neurotoxic plaques is a defining event in Alzheimer's disease (AD)

pathology, making the development of molecular tools to counteract Aβ accumulation a high

priority. Current strategies for Aβ clearance, predominantly based on monoclonal antibodies[50], act

by sequestering Aβ for immune-mediated removal rather than direct peptide degradation[51]. In

contrast, catalytic degradation offers a compelling alternative, as a single enzyme can process

multiple Aβ molecules through direct hydrolysis. Although endogenous proteases like neprilysin

and insulin-degrading enzyme can target Aβ[52,53], their broad specificity presents a risk for

undesirable off-target effects when therapeutically enhanced[54,55]. Our work provides a

fundamentally distinct strategy—fully de novo design of programmable, sequence-specific

proteases for Aβ peptide degradation—enabling the selective hydrolysis of pathogenic sequences

and facilitating efficient monomer clearance. By targeting cleavage within the hydrophobic,

aggregation-prone regions of Aβ, these proteases have the potential to inhibit the formation of

toxic fibrils. Notably, the cleavage site for our most specific design, PP507-S1, is only one residue

away from the α-secretase site responsible for the non-pathogenic processing of APP, showcasing

the potential for precision engineering[56]. As de novo designed enzymes can target different sites,

our multi-enzyme cocktail approach may reveal new possibilities for developing therapeutic

interventions for AD.

Overall, our work establishes a robust and generalizable framework for the de novo design of

programmable, sequence-specific metalloproteases with wide-ranging applications in research and

medicine. Our design approach can be readily adapted to other protease classes, such as serine or

cysteine proteases, opening new possibilities for selective protein manipulation and targeted

protein degradation that are unattainable with naturally occurring enzymes. Looking ahead, the

integration of explicit negative design strategies, machine learning-guided optimization, and

advanced transition-state modeling will likely yield next-generation proteases with even higher

specificity and catalytic efficiency. Ultimately, this technology sets the stage for the rational

development of "designer" proteases for a broad spectrum of diseases and empowers both basic

research and therapeutic innovation by providing unprecedented control over protein function in

living systems.

Data availability

The density maps and structure files have been deposited into the Electron Microscopy Data Bank and the RCSB Protein Data Bank with the following accession codes: EMD-69321 and 23WM (PP507 E110Q); EMD-69322 and 23WN (DP622 E96Q); EMD-69323 and 23WO (DP221 E85Q).

Code availability

The Rosetta macromolecular modelling suite (https://www.rosettacommons.org) is freely available to academic and commercial users. Design protocols and analysis scripts used in this paper are available in the Methods and at https://github.com/LongxingLab/metalloprotease. The code for Proteus2 is currently uploaded to Zenodo at xxx.

Acknowledgements

We extend our sincere thanks to the Dang Lab at Westlake University for their valuable guidance and technical expertise in peptide synthesis. We would like to thank the Mass Spectrometry Facility of Westlake University for help in sample analysis; and the Westlake University HPC Center for computation assistance.

Author contributions

Y.Q., C.W. and L.C. designed the research. Y.Q., C.W. and L.C. developed the design pipeline and made the designs. Y.Q., H.Z. and Y.W. pefromed the experiments. All authors analysed data. L.C. supervised research. Y.Q., C.W. and L.C. wrote the manuscript with the input from the other authors. All authors revised the manuscript.

Competing interests

A provisional patent application will be filed prior to publication.

Funding sources

424   This work was supported by grants from the National Key R&D Program of China

425   (2022YFA1303700), the National Natural Science Foundation of China (32370989, 3257120243),

426   The Zhejiang Province Pioneer Plan (2025C01189), and Westlake Education Foundation to L.C.

Fig. 1: Catalytic mechanism of zinc-dependent metalloproteases and two-step de novo design of clamp-like protease scaffolds. a, The catalytic cycle of zinc-dependent metalloproteases involves a precisely orchestrated sequence of molecular events at the active site. The reaction

initiates as the scissile carbonyl oxygen of the substrate coordinates with the catalytic zinc ion, polarizing the carbonyl and increasing its electrophilicity. Concurrently, a conserved glutamate residue acts as a general base, deprotonating a zinc-bound water molecule to trigger nucleophilic attack on the carbonyl carbon. This generates a transient tetrahedral intermediate, with its negatively charged oxyanion stabilized via hydrogen bonds within the oxyanion hole—typically formed by a tyrosine or histidine. The catalytic glutamate then functions as a general acid, transferring a proton to the amide nitrogen of the leaving group, thereby facilitating intermediate collapse and peptide bond cleavage. The hydrolyzed products subsequently dissociate, and a new water molecule occupies the zinc coordination sphere, restoring the enzyme for further catalytic cycles. **b,** Theozyme models for the Zn(II)-activated water nucleophilic attack on the substrate peptide bond. Both panels illustrate the core catalytic components: the central Zn(II) ion coordinated by three endogenous amino acid residues, the catalytic glutamate acting as a general base, the nucleophilic water molecule, and the substrate scissile bond represented by the P1 and P1' residues. **c,** Computational framework for designing substrate-specific metalloproteases. Generation is initialized with either a minimalist catalytic motif (de novo design) or a largely preserved template topology (pocket design). A pretrained protein language model encodes the sequences of the substrate and catalytic residues into latent embeddings. These embeddings, alongside the structural motif, condition the Proteus2 generative model for substrate-sequence-aware protease design via a two-step encapsulation strategy. **d,** Schematic representation of the two-step encapsulation strategy. The scaffold is assembled in a sequential manner to achieve complete substrate encapsulation, ensuring that the clamp precisely engages the target peptide for sequence-specific recognition.

Fig. 2: Experimental validation of designed metalloproteases targeting Aβ. a, Primary

sequence of Aβ42. The complete amino acid sequence of Aβ42 is shown, with two main

hydrophobic domains highlighted in orange and three selected cleavage sites marked in red. **b**,

Schematic of the proteolysis assay. To facilitate expression and protelytic analysis, Aβ segment

was engineered as a fusion protein (MBP-Aβ_segment-SUMO). The diagram illustrates that

specific cleavage within the Aβ sequence by the designed proteases results in separation of the

MBP and SUMO fragments, which are readily detected via molecular weight shifts in SDS-PAGE

experiments. **c**, Design models and proteolytic screening of designed proteases. Design models of

five designed proteases are displayed alongside SDS-PAGE analysis, which confirms the

successful cleavage of the MBP-Aβ-SUMO substrate.

Fig. 3: Mutagenesis and kinetic characterization of the designed metalloproteases. a, Atomic details of the active site architectures in the design models, highlighting the precise spatial arrangement of key catalytic residues. The coordination of the zinc ion by histidines and

glutamates, the positioning of the general base, and the configuration of the oxyanion hole are

illustrated. **b,** Mutational analysis of catalytic residues. SDS-PAGE analysis of various catalytic

residue mutants is shown, verifying the functional roles of the designed active site residues. **c,**

Kinetic characterization of the designed proteases. Proteolytic kinetics were assessed by measuring

initial reaction velocities at varying substrate concentrations. Data were fitted to the Michaelis-

Menten equation to determine key kinetic parameters, including the Michaelis constant (K_m) and

turnover number (k_{cat}). The calculated k_{cat}/K_m values quantify the catalytic efficiency of the de

novo designed enzymes, demonstrating their substrate recognition and cleavage ability. Data

represent mean ± standard deviation (black dots and error bars).

a Cleavage of different substrates by DP720-S1

d Cleavage of different substrates by DP622-S2

b Cleavage of different substrates by PP507-S1

e Cleavage of different substrates by DP221-S3

c Cleavage of different substrates by PP532-S1

481 **Fig. 4: Proteolytic specificity and substrate selectivity of designed proteases.** The substrate

482 recognition profiles of five designed proteases (**a**, DP720-S1; **b**, PP507-S1; **c**, PP532-S1; **d**,

483 DP622-S2; **e**, DP221-S3) were systematically evaluated using the three structurally distinct MBP-

484   Aβ_segment-SUMO fusion constructs (Substrates 1, 2, and 3). The results demonstrate the

485   sequence and site selectivity of each protease toward their corresponding substrates.

a

PP507 E110Q with substrate   DP622 E96Q with substrate   DP221 E85Q with substrate

S1   S2   S3

b

Aβ 15 - 22: Q K L↓V F F A E   Aβ 34 - 41: L M V G G↓V V I   Aβ 29 - 39: G A I I G↓L M V G G V

c

☐ Designed model   ☐ Cryo-EM structure of PP507 E110Q with substrate   ☐ Cryo-EM structure of DP622 E96Q with substrate   ☐ Cryo-EM structure of DP221 E85Q with substrate

RMSD: 1.88 Å   RMSD: 1.13 Å   RMSD: 1.48 Å

d

E110Q   P1   P1′   4.4 Å   2.7 Å   5.1 Å   2.1 Å   Y192

E96Q   P1′   P1   2.8 Å   2.1 Å   2.8 Å   2.3 Å   H172

E85Q   P1′   P1   4.4 Å   3.2 Å   2.1 Å   2.3 Å   H191

Fig. 5: Cryo-EM structure characterization of designed protease in complex with Aβ42. a, Overall cryo-EM density maps for each of the three designed proteases. To capture the pre-catalytic substrate-bound complex, the catalytic glutamate in each design was mutated to

glutamine (E to Q). **b,** Local cryo-EM density maps highlighting the bound Aβ42 substrate. **c,**

Superimposition of the computational design models (grey) and the experimentally determined

cryo-EM structures (colored), with backbone Cα RMSD values indicated. **d,** Close-up views of

the superimposed active sites. Key catalytic geometries are assessed by measuring the distances

between the substrate P1 backbone carbonyl oxygen, the designed oxyanion hole residue, and the

catalytic zinc ion. Distance labels are shown in black for the design models and in red for the cryo-

EM structures.

Reference

1   Lopez-Otin, C. & Bond, J. S. Proteases: multifunctional enzymes in life and disease. *J Biol*
Chem **283**, 30433-30437 (2008). https://doi.org:10.1074/jbc.R800035200
2   Taylor, R. C., Cullen, S. P. & Martin, S. J. Apoptosis: controlled demolition at the cellular
level. *Nat Rev Mol Cell Biol* **9**, 231-241 (2008). https://doi.org:10.1038/nrm2312
3   Seidah, N. G. & Prat, A. The biology and therapeutic targeting of the proprotein
convertases. *Nat Rev Drug Discov* **11**, 367-383 (2012). https://doi.org:10.1038/nrd3699
4   Turk, B. Targeting proteases: successes, failures and future prospects. *Nat Rev Drug Discov*
5, 785-799 (2006). https://doi.org:10.1038/nrd2092
5   Bonnans, C., Chou, J. & Werb, Z. Remodelling the extracellular matrix in development and
disease. *Nat Rev Mol Cell Biol* **15**, 786-801 (2014). https://doi.org:10.1038/nrm3904
6   Kirk, O., Borchert, T. V. & Fuglsang, C. C. Industrial enzyme applications. *Curr Opin*
Biotechnol **13**, 345-351 (2002). https://doi.org:10.1016/s0958-1669(02)00328-2
7   Li, Q., Yi, L., Marek, P. & Iverson, B. L. Commercial proteases: present and future. *FEBS Lett*
587, 1155-1163 (2013). https://doi.org:10.1016/j.febslet.2012.12.019
8   Perona, J. J. & Craik, C. S. Evolutionary divergence of substrate specificity within the
chymotrypsin-like serine protease fold. *J Biol Chem* **272**, 29987-29990 (1997).
https://doi.org:10.1074/jbc.272.48.29987
9   Hedstrom, L. Serine protease mechanism and specificity. *Chem Rev* **102**, 4501-4524 (2002).
https://doi.org:10.1021/cr000033x
10 Craik, C. S., Page, M. J. & Madison, E. L. Proteases as therapeutics. *Biochem J* **435**, 1-16
(2011). https://doi.org:10.1042/BJ20100965
11 Dyer, R. P. & Weiss, G. A. Making the cut with protease engineering. *Cell Chem Biol* **29**,
177-190 (2022). https://doi.org:10.1016/j.chembiol.2021.12.001
12 Blum, T. R. *et al.* Phage-assisted evolution of botulinum neurotoxin proteases with
reprogrammed specificity. *Science* **371**, 803-810 (2021).
https://doi.org:10.1126/science.abf5972
13 Gao, Z. *et al.* A gasdermin-based life-death evolution system for reprogramming protease
specificity. *Nat Chem Biol* (2025). https://doi.org:10.1038/s41589-025-02063-3
14 Dyer, R. P. *et al.* Reengineering the specificity of the highly selective Clostridium botulinum
protease via directed evolution. *Sci Rep* **12**, 9956 (2022). https://doi.org:10.1038/s41598-
022-13617-z
15 Bode, W., Gomis-Ruth, F. X. & Stockler, W. Astacins, serralysins, snake venom and matrix
metalloproteinases exhibit identical zinc-binding environments (HEXXHXXGXXH and Met-
turn) and topologies and should be grouped into a common family, the 'metzincins'. *FEBS*
Lett **331**, 134-140 (1993). https://doi.org:10.1016/0014-5793(93)80312-i
16 Visse, R. & Nagase, H. Matrix metalloproteinases and tissue inhibitors of
metalloproteinases: structure, function, and biochemistry. *Circ Res* **92**, 827-839 (2003).
https://doi.org:10.1161/01.RES.0000070112.80711.3D
17 Spyroulias, G. A., Galanis, A. S., Pairas, G., Manessi-Zoupa, E. & Cordopatis, P. Structural
features of angiotensin-I converting enzyme catalytic sites: conformational studies in
solution, homology models and comparison with other zinc metallopeptidases. *Curr Top*
Med Chem **4**, 403-429 (2004). https://doi.org:10.2174/1568026043451294

18 Nagase, H. & Woessner, J. F., Jr. Matrix metalloproteinases. *J Biol Chem* **274**, 21491-21494
(1999). https://doi.org:10.1074/jbc.274.31.21491

19 Blobel, C. P. ADAMs: key components in EGFR signalling and development. *Nat Rev Mol*
Cell Biol **6**, 32-43 (2005). https://doi.org:10.1038/nrm1548

20 Bond, J. S. & Beynon, R. J. The astacin family of metalloendopeptidases. *Protein Sci* **4**,
1247-1261 (1995). https://doi.org:10.1002/pro.5560040701

21 Tronrud, D. E., Roderick, S. L. & Matthews, B. W. Structural basis for the action of
thermolysin. *Matrix Suppl* **1**, 107-111 (1992).

22 Miyoshi, S. Extracellular proteolytic enzymes produced by human pathogenic vibrio
species. *Front Microbiol* **4**, 339 (2013). https://doi.org:10.3389/fmicb.2013.00339

23 Varadarajan, N., Rodriguez, S., Hwang, B. Y., Georgiou, G. & Iverson, B. L. Highly active and
selective endopeptidases with programmed substrate specificities. *Nat Chem Biol* **4**, 290-
294 (2008). https://doi.org:10.1038/nchembio.80

24 Vandenbroucke, R. E. & Libert, C. Is there new hope for therapeutic matrix
metalloproteinase inhibition? *Nat Rev Drug Discov* **13**, 904-927 (2014).
https://doi.org:10.1038/nrd4390

25 Yeh, A. H. *et al.* De novo design of luciferases using deep learning. *Nature* **614**, 774-780
(2023). https://doi.org:10.1038/s41586-023-05696-3

26 Ahern, W. *et al.* Atom-level enzyme active site scaffolding using RFdiffusion2. *Nat Methods*
(2025). https://doi.org:10.1038/s41592-025-02975-x

27 Braun, M. *et al.* Computational enzyme design by catalytic motif scaffolding. *Nature*
(2025). https://doi.org:10.1038/s41586-025-09747-9

28 Kim, D. *et al.* Computational design of metallohydrolases. *Nature* (2025).
https://doi.org:10.1038/s41586-025-09746-w

29 Listov, D. *et al.* Complete computational design of high-efficiency Kemp elimination
enzymes. *Nature* **643**, 1421-1427 (2025). https://doi.org:10.1038/s41586-025-09136-2

30 Studer, S. *et al.* Evolution of a highly active and enantiospecific metalloenzyme from short
peptides. *Science* **362**, 1285-1288 (2018). https://doi.org:10.1126/science.aau3744

31 Radzicka, A. & Wolfenden, R. Rates of Uncatalyzed Peptide Bond Hydrolysis in Neutral
Solution and the Transition State Affinities of Proteases. *Journal of the American Chemical*
Society **118**, 6105-6109 (1996). https://doi.org:10.1021/ja954077c

32 Kiss, G., Celebi-Olcum, N., Moretti, R., Baker, D. & Houk, K. N. Computational enzyme
design. *Angew Chem Int Ed Engl* **52**, 5700-5725 (2013).
https://doi.org:10.1002/anie.201204077

33 Tokuriki, N. & Tawfik, D. S. Stability effects of mutations and protein evolvability. *Curr Opin*
Struct Biol **19**, 596-604 (2009). https://doi.org:10.1016/j.sbi.2009.08.003

34 Nestl, B. M. & Hauer, B. Engineering of Flexible Loops in Enzymes. *ACS Catalysis* **4**, 3201-
3211 (2014). https://doi.org:10.1021/cs500325p

35 Barrozo, A., Borstnar, R., Marloie, G. & Kamerlin, S. C. Computational protein engineering:
bridging the gap between rational design and laboratory evolution. *Int J Mol Sci* **13**, 12428-
12460 (2012). https://doi.org:10.3390/ijms131012428

36 Chen, C. Y., Georgiev, I., Anderson, A. C. & Donald, B. R. Computational structure-based
redesign of enzyme activity. *Proc Natl Acad Sci U S A* **106**, 3764-3769 (2009).
https://doi.org:10.1073/pnas.0900266106

37 Pichlo, C. *et al.* Molecular determinants of the mechanism and substrate specificity of
Clostridium difficile proline-proline endopeptidase-1. *J Biol Chem* **294**, 11525-11535
(2019). https://doi.org:10.1074/jbc.RA119.009029
38 Fernandez, D., Russi, S., Vendrell, J., Monod, M. & Pallares, I. A functional and structural
study of the major metalloprotease secreted by the pathogenic fungus Aspergillus
fumigatus. *Acta Crystallogr D Biol Crystallogr* **69**, 1946-1957 (2013).
https://doi.org:10.1107/S0907444913017642
39 Wang, C. *et al.* in *Proceedings of the 41st International Conference on Machine Learning*
Vol. 235 (eds Salakhutdinov Ruslan *et al.*) 51376--51395 (PMLR, Proceedings of Machine
Learning Research, 2024).
40 Suzek, B. E. *et al.* UniRef clusters: a comprehensive and scalable alternative for improving
sequence   similarity   searches.   *Bioinformatics*   **31**,   926-932   (2014).
https://doi.org:10.1093/bioinformatics/btu739
41 Berman, H. M. *et al.* The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242 (2000).
https://doi.org:10.1093/nar/28.1.235
42 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2
43 Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org:10.1038/s41586-024-07487-w
44 Dauparas, J. *et al.* Atomic context-conditioned protein sequence design using
LigandMPNN. *Nat Methods* **22**, 717-723 (2025). https://doi.org:10.1038/s41592-025-
02626-1
45 Selkoe, D. J. & Hardy, J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO*
Mol Med **8**, 595-608 (2016). https://doi.org:10.15252/emmm.201606210
46 Tjernberg, L. O. *et al.* Arrest of beta-amyloid fibril formation by a pentapeptide ligand. *J*
Biol Chem **271**, 8545-8548 (1996). https://doi.org:10.1074/jbc.271.15.8545
47 Jarrett, J. T., Berger, E. P. & Lansbury, P. T., Jr. The carboxy terminus of the beta amyloid
protein is critical for the seeding of amyloid formation: implications for the pathogenesis
of   Alzheimer's   disease.   *Biochemistry*   **32**,   4693-4697   (1993).
https://doi.org:10.1021/bi00069a001
48 Saido, T. & Leissring, M. A. Proteolytic degradation of amyloid beta-protein. *Cold Spring*
Harb Perspect Med **2**, a006379 (2012). https://doi.org:10.1101/cshperspect.a006379
49 Mukherjee, A. *et al.* Insulysin hydrolyzes amyloid beta peptides to products that are
neither neurotoxic nor deposit on amyloid plaques. *J Neurosci* **20**, 8745-8749 (2000).
https://doi.org:10.1523/JNEUROSCI.20-23-08745.2000
50 van Dyck, C. H. *et al.* Lecanemab in Early Alzheimer's Disease. *N Engl J Med* **388**, 9-21
(2023). https://doi.org:10.1056/NEJMoa2212948
51 Sevigny, J. *et al.* The antibody aducanumab reduces Abeta plaques in Alzheimer's disease.
Nature **537**, 50-56 (2016). https://doi.org:10.1038/nature19323
52 Qiu, W. Q. *et al.* Insulin-degrading enzyme regulates extracellular levels of amyloid beta-
protein   by   degradation.   *J   Biol   Chem*   **273**,   32730-32738   (1998).
https://doi.org:10.1074/jbc.273.49.32730
53 Iwata, N. *et al.* Metabolic regulation of brain Abeta by neprilysin. *Science* **292**, 1550-1552
(2001). https://doi.org:10.1126/science.1059946

54 Tang, W. J. Targeting Insulin-Degrading Enzyme to Treat Type 2 Diabetes Mellitus. *Trends*
Endocrinol Metab **27**, 24-34 (2016). https://doi.org:10.1016/j.tem.2015.11.003
55 Turner, A. J., Isaac, R. E. & Coates, D. The neprilysin (NEP) family of zinc
metalloendopeptidases: genomics and function. *Bioessays* **23**, 261-269 (2001).
https://doi.org:10.1002/1521-1878(200103)23:3<261::AID-BIES1036>3.0.CO;2-K
56 Esch, F. S. *et al.* Cleavage of amyloid beta peptide during constitutive processing of its
precursor. *Science* **248**, 1122-1124 (1990). https://doi.org:10.1126/science.2111583
636

a

b

Glu P1 P1′ His
H₂O Zn²⁺
His His Glu

Glu P1 P1′ Tyr
H₂O Zn²⁺
His His Glu

c

De novo protease design

protease   substrate
H E   R   E   H   H H Q K L V F F A E

Sequence Encoding

Pretrained Protein Language Model

H E   R   Y   E   H H Q K L V F F A E

Peptide-binding pocket design

Step 1   Step 2

Latent Embedding   Proteus2

d

Step 0: Catalytic pocket motif   Step 1: Generate first jaw   Step 2: Generate second jaw

General base

Oxyanion hole

a

Aβ 42: 1 DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA 42

S1 S3 S2

b

designed protease → substrate (Aβ, MBP, SUMO) → substrate degradation (MBP, SUMO)

c

site 1: HHQKL↓VFFAED

PP507-S1
kDa | PP507 | S1 | PP507 + S1
→ substrate
→ cleavage product
→ protease
→ cleavage product

PP532-S1
kDa | PP532 | S1 | PP532 + S1
→ substrate
→ cleavage product
→ protease
→ cleavage product

DP720-S1
kDa | DP20 | S1 | DP20 + S1
→ substrate
→ cleavage product
→ protease
→ cleavage product

site 2: GLMVGG↓VVIA

DP622-S2
kDa | DP622 | S2 | DP622 + S2
→ substrate
→ cleavage product
→ protease
→ cleavage product

site 3: GAIIG↓LMVGGV

DP221-S3
kDa | DP221 | S3 | DP221 + S3
→ substrate
→ cleavage product
→ protease
→ cleavage product

a

DP720-S1

DP622-S2

DP221-S3

PP507-S1

PP532-S1

b

DP720 + S1

DP622 + S2

DP221 + S3

PP507 + S1   PP532 + S1

c

Michaelis-Menten kinetic analysis of DP720-S1

k_{cat} = 0.0001 ± 0.00002 s^{-1}
K_M = 28.14 ± 9.92 μM
k_{cat}/K_M = 3.77 M^{-1} s^{-1}

Michaelis-Menten kinetic analysis of DP622-S2

k_{cat} = 0.0009 ± 0.00002 s^{-1}
K_M = 5.36 ± 0.3 μM
k_{cat}/K_M = 160.85 M^{-1} s^{-1}

Michaelis-Menten kinetic analysis of DP221-S3

k_{cat} = 0.0001 ± 0.00001 s^{-1}
K_M = 29.88 ± 4.53 μM
k_{cat}/K_M = 4.16 M^{-1} s^{-1}

Michaelis-Menten kinetic analysis of PP507-S1

k_{cat} = 0.0005 ± 0.0001 s^{-1}
K_M = 17.90 ± 3.06 μM
k_{cat}/K_M = 28.41 M^{-1} s^{-1}

Michaelis-Menten kinetic analysis of PP532-S1

k_{cat} = 0.0008 ± 0.0001 s^{-1}
K_M = 33.96 ± 7.05 μM
k_{cat}/K_M = 24.27 M^{-1} s^{-1}

a Cleavage of different substrates by DP720-S1

b Cleavage of different substrates by PP507-S1

c Cleavage of different substrates by PP532-S1

d Cleavage of different substrates by DP622-S2

e Cleavage of different substrates by DP221-S3

a

PP507 E110Q with substrate   DP622 E96Q with substrate   DP221 E85Q with substrate

S1   S2   S3

b

Aβ 15 - 22: QKL V FFAE   Aβ 34 - 41: LMVGG VVI   Aβ 29 - 39: GAIIG LMVGGV

c

☐ Designed model ■ Cryo-EM structure of   ■ Cryo-EM structure of   ■ Cryo-EM structure of
                    PP507 E110Q with substrate   DP622 E96Q with substrate   DP221 E85Q with substrate

RMSD: 1.88 Å   RMSD: 1.13 Å   RMSD: 1.48 Å

d

E110Q   P1 P1′   E96Q   P1′   E85Q   P1′
                                            P1
         4.4 Å   2.8 Å   4.4 Å
    5.1 Å 2.7 Å   2.1 Å 2.8 Å   3.2 Å 2.3 Å
         2.1 Å   Y192   P1 2.3 Å H172   2.3 Å H191