Plant Kelch phosphatases are Ser/Thr phosphatases involved in cell cycle regulation

Felix Rico-Resendiz[1,*], Oded Pri-Tal[1,*], Pierre Raia[1,*], Andrea Moretti[1,*,$], Houming Chen[1], Jun Yu[2],

Larissa Broger[1], Christelle Fuchs[3], Ludwig A. Hothorn[4], Sylvain Loubéry[3], Michael Hothorn[1,#]

Affiliations

[1]Structural Plant Biology Laboratory, Department of Plant Sciences, University of Geneva, 1211

Geneva, Switzerland.

[2]Department of Molecular and Cellular Biology, University of Geneva, 1211, Geneva, Switzerland.

[3]Plant Imaging Unit, Department of Plant Sciences, University of Geneva, 1211 Geneva,

Switzerland.

[4]retired from Institute of Biostatistics, Leibniz University, 30419, Hannover, Germany.

[$]current address: CSL Behring AG, 3014 Bern, Switzerland.

*These authors contributed equally to this work.

[#]Corresponding author Michael Hothorn **Email:** <u>michael.hothorn@unige.ch</u>

Author Contributions: M.H. and A.M. designed the study. A.M. crystallized proteins, A.M. and

M.H. solved and refined structures, P.R. and M.H. performed structural analyses, P.R. and A.M.

performed phosphatase and kinase assays, J.Y. purified the CDK1 – cyclin B1 – CKS1 complex,

O.P-T., L.B. and F.R. generated transgenic Arabiopsis lines, O.P-T., F.R. and H.C. performed

Arabidopsis phenotyping experiments, O.P-T. and F.R. performed Y2H and GUS assays, O.P-T.

performed the AP-MS experiment, F.R. generated transgenic Marchantia lines, performed

phenotyping experiments, RNA-seq, and together with H.C., C.F. and S.L. performed and analyzed

the SEM, histology and microscopy experiments. L.A.H. performed statistical analyses. M.H. was

responsible for conceptualization, supervision, funding acquisition, and project administration.

M.H., P.R. and F.R. wrote the original draft. All authors reviewed and edited the manuscript.

Competing Interest Statement: The authors declare that they have no competing interests.

Classification: Biological sciences, Plant Biology and Biochemistry

Keywords: protein phosphatase, Kelch domain, brassinosteroid signaling, plant development, cell

cycle, enzyme regulation, *Arabidopsis thaliana*, *Marchantia polymorpha*

Abstract

Brassinosteroids (BRs) are plant steroid hormones sensed by the membrane receptor kinase BRI1. Activation of BRI1 leads to the dephosphorylation of BZR1/BES1 transcription factors. Overexpression of the Kelch phosphatase BRI1 SUPPRESSOR 1 (BSU1) rescued the growth defects of *bri1* mutants. Subsequent studies identified BSU1 as a protein tyrosine phosphatase, which promotes BR signaling by dephosphorylating a phosphotyrosine in the glycogen synthase kinase 3 BIN2. Crystal structures of the BSU1 phosphatase domain now reveal a high degree of structural similarity to protein phosphatase 1 (PP1), a eukaryotic serine/threonine phosphatase. Consistently, BSU1 efficiently dephosphorylated phosphothreonine- and phosphoserine-containing substrate peptides, but showed no detectable activity toward BIN2 and other phosphotyrosine substrates. A catalytically inactive BSU1 phosphatase domain suppresses the growth phenotypes of the Arabidopsis *bri1-5* mutant and binds the BSU1 homologs BSL1-3. *bsu1* and *bsu1 bsl1 bsl2/3* loss-of-function mutants display wild-type-like BR responses, but exhibit stomatal patterning and fertility defects. Importantly, the PP1-like C-terminal tail of BSU1 is phosphorylated at Thr785 by a cyclin-dependent kinase complex. The phosphorylated tail binds to the BSU1 substrate-binding grooves, blocking access to the active site. Mutation of Thr785 to alanine activates BSU1, suggesting that Kelch phosphatases and PP1 share a common regulatory mechanism. Deletion of the *Marchantia polymorpha* Kelch phosphatase MpBSLM results in an undifferentiated cell mass phenotype, associated with the over-activation of a cell cycle reporter. Taken together, our experiments suggest that plant Kelch phosphatases act as PP1-like cell cycle regulators, rather than as tyrosine phosphatases in brassinosteroid signaling. (243 words)

Significance Statement

Brassinosteroid hormones initiate a well-characterized signaling pathways in plants. While the early signaling events at the plasma membrane have been thoroughly examined, the precise sequence of molecular events that comprise the cytoplasmic signaling cascade is less well-understood. Here, the molecular function of the Kelch protein phosphatase BSU1 is re-evaluated. Previous reports have indicated that BSU1 dephosphorylates the GSK3 kinase BIN2 on a central tyrosine residue. However, structural, quantitative enzyme kinetic, and genetic analyses now suggest that BSU1 is a PP1-like serine/threonine phosphatase involved in cell cycle control rather than a tyrosine phosphatase involved in BIN2 de-phosphorylation. The regulation of BSU1 by a cyclin-dependent kinase complex and the role of Kelch phosphatases in cell cycle progression in *Marchantia polymorpha* are experimentally characterized. (121 words)

Introduction

Brassinosteroids (BRs) are a class of steroid hormones (1, 2) that regulate plant growth and development by binding to the membrane receptor kinase BRASSINOSTEROID INSENSITIVE 1 (BRI1) (3–6). BR binding to the BRI1 leucine-rich repeat extracellular domain (7, 8) promotes the dissociation of SOMATIC EMBRYOGENESIS RECEPTOR KINASES (SERKs) from receptor pseudokinases (9, 10), and the formation of a BRI1 – BR – SERK signaling complex (11–15, 6). The ligand-induced interaction of receptor and co-receptor at the cell surface brings their cytoplasmic kinase domains in proximity, triggering transphosphorylation and activation of the BRI1 kinase domain (16–19). The exact sequence of the early cytoplasmic signaling events has not been defined. The inhibitor protein BKI1 dissociates from the BRI1 kinase domain following receptor activation and relocates from the plasma membrane to the cytosol (20–22), where it associates with 14-3-3 proteins (23, 24). The BRI1 kinase domain can associate with different substrates, including the plasma membrane proton pump (25) and several cytoplasmic signaling kinases. BRI1-SIGNALING KINASES (BSKs) (26) form a family of protein kinases / pseudokinases with a conserved kinase core and a C-terminal tetratricopeptide repeat (TPR) domain (27). BSK family members are phosphorylated by BRI1 on a conserved serine residue (Ser230 in BSK1) in the activation segment of the kinase domain upon BR treatment in vivo (26, 28), and constitutively in vitro (26, 29, 24). Consistent with a function in BR signal transduction, *bsk* loss-of-function alleles have reduced BR signaling responses and associated growth phenotypes (26, 29–31). A second set of plasma-membrane associated, cytosolic BRI1 substrates are the protein kinases CONSTITUTIVE DIFFERENTIAL GROWTH 1 (CDG1) and CDL1 (32, 33). Both BSKs and CDG1/CDL1 have been reported to interact with the protein phosphatase BRI SUPPRESSOR 1 (BSU1) (33, 34). BSU1 was initially identified as a dominant suppressor of the *bri1-5* BR-receptor allele in an activation tagging (35) forward genetic screen (36). BSU1 contains an N-terminal Kelch repeat and a C-terminal phosphatase module, a domain architecture only found in plants and some unicellular eukaryotes (37, 38) (*SI Appendix*, Fig. S1). In Arabidopsis, there are three BSU1 homologs (BSL1-3) and over-expression of *BSU1*, *BSL1* and *BSL2* partially rescued the growth and BR signaling defects of both weak and strong BR-receptor alleles (33, 34, 36, 39, 40). *bsu1* T-DNA insertion mutants had no obvious phenotype (36, 37), while *bsl2 bsl3* mutants were small, sterile, and had short roots and twisted leaves (37). No homozygous *bsl1 bsl2 bsl3* mutants could be obtained (37). However, *bsu1 bsl1 BSL2*/*BSL3*-amiRNA plants had strongly reduced BR responses and an associated dwarf growth phenotype (34). At the biochemical level, both BSK1 and CDG1/CDL1 interacted with BSU1 in co-immunoprecipitation assays, in yeast two-hybrid

experiments, and in pull downs (33, 34, 36, 30). CDG1/CDL1 phosphorylated BSU1 Ser764 in

vitro. BSU1 in turn dephosphorylated the glycogen synthase kinase 3 (GSK3)

BRASSINOSTEROID-INSENSITIVE 2 (BIN2) (41) at Tyr200, a GSK3 autophosphorylation site

important for enzyme function (42, 43), in vitro (33, 34). Together, these experiments defined BSU1

as a tyrosine phosphatase that inactivates BIN2, thereby decreasing BIN2-mediated phosphorylation

of the BR transcription factors BZR1 and BES1, ultimately modulating BR-responsive gene

expression (44–50). Outside BR signaling, plant Kelch phosphatases have been characterized as

components of mitogen-activated protein kinase signaling pathways regulating stomatal

development (51–53) or immune signaling (54–56) in Arabidopsis, and in cell cycle progression in

Chlamydomonas reinhardtii (57, 58).

We aim to dissect the molecular mechanism of brassinosteroid signaling from BR sensing at the

plasma membrane to the activation of nuclear transcription factors (59). Here we present the

structure and enzyme mechanism of BSU1 and (re-)investigate its function in BR and cell signaling

in Arabidopsis and in *Marchantia polymorpha*.

Results

<u>BSU1 is a PP1-like Ser/Thr phosphatase</u>

We expressed BSU1 in baculovirus-infected insect cells (*SI Appendix*, Fig. S2) and obtained crystal

structures of its N-terminal Kelch (residues 11-351) and C-terminal phosphatase (residues 450-793)

domains at 2.25 and 2.1 Å resolution, respectively (Fig. 1 *A-C*) (*SI Appendix*, Table S1). The Kelch

domain is composed of a 6-bladed β-propeller (Fig. 1*B*), which shares structural similarity with

many Kelch domain-containing proteins mediating protein-protein interactions, including human E3

ubiquitin ligases (60, 61) (*SI Appendix*, Fig. S3 *A* and *B*). Ser251, which is phosphorylated by the

immune signaling kinase BIK1 (54), maps to a loop connecting blade IV and V (Fig. 1*B*). The C-

terminal phosphatase domain has a canonical protein phosphatase fold (Fig. 1*C*) (*SI Appendix*, Fig.

S3*C*). An extended C-terminal tail folds back into the active site of the enzyme and is well-defined

in our structure (Fig. 1*C*). Ser764, which is phosphorylated by CDG1/CDL1, maps to the C-

terminal tail of BSU1 (Fig. 1*C*) (*SI Appendix*, Fig. S1).

In our structure, the catalytic center of BSU1 coordinates two Zn^{2+} ions (Fig. 1*D*) that co-purified

with the enzyme, as previously reported (62). We mutated BSU1 Asp510 to asparagine and His707

to lysine (63) and obtained a 1.6 Å crystal structure of the putatively inactive enzyme (*SI Appendix*,

Table S1). The mutations disrupt the coordination of metal ions in the catalytic center without

altering the overall structure of the BSU1 core (root mean square deviation, r.m.s.d. is ~0.7 Å

comparing 4,737 corresponding atoms) (Fig. 1*D*). In vitro, BSU1 hydrolyzes the generic

phosphatase substrate *p*-Nitrophenyl Phosphate (*p*-NPP) and has a preference for Mn^{2+} ions when

assayed against *p*-NPP, as previously reported (34) (Fig. 1 *E* and *F*). The BSU1$^{421-793\ D510N\ H707K}$

mutant had no detectable activity in this assay (Fig. 1*E*). Next, we purified full-length BIN2, which

is auto-phosphorylated on Tyr200 when expressed in *E. coli,* as previously reported (34) (*SI*

Appendix, Fig. S2). No dephosphorylation of pTyr200 was observed when BIN2 was incubated with

either the isolated BSU1 phosphatase domain, or with full-length BSU1, even at a 4:1 substrate-to-

enzyme ratio (Fig. 1*G*).

We therefore characterized the substrate specificity of BSU1 in vitro. Structural homology searches

with the program DALI (64) revealed that the BSU1 phosphatase domain most closely resembles

protein phosphatase 1 (PP1) enzymes (65, 66), which are generic Ser/Thr phosphatases (67) (Fig.

1*H)* (*SI Appendix* Fig. S3*C*). Notably, the catalytic centers of BSU1 and human PP1 are virtually

identical (Fig. 1*I*). Optimal substrate peptides for human PP1 have been previously identified using

a high-throughput in vitro dephosphorylation assay (68). Based on this study, we designed three

different substrate peptides for BSU1, harboring either a central pSer, pThr or pTyr residue (*SI*

Appendix, Fig. S4). We performed enzyme kinetic assays and found that wild-type BSU1 but not

the phosphatase-dead BSU1$^{421-793\ D510N\ H707K}$ mutant efficiently dephosphorylated the pThr-containing

peptide, with a catalytic efficiency (k_{cat}/K_M) of $\sim24 \times 10^3$ M^{-1} s^{-1} (Fig. 1 *J* and *K*). BSU1 also

dephosphorylated the pSer-containing peptide (k_{cat}/K_M $\sim4 \times 10^3$ M^{-1} s^{-1}) but had only very low

activity against the pTyr-containing peptide (k_{cat}/K_M $\sim0.05 \times 10^3$ M^{-1} s^{-1}) (Fig. 1 *J* and *K*). In line

with this, the catalytic efficiency for *p*-NPP, a phospho-tyrosine mimicking substrate, is much lower

(k_{cat}/K_M $\sim0.09 \times 10^3$ M^{-1} s^{-1}) than observed for the pThr and pSer-containing substrate peptides (Fig.

1 *E* and *K*). Molecular docking of our synthetic substrate peptides into the Mn^{2+}-bound BSU1

phosphatase core using AlphaFold3 (69) revealed that pThr- and pSer-containing peptides may

adopt well-defined binding poses within the catalytic center of the BSU1 phosphatase domain,

whereas the pTyr-containing peptide was modeled in a distorted conformation to accommodate

access to the active site (*SI Appendix*, Fig. S4).

In agreement with the dephosphorylation experiments performed using full-length BIN2 as

substrate (Fig. 1*G*), BSU1 showed no detectable activity against a short BIN2 substrate peptide

containing pTyr200 (193-KGEANIS(pY)ICSRFYR-207) (Fig. 1 *J* and *K*). Recently, the CYCLIN-

DEPENDENT KINASE B1 (CDKB1) was identified as an in vivo substrate for the

Chlamydomonas reinhardtii Kelch phosphatase BSL1 (58) (*SI Appendix*, Fig. S1). Specifically,

Thr14 and neighboring Tyr15 in CrCDKB1 were identified to be hyperphosphorylated in a *bsl1-1*

mutant phosophoproteome (58). We found that BSU1 was able to de-phosphorylate a synthetic CrCDKB1 peptide containing pThr14 (k_{cat}/K_M ~1×10^3 M^{-1} s^{-1}), but not the pTyr15-containing substrate peptide (k_{cat}/K_M ~0.04×10^3 M^{-1} s^{-1}) (Fig. 1 *J* and *K*). It has been previously reported that the substrate specificity and enzyme activity of animal PP1 are in part determined by the type of metal ions present in the catalytic center (Fig. 1*I*) (70). We therefore assayed BSU1-mediated dephosphorylation of the synthetic pThr- and pTyr-containing substrate peptides in the presence of different metal ions (Fig. 1 *K* and *L*). These assays confirmed BSU1 to represent a Mn^{2+}-dependent Ser/Thr protein phosphatase (Fig. 1*L*). Taken together, the BSU1 phosphatase domain shares strong structural homology with human PP1 and behaves as a specific Ser/Thr phosphatase in vitro.

PP1 phosphatases interact with their protein substrates via distinct, conserved binding surfaces including the RVxF motif binding groove (71, 72). BSU1 contains a hydrophobic RVxF binding groove that is highly similar to human PP1 (Fig. 1*M*) and which in our crystal structures is occupied by the C-terminal Tobacco Etch Virus (TEV) protease recognition site that was included in our protein expression construct (Fig. 1*N*). Thus, BSU1 appears to share the catalytic activity and substrate binding mechanism with canonical PP1 enzymes (73).

<u>Ectopic expression of catalytically inactive BSU1 rescues *bri1-5* mutant phenotypes</u>

Different studies have demonstrated that expression of full-length BSU1 from the *35S* promoter can suppress the growth phenotype of the *bri1-5* BR receptor mutant (33, 34, 36, 39, 40). We used this approach to perform BSU1 structure-function studies, expressing wild-type or mutant fragments of BSU1 fused to a C-terminal 3xFlag tag in the *bri1-5* background (*SI Appendix*, Table S2). We compared the shoot growth phenotypes of the resulting T1 plants to the previously reported *bsu1-1D* allele (36) (Fig. 2*A*). Expression of the isolated BSU1 phosphatase domain (BSU1$^{421-793}$) did partially suppress the *bri1-5* shoot growth phenotype (Fig. 2*A*), while expression of the N-terminal Kelch domain (BSU1$^{1-357}$) further enhanced the stunted growth phenotype of the *bri1-5* mutant (*SI Appendix*, Fig. S5). Surprisingly, the catalytically inactive BSU1 phosphatase domain (BSU1$^{421-793}$ $^{D510N\ H707K}$) (Fig. 1 *E* and *J*) could also suppress the *bri1-5* phenotype (Fig. 2*A*) (*SI Appendix*, Fig. S5). Missense mutations in the RVxF hydrophobic binding groove, such as Cys291 and Leu289 to arginine in human PP1α, have been previously shown to disrupt the interaction of PP1 with protein substrates (74) (Fig. 1*M*). A BSU1 variant, in which we replaced the corresponding Met748 and Ile750 with arginine (BSU1$^{421-793\ M748R\ I750R}$) (Fig. 1*M*), failed to rescue the *bri1-5* mutant growth phenotype, independent of the catalytic activity of the enzyme (Fig. 2*A*).

Our observation that the presence of an intact substrate binding groove allows the BSU1
phosphatase domain to rescue the *bri1-5* phenotype prompted us to investigate if known BR
signaling components may interact with BSU1 when over-expressed in the *bri1-5* mutant. We
performed affinity purification followed by mass spectrometry (AP-MS) experiments with the N-
terminal Kelch domain, the C-terminal phosphatase domain, and the catalytically inactive BSU1[421-
793 D510N H707K], respectively. With the exception of the BSU1 homologs BSL1-3, no known BR
signaling components were recovered. The BSL1-3 proteins were strongly enriched in the BSU1
phosphatase and phosphatase-dead samples, but not in the Kelch domain sample (Fig. 2*B*). Full-
length BSU1 has been previously shown to interact with BSL1-3 in vitro and in planta (40). The
interaction surface on BSU1 has not yet been mapped. We found that the wild-type BSU1
phosphatase domain, but not the BSU1[421-793 M748R I750R] mutant targeting the RVxF binding groove
interacted with BSL2 in yeast two-hybrid assays (Fig. 2*C*). Together, these experiments suggest that
the over-expression of BSU1 suppresses the phenotypes of the BR receptor mutant allele *bri1-5*
through RVxF binding groove-mediated protein-protein interactions rather than via its catalytic
phosphatase activity (Fig. 2*A-C*).
The presence of BSLs in our AP-MS assays led us to reanalyze their role in BR signaling in relation
to BSU1. We used CRISPR/Cas9 gene editing in Col-0 background to generate a *bsu1* knock-out
allele, and higher-order *bsu1 bsl* mutants (*SI Appendix*, Fig. S6). *bsu1* mutants were
indistinguishable from wild-type plants and exhibited normal BR responses (Fig. 2 *D* and *E*), as
previously described (34, 36). *bsu1 bsl1 bsl3* and *bsu1 bsl1 bsl2* triple mutants had significantly
more stomata compared to wild type (*SI Appendix*, Fig. S7), in good agreement with the stomatal
patterning phenotype previously reported for a *bsl1 bsl2 bsl3* T-DNA insertion mutant (52). *bsu1*
bsl1 bsl3 and *bsu1 bsl1 bsl2* mutants showed no significant differences from the wild-type control
in quantitative hypocotyl growth assays performed in the presence and absence of the BR
biosynthesis inhibitor brassinazole (BRZ) (75, 10, 15, 19, 24) (Fig. 2*D*). Addition of the potent
brassinosteroid brassinolide (BL) inhibited primary root growth of *bsu1, bsu1 bsl1 bsl3* and *bsu1*
bsl1 bsl2 mutant plants to wild type-like levels, while the known BR signaling mutants *bsk3-1* (30)
and *bri1-301* (76) were insensitive (Fig. 2*E*). Both *bsu1 bsl1 bsl3* and *bsu1 bsl1 bsl2* mutants had
shorter siliques compared to wild type (*SI Appendix*, Fig. S8*A*), indicative of a fertility defect, as
previously described (37). We generated *bsu1 bsl1 bsl3 bsl2*[+/-] or *bsu1 bsl1 bsl2 bsl3*[+/-] plants, but
could not isolate a homozygous quadruple mutant. In all higher-order *bsu1 bsl* mutants we observed
a severe seed-abortion phenotype, suggesting that the quadruple mutant is likely nonviable and
therefore cannot be evaluated for BR responses (*SI Appendix*, Fig. S8 *B* and *C*). Overall, our

experiments provide no genetic evidence for a function of BSU1 and BSL protein phosphatases as positive regulators of BR signaling, while confirming their role in stomatal patterning and in other, essential cell signaling processes.

Different experimental approaches have defined the BR receptor BRI1 to be an ubiquitously expressed receptor kinase throughout plant development (77–81). However, there are conflicting reports regarding the tissue expression of *BSU1* (36, 37). We generated *proBSU1*::*GUS* and *proBSU1*::*BSU1-GUS* reporter lines and found *BSU1* expression to be low and restricted to the outer leaf margin in cotyledons and true leaves across ~15 independent T2 lines (Fig. 2*F*) (*SI Appendix,* Fig. S9). In contrast, a *proBRI1*::*GUS* reporter line confirmed the ubiquitous expression of *BRI1* (Fig. 2*F*) (*SI Appendix,* Fig. S9). These experiments indicate that the expression patterns of the BR receptor and its previously proposed downstream signaling component BSU1 are largely non-overlapping.

In conclusion, the expression domains of BSU1 and BRI1 only partially overlap (Fig. 2*F*) and *bsu1* and *bsu1 bsl* loss-of-function mutants display wild type-like BR responses (Fig. 2 *D* and *E*). Over-expression of phosphatase-dead BSU1 can rescue the growth phenotypes of the BR receptor allele *bri1-5* (Fig. 2*A*). When over-expressed, BSU1 can interact with BSL1-3, as previously reported (40), and this interaction and the ability to suppress *bri1-5* requires a functional RVxF binding groove (Fig. 2 *B*, *C* and *F*). Thus, the interaction between ectopically expressed BSU1 and BSLs may constitute a key molecular step in reestablishing intracellular BR signaling in the *bri1-5* mutant. In line with this, over-expression of BSL2 can likewise partially suppress the *bri1-5* mutant growth phenotype (Fig. 2*A*). It is thus possible, that the *bsu1-D* allele induces the artificial over-accumulation of Kelch Ser/Thr phosphatase activity (Fig. 1), which in turn could lead to a direct dephosphorylation of pSer/pThr containing BR signaling components. We attempted to identify potential pSer or pThr-containing substrates, by docking known BR signaling components (2) to the BSU1 phosphatase domain using AlphaFold3 (69). The BR transcription factor BZR1 was identified as a high scoring interaction, with an interface predicted Template Modeling (ipTM) score of ~0.9 (Fig. 2*G*). Notably, the previously characterized BZR1[168-178]pS peptide (24), which contains a GSK3 phosphorylation site at Ser173, and which has been previously shown to be phosphorylated by BIN2 (82, 83), resembles a bona fide enzyme – substrate complex when docked to the BSU1 phosphatase domain (Fig. 2*G*). In accordance, BSU1 very efficiently dephosphorylated BZR1[168-178]pS in vitro, with a k_{cat}/K_M of ~59 × 10^3 M^{-1} s^{-1}. (Fig. 2*H*). BSL1-3 could not be evaluated in this assay because we were unable to purify these BSU1 homologs (which share ~65% sequence identity in their phosphatase domains) from either *E. coli* or insect cells, as previously reported for

other PP1 enzymes (73). Taken together, artificial over-accumulation of BSU1 and BSLs may result

in the direct dephosphorylation of pBZR1/pBES1 in the *bsu1-D* mutant, as initially proposed (36).

<u>A PP1-like regulatory tail in BSU1 is phosphorylated by the CDK1 – cyclin B1 – CKS1 complex</u>

Analysis of the crystal structure of the wild-type BSU1 phosphatase domain revealed a well-ordered

C-terminal tail that folds back into the active site of the enzyme (Figs. 1*A* and 3*A*). BSU1 Thr785 is

phosphorylated, and its phosphate moiety is coordinated by Arg548, Arg679 and Tyr731 (Fig. 3*A*),

which are conserved among BSU1, BSLs and human PP1 isoforms (*SI Appendix*, Fig. S1). Despite

its proximity to the catalytic metal center, pThr785 is too far from both the metal-activated water

molecule that mediates the nucleophilic attack of PP1 substrates (66) and the general base His577

(which corresponds to His125 in HsPP1α) to enable catalysis (Fig. 3*A*). Mass spectrometric analysis

of insect cell and plant extracted BSU1 revealed that Thr785 is also phosphorylated in vivo (Fig.

3*B*) (*SI Appendix*, Table S3). Notably, BSU1 and PP1 isoforms have very similar C-terminal tails,

with BSU1 pThr785 mapping to the previously identified PxTPP motif in PP1 (Fig. 3*C*) (84).

Phosphorylation of the corresponding Thr320 in HsPP1α is an important control mechanism for

cell-cycle progression, for example regulating the phosphorylation of the retinoblastoma protein

during G1 to S phase transition (84). It has been previously demonstrated that the cyclin-dependent

kinase 1 (CDK1) – cyclin B1 complex phosphorylates the conserved threonine residue within the

PxTPP motif in human and fission yeast PP1 (85–87). Consistent with this, a recombinantly

expressed and purified CDK1 – cyclin B1 – cyclin-dependent kinase subunit 1 (CKS1) complex

(88) readily phosphorylated the isolated BSU1 phosphatase domain, but not the BSU1[421-793 T785A]

mutant protein (Fig. 3*D*).

Mutation of Thr320 in the C-terminal tail to alanine constitutively activates human PP1α (89) and

phosphorylation of this threonine residue correlates with a reduction of PP1 activity (87, 90).

However, no structure of a PP1 with an ordered C-terminal tail has been resolved thus far (91). Our

structure of the active BSU1 phosphatase domain now reveals that the phosphorylated C-terminus

adopts an inhibitor-like binding mode, triggering rearrangement of the β7-β8 and β11-β12 loops, as

previously described for small molecule and protein inhibitors (92, 93) (Fig. 3 *A* and *E*). We

therefore compared BSU1 with previously resolved PP1 – inhibitor complex structures, identified

with DALI (64) (*SI Appendix*, Fig. S3). We found that part of the BSU1 C-terminal tail (residues

767-778) binds the PP1 C-terminal binding groove, which is also targeted by, for example, the

neuronal PP1 inhibitor protein spinophilin (94) (Fig. 3*F*). In addition, the phosphorylated BSU1 C-

terminal tail occupies the acidic binding groove, adopting a similar conformation as recently

reported for human Inhibitor-3 (62). (Fig. 3*G*). Both the triple Cys-motif of Inhibitor-3 and the [783]PPpTP[786] segment of BSU1 bind across the active site of the enzyme, while two basic residues (Arg788 in BSU1 and Lys59 in Inhibitor-3) target the acidic binding groove (Fig. 3*G*). Spinophilin and Inhibitor-3 have been shown to inhibit the activity of human PP1α by blocking substrate binding grooves and access to the active site (62, 94). By structural analogy, this implies that phosphorylation of Thr785 in the C-terminal tail of BSU1 inhibits both the binding of protein substrates, and their dephosphorylation. Consequently, the BSU1[421-793 T785A] mutant dephosphorylated a synthetic pThr-containing substrate peptide about two times more efficiently than the BSU1[421-793] wild-type phosphatase domain, with k_{cat}/K_M values of ~70 × 10^3 M^{-1} s^{-1} and ~35 × 10^3 M^{-1} s^{-1}, respectively (Fig. 3 *H* and 1*K*).

Together, the structural homology of BSU1 to PP1 enzymes, its phosphorylation by the CDK1 – cyclin B1 – CKS1 complex, and the presence of conserved substrate-binding and regulatory motifs together support a role for plant Kelch phosphatases in cell cycle regulation. In line with this, we observed induction of BSU1 expression and protein accumulation in callus, a tissue formed by rapidly growing and dividing cells (*SI Appendix*, Fig. S10).

<u>The Kelch phosphatase BSLM regulates the Marchantia cell cycle</u>

Our observation that Arabidopsis *bsu1 bsl1 bsl2 bsl3* quadruple mutants are not viable (*SI Appendix*, Fig. S8) prompted us to study the function of Kelch phosphatases in *Marchantia polymorpha*. Marchantia Kelch phosphatases are encoded on the sex chromosome and we named them MpBSLM (male, MpVg00440 https://marchantia.info) and MpBSMF (female, MpUg00300), respectively. During the gametophyte phase MpBSLM appears as a single copy gene in the Tak-1 wild type. *Marchantia polymorpha* lacks a canonical BR receptor and does not respond to brassinosteroids (95). However, Marchantia harbors a BIN2-like GSK3 kinase, MpGSK, and a MpBES1 transcription factor (95, 96). Mp*gsk*[ge] and Mp*bes1*[ge] knock-out lines are both strongly affected in growth and cell differentiation (95, 96).

We generated Mp*bslm*[ge] and Mp*gsk*[ge] knock-out mutants using CRISPR/Cas9 gene editing and an amiRNA-MpBSLM knock-down line (*SI Appendix*, Fig. S11). Mp*bslm*[ge] and Mp*gsk-4*[ge] mutants both had an undifferentiated cell mass phenotype (Fig. 4*A*), similar to what has been previously described for the Mp*gsk-1*[ge] and Mp*gsk-2*[ge] alleles (96). The amiRNA-MpBSLM mutant displayed an intermediate growth phenotype (Fig. 4*A*). The Mp*bslm*[ge] and Mp*gsk-3*[ge] mutants had no recognizable air pores on the thallus surface (Fig. 4*B*). The typical thallus body plan with a dorsal epidermal layer, a central storage layer and a ventral epidermis could not be differentiated in either

the Mp*bslm*[ge] or the Mp*gsk-3*[ge] mutant, respectively (Fig. 4*B*). The ventral and dorsal epidermis and some air chambers were still recognizable in the weaker amiRNA-MpBSLM allele (Fig. 4*B*).

Arabidopsis has four Kelch phosphatases and 10 BIN2-like GSK3/SHAGGY kinases (36, 97). The presence of a single *BSL* and *GSK* gene in Tak-1 and their highly similar loss-of-function phenotypes prompted us to investigate their genetic interaction. We generated a Mp*bslm*[ge] Mp*gsk-4*[ge] double mutant, which displayed a more severe growth defect compared to either the Mp*bslm*[ge] or the Mp*gsk-3*[ge] single mutant, respectively (Fig. 4 *A* and *B*).

We next performed RNA-seq analyses comparing Mp*bslm*[ge] to the Tak-1 wild-type. We found ~500 and ~800 genes to be significantly up- or down-regulated, respectively (*SI Appendix,* Fig. S12 and Table S4). Among the differentially expressed genes, we identified several cell cycle components and cell-cycle regulated genes, including a Marchantia cyclin-dependent kinase and a cyclin D isoform (Fig. 4*C*).

To investigate if the Mp*bslm*[ge] and Mp*gsk-3*[ge] mutants were enriched with cells undergoing the cell cycle we first used 5-Ethynyl-2'-deoxyuridine (EdU) staining to mark cells undergoing DNA duplication during a 3 h time period, as previously described (98). We found EdU staining to be restricted to the meristem of Tak-1 (Fig. 4*D*) (98). In both the Mp*bslm*[ge] and the Mp*gsk-3*[ge] mutant we observed EdU stained cells across the entire thallus surface and no distinct meristematic region could be identified (Fig. 4*D*). A technical limitation of EdU staining is that it captures DNA synthesis only during the labeling window and remains detectable in cells even after DNA replication has been completed. To overcome these issues, we used the previously reported *proMpCYCD*::*Dbox-TdTomato* cell cycle marker (96), which is specifically active during the G1-S phase. We transformed the reporter construct into Tak-1 and our Mp*bslm*[ge] and Mp*gsk-3*[ge] mutants. We observed only a weak TdTomato signal in the Tak-1 wild-type, which was restricted to the meristem (Fig. 4*D*). In contrast, a much stronger TdTomato signal was observed in both the Mp*bslm*[ge] and Mp*gsk-3*[ge] mutants (Fig. 4*D*). In line with the EdU staining results, also the *proMpCYCD*::*Dbox-TdTomato* reporter was expressed throughout the thallus (Fig. 4*D*). In agreement with our biochemical analysis of BSU1 in Arabidopsis (Fig. 3*D*), MpBSLM weakly interacted with MpCDK1 and MpCKS, but not with MpCYCB1 in yeast two-hybrid assays (Fig. 4E). Additionally, we found MpBSLM to weakly interact with the Marchantia retinoblastoma protein MpRBR (Fig. 4*E*), a known PP1 substrate in human cells (84). Together, these experiments suggest a function for MpBSLM in Marchantia cell cycle regulation. Deletion of *MpBSLM* results in cell over-proliferation and uncontrolled cell differentiation. The Mp*gsk-3*[ge] allele has an overall similar phenotype (Fig. 4*A*) and MpBSLM and MpGSK weakly interact in yeast two-hybrid assays

362 (Fig. 4*E*). However, the enhanced growth phenotype of the Mp*bslm*[ge] Mp*gsk-4*[ge] double mutant
363 implies that MpBSLM and MpGSK may not function in a strictly linear signaling pathway (Fig.
364 4*A*).
365
Discussion
The *Arabidopsis thaliana* Kelch phosphatase BSU1 was initially characterized as an okadaic acid-
sensitive, PP1-like Ser/Thr phosphatase (36, 92). Later work positioned BSU1 as a key component
of the intracellular brassinosteroid (BR) signaling pathway. Specifically, BSU1 was proposed to
dephosphorylate the pTyr200 residue of BIN2, thereby regulating its activity toward the
transcription factors BZR1 and BES1 (33, 34). However, based on our structural and enzyme
kinetic analyses, the BSU1 phosphatase domain consistently behaves as PP1-like Ser/Thr
phosphatase and we could not detect dephosphorylation of BIN2 pTyr200, either in the full-length
kinase or in a synthetic peptide containing this site (Fig. 1*G, J–L*).
It has been previously demonstrated that constitutive over-expression of full-length wild-type BSU1
can partially suppress the growth phenotypes of both weak and strong BR receptor alleles (33, 34,
36, 40). Our structure-function experiments now reveal that (i) expression of the isolated Kelch
domain further exacerbates the growth phenotype of the *bri1-5* mutant (*SI Appendix*, Fig. S5), (ii)
the isolated BSU1 phosphatase domain alone is sufficient to suppress this growth defect (Fig. 2*A*)
and (iii) a catalytically inactive BSU1 variant retains the ability to suppress the *bri1-5* phenotype
(Figs. 1*E* and 2*A*). Together, these results, along with our observations that *bsu1* knockout alleles
have no detectable BR signaling defects and that *BSU1* is not broadly expressed, are inconsistent
with a central role for BSU1 in canonical BR signal transduction (2, 34). Instead, they support the
interpretation that *bsu1-1D* and *pro35S*::*BSU1* act as neomorphic alleles in this context. This raises
the question of how over-expressing a catalytically inactive phosphatase core can partially restore
BR signaling in the *bri1-5* allele (Fig. 2*A*). Our findings that (i) a BSU1 variant with a disrupted
RVxF substrate-binding groove fails to suppress the *bri1-5* phenotype (Figs. 1*M* and 2*A*), (ii) BSL1,
BSL2 and BSL3 associate with the overexpressed BSU1 phosphatase domain in *bri1-5* plants (Fig.
2*B*), and (iii) the RVxF binding groove mediates the interaction between BSU1 and BSL2 (Fig. 2*C*),
suggest that BSU1 overexpression in *bri1-5* may influence BSL accumulation, distribution, or sub-
cellular localization, as previously suggested (40). This, in turn, could indirectly alter BR signaling.
In line with this, over-expression of BSL2 also suppresses the *bri1-5* phenotype (Fig. 2*A*). However,
in agreement with an earlier report (37), our newly generated *bsu1 bsl1 bsl3* and *bsu1 bsl1 bsl2*
knock-out mutants had no detectable BR phenotypes, even in highly sensitive and quantitative

hypocotyl growth assays (Fig. 2*D*), which we have previously used to detect very weak BR
signaling defects in accessory BR signaling components such as BIR receptor pseudokinases or 14-
3-3 proteins (10, 24). Importantly, the triple mutants display the previously reported stomatal
patterning defects (52) (SI Appendix, Fig. S7). The fertility defect of *bsu1 bsl1 bsl3 bsl2*[+/-] and *bsu1*
bsl2 bsl3[+/-] mutants limits our genetic analysis of BSL isoforms in BR signaling (*SI Appendix*, Fig.
S8). It remains formally possible that a single *BSL2* or *BSL3* allele can support BR signaling to wild
type-like levels (Fig. 2 *D* and *E*).

While our loss-of-function studies cannot substantiate a role for BSU1 or BSLs in BR signaling,
over-expression of these phosphatases can clearly impact the BR pathway (Fig. 2A and *SI*
Appendix, Fig. S5). One possible mechanism would be that BSU1 and BSL overaccumulation leads
to an increased dephosphorylation of BR signaling components containing pSer and/or pThr sites.
We found that the BSU1 phosphatase domain can efficiently dephosphorylate pBZR[169-182] in vitro, a
fragment containing an important BIN2 phosphorylation site in BZR1 required for 14-3-3 binding
and nuclear-cytoplasmic shuttling (24, 45, 82, 83, 99). Dephosphorylation of the related BES1
transcription factor by BSU1 in vivo and in vitro has been previously reported (36), and
accumulation of dephosphorylated BES1/BZR could rationalize the suppression of the *bri1-5*
mutant phenotype in BSU1 or BSL2 over-expressing lines. An alternative possibility is that a BR-
regulated protein kinase preferentially phosphorylates BSU1, BSL2, or BSU1-BSL complexes in
our *bri1-5* rescue lines (Fig. 2*A*) (40), protein assemblies that would not normally be present in cells
or tissues actively engaged in BR signaling (Fig. 2*F*). Indeed, the recombinant BIN2 kinase can
efficiently trans-phosphorylate BSU1, especially its N-terminal Kelch domain (*SI Appendix*, Fig.
S13). Taken together, several molecular scenarios could account for the enhanced BR signaling
observed in transgenic lines over-accumulating Kelch phosphatases. However, our genetic
experiments and previous studies (37) do not provide compelling evidence for the involvement of
these broadly acting Ser/Thr phosphatases in BR signaling.

While the molecular functions of Arabidopsis BSU1/BSLs in cell signaling (52, 54, 55) warrant
further investigation, a core function for Kelch phosphatases in cell cycle regulation has been
established in Chlamydomonas (57, 58). Specifically, CrBSL1 (*SI Appendix*, Fig. S1) has been
reported to promote mitotic entry by dephosphorylating the cyclin-dependent kinase CDKB1 (58).
We found that the BSU1 phosphatase core is structurally highly similar to human and yeast PP1,
well-established cell cycle regulators that counteract CDK1-mediated phosphorylation events (89,
90). The HsPP1α and BSU1 active sites are virtually identical and BSU1 harbors the known PP1
RVxF, C-terminal, acidic and hydrophobic binding grooves (Figs. 1 *H* and *I*, 3*F*). While the

structure of an apo PP1 enzyme has not been resolved so far (94), our structure of a catalytically

inactive BSU1 reveals the substrate-binding loops in an open conformation (Fig. 3*E*).

In the crystal structure of the active BSU1 phosphatase domain, we detected phosphorylation of the

regulatory C-terminal tail at the previously identified PxTPP motif (84), with the tail occupying the

active site and interacting with both the C-terminal and acidic binding grooves (Fig. 3 *A-C* and *F*).

The phosphorylated C-terminal tail can block both access to the PP1 active site and restrict the

binding of selected substrates and inhibitors (Fig. 3 *F* and *G*). Mutation of the phosphorylated

BSU1 Thr785 to alanine increased the dephosphorylation of a pThr-containing substrate peptide

(Fig. 3*H*), rationalizing previous genetic and biochemical studies (84, 87, 89, 90). Our structural

data further support a recent mechanistic analysis of a PP1 C-terminal tail deletion mutant (100).

The central threonine residue in the PxTPP motif has been previously shown to be phosphorylated

by CDK1 (85, 86), with CDK1 counteracting PP1 activity (101). Consequently, BSU1 Thr785 was

efficiently phosphorylated by the CDK1 – cyclin B1 – CKS1 complex in vitro (Fig. 3*D*). While we

originally identified the pThr785 site in insect cell-expressed BSU1, this site is also found

phosphorylated in Arabidopsis (*SI Appendix*, Table S3), and in CrBSL1 in Chlamydomonas (58),

suggesting that the PP1-like inhibitory mechanism is conserved among plant Kelch phosphatases.

Our structural and biochemical data strongly argue for a function of Kelch phosphatases as cell

cycle regulators, similar to PP1 in yeast and human. Consistent with this notion, knockout of the

single Kelch phosphatase gene in *Marchantia polymorpha* Tak-1 results in a severe phenotype, with

the Mp*bslm*[ge] mutant forming a drastically undifferentiated cell mass (Fig. 4 *A* and *B*). In agreement

with the recent study in Chlamydomonas (58), our work supports a core function for Kelch

phosphatases in algal and plant cell cycle progression, likely via multiple regulatory mechanisms.

Whether Kelch phosphatases share redundant functions with type-one serine/threonine protein

phosphatases (TOPPs) in plants remains to be characterized (102).

Materials and methods

Expression construct cloning

The Kelch domain (residues 1–357; BSU1[1-357]) and the full-length enzyme (residues 1–793; BSU1[1-

793]) from *Arabidopsis thaliana* BSU1 (TAIR-ID: At1g03445, https://www.arabidopsis.org) were

cloned from codon-optimized synthetic genes (Twist Bioscience) into the pBB3 vector for

expression in *Spodoptera frugiperda* Sf9 cells. The constructs carried an N-terminal 10×His-Twin-

StrepII affinity tag followed by a tobacco etch virus protease (TEV)-cleavable site. The phosphatase

domain (residues 421–793; BSU1$^{421\text{-}793}$) was cloned into the pBB40 vector carrying a C-terminal

TEV-cleavable Twin-StrepII-10×His affinity tag (*SI Appendix*, Table S2).

The full-length BIN2 kinase (residues 1–380; BIN2) from *Arabidopsis thaliana* was cloned into the

pMH_HStrx vector from a synthetic gene codon-optimised for expression in *Escherichia coli*. The

vector provides an N-terminal 10×His-Streptavidin II-thioredoxin tag, followed by a TEV-cleavage

site. Point mutations in BSU1$^{421\text{-}793}$ and BIN2 were generated by site-directed mutagenesis and

subcloned into plasmids pBB40 and pMH_HStrx, respectively. The phosphatase-dead BSU1$^{421\text{-}793}$

variant was produced by replacing the Asp510 and His707 residues in the catalytic center with

asparagine and lysine, respectively (BSU1$^{421\text{-}793\ D510N\ H707K}$). The kinase-dead BIN2 variant was

generated by replacing the catalytic Lys69 with arginine (BIN2^{K69R}), and the non-phosphorylatable

BIN2 mutant was obtained by replacing Tyr200 with phenylalanine (BIN2^{Y200F}).

The synthetic genes encoding full-length CYCLIN-DEPENDENT KINASE-1 ISOFORM 1

(residues 1–297; HsCDK1), G2/MITOTIC-SPECIFIC CYCLIN-B1 (residues 1–433; HsCyclin B1)

and CYCLIN-DEPENDENT KINASES REGULATORY SUBUNIT-1 (residues 1–79; HsCKS1)

from *Homo sapiens* were obtained as gene optimized versions for expression in Sf9 cells (Thermo

Fisher Scientific) as previously described (88). HsCDK1, HsCyclin B1 and HsCKS1 were cloned

into the pF1 vector for co-expression forming the CDK1-cyclin B1-CKS1 complex. An 8×His tag

was fused to the C-terminus of HsCDK1 and a TEV-cleavable Twin-StrepII tag was fused to the C-

terminus of HsCyclin B1.

<u>Protein expression and purification</u>

BSU1$^{1\text{-}357}$, BSU1$^{421\text{-}793}$, BSU1$^{421\text{-}793\ D510N\ H707K}$, and BSU1$^{1\text{-}793}$ were expressed in *Spodoptera frugiperda*

Sf9 cells using recombinant baculoviruses. Typically, 10 mL of recombinant P3 baculovirus was

used to infect 200 mL of Sf9 insect cells growing in SF900 media (Gibco) at a cell density of ~2.0

× 10^6 cells/mL. The cells were incubated for 72 h at 28 °C while shaking at 110 rpm. Cells were

harvested by centrifugation at 1,000 × g for 20 min at 4 °C when cell viability was 80–85% and GFP

fluorescence indicated >40% infection rate. The cell pellets were flash-frozen in liquid N$_2$ and

stored at −80 °C in lysis buffer containing 50 mM Bis-Tris HCl (pH 7.5), 300 mM NaCl, 0.5 mM

TCEP, 0.05 mM EDTA, protease inhibitor cocktail tablets (cOmplete EDTA-free; Roche), and

DNase I (Roche). For protein purification, cells were thawed and lysed by sonication (Branson

Sonifier DS450). The lysate was clarified by ultracentrifugation at 50,000 × g for 60 min at 4 °C and

filtered through 0.45 µm filters (Durapore Merck). The supernatant was loaded onto a Ni^{2+} affinity

column (HisTrap HP 5 mL, Cytiva), washed with 50 mM Bis-Tris HCl (pH 7.5), 300 mM NaCl, 0.5

mM TCEP, 0.05 mM EDTA, 20 mM imidazole, and eluted with wash buffer supplemented with

350 mM imidazole (pH 8.0) directly onto a Strep-Tactin XT 4Flow 5 mL column (IBA). The final

elution from the Strep-Tactin column was performed using BXT buffer (IBA). The eluted fractions

were incubated with TEV protease for 16 h at 4 °C to remove the 10×His and Twin-StrepII affinity

tags. The cleaved protein was separated from the tags by a second Ni^{2+} affinity chromatography

step. Further purification was achieved by size-exclusion chromatrography on a HiLoad Superdex

200 16/600 pg column (Cytiva) equilibrated in 20 mM Hepes (pH 7.5) and 150 mM NaCl. The peak

fractions were concentrated to 5–15 mg/mL and immediately used for crystallization experiments or

flash-frozen in liquid N_2 and stored at −80 °C for enzymatic assays.

The expression of the full-length BIN2, BIN2^{K69R} and BIN2^{Y200F} was done in *E. coli* BL21 (DE3)

RIL cells grown in Terrific Broth containing 50 µg/mL kanamycin and 34 µg/mL chloramphenicol.

When the culture reached an $OD_{600 \; nm}$ of ~0.8, protein expression was induced with 0.5 mM

isopropyl β-D-1-thiogalactopyranoside (IPTG) at 18 °C for 16 h. Cells were harvested by

centrifugation and resuspended in a buffer containing 50 mM Bis-Tris HCl (pH 7.5), 300 mM NaCl,

0.5 mM TCEP, and 0.1 mM EDTA, supplemented with a complete protease inhibitor cocktail tablet,

DNase I, and 0.5 mg/mL lysozyme (Roth). Cell lysis was achieved by sonication. The lysate was

then clarified by centrifugation at 50,000 × g for 30 min, and the supernatant was loaded onto a Ni^{2+}

affinity column (HisTrap HP 5 mL, Cytiva) equilibrated in 50 mM Bis-Tris HCl (pH 7.5), 300 mM

NaCl, 0.5 mM TCEP, 0.1 mM EDTA, and 20 mM imidazole. After a washing step with the same

buffer, BIN2 was eluted in buffer supplemented with 250 mM imidazole (pH 8.0). The 10×His-

Streptavidin II-thioredoxin tag was cleaved overnight with TEV protease at 4 °C. The tag-free BIN2

protein was separated from the fusion tag by a second Ni^{2+} affinity chromatography step. Finally,

BIN2 was applied to a HiLoad Superdex 200 16/600 pg column (Cytiva) equilibrated in 20 mM

Hepes (pH 7.5) and 150 mM NaCl. The peak fractions containing BIN2 were concentrated, flash-

frozen in liquid N_2 and stored at −80 °C for enzymatic assays.

The human CDK1 – cyclin B1 – CKS1 complex, was co-expressed in Sf9 cells using baculoviruses

as previously described (88). Typically, 25 ml of recombinant P3 baculoviruses were used to infect

500 ml of Sf9 insect cells at a cell density of roughly 2.0×10^6 cells/mL. The cells were incubated

for 48 h at 27 °C at 110 rpm, harvested at a cell viability rate of 80-85%, flash-frozen in liquid N_2

and stored at −80 °C. The CDK1-cyclin B1-CKS1 complex was purified using StrepTactin

Superflow Cartridge (Qiagen) columns. Lysis and wash buffer contained 50 mM Tris-HCl (pH 8.0),

500 mM NaCl, 10 mM β-mercaptoethanol and 5% (v/v) glycerol. The complex was eluted using

wash buffer supplemented with 2.5 mM desthiobiotin and further loaded onto a HisTrap HP column

(GE Healthcare Life Sciences). The complex was eluted from the HisTrap column in 50 mM Tris-
HCl (pH 8.0), 500 mM NaCl, 10 mM β-mercaptoethanol, 5% (v/v) glycerol, and 250 mM
imidazole. The CDK1-cyclin B1-CKS1 complex was loaded on a Superdex 200 Increase 10/300 GL
column (Cytiva) size exclusion column as a final purification step. Peak fractions were pooled,
concentrated and flash-frozen in liquid nitrogen and stored at −80 °C for enzymatic assays.
531
532 <u>Analytical size-exclusion chromatography</u>
533 Gel filtration experiments were performed using a Superdex 200 Increase 10/300 GL column
534 (Cytiva) pre-equilibrated in 20 mM Hepes pH 7.5, 150 mM NaCl. 80 µl of the respective protein
535 was loaded sequentially onto the column, and elution at 0.75 ml/min was monitored by ultraviolet
536 absorbance at 280 nm. Peak fractions were analyzed by SDS–PAGE gel electrophoresis.
537
<u>Protein crystallization and data collection</u>
Crystals of the BSU1 Kelch domain developed in sitting drops composed of 0.25 µL of protein
solution (BSU1$^{1-357}$ at 15 mg/mL in 20 mM HEPES [pH 7.5], 150 mM NaCl) and 0.25 µL of
crystallization buffer (18% [w/v] PEG 8,000, 5% [v/v] PEG 550 MME, 0.2 M sodium acetate
trihydrate, 0.1 M sodium cacodylate trihydrate [pH 6.5]) suspended over 80 µL of the latter as
reservoir solution. Crystals were transferred to reservoir solution supplemented with 20% (v/v)
glycerol and snap-frozen in liquid N_2. A complete native dataset to 2.25 Å was recorded at beam
line X06DA of the Swiss Light Source, Villigen, Switzerland. Crystals of the BSU1 phosphatase
domain grew in hanging drops composed of 1.5 µL of protein solution (BSU1$^{421-793}$ at 5 mg/mL in
20 mM HEPES [pH 7.5], 150 mM NaCl) and 1.5 µL of crystallization buffer (0.7 M sodium citrate
tribasic dihydrate, 0.1 M Tris [pH 8.0]) using microseeding protocols. Crystals were cryoprotected
by serial transfer into 0.8 M sodium citrate tribasic dihydrate, 0.1 M Tris (pH 8.0), 20% (v/v)
glycerol. A native dataset and a redundant single-wavelength anomalous dispersion (SAD) dataset
close to the Zn K edge (9670.0 eV, three 360° wedges at 0.1° oscillation, with χ set to −15°, 0° and
15°) were collected to 2.1 and 2.7 Å resolution, respectively. Crystals of the BSU1 phosphatase-
dead mutant (BSU1$^{421-793\ D510N\ H707K}$ at 5 mg/mL in 20 mM HEPES [pH 7.5], 150 mM NaCl)
developed in 1.6 M sodium / potassium phosphate (pH 7.0), were snap-frozen in mother liquor
supplemented with 20% (v/v) glycerol, and diffracted to 1.6 Å resolution. Data processing and
scaling was done with XDS (103).
557
<u>Structure solution and refinement</u>

The structure of the BSU1 Kelch domain was solved by molecular replacement as implemented in

in the program MoRDa (https://www.ccp4.ac.uk/morda-automatic-molecular-replacement-

pipeline/). The highest scoring solution comprised one molecule of the Kelch domain of AtNSP1

(104) (PDB-ID pdb_00005gqt) in the asymmetric unit, and was used a starting model for automatic

model building and refinement in ARP/wARP (105). The structure was completed in iterative

rounds of manual model adjustment Coot (106) and restrained refinement in phenix.refine (107).

The anomalous signal in the scaled SAD dataset from BSU1 phosphatase crystals extended to ~4.5

Å resolution (108). 8 consistent Zn^{2+} sites were identified using XPREP/SHELXD (109). Site-

refinement and phasing was done in autoSHARP (110), followed by non-crystallographic symmetry

(NCS) averaging and density modification in phenix.resolve (111). The resulting starting phases

were input in phenix.autobuild (112) and the structure was completed by manual model building in

Coot (106) and restrained refinement in phenix.refine (107). The structure of phosphatase-dead

BSU1 was solved by molecular replacement using the program Phaser (113) and using the refined

BSU1 phosphatase structure as search model. Inspection of the final models with phenix.molprobity

(114) revealed excellent stereochemistry (*SI Appendix*, Table S1). Structural representations were

done with ChimeraX (115).

<u>*p*-NPP phosphatase assay</u>

The phosphatase activity of BSU1[421-793] (or of its catalytically dead variant BSU1[421-793 D510N H707K])

toward *para*-nitrophenyl phosphate (*p*-NPP) was quantified by measuring the absorbance of the

reaction product, *para*-nitrophenol (*p*-NP), at 405 nm using a plate reader (Tecan Infinite M Nano).

Assays were performed in 100 µL reaction mixtures containing 50 mM Hepes (pH 7.5), 200 mM

NaCl, and 2 mM $MnCl_2$ at 37 °C for 2, 4, or 8 min. At each time point, the reaction was quenched

by mixing 50 µL of the reaction mixture with 20 µL of 150 mM EDTA. *p*-NPP substrate

concentrations ranged from 0.5 to 20 mM, while the enzyme concentration was maintained at 5 µM.

The quantity of released product was estimated using a standard curve of $OD_{405\ nm}$ versus *p*-NP

standard concentrations. Enzyme-free blanks at $OD_{405\ nm}$ were measured at each time point to

discard effect of non-enzymatic hydrolysis of *p*-NPP. The linearity of absorbance over time (R^2 >

0.95 for all conditions) confirmed that initial reaction rates were measured. The resulting kinetic

data for BSU1[421-793] were fitted to the Michaelis-Menten equation using RStudio (version 4.2.2) to

determine the V_{max}, K_M, and k_{cat} values. All experiments were performed in triplicates (n = 3).

<u>Dephosphorylation assay of BIN2 at pTyr200</u>

The protein phosphatase activity of BSU1$^{421\text{-}793}$ toward phosphorylated Tyr200 (pTyr200) of BIN2

was assessed by immunoblotting. Assays were conducted in 100 µL reaction mixtures containing 50

mM Hepes (pH 7.5), 200 mM NaCl, and 2 mM MnCl$_2$ at 37 °C for 5, 10, 20, or 30 min. BIN2 and

BIN2^{Y200F} were used at a concentration of 20 µM, resulting in a 4:1 substrate-to-enzyme ratio

relative to either BSU1$^{421\text{-}793}$ or to full-length BSU1$^{1\text{-}793}$ (5 µM). Reactions were quenched at each

time point by mixing 20 µL of the reaction mixture with 5 µL of SDS loading buffer (62.5 mM Tris-

HCl pH 6.8, 30% glycerol, 0.01% [w/v] bromophenol blue), followed by incubation at 95 °C for 5

min. 2 µg of BIN2 and 0.5 µg of BSU1$^{421\text{-}793}$ were resolved on a 10% BIS-Tris acrylamide SDS-

PAGE gel using MES running buffer and transferred to a 0.2 µm PVDF membrane (IB34002,

Thermo Scientific) using a semi-dry protocol (iBlot3, Thermo Scientific). Membranes were blocked

with TBS-Tween (0.1%) containing 5% BSA for 1 hour at room temperature. For pTyr200

detection, membranes were probed for 1 h at room temperature with an anti-phospho-GSK3α/β (Tyr

279/216) monoclonal antibody (clone 5G-2F, mouse, Merck, ref: 05413, lot: 4229392) at a 1:2,000

dilution. Following three 10-min washes with TBS-Tween (0.1%), membranes were incubated for 1

h at room temperature with an anti-mouse-peroxidase polyclonal secondary antibody (goat,

Promega, ref: W402B, lot: 0000169565) at a 1:3,000 dilution. Signal detection was performed using

SuperSignal West Femto Maximum Sensitivity Substrate (34095, Thermo Scientific). The

experiment was performed in triplicate (n = 3).

<u>Malachite green-based phosphatase assay</u>

The protein phosphatase activity of BSU1$^{421\text{-}793}$ (or its catalytically dead variant BSU1$^{421\text{-}793\ D510N}$

^{H707K}) was quantified using the Malachite Green Phosphate Assay Kit (Sigma-Aldrich). The assay

detects the release of inorganic phosphate produced by BSU1$^{421\text{-}793}$ during the dephosphorylation of

three synthetic 13-amino acid peptide substrates, RKLRRK(pT)GKRGSY, RKLRRK(pS)GKRGSY,

and RKLRRK(pY)GKRGSY, which differ solely at the phospho-residue (designed based on Hein et

al. (68); see *SI Appendix* Fig. S4). Synthetic peptides for the putative natural substrates BIN2 (193-

KGEANIS(pY)ICSRFYR-207, BIN2$^{193\text{-}207}$pY), CrCDKB1 (8-EKIGEG(pT)YGKVYK-20,

CrCDKB$^{18\text{-}20}$pT; or 8-EKIGEGT(pY)GKVYK-20, CrCDKB$^{18\text{-}20}$pY) and BZR1 (168-

<u>R</u>LRISN(pS)CPVTP<u>RY</u>-178, BZR1$^{168\text{-}178}$pS) were based on previous reports (33, 34, 58, 24, 82,

83). All peptide were purchased from GenScript biotech. Reactions were performed in 100 µL

mixtures containing 50 mM Hepes (pH 7.5), 200 mM NaCl, and 2 mM MnCl$_2$ at 37 °C. For

RKLRRK(pT)GKRGSY, RKLRRK(pS)GKRGSY, and BZR1$^{168\text{-}178}$pS, 50 nM BSU1$^{421\text{-}793}$ were

incubated with substrate concentrations ranging from 6.25 µM to 1000 µM for 2, 3, or 4 min. For

RKLRRK(pY)GKRGSY, BIN2$^{193\text{-}207}$pY, CrCDKB$^{18\text{-}20}$pT and CrCDKB$^{18\text{-}20}$pY, a higher BSU1$^{421\text{-}793}$

enzyme concentration (2 µM) was used to ensure sufficient Pi release for malachite green reagent

detection. Reactions were quenched by mixing 40 µL of the reaction mixture with 10 µL of

malachite green reagent, and absorbance at 620 nm was measured after 25 min using a plate reader

(Tecan Infinite M Nano). Enzyme-free blanks at OD$_{620\ nm}$ were measured at each time point to

discard effect of non-enzymatic hydrolysis or background inorganic phosphate. The concentration

of released inorganic phosphate was estimated using a standard curve of OD$_{620\ nm}$ versus inorganic

phosphate standards. Kinetic parameters (V_{max}, K_M, and k_{cat}) were calculated by fitting the data to the

Michaelis-Menten equation in RStudio (version 4.2.2), with linear regression analysis confirming

R² > 0.95 for all conditions. All experiments were conducted in triplicate (n = 3).

<u>Generation of transgenic Arabidopsis lines</u>

For *bri1-5* suppression assay the BSU1 (At1g03445) or BSL2 (At1g08420) genomic fragments

were amplified using Phusion High-Fidelity DNA Polymerase (New England Biolabs) according to

manufacturer's protocols and clone into a Golden Gate level 1 plasmid (116) (*SI Appendix*, Table

S2). Point mutations were introduced using Gibson cloning (*SI Appendix*, Table S2). Level 2 binary

plasmids were assembled providing the Cauliflower Mosaic Virus CaMV 35S promoter, the

corresponding BSU1 version, a C-terminal 3xFlag tag, a NOS terminator and FastRed as selection

marker. Final constructs were introduced into *Agrobacterium tumefaciens* strain GV3101 and then

to the *bri1-5* mutant using the floral dip method (117). Transgenic T1 seeds were identified by

mCherry fluorescence with a Zeiss Axio Zoom.V16 stereo microscope (mRFP filter) and a

HXP200C illuminator. The *bsu1*, *bsu1 bsl1 bsl2* and *bsu1 bsl1 bsl3* mutants were generated by

clustered regularly interspaced palindromic repeats (CRISPR/Cas9) gene editing (118). Guide-

RNAs (gRNAs) were designed using the CRISPR-Pv2 website

(http://crispr.hzau.edu.cn/CRISPR2/). Entry vectors containing the gRNAs under the control of the

U6 promoter were generated by annealing and ligating forward and reverse primers. Then, gRNA

entry vectors were used to create the binary vectors containing the *proU6*::*gRNA*, *proAtRPS5*::*Cas9*

and FastRed selection marker using golden gate cloning. Constructs were introduced into

Agrobacterium tumefaciens strain GV3101 and subsequently transformed into *A. thaliana* Col-0

ecotype using the floral dip method (117). Transgenic T1 seeds were identified via their fluorescent

seed coats. Genotyping of CRIPR-Cas9 events was done by PCR followed by Sanger sequencing,

Cas9 was segregated out in the T2 generation. Homozygous single *bsu1* and triple mutants *bsu1*

bsl1 bsl2 and *bsu1 bsl1 bsl3* mutants were isolated in T4 generation by PCR and verified by Sanger-

sequencing (Microsynth, AG). Quadruple segregating lines *bsu1 bsl1 bsl3 bsl2+/-* and *bsu1 bsl1*

bsl2 bsl3+/- were generated, but no quadruple homozygous lines could be obtained.

<u>Phenotypic analysis of BSU1/BSL2 over-expressing lines in *bri1-5*</u>

20-25 transformants per line were selected, sterilized, put on half-strength Murashige and Skoog

($^{1/2}$MS) medium containing 0.8% (w/v) agar plates and stratified at 4 °C for 48 h. Plates were then

moved to a Percival growth chamber with 16 h / 8 h light / dark cycle at 22 °C for 7 d. After that,

seedling were transferred to soil and kept in a controlled environment with 16h / 8h light / dark

regime at 22 °C for 45 d.

<u>Western-blotting</u>

To analyze the protein accumulation of BSU1-3xFlag and BSL2-3xFlag constructs, ~100 mg of

tissue sample were harvested and snap-frozen in liquid N_2 in a 2 mL tube with glass beads. Samples

were ground in a tissue lyzer (MM400, Retsch) and 300 µl of extraction buffer (50 mM Tris [pH

7.5], 150 mM NaCl, 1% [v/v] protease inhibitor cocktail [PE023, Merck]) was added to the tube.

Samples were then gently agitated at 4 °C for 30 min in an orbital shaker (Intelli-Mixer RM-2M,

ELMI). Next, samples were centrifugated at $10,000 \times g$ for 20 min at 4 °C. 150 µL of supernatant

was transferred to a new tube and 30 µl of 6x SDS sample buffer (0.5 M Bis-Tris [pH 6.6], 50%

[v/v] glycerol, 6% [w/v] SDS, 0.5 M DTT with 0.05% [w/v] bromophenol blue) was added.

Samples were boiled at 95 °C for 5 min and 30 µl were loaded and separated on a 10% SDS-PAGE

gel. Proteins were transferred to a 0.2 µm PVDF membrane (IB34002, Thermo Scientific) using a

semi-dry protocol (iBlot3, Thermo Scientific). Membranes were blocked for 1 h with 5% (w/v) milk

powder in TBS-T (50 mM Tris [pH 7.5], 150 mM NaCl, 0.1% [v/v] Tween 20) at room temperature.

Then membranes were incubated at room temperature for 2 h with Anti-FLAG M2-Peroxidase

(HRP) antibody and a dilution of 1:5,000. Membranes were washed three times with TBS-T,

detected using SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Scientific) and

visualized with Amersham Imager (Cytiva).

<u>Affinity purification coupled to mass spectrometry</u>

The immunoprecipitation followed by mass-spectrometry (AP-MS) experiment was performed as

previously described (119). Seedlings of *pro35S::BSU1$^{1-357}$-3xFlag*, *pro35S::BSU1$^{421-793}$-3xFlag* and

pro35S::BSU1$^{421-793\ D510N\ H707K}$-3xFlag were sterilized and plated on $^{1/2}$MS medium (Duchefa). Plates

were stratified at 4 °C for 48 h and transferred to a Percival growth chamber with 16 h / 8 h light /

dark cycle at 22 °C for 12 d. 100 mg of whole seedlings were harvested and snap-frozen in liquid

N_2. Samples were ground in a tissue lyzer (MM400, Retsch) and 300 µl of buffer was added (50mM

Tris [pH 7.5], 150mM NaCl, 0.1% [v/v] Triton X-100, 10% [v/v] glycerol, 1% [v/v] protease

inhibitor cocktail [PE023,Merck]). Tubes were gently agitated for 30 min at 4 °C and then

centrifuged at 10,000 × g for 15 min at 4 °C. The supernatant was transferred to a new tube and

centrifuged again at 10,000 × g for 15 min at 4 °C. Protein concentration was determined using

Bradford method. All samples were adjusted to total protein concentration of 3 mg/mL. 200 µL of

total protein was loaded to 50µl of buffer equilibrated Anti-FLAG magnetic beads (Sigma). Tubes

were gently agitated for 30 min at 4 °C and then washed three times with 400 µL buffer. Proteins

were eluted by adding 100 µL of 0.1 M glycine HCl (pH 3.0), with 5 min incubation. 10 µL were

used for visualization on a silver-stained (Pierce™ Silver Stain for Mass Spectrometry, Thermo

Scientific) SDS-PAGE gel. The remaining elution fractions were dried under speed-vacuum and

proteins were digested with a mix of Trypsin/LysC using an iST kit (PreOmics). Peptides were then

analyzed by nanoLC-ESI-MSMS using an easy-nLC 1000 liquid chromatography system (Thermo

Fisher Scientific) coupled with a Q-Exactive HF Hybrid Quadrupole-Orbitrap Mass Spectrometer

(Thermo Fisher Scientific). spectra were extracted and searched against the *Arabidopsis thaliana*

reference proteome database (Uniprot, https://www.uniprot.org/, 2023-12, 27'448 entries) and an

in-house database of common contaminant using Mascot (Matrix Science, London, UK; version

2.6.2). Trypsin was selected as the enzyme, with one potential missed cleavage. Precursor ion

tolerance was set to 10 ppm and fragment ion tolerance to 0.02 Da. Carbamidomethylation of

cysteine was specified as a fixed modification. Variable amino acid modification was oxidized

methionine, as well as phosphorylation of serine, threonine and tyrosine. Peptide-spectrum matches

were validated using Percolator a target FDR of 0.01 and a Delta Cn of 0.5. For label-free

quantification, features and chromatographic peaks were detected using the "Minora Feature

Detector" Node with the default parameters. PSM and peptides were filtered with a false discovery

rate (FDR) of 1%, and then grouped to proteins with again a FDR of 1% (strict) or 5% (relaxed) and

using peptides with high confidence levels. The ptmRS mode was used for phosphorylation

modifications sites localization and validation. Only sites located with a probability of more than

75% were considered. Both unique and razor peptides were used for quantitation and protein

abundances were calculated as the average of the three most abundant distinct peptide groups. The

abundances were not normalized. "Protein abundance based" option was selected for protein ratio

calculation and associated *p*-values were calculated with an ANOVA test (individual proteins).

 Yeast two-hybrid assays

BSU1$^{421-793}$, BSU1$^{421-793 \text{ M793R I750R}}$, BSL2 and MpBSLM coding sequences were synthesized (Twist

Bioscience). MpCDK1, MpCKS, MpCYCB1, MpGCK and MpRBR were amplified from cDNA

using high fidelity DNA polymerase KOD-plus (TOYOBO, No. KOD-201). All coding sequences

were ligated to P6 and pB29 (Hybrigenics Services) providing an N-terminal GAL4 activating

domain and a C-terminal LexA DNA binding domain, respectively, and the diploid TATA strain

(Hybrigenics Services) was used (*SI Appendix*, Table S2). One colony of yeast was grown overnight

at 30 °C with constant shaking at 200 rpm in 30 mL YPAD liquid media (20 g/L glucose, 20 g/L

bacto-peptone, 10 g/L yeast extract and 0.04 g/L adenine hemisulfate). After 16 h, the culture was

diluted to an OD$_{600 \text{ nm}}$ of 0.2 and grown until the OD$_{600 \text{ nm}}$ reached 0.4-0.6 before being transformed.

Cells were collected at 1,000 × g for 5 min and re-suspended in 25 mL Tris-EDTA buffer (TE).

Cells were centrifuged and re-suspended in 1xTE/LiAc to a final volume of 1.5 mL. 0.25 µg of prey

plasmid, 0.25 µg of bait plasmid were mixed with 10 µl of herring sperm ssDNA (10 mg/ml). 100

µl of yeast cells and 600 ul of PEG/LiAc were added to each transformation, and vortexed for 10 s.

The transformations were incubated 30 min at 30 °C while shaking at 200 rpm before heat shock at

42 °C for 15 min. After being cooled down on ice for 2 min the cells were spun down and re-

suspended in 0.5 mL of TE. 100 µl of transformed cells were plated on SD plates (yeast nitrogen

base with ammonium sulfate and without amino acids 6.7 g/L, glucose 20 g/L, bacto agar 20 g/L)

without leucine and tryptophan and incubated at 30 °C for 3 d. Positive transformants were selected

on SD - LW plates and growth overnight in selective SD liquid media in duplicates at 30 °C while

shaking at 200 rpm. Each sample was diluted to an initial OD$_{600 \text{ nm}}$ of 0.2 and then sequentially

diluted 10x (10x, 100x, 1,000x, 10,000x). 5 µl of each dilution was spotted on SD plates without

leucine and tryptophan and with or without histidine. Plates were incubated at 30 °C for 2 d

(Arabidopsis) or 3 d (Marchantia), respectively.

Seedling hypocotyl growth assay

Seeds were surface sterilized, stratified at 4 °C for 2 d, and plated on $^{1/2}$MS medium containing

0.8% (w/v) agar and supplemented with 1 µM BRZ from a 10 mM stock solution in DMSO (Tokyo

Chemical Industry) or with 0.1% (v/v) DMSO. To induce germination, seeds were exposed to light

for 1 h and then incubated in the dark at 22 °C for 5 d. We then scanned the plates at 600 dots per

inch resolution on a regular flatbed scanner (Perfection V600 photo; EPSON), measured hypocotyl

lengths using Fiji (120) and analyzed the results in R version 4.1.2 (121) using the packages mratios

(122) and multcomp (123). Rather than *p*-values, unadjusted 95% confidence limits for fold-

changes are reported. A mixed-effects model for the ratio of a given line compared to the wild-type

Col-0 is applied, allowing for heterogeneous variances to analyze log-transformed end point

hypocotyl lengths. To evaluate treatment-by-mutant interactions, the 95% two-sided confidence

intervals for the relative inhibition (Col-0: untreated versus BRZ-treated hypocotyl length)/(any

genotype: untreated versus BRZ-treated hypocotyl length) are calculated for the log-transformed

length.

<u>Seedling root growth inhibition assay</u>

Seeds were sterilized, put on $^{1/2}$MS plates supplemented with 1% (w/v) sucrose and stratified at 4 °C

for 48 h. Plates contained either 100 nM brassinolide (OlChemIm) from a 1 mM stock solution in

100% DMSO, or 0.01% (v/v) DMSO. Plates were transferred to a Percival growth chamber with 16

h / 8 h light / dark cycle at 22 °C for 7 d, after-which seedling were scanned (Perfection V600

photo; EPSON) and root length was manually measured in Fiji (120). Multiple comparisons of the

genotypes vs. wild-type (Col-0) were performed using a Dunnett (124) test as implemented in the R

package multcomp (123) (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$).

<u>Stomatal index determination</u>

Seeds were sterilized, put on $^{1/2}$MS plates supplemented with 1% (w/v) sucrose and stratified at 4 °C

for 48 h. 7 d-old seedlings were used to determine stomatal indices, *scrm-D (125)* and *bsk1-2 bsk2-*

2 (126) were used as controls. For confocal imaging, seedlings were incubated in a 10 mg/L

propidium iodide (PI) solution for 10 min and then washed five times with H_2O. Cotyledon abaxial

regions were imaged using a confocal LSM 780 optical microscope (Zeiss). PI staining was

visualized with an excitation wavelength of 563 nm and emission was recorded between 571 nm

and 656 nm, as previously reported (127). Mature stomata over a 0.5 mm by 0.5 mm epidermal

surface area were counted manually using Fiji cell counter (120). A Dunnett test (124) was

performed in the R package multcomp (123) to assess the statistical difference of the different

genotypes compared to the Col-0 control (*$p < 0.05$, ***$p < 0.01$, ***$p < 0.001$).

<u>Quantification silique and seed abortion phenotypes</u>

Col-0, *bsu1 bsl1 bsl3*, *bsu1 bsl1 bsl2*, *bsu1 bsl1 bsl3 bsl2$^{+/-}$* and *bsu1 bsl1 bsl2 bsl3$^{+/-}$* plants were

grown in soil for 5 weeks with a photoperiod of 16 h / 8 h at 22 °C. Representative pictures of

inflorescence branch and siliques were taken (DS126761, Canon) and silique length were measured

using Fiji (120) (n = 20). A Dunnett test (124) in the R package multcomp (123) was used to assess

the statistical difference of the different genotypes compared to the Col-0 control (*p <0.05, ***p <

0.01, ***p < 0.001). Pictures from dissected siliques from the previously mention genotypes were

taken with a binocular Nikon SMZ18 and seed abortion was evaluated on dissected siliques.

<u>β-glucuronidase (GUS) reporter assay</u>

The β-glucuronidase (GUS) reporter activity analysis, 1154 bp upstream of BSU1 ORF were cloned

to drive expression of *proBSU1*::*BSU1-GUS* and *proBSU1*::*GUS*, respectively (*SI Appendix*, Table

S2). For BRI1 construct, 2073bp upstream of the BRI1 ORF (At4g39400) were used to generate the

proBRI1::*GUS* reporter. All constructs were transformed into Ws-2 background. For each construct,

14-17 independent T2 lines were analyzed. Seeds were sterilized and stratified on $^{1/2}$MS plates for

48 h at 4 °C. Plates were then transferred to 22°C with 16 h / 8 h light dark cycle for 5 d, after-

which staining was done for GUS activity in cotyledons. For 4[th] leaf staining, seedlings were

transferred to soil for additional 10 d before staining. Samples were incubated in ice-cold 90% (v/v)

acetone for 20 min and rinsed with rinse solution (0.5 mM $K_4Fe(CN)_6$, 0.5 mM $K_3Fe(CN)_6$, and 50

mM $NaPO_4$ buffer [pH 7.0]). Next, seedlings were vacuum infiltrated for 15 min in staining solution

(0.5 mM $K_4Fe(CN)_6$, 0.5 mM $K_3Fe(CN)_6$, 10 mM EDTA, 0.1% (v/v) Triton X-100, 1 mM X-Gluc,

and 100 mM $NaPO_4$ buffer [pH 7.0]), and incubated at 37 °C for 12 h. Chlorophyll was removed by

incubation in 70% (v/v) ethanol during several washing steps. Pictures were taken with a Nikon

SMZ18 microscope.

<u>Callus induction</u>

Seeds of the *proBSU1*::*GUS*, *proBSU1*::*BSU1-GUS* and *proBRI1*::*GUS* reporter lines were surface

sterilized, stratified at 4 °C for 2 d, and plated on $^{1/2}$MS medium containing 0.8% (w/v) agar. Plates

were transferred to a Percival growth chamber with 16 h / 8 h light / dark cycle at 22 °C. Seven-d-

old seedlings were used as explant to induce callus formation. Cotyledon explants were transferred

to B5 media with 2% (w/v) sucrose and supplemented with 2,4-D (500 μg/L) and kinetin (50 μg/L).

Callus induction was done for 21 d at 22 °C in darkness, as previously reported (128). The GUS

reporter assay was performed as described above.

<u>In vitro radioactive kinase assay</u>

Phosphorylation of BSU1 by human CDK1 bound to cyclin B1 and CKS1 was conducted in 50 μL

mixtures containing kinase buffer (50 mM Bis-Tris HCl, pH 7.5, 5 mM $MgCl_2$, 2 mM DTT), 10 μM

ATP, 10 μCi of [γ-^{32}P]-ATP (6000 Ci/mmol, 10 mCi/mL) (Revvity), 10 nM of the CDK1 – cyclin

B1 – CKS1 complex (88), and 10 µM BSU1$^{421-793}$ substrate, or the BSU1$^{421-793\ T785A}$ mutant protein,

where Thr785 of the PxTPP motif is replaced by alanine. Reactions were incubated at 37 °C for 1, 5

and 10 min, respectively. Reactions were quenched by mixing 15 µL of the reaction mixture with 5

µL of SDS loading buffer (62.5 mM Tris-HCl, pH 6.8, 30% glycerol, 0.01% [w/v] bromophenol

blue) followed by incubation at 95 °C for 5 min. Proteins were resolved on a 4–15% Mini-

PROTEAN TGX Precast Gel (BioRad) using Tris-Glycine running buffer. The gel was then washed

three times for 15 min each with a fixation solution (10% [v/v] acetic acid, 20% [v/v] ethanol),

placed on a sheet of Whatman 3MM filter paper, covered with plastic wrap, and dried at 80 °C for

75 min under vacuum using a conventional gel dryer. The dried gel was exposed for 24 h to a

phosphor screen for signal detection using the Sapphire FL Biomolecular Imager (Azure

biosystems), with laser 638/424BPSP (n = 3).

The protein kinase activity of BIN2 toward BSU1$^{1-357}$, BSU1$^{421-793}$, and BSU1$^{1-793}$ was assessed by

observing the transfer of the radioisotope [γ-^{32}P] from ATP to BSU1 variants. Reactions were

conducted in 50 µL mixtures containing kinase buffer (50 mM Bis-Tris HCl, pH 7.5, 5 mM MgCl$_2$,

2 mM DTT), 10 µM ATP, 10 µCi of [γ-^{32}P]-ATP (6000 Ci/mmol, 10 mCi/mL) (Revvity), 1 µM of

the respective substrate BSU1 variant, and 30 nM BIN2. Reactions were incubated at 37 °C for 2

and 15 min (n = 3).

<u>Determination of in vitro phosphorylation sites</u>

Sample preparation and protein digestion

Protein samples were loaded on a 12% mini polyacrylamide gel, stained by Coomassie Blue. Gel

bands were excised and digested with sequencing-grade trypsin (Promega, Madison, WI, USA,

Product N. V5111) or chymotrypsin (Promega, Prod.N. V1061), as described (129). Extracted

tryptic peptides were dried and resuspended in 0.1% (v/v) formic acid, 2% (v/v) acetonitrile.

Liquid Chromatography-Mass Spectrometry analyses

LC-MS/MS analyses were carried out on a TIMS-TOF Pro (Bruker, Bremen, Germany) mass

spectrometer interfaced through a nanospray ion source ("captive spray") to an Ultimate 3000

RSLCnano HPLC system (Dionex). Peptides were separated on a reversed-phase custom packed 45

cm C18 column (75 µm ID, 100Å, Reprosil Pur 1.9 µm particles, Dr. Maisch, Germany) at a flow

rate of 250 nl/min with a 2-27% acetonitrile gradient in 33 min followed by a ramp to 45% in 5 min

and to 90% in 5 min (total method time: 65 min, all solvents contained 0.1% formic acid). Data-

dependent acquisition was carried out using a standard TIMS PASEF method (130) with ion

accumulation for 100 ms for each the survey MS1 scan and the TIMS-coupled MS2 scans. Duty

cycle was kept at 100%. Up to 10 precursors were targeted per TIMS scan. Precursor isolation was done with a 2 or 3 m/z windows below or above m/z 800, respectively. The minimum threshold intensity for precursor selection was 2500. If the inclusion list allowed it, precursors were targeted more than one time to reach a minimum target total intensity of 20'000. Collision energy was ramped linearly based uniquely on the 1/k0 values from 20 (at 1/k0=0.6) to 59 eV (at 1/k0=1.6). Total duration of a scan cycle including one survey and 10 MS2 TIMS scans was 1.16 s. Precursors could be targeted again in subsequent cycles if their signal increased by a factor 4.0 or more. After selection in one cycle, precursors were excluded from further selection for 60s. Mass resolution in all MS measurements was approximately 35'000.

Data processing

MS data were analyzed using Mascot 2.8 (Matrix Science, London, UK) set up to search the *Arabidopsis thaliana* reference proteome based on the UniProt database, and a custom contaminant database containing the most usual environmental contaminants and enzymes used for digestion (keratins, trypsin, etc). Trypsin (cleavage C-terminal to K,R) , allowing 2 missed cleavages and Chymotrypsin (cleavage C-terminal to Phe, Leu, Trp, Tyr, semispecific) were used as the enzyme definitions. Mascot was searched with a parent ion tolerance of 15 ppm and a fragment ion mass tolerance of 0.04 Da. Carbamidomethylation of cysteine was specified in Mascot as a fixed modification. Protein N-terminal acetylation, methionine oxidation and phosphorylation on Ser, Thr and Tyr were specified as variable modifications.

Data analysis

Scaffold (version Scaffold 5.0.0, Proteome Software Inc., Portland, OR) was used to validate MS/MS based peptide and protein identifications. Peptide identifications were accepted if they could be established at greater than 80.0% probability by the Percolator posterior error probability calculation (131). Protein identifications were accepted if they could be established at greater than 95.0% probability and contained at least 2 identified peptides. Protein probabilities were assigned by the Protein Prophet algorithm (132). All phosphopeptide identifications and phosphosite localizations reported were manually validated.

Marchantia growth conditions

Wild type and CRISPR/Cas9-gene edited *Marchantia polymorpha* plants were Takaragaike-1 (Tak-1) males (133). Plants were asexually maintained and propagated through gemmae or by thallus stubs on ½Gamborg B5 medium (Sigma) adjusted to pH 5.5 with KOH, under constant LED-source

white light (60 μmol/m^2/s) at 22 °C on 90 mm square Petri dishes (Greiner) containing 0.8% (w/v)

plant cell culture agar (Huber lab) as previously reported (134).

<u>Generation of stable transgenic *M. polymorpha* lines</u>

Mp*bslm*^{ge} and Mp*gsk-3*^{ge} loss-of-function mutants were generated by CRISPR/Cas9 gene editing

(118). gRNAs were designed using Casfinder (https://marchantia.info/tools/casfinder/). Primers

were synthesized included forward 5' CTCG- 3' and reverse 5' AAAC- 3' overhangs, respectively

(*SI Appendix*, Table S2). Annealed primers were cloned into a pMpGE_En03 (Addgene #71535),

plasmid digested with BsaI using T4 ligase (NEB). The entry gRNAs were subcloned into binary

vector pMpGE010 (Addgene #71536) containing the Cas9 enzyme and a Hygromycin resistance

gene as selection marker by Gateway Cloning, as described (135). The final plasmids were

transformed into *Agrobacterium tumefaciens* strain GV2260 and plant transformations were done

using the regenerated thallus protocol (133). Transgenic lines were genotyped by PCR using KOD

polymerase (Merck) and by Sanger sequencing, using primers flanking the target regions. We

selected lines bearing insertions or deletions that lead to frame shifts and early stop codons. The

Mp*bslm*^{ge} Mp*gsk-4*^{ge} double mutant was generated by retransforming Mp*bslm*^{ge} with the gRNAs

used to generate the Mp*gsk-3*^{ge} in plasmid pMpGE011 (Addgene #71537) providing a

Chlorsulfuron resistance marker. The artificial micro-RNA was design in the amiRNA Design

Helper tool (available at https://marchantia.info/tools/amir_helper/) and synthesized (Twist

Bioscience). A fragment containing the miRNA was cloned by Golden-Gate into a vector providing

a proMpEF1 promoter, a NOS terminator and hygromycin gene as selection marker. The final

binary plasmid was transform into Tak-1 as described.

<u>Marchantia widefield microscopy</u>

14 d-old Tak-1, amiRNA-MpBSLM, Mp*bslm*^{ge}, Mp*gsk-3*^{ge} and Mp*bslm*^{ge} Mp*gsk-4*^{ge} plants were

observed with an Axio Zoom V16 stereomicroscope (Zeiss) equipped with a 1x PlanNeoFluar NA

0.25 objective, an HXP 200 C fluorescence lamp (Kübler) and an Axiocam 512 colour CCD

camera. Z stacks were acquired; bright-field images were acquired with episcopic white light, and

fluorescence images were acquired with a DAPI filter (excitation 335-383 nm, emission 420-470

nm).

<u>Scanning electron microscopy (SEM)</u>

14 d-old Tak-1, amiRNA-MpBSLM, Mp*bslm*[ge], Mp*gsk-3*[ge] and Mp*bslm*[ge] Mp*gsk-4*[ge] plants were

transferred to aluminum holders on double-sided carbon tape (Electron Microscopy Sciences) and

treated with a JFC-1200 gold sputter coater (JEOL). Imaging was performed with a JSM- 6510LV

scanning electron microscope (JEOL) in high vacuum mode with a spot size of 40 mm, an

acceleration of 15 kV and a working distance of 12 mm.

<u>Marchantia cross section fixation, processing and histology</u>

14 d-old Tak-1 and amiRNA-MpBSLM thalli and Mp*bslm*[ge], Mp*gsk-3*[ge] and Mp*bslm*[ge] Mp*gsk-4*[ge]

thallus stubs were sectioned and fixed in phosphate buffer saline (PBS) pH 7.4 with 4% (v/v)

formaldehyde and 0.2% (v/v) Triton X-100. The fixation was done overnight at 4 °C under

agitation, after vacuum infiltration. Samples were then washed with PBS (2x15 min) and with H_2O

(2x10 min) before undergoing dehydration in a graded ethanol series (ethanol 30%, 50%, 70%, 90%

and 100% [v/v] with 30 min incubation two times until the last bath in ethanol 100% [v/v]

overnight at 4 °C). Technovit7100 (EMS, #14653) was prepared according to the manufacturer's

recommendations by supplementing it with hardener I, and samples were progressively infiltrated

by incubation in 3:1, 1:1 and 1:3 mixes ethanol:Technovit7100 (each time 2 h under agitation at

room temperature), before finally incubating in 100% Technovit7100 for 2 h at room temperature

(after vacuum infiltration) and for another 40 h at 4 °C. Embedding was done in Technovit7100

supplemented with 1/15 Hardener II and 1/25 polyethylene glycol 400; polymerization was done for

30 minutes at room temperature followed by 30 min at 60 °C. Sectioning was performed with a

Histocore AUTOCUT microtome (Leica) using disposable R35 blades and sections of 4 µm were

deposited on SuperFrost slides. Sections were incubated 45 s in 0.2% (w/v) toluidine blue in dH_2O,

extensively rinsed with dH_2O, then incubated one min in xylene and mounted in Pertex. The

sections were observed with a Leica DM6B widefield microscope equipped with a DMC5400

CMOS camera (used with binning 2x2) and a 20x Fluotar NA 0.55 air objective.

<u>Image analysis</u>

For a given dye, all images were acquired and post-processed identically. Image processing was

done using Fiji (120). For toluidine blue pictures, tiling and stitching was done using the Leica LAS

X Navigator tool. For wide-field microscopy images, a gamma of 0.45 was applied using Fiji, then

focus stacking was performed with the Affinity Photo software (version 1.10.6.1665) and Fiji was

finally used to apply a Cyan Hot look-up table to the fluorescence pictures and generate the overlay

between brightfield and fluorescence.

RNA-seq

Tak-1 and Mp*bslm*[ge] plants were grown for 14 d in vitro on ½Gamborg B5 medium (Sigma) adjusted to pH 5.5 with KOH, under constant LED-source white light (60 μmol/m^2/s) at 22 °C. For each biological replicate, 10 plants were pooled and RNA was extracted using the RNeasy plant mini kit (Qiagen). A Qubit fluorometer (Thermofisher) was used to determine RNA concentration, 100 ng per sample were analyzed to check quality control using 2100 Bioanalyzer system (Agilent Technologies). Library preparation and sequencing were performed by the iGE3 Genomic Platform at the Faculty of Medicine, University of Geneva (https://ige3.genomics.unige.ch/). Sequencing was performed with Novaseq 6000 machine from Illumina with 100 bp single-end output. Quality control of the reads and adaptor trimming were done with fastQC and Trimmomatic V6 (136). Genomic and transcript annotation files of the *Marchantia polymorpha* reference genome (137) and annotations were downloaded from MarpolBase (https://marchantia.info/download/MpTak_v6.1/). Read mapping to the reference was done with HISAT2 (138) (v2.2.1 with only the -dta option) and StringTie (139) (v2.2.1 with default options). DESeq2 (140) (v3.17) with default options as implemented in Galaxy server (https://usegalaxy.org/). Gene ontology enrichment analyses were performed in Panther (https://www.pantherdb.org/). DEGs annotation were manually curated to identify putative cell cycle regulated genes.

EdU staining and imaging

EdU staining in Marchantia was performed using the Click-iT EdU (5-ethynyl-2′-deoxyuridine) Alexa Fluor 488 Imaging Kit (Thermo Fisher Scientific), as described (98). 3 d-old thalli of Tak-1 or Mp*bslm*[ge] and Mp*gsk-3*[ge] thallus stubs were grown in $^{1/2}$Gamborg's B5 medium and soaked in half-strength liquid medium supplemented with 10 μM EdU at 22 °C for 3 h under continuous light. The samples were fixed in 50% (v/v) methanol / 10% (v/v) acetic acid (fixative solution) at 4 °C over night. After fixation, the samples were dehydrated through ethanol series of 80%, 90% and 100% (v/v) and a final 90% (v/v) acetone incubation at 80 °C for 5 min each, followed by fixative solution at 22 °C for 30 min. The samples were washed four times with 0.5% (v/v) Triton X-100 in PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na$_2$HPO$_4$, 1.8 mM KH$_2$PO$_4$), incubated with the EdU detection cocktail following the manufacturer's instructions for 30 min in the darkness at room temperature, and rinsed four times with 0.5% (v/v) Triton X-100 in PBS. Finally, the samples were transferred to slides and incubated with chloral hydrate solution (25 g of chloral hydrate (C$_2$H$_3$Cl$_3$O$_2$) in 10 mL H$_2$O) for 30 min. Images were obtained using a Leica DM6B widefield

microscope equipped with a DFC9000GT sCMOS camera with excitation 450-490 nm and
emission 500-550 nm.

<u>Confocal microscopy of Marchantia cell cycle reporter lines</u>
Confocal laser scanning microscopy was done in 3 d-old Tak-1 thallus apical notches or in
Mp*bslm*[ge] and Mp*gsk-3*[ge] thallus stubs transformed with the cell cycle reporter *proMpCYCD*::*Dbox-*
TdTomato (96). Samples were transferred to slides with 10% (v/v) glycerol and image acquisition
was performed on the SP8 microscope (Leica). TdTomato was excited at 552 nm, emission was
recorded between 560 and 575 nm. Z-stack pictures were assembled using max intensity projection
in Fiji (120).

Figure legends

Fig. 1. BSU1 is a PP1-like Ser/Thr phosphatase. (*A*) Schematic representation of *A. thaliana* BSU1,
depicting the Kelch (residues 11–351) and phosphatase (residues 450–793) domains
(https://www.uniprot.org/ Uniprot ID Q9LR78). Structured regions are shown as blocks, putative
linker regions (residues 1–10 and 352–449) as solid line, CDG1 phosphorylation sites are
highlighted in red, the BIK1 phosphorylation site in blue. (*B*) Top-view of the BSU1 Kelch domain.
Shown is a ribbon diagram colored with a rainbow gradient from blue (N-term) to red (C-term).
Kelch motif blades are labeled I to VI, an intrinsically disordered loop is indicated by a dotted line,
the BIK1-mediated Ser251 phosphorylation site is shown alongside (in bonds representation,
marked by a blue circle). (*C*) Ribbon diagram of the BSU1 phosphatase domain (chain A), with α-
helices in blue, β-strands in green, loops in white, and Zn^{2+} ions as gray spheres. The C-terminal tail
is shown in dark gray, and Ser764 is depicted in bonds representation, marked by a red circle. (*D*)
Close-up view of the active sites of wild-type BSU1 (left) and the catalytically inactive variant
BSU1[421-793 D510N H707K] (right). Zn^{2+} ions are shown as green spheres, the distorted trigonal bipyramidal
pentavalent coordination is indicated by dotted lines (in magenta), water molecules are shown as
red spheres. A phased anomalous difference map calculated from data collected at the Zn K-edge
and contoured at 25σ is shown as a green mesh. (*E*) Initial enzyme velocities for BSU1-catalyzed
hydrolysis of para-nitrophenyl phosphate (*p*-NPP), plotted as a function of substrate concentration.
Data were fitted to the Michaelis-Menten equation by nonlinear regression. Extracted kinetic
parameters are summarized in panel *K* (n = 3). (*F*) Metal ion preference of the phosphatase domain
using 2 mM *p*-NPP as substrate. Each metal chloride was tested at a fixed concentration of 2 mM.

(*G*) BIN2 pTyr dephosphorylation time course experiment using 20 µM BIN2 and 5 µM of either

the isolated BSU1 phosphatase domain (top) or the full-length enzyme (bottom). Shown are a

immunoblots and coomassie-stained membranes below as loading controls. The presence of BIN2

pTyr200 was detected using a human GSK3 pTyr antibody (n = 2). (*H*) Structural superposition of

the BSU1 phosphatase domain (cyan C_α trace) and HsPP1γ (in orange, PDB-ID: pdb_00001jk7

(92), r.m.s.d. is ~0.8 Å comparing 245 corresponding C_α atoms). (*I*) Close-up view of the

superimposed BSU1 (in cyan) and HsPP1γ (in orange) active sites. Shown are the catalytic

Zn^{2+}/Mn^{2+} ions (in purple), with coordinating residues depicted in bonds representation. (*J*) Initial

enzyme velocities of BSU1 phosphatase domain (or phosphatase-dead) catalyzed hydrolysis of

synthetic substrate peptides containing either a pThr (left), pSer (center) or pTyr (right), or of the

natural substrate candidates BIN2 and CrCDKB1 (bottom row), using a malachite green-based

phosphatase assay (n = 3). (*K*) Table summaries of the enzyme kinetic parameters derived from the

experiments shown in *E* and *J* (n.d., no activity detected). (*L*) Metal ion preference of the

phosphatase domain vs. synthetic substrate peptides containing either a pThr or pTyr. Assays were

performed using 200 µM of peptide substrate and 50 nM of enzyme in the presence of 2 mM of the

indicated metal chloride. In the case of $ZnCl_2$ + $Fe(II)Cl_2$ a final concentration of 500 µM was used

(n = 3). (*M*) Close-up view of the RVxF peptide binding grooves in BSU1 (in cyan) and HsPP1α (in

orange, PDB-ID: pdb_00003v4y (141), r.m.s.d. is ~0.8 Å comparing 252 corresponding C_α atoms).

Shown is a ribbon diagram of the peptide binding groove with substrate peptide-interacting amino-

acid side-chains shown in bonds representation. (*N*) Molecular surface view of the BSU1 RVxF

peptide binding groove (in cyan) bound to the TEV recognition motif (in yellow, in surface and

bonds representation) of a neighboring molecule in the asymmetric unit.

1041

Fig. 2. A catalytically inactive BSU1 phosphatase domain can suppress the growth phenotype of the

bri1-5 mutant. (*A*) Shoot growth phenotypes of 45 d-old *bri1-5* plants expressing different BSU1

variants, or full-length BSL2 from the *35S* promoter. The Ws-2 wild type, *bri1-5* (142) and the

original *bsu1-1D* (36) allele are shown alongside. BSU1/BSL2 protein expression was assayed by

western blot using an α-Flag antibody, the Ponceau-stained membrane is shown below as loading

control. (*B*) Relative abundance of BLS1, BSL2 and BSL3 in AP-MS experiments with the Flag-

tagged BSU1 Kelch (top), phosphatase (center) or phosphatase-dead (bottom) domains. (*C*) Yeast

co-expressing either the wild-type BSU1 phosphatase domain or the mutant carrying point

mutations in the RVxF binding groove fused to the Gal4-activation domain (Gal4; prey) and full-

length wild-type BSL2 fused to the LexA-binding domain (LexA; bait) were grown on selective SD

medium supplemented with histidine (-LW; co-transformation control) or lacking histidine (−LWH; interaction assay). Shown are serial dilutions from left to right after 2d of incubation. (*D*) Hypocotyl growth assay of 5 d-old dark-grown seedlings in the presence and absence of the BR biosynthesis inhibitor BRZ (75). Representative seedlings are shown in the top panel (scale bar = 0.5 cm). Box plots (bold black line, median; box, interquartile range (IQR); whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile) are in the center panel with raw data shown as individual points (n = 30). Shown below is relative inhibition of hypocotyl growth in the presence of BRZ plotted together with lower and upper confidence intervals, the dotted horizontal line depicts the Col-0 wild-type reference (rel. inhib., relative inhibition). (*E*) Brassinolide (BL)-mediated root growth inhibition assay. Shown are representative 7 d-old seedlings, grown on $^{1/2}$MS plates (mock, top panel) or in the presence of 100 nM BL (+BL, bottom panel). Quantifications are shown below as box plots with raw data indicated as individual points (n = 15) and including Dunnett-type (124), two-sided multiplicity-adjusted *p*-values for the comparison against the Col-0 control (*p < 0.05; **p < 0.01; ***p < 0.001). (*F*) Spatiotemporal expression patterns of BSU1 and BRI1 reporter lines determined by β-glucuronidase activity (GUS) assays. A sequence ~1kb upstream of the BSU1 ORF was used to drive *proBSU1*::*BSU1-GUS* and *proBSU1*::*GUS*, respectively. A ~2kb upstream of BRI1 ORF was used to drive *proBRI1*::*GUS*. 14-17 independent T2 lines for each construct were stained for GUS activity, either using 5 d-old cotyledons (top), or the 4th leaf of 15 d-old soil-grown plants. Shown are distribution star plots with expression patterns divided into categories. (*G*) AlphaFold3-predicted complex structure of BSU1$^{421-793}$ (ribbon diagram) bound to Mn^{2+} ions (purple spheres) and BZR1$^{168-178}$pS (in orange). Predicted salt bridges between the phosphate group and BSU1 residues are depicted as dotted lines (in green). AlphaFold3 pTM and ipTM values are reported alongside. (*H*) Initial enzyme velocities of BSU1 phosphatase domain catalyzed hydrolysis of the synthetic BZR1$^{168-178}$pS substrate peptide and including enzyme kinetic parameters (n = 3).

Fig. 3. The C-terminal tail of BSU1 is phoshorylated by the CDK1 – cyclin B1 – CKS1 complex. (*A*) Molecular surface view of the BSU1 phosphatase domain (residues 450-755), the C-terminal tail (residues 756-793) is shown as a ribbon diagram (in orange), pThr785 is inserted into the active site (in bonds representation). The relative positions of two substrate-binding loops are indicated. The inset provides a close-up view of the BSU1 active site with two divalent metal ions and the catalytic water molecule shown as purple and red spheres, respectively. Residues contacting pThr785 (in orange) are highlighted in bonds representation (in blue), hydrogen bonds are shown as

green dotted lines. The closest distances between the phosphate moiety of pThr785 and the catalytic water molecule or the general base His577 are indicated by yellow dotted lines. (*B*) MS1 spectrum of the phosphorylated BSU1$^{755\text{-}788}$ peptide. (*C*) Multiple sequence alignment of the C-terminal tail of BSU1, BSL1-3, two PP1-like TOPP phosphatases from *Arabidopsis thaliana*, and human and fission yeast PP1. The position of the PxTPP motif containing the pThr site is indicated (in red). (*D*) Autoradiograph of an in vitro transphosphorylation time course assay. 10 nM of the human CDK1 – cyclin B1 - CKS1 complex were incubated with 10 µM BSU1$^{421\text{-}793}$ or with BSU1$^{421\text{-}793\ T785A}$, where Thr785 in the PxTPP motif was replaced by an alanine, at 37 °C for the indicated time. (*E*) Structural superposition of BSU1$^{421\text{-}793}$ (ribbon diagram with α-helices shown in blue, β-strands in green and loops in white) and BSU1$^{421\text{-}793\ D510N\ H707K}$ (in red, r.m.s.d. is ~0.4 Å and 2.7 Å comparing 295 and 311 corresponding C$_\alpha$ atoms, respectively). Loop regions undergoing conformational changes (in red) upon binding of the phosphorylated C-terminal tail in BSU1$^{421\text{-}793}$ (in orange) are labeled and indicated by red arrows. (*F*) Structural superposition of BSU1$^{421\text{-}793}$ phosphatase domain (residues 450-755, gray molecular surface; residues 756-793, orange ribbon diagram), the HsPP1α – spinophilin complex (in blue, spinophilin residues 425-474, r.m.s.d. is ~0.8 Å comparing 228 corresponding C$_\alpha$ atoms, PDB-ID: pdb_00003egg) (94), and the HsPP1α – Inhibitor-3 complex (in magenta, Inhibitor-3 residues 39-62, r.m.s.d. is ~0.8 Å comparing 245 corresponding C$_\alpha$ atoms, PDB-ID: pdb_00008dwk) (62). The positions of the C-terminal (**C**) , acidic (**A**) and hydrophobic (**H**) binding grooves in PP1 enzymes are indicated (73). (*G*) Close-up view of the BSU1 substrate binding pocket (gray molecular surface) shown together with the phosphorylated C-terminal tail (in orange) and the C-terminus of Inhibitor-3 (in magenta, residues 59-62 shown in bonds representation, PDB-ID: pdb_00008dwk) (62). (*H*) Initial enzyme velocities of the BSU1 phosphatase domain or of the BSU1 Thr785Ala mutant vs a synthetic pThr-containing substrate peptide, and including the derived enzyme kinetic parameters (n = 3).

Fig. 4. The Kelch phosphatase BSLM is a cell cycle regulator in Marchantia. (*A*) Representative thallus phenotypes of 14 d-old Tak-1 and amiRNA-MpBSLM plants grown from gemma, and Mp*bslm*^{ge}, Mp*gsk-3*^{ge} and Mp*bslm*^{ge} Mp*gsk-4*^{ge} propagated from thallus stubs (scale bar = 1 cm). (*B*) Ultrastructural phenotypes of 14 d-old plants. On top, Bright Field + Fluorescence (BF+F) thallus views showing the air pore distribution across the genotypes from A (pore autofluorescence in blue, scale bar = 500 µm). Center: Scanning Electron Microscopy (SEM) views of selected thallus areas (scale bar = 200 µm). Bottom: Technovit7100-fixed histology thallus cross sections stained with toluidine blue (scale bar = 10 µm). (*C*) Heat map of selected up regulated genes putatively involved

in cell cycle regulation, identified by RNAseq comparing 14 d-old Tak-1 and Mp*bslm*[ge], $\log_2$FC > 1 and FDR < 0.01. (*D*) Top: Combined bright-field view (in gray scale) and EdU staining (in green) of 3 d-old thalli (scale bar = 200µm). Bottom: Expression of the *proMpCYCD*::*Dbox-TdTomato* cell cycle reporter (in magenta) in Tak-1, and in Mp*bslm*[ge] or Mp*gsk-3*[ge] double transgenic lines (scale bar = 50 µm). (*E*) Yeast co-expressing MpBSLM fused to the Gal4-activation domain (Gal4; prey) and MpCDK1 (Mp2g04270), MpCKS (Mp1g16700), MpCYCB1 (Mp5g10030), MpGSK (Mp7g04170) or MpRBR (Mp8g18830) fused to the LexA-binding domain (LexA; bait) were grown on selective SD medium supplemented with histidine (-LW; co-transformation control) or lacking histidine (−LWH; interaction assay). Shown are serial dilutions from left to right after 3 d of incubation.

Data, Materials, and Software Availability

The atomic coordinates and structure factors have been deposited with the Protein Data Bank with accession codes pdb_00009TP5 (BSU1[1-357] Kelch), pdb_00009TP7 (BSU1[421-793] phosphatase domain) and pdb_00009TP8 (BSU1[421-793] phosphatase-dead mutant). The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD072713. Raw RNA-seq reads have been deposited with the NCBI sequence read archive (SRA; https://submit.ncbi.nlm.nih.gov/sra) with identifier PRJNA1381677.

Acknowledgments

We thank Martin Bayer for providing *bsk1 bsk2* and *scrm-D* mutant seeds, and Andreas Boland, Silke Hauf, Martin Bayer, Roman Ulm and Philippe Rieu for their helpful comments on the manuscript and Zhiyong Wang for suggesting the BIN2 and CrBSL1 peptide substrates. We thank the University of Lausanne Protein Analysis Facility and the Proteomics Core Facility, Centre Medical Universitaire (CMU), University of Geneva for mass spectrometry analysis, and the staff of the Swiss Light Source (SLS) for technical assistance during data collection. This work was supported by the Swiss National Science Foundation grant 310030_205201 and by an International Research Scholar Award 55008733 by the Howard Hughes Medical Institute (to M.H.).

References

1. M. D. Grove, *et al.*, Brassinolide, a plant growth-promoting steroid isolated from Brassica napus pollen. *Nature* **281**, 216–217 (1979).

2.   N. Vukašinović, T. M. Nolan, E. Russinova, Unlocking the potential of brassinosteroids: A path to precision plant engineering. *Science* **390**, eadu9798 (2025).

3.   S. D. Clouse, M. Langford, T. C. McMorris, A brassinosteroid-insensitive mutant in Arabidopsis thaliana exhibits multiple defects in growth and development. *Plant Physiol* **111**, 671–678 (1996).

4.   J. Li, J. Chory, A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* **90**, 929–938 (1997).

5.   Z. Y. Wang, H. Seto, S. Fujioka, S. Yoshida, J. Chory, BRI1 is a critical component of a plasma-membrane receptor for plant steroids. *Nature* **410**, 380–383 (2001).

6.   A. Caregnato, *et al.*, A mechanistic framework for the recognition of chemically diverse brassinosteroids by BRI1-family receptor kinases. [Preprint] (2025). Available at: https://www.biorxiv.org/content/10.1101/2025.08.08.669299v2 [Accessed 2 December 2025].

7.   M. Hothorn, *et al.*, Structural basis of steroid hormone perception by the receptor kinase BRI1. *Nature* **474**, 467–471 (2011).

8.   J. She, *et al.*, Structural insight into brassinosteroid perception by BRI1. *Nature* **474**, 472–476 (2011).

9.   J. Imkampe, *et al.*, The Arabidopsis Leucine-rich Repeat Receptor Kinase BIR3 Negatively Regulates BAK1 Receptor Complex Formation and Stabilizes BAK1. *The Plant Cell Online* tpc.00376.2017 (2017). https://doi.org/10.1105/tpc.17.00376.

10.   U. Hohmann, J. Nicolet, A. Moretti, L. A. Hothorn, M. Hothorn, The SERK3 elongated allele defines a role for BIR ectodomains in brassinosteroid signalling. *Nat Plants* **4**, 345–351 (2018).

11.   J. Li, *et al.*, BAK1, an Arabidopsis LRR Receptor-like Protein Kinase, Interacts with BRI1 and Modulates Brassinosteroid Signaling. *Cell* **110**, 213–222 (2002).

12.   K. H. Nam, J. Li, BRI1/BAK1, a Receptor Kinase Pair Mediating Brassinosteroid Signaling. *Cell* **110**, 203–212 (2002).

13.   X. Gou, *et al.*, Genetic evidence for an indispensable role of somatic embryogenesis receptor kinases in brassinosteroid signaling. *PLoS Genet.* **8**, e1002452 (2012).

14.   J. Santiago, C. Henzler, M. Hothorn, Molecular mechanism for plant steroid receptor activation by somatic embryogenesis co-receptor kinases. *Science* **341**, 889–892 (2013).

15.   U. Hohmann, *et al.*, Mechanistic basis for the activation of plant membrane receptor kinases by SERK-family co-receptors. *Proc Natl Acad Sci U S A* **115**, 3488–3493 (2018).

16.   X. Wang, *et al.*, Sequential Transphosphorylation of the BRI1/BAK1 Receptor Kinase Complex Impacts Early Events in Brassinosteroid Signaling. *Developmental Cell* **15**, 220–235 (2008).

17.   L. Yan, *et al.*, Structural basis for the impact of phosphorylation on the activation of plant receptor-like kinase BAK1. *Cell Res* **22**, 1304–1308 (2012).

18. D. Bojar, *et al.*, Crystal structures of the phosphorylated BRI1 kinase domain and implications for brassinosteroid signal initiation. *Plant J* **78**, 31–43 (2014).

19. U. Hohmann, *et al.*, Constitutive Activation of Leucine-Rich Repeat Receptor Kinase Signaling Pathways by BAK1-INTERACTING RECEPTOR-LIKE KINASE3 Chimera. *Plant Cell* **32**, 3311–3323 (2020).

20. X. Wang, J. Chory, Brassinosteroids Regulate Dissociation of BKI1, a Negative Regulator of BRI1 Signaling, from the Plasma Membrane. *Science* **313**, 1118–1122 (2006).

21. Y. Jaillais, *et al.*, Tyrosine phosphorylation controls brassinosteroid receptor activation by triggering membrane release of its kinase inhibitor. *Genes Dev* **25**, 232–237 (2011).

22. J. Wang, *et al.*, Structural insights into the negative regulation of BRI1 signaling by BRI1-interacting protein BKI1. *Cell Res* **24**, 1328–1341 (2014).

23. H. Wang, *et al.*, Dual role of BKI1 and 14-3-3 s in brassinosteroid signaling to link receptor with transcription factors. *Dev Cell* **21**, 825–834 (2011).

24. E. Obergfell, U. Hohmann, A. Moretti, H. Chen, M. Hothorn, Mechanistic Insights into the Function of 14-3-3 Proteins as Negative Regulators of Brassinosteroid Signaling in Arabidopsis. *Plant Cell Physiol* **65**, 1674–1688 (2024).

25. K. Caesar, *et al.*, A fast brassinolide-regulated response pathway in the plasma membrane of Arabidopsis thaliana. *Plant J* **66**, 528–540 (2011).

26. W. Tang, *et al.*, BSKs mediate signal transduction from the receptor kinase BRI1 in Arabidopsis. *Science* **321**, 557–560 (2008).

27. C. Grütter, S. Sreeramulu, G. Sessa, D. Rauh, Structural characterization of the RLCK family member BSK8: a pseudokinase with an unprecedented architecture. *J. Mol. Biol.* **425**, 4455–4467 (2013).

28. T. Niittylä, A. T. Fuglsang, M. G. Palmgren, W. B. Frommer, W. X. Schulze, Temporal analysis of sucrose-induced phosphorylation changes in plasma membrane proteins of Arabidopsis. *Mol Cell Proteomics* **6**, 1711–1726 (2007).

29. S. Sreeramulu, *et al.*, BSKs are partially redundant positive regulators of brassinosteroid signaling in Arabidopsis. *The Plant Journal* **74**, 905–919 (2013).

30. H. Ren, *et al.*, BRASSINOSTEROID-SIGNALING KINASE 3, a plasma membrane-associated scaffold protein involved in early brassinosteroid signaling. *PLoS Genet* **15**, e1007904 (2019).

31. J. Yan, *et al.*, BSK family kinases are essential for brassinosteroid signaling and suppression of adventitious rooting by repressing the expression of LBD16. *New Phytol* **250**, 283–297 (2026).

32. H. Muto, N. Yabe, T. Asami, K. Hasunuma, K. T. Yamamoto, Overexpression of constitutive differential growth 1 gene, which encodes a RLCKVII-subfamily protein kinase, causes abnormal differential and elongation growth after organ differentiation in Arabidopsis. *Plant Physiol* **136**, 3124–3133 (2004).

33. T.-W. Kim, S. Guan, A. L. Burlingame, Z.-Y. Wang, The CDG1 Kinase Mediates Brassinosteroid Signal Transduction from BRI1 Receptor Kinase to BSU1 Phosphatase and GSK3-like Kinase BIN2. *Molecular Cell* **43**, 561–571 (2011).

34. T.-W. Kim, *et al.*, Brassinosteroid signal transduction from cell-surface receptor kinases to nuclear transcription factors. *Nat. Cell Biol.* **11**, 1254–1260 (2009).

35. D. Weigel, *et al.*, Activation tagging in Arabidopsis. *Plant Physiol* **122**, 1003–1013 (2000).

36. S. Mora-García, *et al.*, Nuclear protein phosphatases with Kelch-repeat domains modulate the response to brassinosteroids in Arabidopsis. *Genes Dev.* **18**, 448–460 (2004).

37. G. A. Maselli, *et al.*, Revisiting the evolutionary history and roles of protein phosphatases with Kelch-like domains in plants. *Plant Physiol.* **164**, 1527–1541 (2014).

38. N. Philip, H. J. Vaikkinen, L. Tetley, A. P. Waters, A unique Kelch domain phosphatase in Plasmodium regulates ookinete morphology, motility and invasion. *PLoS One* **7**, e44617 (2012).

39. E.-J. Kim, S.-H. Lee, C.-H. Park, T.-W. Kim, Functional Role of BSL1 Subcellular Localization in Brassinosteroid Signaling. *J. Plant Biol.* **61**, 40–49 (2018).

40. E.-J. Kim, *et al.*, Oligomerization between BSU1 Family Members Potentiates Brassinosteroid Signaling in Arabidopsis. *Mol Plant* **9**, 178–181 (2016).

41. J. Li, K. H. Nam, Regulation of brassinosteroid signaling by a GSK3/SHAGGY-like kinase. *Science* **295**, 1299–1301 (2002).

42. K. Hughes, E. Nikolakaki, S. E. Plyte, N. F. Totty, J. R. Woodgett, Modulation of the glycogen synthase kinase-3 family by tyrosine phosphorylation. *EMBO J* **12**, 803–808 (1993).

43. A. Cole, S. Frame, P. Cohen, Further evidence that the tyrosine phosphorylation of glycogen synthase kinase-3 (GSK3) in mammalian cells is an autophosphorylation event. *Biochem J* **377**, 249–255 (2004).

44. Z. Y. Wang, *et al.*, Nuclear-localized BZR1 mediates brassinosteroid-induced growth and feedback suppression of brassinosteroid biosynthesis. *Dev. Cell* **2**, 505–513 (2002).

45. Y. Yin, *et al.*, BES1 accumulates in the nucleus in response to brassinosteroids to regulate gene expression and promote stem elongation. *Cell* **109**, 181–191 (2002).

46. J. Zhao, *et al.*, Two putative BIN2 substrates are nuclear components of brassinosteroid signaling. *Plant Physiol.* **130**, 1221–1229 (2002).

47. Y. Yin, *et al.*, A new class of transcription factors mediates brassinosteroid-regulated gene expression in Arabidopsis. *Cell* **120**, 249–259 (2005).

48. J.-X. He, *et al.*, BZR1 is a transcriptional repressor with dual roles in brassinosteroid homeostasis and growth responses. *Science* **307**, 1634–1638 (2005).

49. G. Vert, J. Chory, Downstream nuclear events in brassinosteroid signalling. *Nature* **441**, 96–100 (2006).

50. L.-G. Chen, *et al.*, BZR1 Family Transcription Factors Function Redundantly and Indispensably in BR Signaling but Exhibit BRI1-Independent Function in Regulating Anther Development in Arabidopsis. *Mol Plant* **12**, 1408–1415 (2019).

51. T.-W. Kim, M. Michniewicz, D. C. Bergmann, Z.-Y. Wang, Brassinosteroid regulates stomatal development by GSK3-mediated inhibition of a MAPK pathway. *Nature* **482**, 419–422 (2012).

52. X. Guo, X. Ding, J. Dong, Dichotomy of the BSL phosphatase signaling spatially regulates MAPK components in stomatal fate determination. *Nat Commun* **13**, 2438 (2022).

53. X. Guo, C. H. Park, Z.-Y. Wang, B. E. Nickels, J. Dong, A spatiotemporal molecular switch governs plant asymmetric cell division. *Nat Plants* **7**, 667–680 (2021).

54. C. H. Park, *et al.*, Deconvoluting signals downstream of growth and immune receptor kinases by phosphocodes of the BSU1 family phosphatases. *Nat Plants* **8**, 646–655 (2022).

55. D. G. O. Saunders, *et al.*, Host protein BSL1 associates with Phytophthora infestans RXLR effector AVR2 and the Solanum demissum Immune receptor R2 to mediate disease resistance. *Plant Cell* **24**, 3420–3434 (2012).

56. D. Turnbull, *et al.*, AVR2 Targets BSL Family Members, Which Act as Susceptibility Factors to Suppress Host Immunity. *Plant Physiol* **180**, 571–581 (2019).

57. F. Tulin, F. R. Cross, Cyclin-Dependent Kinase Regulation of Diurnal Transcription in Chlamydomonas. *Plant Cell* **27**, 2727–2742 (2015).

58. F. Tulin, *et al.*, Mitotic entry is controlled by the plant-specific phosphatase BSL1 and cyclin-dependent kinase B. *Nat Plants* **11**, 2395–2408 (2025).

59. T. M. Nolan, N. Vukašinović, D. Liu, E. Russinova, Y. Yin, Brassinosteroids: Multidimensional Regulators of Plant Growth, Development, and Stress Responses. *Plant Cell* **32**, 295–318 (2020).

60. S. Dharmaiah, *et al.*, Structural basis for LZTR1 recognition of RAS GTPases for degradation. *Science* **389**, 1112–1117 (2025).

61. D. C. Scott, *et al.*, Structural basis for C-degron selectivity across KLHDCX family E3 ubiquitin ligases. *Nat Commun* **15**, 9899 (2024).

62. G. Srivastava, M. S. Choy, N. Bolik-Coulon, R. Page, W. Peti, Inhibitor-3 inhibits Protein Phosphatase 1 via a metal binding dynamic protein-protein interaction. *Nat Commun* **14**, 1798 (2023).

63. J. Zhang, Z. Zhang, K. Brew, E. Y. Lee, Mutational analysis of the catalytic subunit of muscle protein phosphatase-1. *Biochemistry* **35**, 6276–6282 (1996).

64. L. Holm, P. Rosenström, Dali server: conservation mapping in 3D. *Nucl. Acids Res.* **38**, W545–W549 (2010).

65. J. Goldberg, *et al.*, Three-dimensional structure of the catalytic subunit of protein serine/threonine phosphatase-1. *Nature* **376**, 745–753 (1995).

66. M. P. Egloff, P. T. Cohen, P. Reinemer, D. Barford, Crystal structure of the catalytic subunit of human protein phosphatase 1 and its complex with tungstate. *J Mol Biol* **254**, 942–959 (1995).

67. M. Bollen, W. Peti, M. J. Ragusa, M. Beullens, The extended PP1 toolkit: designed to create specificity. *Trends Biochem Sci* **35**, 450–458 (2010).

68. J. B. Hein, *et al.*, Phosphatase specificity principles uncovered by MRBLE:Dephos and global substrate identification. *Mol Syst Biol* **19**, e11782 (2023).

69. J. Abramson, *et al.*, Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493–500 (2024).

70. E. Heroes, *et al.*, Metals in the active site of native protein phosphatase-1. *J Inorg Biochem* **149**, 1–5 (2015).

71. M. P. Egloff, *et al.*, Structural basis for the recognition of regulatory subunits by the catalytic subunit of protein phosphatase 1. *EMBO J* **16**, 1876–1887 (1997).

72. M. Terrak, F. Kerff, K. Langsetmo, T. Tao, R. Dominguez, Structural basis of protein phosphatase 1 regulation. *Nature* **429**, 780–784 (2004).

73. W. Peti, A. C. Nairn, R. Page, Structural Basis for Protein Phosphatase 1 Regulation and Specificity. *FEBS J* **280**, 596–611 (2013).

74. J. A. Gibbons, D. C. Weiser, S. Shenolikar, Importance of a surface hydrophobic pocket on protein phosphatase-1 catalytic subunit in recognizing cellular regulators. *J Biol Chem* **280**, 15903–15911 (2005).

75. T. Asami, *et al.*, Characterization of brassinazole, a triazole-type brassinosteroid biosynthesis inhibitor. *Plant Physiol.* **123**, 93–100 (2000).

76. X. Zhang, *et al.*, A Temperature-Sensitive Misfolded bri1-301 Receptor Requires Its Kinase Activity to Promote Growth. *Plant Physiology* **178**, 1704–1719 (2018).

77. D. M. Friedrichsen, C. A. Joazeiro, J. Li, T. Hunter, J. Chory, Brassinosteroid-insensitive-1 is a ubiquitously expressed leucine-rich repeat receptor serine/threonine kinase. *Plant Physiol* **123**, 1247–1256 (2000).

78. E. Russinova, *et al.*, Heterodimerization and Endocytosis of Arabidopsis Brassinosteroid Receptors BRI1 and AtSERK3 (BAK1). *Plant Cell* **16**, 3216–3229 (2004).

79. T. M. Nolan, *et al.*, Brassinosteroid gene regulatory networks at cellular resolution in the Arabidopsis root. *Science* **379**, eadf4721 (2023).

80. N. Blanco-Touriñán, *et al.*, The brassinosteroid receptor gene BRI1 safeguards cell-autonomous brassinosteroid signaling across tissues. *Sci Adv* **10**, eadq3352 (2024).

81. A. Caño-Delgado, *et al.*, BRL1 and BRL3 are novel brassinosteroid receptors that function in vascular differentiation in Arabidopsis. *Development* **131**, 5341–5351 (2004).

82. S. S. Gampala, *et al.*, An essential role for 14-3-3 proteins in brassinosteroid signal transduction in Arabidopsis. *Dev. Cell* **13**, 177–189 (2007).

83. H. Ryu, *et al.*, Nucleocytoplasmic shuttling of BZR1 mediated by phosphorylation is essential in Arabidopsis brassinosteroid signaling. *Plant Cell* **19**, 2749–2762 (2007).

84. C. W. Y. Liu, *et al.*, Inhibitory Phosphorylation of PP1α Catalytic Subunit during the G1/S Transition *. *Journal of Biological Chemistry* **274**, 29470–29475 (1999).

85. M. Dohadwala, *et al.*, Phosphorylation and inactivation of protein phosphatase 1 by cyclin-dependent kinases. *Proc Natl Acad Sci U S A* **91**, 6408–6412 (1994).

86. H. Yamano, K. Ishii, M. Yanagida, Phosphorylation of dis2 protein phosphatase at the C-terminal cdc2 consensus and its potential role in cell cycle regulation. *EMBO J* **13**, 5310–5318 (1994).

87. Y. G. Kwon, S. Y. Lee, Y. Choi, P. Greengard, A. C. Nairn, Cell cycle-dependent phosphorylation of mammalian protein phosphatase 1 by cdc2 kinase. *Proc Natl Acad Sci U S A* **94**, 2168–2173 (1997).

88. J. Yu, *et al.*, Structural basis of human separase regulation by securin and CDK1-cyclin B1. *Nature* **596**, 138–142 (2021).

89. N. Berndt, M. Dohadwala, C. W. Liu, Constitutively active protein phosphatase 1alpha causes Rb-dependent G1 arrest in human cancer cells. *Curr Biol* **7**, 375–386 (1997).

90. J. Q. Wu, *et al.*, PP1-mediated dephosphorylation of phosphoproteins at mitotic exit is controlled by inhibitor-1 and PP1 phosphorylation. *Nat Cell Biol* **11**, 644–651 (2009).

91. F. Salvi, *et al.*, Towards Dissecting the Mechanism of Protein Phosphatase-1 Inhibition by Its C-Terminal Phosphorylation. *Chembiochem* **22**, 834–838 (2021).

92. J. T. Maynes, *et al.*, Crystal structure of the tumor-promoter okadaic acid bound to protein phosphatase-1. *J Biol Chem* **276**, 44078–44082 (2001).

93. J. H. Connor, T. Kleeman, S. Barik, R. E. Honkanen, S. Shenolikar, Importance of the beta12-beta13 loop in protein phosphatase-1 catalytic subunit for inhibition by toxins and mammalian protein inhibitors. *J Biol Chem* **274**, 22366–22372 (1999).

94. M. J. Ragusa, *et al.*, Spinophilin directs protein phosphatase 1 specificity by blocking substrate binding sites. *Nat Struct Mol Biol* **17**, 459–464 (2010).

95. M. A. Mecchia, *et al.*, The BES1/BZR1-family transcription factor MpBES1 regulates cell division and differentiation in Marchantia polymorpha. *Curr Biol* **31**, 4860-4869.e8 (2021).

96. T. Furuya, *et al.*, A glycogen synthase kinase 3-like kinase MpGSK regulates cell differentiation in Marchantia polymorpha. *Plant Biotechnol (Tokyo)* **39**, 65–72 (2022).

97. Y. Saidi, T. J. Hearn, J. C. Coates, Function and evolution of "green" GSK3/Shaggy-like kinases. *Trends Plant Sci* **17**, 39–46 (2012).

98. T. Furuya, *et al.*, ANGUSTIFOLIA contributes to the regulation of three-dimensional morphogenesis in the liverwort Marchantia polymorpha. *Development* **145**, dev161398 (2018).

99. J.-X. He, J. M. Gendron, Y. Yang, J. Li, Z.-Y. Wang, The GSK3-like kinase BIN2 phosphorylates and destabilizes BZR1, a positive regulator of the brassinosteroid signaling pathway in Arabidopsis. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 10185–10190 (2002).

100. S. Lemaire, *et al.*, PPP1R2 stimulates protein phosphatase-1 through stabilisation of dynamic subunit interactions. *Nat Commun* **15**, 9822 (2024).

101. S. Rogers, *et al.*, PP1 initiates the dephosphorylation of MASTL, triggering mitotic exit and bistability in human cells. *J Cell Sci* **129**, 1340–1354 (2016).

102. Q. Lin, J. Li, R. D. Smith, J. C. Walker, Molecular cloning and chromosomal mapping of type one serine/threonine protein phosphatases in Arabidopsis thaliana. *Plant Mol Biol* **37**, 471–481 (1998).

103. W. Kabsch, Automatic processing of rotation diffraction data from crystals of initially unknown symmetry land cell constants. *Journal of Applied Crystallography* **26**, 795–800 (1993).

104. W. Zhang, *et al.*, Crystal structure of the nitrile-specifier protein NSP1 from Arabidopsis thaliana. *Biochem Biophys Res Commun* **488**, 147–152 (2017).

105. A. Perrakis, M. Harkiolaki, K. S. Wilson, V. S. Lamzin, ARP/wARP and molecular replacement. *Acta Crystallogr D Biol Crystallogr* **57**, 1445–1450 (2001).

106. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Cryst D* **60**, 2126–2132 (2004).

107. P. V. Afonine, *et al.*, Towards automated crystallographic structure refinement with phenix.refine. *Acta Cryst D* **68**, 352–367 (2012).

108. P. H. Zwart, R. W. Grosse-Kunstleve, P. D. Adams, Xtriage and Fest: automatic assessment of X-ray data and substructure structure factor estimation. *CCP4 newsletter* (2005).

109. G. M. Sheldrick, A short history of SHELX. *Acta Crystallogr., A, Found. Crystallogr* **64**, 112–122 (2008).

110. C. Vonrhein, E. Blanc, P. Roversi, G. Bricogne, Automated structure solution with autoSHARP. *Methods Mol Biol* **364**, 215–230 (2007).

111. T. C. Terwilliger, SOLVE and RESOLVE: automated structure solution and density modification. *Meth. Enzymol.* **374**, 22–37 (2003).

112. T. C. Terwilliger, *et al.*, Iterative model building, structure refinement and density modification with the PHENIX AutoBuild wizard. *Acta Crystallogr. D Biol. Crystallogr* **64**, 61–69 (2008).

113. A. J. McCoy, *et al.*, Phaser crystallographic software. *J Appl Cryst* **40**, 658–674 (2007).

114. I. W. Davis, *et al.*, MolProbity: all-atom contacts and structure validation for proteins and nucleic acids. *Nucleic Acids Res* **35**, W375-383 (2007).

115. E. C. Meng, *et al.*, UCSF ChimeraX: Tools for structure building and analysis. *Protein Science* **32**, e4792 (2023).

116. D. Chiasson, *et al.*, A unified multi-kingdom Golden Gate cloning platform. *Sci Rep* **9**, 10131 (2019).

117. S. J. Clough, A. F. Bent, Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J.* **16**, 735–743 (1998).

118. M. Jinek, *et al.*, A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821 (2012).

119. M. K. Singh, *et al.*, A single class of ARF GTPase activated by several pathway-specific ARF-GEFs regulates essential membrane traffic in Arabidopsis. *PLoS Genet* **14**, e1007795 (2018).

120. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

121. R Core Team, *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2013* (ISBN 3-900051-07-0, 2014).

122. A. Kitsche, L. A. Hothorn, Testing for qualitative interaction using ratios of treatment differences. *Stat Med* **33**, 1477–1489 (2014).

123. T. Hothorn, F. Bretz, P. Westfall, Simultaneous inference in general parametric models. *Biom J* **50**, 346–363 (2008).

124. C. W. Dunnett, A Multiple Comparison Procedure for Comparing Several Treatments with a Control. *Journal of the American Statistical Association* **50**, 1096–1121 (1955).

125. M. M. Kanaoka, *et al.*, SCREAM/ICE1 and SCREAM2 Specify Three Cell-State Transitional Steps Leading to Arabidopsis Stomatal Differentiation. *Plant Cell* **20**, 1775–1785 (2008).

126. A. Neu, *et al.*, Constitutive signaling activity of a receptor-associated protein links fertilization with embryonic patterning in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* **116**, 5795–5804 (2019).

127. H. Chen, *et al.*, Phosphorylation-dependent activation of the bHLH transcription factor ICE1/SCRM promotes polarization of the Arabidopsis zygote. *New Phytol* **245**, 1029–1039 (2025).

128. J. Mergner, *et al.*, Mass-spectrometry-based draft of the Arabidopsis proteome. *Nature* **579**, 409–414 (2020).

129. A. Shevchenko, H. Tomas, J. Havlis, J. V. Olsen, M. Mann, In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nat Protoc* **1**, 2856–2860 (2006).

130. F. Meier, *et al.*, Online Parallel Accumulation-Serial Fragmentation (PASEF) with a Novel Trapped Ion Mobility Mass Spectrometer. *Mol Cell Proteomics* **17**, 2534–2545 (2018).

131. L. Käll, J. D. Storey, W. S. Noble, Non-parametric estimation of posterior error probabilities associated with peptides identified by tandem mass spectrometry. *Bioinformatics* **24**, i42-48 (2008).

132. A. I. Nesvizhskii, A. Keller, E. Kolker, R. Aebersold, A statistical model for identifying proteins by tandem mass spectrometry. *Anal Chem* **75**, 4646–4658 (2003).

133. K. Ishizaki, S. Chiyoda, K. T. Yamato, T. Kohchi, Agrobacterium-Mediated Transformation of the Haploid Liverwort Marchantia polymorpha L., an Emerging Model for Plant Biology. *Plant and Cell Physiology* **49**, 1084–1091 (2008).

134. F. Laurent, *et al.*, Inositol pyrophosphate catabolism by three families of phosphatases regulates plant growth and development. *PLOS Genetics* **20**, e1011468 (2024).

135. S. S. Sugano, *et al.*, Efficient CRISPR/Cas9-based genome editing and its application to conditional genetic analysis in Marchantia polymorpha. *PLoS One* **13**, e0205117 (2018).

136. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

137. J. L. Bowman, *et al.*, Insights into Land Plant Evolution Garnered from the Marchantia polymorpha Genome. *Cell* **171**, 287-304.e15 (2017).

138. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

139. M. Pertea, D. Kim, G. M. Pertea, J. T. Leek, S. L. Salzberg, Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11**, 1650–1667 (2016).

140. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

141. N. O'Connell, *et al.*, The molecular basis for substrate specificity of the nuclear NIPP1:PP1 holoenzyme. *Structure* **20**, 1746–1756 (2012).

142. T. Noguchi, *et al.*, Brassinosteroid-insensitive dwarf mutants of Arabidopsis accumulate brassinosteroids. *Plant Physiol* **121**, 743–752 (1999).

143. W. Kabsch, C. Sander, Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **22**, 2577–2637 (1983).

144. W. Kabsch, Automatic processing of rotation diffraction data from crystals of initially unknown symmetry and cell constants. *Journal of Applied Crystallography* **26**, 795–800 (1993).

145. G. Bricogne, C. Vonrhein, C. Flensburg, M. Schiltz, W. Paciorek, Generation, representation and flow of phase information in structure determination: recent developments in and around SHARP 2.0. *Acta Crystallogr. D Biol. Crystallogr* **59**, 2023–2030 (2003).

Fig. 1. BSU1 is a PP1-like Ser/Thr phosphatase. (*A*) Schematic representation of *A. thaliana* BSU1, depicting the Kelch (residues 11–351) and phosphatase (residues 450–793) domains (https:// www.uniprot.org/ Uniprot ID Q9LR78). Structured regions are shown as blocks, putative linker regions (residues 1–10 and 352–449) as solid line, CDG1 phosphorylation sites are highlighted in red, the BIK1 phosphorylation site in blue. (*B*) Top-view of the BSU1 Kelch domain. Shown is a ribbon diagram colored with a rainbow gradient from blue (N-term) to red (C-term). Kelch motif blades are labeled I to VI, an intrinsically disordered loop is indicated by a dotted line, the BIK1-mediated Ser251 phosphorylation site is shown alongside (in bonds representation, marked by a blue circle). (*C*) Ribbon diagram of the BSU1 phosphatase domain (chain A), with α-helices in blue, β-strands in green, loops in white, and Zn^{2+} ions as gray spheres. The C-terminal tail is shown in dark gray, and Ser764 is depicted in bonds representation, marked by a red circle. (*D*) Close-up view of the active sites of wild-type BSU1 (left) and the catalytically inactive variant BSU1$^{421-793 \; D510N \; H707K}$ (right). Zn^{2+} ions are shown as green spheres, the distorted trigonal bipyramidal pentavalent coordination is indicated by dotted lines (in magenta), water molecules are shown as red spheres. A phased anomalous difference map calculated from data collected at the Zn K-edge and contoured at 25 σ is shown as a green mesh. (E) Initial enzyme velocities for BSU1-catalyzed hydrolysis of para-nitrophenyl phosphate (*p*-NPP), plotted as a function of substrate concentration. Data were fitted to the Michaelis-Menten equation by nonlinear regression. Extracted kinetic parameters are summarized in panel K (n = 3). (F) Metal ion preference of the phosphatase domain using 2 mM *p*-NPP as substrate. Each metal chloride was tested at a fixed concentration of 2 mM. (G) BIN2 pTyr dephosphorylation time course experiment using 20 μM BIN2 and 5 μM of either the isolated BSU1 phosphatase domain (top) or the full-length enzyme (bottom). Shown are a immunoblots and coomassie-stained membranes below as loading controls. The presence of BIN2 pTyr200 was detected using a human GSK3 pTyr antibody (n = 2). (H) Structural superposition of the BSU1 phosphatase domain (cyan Cα trace) and HsPP1γ (in orange, PDB-ID: pdb_00001jk7 (92), r.m.s.d. is ~0.8 Å comparing 245 corresponding Cα atoms). (*I*) Close-up view of the superimposed BSU1 (in cyan) and HsPP1γ (in orange) active sites. Shown are the catalytic Zn^{2+}/Mn^{2+} ions (in purple), with coordinating residues depicted in bonds representation. (*J*) Initial enzyme velocities of BSU1 phosphatase domain (or phosphatase-dead) catalyzed hydrolysis of synthetic substrate peptides containing either a pThr (left), pSer (center) or pTyr (right), or of the natural substrate candidates BIN2 and CrCDKB1 (bottom row), using a malachite green-based phosphatase assay (n = 3). (*K*) Table summaries of the enzyme kinetic parameters derived from the experiments shown in E and J (n.d., no activity detected). (*L*) Metal ion preference of the phosphatase domain vs. synthetic substrate peptides containing either a pThr or pTyr. Assays were performed using 200 μM of peptide substrate and 50 nM of enzyme in the presence of 2 mM of the indicated metal chloride. In the case of ZnCl₂ + Fe(II)Cl₂ a final concentration of 500 μM was used (n = 3). (*M*) Close-up view of the RVxF peptide binding grooves in BSU1 (in cyan) and HsPP1α (in orange, PDB-ID: pdb_00003v4y (141), r.m.s.d. is ~0.8 Å comparing 252 corresponding Cα atoms). Shown is a ribbon diagram of the peptide binding groove with substrate peptide-interacting amino-acid side-chains shown in bonds representation. (*N*) Molecular surface view of the BSU1 RVxF peptide binding groove (in cyan) bound to the TEV recognition motif (in yellow, in surface and bonds representation) of a neighboring molecule in the asymmetric unit.

A

pro35S::CDS-3xFlag / bri1-5

Ws-2 bri1-5 bsu1-1D BSU1^421-793 BSU1^421-793 D510N H707K BSU1^421-793 M748R I750R BSU1^421-793 D510N H707K M748R I750R BSL2

Ws-2 | Ws-2 | Ws-2 | Ws-2 | Ws-2

115-
70-
50-
30-

B

	normalized protein abundances (millions) ±SE		
pro35S::CDS-3xFlag / bri1-5	BSL1	BSL2	BSL3
BSU1^1-357	0.12±0.02	0.14±0.03	0.38±0.09
BSU1^421-793	250.21±12.06	47.16±1.26	74.81±3.66
BSU1^421-793 D510N H707K	2.60±0.09	17.12±0.19	23.14±0.74

C

Gal4 empty - LexA empty
Gal4 empty - LexA AtBSL2
Gal4 AtBSU1^421-793 - LexA empty
Gal4 AtBSU1^421-793 - LexA AtBSL2
Gal4 AtBSU1^421-793 M748R I750R - LexA empty
Gal4 AtBSU1^421-793 M748R I750R - LexA AtBSL2

SD -LW   SD -LWH

D

mock

+BRZ

mock
+BRZ

rel. inhib.

Col-0 bsk3-1 bri1-301 bsu1 bsu1 bsl1 bsl3 bsu1 bsl1 bsl2

E

mock

+BL

Col-0 bsk3-1 bri1-301 bsu1 bsu1 bsl1 bsl3 bsu1 bsl1 bsl2

*** *** * *** * ***

F

100%
50%
0%

100%
50%
0%

proBSU1::BSU1-GUS
proBSU1::GUS
proBRI1::GUS

G

H629 D544 D510
H577
N576
H707
pTM 0.83
ipTM 0.91
H512
Y731
R679
BZR1
N
R548 pS173

H

V_max = 14.9 µmol min⁻¹ mg⁻¹
K_M = 0.16 mM
k_cat = 9.5 s⁻¹

RLRISN(pS)CPVTPRY (µM)

Fig. 2. A catalytically inactive BSU1 phosphatase domain can suppress the growth phenotype of the *bri1-5* mutant. (*A*) Inflorescence phenotypes of 45 d-old *bri1-5* plants expressing different BSU1 variants, or full-length BSL2 from the 35S promoter. The Ws-2 wild type, *bri1-5* (136) and the original *bsu1-1D* (35) allele are shown alongside. BSU1/BSL2 protein expression was assayed by western blot using an α-Flag antibody, the Ponceau-stained membrane is shown below as loading control. (*B*) Relative abundance of BLS1, BSL2 and BSL3 in AP-MS experiments with the Flag-tagged BSU1 Kelch (top), phosphatase (center) or phosphatase-dead (bottom) domains. (*C*) Yeast co-expressing either the wild-type BSU1 phosphatase domain or the mutant carrying point mutations in the RVxF binding groove fused to the Gal4-activation domain (Gal4; prey) and full-length wild-type BSL2 fused to the LexA-binding domain (LexA; bait) were grown on selective SD medium supplemented with histidine (-LW; co-transformation control) or lacking histidine (−LWH; interaction assay). Shown are serial dilutions from left to right after 2d of incubation. (*D*) Hypocotyl growth assay of 5 d-old dark-grown seedlings in the presence and absence of the BR biosynthesis inhibitor BRZ (73). Representative seedlings are shown in the top panel (scale bar = 0.5cm). Box plots (bold black line, median; box, interquartile range (IQR); whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile) are in the center panel with raw data shown as individual points (n=30). Shown below is relative inhibition of hypocotyl growth in the presence of BRZ plotted together with lower and upper confidence intervals, the dotted horizontal line depicts the Col-0 wild-type reference (rel. inhib., relative inhibition). (*E*) Brassinolide (BL)-mediated root growth inhibition assay. Shown are representative 7 d-old seedlings, grown on $^{1/2}$MS plates (mock, top panel) or in the presence of 100 nM BL (+BL, bottom panel). Quantifications are shown below as box plots with raw data indicated as individual points (n=15) and including Dunnett-type (119), two-sided multiplicity-adjusted *p*-values for the comparison against the Col-0 control (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). (*F*) Spatiotemporal expression patterns of BSU1 and BRI1 reporter lines determined by β-glucuronidase activity (GUS) assays. A sequence ~1kb upstream of the BSU1 ORF was used to drive *proBSU1*::BSU1-GUS and *proBSU1*::*GUS*, respectively. A ~2kb upstream of BRI1 ORF was used to drive *proBRI1*::*GUS*. 14-17 independent T2 lines for each construct were stained for GUS activity, either using 5 d-old cotyledons (top), or the 4th leaf of 15 d-old soil-grown plants. Shown are distribution star plots with expression patterns divided into categories. (*G*) AlphaFold3-predicted complex structure of $BSU1^{421-793}$ (ribbon diagram) bound to Mn^{2+} ions (purple spheres) and $pBZR1^{168-178}pS$ (in orange). Predicted salt bridges between the phosphate group and BSU1 residues are depicted as dotted lines (in green). AlphaFold3 pTM and ipTM values are reported alongside. (*H*) Initial enzyme velocities of BSU1 phosphatase domain catalyzed hydrolysis of the synthetic $pBZR1^{168-178}pS$ substrate peptide and including enzyme kinetic parameters (n = 3).

Fig. 3. The C-terminal tail of BSU1 is phosphorylated by the CDK1 – cyclin B1 – CKS1 complex. (*A*) Molecular surface view of the BSU1 phosphatase domain (residues 450-755), the C-terminal tail (residues 756-793) is shown as a ribbon diagram (in orange), pThr785 is inserted into the active site (in bonds representation). The relative positions of two substrate-binding loops are indicated. The inset provides a close-up view of the BSU1 active site with two divalent metal ions and the catalytic water molecule shown as purple and red spheres, respectively. Residues contacting pThr785 (in orange) are highlighted in bonds representation (in blue), hydrogen bonds are shown as green dotted lines. The closest distances between the phosphate moiety of pThr785 and the catalytic water molecule or the general base His577 are indicated by yellow dotted lines. (*B*) MS1 spectrum of the phosphorylated BSU1[755-788] peptide. (*C*) Multiple sequence alignment of the C-terminal tail of BSU1, BSL1-3, two PP1-like TOPP phosphatases from *Arabidopsis thaliana*, and human and fission yeast PP1. The position of the PxTPP motif containing the pThr site is indicated (in red). (*D*) Autoradiograph of an in vitro transphosphorylation time course assay. 10 nM of the human CDK1 – cyclin B1 - CKS1 complex were incubated with 10 µM BSU1421-793 or with BSU1421-793 T785A, where Thr785 in the PxTPP motif was replaced by an alanine, at 37 °C for the indicated time. (*E*) Structural superposition of BSU1[421-793] (ribbon diagram with α-helices shown in blue, β-strands in green and loops in white) and BSU1[421-793' D510N H707K] (in red, r.m.s.d. is ~0.4 Å and 2.7 Å comparing 295 and 311 corresponding Cα atoms, respectively). Loop regions undergoing conformational changes (in red) upon binding of the phosphorylated C-terminal tail in BSU1[421-793] (in orange) are labeled and indicated by red arrows. (*F*) Structural superposition of BSU1[421-793] phosphatase domain (residues 450-755, gray molecular surface; residues 756-793, orange ribbon diagram), the HsPP1α – spinophilin complex (in blue, spinophilin residues 425-474, r.m.s.d. is ~0.8 Å comparing 228 corresponding Cα atoms, PDB-ID: pdb_00003egg) (94), and the HsPP1α – Inhibitor-3 complex (in magenta, Inhibitor-3 residues 39-62, r.m.s.d. is ~0.8 Å comparing 245 corresponding Cα atoms, PDB-ID: pdb_00008dwk) (62). The positions of the C-terminal (C) , acidic (A) and hydrophobic (H) binding grooves in PP1 enzymes are indicated (73). (*G*) Close-up view of the BSU1 substrate binding pocket (gray molecular surface) shown together with the phosphorylated C-terminal tail (in orange) and the C-terminus of Inhibitor-3 (in magenta, residues 59-62 shown in bonds representation, PDB-ID: pdb_00008dwk) (62). (*H*) Initial enzyme velocities of the BSU1 phosphatase domain or of the BSU1 Thr785Ala mutant vs a synthetic pThr-containing substrate peptide, and including the derived enzyme kinetic parameters (n = 3).

Fig. 4. The Kelch phosphatase BSLM is a cell cycle regulator in Marchantia. (*A*) Representative thallus phenotypes of 14 d-old Tak-1 and amiRNA-MpBSLM plants grown from gemma, and Mp*bslm*ge, Mp*gsk-3*ge and Mp*bslm*ge Mp*gsk-4*ge propagated from thallus stubs (scale bar = 1 cm). (*B*) Ultrastructural phenotypes of 14 d-old plants. On top, Bright Field + Fluorescence (BF+F) thallus views showing the air pore distribution across the genotypes from A (pore autofluorescence in blue, scale bar = 500 μm). Center: Scanning Electron Microscopy (SEM) views of selected thallus areas (scale bar = 200 μm). Bottom: Technovit7100-fixed histology thallus cross sections stained with toluidine blue (scale bar = 10 μm). (*C*) Heat map of selected up regulated genes putatively involved in cell cycle regulation, identified by RNAseq comparing 14 d-old Tak-1 and Mp*bslm*ge, log$_2$FC > 1 and FDR < 0.01. (*D*) Top: Combined bright-field view (in gray scale) and EdU staining (in green) of 3 d-old thalli (scale bar = 200μm). Bottom: Expression of the *proMpCYCD::Dbox-TdTomato* cell cycle reporter (in magenta) in Tak-1, and in Mp*bslm*ge or Mp*gsk-3*ge double transgenic lines (scale bar = 50 μm). (*E*) Yeast co-expressing MpBSLM fused to the Gal4-activation domain (Gal4; prey) and MpCDK1 (Mp2g04270), MpCKS (Mp1g16700), MpCYCB1 (Mp5g10030), MpGSK (Mp7g04170) or MpRBR (Mp8g18830) fused to the LexA-binding domain (LexA; bait) were grown on selective SD medium supplemented with histidine (-LW; co-transformation control) or lacking histidine (−LWH; interaction assay). Shown are serial dilutions from left to right after 3 d of incubation.

Supporting Information for

Plant Kelch phosphatases are Ser/Thr phosphatases involved in cell cycle regulation

Felix Rico-Resendiz[1,*], Oded Pri-Tal[1,*], Pierre Raia[1,*], Andrea Moretti[1,*,$], Houming Chen[1], Jun Yu[2], Larissa Broger[1], Christelle Fuchs[3], Ludwig A. Hothorn[4], Sylvain Loubéry[3], Michael Hothorn[1,#]

Affiliations
[1]Structural Plant Biology Laboratory, Department of Plant Sciences, University of Geneva, 1211 Geneva, Switzerland.
[2]Department of Molecular and Cellular Biology, University of Geneva, 1211, Geneva, Switzerland.
[3]Plant Imaging Unit, Department of Plant Sciences, University of Geneva, 1211 Geneva, Switzerland.
[4]retired from Institute of Biostatistics, Leibniz University, 30419, Hannover, Germany.
[$]current address: CSL Behring AG, 3014 Bern, Switzerland.
*These authors contributed equally to this work.

[#]Corresponding author Michael Hothorn Email: michael.hothorn@unige.ch

Fig. S1. Structure-based sequence alignment of BSU1-related protein phosphatases. Shown is a multiple sequence alignment (*A. thaliana* BSU1, https://uniprot.org UniProt-ID: Q9LR78; *A. thaliana* BSL1, Q8L7U5; *O. sativa* BSL1, Q60EX6; *A. thaliana* BSL2, Q9SJF0; *A. thaliana* BSL3, Q9SHS7-1; *O. sativa* BSL2, Q2QM47; *M. polymorpha* BSLM, A0A2R6VWR4; *C. reinhardtii*, phytozome.org (v.5.6) identifier Cre01.g050850; *A. thaliana* PP1/TOPP3, P48483; *H. sapiens* PP1α, P62136; *H. sapiens* PP1γ, P36873; *H. sapiens* PP1β, P62140) and including a BSU1 secondary structure assignment (α-helices in gray, β-strands in black) calculated with the program DSSP (143). Conserved residues among plant Kelch-phosphatases are highlighted in light red, residues also conserved among PP1-type Ser/Thr phosphatases are shown in dark red, residues involved in metal ion coordination are denoted by a red *, the position of the D510N and H707K mutations are highlighted in blue. Phosphorylation sites are highlighted by a colored circle in either red (CDG1-mediated phosphorylation), magenta (CDK1-mediated phosphorylation) or gray (BIN2-mediated phosphorylation).

Fig. S2. Purity of the BSU1 and BIN2 preparations used in this study. (*A*) Coomassie-stained SDS-PAGE analysis of the different wild-type and mutant fragments of BSU1 and BIN2. (*B*) Analytical size exclusion chromatography analyses of wild-type and mutant BSU1 and BIN2 fragments. 80 µL of sample were loaded to a Superdex 200 increase HR10/30 column and sample elution at 0.75 mL/min was monitored by UV absorption at 280 nm. The red and blue vertical lines indicate the characteristic elution volumes for monomeric wild-type BSU1$^{421-793}$ and full-length BIN2 proteins, respectively.

A

```
PDB      Z>30  rmsd  %id  Description

9mey-B   38.7  2.0   23   KELCH DOMAIN LEUCINE ZIPPER-LIKE TRANSCRIPTION REGULATOR 1 (Homo sapiens)

[...]

8uxs-A   36.6  1.8   21   KELCH DOMAIN-CONTAINING PROTEIN 2 (Homo sapiens)

[...]

5gqt-A   36.5  2.3   24   KELCH DOMAIN NITRILE-SPECIFIER PROTEIN 1 (Arabidopsis thaliana)

[...]

5gq0-B   36.1  2.3   22   EPITHIOSPECIFIER PROTEIN (Arabidopsis thaliana)

5a10-A   35.8  2.2   21   THIOCYANATE FORMING PROTEIN (Arabidopsis thaliana)

9d8p-A   35.7  2.2   22   KELCH DOMAIN-CONTAINING PROTEIN 10 (Homo sapiens)

9d1z-A   35.6  2.1   22   KELCH DOMAIN-CONTAINING PROTEIN 3 (Homo sapiens)

[...]

5yy8-A   33.1  2.0   20   KELCH DOMAIN INFLUENZA VIRUS NS1A-BINDING PROTEIN (Homo sapiens)

[...]

8ttq-B   30.6  2.3   20   MUSKELIN (Homo sapiens)

6hrl-A   30.1  1.8   21   KELCH-LIKE PROTEIN 17 (Homo sapiens)

[...]
```

B

C

```
PDB      Z>40  rmsd  %id  Description

1jk7-A   43.3  1.4   43   SERINE/THREONINE PROTEIN PHOSPHATASE PP1-GAMMA (Homo sapiens)

8pq7-C   43.2  1.2   44   SERINE/THREONINE-PROTEIN PHOSPHATASE PP1c (Solanum tuberosum)

3v4y-C   43.2  1.5   43   SERINE/THREONINE-PROTEIN PHOSPHATASE PP1-ALPHA (Homo sapiens)

[...]

5jpe-A   42.9  1.6   40   SERINE/THREONINE-PROTEIN PHOSPHATASE PPZ1 (Candida albicans)

[...]

7txh-F   42.7  1.5   43   SERINE/THREONINE-PROTEIN PHOSPHATASE PP1c (Homo sapiens)

[...]

8a8f-A   42.7  1.5   43   SERINE/THREONINE-PROTEIN PHOSPHATASE PP1-2 (Saccharomyces cerevisiae)

[...]

1s70-A   42.6  2.1   42   SERINE/THREONINE PROTEIN PHOSPHATASE PP1-DELTA (Homo sapiens)

[...]

4i5n-F   41.7  1.6   36   SERINE/THREONINE-PROTEIN PHOSPHATASE 2A (Homo sapiens)

[...]

1s95-B   41.6  2.0   36   SERINE/THREONINE PROTEIN PHOSPHATASE 5 (Homo sapiens)

[...]

2p6b-A   40.3  1.9   35   SERINE/THREONINE-PROTEIN PHOSPHATASE 2B CALMODULIN-DEPENDENT CALCINEURIN A (Homo sapiens)

[...]

4i11-A   40.2  2.1   34   SERINE/THREONINE-PROTEIN PHOSPHATASE 2B CALMODULIN-DEPENDENT CALCINEURIN A (Rattus norvegicus)

[...]
```

Fig. S3. BSU1 shares structural homology with Kelch domain-containing proteins and with PP1-like Ser/Thr phosphatases. (*A*) Selected top hits from the DALI webserver (http://ekhidna2.biocenter.helsinki.fi/dali/) (64) using the isolated BSU1[11-351] Kelch domain as search model. (*B*) Structural superposition of the BSU1 Kelch domain (cyan C_α trace) with the Kelch domain of human Leucine Zipper-like Transcription Regulator 1 (in orange, PDB-ID: pdb_00009mey, r.m.s.d. ~1 Å comparing 184 corresponding C_α atoms) (60). (*C*) Selected DALI hits for the BSU1[450-793] phosphatase domain.

Fig. S4. Design and molecular docking of BSU1 synthetic substrate peptides. (*A*) Amino acid sequence of the synthetic peptide used in BSU1 phosphatase substrate specificity assays. Basic, acidic, and hydrophobic residues are highlighted in blue, red, and yellow, respectively. The central phosphorylated residue (pThr, pSer, or pTyr) is denoted as pX. Residue positions relative to pX are indicated below, with negative and positive numbers representing upstream and downstream residues, respectively. Based on the analysis by Hein et al. (68), PP1 phosphatases exhibit a preference for basic residues at positions −6, −5, −3, −2, −1, +2, and +3, while there is no amino acid preference at position +1, except for the exclusion of proline when pSer is present. Positions +4, +5, and +6 show no preference; thus, a Gly-Ser-Tyr linker was incorporated at the C-terminus to accurately estimate the molar concentration of the peptide in solution, using the molar extinction coefficients at $A_{280\text{ nm}}$. (*B-D*) AlphaFold3-predicted complex structures of BSU1$^{421-793}$ bound to Mn^{2+} ions and three distinct phosphopeptides: the pThr-containing peptide in *B*, the pSer-peptide in *C*, and the pTyr-peptide in *D*, as implemented in https://alphafoldserver.com/ (69). The upper panels provide a close-up view of BSU1 active site in ribbon representation. The experimental crystal structure of BSU1$^{421-793}$ is shown in white, the AlphaFold3 docking models are in blue. Phosphopeptides are colored in yellow (pThr), in orange (pSer) and in dark gray (pTyr). Predicted salt bridges between the phosphate group and BSU1 residues are depicted as dotted lines (in green). The lower panels present an overview of the AlphaFold3 docking models as C$_\alpha$ chain traces, colored by predicted local distance difference test (pLDDT) score (very high, dark blue, pLDDT > 90; high, cyan, 90 > pLDDT > 70; low, yellow, 70 > pLDDT > 50; very low, red, pLDDT < 50) and visualized using Py2DMol (https://py2dmol.solab.org). AlphaFold3 pTM and ipTM values are reported alongside (69). (*E*) Molecular surface representation of BSU1$^{421-793}$ (in blue) bound to the different phosphopeptides, colored as in panels *B-D*. Note that the pTyr-containing peptide can only be accommodated in the active site adopting a highly unusual twisted conformation.

Fig. S5. The isolated BSU1 Kelch domain further enhances the growth phenotype of the *bri1-5* mutant. Shown are shoot growth phenotypes of 45 d-old *bri1-5* plants expressing BSU1$^{1-357}$ (Kelch domain), BSU1$^{421-793}$ (phosphatase domain) or BSU1$^{421-793\ D510N\ H707K}$ (phosphatase-dead) variants from the 35S promoter. Shown are T1 generation plants, the Ws-2 wild type, *bri1-5* (142) and the *bsu1-1D* (36) allele. BSU1 protein expression was assayed by western blot using an α-Flag antibody, the Ponceau-stained membranes are shown on the left as loading controls.

Fig. S6. Arabidopsis *bsu1* and *bsl1-3* mutant CRISPR/Cas9 gene editing events. (*A*) Genomic representations of BSU1 (TAIR-ID: At1g03445), BSL1 (At4g03080), BSL2 (At1g08420) and BSL3 (At2g27210) genes with exons in yellow and introns shown as black lines. sgRNA targeting regions are represented with red arrowheads. (*B*) CRISPR events are shown below the corresponding genotype, as confirmed by Sanger sequencing. The altered DNA sequence and the stop codon are shown in red, the corresponding amino acid sequence is shown in black (wild type) and blue (altered sequence by the frame shift editing event).

Fig. S7. Deletion of multiple Kelch phosphatase triggers stomata over-proliferation. (*A*) Representative confocal images of 7 d-old abaxial cotyledon regions for Col-0, *bsk1-2 bsk2-2* (126), *scrm-D* (125), *bsu1* and different *bsu1 bsl* mutant combinations, stained with PI (in magenta). (scale bar = 200 μm). (*B*) Stomatal index quantification represented in box plots (n = 6; bold black line, median; box, inter-quartile range (IQR); whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile, raw data shown as individual points) and including Dunnett-type (124) two-sided multiplicity-adjusted *p*-values for the comparison against the Col-0 control (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). The stomatal index of the strong *scrm-D* allele could not be determined reliably.

A

B

Genotype	Seeds		
	Total	Aborted	Rate aborted
Col-0	245	40	16.32 %
bsu1 bsl1 bsl3	203	57	28.07 %
bsu1 bsl1 bsl2	172	127	73.83 %
bsu1 bsl1 bsl3 bsl2$^{+/-}$	215	67	31.16 %
bsu1 bsl1 bsl2 bsl3$^{+/-}$	174	140	80.45 %

C

Col-0   *bsu1 bsl1 bsl3*   *bsu1 bsl1 bsl2*   *bsu1 bsl1 bsl3 bsl2$^{+/-}$*   *bsu1 bsl1 bsl2 bsl3$^{+/-}$*

Fig. S8. Higher order *bsu1 bsl* mutants have fertility defects. (*A*) Quantification of silique length in different *bsu1 bsl* mutant combinations with representative siliques shown on top (scale bar = 0.5 cm) and quantifications below (n = 20). Box plot bold black line, median; box, inter-quartile range (IQR); whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile, raw data shown as individual points. Dunnet-type (124), two-sided multiplicity-adjusted *p*-values for the comparison against the Col-0 control are shown alongside (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). (*B*) Quantification of seed abortion in Col-0 and different *bsu1 bsl* mutant combinations. The table summarizes the total number of seed and aborted seeds. The abortion rate is expressed as percentage of wild type seeds. Shown below in (*C*) are representative shoot branches of the different genotypes on the top (scale bar = 1 cm) and individual opened siliques below (scale bar = 0.2 cm). Aborted seeds are denoted by white asterisks.

Fig. S9. The expression patterns of BSU1 and BRI1 are not overlapping. Shown are spatiotemporal expression patterns of BSU1 and BRI1 reporter lines determined by β-glucuronidase activity (GUS) assays. A sequence ~1kb upstream of the BSU1 ORF was used to drive *proBSU1::BSU1-GUS* and *proBSU1::GUS*, respectively. A ~2kb upstream of BRI1 ORF was used to drive *proBRI1::GUS*. 14-17 independent T2 lines for each construct were stained for GUS activity, either using 5 d-old cotyledons (*A*), or the 4th leaf of 15 d-old soil-grown plants (*B*). Examples for the different categories summarized in the star plots in Fig. 2*F* are represented.

Fig. S10. BSU1 expression pattern during callus induction. Promoter β-glucuronidase (GUS) reporter assay for 3-week-old callus induced from Col-0, *proBSU1::GUS* and *proBSU1::BSU1-GUS* plants. Representative pictures of the induced calli (top panel) and stained calli (bottom panel) are shown for all genotypes (scale bar = 0.5 cm).

Fig. S11. Marchantia Mp*bslm*[ge] and Mp*gsk*[ge] mutant CRISPR/Cas9 gene editing events. (*A*) Genomic representations of MpBSLM (MpVg00440, https://marchantia.info) and MpGSK (Mp7g04170) genes with exons in yellow and introns shown as black lines. SgRNA targeting regions are represented with red arrowheads. (*B*) CRISPR events are shown below the corresponding genotype, as confirmed by Sanger sequencing. The altered DNA sequence and the stop codon are shown in red, the corresponding amino acid sequence is shown in black (wild type) and blue (altered sequence by the frame-shift editing event).

Fig. S12. Summary of a RNA-seq experiment comparing 14 d-old Mp*bslm*^{ge} mutant plants with the Tak-1 wild type. (*A*) Volcano plots showing differentially expressed genes (DEGs) (Log$_2$FC > 1, in red and Log$_2$FC < -1, in blue) of Mp*bslm*^{ge} vs. Tak-1. Putative cell cycle-regulated genes discussed in the text are indicated in black (*B*) Manually curated gene-ontology classification of DEGs. Up-regulated genes with |log$_2$(FC)|>1 and $p < 0.05$ were considered differentially expressed.

Fig. S13. BIN2 phosphorylates BSU1 in vitro. (*A*) Autoradiograph of an in vitro kinase assay of BIN2 vs. the BSU1 Kelch domain (BSU1[1-357]), BSU1 phosphatase domain (BSU1[421-793]), or full-length BSU1 (BSU1[1-793]). BIN2[K69R] is a kinase-dead mutant (41). (*B*) Schematic representation of BSU1, depicting the Kelch domain (residues 11–351), phosphatase domain (residues 450–767), and C-terminal tail (residues 768–793). Structured regions are shown as blocks, and putative linker regions (residues 1–10 and 352–449) as solid lines. Phosphorylated residues targeted by BIN2 are shown in gray. Below, the MS1 spectra of the phosphorylated BSU1[243-260], BSU1[421-441], and BSU1[141-162] peptides are shown from top to bottom.

Table S1. Crystallographic data collection and refinement statistics.

	BSU1[1-357] Kelch	BSU1[421-793] SAD	BSU1[421-793] native	BSU1[421-793 D510N H707K] phosphatase-dead
PDB-ID	pdb_00009TP5		pdb_00009TP7	pdb_00009TP8
Data collection				
Wavelength (Å)	0.999874	1.282151	1.000029	1.000031
Space group	$P\,4_3\,2_1\,2$	$C\,2$	$C\,2$	$C\,2$
Cell dimensions				
a, b, c (Å)	77.31, 77.31, 134.32	226.07, 97.64, 157.79	226.41, 97.79, 157.80	139.76, 44.05, 79.24
α, β, γ (°)	90, 90, 90	90, 129.23, 90	90, 129.41, 90	90, 121.07, 90
Resolution (Å)	42.40 – 2.25 (2.39– 2.25)	45.69 – 2.70 (2.77 – 2.70)	45.38 – 2.10 (2.22 – 2.10)	41.34 – 1.60 (1.69 – 1.60)
R_{meas}[#]	0.225 (3.0)	0.196 (2.93)	0.147 (1.87)	0.079 (1.37)
CC(1/2)[#]	1.0 (0.30)	1.0 (0.59)	1.0 (0.56)	1.0 (0.50)
$I/\sigma\,I$[#]	12.72 (1.0)	9.4 (1.0)	11.44 (1.20)	11.7 (1.1)
Completeness (%)[#]	100 (100)	99.6 (99.6)	99.5 (99.2)	99.0 (98.6)
Redundancy[#]	13.9 (13.5)	10.4 (10.6)	6.8 (6.8)	3.4 (3.3)
Wilson B-factor[#]	51.3	65.1	43.1	29.5
Phasing				
No. of Zn^{2+} sites		8		
Figure of merit[^]		0.39 / 0.82		
Refinement				
Resolution (Å)	42.40 – 2.25		45.38 – 2.10	41.34 – 1.60
No. reflections	36,873		155,227	106,610
R_{work} / R_{free}[*]	0.22 / 0.25		0.20 / 0.23	0.16 / 0.18
No. atoms				
protein	4,894		32,221	5,029
solvent	43		870	139
ligand	44		68	77
Res. B-factors[*]				
protein	70.1		58.7	33.43
solvent	49.2		49.0	38.3
ligand	76.9		76.5	54.6
R.m.s deviations[*]				
bond lengths (Å)	0.002		0.002	0.015
bond angles (°)	0.49		0.53	1.44
Ramachandran plot[$]:				
most favored regions (%)	96.62		95.58	96.09
outliers (%)	0		0	0
MolProbity score[$]	1.13		1.51	1.64

[#]as defined in XDS (144)
[^]FOM before and after NCS averaging and density modification with phenix.resolve (145, 111)
[*]as defined in phenix.refine (107)
[$]as defined in Molprobity (114)