A Novel *miR-4745*-KLC2 Axis Regulates Cancer Stem Cell Traits in Colorectal Cancer

Kuldeep Lahry[1,2#], Pauline Barraja[2#], Julie Ripoll[3], Océane Bouvet[2], Guillaume Pinna[4], Amandine Bastide[1,2], Stéphanie Mateus[1], Lisa Tellier[1], Kevin Bigot[1], Christy Matta[1], Elodie Pothin[2], Lucile Bansard[2], Cécile Dejou[1], Anne-Alicia Gonzalez[5], Eric Rivals[3], Julie Pannequin[2], Jean-Marc Pascussi[2*], Alexandre David[1,2*] and Chris Planque[1,2*]

[1] IRCM, Université de Montpellier, INSERM, ICM, Montpellier, France.

[2] IGF, Université de Montpellier, CNRS, INSERM, Montpellier, France.

[3] LIRMM, Université de Montpellier, CNRS, Montpellier, France.

[4] Plateforme ARN interférence (PARi), Université Paris Cité & Université Paris-Saclay, INSERM, iRCM/IBFJ CEA, Stabilité Génétique Cellules Souches et Radiations, Fontenay-aux-Roses, France.

[5] MGX-Montpellier GenomiX, Université de Montpellier, CNRS, INSERM, Montpellier, France.

* Corresponding authors:

chris.planque@inserm.fr;

alexandre.david@inserm.fr;

jean-marc.pascussi@inserm.fr;

[#], These authors contributed equally to this work.

[*], These authors jointly supervised this work.

***Running title**: miR-4745 Regulates Colorectal Cancer Stem Cell Phenotype*

ABSTRACT

Advanced colorectal cancer (CRC) treatments often fail due to chemoresistance and tumor recurrence, primarily driven by cancer stem cells (CSCs), which play a significant role in therapy ineffectiveness. MicroRNAs (miRNAs) play a pivotal role in regulating numerous biological functions within CSCs. Through a gain-of-function miRNA screening, we identified miRNAs that regulate the ALDEFLUOR-positive CSC population in colon cancer, with *miR-4745* emerging as a promising inhibitor. Overexpression of *miR-4745* reduced CSC self-renewal, diminished chemoresistance, and prevented CSC enrichment after chemotherapy in patient-derived CRC models. Mechanistically, *miR-4745* directly targets the 3'UTR of kinesin-light chain 2 (*KLC2*) mRNA, a member of the kinesin superfamily associated with poor clinical outcomes across various cancers. siRNA-mediated *KLC2* downregulation reduced ALDH-positive CSCs and suppressed sphere formation, while *KLC2* overexpression promoted CSC traits. Additionally, we observed an inverse correlation between *KLC2* and *miR-4745* mRNA levels in different colon cancer cell culture settings. Notably, *KLC2* mRNA levels increased with local CRC progression and were correlated with poor overall survival in a cohort of over 1 000 stage I to IV colorectal cancer patients. Our study uncovers a novel *miR-4745*-KLC2 axis that regulates CSC properties and chemoresistance in CRC, offering a promising strategy to prevent relapse and improve clinical outcomes for patients.

Keywords: *miR-4745*, *KLC2*, colorectal cancer, cancer stem cell, chemoresistance

BACKGROUND

Colorectal cancer (CRC) is the most common cancer in men and the second most common in women worldwide (WHO). In the United States, colorectal cancer ranks as the third most common cancer diagnosis and the third leading cause of cancer-related mortality among both men and women (1). According to projections by the World Health Organization, the global impact of colorectal cancer is anticipated to grow substantially by 2040, with an estimated 63% rise in new cases and a 73% increase in yearly deaths. For localized CRC, surgery remains the standard of care, while advanced CRC is managed through a multimodal approach combining chemotherapy, targeted therapies, and immunotherapy. The primary chemotherapeutic agents include 5-fluorouracil (5-FU), irinotecan, and oxaliplatin, often administered in various combinations. However, these treatments fail in 90% of metastatic cases, primarily due to drug resistance and tumor recurrence driven by cancer stem cells (CSCs), a highly resilient subset of tumor cells (2). CSCs are defined by their expression of stem cell factors that confer stemness properties, including the ability to self-renew. Moreover, CSCs can initiate and sustain tumor growth in serial transplantation assays (3) . These cells exhibit increased resistance to treatments compared to other tumor cells (4,5) and often become more prevalent in CRC tumors following chemotherapy (6,7).

Considering the significant impact of CSCs on therapeutic failure in CRC patients, it is essential to develop and implement treatment approaches that increase the responsiveness of these cells to current therapies, thereby improving patient outcomes. However, the precise mechanisms driving the heightened therapeutic resistance of CSCs remain poorly understood. This resistance is multifactorial, arising from factors such as increased resistance to DNA damage, reduced apoptosis, enhanced autophagy, slow proliferation, and the activation of the WNT/β-catenin, Hedgehog, and Notch signaling pathways. Additionally, CSCs often express multidrug transporters such as ATP-binding cassette G2 (ABCG2) (8) and drug-metabolizing

enzymes like aldehyde dehydrogenase 1A1 (ALDH1A1) (9). To identify CSCs in solid

tumors, including CRC, flow cytometry techniques have been employed, targeting ABCG2

expression (Side Population, SP) and ALDH activity (ALDEFLUOR-positive cells) (10).

Both SP cells and ALDH-positive cells exhibit increased resistance to cytotoxic drugs *in vitro*

and *in vivo,* and their populations are enriched following chemotherapy (6).

Micro-RNAs (miRNAs) are small non-coding RNA molecules, typically 20-25 nucleotides in

length, that regulate gene expression by binding to the 3' untranslated region (3'UTR) of

messenger RNA (mRNA), resulting in either translation inhibition or mRNA degradation. In

cancer, miRNA expression is often dysregulated, and increasing evidence indicates that

miRNAs can function as oncogenes or tumor suppressors. They play a crucial role in

regulating gene expression related to various cancer-related processes, including tumor

initiation, progression (11,12), and drug resistance (13).

Interestingly, miRNAs are increasingly recognized as pivotal regulators in the complex

control of CSC fate. Multiple studies have highlighted their role in regulating CSC self-

renewal across various cancers, including prostate cancer (14), breast cancer (15), and colon

cancer (16). In colorectal CSCs specifically, overexpression of *miR-451* has been shown to

reduce tumorigenicity, self-renewal, and resistance to irinotecan by downregulating the

ABCB1 drug transporter (17). Moreover, *miR-137* suppresses the tumorigenic potential of

colorectal cancer stem cells by directly targeting Musashi-1 (18), whereas *miR-140*

contributes to resistance against methotrexate and 5-fluorouracil through its regulation of

histone deacetylase 4 (*HDAC4*) (19).

We recently reported that *miR-148a* overexpression suppresses Pregnane X Receptor (PXR)

signaling, which is crucial for the expression of chemoresistance genes such as *ALDH1A1* and

ABCG2 in CSCs. Our study further demonstrated that niclosamide induces *miR-148a*, thereby

inhibiting PXR and preventing CSC chemoresistance and tumor recurrence, both *in vitro* and

in vivo (16).

Given the critical role of CSCs in CRC chemoresistance and the impact of miRNAs on the

CSC phenotype, we conducted a gain-of-function screen to identify miRNAs that regulate the

ALDH-positive CSC population in CRC cells. This screen identified *miR-4745* as a novel and

potent inhibitor of ALDH activity. Ectopic overexpression of *miR-4745* diminished the self-

renewal and chemoresistance properties of colon CSCs and prevented their enrichment

following chemotherapy. Subsequent analysis revealed that kinesin light chain 2 (*KLC2*)

mRNA is a direct target of *miR-4745*. This microRNA binds to the 3′ untranslated region

(3′UTR) of *KLC2* mRNA, resulting in decreased expression of the target transcript.

Additionally, we observed an increase in *KLC2* expression with advancing cancer stages in

colon cancer tissue samples.

These findings position *miR-4745* as a promising therapeutic candidate for reducing

recurrence and drug resistance in CRC, while also uncovering a novel role for KLC2 in

regulating colon CSCs.

METHODS

Cell lines and patient-derived tumor cell culture

CRC cell lines ((SW480 (RRID:CVCL_0546), SW620 (RRID:CVCL_0547), HT29

(RRID:CVCL_A8EZ), LS174T (RRID:CVCL_1384), HCT116 (RRID:CVCL_0291), LoVo

(RRID:CVCL_0399), and Colo205 (RRID:CVCL_0218)) were obtained from ATCC and

cultured either in DMEM or RPMI medium (Gibco) supplemented with 10% FBS, or were

cultured as spheroids in M11-defined medium using either ultra-low attachment flasks from

Corning or flasks coated with Poly 2-hydroxyethyl methacrylate. M11 medium consisted of

DMEM/F12 (1:1) (Gibco), supplemented with N2, 3 mM Glutamine, 0.6% Glucose, 4µg/ml

insulin (Sigma-Aldrich), 100U/ml Penicillin G, 100ug/ml Streptomycin, 10ng/ml hBasic-FGF

(R&D Systems), and 20ng/ml hEGF (R&D Systems). Patient-derived colon cancer cells

(CPP1, CPP6, CPP19, CPP25, CPP36, and CPP44) were established from CRC biopsies,

while two circulating tumor cell lines (CTC44 and CTC45) were isolated from the blood of a

patient with advanced metastatic colorectal cancer. These biological samples were provided

by CHU-Carémeau (Nîmes, France) as part of a previously reported study (20,21), registered

under ClinicalTrials.gov Identifier#NCT01577511 (First submitted: 2012-04-12; Last update

posted: 2021-06-21). The study protocol was approved by the French Ethics Committee

(Comité de Protection des Personnes, Sud Méditerrannée III). All relevant ethical regulations

for research involving human participants were followed with written informed consent

obtained from all patients. Cell cultures were maintained at 37°C in a humidified atmosphere

with 5% CO_2.

Colon tumor tissue samples

RNA extracted from colon tumor tissue samples was obtained as part of the Clinical and

Biological Database BCBCOLON (Institut du Cancer de Montpellier—Val d'Aurelle, France)

registered under ClinicalTrials.gov Identifier #NCT03976960 (First submitted: 2019-06-05;

Last update posted: 2025-02-12). Tissue samples were obtained from 41 patients at various

stages of colon cancer: adenoma (n = 3), primary adenocarcinoma at stages I (n = 8), II (n =

11), III (n = 5), IV (n = 8) and metastases (n = 6). All patients were chemonaive.

The tumor samples were collected in accordance with French laws under the supervision of a

designated investigator and registered with the French Ministry of Higher Education and

Research (declaration number DC-2008–695). The research protocol received approval from

the French Ethics Committee (CPP Sud Méditerranée III, reference number 2014.02.04) and

was also authorized by the institutional translational research board (ICM-CORT-2018-28).

All relevant ethical regulations for research involving human participants were followed, and

written informed consent was obtained from all patients.

Plasmid constructions

The dual sensor plasmid pmiR-sensor contains a bidirectional promoter that drives the

expression of both AcGFP (a green fluorescent protein from *Aequorea coerulescens*) and the

red fluorescent protein DsRed1 (22). In the 3'UTR of the GFP transcript, we inserted either a

full-length 3'UTR of *KLC2* (at the *Xba*I restriction site) or two tandem copies of predicted

miR-4745 binding sites (between the *Apa*I and *Xba*I restriction sites). The sequences of the

cloned regions are provided in **Supplementary Table S1**. GFP expression levels serve as a

direct indicator of *miR-4745* activity, while DsRed1 is used for normalization. This dual

expression system minimizes the effects of variable expression levels and nonspecific

transcriptional regulation, allowing for ratiometric measurements (AcGFP/DsRed1). To create

the pmiR-KLC2-3'UTR mutant #2, site directed mutagenesis was performed using the

primers KLC2-3'UTR Mut2 Fp (CGGTACTACCCAAACCTCCCCTCGTCCCTCTTC) and

KLC2-3'UTR Mut2 Rp (GACGAGGGGAGGTTTGGGTAGTACCGTGGACTG). The

clones were confirmed via Sanger sequencing at Eurofins using the AcGFP Fp primer

(CAACATCGAGGATGGCAG).

To create the KLC2-overexpressing plasmid (oe-KLC2), the *KLC2* coding sequence, lacking

the 3'UTR *miR-4745* targeting sequence, was amplified from the pDONOR-KLC2 plasmid

(Montpellier Genomic Collection MGC Facility) and inserted into the pCDNA3-3xFlag Cter

plasmid at the *Hind*III and *Xba*I restriction sites.

Transfection assay and oligonucleotides

The mirVana™ miRNA mimics, including *miR-4745-3p* (ref: 4464066, ID: MC21191), *miR-*

497-3p (ref: 4464066, ID: MC12884), *miR-4723-5p* (ref: 4464066, ID: MC21507), *miR-4784*

(ref: 4464066, ID: MC22683), and the mirVana microRNA Mimic Negative Control #1 (ref:

4464058), as well as the anti-miR™ miRNA inhibitors, including anti-*miR4745-3p* (ref:

AM17000, ID: AM21191) and the anti-miR™ miRNA Inhibitor Negative Control #1 (ref:

AM17010) were purchased from Life Technologies. siRNA duplexes specific to *KLC2*,

HEY2, *CCND3*, and *miR31HG* were purchased from Genecust, and their corresponding

sequences are listed in **Supplementary Table S2**. Transfection of 50nM miRNA or siRNA

duplexes was performed using Lipofectamine RNAiMax (Invitrogen) according to the

manufacturer's instructions. Additionally, transfection of 200 000 cells in a 6-well plate with

2µg of plasmid DNA was carried out using Lipofectamine 2000 (Invitrogen) for 72 hours,

following the manufacturer's protocol. For co-transfection experiments involving both a

miRNA mimic and plasmid DNA, Lipofectamine 2000 was also used.

miRNA extraction and quantification by real-time RT-PCR

miRNA was purified using the miRCURY™ RNA isolation kit. First-strand cDNA synthesis

was performed with the miRCURY LNA™ RT Kit (Qiagen), followed by real-time PCR

amplification using the miRCURY LNA™ SYBR Green PCR Kit (Qiagen). Each sample was

amplified in triplicate on an LC480 Real-Time PCR System (Roche). The expression levels of

miR-4745-3p were calculated using the $2^{-\Delta\Delta Ct}$ method, with *miR-103a* used as the

normalization control.

RNA extraction and quantification by real-time PCR

198   Total RNA was extracted using the RNeasy Mini Kit (Qiagen) and treated with DNAse I to

199   remove genomic DNA contamination. First-strand cDNA was synthesized using Superscript

II (Invitrogen) and random hexamers, followed by real-time PCR for gene expression

analysis. Each cDNA sample was analyzed in triplicate using SYBR Green chemistry

(Roche) on a LightCycler 480 Real-Time PCR System (Roche). *GAPDH*, *β-actin*, *RPLO*, *18S*

ribosomal RNA, and *RPL13* were used as endogenous controls. Gene expression fold changes

were determined using the comparative Ct approach, specifically the $2^{-\Delta\Delta Ct}$ method. Primer

sequences are provided in **Supplementary Table S3**.

mRNA purification

mRNA was isolated from total RNA by performing two successive purifications with the

GeneElute™ mRNA Purification Kit (Sigma), following the protocol provided by the

manufacturer.

Western blot

Cells were lysed using RIPA buffer supplemented with protease inhibitors (Roche) for total

protein extraction. For western blotting, samples were separated on a 10% SDS-PAGE gel

and transferred to a nitrocellulose membrane (Amersham). The antibodies utilized in this

study included GAPDH (sc-32233, dilution 1:5,000; Santa Cruz), β-ACTIN (A-5441, dilution

1:10,000; Sigma-Aldrich), and KLC2 (17668-1-AP, dilution 1:1,000; Proteintech). Band

intensities were quantified using Image Lab software (Bio-Rad).

Proliferation assay

One thousand cells were seeded in triplicate in a 96-well plate and incubated for 24, 48, 72,

and 96 hours. Cells were subsequently fixed in 10% trichloroacetic acid (TCA) at 4 °C for

a minimum duration of 2 hours. After three washes with Milli-Q water, cells were stained

with 0.4% Sulforhodamine B solution for 30 minutes at room temperature, followed by three

225 additional washes with MilliQ water. Following resuspension in 10 mM Tris-Base,

226 absorbance was recorded at a wavelength of 562 nm.

Staining for flow cytometry and fluorescence-activated cell sorting (FACS)

CD44v6-APC (clone REA706, 130-111-238; Macs Miltenyi) antibody was incubated with

100 000 cells in PBS containing 5% FBS for 20 min at 4°C. The Aldefluor assay (Stem Cell

Technologies) was performed according to the manufacturer's instructions. ALDH-positive

and ALDH-negative cells were identified by comparing the same sample treated with and

without the ALDH inhibitor diethylaminobenzaldehyde (DEAB). The flow cytometry gating

strategy for Aldefluor-stained samples involved initial staining using the ALEFLUOR assay,

followed by SYTOX Blue Dead Cell Stain (Invitrogen). A stepwise gating strategy was

employed to identify the primary cell population (SSC vs. FSC), isolate single cells (SSC-A

vs. SSC-W), and select viable cells based on SYTOX Blue exclusion (SSC-A vs. SYTOX

Blue fluorescence). Dead cells were excluded based on light scatter and SYTOX Blue

Staining. DEAB-treated cells served as a negative control to set the gating threshold for

identifying ALDH-positive subpopulation in test samples without DEAB. Cells were sorted

using a FACSAria II instrument (BD), and the resulting data were processed and analyzed

with Flowing Software version 2.5.1 (http://flowingsoftware.btk.fi/). ALDH activity was also

assessed using the MACSQUANT analyzer (Miltenyi) with the same protocol.

For sphere formation assay experiments, live cells were sorted using FACSAria II (BD) based

on negative labeling with SYTOX Blue or 7-AAD.

To experimentally validate the binding of *miR-4745* to the *KLC2* mRNA transcript, we

employed a fluorescence-based dual reporter assay system using the pmiR-sensor plasmid.

CPP1 cells were co-transfected with the pmiR-sensor plasmid containing the *KLC2* 3'UTR,

and the *miR-4745-3p* mimic. GFP and DsRed1 expression levels were subsequently measured

using a CytoFLEX LX flow cytometer (Beckman Coulter) at the MRI facility in Montpellier.

In vitro cytotoxic treatment

253 Cells were plated at 200 000 cells per well in DMEM with 10% FBS in 6-well plates. After 24

254 hours, cells were exposed to FIRI (1X = 50μM 5-FU + 500nM SN38) or vehicle (0.1%

255 DMSO), with 3 wells per condition. Following 72 hours of treatment, cells were washed twice

256 with ice-cold PBS before RNA extraction.

In vitro chemosensitivity assays

259 Cells were plated at 2 000 cells per well in DMEM with 10% FBS in 96-well plates. After 24

260 hours, they were treated with chemotherapeutic agents for 48 to 72 hours. Cell viability was

261 evaluated using the CellTiter-Glo assay (Promega), and half-maximal effective concentration

262 (EC_{50}) values were determined using Prism9 software.

Sphere formation assays

265 The percentage of cells forming spheres was determined by plating 100 or 200 live cells per

266 well in M11 medium in ultra-low attachment 96-well plates. Live cells were sorted based on

267 negative labeling with SYTOX Blue or 7-AAD. Spheres with a diameter exceeding 50 μm

268 were counted.

Polysome fractionation

This procedure was carried out as previously described (23). Briefly, 4 x 10^6 CPP1 cells were

seeded in Petri dishes (150 mm) and transfected 24 hours later with either the *miR-4745-3p*

mimic or the mirVana microRNA Mimic Negative Control #1 (miCTRL). Following 48 hours

of transfection, cells were exposed to 20 µg/ml emetine for 5 minutes at 37°C. Subsequently,

they were rinsed twice with ice-cold PBS and harvested by scraping into ice-cold PBS. The

cell suspension was centrifuged, resuspended in 1 mL of polysome lysis buffer, and

homogenized by vigorous shaking with 1.4 mm ceramic spheres (Lysing matrix D MPBio)

using a FastPrep machine (MPBio). The lysates were subsequently centrifuged at maximum

speed for 10 minutes at 4°C.

The resulting supernatant (cytosolic fraction) was loaded onto a 15-50% sucrose gradient and

centrifuged at 35 000 rpm for 2.5 hours at 4°C using a SW41 rotor (Beckman Coulter).

Polysomes were separated and monitored in real-time using an optical density (OD) 254 nm

UV spectrometer. Gradient fractionation was carried out using an ISCO density gradient

fractionation system (Lincoln, NE), while continuously monitoring absorbance at 254 nm

across the gradient depth.

Transcriptome and Translatome analyses

Total RNA was isolated using TRIzol reagent (Invitrogen) following the manufacturer's

instructions. mRNA associated with 1 to 3 ribosomes ("Light polysomes") and mRNA

associated with more than 3 ribosomes ("Heavy polysomes") were extracted using TRIzol LS

(Invitrogen) according to the manufacturer's instructions.

We prepared RNA-seq libraries using the NuGEN Universal Plus mRNA-seq kit (Tecan,

Redwood City, CA, USA) according to the manufacturer's instructions. Briefly,

polyadenylated RNAs were selected using oligo-dT magnetic beads and chemically

fragmented. First strand cDNAs were synthesized using random primers in the presence of

actinomycin D to prevent spurious DNA-dependent synthesis. After second strand synthesis

and end-repair, adapters including sequences used later in PCR amplification were ligated.

The strand to be sequenced was purified and PCR amplified for 15 cycles. PCR products were

purified using AMPure XP Beads (Beckman Coulter Genomics, Brea, CA, USA). The size

distribution of the resulting libraries was monitored using a Fragment Analyzer (Agilent

Technologies, Santa Clara, CA, USA) and the libraries were quantified using the KAPA

Library quantification kit (Roche, Basel, Switzerland).

The libraries were denatured with NaOH, neutralized with Tris-HCl, and diluted to 300 pM.

Clustering and sequencing were performed on a NovaSeq 6000 (Illumina, San Diego, CA,

USA) using the single read 100nt protocol on 2 lanes of a flow cell S2.

Image analyses and base calling were performed in real time by the Real-Time Analysis

software (Illumina). Demultiplexing was performed using Illumina's conversion software

(bcl2fastq 2.20).

Bioinformatic analysis of transcriptome and translatome sequencing data

We describe the pipeline used to analyze RNA-seq data from the transcriptome (cytosolic

fraction) and translatome (light and heavy polysomal fractions) under two conditions: control

condition (CTL) and *miR-4745* overexpression (*miR-4745*). We refer to these three libraries as

Cyto, Heavy and Light, and to the conditions as CTL and *miR-4745*. For each library, read

quality was assessed using FastQC (v0.11.9, Babraham Institute, Cambridge, UK) and

Illumina's Sequencing Analysis Viewer (software SAV). FastqScreen (v0.15.1) was used to

identify potential contamination. Reads were filtered using a quality threshold of Q35, and 12

bases at the 5'end were trimmed with Cutadapt v3.2. High-quality reads were then aligned to

the Homo sapiens reference genome GRCh38 (primary assembly release 104) using STAR

v2.7.9a (24), and annotated with the corresponding Ensembl annotation. Quantification of

mapped reads was performed using featureCounts from subread v2.0.1 (25), and features with

zero counts across all fractions were removed.

Statistical analyses were conducted using DESeq2 v1.30.1 (26) to identify differentially

expressed genes (DEGs) between the two conditions for each fraction, with statistics

computed using the Wald test. P-value adjustment for multiple testing was performed using

the Benjamini-Hochberg method (27) to control the false discovery rate. Only genes with an

adjusted p-value below 0.05 and a log fold change greater than one were selected for further

analyses. Results from the cytosolic fraction were combined with those from the polysomal

fraction to determine whether genes were regulated at the transcriptional level, translational

level, or both. DEGs from pairwise comparisons (e.g., CTL-Heavy vs. *miR4745*-Heavy) were

categorized using Pandas v1.1.5 into seven regulatory categories: no significant variation

(background), regulation at the transcript level only (Transcription UP or Transcription

DOWN), regulation at the mRNA translation level only (Translation UP or Translation

DOWN), or regulation at both transcript and translation levels (total mRNA abundance UP or

total mRNA abundance DOWN). Scatter plots illustrating gene variation and regulatory

categories were generated using matplotlib v3.3.0 (28).

miRNome-wide screening coupled with a miniaturized ALDEFLUOR assay

We conducted a miRNome-wide screening in the SW620 cell line using a miRNA

overexpression (OE) library. The OE library contained miRNA mimics of endogenous

miRNA duplexes (miRIDIAN® microRNA Mimics Library, Dharmacon/Horizon Discovery),

designed to mimic 2 048 human miRNAs listed in miRBase v.19.0, as detailed previously

(15). For screening, an automated reverse transfection method targeting pre-miRs was

established using a robotic workstation fitted with a 96-well head probe (Nimbus, Hamilton).

In short, pre-miRs were combined with Lipofectamine RNAiMAX (Life Technologies) and

applied to collagen-coated, clear-bottom, black-walled 96-well culture plates (Costar

Cat#3904). After 15 minutes of complexation, SW620 cells were seeded onto the lipoplexes

(12 500 cells per well; final [pre-miR] = 25nM).

Three different controls were used in the OE screen: a commercial mixture of cytotoxic

siRNAs as a positive transfection control (AllStars Maximal Death Control, Qiagen), a

miRNA mimic against *miR-148a-3p* (phenotypic positive control, previously shown to

downregulate the relative CSC proportion, Dharmacon/Horizon Discovery) and a mimic

negative control (miRIDIAN® microRNA Mimic Negative Control, Dharmacon/Horizon

Discovery). Three days post-transfection, the relative CSC proportion in SW620 cells treated

with miRNA mimics was assessed using a previously described adaptation of the

ALDEFLUOR assay (Stem Cell Technologies) for image acquisition and analysis in a 96-

well format (29,30).   Briefly, after removing the culture supernatants, 40 µL of an

ALDEFLUOR/Hoechst mixture (5 µL of BAAA substrate per mL of ALDEFLUOR buffer +

Hoechst 33342 [Sigma] at 50 µg/mL) were added to the culture wells. All plates included one

control well that received N, N-diethylaminobenzaldehyde (DEAB), an inhibitor of ALDH.

After 45 minutes of incubation at 37°C, the mixture was replaced with 50µL of ice-cold

ALDEFLUOR buffer, and the plates were immediately imaged using a High Content Imaging

device (Operetta HCS epifluorescence microscope, Perkin Elmer). During image acquisition,

the plates were kept at 4°C. For each well, nine fields were captured at 10X magnification

using two fluorescence channels: green for ALDEFLUOR (excitation: 470±10 nm; emission:

525±25 nm) and blue for Hoechst 33342 (ex: 380±20 nm; em: 445±35 nm). An automated

algorithm was developed using Harmony 3.0 (Perkin Elmer) to quantify the relative CSC

proportion. In summary, nuclear Regions of Interest (ROIs) were identified within the

Hoechst channel, and the mean ALDEFLUOR signal, corrected for background, was

quantified. Cells were defined as ALDH-positive when this signal exceeded the one

quantified in the DEAB condition. The relative CSC proportion was computed as the number

of ALDH-positive cells relative to the total number of cells.

Data Preprocessing and Hit Selection

For each culture plate, the total cell amount and the percentage of ALDH-positive cells

measured in sample wells were first normalized to the average values of the corresponding

negative control (NEG) wells. The normalized results were labeled as *Relative Total Cell*

Amount and *Relative % ALDH-positive cells*. A time dependent decay of the ALDEFLUOR

signal was observed, leading to a slight decrease in the *Relative % ALDH-positive cells*

measured over the course of plate acquisitions. To account for and adjust this decay, a non-

linear polynomial regression model was established for each plate to describe the relationship

between the median *Relative % ALDH-positive cells* per column and the column index. For

each column index, a correction factor was computed by dividing the plate's median *Relative*

% ALDH-positive cells by the predicted value from the model at the column index. These

correction factors were subsequently applied to each column to normalize the individual

Relative % ALDH-positive cell measurements. The corrected results were labeled as

Corrected Relative % ALDH-positive cells. Analysis of the *Corrected Relative % ALDH-*

positive cells revealed a non Gaussian, long tailed distribution in the sample population. To

normalize this distribution, a Box–Cox transformation was applied to the population. The

optimal λ coefficient for the Box–Cox transformation was determined by fitting a linear

regression to quantile to quantile (QQ) plots, constructed from quantiles of the Box–Cox

transformed *Corrected Relative % ALDH-positive cells* distribution plotted against quantiles

of a corresponding theoretical Gaussian distribution. An optimal $\lambda = 0.7$ was determined to

achieve the best linear fit. The normality of the Box–Cox transformed distribution was

confirmed using a Kolmogorov–Smirnov test ($p < 0.05$), and the transformed results were

labeled as *Box–Cox Corrected Relative % ALDH-positive cells.* The *Box–Cox Corrected*

Relative % ALDH-positive cells sample results were then statistically scored using a Z Score

method. miRNA mimics were selected as hits if their absolute Z Score was greater than 3.

Statistical analysis

For each experiment, data are shown as mean ± S.E.M of n independent experiments (n =

number of independent experiments). GraphPad Prism9 software was used for data analysis.

The Mann Whitney test was used to analyse the difference between two groups of quantitative

variables with alpha value set at 5%. *p<0.05; **p<0.005; ***p<0.001.

RESULTS

A miRnome-wide high content screen identifies *miR-4745* as a negative regulator of the ALDH-positive CRC population.

We, and others, previously reported that the brightest subpopulation of CRC cells, identified using ALDEFLUOR^TM assay, a reporter for cellular ALDH enzymatic activity, is highly enriched in CSCs (10,21,29). These cells exhibit enhanced spheroid-forming efficiency and greater resistance to FIRI treatment. To investigate miRNAs involved in regulating colon CSCs, we conducted a miRNome-wide gain-of-function screen on a miniaturized ALDEFLUOR-based detection assay. This involved the systemic testing of a miRNA mimics library in the SW620 cell line (**Figure 1A**). SW620 was selected as the primary screening platform because it combines high technical suitability for automated high-content imaging with a biologically aggressive, metastatic phenotype characterized by a robust and stable ALDH+ cancer stem cell subpopulation. After 72 hours of miRNA transfection, CSCs were detected and quantified by measuring ALDEFLUOR-positive cells through high-content epifluorescence microscopy. The process involved identifying and counting cell nuclei stained with Hoechst dye and measuring cellular ALDEFLUOR intensities. ALDEFLUOR-positive cells were defined as those with intensities above the average intensity of cells treated with the ALDH inhibitor DEAB. The PARi platform optimized and miniaturized the ALDEFLUOR assay for 96-well plates, and developed automated procedures for image acquisition and analysis, as previously described for breast cancer cells (15). The screen revealed multiple miRNAs that significantly decreased ALDEFLUOR activity in SW620 cells (**Figure 1B; Supplementary Table S4**).

Among the top miRNAs identified, miR-*137* (31), *miR-202-5p* (32) and *miR-4443* (33), have been previously reported to regulate multiple pathways involved in CSC maintenance and function, supporting the validity of our screening results. Subsequently, miRNAs were

selected based on their relevance to CRC, CSC biology, and key phenotypic properties such as chemoresistance, self-renewal, and tumor initiation, with a particular focus on miRNAs that have been less extensively characterized but show promising potential.

We validated the activity of the selected miRNAs across three independent and complementary cellular models: the established SW620 cell line, patient-derived CRC cells from a primary tumor (CPP1), and patient-derived circulating tumor cell (CTC) lines (CTC44). CPP1 and CTC44 were specifically selected because they retain native genetic alterations and phenotypic heterogeneity that are often lost in long-established, immortalized cell lines. The inclusion of CTC44 is particularly significant as it provides a clinically relevant model of metastatic disease that retains the stem-like and chemoresistant features reflective of the original patient tumor. ALDEFLUOR activity was quantified by flow cytometry. Consistent with the screening findings, overexpression of *miR-4745*, *miR-497*, *miR-4723*, and *miR-4784* led to a significant decrease in the percentage of ALDH-positive cells relative to cells transfected with the negative control miRNA mimic (miCTRL) in all three cell lines (**Figure 1C**). This reduction in ALDH-positive cells correlated with decreased ALDH1A1 mRNA expression (**Figure 1D**). Among these candidates, *miR-4745* emerged as the most potent negative regulator of both the ALDH-positive cell population and ALDH1A1 mRNA expression, prompting further investigation into its effects.

To confirm that the reduction in ALDEFLUOR activity in CRC cells was specifically mediated by *miR-4745*, we used antagomiR-4745, a chemically modified RNA molecule designed to inhibit endogenous *miR-4745*. CPP1 and SW620 cells were co-transfected with either antagomiR-4745 (anti-*miR-4745*) to block miR-4745 activity, or a negative control (anti-miCTRL), along with *miR-4745* or miCTRL mimics. As shown in **Figure 1E**, the *miR-4745* mimic significantly decreased ALDEFLUOR activity in cells co-transfected with anti-miCTRL, but this effect was abolished in cells co-transfected with antagomiR-4745. This

confirms that the reduction in ALDEFLUOR activity was specifically due to *miR-4745*, as

blocking its activity with antagomiR negated the effect.

Finally, we assessed *miR-4745* expression in CSCs enriched from CRC cells using

ALDEFLUOR cell sorting. We found that *miR-4745* expression was significantly lower in

CSC populations (ALDH-positive cells) compared to non-CSC populations (ALDH-negative

cells) sorted from SW620, CPP1, HT29, and CTC44 cell lines respectively (**Figure 1F**).

These findings confirm that *miR-4745* expression or activity is downregulated in CSCs.

Overexpression of *miR-4745* represses the CSC phenotype *in vitro*.

To assess the functional effects of *miR-4745* modulation on CSC fate, CRC cell models,

including patient-derived CTC lines representing metastatic and therapy resistant disease

states, were transfected with a *miR-4745-3p* mimic. Initially, we confirmed that *miR-4745*

overexpression does not affect cell growth, as cells transfected with the *miR-4745* mimic

exhibited growth rates similar to those transfected with the control miRNA (miCTRL)

(**Supplementary Figure 1A**). This indicates that any observed effects of *miR-4745* on the

CSC phenotype are not attributable to changes in cell proliferation.

Subsequently, we assessed the presence of CD44v6-positive cells, a membrane marker linked

to cancer stem cells and colorectal cancer metastasis (34), in CPP1 and CTC44 cells following

miR-4745 overexpression. We observed that *miR-4745* overexpression significantly reduced

the percentage of CD44v6-positive cells (**Figure 2A**) and decreased CD44v6 mRNA

expression (**Supplementary Figure 1B**) compared to cells transfected with the control

miRNA (miCTRL).

We further evaluated the impact of *miR-4745* on the self-renewal capacity of CRC cells by

assessing their ability to form tumorspheres. As shown in **Figure 2B** and **Supplementary**

Figure 1C, cells transfected with the *miR-4745* mimic exhibited significantly lower spheroid-

forming efficiency compared to those transfected with the negative control (miCTRL). To counteract the effects of *miR-4745* on sphere formation, we used antagomiR-4745. As depicted in **Supplementary Figure 1C**, the *miR-4745* mimic reduced the number of spheres in CPP1 and SW620 cells co-transfected with the antagomiR control sequence (anti-miCTRL + *miR4745*), but this effect was abolished in cells co-transfected with the *miR-4745* inhibitor (anti-*miR4745* + *miR4745*). Notably, the inhibitory effect of *miR-4745* extended beyond primary spheres to subsequent generations (**Figure 2C**), emphasizing its role in diminishing CSC self-renewal ability.

Next, we extended our investigation to chemoresistance, another hallmark of CSCs. We examined the effect of *miR-4745* overexpression on the survival of ALDH-positive patient-derived CRC cells treated *in vitro* with a standard CRC chemotherapy cocktail (FIRI = 5-fluorouracil [5-FU] + SN38, the active metabolite of irinotecan). As shown in **Figure 2D** and **Supplementary Figure 1D**, *miR-4745* transfection significantly decreased the survival of ALDH-positive cells after 48 hours of FIRI treatment, resulting in a four-fold decrease in EC_{50} (half maximal effective concentration) for CPP1 and CTC44 cells compared to those transfected with miCTRL ($p < 0.05$). Furthermore, **Figure 2E** and **Supplementary Figure 1E** illustrate that, although the proportion of ALDH-positive cells increased significantly following chemotherapy in cells transfected with miCTRL, this enrichment was notably reduced in cells transfected with *miR-4745*. These findings suggest that *miR-4745* sensitizes CSCs to chemotherapy. Taken together, these findings suggest that elevated *miR-4745* expression influences the cancer stem cell phenotype.

***KLC2* is a potential target gene of *miR-4745*.**

Having established *miR-4745* as a regulator of CSC traits, our next objective was to identify its molecular targets. Given that miRNAs can modulate gene expression by degrading target

mRNAs or inhibiting their translation, we conducted both transcriptome and translatome analyses in CPP1 cells overexpressing *miR-4745*. For transcriptome analysis, total RNAs were isolated from cytoplasmic lysates. For translatome analysis, mRNAs were isolated from polysome fractions of the same lysates, separated by sucrose gradient ultracentrifugation.

Overexpression of *miR-4745* did not alter the polysome profile in CPP1 cells, indicating that global protein synthesis remains unaffected (**Figure 3A**). To further investigate changes in translation, we sequenced mRNAs from both poorly translated fractions (light fractions: 1 to 3 ribosomes per mRNA) and highly translated fractions (heavy fractions: more than 3 ribosomes per mRNA) of the polysome profile. Overexpression of *miR-4745* in CPP1 cells resulted in selective modulation of the transcriptome, with 25 transcripts significantly upregulated and 20 downregulated (adjusted p-value < 0.05 and a fold change > 2 or < 0.5), highlighting a focused regulatory effect. Similarly, *miR-4745* overexpression induced targeted changes in mRNA translation, with 23 highly translated mRNAs upregulated and 17 downregulated, suggesting a refined role in translational control (**Figure 3B and Supplementary Tables S5 & S6**).

We then validated the RNA sequencing results by analyzing the expression of two of the most downregulated genes (*KLC2* [Kinesin Light Chain 2] and *HEY2* [Hes Related Family BHLH Transcription Factor With YRPW Motif 2]) and two of the most upregulated genes (*CCND3* [Cyclin D3] and *MIR31HG* [miR-31 host gene]) at both transcriptional and translational levels, following transient overexpression of *miR-4745* in SW620 and CPP1 cell lines (**Figure 3C**).

Within each category of significantly differentially expressed genes (DEGs) (see section *Bioinformatic analysis of transcriptome and translatome sequencing data* in Materials and Methods), we searched for *miR-4745* target sites using TargetScan, TargetMiner, and the miRDB database. Since *miR-4745* target sites are preferentially located in the 3' UTR of

messenger RNAs, we computed the intersection of DEGs in each category with the set of

predicted *miR-4745* target genes from each tool or database. As shown in the Venn diagram in

Figure 3D, only one differentially expressed gene was predicted by all three approaches to

have a *miR-4745* target site in its 3' UTR. This gene was *KLC2*, which is regulated at both

mRNA and protein levels. The downregulation of KLC2 at the protein level with *miR-4745*

overexpression was confirmed by immunoblotting in CPP1 and SW620 cells (**Figure 3E**).

***MicroRNA-4745* targets *KLC2* mRNA by binding to a specific sequence within its**

3'untranslated region (3'UTR).

To experimentally confirm the binding of *miR-4745* to *KLC2* mRNA, we employed a

fluorescence-based dual reporter assay system using a pmiR-sensor plasmid (16). This

plasmid features a bidirectional promoter that drives the simultaneous expression of GFP

from *Aequorea coerulescens* (AcGFP) and the red fluorescent protein DsRed1 (22). The

GFP's 3'UTR was modified to include the full-length 3'UTR of *KLC2* (*KLC2*-3'UTR), which

contains a binding site for *miR-4745* (**Figure 4A**). Thus, the level of GFP reflects *miR-4745*

activity, while DsRed1 expression normalizes for expression variability and nonspecific

transcriptional regulation, providing a ratiometric measure (GFP/DsRed).

CPP1 cells were co-transfected with the miR-sensor plasmid containing the *KLC2* 3'UTR

(pmiR-*KLC2*-3'UTR), with or without a *miR-4745-3p* mimic. GFP and DsRed expression

levels were then measured using flow cytometry (**Figures 4B** to **4E**). As shown in **Figure 4E**,

co-transfection with the *miR-4745-3p* mimic specifically led to a decrease in GFP expression

in cells transfected with the pmiR-*KLC2*-3'UTR compared to other conditions, indicating that

miR-4745-3p binds to the *KLC2* 3'UTR.

To pinpoint the exact binding location of *miR-4745* within the *KLC2* 3'UTR, we used

TargetScan, an online database that predicts miRNA binding sites in mammalian mRNAs

transcripts. This analysis identified three 7-mer binding sites (seed sequence - CCGGGCC)

within the *KLC2* 3'UTR that match the seed region (position 2-8 of the mature miRNA) of

miR-4745 (**Supplementary Figure 2**). The locations were identified as loci 1, 2, and 3. Two

copies of each locus were cloned into the 3'UTR of GFP in the pmiR-sensor plasmid, and

mutant versions of these loci (CGGGCC to CAAACC) were generated to disrupt *miR-4745*

binding. As shown in **Figures 4C** and **4D**, GFP intensity significantly decreased when the

wild-type locus 2 construct was co-transfected with *miR-4745*, but this effect was rescued

when the binding site at locus 2 was mutated. No significant effects were observed at the

other loci (**Figure 4D**). This data confirms that *miR-4745* specifically binds to locus 2 within

the *KLC2* 3'UTR and downregulates *KLC2* expression. Furthermore, mutating the same locus

in the pmiR-KLC2-3'UTR plasmid also rescued the effect of the *miR-4745* mimic (**Figure

4E**). An overall decrease in fluorescence was seen when the full-length *KLC2* 3'UTR was

cloned into the 3'UTR of GFP compared to the pmiR-sensor, which may be due to its larger

size (~1 kb). Overall, these findings validate the specific binding of *miR-4745* at locus 2

within the *KLC2* 3'UTR, leading to the downregulation of *KLC2* expression.

Downregulation of *KLC2* reduces the colorectal CSC phenotype.

Our research has established that *miR-4745* modulates *KLC2* expression, suggesting that the

miR-4745-KLC2 axis may be involved in the CSC phenotype. To elucidate the direct role of

KLC2 in this process, CPP1 and SW620 cells were transfected with siRNA targeting *KLC2*

(si*KLC2*) (**Supplementary Figures 3A** and **3B**), and ALDEFLUOR activity was assessed.

Consistent with the effects observed with the *miR-4745* mimic, the knockdown of *KLC2*

resulted in a significant reduction in the ALDH-positive cell population in both CPP1 and

SW620 cells compared to controls transfected with the non-targeting siRNA (siCTRL)

(**Figure 5A**).

To evaluate the proportion of CSCs within the tumor population, we performed a sphere formation assay. As illustrated in **Figure 5B**, *KLC2* knockdown reduced the ability of CPP1 and SW620 cells to form tumorspheres *in vitro* by approximately 40% to 50%, with no change in cell proliferation (**Supplementary Figure 3C**). Conversely, transient overexpression of *KLC2* in CRC cells (**Supplementary Figure 3D**) enhanced the stem cell phenotype, as evidenced by increased ALDEFLUOR activity and improved tumorsphere formation capacity *in vitro* (**Figures 5C** and **5D**).

To demonstrate that the effect of *miR-4745* on the CSC phenotype is mediated through *KLC2*, SW620 and CPP1 cells were co-transfected with either a *KLC2* overexpression plasmid construct lacking the 3'UTR *miR-4745* binding site (oeKLC2) or a control plasmid (oeCTRL), along with *miR-4745* or miCTRL mimics. While ALDEFLUOR activity decreased in cells co-transfected with the control plasmid and *miR4745* mimic, the inhibitory effect of *miR-4745* on ALDH-positive cells was counteracted in cells co-transfected with the cDNA construct lacking the 3'UTR *miR-4745* binding site (**Figure 5E**). This rescue experiment supports the hypothesis that *KLC2* downregulation is a crucial mechanism through which *miR-4745* exerts its effects.

miR-4745 negatively correlates with *KLC2* expression in CRC cells

Given our demonstration of the direct post-transcriptional regulation of *KLC2* by *miR-4745* in colon cancer cells, we expected to observe an inverse relationship between *miR-4745* and *KLC2* levels in colorectal cancer cells. RT-qPCR analysis revealed a significant negative correlation between *KLC2* mRNA and *miR-4745* (r = -0.716, p = 0.0027) across a panel of CRC cell lines (SW480, SW620, HT29, LS174T, HCT116, Lovo, and Colo205), as well as in patient-derived CRC cells from primary tumors (CPP1, CPP25, CPP44), metastatic tumors (CPP6, CPP19, CPP36), and circulating tumor cells (CTC44, CTC45) (**Figure 6A**).

Furthermore, *miR-4745* expression was significantly lower in CSCs enriched from CRC cells

treated with chemotherapy (FIRI = 5-FU + SN38) compared to non-CSCs. This pattern was

consistent in both ATCC-derived CRC cells (HT29) and patient-derived CRC cells (CPP1,

CTC44, CTC45) under CSC-enriching conditions (**Figure 6B**). In contrast, *KLC2* mRNA and

protein levels significantly increased following FIRI treatment, exhibiting an expression

pattern opposite to that of *miR-4745* (**Figures 6B** and **6C**).

Additionally, *KLC2* transcript levels were elevated in ALDH-positive cells across all four

CRC lines tested, suggesting that the downregulation of *miR-4745* contributes to the

preferential expression of *KLC2* in these cells (**Figure 6D**).

Differential expression of *miR-4745* and *KLC2* in human CRC samples.

To further explore the expression patterns of *miR-4745* and *KLC2* in human colorectal cancer

samples, we analyzed data from The Cancer Genome Atlas (TCGA) and dbDEMC databases.

In the TCGA dataset, *miR-4745-3p* expression was found to be marginally lower in colon

adenocarcinoma samples compared to non-cancerous tissues (n=103 vs. n=3, p = 0.06),

although this difference was not statistically significant, possibly due to the small size of the

non-cancerous group. Conversely, analysis of the dbDEMC dataset demonstrated a significant

decrease in *miR-4745-3p* expression in colon adenocarcinoma samples (n=444) relative to

non-cancerous controls (n=8, p = 0.00125). Additionally, we measured *KLC2* mRNA

expression in colon cancer tissues from 41 patients across different stages of CRC: adenoma

(n=3), stage I (n=8), stage II (n=11), stage III (n=5), stage IV (n=8), and metastasis (n=6).

KLC2 expression showed a significant increase with early local progression of CRC,

suggesting its involvement in the early stages of tumor development (**Figure 6E**). However, a

reduction in *KLC2* mRNA levels was observed in stage IV and metastatic samples, where

cancer cells had spread from their primary site, suggesting that the gene may be subject to

microenvironment-dependent regulation or reprogramming in advanced disease stages (35,36)

(**Figure 6E**).

In addition, we sought to determine whether *KLC2* expression correlates with clinical

outcomes in colorectal cancer patients. We performed a Kaplan-Meier analysis of *KLC2*

mRNA expression in a cohort of stage I to IV CRC patients (37). Our analysis revealed that

patients with tumors exhibiting high *KLC2* expression had a significantly lower probability of

overall survival, indicating that elevated *KLC2* expression is associated with poor prognosis in

CRC patients (n=1 061; p=0.031, **Figure 6F**). Taken together, these data demonstrate that the

miR-4745-KLC2 axis may serve as a prognostic indicator for overall survival in CRC.

DISCUSSION

Colorectal cancer remains a major clinical challenge due to its high rates of chemoresistance

and tumor recurrence, primarily driven by CSCs. Among potential therapeutic targets,

miRNAs have gained significant attention for their pivotal role in regulating the CSC

phenotype (38), offering new opportunities for targeting these cells in cancer therapy. In this

study, we identified several miRNAs of interest that regulate the CSC phenotype, among

which *miR-4745* was selected for further characterization. We demonstrated that *miR-4745*

regulates CSC traits such as sphere-forming ability, self-renewal and chemoresistance.

Mechanistically we identified the 3' UTR of *KLC2* as a direct target of *miR-4745* and a key

intermediate in the molecular mechanism controlling CSC traits.

Our bioinformatic analysis revealed that, in addition to its role in regulating *KLC2*, *miR-4745*

indirectly downregulates the expression of genes that promote key hallmarks of cancer stem

cells, including self-renewal (e.g. *SQSTM1* (39) and SPP1 (40)), invasion and metastasis (e.g.

MMP15 (41). This occurs in part through the activation of signaling pathways such as the

Notch pathway (e.g. *HEY2,* (42)) or the ERK1/2 pathway (e.g. *GDF15*, (43), the modulation

of the cellular stress response (e.g. *CYP2E1* (44)), and drug resistance (e.g., *LGMN* (45) and

UCP2 (46)). Importantly, we observed that *miR-4745* controls the expression of genes

previously reported to negatively impact disease-free survival in colon cancer patients,

including *GDF15*, *UCP2*, and *SPP1*. These data strongly suggest that, in addition to *KLC2*,

miR-4745 downregulates a network of critical downstream targets that may play a role in

regulating the CSC phenotype.

KLC2, a light chain isoform of Kinesin-1, plays a critical role in the intracellular transport of

various cargoes along microtubules. Mutations in the *KLC2* gene are associated with several

neurological disorders, including spastic paraplegia, optic atrophy, and neuropathy (47).

Notably, *KLC2* overexpression has been shown to promote radioresistance in non-small cell

lung cancer (NSCLC), resulting in increased tumor volume and reduced survival in tumor-

bearing mice. This suggests a potential role for *KLC2* in the survival of CSCs following

radiotherapy. Additionally, elevated *KLC2* levels are correlated with decreased p53

phosphorylation and poor prognosis in NSCLC patients (48). *KLC2* is also a target of *miR-*

125b, a microRNA involved in cancer cell migration and metastasis (49). Nonetheless, the

target genes of *miR-125b* differ among various cancer types, in which its expression is

frequently dysregulated. One known binding partner of *KLC2* is *SMAD2*, which is crucial for

TGFβ signal transduction, leading to various effects during tumorigenesis. Further studies on

the downstream pathways of *KLC2* could help uncover the molecular mechanisms behind its

impact on CSC survival and its clinicopathological significance.

We observed a clear inverse correlation between *miR-4745* and *KLC2* mRNA levels across

various stages of CRC cell lines, substantiating the existence of a *miR-4745/KLC2* axis.

Furthermore, *KLC2* expression progressively increased with CRC advancement, peaking in

stage III patient samples, suggesting it may serve as a potential local driver of CRC

progression. However, further investigation is needed to understand the underlying reasons

for the reduction in *KLC2* expression in stage IV tissues and metastatic sites.

Our study emphasizes the promise of targeting the *miR-4745/KLC2* pathway as a therapeutic

strategy for colorectal cancer. To ensure that the observed effects on colorectal CSC traits

were biologically relevant, and not simply the result of supraphysiological levels of *miR-*

4745-3p, we quantified its expression by RT-qPCR in two CRC cell models transfected with

decreasing concentrations of *miR-4745-3p* mimic. Although the effects were attenuated

compared to transfection with 50 nM mimic, we still observed a measurable impact on the

cells' sphere-forming capacity at concentrations as low as 0.001 nM (**Supplementary Figure**

4B). Notably, this concentration aligns with the endogenous expression level of *miR-4745-3p*

observed under control mimic conditions (**Supplementary Figure 4A**). These results suggest

that the *miR-4745/KLC2* regulatory axis remains biologically relevant in CRC, even if its

impact under physiological conditions may be moderate. Although *miR-4745-3p* exhibits low

endogenous expression, it is likely that its regulation is context-dependent or limited to

particular CRC subtypes or stem-like cell populations, analogous to other key miRNAs that

play roles in stemness or stress response mechanisms.

Modulating the *miR-4745/KLC2* pathway may help reprogram CSC behavior, offering a

strategy to overcome chemoresistance. For example, upregulating *miR-4745* or directly

targeting *KLC2* could sensitize CSCs to chemotherapy, potentially reducing the likelihood of

tumor recurrence and improving patient outcomes. While miRNA-based therapies faces

challenges, particularly in achieving targeted and efficient systemic delivery, innovative

technologies, including nanocapsules, nanocarriers, micro/nanoparticles, liposomes and

PEGylated vesicles have been developed to address these limitations (50). Another promising

alternative to modulate the *miR-4745/KLC2* axis could involve pharmacological agents that

act as endogenous inducers or activators of *miR-4745*. In our previous study, we successfully

induced *miR-148a-3p* expression with the anthelmintic drug niclosamide (16), demonstrating

a practical strategy to enhance miRNA activity *in vivo* and raising the possibility of

repurposing niclosamide as an adjuvant strategy to reduce CSC-mediated chemoresistance.

Furthermore, assessing *miR-4745* expression in colorectal tumor samples, including those

from chemotherapy-treated patients, could provide deeper insights into its role in therapy

resistance. As mentioned, directly targeting *KLC2* through the development of a specific

inhibitor or using PROTAC technology might represent another strategy to regulate the *miR-*

4745/KLC2 axis.

Finally, our miRnome-wide high content screening efforts identified other miRNAs, including

miR-497, *miR-4723*, and *miR-4784*, which also contribute to reducing the CSC phenotype.

Further investigation into the molecular mechanisms of these miRNAs could provide broader

insights into CSC regulation and potential therapeutic strategies.

CONCLUSIONS

In conclusion, our study identifies *miR-4745* as a novel and functionally significant regulator

of colorectal CSC properties, acting in part through the direct repression of the kinesin light

chain protein *KLC2*. We demonstrate that *miR-4745* impairs key CSC traits, including self-

renewal, chemoresistance, and sphere-forming capacity, and delineate a broader regulatory

network involving multiple downstream effectors implicated in CSC maintenance, tumor

progression, and adverse clinical outcomes. The observed inverse correlation between *miR-*

4745 and *KLC2* expression across CRC stages reinforces the relevance of this regulatory axis

in colorectal tumorigenesis. Although endogenously expressed at low levels, *miR-4745-3p*

exhibits biologically meaningful activity within a physiologically relevant range, suggesting

its potential as a context-specific modulator of CSC behavior. These findings provide a

compelling rationale for therapeutic strategies aimed at restoring *miR-4745* expression or

pharmacologically inhibiting *KLC2* to overcome CSC-driven chemoresistance. Finally, the

identification of additional CSC-targeting miRNAs through our high-content screening

broadens the scope for future combinatorial approaches. Altogether, our results highlight the

therapeutic potential of miRNA-based interventions in targeting colorectal CSCs and

improving patient outcomes.

ABBREVIATIONS

3' UTR : 3' Untranslated region

5-FU : 5-fluorouracil

ABCB1 : ATP-Binding Cassette Subfamily B Member 1

ABCG2 : ATP-Binding Cassette G2

AcGFP : *Aequora coerulescens* Green fluorescent protein

ALDH1A1 : Aldehyde Dehydrogenase 1A1

ATCC : American Type Culture Collection

CCND3 : Cyclin D3

CRC : Colorectal cancer

CSCs : Cancer stem cells

CYP2E1 : Cytochrome P450 2E1

dbDEMC : database of Differentially Expressed MiRNAs in human Cancers

DEAB : diethylaminobenzaldehyde

DEG : Differentially expressed genes

DMEM : Dulbecco's Modified Eagle Medium

DS-Red : *Discosoma sp.* red fluorescent protein

EGF: Epidermal growth factor

ERK1/2 : Extracellular Signal Regulated Kinase 1/2

FBS : Fetal bovine serum

FGF: Fibroblast growth factor

FSC : Forward Scatter

GAPDH : Glyceraldehyde-3-phosphate dehydrogenase

GDF15: Growth differentiation factor 15

GO : Gene Ontology

HDAC4 : Histone Deacetylase 4

HEY2 : Hes Related Family BHLH Transcription Factor With YRPW Motif 2

KLC2 : Kinesin-light chain 2

LGMN : Legumain

mRNA : messenger RNA

miRNA : micro-RNA

MIR31HG : miR-31 Host Gene

MMP15 : Matrix metalloproteinase 15

NSCLC : Non-Small Cell Lung Cancer

OE : overexpression

PXR : Pregnane X Receptor

QQ : quantile-to-quantile

RNA-seq : RNA sequencing

RPLO : Ribosomal protein L0

RPL13 : Ribosomal protein L13

RPMI : Roswell Park Memorial Institute Medium

siRNA : small interfering RNA

SMAD2 : SMAD Family Member 2

SPP1 : Secreted Phosphoprotein 1

SQSTM1 : Sequestosome 1

SSC : Side Scatter

TCGA: The Cancer Genome Atlas

TGF-β : Transforming growth factor-β

UCP2 : Uncoupling Protein 2

WHO : World Health Organization

COMPETING INTERESTS

The authors declare no potential conflicts of interest.

AUTHORS' CONTRIBUTIONS

C.P., A.D., J.M.P., and J.P. designed the research; C.P., A.D, K.L., J.M.P, and J.P. designed

the experiments; P.B., K.L., J.R., O.B., G.P., A.B., S.M., C.M., E.P., L.B., C.D., A.A.G., and

C.P. performed the experiments; P.B., K.L., J.R., O.B., G.P., A.B., S.M., C.M., E.P., L.B.,

C.D., A.A.G., E.R., J.P., J.M.P., A.D., and C.P. analyzed the data; C.P. K.L., and A.D. wrote

the manuscript.

ACKNOWLEDGMENTS

We thank C. Duperray and F. Leccia (IRMB, Montpellier) from the Montpellier RIO Imaging

platform for flow cytometry experiments.

FUNDING DECLARATION

MGX acknowledges financial support from France Génomique National infrastructure,

funded as part of "Investissement d'Avenir" program managed by Agence Nationale pour la

Recherche (contrat ANR-10-INBS-09). This work was supported by grants from the Ligue

Contre le Cancer, INCa, Association pour la Recherche sur le Cancer, Cancéropôle GSO, and

Siric of Montpellier Cancer (France).

DECLARATION

We confirm that all research involving human subjects, human material, or human data in our

study was conducted in full accordance with the ethical principles outlined in the Declaration

of Helsinki.

Cell lines and patient-derived tumor cell cultures:

Patient-derived colon cancer cell lines (CPP1, CPP6, CPP19, CPP25, CPP36, and CPP44) and

circulating tumor cell lines (CTC44 and CTC45) were established from colorectal cancer

biopsies and peripheral blood samples collected at CHU-Carémeau (Nîmes, France) as part of

a previously clinical study (20,21), registered on ClinicalTrials.gov (Identifier

NCT01577511 ; Principal Investigator: Dr. Jean-François Bourgaux ; First submitted: 2012-

04-12; First submitted that met QC criteria : 2012-04-13 ; First posted (estimated) : 2012-04-

16 ; Last Update submitted that met QC criteria : 2021-06-18 ; Last update posted: 2021-06-

21; Last verified : 2020-01 ; **https://clinicaltrials.gov/study/NCT01577511**).

The study protocol was approved by the French Ethics Committee "Comité de Protection des

Personnes Sud Méditerranée III (authorization number 2011-A01141-40)," which operates in

full compliance with the Declaration of Helsinki. Written informed consent was obtained

from all participants involved in the study, in accordance with all relevant ethical regulations.

Commercial CRC cell lines used in this study - SW480 (RRID: CVCL_0546), SW620

(RRID:CVCL_0547), HT29 (RRID:CVCL_A8EZ), LS174T (RRID:CVCL_1384), HCT116

(RRID:CVCL_0291), LoVo (RRID:CVCL_0399), and Colo205 (RRID:CVCL_0218) were

obtained from ATCC and used in accordance with standard ethical practices and institutional

guidelines.

Colon tumor tissue samples:

RNA extracted from 41 colon tumor tissue samples was obtained through the Clinical and

Biological Database BCBCOLON (Institut du Cancer de Montpellier—Val d'Aurelle, France)

registered at ClinicalTrials.gov (Identifier #NCT03976960 ; Principal Investigator: Dr. Eric

Assenat ; First submitted: 2019-06-05; First submitted that met QC criteria : 2019-06-05 ;

First posted : 2019-06-06 ; Last Update submitted that met QC criteria : 2025-02-11 ; Last

update   posted: 2025-02-12;   Last   verified :   2025-02 ;

https://clinicaltrials.gov/study/NCT03976960).

These samples were collected in compliance with French legislation and declared to the

French Ministry of Higher Education and Research (Declaration No. DC-2008–695). The

study received approval from the French Ethics Committee (CPP Sud Méditerrannée III,

Ref#2014.02.04) and the ICM Translational Research Committee (ICM-CORT-2018-28). All

participants provided written consent, and all relevant ethical regulations were strictly

followed.

FIGURE LEGENDS

Figure 1: miRnome-wide High Content Screen identifies *miR-4745* as a negative

regulator of the ALDH-positive CRC population.

(a) Experimental design for the miRnome-wide gain-of-function screen employing pre-miRs

as miRNA mimics in a scaled-down ALDEFLUOR detection assay. Following the individual

transfection of miRNA mimics from a human miRNome-wide library into SW620 colon

cancer cells, high-content screening was employed to measure the proportion of CSC

(ALDEFLUOR-positive cells) in cells transfected with miRNA mimics or miRVana

microRNA Mimic Negative Control #1 (miCTRL). **(b)** MA-plot of the high-content screening

data, showing the Z-score of Aldefluor activity (mean of three replicates) in SW620 cells

transfected with miRNA mimics from the human miRNome-wide library. The Z-scores are

plotted against the plate well index, indicating the specific position of each miRNA mimic on

the plate. **(c)** Percentage of Aldefluor-positive cells (% ALDH$^+$ cells) in patient-derived CPP1

and SW620 colon cancer cells or circulating tumor cell line (CTC44) transfected for 72 hours

with *miR-4745*, *miR-497*, *miR-4723*, *miR-4784* or miCTRL mimics. Data are expressed as

mean ± SEM (n = 6). **(d)** RT-qPCR analysis of *ALDH1A1* mRNA expression in patient-

derived CPP1 and SW620 colon cancer cells or circulating tumor cell line (CTC44)

transfected for 72 hours with *miR-4745*, *miR-497*, *miR-4723*, *miR-4784* or miCTRL mimics.

Data are expressed as mean ± SEM (n=3). Fold change is calculated relative to cells

transfected with negative control mimic. **(e)** Percentage of Aldefluor-positive cells (%

ALDH$^+$ cells) in CPP1 or SW620 cells co-transfected for 72 hours with either anti-miR-4745

or anti-miCTRL, along with *miR-4745* or miCTRL mimics. Data are expressed as mean ±

SEM (n = 4). **(f)** Expression levels of *miR-4745-3p* in patient-derived CPP1, SW620, and

HT29 colon cancer cells, as well as in the circulating tumor cell line (CTC44), following cell

sorting based on Aldefluor activity (ALDH-negative and ALDH-positive populations). Data

are presented as mean ± SEM (n = 3) and are expressed as fold change relative to the ALDH-

negative population. *, p < 0.05, **, p < 0.005, ***, p < 0.001, ns = not significant, Mann

Whitney test.

Figure 2: Overexpression of *miR-4745* represses the CSC phenotype *in vitro*.

(a) Quantification of the CD44v6-APC-positive population in CPP1 (n=7) and CTC44 (n=5)

cells transfected with *miR-4745-3p* (*miR4745*) or miRVana microRNA Mimic Negative

Control #1 (miCTRL) mimics (right panel). Percentage of CD44v6-APC positive CPP1 cells

is indicated in inset boxes for a representative experiment (left panel). **(b)** Representative

images of tumorspheres (upper panel) and the percentage (lower panel) of sphere-forming

cells in patient-derived CPP1 and circulating tumor (CTC44) colon cancer cells transfected

with *miR-4745-3p* (*miR4745*) or miCTRL mimics. Data from individual replicates (n=15 per

experiment) are presented, along with the mean ± SEM from six independent experiments. **(c)** Percentage of sphere-forming cells in patient-derived CPP1 colon cancer cells transfected with *miR4745* or negative control (miCTRL) mimics and, maintained as first-generation colonospheres (S1) or after several passages (S2, S3). Data from individual replicates (n=12 per experiment) are presented, along with the mean ± SEM from three independent experiments. **(d)** Percentage of surviving cancer stem cells (Aldefluor-positive) measured 48 hours following treatment with specified concentrations of FIRI (1X = 50 µM 5-FU + 500 nM SN38) *in vitro*. Sorted CPP1 Aldefluor-positive cells were first transfected with 50 nM *miR-4745-3p* (*miR-4745*) or miRVana microRNA Mimic Negative Control #1 (miCTRL) mimics, 24 hours prior to FIRI treatment. Data are expressed as mean ± SEM (n = 3). **(e)** Percentage of Aldefluor-positive cells (% ALDH$^+$ cells) in CPP1 cells transfected with *miR-4745* or miCTRL mimics and then exposed for 72 hours to chemotherapy (Firi=5µM 5-FU + 50nM SN38). Data are expressed as mean ± SEM (n=6). *, $p<0.05$; **, $p<0.005$, ***, $p < 0.001$, Mann Whitney test.

Figure 3: *KLC2* is a potential target gene of *miR-4745*.

(a) Representative polysome profiling of CPP1 cells transfected with *miR-4745-3p* (*miR4745*) or miCTRL mimics for 72 hours. The profiles show peaks corresponding to the small (40S) and large (60S) ribosomal subunits, free fractions, light fractions (1 to 3 ribosomes per mRNA) and heavy fractions (more than 3 ribosomes per mRNA). This experiment was repeated three times. **(b)** Scatter plot showing the differential distribution of the log2 fold change of RNA from heavy fractions (y-axis) versus the log2 fold change of RNA from free fractions (x-axis) in CPP1 cells transfected with *miR-4745* or miCTRL mimics. **(c)** mRNA expression levels of potential *miR-4745* target genes were assessed in CPP1 and SW620 cells transfected with *miR-4745-3p* (*miR4745*) or miRVana microRNA Mimic Negative

Control #1 (miCTRL) mimics for 72 hours. Data are presented as mean ± SEM (n = 3) and are expressed as fold change relative to cells transfected with the miCTRL mimic. (**d**) Venn diagram showing the intersection of the sets of *miR-4745* target genes predicted by TargetScan (blue), TargetMiner (yellow), miRDB (red), and the set of differentially expressed genes (DEGs) regulated at both transcriptional and translational levels (green - UP or DOWN). The intersection of all four sets contains a single gene, *KLC2*. (**e**) Representative Western blot analysis (upper panel) and quantification (lower panel) of KLC2 protein levels in patient-derived CPP1 and SW620 colon cancer cells transfected for 72 hours with either *miR-4745-3p* (*miR4745*) or miCTRL mimics. Data are presented as mean ± SEM (n=4). ***, p<0.001, Mann Whitney test.

Figure 4: *MicroRNA-4745* targets *KLC2* mRNA by binding to a specific sequence in its 3'untranslated region (3'UTR).

(**a**) The schematic illustration depicts the dual-reporter pmiR-sensor plasmid, in which the full- length *KLC2*-3'UTR or one of its three predicted binding sites (loci #1, #2, and #3) for *miR-4745*, is inserted into the 3'UTR of AcGFP (GFP). (**b**) Flow cytometry was used to measure the fluorescence intensities of GFP and DsRed in DsRed-positive cells. (**c**) Flow cytometry analysis (count versus GFP) was conducted on CPP1 cells co-transfected with a pmiR-sensor plasmid containing two copies of either the wild-type (WT) or mutated (Mutant) predicted binding site within the KLC2-3'UTR (pmiR-locus #2 WT or mutant), along with a *miR-4745-3p* mimic. (**d**) The GFP/DsRed ratio (geometric mean of intensities) was analyzed for CPP1 cells co-transfected with the pmiR-sensor plasmid containing two copies of the predicted binding sites within the KLC2-3'UTR or their mutant versions (sequences listed in **Supplementary Table S4**), along with a *miR-4745-3p* mimic. (**e**) The GFP/DsRed ratio (geometric mean of intensities) was analyzed for CPP1 cells co-transfected with the pmiR-

sensor plasmid containing the full-length KLC2-3'UTR or its mutated version at locus #2,

along with a *miR-4745-3p* mimic. Data are expressed as mean ± SEM (n=3). **, p<0.005,

Student's t-test.

Figure 5: *KLC2* inactivation inhibits colorectal CSC phenotype.

(a) Percentage of Aldefluor-positive cells (% ALDH$^+$ cells) in CPP1 or SW620 cells

transfected for 72 hours with 50nM of control (siCTRL) or KLC2-specific (*siKLC2*) siRNA.

Data are expressed as mean ± SEM (n = 9 for CPP1 and n = 5 for SW620). **(b)** Percentage of

sphere-forming cells among sorted live (Sytox Blue-negative) CPP1 and SW620 cells

transfected for 72 hours with 50nM of control (siCTRL) or KLC2-specific (siKLC2) siRNA.

Data are expressed as mean ± SEM (n=3 experiments with 15 individual replicates per

experiment). **(c)** Percentage of Aldefluor-positive cells (% ALDH$^+$ cells) in CPP1 or SW620

cells transfected for 72 hours with either a control plasmid (oe-CTRL) or a plasmid

overexpressing *KLC2* (oeKLC2). Data are expressed as mean ± SEM, n = 5. **(d)** Percentage of

sphere-forming cells among sorted live (Sytox Blue-negative) CPP1 and SW620 cells

transfected for 72 hours with either a control plasmid (oe-CTRL) or a plasmid overexpressing

KLC2 (oeKLC2). Data are presented as mean ± SEM, n = 3 (10 individual replicate values

per experiment). **(e)** Percentage of Aldefluor-positive cells (% ALDH$^+$ cells) in CPP1 or

SW620 cells co-transfected for 72 hours with either a control plasmid (oe-CTRL) or a

plasmid overexpressing *KLC2* lacking the 3'UTR *miR-4745*-targeting sequence (oeKLC2),

along with miR-4745 or miCTRL mimics. Data are presented as mean ± SEM (n = 4). *, p <

0.05; **, p < 0.005; ***, p < 0.001; ns = not significant, Mann Whitney test.

Figure 6: *KLC2* is inversely correlated with *miR-4745-3p* in CRC and is upregulated

under chemotherapy treatment and with local CRC progression.

(a) Inverse correlation between *KLC2* and *miR-4745-3p* expression in CRC cell lines obtained from ATCC (SW480, SW620, HT29, LS174T, HCT116, Lovo, and Colo205, black squares) and patient-derived CRC cells established in the IGF laboratory, including those from primary tumors (CPP1, CPP25, CPP44), metastatic tumors (CPP6, CPP19, CPP36), and blood (circulating tumor cells, CTC44, CTC45, red squares). Expression levels were measured by RT-qPCR. *KLC2* expression (mean of each technical replicate) was normalized to RPL0 (Ribosomal protein L0), while *miR-4745-3p* expression was normalized to *miR-103a-3p*. **(b)** *miR-4745-3p* (upper panel) and *KLC2* (lower panel) mRNA expression in HT29, CPP1, CTC44, and CTC45 colon cancer cells after 72 hours of chemotherapy treatment (Firi = 5 µM 5-FU + 50 nM SN38). Data are presented as mean ± SEM (n=3), showing fold change compared to untreated cells (NT). **(c)** Representative Western blot analysis (upper panel) and quantification (lower panel) of KLC2 protein expression levels in HT29, CPP1, CTC44, and CTC45 colon cancer cells after 72 hours of chemotherapy treatment (Firi = 5 µM 5-FU + 50 nM SN38). Data are presented as mean ± SEM (n=4), showing fold change compared to untreated cells (NT). **(d)** Expression levels of *KLC2* in patient-derived CPP1, SW620, and HT29 colon cancer cells, as well as in the circulating tumor cell line (CTC44), following cell sorting based on Aldefluor activity (ALDH-negative and ALDH-positive populations). Data are presented as mean ± SEM (n = 3) and are expressed as fold change relative to the ALDH-negative population. **(e)** *KLC2* mRNA expression in colon cancer tissue samples from 41 patients at various stages of cancer: adenoma (n = 3), primary adenocarcinoma at stages I (n = 8), II (n = 11), III (n = 5), IV (n = 8) and metastases (n = 6). Each dot represents an individual value. *KLC2* mRNA expression was normalized to actin. *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$, Mann Whitney test (**b** to **d**) or two-way ANOVA (Tukey's multiple comparisons test, **e**). **(f)** Kaplan-Meier estimates of overall survival probability based on *KLC2* expression levels in a cohort of 1 061 stage I to IV colorectal cancer patients

(https://kmplot.com/analysis/index.php?p=service&cancer=colon). Patients with low *KLC2*

expression are depicted in black, while those with high *KLC2* expression are shown in red.

The p-value from the likelihood ratio test is indicated.

REFERENCES

1. Siegel RL, Wagle NS, Cercek A, Smith RA, Jemal A. Colorectal cancer statistics, 2023.
CA: A Cancer Journal for Clinicians. 2023;73(3):233□54.

2. Adorno-Cruz V, Kibria G, Liu X, Doherty M, Junk DJ, Guan D, et al. Cancer stem cells:
targeting the roots of cancer, seeds of metastasis, and sources of therapy resistance.
Cancer Res. 15 mars 2015;75(6):924□9.

3. Ricci-Vitiani L, Lombardi DG, Pilozzi E, Biffoni M, Todaro M, Peschle C, et al.
Identification and expansion of human colon-cancer-initiating cells. Nature. 4 janv
2007;445(7123):111□5.

4. Kreso A, O'Brien CA, van Galen P, Gan OI, Notta F, Brown AMK, et al. Variable clonal
repopulation dynamics influence chemotherapy response in colorectal cancer. Science. 1
févr 2013;339(6119):543□8.

5. Todaro M, Alea MP, Di Stefano AB, Cammareri P, Vermeulen L, Iovino F, et al. Colon
cancer stem cells dictate tumor growth and resist cell death by production of interleukin-
4. Cell Stem Cell. 11 oct 2007;1(4):389□402.

6. Dylla SJ, Beviglia L, Park IK, Chartier C, Raval J, Ngan L, et al. Colorectal cancer stem
cells are enriched in xenogeneic tumors following chemotherapy. PLoS One. 18 juin
2008;3(6):e2428.

7. Merlos-Suárez A, Barriga FM, Jung P, Iglesias M, Céspedes MV, Rossell D, et al. The
intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease
relapse. Cell Stem Cell. 6 mai 2011;8(5):511□24.

8. Haraguchi N, Utsunomiya T, Inoue H, Tanaka F, Mimori K, Barnard GF, et al.
Characterization of a side population of cancer cells from human gastrointestinal system.
Stem Cells. mars 2006;24(3):506□13.

9. Huang EH, Hynes MJ, Zhang T, Ginestier C, Dontu G, Appelman H, et al. Aldehyde
dehydrogenase 1 is a marker for normal and malignant human colonic stem cells (SC) and
tracks SC overpopulation during colon tumorigenesis. Cancer Res. 15 avr
2009;69(8):3382□9.

10. Shenoy A, Butterworth E, Huang EH. ALDH as a marker for enriching tumorigenic
1025   human colonic stem cells. Methods Mol Biol. 2012;916:373□85.

1026   11. Fang Y, Sun B, Xiang J, Chen Z. MiR-301a promotes colorectal cancer cell growth and
1027   invasion by directly targeting SOCS6. Cell Physiol Biochem. 2015;35(1):227□36.

1028   12. Xu K, Liu X, Mao X, Xue L, Wang R, Chen L, et al. MicroRNA-149 suppresses
1029   colorectal cancer cell migration and invasion by directly targeting forkhead box
1030   transcription factor FOXM1. Cell Physiol Biochem. 2015;35(2):499□515.

1031   13. Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor
1032   suppressors. Dev Biol. 1 févr 2007;302(1):1□12.

1033   14. Liu C, Kelnar K, Liu B, Chen X, Calhoun-Davis T, Li H, et al. The microRNA miR-34a
1034   inhibits prostate cancer stem cells and metastasis by directly repressing CD44. Nat Med.
1035   févr 2011;17(2):211□5.

1036   15. El Helou R, Pinna G, Cabaud O, Wicinski J, Bhajun R, Guyon L, et al. miR-600 Acts as a
1037   Bimodal Switch that Regulates Breast Cancer Stem Cell Fate through WNT Signaling.
1038   Cell Rep. 28 févr 2017;18(9):2256□68.

1039   16. Bansard L, Bouvet O, Moutin E, Le Gall G, Giammona A, Pothin E, et al. Niclosamide
1040   induces miR-148a to inhibit PXR and sensitize colon cancer stem cells to chemotherapy.
1041   Stem Cell Reports. 12 avr 2022;17(4):835□48.

1042   17. Bitarte N, Bandres E, Boni V, Zarate R, Rodriguez J, Gonzalez-Huarriz M, et al.
1043   MicroRNA-451 is involved in the self-renewal, tumorigenicity, and chemoresistance of
1044   colorectal cancer stem cells. Stem Cells. nov 2011;29(11):1661□71.

1045   18. Smith AR, Marquez RT, Tsao WC, Pathak S, Roy A, Ping J, et al. Tumor suppressive
1046   microRNA-137 negatively regulates Musashi-1 and colorectal cancer progression.
1047   Oncotarget. 20 mai 2015;6(14):12558□73.

1048   19. Song B, Wang Y, Xi Y, Kudo K, Bruheim S, Botchkina GI, et al. Mechanism of
1049   chemoresistance mediated by miR-140 in human osteosarcoma and colon cancer cells.
1050   Oncogene. 19 nov 2009;28(46):4065□74.

1051   20. Grillet F, Bayet E, Villeronce O, Zappia L, Lagerqvist EL, Lunke S, et al. Circulating
1052   tumour cells from patients with colorectal cancer have cancer stem cell hallmarks in ex
1053   vivo culture. Gut. oct 2017;66(10):1802□10.

1054   21. Planque C, Rajabi F, Grillet F, Finetti P, Bertucci F, Gironella M, et al. Pregnane X-
1055   receptor promotes stem cell-mediated colon cancer relapse. Oncotarget. 18 juill
1056   2016;7(35):56558□73.

1057   22. Magill ST, Cambronne XA, Luikart BW, Lioy DT, Leighton BH, Westbrook GL, et al.
1058   microRNA-132 regulates dendritic growth and arborization of newborn neurons in the
1059   adult hippocampus. Proc Natl Acad Sci U S A. 23 nov 2010;107(47):20382□7.

1060   23. David A, Netzer N, Strader MB, Das SR, Chen CY, Gibbs J, et al. RNA binding targets
1061   aminoacyl-tRNA synthetases to translating ribosomes. J Biol Chem. 10 juin
1062   2011;286(23):20688□700.

24. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 1 janv 2013;29(1):15□21.

25. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 1 avr 2014;30(7):923□30.

26. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

27. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological). 1 janv 1995;57(1):289□300.

28. Hunter JD. Matplotlib: A 2D Graphics Environment. Computing in Science &
Engineering. mai 2007;9(3):90□5.

29. Charafe-Jauffret E, Ginestier C, Bertucci F, Cabaud O, Wicinski J, Finetti P, et al.
ALDH1-positive cancer stem cells predict engraftment of primary breast tumors and are
governed by a common stem cell program. Cancer Res. 15 déc 2013;73(24):7290□300.

30. Pinna G, Vandamme M, Rouault C, Charafe-Jauffret E, Ginestier C. Genome-wide RNA
interference screen in cancer stem cells. Methods Cell Biol. 2022;170:81□99.

31. Sakaguchi M, Hisamori S, Oshima N, Sato F, Shimono Y, Sakai Y. miR-137 Regulates
the Tumorigenicity of Colon Cancer Stem Cells through the Inhibition of DCLK1. Mol
Cancer Res. avr 2016;14(4):354□62.

32. Ke SB, Qiu H, Chen JM, Shi W, Chen YS. MicroRNA-202-5p functions as a tumor
suppressor in colorectal carcinoma by directly targeting SMARCC1. Gene. 15 nov
2018;676:329□35.

33. Meerson A, Yehuda H. Leptin and insulin up-regulate miR-4443 to suppress NCOA1 and
TRAF4, and decrease the invasiveness of human colon cancer cells. BMC Cancer. 14 nov
2016;16(1):882.

34. Todaro M, Gaggianesi M, Catalano V, Benfante A, Iovino F, Biffoni M, et al. CD44v6 is
a marker of constitutive and reprogrammed cancer stem cells driving colon cancer
metastasis. Cell Stem Cell. 6 mars 2014;14(3):342□56.

35. Strainiene E, Binkis M, Urnikyte S, Stankevicius V, Sasnauskiene A, Kundrotas G, et al.
Microenvironment dependent gene expression signatures in reprogrammed human colon
normal and cancer cell lines. BMC Cancer. 27 févr 2018;18(1):222.

36. Wang L, Jiang Z, Huang D, Duan J, Huang C, Sullivan S, et al. JAK/STAT3 regulated
global gene expression dynamics during late-stage reprogramming process. BMC
Genomics. 6 mars 2018;19(1):183.

37. Győrffy B. Integrated analysis of public datasets for the discovery and validation of
survival-associated genes in solid tumors. Innovation (Camb). 6 mai 2024;5(3):100625.

38.   Chao HM, Wang TW, Chern E, Hsu SH. Regulatory RNAs, microRNA, long-non coding
RNA and circular RNA roles in colorectal cancer stem cells. World J Gastrointest Oncol.
15 avr 2022;14(4):748‑64.

39.   Xu LZ, Li SS, Zhou W, Kang ZJ, Zhang QX, Kamran M, et al. p62/SQSTM1 enhances
breast cancer stem-like properties by stabilizing MYC mRNA. Oncogene. 19 janv
2017;36(3):304‑17.

40.   Nallasamy P, Nimmakayala RK, Karmakar S, Leon F, Seshacharyulu P, Lakshmanan I, et
al. Pancreatic Tumor Microenvironment Factor Promotes Cancer Stemness via SPP1-
CD44 Axis. Gastroenterology. déc 2021;161(6):1998-2013.e7.

41.   Mustafa S, Koran S, AlOmair L. Insights Into the Role of Matrix Metalloproteinases in
Cancer and its Various Therapeutic Aspects: A Review. Front Mol Biosci. 29 sept
2022;9:896099.

42.   Liu Z, Sanders AJ, Liang G, Song E, Jiang WG, Gong C. Hey Factors at the Crossroad of
Tumorigenesis and Clinical Therapeutic Modulation of Hey for Anticancer Treatment.
Mol Cancer Ther. mai 2017;16(5):775‑86.

43.   Sasahara A, Tominaga K, Nishimura T, Yano M, Kiyokawa E, Noguchi M, et al. An
autocrine/paracrine circuit of growth differentiation factor (GDF) 15 has a role for
maintenance of breast cancer stem-like cells. Oncotarget. 11 févr 2017;8(15):24869‑81.

44.   Leung T, Rajendran R, Singh S, Garva R, Krstic-Demonacos M, Demonacos C.
Cytochrome P450 2E1 (CYP2E1) regulates the response to oxidative stress and migration
of breast cancer cells. Breast Cancer Res. 8 nov 2013;15(6):R107.

45.   Pei X, Zhang SL, Qiu BQ, Zhang PF, Liu TS, Wang Y. Cancer Cell Secreted Legumain
Promotes Gastric Cancer Resistance to Anti-PD-1 Immunotherapy by Enhancing
Macrophage M2 Polarization. Pharmaceuticals (Basel). 16 juill 2024;17(7):951.

46.   Yu J, Shi L, Lin W, Lu B, Zhao Y. UCP2 promotes proliferation and chemoresistance
through regulating the NF-κB/β-catenin axis and mitochondrial ROS in gallbladder
cancer. Biochem Pharmacol. févr 2020;172:113745.

47.   Melo US, Macedo-Souza LI, Figueiredo T, Muotri AR, Gleeson JG, Coux G, et al.
Overexpression of KLC2 due to a homozygous deletion in the non-coding region causes
SPOAN syndrome. Hum Mol Genet. 15 déc 2015;24(24):6877‑85.

48.   Qiao S, Jiang Y, Li N, Zhu X. The kinesin light chain‑2, a target of mRNA stabilizing
protein HuR, inhibits p53 protein phosphorylation to promote radioresistance in NSCLC.
Thorac Cancer. 13 avr 2023;14(16):1440‑50.

49.   Wang M, Zhu X, Sha Z, Li N, Li D, Chen L. High expression of kinesin light chain-2, a
novel target of miR-125b, is associated with poor clinical outcome of elderly non-small-
cell lung cancer patients. Br J Cancer. 3 mars 2015;112(5):874‑82.

50.   Charbe NB, Amnerkar ND, Ramesh B, Tambuwala MM, Bakshi HA, Aljabali AAA, et
al. Small interfering RNA for cancer treatment: overcoming hurdles in delivery. Acta
Pharm Sin B. nov 2020;10(11):2075‑109.

1138

A

hsa-*miR-4745* :   3' CUGGCACUCUGCAGC GGCCCGG U

KLC2 mRNA 3'UTR : 5' ...AGUCCACGGUACUAC CCGGGCC U

Ψ

5' LTR   DsRedx1   Bi-CMV   AcGFP   *miR-4745*   3' SIN LTR

KLC2-3'UTR/locus 1/2/3

B

Transfected cells

FSC (Lin)

DsRed (Log)

Count

GFP (Log)

C

pmiR-locus 2 WT   pmiR-locus 2 WT + *miR-4745*   pmiR-locus 2 Mut   pmiR-locus 2 Mut + *miR-4745*

Count

GFP (Log)

GFP (Log)

D

GFP/DsRed ratio

ns   ns   ns   **   ns   ns   ns

0.45

0.30

0.15

0.00

miR4745   - +   - +   - +   - +   - +   - +   - +

WT   Mut   WT   Mut   WT   Mut

pmiR-sensor   pmiR-locus 1   pmiR-locus 2   pmiR-locus 3

E

GFP/DsRed ratio

ns   **   ns

0.50
0.45
0.40
0.35
0.30
0.15
0.10
0.05
0.00

miR4745   - +   - +   - +

WT   Mut locus 2

pmiR-sensor   pmiR- *KLC2*-3'UTR

A

CPP1

B

C

D

E

CTC44

Supplementary Figure 2: Overexpression of *miR-4745* represses the CSC phenotype in vitro.

(a) Cell proliferation was assessed daily for 4 days using the sulforhodamine B colorimetric assay in patient-derived CPP1 cells co-transfected for 72 hours with either anti-*miR-4745* or anti-miCTRL, along with *miR-4745* or miCTRL mimics. **(b)** CD44v6 mRNA expression in patient-derived CPP1 colon cancer cells and the circulating tumor cell line (CTC44), transfected with *miR-4745* or a negative control mimic (miCTRL). Data are presented as mean ± SEM (n=3). Results are shown as fold change relative to cells transfected with the negative control mimic. **(c)** Percentage of sphere-forming cells in patient-derived CPP1 and SW620 colon cancer cells co-transfected for 72 hours with either anti-*miR-4745* or anti-miCTRL, along with *miR-4745* or miCTRL mimics. Individual replicate values (n=15 per experiment) are plotted along with the mean ± SEM of three independent experiments. **(d)** Percentage of surviving CSCs (Aldefluor-positive cells) 48 hours after exposure to the indicated concentrations of FIRI (1X = 50μM 5-FU + 500nM SN38) in vitro. Sorted CTC44 Aldefluor-positive cells were transfected with 50nM *miR-4745-3p* (miR-4745) or miRVana microRNA Mimic Negative Control #1 (miCTRL) mimics, 24hours before FIRI treatment. Data are presented as mean ± SEM (n = 3).

(e) Percentage of Aldefluor-positive cells (% ALDH$^+$ cells) in CTC44 cells transfected with miR-4745 or miCTRL mimics and then exposed to chemotherapy (FIRI = 5μM 5-FU + 50nM SN38) for 72 hours. Data are presented as mean ± SEM (n = 6). *, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.001$; ns = not significant, Mann-Whitney test.

TCTAGAtgctgaaggggcagccagtcaccagagcgcccacctggcacacccc
ctcaccccagccctgcgcatgggcctgctgcttgtcccgcctgtctctcccac
agcccctgtcttttctgttcaatctcagggtaaccttctcccttgtcatctca
gcctgagccctggaggctgggcctgcccactccagctccatcccttatttatt
ccttccagcagggccctcttccctaggtt**ggggcc**agcaggaggtgccggctg
gagtctccaccatagactcagtggcctggcctccccagaccccagagccaaga
acactaagcactcgccggcccttcggcaccctcgccctccctcccgactcaac
ccggccgttgcttctgtatatagagaaataagttattggccgcgcgcctccct
tcagtccacggtactacc**ggggcc**tcccctcgtccctcttctagtggtaccgc
ccaggccttaatcacccccattccgtgcggtggtatctcccaggctctacatt
ctcgggagcggcgcctcccaagggggtcctgggaccttctcgcgctcctcctg
gcctctgagggatgcgtcctacccgcgccatcgccccgtggcccaggacgggg
acctccccttagtccgtcctcccaccgc**ggggcc**ctgccccgcatcccggcct
tatgcactgcccctcccacccggccccgcccaggcacggccgaccccgccccg
ggcaccgcccaccgagccatcctgcctcgcctccccccacgcctgcagcttct
cgcgaggggcggcgacggtcccctggtggcaggaggggctccccctgttgcgg
gtgaggcggctgctctcatattttcagatgttgctgtagaaataaagacggtt
taaatctga**TCTAGA**

28
29 **Supplementary Figure 4 :** *KLC2* **3'UTR sequence**

30 *KLC2* 3'UTR sequence with XbaI sites (highlighted at the ends) cloned into the pmiR-sensor

31 plasmid. The three predicted binding sites for *miR-4745* are highlighted in red.

A

B

C

D

Supplementary Figure 5: *KLC2* **inactivation inhibits colorectal CSC phenotype.**

(a) mRNA expression levels of *KLC2*, *HEY2*, *CCND3*, and *miR31HG* in CPP1 and SW620 cells transfected for 72 hours with 50nM control siRNA (siCTRL) or specific siRNA targeting *KLC2* (siKLC2), *HEY2* (siHEY2), *CCND3* (siCCND3), or *miR31HG* (simiR31HG). Data are presented as mean ± SEM (n=3) and are expressed as fold change relative to siCTRL-transfected cells. **(b)** Representative Western blot analysis (left panel) and quantification (right panel) of KLC2 protein levels in patient-derived CPP1 and SW620 colon cancer cells transfected for 72 hours with 50nM control siRNA (siCTRL) or siRNA targeting KLC2 (siKLC2). Data are presented as mean +- SEM (n=4). **(c)** Cell proliferation was assessed daily for 4 days using the sulforhodamine B colorimetric assay in patient-derived CPP1 cells transfected for 72 hours with 50nM control siRNA (siCTRL) or specific siRNAs targeting *KLC2* (siKLC2), *HEY2* (siHEY2), *CCND3* (siCCND3), or *miR31HG* (simiR31HG). Data are presented as mean +/- SEM (n=3). ***$p < 0.001$; Mann-Whitney test. **(d)** Western blot analysis of KLC2 protein levels (both endogenous KLC2 and KLC2-3x Flag) in patient-derived CPP1 and SW620 colon cancer cells transfected with a KLC2-overexpressing plasmid (oe-KLC2) or an empty vector (oe-CTRL). NT refers to non-transfected cells.

Supplementary Figure 6: *miR-4745-3p* mimic induces dose-dependent changes in expression levels and sphere-forming capacity in CRC cell lines.

(a) Relative expression levels of *miR-4745-3p* in SW620 colon cancer cells and in the circulating tumor cell line CTC45, transfected for 72 hours with varying concentrations of *miR-4745-3p* or control (miCTRL) mimics. Results are expressed as fold change relative to miCTRL-transfected cells and presented as mean ± SEM (n = 3). **(b)** Sphere-forming efficiency of SW620 and CTC45 cells transfected for 72 hours with varying concentrations of miR-4745-3p or miCTRL mimics. Data represent the percentage of sphere-forming cells (mean ± SEM; n = 3 independent experiments, 10 replicates per experiment). Statistical significance was assessed using one-way ANOVA: *, $p < 0.05$, **, $p < 0.005$, ***, $p < 0.001$, ns = not significant.

Supplementary Table S1: *KLC2* 3'-UTR regions cloned into the pmiR-sensor plasmid. The underlined region indicates the predicted binding sequence for the seed region of *miR-4745*.

Clones	Sequence (5′- 3′)
Clone 1	CCAGCAAGGCCCTCTTCCCTAGGTT**CGGGCC**AGCAGGAGGTGCCGGCTGGAGTCTCCCAGCAAGGCCCTCTTCCCTAGGTT**CGGGCC**AGCAGGAGGTGCCGGCTGGAGTCTC
Clone 1 Mutant	CCAGCAAGGCCCTCTTCCCTAGGTT**CAAACC**AGCAGGAGGTGCCGGCTGGAGTCTCCCAGCAAGGCCCTCTTCCCTAGGTT**CAAACC**AGCAGGAGGTGCCGGCTGGAGTCTC
Clone 2	CCTCCCTTCAGTCCACGGTACTACC**CGGGCC**TCCCCTCGTCCCTCTTCTAGTGGTACCTCCCTTCAGTCCACGGTACTACC**CGGGCC**TCCCCTCGTCCCTCTTCTAGTGGTA
Clone 2 Mutant	CCTCCCTTCAGTCCACGGTACTACC**CAAACC**TCCCCTCGTCCCTCTTCTAGTGGTACCTCCCTTCAGTCCACGGTACTA

	CC**CAAACC**TCCCCTCGTCCCTCTTCTAGTGGTA
Clone 3	TCCCCTTAGTCCGTCCTCCCACCGC**CGGGCC**GTGCCCCG CATCCCGGCCTTATGCATCCCCTTAGTCCGTCCTCCCACC GC**CGGGCCG**TGCCCCGCATCCCGGCCTTATGCA
Clone 3 Mutant	TCCCCTTAGTCCGTCCTCCCACCGC**CAAACC**GTGCCCCG CATCCCGGCCTTATGCATCCCCTTAGTCCGTCCTCCCACC GC**CAAACC**GTGCCCCGCATCCCGGCCTTATGCA

88 **Supplementary Table S2: List of siRNA duplexes targeting *KLC2*, *HEY2*, *CCND3*, and**

89 *miR31HG.*

siRNA	Sequences
si-KLC2	5' GGAGGUGGAAUAUUACUAUTT 3' 3' AUAGUAAUAUUCCACCUCCTT 5'
si-HEY2	5' CGGGAUCGGAUAAAUAACATT 3' 3' UGUUAUUUAUCCGAUCCCGTT 5'
si-MiR31HG	5' CCCUCCUUCUAGAAUAUAUTT 3' 3' AUAUAUUCUAGAAGGAGGGTT 5'
si-CCND3	5' CUGCUGUGAUUGCACAUGATT 3' 3' UCAUGUGCAAUCACAGCAGTT 5'

92 **Supplementary Table S3: Primer sequences.**

Genes	Primer Sequences (5'-3')
KLC2	Forward: AGGGGATGTGTCTGGTCAG Reverse: CCAGGTTGTTTAGTGTCGCAG
HEY2	Forward: TGTTTGGTTCAAGGCAGCTC Reverse: TGATGCACTGCTGGATGGTA
MiR31HG	Forward: CGCTTCTGTCCTCCTACTCG Reverse: ACAAGCAGACCCTTGGAATG
CCND3	Forward: TACCCGCCATCCATGATCG

		Reverse: AGGCAGTCCACTTCAGTGC
	ALDH1A1	Forward: TCCTGGTTATGGGCCTACAG Reverse: CAACAGCATTGTCCAAGTCG
	CD44v6	Forward: GGAACAGTGGTTTGGCAACAG Reverse: TTGGGTGTTTGGCGATATCC
	β-ACTIN	Forward: AGCACGGCATCGTCACCAACT Reverse: TGGCTGGGGTGTTGAAGGTCT
	GAPDH	Forward: AATTGAGCCCGCAGCCTCCC Reverse: CCAGGCGCCCAATACGACCA
	RPLO	Forward: TCGACAATGGCAGCATCTAC Reverse: GCCTTGACCTTTTCAGCAAG
	RPL13	Forward: CATAGGAAGCTGGGAGCAAG Reverse: GCCCTCCAATCAGTCTTCTG
	18S	Forward: GTAACCCGTTGAACCCCATT Reverse: CCATCCAATCGGTAGTAGCG

Supplementary Table S4: List of miRNAs identified from human miRNome-wide screening using a miRNA overexpression (OE) library that deregulate ALDEFLUOR activity in the SW620 cell line model (Excel file).

Supplementary Table S5: List of transcripts that are downregulated or upregulated in the transcriptome analysis following *miR-4745* overexpression in CPP1 cells.

Supplementary Table S6: List of transcripts that are downregulated or upregulated in the translatome analysis, specifically in the highly translated fractions (heavy fractions with more than 3 ribosomes per mRNA).