CellWHISPER disentangles direct cell–cell communication from structural proximity

Anurendra Kumar[1], Felix Rivera[2], Bhavay Aggarwal[2], Nicholas Zhang[2], Ahmet Coskun[2], Saurabh Sinha[2,3*]

1 College of Computing, Georgia Institute of Technology, Atlanta, GA, 30332, USA
2 The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology, Atlanta, GA, 30332, USA
3 H. Milton Stewart School of Industrial & Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30318, USA

* Corresponding Authors
Corresponding Authors-
1. Saurabh Sinha, Email- saurabh.sinha@bme.gatech.edu

15   *Abstract*

16
Direct, contact-dependent cell–cell communication shapes tissue physiology and disease. Spatial
transcriptomics offers a unique opportunity to infer such communication mechanisms, but the few
existing tools for such inference are prone to high error rates due to spatial structure in distribution
of cell types and gene expression. We present CellWHISPER, a statistical framework for inferring
contact-mediated cell–cell communication, including gap junctions and ligand-receptor
mechanisms, from single-cell resolution spatial data. CellWHISPER achieves strict error control
via a specialized hypothesis-testing procedure that accounts for the confounding effects of cell-
type-specific expression and spatial organization. Its use of an exact closed-form significance
score obviates expensive permutation testing and enables scalable analysis of large tissues an
extensive signaling compendia. To uncover salient communication patterns, CellWHISPER also
includes a latent variable model that distills recurrent patterns into maps of mutual preferences
among cell types and signaling genes. Applied to mouse brain STEREO-seq and Xenium
datasets, CellWHISPER identifies cell type-specific gap-junction coupling and yields a
comprehensive connexin-interaction map. We experimentally confirmed Connexin 43-mediated
interfaces among astrocytes, microglia and endothelial cells, which were among the highest-
ranked predictions; in contrast, existing approaches fail to prioritize these confirmed interactions.
Differential analysis of an Alzheimer's disease model versus wild-type identifies preserved
astroglial coupling but increased microglia-associated gap-junction communication. In summary,
CellWHISPER disentangles direct cell–cell communication from structural proximity, enabling
statistically robust and computationally scalable inference from spatial transcriptomics, leading to
the discovery of tissue- and disease-associated contact-mediated signaling programs.
38
39

INTRODUCTION

Cell signaling plays a pivotal role in driving cellular heterogeneity, differentiation, and

pathogenesis within tissues. Rapid proliferation of single-cell transcriptomics technologies has led

to several computational tools for discovering cell-cell communication (CCC) mechanisms[1-4].

These tools, such as CellChat[5], CellPhoneDB[6] and NicheNet[7], typically reveal communication

between specific cell types via curated ligand-receptor (LR) pairs. Broadly, CCC can be

categorized into paracrine, autocrine, juxtacrine, and endocrine signaling[1]. Existing CCC methods

focus on paracrine and autocrine signaling mediated by LR interactions, whereas direct juxtacrine

signaling, particularly via gap junction communication (GJC), remains largely unexplored. Rapidly

emerging spatial transcriptomics (ST) technologies including Stereoseq, VisiumHD, Nanostring

and Xenium, provide a transformative opportunity to bridge this gap by combining gene

expression with spatial proximity, enabling investigation of direct (contact-mediated)

communication between cells. However, computational inference of direct CCC poses distinct

statistical challenges: spatial organization of cell-types and heterogeneous gene expression act

as strong confounders that can inflate apparent evidence for signaling if not explicitly controlled

(**Fig. 1a**).

Direct cell-cell signaling operates primarily via two mechanisms: GJC[8] and LR interactions. Gap

junctions are intercellular channels (**Supp. Fig. 7**) that enable direct cell-cell transfer of ions,

metabolites, second messengers, and regulatory molecules[9]. GJC-mediated inter-cellular

networks play vital roles in normal physiology[10-12] and are implicated in diverse pathological

conditions[13-15]. Gap junction channels are formed by connexin proteins, assembled as hexamers

of homotypic or heterotypic connexins. Over 20 connexins have been identified in mammals, and

the existence of a compatibility "code" governing gap junction formation is an important open

problem[16,17]. Similarly, direct LR pathways, such as Notch-Delta[18] and Eph-Ephrin[19] regulate

differentiation, embryogenesis, and immune cell communication[20], motivating the search for

additional instances of direct LR signaling in varying contexts[21].

Several existing tools allow CCC inference from single-cell transcriptomics, but very few of these

(CellChatv2[22], SpaTalk[24], and COMMOT[25]) exploit inter-cellular proximity information available in

ST data, a key requirement for inferring *contact-dependent* communication. We found these ST-

based inference tools to exhibit surprisingly high false positive errors, motivating this study.

Specifically, we performed a randomized control by permuting cell locations within each cell type,

preserving cell-type-specific spatial organization and LR expression while destroying spatial

proximity between ligand- and receptor-expressing cells. In this control, evidence for true direct

CCC should be eliminated. Strikingly, the above-mentioned ST-based CCC inference tools

reported comparable numbers of direct CCC mechanisms in randomized and original datasets,

indicating insufficient statistical calibration for spatially constrained inference. This finding is

consistent with published studies debating the error rates of existing tools [3,26] and suggestions of

inflated false positives[1,4,27]. It motivated us to devise a statistical procedure for CCC inference that

explicitly accounts for gene-expression heterogeneity, spatial organization, and statistical

dependencies induced by shared cellular neighborhoods.

We present CellWHISPER (Workflow for HIgh-precision Spatial Proximity-mediated cEll-cell

inteRactions), a toolkit to infer direct CCC from single-cell ST data. CellWHISPER makes three

core computational contributions: (i) a proximity-constrained statistical test with a confounder-

aware null that improves specificity for direct CCC, (ii) an analytically derived exact null (closed-

form mean/variance of the CCC-measuring statistic) that avoids explicit permutations and enables

scalable inference across large tissues and signaling compendia, and (iii) a latent variable model

that analyzes the full tensor of tested CCC mechanisms to capture its higher-order interpretable

structures without first collapsing the tensor into information-losing matrices (as is currently done).

Applying CellWHISPER to mouse brain datasets from STEREO-seq[28] and Xenium reveals known

and novel GJC, reporting the first comprehensive brain "connexin code"[16,29]. We experimentally

confirmed Connexin 43-mediated interfaces among astrocytes, microglia and endothelial cells,

which were among the highest-ranked predictions; in contrast, existing approaches fail to prioritize

these confirmed interactions. LR analysis recapitulates established CCC programs such as

mossy fiber pathway while differential analyses identify communication mechanisms associated

with Alzheimer's Disease. In summary, CellWHISPER enables rigorous, scalable inference of

direct inter-cellular communication, revealing tissue- and disease-specific signaling programs.

RESULTS

Overview of CellWHISPER

CellWHISPER infers direct cell-cell communication (CCC) involving a "quadruplet" of two cell types and two signaling genes. Given ST data and cell-type labels, it tests whether a cell-type pair shows evidence of direct communication via a signaling gene pair[3] (connexins or LR pairs), returning a list of cell type-signaling gene quadruplets with z-scores quantifying significance (**Fig. 1b,c,** Methods). A latent variable model (LVM) summarizes the resulting quadruplet set into interpretable modules and preference maps, and an optional differential analysis identifies condition-specific mechanisms (Methods, **Fig. 1c**).

At the core is a statistical test for whether cell types CT_i and CT_j communicate via genes SG_k and SG_l. CellWHISPER binarizes SG expression using a quantile-based threshold. Spatial information is represented by a k-nearest neighbors (KNN) graph[30], from which a subgraph ("whisper network") is derived to include edges between neighboring cells of types CT_i and CT_j expressing genes SG_k and SG_l respectively (Fig. 1b). The whisper network edge count (N_{ijkl}) quantifies the evidence for this quadruplet in the tissue sample. To assess significance, CellWHISPER shuffles locations among cells of same type (preserving cell-type expression and spatial distributions), recomputes N_{ijkl} and uses the resulting null to compute a z-score. For scalability, CellWHISPER derives an analytically computed exact null for the within-cell-type randomization scheme, avoiding explicit shuffling (Methods). This exact null utilizes closed-form expressions for the expectation and variance of the whisper network edge count, explicitly accounting for dependence induced by shared cells in the spatial graph. As a result, CellWHISPER computes calibrated z-scores and enables orders-of-magnitude improvements in computational efficiency relative to permutation-based approach (**Fig. 1e,f**). In practice, this enables direct CCC inference on datasets comprising tens of thousands of cells and thousands of signaling gene pairs on standard hardware, without exceeding memory limits or incurring prohibitive runtimes. To assess statistical control for direct CCC, we applied the within-type randomization control described above. On downsampled Stereo-seq data (~5,000 cells, 28 cell types, 367 LR pairs) competing tools (CellChat v2[22], COMMOT[25], SpaTalk[24]) reported similar numbers of significant quadruplets on real and randomized data whereas CellWHISPER reported far fewer quadruplets on randomized data, yielding an empirically estimated FPR <5%, underscoring its high specificity (**Fig. 1d,** Methods).

Figure 1. Overview of CellWHISPER. (**a**) **Cell-type confounding in ligand–receptor communication inference.** When ligand L is expressed broadly across cell type CT1 and receptor R across CT2, spatial proximity between the two cell types produces apparent L–R co-expression at their interface (left; yellow zone) even without specific L–R signaling. When the same number of high-expressing cells are enriched specifically at the CT1–CT2 interface, the co-expression reflects genuine signaling (right; yellow zone). Both scenarios appear identical to existing methods, which report significant communication in both cases. CellWHISPER evaluates co-expression against a null that randomizes cell identities within each cell type, preserving cell-type-specific expression distributions and spatial organization, and correctly distinguishes the two. (**b**) **Statistical testing for a CT-SG quadruplet**: For a given cell type pair (ct_i, ct_j) and signaling gene pair (g_k, g_l), CellWHISPER constructs a "whisper network", where each edge represents a pair of

adjacent cells of types ct_i, ct_j expressing genes g_k, g_l respectively, and the count of such "whisper edges"

serves as the test statistic. Under the null model, cell locations are randomized within each cell type,

generating a background distribution of edge counts, which is used to assess statistical significance. (**c**)

CellWHISPER framework: CellWHISPER takes single-cell resolution ST data as input, along with cell type

(CT) labels and signaling gene pairs, which may be ligand-receptor (LR) gene pairs or gap junction (GJ)

gene pairs. It performs a statistical test (panel b) to compute a z-score for each cell type pair and signaling

gene pair (a CT-SG quadruplet), reporting all quadruplets with significant z-scores. A latent variable model

can then analyze the full complement of quadruplet-level results to infer higher-order communication

patterns such as CT pairs that are similar in their SG usage (CT-CT similarity matrix), SG pairs that mediate

communication between similar cell types (SG-SG similarity matrix), and quantify whether a certain cell

types tends to communicate using a specific SG (CT-SG preference matrix). CellWHISPER can also be

applied across conditions for differential analysis, identifying CT-SG quadruplets specific to one condition

versus the other. (**d**) **Randomized control reveals poor specificity**: Shown is the average number of

communicating cell-type pairs inferred per ligand–receptor by four different methods, on real versus

randomized data (cell identities permuted within each cell type, preserving expression distributions and cell-

type spatial organization while disrupting cross-type proximity). Competing methods show limited

separation between real and randomized data, whereas CellWHISPER yields substantially fewer calls

under randomization, indicating significantly improved specificity for direct CCC. We summarize this as an

empirical FPR, defined as the number of significant CT–SG quadruplets detected on randomized data

divided by the number detected on real data under fixed thresholds (CellWHISPER: <5%). **Benchmarking**

was performed on a downsampled hippocampal subset (~5,000 cells) due to computational

constraints of competing methods (Methods, Supp. Fig. 5). (**e**) **Runtime and scalability.** Wall-clock

time to evaluate all cell-type pairs for a single signaling-gene pair on datasets of increasing size (~5,000;

~10,000; and ~50,000 cells), comparing CellWHISPER with the analytically derived exact null versus

permutation testing (n = 500), and three competing tools (CellChat, COMMOT, SpaTalk) run with

recommended settings. CellWHISPER's analytical null achieves ~100× speedup over permutation on

~5,000 cells and substantially larger speedups at ~10,000 and ~50,000 cells, enabling practical screening

on full-resolution tissues. SpaTalk did not complete within the allotted runtime at one or more dataset sizes

(shown as timeouts). (**f**) **Consistency between analytical and permutation-based significance**. Z-

scores computed using the analytical null closely match those from permutation testing across CT–SG

tests, confirming that the analytical formulation reproduces the permutation-based null while avoiding

explicit shuffling. (**g**) **Whisper network specificity for connexins:** For each homotypic connexin gene,

an aggregated whisper network was constructed as the union of whisper networks across all cell-type pairs

in the Stereo-seq mouse brain dataset. Edge counts for connexin networks were significantly higher than

those for networks constructed from randomly selected genes (n = 5000) (Mann–Whitney U test p = 4.6 ×

10e-8). All connexins yielded $\geq$8,878 edges, whereas only 0.4% of random genes reached this threshold.

We focused initial validation on one of the best studied mechanisms of direct CCC: gap junctions

(GJ). Gap junction communication provides a stringent stress test for direct CCC inference

because they are purely contact-mediated and therefore particularly sensitive to spatial and

expression confounders. GJ form when connexin protein in adjacent cells dimerize to establish

intercellular channels[31] (Fig. 1a). To evaluate the plausibility of using mRNA levels as proxies for

connexin-mediated communication, we performed targeted spatial profiling in an induced

pluripotent stem cell model[12] and observed concordance between *Gja1* mRNA and Cx43 protein

abundance, as well as substantial overlap between mRNA- and protein-derived whisper networks

(**Supp. Figure 1, Methods**). In a Stereo-seq mouse brain dataset[28] (Methods), aggregated

whisper networks for connexin pairs were significantly larger than those from random genes (**Fig.**

1g, Mann-Whitney U test p-value 4.6e-8, Methods): each of 10 connexins yielded a whisper

network of $\geq$8878 edges, while only 22/5000 (0.4%) random genes met this criterion (**Supp.**

Figure 2).

CellWHISPER identifies known and novel mechanisms of gap junction communication

(GJC) in the mouse brain

To demonstrate the utility of CellWHISPER, we first applied it to discover GJC in mouse brain.

From a computational perspective, GJC provides a stringent benchmark for direct CCC inference

because it is purely contact-mediated and therefore maximally sensitive to spatial and expression

confounders. GJC is believed to be widespread in the brain[32] and has been implicated in

neurodevelopment and diseases [33-35]. Certain combinations of cell types and connexins are

preferentially involved in brain GJC [31,36] and we sought to explore such a "connexin code"[16,29]. We

analyzed a STEREO-seq mouse hemibrain dataset[28], comprising of ~50,000 cells across 28 cell

types (CTs), using ten highly variable connexins as signaling genes (SG). (All 100 pairs of these

genes were included in the SG compendium). CellWHISPER identified 1726 significant CT-SG

quadruplets (z-score > 3, whisper network size > 30, **Fig. 2a**, **Supp. Data 1**) at an empirical FPR

<1% estimated from randomized controls (**Supp. Figure 3**). This demonstrates that well-

controlled inference over large quadruplet spaces is feasible at tissue scale (Fig 1e). Predictions

were dominated by inter-glial GJC mechanisms (**Fig. 2b**).

Figure 2. CellWHISPER Identifies Direct Gap Junction Communication in the Mouse Brain. (a) Gap
junction communication identified in mouse brain: Sankey plot of the top 150 significant CT-SG
quadruplets (a cell type pair communicating via a gap junction gene pair) ranked by whisper network size
identified in Stereo-seq data (~50K cells). For visual ease, "Astr" represents all astrocyte subtypes (Astr1-
5), "IN" represents all interneuron subtypes and "EX L2-6" represents all excitatory neuron layers. ("Olig" =
oligodendrocyte, "Micro" = microglia.) Lines indicate CellWHISPER-predicted communication involving
specific cell type pairs (or aggregated groups such as Astr) and specific gap junction genes: red for glial-
glial communication, blue for neuronal-neuronal communication, and pink for neuroglial communication.
Line thickness reflects the magnitude of communication for each CT–SG pair, while node height reflects
the total communication associated with that cell type or signaling gene. (**b**) Pie chart of the three major
classes of gap junction communication in the top 150 predicted CT-SG quadruplets shown in (a). (**c**)
Oligodendrocyte self-communication. Whisper network of oligodendrocytes communicating via
homotypic Cx32 gap junctions. Blue edges represent whisper network connections; red shading denotes
kernel density smoothing at edge centers for visual clarity. (**d**) **Statistical significance of whisper edges.**
Histogram of edge counts from randomized whisper networks (cell locations permuted within cell type), with

fitted normal distribution (blue line). The red dashed line marks the observed number of edges in the whisper
network shown in (c) which is significantly higher than expected under the null. Y-axis shows the probability
density of randomized edge counts. (**e**) **Heterotypic astroglial communication.** Whisper network of
astrocytes predicted to communicate with oligodendrocytes via heterotypic gap junction formed by Cx43
(astrocytic side) and Cx47 (oligodendrocytic side). (**f**) Cell types predicted to communicate using homotypic
Cx36 gap junctions. Blue entries in heatmap denote significant cell type pairs (z score>3, whisper edge
count > 30) found by CellWHISPER. Neuronal communication is predicted more frequently compared to
glial communication. (**g**) Glial cell types predicted to communicate using Cx43-Cx43 gap junctions. Only
glial communication is shown since it is predicted more frequently compared to neuronal communication
(Supp. Fig. 6) (**h–j**) Astrocytic communication enriched in hippocampus. Whisper networks showing (h)
astrocyte–astrocyte, (i) astrocyte–endothelial, and (j) astrocyte–microglia interactions mediated by
homotypic Cx43 (GJA1) gap junctions

We examined the highest-confidence mechanisms from two complementary perspectives.
Ranking quadruplets by whisper network size revealed that the top six represent inter-
oligodendrocyte GJC via combinations of Cx47, Cx32, and Cx29 (**Supp. Data 1, Supp. Data 2**)
consistent with established roles of Cx47 and Cx32 in oligodendrocytic coupling[37,38]. **Fig. 2c**
illustrates homotypic Cx32 whisper network between oligodendrocytes, comprising 4,283 edges,
far exceeding chance expectation (**Fig. 2d**, z-score = 13.09). The top predictions also included
Cx29-mediated oligodendrocytic GJC. Although Cx29 transcripts are enriched in
oligodendrocytes[39], prior studies indicate that it primarily localizes to the adaxonal membrane
rather than forming functional gap junctions[40], highlighting a known limitation of transcript-based
inference when subcellular protein localization is critical. The next highest-ranking quadruplets
involve astrocyte-astrocyte GJC via Cx43, Cx30, and Cx26, connexins known to couple
astrocytes into an astrocytic syncytium essential for homeostasis and adult neurogenesis[17,41-44].
We also identified high-confidence astrocyte-oligodendrocyte (A/O) communication via Cx43 on
the astrocyte side and Cx47, Cx32 or Cx29 on the oligodendrocytic side. **Fig. 2e** illustrates *Astr3-*
oligodendrocyte communication via Cx43-Cx47, consistent with prior reports supporting
functional A/O GJ coupling[45], though there is some unresolved debate about their functionality[44].

Examining CellWHISPER results ranked by z-score rather than network size (**Supp. Data 3**)
highlights additional mechanisms supported by fewer communicating cell pairs, which may thus
be less well-studied in prior work. Among these, we identified GJC between excitatory neurons
and other cell types, including thalamic excitatory neurons, oligodendrocytes, and astrocytes
(**Supp. Data 3**). Maubecin et al.[46] provided direct evidence of functional neuro-glial coupling using

heterotypic gap junctions in rodents. Gap junctions between excitatory neurons are known to be

primarily mediated by Cx36 (Gjd2)[47,48] and extensively studied in rodent models[49-52]. As shown in

Fig. 2f, excitatory neurons are a major site of homotypic Cx36-mediated communication predicted

by CellWHISPER.

CellWHISPER identifies homotypic Cx43 gap junctions as a mechanism underlying

astrocytic syncytium

Cx43 is the most studied connexin, with key roles in differentiation and homeostatic regulation[53-

55]. We examined the complete map of cell type pairs predicted by CellWHISPER to participate in

Cx43-mediated GJC (**Fig. 2g**), finding a predominance of glial over neuronal communication. This

analysis provides a focused validation setting, as Cx43-mediated interactions are well

characterized. Astrocyte subpopulation Astr1 is predicted to communicate with other astrocytes,

as well as endothelial, microglia, oligodendrocytes and ependymal cells. This aligns with known

biology: Cx43 mediates astrocyte-oligodendrocyte-ependymal cell interactions forming an

isopotential syncytium [54,56-58]. Predicted astroglial interactions exhibited a higher network density

in the hippocampus (**Figs. 2h-j**). Inter-astrocyte GJC (z-score 18.13, **Fig. 2h**) in the hippocampus

has been documented to be Cx43-mediated. Astrocyte-endothelial GJC (z-score 10.94, **Fig. 2i**)

has been implicated in maintaining blood–brain barrier integrity[59-61]. Additionally, CellWHISPER

predicted Cx43-mediated communication between microglia and astrocytes (**Fig. 2j**) as well other

microglia. Indeed, recent studies suggest that microglia upregulate Cx43 under pathological or

inflammatory conditions and form functional gap junctions – with each other and with astrocytes[62-

64].

Figure 3. Spatially resolved analysis of Cx43-mediated cellular crosstalk across brain regions. (**a**)

Schematic of gap junction–mediated interactions between astrocytes (GFAP), microglia (IBA1), and

endothelial cells (CD31) via Cx43 (created with BioRender.com). (**b**) Immunofluorescence of coronal mouse

brain section stained for Cx43, GFAP, IBA1, and CD31. Hippocampal boundaries were defined using the

Allen Mouse Brain Atlas. Regions of interest (ROI) within hippocampus (Hippo1, Hippo2) and cortex were

outlined for higher-resolution imaging. Scale bar, 100 μm. (**c**) Single-cell–resolution images of hippocampal

ROIs showing colocalization of astrocytes and microglia with Cx43. (**d**) Quantification of Cx43 colocalization

across hippocampus and cortex. We selected 10 regions (478 × 262 pixels each) within and outside the

hippocampus and quantified Cx43 colocalization with astrocytes, endothelial cells, and microglia.

Colocalization was defined as instances where Cx43 puncta were positioned between or overlapping the

indicated cell types, yielding 30 values (one per cell type per region) and a total of 333 interactions. Two-

way ANOVA revealed a significant enrichment of astrocyte–Cx43 interactions in the hippocampus

compared to the cortex (p = 0.0002).

To validate these computationally inferred predictions, we performed immunofluorescent staining

on mouse coronal brain sections for Cx43 together with markers of astrocytes (GFAP), endothelial

cells (CD31), and microglia (IBA1) (**Fig. 3a, b**; Methods). All channels showed robust signal

across brain regions (**Fig. 3b**). Distinct Cx43 co-localization was evident at astrocyte–astrocyte,

astrocyte–endothelial, astrocyte–microglial, and microglial–microglial interfaces (Fig 3b,c).

Quantitative analysis showed higher co-localization of Cx43 with cell-type markers in the

hippocampus compared to cortex (**Fig. 3d**), with astrocytic contacts enriched >5-fold.

Representative examples of colocalization of astrocytes, microglia and Cx43 in the hippocampus

are shown in **Fig. 3c**. Notably, these experimentally supported Cx43 interfaces (Astrocyte–

Astrocyte, Astrocyte–Endothelium, and Astrocyte–Microglia) are prioritized among the highest-

ranked CellWHISPER predictions, enabling targeted experimental follow-up. By contrast,

CellChat v2 (the only ST-based tool that scales to this data set) assigns extremal significance

scores (p-value = 0) to a large set of candidate cell-type pairs (~150 pairs), including these

interfaces, limiting its usefulness for prioritizing specific mechanisms. Together, these

observations support the ability of CellWHISPER to recover known, region-specific contact-

mediated communication patterns using a rigorous confounder-aware framework.

CellWHISPER identifies direct ligand-receptor (LR) communication

Direct LR communication plays a fundamental role in organizing local tissue architecture[21]. We

next used LR interactions as a test of generality of the CellWHISPER analytic framework. Unlike

gap junctions, LR signaling spans diverse functional classes (adhesion, guidance, immune and

ECM), while still requiring spatial proximity for direct interactions. To identify such interactions in

the mouse brain, we re-analyzed STEREO-seq data using 367 curated LR pairs from the CellChat

database[5] (Methods). CellWHISPER identified 4743 significant CT–SG quadruplets (z-score > 4,

 whisper network size > 30) (**Fig. 4a, Supp. Data 4**), with an empirically estimated FPR < 1%,

spanning inter-glial, neuron-glia and neuron-neuron communication (**Fig. 4b**).

Figure 4. CellWHISPER Identifies direct Ligand-Receptor communication in the mouse brain. (a)
Ligand–receptor communication mechanisms. Sankey plot showing the top 25 significant CT–SG
quadruplets **ranked by z-score** (z-score > 4, whisper network size > 30) identified using 367 curated
ligand–receptor pairs. Randomized control analysis estimated a false positive rate (FPR) < 1% for this
compendium. **(b)** Communication classes. Distribution of major interaction types, including neuron–neuron,
neuron–glia, and inter-glial communication. **(c–e)** Six representative whisper networks (two per panel,
left/right). Selected ligand–receptor–mediated cell–cell interactions illustrating key modes of direct
communication: (c) excitatory CA3 neurons (EX CA) and dentate gyrus granule neurons (GN DG)
communicating via Efnb3-Epha4 (right) and tight junction Ocln (left); (d) excitatory neuron-oligodendrocyte
interactions involving *Cldn* (right) and semaplexin family member (left); and (e) meningeal- astrocyte
communication via extracellular matrix components (*Lama2*, *Col4a1–6*, *Adgrg6*).

The highest confidence predictions revealed communication between CA3 excitatory neurons
(EX CA) and dentate gyrus granule neurons (GN DG) mediated by LR pairs of the mossy fiber
pathway[65] (**Supp. Data 5**). These include multiple ephrin/Eph family members, C1ql2/3-Adgrb3
and homotypic interactions of occludin, protocadherin and claudin (**Supp. Data 5, Fig. 4c**).
Ephrin–Eph and C1ql2/3–Adgrb3 (BAI3) interactions, both implicated in mossy fiber–CA3
synapse organization[66-68], ranked among the highest-confidence predictions, alongside multiple

adhesion-related pairs (including claudins/occludin). These findings are consistent with known

hippocampal wiring programs[69] and support the biological plausibility of CellWHISPER's

proximity-constrained LR inference.

CellWHISPER also predicted LR-mediated communication between oligodendrocytes and

excitatory neurons, involving homotypic MAG, claudin and semaplexin interactions (**Supp. Data**

6, Fig. 4d), consistent with known roles for these mechanisms in axon–glia recognition, myelin

stability[70], and activity-dependent modulation of myelination[71]. Additionally, we observed

interactions between astrocytes and meningeal cells, notably involving extracellular matrix

components (ECM) such as Lama2, type IV collagens (Col4a1–6), Adgrg6 (**Supp. Data 7, Fig.**

4e). These examples illustrate that CellWHISPER supports discovery across multiple LR

functional categories while maintaining stringent specificity for direct CCC.

Latent Variable Modeling (LVM) infers general patterns of communication mechanisms

We used CellWHISPER's "LVM" to infer higher-order patterns among significant quadruplets

(Methods). Unlike approaches that first collapse quadruplet-level information into CT x SG or CT

x CT matrices[5,6], an aggregation step that can be information-losing; CellWHISPER directly

models quadruplets to characterize communication profiles of cell types and signaling genes.

Specifically, the LVM models the presence or absence of statistically supported CT–SG

quadruplets using a shared low-dimensional embedding space for cell types and signaling genes

(Methods). Under this model, a quadruplet is more likely when each cell type exhibits higher

affinity for its corresponding signaling gene, yielding embedding geometry that captures CCC

preferences. The learned embeddings yield (1) CT–SG preferences, (2) CT–CT similarity, and

(3) SG–SG similarity. Model parameters are estimated via constrained log-likelihood

maximization. **Fig. 5b** shows that the optimization can accurately infer cell type similarity on

simulated data (Methods).

Applying LVM to GJ results in mouse brain Stereo-seq data revealed five dominant cell type

modules in CT-CT similarity map (**Fig. 5a, Supp. Data 8**) whereas a baseline method performing

naïve aggregation of quadruplet tensor did not reveal modular structure (**Supp. Figure 4**). LVM

also identified four GJ modules in SG-SG similarity map (**Fig. 5d**, Supp. Data 8). Cell type

modules showed preferences for specific GJ modules in the CT-SG preference matrix (**Fig. 5c**).

For example, excitatory neuron subtypes in cell type modules M1/M2 (EX Mb, EX L2/3, EX CA)

prefer GJ module M4 consisting of Gjd2 (Fig. 5c,d). Notably, DA neurons (excitatory or inhibitory

Figure 5. LVM infers module of signaling cell types and signaling genes. (**a**) Cell type similarity matrix. Cosine similarity of latent cell-type embeddings derived from the LVM routine of CellWHISPER, applied to gap junction communication in the Stereo-seq mouse brain dataset. Five dominant modules of cell types emerge, reflecting recurrent patterns of gap junction communication across the tissue. (**b**) Optimization. Constraint optimization within the latent variable model (Methods) identifies underlying cell type pair similarity, resolving higher-order structure in the data. (**c**) Cell type–gap junction preferences. Heatmap showing associations between cell type modules and gap junction gene modules. Distinct modules exhibit preferential coupling, e.g., Module M1 and M2 are enriched in excitatory neurons and prefer Gjd2 (GJ module M4). DA neurons and IN S+ prefer GJ module M2. **(d)** Cosine similarity of latent embeddings across gap junction genes, revealing four dominant connexin modules. (**e**) Cell-type similarity matrix for ligand–receptor communication. Cosine similarity of latent cell-type embeddings derived from CellWHISPER applied to ligand–receptor interactions (Stereo-seq mouse brain dataset). Four dominant cell-type modules emerge. Excitatory neurons in cortical layers and hippocampal CA region cluster together (M2), whereas excitatory neurons in thalamus is in module M4. (**f**) Communication overlap between excitatory neurons. Quantification of overlap in ligand–receptor communication patterns (shared CT–ligand–receptor triplets) shows that excitatory neurons in cortical layers 2/3 are more similar to CA excitatory neurons than to thalamic excitatory neurons, consistent with known laminar and neuromodulatory properties of neocortical and hippocampal neurons. (g–h) Cell type–ligand and cell type–receptor module specificity. Sankey plots showing preferential associations between cell-type modules and ligand modules (g), and between cell-type modules and receptor modules (h). Links were derived by averaging CT–SG preference scores within each module. (**i**) Ligand similarity map. Cosine similarity of ligand embeddings reveals five dominant modules (Supp. Data 9). Module M1 consists mainly of genes related to cell adhesion (e.g., PECAM1, CDH1/5, ICAM1), extracellular matrix components (COL4A1–6, LAMA2), and immune modulators (H2-Aa, H2-Eb1, H2-Ab1, CD52, CD55, CD86, PTPRC). Another ligand module (M5, Supp.

Data 9) is enriched for axon guidance and synaptic signaling genes , including protocadherins, ephrins,

neurexins, semaphorins, Lrrc and Flrt family (**j**) Receptor similarity map. Cosine similarity of receptor

embeddings identifies five receptor modules(Supp. Data 9).

depending on context[72]), clustered with inhibitory Sst+ interneurons[73] in cell type module M3 (**Fig.**

5a, Supp. Data 8) and shared preference for GJ module M2 (**Fig. 5c**), consistent with dye-coupling

evidence of DA–interneuron electrical synapses[74] and suggesting candidate GJ mediating these

synapses.

Next, we applied LVM to LR interactions (Fig. 4), identifying four dominant cell-type modules (**Fig.**

5e, Supp. Data 9). Excitatory neurons in cortical layers and CA region co-cluster, whereas

thalamic excitatory neurons lie in different module (**Fig. 5e,f**). This aligns with known features of

excitatory neurons in the neocortex and hippocampus (of which CA is part), including laminar

structure[75,76] and shared neuromodulatory response[77], different from excitatory neurons in

thalamic neurons[78]. Analysis of ligand similarity map (**Fig. 5i**) identified five dominant modules, of

which one (M1) is enriched for cell adhesion, ECM and immune modulators (**Supp. Data 9**). This

ligand module preferentially associates with cell type module M1 (**Fig. 5g**), enriched for glial cell

types located near the brain periphery, such as ependymal cells, meninges, and smooth muscle

cells (Fig. 5e), suggesting a specialized glial–meningeal signaling axis potentially involved in

barrier maintenance and neuroimmune communication. Another ligand module (M5, Supp. Data

9) is enriched for axon guidance and synaptic signaling genes[79,80], and prefers cell type module

M2 (**Fig. 5g**), enriched in excitatory neurons in the cortical layers and hippocampus, consistent

with their known role in neuronal cell-cell communication[81]. Similarly, we found five dominant

modules in receptor similarity map (**Fig. 5j**, Supp Data 9) having preference for specific cell type

modules (**Fig. 5h**). Together, LVM captures dominant communication patterns across GJC and

LR signaling.

Differential analysis reveals Alzheimer's disease-associated GJC mechanisms

Gap junction-mediated communication (GJC) supports central nervous system homeostasis and

is implicated in neurodegenerative diseases, including Alzheimer's disease (AD)[31,82,83]. Under

normal conditions, GJC supports metabolic and synaptic function, especially through astrocytic

networks[63,84] whereas in AD, aberrant gap junctions and hemichannel activity contributes to

neuroinflammation and neurodegeneration[31,82,83] [31]. From a computational perspective, this

analysis tests whether CellWHISPER can identify condition-associated rewiring using the same

statistically well-controlled, proximity-constrained quadruplet framework, while limiting false
positives arising from differences in cell-type composition, spatial organization, or expression. To
systematically interrogate these dynamics, we used CellWHISPER to decode GJC networks
altered in AD.

Figure 6. Differential Analysis infers conserved and condition-specific gap junction communication in Alzheimer's disease (AD). (**a**) Scatter plot of 60 high-confidence CT–SG quadruplets (z-score > 4, whisper network size > 30 in at least one condition) showing their z-scores in wild-type (WT) and TgCRND8 (AD) Xenium datasets for Gja1, Gjb2, and Gjc3. Quadruplets are classified as shared, condition-specific, or indeterminate. (**b**) Bar plot showing the frequency of each cell type across the 60 quadruplets. Astrocytes and oligodendrocytes are most enriched in shared interactions, microglia dominate AD-specific interactions, and astrocytes are more frequent in WT-specific interactions. (**c**) Whisper networks of Astr1–Olig1 communication via homotypic Cx43 in WT (left) and AD (right), showing preserved symmetry between conditions. (**d**) Selected condition-specific quadruplets, showing z-scores in WT versus AD. Top three WT-specific and AD-specific interactions were chosen based on whisper network size. (**e**) Whisper network of Astr1–Ex th1 communication via homotypic Cx43 in WT. (**f**) Whisper network of Smc1–Smc1 (smooth muscle cell self-communication) via homotypic Cx43 in AD, highlighting potential vascular remodeling. (**g**) Aggregated microglial whisper network in AD, constructed by taking the union of all AD-specific quadruplets involving microglia, revealing enrichment in thalamus and hypothalamus regions.

We analyzed Xenium mouse brain data from wild-type and TgCRND8 mice[85] (aged >13 months,

Methods) using three annotated GJ genes. After cross-sample integration to harmonize cell-type

labels (Methods), CellWHISPER's Differential Analysis (DA) function quantified quadruplets per

condition, and classified them as shared, condition-specific or indeterminate (Methods). This

revealed 60 high-confidence quadruplets (z-score > 4, whisper network size > 30, **Fig. 6a, Supp.**

Data 10). The twenty shared quadruplets are enriched for astrocytic (13/20) and oligodendrocytic

communication (9/20) (**Fig. 6b**) including Astr1-Olig1 homotypic Cx43 (**Fig. 6c**). Despite known

Cx43 upregulation near Aβ plaques, prior work suggests structural preservation of astrocytic GJC

in AD brain[86]; consistent with this, CellWHISPER identifies conserved GJC mechanisms despite

change in signaling gene expression.

DA further identified 15 TgCRND8-specific and 14 wild-type-specific quadruplets (Fig. 6a,d,

Supp. 10), including reduced astrocyte-thalamic excitatory neuron GJC via homotypic GJA1 (**Fig.**

6e). The corresponding whisper network (Fig. 6e) shows enrichment of this communication in the

WT thalamus. In contrast, TgCRND8-specific mechanisms included increased smooth muscle

cells homotypic GJA1 (Fig. 6d,f), potentially reflecting vascular remodeling[87,88]. TgCRND8-

specific mechanisms were enriched for microglial interactions (5/17 vs. 0/23 in WT-specific, p-

value < 0.01, two-tailed proportions test), and WT-specific mechanisms to be marginally enriched

for astrocytes (6/23 vs. 2/17 in TgCRND8) (**Fig. 6b**). These trends align with existing literature:

astrocyte networks maintain homeostasis in healthy brain[89], while microglial GJC becomes more

prominent during AD neuroinflammation[31,90,91]. Aggregated microglial whisper network (**Fig. 6g**)

showed enrichment in thalamus and hypothalamus. Connexin alterations have been reported in

thalamic regions in AD[82] while microglial changes in thalamus are linked to cognitive impairment

after cortical brain injury[92].

DISCUSSION

Direct cell–cell communication shapes tissue homeostasis and disease, yet existing spatial

transcriptomics tools conflate genuine signaling with cell type co-localization, limiting their utility

for rigorous inference. CellWHISPER addresses this by explicitly accounting for confounding

effects of gene-expression and spatial organization, CellWHISPER markedly reduces FPR

relative to competing tools. Two additional computational features broaden practicality and

interpretability: an analytical exact null that avoids explicit permutation testing and enables

scalable inference across large tissues and signaling compendia, and a quadruplet-native latent

variable model that extracts higher-order structure without collapsing interactions into information-
losing pairwise summaries. A differential analysis module further extends the same testing
framework to multi-condition settings to identify shared versus condition-associated mechanisms.
The framework is agnostic to the signaling compendium (connexins, curated LR pairs, or user-
defined contact genes) and scales to tens of thousands of cells, enabling routine application to
emerging high-resolution spatial platforms.

We demonstrated CellWHISPER on mouse brain spatial datasets as validation settings spanning
two major classes of direct CCC: gap junctions and contact-dependent ligand–receptor
interactions. CellWHISPER recovered canonical glial networks and generated a large-scale map
of predicted connexin-mediated interactions ("connexin code"). Immunostaining validated Cx43
interactions, demonstrating its colocalization at astrocyte–astrocyte/endothelial/microglial
interfaces. LR analysis uncovered adhesion/synaptic programs, including the mossy fiber
pathway and a glial–meningeal signaling axis. Differential analysis of an AD mouse model
identified a shift toward microglial-associated GJC and region-specific remodeling in the thalamus
and vasculature, consistent with emerging models of neuroinflammatory progression.

Several limitations motivate future work. CellWHISPER inference relies on mRNA as a proxy for
membrane proteins, and post-transcriptional regulation or subcellular localization can yield false
positives . It prioritizes specificity through a confounder-aware null model, which may come at the
cost of sensitivity. In contrast, CellWHISPER excels at detecting communication events mediated
by a smaller number of cell pairs. Extensions to lower-resolution platforms (e.g. Visium) may be
possible via deconvolution approaches, and integrating pathway information could improve
robustness of whisper networks. Finally, segmentation errors in ST[93] may propagate to CCC
inference, a limitation shared across current methods.

In summary, CellWHISPER establishes a robust paradigm for inferring direct signaling from ST
data, combining stringent statistical control with interpretable latent structure, to enable
mechanistic dissection of tissue organization and for identifying disease-specific communication
in development, homeostasis, and disease.

527
528
529
530 **METHODS**

Overview of CellWHISPER

CellWHISPER infers direct cell–cell communication (CCC) from spatial transcriptomics (ST) data. The input consists of gene expression profiles, spatial coordinates, and cell-type annotations for individual cells, together with a precompiled compendium of signaling gene (SG) pairs, such as connexins or ligand–receptor (LR) pairs. SG compendia can be obtained from existing resources or provided by the user as a custom list.

For each candidate combination of a cell-type pair (ct_i, ct_j) and a signaling gene pair (sg_k, sg_l), CellWHISPER tests whether the two cell types exhibit evidence of direct communication mediated by the signaling genes (Fig. 1b). Statistically supported interactions are reported as CT–SG quadruplets, each associated with a z-score quantifying significance (Fig. 1c). CellWHISPER includes three analysis modules (Fig. 1c). (i) A statistical testing module identifies significant CT–SG quadruplets. (ii) A latent variable model (LVM) summarizes the collection of significant quadruplets into interpretable preference and similarity maps, including CT–CT, CT–SG, and SG–SG relationships.

(iii) An optional differential analysis (DA) module identifies condition-specific or shared communication mechanisms across biological conditions.

Whisper network construction

Gene expression was binarized using a quantile-based threshold (75th percentile by default), yielding binary indicators of whether a signaling gene is expressed in a given cell.

Spatial proximity between cells was represented using a k-nearest neighbors (KNN) graph constructed from cell coordinates (k = 5 unless otherwise specified). This graph defines the set of potential physical contacts used for CCC inference. For datasets with irregular tissue geometry, distance-based thresholds should be used instead of fixed neighbor counts.

For each CT–SG quadruplet (ct_i, ct_j, sg_k, sg_l), a whisper network was derived by retaining only edges connecting neighboring cells of types ct_i and ct_j that express sg_k and sg_l, respectively (Fig. 1b). The number of edges in this subgraph, denoted N_{ijkl}, quantifies spatially supported evidence for the corresponding communication mechanism.

Statistical test for CT–SG quadruplets

To assess statistical significance, CellWHISPER constructs a null model by permuting cell identities within each cell type. This preserves cell-type–specific gene expression distributions and spatial organization while disrupting spatial proximity between signaling cells of different

types. The whisper network is recomputed under the null, yielding a null distribution for N_{ijkl}, from

which a z-score is computed.

To enable scalability to large datasets, CellWHISPER derives an analytical approximation to the

null distribution, avoiding explicit permutations while preserving control of confounding effects

arising from expression heterogeneity and spatial cell-type structure.

Analytical approach for Statistical test of a quadruplet:

Let Z_u be a Boolean random variable indicating the presence of a whisper edge for a proximal

cell pair corresponding to a quadruplet i, j, k, l. Our objective is to estimate the expectation and

variance of the total number of such whisper edges: $E(\sum_u Z_u), Var(\sum_u Z_u)$.The probability that a

specific edge u is present (i.e. $Z_u = 1$) is modeled as the product of probabilities that gene k is

expressed in cell of type i and gene l is expressed in a cell of type j. Gene expression is

determined by a global percentile threshold across all cells. Thus:

$$P(Z_u = 1) = p_i p_j$$

Where p_i is the probability that a cell of type i expresses gene k, and p_j is the probability that a

cell of type j expresses gene l.

If Z_u are Independent (i.i.d.):

Assuming the Z_u are i.i.d. Bernoulli variables, the total follows a Binomial distribution with N_u

WHISPER edges:

$$E\left(\sum_u Z_u\right) = \sum_u P(Z_u) = N_u P(Z_u = 1) = N_u p_i p_j$$

$$Var\left(\sum_u Z_u\right) = N_u P(Z_u = 1)\left(1 - P(Z_u = 1)\right) = N_u p_i p_j (1 - p_i p_j)$$

Accounting for Dependence Between Edges:

However, the Z_u variables are not independent because edges may share a common cell. While

the expectation remains unchanged due to linearity, the variance must account for dependencies:

$$Var\left(\sum_u Z_u\right) = E\left(\left(\sum_u Z_u\right)^2\right) - \left(E\left(\sum_u Z_u\right)\right)^2$$

593
$$= E\left(\left(\sum_u Z_u\right)^2\right) - N_u^2 p_i^2 p_j^2$$

594 To expand the first term,

595
$$E\left(\left(\sum_u Z_u\right)^2\right) = E\left(\sum_u Z_u^2\right) - E\left(\sum_{u,v} Z_u Z_v\right)$$

Since, $Z_u \in [\{0,1\}]$, $Z_u^2 = Z_u$, so the first sum reduces to $E(\sum_u Z_u) = N_u p_i p_j$. The second term

depends on the structure of edge overlap and has the following cases (assuming $i \neq j$):

• Case1: No shared cells between u and v , therefore Z_u, Z_v are independent, $E(Z_u Z_v) =$

$p_i^2 p_j^2$

• Case 2: Shared cell of type j but distinct cells of type i, therefore, $E(Z_u Z_v) = p_i p_j^2$

• Case 3: Shared cell of type i but distinct cells of type j, therefore $E(Z_u Z_v) = p_i^2 p_j$

If $i = j$, cases 2 and 3 collapse into a single case due to symmetry.

Stereoseq Data and Analysis

Mouse brain STEREO-seq data[28] were obtained from the Spatial Omics Database[94]. After quality

control filtering, the dataset comprised 50,140 cells and 18,527 genes, including 10 gap junction

(connexin) genes: *Gjd2* (Cx36), *Gja5* (Cx40), *Gjc1* (Cx45), *Gja4* (Cx37), *Gjb2* (Cx26), *Gja1*

(Cx43), *Gjb6* (Cx30), *Gjb1* (Cx32), *Gjc2* (Cx47), and *Gjc3* (Cx29). We used the authors' annotated

cell-type labels, encompassing 28 distinct cell types. Abbreviations include: CA3, cornu ammonis

area 3; Mb, midbrain; EX, excitatory glutamatergic neuron; IN, GABAergic interneuron; DA,

dopaminergic neuron; GN DG, granule cell of dentate gyrus; Astr, astrocyte; Micro, microglia;

OPC, oligodendrocyte precursor cell; Oligo, oligodendrocyte; SMC, smooth muscle cell; Ery,

erythrocyte; and Endo, endothelial cell.

Gene expression matrices were normalized and subsequently denoised using MAGIC imputation

(t=2) to recover continuous expression profiles. To construct the *whisper network*, a gene was

considered "expressed" in a given cell if its expression exceeded the 75th percentile of that gene's

expression across all cells. For gap-junction analysis, we evaluated all possible connexin–

connexin combinations. Although several connexins are known to preferentially form specific

homo- or heterotypic channels, experimental evidence for many pairings remains incomplete or

context-dependent; thus, interaction patterns were inferred empirically from the data.

Ligand–receptor database

We used the CellChat v2[22] ligand–receptor compendium, focusing specifically on pairs annotated as cell–cell contact interactions. After intersecting this database with the gene set retained from the STEREO-seq dataset, 367 ligand–receptor pairs were included for downstream analyses.

Randomization and False Positive Rate estimation:

We performed randomization by permuting cell IDs within each cell type. This procedure preserves the marginal spatial distribution and gene expression profiles specific to each cell type. False positive rate (FPR) is obtained by comparing the number of detected pairs obtained on randomized data with number of detected pairs on real data.

Benchmarking

We could not run Spatalk on full mouse hemibrain dataset containing 50K cells. Therefore, we selected part of hippocampal region (Supp. Fig. 5) containing 5K cells for benchmarking analysis in Fig. 1d. We randomized by permuting cell IDs within each cell type.

All comparison methods were downloaded from their official repositories and executed using recommended parameters. For CellChat v2, we used the computeCommunProb function with default parameters (type = "triMean", trim = 0.1, interaction.range = 250, contact.knn.k = 5), followed by filterCommunication with a minimum of 20 cells per cluster. For COMMOT, we applied commot.tl.cluster_communication to compute cell–cell communication for each ligand–receptor pair. For SpaTalk, we used the default parameters.

Connexin mRNA-protein concordance

CellWHISPER relies on cellular mRNA abundance as a proxy for membrane-bound connexin proteins. To evaluate this assumption, we performed targeted spatial profiling of connexin 43 (Cx43) protein and its corresponding mRNA (Gja1) at single-molecule resolution in an induced pluripotent stem cell (iPSC) model, where gap junction communication is critical for spatial patterning.

We observed a significant correlation between Gja1 mRNA expression and average cytosolic Cx43 protein intensity at the single-cell level (Spearman's R = 0.47, $p = 5 \times 10^{-20}$; Supplementary Fig. 1). Consistently, whisper networks inferred from Gja1 mRNA showed strong overlap with those inferred from Cx43 protein measurements (Fig. 1e, Supp. Fig. 1), supporting the use of mRNA as a proxy for connexin-mediated communication in this context.

Latent Variable Model (LVM):

LVM stipulates that the binary variable $I(ct_i, ct_j, g_k, g_l)$ indicating whether the CT-SG quadruplet ct_i, ct_j, g_k, g_l has statistical support in the data follows a Bernoulli distribution:

$$I(ct_i, ct_j, g_k, g_l) \sim Ber\ (p_{ijkl})$$

where u_i, u_j denote to-be-learnt low-dimensional representations (embeddings) for CTs ct_i, ct_j and v_k, v_l denote embeddings for SGs g_k, g_l, respectively, all embeddings being in a common latent space. The dot product $u_i. v_k$ captures the general propensity of ct_i for participating in CCC using g_k. Thus, the model posits that the quadruplet ct_i, ct_j, g_k, g_l is more likely if there is a higher tendency for ct_i communicating via g_k and ct_j communicating via g_l.

Using this formulation CellWHISPER learn embeddings that enable quantification of (1) cell type-signaling gene preferences ($u_i. v_k$), (2) cell type similarities ($u_i. u_j$), i.e., which pairs of CTs have similar communication mechanisms, and (3) signaling gene similarities ($v_k. v_l$), i.e., which SGs are used by the same cell types. It uses cosine similarity rather than dot products between embeddings to emphasize specificity and mitigate abundance bias inherent in dot product.

In the null model, p_{ijkl} is assumed to be constant for all indices i, j, k, l. In contrast, the alternate model factorizes p_{ijkl} into two terms:

- the propensity of cell type i to communicate via signaling gene k, and
- the propensity of cell type j to communicate via signaling gene l

Formally, this is expressed as:

$$p_{ijkl} = (u_i. v_k)(u_j. v_l)$$

where u_i and u_j are low dimensional embeddings for cell types i, j, and v_k, v_l are embeddings for signaling genes k and l. All embeddings lie in a shared latent space, and we enforce the constraints:

$$0 < u_{ir} < 1, \qquad 0 < v_{ir} < 1, \qquad ||u_{ir}||_1 < 1, ||v_{kr}||_1 < 1$$

691 This formulation is invariant under simultaneous permutation of both CT and SG pairs, i.e.

692
$$I\big(ct_i, ct_j, g_k, g_l\big) = I\big(ct_j, ct_i, g_l, g_k\big) \Rightarrow p_{ijkl} = p_{jilk}$$

We optimize the model using constrained log-likelihood maximization, implemented via
`scipy.optimize` using the SLQSP method. The constraint is posed using the L1 norm of the
embeddings that ensures that the problem is a linear optimization and probabilities $0 < p_{ijkl} < 1$.

To confirm identifiability and recovery, we simulated Gap Junction Communication results with
the same orders as the STEREO-seq analysis (29 cell types, 10 connexins, d=3). Ground-truth
embeddings were drawn from numpy.random.rand with unit norm cell type and signaling gene
embeddings. The optimizer recovered CT–CT similarity, SG–SG similarity, and CT–SG
preference structure with high agreement (Fig. 5b).

For ligand–receptor analysis, only a subset of ligand–receptor pairs are biologically plausible. To
ensure model learnability, we explicitly set the probability $p_{ijkl} = 0$ for incompatible ligand–
receptor combinations. For computational feasibility, we randomly subsample a small number of
such incompatible pairs (2000 in our experiments) to include as negative examples during
optimization. This approach enables efficient training while maintaining biological specificity in the
learned embeddings.

Differential Analysis:
We performed Differential Analysis (DA) at the terminal time point at which amyloid pathology is
well established. Wild-type (WT) and TgCRND8 (AD) Xenium datasets were jointly integrated
using residualPCA to achieve conservative alignment across conditions while preserving
biological differences. A joint embedding was constructed and cell-type clustering was performed
on the integrated space yielding 25 clusters.

Label transfer from reference. Cell-type labels were transferred from the annotated STEREO-seq
reference (analyzed above). For this step, the WT, AD, and STEREO-seq datasets were co-
embedded with reciprocal PCA[95]. For each query cell (WT or AD), we identified its k = 7 nearest
neighbors in the reference and recorded their labels. For each cluster in the joint WT+AD

embedding, neighbor votes were aggregated across all member cells and normalized by the total
number of votes in that cluster to yield a label-probability vector. The cluster label was assigned
as the cell type with the maximum normalized vote. Clusters containing multiple subpopulations
of the same broad class (e.g., oligodendrocytes) were further subdivided and relabeled as
subtypes (e.g., Olig-1, Olig-2).

DA quantification. Using these harmonized annotations, CellWHISPER quantified CT–SG
quadruplets separately in WT and AD datasets. Quadruplets were retained if they satisfied
significance thresholds (z-score > 4 and whisper-network size > 30). This procedure yielded 60
high-confidence quadruplets across the two conditions. A quadruplet was classified as shared if
both WT and AD satisfied these thresholds, condition-specific if one condition satisfied the
thresholds while the other did not (z-score < 3 or whisper-network size < 20), and indeterminate
otherwise.

Experimental details for GJA1 mRNA-protein correlation
iPSC culture
iPSCs were obtained and cultured according to Culture and Freezing Methods for WTC Derived
AICS hiPSC Lines: https://www.coriell.org/0/PDF/Allen/iPSC/AICS_SOP_WTC_CellCulture.pdf.
Coverslips (22 × 22 mm) were placed flat in sterile 6-well plates and coated with growth factor–
reduced, phenol-red–free Matrigel (Corning, cat. no. 356231) at a final protein concentration of
0.337 mg/mL. Briefly, Matrigel aliquots thawed at 4 °C were diluted on ice in cold DMEM/F12 to
1.5 mL per well, mixed thoroughly with pre-chilled pipettes to ensure homogeneity, and
immediately dispensed to fully cover each coverslip. Plates were incubated at room temperature
for 1 h, after which excess coating solution was aspirated and wells were gently rinsed with room-
temperature mTeSR1 medium supplemented with 1 μM Y-27632 (ROCK inhibitor) before cell
seeding. Coated coverslips were used for experiments within 6 h or, if needed, sealed with
Parafilm and stored at 4 °C for up to 2 weeks. Once cells were confluent, they were fixed in 4%
formaldehyde for staining.

iPSC RNA and protein multiplexing
Fixed cells on coverslip with permeabilized with 70% EtOH for 30 mins at -20 C. Cycle 1 RNA
was labeled using Molecular Instruments hybridized chain reaction RNA fluorescence in situ
hybridzation (HCR RNA FISH) v3.0 probes: Human Cx43 B3 555, Human MALAT1 B2 647. All
probes were amplified 75 mins at room temperature. Fluorescence images were acquired on

adapted Nikon Ti 2000-U microscope with appropriate filters. Then, samples were blocked and
incubated with Cx43 Ab (Invitrogen 71-0700, 1;150) overnight at 4 C followed by secondary
antibody staining (anti-rabbit 488) for 1 hr at room temperature. This Cycle 2 Cx43 Ab was imaged
again. DAPI counterstain was used to register the images from both cycles. Images were
segmented with Cellpose and quantified for downstream analysis.
763
Immunofluorescence staining
Mouse brain sections were fixed in 4% paraformaldehyde (PFA) and sectioned using Slides were
deparaffinized and rehydrated by subsequent rinses in Xylene, 100% EtOH, 95% EtOH, 80%
EtOH, 70% EtOH and deionized water (diH$_2$O). Then we performed antigen retrieval by placing
slides in a Coplin jar with 10% sodium citrate buffer in 1x phosphate-buffered saline (PBS). Coplin
jars were placed in a pressure cooker with 300 mL of tap water and set on high pressure for 15
minutes. The slides were let to cool for 20 minutes and then washed with diH$_2$O. They were then
blocked using 50 mg/mL bovine serum albumin (BSA) in 1x PBS. They were then washed with
.1% Tween 20 in 1x PBS (PBS-T). Primary antibodies were diluted based on manufacturers'
recommendation in 5% BSA and left to incubate on slides at 4°C, overnight. A list of antibodies
and their dilutions are below. Slides were washed the next day with PBS-T. For unconjugated
primary antibodies, slides were washed with three times with PBS-T for 2 minutes and incubated
in 1:5000 <u>secondary antibody</u> in PBS for 1 hour at room temperature. Slides were washed were
washed three times with PBS-T and counterstained by a 1:500 Dapi in 1x PBS solution for 10
minutes. Slides were washed with PBS-T for 2 minutes and nuclear staining was done using
1:2000 DAPI (Thermo Scientific, Ref:62248). Slides were mounted using 10% glycerol in 1x PBS.
Coverslips were placed over top of the slides, and the edges were coated with nail polish to
prevent dehydration. Slides were imaged using the Keyence BZ-X800 fluorescence microscope.
782
Image preprocessing and quantification of colocalization
Images of cells in tissue all underwent the following preprocessing before further analysis. Each
image contains multiple channels. The maximum intensity projection of each channel was then
generated. The projected image was the final image used for cellular counting analysis. For the
DAPI channel, the maximum intensity projection was generated from the raw image. For
visualization consistency, brightness and contrast of each fluorescent channel were adjusted
uniformly across all images using the "Brightness/Contrast" tool without altering the underlying
pixel intensity values. To measure intercellular colocalization, the straight-line tool was used to
draw a line between the centers of identified cells of interest (e.g., nuclei or marker-specific

regions). The length of each line was recorded using the "Measure" function, which reports distances in calibrated units (µm) based on the microscope's pixel-to-micrometer conversion. At least 629 cells per field were analyzed, and distances were exported for downstream statistical analysis. Previous steps were conducted in ImageJ/Fiji (National Institutes of Health). Statistical significance was calculated using a regular two-way ANOVA in *BioRender Graph*. Statistical levels are provided as follows: ns for $0.05 < p \leq 1$, * for $0.01 < p \leq 0.05$, ** for $0.001 < p \leq 0.01$, *** for $0.0001 < p < 0.001$, and **** for $p \leq 0.0001$.

Code Availability

The code is available at https://github.com/anurendra/CellWHISPER.

Data Availability

Mouse brain STEREO-seq data[28] were obtained from the Spatial Omics Database[94]. Our iPSC-derived and mouse brain immunostaining data will be made available upon publication. Xenium *TgCRND8* Alzheimer's-disease mouse-model data were obtained from wild-type and *TgCRND8* mice[85], as provided by 10x Genomics. Only the final time point (age > 13 months) was used for analysis.

REFERENCES

1. Armingol, E., Officer, A., Harismendy, O. & Lewis, N.E. Deciphering cell–cell interactions and communication from gene expression. *Nature Reviews Genetics* **22**, 71-88 (2021).

2. Wilk, A.J., Shalek, A.K., Holmes, S. & Blish, C.A. Comparative analysis of cell–cell communication at single-cell resolution. *Nature biotechnology* **42**, 470-483 (2024).

3. Dimitrov, D. *et al.* LIANA+ provides an all-in-one framework for cell–cell communication inference. *Nature Cell Biology* **26**, 1613-1622 (2024).

4. Dimitrov, D. *et al.* Comparison of methods and resources for cell-cell communication inference from single-cell RNA-Seq data. *Nature communications* **13**, 3224 (2022).

5. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature communications* **12**, 1088 (2021).

6. Efremova, M., Vento-Tormo, M., Teichmann, S.A. & Vento-Tormo, R. CellPhoneDB: inferring cell–cell communication from combined expression of multi-subunit ligand–receptor complexes. *Nature protocols* **15**, 1484-1506 (2020).

7. Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication by linking ligands to target genes. *Nature methods* **17**, 159-162 (2020).

8. Hervé, J.-C. & Derangeon, M. Gap-junction-mediated cell-to-cell communication. *Cell and tissue research* **352**, 21-31 (2013).

9. Goodenough, D.A. & Paul, D.L. Gap junctions. *Cold Spring Harbor perspectives in biology* **1**, a002576 (2009).

10. Battefeld, A., Klooster, J. & Kole, M.H. Myelinating satellite oligodendrocytes are integrated in a glial syncytium constraining neuronal high-frequency activity. *Nature Communications* **7**, 1-13 (2016).

11. Rash, J.E. Molecular disruptions of the panglial syncytium block potassium siphoning and axonal saltatory conduction: pertinence to neuromyelitis optica and other demyelinating diseases of the central nervous system. *Neuroscience* **168**, 982-1008 (2010).

12. Glen, C.M., McDevitt, T.C. & Kemp, M.L. Dynamic intercellular transport modulates the spatial patterning of differentiation during early neural commitment. *Nature communications* **9**, 4111 (2018).

13. Harris, A. & Locke, D. *Connexins: a guide*, (Springer Science & Business Media, 2008).

14. Nakase, T. & Naus, C.C. Gap junctions and neurological disorders of the central nervous system. *Biochimica Et Biophysica Acta (BBA)-Biomembranes* **1662**, 149-158 (2004).

15. Jongsma, H.J. & Wilders, R. Gap junctions in cardiovascular disease. *Circulation research* **86**, 1193-1197 (2000).

16. Koval, M., Molina, S.A. & Burt, J.M. Mix and match: investigating heteromeric and heterotypic gap junction channels in model systems and native tissues. *FEBS letters* **588**, 1193-1204 (2014).

17. Stephan, J., Eitelmann, S. & Zhou, M. Approaches to study gap junctional coupling. *Frontiers in Cellular Neuroscience* **15**, 640406 (2021).

18. Zhou, B. *et al.* Notch signaling pathway: architecture, disease, and therapeutics. *Signal transduction and targeted therapy* **7**, 95 (2022).

19. Kania, A. & Klein, R. Mechanisms of ephrin–Eph signalling in development, physiology and disease. *Nature reviews Molecular cell biology* **17**, 240-256 (2016).

20. Nakandakari-Higa, S. *et al.* Universal recording of immune cell interactions in vivo. *Nature* **627**, 399-406 (2024).

21. Mattes, B. & Scholpp, S. Emerging role of contact-mediated cell communication in tissue development and diseases. *Histochemistry and cell biology* **150**, 431-442 (2018).

22. Jin, S., Plikus, M.V. & Nie, Q. CellChat for systematic analysis of cell–cell
communication from single-cell transcriptomics. *Nature Protocols* **20**, 180-219 (2025).
23. Troulé, K. *et al.* CellPhoneDB v5: inferring cell–cell communication from single-cell
multiomics data. *Nature Protocols*, 1-29 (2025).
24. Shao, X. *et al.* Knowledge-graph-based cell-cell communication inference for spatially
resolved transcriptomic data with SpaTalk. *Nature communications* **13**, 4429 (2022).
25. Cang, Z. *et al.* Screening cell–cell communication in spatial transcriptomics via collective
optimal transport. *Nature methods* **20**, 218-228 (2023).
26. Cesaro, G. *et al.* Advances and challenges in cell–cell communication inference: a
comprehensive review of tools, resources, and future directions. *Briefings in*
Bioinformatics **26**, bbaf280 (2025).
27. Zohora, F.T. *et al.* NEST: Spatially-mapped cell-cell communication patterns using a
deep learning-based attention mechanism. *bioRxiv*, 2024.03. 19.585796 (2024).
28. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA
nanoball-patterned arrays. *Cell* **185**, 1777-1792. e21 (2022).
29. Evans, W.H. & Martin, P.E. Gap junctions: structure and function. *Molecular membrane*
biology **19**, 121-136 (2002).
30. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nature methods*
19, 171-178 (2022).
31. Takeuchi, H. & Suzumura, A. Gap junctions and hemichannels composed of connexins:
potential therapeutic targets for neurodegenerative diseases. *Frontiers in cellular*
neuroscience **8**, 189 (2014).
32. Söhl, G., Maxeiner, S. & Willecke, K. Expression and functions of neuronal gap
junctions. *Nature reviews neuroscience* **6**, 191-200 (2005).
33. Dere, E. & Zlomuzica, A. The role of gap junctions in the brain in health and disease.
Neuroscience & Biobehavioral Reviews **36**, 206-217 (2012).
34. Dong, A., Liu, S. & Li, Y. Gap junctions in the nervous system: probing functional
connections using new imaging approaches. *Frontiers in Cellular Neuroscience* **12**, 320
(2018).
35. Kajiwara, Y. *et al.* GJA1 (connexin43) is a key regulator of Alzheimer's disease
pathogenesis. *Acta neuropathologica communications* **6**, 1-20 (2018).
36. Sánchez, O.F. *et al.* Role of connexins 30, 36, and 43 in brain tumors,
neurodegenerative diseases, and neuroprotection. *Cells* **9**, 846 (2020).
37. Nualart-Marti, A., Solsona, C. & Fields, R.D. Gap junction communication in myelinating
glia. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **1828**, 69-78 (2013).
38. Papaneophytou, C., Georgiou, E. & Kleopa, K.A. The role of oligodendrocyte gap
junctions in neuroinflammation. *Channels* **13**, 247-263 (2019).
39. Nagy, J.I., Ionescu, A.V., Lynn, B.D. & Rash, J.E. Connexin29 and connexin32 at
oligodendrocyte and astrocyte gap junctions and in myelin of the mouse central nervous
system. *Journal of Comparative Neurology* **464**, 356-370 (2003).
40. Kleopa, K.A., Orthmann, J.L., Enriquez, A., Paul, D.L. & Scherer, S.S. Unique
distributions of the gap junction proteins connexin29, connexin32, and connexin47 in
oligodendrocytes. *Glia* **47**, 346-357 (2004).
41. Orthmann-Murphy, J.L., Abrams, C.K. & Scherer, S.S. Gap junctions couple astrocytes
and oligodendrocytes. *Journal of Molecular Neuroscience* **35**, 101-116 (2008).
42. Xing, L., Yang, T., Cui, S. & Chen, G. Connexin hemichannels in astrocytes: role in CNS
disorders. *Frontiers in molecular neuroscience* **12**, 23 (2019).
43. Li, M. *et al.* Astrocytic Gap Junctions protein Cx43/Cx30 modulate EAAT1 and glutamate
to mediate cerebral ischemia–reperfusion injury. *Brain Research* **1847**, 149306 (2025).

44.   Nagy, J., Ionescu, A.V., Lynn, B. & Rash, J. Coupling of astrocyte connexins Cx26,
926   Cx30, Cx43 to oligodendrocyte Cx29, Cx32, Cx47: Implications from normal and
927   connexin32 knockout mice. *Glia* **44**, 205-218 (2003).
928   45.   Orthmann-Murphy, J.L., Freidin, M., Fischer, E., Scherer, S.S. & Abrams, C.K. Two
929   distinct heterotypic channels mediate gap junction coupling between astrocyte and
930   oligodendrocyte connexins. *Journal of Neuroscience* **27**, 13949-13957 (2007).
931   46.   Alvarez-Maubecin, V., García-Hernández, F., Williams, J.T. & Van Bockstaele, E.J.
932   Functional coupling between neurons and glia. *Journal of Neuroscience* **20**, 4091-4098
933   (2000).
934   47.   Srinivas, M. *et al.* Functional properties of channels formed by the neuronal gap junction
935   protein connexin36. *Journal of Neuroscience* **19**, 9848-9855 (1999).
936   48.   Lee, S.-N. *et al.* Cryo-EM structures of human Cx36/GJD2 neuronal gap junction
937   channel. *Nature communications* **14**, 1347 (2023).
938   49.   Nagy, J. & Rash, J. Cx36, Cx43 and Cx45 in mouse and rat cerebellar cortex: Species-
939   specific expression, compensation in Cx36 null mice and co-localization in neurons vs.
940   glia. *European Journal of Neuroscience* **46**, 1790-1804 (2017).
941   50.   Feigenspan, A. *et al.* Expression of connexin36 in cone pedicles and OFF-cone bipolar
942   cells of the mouse retina. *Journal of Neuroscience* **24**, 3325-3334 (2004).
943   51.   Van Bockstaele, E., Garcia-Hernandez, F., Fox, K., Alvarez, V. & Williams, J. Expression
944   of connexins during development and following manipulation of afferent input in the rat
945   locus coeruleus. *Neurochemistry international* **45**, 421-428 (2004).
946   52.   Kay, C.W., Ursu, D., Sher, E. & King, A.E. The role of Cx36 and Cx43 in 4-
947   aminopyridine-induced rhythmic activity in the spinal nociceptive dorsal horn: an
948   electrophysiological study in vitro. *Physiological Reports* **4**, e12852 (2016).
949   53.   Zhou, M., Zhong, S. & Verkhratsky, A. Astrocyte syncytium: from neonatal genesis to
950   aging degeneration. *Neural regeneration research* **19**, 395-396 (2024).
951   54.   Li, Z.A. & Zhou, M. Astrocyte syncytium–a biopower grid system in the brain. *Journal of*
952   *integrative neuroscience* **23**, 71 (2024).
953   55.   Ma, B. *et al.* Gap junction coupling confers isopotentiality on astrocyte syncytium. *Glia*
954   **64**, 214-226 (2016).
955   56.   Serra, R. & Simard, J.M. Adherens, tight, and gap junctions in ependymal cells: A
956   systematic review of their contribution to CSF-brain barrier. *Frontiers in neurology* **14**,
957   1092205 (2023).
958   57.   Fabbiani, G. *et al.* Connexin signaling is involved in the reactivation of a latent stem cell
959   niche after spinal cord injury. *Journal of Neuroscience* **40**, 2246-2258 (2020).
960   58.   Bouillé, C., Mesnil, M., Barriere, H. & Gabrion, J. Gap junctional intercellular
961   communication between cultured ependymal cells, revealed by lucifer yellow CH transfer
962   and freeze-fracture. *Glia* **4**, 25-36 (1991).
963   59.   De Bock, M. *et al.* Targeting gliovascular connexins prevents inflammatory blood-brain
964   barrier leakage and astrogliosis. *Jci Insight* **7**, e135263 (2022).
965   60.   Jiang, H., Zhang, Y., Wang, Z.-Z. & Chen, N.-H. Connexin 43: an interface connecting
966   neuroinflammation to depression. *Molecules* **28**, 1820 (2023).
967   61.   Johnson, A.M. *et al.* Connexin 43 gap junctions contribute to brain endothelial barrier
968   hyperpermeability in familial cerebral cavernous malformations type III by modulating
969   tight junction structure. *The FASEB Journal* **32**, 2615 (2018).
970   62.   Eugenín, E.A. *et al.* Microglia at brain stab wounds express connexin 43 and in vitro
971   form functional gap junctions after treatment with interferon-γ and tumor necrosis factor-
972   α. *Proceedings of the National Academy of Sciences* **98**, 4190-4195 (2001).
973   63.   Nakase, T., Fushiki, S. & Naus, C.C. Astrocytic gap junctions composed of connexin 43
974   reduce apoptotic neuronal damage in cerebral ischemia. *Stroke* **34**, 1987-1993 (2003).

975	64.	Abudara, V. *et al.* Activated microglia impairs neuroglial interaction by opening C x43
976		hemichannels in hippocampal astrocytes. *Glia* **63**, 795-811 (2015).
977	65.	Henze, D.A., Urban, N.N. & Barrionuevo, G. The multifarious hippocampal mossy fiber
978		pathway: a review. *Neuroscience* **98**, 407-427 (2000).
979	66.	Triplett, J.W. & Feldheim, D.A. Eph and ephrin signaling in the formation of topographic
980		maps. in *Seminars in cell & developmental biology* Vol. 23 7-15 (Elsevier, 2012).
981	67.	Yang, J.S., Wei, H.X., Chen, P.P. & Wu, G. Roles of Eph/ephrin bidirectional signaling in
982		central nervous system injury and recovery. *Experimental and therapeutic medicine* **15**,
983		2219-2227 (2018).
984	68.	Martinelli, D.C. *et al.* Expression of C1ql3 in discrete neuronal populations controls
985		efferent synapse numbers and diverse behaviors. *Neuron* **91**, 1034-1051 (2016).
986	69.	Newrzella, D. *et al.* The functional genome of CA1 and CA3 neurons under native
987		conditions and in response to ischemia. *BMC genomics* **8**, 1-18 (2007).
988	70.	Quarles, R.H. Myelin-associated glycoprotein (MAG): past, present and beyond. *Journal*
989		*of neurochemistry* **100**, 1431-1448 (2007).
990	71.	Gow, A. *et al.* CNS myelin and sertoli cell tight junction strands are absent in
991		Osp/claudin-11 null mice. *Cell* **99**, 649-659 (1999).
992	72.	Hutchison, M. *et al.* Genetic inhibition of neurotransmission reveals role of glutamatergic
993		input to dopamine neurons in high-effort behavior. *Molecular psychiatry* **23**, 1213-1225
994		(2018).
995	73.	Song, Y.-H., Yoon, J. & Lee, S.-H. The role of neuropeptide somatostatin in the brain
996		and its application in treating neurological disorders. *Experimental & Molecular Medicine*
997		**53**, 328-338 (2021).
998	74.	Vandecasteele, M., Glowinski, J. & Venance, L. Electrical synapses between
999		dopaminergic neurons of the substantia nigra pars compacta. *Journal of Neuroscience*
1000		**25**, 291-298 (2005).
1001	75.	Radnikow, G. & Feldmeyer, D. Layer-and cell type-specific modulation of excitatory
1002		neuronal activity in the neocortex. *Frontiers in neuroanatomy* **12**, 1 (2018).
1003	76.	Yan, J. & Gao, L. Organizing principles of the prefrontal cortex from a single-neuron
1004		projectome. Vol. 25 408-409 (NATURE PORTFOLIO HEIDELBERGER PLATZ 3,
1005		BERLIN, 14197, GERMANY, 2022).
1006	77.	Gulledge, A.T. & Kawaguchi, Y. Phasic cholinergic signaling in the hippocampus:
1007		functional homology with the neocortex? *Hippocampus* **17**, 327-332 (2007).
1008	78.	Sherman, S.M. Thalamus plays a central role in ongoing cortical functioning. *Nature*
1009		*neuroscience* **19**, 533-541 (2016).
1010	79.	Yuasa-Kawada, J., Kinoshita-Kawada, M., Tsuboi, Y. & Wu, J.Y. Neuronal guidance
1011		genes in health and diseases. *Protein & cell* **14**, 238-261 (2023).
1012	80.	Südhof, T.C. Synaptic neurexin complexes: a molecular code for the logic of neural
1013		circuits. *Cell* **171**, 745-769 (2017).
1014	81.	Lv, X. *et al.* Patterned cPCDH expression regulates the fine organization of the
1015		neocortex. *Nature* **612**, 503-511 (2022).
1016	82.	Angeli, S. *et al.* Altered expression of glial gap junction proteins Cx43, Cx30, and Cx47
1017		in the 5XFAD model of Alzheimer's disease. *Frontiers in Neuroscience* **14**, 582934
1018		(2020).
1019	83.	Maulik, M. *et al.* Amyloid-β regulates gap junction protein connexin 43 trafficking in
1020		cultured primary astrocytes. *Journal of Biological Chemistry* **295**, 15097-15111 (2020).
1021	84.	Pechlivanidou, M. *et al.* Glial gap junction pathology in the spinal cord of the 5xFAD
1022		mouse model of early-onset Alzheimer's disease. *International journal of molecular*
1023		*sciences* **23**, 15597 (2022).
1024	85.	Genomics, x. Xenium In Situ Analysis of Alzheimer's Disease Mouse Model Brain
1025		Coronal Sections from One Hemisphere Over a Time Course. (2024).

86. Yi, C. *et al.* Astroglial connexin43 contributes to neuronal suffering in a mouse model of
Alzheimer's disease. *Cell Death & Differentiation* **23**, 1691-1701 (2016).
87. Kozberg, M.G. *et al.* Blood–brain barrier leakage and perivascular inflammation in
cerebral amyloid angiopathy. *Brain Communications* **4**, fcac245 (2022).
88. Figueroa, X.F. & Duling, B.R. Gap junctions in the control of vascular function.
Antioxidants & redox signaling **11**, 251-266 (2009).
89. Huang, X. *et al.* Astroglial connexins in neurodegenerative diseases. *Frontiers in*
molecular neuroscience **14**, 657514 (2021).
90. Caruso, G., Di Pietro, L. & Caraci, F. Gap junctions and connexins in microglia-related
oxidative stress and neuroinflammation: perspectives for drug discovery. *Biomolecules*
13, 505 (2023).
91. Anwar, M.M. *et al.* Assessing the role of primary healthy microglia and gap junction
blocker in hindering Alzheimer's disease neuroinflammatory type: early approaches for
therapeutic intervention. *Frontiers in Neuroscience* **16**, 1041461 (2023).
92. Matoba, K. *et al.* FcγR-and CD9-dependent synapse engulfing microglia in the thalamus
drives cognitive impairment following cortical brain injury. *bioRxiv*, 2024.09. 19.609743
(2024).
93. Yu, H. *et al.* Segmentation Matters: Cell Segmentation Challenge in Spatial
Transcriptomics. *NEURON-D-25-01846*.
94. Yuan, Z. *et al.* SODB facilitates comprehensive exploration of spatial omics data. *Nature*
Methods **20**, 387-399 (2023).
95. Stuart, T. *et al.* Comprehensive integration of single-cell data. *cell* **177**, 1888-1902. e21
(2019).
1049