CELLWHISPER: INFERENCE OF DIRECT CELL-CELL COMMUNICATION FROM SPATIAL TRANSCRIPTOMICS

Anurendra Kumar[1], Felix Rivera[2], Bhavay Aggarwal[2], Nicholas Zhang[2], Ahmet Coskun[2], Saurabh Sinha[2,3]*

1 College of Computing, Georgia Institute of Technology, Atlanta, GA, 30332, USA
2 The Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology, Atlanta, GA, 30332, USA
3 H. Milton Stewart School of Industrial & Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30318, USA

* Corresponding Authors
Corresponding Authors-
1. Saurabh Sinha, Email- saurabh.sinha@bme.gatech.edu

17   *Abstract*

Cell–cell communication underlies tissue function, and several computational tools aim to recover
such communication from spatial transcriptomics data. However, these tools are prone to
confounders such as cell-level gene expression and spatial distribution of cell types. Moreover,
existing methods have largely focused on ligand–receptor signaling, leaving gap junction (GJ)
communication poorly charted. We present CellWHISPER, a statistically grounded method for
inferring direct communication mechanisms in the form of cell-type–signaling-gene quadruplets.
CellWHISPER employs an analytical null model that conditions on spatial organization and
expression profiles of cell types, yielding calibrated z-scores that provide improved false-positive
control. To uncover salient communication patterns, CellWHISPER also includes a latent variable
model that distils recurrent patterns into maps of mutual preferences among cell types and
signaling genes. Applied to mouse brain STEREO-seq and Xenium data, CellWHISPER
recapitulates established glial coupling, identifies novel candidates such as excitatory neuron–
astrocyte GJ interactions, and highlights conserved and condition-specific mechanisms.
Differential analysis in an Alzheimer's disease model revealed preserved astroglial Connexin43
networks and increased microglia-associated GJ communication. Together, these results
establish CellWHISPER as a confounder-aware, scalable method for mapping direct intercellular
communication from spatial transcriptomics data, enabling systematic discovery of cellular
crosstalk.

39   INTRODUCTION

Cell signaling plays a pivotal role in driving cellular heterogeneity, differentiation, and

pathogenesis within tissues. Rapid proliferation of single-cell transcriptomics technologies has led

to several computational tools for discovering cell-cell communication (CCC) mechanisms[1-4].

These tools, such as CellChat[5], CellPhoneDB[6] and NicheNet[7], typically reveal communication

between specific cell types via curated ligand-receptor (LR) pairs. Broadly, CCC can be

categorized into paracrine, autocrine, juxtacrine, and endocrine signaling[1]. Existing CCC methods

focus on paracrine and autocrine signaling mediated by LR interactions, whereas direct juxtacrine

signaling—particularly via gap junction communication (GJC)—remains largely unexplored.

Rapidly emerging spatial transcriptomics (ST) technologies including Stereoseq, VisiumHD,

Nanostring and Xenium, provide a transformative opportunity to bridge this gap by combining

gene expression with spatial proximity, enabling investigation of direct (contact-mediated)

interactions between cells.

Direct cell-cell signaling operates primarily via two mechanisms: GJC[8] and LR interactions. Gap

junctions are intercellular channels (**Fig. 1a**) that enable direct cell-cell transfer of ions,

metabolites, second messengers, and regulatory molecules[9]. GJC-mediated inter-cellular

networks play vital roles in normal physiology[10-12] and are implicated in diverse pathological

conditions[13-15]. Gap junction channels are formed by connexin proteins, assembled as hexamers

of homotypic or heterotypic connexins. Over 20 connexins have been identified in mammals, and

the existence of a compatibility "code" governing gap junction formation is an important open

problem[16,17]. Similarly, direct LR pathways, such as Notch-Delta[18] and Eph-Ephrin[19] regulate

differentiation, embryogenesis, and　immune cell communication[20], motivating the search for

additional instances of direct LR signaling in varying contexts[21].

Several recently published tools exploit ST data for CCC inference, including CellChatv2[22],

CellPhoneDBv5[23], SpaTalk[24], and COMMOT[25].　However, their error rates remain debated[3,26],

with studies suggesting inflated false positives[1,4,27]. To examine this issue for *direct* CCC, we

performed a simple randomized control by permuting cell locations within each cell type,

preserving cell-type-specific spatial organization and LR expression while destroying spatial

proximity between ligand- and receptor-expressing cells. Such a randomization should eliminate

evidence for direct CCC. Surprisingly, leading CCC tools reported similar numbers of CCC

mechanisms in the randomized and original datasets, suggesting a high false positive rate (FPR)

for direct CCC inference. Motivated by this, we focus on improving the specificity (i.e., reducing

FPR) of direct CCC detection. This motivated the development of a more stringent, confounder-

aware null model.

We present CellWHISPER (Workflow for HIgh-precision Spatial Proximity-mediated cEll-cell

inteRactions), a toolkit to infer direct CCC from single-cell ST data. CellWHISPER overcomes

spatial and expression confounders through a strong analytical null model that enables scalable

inference while maintaining low FPR. A latent variable model identifies persistent patterns across

CCC mechanisms, yielding interpretable maps of cell-type and signaling-gene preferences.

Applying CellWHISPER to mouse brain datasets from STEREO-seq[28] and Xenium reveals known

and novel GJC, reporting the first comprehensive brain "connexin code"[16,29]. LR analysis

recapitulates established CCC programs such as mossy fiber pathway while differential analyses

to identifies communication mechanisms associated with Alzheimer's Disease. In summary,

CellWHISPER provides a scalable and rigorous toolkit for mapping direct inter-cellular

communication mechanisms at the cell type-level.

RESULTS

Overview of CellWHISPER

CellWHISPER infers direct cell-cell communication (CCC) involving a "quadruplet" of two cell

types and two signaling genes. Given ST data and cell-type labels, it tests whether a cell-type

pair shows evidence of direct communication via a signaling gene pair[3] (connexins or LR pairs),

returning a list of cell type-signaling gene quadruplets with z-scores quantifying significance (**Fig.**

1b,d, Methods). A latent variable model (LVM) summarizes the resulting quadruplet set into

interpretable modules and preference maps, and an optional differential analysis identifies

condition-specific mechanisms (Methods, **Fig. 1d**).

At the core is a statistical test for whether cell types CT_i and CT_j communicate via genes SG_k and

SG_l. CellWHISPER binarizes SG expression using a quantile-based threshold. Spatial information

is represented by a k-nearest neighbors (KNN) graph[30], from which a subgraph ("whisper

network") is derived to include edges between neighboring cells of types CT_i and CT_j expressing

genes SG_k and SG_l respectively (Fig. 1b). The whisper network edge count (N_{ijkl}) quantifies the

evidence for this quadruplet in the tissue sample. To assess significance, CellWHISPER shuffles

locations among cells of same type (preserving cell-type expression and spatial distributions),

recomputes N_{ijkl} and uses the resulting null to compute a z-score. For scalability, CellWHISPER

derives an analytical null distribution that avoids explicit shuffling (Methods).

We focused initial validation on one of the best studied mechanisms of direct CCC: gap junctions

(GJ). GJ form when connexin protein in adjacent cells dimerize to establish intercellular

channels[31] (Fig. 1a). To evaluate the plausibility of using mRNA levels as proxies for connexin-

mediated communication, we performed targeted spatial profiling in an induced pluripotent stem

cell model[12] and observed concordance between *Gja1* mRNA and Cx43 protein abundance, as

well as substantial overlap between mRNA- and protein-derived whisper networks (**Fig. 1e, Supp.**

Figure 1, Methods). In a Stereo-seq mouse brain dataset[28] (Methods), aggregated whisper

networks for connexin pairs were significantly larger than those from random genes (**Fig. 1f**,

Mann-Whitney U test p-value 4.6e-8, Methods): each of 10 connexins yielded a whisper network

of ≥8878 edges, while only 22/5000 (0.4%) random genes met this criterion (**Supp. Figure 2**).

Figure 1. Overview of CellWHISPER. (a) Schematic of Gap junction communication: Connexin proteins dock at the membranes of adjacent cells, potentially of different cell types (CT_i, CT_j) to form intercellular channels called "gap junctions", enabling direct communication via transport of small molecules **such as ions, metabolites, and second messengers**. The connexin proteins (Cx_k, Cx_l) may be homotypic or heterotypic. (**b**) **Statistical testing for a CT-SG quadruplet**: For a given cell type pair (ct_i, ct_j) and signaling gene pair (g_k, g_l), CellWHISPER constructs a "whisper network", where each edge represents a pair of adjacent cells of types ct_i, ct_j expressing genes g_k, g_l respectively, and the count of such "whisper edges" serves as the test statistic. Under the null model, cell locations are randomized within each cell type, generating a background distribution of edge counts, which is used to assess statistical significance. (**c**) **Benchmarking against existing methods**: Shown is the number of cell-type pairs inferred as communicating via each ligand–receptor on average, according to four different methods, on real and randomized data. CellChat v2, COMMOT, and SpaTalk predict similar numbers of interactions on real (blue) and randomized (orange) data, indicating poor specificity and false positive rates (FPR) >90%. In contrast, CellWHISPER produced markedly fewer interactions on randomized data compared to real data, suggesting FPR < 5%. **Benchmarking was performed on a downsampled hippocampal subset (~5,000 cells) due to computational constraints of competing methods** (Methods, Supp. Fig. 5). (**d**) **CellWHISPER framework**: CellWHISPER takes single-cell resolution ST data as input, along with cell type (CT) labels and signaling gene pairs, which may be ligand-receptor (LR) gene pairs or gap junction (GJ) gene pairs. It performs a statistical test (panel b) to compute a z-score for each cell type pair and signaling

gene pair (a CT-SG quadruplet), reporting all quadruplets with significant z-scores. A latent variable model

can then analyze the full complement of quadruplet-level results to infer higher-order communication

patterns such as CT pairs that are similar in their SG usage (CT-CT similarity matrix), SG pairs that mediate

communication between similar cell types (SG-SG similarity matrix), and quantify whether a certain cell

types tends to communicate using a specific SG (CT-SG preference matrix). CellWHISPER can also be

applied across conditions for differential analysis, identifying CT-SG quadruplets specific to one condition

versus the other. (**e**) **Gap junction gene-protein concordance**: Gja1 mRNA expression correlates with

cytosolic concentration of its encoded connexin protein Cx43 (Spearman correlation = 0.47), as measured

in our induced pluripotent stem cell (iPSC)- derived cultures using targeted spatial profiling. (**f**) **Whisper**

network specificity for connexins: For each homotypic connexin gene, an aggregated whisper network

was constructed as the union of whisper networks across all cell-type pairs in the Stereo-seq mouse brain

dataset. Edge counts for connexin networks were significantly higher than those for networks constructed

from randomly selected genes (n = 5000) (Mann–Whitney U test p = 4.6×10^{-8}). All connexins yielded $\geq$

8,878 edges, whereas only 0.4% of random genes reached this threshold.

Finally, we performed the randomized control of permuting locations within each type. On

downsampled Stereo-seq data (~5,000 cells, 28 cell types, 367 LR pairs) competing tools

(CellChat v2[22], COMMOT[25], SpaTalk[24]) reported similar numbers of significant quadruplets on

real and randomized data whereas CellWHISPER reported far fewer quadruplets on randomized

data, yielding an empirically estimated FPR <5%, underscoring its high specificity (**Fig. 1c,**

Methods).

CellWHISPER identifies known and novel mechanisms of gap junction communication

(GJC) in the mouse brain

To demonstrate the utility of CellWHISPER, we first applied it to discover GJC in mouse brain.

GJC is believed to be widespread in the brain[32] and has been implicated in neurodevelopment

and diseases [33-35]. Certain combinations of cell types and connexins are preferentially involved in

brain GJC [31,36] and we sought to explore such a "connexin code"[16,29]. We analyzed a STEREO-

seq mouse hemibrain dataset[28], comprising of ~50,000 cells of 28 cell types (CTs), using ten

highly variable connexins as signaling genes (SG). (All 100 pairs of these genes were included

in the SG compendium). CellWHISPER identified 1726 significant CT-SG quadruplets (z-score >

3, whisper network size > 30, **Fig. 2a**, **Supp. Data 1**) at an empirical FPR <1% estimated from

randomized controls (**Supp. Figure 3**). These predictions were dominated by inter-glial GJC

mechanisms (**Fig. 2b**).

Figure 2. CellWHISPER Identifies Direct Gap Junction Communication in the Mouse Brain. (a) Gap junction communication identified in mouse brain: Sankey plot of the top 150 significant CT-SG quadruplets (a cell type pair communicating via a gap junction gene pair) ranked by whisper network size identified in Stereo-seq data (~50K cells). For visual ease, "Astr" represents all astrocyte subtypes (Astr1-5), "IN" represents all interneuron subtypes and "EX L2-6" represents all excitatory neuron layers. ("Olig" = oligodendrocyte, "Micro" = microglia.) Lines indicate CellWHISPER-predicted communication involving specific cell type pairs (or aggregated groups such as Astr) and specific gap junction genes: red for glial-glial communication, blue for neuronal-neuronal communication, and pink for neuroglial communication. Line thickness reflects the magnitude of communication for each CT–SG pair, while node height reflects the total communication associated with that cell type or signaling gene. **(b)** Pie chart of the three major classes of gap junction communication in the top 150 predicted CT-SG quadruplets shown in (a). **(c)**

Oligodendrocyte self-communication. Whisper network of oligodendrocytes communicating via
homotypic Cx32 gap junctions. Blue edges represent whisper network connections; red shading denotes
kernel density smoothing at edge centers for visual clarity. (**d**) **Statistical significance of whisper edges.**
Histogram of edge counts from randomized whisper networks (cell locations permuted within cell type), with
fitted normal distribution (blue line). The red dashed line marks the observed number of edges in the whisper
network shown in (c) which is significantly higher than expected under the null. Y-axis shows the probability
density of randomized edge counts. (**e**) **Heterotypic astroglial communication.** Whisper network of
astrocytes predicted to communicate with oligodendrocytes via heterotypic gap junction formed by Cx43
(astrocytic side) and Cx47 (oligodendrocytic side). (**f**) Cell types predicted to communicate using homotypic
Cx36 gap junctions. Blue entries in heatmap denote significant cell type pairs (z score>3, whisper edge
count > 30) found by CellWHISPER. Neuronal communication is predicted more frequently compared to
glial communication. (**g**) Glial cell types predicted to communicate using Cx43-Cx43 gap junctions. Only
glial communication is shown since it is predicted more frequently compared to neuronal communication
(Supp. Fig. 6) (**h–j**) Astrocytic communication enriched in hippocampus. Whisper networks showing (h)
astrocyte–astrocyte, (i) astrocyte–endothelial, and (j) astrocyte–microglia interactions mediated by
homotypic Cx43 (GJA1) gap junctions.

We examined the highest-confidence mechanisms from two complementary perspectives.
Ranking quadruplets by whisper network size revealed that the top six represent inter-
oligodendrocyte GJC via combinations of Cx47, Cx32, and Cx29 (**Supp. Data 1, Supp. Data 2**)
consistent with established roles of Cx47 and Cx32 in oligodendrocytic coupling[37,38]. **Fig. 2c**
illustrates homotypic Cx32 whisper network between oligodendrocytes, comprising 4,283 edges,
far exceeding chance expectation (**Fig. 2d**, z-score = 13.09). The top predictions also included
Cx29-mediated oligodendrocytic GJC. Although Cx29 is frequently associated with
oligodendrocytes[39], prior studies indicate that it primarily localizes to the adaxonal membrane
rather than forming functional gap junctions[40]. This likely represents a false-positive prediction
and underscores a limitation of transcript-based inference when subcellular protein localization is
critical. The next highest-ranking quadruplets involve astrocyte-astrocyte GJC via Cx43, Cx30,
and Cx26, connexins known to couple astrocytes into an astrocytic syncytium essential for
homeostasis and adult neurogenesis[17,41-44]. We also identified high-confidence astrocyte-
oligodendrocyte (A/O) communication via Cx43 on the astrocyte side and Cx47, Cx32 or Cx29 on
the oligodendrocytic side. **Fig. 2e** illustrates *Astr3*-oligodendrocyte communication via Cx43-
Cx47, consistent with prior reports supporting functional A/O GJ coupling[45], though there is some
unresolved debate about their functionality[44].

Examining CellWHISPER results ranked by z-score rather than network size (**Supp. Data 3**)
highlights additional mechanisms supported by fewer communicating cell pairs, which may thus
be less well-studied in prior work. Among these, we identified GJC between excitatory neurons
and other cell types, including thalamic excitatory neurons, oligodendrocytes, and astrocytes
(**Supp. Data 3**). Maubecin et al.[46] provided direct evidence of functional neuro-glial coupling using
heterotypic gap junctions in rodents. Gap junctions between excitatory neurons are known to be
primarily mediated by Cx36 (Gjd2)[47,48] and extensively studied in rodent models[49-52]. As shown in
Fig. 2f, excitatory neurons are a major site of homotypic Cx36-mediated communication predicted
by CellWHISPER.
231
CellWHISPER identifies homotypic Cx43 gap junctions as a mechanism underlying
astrocytic syncytium
Cx43 is the most studied connexin, with key roles in differentiation and homeostatic regulation[53-
55]. We examined the complete map of cell type pairs predicted by CellWHISPER to participate in
Cx43-mediated GJC (**Fig. 2g**), finding a predominance of glial over neuronal communication. For
instance, astrocyte subpopulation Astr1 is predicted to communicate with other astrocytes, as well
as endothelial, microglia, oligodendrocytes and ependymal cells. This aligns with known biology:
Cx43 mediates astrocyte-oligodendrocyte-ependymal cell interactions forming an isopotential
syncytium [54,56-58]. Predicted astroglial interactions exhibited a higher network density in the
hippocampus (**Figs. 2h-j**). Inter-astrocyte GJC (z-score 18.13, **Fig. 2h**) in the hippocampus has
been documented to be Cx43-mediated. Astrocyte-endothelial GJC (z-score 10.94, **Fig. 2i**) has
been implicated in maintaining blood–brain barrier integrity[59-61]. Additionally, CellWHISPER
predicted Cx43-mediated communication between microglia and astrocytes (**Fig. 2j**) as well other
microglia. Indeed, recent studies suggest that microglia upregulate Cx43 under pathological or
inflammatory conditions and form functional gap junctions – with each other and with astrocytes[62-
64].
248
To validate these predictions, we performed immunofluorescent staining on mouse coronal brain
sections for Cx43 together with markers of astrocytes (GFAP), endothelial cells (CD31), and
microglia (IBA1) (**Fig. 3a, b**; Methods). All channels showed robust signal across brain regions
(**Fig. 3b**). Distinct Cx43 co-localization was evident at astrocyte–astrocyte, astrocyte–endothelial,
astrocyte–microglial, and microglial–microglial interfaces. Regions of interest were identified
using the Allen Mouse Brain Atlas[65], focusing on the hippocampus and cerebral cortex (outside
hippocampus). Quantitative analysis showed higher co-localization of Cx43 with cell-type markers

Figure 3. Spatially resolved analysis of Cx43-mediated cellular crosstalk across brain regions. (**a**)
Schematic of gap junction–mediated interactions between astrocytes (GFAP), microglia (IBA1), and
endothelial cells (CD31) via Cx43 (created with BioRender.com). (**b**) Immunofluorescence of coronal mouse
brain section stained for Cx43, GFAP, IBA1, and CD31. Hippocampal boundaries were defined using the
Allen Mouse Brain Atlas. Regions of interest (ROI) within hippocampus (Hippo1, Hippo2) and cortex were
outlined for higher-resolution imaging. Scale bar, 100 μm. (**c**) Single-cell–resolution images of hippocampal
ROIs showing colocalization of astrocytes and microglia with Cx43. (**d**) Quantification of Cx43 colocalization
across hippocampus and cortex. We selected 10 regions (478 × 262 pixels each) within and outside the
hippocampus and quantified Cx43 colocalization with astrocytes, endothelial cells, and microglia.
Colocalization was defined as instances where Cx43 puncta were positioned between or overlapping the
indicated cell types, yielding 30 values (one per cell type per region) and a total of 333 interactions. Two-
way ANOVA revealed a significant enrichment of astrocyte–Cx43 interactions in the hippocampus
compared to the cortex (p = 0.0002).

in the hippocampus compared to cortex (**Fig. 3d**), with astrocytic contacts enriched >5-fold.
Representative examples of colocalization of astrocytes, microglia and Cx43 in the hippocampus
are shown in **Fig. 3c**. Given the hippocampus' central role in learning, memory consolidation, and
emotional regulation, these observations point to a potential role of Cx43-mediated coupling in
coordinating blood–brain barrier regulation and neuroinflammatory signaling.

CellWHISPER identifies direct ligand-receptor (LR) communication
Direct LR communication plays a fundamental role in organizing local tissue architecture[21]. To
identify such interactions in the mouse brain, we re-analyzed STEREO-seq data using 367
curated LR pairs from the CellChat database[5] (Methods). CellWHISPER identified 4743
significant CT–SG quadruplets (z-score > 4, whisper network size > 30) (**Fig. 4a, Supp. Data 4**),
with an empirically estimated FPR < 1%, spanning inter-glial, neuron-glia and neuron-neuron
communication (**Fig. 4b**).

The highest confidence predictions revealed communication between CA3 excitatory neurons
(EX CA) and dentate gyrus granule neurons (GN DG) mediated by LR pairs of the mossy fiber
pathway[66] (**Supp. Data 5**). These include multiple ephrin/Eph family members, C1ql2/3-Adgrb3
and homotypic interactions of occludin, protocadherin and claudin (**Supp. Data 5, Fig. 4c**).
Ephrin–Eph findings are consistent with known hippocampal gradients that mediate excitatory
connectivity and synaptic adhesion[67,68]. C1ql2/3 act as organizers of mossy fiber–CA3 synapses

Figure 4. CellWHISPER Identifies direct Ligand-Receptor communication in the mouse brain. (a)
Ligand–receptor communication mechanisms. Sankey plot showing the top 25 significant CT–SG
quadruplets **ranked by z-score** (z-score > 4, whisper network size > 30) identified using 367 curated
ligand–receptor pairs. Randomized control analysis estimated a false positive rate (FPR) < 1% for this
compendium. **(b)** Communication classes. Distribution of major interaction types, including neuron–neuron,
neuron–glia, and inter-glial communication. **(c–e)** Six representative whisper networks (two per panel,
left/right). Selected ligand–receptor–mediated cell–cell interactions illustrating key modes of direct
communication: (c) excitatory CA3 neurons (EX CA) and dentate gyrus granule neurons (GN DG)
communicating via Efnb3-Epha4 (right) and tight junction Ocln (left); (d) excitatory neuron-oligodendrocyte
interactions involving *Cldn* (right) and semaplexin family member (left); and (e) meningeal- astrocyte
communication via extracellular matrix components (*Lama2*, *Col4a1–6*, *Adgrg6*).

by binding ADGRB3 (BAI3) to regulate excitatory synapse formation through bidirectional trans-
synaptic signaling[69]. Claudins and occludin, classically tight-junction components, are enriched in
hippocampal neurons and associate with synaptic scaffolding proteins[70], suggesting adhesive
function at synapses. Together, these interactions suggest a multilayered molecular basis for
mossy fiber circuit organization.

CellWHISPER also predicted LR-mediated communication between oligodendrocytes and
excitatory neurons, involving homotypic MAG**,** claudin and semaplexin interactions (**Supp. Data**
6, Fig. 4d), consistent with known roles for these mechanisms in axon–glia recognition, myelin
stability[71], and activity-dependent modulation of myelination[72]. Additionally, we observed
interactions between astrocytes and meningeal cells, notably involving extracellular matrix
components (ECM) such as Lama2, type IV collagens (Col4a1–6), Adgrg6 (**Supp. Data 7, Fig.**
4e).

Latent Variable Modeling (LVM) infers general patterns of communication mechanisms
We used CellWHISPER's "LVM" to infer higher-order patterns among significant quadruplets
(Methods). Unlike approaches that first collapse quadruplet-level information into CT x SG or CT
x CT matrices[5,6], CellWHISPER directly models quadruplets, learning a shared low-dimensional
embedding space for cell types and signaling genes, such that proximity in the embedding space
reflects CCC preferences. The learned embeddings yield (1) CT–SG preferences, (2) CT–CT
similarity, and (3) SG–SG similarity. Model parameters are estimated via constrained log-
likelihood maximization. **Fig. 5b** shows that the optimization can accurately infer cell type
similarity on simulated data (Methods).

328
329 **Figure 5. LVM infers module of signaling cell types and signaling genes.** (**a**) Cell type similarity matrix.
330 Cosine similarity of latent cell-type embeddings derived from the LVM routine of CellWHISPER, applied to
331 gap junction communication in the Stereo-seq mouse brain dataset. Five dominant modules of cell types
332 emerge, reflecting recurrent patterns of gap junction communication across the tissue. (**b**) Optimization.

Constraint optimization within the latent variable model (Methods) identifies underlying cell type pair similarity, resolving higher-order structure in the data. (**c**) Cell type–gap junction preferences. Heatmap showing associations between cell type modules and gap junction gene modules. Distinct modules exhibit preferential coupling, e.g., Module M1 and M2 are enriched in excitatory neurons and prefer Gjd2 (GJ module M4). DA neurons and IN S+ prefer GJ module M2. (**d**) Cosine similarity of latent embeddings across gap junction genes, revealing four dominant connexin modules. (**e**) Cell-type similarity matrix for ligand–receptor communication. Cosine similarity of latent cell-type embeddings derived from CellWHISPER applied to ligand–receptor interactions (Stereo-seq mouse brain dataset). Four dominant cell-type modules emerge. Excitatory neurons in cortical layers and hippocampal CA region cluster together (M2), whereas excitatory neurons in thalamus is in module M4. (**f**) Communication overlap between excitatory neurons. Quantification of overlap in ligand–receptor communication patterns (shared CT– ligand–receptor triplets) shows that excitatory neurons in cortical layers 2/3 are more similar to CA excitatory neurons than to thalamic excitatory neurons, consistent with known laminar and neuromodulatory properties of neocortical and hippocampal neurons. (g–h) Cell type–ligand and cell type–receptor module specificity. Sankey plots showing preferential associations between cell-type modules and ligand modules (g), and between cell-type modules and receptor modules (h). Links were derived by averaging CT–SG preference scores within each module. (**i**) Ligand similarity map. Cosine similarity of ligand embeddings reveals five dominant modules (Supp. Data 9). Module M1 consists mainly of genes related to cell adhesion (e.g., PECAM1, CDH1/5, ICAM1), extracellular matrix components (COL4A1–6, LAMA2), and immune modulators (H2-Aa, H2-Eb1, H2-Ab1, CD52, CD55, CD86, PTPRC). Another ligand module (M5, Supp. Data 9) is enriched for axon guidance and synaptic signaling genes , including protocadherins, ephrins, neurexins, semaphorins, Lrrc and Flrt family (**j**) Receptor similarity map. Cosine similarity of receptor embeddings identifies five receptor modules(Supp. Data 9).

Applying LVM to GJ results in mouse brain Stereo-seq data revealed five dominant cell type modules in CT-CT similarity map (**Fig. 5a, Supp. Data 8**) whereas a baseline method performing naïve aggregation of quadruplet tensor did not reveal modular structure (**Supp. Figure 4**). LVM also identified four GJ modules in SG-SG similarity map (**Fig. 5d**, Supp. Data 8). Cell type modules showed preferences for specific GJ modules in the CT-SG preference matrix (**Fig. 5c**). For example, excitatory neuron subtypes in cell type modules M1/M2 (EX Mb, EX L2/3, EX CA) prefer GJ module M4 consisting of Gjd2 (Fig. 5c,d). Notably, DA neurons (excitatory or inhibitory depending on context[73]), clustered with inhibitory Sst+ interneurons[74] in cell type module M3 (**Fig. 5a**, Supp. Data 8) and shared preference for GJ module M2 (**Fig. 5c**), consistent with dye-coupling evidence of DA–interneuron electrical synapses[75] and suggesting candidate GJ mediating these synapses.

Next, we applied LVM to LR interactions (Fig. 4), identifying four dominant cell-type modules (**Fig.**
5e, Supp. Data 9). Excitatory neurons in cortical layers and CA region co-cluster, whereas
thalamic excitatory neurons lie in different module (**Fig. 5e,f**). This aligns with known features of
excitatory neurons in the neocortex and hippocampus (of which CA is part), including laminar
structure[76,77] and shared neuromodulatory response[78], different from excitatory neurons in
thalamic neurons[79]. Analysis of ligand similarity map (**Fig. 5i**) identified five dominant modules, of
which one (M1) is enriched for cell adhesion, ECM and immune modulators (**Supp. Data 9**). This
ligand module preferentially associates with cell type module M1 (**Fig. 5g**), enriched for glial cell
types located near the brain periphery, such as ependymal cells, meninges, and smooth muscle
cells (Fig. 5e), suggesting a specialized glial–meningeal signaling axis potentially involved in
barrier maintenance and neuroimmune communication. Another ligand module (M5, Supp. Data
9) is enriched for axon guidance and synaptic signaling genes[80,81], and prefers cell type module
M2 (**Fig. 5g**), enriched in excitatory neurons in the cortical layers and hippocampus, consistent
with their known role in neuronal cell-cell communication[82]. Similarly, we found five dominant
modules in receptor similarity map (**Fig. 5j**, Supp Data 9) having preference for specific cell type
modules (**Fig. 5h**). Together, LVM captures dominant communication patterns across GJC and
LR signaling.
386
Differential analysis reveals Alzheimer's disease-associated GJC mechanisms
Gap junction-mediated communication (GJC) supports central nervous system homeostasis and
is implicated in neurodegenerative diseases, including Alzheimer's disease (AD)[31,83,84]. Under
normal conditions, GJC supports metabolic and synaptic function, especially through astrocytic
networks[63,85] whereas in AD, aberrant gap junctions and hemichannel activity contributes to
neuroinflammation and neurodegeneration[31,83,84 31]. To systematically interrogate these dynamics,
we used CellWHISPER to decode GJC networks altered in AD.
394
We analyzed Xenium mouse brain data from wild-type and TgCRND8 mice[86] (aged >13 months,
Methods) using three annotated GJ genes. After cross-sample integration to harmonize cell-type
labels (Methods), CellWHISPER's Diffrential Analysis (DA) function quantified quadruplets per
condition, and classified them as shared, condition-specific or indeterminate (Methods). This
revealed 60 high-confidence quadruplets (z-score > 4, whisper network size > 30, **Fig. 6a, Supp.**
Data 10). The twenty shared quadruplets are enriched for astrocytic (13/20) and oligodendrocytic
communication (9/20) (**Fig. 6b**) including Astr1-Olig1 homotypic Cx43 (**Fig. 6c**). Despite known
Cx43 upregulation near Aβ plaques, prior work suggests structural preservation of astrocytic GJC

in AD brain[87]; consistent with this, CellWHISPER identifies conserved GJC mechanisms despite

change in signaling gene expression.

Figure 6. Differential Analysis infers conserved and condition-specific gap junction communication

in Alzheimer's disease (AD). (**a**) Scatter plot of 60 high-confidence CT–SG quadruplets (z-score > 4,

whisper network size > 30 in at least one condition) showing their z-scores in wild-type (WT) and TgCRND8

(AD) Xenium datasets for Gja1, Gjb2, and Gjc3. Quadruplets are classified as shared, condition-specific,

or indeterminate. (**b**) Bar plot showing the frequency of each cell type across the 60 quadruplets. Astrocytes

and oligodendrocytes are most enriched in shared interactions, microglia dominate AD-specific interactions,

and astrocytes are more frequent in WT-specific interactions. (**c**) Whisper networks of Astr1–Olig1

communication via homotypic Cx43 in WT (left) and AD (right), showing preserved symmetry between

conditions. (**d**) Selected condition-specific quadruplets, showing z-scores in WT versus AD. Top three WT-

specific and AD-specific interactions were chosen based on whisper network size. (**e**) Whisper network of

Astr1–Ex th1 communication via homotypic Cx43 in WT. (**f**) Whisper network of Smc1–Smc1 (smooth

muscle cell self-communication) via homotypic Cx43 in AD, highlighting potential vascular remodeling. (**g**)

Aggregated microglial whisper network in AD, constructed by taking the union of all AD-specific quadruplets

involving microglia, revealing enrichment in thalamus and hypothalamus regions.

DA further identified 15 TgCRND8-specific and 14 wild-type-specific quadruplets (Fig. 6a,d,

Supp. 10), including reduced astrocyte-thalamic excitatory neuron GJC via homotypic GJA1 (**Fig.**

6e). In contrast, TgCRND8-specific mechanisms included increased smooth muscle cells

homotypic GJA1 (Fig. 6d,f), potentially reflecting vascular remodeling[88,89]. TgCRND8-specific

mechanisms were enriched for microglial interactions (5/17 vs. 0/23 in WT-specific, p-value <

0.01, two-tailed proportions test), and WT-specific mechanisms to be marginally enriched for

astrocytes (6/23 vs. 2/17 in TgCRND8) (**Fig. 6b**). These trends align with existing literature:

astrocyte networks maintain homeostasis in healthy brain[90], while microglial GJC becomes more

prominent during AD neuroinflammation[31,91,92]. Aggregated microglial whisper network (**Fig. 6g**)

showed enrichment in thalamus and hypothalamus. Connexin alterations have been reported in

thalamic regions in AD[83] while microglial changes in thalamus are linked to cognitive impairment

after cortical brain injury[93].

DISCUSSION

CellWHISPER provides a statistically rigorous framework for decoding direct cell-cell

communication from spatial transcriptomics. By explicitly accounting for confounding effects of

gene-expression and spatial organization, CellWHISPER markedly reduces FPR relative to

competing tools. Its LVM gleans higher order trends by operating directly on CT-SG quadruplets,

while differential analysis allows identification of condition-specific mechanisms.

Applying CellWHISPER to mouse brain ST data revealed the first large-scale map of connexin

code, recapitulating canonical glial networks and generating testable hypotheses.

Immunostaining validated Cx43 interactions, demonstrating its colocalization at astrocyte–

astrocyte/endothelial/microglial interfaces. LR analysis uncovered adhesion/synaptic programs,

including the mossy fiber pathway and a glial–meningeal signaling axis. Differential analysis of an

AD mouse model identified a shift toward microglial-associated GJC and region-specific

remodeling in the thalamus and vasculature, consistent with emerging models of

neuroinflammatory progression.

Several limitations motivate future work. CellWHISPER inference relies on mRNA as a proxy for

membrane proteins, and post-transcriptional regulation or subcellular localization can yield false

positives. It prioritizes specificity through a confounder-aware null model, which may come at the

cost of sensitivity. In contrast, CellWHISPER excels at detecting communication events mediated

by a smaller number of cell pairs. Extensions to lower-resolution platforms (e.g. Visium) may be

possible via deconvolution approaches, and integrating pathway information could improve
robustness of whisper networks. Finally, segmentation errors in ST[94] may propagate to CCC
inference, a limitation shared across current methods.

In summary, CellWHISPER establishes a robust paradigm for inferring direct signaling from ST
data, combining stringent statistical control with interpretable latent structure, to enable
mechanistic dissection of tissue organization and for identifying disease-specific communication
in development, homeostasis, and disease.

METHODS

Overview of CellWHISPER

CellWHISPER infers direct cell–cell communication (CCC) from spatial transcriptomics (ST) data.
The input consists of gene expression profiles, spatial coordinates, and cell-type annotations for
individual cells, together with a precompiled compendium of signaling gene (SG) pairs, such as
connexins or ligand–receptor (LR) pairs. SG compendia can be obtained from existing resources
or provided by the user as a custom list.

For each candidate combination of a cell-type pair (ct_i, ct_j) and a signaling gene pair (sg_k, sg_l),
CellWHISPER tests whether the two cell types exhibit evidence of direct communication mediated
by the signaling genes (Fig. 1b). Statistically supported interactions are reported as CT–SG
quadruplets, each associated with a z-score quantifying significance (Fig. 1d). CellWHISPER
includes three analysis modules (Fig. 1d).
(i) A statistical testing module identifies significant CT–SG quadruplets.
(ii) A latent variable model (LVM) summarizes the collection of significant quadruplets into
interpretable preference and similarity maps, including CT–CT, CT–SG, and SG–SG
relationships.
(iii) An optional differential analysis (DA) module identifies condition-specific or shared
communication mechanisms across biological conditions.

Whisper network construction

Gene expression was binarized using a quantile-based threshold (75th percentile by default),
yielding binary indicators of whether a signaling gene is expressed in a given cell.

Spatial proximity between cells was represented using a k-nearest neighbors (KNN) graph
constructed from cell coordinates (k = 5 unless otherwise specified). This graph defines the set

of potential physical contacts used for CCC inference. For datasets with irregular tissue geometry,

distance-based thresholds should be used instead of fixed neighbor counts.

For each CT–SG quadruplet (ct_i, ct_j, sg_k, sg_l), a whisper network was derived by retaining only

edges connecting neighboring cells of types ct_i and ct_j that express sg_k and sg_l, respectively

(Fig. 1b). The number of edges in this subgraph, denoted N_{ijkl}, quantifies spatially supported

evidence for the corresponding communication mechanism.

Statistical test for CT–SG quadruplets

To assess statistical significance, CellWHISPER constructs a null model by permuting cell

identities within each cell type. This preserves cell-type–specific gene expression distributions

and spatial organization while disrupting spatial proximity between signaling cells of different

types. The whisper network is recomputed under the null, yielding a null distribution for N_{ijkl}, from

which a z-score is computed.

To enable scalability to large datasets, CellWHISPER derives an analytical approximation to the

null distribution, avoiding explicit permutations while preserving control of confounding effects

arising from expression heterogeneity and spatial cell-type structure.

Analytical approach for Statistical test of a quadruplet:

Let Z_u be a Boolean random variable indicating the presence of a whisper edge for a proximal

cell pair corresponding to a quadruplet i, j, k, l. Our objective is to estimate the expectation and

variance of the total number of such whisper edges: $E(\sum_u Z_u), Var(\sum_u Z_u)$.The probability that a

specific edge u is present (i.e. $Z_u = 1$) is modeled as the product of probabilities that gene k is

expressed in cell of type i and gene l is expressed in a cell of type j. Gene expression is

determined by a global percentile threshold across all cells. Thus:

$$P(Z_u = 1) = p_i p_j$$

Where p_i is the probability that a cell of type i expresses gene k, and p_j is the probability that a

cell of type j expresses gene l.

If Z_u are Independent (i.i.d.):

Assuming the Z_u are i.i.d. Bernoulli variables, the total follows a Binomial distribution with N_u

WHISPER edges:

521
$$E\left(\sum_u Z_u\right) = \sum_u P(Z_u) = N_u P(Z_u = 1) = N_u p_i p_j$$

522
$$Var\left(\sum_u Z_u\right) = N_u P(Z_u = 1)(1 - P(Z_u = 1)) = N_u p_i p_j (1 - p_i p_j)$$

524 **Accounting for Dependence Between Edges:**

525 However, the Z_u variables are not independent because edges may share a common cell. While

526 the expectation remains unchanged due to linearity, the variance must account for dependencies:

527
$$Var\left(\sum_u Z_u\right) = E\left((\sum_u Z_u)^2\right) - \left(E\left(\sum_u Z_u\right)\right)^2$$

528
$$= E\left((\sum_u Z_u)^2\right) - N_u^2 p_i^2 p_j^2$$

529 To expand the first term,

530
$$E\left((\sum_u Z_u)^2\right) = E\left(\sum_u Z_u^2\right) - E(\sum_{u,v} Z_u Z_v)$$

531 Since, $Z_u \in [\{0,1\}]$, $Z_u^2 = Z_u$, so the first sum reduces to $E(\sum_u Z_u) = N_u p_i p_j$. The second term

532 depends on the structure of edge overlap and has the following cases (assuming $i \neq j$):

- Case1: No shared cells between u and v , therefore Z_u, Z_v are independent, $E(Z_u Z_v) =$ 535 $p_i^2 p_j^2$

- Case 2: Shared cell of type j but distinct cells of type i, therefore, $E(Z_u Z_v) = p_i p_j^2$

- Case 3: Shared cell of type i but distinct cells of type j, therefore $E(Z_u Z_v) = p_i^2 p_j$

If $i = j$, cases 2 and 3 collapse into a single case due to symmetry.

Stereoseq Data and Analysis

Mouse brain STEREO-seq data[28] were obtained from the Spatial Omics Database[95]. After quality

control filtering, the dataset comprised 50,140 cells and 18,527 genes, including 10 gap junction

(connexin) genes: *Gjd2* (Cx36), *Gja5* (Cx40), *Gjc1* (Cx45), *Gja4* (Cx37), *Gjb2* (Cx26), *Gja1*

(Cx43), *Gjb6* (Cx30), *Gjb1* (Cx32), *Gjc2* (Cx47), and *Gjc3* (Cx29). We used the authors' annotated

cell-type labels, encompassing 28 distinct cell types. Abbreviations include: CA3, cornu ammonis

area 3; Mb, midbrain; EX, excitatory glutamatergic neuron; IN, GABAergic interneuron; DA,

dopaminergic neuron; GN DG, granule cell of dentate gyrus; Astr, astrocyte; Micro, microglia;

OPC, oligodendrocyte precursor cell; Oligo, oligodendrocyte; SMC, smooth muscle cell; Ery,

erythrocyte; and Endo, endothelial cell.

Gene expression matrices were normalized and subsequently denoised using MAGIC imputation

(t=2) to recover continuous expression profiles. To construct the *whisper network*, a gene was

considered "expressed" in a given cell if its expression exceeded the 75th percentile of that gene's

expression across all cells. For gap-junction analysis, we evaluated all possible connexin–

connexin combinations. Although several connexins are known to preferentially form specific

homo- or heterotypic channels, experimental evidence for many pairings remains incomplete or

context-dependent; thus, interaction patterns were inferred empirically from the data.

Ligand–receptor database

We used the CellChat v2[22] ligand–receptor compendium, focusing specifically on pairs annotated

as cell–cell contact interactions. After intersecting this database with the gene set retained from

the STEREO-seq dataset, 367 ligand–receptor pairs were included for downstream analyses.

Randomization and False Positive Rate estimation:

We performed randomization by permuting cell IDs within each cell type. This procedure

preserves the marginal spatial distribution and gene expression profiles specific to each cell type.

False positive rate (FPR) is obtained by comparing the number of detected pairs obtained on

randomized data with number of detected pairs on real data.

Benchmarking

We couldn't run Commot and Spatalk on full mouse hemibrain dataset containing 50K cells.

Therefore, we selected part of hippocampal region (Supp. Fig. 5) containing 5K cells for

benchmarking analysis in Fig. 1c. We randomized by permuting cell IDs within each cell type.

All comparison methods were downloaded from their official repositories and executed using

recommended parameters. For CellChat v2, we used the computeCommunProb function with

default parameters (type = "triMean", trim = 0.1, interaction.range = 250, contact.knn.k = 5),

followed by filterCommunication with a minimum of 20 cells per cluster. For COMMOT, we applied

commot.tl.cluster_communication to compute cell–cell communication for each ligand–receptor

pair. For SpaTalk, we used the default parameters.

Connexin mRNA-protein concordance

CellWHISPER relies on cellular mRNA abundance as a proxy for membrane-bound connexin

proteins. To evaluate this assumption, we performed targeted spatial profiling of connexin 43

(Cx43) protein and its corresponding mRNA (Gja1) at single-molecule resolution in an induced

pluripotent stem cell (iPSC) model, where gap junction communication is critical for spatial

patterning.

We observed a significant correlation between Gja1 mRNA expression and average cytosolic

Cx43 protein intensity at the single-cell level (Spearman's R = 0.47, p = 5 × 10^{-20}; Fig. 1e).

Consistently, whisper networks inferred from Gja1 mRNA showed strong overlap with those

inferred from Cx43 protein measurements (Fig. 1e, Supp. Fig. 1), supporting the use of mRNA as

a proxy for connexin-mediated communication in this context.

Latent Variable Model (LVM):

LVM stipulates that the binary variable $I(ct_i, ct_j, g_k, g_l)$ indicating whether the CT-SG quadruplet

ct_i, ct_j, g_k, g_l has statistical support in the data follows a Bernoulli distribution:

$$I(ct_i, ct_j, g_k, g_l) \sim Ber\,(p_{ijkl})$$

where u_i , u_j denote to-be-learnt low-dimensional representations (embeddings) for CTs ct_i , ct_j

and v_k ,v_l denote embeddings for SGs g_k, g_l, respectively, all embeddings being in a common

latent space. The dot product $u_i.v_k$ captures the general propensity of ct_i for participating in CCC

using g_k. Thus, the model posits that the quadruplet ct_i, ct_j, g_k, g_l is more likely if there is a higher

tendency for ct_i communicating via g_k and ct_j communicating via g_l.

Using this formulation CellWHISPER learn embeddings that enable quantification of (1) cell type-

signaling gene preferences ($u_i.v_k$), (2) cell type similarities ($u_i.u_j$), i.e., which pairs of CTs have

similar communication mechanisms, and (3) signaling gene similarities ($v_k.v_l$), i.e., which SGs

are used by the same cell types. It uses cosine similarity rather than dot products between

embeddings to emphasize specificity and mitigate abundance bias inherent in dot product.

In the null model, p_{ijkl} is assumed to be constant for all indices i, j, k, l. In contrast, the alternate

model factorizes p_{ijkl} into two terms:

•   the propensity of cell type i to communicate via signaling gene k, and

•   the propensity of cell type j to communicate via signaling gene l

Formally, this is expressed as:

619
$$p_{ijkl} = (u_i . v_k)(u_j . v_l)$$

where u_i and u_j are low dimensional embeddings for cell types i, j, and v_k, v_l are embeddings

for signaling genes k and l. All embeddings lie in a shared latent space, and we enforce the

constraints:

$$0 < u_{ir} < 1, \qquad 0 < v_{ir} < 1, \qquad \left\|u_{ir}\right\|_1 < 1, \left\|v_{kr}\right\|_1 < 1$$

This formulation is invariant under simultaneous permutation of both CT and SG pairs, i.e.

$$I\big(ct_i, ct_j, g_k, g_l\big) = I\big(ct_j, ct_i, g_l, g_k\big) \Rightarrow p_{ijkl} = p_{jilk}$$

We optimize the model using constrained log-likelihood maximization, implemented via

`scipy.optimize` using the SLQSP method. The constraint is posed using the L1 norm of the

embeddings that ensures that the problem is a linear optimization and probabilities $0 < p_{ijkl} < 1$.

To confirm identifiability and recovery, we simulated Gap Junction Communication results with

the same orders as the STEREO-seq analysis (29 cell types, 10 connexins, d=3). Ground-truth

embeddings were drawn from numpy.random.rand with unit norm cell type and signaling gene

embeddings. The optimizer recovered CT–CT similarity, SG–SG similarity, and CT–SG

preference structure with high agreement (Fig. 5b).

For ligand–receptor analysis, only a subset of ligand–receptor pairs are biologically plausible. To

ensure model learnability, we explicitly set the probability $p_{ijkl} = 0$ for incompatible ligand–

receptor combinations. For computational feasibility, we randomly subsample a small number of

such incompatible pairs (2000 in our experiments) to include as negative examples during

optimization. This approach enables efficient training while maintaining biological specificity in the

learned embeddings.

Differential Analysis:

We performed Differential Analysis (DA) at the terminal time point at which amyloid pathology is well established. Wild-type (WT) and TgCRND8 (AD) Xenium datasets were jointly integrated using residualPCA to achieve conservative alignment across conditions while preserving biological differences. A joint embedding was constructed and cell-type clustering was performed on the integrated space yielding 25 clusters.

Label transfer from reference. Cell-type labels were transferred from the annotated STEREO-seq reference (analyzed above). For this step, the WT, AD, and STEREO-seq datasets were co-embedded with reciprocal PCA[96]. For each query cell (WT or AD), we identified its k = 7 nearest neighbors in the reference and recorded their labels. For each cluster in the joint WT+AD embedding, neighbor votes were aggregated across all member cells and normalized by the total number of votes in that cluster to yield a label-probability vector. The cluster label was assigned as the cell type with the maximum normalized vote. Clusters containing multiple subpopulations of the same broad class (e.g., oligodendrocytes) were further subdivided and relabeled as subtypes (e.g., Olig-1, Olig-2).

DA quantification. Using these harmonized annotations, CellWHISPER quantified CT–SG quadruplets separately in WT and AD datasets. Quadruplets were retained if they satisfied significance thresholds (z-score > 4 and whisper-network size > 30). This procedure yielded 60 high-confidence quadruplets across the two conditions. A quadruplet was classified as shared if both WT and AD satisfied these thresholds, condition-specific if one condition satisfied the thresholds while the other did not (z-score < 3 or whisper-network size < 20), and indeterminate otherwise.

Experimental details for GJA1 mRNA-protein correlation
iPSC culture
iPSCs were obtained and cultured according to Culture and Freezing Methods for WTC Derived AICS hiPSC Lines: https://www.coriell.org/0/PDF/Allen/iPSC/AICS_SOP_WTC_CellCulture.pdf. Coverslips (22 × 22 mm) were placed flat in sterile 6-well plates and coated with growth factor–reduced, phenol-red–free Matrigel (Corning, cat. no. 356231) at a final protein concentration of 0.337 mg/mL. Briefly, Matrigel aliquots thawed at 4 °C were diluted on ice in cold DMEM/F12 to 1.5 mL per well, mixed thoroughly with pre-chilled pipettes to ensure homogeneity, and

immediately dispensed to fully cover each coverslip. Plates were incubated at room temperature

for 1 h, after which excess coating solution was aspirated and wells were gently rinsed with room-

temperature mTeSR1 medium supplemented with 1 µM Y-27632 (ROCK inhibitor) before cell

seeding. Coated coverslips were used for experiments within 6 h or, if needed, sealed with

Parafilm and stored at 4 °C for up to 2 weeks. Once cells were confluent, they were fixed in 4%

formaldehyde for staining.

iPSC RNA and protein multiplexing

Fixed cells on coverslip with permeabilized with 70% EtOH for 30 mins at -20 C. Cycle 1 RNA

was labeled using Molecular Instruments hybridized chain reaction RNA fluorescence in situ

hybridzation (HCR RNA FISH) v3.0 probes: Human Cx43 B3 555, Human MALAT1 B2 647. All

probes were amplified 75 mins at room temperature. Fluorescence images were acquired on

adapted Nikon Ti 2000-U microscope with appropriate filters. Then, samples were blocked and

incubated with Cx43 Ab (Invitrogen 71-0700, 1;150) overnight at 4 C followed by secondary

antibody staining (anti-rabbit 488) for 1 hr at room temperature. This Cycle 2 Cx43 Ab was imaged

again. DAPI counterstain was used to register the images from both cycles. Images were

segmented with Cellpose and quantified for downstream analysis.

Immunofluorescence staining

Mouse brain sections were fixed in 4% paraformaldehyde (PFA) and sectioned using Slides were

deparaffinized and rehydrated by subsequent rinses in Xylene, 100% EtOH, 95% EtOH, 80%

EtOH, 70% EtOH and deionized water (diH$_2$O). Then we performed antigen retrieval by placing

slides in a Coplin jar with 10% sodium citrate buffer in 1x phosphate-buffered saline (PBS). Coplin

jars were placed in a pressure cooker with 300 mL of tap water and set on high pressure for 15

minutes. The slides were let to cool for 20 minutes and then washed with diH$_2$O. They were then

blocked using 50 mg/mL bovine serum albumin (BSA) in 1x PBS. They were then washed with

.1% Tween 20 in 1x PBS (PBS-T). Primary antibodies were diluted based on manufacturers'

recommendation in 5% BSA and left to incubate on slides at 4°C, overnight. A list of antibodies

and their dilutions are below. Slides were washed the next day with PBS-T. For unconjugated

primary antibodies, slides were washed with three times with PBS-T for 2 minutes and incubated

in 1:5000 <u>secondary antibody</u> in PBS for 1 hour at room temperature. Slides were washed were

washed three times with PBS-T and counterstained by a 1:500 Dapi in 1x PBS solution for 10

minutes. Slides were washed with PBS-T for 2 minutes and nuclear staining was done using

1:2000 DAPI (Thermo Scientific, Ref:62248). Slides were mounted using 10% glycerol in 1x PBS.

Coverslips were placed over top of the slides, and the edges were coated with nail polish to
prevent dehydration. Slides were imaged using the Keyence BZ-X800 fluorescence microscope.
718

Image preprocessing and quantification of colocalization
Images of cells in tissue all underwent the following preprocessing before further analysis. Each
image contains multiple channels. The maximum intensity projection of each channel was then
generated. The projected image was the final image used for cellular counting analysis. For the
DAPI channel, the maximum intensity projection was generated from the raw image. For
visualization consistency, brightness and contrast of each fluorescent channel were adjusted
uniformly across all images using the "Brightness/Contrast" tool without altering the underlying
pixel intensity values. To measure intercellular colocalization, the straight-line tool was used to
draw a line between the centers of identified cells of interest (e.g., nuclei or marker-specific
regions). The length of each line was recorded using the "Measure" function, which reports
distances in calibrated units (µm) based on the microscope's pixel-to-micrometer conversion. At
least 629 cells per field were analyzed, and distances were exported for downstream statistical
analysis. Previous steps were conducted in ImageJ/Fiji (National Institutes of Health). Statistical
significance was calculated using a regular two-way ANOVA in *BioRender Graph*. Statistical
levels are provided as follows: ns for $0.05 < p \leq 1$, * for $0.01 < p \leq 0.05$, ** for $0.001 < p \leq 0.01$, ***
for $0.0001 < p < 0.001$, and **** for $p \leq 0.0001$.

Code Availability
The code is available at https://github.com/anurendra/CellWHISPER.

Data Availability
Mouse brain STEREO-seq data[28] were obtained from the Spatial Omics Database[95]. Our iPSC-
derived and mouse brain immunostaining data will be made available upon publication. Xenium
TgCRND8 Alzheimer's-disease mouse-model data were obtained from wild-type and *TgCRND8*
mice[86], as provided by 10x Genomics. Only the final time point (age > 13 months) was used for
analysis.

786   **REFERENCES**

787
788
1.   Armingol, E., Officer, A., Harismendy, O. & Lewis, N.E. Deciphering cell–cell interactions
and communication from gene expression. *Nature Reviews Genetics* **22**, 71-88 (2021).
2.   Wilk, A.J., Shalek, A.K., Holmes, S. & Blish, C.A. Comparative analysis of cell–cell
communication at single-cell resolution. *Nature biotechnology* **42**, 470-483 (2024).
3.   Dimitrov, D. *et al.* LIANA+ provides an all-in-one framework for cell–cell communication
inference. *Nature Cell Biology* **26**, 1613-1622 (2024).
4.   Dimitrov, D. *et al.* Comparison of methods and resources for cell-cell communication
inference from single-cell RNA-Seq data. *Nature communications* **13**, 3224 (2022).
5.   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature*
communications **12**, 1088 (2021).
6.   Efremova, M., Vento-Tormo, M., Teichmann, S.A. & Vento-Tormo, R. CellPhoneDB:
inferring cell–cell communication from combined expression of multi-subunit ligand–
receptor complexes. *Nature protocols* **15**, 1484-1506 (2020).
7.   Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication
by linking ligands to target genes. *Nature methods* **17**, 159-162 (2020).
8.   Hervé, J.-C. & Derangeon, M. Gap-junction-mediated cell-to-cell communication. *Cell*
and tissue research **352**, 21-31 (2013).
9.   Goodenough, D.A. & Paul, D.L. Gap junctions. *Cold Spring Harbor perspectives in*
biology **1**, a002576 (2009).
10.   Battefeld, A., Klooster, J. & Kole, M.H. Myelinating satellite oligodendrocytes are
integrated in a glial syncytium constraining neuronal high-frequency activity. *Nature*
Communications **7**, 1-13 (2016).
11.   Rash, J.E. Molecular disruptions of the panglial syncytium block potassium siphoning
and axonal saltatory conduction: pertinence to neuromyelitis optica and other
demyelinating diseases of the central nervous system. *Neuroscience* **168**, 982-1008
(2010).
12.   Glen, C.M., McDevitt, T.C. & Kemp, M.L. Dynamic intercellular transport modulates the
spatial patterning of differentiation during early neural commitment. *Nature*
communications **9**, 4111 (2018).
13.   Harris, A. & Locke, D. *Connexins: a guide*, (Springer Science & Business Media, 2008).
14.   Nakase, T. & Naus, C.C. Gap junctions and neurological disorders of the central nervous
system. *Biochimica Et Biophysica Acta (BBA)-Biomembranes* **1662**, 149-158 (2004).
15.   Jongsma, H.J. & Wilders, R. Gap junctions in cardiovascular disease. *Circulation*
research **86**, 1193-1197 (2000).
16.   Koval, M., Molina, S.A. & Burt, J.M. Mix and match: investigating heteromeric and
heterotypic gap junction channels in model systems and native tissues. *FEBS letters*
588, 1193-1204 (2014).
17.   Stephan, J., Eitelmann, S. & Zhou, M. Approaches to study gap junctional coupling.
Frontiers in Cellular Neuroscience **15**, 640406 (2021).
18.   Zhou, B. *et al.* Notch signaling pathway: architecture, disease, and therapeutics. *Signal*
transduction and targeted therapy **7**, 95 (2022).
19.   Kania, A. & Klein, R. Mechanisms of ephrin–Eph signalling in development, physiology
and disease. *Nature reviews Molecular cell biology* **17**, 240-256 (2016).

20. Nakandakari-Higa, S. *et al.* Universal recording of immune cell interactions in vivo.
Nature **627**, 399-406 (2024).
21. Mattes, B. & Scholpp, S. Emerging role of contact-mediated cell communication in tissue
development and diseases. *Histochemistry and cell biology* **150**, 431-442 (2018).
22. Jin, S., Plikus, M.V. & Nie, Q. CellChat for systematic analysis of cell–cell
communication from single-cell transcriptomics. *Nature Protocols* **20**, 180-219 (2025).
23. Troulé, K. *et al.* CellPhoneDB v5: inferring cell–cell communication from single-cell
multiomics data. *Nature Protocols*, 1-29 (2025).
24. Shao, X. *et al.* Knowledge-graph-based cell-cell communication inference for spatially
resolved transcriptomic data with SpaTalk. *Nature communications* **13**, 4429 (2022).
25. Cang, Z. *et al.* Screening cell–cell communication in spatial transcriptomics via collective
optimal transport. *Nature methods* **20**, 218-228 (2023).
26. Cesaro, G. *et al.* Advances and challenges in cell–cell communication inference: a
comprehensive review of tools, resources, and future directions. *Briefings in*
Bioinformatics **26**, bbaf280 (2025).
27. Zohora, F.T. *et al.* NEST: Spatially-mapped cell-cell communication patterns using a
deep learning-based attention mechanism. *bioRxiv*, 2024.03. 19.585796 (2024).
28. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA
nanoball-patterned arrays. *Cell* **185**, 1777-1792. e21 (2022).
29. Evans, W.H. & Martin, P.E. Gap junctions: structure and function. *Molecular membrane*
biology **19**, 121-136 (2002).
30. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nature methods*
19, 171-178 (2022).
31. Takeuchi, H. & Suzumura, A. Gap junctions and hemichannels composed of connexins:
potential therapeutic targets for neurodegenerative diseases. *Frontiers in cellular*
neuroscience **8**, 189 (2014).
32. Söhl, G., Maxeiner, S. & Willecke, K. Expression and functions of neuronal gap
junctions. *Nature reviews neuroscience* **6**, 191-200 (2005).
33. Dere, E. & Zlomuzica, A. The role of gap junctions in the brain in health and disease.
Neuroscience & Biobehavioral Reviews **36**, 206-217 (2012).
34. Dong, A., Liu, S. & Li, Y. Gap junctions in the nervous system: probing functional
connections using new imaging approaches. *Frontiers in Cellular Neuroscience* **12**, 320
(2018).
35. Kajiwara, Y. *et al.* GJA1 (connexin43) is a key regulator of Alzheimer's disease
pathogenesis. *Acta neuropathologica communications* **6**, 1-20 (2018).
36. Sánchez, O.F. *et al.* Role of connexins 30, 36, and 43 in brain tumors,
neurodegenerative diseases, and neuroprotection. *Cells* **9**, 846 (2020).
37. Nualart-Marti, A., Solsona, C. & Fields, R.D. Gap junction communication in myelinating
glia. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **1828**, 69-78 (2013).
38. Papaneophytou, C., Georgiou, E. & Kleopa, K.A. The role of oligodendrocyte gap
junctions in neuroinflammation. *Channels* **13**, 247-263 (2019).
39. Nagy, J.I., Ionescu, A.V., Lynn, B.D. & Rash, J.E. Connexin29 and connexin32 at
oligodendrocyte and astrocyte gap junctions and in myelin of the mouse central nervous
system. *Journal of Comparative Neurology* **464**, 356-370 (2003).
40. Kleopa, K.A., Orthmann, J.L., Enriquez, A., Paul, D.L. & Scherer, S.S. Unique
distributions of the gap junction proteins connexin29, connexin32, and connexin47 in
oligodendrocytes. *Glia* **47**, 346-357 (2004).
41. Orthmann-Murphy, J.L., Abrams, C.K. & Scherer, S.S. Gap junctions couple astrocytes
and oligodendrocytes. *Journal of Molecular Neuroscience* **35**, 101-116 (2008).
42. Xing, L., Yang, T., Cui, S. & Chen, G. Connexin hemichannels in astrocytes: role in CNS
disorders. *Frontiers in molecular neuroscience* **12**, 23 (2019).

43. Li, M. *et al.* Astrocytic Gap Junctions protein Cx43/Cx30 modulate EAAT1 and glutamate
to mediate cerebral ischemia–reperfusion injury. *Brain Research* **1847**, 149306 (2025).

44. Nagy, J., Ionescu, A.V., Lynn, B. & Rash, J. Coupling of astrocyte connexins Cx26,
Cx30, Cx43 to oligodendrocyte Cx29, Cx32, Cx47: Implications from normal and
connexin32 knockout mice. *Glia* **44**, 205-218 (2003).

45. Orthmann-Murphy, J.L., Freidin, M., Fischer, E., Scherer, S.S. & Abrams, C.K. Two
distinct heterotypic channels mediate gap junction coupling between astrocyte and
oligodendrocyte connexins. *Journal of Neuroscience* **27**, 13949-13957 (2007).

46. Alvarez-Maubecin, V., García-Hernández, F., Williams, J.T. & Van Bockstaele, E.J.
Functional coupling between neurons and glia. *Journal of Neuroscience* **20**, 4091-4098
(2000).

47. Srinivas, M. *et al.* Functional properties of channels formed by the neuronal gap junction
protein connexin36. *Journal of Neuroscience* **19**, 9848-9855 (1999).

48. Lee, S.-N. *et al.* Cryo-EM structures of human Cx36/GJD2 neuronal gap junction
channel. *Nature communications* **14**, 1347 (2023).

49. Nagy, J. & Rash, J. Cx36, Cx43 and Cx45 in mouse and rat cerebellar cortex: Species-
specific expression, compensation in Cx36 null mice and co-localization in neurons vs.
glia. *European Journal of Neuroscience* **46**, 1790-1804 (2017).

50. Feigenspan, A. *et al.* Expression of connexin36 in cone pedicles and OFF-cone bipolar
cells of the mouse retina. *Journal of Neuroscience* **24**, 3325-3334 (2004).

51. Van Bockstaele, E., Garcia-Hernandez, F., Fox, K., Alvarez, V. & Williams, J. Expression
of connexins during development and following manipulation of afferent input in the rat
locus coeruleus. *Neurochemistry international* **45**, 421-428 (2004).

52. Kay, C.W., Ursu, D., Sher, E. & King, A.E. The role of Cx36 and Cx43 in 4-
aminopyridine-induced rhythmic activity in the spinal nociceptive dorsal horn: an
electrophysiological study in vitro. *Physiological Reports* **4**, e12852 (2016).

53. Zhou, M., Zhong, S. & Verkhratsky, A. Astrocyte syncytium: from neonatal genesis to
aging degeneration. *Neural regeneration research* **19**, 395-396 (2024).

54. Li, Z.A. & Zhou, M. Astrocyte syncytium–a biopower grid system in the brain. *Journal of
integrative neuroscience* **23**, 71 (2024).

55. Ma, B. *et al.* Gap junction coupling confers isopotentiality on astrocyte syncytium. *Glia*
64, 214-226 (2016).

56. Serra, R. & Simard, J.M. Adherens, tight, and gap junctions in ependymal cells: A
systematic review of their contribution to CSF-brain barrier. *Frontiers in neurology* **14**,
1092205 (2023).

57. Fabbiani, G. *et al.* Connexin signaling is involved in the reactivation of a latent stem cell
niche after spinal cord injury. *Journal of Neuroscience* **40**, 2246-2258 (2020).

58. Bouillé, C., Mesnil, M., Barriere, H. & Gabrion, J. Gap junctional intercellular
communication between cultured ependymal cells, revealed by lucifer yellow CH transfer
and freeze-fracture. *Glia* **4**, 25-36 (1991).

59. De Bock, M. *et al.* Targeting gliovascular connexins prevents inflammatory blood-brain
barrier leakage and astrogliosis. *Jci Insight* **7**, e135263 (2022).

60. Jiang, H., Zhang, Y., Wang, Z.-Z. & Chen, N.-H. Connexin 43: an interface connecting
neuroinflammation to depression. *Molecules* **28**, 1820 (2023).

61. Johnson, A.M. *et al.* Connexin 43 gap junctions contribute to brain endothelial barrier
hyperpermeability in familial cerebral cavernous malformations type III by modulating
tight junction structure. *The FASEB Journal* **32**, 2615 (2018).

62. Eugenín, E.A. *et al.* Microglia at brain stab wounds express connexin 43 and in vitro
form functional gap junctions after treatment with interferon-γ and tumor necrosis factor-
α. *Proceedings of the National Academy of Sciences* **98**, 4190-4195 (2001).

63. Nakase, T., Fushiki, S. & Naus, C.C. Astrocytic gap junctions composed of connexin 43
reduce apoptotic neuronal damage in cerebral ischemia. *Stroke* **34**, 1987-1993 (2003).

64. Abudara, V. *et al.* Activated microglia impairs neuroglial interaction by opening C x43
hemichannels in hippocampal astrocytes. *Glia* **63**, 795-811 (2015).

65. Hawrylycz, M. *et al.* The allen brain atlas. *Springer handbook of bio-/neuroinformatics*,
1111-1126 (2014).

66. Henze, D.A., Urban, N.N. & Barrionuevo, G. The multifarious hippocampal mossy fiber
pathway: a review. *Neuroscience* **98**, 407-427 (2000).

67. Triplett, J.W. & Feldheim, D.A. Eph and ephrin signaling in the formation of topographic
maps. in *Seminars in cell & developmental biology* Vol. 23 7-15 (Elsevier, 2012).

68. Yang, J.S., Wei, H.X., Chen, P.P. & Wu, G. Roles of Eph/ephrin bidirectional signaling in
central nervous system injury and recovery. *Experimental and therapeutic medicine* **15**,
2219-2227 (2018).

69. Martinelli, D.C. *et al.* Expression of C1ql3 in discrete neuronal populations controls
efferent synapse numbers and diverse behaviors. *Neuron* **91**, 1034-1051 (2016).

70. Newrzella, D. *et al.* The functional genome of CA1 and CA3 neurons under native
conditions and in response to ischemia. *BMC genomics* **8**, 1-18 (2007).

71. Quarles, R.H. Myelin-associated glycoprotein (MAG): past, present and beyond. *Journal*
of neurochemistry **100**, 1431-1448 (2007).

72. Gow, A. *et al.* CNS myelin and sertoli cell tight junction strands are absent in
Osp/claudin-11 null mice. *Cell* **99**, 649-659 (1999).

73. Hutchison, M. *et al.* Genetic inhibition of neurotransmission reveals role of glutamatergic
input to dopamine neurons in high-effort behavior. *Molecular psychiatry* **23**, 1213-1225
(2018).

74. Song, Y.-H., Yoon, J. & Lee, S.-H. The role of neuropeptide somatostatin in the brain
and its application in treating neurological disorders. *Experimental & Molecular Medicine*
53, 328-338 (2021).

75. Vandecasteele, M., Glowinski, J. & Venance, L. Electrical synapses between
dopaminergic neurons of the substantia nigra pars compacta. *Journal of Neuroscience*
25, 291-298 (2005).

76. Radnikow, G. & Feldmeyer, D. Layer-and cell type-specific modulation of excitatory
neuronal activity in the neocortex. *Frontiers in neuroanatomy* **12**, 1 (2018).

77. Yan, J. & Gao, L. Organizing principles of the prefrontal cortex from a single-neuron
projectome. Vol. 25 408-409 (NATURE PORTFOLIO HEIDELBERGER PLATZ 3,
BERLIN, 14197, GERMANY, 2022).

78. Gulledge, A.T. & Kawaguchi, Y. Phasic cholinergic signaling in the hippocampus:
functional homology with the neocortex? *Hippocampus* **17**, 327-332 (2007).

79. Sherman, S.M. Thalamus plays a central role in ongoing cortical functioning. *Nature*
neuroscience **19**, 533-541 (2016).

80. Yuasa-Kawada, J., Kinoshita-Kawada, M., Tsuboi, Y. & Wu, J.Y. Neuronal guidance
genes in health and diseases. *Protein & cell* **14**, 238-261 (2023).

81. Südhof, T.C. Synaptic neurexin complexes: a molecular code for the logic of neural
circuits. *Cell* **171**, 745-769 (2017).

82. Lv, X. *et al.* Patterned cPCDH expression regulates the fine organization of the
neocortex. *Nature* **612**, 503-511 (2022).

83. Angeli, S. *et al.* Altered expression of glial gap junction proteins Cx43, Cx30, and Cx47
in the 5XFAD model of Alzheimer's disease. *Frontiers in Neuroscience* **14**, 582934
(2020).

84. Maulik, M. *et al.* Amyloid-β regulates gap junction protein connexin 43 trafficking in
cultured primary astrocytes. *Journal of Biological Chemistry* **295**, 15097-15111 (2020).

85.   Pechlivanidou, M. *et al.* Glial gap junction pathology in the spinal cord of the 5xFAD
mouse model of early-onset Alzheimer's disease. *International journal of molecular*
sciences **23**, 15597 (2022).

86.   Genomics, x. Xenium In Situ Analysis of Alzheimer's Disease Mouse Model Brain
Coronal Sections from One Hemisphere Over a Time Course. (2024).

87.   Yi, C. *et al.* Astroglial connexin43 contributes to neuronal suffering in a mouse model of
Alzheimer's disease. *Cell Death & Differentiation* **23**, 1691-1701 (2016).

88.   Kozberg, M.G. *et al.* Blood–brain barrier leakage and perivascular inflammation in
cerebral amyloid angiopathy. *Brain Communications* **4**, fcac245 (2022).

89.   Figueroa, X.F. & Duling, B.R. Gap junctions in the control of vascular function.
Antioxidants & redox signaling **11**, 251-266 (2009).

90.   Huang, X. *et al.* Astroglial connexins in neurodegenerative diseases. *Frontiers in*
molecular neuroscience **14**, 657514 (2021).

91.   Caruso, G., Di Pietro, L. & Caraci, F. Gap junctions and connexins in microglia-related
oxidative stress and neuroinflammation: perspectives for drug discovery. *Biomolecules*
13, 505 (2023).

92.   Anwar, M.M. *et al.* Assessing the role of primary healthy microglia and gap junction
blocker in hindering Alzheimer's disease neuroinflammatory type: early approaches for
therapeutic intervention. *Frontiers in Neuroscience* **16**, 1041461 (2023).

93.   Matoba, K. *et al.* FcγR-and CD9-dependent synapse engulfing microglia in the thalamus
drives cognitive impairment following cortical brain injury. *bioRxiv*, 2024.09. 19.609743
(2024).

94.   Yu, H. *et al.* Segmentation Matters: Cell Segmentation Challenge in Spatial
Transcriptomics. *NEURON-D-25-01846*.

95.   Yuan, Z. *et al.* SODB facilitates comprehensive exploration of spatial omics data. *Nature*
Methods **20**, 387-399 (2023).

96.   Stuart, T. *et al.* Comprehensive integration of single-cell data. *cell* **177**, 1888-1902. e21
(2019).

1011