Transient species can increase resident metapopulation size by modifying
local conditions and regional connectivity

Manuscript elements: Table 1, Figure 1, Figure 2, Figure 3, Figure 4, Figure 5, Supplementary
material (Table S1, Table S2, Figure S1, Figure S2, Figure S3, Figure S4)

Word count: Main text (5720)

Keywords: asymmetric dispersal, metapopulation size, source-sink dynamics, ideal free
distribution

Manuscript type: Major article

Abstract

Populations are regulated by processes operating at both local and regional scales. As a result, species can interact not only within local communities but also through regional-scale processes. However, the feedback between local and regional processes of species interactions in shaping metapopulation dynamics have rarely been tested experimentally. Using nectar microbes as a model system, we examine how a bacterial species dispersing to a sink habitat, sustained by external supply, can interact with a resident yeast species directly by modifying local nectar chemistry and indirectly by altering metapopulation connectivity. We show that the external supply of bacteria increases resident yeast densities, potentially through the provision of resource subsidies, and that this benefit is enhanced when the dispersal rate of yeast is reduced in patches supplied with bacteria. Analysis of a metapopulation model indicates that the benefit of dispersal modification on metapopulation size occurs only when resource availability differs across patches–supporting our interpretation that bacterial supply provides resource subsidies. Taken together, our results underscore the role of dispersal in shaping how metapopulations experience density-dependence across heterogeneous landscapes, highlight the relevance of material flows across ecosystem boundaries, and suggest a plausible mechanism in which transient species may leave lasting impacts on ecological communities.

INTRODUCTION

Many natural populations are spatially subdivided into a network of local populations connected by

dispersal, *i.e.*, a metapopulation. Although originally introduced as a spatially implicit model that tracks

the number (or fraction) of patches occupied by a single species, metapopulation theory has developed

into a more general theory that can accommodate multiple species, explicit local dynamics, and spatial

structure (e.g., Hanski, 1998; Tilman, 1994; Leibold et al. 2004). Decades of theoretical work has shown

that dispersal does more than merely shuffle individuals across space (Hastings, 1982; Holt, 1985; Karlin

& McGregor, 1972; Levin, 1974; Levins, 1969). Even under the simplest scenarios, dispersal of

organisms can have non-intuitive effects on local and regional patterns of biodiversity (Hanski, 1998;

Huffaker, 1958; Levin, 1974; Levins, 1969). For instance, dispersal allows metapopulations to persist

through patch dynamics even when local populations are certain to go extinct (Brown & Kodric-Brown,

1977). Dispersal can even enable metapopulations to persist when average local growth rates are negative,

as occasional periods of positive growth inflate overall population size (Kortessis et al., 2025; Matthews

& Gonzalez, 2007). Furthermore, dispersal can stabilize or destabilize local populations (Bascompte &

Solé, 1994; Doebeli, 1995; Huffaker, 1958; Pascual, 1993) or allow intePracting species (or genotypes) to

coexist (Hassell et al., 1991; Karlin & McGregor, 1972; Levin, 1974; Solé et al., 1992). Recent work

demonstrates that dispersal can even increase the total number of individuals across the metapopulation

(i.e., metapopulation size) relative to the sum of its local carrying capacities without dispersal if local

conditions are spatially heterogeneous (Arditi et al., 2015; DeAngelis et al., 2016; He et al., 2025; Zhang

et al., 2017). In sum, these results indicate that the dynamics of a regional metapopulation are rarely the

simple sum of its constituent populations' dynamics considered in isolation and instead often depend on

how dispersal links them.

Despite progress made in understanding how dispersal structures ecological communities,

relatively less is known about the ecological consequences when the dynamics of dispersal is itself

coupled to local dynamics. In natural ecological settings, non-random (directed or biased) dispersal is

more likely the norm than the exception (Amarasekare, 2010; Clobert et al., 2009; McPeek et al., 2024;

Vuilleumier & Possingham, 2006). While directionality in movement can arise naturally via physical

forces such as wind, currents, and gravity (Germain et al., 2019; Mari et al., 2014; Okubo & Levin, 2001),

it can also result from biotic mechanisms such as habitat selection (Clobert et al., 2009). Although it

might seem that directed dispersal via habitat selection is unique to mobile animals with cognitive

capacities, organisms with limited motility can also disperse non-randomly via mechanisms such as

phoresy (or zoochory in plants), where their interactions with animal dispersal vectors can dictate both the

rate and direction of movement (Bartlow & Agosta, 2021).Given that dispersal can modify population

size and that species can influence each other's dispersal, a natural question to ask is whether species

interactions can be mediated from purely regional processes. For example, what happens when multiple

phoretic species share the same vectors? Could shared dispersal vectors mediate interactions between

phoretic organisms? If vectors are limited, could competition for them shape the relative abundances of

phoretic species at the metapopulation level (Nell et al., 2025)? To date, few studies have tackled these

questions, even though doing so could yield important insights into metapopulation dynamics in natural

ecological settings.

Recognizing that dispersal can be an integral component of how species interact could reveal

cryptic interactions that shape ecological communities. This possibility is particularly relevant for species

that form sink populations—species whose persistence is determined less by local conditions than by

immigration from adjacent habitats. Because such species cannot maintain growth locally, their potential

impacts on 'resident' species—i.e., species capable of persisting without immigration—are often

overlooked (Amor et al., 2020; Mallon et al., 2018). However, previous studies suggest that sink

populations are not only prevalent but can also impact resident species (Amarasekare & Nisbet, 2001;

Holt et al., 2003; Long et al., 2007; Mallon et al., 2018; Pulliam, 1988). The primary mechanism by

which this impact occurs is through the sheer abundance by which immigration introduces the species

into a local community, amplifying their total impact despite their otherwise small per-capita effects on

residents. For instance, a species occupying a sink habitat can reach sufficiently high densities via

immigration and deplete resources to levels that depress the abundance of, or even competitively exclude,

an otherwise locally superior competitor (Amarasekare & Nisbet, 2001; Long et al., 2007). Notably, these

studies assume that sink populations impact residents purely through local interactions. However, if

dispersal can be shaped by other species, sink populations may also influence residents by altering their

dispersal patterns across the metapopulation. Despite evidence showing that shared dispersal vectors can

reshape dispersal patterns among competing species (e.g., Muller-Landau et al., 2008; Xiao & Zhang,

2016), their long-term demographic consequences remain unknown.

A system in which dispersal vectors modulate species interactions is the floral microbiome,

making it an excellent model system for examining the potential feedback between local and regional

processes. Floral nectar often contains microbial communities (Vannette, 2020) that are dispersed by

pollinators (Belisle et al., 2012; Herrera et al., 2009; Vannette & Fukami, 2017). The identity of

pollinators can be a strong determinant of microbial composition, suggesting that their abundance and

foraging patterns can shape the spatial distribution of microbial communities in floral nectar (Belisle et

al., 2012; Morris et al., 2020; Vannette et al., 2021; Vannette & Fukami, 2017). As is true for many

microbial systems (Mallon et al., 2018), floral nectar is likely to contain many transient invaders given

that nectar microbes are often a subset of the microbial communities found on pollinators (Belisle et al.,

2012; Herrera et al., 2009; Vannette & Fukami, 2017). Importantly, the microbial activity within nectar

can alter pollinator foraging behavior by modifying nectar chemistry or producing volatiles that attract or

repel certain pollinators (Chappell et al., 2022; Schaeffer et al., 2019; Vannette et al., 2013). Local

modification of nectar chemistry may thus bias dispersal towards or away from certain flowers. In other

words, these microbes participate in their own dispersal despite not being able to move on their own.

Considered together, these processes create a feedback loop between local dynamics and regional

dispersal via pollinator foraging (Nell et al., 2025), providing an additional pathway in which sink

populations could interact with residents at the metapopulation level.

Here, using nectar microbes isolated from the hummingbird-pollinated sticky monkeyflower in

California (*Diplacus aurantiacus*), we conducted a laboratory experiment to investigate how a bacterial

species invading a sink habitat can impact the metapopulation dynamics of a resident yeast species. To

avoid ambiguity, we refer to dispersal from external sources as invasion (or supply) and reserve dispersal

for movement among patches within the metapopulation (external vs. internal dispersal sensu Fukami

2005). Previous work in this system has shown that the bacteria species reduces nectar pH, which in turn

suppresses yeast growth and reduces pollinators' nectar consumption (Chappell et al., 2022). In a pilot

experiment, we found that the bacteria species cannot sustain a viable population under our experimental

conditions but can nevertheless reduce nectar pH, making it well suited for examining the effects of

species invading a sink habitat on resident yeast species. Specifically, we ask how the yeast

metapopulation responds to changes in local nectar chemistry (i.e., reduced pH) and patch (*i.e.*, local

population) connectivity mediated by pollinator foraging preferences in the presence of bacterial sink

populations. First, we test the hypothesis that reduction of nectar pH by the external supply of bacteria

reduces yeast metapopulation size by modifying nectar pH to the detriment of yeast population growth

(*i.e.*, local effects). Second, we test the hypothesis that pollinator preference biases dispersal away from

low-pH patches, thus altering metapopulation size of yeast by altering its metapopulation connectivity

(*i.e.*, local-regional interaction effects). Finally, we analyze a simple two-patch metapopulation model to

verify and complement inferences drawn from our experimental results.

METHODS

 Overview of study system

To investigate the impact of a species invading a sink habitat on the metapopulation dynamics of a

resident species, we conducted experiments using two nectar microbes: the bacterium *Acinetobacter*

nectaris (strain FNA17) and the yeast *Metschnikowia reukaufii* (strain MR1). These microbes naturally

inhabit the floral nectar of the sticky monkeyflower, *Diplacus aurantiacus,* and the specific strains we

used were isolated from flowers in Jasper Ridge Biological Preserve ('Ootchamin 'Ooyakma) located in

the Santa Cruz Mountains of California. *Diplacus aurantaicaus* is a shrub native to coastal Oregon and

California and is commonly pollinated by hummingbirds (Belisle et al., 2012). Instead of dispersing

between flowers themselves, these microbes "hitchhike" onto pollinators to move between flowers. When

microbes arrive at a new flower, they proliferate by metabolising nectar resources and in so doing, modify

nectar chemistry and set in motion the diverse ecological interactions that occur in these microscopic

ecosystems (e.g., microbe-microbe interactions, plant-pollinator mutualisms; Chappell & Fukami, 2018;

Quevedo-Caraballo et al., 2025). Its relative simplicity and tractability allows this study system to be

replicated in laboratory settings with artificial nectar in microwell plates as flowers and pipettes to

simulate pollinator-assisted dispersal (Chappell et al., 2022; Grainger et al., 2019; Letten et al., 2018;

Tucker & Fukami, 2014; Vannette & Fukami, 2014). Following these prior studies, we adapted this

experimental microcosm for the purpose of our study.

Experimental setup

We conducted an experiment that factorially manipulated 'bacterial supply' (i.e., the number of habitat

patches in each metapopulation that would receive bacteria from an external source, dispersal bias, and

dispersal frequency (i.e., the number of dispersal events within a growth cycle) to test how *A. nectaris*

bacteria-driven pH changes affect *M. reukaufii* yeast metapopulation by modifying local nectar conditions

and metapopulation connectivity through dispersal bias (Fig. 1). Note that our hypothesis is concerned

with how patches are wired (*i.e.*, dispersal bias) and not the overall frequency of dispersal events *per se*.

Thus, we included two levels of dispersal frequency to ensure a meaningful difference between the levels

of the dispersal bias treatment is realized, rather than being a variable that is itself of interest. Our

experimental metapopulations consisted of eight wells (i.e., patches) on a 96-well PCR plate, connected

by dispersal. Each metapopulation was assigned to one of 12 treatment groups defined by the fully

factorial combination (Fig. 1A) of bacterial supply (three levels: 0, 3, or 6), dispersal bias (two levels:

random or bias), and dispersal frequency (two levels: 3 or 6), with each treatment replicated six times (12

treatments x 6 replicates = 72 metapopulations). Each replicate metapopulation was initiated by seeding

patches with approximately 100-150 viable cells of *M. reukaufii* (see Supplementary Methods) and 100 μl

of sterile artificial nectar. Artificial nectar was prepared as specified by Chappell et al. (2022) and the full

recipe is provided in Supplementary Methods. Plates were sealed with a sterilized air-permeable

membrane and incubated at 26°C for two days before beginning the experiment. After initiation, each

metapopulation underwent a total of five two-day growth-dilution-cycles where each cycle included the

following order of events: 1) pH sampling + dispersal, 2) resource replenishment, 3) community census,

4) bacterial supply, and 5) a two-day incubation period (Fig. 1B):

Event 1: Dispersal. To simulate pollinator-assisted dispersal, we first sampled the pH of local patches,

generated a sequence of dispersal events, then pipetted nectar between patches according to the order of

generated sequences. Specifically, for the bias treatment, 10 μl of nectar from each patch were pipetted

onto pH indicator strips. Once pH readings developed, we recorded the pH values and used them as

weights (representing pollinator preferences) to generate a new sequence of dispersal events between

donor-recipient pairs. Specifically, donor and recipient patches were selected by sampling two patches

without replacement, with a weighted distribution over the eight patches (A–H), where selection

probabilities are proportional to their respective pH values. The weight of each patch (ω_i) was calculated

using the Monod equation with a half-saturation constant set at 3, which is a pH value that lies between

nectar conditioned by bacteria (pH = 2) and yeast (pH = 4; Chappell et al 2022). More precisely, the

weight of patch i is $\omega_i = pH_i / (3 + pH_i)$ and the probability of selecting it is obtained by normalizing

these weights, $P_i = \omega_i / \sum_{j=1}^{8} \omega_j$. This setup reflects the field observation that low-pH patches are

less likely to be visited by pollinators and, consequently, microbes are less likely to disperse into or out of

them (Vannette et al. 2013; Chappell et al. 2022). For the random treatment, the pH sampling step was

skipped (but still discarding 10 µl of nectar from each patch) because dispersal did not depend on pH.

Instead, dispersal sequences from random dispersal treatments were done in a similar manner except that

the probabilities were always equal across patches (*i.e.*, independent of pH); the sequence for background

dispersal was identical for all 72 metapopulations in the random dispersal treatment. For both the biased

and random dispersal treatment, each dispersal event was achieved by pipetting 3 µl of nectar from a

donor patch to a recipient patch. Background dispersal was conducted to ensure that patches within a

replicate metapopulation are connected within the experimental duration. The number of dispersal events

per growth cycle was determined by the dispersal frequency treatment, either 3 or 6, which was

manipulated factorially with dispersal bias. In other words, each replicate underwent 3 or 6 sequential

dispersal events (in addition to background dispersal) within a growth-dilution cycle.

Event 2: Resource replenishment. Once the dispersal events were completed, all patches were diluted

by transferring 5 µl of nectar into 95 µl of fresh, sterilized media on new 96-well plates to simulate

resource replenishment (i.e., nectar production in flowers). We used new sterile 96-well plates, instead of

removing and adding nectar from the original, to minimize contamination.

Event 3: Sampling. Immediately after dilution, we sampled each local community by plating 5 µl of the

newly diluted nectar onto yeast malt agar (YMA) for yeast and 5 µl onto tryptic soy agar (TSA) for

bacteria, and then enumerated the resulting colony-forming units (CFUs) after >3 days of incubation at

room temperature as a measure of microbial density (following Peay et al., 2012 and Vannette et al.,

2013). Due to time constraints associated with simultaneously conducting the experiment and counting

CFUs, we took images of the agar plates and enumerated CFUs after completion of the experiment (see

Appendix for sample images). In a few cases, colonies were too numerous to count so we set their counts

to 200, a value just above the maximum from samples that were countable. To ensure this artificial cutoff

did not qualitatively affect our results, we repeated our analyses with these samples removed (see

'Potential artifacts of experimental design' below).

Event 4: Bacterial supply. Following sampling, bacterial cells were added to patches according to the

bacterial supply treatment. Specifically, bacterial cells were either added to no patches (bacterial supply =

0), patches A to C (bacterial supply = 3), or patches A to F (bacterial supply = 6). We chose to add

bacteria to only a subset of patches within the metapopulation because this allows spatial heterogeneities

in nectar pH and thus creates variation in dispersal probabilities under the dispersal bias treatment.

Although doing so is not required, we chose to fix the patches in which bacteria were added (as opposed

to randomly selecting patches to inoculate) to minimize experimental error. Approximately 20,000

bacterial cells (~200 cells/μl nectar) were inoculated to each recipient patch (see Supplementary

Methods). Bacterial cells were supplied after sampling because in our pilot trials, this amount of bacteria

was necessary to reduce pH (after 2 days) but would be too dense to count if plated directly after

inoculation. Thus, the bacteria that we sampled came from the remaining bacteria inoculated two days

prior.

Event 5: Incubation. Following bacterial supply, 96-well plates were sealed with sterilized, air-

permeable membranes and incubated at 26°C for 48 hours before beginning the next cycle.

224   *Data analysis*

Yeast metapopulation size

To assess treatment effects on yeast metapopulation size, we defined metapopulation size as the sum of

local patch densities within a metapopulation replicate and modelled it as a response variable in a

negative binomial generalized linear mixed model (GLMM). This model included dispersal frequency

and bacterial supply as continuous predictors, dispersal bias as a categorical variable, and all two-way

interactions as fixed effects, with replicate and time as random effects. We included all two-way

interactions because they aligned with our hypotheses and excluded the three-way interaction term

because including it did not improve model fit. Because initial statistical fits with Poisson-distributed

errors showed violation of homoscedasticity, we implemented a forward model selection to account for

potential variance structure in our data. Specifically, we sequentially added treatment terms as predictors

to the dispersion component of the negative binomial GLMM and only kept terms that significantly

improved model fit (as evaluated by χ^2 tests; Table S1). This led to the selection of a model that included

bacterial supply as a significant predictor of the variance in yeast metapopulation size (Table 1; ΔAIC = -

34.8, P < 0.001) which satisfied the assumption of residual homoscedasticity. Furthermore, we also

assessed the correlation between local densities of yeast and bacteria to provide additional evidence for

the local effects of bacteria supply on yeast population size using the `cor.test()` function in base R.

Statistical models were fitted in R using the 'glmmTMB' and model assumptions were evaluated using

the 'DHARMa' package.

Potential artefacts of experimental design

We conducted additional tests to confirm that our experimental results were not contingent on the choices

of our experimental design. We describe three specific tests here in brief, and provide more detailed

descriptions in the Supplementary Methods, with results shown in Fig. S3 and Fig. S4. First, because a

few samples formed colonies that were too dense to count (35 samples out of 354 total; denoted 'Inf' in

our raw data), we included them in our statistical analyses by setting their counts to 200. To ensure that

this decision did not artificially bias our results, we refitted the same statistical models with these samples

removed and found no qualitative differences (Table S1-2; Fig. S2). Second, we verified the extent to

which our experimental treatments modified the dispersal in patches supplied with bacteria relative to

those without. For example, if treatments were successful, then bacteria supplied patches should be less

likely to be selected for dispersal relative to patches with no supply under the dispersal bias treatment. To

test this, we measured the differences in dispersal propensies (i.e., frequency that a patch type is selected

for dispersal) between dispersal treatments and tested whether these differences differed by patch type. A

GLMM revealed marginally significant difference in the dispersal propensities between patch types (P =

0.05; Fig S3). Lastly, we confirmed that the position of microplate wells had no effect on yeast densities

by comparing the densities between wells within replicate after controlling for treatment effects via

bootstrapping.

Metapopulation model

We complemented our experiment with the analysis of a two-patch metapopulation model. As described

in our Results, we unexpectedly found that the densities of the resident species *M. reukaufii* were

positively correlated with the densities of the bacterial species, *A. nectaris* (Fig. 3). Given this finding, we

explored the possibility that, in addition to affecting regional connectivity by modifying nectar pH,

immigration of *A. nectaris* could also impact the resident species locally by bringing in resource subsidies

that increase the local carrying capacity of yeast. To explore this possibility in a model and see if patterns

resemble those seen in our experiment, we analyzed a two-patch metapopulation model and asked how

dispersal bias (induced by pH modification) and local resource subsidies (via external supply of bacteria)

jointly affect yeast metapopulation size.

To simplify analysis, we did not explicitly model the densities of the bacterial species and simply

assumed that its effects are reflected in demographic and dispersal parameters of the resident yeast. Our

general model takes the form:

278
$$\frac{dN_B}{dt} = rN_B\left(1 - \frac{N_B}{\beta K}\right) - D(1 + \delta)N_B + D(1 - \delta)N_N$$

279
$$\frac{dN_N}{dt} = rN_N\left(1 - \frac{N_N}{K}\right) - D(1 - \delta)N_N + D(1 + \delta)N_B$$

where N_B and N_N are the densities of the resident yeast in patch B (bacteria-containing) and patch N (no-

bacteria), respectively. The parameters r and K represent the local intrinsic growth rate and baseline

carrying capacity (*i.e.*, the carrying capacity in the absence of enhancement by bacterial subsidies), while

D denotes the average per-capita dispersal rate across the metapopulation. The parameters β and δ control

asymmetries in carrying capacity and per-capita dispersal rate between the two types of patches.

Specifically, β represents the enhancement of carrying capacity in the bacteria-containing patch, whereas

δ represents asymmetry in per-capita dispersal rates between patches, with $\delta < 0$ indicating reduced patch

B per-capita dispersal rate relative to patch N (*i.e.*, dispersal bias treatment in experiment). Importantly, β

introduces asymmetry in carrying capacity between local patches while also increasing the total carrying

capacity at the metapopulation-level, whereas δ modifies the difference (or asymmetry) in per-capita

dispersal rates between patches without altering average per-capita dispersal rate across patches.

This parameterization mirrors our experimental setup. In the experiment, bacterial supply

increases the resources available in bacteria-containing patches (and therefore the metapopulation as a

whole) and thereby increases yeast density through demographic feedback. In contrast, the dispersal bias

treatment controls the likelihood that a bacteria-containing patch is selected for dispersal relative to a no-

bacteria patch. Specifically, because each bacteria-containing patch is sampled with weight determined by

its pH (*i.e.*, ω_B), the probability of sampling a bacteria-containing patch from the experimental

metapopulation is proportional to ω_B times the number of bacteria-containing patches (i.e., $\omega_B * n_B$),

which aligns with the $D(1 + \delta) * N_B$ term in the model. Accordingly, if bacteria do not reduce pH, then

$\omega_B = \omega_N$ (equivalent to the random dispersal treatment) and the probability of selecting a given patch

type for dispersal is proportional to the frequency of that type in the metapopulation (*i.e.*, determined

strictly by bacterial supply). By reducing the system to two patches, the model assumes that

metapopulation differences arise primarily from dispersal between patch types ($N_B \rightarrow N_N$ or $N_N \rightarrow N_B$) and

not within ($N_B \rightarrow N_B$ or $N_N \rightarrow N_N$) (see Discussion for limitations).

With this model, we evaluated how and examined why metapopulation size varies as a function

of parameter values. For simplicity and given the non-significant effect of dispersal frequency (see

Results), we assume $r = K = D = 1$, which reduces the number of free parameters to the two aligned with

our hypotheses: bacterial supply (β) and dispersal bias (δ). With this reduced model, we computed the

metapopulation size at varying levels of β *and* δ by solving for its steady-state solution numerically in R

and taking the sum of equilibrium densities (denoted N_B^* and N_N^*) across both patches. To obtain a

heuristic understanding of why metapopulation size varies at all, we derived analytically the condition

under which metapopulation size would be maximized under our model, as predicted by the theory of

ideal free distribution (IFD; (Fretwell & Lucas, 1969; Holt 1985). In brief, IFD predicts that mean fitness

(or average per-capita growth rate) is maximized when individuals distribute themselves such that fitness

is equalized across patches. At equilibrium, this corresponds to the condition where the density in each

patch equals its local carrying capacity, since any deviation would imply unequal fitness: one patch would

be underfilled (positive growth) while the other would be overfilled (negative growth). One trivial case in

which this condition holds is when $D = 0$. However, since we assume dispersal ($D = 1$), the remaining

condition that satisfies IFD occurs when the number of migrants in each patch cancels out: $(1 + \delta)N_B^* =$

$(1 - \delta)N_N^*$. By substituting the local carrying capacities β and 1 for the equilibrium densities N_B^* and N_N^*,

respectively, and re-arranging yields the condition: $\delta = (1 - \beta) / (1 + \beta)$. We verify that this condition

corresponds to the maximization of metapopulation size by comparing it to simulations.

325   **RESULTS**

327   *Effects of bacterial supply, dispersal bias, and dispersal frequency on yeast metapopulation size*

Bacterial supply significantly modified overall yeast metapopulation size ($\chi^2 = 6.103$, $P = 0.013$; Table 1;

Fig. 2&3a) and the magnitude of this effect depended on dispersal bias ($\chi^2 = 6.162$, $P=0.013$; Table 1).

Specifically, bacterial supply increased yeast metapopulation size under biased dispersal, but not under

random dispersal (Fig. 3c). In contrast, neither dispersal frequency ($\chi^2 = 1.053, P = 0.305$) nor dispersal

bias ($\chi^2 = 1.626, P = 0.202$) alone had significant main effects on yeast metapopulation size (Table 1).

We also found no significant interaction between bacterial supply and dispersal frequency ($\chi^2 = 3.419$; P

$= 0.064$; Table 1). The non-significant main effect of dispersal bias is expected, as the experimental

design ensures that dispersal bias is only functionally consequential when bacterial sink populations are

present (i.e., when bacterial supply > 0). Only under these conditions would pH values differ between

patches and asymmetric dispersal be possible under dispersal bias. Thus, the non-significant effect of

dispersal bias in isolation reflects a sensible experimental setup. Lastly, local yeast densities also

positively correlated with local bacteria densities (ρ =0.149, P<0.01; Fig. 5), an unexpected result that

suggests that bacteria have a net positive (rather than purely competitive) effect on yeast.

342   *Theoretical predictions of metapopulation model*

Model predictions agree qualitatively with our experimental results that on average, bacterial supply

increases metapopulation size and that this effect is greater when per-capita dispersal rate is lower in the

bacteria-containing patches (Fig. 5). Note that because we assume $r = K = D = 1$ for analytical tractability,

comparisons between model predictions and experiment are thus based on relative effects, as the

theoretical and experimental system may differ in their baseline parameter values (*e.g.*, experiment $K \neq$

1). Specifically, our model predicts that increasing the carrying capacity of yeast in the bacteria-

containing patch ($\beta > 1$) increases the overall metapopulation size (Fig. 5), similar to how bacterial

supply had an overall positive effect on yeast metapopulation size across treatments in our experiment

(Fig. 3B; average across treatments). Furthermore, the effect of lowering per-capita dispersal rate in the

bacteria-containing patch relative to the no-bacteria patch (*i.e.*, decreasing δ, analogous to dispersal bias

in the experiment) reduced metapopulation size when carrying capacities are similar, but increased it as

the carrying capacity of the bacteria-containing patch increased ($\beta \gg 1$; Fig. 5a). In other words,

dispersal bias increased metapopulation size relative to symmetric random dispersal only when the

bacteria-containing patch had sufficiently more resources, consistent with the significant interaction term

between bacterial supply and dispersal bias observed experimentally (predicted lines cross in Fig. 3A).

Lastly, we found that metapopulation size is maximized when the total number of migrants in each patch

are equalized, in accordance with the ideal free distribution (unpacked in Discussion; Fig. 5B).

DISCUSSION

Our study demonstrated that an invader with negative growth rates in all patches could still influence the metapopulation of a resident species by modifying both local habitat conditions and regional connectivity. In particular, we found that addition of an externally-sustained bacterial species increased the resident yeast metapopulation size, and that this effect was larger when they also reduced dispersal rates in the patches in which they occurred. Furthermore, our theoretical results align with experiments, and suggest that the mechanism by which dispersal bias increases yeast metapopulation size is through equalizing per-capita growth rates across patches, thereby bringing the metapopulation size closer to the theoretical maximum defined by the ideal free distribution. Taken together, these results demonstrate that transient invaders can impose demographic consequences on resident species through the interaction between local and regional processes. Below, we connect our results to metapopulation theory, discuss the biological mechanisms that may have generated these patterns, place our findings within broader ecological frameworks, and discuss key study limitations as well as opportunities for future research.

Dispersal and metapopulation size

Our study provides a rare empirical demonstration of how biotically-induced dispersal asymmetries can alter metapopulation size. Furthermore, the analysis of our theoretical model connects our experimental results to well-known effects of dispersal in spatially heterogeneous landscapes. In particular, theory shows that when carrying capacities vary spatially, dispersal typically reduces metapopulation size because productive patches export more migrants than they receive, causing greater overfilling of low-capacity patches than underfilling of high-capacity patches and thus stronger overall density dependence (Holt, 1985; McPeek & Holt, 1992; Pacala & Roughgarden, 1982). This demographic cost of dispersal underlies the "slower disperser wins" phenomenon (Dockery et al., 1998; Hastings, 1983) and explains

384   how a locally inferior competitor can coexist with a competitively superior species with faster dispersal

385   rates (Abrams & Wilson, 2004; Amarasekare & Nisbet, 2001; Pacala & Roughgarden, 1982). With

386   respect to our experiment, the demographic cost of dispersal in a landscape with heterogeneous carrying

387   capacities also explains why the positive effect of bacterial supply on metapopulation size was absent

388   when dispersal was random (Fig. 3A).

390   In agreement with McPeek & Holt (1992), our study shows that the detriments of dispersal on

metapopulation size can be nullified if differences in local per-capita dispersal rates offset differences in

local carrying capacities. The degree of this offset can be interpreted as a deviation from the ideal free

distribution, the condition under which local growth rates are equalized and metapopulation size is

maximized (Fig. 5B). Our formulation of dispersal slightly differs from that of McPeek & Holt (1992);

rather than assigning each patch an independent dispersal parameter, our model contains a single

parameter that controls the asymmetry in per-capita dispersal between patches. This formulation reflects

dispersal asymmetry arising from the foraging preferences of the dispersal vector (i.e., pollinators), such

that a single parameter captures a metapopulation-level process that generates patch-specific differences

in per-capita dispersal rates. More broadly, our results also align with empirical studies showing that

movement between habitats can modify demographic parameters (Donahue et al., 2003; Fox, 2007),

while extending this idea to the modification of dispersal parameters themselves.

403   *Biological mechanisms: Why does bacterial supply increase yeast densities?*

Our experimental results appear to contradict previous work showing that the reduction in pH by bacteria

inhibits yeast growth (Chappell et al., 2022; Figs. 3-4 & S1). We posit that experimental differences may

have contributed to the apparent disagreements. In the present experiment, densities of yeast and bacteria

were approximately the same at the beginning of each dilution cycle (~100-200 cells/µL; Fig. 1, see

Methods). Thus, nectar acidification by the bacteria occurred concurrently with yeast population growth,

possibly preventing pH from dropping fast enough to suppress yeast growth. However, this explanation is

only partial because it does not explain the observed positive effect of bacterial supply on yeast densities.

We posit that the addition of bacterial cells may have also introduced resource subsidies that enabled

yeast to reach higher densities. Although each inoculation event contributed only ~1% of the nectar

volume, additional inputs may have arisen from metabolites released by bacterial cells or from cell lysis,

beyond those present in the nectar medium used to suspend cells prior to inoculation. While this

mechanism has not been confirmed experimentally, it is consistent with reports of *Acinetobacter* spp.

releasing metabolic by-products in low-nitrogen nectar environments (Álvarez-Pérez et al., 2019; Kato et

al., 2025; Morales-Poole et al., 2023). Because many microbes secrete metabolites (Douglas, 2020) and

deceased cells can impact local communities (Hao et al., 2026), future work can identify the biochemical

pathways mediating these interactions.

Meta-ecosystems and nectar microbiome

The nutrient-subsidy mechanism we propose may reflect an experimental artefact, but the process may be

more general and relevant across ecosystems. Our results align with core ideas in meta-ecosystem

ecology, which emphasizes how flows of matter and energy shape ecosystem dynamics (Gounand et al.,

2018; Loreau et al., 2003; Massol et al., 2017; Polis et al., 1997). By focusing on connections across

ecosystem boundaries, this framework considers a broader set of spatial flows than traditional

metapopulation approaches would, including the movement of inorganic nutrients, organic matter, and

other non-living biotic materials in addition to living organisms. In nectar microbes, for instance, floral

nectar represents a primary habitat but not the only one they occupy. Nectar yeast may overwinter in bee

nests (Pozo et al., 2018), while soil and plant surfaces can also serve as refugia for other microbes before

they recolonize flowers in the following season (Vannette, 2020). Together, these examples highlight that

dispersal moves not just organisms but also the energy and nutrients associated with them from previous

places, prompting questions about the extent to which these cross-ecosystem fluxes affect local microbial

communities.

Experimental caveats and theoretical assumptions

Our experimental and theoretical approaches necessarily involve several caveats and simplifying

assumptions. For tractability, our two-patch model formalism ignores dispersal among identical habitats,

implicitly assuming that such movements have negligible effects on metapopulation dynamics. While this

may be a reasonable assumption within an idealized experimental setting like ours, movement between

identical habitats is likely to be important in real-world systems with extinction-recolonization dynamics

(*e.g.*, Brown & Kodric-Brown, 1977). With respect to our experiment, bacterial cells were always

inoculated into the same subset of patches, raising the possibility that our results reflect spatial placement

(e.g., edge effects on the PCR plate) rather than the effects of experimental treatments. However,

supplementary analyses found no evidence for location-specific effects (Appendix S4). Moreover,

although the factorial design of our experiment allowed us to isolate the effects of dispersal frequency and

dispersal bias, these factors likely covary in nature. For instance, pollinators may reduce visitation to

regions dominated by bacteria, thereby also lowering visitation to any yeast-dominated patches within

those regions. Lastly, both our experiment and model assumed conservative dispersal, whereas nectar

consumption by pollinators implies that only a fraction of microbes and nectar material are transferred

among flowers (*i.e*., dispersal has no fitness costs). Although this assumption is unlikely to hold in natural

systems, the rapid life cycles of nectar microbes suggests that even small numbers of immigrants may be

sufficient to modify local community structure.

Conclusion

This study experimentally demonstrated that a species invading a sink habitat can influence a resident metapopulation by modifying local conditions and altering metapopulation connectivity. More broadly, our results highlight that dispersing organisms can carry resources and nutrients acquired from previous habitats, allowing them to interact with resident species—even after death—by modifying the resource landscape that those species experience. Our experiments thus demonstrate the possibility that transient invaders may leave lasting ecological effects that persist well beyond their extinction. Given the prevalence of sink populations (Dias, 1996; Pulliam, 1988) in general and of unsuccessful microbial invasions in particular (Amor et al., 2020; Mallon et al., 2018), we suggest that it is worth investigating the extent to which these transient invaders structure local communities and metapopulations in natural ecosystems.

TABLES

Table 1. ANOVA table for the effects of treatments on metapopulation size and variance (n = 354)

Component	Term	χ^2	P
conditional	**Bacterial supply**	**6.10**	**0.01**
conditional	Bias	1.62	0.20
conditional	Frequency	1.02	0.31
conditional	**Bacterial supply:Bias**	**6.22**	**0.01**
conditional	Bacterial supply:Frequency	3.42	0.06
conditional	Bias:Frequency	1.12	0.29
dispersion	**Bacterial supply**	**4.20**	**0.04**

FIGURES

Figure 1. Overview of experimental setup. (**A**) Fully crossed design: dispersal bias (2) × dispersal frequency (2) × bacterial supply (3), each with six replicates. Each metapopulation consisted of 8 patches (PCR plate wells) connected by dispersal and underwent four 2-day growth–dilution cycles. (**B**) At the start of each cycle, patch pH was measured and used to generate donor–recipient pairs (see Methods). Dispersal was performed by transferring 3 µl of nectar from donor to recipient (red arrows). Patches were then diluted (5 µl into 95 µl fresh nectar) to emulate resource renewal, and samples were plated (TSA for bacteria, YMA for yeast) to estimate densities (CFUs). Bacteria (~20,000 cells) were added according to treatment, and plates were sealed and incubated at 26°C for 2 days.

Figure 2. Time-series data of bacteria and yeast densities at local and regional scales. Lines show the
time-series of the microbial densities of a single replicate metapopulation. The densities of bacteria
(strain: FNA17) and yeast (strain: MR1) are colored in purple and yellow, respectively. Solid lines (with
points) show the metapopulation size ($\sum N_i$) while dashed lines show the average density of a single patch
($\bar{N}$) within a replicate metapopulation. Each facet contains the 6 replicates of a single treatment specified
by the facet labels.

Figure 3. Effects of bacterial supply and dispersal bias on yeast metapopulation size and variance.
(**A**) Estimated treatment effects on the metapopulation-level colony forming units of yeast per μl, while fitted lines and 95% confidence intervals are from GLMM estimates. Points in (**B**) show the estimated marginal effect of bacterial supply on yeast CFUs at different levels of dispersal bias and evaluated at the mean dispersal frequency (4.5) with error bars indicating SE.

Figure 4. Association between patch-level yeast and bacteria densities. Points indicate the number of colony forming units (CFU) of bacteria and yeast per μl in a single patch of a metapopulation. Line of best fit was drawn to aid visualization and ρ is the Pearson correlation coefficient ($p < 0.01$). Points clustered around 200 CFUs represent cases where CFUs were too dense to count; see Methods.

Figure 5. Theoretical predictions of metapopulation size with heterogeneous per-capita dispersal rates and carrying capacities. Metapopulation size is the total equilibrium density across both patches $(N_B^* + N_N^*)$. (**A**) Metapopulation size increases with patch B carrying capacity (β), but more strongly when per-capita dispersal from patch B is lower (blue relative to grey curve). (**B**) As β increases, equalizing total migrants between patches requires lower per-capita dispersal from patch B ($\delta < 0$). The black curve shows the condition $(1 + \delta)N_B^* = (1 - \delta)N_N^*$. Above and below the curve, patch B and patch N, respectively, contribute more migrants at equilibrium. Points indicate the value of δ that maximizes metapopulation size for a given β (point size proportional to metapopulation size). In both panels, $r = K = D = 1$.

References

Abrams, P. A., & Wilson, W. G. (2004). Coexistence of competitors in metacommunities due to spatial
variation in resource growth rates; does R * predict the outcome of competition? Ecology Letters,
7(10), 929–940. https://doi.org/10.1111/j.1461-0248.2004.00644.x

Álvarez-Pérez, S., Lievens, B., & Fukami, T. (2019). Yeast–Bacterium Interactions: The Next Frontier in
Nectar Research. Trends in Plant Science, 24(5), 393–401.
https://doi.org/10.1016/j.tplants.2019.01.012

Amarasekare, P. (2010). Effect of non-random dispersal strategies on spatial coexistence mechanisms.
Journal of Animal Ecology, 79(1), 282–293. https://doi.org/10.1111/j.1365-2656.2009.01607.x

Amarasekare, P., & Nisbet, R. M. (2001). Spatial Heterogeneity, Source-Sink Dynamics, and the Local
Coexistence of Competing Species. The American Naturalist, 158(6), 572–584.
https://doi.org/10.1086/323586

Amor, D. R., Ratzke, C., & Gore, J. (2020). Transient invaders can induce shifts between alternative
stable states of microbial communities. Science Advances, 6(8), eaay8676.
https://doi.org/10.1126/sciadv.aay8676

Arditi, R., Lobry, C., & Sari, T. (2015). Is dispersal always beneficial to carrying capacity? New insights
from the multi-patch logistic equation. Theoretical Population Biology, 106, 45–59.
https://doi.org/10.1016/j.tpb.2015.10.001

Bartlow, A. W., & Agosta, S. J. (2021). Phoresy in animals: Review and synthesis of a common but
understudied mode of dispersal. Biological Reviews, 96(1), 223–246.
https://doi.org/10.1111/brv.12654

Bascompte, J., & Solé, R. V. (1994). Spatially Induced Bifurcations in Single-Species Population
Dynamics. Journal of Animal Ecology, 63(2), 256–264. https://doi.org/10.2307/5544

Belisle, M., Peay, K. G., & Fukami, T. (2012). Flowers as Islands: Spatial Distribution of Nectar-
Inhabiting Microfungi among Plants of Mimulus aurantiacus, a Hummingbird-Pollinated Shrub.
Microbial Ecology, 63(4), 711–718. https://doi.org/10.1007/s00248-011-9975-8

Brown, J. H., & Kodric-Brown, A. (1977). Turnover Rates in Insular Biogeography: Effect of
Immigration on Extinction. Ecology, 58(2), 445–449. https://doi.org/10.2307/1935620

Chappell, C. R., Dhami, M. K., Bitter, M. C., Czech, L., Herrera Paredes, S., Barrie, F. B., Calderón, Y.,
Eritano, K., Golden, L.-A., Hekmat-Scafe, D., Hsu, V., Kieschnick, C., Malladi, S., Rush, N., &
Fukami, T. (2022). Wide-ranging consequences of priority effects governed by an overarching
factor. eLife, 11, e79647. https://doi.org/10.7554/eLife.79647

Chappell, C. R., & Fukami, T. (2018). Nectar yeasts: A natural microcosm for ecology. Yeast, 35(6),
417–423. https://doi.org/10.1002/yea.3311

Clobert, J., Le Galliard, J.-F., Cote, J., Meylan, S., & Massot, M. (2009). Informed dispersal,
heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations.
Ecology Letters, 12(3), 197–209. https://doi.org/10.1111/j.1461-0248.2008.01267.x

DeAngelis, D. L., Ni, W.-M., & Zhang, B. (2016). Dispersal and spatial heterogeneity: Single species.
Journal of Mathematical Biology, 72(1), 239–254. https://doi.org/10.1007/s00285-015-0879-y

Dias, P. C. (1996). Sources and sinks in population biology. Trends in Ecology & Evolution, 11(8), 326–
330. https://doi.org/10.1016/0169-5347(96)10037-9

Dockery, J., Hutson, V., Mischaikow, K., & Pernarowski, M. (1998). The evolution of slow dispersal rates: A reaction diffusion model. Journal of Mathematical Biology, 37(1), 61–83. https://doi.org/10.1007/s002850050120

Doebeli, M. (1995). Dispersal and Dynamics. Theoretical Population Biology, 47(1), 82–106. https://doi.org/10.1006/tpbi.1995.1004

Donahue, M. J., Holyoak, M., & Feng, C. (2003). Patterns of Dispersal and Dynamics among Habitat Patches Varying in Quality. The American Naturalist, 162(3), 302–317. https://doi.org/10.1086/377185

Douglas, A. E. (2020). The microbial exometabolome: Ecological resource and architect of microbial communities. Philosophical Transactions of the Royal Society B: Biological Sciences, 375(1798), 20190250. https://doi.org/10.1098/rstb.2019.0250

Fox, J. W. (2007). Testing the Mechanisms by Which Source-Sink Dynamics Alter Competitive Outcomes in a Model System. The American Naturalist, 170(3), 396–408. https://doi.org/10.1086/519855

Fretwell, S. D., & Lucas, H. L. (1969). On territorial behavior and other factors influencing habitat distribution in birds. Acta Biotheoretica, 19(1), 16–36. https://doi.org/10.1007/BF01601953

Fukami, T. (2005). Integrating internal and external dispersal in metacommunity assembly: Preliminary theoretical analyses. Ecological Research, 20(6), 623–631. https://doi.org/10.1007/s11284-005-0092-3

Germain, R. M., Jones, N. T., & Grainger, T. N. (2019). Cryptic dispersal networks shape biodiversity in an invaded landscape. Ecology, 100(8), e02738. https://doi.org/10.1002/ecy.2738

Gounand, I., Harvey, E., Little, C. J., & Altermatt, F. (2018). Meta-Ecosystems 2.0: Rooting the Theory into the Field. Trends in Ecology & Evolution, 33(1), 36–46. https://doi.org/10.1016/j.tree.2017.10.006

Grainger, T. N., Letten, A. D., Gilbert, B., & Fukami, T. (2019). Applying modern coexistence theory to priority effects. Proceedings of the National Academy of Sciences, 116(13), 6205–6210. https://doi.org/10.1073/pnas.1803122116

Hanski, I. (1998). Metapopulation dynamics. Nature, 396(6706), 41–49. https://doi.org/10.1038/23876

Hao, Y.-Q., Li, B.-H., Chen, J.-Y., Shu, W.-S., & Zhao, X.-F. (2026). Bacterial necromass recycling promotes diversity maintenance in bacterial communities via resource partitioning. Nature Ecology & Evolution, 1–14. https://doi.org/10.1038/s41559-025-02967-2

Hassell, M. P., Comins, H. N., & Mayt, R. M. (1991). Spatial structure and chaos in insect population dynamics. Nature, 353(6341), 255–258. https://doi.org/10.1038/353255a0

Hastings, A. (1982). Dynamics of a single species in a spatially varying environment: The stabilizing role of high dispersal rates. Journal of Mathematical Biology, 16(1), 49–55. https://doi.org/10.1007/BF00275160

Hastings, A. (1983). Can spatial variation alone lead to selection for dispersal? Theoretical Population Biology, 24(3), 244–251. https://doi.org/10.1016/0040-5809(83)90027-8

He, X., Ni, W.-M., Ye, Z., & Zhang, B. (2025). Total population for a resource-limited single consumer model. Journal of Mathematical Biology, 90(2), 20. https://doi.org/10.1007/s00285-025-02186-0

Herrera, C. M., Canto, A., Pozo, M. I., & Bazaga, P. (2009). Inhospitable sweetness: Nectar filtering of pollinator-borne inocula leads to impoverished, phylogenetically clustered yeast communities. Proceedings of the Royal Society B: Biological Sciences, 277(1682), 747–754. https://doi.org/10.1098/rspb.2009.1485

Holt, R. D. (1985). Population dynamics in two-patch environments: Some anomalous consequences of an optimal habitat distribution. Theoretical Population Biology, 28(2), 181–208. https://doi.org/10.1016/0040-5809(85)90027-9

Holt, R. D., Barfield, M., & Gonzalez, A. (2003). Impacts of environmental variability in open populations and communities: "Inflation" in sink environments. Theoretical Population Biology, Understanding the Role of Environmental Vatiation in Population and Community Dynamics, 64(3), 315–330. https://doi.org/10.1016/S0040-5809(03)00087-X

Huffaker, C. B. (1958). EXPERIMENTAL STUDIES ON PREDATION: DISPERSION FACTORS AND PREDATOR-PREY OSCILLATIONS. HILGARDIA, 27(14).

Karlin, S., & McGregor, J. (1972). Polymorphisms for genetic and ecological systems with weak coupling. Theoretical Population Biology, 3(2), 210–238. https://doi.org/10.1016/0040-5809(72)90027-5

Kato, Y., Miura, H., Takayama, S., & Fujii, S. (2025). Glucose-dependent acidification strategy by nectar-dwelling bacteria mediates pollen tube burst. Research Square. https://doi.org/10.21203/rs.3.rs-7063044/v1

Kortessis, N., Glass, G., Gonzalez, A., Ruktanonchai, N. W., Simon, M. W., Singer, B., & Holt, R. D. (2025). Metapopulations, the Inflationary Effect, and Consequences for Public Health. The American Naturalist, 205(3), 342–359. https://doi.org/10.1086/733896

Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D., Shurin, J. B., Law, R., Tilman, D., Loreau, M., & Gonzalez, A. (2004). The metacommunity concept: A framework for multi-scale community ecology. Ecology Letters, 7(7), 601–613. https://doi.org/10.1111/j.1461-0248.2004.00608.x

Letten, A. D., Dhami, M. K., Ke, P.-J., & Fukami, T. (2018). Species coexistence through simultaneous fluctuation-dependent mechanisms. Proceedings of the National Academy of Sciences, 115(26), 6745–6750. https://doi.org/10.1073/pnas.1801846115

Levin, S. A. (1974). Dispersion and Population Interactions. The American Naturalist, 108(960), 207–228.

Levins, R. (1969). Some Demographic and Genetic Consequences of Environmental Heterogeneity for Biological Control1. Bulletin of the Entomological Society of America, 15(3), 237–240. https://doi.org/10.1093/besa/15.3.237

Long, Z. T., Petchey, O. L., & Holt, R. D. (2007). The effects of immigration and environmental variability on the persistence of an inferior competitor. Ecology Letters, 10(7), 574–585. https://doi.org/10.1111/j.1461-0248.2007.01049.x

Loreau, M., Mouquet, N., & Holt, R. D. (2003). Meta-ecosystems: A theoretical framework for a spatial ecosystem ecology. Ecology Letters, 6(8), 673–679. https://doi.org/10.1046/j.1461-0248.2003.00483.x

Lou, Y. (2006). On the effects of migration and spatial heterogeneity on single and multiple species. Journal of Differential Equations, 223(2), 400–426. https://doi.org/10.1016/j.jde.2005.05.010

Mallon, C. A., Le Roux, X., van Doorn, G. S., Dini-Andreote, F., Poly, F., & Salles, J. F. (2018). The impact of failure: Unsuccessful bacterial invasions steer the soil microbial community away from the invader's niche. The ISME Journal, 12(3), 728–741. https://doi.org/10.1038/s41396-017-0003-y

Mari, L., Casagrandi, R., Bertuzzo, E., Rinaldo, A., & Gatto, M. (2014). Metapopulation persistence and species spread in river networks. Ecology Letters, 17(4), 426–434. https://doi.org/10.1111/ele.12242

Massol, F., Altermatt, F., Gounand, I., Gravel, D., Leibold, M. A., & Mouquet, N. (2017). How life-history traits affect ecosystem properties: Effects of dispersal in meta-ecosystems. Oikos, 126(4), 532–546. https://doi.org/10.1111/oik.03893

Matthews, D. P., & Gonzalez, A. (2007). The Inflationary Effects of Environmental Fluctuations Ensure the Persistence of Sink Metapopulations. Ecology, 88(11), 2848–2856. https://doi.org/10.1890/06-1107.1

McPeek, M. A., & Holt, R. D. (1992). The Evolution of Dispersal in Spatially and Temporally Varying Environments. The American Naturalist, 140(6), 1010–1027.

McPeek, M. A., Resetarits, W. J., Jr, & Holt, R. D. (2024). The evolution of passive dispersal versus habitat selection have differing emergent consequences in metacommunities. Philosophical Transactions of the Royal Society B: Biological Sciences, 379(1907), 20230126. https://doi.org/10.1098/rstb.2023.0126

Morales-Poole, J. R., de Vega, C., Tsuji, K., Jacquemyn, H., Junker, R. R., Herrera, C. M., Michiels, C., Lievens, B., & Álvarez-Pérez, S. (2023). Sugar Concentration, Nitrogen Availability, and Phylogenetic Factors Determine the Ability of Acinetobacter spp. And Rosenbergiella spp. To Grow in Floral Nectar. Microbial Ecology, 86(1), 377–391. https://doi.org/10.1007/s00248-022-02088-4

Morris, M. M., Frixione, N. J., Burkert, A. C., Dinsdale, E. A., & Vannette, R. L. (2020). Microbial abundance, composition, and function in nectar are shaped by flower visitor identity. FEMS Microbiology Ecology, 96(3), fiaa003. https://doi.org/10.1093/femsec/fiaa003

Muller-Landau, H. C., Wright, S. J., Calderón, O., Condit, R., & Hubbell, S. P. (2008). Interspecific variation in primary seed dispersal in a tropical forest. Journal of Ecology, 96(4), 653–667. https://doi.org/10.1111/j.1365-2745.2008.01399.x

Nell, L. A., Klausmeier, C. A., & Fukami, T. (2025). Regional species coexistence despite local priority effects: The overlooked role of dispersal–community feedback (p. 2025.03.31.646410). bioRxiv. https://doi.org/10.1101/2025.03.31.646410

Okubo, A., & Levin, S. A. (2001). Diffusion and Ecological Problems: Modern Perspectives (Vol. 14). Springer. https://doi.org/10.1007/978-1-4757-4978-6

Pacala, S. W., & Roughgarden, J. (1982). Spatial heterogeneity and interspecific competition. Theoretical Population Biology, 21(1), 92–113. https://doi.org/10.1016/0040-5809(82)90008-9

Pascual, M. (1993). Diffusion-Induced Chaos in a Spatial Predator—Prey System. Proceedings: Biological Sciences, 251(1330), 1–7.

Peay, K. G., Belisle, M., & Fukami, T. (2012). Phylogenetic relatedness predicts priority effects in nectar yeast communities. Proceedings: Biological Sciences, 279(1729), 749–758.

Pozo, M. I., Bartlewicz, J., van Oystaeyen, A., Benavente, A., van Kemenade, G., Wäckers, F., & Jacquemyn, H. (2018). Surviving in the absence of flowers: Do nectar yeasts rely on overwintering bumblebee queens to complete their annual life cycle? FEMS Microbiology Ecology, 94(12), fiy196. https://doi.org/10.1093/femsec/fiy196

Pulliam, H. R. (1988). Sources, Sinks, and Population Regulation. The American Naturalist, 132(5), 652–
661.

Quevedo-Caraballo, S., de Vega, C., Lievens, B., Fukami, T., & Álvarez-Pérez, S. (2025). Tiny but
mighty? Overview of a decade of research on nectar bacteria. New Phytologist, 245(5), 1897–1910.
https://doi.org/10.1111/nph.20369

Schaeffer, R. N., Rering, C. C., Maalouf, I., Beck, J. J., & Vannette, R. L. (2019). Microbial metabolites
elicit distinct olfactory and gustatory preferences in bumblebees. Biology Letters, 15(7), 20190132.
https://doi.org/10.1098/rsbl.2019.0132

Solé, R. V., Bascompte, J., & Valls, J. (1992). Stability and complexity of spatially extended two-species
competition. Journal of Theoretical Biology, 159(4), 469–480. https://doi.org/10.1016/S0022-
5193(05)80691-5

Tilman, D. (1994). Competition and Biodiversity in Spatially Structured Habitats. Ecology, 75(1), 2–16.
https://doi.org/10.2307/1939377

Tucker, C. M., & Fukami, T. (2014). Environmental variability counteracts priority effects to facilitate
species coexistence: Evidence from nectar microbes. Proceedings of the Royal Society B:
Biological Sciences, 281(1778), 20132637. https://doi.org/10.1098/rspb.2013.2637

Turchin, P. (1990). Rarity of density dependence or population regulation with lags? Nature, 344(6267),
660–663. https://doi.org/10.1038/344660a0

Vannette, R. L. (2020). The Floral Microbiome: Plant, Pollinator, and Microbial Perspectives. Annual
Review of Ecology, Evolution, and Systematics, 51(Volume 51, 2020), 363–386.
https://doi.org/10.1146/annurev-ecolsys-011720-013401

Vannette, R. L., & Fukami, T. (2014). Historical contingency in species interactions: Towards niche-
based predictions. Ecology Letters, 17(1), 115–124. https://doi.org/10.1111/ele.12204

Vannette, R. L., & Fukami, T. (2017). Dispersal enhances beta diversity in nectar microbes. Ecology
Letters, 20(7), 901–910. https://doi.org/10.1111/ele.12787

Vannette, R. L., Gauthier, M.-P. L., & Fukami, T. (2013). Nectar bacteria, but not yeast, weaken a plant–
pollinator mutualism. Proceedings of the Royal Society B: Biological Sciences, 280(1752),
20122601. https://doi.org/10.1098/rspb.2012.2601

Vannette, R. L., McMunn, M. S., Hall, G. W., Mueller, T. G., Munkres, I., & Perry, D. (2021). Culturable
bacteria are more common than fungi in floral nectar and are more easily dispersed by thrips, a
ubiquitous flower visitor. FEMS Microbiology Ecology, 97(12), fiab150.
https://doi.org/10.1093/femsec/fiab150

Vuilleumier, S., & Possingham, H. P. (2006). Does colonization asymmetry matter in metapopulations?
Proceedings of the Royal Society B: Biological Sciences, 273(1594), 1637–1642.
https://doi.org/10.1098/rspb.2006.3469

Xiao, Z., & Zhang, Z. (2016). Contrasting patterns of short-term indirect seed–seed interactions mediated
by scatter-hoarding rodents. Journal of Animal Ecology, 85(5), 1370–1377.
https://doi.org/10.1111/1365-2656.12542

Zhang, B., Kula, A., Mack, K. M. L., Zhai, L., Ryce, A. L., Ni, W.-M., DeAngelis, D. L., & Van Dyken,
J. D. (2017). Carrying capacity in a heterogeneous environment with habitat connectivity. Ecology
Letters, 20(9), 1118–1128. https://doi.org/10.1111/ele.12807

732