Reciprocal Environmental Decision Support (REDS): better tailored advice in return for data

Ben Kenward[1,2], Nick Casey[1,3], Perline Bastid[1], Francis Buner[4], Ishi Buffam[5], Julie Ewald[1,4], Robert Kenward[1,6]

Institutions and addresses

[1]European Sustainable Use Group, Rue de la Science 10, Bruxelles 1000, Belgium

[2]Centre for Psychological Research, Oxford Brookes University, Headington Campus, Oxford, OX3 0BP, UK

[3]Anatrack Ltd, 52 Furzebrook Road, Wareham, Dorset, BH20 5AX, UK

[4]Game & Wildlife Conservation Trust, Burgate Manor, Fordingbridge, Hampshire, SP6 1EF, UK

[5]Department of Landscape Architecture, Planning and Management, Swedish University of Agricultural Sciences, Alnarp, Sweden

[6]UK Centre for Ecology and Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

Corresponding author

Ben Kenward
ben.kenward@wolfson.oxon.org

Data availability statement

Data is available via Supporting Information.

Acknowledgements

Thanks to Nicholas Aebischer and Johan Vegelius for useful discussions, to Prolific-recruited participants who went the extra mile, and to Bridget Kenward for much practical assistance. This study was funded by the EU Horizon grant PRO-COAST (Project ID 101082327).

Author contributions

Conceptualisation: BK, NC, JE, RK; Data curation: BK, NC; Analysis: BK; Funding acquisition: JE, RK; Methodology: all authors; Software: NC, BK; Writing (original draft): BK; Writing (review): all authors.

Conflict of interest statement

Ben Kenward, Nick Casey, and Robert Kenward are co-directors of Anatrack Ltd, a company that assisted in the development of technical systems used in the study, and Nick Casey is an employee. Following a decision at the 2025 board of directors meeting, the company is in the process of restructuring to a non-profit organisation.

Lay Summary

A summary for non-specialists is available at https://bit.ly/reds_brochure.

Reciprocal Environmental Decision Support (REDS): better tailored advice in return for data

Abstract

1. Environmental Decision Support systems use models to make tailored predictions about local circumstances based on user input. Citizen Science systems accept contributions about local circumstances to improve models. Each system type emphasises data flow in one direction: system to user or user to system, respectively.

2. We introduce a new system type that synergises data flows in both directions. Reciprocal Environmental Decision Support (REDS) systems process user-contributed information to (1) provide tailored predictions and (2) improve the model that generates the advice. We assessed existing decision support systems and found no REDS system for ecosystem management.

3. For our proof-of-concept REDS system, Garden Advice, we began with an initial Bayesian species-habitat association model (for the House Sparrow *Passer domesticus*) and solicited data from UK residents about habitat structure and sparrow observations in domestic gardens. The model made predictions about sparrow presence, and effects of planned garden changes, while updating parameter estimates based on contributed data, generally becoming more confident. One salient parameter update likely reflects reality (a positive association with grass), while another (a reduced association with roof proximity) likely reflects observation bias.

4. Cross-validated comparison indicated that the final model predicted the solicited data better than the original model. Thus, untrained observers can provide data of sufficient quality to refine a model already parameterised with trained observer data. Notwithstanding important questions about distinguishing observation bias from ecologically meaningful information, our system demonstrates that important synergies can be obtained from the REDS approach. Later users of the system obtained better advice thanks to contributions from earlier users and automated Bayesian updating.

Keywords

Reciprocal environmental decision support; REDS; environmental decision support; EDS; Citizen Science; House Sparrow *Passer domesticus*; ecological modelling; Bayesian updating.

Word counts

262 (abstract); 6,691 (all other text)

Introduction

Introducing Reciprocal Environmental Decision Support (REDS)

A salient feature of many information systems is that query input is not discarded once query output has been returned, because this data has value (e.g., Bondia-Barceló et al., 2016). Such use is often of ambiguous social benefit (for example in advertising, Frick et al., 2021; Martin & Murphy, 2017), but examples of social good include web search data revealing trends in public consciousness of environmental issues (Nghiem et al., 2016), and enabling epidemics to be rapidly tracked (Ginsberg et al., 2009).

Environmental decision support (EDS) systems typically require complex query input, because optimised predictions about environmental consequences of decisions require the situation to be described in some detail. Frequently, land managers collate and input detailed information that is processed to produce output that can support decisions but is not put to further use. As one example illustrating this general point, the Cool Farm Tool uses extensive farm activity data to estimate farm carbon footprint and to support decisions by analysing hypothetical activity changes; it has been used for over 125,000 assessments (Cool Farm, 2025a; Kayatz et al., 2019). This implies a rich central database, but as they note, "the Cool Farm Alliance has not used any data stored in the tool" for purposes except returning assessment results (Cool Farm, 2025b).

The richness of EDS query input data, along with the lack of evidence for its use (beyond returning query output), implies that important opportunities are being missed. The typically multivariate nature of this data implies that it could parameterise predictive models that themselves could be useful within EDS. This would begin to bridge the gap between EDS and Citizen Science (CS). The current work introduces a new type of system that combines the two (see also conceptual groundwork in Kenward et al., 2013; Papathanasiou & Kenward, 2014). Whereas CS emphasises the one-way data flow from citizen to system, and EDS emphasises the one-way data flow from system to citizen, Reciprocal Environmental Decision Support (REDS) represents a novel system type which closes this loop (Figure 1).

Information loop closure has the theoretical potential to revolutionise the quality of data, models, and decisions, through iterative mutual improvement via feedback loops (Wiener, 1948). Here, we empirically demonstrate a REDS system that uses input not only to generate model predictions that can support decision making, but also to improve the model itself. We illustrate this by implementing a habitat association model for an animal indicator-species that makes predictions about species occurrence based on user-provided habitat descriptions, with model updating enabled by user-provided animal observations.

67   *Figure 1*. Information flows in EDS, CS, and REDS.

Environmental decision support

We define EDS systems as those with the primary purpose of supporting environmental management decisions and that provide model-generated predictions about environmentally relevant consequences of possible actions. Non-predictive databases can also support environmental decisions, but in line with previous literature requiring EDS systems to feature "analysis" or "models" or to be "intelligent" (McIntosh et al., 2011, p. 1391), we exclude such systems from our definition. REDS systems are EDS systems that are reciprocal; not only is user knowledge updated by the model predictions, but model predictions are updated by user observations (Figure 1).

We define a REDS system as self-contained (though network-accessed) and with model updating on a time scale of a few days at most after data contribution. Otherwise, any EDS system informed by users in an indirect way (e.g., by a CS study reported in the literature) might be classified as REDS. Indirect and delayed closure of the information loop cannot lead to dynamic iterative updating, and is less likely to lead to productive synergies between CS and EDS.

Table 1 lists example systems and evaluates them as REDS candidates. Some systems feature reciprocal sharing of environmentally relevant data but are not REDS (or EDS) because there is no model, for example the PurpleAir system for sharing air quality measures (PurpleAir, 2025; Robinson et al., 2023). The species-mapping functionality of iNaturalist does have a model that is reciprocally updated: users report species observations that update a model predicting occurrence at unobserved locations (Cole et al., 2023). The data flow and model updating is REDS-like, but iNaturalist is not an EDS. The species occurrence models are based on spatial coordinates at very coarse scale, and (as implemented) do not involve environmental conditions predictors, so the models are not suitable to support the what-if analyses needed for EDS.

Google Maps route-planning is presented as supporting environmental decisions, as it offers information on the relative energy efficiency of different routes. Further, it is reciprocal, as measurements of traffic speed from users update its models of traffic conditions which in turn inform route recommendations. Thus, Google Maps route-planning is the only existing REDS of which we are aware (although perhaps not a successful implementation, Burris et al., 2023), and it stretches the definition, as it does not assist ecosystem management directly.

97   Table 1: Data flows and REDS status for example candidate systems

	User → system	System → user			Reciprocal	Ecosystem
	Data	Data	Model predictions	EDS	model update	management
PurpleAir	Air quality	Air quality	-	No	No	No
iNaturalist	Species occurrence	Species occurrence	Location → species occurrence	No	Yes	No
Google Maps	Traffic speed	Traffic speed	Route choice → energy efficiency	Yes	Yes	No
InVEST	Diverse variables	-	Diverse variables → ecosystem services	Yes	No	Yes
Garden Advice	Habitat, species occurrence	-	Habitat → species occurrence	Yes	Yes	Yes

Note: Ecosystem management denotes whether there are model predictions based on environmental condition
variables, as opposed to (for example) purely spatial variables, which cannot be managed.

None of the above systems feature models where the inputs include environmental conditions,
although such models are key for most EDS systems (e.g., the InVEST suite of models, Natural
Capital Project, 2025), because ecosystem management involves manipulation of environmental
conditions. Following a systematic review (see Supporting Information), we found no existing
REDS systems with models based on environmental conditions. Reciprocal model updating is
common for species identification, but not for EDS. We therefore introduce Garden Advice as
the first REDS suitable for ecosystem management.

The Garden Advice REDS system and our associated research questions

Garden Advice (GA) records (amongst other things) habitat descriptions and House Sparrow
Passer domesticus observations in domestic gardens across the UK. We chose a sparrow habitat
association model for testing REDS-related research questions for multiple reasons. House
Sparrows are in global decline, for complex reasons, and are frequently studied, so are a useful
indicator species (Hanson et al., 2020). Despite their decline they are relatively common in UK
domestic gardens, thanks to their association with humans (De Laet & Summers-Smith, 2007).
UK domestic gardens are an excellent arena for citizen science, thanks to the popularity of
gardening (RHS, 2025; Sandhaus et al., 2019; Sharma et al., 2019). Importantly, a raw dataset is
available for sparrow habitat association in UK domestic gardens, thanks to the Glasgow House
Sparrow Project (GHSP; Matthiopoulos et al., 2019), enabling an initial model.

Our primary research question was whether we could successfully implement a functioning
REDS system, defined as confirmation of the following preregistered hypothesis: after updating
the initial model with user contributed data, the model will better predict the user data. In
principle, this is a modest criterion, because if there is information in the user data absent from
the initial model, then with enough data, logically the criterion must be met. In practice, when
soliciting internet data from untrained users across an appreciable geographic range, there are
reasons why these conditions may not apply, rendering the question non-trivial. Our focus is on
predicting newly user-contributed data rather than the initial data, because user data is likely
more representative of house sparrows across the UK (all initial data is from Glasgow). A
secondary purpose of this report is to describe user advice-seeking behaviours within Garden
Advice, to investigate how those managing land might use a REDS system.

Our methods include no independent verification of accuracy of citizen contributions. This raises
concerns about citizen science data quality (Aceves-Bueno et al., 2017; Backstrom et al., 2024;

Johnston et al., 2023; Kosmala et al., 2016), questionable in similar contexts without additional
verification (Williams et al., 2015). We return to this issue in the light of our results.

Methods

Declarations

Extensive additional detail concerning all methods is provided in Supporting Information. Large
Language Models were sometimes used to check natural language style and to assist with writing
of code. The procedure received ethical approval from the [ANONYMISED] human-participant
research-ethics internal assessment procedure. Participants gave fully informed consent. We
preregistered a specific method to test the preregistered hypothesis, which we followed except
where unforeseen circumstances intervened, as noted where relevant.

Participants

Prolific, a service for social scientists to recruit paid participants, known for good quality data in
comparison to other similar services (Albert & Smilek, 2023; Peer et al., 2022), was used to
commission 71 UK residents to describe their gardens using Garden Advice in May 2025. An
additional 68 UK residents made initial attempts to participate, but could not, for example due to
misidentifying birds in a screening questionnaire. All garden data obtained during the study
period that could be included in model updating was included. A small number of broad
requirements for participation (e.g., having an upper secondary education) were implemented in
Prolific to optimise participant ability and motivation to succeed at mapping a domestic garden.
Most participants were middle-aged; most were in full-time work; most had university degrees.

Garden Advice app

The system provides user-friendly methods for citizens to map habitats in domestic gardens and
to specify areas where they typically observe House Sparrows. The system is REDS because it
provides users with scores indicating the likelihoods of observing sparrows in gardens they
describe, and updates the model that generates these scores with the user-provided data. The
system also indicates how planned garden changes would impact the sparrow scores. Thus, the
system is designed to be capable of providing better EDS, the more it is used – the hallmark of
the REDS approach. Garden Advice can be accessed at https://garden.earthadvice.org (those
providing non-genuine data for testing purposes are asked to enable testing mode).

In-app instructions guided the current users to (1) find their garden on the base map, (2) draw a
boundary around it, (3) add habitat patches until the bounded area was filled, (4) certify that the
map was accurate (with direction to editing functions if not), (5) answer whether they had ever
seen a House Sparrow in their garden, (6) if they answered yes, draw polygons around all
specific areas where sparrows have been seen. All data was contributed without any non-
automated assistance to users.

Garden Advice also incorporates a carbon model estimating how much carbon gardens sequester.
It provides these estimates in the form of hours of human life saved, by converting carbon to
human mortality via the thousand-ton rule (see Pearce & Parncutt, 2023 for a presentation of this
rule, and Bressler, 2021 for an example model from which it is derived). Although not part of the
REDS functionality, this is an important part of Garden Advice because it attracts users (this was
how we developed the system), and because framing carbon sources and sinks in terms of
countable human deaths and lives saved is a useful messaging tool (Kenward, 2024).

The sparrow habitat association model

Variables

Model variables were chosen for compatibility with the publicly available Glasgow House
Sparrow Project dataset (GHSP), consisting of observations from trained surveyors at 32
individual colony areas in and around the city of Glasgow (Matthiopoulos et al., 2019). Each
data point represents a 2 m^2 map grid-cell where a sparrow was (1) or was not (0) observed.
Predictor variables are proportion of cell covered by each of six habitat types (tree, bush, hedge,
grass, roof, and artificial surface) plus two distance variables representing the distance from the
cell to the nearest hedge or roof. Garden Advice (GA) allows mapping of two additional habitat
types anticipated to be necessary for participants to fully map their gardens: water (ignored in the
current model) and cultivated beds (treated as bushes).

The GA binary dependent variable is 1 if any sparrow observation area overlaps with the cell,
otherwise 0. Utilised GHSP distances were approximate reconstructions because available GHSP
distances were z-transformed before release; GA roof distances are calculated using existing map
data as well as user data; GA hedge distances in gardens with no hedges are imputed as the
maximum observed value (96.7 m). For presentation, distance variables are converted to
proximity variables by reversing the sign, so that effects have a more intuitive direction.

Model specification and updating

The model is a Bayesian binary logistic regression, with R syntax:

sparrow ~ hedge + grass + tree + roof + artificial + bush + hedgeProximity + roofProximity +
(1 | location) +
(0 + hedge + grass + tree + roof + artificial + bush + hedgeProximity + roofProximity || location)

In other words, there are eight fixed predictors, an overall intercept, a random intercept for each
observed location (GHSP colony or GA garden), and random slopes for each fixed predictor and
location (uncorrelated to ease model fit). The random slopes control for intra-location spatial
autocorrelation, as well as capturing any inter-location heterogeneity in habitat associations.

The model was initially fit to all GHSP data and incorporated into the GA system. It was then
updated with the new GA data every day during the study period, entirely automatically except
for manual addition of certain data contributions to an exclusion list (Table S3).

GHSP data was sampled in case-control format – each colony area was observed until a sparrow
was registered at 40 grid cells, then 40 grid cells where no sparrow was observed were randomly
selected, and habitat was described for the 80 cells. GA data, on the other hand, comprises full
presence-absence information, enabling prevalence calculations. We used a prevalence-based
offset to make the data sets compatible. Information about offsetting in the Supporting
Information is important because it involves a deviation from preregistration, for the sake of
improved compatibility when comparing models.

Preregistration stated that models for leave-one-group-out cross-validation (see below) would be
fitted using the standard Markov chain Monte Carlo (MCMC) approach, but in practice these
models ran too slowly so we switched to Integrated nested Laplace approximation (INLA).
Priors were the default in all cases (as preregistered), except for random slope variance
hyperparameters in the INLA models, where initial poor fit, combined with re-consideration of
appropriate prior beliefs, led to a specification of a comparatively high variance prior.

Model predictions and their evaluation

Random components of any version of the models explain large amounts of variation. However,
this variation cannot be used for predictions at novel locations, and we therefore only use fixed
model components for making predictions. For evaluating prediction accuracy, we preregistered
the use of Brier scores (equivalent to mean squared error) and the standard decomposition into
two primary components (Merkle & Hartman, 2018; Siegert, 2017). Discrimination (also known
as resolution) measures how well separated the forecast probabilities are for observed presences
versus observed absences (higher scores are better). Miscalibration (also known as calibration)
measures how far the forecast probabilities deviate from the observed frequencies on average
(lower scores are better). We ignore the third decomposition, uncertainty, which measures
uncertainty inherent to the data.

Our primary model comparison examines the relative abilities of the initial GHSP-data-only
model, and the model updated with GA data, to predict GA data. This is because the GA data is
assumed closer to the population we aim to represent with this model – but we also explore other
predictions and discuss this assumption. When comparing model performance metrics, we
generate them using leave-one-group-out analysis (Vehtari, 2025), a hierarchical leave-one-out
cross-validation where a model is never evaluated for predicting data it is trained with.

Results

Data description

Gardens were distributed throughout the United Kingdom (Figure 2). Descriptive statistics for
model variables are in Table S4 (additional details of all results are in Supporting Information).
Key figures include that sparrows were observed in 85% of GA gardens and in mean 22% of GA
map grid cells. Examples of user-created maps are in Figure 3.

238
Figure 2. Locations of the 71 Garden Advice (GA) gardens within the UK, snapped to a 5 km
grid for privacy. Rural / urban classification data (ONS Geography, 2025) covers England and
Wales only and is sometimes unavailable for other reasons.

Figure 3. Six user-provided GA garden maps, selected as those nearest the duodeciles (0%,
20%, … 100%) of the Brier score discrimination decomposition distribution, to provide
examples of where sparrows are poorly and well predicted by the final model. Dotted edges
indicate buildings not drawn by users but automatically imported to calculate roof proximity.

Effect of model updating on model parameters

Figure 4 shows how fixed model parameters were updated (and indicates very good convergence
between INLA and MCMC models). For trees, bushes, and hedges (cover and proximity), the
model became more confident of more positive effects. The effect size differences were small,
but the model updated from being only weakly confident to being very confident in positive
effects. There was a salient change for grass, where the model updated from being weakly

confident in a negative effect to slightly more confident in a positive effect (with a 91%
probability this represents a real difference[1]).

For roofs and artificial surfaces, on the other hand, the model decreased its estimates of impact
(positive and negative, respectively). Here, the most salient change was for roof proximity, with
a 98% probability that this represents a real difference. This change also implied a reduction in
confidence about the effect (unique to this parameter).

Figure 4. Posterior distributions for model parameters before and after updating of GHSP-data
model with GA data. Variable units are metres or cell proportion covered, as appropriate.

---

[1] The probability that a point from the first posterior differs from a point in the second posterior in the same direction as the observed difference in medians.

Effect of model updating on predictive power

Updating with GA data improved the model's overall ability to predict the GA data, but decreased its ability to predict the GHSP data, and thus its ability to predict all data together was unchanged (Figure 5). The trends for both Brier decompositions are in the same direction as the overall Brier score, and due to greater uncertainty when decomposing (credible intervals in Figure 5), it's not possible to determine from this analysis whether the overall improvement relates primarily to discrimination or miscalibration or both. The only relevant preregistered hypothesis, which predicted improvements in overall Brier score and its decompositions when predicting GA data, was thus confirmed in essence.[2]

Figure 5. Model performance (leave-one-out cross-validated) before and after the GHSP-only model was updated with GA data. Vertical scale direction is set so upwards always represents improvement. For Brier score and decompositions (left hand panels), small points and lines show performance for individual gardens and colonies, while large points and lines show mean performance. Change scores (right-hand panels) are median and 95% credible intervals.

For the current EDS application, thresholding model output to produce a presence-absence prediction is unnecessary, but in many contexts, decision making requires a best guess. We therefore explored thresholded presence-absence predictions for GA data, finding at least small improvements for the updated model, depending on threshold selection approach.

---

[2] There was one additional preregistered hypothesis, that the updated model would explain variance; this hypothesis is ignored as it is trivial given that the primary hypothesis is confirmed.

Effect consistency

An aspect of model prediction particularly relevant in decision-making contexts is effect consistency, defined as the estimated probability that a random location will display an effect in the predicted direction. This is correlated with but separable from the estimated fixed effect, because it also depends on inter-location effect heterogeneity. Figure 6 illustrates good confidence in most vegetation-related effects and effect consistencies – for example, trees are predicted to have a positive effect in approximately 80% of gardens. There is some weak evidence that the consistency of hedge effects might be slightly greater than expected, given their fixed effect sizes, although trees and bushes are estimated to be most beneficial. The equivalent figure for the GHSP-only model (Figure S2) indicates that GHSP data by itself does not enable confident predictions except for those based on roof proximity and artificial surface.

Decision-support-seeking behaviours

Participants were paid small sums to follow specific instructions within the GA system, necessary to provide data that could support model updating. They were not instructed to obtain environmental decision support by running models to see their garden's scores for sparrows or carbon. However, 56% examined their sparrow scores, 59% examined their carbon scores, 76% examined at least one of those, 18% planned changes to their garden using planning mode, and 11% examined how these changes affected scores.

300   Figure 6. Effect consistency and standardised effect size across predictors for GA-updated
301   MCMC model. Effect sizes are standardized by multiplying regression parameters by the
302   pooled within-location standard deviation for each predictor.

Discussion

Observed model improvements

The REDS approach was successfully demonstrated in a proof-of-concept system, Garden
Advice (GA). Users contributed habitat and species occurrence data, refining the decision
support available to them by increasing the accuracy of the model that generated predictions. The
Brier score improvement of a little over 0.01 corresponds to probability predictions improving by
a little over 0.1 on average – a non-trivial improvement – and threshold approaches also revealed
meaningful improvements. As discussed in Supporting Information, the Brier score improvement
almost certainly represents an improvement in discrimination, because our model offsetting
approach makes calibration improvement unlikely.

Whether these improvements reflect reality depends on the veracity of the contributed data, but
literature suggests that the most marked parameter change represents real ecological information.
Weak confidence in a negative association for grass shifted to weak confidence in a positive
association. Consistent evidence includes anecdotal (Bradbury, 2024; Teo, 2023) and systematic
(Vincent, 2005) observations of House Sparrows indicating that although grass is not a preferred

foraging habitat, they can obtain seeds and invertebrates there. In Spain, reduction of grass in
urban parks has been associated with House Sparrow decline (Bernat-Ponce et al., 2020).

Other studies indicate negative associations with grass, however (Havlíček, 2021; Weir, 2015).
One likely resolution to this conflicting evidence relates to temporal variation. House Sparrow
preference for grass has been observed to fluctuate month-on-month and year-on-year (Vincent,
2005), so grass may be preferred when temporary conditions increase its relative benefits. All
data here was from May 2025, but REDS systems are well suited to provide data over time and
across circumstances to support a richer model that could address issues of seasonality and other
reasons for heterogenous effects.

Of the remaining changes in model parameters, most represented increased confidence in slightly
larger positive effects. This was the case for trees, bushes, and hedges (cover and proximity), and
this is also supported by the general literature. There were weak indications that the effects of
hedges may be more consistent across locations than would be predicted by effect size alone,
according with previous observations of the great importance of hedgerows for birds and
reinforcing the importance of conserving hedges (Glasgow House Sparrow Project, n.d.; Hinsley
& Bellamy, 2000). Such effect consistency estimates are only available when sampling large
numbers of independent locations, as they are transformations of inter-location effect
heterogeneity. REDS and other citizen science systems are well suited to collecting such data
(Garretson et al., 2023). Effect consistency is important for EDS as it is appropriate for decision
support to emphasise consistent effects, independently of effect size, because consistent effects
are robust to local idiosyncrasies. We are unaware of previous uses of our specific method for
intuitively presenting effect heterogeneity, by transforming to the percentage of locations where
the effect is expected to apply. This presentation method could be generally beneficial for
decision support.

The GHSP-only model's initial strong confidence in the association of sparrows with roof
proximity was removed by updating the model with GA data. This is unlikely to be realistic,
given the close affiliation of House Sparrows with human dwellings (Shaw et al., 2008). This
discrepancy highlights a key issue: the accuracy of GA user observations was not evaluated, and
the result likely reflects observation bias. Birds on or near roofs are harder to see, especially for
those observing from indoors. Another study that solicited non-expert animal observations from
UK gardens found that although participants could provide accurate habitat descriptions,
hedgehog presence-absence reports were uncorrelated with objective data (Williams et al., 2015).
Hedgehogs are harder than sparrows for humans to detect, but this nevertheless provides a
sobering reminder of the necessity for independent confirmation of citizen science data.

REDS – broader implications

Reviews indicate that citizen science data is often reliable, and measures such as additional
training provide benefits (Aceves-Bueno et al., 2017; Backstrom et al., 2024; Johnston et al.,
2023; Kosmala et al., 2016). Further, although our study involved data solicitation from a wide
public, in practice EDS systems are often used by land managers such as farmers, who might
provide more accurate observations. Nevertheless, the next stage in REDS concept validation
must involve independent verification of data quality. Such model validation is one of the factors
most associated with EDS system uptake (Walling & Vaneeckhaute, 2020).

EDS systems predict the effects of potential changes to local conditions so that management can
be planned. However, standard EDS relies on static models that may not optimally reflect the

reality experienced by current users. There are many reasons why relations between
environmental variables may not be static, because they may be moderated by additional
variables which are themselves changing (such as climate, Bueno de Mesquita et al., 2021).
REDS systems can better remain up to date, in essence informing users "if you change your
conditions in such a way, your situation will more closely resemble the situation recently
reported by certain other users, so your observations will become more like theirs." A concrete
example from this study is that, thanks to the provision of user data that updated the model to
separate grass from artificial surfaces, by the end of the study period the system advised that
lawns are better for sparrows than decking or paving. Although our users arrived at the system
because of payment, and were never instructed to use the EDS functions, most of them did
request model predictions, so later users did benefit from this improved advice.

The REDS design concept joins that of the digital twin as a system using modern technology to
close an information loop to improve model predictions (De Koning, 2025). REDS systems
could also be digital twins, although REDS does not necessarily involve real-time modelling of
an ongoing phenomenon – the defining feature of digital twins – and digital twins don't
necessarily directly support environmental decision making. Despite recent reviews of digital
twins in environmental science (Hazeleger et al., 2024; Trantas et al., 2023), we are unaware of
any digital twin system yet operating to offer practical decision support for ecosystem managers.
These reviews list appreciable practical challenges for the digital twin approach – perhaps in
some contexts, the less technically ambitious REDS approach could offer some of the same
benefits more straightforwardly.

Future REDS systems would benefit from more complex models. Habitat association models do
not consider causal mechanisms and do not necessarily predict long-term species health, because
local observations may not predict longer term demographic changes (Bacon et al., 2017;
Matthiopoulos et al., 2019; Regos et al., 2021). That said, reviews indicate that although there
are important exceptions, the availability of preferred habitats does tend to predict species
abundance (Boyce et al., 2016; Weber et al., 2017). In principle, the REDS citizen science
approach could also track demographic change, as one route to support more complex and
accurate models.

Further development of the REDS concept will benefit from closer examination of the potential
social synergies generated by combining citizen science and EDS. There is always potential for
useful synergy when closing an information loop, allowing feedback mechanisms to operate
(Wiener, 1948). For REDS, some of these synergies might relate to the strong human motivation
towards reciprocity (Bowles & Gintis, 2011). It is likely that public data sharing that already
exists on platforms such as iNaturalist and PurpleAir is motivated in large part in this way
(Metzger & Runge, 2023; Pai & Tsai, 2016).

Examples of existing initiatives that could become REDS include membership-organisation
citizen-science initiatives, such as the British Trust for Ornithology's Garden Birdwatch
(Morrison et al., 2014), schemes associated with the Bumblebee Conservation Trust (Sharma et
al., 2019), or the Game & Wildlife Conservation Trust's Partridge Count Scheme (Ewald et al.,
2009). Participation could be increased, and relevant environmental management behaviour
could be improved, as citizens receive tailored advice as an immediate and integral part of
reporting their contributions.

Additional motivation for engagement may come from participants more directly seeing how
individual contributions improve centralised knowledge. Stakeholder engagement is a key
determinant of EDS system success (Walling & Vaneeckhaute, 2020). Provided that careful
consideration is given to the structure of the initial model, REDS can save time for scientists
operating such schemes, as analysing and reporting the data is an automated and integrated part
of the data-collection process itself. Like digital twins, the REDS concept is at an early stage of
development, and the next stages must also focus on how this can be achieved without
sacrificing the quality of analysis and conclusions.

In summary, the sense that one is helping at the same time as being helped could be a strong
factor for further activating the synergies between EDS and citizen science that this study
indicates REDS can provide. These synergies offer positive feedback loops potentially
generating a virtuous circle of greater user engagement, data provision, advice quality, and
advice uptake.

References

Aceves-Bueno, E., Adeleye, A. S., Feraud, M., Huang, Y., Tao, M., Yang, Y., & Anderson, S. E.
(2017). The Accuracy of Citizen Science Data: A Quantitative Review. *The Bulletin of*
the Ecological Society of America, *98*(4), 278-290.
Albert, D. A., & Smilek, D. (2023). Comparing attentional disengagement between Prolific and
MTurk samples. *Scientific Reports*, *13*(1), 20574.
Backstrom, L. J., Callaghan, C. T., Leseberg, N. P., Sanderson, C., Fuller, R. A., & Watson, J. E.
M. (2024). Assessing adequacy of citizen science datasets for biodiversity monitoring.
Ecology and Evolution, *14*(2), e10857.
Bacon, L., Hingrat, Y., Jiguet, F., Monnet, A.-C., Sarrazin, F., & Robert, A. (2017). Habitat
suitability and demography, a time-dependent relationship. *Ecology and Evolution*, *7*(7),
2214-2222.
Bernat-Ponce, E., Gil-Delgado, J. A., & López-Iborra, G. M. (2020). Replacement of semi-
natural cover with artificial substrates in urban parks causes a decline of house sparrows
Passer domesticus in Mediterranean towns. *Urban Ecosystems*, *23*(3), 471-481.
Bondia-Barceló, J., Castellà-Roca, J., & Viejo, A. (2016). Building Privacy-Preserving Search
Engine Query Logs for Data Monetization. 2016 Intl IEEE Conferences on Ubiquitous
Intelligence & Computing, Advanced and Trusted Computing, Scalable Computing and
Communications, Cloud and Big Data Computing, Internet of People, and Smart World
Congress (UIC/ATC/ScalCom/CBDCom/IoP/SmartWorld),
Bowles, S., & Gintis, H. (2011). *A cooperative species: Human reciprocity and its evolution*.
Princeton University Press.
Boyce, M. S., Johnson, C. J., Merrill, E. H., Nielsen, S. E., Solberg, E. J., & van Moorter, B.
(2016). REVIEW: Can habitat selection predict abundance? *Journal of Animal Ecology*,
85(1), 11-20.
Bradbury, K. (2024). *The house sparrow (Passer domesticus)*. https://www.encounter-
nature.com/features/the-house-sparrow-passer-domesticus
Bressler, R. D. (2021). The mortality cost of carbon. *Nature Communications*, *12*(1), 4467.
Bueno de Mesquita, C. P., White, C. T., Farrer, E. C., Hallett, L. M., & Suding, K. N. (2021).
Taking climate change into account: Non-stationarity in climate drivers of ecological
response. *Journal of Ecology*, *109*(3), 1491-1500.

Burris, M., Khan, M., & Johnson, J. (2023). *"Is That Route Really the Most Fuel-Efficient?"* (National Institute For Congestion Reduction Research Reports, Issue 17). https://doi.org/10.5038/CUTR-NICR-Y3-4-8

Cole, E., Van Horn, G., Lange, C., Shepard, A., Leary, P., Perona, P., Loarie, S., & Mac Aodha, O. (2023). *Spatial Implicit Neural Representations for Global-Scale Species Mapping* Proceedings of the 40th International Conference on Machine Learning, Proceedings of Machine Learning Research. https://proceedings.mlr.press/v202/cole23a.html

Cool Farm. (2025a). *Cool Farm*. Retrieved 2025-10-03 from https://coolfarm.org/

Cool Farm. (2025b). *Cool Farm: Frequently Asked Questions*. Retrieved 2025-10-03 from https://coolfarm.org/frequently-asked-questions/

De Koning, K. (2025). The crane radar: Development and deployment of an operational eco-digital twin. *Ecological Informatics*, *85*, 102938.

De Laet, J., & Summers-Smith, J. D. (2007). The status of the urban house sparrow Passer domesticus in north-western Europe: a review. *Journal of Ornithology*, *148*(2), 275-278.

Ewald, J. A., Kingdon, N. G., & Santin-Janin, H. (2009). The GWCT Partridge Count Scheme: a volunteer-based monitoring and conservation promotion scheme. In S. B. Cederbaum, B. C. Faircloth, T. M. Terhune, J. J. Thompson, & J. P. Carroll (Eds.), *Gamebird 2006: Quail VI and Perdix XII* (pp. 27-37). Warnell School of Forestry and Natural Resources.

Frick, V., Matthies, E., Thøgersen, J., & Santarius, T. (2021). Do online environments promote sufficiency or overconsumption? Online advertisement and social media effects on clothing, digital devices, and air travel consumption. *Journal of Consumer Behaviour*, *20*(2), 288-308.

Garretson, A., Cuddy, T., Duffy, A. G., iNaturalist Citizen, S., & Forkner, R. E. (2023). Citizen science data reveal regional heterogeneity in phenological response to climate in the large milkweed bug, Oncopeltus fasciatus. *Ecology and Evolution*, *13*(7), e10213.

Ginsberg, J., Mohebbi, M. H., Patel, R. S., Brammer, L., Smolinski, M. S., & Brilliant, L. (2009). Detecting influenza epidemics using search engine query data. *Nature*, *457*(7232), 1012-1014.

Glasgow House Sparrow Project. (n.d.). *Help your sparrows*. Retrieved 2025-12-15 from https://www.housesparrowscience.com/help-your-sparrows/

Hanson, H. E., Mathews, N. S., Hauber, M. E., & Martin, L. B. (2020). The house sparrow in the service of basic and applied biology. *Elife*, *9*.

Havlíček, J. (2021). *The breeding and foraging ecology of the House Sparrow in rural and urban environments* [PhD Thesis, University of South Bohemia in České Budějovice].

Hazeleger, W., Aerts, J. P. M., Bauer, P., Bierkens, M. F. P., Camps-Valls, G., Dekker, M. M., Doblas-Reyes, F. J., Eyring, V., Finkenauer, C., Grundner, A., Hachinger, S., Hall, D. M., Hartmann, T., Iglesias-Suarez, F., Janssens, M., Jones, E. R., Kölling, T., Lees, M., Lhermitte, S., . . . Vossepoel, F. C. (2024). Digital twins of the Earth with and for humans. *Communications Earth & Environment*, *5*(1), 463.

Hinsley, S. A., & Bellamy, P. E. (2000). The influence of hedge structure, management and landscape context on the value of hedgerows to birds: A review. *Journal of Environmental Management*, *60*(1), 33-49.

Johnston, A., Matechou, E., & Dennis, E. B. (2023). Outstanding challenges and future directions for biodiversity monitoring using citizen science data. *Methods in Ecology and Evolution*, *14*(1), 103-116.

Kayatz, B., Baroni, G., Hillier, J., Lüdtke, S., Heathcote, R., Malin, D., van Tonder, C., Kuster, B., Freese, D., Hüttl, R., & Wattenbach, M. (2019). Cool Farm Tool Water: A global on-line tool to assess water use in crop production. *Journal of Cleaner Production*, *207*, 1163-1179.

Kenward, B. (2024). *The 1000-ton rule, which emphasises the mortality cost of carbon, can be an effective persuasion tool, including for use by activists.* https://www.benkenward.com/articles/the_1000_ton_rule_can_be_an_effective_persuasion_tool.pdf

Kenward, R., Papathanasiou, J., Manos, B., & Arampatzis, E. (Eds.). (2013). *Transactional Environmental Support System Design: Global Solutions*. IGI Global.

Kosmala, M., Wiggins, A., Swanson, A., & Simmons, B. (2016). Assessing data quality in citizen science. *Frontiers in Ecology and the Environment*, *14*(10), 551-560.

Martin, K. D., & Murphy, P. E. (2017). The role of data privacy in marketing. *Journal of the Academy of Marketing Science*, *45*(2), 135-155.

Matthiopoulos, J., Field, C., & MacLeod, R. (2019). Predicting population change from models based on habitat availability and utilization. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1901), 20182911.

McIntosh, B. S., Ascough, J. C., Twery, M., Chew, J., Elmahdi, A., Haase, D., Harou, J. J., Hepting, D., Cuddy, S., Jakeman, A. J., Chen, S., Kassahun, A., Lautenbach, S., Matthews, K., Merritt, W., Quinn, N. W. T., Rodriguez-Roda, I., Sieber, S., Stavenga, M., . . . Voinov, A. (2011). Environmental decision support systems (EDSS) development – Challenges and best practices. *Environmental Modelling & Software*, *26*(12), 1389-1402.

Merkle, E. C., & Hartman, R. (2018). Weighted Brier score decompositions for topically heterogenous forecasting tournaments. *Judgment and Decision Making*, *13*(2), 185-201.

Metzger, F. M., & Runge, G. (2023). Reciprocity of Data Sharing Infrastructures: A Conceptual Norms Framework. In A. Roßnagel, M. Friedewald, C. L. Geminn, M. Karaboga, & S. Schindler (Eds.), *Data Sharing – Datenkapitalismus by Default?* https://doi.org/10.24406/publica-2072

Morrison, C. A., Robinson, R. A., Leech, D. I., Dadam, D., & Toms, M. P. (2014). Using citizen science to investigate the role of productivity in House Sparrow Passer domesticus population trends. *Bird Study*, *61*(1), 91-100.

Natural Capital Project. (2025). InVEST 3.16.2: Stanford University, University of Minnesota, Chinese Academy of Sciences, The Nature Conservancy, World Wildlife Fund, Stockholm Resilience Centre and the Royal Swedish Academy of Sciences. https://doi.org/10.60793/natcap-invest-3.16.2

Nghiem, L. T. P., Papworth, S. K., Lim, F. K. S., & Carrasco, L. R. (2016). Analysis of the Capacity of Google Trends to Measure Interest in Conservation Topics and the Role of Online News. *PLOS ONE*, *11*(3), e0152802.

ONS Geography. (2025). *Rural Urban Classification (2021) of Output Areas in EW*. Retrieved 2025-05-19 from https://geoportal.statistics.gov.uk/datasets/ons::rural-urban-classification-2021-of-output-areas-in-ew/about

Pai, P., & Tsai, H.-T. (2016). Reciprocity norms and information-sharing behavior in online consumption communities: An empirical investigation of antecedents and moderators. *Information & Management*, *53*(1), 38-52.

Papathanasiou, J., & Kenward, R. (2014). Design of a data-driven environmental decision support system and testing of stakeholder data-collection. *Environmental Modelling & Software*, *55*, 92-106.

Pearce, J. M., & Parncutt, R. (2023). Quantifying Global Greenhouse Gas Emissions in Human Deaths to Guide Energy Policy. *Energies*, *16*(16).

Peer, E., Rothschild, D., Gordon, A., Evernden, Z., & Damer, E. (2022). Data quality of platforms and panels for online behavioral research. *Behavior Research Methods*, *54*(4), 1643-1662.

PurpleAir. (2025). *PurpleAir*. Retrieved 2025-10-03 from https://purpleair.com/

Regos, A., Tapia, L., Gil-Carrera, A., & Domínguez, J. (2021). Caution Is Needed When Using Niche Models to Infer Changes in Species Abundance: The Case of Two Sympatric Raptor Populations. *Animals*, *11*(7), 2020.

RHS. (2025). *RHS State of Gardening Report 2025*. https://www.rhs.org.uk/about-us/what-we-do/rhs-state-of-gardening

Robinson, D. L., Goodman, N., & Vardoulakis, S. (2023). Five Years of Accurate PM2.5 Measurements Demonstrate the Value of Low-Cost PurpleAir Monitors in Areas Affected by Woodsmoke. *International Journal of Environmental Research and Public Health*, *20*(23).

Sandhaus, S., Kaufmann, D., & Ramirez-Andreotta, M. (2019). Public participation, trust and data sharing: gardens as hubs for citizen science and environmental health literacy efforts. *International Journal of Science Education, Part B*, *9*(1), 54-71.

Sharma, N., Greaves, S., Siddharthan, A., Anderson, H. B., Robinson, A.-M., Colucci-Gray, L., Wibowo, A. T., Bostock, H., Salisbury, A., & Roberts, S. (2019). From citizen science to citizen action: analysing the potential for a digital platform to cultivate attachments to nature. *Journal of Science Communication*, *18*(01), A07.

Shaw, L. M., Chamberlain, D., & Evans, M. (2008). The House Sparrow Passer domesticus in urban areas: reviewing a possible link between post-decline distribution and human socioeconomic status. *Journal of Ornithology*, *149*(3), 293-299.

Siegert, S. (2017). Simplifying and generalising Murphy's Brier score decomposition. *Quarterly Journal of the Royal Meteorological Society*, *143*(703), 1178-1183.

Teo, L. W. (2023). *House Sparrows dig and chew on grass stolons*. https://besgroup.org/2023/05/18/house-sparrows-dig-chew-grass-stolons/

Trantas, A., Plug, R., Pileggi, P., & Lazovik, E. (2023). Digital twin challenges in biodiversity modelling. *Ecological Informatics*, *78*, 102357.

Vehtari, A. (2025). *Can cross-validation be used for hierarchical / multilevel models?* https://users.aalto.fi/~ave/CV-FAQ.html#8_Can_cross-validation_be_used_for_hierarchical__multilevel_models

Vincent, K. (2005). *Investigating the causes of the decline of the urban house sparrow Passer domesticus in Britain* [PhD Thesis, De Montfort University, Leicester].

Walling, E., & Vaneeckhaute, C. (2020). Developing successful environmental decision support systems: Challenges and best practices. *Journal of Environmental Management*, *264*, 110513.

Weber, M. M., Stevens, R. D., Diniz-Filho, J. A. F., & Grelle, C. E. V. (2017). Is there a correlation between abundance and environmental suitability derived from ecological niche modelling? A meta-analysis. *Ecography*, *40*(7), 817-828.

Weir, J. E. S. (2015). *Urban green space management for invertebrates and house sparrows*
[PhD Thesis, Imperial College London, Department of Ecology and Evolution].
Wiener, N. (1948). *Cybernetics: Or Control and Communication in the Animal and the Machine*.
MIT University Press.
Williams, R. L., Stafford, R., & Goodenough, A. E. (2015). Biodiversity in urban gardens:
Assessing the accuracy of citizen science data on garden hedgehogs. *Urban Ecosystems*,
18(3), 819-833.