Ex Vivo Assay for Organ-Specific Cancer Cell Invasion

François Tyckaert[1a], Paul Frieso Göddertz[1a], Maria Reichhold[1], Bettina Sarg[2], Klaus Faserl[2], Pere Patón González[1], Felix Eichin[3], Andreas Villunger[3], Steffen Ormanns[4], Stefan Redl[5], Julia Hofmann[6], Theresa Hautz[6], Francesco Baschieri[1b]

[1] Institute of Pathophysiology, Biocenter, Medical University of Innsbruck, Innsbruck, Austria.
[2] Medical University of Innsbruck, CCB-Biocenter, Institute of Medical Biochemistry, Protein Core Facility, Innsbruck, Austria.
[3] Institute for Developmental Immunology, Biocenter, Medical University of Innsbruck, Innsbruck, Austria.
[4] Institute of General Pathology, Medical University Innsbruck, Innsbruck, Austria.
[5] Institute of Neuroanatomy, Medical University of Innsbruck, Innsbruck, Austria.
[6] Department of Visceral, Transplant and Thoracic Surgery, Center of Operative Medicine, organLife Laboratory and D. Swarovski Research Laboratory, Medical University of Innsbruck, Innsbruck, Austria.
[a] These authors contributed equally
[b] Correspondence to francesco.baschieri@i-med.ac.at - Medical University Innsbruck - CCB - Innrain 80 - 6020 Innsbruck, Austria - +43 512 9003 70970

Abstract

Metastasis is the leading cause of cancer-related mortality, yet experimental models inadequately recapitulate the tissue-specific microenvironments that shape metastatic dissemination. In vivo systems provide physiological relevance but are poorly suited for mechanistic studies and screening, whereas conventional in vitro assays lack the organ-specific extracellular matrix (ECM) context that critically influences invasive behavior. To address this gap, an ex vivo method is established that balances biological relevance with scalability, affordability, and ease of use.

Mild detergent decellularization of mouse organs followed by vibratome slicing generates optically transparent lung, liver, and intestine scaffolds that preserve native ECM architecture, mechanics, and composition. These organ-derived matrices are readily integrated into standard microfluidic channels and analyzed using conventional fluorescence microscopy, enabling quantitative assessment of cancer cell invasion without specialized infrastructure.

Benchmarking with breast cancer cell lines of well-defined invasive capacity confirms the robustness and biological relevance of the platform. Non-invasive MCF7 cells fail to infiltrate any scaffold, whereas highly invasive MDA-MB-231 cells display organ-specific invasion, preferentially penetrating lung and liver ECM while showing virtually no invasion of intestinal scaffolds, consistent with clinically observed metastatic tropism. Quantitative invasion rates closely match values reported in vivo by intravital microscopy.

Overall, this ex vivo system provides an accessible and scalable platform to study ECM-driven determinants of metastatic invasion while reducing reliance on animal models.

Keywords: Cancer; Metastasis; Invasion; Extracellular matrix; Tissue Decellularization

1. Introduction

Cells within our body are embedded in a complex meshwork of proteins known as the extracellular matrix (ECM) [1], [2], [3]. Although fibrillar collagens constitute the bulk of the ECM, with collagen I alone accounting for up to 12–17% of total mouse protein mass [4], the composition, architecture, and mechanical properties of the ECM vary markedly between organs, reflecting their distinct physiological requirements [3], [5].

Beyond its structural role, the architecture and mechanical properties of the ECM guide cell movement, determine regions of tissue growth, and even dictate where cellular extrusion occurs [6], [7], [8], [9], [10]. This regulation is mediated by a finely tuned interaction between ECM components and cell adhesion molecules (CAMs), such as integrins, on the cell surface [11].

The importance of cell–ECM interactions has long been recognized in cancer biology. More than a century ago, Paget's "seed and soil" hypothesis proposed that metastatic dissemination depends on the compatibility between disseminating cancer cells and the microenvironment of target organs. Consistent with this idea, clinical and experimental evidence demonstrates that metastases arising from distinct primary tumors exhibit reproducible organ-specific dissemination patterns, a phenomenon known as organotropism [12], [13], [14]. While soluble factors and stromal cells contribute to metastatic niche formation, the ECM represents a fundamental and relatively stable component of the metastatic soil.

Metastasis is the leading cause of cancer-related mortality, and improving our understanding of this process remains a central challenge in efforts to enhance patient outcomes. Despite its importance, ECM-driven organotropism remains difficult to investigate experimentally. *In vivo* mouse models offer physiological relevance but are time-consuming, costly, and challenging to dissect mechanistically. Conversely, conventional *in vitro* assays are easier to interpret but fail to recapitulate the tissue-specific ECM context that critically shapes metastatic behavior, contributing to poor predictive power in preclinical studies.

There is a pressing need for improved *in vitro* or *ex vivo* models that better mimic the complexity of metastatic processes preserving the complexity of organ-specific ECMs. To address this gap, we developed an *ex vivo* invasion assay based on decellularized mouse organs. By combining mild detergent-based decellularization with vibratome slicing, we generate thin, optically transparent tissue sections that retain native ECM architecture, mechanics, and composition. These scaffolds can be integrated into standard microfluidic channels and imaged using conventional microscopy, enabling quantitative and comparative analysis of cancer cell invasion across distinct organ-derived ECMs.

2. Materials and methods

2.1. Tissue decellularization

Immediately after collection, organs from healthy, 6-10 week old, male BL6N mice were quickly washed in PBS to eliminate excess blood. Importantly, the mice had been euthanized for unrelated procedures, and only surplus organs were used in this study. Liver lobes were separated, and intestine lumens (duodenums) were flushed multiple times with PBS. Organs were then immediately submerged in 50 ml of PBS containing 0.1% SDS and incubated rotating at room temperature (RT). The solution was changed every two days until full decellularization of the organs, which could be evaluated empirically by observing the organs through a source of light to assess their transparency. Approximately, 15 days were needed to decellularize liver and lungs, while gut decellularization was achieved in 7 days.
SDS being an ionic detergent, a 3h wash with the non-ionic detergent Triton X-100 (0.05%) was used to wash away SDS, followed by 4 days washing in 50 ml H2O, with H2O changed daily.
Organs were then stored in PBS until slicing. Quantification of residual SDS was performed using a colorimetric assay based on a carbocyianine dye (Stains all - Sigma-Alfrich - Cat. nr. E9379-1G) as previously described [15].

2.2. Tissue slicing

Decellularized organs were then included in low temperature melting agarose at 1% for liver and lung, and 3% for intestine. Organs were sliced with the help of an Alabama R&D Tissue Slicer (Alabama R&D, Munford, AL, USA) using stainless steel razor blades (Personna Medical, Stainton, VA, USA) in PBS. Native organs were sliced in ice cold PBS in the presence of proteases inhibitors. Slices of 500 µm were recovered in PBS and incubated at 37°C for 30 min to eliminate agarose residues, then stored at 4°C in PBS. Of note, as organs are likely to experience some minor volume changes, if the final volume of the slices is important, slicing has to be performed on already decellularized organs.

2.3. Stiffness measurements

A Chiaro Nanoindenter (Optics11 Life) was used to measure the mechanical properties of the tissues. Vibratome cut slices of tissues were fixed on a plastic tissue culture dish with the help of super-glue, then submerged into PBS. For fresh tissues, proteases inhibitors and propidium iodide were added to the PBS to prevent and monitor cellular damage. Fresh tissues were analyzed within 8 hours of collection, a time window during which minimal to no cell death was observed. Measurements were performed with a rounded-tip probe with 27.5 µm diameter and stiffness of 0.5 N/m. Indentations of 3 µm were performed with an approach speed of 3000 nm/s. For each tissue slice, at least 36 measurements spaced of 100 µm were acquired. Curves were fitted using the Prova software from Optics11 life. The upper 1000 nm of indentation were fitted using Hertzian contact model and assuming Poisson's ratio of 0.5 for the calculation of the effective Young's elastic modulus.

2.4. Histology

Native or decellularized organs were fixed with PFA 4% for 24h and embedded in paraffin. Four micrometer thick sections of paraffin embedded tissues were stained using automated routine hematoxylin-eosin staining (Sakura Prisma, Sakura Finetek Europe, Umkirch, Germany), the connective tissue stainings van Gieson (vG), as well as chromotrop-anilin blue (CAB) on an automated special staining device (Ventana BenchMark, Roche, Tucson, AZ, USA) according to the manufacturer's instructions. Slides were scanned using a Leica Aperio GT450 DX. Acquired images were cropped and further processed with the QuPath software and FIJI.

2.5. Scanning Electron Microscopy

Decellularized organs were fixed in 2.5% GA and 2% PFA diluted in sodium cacodylate buffer (0.1M, pH = 7.4) for several days and postfixed in 4% Glutaraldehyde in 0.1M PB, pH 7.4 overnight. After washing with PB, samples were cryoprotected overnight in 30% sucrose, embedded in OCT and the surface was trimmed and cut using a Leica Cryocut CM3050 S. After washing in 0.1M PB, samples were postfixed in 1% OsO_4 in A.d., dehydrated in a standard ethanol series and critical point dried using a BAL-TEC CPD 030 critical point dryer, mounted with carbon adhesive tape and Leit-C on aluminium stubs and sputtered with 10 nm gold using a BAL-TEC MED 020 coating system. Scanning electron microscopic images were taken with a Tescan-Clara, at the Zoology department, University of Innsbruck.

2.6. Proteomics

Samples were dried in a SpeedVac. Each individual sample was recovered in 40 µL of 100 mM ammonium bicarbonate (NH_4HCO_3), 1% deoxycholate (DOC) and 50 mM dithiothreitol and incubated at 56°C for 1 h. The samples were alkylated with 5 µl of 500 mM IAA solution for 30 min at RT in the dark. Samples were then dried again in a SpeedVac, resuspended in 50 µL of trypsin solution (10 ng/µL) in 50 mM NH_4HCO_3 and incubated overnight at 37°C [16]. By adding 10 µl concentrated formic acid samples were acidified (pH lower than 2) to precipitate deoxycholate. Afterwards, they were centrifuged for 10 min at 36000g and transferred to a 200 µl low binding Eppendorf tube.

For LC-MS analysis an UltiMate 3000 nano-HPLC system coupled to an Orbitrap Eclipse mass spectrometer (Thermo Scientific, Bremen, Germany) was used. The peptides were separated on a homemade frit-less fused-silica micro-capillary column (75 µm i.d. x 360 µm o.d. x 20.5 cm length) packed with 2.4 µm reversed-phase C18 material (Reprosil). Solvents for HPLC were 0.1% formic acid (solvent A) and 0.1% formic acid in 85% acetonitrile (solvent B). The gradient profile was as follows: 0-4 min, 4% B; 4-117 min, 30% B; 117-127 min, 100% B. The flow rate was 250 nL/min [17].

The Orbitrap Eclipse mass spectrometer was operating in the data dependent mode with a cycle time of one second. Survey full scan MS spectra were acquired from 375 to 1500 m/z at a resolution of 60 000, a maximum injection time (IT) of 118 ms and automatic gain control (AGC) target 400 000. The MS2 Spectrum was measured in the Orbitrap analyzer at a resolution of 15000 with an isolation window of 1.2 mass-to-charge ratio (m/z), a maximum IT of 22 ms, and AGC target of 50 000. The selected isotope patterns were fragmented by higher-energy collisional dissociation with normalized collision energy of 30%.

Data analysis was performed using Proteome Discoverer 3.1 (Thermo Scientific) with Sequest search engine. The raw files were searched against UniProt mouse-2024_UP000000589_10090 database. Precursor and fragment mass tolerance was set to 10 ppm and 0.02 Da, respectively, and up to two missed cleavages were allowed. Carbamidomethylation of cysteine was set as static modification, oxidation of Methionine, Galactosyl and Glucosylgalactosyl were set as variable modifications. Peptide identifications were filtered at 1% false discovery rate. For label free quantification the protein abundances were normalized to total protein amount.

Three biological replicates were analyzed for each decellularized organ. Only proteins detected in all three replicates were retained for further analysis. To generate organ-specific protein lists, we additionally applied a threshold of ≥0.01% of the total protein mass. Gene ontology (GO) analysis was then performed online using String database version 12.0 with the following parameters: False Discovery Rate (FDR) ≤ 1E-5; Signal ≥ 2; Strength ≥ 1; Minimum count = 10.

2.7. Cell culture and in vitro testing

MDA-MB-231 cells (a gift from G. Montagnac, Inst. Gustave Roussy, Villejuif, France; ATCC HTB-26). MCF7 and HEK293T cells were obtained from H. Farhan lab. All cell lines were grown in DMEM Glutamax supplemented with 10% FCS and 100 U ml–1 penicillin/streptomycin at 37°C in 5% CO2. All cell lines were regularly tested by PCR for the absence of mycoplasma contamination. For channel invasion experiments, Primocin (Invivogen - Cat. ant-pm-05) was also added to the medium at the final concentration of 100µg/ml. Lentiviruses were produced in HEK293T cells using the packaging plasmid psPAX2 and the VSV-G envelope expressing plasmid pMD2.G according to the Trono lab protocol. Stable cell lines expressing empty pLVTHM (a gift from Didier Trono, Addgene plasmid #12247) were obtained by lentiviral transduction, followed by sorting via fluorescence-activated cell sorting.

2.8. 2D cell migration

For wound healing, Ibidi 2 well culture inserts (cat. Nr. 80209) were used. Briefly, 30.000 cells per well were seeded on uncoated plates or on collagen I coated (50 µg/ml) plates. Upon cell adhesion the inserts were removed, and cell migration across the wound was imaged (1 image every 20 min). Imaging was performed on an epifluorescence microscope (Leica Microsystems Ltd., Wetzlar, Germany) through a 10✕ (Fluotar N.A. 0.32 PH1) objective. The microscope was equipped with an LED5 lamp from Leica, a filter cube DFT51010, an external wheel EFW for DMi8, and an sCMOS K5 camera. Wound healing closure was analyzed using FIJI.

2.9. Gelatin digestion

For Gelatin digestion, glass coverslips were activated with the help of a plasma cleaner (Harrick Plasma PDC-002-CE). Activated coverslips were incubated for 10 min with Gelatin Oregon Green™ 488 Conjugate (Invitrogen – Cat. Nr. G13186) diluted at 1 mg/ml. Coverslips were then rinsed in PBS, and glutaraldehyde was used to crosslink gelatin, followed by quenching with Sodium Borohydride. Cells were plated and incubated at 37°C for 24h. Then, cells were fixed and stained with Phalloidin iFluor 555 (Abcam - Cat. Nr. ab176756). Images were acquired using a 60✕ TIRF objective (Plan-APOCHROMAT 60✕/1.49 Oil, Nikon) mounted on an inverted

microscope (Eclipse Ti2-E; Nikon) with a spinning disc confocal unit (X-Light V2 - 60 µm pinhole), a sCMOD camera (Photometrics Prime 95B), and Lumencor Spectra III LED light source (380 –750   nm), controlled via the NIS-Elements   software (Nikon). Acquired images were analyzed with FIJI software.

2.10. Transwell invasion assays

Fluoroblok™ inserts with 8 µm membrane pores, 24 well plate format (BD Falcon – Ref. 351152) were coated with 50 µl Geltrex (Gibco – Life Technologies – Cat. Nr. A14132-02) diluted in DMEM with with 10% FCS to a final concentration of 1.2–1.8 mg/ml. Geltrex was let to polymerize for 1h at 37°C. 600 µl of DMEM supplemented with 10% FCS and 100 U ml–1 penicillin/streptomycin were added in each well, while 1E5 cells were resuspended in a volume of 300 µl of medium with 10% FCS and plated inside the coated invasion insert. Cells were then allowed to invade at 37°C for 48h. Then, cells were fixed with PFA and stained for DAPI. Imaging was performed on an epifluorescence microscope (Leica Microsystems Ltd., Wetzlar, Germany) through 63× [Numerical Aperture (N.A.) 1.40], 20× (HC N.A. 0.40 PH1), 10× (Fluotar N.A. 0.32 PH1), or 5× (PLAN 5/N.A. 0.12 PHO) objectives. The microscope was equipped with an LED5 lamp from Leica, a filter cube DFT51010, an external wheel EFW for DMi8, and an sCMOS K5 camera. A PeCon chamber was installed on the microscope to perform live imaging at 37°C with 5% CO2. The microscope was steered by the Leica dedicated LasX software with Navigator, to obtain mosaic stitched images. Using this function, the entire lower side of the Fluoroblok™ was imaged and cells having invaded were counted using FIJI.

2.11. Inclusion of organs into microchannels

Ibidi sticky-Slide VI 0.4 (Cat. Nr. 80608) and polymer Coverslips for sticky-Slides (Cat. Nr. 10813) were used to host decellularized tissue slices. Channels were. According to manufacturer's instructions, channel thickness is 400 µm ± 130 µm. Organ slices were incubated in FCS for 1h prior to being included in the open sticky slide, one slice per channel, paying attention not to completely block the channel. Channels were then sealed with the help of the Ibidi Adapter for Sticky Slides (Cat. Nr. 80043) and a clamp. DMEM supplemented with 100 U ml–1 penicillin/streptomycin and Primocin (Invivogen - Cat. Nr. ant-pm-05) at 100 µg/ml was added to the channels, and organs were let equilibrate in medium for at least 3h at 37°C in 5% CO_2. 10E5 cells were resuspended in 30µm DMEM with antibiotics and injected in the channels. Slides with organs and cells were let under agitation at RT for 1h, then non adherent cells were washed away and 100-150 µl medium with antibiotics and FCS were added in each channel. Slides were then incubated at 37°C in 5% CO_2 for the whole duration of the experiment, replacing medium every other day.

2.12. Immunostaining and imaging

PFA was added in the channels for 10 min, followed by washes with PBS. Cleaved collagen antibody (ImmunoGlobe - Cat. Nr. 0217-025) was diluted 1:100 in PBS Tx 0.3% BSA 1%, added to the channels, and incubated ON at 4°C. Secondary antibodies, as well as phalloidin and DAPI were then used prior to performing imaging.

Imaging was performed on a Leica Sp8 confocal microscope equipped with a fast resonant scanner (8000 Hz), an incubation chamber, two hybrid and three PMT detectors, using a white laser to excite all the fluorophores at the required wavelength. Imaging was performed using a 63× glycerol immersion objective (1.4 NA) or a 20x glycerol immersion objective (0.75 NA) and the 633 nm laser was used to obtain interference reflection microscopy (IRM) images. 3D reconstructions of images were prepared using Imaris Version 11.0.

2.13. Statistical analysis

Images were analyzed in FIJI. All images were background subtracted prior to performing any analysis. Graphics were prepared with Prism V10.0 and statistical analyses were performed using Prism V10.2.3. Box-plot are built as follows: The line within the box corresponds to the mean value of at least three independent experiments (n indicated in each figure legend), boxes indicate interquartile ranges (Q1 to Q3), whiskers extend to min and max values, and, eventually, outliers are represented as points outside the box-whiskers. Empty circles on top of the graphs represent the mean of each individual experiment unless otherwise stated.

3. Results

3.1. Decellularization of mouse organs

Several decellularization strategies exist, but detergent-based methods are generally considered the most effective strategy to remove cells from tissues [18]. Recently, it was shown that using low concentration of SDS allowed to better preserve tissue structure [19]. Given the importance of mechanical and structural properties of the ECM in cancer progression [20], we hence employed SDS as a decellularizing agent.

Organs from overnumbered adult, wild type mice were hence decellularized using SDS at low concentration. Efficient decellularization was visually confirmed by loss of pigmentation and transparency, making it possible to see blood vessels with the naked eye (Fig. 1A). To ensure full decellularization, DAPI stainings were performed on sections of organs, and fluorescence quantification confirmed the absence of nuclei from decellularized organ slices (Suppl. Fig. 1A-B). Additionally, hematoxylin and eosin stainings were performed on paraffin embedded slides to search for remaining cells. Visual inspection of the slices confirmed that full decellularization was achieved in all tissues (Fig. 1B). At the macroscopic level, decellularized organs retained their overall shape, suggesting that the decellularization procedure did not markedly affect tissue topography or gross mechanical properties. To further assess structural preservation, we imaged decellularized organs by scanning electron microscopy (SEM), which revealed intact microanatomical features characteristic of each tissue type (Fig. 1C). For example, we could detect the Glisson's triad in the liver, bronchial structures in the lung, and intact blood vessels within the gut submucosa (Fig. 1D). Consistent with these observations, histological stainings for collagen and elastin performed on native and decellularized tissues confirmed that the protocol did not substantially alter the underlying tissue architecture (Suppl. Fig. 2A-B).

Finally, as SDS is toxic to cells, optimal removal of SDS was assessed with a colorimetric assay previously described [15] (Suppl. Fig. 2C). Of note, residual SDS below 0.005% is non-toxic to human cells [21], and the washing protocol used achieved this level in all organs after just two washes.

3.2. Mechanical tissue properties are maintained after decellularization

We next assessed whether decellularization altered the mechanical properties of the tissues. Elastic modulus measurements were performed using a Chiaro nanoindenter (Optics11 life) on both fresh and decellularized tissue slices of lungs, liver, and gut. In this system, an optical fiber detects the phase shift caused by cantilever deflection during indentation, allowing precise measurement of tissue mechanical properties.

Slices of 500 µm of thickness were immobilized on the bottom of a petri dish and indentation measurements were performed with a round tip as described in the materials and methods. Importantly, all the tissues were analyzed using the same settings, allowing for direct comparison of the results. Native tissues were imaged in the presence of protease inhibitors to reduce tissue damage, and propidium iodide was added to the imaging medium to visually confirm the absence of damaged cells from the measured areas. Measurements were performed over areas of 600 µm x 600 µm. No significant differences were found between native and decellularized tissue for any of the organs tested (Fig. 2).

These data confirm that our detergent-based decellularization procedure did not significantly alter tissue stiffness, making the decellularized scaffolds suitable for the study of mechanosensitive processes such as cell invasion.

3.3. Composition of decellularized organs

To evaluate the molecular integrity of the ECM, we performed label-free, quantitative proteomics on decellularized tissues (Suppl. Table 1). A core set of ECM proteins was consistently detected across all samples, along with distinct tissue-specific components. Notably, proteins known to be enriched in each organ, such as Collagen IV in liver, Thrombospondin and Elastin in lung, and Mucins in the gut [3], [22], were reliably identified by this approach (Fig. 3A, Suppl. Table 1). Gene Ontology (GO) analysis revealed ECM and ECM-associated functions among the top three molecular categories in all tissues examined (Fig. 3B, Suppl. Table 1). In addition, the UNIPROT keywords "basement membrane" and "collagen" were enriched with high confidence across all decellularized scaffolds (Suppl. Table 1).

Together, these findings demonstrate that our protocol preserves not only tissue architecture and mechanical properties but also the biochemical composition of each organ's ECM.

3.4. In Vitro invasion assays confirm intrinsic cell line differences

To benchmark our *ex vivo* model, we first tested the invasive potential of MDA-MB-231 and MCF7 breast cancer cells in conventional 2D and 3D *in vitro* assays. In 2D wound healing, both cell lines displayed the ability to move, but MDA-MB-231 were significantly faster than MCF7 both on uncoated substrates and substrates coated with collagen I (Fig. 4A-B). Invasion through artificial ECM was assessed using fluorescence blocking transwells. In agreement with the literature, MDA-MB-231 were strongly invasive, contrary to MCF7 (Fig. 4C). The ability to

degrade the matrix is necessary to invade through dense matrices. We therefore measured the degradation capacity of the two cell lines by plating them on coverslips coated with fluorescent collagen and allowing them to degrade the substrate overnight. Again, in agreement with the literature, MCF7 did not display any degradative capacity, while MDA-MB-231 degraded the fluorescent collagen in regions resembling focal adhesions (Fig. 4D-E).

These results confirm the differential invasive behavior of the two cell lines and provide a reference for their behavior in the organ-derived scaffolds.

3.5. Breast cancer cell invasion is organ-specific in decellularized scaffolds

For *ex vivo* invasion assays, organ slices were inserted into ibidi channels. Thanks to the precision of the cutting process [23], we were able to place the slices into the channels without the need for glue or excessive compression. GFP-expressing cancer cells were added and allowed to invade for 7 days (Suppl. Fig. 3A). Although some autofluorescence is always present when imaging tissues, the decellularization protocol renders tissues virtually transparent and makes imaging possible using a standard fluorescence microscope without requiring fixation or staining (Suppl. Movie 1). Visual inspection of all organs, along with live imaging of lung slices, confirmed that both cell lines adhered to the scaffold surfaces from the start of the assay, ensuring that differences in invasion were not due to attachment efficiency (Suppl. Movie 1).

After 7 days of invasion, cells and tissues were fixed and subjected to immunostaining. Consistent with epifluorescence imaging, confocal microscopy confirmed that MCF7 cells failed to infiltrate any of the three tissues tested (Fig. 5A-C). In contrast, MDA-MB-231 cells invaded the lungs and liver, whereas their penetration in the intestine was negligible (Fig. 5A-C; Suppl. Movies 2 and 3). Both single cell invasion and collective invasion were detected. Notably, localized collagen degradation, indicative of metalloprotease-dependent mesenchymal invasion, was readily detected in association with single invading cells (Fig. 5A-B).

Quantitative analysis of invasion depth revealed a trend toward deeper penetration in liver compared to lungs, although this difference did not reach statistical significance (Fig. 5D and Suppl. Fig. 3B). Penetration distances could be converted into an average effective invasion speed, yielding values of approximately 27–30 µm/day in liver and 12–27 µm/day in lung (Fig. 5E).

Overall, these findings demonstrate that decellularized organ scaffolds quantitatively recapitulate clinically observed metastatic tropism, with breast cancer cells preferentially invading lung and liver over intestine tissue.

4. Discussion and conclusion

In summary, we established an *ex vivo* invasion platform based on decellularized organ scaffolds that enables quantitative investigation of cancer cell invasion within physiologically relevant extracellular matrices. This platform is designed to bridge the gap between reductionist *in vitro* assays and the structural and biochemical complexity of *in vivo* metastasis models, providing a practical and biologically meaningful tool for preclinical research.

Because ECM properties vary by organ and strongly influence metastatic behavior, incorporating native ECM is essential for modeling organ-specific invasion [14]. Mechanical, structural, and biochemical analyses confirmed that the decellularization protocol preserved tissue mechanics and retained key organ-specific ECM components. Using this platform, we distinguished invasive from non-invasive cancer cells in agreement with established assays, while also observing organ-dependent differences in invasion that are not captured by conventional models. In particular, MDA-MB-231 cells invaded lung and liver scaffolds but showed minimal penetration into intestine ECM, consistent with reported patterns of metastatic organotropism in triple-negative breast cancer [24]. Invasion rates measured in the scaffolds were within the range reported in vivo by intravital microscopy [25], supporting the relevance of the system for quantitative invasion studies.

The system is fully compatible with live and fixed imaging, immunostaining, and three-dimensional reconstruction, enabling detailed spatial and temporal analysis of invasion modes. The preservation of tissue orientation throughout the assay further facilitates correlative live-to-fixed studies. The workflow is simple to implement, cost-effective, and scalable, making it well suited for comparative studies and medium-throughput screening applications.

While this system does not recapitulate all aspects of the *in vivo* metastatic niche, such as immune surveillance, active stromal remodeling, or vascular perfusion, it isolates the contribution of the native ECM, allowing its role in organ-specific invasion to be interrogated with unprecedented clarity. As such, it complements *in vivo* models, and provides a valuable refinement strategy in line with ongoing efforts to reduce animal experimentation.

Although demonstrated here using mouse tissues, the approach is readily adaptable to other sources, including human biopsies. The use of thin slices allows multiplex analyses from limited material. By preserving native ECM structure and mechanics while remaining simple to implement, this platform represents a scalable complement and, in some contexts, a potential alternative to traditional *in vivo* metastasis models.

CRediT authorship contribution statement

FT: conceptualization, data curation, investigation, methodology, validation, visualization, review and editing. **PFG**: investigation, methodology, visualization, review and editing. **MR**: methodology, investigation. **BS**: methodology, investigation. **KF:** methodology, investigation. **PPG**: data curation, formal analysis. **FE**: resources. **AV**: resources. **SO**: investigation, resources. **SR**: investigation, resources. **JH**: investigation, methodology, resources. **TH**: methodology, resources. **FB**: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, supervision, visualization, writing original draft

Conflict of interest

All authors declare that they have no competing interests.

Acknowledgements

We thank the Biooptics facility and the Proteomics facility of Medical University Innsbruck for technical support throughout the research project. We thank Dr. Anna Seybold from the electron microscopy facility of the University Innsbruck for technical support. Open Access Funding provided by the Medical University of Innsbruck. This work was supported by the Austrian Science Fund (FWF Grant-DOI: 10.55776/PAT4730323) and by a MUI installation grant given to FB.

References

[1]   A. L. Berrier and K. M. Yamada, "Cell-matrix adhesion," *J Cell Physiol*, vol. 213, no. 3, pp. 565–573, Dec. 2007, doi: 10.1002/JCP.21237;CTYPE:STRING:JOURNAL.

[2]   R. O. Hynes, "The extracellular matrix: not just pretty fibrils.," *Science*, vol. 326, no. 5957, pp. 1216–9, Nov. 2009, doi: 10.1126/science.1176009.

[3]   A. Naba, K. R. Clauser, S. Hoersch, H. Liu, S. A. Carr, and R. O. Hynes, "The Matrisome: In Silico Definition and In Vivo Characterization by Proteomics of Normal and Tumor Extracellular Matrices," *Molecular & Cellular Proteomics*, vol. 11, no. 4, p. M111.014647, Apr. 2012, doi: 10.1074/MCP.M111.014647.

[4]   K. Tarnutzer, D. Siva Sankar, J. Dengjel, and C. Y. Ewald, "Collagen constitutes about 12% in females and 17% in males of the total protein in mice," *Sci Rep*, vol. 13, no. 1, pp. 1–12, Dec. 2023, doi: 10.1038/S41598-023-31566-Z;SUBJMETA=45,612,631,750,79,80;KWRD=EXTRACELLULAR+MATRIX,PROTEINS.

[5]   J. K. Mouw, G. Ou, and V. M. Weaver, "Extracellular matrix assembly: a multiscale deconstruction," *Nature Reviews Molecular Cell Biology 2014 15:12*, vol. 15, no. 12, pp. 771–785, Nov. 2014, doi: 10.1038/nrm3902.

[6]   K. M. Yamada and M. Sixt, "Mechanisms of 3D cell migration," *Nature Reviews Molecular Cell Biology 2019 20:12*, vol. 20, no. 12, pp. 738–752, Oct. 2019, doi: 10.1038/s41580-019-0172-9.

[7]   A. P. Le, J. F. Rupprecht, R. M. Mège, Y. Toyama, C. T. Lim, and B. Ladoux, "Adhesion-mediated heterogeneous actin organization governs apoptotic cell extrusion," *Nature Communications 2021 12:1*, vol. 12, no. 1, pp. 1–18, Jan. 2021, doi: 10.1038/s41467-020-20563-9.

[8]   S. Melo *et al.*, "The ECM and tissue architecture are major determinants of early invasion mediated by E-cadherin dysfunction," *Communications Biology 2023 6:1*, vol. 6, no. 1, pp. 1–11, Nov. 2023, doi: 10.1038/s42003-023-05482-x.

[9]   M. J. Jones and M. C. Jones, "Cell cycle control by cell-matrix interactions," *Curr Opin Cell Biol*, vol. 86, p. 102288, Feb. 2024, doi: 10.1016/J.CEB.2023.102288.

[10] A. Naba, "Mechanisms of assembly and remodelling of the extracellular matrix," *Nature Reviews Molecular Cell Biology 2024 25:11*, vol. 25, no. 11, pp. 865–885, Sep. 2024, doi: 10.1038/s41580-024-00767-3.

[11] M. R. Chastney, J. R. W. Conway, and J. Ivaska, "Integrin adhesion complexes," *Current Biology*, vol. 31, no. 10, pp. R536–R542, May 2021, doi: 10.1016/J.CUB.2021.01.038.

[12] A. Hoshino *et al.*, "Tumour exosome integrins determine organotropic metastasis," *Nature*, vol. 527, no. 7578, pp. 329–335, Nov. 2015, doi: 10.1038/NATURE15756;SUBJMETA=304,322,327,631,67,80;KWRD=CANCER+MICROENV IRONMENT,MECHANISMS+OF+DISEASE,METASTASIS.

[13] H. Peinado *et al.*, "Pre-metastatic niches: organ-specific homes for metastases," *Nature Reviews Cancer 2017 17:5*, vol. 17, no. 5, pp. 302–317, Mar. 2017, doi: 10.1038/nrc.2017.6.

[14] Y. Gao, I. Bado, H. Wang, W. Zhang, J. M. Rosen, and X. H. F. Zhang, "Metastasis Organotropism: Redefining the Congenial Soil," *Dev Cell*, vol. 49, no. 3, pp. 375–391, May 2019, doi: 10.1016/J.DEVCEL.2019.04.012.

[15] W. Godefroy *et al.*, "Development and qualification of clinical grade decellularized and cryopreserved human esophagi," *Sci Rep*, vol. 13, no. 1, pp. 1–17, Dec. 2023, doi: 10.1038/S41598-023-45610-5;SUBJMETA=54,61,631,994;KWRD=BIOMATERIALS,TISSUES.

[16] J. M. Wallis *et al.*, "Comparative Assessment of Detergent-Based Protocols for Mouse Lung De-Cellularization and Re-Cellularization," *https://home.liebertpub.com/tec*, vol. 18, no. 6, pp. 420–432, Jan. 2012, doi: 10.1089/TEN.TEC.2011.0567.

[17] K. Faserl, A. J. Chetwynd, I. Lynch, J. A. Thorn, and H. H. Lindner, "Corona Isolation Method Matters: Capillary Electrophoresis Mass Spectrometry Based Comparison of Protein Corona Compositions Following On-Particle versus In-Solution or In-Gel Digestion," *Nanomaterials 2019, Vol. 9, Page 898*, vol. 9, no. 6, p. 898, Jun. 2019, doi: 10.3390/NANO9060898.

[18] P. M. Crapo, T. W. Gilbert, and S. F. Badylak, "An overview of tissue and whole organ decellularization processes," *Biomaterials*, vol. 32, no. 12, pp. 3233–3243, Apr. 2011, doi: 10.1016/J.BIOMATERIALS.2011.01.057.

[19] A. L. Wishart *et al.*, "Decellularized extracellular matrix scaffolds identify full-length collagen VI as a driver of breast cancer cell invasion in obesity and metastasis," *Sci Adv*, vol. 6, no. 43, Oct. 2020, doi: 10.1126/SCIADV.ABC3175/SUPPL_FILE/ABC3175_SM.PDF.

[20] M. J. Oudin and V. M. Weaver, "Physical and Chemical Gradients in the Tumor Microenvironment Regulate Tumor Cell Invasion, Migration, and Metastasis.," *Cold Spring Harb Symp Quant Biol*, vol. 81, pp. 189–205, Jan. 2016, doi: 10.1101/sqb.2016.81.030817.

[21] S. Cebotari *et al.*, "Detergent decellularization of heart valves for tissue engineering: Toxicological effects of residual detergents on human endothelial cells," *Artif Organs*, vol. 34, no. 3, pp. 206–210, Mar. 2010, doi: 10.1111/J.1525-1594.2009.00796.X;WGROUP:STRING:PUBLICATION.

[22] A. M. Roy, R. Iyer, and S. Chakraborty, "The extracellular matrix in hepatocellular carcinoma: Mechanisms and therapeutic vulnerability," *Cell Rep Med*, vol. 4, no. 9, p. 101170, Sep. 2023, doi: 10.1016/J.XCRM.2023.101170.

[23] M. Zimmermann *et al.*, "Improved reproducibility in preparing precision-cut liver tissue slices," *Cytotechnology 2009 61:3*, vol. 61, no. 3, pp. 145–152, Dec. 2009, doi: 10.1007/S10616-009-9246-4.

[24]   C. Bartmann *et al.*, "Pattern of metastatic spread and subcategories of breast cancer," *Arch Gynecol Obstet*, vol. 295, no. 1, pp. 211–223, Jan. 2017, doi: 10.1007/S00404-016-4225-4/TABLES/3.

[25]   B. Weigelin, G.-J. Bakker, and P. Friedl, "Intravital third harmonic generation microscopy of collective melanoma cell invasion," *Intravital*, vol. 1, no. 1, pp. 32–43, Jul. 2012, doi: 10.4161/INTV.21223.

Figure Legends

Figure 1 – Preparation and characterization of decellularized organs

(A) Representative images of native and decellularized organs. **(B)** Hematoxylin and eosin staining of native and decellularized organs **(C)** Representative SEM images of decellularized organs. **(D)** Close-up of organ-specific structural features (from left to right: Glisson's triad for the liver, bronchi in the lung, and blood vessels included in the intestine submucosa).

Figure 2 – Mechanical characterization of sliced organs

Measures of the Effective Young Modulus of slices of native or decellularized organs. 36 measure per sample were performed and violin plots were built using the individual measures. Empty circles represent the average Effective Young Modulus of every organ measured. At least 3 slices, each corresponding to a different organ, were measured in each category. Results were analyzed by ANOVA Tukey's multiple comparisons test (p values are indicated for non significant differences; **** $p < 0.0001$).

Figure 3 – Proteomics analysis on decellularized organs

(A) Decellularized organs were subjected to mass spectrometry. The number of organ-specific proteins, as well as the number of proteins shared between different organs is displayed. **(B)** Gene Ontology enrichment of the proteins identified in each decellularized organs with the (top 5) significant molecular functions listed.

Figure 4 – In vitro characterization of breast cancer cell lines

(A) Cells were seeded to confluence in ibidi wound healing inserts. Inserts were removed and cells were imaged every 2h for a total of 8h. 5-6 wounds per cell type from 2 independent experiments were quantified and results were analyzed by by Two-way ANOVA with multiple comparisons test (* $p < 0.05$; *** $p < 0.001$). **(B)** Representative images of wounds at the beginning (o hours) and at the end (8 hours) of the videos. **(C)** Cells were allowed to invade through GelTrex for 48h. Then, invading cells were labeled with DAPI, imaged and counted. Three independent experiments, each containing 2 technical replicates per condition were performed and analyzed by Student's T-test (* $p < 0.05$). **(D)** Cells were seeded on fluorescent gelatin-coated coverslips and allowed to adhere for 24h. Then, cells were fixed and stained for actin (phalloidin) and nuclei (DAPI). The digested area underneath the cells (regions devoid of

gelatin) was measured as compared to the total cell area (segmented based on the actin cytoskeleton) to obtain percentages of digested areas. 167 and 80 cells respectively for MCF7 and MDA-MB-231 from 3 independent experiments were analyzed by unpaired T-test (** $p <$ 0.01).

Figure 5 – Ex vivo invasion of breast cancer cells through decellularized organs

(A) Representative mid-section of organs slices after 7 days of incubation with cancer cells (red). Interference reflection microscopy was used to visualize fibrillar ECM (cyan) and cleaved collagen I was immunostained (yellow). **(B)** Maximum projections of 100 µm thick stacks representative of the behavior of each distinct cell line in every organ. **(C)** 3D rendering of cells in different organs. **(D)** The distance of the invasive front of MDA-MB-231 cells was quantified in the indicated organs across three independent experiments. Violin plots show invasion distances measured at multiple points along individual invasive foci for each organ. Empty circles indicate the mean invasion distance per organ. **(E)** A theoretical invasion speed of MDA-MB-231 in the indicated organs was calculated by dividing the invaded distance as in (D) by the number of invasion days.

Supplementary Material

Supplementary Movie 1

MCF7 (left) or MDA-MB-231 (right) cells were allowed to adhere to lung slices for 24h, then imaged on an epifluorescence inverted microscope at the frequency of 1 image every 10 min for 15h. Cells are GFP-positive, and decellularized organs are visualized by phase contrast. Scale bars: 50 µm.

Supplementary Movie 2

3D rendering of MDA-MB-231 cells adhering on the external surface of the intestine without invading. Red: cells. Cyan: Fibrillar ECM imaged by interference reflection microscopy.

Supplementary Movie 3

3D rendering of MDA-MB-231 cells invading into the liver. Red: cells. Cyan: Fibrillar ECM imaged by interference reflection microscopy.

Supplementary Table 1

Proteomics analysis and permalinks to Gene Ontology Enrichment Analyses.

Suppl. Fig. 1 – Characterization of the decellularization protocol

(A) Representative images of native or decellularized organs stained with DAPI. **(B)** DAPI fluorescence of images as in (A) was quantified over a surface of at least 4 mm^2 for each organ. Results were analyzed by non-parametric LSD Fisher's test (* $p < 0.05$; **** $p < 0.0001$).

Suppl. Fig. 2 – Histology of native and decellularized organs

(A) Connective tissue in the different organs was visualized by Elastic van Gieson staining. **(B)** Collagen was visualized in different organs by CAB staining. **(C)** Residual SDS in the solution where decellularized organs were washed and stored was quantified in three to five independent experiments. The blue dashed line corresponds to the maximum tolerated levels of residual SDS for recellularization purposes.

Suppl. Fig. 3 – Schematic of the ex-vivo system's setup

(A) Schematic of the protocol to place decellularized organ slices in imaging microchannels divided into the two phases of preparation and invasion experiment. **(B)** Cancer cell invasion was quantified as illustrated in the schematic: First, a cross-sectional image of each tissue slice was captured to visualize invasive foci. For each invaded region, a higher-resolution volumetric image was then acquired. Within these images, the distance between invading cancer cells (red) and the original organ surface (dashed black line) was measured by drawing perpendicular lines (yellow). The mean of these measurements was used to define the penetration depth of cancer cells within each invaded region across all organ scaffolds.

Figure 1

A Liver Lung Intestine

1 cm

B Native Decellularized

200 µm

C

200 µm

D

50 µm

Figure 2

Figure 3

A

B

Figure 4

A

B

MCF7

MDA MB 231

0 hours   8 hours

100 µm

C

MCF7 collagen
MCF7 no collagen
MDA-MB-231 collagen
MDA-MB-231 no collagen

D

MCF7

MDA-MB-231

Actin Gelatin DAPI

10 µm

E

Figure 5

A

Liver — 500 µm

Lung — 500 µm

Intestine — 500 µm

Cells
ECM
Cleaved collagen

B

MDA-MB-231 — 50 µm

MCF7 — 50 µm

Cells ECM Cleaved collagen

C

Liver - MDA-MB-231 — 30 µm

Liver - MCF7 — 40 µm

Lung - MDA-MB-231 — 30 µm

Intestine - MDA-MB-231 — 50 µm

Cells ECM

D Invasion

Dist. from tissue surface (µm)

Liver   Lung

E Theoretical Speed

Speed (µm/day)

Liver   Lung