Seabird diet and transmission routes mediate the increase in pathogen prevalence with human disturbance

Shabnam Shadloo[1], Amy Wilson[2], Marie Auger-Méthé[1,3],

1 Institute for the Oceans and Fisheries, University of British Columbia, Vancouver, British Columbia, Canada

2 Department of Forest and Conservation Sciences, University of British Columbia

3 Department of Statistics, University of British Columbia

* s.shadloo@oceans.ubc.ca

1 Abstract

Human disturbances can alter patterns of wildlife infection, but their effects are likely to differ between

terrestrial and marine environments. Seabirds range across land and sea and are therefore exposed to

diverse human pressures such as urbanization, agriculture, and landfills, as well as fisheries, shipping, and

marine pollution. However, most literature on wildlife infection has focused on terrestrial hosts or single

type of disturbance (e.g., urbanization). Studying seabirds, which exploit diverse diets and habitats, offers

a unique opportunity to disentangle how disturbances at sea and on land, as well as diets, and transmission

routes, can change pathogen prevalence at a global scale. We conducted a global literature review and

extracted datasets on seabird infection status for a range of pathogens, seabirds diets (piscivore,

invertivore, generalist), and pathogen transmission routes (contact-based, vector-borne, intermediate host)

to link pathogen prevalence to human disturbances at sea and on land. Our results showed that the

association between human disturbances and pathogen prevalence was mediated by species' diets and

differed across transmission routes. Across gradients of human disturbance at sea, pathogen prevalence in

piscivorous seabirds increased from near zero to over 0.7, particularly for contact-based pathogens.

Land-based disturbance was also associated with higher pathogen prevalence in generalists across

transmission routes, whereas invertivores showed a negative association with terrestrial disturbance. These

results indicate that piscivores are most vulnerable in heavily disturbed marine environments, while

generalists face greater risk on highly disturbed lands, suggesting that effective disease mitigation will

require targeted management actions in both environments.

Keywords: Wildlife health, Anthropogenic pressure, Disease ecology, Marine environment, Terrestrial

environment

2   Introduction

Disease outbreaks in wildlife occur through intrinsic and extrinsic factors and can be accelerated by human activities [1, 2]. These outbreaks pose serious threats to species by leading to higher mortality, reduced reproductive success, population decline, local extinction, and biodiversity loss [3–6]. For example, viral and bacterial diseases, including avian cholera (*Pasteurella multocida*), have caused significant mortality in aquatic birds and can lead to rapid population decline and even local extinction [7–9]. Recently, highly pathogenic avian influenza panzootic decimated seabird populations in North and South America, Europe, and Antarctica [10, 11]. Over three billion people live in areas with frequent contact with wildlife, further increasing the risk of disease transmission [12]. Urbanization creates opportunities for interactions between wildlife, livestock, and humans, modifying pathogen transmission through host density, movement, and immunity, as well as abiotic factors such as pollution and climate change [13]. Seabirds are especially vulnerable wildlife species. They are effective reservoirs for pathogens as they live long, breed in dense colonies, and have diverse diets. They also travel long distances, sometimes across continents, crossing different gradients of human impact [4, 14–16]. Half of seabird species are listed as globally threatened or near threatened by the IUCN [17]. This situation may worsen due to human-related factors such as climate change, pollution, and the dilution effect due to loss of biodiversity [2, 18–20].

Anthropogenic factors on terrestrial environments, such as urban development, alter seabirds' environments and behaviour, influencing their interactions with pathogens [13, 21, 22]. Human activities concentrate food resources and pathogen hosts in new ways. For instance, open-air landfills and agricultural sites provide abundant, predictable food that attracts large numbers of opportunistic seabirds. However, these dense foraging aggregations become hotspots for pathogen transmission, since many diseases often rely on the spatial overlap of hosts and vectors [2, 13, 23–27]. For example, birds in urbanized landscapes experience higher exposure to mosquito-borne viruses such as West Nile virus [28]. We predict that higher human pressure on terrestrial environments could lead to increased prevalence of pathogens, due to more overlap with intermediate hosts and vectors, and aggregation of species in anthropogenic food sources and contaminated habitats.

Human pressures on marine environments can have a similar impact on seabirds. Marine pollution can

introduce new pathogens to seabirds. For example, seabirds foraging near sewage outfalls or wastewater treatment plants often ingest pathogens and antimicrobial-resistant microbes originating from human waste [29]. Another example is the spread of *Toxoplasma gondii* to marine environments through urban runoffs, leading to infection in seabirds [30]. Fishing operations produce abundant waste that can be opportunistically exploited by seabirds [31, 32]. Scavenging seabirds congregate in large numbers around fishing vessels or fish processing outfalls and show a strong tendency to repeatedly return to the same feeding locations [31]. This aggregation around fishing vessels and processing facilities can increase interspecies contact and pathogen exchange. Bycatch can also indirectly increase susceptibility to novel pathogens, especially when it leads to large population declines associated with reduced genetic diversity and resilience [33]. Many pelagic and polar seabirds may rarely visit cities, but can be affected by marine-based impacts such as bioaccumulation of contaminants and climate-driven shifts in marine pathogens. For example, penguins are exposed to persistent organic pollutants, which can impair their immune function [34]. Many offshore seabirds, such as petrels and shearwaters, chronically ingest microplastics, which can increase the prevalence of pathogenic bacteria [35, 36]. Climate change is also shifting the distribution of some pathogens towards higher altitudes, mainly threatening naive species [37–39]. Accordingly, we predict that pathogens would occur more in seabirds in waters with higher cumulative human disturbance.

Seabirds' foraging ecology can also shape how human activities influence the prevalence of pathogens that seabirds encounter. Their foraging ecology can determine movement and habitat use, drawing some species into cities where they access abundant food, while leading others to avoid urban environments. Generalist seabirds are more likely to be exposed to a broader array of pathogens, particularly in anthropogenic landscapes [40]. For example, colonies of European herring gulls (*Larus argentatus*) foraging across varied habitats exhibited greater microbial pathogen diversity than those with more specialized diets [41]. However, dietary specialization does not eliminate the risk of pathogens. Foodborne pathogens such as Vibrio spp. can spread through prey species, such as shellfish, fish, zooplankton, and crustaceans [41, 42]. We predict that human-related factors would interact with seabird diet to alter the pathogen prevalence.

The transmission routes used by pathogens can determine which factors accelerate their transmission

and how human activities influence their prevalence in seabirds. Transmission occurs through vertical

(passed directly from parent to offspring) and/or horizontal routes [43]. For pathogens to transmit through

horizontal routes, suitable ecological and environmental conditions are required, such as direct contact with

infected individuals, contaminated environments, or the presence of vectors or intermediate hosts [44, 45].

Because human disturbances can modify these conditions, for example, by concentrating animals at

landfills, altering prey communities, increasing pollution, and altering food and water-borne pathogens

dynamics, the potential effects of humans on pathogen prevalence depend on the transmission route.

Numerous studies have examined pathogen prevalence in wildlife species. Much of this research has

focused on the relationship between pathogens and wild mammals [46, 47], species' habitat use [40], or has

concentrated on specific groups of pathogens [48–51]. Among studies on wild birds [52, 53], the impacts of

human disturbances were not assessed. Some studies have documented how urbanization affects pathogen

prevalence; however, they have primarily focused on terrestrial species and have rarely considered species'

diets [2, 54]. Predictions derived from such studies are unlikely to fully apply to seabirds, as they encounter

different opportunities and constraints for pathogen transmission at sea and on land due to their unique

ecology. Given the recent emergence of global wildlife pandemics, such as highly pathogenic avian

influenza, linking human impacts to pathogen prevalence in seabirds can provide valuable insights for their

conservation. We aim to fill the gap by addressing three questions: (1) whether the prevalence of

pathogens in seabirds increases with human impact on marine and terrestrial environments, (2) if this

association is influenced by an interaction with seabirds' diets, and (3) whether pathogen associations with

human disturbances vary by transmission route.

3 Materials and Methods

3.1 Literature review

98 We conducted a comprehensive literature search in Web of Science from 2000 to 2024 to identify English

99 publications on seabirds and pathogens. The search terms were (seabird* OR gull* OR penguin* OR

100 albatross* OR petrel* OR fulmar* OR shearwater* OR murre* OR tern* OR skua* OR puffin* OR

101 frigatebird* OR auklet* OR guillemot* OR kittiwake* OR cormorant* OR shag* OR booby* OR

tropicbird* OR auk* OR pelican* OR gannet*) AND (pathogen* OR disease* OR infection* OR parasite* OR virus* OR bacteria OR helminth* OR protozoa*). We restricted the search to keywords appearing in titles and abstracts to enhance relevance. We then screened titles and abstracts and excluded the ones that were irrelevant. To minimize bias, we excluded experimental studies and samples from captive birds, rehabilitation centers, and carcasses, as well as studies where the sampling effort relied on clinical signs and diagnoses were based on symptoms rather than laboratory tests. Additionally, we removed research on pooled samples, eggs, and studies that collected data from fecal samples from nests or the ground, rather than from specified individual birds, because the number of individuals was unknown in such cases. We excluded studies focused on vectors, rather than pathogens, and removed studies on shorebirds based on two criteria: species belonging to the families Scolopacidae (e.g., sandpipers), Charadriidae (e.g., plovers), Haematopodidae (e.g., oystercatchers), and Jacanidae (jacanas), and species that never occupy marine neritic or oceanic habitats at any stage of their life cycle (breeding or non-breeding) according to IUCN Red List [55]. We also removed datasets used in more than one study. In the final step, we excluded samples for which we could not obtain covariate values (primarily studies located far inland and thus lacking marine environment information).

From each study, we collected data on sample size, number of positive cases, geographic location (x and y coordinates if mentioned or coordinates extracted from Google Earth based on location name), pathogen, and detection method. If any of this data was missing in a study, we excluded that study as well. We categorized the methods for pathogen detection into two groups: 1) isolation, including methods such as DNA extraction, blood smears, physical examination, antigen detection, microscopy, culturing bacteria, and histological analysis, and 2) exposure, which includes different methods of detecting antibodies, such as Enzyme-Linked Immunosorbent Assay (ELISA) and Microscopic Agglutination Test (MAT).

For each sample, we recorded the counts of infected and uninfected individuals, the sampling year(s), geographic location, pathogen, and host species. Many studies contributed multiple samples because they reported several pathogens, species, locations, or, in a few cases, distinct sampling periods. So the number of samples in our dataset exceeds the number of studies (Fig. S2 and S5). The literature search data are available in the Supplement Materials: Pathogen Prevalence in Seabirds Dataset.

3.2 Predictor variables

We used two spatial layers that quantify overall human disturbances that may affect pathogen prevalence: the Human Footprint Index for terrestrial environments (hereafter H.Land; [56]) and the Cumulative Human Impact on Marine Ecosystems dataset (hereafter H.Sea; [57]). The H.Land layer combines eight layers of human pressure, including urban development, population density, nighttime lights, croplands, pasturelands, roads, railways, and navigable waterways. For 81 study locations that fell outside the H.Land layer (e.g., some remote small islands), we manually assigned scores by examining the corresponding areas using satellite imagery, following the methodology outlined in the original dataset [56, 58]. The H.Sea layer aggregates human impacts on global oceans from 19 anthropogenic stressors across four categories: land-based pollution (such as nutrient runoff); ocean-based activities (such as fishing); climate change; and invasive species. We created a buffer of 50 km around each data point and extracted the mean value for each covariate (H.Land and H.Sea) in each buffer. To simplify calculations and facilitate more explicit comparisons among predictors, we standardized these predictors using z-transformation [59]. According to the Pearson correlation coefficient, H.Land and H.Sea were not strongly colinear in our dataset (r = 0.40).

To understand the association between diet and its interaction with human impact, we extracted diet data from the EltonTraits 1.0 dataset [60]. We reclassified these diets as predominantly piscivore, predominantly invertivore, or generalist (Table 1). Species falling in the third category did not have a dominant diet except for two species that had unique diets compared to others: dolphin gull *Leucophaeus scoresbii* (dominantly scavenger) and brown skua *Stercorarius antarcticus* (mainly feeds on mammals and birds).

Table 1. Definition of each diet category extracted from EltonTraits 1.0 dataset [60].

Diet	Definition
Piscivore	$\geq 50\%$ of the diet is fish.
Invertivores	$\geq 50\%$ of the diet is invertebrates: aquatic invertebrates, shrimp, krill, squid, crustaceans, molluscs, cephalopods, polychaetes, gastropods, orthoptera, terrestrial invertebrates, ground insects, insect larvae, worms, orthopterans, and flying insects.
Generalists	None of the species-specific diets reach 50%, except for dolphin gull and brown skua.

3.3 Extracting transmission route

To understand how the relationship between human disturbance and pathogen prevalence may be altered by its transmission route, we obtained data on pathogen transmission routes from vatious sources as we could not find one comprehensive database for pathogens transmission routes related to birds. We used the Global Mammal Parasite Database (GMPD), Centers for Disease Control and Prevention (CDC), and published articles related to pathogens not found in these two databases [61, 62] (see S6). We classified these mechanisms into three categories: close and non-close contact (hereafter, contact-based), vector-borne, and intermediate host [44, 45]. Close-contact pathogens spread through direct interactions (e.g., touching and biting) between individuals, while non-close contact pathogens spread indirectly through contaminated objects (e.g., surfaces) or environments (e.g., water, food sources, or soil) [62]. In this study, we combined these two groups because many viruses and bacteria in our dataset can be transmitted via both routes. Vector-borne pathogens rely on living carriers, typically arthropods such as mosquitoes, flies, and ticks [62]. The final category, intermediate host, contains pathogens that require a secondary host to complete part of their life cycle, such as growth or asexual reproduction, before passing to their definitive host. In our dataset, some pathogens belong to more than one category [63].

3.4 Statistical analysis

Our analyses focused on identifying the factors that explain pathogen prevalence, defined as the number of infected individuals in each sample, bounded by the number of individuals tested for infection. Since many samples did not contain infected individuals, we used generalized linear mixed model (GLMM) with a zero-inflated binomial distribution (ZIB) using the R package glmmTMB [64–66]. To account for the variability that our fixed effects cannot explain, we included two random effects. The first is the seabird species nested within genus, since the evolutionary history and genetic relatedness can influence a species' susceptibility or resistance to a disease [67]. For example, birds in the order Passeriformes are more susceptible to avian malaria compared to other bird groups [68]. Then, we incorporated a random effect for study ID to control for variation specific to each study.

We modeled pathogen prevalence (Y_i) as a function of H.Land, H.Sea, and diet, and allowed excessive zeros to depend on the method of pathogen detection as

$$
\begin{aligned}
&Y_i \sim \text{ZIB}(n_i, p_i, \pi_i), \\
&Y_i =
\begin{cases}
0 & \text{with probability } p_i + (1 - p_i)(1 - \pi_i)^{n_i}, \\
k & \text{with probability } (1 - p_i)\binom{n_i}{k}\pi_i^{n_i}(1 - \pi_i)^{n_i - k}, \quad k = 1, \ldots, n_i,
\end{cases}
\end{aligned}
\tag{1}
$$

where i is a sampling unit defined with unique pathogen, seabird species, study, and location, p_i is the probability of a zero because infection was not possible and/or not detectable (e.g., no exposure, unsuitable host, limits of the detection method; considered as the zero inflation), n_i is the number of individuals in sample i, and π_i is the probability of the individual being infected if infection is possible (i.e., probability of success) in a single trial [66].

The probability of individual infection is modelled as:

$$\text{logit}(\pi_i) = \beta_0 + \beta_1 \text{diet:piscivore}_i + \beta_2 \text{diet:generalist}_i + \beta_3 \text{H.Sea}_i + \beta_4 \text{H.Land}_i$$
$$+ \beta_5(\text{diet:piscivore}_i \times \text{H.Sea}_i) + \beta_6(\text{diet:piscivore}_i \times \text{H.Land}_i)$$
$$+ \beta_7(\text{diet:generalist}_i \times \text{H.Sea}_i) + \beta_8(\text{diet:generalist}_i \times \text{H.Land}_i)$$
$$+ v_{\text{study}[i]} + v_{\text{SWG}[i]}, \tag{2}$$

$$v_{\text{study}[i]} \sim \mathcal{N}(0, \sigma^2_{\text{study}}), \tag{3}$$
$$v_{\text{SWG}[i]} \sim \mathcal{N}(0, \sigma^2_{\text{SWG}}), \tag{4}$$

where $\beta_{predictor}$ is the coefficient for each predictor, $v_{\text{study}[i]}$ is random effect of each study for sample i, and $v_{\text{SBSwG}[i]}$ is random effect of seabird species within genus for sample i. While H.Sea_i and H.Land_i are continuous predictors, diet:piscivore_i and diet:generalist_i are dummy variables that identify whether the species in sample unit i fall in each of these diets. The probability of a structural zero that is not defined in the binomial distribution is modelled as

$$\text{logit}(p_i) = \alpha_0 + \alpha_1 \text{method}_i, \tag{5}$$

where p_i is the probability that sample i is zero, α_0 is the intercept, and α_1 is the coefficient for the method of pathogen detection.

We conducted two sets of GLMMs to investigate differences between pathogen transmission routes. The first set included all available data, regardless of the pathogen transmission route (hereafter, the general model set). The second set comprised separate models focusing on each of the three transmission route categories (hereafter, contact-based, vector-borne, and intermediate host).

For model selection, we used the Bayesian Information Criterion (BIC) to identify the best-fitting models [69]. To perform residual diagnostics, we applied simulation-based methods available in the

DHARMa package [70]. When the BIC-selected model showed any association with H.Sea or H.Land, we used parametric bootstrap likelihood-ratio tests to assess whether the model including H.Sea and H.Land significantly improved model fit relative to a nested model without these two covariates (for more detail, see section Parametric bootstrap likelihood ratio).

4 Results

Our literature-review dataset spans all oceans and includes samples from 5 orders, 35 seabird genera, and 65 speccies (Fig 1 and S5). The first three selected models by BIC (i.e., general, contact-based, and vector-borne models) retained both H.Sea and H.Land (Tables S1, S2, and S3), and parametric bootstrap likelihood-ratio tests showed adding human-related predictor can significantly improve the fit for the three models (general: Δ deviance = 78.2.9, p_{value} = 0.001; contact-based: Δ deviance = 101.2, p_{value} = 0.001; vector-borne: Δ deviance = 80.8, p_{value} = 0.001). The intermediate-host model showed no clear association with any predictors. DHARMa's outlier test detected no extreme residuals, and the residual diagnostics indicated small but significant deviations for general, contact-based, and vector-borne models, and a poor fit for the intermediate-host model (Fig S3).

Fig 1. Global distribution of the samples and seabird orders. Each dot marks one independent dataset sample from published studies. The colour of each dot shows the pathogen prevalence reported in samples across a gradient of human impact on marine and terrestrial environments. The pie chart shows seabirds orders in the study. Our dataset contained 5 orders of seabirds with the most frequent being Charadriiformes.

₂₀₉   Associations between pathogen prevalence and human impact varied with diet and transmission route

₂₁₀   (Fig 2 panel A; Table 2). The best model from the general set indicates that the prevalence in piscivores

₂₁₁   increased with both H.Sea and H.Land, and that the prevalence in generalists increased with H.Land.

₂₁₂   Invertivores, in contrast, showed a very small decline with increasing H.Land.

A) General model

B) Contact-based model

C) Vector-borne model

Human disturbance at sea · Human disturbance on land

Fig 2. Pathogen prevalence as a function of H.Sea and H.Land and their interaction with 95% confidence intervals. The figures are A) general, B) contact-based, and C) vector-borne models. The main effects of H.Sea and H.Land give the slope for invertivores, and the interaction terms for generalists and piscivores show how their slopes differ from those of invertivores. Pink rectangles mark parameters whose 95% CIs do not include zero.

Table 2. Parameters in three BIC-selected models (General, Contact-based, Vector-borne). Estimates are on the logit scale. Here, FE and RE are fixed effects and random effects, respectively. The p-values in the first column are the result of the parametric bootstrap likelihood ratio test. The intercept corresponds to the reference diet (invertivore).

Model (p-value)	Term	Variable	estimate	Standard error	p-value
General (0.001)	FE	intercept (invertivore)	$\beta_0 = -4.75$	0.84	< 0.01
	FE	diet:piscivore	$\beta_1 = 3.2$	0.89	< 0.01
	FE	diet:generalist	$\beta_2 = 2.21$	0.88	0.01
	FE	H.Sea (invertivore)	$\beta_3 = -0.23$	0.24	0.34
	FE	H.Land (invertivore)	$\beta_4 = -1.93$	0.49	< 0.01
	FE	diet:piscivore × H.Sea	$\beta_5 = 1.02$	0.26	< 0.01
	FE	diet:piscivore × H.Land	$\beta_6 = 2.26$	0.50	< 0.01
	FE	diet:generalist × H.Sea	$\beta_7 = 0.26$	0.31	0.39
	FE	diet:generalist × H.Land	$\beta_8 = 2.59$	0.57	< 0.01
	RE	Species within genus	$v_{\text{SWG}} = 1.68$		
	RE	Genus	$v_{genus} = 1.92$		
	RE	Study	$v_{study} = 7.48$		
Contact-based (0.001)	FE	intercept (invertivore)	$\beta_0 = -2.55$	0.85	< 0.01
	FE	diet:piscivore	$\beta_2 = 1.39$	0.99	0.16
	FE	diet:generalist	$\beta_1 = 0.22$	0.92	0.81
	FE	H.Sea (invertivore)	$\beta_3 = -0.15$	0.24	0.51
	FE	H.Land (invertivore)	$\beta_4 = -2.07$	0.49	< 0.01
	FE	diet:piscivore × H.Sea	$\beta_5 = 1.04$	0.29	< 0.01
	FE	diet:piscivore × H.Land	$\beta_6 = 1.88$	0.50	< 0.01
	FE	diet:generalist × H.Sea	$\beta_7 = -0.54$	0.80	0.50
	FE	diet:generalist × H.Land	$\beta_8 = 2.68$	0.77	< 0.01
	RE	Species within genus	$v_{\text{SWG}} = 0.97$		
	RE	Genus	$v_{genus} = 1.67$		
	RE	Study	$v_{study} = 5.37$		
Vector-borne (0.001)	FE	intercept	$\beta_3 = -5.31$	1.51	< 0.01
	FE	H.Sea	$\beta_3 = 0.51$	0.12	< 0.01
	FE	H.Land	$\beta_4 = 0.82$	0.12	< 0.01
	RE	Species within genus	$v_{\text{SWG}} = 0.62$	0.38	
	RE	Genus	$v_{genus} = 3.38$		
	RE	Study	$v_{study} = 2.44$		

213   Best models from sets focused on specific transmission routes differed from the general results,

214   indicating that transmission routes may alter the effect of human disturbances on pathogen prevalence (2

panels B nad C). As for the contact-based model, prevalence in piscivores increased with H.Sea, prevalence in generalists increased with H.Land, and prevalence of inventories decreased with H.Land. In contrast to the general model, pathogen prevalence in piscivores decreased with H.Land. However, the difference between the H.Land effect on piscivores and the baseline diet (invertivores) was positive. The prevalence of vector-borne pathogens was generally low, and showed small positive associations with both H.Sea and H.Land, regardless of diets (Fig 2). Other coefficients were not significant (their 95% confidence interval crossed zero; Fig S4). In the intermediate model, no covariate showed a significant association with prevalence. The intermediate dataset consisted of 85 samples from 4 studies. The small dataset may be why our models did not find any patterns.

To assess whether some seabird groups had higher overall pathogen prevalence than the average, we looked at the predicted odds ratio of pathogen prevalence for each genus. While most genera do not indicate clear deviation from the overall rate, two genera of seabirds (i.e., Pachyptila and Fregata) had odds ratios entirely above 1 in the selected vector-borne model (Fig S5).

5   Discussion

We found that pathogen prevalence in seabirds generally increased with human disturbance at sea (H.Sea) and on land (H.Land). Vector-borne pathogen prevalence rose with both H.Sea and H.Land across all diets, indicating a consistent positive association with disturbance. Diet and transmission route modified these patterns for other pathogen types. Piscivores showed particularly strong increases in prevalence with H.Sea, especially for contact-based pathogens (over 0.75 increase). Generalists had a higher prevalence with H.Land. In contrast, at high levels of H.Land, piscivores and invertivores showed lower prevalence of contact-based pathogens.

Human pressures can alter pathogen risks in wildlife through ecological and physiological mechanisms (e.g., [46,71]). Anthropogenic food leads to different disease outcomes in wildlife, often amplifying contact-based microparasites (e.g., many bacteria and viruses) by aggregating hosts at landfills, feeders, crops, or refuse [71]. Our results for generalist seabirds showed the same trend, with increases in pathogen prevalence with human disturbance on land. However, some studies showed contradictory results for

macroparasites with complex life cycles. They suggest food provided by humans can shift away from infected intermediate hosts [46, 71]. Some studies did not show a consistent positive or negative effect of urbanisation on bird and mammal parasites. They argued that differences in prevalence can be a result of parasite types, and their transmission modes, and can increase with pollutants, chronic stress, and weak immune defences [2, 53, 72–75]. Our findings are consistent with this idea, as we also observed that the relationship between human disturbance and pathogen prevalence depends on the pathogen type and its transmission mode.

The increase in pathogen prevalence in piscivores in high human disturbance at sea (H.Sea) likely reflects their foraging ecology. Piscivores often are exposed to pathogens through their prey, as many fish serve as tolerant reservoirs for parasites [76]. Fish host many parasites to reach infective stages, so predation on fish efficiently passes on these parasites up the food chain [77]. Many marine helminths are generalists that move easily across species through food chains. At the same time, their long-lived larval stages persist in intermediate or paratenic hosts and keep infection pressure high [77]. Piscivorous seabirds often bio-accumulate contaminants as well, which can disrupt their endocrine and immune systems [78–81], increasing the risk of infection. For instance, organochlorine loads correlate with immune suppression in Caspian terns (*Sterna caspia*) and toxic pollution and heavy metals such as polychlorinated biphenyls and mercury are correlated with higher prevalence of avian influenza in waterfowls [82, 83]. The reduction in pathogen prevalence with increased disturbance on land (H.Land) association for contact-based pathogens for piscivores may result from their different behaviour in urban areas. Many seabirds are offshore species that breed on remote islets in dense colonies. However, some species are increasingly breeding in human-dominated landscapes, forming more dispersed colonies (e.g., on rooftops; [84–87]). Moreover, unlike generalists, there is limited evidence of strictly piscivorous seabirds aggregating on landfills and poultry facilities, where contact-based pathogens such as avian influenza can be transmitted. For example, a large-scale survey in the Netherlands (2006–2009) compared over 6000 birds across an area with varying gradients of urbanization, and found that avian influenza prevalence was higher in highly urbanized sites in comparison to low-urbanized sites [88]. However, piscivorous seabirds (e.g., cormorants) inhabiting urban areas aggregate around anthropogenic fish-related resources, such as aquacultural sites and fishing piers (e.g., [89, 90]). Because our H.Land layer did not include aquaculture, it may underrepresent an important

pathway of pathogen exposure for these species.

Generalist seabirds in cities frequently use landfills, agricultural areas, and other anthropogenic food

sources [41, 91, 92]. Such locations can provide an opportunity for pathogen carriers to be in proximity with

susceptible individuals [93, 94]. For example, ibises in Florida and gulls in the Iberian coast showed a

higher prevalence of Salmonelosis and Campylobacter, respectively, in more developed areas originating

from urban parks and human wastes [95–97]. In addition, this behaviour can compromise their immune

systems through poor-quality food (high in fat and low in protein) [71, 98–100]. In an experimental study,

mice with an anthropogenic diet showed lower ability to kill *Salmonella paratyphi* compared to the ones

with a natural diet [101].

Invertivores also showed a lower prevalence in highly disturbed lands. Invertivore species in our study

are mostly offshore and have limited exposure to land-based pathogens, especially those transmitted

through contact [102]. And similar to piscivorous seabirds, they often do not aggregate in highly

contaminated sites on land. It is important to note that the accumulated H.Land we used did not account

for sewage impact, which may lead to an underrepresentation of human disturbances in such locations.

Invertivores foraging on estuarine mudflats contaminated by sewage can show a higher prevalence of

pathogens. For example, Olrog's gull feeding on benthic prey and crabs are exposed to pathogens

originating from human disturbances [103].

Our vector-borne category aggregates pathogens transmitted by a range of arthropods, such as ticks,

fleas, louse flies, and mosquitoes. We found a consistent, though slight, positive association between human

impact and diets. Climate change, warmer microclimates (e.g., in cities), and higher humidity can increase

the number and survival of vectors such as mosquitoes and ticks [2, 104–106]. For example, warmer winters

in the Arctic are associated with higher prevalence and intensity of *Ixodes uriae* ticks in seabird colonies [9].

Recent work suggests an expansion or a change in the distribution of *I. uriae* towards higher latitude due

to warmer winters in the Arctic [37]. Mosquitoes (Culex spp.) are also well adapted to man-made water

bodies (e.g., treatment plants and sewers; [28]). The abundance of mosquitoes can lead to increased

transmission of vector-borne pathogens, such as West Nile virus in avian species [28]. Also, simplifying bird

communities and favouring competent reservoir hosts may increase the prevalence of some vector-borne

viruses [37].

In our study, thin-billed prion (*Pachyptila belcheri*) and species within the genus Fregata (frigatebirds) showed higher odds of prevalence of vector-borne pathogens. A study showed that Haemoproteus *iwa* infections are particularly common in frigatebirds relative to other seabirds, and are transmitted by host-specialist hippoboscid flies *Olfersia spinifera* that are abundant on frigatebirds throughout their range in comparison to other species [107]. This result is in line with a review on seabirds [48]. They argued that since frigatebirds breed in warm, nearshore tropical environments where vectors such as louse flies (Hippoboscidae) and biting midges (Ceratopogonidae) are abundant, they can experience a high exposure to such parasites as well [48]. The same study also linked increased prevalence of another vector-borne pathogen, Plasmodium, to traits that prolong exposure at the nest, especially burrow nesting and longer fledging periods. *P. Belcheri* is a burrow-nester with a relatively long fledging period of about 50 days [108, 109].

Many seabird species shift their diet between breeding and non-breeding seasons. Quantifying diet across the full annual cycle is difficult, especially during the non-breeding period, and most traditional diet studies are limited to the chick-rearing season [110–112]. Therefore, our use of three coarse diet categories can introduce uncertainty. However, in the BIC model selection for general and contact-based pathogens, the best-supported models retained the interaction between diet and human disturbance, indicating that this covariate still captures significant variation in pathogen prevalence despite these limitations. Models without diet as a fixed effect had higher BIC values for these two models (Tables S1 to S4). The best vector-borne model result, however, did not include diet.

Our literature review identified important data gaps. We found a few studies on seabird pathogens with intermediate hosts that met our criteria. As a result, it was difficult to assess whether the lack of relationships between diet and human impacts was real or due to the small sample size. Pathogens requiring intermediate hosts can be influenced by human-modified environments. For instance, the prevalence of *T. gondii* in mammalian wildlife depends on their diet and habitat and increases with human population density [113, 114]. We also excluded many papers that lacked the necessary details about their data, such as fine-scale locations or counts of infected and uninfected individuals. We recommend future research to focus more on intermediate-host pathogens in seabirds and publish more detailed data, including diagnostic protocols, locations, sample sizes, and raw counts, to enable comparisons and

meta-analyses.

Overall, we saw that human impacts can increase the prevalence of pathogens in seabirds. One of the clearest patterns is related to piscivores and human disturance at sea. Our results showed that, through the low to high gradients of H.Sea, the prevalence of pathogens in seabirds can increase from nearly zero to around 70% and even higher in contact-based pathogens. Generalists exposed to high human disturbance on land have a higher rate of contact-transmitted pathogens. It is essential to reduce human pressure by lowering contaminants and enhancing waste treatment in the sea, as well as improving waste management on land, with an understanding of how seabirds use highly disturbed landscape on land and sea. For example, landfill management and restricting access to seairds has shown an increase in their stress levels in the short term and controversial results due to losing a reliable food (e.g., [115–117]). Identifying which combinations of diet and human disturbance drive the highest pathogen prevalence can be key to prioritizing monitoring and management efforts for seabird populations.

6 Acknowledgement

We are deeply grateful to the S$\underline{\text{T}}$ÁUT$\underline{\text{W}}$ (Tsawout) and W̱SIḴEM (Tseycum) for giving us access to X̱O, X̱DEȽ (Mandarte) Island. We thank Mark Hipfner, Peter Arcese, and Jade Banstraa for their assistance and collaboration throughout the study. This work was supported by Natural Sciences and Engineering Research Council of Canada (NSERC), the Canada Research Chairs program, BC Knowledge Development fund, Canada Foundation for Innovation's John R.Evans Leaders Fund, Cecil and Kathleen Morrow Scholarship, Environment and Climate Change Canada (ECCC), Fisheries and Oceans Canada (DFO), and the Four Year Fellowship (4YF).

References

1. Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife–threats to biodiversity and human health. Science. 2000;287(5452):443-9.

2. Bradley CA, Altizer S. Urbanization and the ecology of wildlife diseases. Trends in ecology & evolution. 2007;22(2):95-102.

3. Brearley G, Rhodes J, Bradley A, Baxter G, Seabrook L, Lunney D, et al. Wildlife disease prevalence in human-modified landscapes. Biological Reviews. 2013;88(2):427-42.

4. Khan JS, Provencher JF, Forbes MR, Mallory ML, Lebarbenchon C, McCoy KD. Parasites of seabirds: A survey of effects and ecological implications. In: Advances in Marine Biology. vol. 82;. ISSN: 21625875. doi:10.1016/bs.amb.2019.02.001.

5. Sanz-Aguilar A, Payo-Payo A, Rotger A, Yousfi L, Moutailler S, Beck C, et al. Infestation of small seabirds by *Ornithodoros maritimus* ticks: Effects on chick body condition, reproduction and associated infectious agents. Ticks and Tick-borne diseases. 2020;11(1):101281.

6. Camphuysen C, Gear S, Furness R. Avian influenza leads to mass mortality of adult Great Skuas in Foula in summer 2022. Scottish Birds. 2022;42:312-23.

7. Newman SH, Chmura A, Converse K, Kilpatrick AM, Patel N, Lammers E, et al. Aquatic bird disease and mortality as an indicator of changing ecosystem health. Marine Ecology Progress Series;352. doi:10.3354/meps07076.

8. Iverson SA, Gilchrist HG, Soos C, Buttler II, Harms NJ, Forbes MR. Injecting epidemiology into population viability analysis: avian cholera transmission dynamics at an arctic seabird colony. Journal of Animal Ecology. 2016;85(6):1481-90.

9. Descamps S, Jenouvrier S, Gilchrist HG, Forbes MR. Avian cholera, a threat to the viability of an Arctic seabird colony? PloS One. 2012;7(2):e29659.

10. McPhail GM, Collins SM, Burt TV, Careen NG, Doiron PB, Avery-Gomm S, et al. Geographic, ecological, and temporal patterns of seabird mortality during the 2022 HPAI H5N1 outbreak on the island of Newfoundland. Canadian Journal of Zoology. 2024;103:1-12.

11. Grémillet D, Ponchon A, Provost P, Gamble A, Abed-Zahar M, Bernard A, et al. Strong breeding colony fidelity in northern gannets following high pathogenicity avian influenza virus (HPAIV) outbreak. Biological Conservation. 2023;286:110269.

12. Simkin RD, Han BA, Radeloff VC, LaDeau S, Schug F, Seto KC. Zoonotic Host Richness in the Global Wildland–Urban Interface. Global Change Biology. 2025;31(2):e70039.

13. Hassell JM, Begon M, Ward MJ, Fèvre EM. Urbanization and disease emergence: dynamics at the wildlife–livestock–human interface. Trends in ecology & evolution. 2017;32(1):55-67.

14. Dietrich M, Gómez-Díaz E, McCoy KD. Worldwide distribution and diversity of seabird ticks: implications for the ecology and epidemiology of tick-borne pathogens. Vector borne and zoonotic diseases. 2011;11(5):453-70.

15. McCoy KD, Dietrich M, Jaeger A, Wilkinson DA, Bastien M, Lagadec E, et al. The role of seabirds of the Iles Eparses as reservoirs and disseminators of parasites and pathogens. Acta Oecologica. 2016;72:98-109.

16. Dhama K, Mahendran M, Tomar S. Pathogens transmitted by migratory birds: threat perceptions to poultry health and production. International Journal of Poultry Science. 2008;7(6):516-25.

17. Dias MP, Martin R, Pearmain EJ, Burfield IJ, Small C, Phillips RA, et al. Threats to seabirds: a global assessment. Biological conservation. 2019;237:525-37.

18. Cohen JM, Sauer EL, Santiago O, Spencer S, Rohr JR. Divergent impacts of warming weather on wildlife disease risk across climates. Science. 2020;370(6519):eabb1702.

19. Grimaldi WW, Seddon PJ, Lyver PO, Nakagawa S, Tompkins DM. Infectious diseases of Antarctic penguins: current status and future threats. Polar Biology. 2015;38(5):591-606.

20. Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, et al. Impacts of biodiversity on the emergence and transmission of infectious diseases. Nature. 2010;468(7324):647-52.

21. Matos AMRNd, Domit C, Bracarense APFRL. Seabirds: studies with parasitofauna and potential indicator for environmental anthropogenic impacts. Semina: Ciências Agrárias;41(4):1439-50. Number: 4. doi:10.5433/1679-0359.2020v41n4p1439.

22. Morse SS, Mazet JA, Woolhouse M, Parrish CR, Carroll D, Karesh WB, et al. Prediction and prevention of the next pandemic zoonosis. The Lancet. 2012;380(9857):1956-65.

23. Carlberg RA, Karris G, Verma M, Foufopoulos J. Food versus Disturbance: Contradictory Effects of Human Activities on an Opportunistic Seabird Breeding in an Oligotrophic Marine System. Diversity;14(6). doi:10.3390/d14060421.

24. Sadeghi S, Nikaeen M, Mohammadi F, Hossein Nafez A, Gholipour S, Shamsizadeh Z, et al. Microbial characteristics of municipal solid waste compost: Occupational and public health risks from surface applied compost. Waste Management;144. doi:10.1016/j.wasman.2022.03.012.

25. López-Calderón C, Martín-Vélez V, Blas J, Höfle U, Sánchez MI, Flack A, et al. White stork movements reveal the ecological connectivity between landfills and different habitats. Movement Ecology. 2023;11(1):18.

26. Altizer S, Bartel R, Han BA. Animal migration and infectious disease risk. Science. 2011;331(6015):296-302.

27. Plowright RK, Foley P, Field HE, Dobson AP, Foley JE, Eby P, et al. Urban habituation, ecological connectivity and epidemic dampening: the emergence of Hendra virus from flying foxes (Pteropus spp.). Proceedings of the Royal Society B: Biological Sciences. 2011;278(1725):3703-12.

28. Bradley CA, Gibbs SE, Altizer S. Urban land use predicts West Nile virus exposure in songbirds. Ecological Applications. 2008;18(5):1083-92.

29. Hubálek Z. Pathogenic microorganisms associated with gulls and terns (Laridae). Journal of Vertebrate Biology. 2021;70(3):21009-1.

30. Poulle ML, Le Corre M, Bastien M, Gedda E, Feare C, Jaeger A, et al. Exposure of pelagic seabirds to *Toxoplasma gondii* in the Western Indian Ocean points to an open sea dispersal of this terrestrial parasite. PLoS One. 2021;16(8):e0255664.

31. Bartumeus F, Giuggioli L, Louzao M, Bretagnolle V, Oro D, Levin SA. Fishery discards impact on seabird movement patterns at regional scales. Current Biology. 2010;20(3):215-22.

32. Cianchetti-Benedetti M, Dell'Omo G, Russo T, Catoni C, Quillfeldt P. Interactions between commercial fishing vessels and a pelagic seabird in the southern Mediterranean Sea. BMC Ecology. 2018;18:1-10.

33. Cortés-Hinojosa G. Marine bird of Neotropics, what we know, and we should know of diseases in a changing world. In: Ecology of Wildlife Diseases in the Neotropics. Springer; 2024. p. 121-41.

34. Jara-Carrasco S, González M, González-Acuña D, Chiang G, Celis J, Espejo W, et al. Potential immunohaematological effects of persistent organic pollutants on chinstrap penguin. Antarctic Science. 2015;27(4):373-81.

35. Fackelmann G, Pham CK, Rodríguez Y, Mallory ML, Provencher JF, Baak JE, et al. Current levels of microplastic pollution impact wild seabird gut microbiomes. Nature Ecology & Evolution. 2023;7(5):698-706.

36. de Jersey AM, Lavers JL, Bond AL, Wilson R, Zosky GR, Rivers-Auty J. Seabirds in crisis: Plastic ingestion induces proteomic signatures of multiorgan failure and neurodegeneration. Science Advances. 2025;11(11):eads0834.

37. McCoy KD, Toty C, Dupraz M, Tornos J, Gamble A, Garnier R, et al. Climate change in the Arctic: Testing the poleward expansion of ticks and tick-borne diseases. Global Change Biology. 2023;29(7):1729-40.

38. Van Hemert C, Pearce JM, Handel CM. Wildlife health in a rapidly changing North: focus on avian disease. Frontiers in Ecology and the Environment. 2014;12(10):548-56.

39. Bradley MJ, Kutz SJ, Jenkins E, O'Hara TM. The potential impact of climate change on infectious diseases of Arctic fauna. International Journal of Circumpolar Health. 2005;64(5):468-77.

40. Gutiérrez JS, Rakhimberdiev E, Piersma T, Thieltges DW. Migration and parasitism: habitat use, not migration distance, influences helminth species richness in Charadriiform birds. Journal of Biogeography;44(5):1137-47. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/jbi.12956. doi:10.1111/jbi.12956.

41. Fuirst M, Veit RR, Hahn M, Dheilly N, Thorne LH. Effects of urbanization on the foraging ecology and microbiota of the generalist seabird Larus argentatus. PLoS One. 2018;13(12):e0209200.

42. Holzman DC. Marine and Coastal Science: Where Space and Ocean Meet. Environmental Health Perspectives;115(5):A243. Publisher: National Institute of Environmental Health Sciences. doi:10.1289/EHP.115-A243A.

43. Carr KS, Bronze MS. Vertical Transplacental Infections. StatPearls Publishing, National Center for Biotechnology Information; 2023. Accessed: 2025-05-12. https://www.ncbi.nlm.nih.gov/books/NBK604195/.

44. Pedersen AB, Altizer S, Poss M, Cunningham AA, Nunn CL. Patterns of host specificity and transmission among parasites of wild primates. International Journal for Parasitology. 2005;35(6):647-57.

45. Thrusfield M. Veterinary epidemiology. John Wiley & Sons; 2013.

46. Werner CS, Nunn CL. Effect of urban habitat use on parasitism in mammals: a meta-analysis. Proceedings of the Royal Society B. 2020;287(1927):20200397.

47. Albery GF, Carlson CJ, Cohen LE, Eskew EA, Gibb R, Ryan SJ, et al. Urban-adapted mammal species have more known pathogens. Nature Ecology & Evolution. 2022;6(6):794-801.

48. Quillfeldt P, Arriero E, Martinez J, Masello JF, Merino S. Prevalence of blood parasites in seabirds-a review. Frontiers in Zoology. 2011;8(1):26.

49. Wilson A, Lapen D, Mitchell G, Provencher J, Wilson S. Interaction of diet and habitat predicts *Toxoplasma gondii* infection rates in wild birds at a global scale. Global Ecology and Biogeography;29(7). doi:10.1111/geb.13096.

50. Wilson A, Wilson S, Alavi N, Lapen D. Human density is associated with the increased prevalence of a generalist zoonotic parasite in mammalian wildlife. Proceedings of the Royal Society B: Biological Sciences;288(1961). doi:10.1098/rspb.2021.1724.

51. Wilson A, Lapen DR, Provencher JF, Wilson S. The role of species ecology in predicting *Toxoplasma gondii* prevalence in wild and domesticated mammals globally. PLOS Pathogens;20(1):e1011908. doi:10.1371/journal.ppat.1011908.

52. Benskin CMH, Wilson K, Jones K, Hartley IR. Bacterial pathogens in wild birds: a review of the frequency and effects of infection. Biological Reviews. 2009;84(3):349-73.

53. Delgado-V CA, French K. Parasite–bird interactions in urban areas: Current evidence and emerging questions. Landscape and urban planning. 2012;105(1-2):5-14.

54. Murray MH, Sánchez CA, Becker DJ, Byers KA, Worsley-Tonks KEL, Craft ME. City sicker? A meta-analysis of wildlife health and urbanization. Frontiers in Ecology and the Environment;17(10). doi:10.1002/fee.2126.

55. International Union for Conservation of Nature. The IUCN Red List of Threatened Species; 2025. Accessed on 29 May 2025. Available from: `https://www.iucnredlist.org`.

56. Mu H, Li X, Wen Y, Huang J, Du P, Su W, et al. A global record of annual terrestrial Human Footprint dataset from 2000 to 2018. Scientific Data;9(1). doi:10.1038/s41597-022-01284-8.

57. Halpern BS, Frazier M, Afflerbach J, Lowndes JS, Micheli F, O'Hara C, et al. Recent pace of change in human impact on the world's ocean. Scientific Reports;9(1):11609. Publisher: Nature Publishing Group. doi:10.1038/s41598-019-47201-9.

58. Venter O, Sanderson EW, Magrach A, Allan JR, Beher J, Jones KR, et al. Global terrestrial Human Footprint maps for 1993 and 2009. Scientific Data;3. doi:10.1038/sdata.2016.67.

59. Beck JJ, Li D, Johnson SE, Rogers D, Cameron KM, Sytsma KJ, et al. Functional traits mediate individualistic species-environment distributions at broad spatial scales while fine-scale species associations remain unpredictable. American Journal of Botany. 2022;109(12):1991-2005.

60. Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W. EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. Ecology;95(7). doi:10.1890/13-1917.1.

61. Stephens PR, Pappalardo P, Huang S, Byers JE, Farrell MJ, Gehman A, et al.. Global mammal parasite database version 2.0. Wiley Online Library; 2017.

62. Centers for Disease Control and Prevention. CDC website; 2025. Accessed: 2025-09-04. https://www.cdc.gov.

63. Onstad D, Fuxa J, Humber R, Oestergaard J, Shapiro-Ilan D, Gouli V, et al. An abridged glossary of terms used in invertebrate pathology. Society for Invertebrate Pathology. 2006.

64. Britt CL, Rocque M, Zimmerman GM. The analysis of bounded count data in criminology. Journal of Quantitative Criminology. 2018;34(2):591-607.

65. Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, et al. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. The R Journal. 2017;9(2):378-400. Available from: https://journal.r-project.org/archive/2017/RJ-2017-066/.

66. Hall DB. Zero-inflated Poisson and binomial regression with random effects: a case study. Biometrics. 2000;56(4):1030-9.

67. Ilgunas M, Palinauskas V, Platonova E, Iezhova T, Valkiunas G. The experimental study on susceptibility of common European songbirds to Plasmodium elongatum (lineage pGRW6), a widespread avian malaria parasite. Malaria Journal;18(1). doi:10.1186/s12936-019-2926-4.

68. Robles-Fernández L, Santiago-Alarcon D, Lira-Noriega A. Wildlife susceptibility to infectious diseases at global scales. Proceedings of the National Academy of Sciences of the United States of America;119(35). doi:10.1073/pnas.2122851119.

69. Brewer MJ, Butler A, Cooksley SL. The relative performance of AIC, AICC and BIC in the presence of unobserved heterogeneity. Methods in Ecology and Evolution. 2016;7(6):679-92.

70. Hartig F, Hartig MF. Package 'dharma'. R package. 2017;531:532.

71. Becker DJ, Streicker DG, Altizer S. Linking anthropogenic resources to wildlife–pathogen dynamics: a review and meta-analysis. Ecology Letters. 2015;18(5):483-95.

72. Jacobson KC, Arkoosh MR, Kagley AN, Clemons ER, Collier TK, Casillas E. Cumulative effects of natural and anthropogenic stress on immune function and disease resistance in juvenile Chinook salmon. Journal of Aquatic Animal Health. 2003;15(1):1-12.

73. Martin LB, Hopkins WA, Mydlarz LD, Rohr JR. The effects of anthropogenic global changes on immune functions and disease resistance. Annals of the New York Academy of Sciences. 2010;1195(1):129-48.

74. Rollins-Smith LA. Amphibian immunity–stress, disease, and climate change. Developmental & Comparative Immunology. 2017;66:111-9.

75. Palmer CV. Immunity and the coral crisis. Communications Biology. 2018;1(1):91.

76. Novotny L, Dvorska L, Lorencova A, Beran V, Pavlik I, et al. Fish: a potential source of bacterial pathogens for human beings. Veterinární Medicína. 2004;49(9):343-58.

77. Marcogliese D. Food webs and the transmission of parasites to marine fish. Parasitology. 2002;124(7):83-99.

78. Carravieri A, Burthe SJ, de La Vega C, Yonehara Y, Daunt F, Newell MA, et al. Interactions between environmental contaminants and gastrointestinal parasites: novel insights from an integrative approach in a marine predator. Environmental Science & Technology. 2020;54(14):8938-48.

79. Bracey AM, Etterson MA, Strand FC, Matteson SW, Niemi GJ, Cuthbert FJ, et al. Foraging ecology differentiates life stages and mercury exposure in common terns (Sterna hirundo). Integrated Environmental Assessment and Management. 2020;17(2):398-410.

80. Sunderland EM, Hu XC, Dassuncao C, Tokranov AK, Wagner CC, Allen JG. A review of the pathways of human exposure to poly-and perfluoroalkyl substances (PFASs) and present understanding of health effects. Journal of Exposure Science & Environmental Epidemiology. 2019;29(2):131-47.

81. Whitney MC, Cristol DA. Impacts of sublethal mercury exposure on birds: a detailed review. Reviews of Environmental Contamination and Toxicology Volume 244. 2017:113-63.

82. Grasman KA, Fox GA. Associations between altered immune function and organochlorine contamination in young Caspian terns (*Sterna caspia*) from Lake Huron, 1997–1999. Ecotoxicology. 2001;10(2):101-14.

83. Teitelbaum CS, Ackerman JT, Hill MA, Satter JM, Casazza ML, De La Cruz SE, et al. Avian influenza antibody prevalence increases with mercury contamination in wild waterfowl. Proceedings of the Royal Society B. 2022;289(1982):20221312.

84. Idle JL, Wilhite CJ, Harmon KC, Friswold B, Price MR. Wedge-tailed Shearwater (*Ardenna pacifica*) nesting success in human-dominated coastal environments. PeerJ. 2021;9:e12096.

85. Macfarlane AE. Roof-nesting by common terns. Wilson Bulletin. 1977;89(3):22.

86. Clancy C, Ward K. Auto-rewilding in post-industrial cities: the case of inland cormorants in urban Britain. Conservation and Society. 2020;18(2):126-36.

87. Pichegru L, Sherley R, Malan T, Barham B, Ludynia K, Geldenhuys D, et al. Decades of artificial nests towards African penguin conservation—Have they made a difference? Ecological Solutions and Evidence. 2024;5(4):e12388.

88. Verhagen JH, Munster VJ, Majoor F, Lexmond P, Vuong O, Stumpel JB, et al. Avian influenza a virus in wild birds in highly urbanized areas. PLoS One. 2012;7(6):e38256.

89. Burr PC, Dorr BS, Avery JL, Street GM, Strickland BK. Long term changes in aquaculture influence migration, regional abundance, and distribution of an avian species. PloS One. 2023;18(4):e0284265.

90. Thomas FL, Forys EA. The Role of Fishing Piers in Brown Pelican (*Pelecanus occidentalis*) Entanglement. Animals. 2022;12(18):2352.

91. Ceia FR, Silva NC, Paiva VH, Morais L, Serrão EA, Ramos JA. Gulls as indicators of environmental changes in the North Atlantic: A long-term study on Berlenga Island, Western Portugal. Diversity. 2023;15(11):1148.

92. Abell R, Thieme ML, Revenga C, Bryer M, Kottelat M, Bogutskaya N, et al. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. BioScience;58(5):403-14. Publisher: American Institute of Biological Sciences.

93. Miller S, Zieger U, Ganser C, Satterlee SA, Bankovich B, Amadi V, et al. Influence of land use and climate on Salmonella carrier status in the small Indian mongoose (*Herpestes auropunctatus*) in Grenada, West Indies. Journal of Wildlife Diseases. 2015;51(1):60-8.

94. Fleischer R, Eibner GJ, Schwensow NI, Pirzer F, Paraskevopoulou S, Mayer G, et al. Immunogenetic-pathogen networks shrink in Tome's spiny rat, a generalist rodent inhabiting disturbed landscapes. Communications Biology. 2024;7(1):169.

95. Hernandez SM, Welch CN, Peters VE, Lipp EK, Curry S, Yabsley MJ, et al. Urbanized white ibises (Eudocimus albus) as carriers of Salmonella enterica of significance to public health and wildlife. PLoS One. 2016;11(10):e0164402.

96. Seixas JS, Hernandez SM, Christie KF, Norfolk WA, Rozier RS, Kurimo-Beechuk E, et al. Salmonella infections in white ibis nestlings in urban areas as a result of environmental contamination reveal a potential health risk to humans and animals. Journal of the American Veterinary Medical Association. 2025;1(aop):1-8.

97. Ramos R, Cerdà-Cuéllar M, Ramírez F, Jover L, Ruiz X. Influence of refuse sites on the prevalence of Campylobacter spp. and Salmonella serovars in seagulls. Applied and Environmental Microbiology. 2010;76(9):3052-6.

98. Van Heugten E, Coffey M, Spears J. Effects of immune challenge, dietary energy density, and source of energy on performance and immunity in weanling pigs. Journal of Animal Science. 1996;74(10):2431-40.

99. Coop RL, Kyriazakis I. Influence of host nutrition on the development and consequences of nematode parasitism in ruminants. Trends in Parasitology. 2001;17(7):325-30.

100. Ezenwa VO. Interactions among host diet, nutritional status and gastrointestinal parasite infection in wild bovids. International Journal for Parasitology. 2004;34(4):535-42.

101. Cummings CR, Hernandez SM, Murray M, Ellison T, Adams HC, Cooper RE, et al. Effects of an anthropogenic diet on indicators of physiological challenge and immunity of white ibis nestlings raised in captivity. Ecology and Evolution. 2020;10(15):8416-28.

102. BirdLife International. IUCN Red List for birds; 2025. Accessed 2025-11-05. Available from: `https://datazone.birdlife.org/`.

103. La Sala LF, Petracci PF, Randazzo V, Fernández-Miyakawa ME. Enteric bacteria in Olrog's gull (*Larus atlanticus*) and kelp gull (Larus dominicanus) from the Bahia Blanca Estuary, Argentina. El hornero. 2013;28(2):59-64.

104. Descamps S. Winter temperature affects the prevalence of ticks in an Arctic seabird. PLoS One. 2013;8(6):e65374.

105. Epstein PR. West Nile virus and the climate. Journal of Urban Health. 2001;78(2):367-71.

106. Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West nile virus. The Lancet infectious diseases. 2002;2(9):519-29.

107. Bastien M, Jaeger A, Le Corre M, Tortosa P, Lebarbenchon C. Haemoproteus iwa in great frigatebirds (*Fregata minor*) in the islands of the Western Indian Ocean. PLoS One. 2014;9(5):e97185.

108. Quillfeldt P, Strange IJ, Segelbacher G, Masello JF. Male and female contributions to provisioning rates of thin-billed prions, *Pachyptila belcheri*, in the South Atlantic. Journal of Ornithology. 2007;148(3):367-72.

109. Strange IJ. The Thin-billed Prion, *Pachyptila belcheri*, at New Island, Falkand Islands [breeding sites; nest sites; behaviour; homing; predation, chick mortality]. Gerfaut. 1980;70.

110. Einoder LD. A review of the use of seabirds as indicators in fisheries and ecosystem management. Fisheries Research. 2009;95(1):6-13.

111. Williams CT, Buck CL, Sears J, Kitaysky AS. Effects of nutritional restriction on nitrogen and carbon stable isotopes in growing seabirds. Oecologia. 2007;153(1):11-8.

112. Hedd A, Fifield DA, Burke CM, Montevecchi WA, Tranquilla LM, Regular PM, et al. Seasonal shift in the foraging niche of Atlantic puffins Fratercula arctica revealed by stable isotope (δ15N and δ13C) analyses. Aquatic Biology. 2010;9(1):13-22.

113. Wilson A, Wilson S, Alavi N, Lapen DR. Human density is associated with the increased prevalence of a generalist zoonotic parasite in mammalian wildlife. Proceedings of the Royal Society B. 2021;288(1961):20211724.

114. Wilson A, Lapen DR, Provencher JF, Wilson S. The role of species ecology in predicting *Toxoplasma gondii* prevalence in wild and domesticated mammals globally. PLoS Pathogens. 2024;20(1):e1011908.

115. Pereira JM, Ramos JA, Domingues A, Almeida A, Marçalo A, Cascão C, et al. Experimental anthropogenic food restrictions drive short-term foraging and immuno-haematological changes in sympatric breeding gulls. Science of the Total Environment. 2025;1003:180672.

116. Zorrozua N, Aldalur A, Herrero A, Beñat D, Delgado S, Sanpera C, et al. Breeding Yellow-legged Gulls increase consumption of terrestrial prey after landfill closure. Ibis. 2020;162(1):50-62. doi:10.1111/ibi.12701.

117. Delgado S, Herrero A, Galarza A, Aldalur A, Zorrozua N, Arizaga J. Demographic impact of landfill closure on a resident opportunistic gull. Population Ecology. 2021;63(3):238-46. doi:10.1002/1438-390X.12083.

118. Warmuth VM, Metzler D, Zamora-Gutierrez V. Human disturbance increases coronavirus prevalence in bats. Science Advances. 2023;9(13):eadd0688.

119. North BV, Curtis D, Sham PC. A note on the calculation of empirical P values from Monte Carlo procedures. The American Journal of Human Genetics. 2002;71(2):439-41.

120. Martín-Vélez V, Navarro J, Figuerola J, Aymí R, Sabaté S, Planell R, et al. A spatial analysis of urban gulls contribution to the potential spread of zoonotic and antibiotic-resistant bacteria. Science of The Total Environment. 2024 Feb;912:168762. doi:10.1016/j.scitotenv.2023.168762.

121. Kay E, Young MJ, Muller C, Howe L, Roe W, Gartrell BD. Prevalence and pathogen load of Eimeria in wild yellow-eyed penguins *(Megadyptes antipodes)* and the morphologic characterization of a novel Eimeria species. Journal of Wildlife Diseases. 2022 Nov;58(4). doi:10.7589/jwd-d-21-00146.

122. Gallo Vaulet L, Vanstreels RET, Gallo L, Entrocassi AC, Peker L, Blanco GS, et al. Chlamydiaceae-Like Bacterium in Wild Magellanic Penguins *(Spheniscus magellanicus)*. Diversity. 2022 Sep;14(9):746. doi:10.3390/d14090746.

123. Poulle ML, Le Corre M, Bastien M, Gedda E, Feare C, Jaeger A, et al. Exposure of pelagic seabirds to *Toxoplasma gondii* in the Western Indian Ocean points to an open sea dispersal of this terrestrial parasite. PLoS One. 2021 Aug;16(8):e0255664. doi:10.1371/journal.pone.0255664.

124. Odoi JO, Sugiyama M, Kitamura Y, Sudo A, Omatsu T, Asai T. Prevalence of antimicrobial resistance in bacteria isolated from Great Cormorants *(Phalacrocorax carbo hanedae)* in Japan. Journal of Veterinary Medical Science. 2021;83(8):1191-5.

125. Gamble A, Bazire R, Delord K, Barbraud C, Jaeger A, Gantelet H, et al. Predator and scavenger movements among and within endangered seabird colonies: Opportunities for pathogen spread. Journal of Applied Ecology. 2019 Dec;57(2):367–378. doi:10.1111/1365-2664.13531.

126. Verdugo C, Pinto A, Ariyama N, Moroni M, Hernandez C. Molecular identification of avian viruses in neotropic cormorants (*Phalacrocorax brasilianus*) in Chile. Journal of Wildlife Diseases. 2019;55(1):105-12.

127. Navarro J, Grémillet D, Afán I, Miranda F, Bouten W, Forero MG, et al. Pathogen transmission risk by opportunistic gulls moving across human landscapes. Scientific Reports. 2019 Jul;9(1). doi:10.1038/s41598-019-46326-1.

128. Espinaze MPA, Hui C, Waller L, Dreyer F, Matthee S. Parasite diversity associated with African penguins *(Spheniscus demersus)* and the effect of host and environmental factors. Parasitology. 2019 Feb;146(6):791–804. doi:10.1017/s0031182018002159.

129. Pedro R, Claudio N, Elena C, Claudio V. Low occurrence of hemosporidian parasites in the Neotropic cormorant (*Phalacrocorax brasilianus*) in Chile. Parasitology Research. 2018 Nov;118(1):325–333. doi:10.1007/s00436-018-6146-6.

130. Campioni L, Martínez-de la Puente J, Figuerola J, Granadeiro JP, Silva MC, Catry P. Absence of haemosporidian parasite infections in the long-lived Cory's shearwater: evidence from molecular analyses and review of the literature. Parasitology Research. 2017 Nov;117(1):323–329. doi:10.1007/s00436-017-5676-7.

131. de Souza Petersen Elisa, de Araujo Jansen, Lucas K, M SM, Tatiana O, M TL, et al. First detection of avian influenza virus (H4N7) in giant petrel monitored by geolocators in the Antarctic region. Marine Biology. 2017;164(4). doi:10.1007/s00227-017-3086-0.

132. Vanstreels RET, Uhart M, Rago V, Hurtado R, Epiphanio S, Catao-Dias JL. Do blood parasites infect Magellanic penguins *(Spheniscus magellanicus)* in the wild? Prospective investigation and climatogeographic considerations. Parasitology. 2017;144(5):698-705.

133. Barriga GP, Boric-Bargetto D, San Martin MC, Neira V, van Bakel H, Thompsom M, et al. Avian Influenza Virus H5 Strain with North American and Eurasian Lineage Genes in an Antarctic Penguin. Emerging Infectious Diseases. 2016 Dec;22(12):2221–2223. doi:10.3201/eid2212.161076.

134. Zagalska-Neubauer M, Bensch S. High prevalence of Leucocytozoon parasites in fresh water breeding gulls. Journal of Ornithology. 2015 Sep;157(2):525–532. doi:10.1007/s10336-015-1291-5.

135. Kim HK, Choi CY, Jeong MS, Kang HY, Lee WS. Regurgitation of the koilin layer in chinstrap penguins (*Pygoscelis antarcticus*) and its association with gastric parasites. Polar Research. 2016 Jan;35(1):25966. doi:10.3402/polar.v35.25966.

136. Lebarbenchon C, Jaeger A, Feare C, Bastien M, Dietrich M, Larose C, et al. Influenza A virus on oceanic islands: Host and viral diversity in seabirds in the western Indian Ocean. PLoS Pathogens. 2015 May;11(5):e1004925. doi:10.1371/journal.ppat.1004925.

137. Hammouda A, Pearce-Duvet J, Boulinier T, Selmi S. Egg sampling as a possible alternative to blood sampling when monitoring the exposure of yellow-legged gulls *(Larus michahellis)* to avian influenza viruses. Avian Pathology. 2014 Nov;43(6):547–551. doi:10.1080/03079457.2014.972340.

138. Abad FX, Busquets N, Sanchez A, Ryan PG, Majó N, Gonzalez-Solís J. Serological and virological surveys of the influenza A viruses in Antarctic and sub-Antarctic penguins. Antarctic Science. 2013 Mar;25(2):339–344. doi:10.1017/s0954102012001228.

139. Rivera WA, Husic JS, Gaylets CE, Mosher RH, Palestis BG, Houlihan AJ. Carriage of bacteria and protozoa in the intestinal tract of common tern chicks. Waterbirds. 2012 Sep;35(3):490–494. doi:10.1675/063.035.0314.

140. Levin II, Parker PG. Prevalence of Hemoproteus iwa in Galapagos Great Frigatebirds *(Fregata minor)* and Their Obligate Fly Ectoparasite *(Olfersia spinifera).* Journal of Parasitology. 2012 Oct;98(5):924–929. doi:10.1645/ge-3027.1.

141. Levin II, Outlaw DC, Vargas FH, Parker PG. Plasmodium blood parasite found in endangered Galapagos penguins *(Spheniscus mendiculus).* Biological Conservation. 2009 Dec;142(12):3191–3195. doi:10.1016/j.biocon.2009.06.017.

142. Merino S, Hennicke J, Martínez J, Ludynia K, Torres R, Work TM, et al. Infection by Haemoproteus Parasites in Four Species of Frigatebirds and the Description of a New Species of Haemoproteus (*Haemosporida: Haemoproteidae*). Journal of Parasitology. 2012 Apr;98(2):388–397. doi:10.1645/ge-2415.1.

143. Krams I, Suraka V, Rattiste K, Āboliņš Ābols M, Krama T, Rantala MJ, et al. Comparative analysis reveals a possible immunity-related absence of blood parasites in Common Gulls *(Larus canus)* and Black-headed Gulls *(Chroicocephalus ridibundus).* Journal of Ornithology. 2012 May;153(4):1245–1252. doi:10.1007/s10336-012-0859-6.

144. Lobato E, Pearce-Duvet J, Staszewski V, Gómez-Díaz E, González-Solís J, Kitaysky A, et al. Seabirds and the Circulation of Lyme Borreliosis Bacteria in the North Pacific. Vector-Borne and Zoonotic Diseases. 2011 Dec;11(12):1521–1527. doi:10.1089/vbz.2010.0267.

145. Quillfeldt P, Martínez J, Hennicke J, Ludynia K, Gladbach A, Masello JF, et al. Hemosporidian blood parasites in seabirds—a comparative genetic study of species from Antarctic to tropical habitats. Naturwissenschaften. 2010 Jul;97(9):809–817. doi:10.1007/s00114-010-0698-3.

146. Wojczulanis-Jakubas K, Svoboda A, Kruszewicz A, Johnsen A. No Evidence of Blood Parasites in Little Auks *(Alle alle)* Breeding on Svalbard. Journal of Wildlife Diseases. 2010 Apr;46(2):574–578. doi:10.7589/0090-3558-46.2.574.

147. Staszewski V, McCoy KD, Boulinier T. Variable exposure and immunological response to Lyme disease Borrelia among North Atlantic seabird species. Proceedings of the Royal Society B: Biological Sciences. 2008 Jun;275(1647):2101–2109. doi:10.1098/rspb.2008.0515.

148. Mendes L, Piersma T, Lecoq M, Spaans B, E Ricklefs R. Disease-limited distributions? Contrasts in the prevalence of avian malaria in shorebird species using marine and freshwater habitats. Oikos. 2005 Mar;109(2):396–404. doi:10.1111/j.0030-1299.2005.13509.x.

149. Fouchier RAM, Munster V, Wallensten A, Bestebroer TM, Herfst S, Smith D, et al. Characterization of a Novel Influenza A Virus Hemagglutinin Subtype (H16) Obtained from black-headed gulls. Journal of Virology. 2005 Mar;79(5):2814–2822. doi:10.1128/jvi.79.5.2814-2822.2005.

150. Wallensten A, Munster VJ, Elmberg J, Osterhaus ADME, Fouchier RAM, Olsen B. Multiple gene segment reassortment between Eurasian and American lineages of influenza A virus (H6N2) in guillemot *(Uria aalge)*. Archives of Virology. 2005 May;150(8):1685–1692.

151. Hollmén T, Franson JC, Docherty DE, Kilpi M, Hario M, Creekmore LH, et al. Infectious bursal disease virus antibodies in eider ducks and herring gulls. The Condor. 2000 Aug;102(3):688–691. doi:10.1093/condor/102.3.688.

152. Jimenez-Uzcategui G, Velez A, Vega P, Buendia V, Montenegro-Benalcázar V, Sevilla C, et al. No evidence for high pathogenicity avian influenza in Waved Albatross (*Pvhoebastria irrorata*). Marine Ornithology. 2024;52(2):349-53.

153. Bregnballe T, Herrmann C, Globig A, Günther A, Staubach C, Heise JN, et al. Outbreaks of highly pathogenic avian influenza (HPAI) epidemics in Baltic Great Cormorant *Phalacrocorax carbo* colonies in 2021 and 2022. Bird Study. 2024 Sep;71(4):353–366. doi:10.1080/00063657.2024.2399168.

154. Ushine N, Ozawa M, Nakayama SM, Ishizuka M, Kato T, Hayama Si. Evaluation of the effect of Pb pollution on avian influenza virus-specific antibody production in black-headed gulls (*Chroicocephalus ridibundus*). Animals. 2023;13(14):2338.

155. Lobato E, Pearce-Duvet J, Staszewski V, Gomez-Diaz E, González-Solís J, Kitaysky A, et al. Seabirds and the circulation of Lyme borreliosis bacteria in the North Pacific. Vector-Borne and Zoonotic Diseases. 2011;11(12):1521-7.

156. Zhao Y, Aarnink A, Cambra-Lopez M, Fabri T. Shedding and Emission of Airborne Viral Microorganisms from Animal Houses. In: 2012 IX International Livestock Environment Symposium (ILES IX). American Society of Agricultural and Biological Engineers; 2012. p. 3.

157. Hernández AC. Poultry and avian diseases. Encyclopedia of Agriculture and Food Systems. 2014:504.

158. González-Acuña DA, Moreno L, Wille M, Herrmann B, Kinsella MJ, Palma RL. Parasites of chinstrap penguins (*Pygoscelis antarctica*) from three localities in the Antarctic Peninsula and a review of their parasitic fauna. Polar Biology. 2021;44(11):2099-105.

159. Murphy JR. Corynebacterium diphtheriae. Medical Microbiology 4th edition. 1996.

160. Clifton IJ, Peckham DG. Defining routes of airborne transmission of Pseudomonas aeruginosa in people with cystic fibrosis. Expert Review of Respiratory Medicine. 2010;4(4):519-29.

161. Battaglia M, Garrett-Sinha LA. Staphylococcus xylosus and Staphylococcus aureus as commensals and pathogens on murine skin. Laboratory Animal Research. 2023;39(1):18.

162. Lanao AE, Chakraborty RK, Pearson-Shaver AL. Mycoplasma infections. In: StatPearls [Internet]. StatPearls Publishing; 2023. .

163. Wong P, Anderson R. The transmission and development of Cosmocephalus obvelatus (Nematoda: Acuarioidea) of gulls (Laridae). Canadian Journal of Zoology. 1982;60(6):1426-40.

164. Chai JY, Jung BK. Fishborne zoonotic heterophyid infections: an update. Food and Waterborne Parasitology. 2017;8:33-63.

165. Gonchar A, Jouet D, Skírnisson K, Krupenko D, Galaktionov KV. Transatlantic discovery of Notocotylus atlanticus (Digenea: Notocotylidae) based on life cycle data. Parasitology Research. 2019;118(5):1445-56.

166. Hoberg EP. Evolution and historical biogeography of a parasite–host assemblage: Alcataenia spp.(Cyclophyllidea: Dilepididae) in Alcidae (Charadriiformes). Canadian Journal of Zoology. 1986;64(11):2576-89.

167. Blakeslee AM, Keogh CL, Fowler AE, Griffen BD. Assessing the effects of trematode infection on invasive green crabs in eastern North America. PLoS One. 2015;10(6):e0128674.

168. Sewoyo PS, Putra IPC, Nainggolan WM. Pathology of proventricular tetrameriasis in a free-range chicken. ARSHI Veterinary Letters. 2024;8(3):47-8.

169. Traversa D, Di Cesare A, Conboy G. Canine and feline cardiopulmonary parasitic nematodes in Europe: emerging and underestimated. Parasites & vectors. 2010;3(1):62.

170. Gray GC. Adenovirus transmission—worthy of our attention. The Journal of infectious diseases. 2006;194(7):871-3.

171. Wozniakowski G, Samorek-Salamonowicz E. Animal herpesviruses and their zoonotic potential for cross-species infection. Annals of Agricultural and Environmental Medicine. 2015;22(2).

172. Woźniakowski G, Frant M, Mamczur A. Avian reticuloendotheliosis in chickens–an update on disease occurrence and clinical course. Journal of Veterinary Research. 2018;62(3):257.

173. Sapiro AL, Hayes BM, Volk RF, Zhang JY, Brooks DM, Martyn C, et al. Longitudinal map of transcriptome changes in the Lyme pathogen Borrelia burgdorferi during tick-borne transmission. Elife. 2023;12:RP86636.

174. Cheong HC, Lee CYQ, Cheok YY, Tan GMY, Looi CY, Wong WF. Chlamydiaceae: diseases in primary hosts and zoonosis. Microorganisms. 2019;7(5):146.

175. Du XY, Zhang PF, Gong SR, Liang YS, Huang YH, Li HS, et al. Discovery of a novel circulation route of free-living Serratiasymbiotica mediated by predatory ladybird beetles. FEMS Microbiology Ecology. 2023;99(11):fiad133.

Supplementary Material

A Materials and Methods

A.1 Parametric bootstrap likelihood ratio

To test the significance of each alternative model, which includes the H.Sea and H.Land, we created a null model including only diet as a fixed effect, except for vector-borne model whose selected model, and therefore the null model, did no include diet. We used this fitted model to simulate 1,000 new datasets, and then for each simulated dataset, we refitted both null and alternative models fitted based on the BIC-selected model and computed the corresponding likelihood ratios. We calculated the p-value as $(k + 1)/1001$, where k is the number of simulated datasets in which the likelihood ratio was equal to or greater than that observed in the original dataset [118, 119].

genus

- Larus (0.222)
- Spheniscus (0.12)
- Pygoscelis (0.106)
- Phalacrocorax (0.057)
- Phaethon (0.049)
- Fregata (0.045)
- Hydrobates (0.035)
- Sula (0.033)
- Uria (0.03)
- Anous (0.028)
- Rissa (0.028)
- Chroicocephalus (0.022)
- Onychoprion (0.022)
- Puffinus (0.02)
- Aethia (0.016)
- Fratercula (0.016)
- Catharacta (0.014)
- Leucophaeus (0.014)
- Sterna (0.014)
- Eudyptes (0.012)
- Ardenna (0.01)
- Calonectris (0.01)
- Stercorarius (0.01)
- Aptenodytes (0.008)
- Thalasseus (0.008)
- Alle (0.006)
- Oceanites (0.006)
- Pachyptila (0.006)
- Papasula (0.006)
- Pterodroma (0.006)
- Macronectes (0.004)
- Megadyptes (0.004)
- Phalaropus (0.004)
- Phoebastria (0.004)
- Leucocarbo (0.002)

Fig S1. There are 35 genera of seabirds in our study. Gulls have the highest proportion (Larus 22%), followed by penguins (Spheniscus 11% and Pygoscelis 9%) and cormorants (Phalacrocorax 6%); all other genera each contributed 5% or less.

Table S1. The results of general BIC-selected model

Model #	Model (covariates)	BIC	ΔBIC
1	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn	5254.6	0.00
2	diet + H.Lnd + H.Mrn + diet:H.Lnd	5260.3	5.75
3	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn + zi(mtC)*	2560.6	6.05
4	diet + H.Lnd + H.Mrn + diet:H.Mrn	5265.0	10.44
5	diet + H.Lnd + H.Mrn + diet:H.Mrn + zi(mtC)	5271.1	16.51
6	diet + H.Lnd + H.Mrn + diet:H.Lnd + zi(mtC)	5271.9	17.34
7	diet + H.Mrn + diet:H.Mrn	5284.7	30.17
8	diet + H.Mrn + diet:H.Mrn + zi(mtC)	5289.8	35.23
9	diet + H.Lnd + diet:H.Lnd + zi(mtC)	5303.4	48.78
10	H.Lnd	5368.9	114.28
11	(null)	5368.6	115.01
12	H.Marine	5373.2	118.59
13	diet + H.Lnd	5373.9	119.36
14	H.Lnd + H.Mrn	5374.1	119.56
15	diet	5374.3	119.73
16	H.Land + zi(mtC)	5375.0	120.47
17	diet + H.Marine + zi(mtC)	5375.7	121.14
18	diet + H.Lnd + H.Mrn	5378.0	123.43
19	H.MArine	5379.3	124.68
20	diet + H.Land + zi(mtC)	5379.3	124.70
21	H.Land + H.Marine + zi(mtC)	5380.1	125.56
22	diet + zi(mtC)	5380.3	125.73
23	H.Marine + zi(mtC)	5380.5	125.89
24	diet + H.Lnd + H.Mrn	5384.1	129.56
25	diet + H.Lnd + zi(mtC)	5385.4	130.86
26	diet + H.Lnd + diet:H.Lnd	-	-

All models include conditional, dispersion, and zero-inflation intercepts.
*zi(mtC) is the method of pathogen detection in each sample in the zero-inflation.

Table S2. The results of contact-based BIC-selected model

Model #	Model (covariates)	BIC	ΔBIC
1	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn	1592.5	0.00
2	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn + zi(mtC)	1594.5	2.02
3	diet + H.Mrn + diet:H.Mrn	1599.1	6.56
4	diet + H.Lnd + H.Mrn + diet:H.Mrn	1599.2	6.65
5	diet + H.Lnd + H.Mrn + diet:H.Lnd	1601.0	8.46
6	diet + H.Lnd + H.Mrn + diet:H.Lnd + zi(mtC)	1601.1	8.58
7	diet + H.Lnd + H.Mrn + diet:H.Mrn + zi(mtC)	1601.2	8.67
8	diet + H.Mrn + diet:H.Mrn + zi(mtC)	1601.5	8.98
9	diet + H.Lnd + diet:H.Lnd	1607.6	15.07
10	diet + H.Lnd + diet:H.Lnd + zi(mtC)	1610.3	17.79
11	H.Lnd	1641.4	48.92
12	H.Lnd + zi(mtC)	1643.6	51.09
13	H.Lnd + H.Mrn	1645.6	53.07
14	H.Lnd + H.Mrn + zi(mtC)	1648.1	55.56
15	H.Mrn +	1651.1	58.60
16	diet + H.Lnd	1652.1	59.57
17	(null)	1652.5	16.01
18	diet + H.Lnd + zi(mtC)	1654.2	61.73
19	H.Mrn	1654.7	62.22
20	zi(mtC)	1655.4	62.93
21	diet + H.Lnd + H.Mrn + zi(mtC)	1656.2	63.69
22	diet + H.Lnd + H.Mrn + zi(mtC)	1658.7	66.19
23	diet + H.Mrn	1661.7	69.24
24	diet	1663.2	70.69
25	diet + H.Mrn + zi(mtC)	1665.4	72.87
26	diet + zi(mtC)	-	-

All models include conditional, dispersion, and zero-inflation intercepts.

Table S3. The results of vector-borne BIC-selected model

Model #	Model (covariates)	BIC	ΔBIC
1	H.Lnd + H.Mrn	606.3	0.00
2	H.Lnd + H.Mrn + zi(mtC)	611.6	5.25
3	diet + H.Lnd + H.Mrn + diet:H.Mrn	613.8	7.45
4	diet + H.Lnd + H.Mrn	614.2	7.91
5	diet + H.Lnd + H.Mrn + diet:H.Mrn + zi(mtC)	619.0	12.68
6	diet + H.Lnd + H.Mrn + zi(mtC)	619.5	13.15
7	diet + H.Lnd + H.Mrn + diet:H.Lnd	621.7	15.38
8	H.Lnd	622.5	16.16
9	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn	624.0	17.72
10	diet + H.Lnd + H.Mrn + diet:H.Lnd + zi(mtC)	626.9	20.62
11	H.Lnd + zi(mtC)	627.7	21.41
12	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn + zi(mtC)	629.3	22.96
13	diet + H.lnd	629.9	23.59
14	diet + H.Lnd + zi(mtC)	635.1	28.84
15	diet + H.Lnd + diet:H.lnd	639.7	33.43
16	diet + H.Lnd + diet:H.lnd + zi(mtC)	645.0	38.68
17	H.Mrn	655.6	49.25
18	H.Mrn + zi(mtC)	660.8	54.49
19	diet + H.Mrn	663.8	57.49
20	diet + H.Mrn + diet:H.Mrn	668.9	63.56
21	diet + H.Mrn + zi(mtC)	669.0	62.72
22	diet + H.Mrn + diet:H.Mrn + zi(mtC)	674.1	67.81
23	(null)	676.7	70.39
24	zi(mtC)	681.9	75.63
25	diet	684.5	78.17
26	diet + zi(mtC)	689.7	83.42

All models include conditional, dispersion, and zero-inflation intercepts.

Table S4. The results of Intermediate host BIC-selected model

Model #	Model (covariates)	BIC	ΔBIC
1	(null)		0.00
2	zi(mtC)	3258.8	3.03
3	H.Mrn	3259.3	3.56
4	H.Lnd	3259.4	4.19
5	diet	3261.9	6.17
6	diet + H.Lnd + zi(mtC)	3262.5	6.75
7	H.Mrn + zi(mtC)	3262.9	7.12
8	H.Mrn + H.Lnd	3263.4	7.68
9	H.Lnd + H.Mrn	3264.2	8.48
10	diet + zi(mtC)	3264.9	9.19
11	diet + H.Lnd + H.Mrn	3266.0	10.24
12	diet + H.Lnd + H.Mrn + zi(mtC)	3266.6	10.89
13	diet + H.Mrn	3268.0	12.30
14	diet + H.Lnd + H.Mrn + zi(mtC)	635.1	28.84
15	diet + H.Mrn + diet:H.Mrn	3268.2	12.49
16	diet + H.Lnd + zi(mtC)	3268.4	12.68
17	diet + H.Lnd + diet:H.Lnd	3268.5	12.75
18	diet + H.Lnd + diet:H:Lnd + zi(mtC)	3270.6	14.87
19	diet + H.Lnd + H.Mrn + zi(mtC)	3271.7	16.01
20	diet + H.Mrn + diet:H.Mrn + zi(mtC)	3271.8	16.02
21	diet + H.Lnd + H.Mrn + diet:H.Lnd	3272.0	16.31
22	diet + H.Lnd + H.Mrn + diet:H.Lnd + diet:H.Mrn	3272.0	16.31
23	diet + H.Lnd + H.Mrn + diet:H.Mrn	3272.6	16.31
24	diet + H.Lnd + H.Mrn + diet:H.Lnd	3275.0	19.22
25	diet + H.Lnd + H.Mrn + diet.H.Mrn + zi(mtC)	3275.0	19.22
26	diet + H.Lnd + H.Mrn + diet:H.Mrn + zi(mtC)	3275.0	19.22

All models include conditional, dispersion, and zero-inflation intercepts.

Fig S2. PRISMA flow diagram. We excluded studies lacking details on sample size, location, seabird species, or pathogen species. We also excluded study designs likely to bias our results, including experimental studies, samples from captive birds, rehabilitation centers, or carcasses, and studies where the number of individuals was not clear, and where sampling relied on clinical signs and diagnoses were based on symptoms rather than laboratory tests, and studies with sampling locations more than 50 km from the coastline. We collected 492 datasets from 54 studies, comprising 35 seabird genera and 65 species.

Fig S3. Residual diagnostics for prevalence models. DHARMa Q–Q plots of simulated residuals. Each panel reports the Kolmogorov–Smirnov tests. The general, contact-based, and vector-borne models show a small deviation, however the deviation for intermediate host model is large.

Fig S4. Predicted pathogen prevalence in seabirds by diet and human disturbance with 95% confidence interval.

Fig S5. Odds ratio of seabird genera being infected with pathogens. Only two genera of Pachyptila and Fregata showed odds ratio over 1 with confidence interval not crossing 1 in vector-borne model. It means that these genera are more susceptible to vector-borne pathogens.

Table S5. List of papers used in the meta-analysis

Title	Reference
A spatial analysis of urban gulls contribution to the potential spread of zoonotic and antibiotic-resistant bacteria	[120]
Prevalence and pathogen load of Eimeria in wild yellow-eyed penguins *(Megadyptes antipodes)* and the morphologic characterization of a novel Eimeria species	[121]
Chlamydiaceae-Like Bacterium in Wild Magellanic Penguins *(Spheniscus magellanicus)*	[122]
Exposure of pelagic seabirds to *Toxoplasma gondii* in the Western Indian Ocean points to an open sea dispersal of this terrestrial parasite	[123]
Prevalence of antimicrobial resistance in bacteria isolated from Great Cormorants *(Phalacrocorax carbo hanedae)* in Japan	[124]
Predator and scavenger movements among and within endangered seabird colonies: Opportunities for pathogen spread	[125]
Molecular identification of avian viruses in neotropic cormorants (*Phalacrocorax brasilianus*) in Chile	[126]
Pathogen transmission risk by opportunistic gulls moving across human landscapes	[127]
Parasite diversity associated with African penguins *(Spheniscus demersus)* and the effect of host and environmental factors	[128]
Low occurrence of hemosporidian parasites in the Neotropic cormorant (*Phalacrocorax brasilianus*) in Chile	[129]

Absence of haemosporidian parasite infections in the long-lived Cory's shearwater: evidence from molecular analyses and review of the literature	[130]
First detection of avian influenza virus (H4N7) in giant petrel monitored by geolocators in the Antarctic region	[131]
Do blood parasites infect Magellanic penguins *(Spheniscus magellanicus)* in the wild? Prospective investigation and climatogeographic considerations	[132]
Avian Influenza Virus H5 Strain with North American and Eurasian Lineage Genes in an Antarctic Penguin	[133]
High prevalence of Leucocytozoon parasites in fresh water breeding gulls	[134]
Regurgitation of the koilin layer in chinstrap penguins (*Pygoscelis antarcticus*) and its association with gastric parasites	[135]
Influenza A virus on oceanic islands: Host and viral diversity in seabirds in the western Indian Ocean	[136]
Egg sampling as a possible alternative to blood sampling when monitoring the exposure of yellow-legged gulls *(Larus michahellis)* to avian influenza viruses	[137]
Serological and virological surveys of the influenza A viruses in Antarctic and sub-Antarctic penguins	[138]
Carriage of bacteria and protozoa in the intestinal tract of common tern chicks	[139]
Prevalence of Hemoproteus iwa in Galapagos Great Frigatebirds *(Fregata minor)* and Their Obligate Fly Ectoparasite *(Olfersia spinifera)*	[140]

Plasmodium blood parasite found in endangered Galapagos penguins *(Spheniscus mendiculus)*	[141]
Infection by Haemoproteus Parasites in Four Species of Frigatebirds and the Description of a New Species of Haemoproteus *(Haemosporida: Haemoproteidae)*	[142]
Comparative analysis reveals a possible immunity-related absence of blood parasites in Common Gulls *(Larus canus)* and Black-headed Gulls *(Chroicocephalus ridibundus)*	[143]
Seabirds and the Circulation of Lyme Borreliosis Bacteria in the North Pacific	[144]
Hemosporidian blood parasites in seabirds—a comparative genetic study of species from Antarctic to tropical habitats	[145]
No Evidence of Blood Parasites in Little Auks *(Alle alle)* Breeding on Svalbard	[146]
Variable exposure and immunological response to Lyme disease Borrelia among North Atlantic seabird species	[147]
Disease-limited distributions? Contrasts in the prevalence of avian malaria in shorebird species using marine and freshwater habitats	[148]
Characterization of a Novel Influenza A Virus Hemagglutinin Subtype (H16) Obtained from black-headed gulls	[149]
Multiple gene segment reassortment between Eurasian and American lineages of influenza A virus (H6N2) in guillemot *(Uria aalge)*	[150]
Infectious bursal disease virus antibodies in eider ducks and herring gulls	[151]

No evidence for high pathogenicity avian influenza in Waved Albatross (*Pvhoebastria irrorata*)	[152]
Outbreaks of highly pathogenic avian influenza (HPAI) epidemics in Baltic Great Cormorant *Phalacrocorax carbo* colonies in 2021 and 2022	[153]
Evaluation of the effect of Pb pollution on avian influenza virus-specific antibody production in black-headed gulls (*Chroicocephalus ridibundus*)	[154]
Seabirds and the circulation of Lyme borreliosis bacteria in the North Pacific	[155]

Table S6. Resources for transmission routes of the pathogens not mentioned in GMPD or CDC

Title	Reference
Shedding and Emission of Airborne Viral Microorganisms from Animal Houses	[156]
Poultry and avian diseases	[157]
Parasites of chinstrap penguins (*Pygoscelis antarctica*) from three localities in the Antarctic Peninsula and a review of their parasitic fauna	[158]
Corynebacterium diphtheriae	[159]
Defining routes of airborne transmission of Pseudomonas aeruginosa in people with cystic fibrosis	[160]
Staphylococcus xylosus and Staphylococcus aureus as commensals and pathogens on murine skin	[161]
Mycoplasma infections	[162]
The transmission and development of Cosmocephalus obvelatus (Nematoda: Acuarioidea) of gulls (Laridae)	[163]
Fishborne zoonotic heterophyid infections: an update	[164]
Transatlantic discovery of Notocotylus atlanticus (Digenea: Notocotylidae) based on life cycle data	[165]
Evolution and historical biogeography of a parasite–host assemblage: Alcataenia spp.(Cyclophyllidea: Dilepididae) in Alcidae (Charadriiformes)	[166]
Assessing the effects of trematode infection on invasive green crabs in eastern North America	[167]
Pathology of proventricular tetrameriasis in a free-range chicken	[168]
Canine and feline cardiopulmonary parasitic nematodes in Europe: emerging and underestimated	[169]
Adenovirus transmission—worthy of our attention	[170]

Animal herpesviruses and their zoonotic potential for cross-species infection	[171]
Avian reticuloendotheliosis in chickens–an update on disease occurrence and clinical course	[172]
Longitudinal map of transcriptome changes in the Lyme pathogen Borrelia burgdorferi during tick-borne transmission	[173]
Chlamydiaceae: diseases in primary hosts and zoonosis	[174]
Discovery of a novel circulation route of free-living Serratiasymbiotica mediated by predatory ladybird beetles	[175]