Single-cell characterization of skin response to a bite by West Nile virus-infected

mosquito reveals fibroblast-mediated barrier to transmission

Hacène Medkour[1*], Federico Bocci[2*], Norman Schneider[1], Idalba Serrato-Pomar[1],

Félix Rey-Cadilhac[1], Yingzi Liu[3], Elliott F. Miot[1], Axel A. Almet[4], Marine Louarn[5],

Nicolas Aubourg[5], Wilfried Saron[6], Ghizlane Maarifi[7,11,12], Dorothée Missé[1], Qing

Nie[3,4], Duncan R. Smith[8], Sébastien Nisole[7,10], Ashley St John[6], Charles Dutertre[5,9],

Maksim V. Plikus[3,#,¶], Julien Pompon[1,8,#,¶]

1 MIVEGEC, University Montpellier, IRD, CNRS, Montpellier, France

2 Radboud institute for molecular life sciences, Radboud University, Nijmegen,

Netherlands

3 Department of Developmental and Cell Biology, University of California Irvine, CA,

USA

4 Department of Mathematics, University of California Irvine, Irvine, CA, USA

5 INSERM U955, Institut Mondor de Recherche Biomedicale, Université Paris-Est

Créteil, Créteil, France

6 Program in Emerging Infectious Diseases, Duke-NUS Medical School, Singapore.

7 Institut de Recherche en Infectiologie de Montpellier (IRIM), Univ Montpellier, CNRS,

INSERM, Montpellier, France

8 Institute of Molecular Biosciences, Mahidol University, Nakhon Pathom, Thailand.

9 Experimental and Translational Pathology Platform, AMMICa, INSERM, Gustave

Roussy, Paris-Saclay University, Villejuif, France

10 Centre Armand-Frappier Santé Biotechnologie, Institut National de la Recherche

Scientifique, Laval, Québec, Canada

11 CIRAD, UMR INTERTRYP, Montpellier, France

12 INTERTRYP, Univ. Montpellier, CIRAD, IRD, Montpellier, France

* Authors contributed equally.

# Authors contributed equally.

¶ Corresponding author: Julien.pompon@ird.fr; plikus@uci.fr

ABSTRACT

Cutaneous events at the mosquito bite site that determine orthoflavivirus transmission efficiency remain largely uncharacterized. Here, we report single-cell RNA-sequencing of skin from immunocompetent mice bitten by West Nile virus-infected mosquitoes, capturing early response at a critical transmission bottleneck. Fibroblasts were the dominant cells exposed to infectious saliva. While neutrophils and *Folr2*+ macrophages diminished, monocyte-derived and antigen-presenting macrophages, and lymphocytes became more abundant. Cell-cell communication analysis revealed the central roles of fibroblasts and myeloid cells in induced immune-related signaling. Transcriptional profiling defined cell type-specific responses to infectious bite integrating immune modulation, skin repair and metabolic remodeling. Using *in vivo* gene silencing, we demonstrated that fibroblast-expressed LRRC15 (leucine rich repeat-containing 15) functions as a cutaneous restriction factor, limiting viral replication in skin, viral dissemination to draining lymph nodes, and disease severity. Collectively, our analyses provide cellular and molecular understanding of bite-initiated arboviral transmission, establishing skin-resident fibroblasts as frontline defender cells.

Key words: Arbovirus, Transmission, skin, cutaneous, Orthoflavivirus, single-cell transcriptomics, mosquito bite, immunity, dermal fibroblasts, LRRC15, antiviral restriction factors, cell-cell communication, mouse model.

HIGHLIGHTS

* Single-cell profiling captures early cutaneous response to West Nile virus-infected mosquito bites with cell type resolution.

* Skin-resident fibroblasts are the dominant frontline cells exposed to infectious mosquito saliva.

* Infectious bite reconfigures cutaneous immune cells, and activates multi-directional signaling between immune and structural skin cell types.

* Bite-induced fibroblast-expressed LRRC15 restricts viral transmission and attenuates disease severity.

INTRODUCTION

Mosquito-borne orthoflaviviruses—including West Nile virus (WNV), dengue virus (DENV) and Zika virus (ZIKV)—cause recurrent outbreaks worldwide, infecting an estimated half a billion people each year and imposing considerable health and economic burden[1,2]. Existing vaccines remain limited and raise safety concerns, complicating both their design and clinical evaluation[3–6]. No antiviral drugs are licensed to-date and therapeutic approaches for acute infection are constrained by an extremely narrow treatment window[7,8]. These challenges underscore the urgent need for novel intervention strategies for mosquito-transmitted infectious diseases.

Bite-initiated skin infection is a critical vulnerability in viral transmission, representing a potential therapeutic target[9,10]. During probing, virus-laden mosquito saliva is injected into the skin, initiating viral transmission through a coordinated molecular and cellular response[11–13]. Saliva and bite-triggered mechanical injury induce local tissue inflammation and increase vascular permeability, which together promote recruitment of neutrophils. These neutrophils release chemoattractants that then drive tissue infiltration by monocytes and the activation or repositioning of resident dendritic cells (DC) and macrophages[14–21]. These infiltrating myeloid cells—themselves being highly susceptible to orthoflavivirus infection—fulfill their defensive immune function as antigen-presenting cells while concurrently carrying viruses to draining lymph nodes (LNs), and as such, enable systemic dissemination and disease transmission[16,17,21–25]. Bite-induced skin edema further facilitates initial virus replication by restricting virus dispersion and maintaining saliva-delivered inoculum[11,17,19]. This sequence of events unfolds with remarkable speed: transcriptional up-regulation of neutrophil-attracting chemokines and neutrophil influx occurs within 4 hours post-bite, myeloid cell infiltration by 6th hour[16,19] and viral spread to LNs within 12-24 hours[11,25–

[27]. Consistently, excision of the bitten skin site at 4 hours or therapeutic activation of

antiviral defenses before 5th hour post-bite abolishes transmission[21,28].

Structural cells of the skin also contribute to these early events. Major structural

skin cells—keratinocytes in the epidermis and ectodermal appendages, as well as

fibroblasts in the dermis[29]—are likely among the first cells to encounter infectious

mosquito saliva and their permissiveness to multiple orthoflaviviruses may influence

disease transmission[20,30–33]. Upon infection, structural cells activate innate immune

responses[32,34–37], including interferon (IFN) pathways, which limit viral replication[33].

Moreover, infected fibroblasts promote DCs' viral resistance and T cell recruitment[31,38],

highlighting the importance of cell-cell communication in the cutaneous response to

infectious bites.

Despite these insights, prevailing mechanistic models of bite-initiated viral

transmission rely largely on indirect and approximate experimental systems—such as

ex vivo virus incubation of skin explants or dissociated skin cells, artificial needle-based

injection of viruses either without or with saliva or with mosquito salivary gland extracts

into immunodeficient mice, uninfected bites followed by viral needle-based inoculation

or uninfected bite alone[25,39–41]. However, the well-documented impact of mosquito bite

on viral transmission kinetics, tissue tropism, cutaneous gene expression and disease

severity[17,42,43] calls for the use of an *in vivo* transmission model to study the early

events of mosquito-borne viral transmission. Here, for the first time we applied single-

cell RNA-sequencing (scRNA-seq) to skin of an immunocompetent mouse model

exposed to infectious mosquito bites to characterize the early molecular and cellular

responses to viral transmission. Our study identified the specific cell types that first

encounter infectious saliva, delineated the rapid reorganization of the cellular

landscape of the skin, mapped induced cell-cell communications and defined cell

subtype-specific transcriptomic responses. Finally, we revealed a fibroblast-mediated

bite-induced antiviral program that restricts viral transmission.

RESULTS

Single-cell characterization of mouse skin bitten by infectious mosquitoes

To conduct informative scRNA-seq evaluation of infectious mosquito-bitten mouse

skin, we first characterized the *in vivo* bite-initiated transmission dynamics in wild-type

(WT) mice. Multiple mosquitoes infected with WNV by intrathoracic inoculation—to

standardize the inoculum—were allowed to bite immunocompetent WT mice (Fig.

S1A). All mosquitoes had infected salivary glands (Fig. S1B,C) and their bites induced

viremia (peaking at 4 days post-bite), clinical signs of infection, prominent weight loss

and mortality in experimental mice (Fig. S1D-G; Dataset S1A,B), thereby validating

viral transmission. To define an optimal time point for the scRNA-seq assay, we next

characterized viral infection kinetics in the bitten skin. We developed a method for

precise identification of the bite site: individually caged mosquitoes (as described in[13])

(Fig. S2A,B) were placed on shaved mouse abdomen and access to the skin was

further directed by applying tape perforated with four 2 mm-diameter openings (Fig.

S2C,D). Visually identified bite sites were marked and collected from 10 minutes to 24

hours post-bite (Fig. S2E). Because the mosquito proboscis penetrates < 2 mm in any

direction into mouse skin[13], we collected 2-mm-diameter skin biopsies and used them

to quantify WNV genomic RNA (gRNA) (Fig. 1A). At 10 minutes post-bite, before virus

amplification[44,45], $\sim 10^4$ copies of gRNA were expectorated into the skin. This gRNA

level remained stable until 3 hours post-bite (hpb), then declined at 6-to-9 hpb—

matching the time of viral genome replication onset[44,45]—before increasing to $\sim 9.6 \times$

10^4 copies at 24 hpb, indicating virus multiplication. To verify dissemination, WNV

gRNA was quantified in draining LNs (Fig. 1B). Viral gRNA was detected as early as

10 minutes post-bite (geometric mean, 345 gRNA copies per LN), remained stable until

9 hpb, and then sharply increased to ~4.1 x 10^4 copies at 24 hpb, validating systemic

infection. Based on these kinetics, we selected 6 hpb for cutaneous scRNA-seq

analysis, a time point corresponding to the bottleneck phase characterized by transient

reduction in cutaneous viral load and the absence of detectable viral expansion in

draining LNs.

To perform scRNA-seq on skin at 6 hpb (Fig. 1C), we collected 2-mm biopsies

centered on the bite sites to enrich for cells directly exposed to infectious saliva and

processed tissues for single-cell isolation and capture. To minimize inter-sample

variability and to reduce animal use, each mouse received between 3 and 5 bites,

yielding multiple skin biopsies per individual. Multiple bites did not alter local cutaneous

viral load at 24 hpb (Fig. S1H) but increased the viral inoculum reaching draining LNs

(Fig. S1I). We optimized skin cell dissociation using enzymatic digestion combined with

needle-assisted mechanical disruption to improve recovery of viable cells, including

dermal DCs (Fig. S3A-C). Skin from non-bitten WT mice (hereafter referred to as

resting skin) was identically processed and served as a control. After sequencing,

genome mapping, quality controls and bioinformatic sample integration (Fig. S4A-E),

we obtained > 6,000 high quality sequenced cells per experimental condition (Fig. 1D).

Unsupervised clustering identified eight cell types (Fig. 1E), each defined by a

distinct transcriptional signature (Fig. 1F; Dataset S2), including established cell type-

specific marker genes (Fig. S5A-H). Fibroblasts were the predominant cell population

(~62.5%), followed by keratinocytes (~15%) and immune cells of the myeloid (~9.4%)

and lymphoid (~7.3%) lineages. Lymphatic (~3.1%) and vascular (~1.6%) endothelial

cells (EC), that delineate lymphatic and blood vessels, respectively, were also

detected, along with vascular smooth muscle cells (SMC) (~0.5%), which surround

larger blood vessels to regulate blood flow. A minor proportion of melanocytes was

also present (~0.4%). These cellular compositions were consistent with prior

cutaneous scRNA-seq datasets[46].

To refine cell annotations, we subclustered the four major cell types based on

subtype-enriched genes. Fibroblasts segregated into eight subtypes (Fig. 1G; Fig.

S6A, B; Dataset S3), listed in the order of decreasing abundance: *Ccl19*+ fibroblasts

(~46% of all fibroblasts), which promote tertiary lymphoid structures by recruiting

lymphocytes and enhancing local adaptive immune response[47,48]; dermal white

adipose tissue (dWAT) fibroblast cells (~28.2%), which reside beneath the dermis and

contribute to wound healing, immune defenses and skin homeostasis[49]; *Pi16*+

fibroblasts (~7.3%), which form perivascular sheaths that regulates vascular stability,

immune cell extravasation and stromal organization through extracellular matrix (ECM)

production and expression of adhesion molecules and chemokines[50–53]; dermal sheath

fibroblasts (~6.4%), which create a contractile niche layer around hair follicles,

anchoring them within the skin and participating in dermal repair[54,55]; *Crabp1*+

fibroblasts (~4.3%), which modulate retinoic acid signaling, regulating ECM and dermal

homeostasis[56,57]; telogen (~3.1%) and anagen (~2.5%) dermal papilla (DP) fibroblasts,

which constitute the principal hair follicle signaling niche cells and reside at its base,

with telogen DP cells maintaining follicle's quiescence, and anagen DP cells driving

active hair growth[58]; and proliferating fibroblasts (~1.1%), which actively divide to

support cutaneous tissue growth, repair and remodeling[59]. Epidermal and ectodermal

appendage cells segregated into nine subtypes (Fig. 1H; Fig. S6C, D; Dataset S3):

germinative layer cells (~28.5% of all keratinocytes)[60], that reside at the base of

growing hair follicles where they actively divide, contributing to the steady-state hair

growth[61]; basal keratinocytes (~14.8%), which form the deepest epidermal layer and

maintain epidermal attachment[62]; suprabasal keratinocytes (~12.6%), which form

upper layers of the epidermis where they differentiate to form water-tight barrier[61];

sebaceous gland cells (~11.4%), which produce lipid-rich secretome to lubricate

epidermis and hairs, reinforce epidermal barrier and modulate local immune and

antibacterial defenses[63]; hair follicle stem cells (HFSCs) (~8.7%), which reside in the

so-called bulge region of the follicle, and critically sustain cyclical hair growth and

contribute to epidermal repair after wounding[61]; upper hair follicle keratinocytes

(~8.3%), which form a protective epithelium at the boundary with the epidermis[61];

cortex/cuticle cells (~8.1%) and medulla cells (~2.1%), which represent differentiated

cell types of the hair shaft; and inner root sheath (IRS) keratinocytes (~6.6%), which

form a protective sleeve around the growing hair shaft[64]. Among myeloid cells (Fig. 1I;

Fig. S6E, F; Dataset S3), we identified multiple populations of macrophages: *Spp1+*

macrophages (~21.6% of myeloid cells), which are associated with sustained

inflammation, extracellular matrix and immune response modulation[65]; *Folr2+*

macrophages (~15.76%), which are perivascular tissue-resident macrophages with the

capacity to activate CD8$^+$ T cells[66]; monocyte-derived macrophages (moMacro;

~12.8%), which arise from recruited circulating monocytes and provide a rapidly

expandable population that drives inflammatory responses and pathogen clearance[67]

and antigen-presenting macrophages (~5.95%) with an activated MHC-II program as

indicated by *Ciita* high expression[68]. We also observed several DC populations: cDC2

(~18.0%) and cDC1 (~5.8%) which activate CD4$^+$ and CD8$^+$ T cells, respectively[69]; and

migratory DCs (~3.0%) with increased *Ccr7* expression[70,71]. Finally in the myeloid

compartment, we identified monocytes (~10.5%), which have functions in

phagocytosis and immune modulation[72]; proliferative myeloids (~3.5%) with an

activated program of cell proliferation (i.e., *Ccna2, Spc24, Ncapg, Ncapd2,* and

Pimreg); and neutrophils (~3.2%), which shape early inflammatory response[73]. Among

lymphoid cells (Fig. 1J; Fig. S6G, H; Dataset S3), subclustering distinguished: a mixed

populations of innate lymphoid cells (ILC), which initiate inflammation through early

cytokine secretion[74,75] , and T cells (~46% of all lymphoid cells); dendritic epidermal T

cells (DETC) (~33.2%), which are skin-resident γδ T cells with roles in immune

surveillance and tissue homeostasis[76]; and quiescent-like ILC (~20.1%), which can

differentiate into active ILCs[77,78]. Altogether, these data establish a robust scRNA-seq

framework to investigate mosquito bite-initiated viral infection in the skin at single-cell

resolution.

Fibroblasts are the frontline infectious saliva-exposed cells

As mosquito RNA is secreted into the saliva[79,80], we leveraged it as a molecular proxy

to identify skin cell types that directly interact with infectious saliva components. Non-

polyadenylated viral reads could not be detected with the scRNAseq technology. Upon

data mapping, we aligned scRNA-seq reads against the mosquito genome and

identified nearly 300 mosquito-derived reads in bitten skin (after quality controls),

whereas none were found in resting control skin. Detected mosquito reads mapped to

six genes, two of which—*HISTONE H4* and an uncharacterized locus

(*LOC23687614*)—accounted for the large majority (Fig. 2A). Mosquito RNA was

detected in 3.9% of all sequenced cells (Fig. 2B), providing an estimate of the limited

subset of skin cells directly exposed to mosquito saliva, even with our narrow skin

collection method. All annotated skin cell types, except melanocytes, contained

mosquito RNA; however, fibroblasts, vascular SMC and keratinocytes exhibited the

highest proportion of positive cells (~ 4%), with fibroblasts being the most exposed in

absolute numbers (Fig. 2C). Moreover, only fibroblasts, keratinocytes and myeloid

cells contained more than one mosquito read per cell (Fig. S7A-H), supporting their

active interaction with mosquito saliva, such as through endocytosis[81]. Intriguingly,

among fibroblasts, the distribution of mosquito RNA did not match subtype abundance:

proliferating (i.e., the least prevalent subtype) and dWAT fibroblasts exhibited the

highest frequencies of mosquito RNA positive cells, followed by *Ccl19*+ (i.e., the most

prevalent subtype) and *Crapb1*+ fibroblasts (Fig. 2D). This suggests preferential

exposure and/or uptake by specific fibroblast subsets.

Next, we characterized transcriptional response to infectious mosquito saliva by

comparing gene expression patterns between mosquito RNA-positive and -negative

fibroblasts, identifying 20 significantly upregulated genes based on expression fold-

change > 1 (Fig. 2E, Dataset S4). Other non-fibroblast cell types contained too few

mosquito RNA-positive cells to support statistically robust analyses. In fibroblasts, four

differentially-expressed genes (DEGs) are associated with immune-related functions:

Fxyd3 enhances Il-17A signalling, promoting cutaneous inflammation[82]; *H2-Eb1*

encodes an MHC-II protein that drives adaptive immune activation[83]; *Dusp22*

modulates T cell activation through the JNK pathway[84]; and *Steap4* responds to

inflammation-induced oxidative stress[85]. Four DEGs are related to metabolic

pathways, inducing steroid biosynthesis (*Dhcr24*[86] and *Fdft1*[87]) and energy-producing

glycolysis (*Acss2*[88] and *Pfkfb3*[89]). Three DEGs are associated with skin repair and

tissue remodeling: *Dmkn* promotes cutaneous barrier formation[90]; *Efnb1* is an

adhesion molecule with an established role in promoting angiogenesis[91]; and *Dmpk*

drives myoblast differentiation[92]. Three DEGs promote cell motility: *Mylk* and *Myo18A*

alter cytoskeleton[93,94], and *Uchl3* influences epithelial-mesenchymal transition[95]. Five

DEGs are involved in general cellular homeostasis: *Lvrn* is implicated in ECM

remodeling[96]; *Tpd52* and *Rbp1* in promoting cell proliferation[97,98]; *Carmn*, a long non-

coding RNA, in regulating cell differentiation[99]; and *Vps37B* in endosomal sorting

through the ESCRT pathway[100]. Collectively, these scRNA-seq analyses provide the

first single-cell-level identification of cutaneous cell types exposed to mosquito saliva

and highlight fibroblasts, a structural cell type, as primary responders during the

earliest stages of vector-host interactions.

Myeloid and lymphoid immune cells are remodeled at the bite site

To characterize cutaneous cellular dynamics in response to infectious bites, we

compared cellular composition of bitten *vs.* resting skin in our scRNA-seq dataset via

permutation testing. At the cell type level, we observed a reduction in melanocyte and

keratinocyte populations, whereas lymphocytes, vascular SMCs, vascular and

lymphatic ECs expanded following infectious bites (Fig. 3A-C). Among fibroblast

subtypes, telogen DP cells were markedly enriched upon infectious bite (Fig. 3D-F).

Among keratinocytes and ectodermal appendage cells, populations of germinative

layer, medulla, cortex and IRS strongly contracted, while sebaceous and suprabasal

keratinocyte populations expanded (Fig. 3G-I). Among myeloid cells, abundances of

Folr2 macrophages and neutrophils were reduced, whereas monocyte-derived and

antigen-presenting macrophage populations were strongly increased (Fig. 3J-L).

Additionally, monocyte and proliferative myeloid compartments were marginally

enhanced from 8.2 to 12.5% and from 2.8 to 4%, respectively. Interestingly, population

sizes of migratory DCs remained unchanged. Among lymphocytes, quiescent-like ILCs

and DETC were strongly recruited to the bite site, increasing by about 4-fold (Fig. 3M-

O). Collectively, while the observed hair follicle cell population differences might reflect

natural hair cycle heterogeneity between animals[101], differences in the other cell types

reveal rapid and extensive multicellular remodeling of the skin in response to an

infectious bite, establishing the early recruitment of multiple types of innate and

adaptive immune cells.

Fibroblasts and myeloid cells orchestrate immune signaling response to

infectious bite

To investigate how an infectious bite reshapes microenvironmental signaling in the

skin, we inferred cell-cell communication (CCC) activity between cell subtypes from

the scRNA-seq dataset using CellChat[102,103]. CellChat database annotates ligands,

receptors, and signaling agonists/antagonists across CCC pathways. First, we

assessed global shifts in CCC of the skin by aggregating all ligand-receptor interactions

significantly present in the data (Fig. 4A). This overview revealed that infectious bite

enhanced communication between fibroblast subtypes, while reducing signaling from

fibroblasts to keratinocytes and between keratinocyte subtypes. Among fibroblasts,

Crabp1+ cells exhibited the strongest activation, both as signal senders and receivers.

Among immune cells, communication across myeloid subtypes was moderately

increased, especially between neutrophils, monocytes and macrophages. Next, we

quantified the activity of individual CCC pathways. Of 62 pathways that were detected

as active in either sample, 24 were upregulated and 11 downregulated by infectious

bite (Fig. 4B). Notably, upregulated pathways were predominantly immune-related (18

out 24), whereas downregulated pathways were mostly associated with skin repair (7

out of 11).

We mapped the activity of regulated pathways to specific cell subtypes (Fig. 4C)

and detailed their contributions as signal senders, receivers, mediators and/or

influencers (Fig. S8). Focusing on immunity, we integrated the upregulated interactions

to delineate global immune CCC patterns (Fig. 4E). Fibroblasts emerged as active

communicators, engaging six CCC pathways toward keratinocytes and myeloid cells.

Toward keratinocytes, most fibroblast subtypes induced pro-inflammatory SEMA5

signaling[104], while dWAT subtypes triggered CADM-mediated immune cell

adhesion[105]. Chemotactic CHEMERIN signaling[106] was broadly promoted in *Folr2*+

macrophages by multiple fibroblast subtypes. In contrast, dWAT and *Crabp1*+

fibroblasts activated pro-inflammatory CADM[107] and hematopoietic-activating FLT3[108]

signaling, respectively, in cDC1. Additionally, dWAT fibroblasts induced

COMPLEMENT signaling in *Spp1*+ macrophages. Keratinocytes coordinated three

CCC pathways, directing signals toward keratinocytes, myeloid cells and NK cells.

Reciprocal SEMA5-mediated inflammatory signaling occurred between keratinocyte

subtypes, while upper hair follicle keratinocytes stimulated MHC-I-dependent antigen-

presentation[109] in DETC. In contrast, suprabasal keratinocytes promoted anti-

inflammatory LXA4 signaling in neutrophils[110] and upper hair follicle induced anti-

inflammatory CD200[111] in cDC2, *Folr2*+ and proliferative macrophages. The infectious

bite provoked the most extensive reorganization of intercellular communication within

the myeloid compartment, activating eight distinct pathways targeting fibroblasts,

myeloid subtypes and lymphocytes. Monocytes induced both immuno-modulatory

TGFβ signaling[112] and immuno-inhibitory SIRP signaling[113] in *Crabp1*+ fibroblasts,

while cooperatively stimulating anti-viral BST2 signaling[114] with *Spp1*+ macrophages

across myeloid subtypes. In return, *Spp1*+ macrophages promoted anti-inflammatory

LAIR1 signaling[115] in monocytes, highlighting layered regulatory mechanisms.

Neutrophils stimulated CSF-dependent phagocytosis[116] in *Folr2*+ macrophages, which

induced phagocytosis-promoting SN signaling[117] across lymphocyte populations.

Conversely, neutrophils and migratory DCs co-activated immuno-modulatory PD-L1

and PD-L2 pathways[118] in quiescent-like ILCs. Taken together, these CellChat

analyses illustrate how infectious bite rapidly reprograms the cutaneous signaling

networks, dampening tissue repair pathways while amplifying pro- and anti-immune-

directed communication across stromal, epithelial and immune compartments.

Cell subtype-specific responses to infectious bite integrate immune, skin repair

and metabolic transcriptional programs

To assess cell subtype-specific responses to infectious bite, we identified DEGs

between infectious and resting conditions within each cell subtype using a conservative

fold-change threshold of ≥ 4, revealing 2,953 DEGs (Fig. 5A; Dataset S5).

Keratinocytes and ectodermal appendages showed the most extensive transcriptional

reprogramming (986 DEGs: 699 up, 287 down), with basal, cortex/cuticle, germinative

layer, and IRS cells exhibiting the largest responses (>162 DEGs each), whereas

HFSCs, medulla, sebaceous gland, and upper hair follicle cells had more modest

responses (<66 DEGs each). Skin repair-associated genes were predominantly

regulated, followed by immune-related genes, which accounted for more than 23.3%

of DEGs in cortex/cuticle, germinative, HFSC, and IRS cells, with metabolic genes also

being significantly modulated (Fig. S9A). Fibroblast subtypes featured fewer overall

DEGs (717 DEGs: 289 up, 428 down) (Fig. 5A; Dataset S5), albeit $Ccl19^+$, $Crabp1^+$,

dermal sheath, dWAT, and $Pi16^+$ fibroblasts each had between 94 and 124 DEGs. Skin

repair-associated genes were most regulated, followed by immune- and metabolic-

related genes (Fig. S9A). Among myeloid cells featuring 745 DEGs (193 up, 552 down)

(Fig. 5A; Dataset S5), moMacrophages were the most responsive (274 DEGs),

followed by antigen-presenting macrophages (148 DEGs), while other subtypes

exhibited a moderate response (<61). Interestingly, while skin repair-associated genes

were also highly regulated across myeloid cell subtypes, immune-related genes were

the most regulated functional category in migratory DCs and present in other subtypes

(Fig. S9A). Metabolic genes were modulated across all myeloid cell subtypes. Among

lymphocytes, DEGs were predominantly downregulated (353 DEGs: 78 up, 275 down)

and fairly homogeneous across DETCs, quiescent-like ILCs and ILC/T cells (125, 122,

and 106 DEGs, respectively) (Fig. 5A; Dataset S5). Here too, skin repair-associated

genes were predominant, followed by immune- and metabolic-related genes (Fig.

S9A). Finally, lymphatic and vascular ECs and SMCs showed moderate responses

(76, 54, and 22 DEGs, respectively) (Fig. 5A; Dataset S5), mainly involving skin repair,

with minor immune- and metabolism-related regulation.

To refine characterization of transcriptional remodeling across cell subtypes, we

performed GO analyses on within-subtype upregulated DEGs using stringent

thresholds (>10% DEGs; $p < 0.01$), presenting results in a decreasing order of cell

abundance. Among fibroblasts (Fig. 5B), $Ccl19^+$ cells showed robust activation of fatty-

acid biosynthesis, accompanied by modulation of macrophage activation and induction

of the pro-angiogenic program. dWAT fibroblasts predominantly upregulated immune-

related processes, including several lymphoid-targeting chemoattractants and

phagocytosis-related genes. $Pi16^+$ and dermal sheath fibroblasts exhibited strong

fatty-acid biosynthetic signatures and moderate chemotactic activity—directed toward

T cells and macrophages in $Pi16^+$ cells, or toward leukocytes in dermal sheath cells—

and both contributed modestly to skin repair programs. In contrast, $Crabp1^+$ fibroblasts

strongly induced skin repair pathways, particularly those linked to angiogenesis and

dermal development, while moderately upregulating prostaglandin biosynthesis genes.

Anagen DP fibroblasts preferentially activated T cell-related pathways, whereas

telogen DP fibroblasts elicited a more modest neutrophil-attracting signature (Fig.

S9B). Proliferating fibroblasts uniquely upregulated endothelial development pathway

genes.

Among keratinocytes and ectodermal appendage cells (Fig. 5C), germinative

cells markedly elicited granulocyte and neutrophil chemoattraction and antimicrobial

humoral responses, while also contributing to ECM-driven and cell migratory

processes associated with skin repair, but with minimal engagement of metabolic

pathways. Basal keratinocytes predominantly activated skin-repair programs, including

wound healing and skin development pathways, while minimally modulating immune

or metabolic functions. Sebaceous gland cells displayed an immune-centric response

focused on B-cell activation, together with the induction of tissue growth associated

pathways. Both cortex/cuticle cells and IRS primarily upregulated skin repair programs,

with moderate contributions to immune chemoattraction (Fig. S9C). HFSCs and

medulla cells each displayed only a single enriched biological process.

Among myeloid cells (Fig. 5D), $Folr2^+$ macrophages exhibited strong activation

of chemoattraction-related GO terms, accompanied by a moderate induction of

myeloid differentiation and skin development. $Spp1^+$ macrophages contributed to T cell

differentiation, chemotaxis and angiogenesis. Monocytes showed enrichment of GO

terms related to myeloid chemoattraction, keratinocyte proliferation and prostaglandin

metabolism. cDC2 had major involvement in sterol-related metabolism, with a modest

implication in B cell activation. Neutrophils uniquely displayed T cell receptor complex

enrichment (Fig. S9D). Among lymphocytes (Fig. 5E), whereas DETC cells lacked

significant GO enrichment, ILC/T cells did not upregulate classical immune-response

pathways but, instead, prominently activated eicosanoid biosynthesis genes.

Quiescent-like ILCs, in contrast, activated chemotactic programs targeting DCs,

neutrophils, and monocytes, and contributed moderately to skin development

processes.

Finally, lymphatic ECs engaged a mix of immune, metabolic, and tissue repair

responses; vascular ECs displayed modest immune-related signature; and vascular

SMCs showed limited immune- and tissue repair-related activation (Fig. S9E-G).

Together, these GO analyzes indicate that infectious mosquito bites trigger a

multifaceted, cell subtype-specific transcriptional response integrating skin repair,

immune activation, and metabolic adaptation across diverse tissue compartments,

revealing specialized functions for distinct cell populations.

Fibroblast response to infectious bite reduces viral transmission

To determine how the above described cutaneous transcriptional responses contribute

to viral transmission, we selected three antiviral DEGs in fibroblasts; *Saa2* primarily

upregulated in Pi16+ fibroblasts and three other subtypes induces anti-viral IFN

response[119], *Defb8* upregulated in DP*, Ccl19*+ and dWAT fibroblasts directly acts on

viruses and *Lrrc15* selectively induced in Crabp1+ fibroblasts has demonstrated

function against viruses[120,121] (Fig. 6A; Fig. S10A-C). We first assessed their regulation

following *in vitro* exposure to virus, saliva (mimicking the bite), or both in the lineage-

matched human cell models: human foreskin fibroblasts (HFF1) (Fig. 6B). Control

experiments confirmed that the dye used for saliva collection[44] and saliva

supplementation alone did not alter cell viability or infection rates (Fig. S10B, C). At 6

hours post-infection, only *LRRC15* was modulated by virus in a manner concordant

with the *in vivo* regulation in mouse skin (Fig. 6C), whereas the remaining genes

displayed inverse patterns (Fig. S10D). Next, we depleted *LRRC15* in HFF1 to

interrogate its function on WNV infection (Fig. 6D). Notably, siRNA-mediated reduction

of LRRC15 mRNA and protein levels (Fig. S10E, F) triggered a strong viral

enhancement (Fig. 6E), identifying LRRC15 as a potent restriction factor.

Next, we developed an *in vivo* cutaneous siRNA-mediated silencing approach

in mice to investigate the effect of *Lrrc15* on transmission (Fig. 6F). Forty-eight hours

after siRNA injection, when cutaneous levels of *Lrrc15* transcript (Fig. 6G) and protein

(Fig. 6H, I) were reduced, skin site was exposed to infectious bites. Local Lrrc15

depletion enhanced viral load in both skin (Fig. 6J) and draining LNs (Fig. 6K) at 24

hpb. Furthermore, while RNAemia at 4 days post-bite confirmed transmission (Fig.

S10G), *Lrrc15* silencing exacerbated weight loss (Fig. 6L), disease severity (Fig. 6M;

Dataset S1) and animal mortality (Fig. 6N). Collectively, these findings demonstrate

that fibroblast-expressed *Lrrc15* functions as a critical antiviral factor restraining early

mosquito-borne infection and underscore the essential contribution of dermal fibroblast

responses to arbovirus transmission.

Discussion

The initial cutaneous response to an infectious mosquito bite is a critical determinant

of arbovirus transmission success[41]. To dissect the cellular and molecular architecture

of this decisive early event, we developed a scRNA-seq protocol targeting a minimal

skin area at the site of WNV-infected mosquito bites in immunocompetent mice,

precisely during the transmission bottleneck phase. Through integrated bioinformatic

analyses, we defined the skin cell composition, identified mosquito saliva-exposed

cells, inferred cell-cell communication networks, and characterized cell type-specific

transcriptional programs, providing a high-resolution map of the earliest stages of

arbovirus transmission. These analyses reveal distinct, subtype-specific cellular roles

and underscore the essential contribution of non-immune, structural skin cells—

particularly fibroblasts—to infection establishment. Using skin-localized gene depletion, we demonstrate that fibroblast-intrinsic responses are key determinants of transmission efficiency and downstream pathogenesis. By combining high-resolution transcriptional profiling with functional perturbation approaches, this study provides an integrated, multiscale view of the cross-cellular response to an infectious mosquito bite, illuminating the molecular and cellular dynamics that govern arbovirus transmission.

The orchestration of myeloid cell response in the skin is a critical driver of arbovirus transmission[41,122]. This response is first initiated by a rapid influx of neutrophils, detectable as early as 3 hours after an uninfectious bite in mice and 4 hours in humans, followed by a decline in neutrophil abundance by 5th hour post-bite in mice[17,19]. Consistent with this kinetic profile, we observed a reduced neutrophil population at 6 hours post-infectious bite, indicative of ongoing resolution of neutrophil-driven inflammation[123,124], further supported by the anti-inflammatory LXA4 signalling[110] toward neutrophils. Rapid attenuation of non-infectious bite-induced inflammation has similarly been described in human skin through the differentiation of anti-inflammatory M2-like macrophages[19,125]. In our study, *Folr2*+ macrophage abundance was reduced following infectious bite, potentially resulting from inflammation resolution[126]. Despite these resolution-associated signatures, inflammation remained active at 6 hours post-infectious bite, as reflected by sustained TNF expression in neutrophils, multiple chemokine expressions in cDC2, *Folr2*+ macrophages and *Spp1*+ macrophages (Dataset S5), enrichment of GO terms associated with chemoattraction, and pro-inflammatory signaling. We also observed a second wave of myeloid cell recruitment[19,41], comprising monocyte-derived, antigen-presenting macrophages, monocytes and proliferative myeloids, although at non-

significant levels for the two latter. While all myeloid populations displayed a mixed

transcriptional response to infectious bite characterized by concurrent up- and down-

regulation of cytokines, coexisting pro- and anti-inflammatory signaling pathways, and

GO terms linked to leukocyte chemoattraction, migratory DCs had a clear signature of

induced anti-viral interferon response (IFIT1, IRF7, ISG15) (Dataset S5). Our results

detail with molecular resolution the dynamic and balanced nature of myeloid responses

at early stages of bite-initiated transmission, delineating a tightly regulated cutaneous

innate immune landscape, in which inflammatory activation and resolution programs

are simultaneously engaged.

Within the lymphoid compartment, we observed a robust influx of DETCs,

skin-resident sentinels that mount early defense responses through rapid release of

inflammatory mediators upon activation [76]. Consistent with a role in viral transmission,

diverse viral infections similarly recruit DETCs, which subsequently orchestrate

myeloid-cell infiltration[127] and modulate early infection dynamics[128]. Infectious

mosquito bite also expanded a population of quiescent-like ILCs, suggesting the

formation of a poised reservoir of tissue-resident precursors capable of rapidly

differentiating into effector ILCs upon exposure to inflammatory or viral cues[77,78].

Alternatively, their accumulation may reflect active immunomodulation by saliva

components, as indicated by the co-induction of anti-inflammatory PD-L1 and PD-L2

pathways[118]. Despite an overall bias toward transcriptional restraint, both DETCs and

quiescent-like ILCs remained responsive to infectious bite, showing bidirectional

regulation of inflammatory chemokines, interleukins, and immunoregulatory pathways.

In contrast, the mixed ILC/T-cell compartment exhibited stronger transcriptional

activation and, along with other lymphocyte subsets, was preferentially targeted by SN,

which promotes phagocytic interactions[117]. Collectively, extensive recruitment and

activation of lymphoid cells at the bite site indicate their potential and previously

underappreciated role in shaping early events that influence viral transmission.

Our untargeted approach details the contribution of structural cells, which have

the capacity to mount an innate immune response[59,129], in actively shaping the broader

cutaneous immune response to an infectious bite. All fibroblast and keratinocyte

subtypes showed enrichment for inflammatory genes and chemoattraction pathways

(Dataset S5), with dWAT fibroblasts and germinative keratinocytes exhibiting a

particularly strong immune-associated transcriptional profile. Structural cells also

displayed direct anti-viral features including defensins[130] in *Ccl19*+ and DP fibroblasts,

and in basal, germinative, IRS, cortex/cuticle keratinocytes; cytotoxic mediators

(granzymes and perforin) in dWAT and *Pi16*+ fibroblasts, and in germinative

keratinocytes; pattern-recognition receptors (C-type lectins) in germinative

keratinocytes; and multiple interferon-inducible genes (Dataset S5). CCC analyses

suggest that fibroblasts and keratinocytes engage with myeloid cells, activating

pathways linked to pathogen control (CADM[107], MHC-I[109], complement[131]), immune

cell proliferation (FLT3[132]), recruitment (chemerin[106], SEMA5[133]), and regulation of

inflammation (LXA4[134], CD200[135]). Together, these data indicate that structural skin

cells adopt immune-active states following infectious bite, contributing to early innate

defense and immune orchestration.

A major finding of our study is the *in vivo* identification of an anti-transmission

response mediated by fibroblasts, revealing fibroblast-expressed LRRC15 as a potent

restricting factor for WNV. Being predominantly expressed in the skin[120,136], *LRRC15*

encodes a membrane protein related to Toll-like receptors[120], a family of pattern-

recognition receptors that initiate innate immune signaling[137,138]. Consistent with this

function, LRRC15 induces a robust antiviral program in fibroblasts and limits infection

by SARS-CoV-2 and adenovirus[120,121], supporting a broad anti-viral function in line with

our findings. Using localized depletion of Lrrc15 in mouse skin, we demonstrate that

Lrrc15 restricts viral transmission and attenuates disease severity. Underscoring its

potential relevance to viral transmission in humans, meta-analysis of single-cell skin

data part of Human Cell Atlas (HCA) project, which consists of 463,503 cells integrated

from 24 published studies, revealed prominent *LRRC15* expression primarily restricted

to 'secretory' and 'mesenchymal' fibroblast types enriched in upper dermis and around

hair follicles, respectively and present across multiple body locations (Fig. S11A-I). A

central unresolved question in vector-borne viral transmission concerns the identity of

the initial target cells following exposure to infectious saliva. Our results demonstrate

that productive infection of fibroblasts is required for transmission and, together with

their pronounced exposure to bite-delivered infectious saliva, support the conclusion

that fibroblasts constitute the initial cellular targets. Although virus is detectable in

draining LNs as early as 10 minutes post-bite, consistent with reported rapid lymphatic

viral dissemination[25,139], restricting local infection in skin fibroblasts markedly reduces

LN seeding and disease outcome, establishing fibroblasts as critical regulators of early

viral spread and transmission.

Limitations of the study

Despite numerous insights that our study offers, it has several limitations. First, our

analysis provides a snapshot of the skin response at 6 hpb, capturing early dynamics

but not the full temporal trajectory of infection, immune activation, and tissue

remodeling. Second, although mice provide a tractable and highly susceptible model

for WNV, inter-species differences in skin architecture and immune composition may

limit direct translation to humans, albeit prominent fibroblast-specific expression of

LRRC15 in skin HCA dataset suggests translatability. Similarly, our study was

conducted with one mosquito vector species and require further experimentation to

expand our findings to other vector species. Finally, as our work centers on WNV, it

remains uncertain whether the identified fibroblast responses, including the restricting

function of LRRC15, represent a generalizable mechanism across arboviruses.

Material and methods

Cells. African green monkey kidney Vero (CCL-81), HFF-1 (SCRC-1041), U937 (CRL-1593.2), and Jurkat (TIB-152) cells were obtained from ATCC. Vero and HFF-1 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Gibco). U937 and Jurkat were maintained in Roswell Park Memorial Institute (RPMI) (Gibco) medium. All media were supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Eurobio), except for HFF-1 which were grown with 15% FBS, and 1% penicillin/streptomycin mix (Invitrogen). Cells were incubated at 37°C with 5% CO_2. All cell lines were tested annually for mycoplasma contamination using specific primers and consistently tested negative.

Viruses. West Nile virus (WNV) strain IS98-ST1 (or Stork 98) was isolated from a stork in Israel in 1998 [140] and obtained from Dr. Philippe Desprès, Centre de Ressources Biologiques, Institut Pasteur, Paris. Virus was propagated in C6/36 cells with 2% FBS, titrated using plaque assay with Vero cells as described previously [141], and stored at -70°C.

Mice. Six- to eight-week-old male C57BL/6J mice were obtained from Charles River (France) and housed in ventilated NexGen Mouse 500 cages (Allentown; Serial number: 1304A0078) within the biosafety level 3 (BSL-3) animal facility at MIVEGEC-IRD, Montpellier, France. They were maintained under a 17 h:7 h light/dark cycle at 53–57% humidity and a temperature of 20–24°C, with *ad libitum* access to sterilized water and an irradiation-sterilized mouse diet (A03, SAFE, France). To allow acclimatization, mice were used for experiments one week after arrival. All efforts were

made to minimize pain and stress. Animal protocols were approved by the APAFiS

national ethical committee (permission numbers: DAP31273, DAP43466).

Mosquitoes. The *Aedes aegypti* BORA colony, originally collected from Bora-Bora

Island in 1980 [142], was maintained at the VectoPole insectary, MIVEGEC. Eggs were

hatched in deionized water, and larvae were reared at 26°C under a 12 h:12 h light-

dark cycle, receiving ground fish food (TetraMin, Tetra) until pupation. Adult

mosquitoes were housed in Bioquip cages at 28°C with 70% relative humidity, a

14h:10h light-dark cycle, and continuous access to a 10% sugar water solution.

Mosquito inoculation. Cold-anesthetized female *Ae. aegypti* (3 to 5 day-old) were

intrathoracically inoculated with 69 nl of RPMI containing 100 particle forming unit

(PFU) of WNV using a Nanoject II (Drummond Scientific Company) and needles made

from 1.14 mm O.D. glass capillaries (1.75″ length, Drummond). Control mosquitoes

were injected with the same volume of RPMI. After inoculation, mosquitoes were

maintained under standard rearing conditions for 9 days before biting.

Biting of mice. Mouse belly was shaved with a hair trimmer (VITIVA MINI, BIOSEB)

one day before challenge. On the day of mosquito bite challenge, mice were

anesthetized by intraperitoneal (IP) injection of 0.2 ml/mouse of ketamine (10 mg/ml,

Imalgène 1000, Boehringer Ingelheim Animal Health) and xylazine (1 mg/ml, Rompun

2%, Elanco GmbH). A tape perforated with 2 mm openings was applied on trimmed

skin to control mosquito bite location. 16h-starved mosquitoes were placed in small

3D-printed cages (Fig. S2) covered with mosquito net to allow biting. Mice were

exposed to one to five encaged WNV-infected mosquitoes. Mosquito blood feeding

was visually recorded and confirmed by the presence of blood in this abdomen. Only

blood-fed mosquitoes were considered to confirm biting. Bite sites identified thanks to

the openings in tape were marked with a permanent pen (Fig. S2).

Tissue collection. 2 mm skin biopsies were collected from the bite sites at 10 min, 1

h, 3 h, 6 h, 9 h, and 24 h post-bite using biopsy punches (Kai Medical) on mice

anesthetized by IP injection of 0.2 ml/mouse of ketamine and xylazine. On the same

mice, inguinal lymph nodes (LNs) and blood were collected.

Absolute quantification of WNV (+) gRNA. Total RNA was extracted from skin

biopsies, LNs, mosquitoes, SG or cells using EZNA RNA extraction kit I (OMEGA) or

from blood samples using QIAamp Viral RNA Mini Kit (Qiagen). Absolute quantification

of positive strand gRNA was performed through one-step RT-qPCR. Total reaction

volume was 10 μl and contained 5 μl of iTaq Universal SYBR green one-step kit (Bio-

Rad), 300 nM of forward and reverse primers [44] (Table S5) and 2 μl of RNA extract.

The reaction was performed in AriaMx Real-time PCR System (Agilent) with the

following thermal profile: 50°C for 10 min, 95°C for 1 min and 40 cycles of 95°C for 10

sec and 60°C for 25 sec, followed by a melting curve analysis. An absolute standard

curve for DENV and WNV gRNA was generated by amplifying the qPCR target using

primers detailed in Table S5 as previously performed [143,144]. Using the gRNA

templates, the limit of detection (LoD) at 95% was determined by calculating fractions

of detected samples in six replicates of serial dilutions [145].

Mouse symptom evaluation. After mosquito biting, mice were weighed daily to

calculate the percent of weight loss. Daily clinical examinations were conducted and a

clinical score (CS) ranging from 0 to 5 was assigned to each mouse following criteria

from a previous study [146] where CS of 0 was assigned to healthy mice; CS of 1 for

mice with ruffled fur, lethargy, hunched posture, no paresis, normal gait; CS of 2 for

mice with altered gait, limited movement in 1 hind limb; CS of 3 for lack of movement,

paralysis in 1 or both hind limbs, and CS of 4 for moribund mice. A CS of 5 indicated

mortality (Table S1). Mice were euthanized if they displayed neurological symptoms,

severe distress, or weight loss exceeding 20%. At day 15, mice were euthanized under

anesthesia.

Skin cell dissociation. Skin biopsies were minced with a scissor and incubated at

37°C for 60 min in 1 ml dissociation buffer containing 5 ml RPMI 1640 without Ca^{2+} or

Mg^{2+} (No FBS or EDTA), 23.2 µl DNAse 1 (10U/µL) (Sigma-Aldrich), 50 µl Liberase

(25mg/mL) (Sigma-Aldrich), 116 µl of 1M HEPES (Gibco), 116 µl of 100 mM sodium

pyruvate (Gibco), 0.0025 g Hyaluronidase (Worthington), 500 µl of Dispase:

Collagenase (1mg/ml) (Sigma-Aldrich). The dissociation was stopped by adding 10 µl

of 0.5 M EDTA (Invitrogen) and 400 µl of FBS. Cell aggregates were mechanically

dissociated by (i) vortexing for 30 sec, (ii) passing through 19G needle (TERUMO)

several times (~10 times), or (iii) using GentleMACS (MACS) dissociator with

m_imptumor_01 program. Cells were filtered through 70 µm (PluriSelect) and 40 µm

cell strainers before centrifugation at 500 g for 10 min at 4°C (low break speed) and

resuspension in appropriate volume of PBS for flow cytometry or 0.04% BSA-RPMI for

scRNA-seq.

Flow cytometry. Three pools of 25 2-mm skin biopsies were collected from the

trimmed abdomen of unbitten mice. After cell dissociation, cell surface staining of skin

cells was performed in PBS buffer containing 1% FCS for 30 min at 4°C with the

following antibodies: MHC Class II (I-A/I-E), monoclonal Antibody APC-eFluor780

(eBioscience), PE/Cyanine7 anti-mouse CD103 Antibody (Biolegend), APC anti-

mouse/human CD207 (Langerin) Antibody (Biolegend). Cells were subsequently

washed with PBS, pelleted by centrifugation at 500 x g for 5 min at 4°C, resuspended

in PBS and analyzed on an LSR Fortessa (BD Biosciences) flow cytometer. Results

were analyzed with FlowJo (BD Biosciences). Antigen-presenting cells (MHC-II+)

present in skin cell suspensions were analyzed for CD103 and CD207 expression to

evaluate the proportion of Langerhans cells (CD207+) and dermal dendritic cells

(CD103+).

5′-end single cell RNA-sequencing. 2-mm skin biopsies corresponding to the areas

bitten by WNV-infected mosquitoes were collected at 6 h post biting. Each mouse was

bitten by three to five mosquitoes. Twenty-two biopsies collected from eight animals

were pooled. The same number of skin biopsies were collected from unbitten mice that

underwent the same procedures as control (resting skin). Skin biopsies were cell

dissociated using enzymatic digestion and needle-assisted mechanical dissociation

and pelleted in 0.04% UltraPure BSA solution (Thermofisher Scientific). Dead cells

were removed using the Dead Cell Removal Kit (Miltenyi Biotec) with MS columns.

Live cells were resuspended in 2% UltraPure BSA, counted using Luna automated cell

counter (Logos), and concentrated to ~1,000 cell/µl. GEM generation, barcoding, post

GEM-RT cleanup, cDNA amplification, and cDNA library construction were performed

using the Chromium Single-Cell 5′ Reagent version 2 kit (10x Genomics). Libraries

were sequenced on an Illumina HiSeq4000 platform (Illumina). Mouse challenge, cell

preparation, GEM generation were performed at the BSL3 Vectopole of MIVEGEC,

IRD, Montpellier; barcoding, post GEM-RT cleanup, cDNA amplification, library

preparation in molecular biology laboratories at MIVEGEC; quality control and

sequencing at the Genomics Research and Technology Hub of UCI. Multiple repeats

were conducted and two repeats for infectious bite and one for resting skin were

successful.

Data processing and quality control for 5′-end transcripts. Raw counts were

aligned against the *Mus musculus* (mm39), *Ae. aegypti* (AaegL5.0) [147], and West Nile

Virus [148] genomes using 10X Genomics Cell Ranger v7.1.0. Data was preprocessed

and integrated with Seurat [149]. First, cells were removed if total RNA count was low

(nCount_RNA < 500), number of expressed features was too low or too high

(nFeature_RNA < 500 and nFeature_RNA > 6000), and mitochondrial gene count was

high (percent.mt > 10). Second, the samples were normalized, clustered, and projected

onto low-dimensional embedding (UMAP) following the standard Seurat workflow with

default parameters. The first 20 PCA dimensions were used for computation of

neighbors and low-dimensional embedding. Third, the samples (1 resting, 2 infectious

bite) were integrated by selecting integration features (*SelectIntegrationFeatures()*)

and anchors (*FindIntegrationAnchors()*), resulting in the integrated dataset

(*IntegrateData()*). Finally, the integrated data was rescaled and UMAP representation

was computed again. The two infectious bite samples were merged into a single

sample for downstream analysis.

Clustering, annotation, and proportion testing of 5'end transcripts. Cell types

were identified by computing cell neighbors using the top 15 PCA dimensions, followed

by clustering with resolution = 0.02, resulting in n = 8 clusters. This resolution threshold

was identified by trial-and-error and based on the clear separation between the cell

types in the UMAP space. Clusters were annotated based on the expression of marker

genes identified with the *FindAllMarkers()* Seurat function with parameters (only.pos =

TRUE, min.pct = 0.8, logfc.threshold = 0.5) based on Wilcoxon rank sum test and

adjusted p-values with Bonferroni correction. Based on annotation at cell type level,

fibroblasts, keratinocytes, lymphoid cells and myeloid cells were selected for individual

sub-clustering. For each cell type, multiple clustering resolution parameters were

tested to identify cell subtypes, and the final parameter value was set based on the

interpretability of the resulting clusters (fibroblasts: 0.2; lymphoid: 0.8; myeloid: 0.9;

keratinocytes: 0.3), whereby gene markers for each subtype were identified following

the same procedure explained above. Proportion tests for cell types and subtypes

between Resting and Infectious bite were performed using the scProportionTest R

package [150] by applying a permutation test that associates a p-value to the cell fraction

fold-change [150].

Identification of mosquito RNA-containing cells. Based on alignment against the

Ae. aegypti genome (AaegL5.0) [147], cells were categorized as mosquito-positive or

mosquito-negative based on detection of one or more mosquito RNA reads, thus

enabling the quantification of mosquito-positive cells per cell type and per subtype (in

the case of fibroblasts). Further classification based on the number of mosquito RNA

reads was not conducted as the majority of cells expressed either none or a single

mosquito RNA reads.

Identification of differentially expressed genes (DEGs). DEGs between mosquito

RNA-positive and -negative cells within fibroblasts, and between resting and infectious

samples within cell subtypes were identified with the *findmarkers()* function from Seurat. Given the sparsity of several genes with high fold-change in the infectious bite, we implemented the MAST statistical testing that specifically considers zero-inflation in scRNA-seq data [151] by setting *test.use="MAST"*. This statistical test was performed for all main cell types as well as on the subtypes for fibroblasts, keratinocytes, lymphoid cells and myeloid cells. DEGs were defined based on adjusted p value < 0.001 and a |log2 fold-change| > 2. DEGs were functionally categorized into immunity (including inflammation, innate immunity, adaptive immunity), skin regeneration (including cell motility, cell proliferation, angiogenesis, tissue development, ECM), metabolism, RTT (replication, transcription and translation), Redox–stress–mitochondria, non-coding RNA, or divers (for genes not related to the above functions or with unknown functions) based on GO biological processes retrieved from Ensembl database (ensembl.org/biomart/martview/).

Gene Ontology (GO) enrichment analysis. GO enrichment analysis was performed on the within cell subtypes upregulated DEGs with the gseapy package [152] using the 'GO_Biological_Process_2025' gene sets. Significant biological processes were filtered based on GO term presence in > 10% of within-subtype DEGs and p-value < 0.01 (i.e., log10 (1/FDR) > 2). Combined score represents the product of the p-value and the z-score of the deviation from expected rank, providing a comprehensive measure of enrichment strength.

Cell-cell communication analyses. The data was analyzed using the CellChat package[153]. First, the individual samples (Rest, Infectious bite) were converted into CellChat data objects following the standard procedure explained in CellChat tutorial.

Cell-cell communication analysis was also performed using CellChat's default setting,

except for setting a minimum number of cells min.cells = 5 per cell type to filter out

weak cell-cell communication pathways (i.e., the pathway must be detected in at least

5 cells per cell type). Then, the samples were integrated, and the cell-cell

communication strength and pathways were identified and plotted using the

netVisual_heatmap and Ranknet functions as well as custom script.

Collection of mosquito saliva. Bulk saliva collection was based on previously

described protocols [154,155]. Uninfected female mosquitoes were let to feed on a

Hemotek feeding system (Hemotek) containing 3 ml of RPMI and 1% 25 mM

Erioglaucine (SigmaAldrich). The number of mosquitoes that salivated was estimated

by visually counting individuals with blue abdomens, indicative of feeding and

salivation. Salivation media was collected after 1h.

Evaluation of cell viability. 10^5 HFF-1, U-937 or Jurkat were supplemented or not

with 1% Erioglaucine. After 6h, cell viability was estimated using CyQUANT NF Cell

Proliferation Assay Kit (Invitrogen).

Cell infection. 10^5 HFF-1, U-937 or Jurkat cells were seeded in 48-well plates. 24h

later, cells were treated with WNV at MOI = 1, with or without the addition of saliva

collected from 5 mosquitoes (4.5 µl of solution corresponding to saliva from one

mosquito), or with mosquito saliva alone. Control cells received the same volume of

saliva collection medium. At 6 h post treatments, cells were collected in TRK lysis

buffer.

Relative gene and WNV gRNA quantification. Total RNA from cells was extracted using EZNA Total RNA kit I. Relative gene expression was quantified with one-step RT-qPCR using iTaq Universal SYBR green one-step kit (Bio-Rad) with the corresponding primers (Table S1). Amplification was conducted in AriaMx using the same thermal profile as for WNV (+) gRNA quantification. Expression of three housekeeping genes (i.e., *RPL13A, ACTB,* and *GAPDH*) was used for normalization (Table S5). Relative expression was calculated using the 2-ΔΔCt method.

siRNA silencing. 10^5 U-937 or HFF-1 cells were transfected with 12 pmol of multiplex flexitube siRNAs (Qiagen) using Lipofectamine RNAiMax (ThermoFisherScientific). siRNA GS131578 against *Lrrc15* was used in HFF1 cells; siRNA GS10437 against *ifi30* and siRNA GS3123 against *Hla-drb1* were used in U-937 cells (Table S1). Allstars multiplex siRNAs (Qiagen) was transfected as control. 48 h after transfection, cells were infected with WNV as described above. At 36 hpi, cells were collected in TRK lysis buffer for RNA quantification or in RIPA lysis buffers (ThermoFisherScientific) protein analysis.

siRNA delivery in skin. One day prior siRNA injection, mouse hairs on the lower back were shaved with the hair trimmer (VITIVA MINI, BIOSEB). Anesthetized mice were intradermally inoculated, using CellTram 4r Oil microinjector (Calibre Scientific), with 30 pmol in 3 µl of siRNA GS74488 against *Lrrc15* or Allstars multiplex siRNAs (Qiagen) mixed with 5 µl of Lipofectamine RNAiMax, completed to 10 µl with water. Each mouse was injected at three different sites around the planned biting site. The inoculation sites were identified by a permanent pen.

Western blot on cells and skin. 48 h post siRNA transfection, cells and skin were

collected. Cells were washed twice with PBS, scrapped in 70 μl of RIPA containing 1X

protease inhibitors (Sigma-Aldrich) and the supernatant was collected after

centrifugation at 12,000 g for 1 min. For each mouse, ~5 mm diameter biopsies were

collected from the three siRNA injected-bitten skin in 200 μl 1X RIPA lysis buffer, micro-

dissected with scissors, homogenized using Fast-prep bead bitter (MP) with glass

beads, and supernatant was collected after centrifugation at 12,000 g for 1 min.

Normalized protein quantities were separated under denaturing conditions in 10%

polyacrylamide gel and transferred onto 0.2 μm Nitrocellulose membrane using

TransBlot system (BioRad). Staining was conducted with 1:1,000 and 1:500 rabbit anti-

LRRC15 (50546, Cell Signaling) for human cells and mouse skin, respectively, and

1:400 anti-Actin (MA5-11869, Invitrogen) in 0.1 % Tween-20 1% BSA in PBS.

Secondary staining with 1:2,000 of goat anti-rabbit (7074P2, Cell Signaling) and goat

anti-mouse (7076P2, Cell Signaling) was similarly performed. Pictures were taken with

ChemiDoc (BioRad) using SuperSignal West Pico Chemiluminescent substrate

(ThermoFisherScientific). Actin expression was used for normalization.

Quantification of siRNA silencing effect on skin infection and symptoms. 48 h

post siRNA injection, mice were bitten at the siRNA injection sites by one WNV-infected

mosquito. Each mouse was bitten by three mosquitoes at three different sites

previously injected with siRNA. At 24h post-bite, skin biopsies from the bite sites and

inguinal LNs were collected and homogenized with Fast-prep bead bitter (MP) with

glass beads in 350 μl of TRK lysis buffer (EZNA). After RNA extraction with EZNA Total

RNA extraction kit I, WNV gRNA was quantified. Additional mice were monitored for

symptoms, weight loss and survival. At 4 days post bite, blood samples were collected

via mandibular puncture and sample volumes were estimated by pipetting before RNA

extraction using EZNA total RNA extraction kit I and WNV gRNA quantification. Mice

were euthanized if they displayed neurological symptoms, severe distress, or weight

loss exceeding 20%. Mice were euthanized under anesthesia at day 14.

Skin Human Cell Atlas (HCA) data meta-analysis. We re-analyzed the skin

component of the HCA, which is an integrated single-cell dataset from 24 independent

single-cell datasets of normal, non-diseased human skin sampled across different

ages, sexes and genders, and anatomical body sites. Skin HCA contains two levels of

cell annotations. The first level of annotations identifies all skin-resident cell lineages.

The second level identifies lineage-specific subclusters. We analyzed *LRRC15*

expression across all lineages, before restricting analysis to fibroblast subclusters.

Before analyzing *LRRC15* expression in fibroblasts across anatomical regions, we

removed cells sampled from anatomical regions for which there were fewer than 500

fibroblasts, and cells from anatomical regions of unknown origin. This step of filtering

was to ensure that observations of low *LRRC15* expression was a true biological

signal, rather than due to low fibroblast numbers.

Spatial mapping of skin HCA to Xenium data. Prior to spatial projection analysis,

we optimized SpaceRanger-generated cell segmentation of the raw spatial Xenium

data using ProSeg [156]. We then performed quality control of individual samples using

Spotsweeper-py [157], removing cells with excessively low total counts, excessively low

numbers of unique transcripts, and an excessively high proportion of gene transcription

due to mitochondrial genes. We then normalized cell counts by median library size

before log-transforming the normalized counts (adding a pseudocount of 1). When

analyzing the Scalp sample, we used ResolVI [158] to cluster the data initially and remove

adipocytes, as this cell lineage is absent from single-cell skin HCA dataset.

We used cell2location [159] to project HCA fibroblast populations to their likely

spatial location in Xenium spatial transcriptomics samples. We assumed that each

Xenium cell mapped exactly to one cell type. As cell2location was originally developed

for the first generation 10X Visium spatial sequencing platform, which contains

multicellular spots, cell2location output estimates the expected "number" of cells from

each cell annotation label at each spot. Therefore, we annotated Xenium cells as the

cell type annotation with the highest number of cells predicted by cell2location. Briefly,

we performed two levels of spatial projection using the HCA and the Xenium samples.

First, we trained cell2location using the global cell lineage annotations to predict which

Xenium cells were likely fibroblasts. Then, we trained a second cell2location model

using only HCA fibroblast subcluster annotations and projected these annotations onto

the identified Xenium fibroblast cells.

Statistical analysis. Differences in log-transformed gRNA copies and relative gene

expression were tested with one-tailed T-test or post-hoc Dunnett's test. Differences

in survival were tested with Kaplan-Meier survival analysis with Gehan-Breslow-

Wilcoxon test. The statistical analyses were performed using Prism 8.0.2 (GraphPad).

Clinical scores and weight loss were analyzed using generalized linear mixed-effects

models (GLMMs). Clinical scores were modeled assuming a Poisson error distribution

with a log link function. Weight loss were modeled using an inverse Gaussian error

distribution with a log link. Both models included dpi, condition (siNeg vs. siLRRC15),

and their interaction as fixed effects, with mouse specified as a random intercept.

Analyses were performed in R (version 4.3.3) using the glmmTMB package (version

1.1.9). Model adequacy was evaluated using simulation-based diagnostics

implemented in the DHARMa package (version 0.4.7). No evidence of overdispersion

was detected for either the clinical score model (dispersion = 0.579, p = 0.496) or the

weight loss model (dispersion = 1.038, p = 0.728).

Data and Code Availability

scRNA-seq data have been deposited at https://zenodo.org/records/17953412 and are

publicly available as of the date of publication. Any additional information required to

reanalyze the data reported in this paper is available from the Lead Contact upon

request.

Acknowledgments

We thank all members of the Pompon's Laboratory at MIVEGEC, IRD, for their

valuable suggestions and support. We are especially grateful to VectoPole team in

Montpellier, particularly Bethsabée Scheid and Carole Ginibre, for providing mosquito

eggs, and to Nathalie Barougier, Pascal Bautinaud and Drs. Sylvie Cornelie and Idris

Mhaidi for their assistance with mouse management. We thank Dany Sevrac and the

MGX-Montpellier GenomiX platfrom, part of the French National Infrastructure at the

Montpellier Biocampus, for providing the Chromium system and technical support. We

acknowledge the imaging facility MRI, member of the national infrastructure France-

BioImaging (https://ror. org/01y7vt929) supported by the French National Research

Agency (ANR-24-INBS-0005 FBI BIOGEN). Finally, we are grateful to Melanie Oakes

and the Genomics Research and Technology Hub at the University of California, Irvine

(UCI), USA, for sequencing support.

Funding

Support for this research came from fellowships from the Fondation pour la Recherche

Médicale to H.M. (SPF202110013925) and to E.F.M. (ARF202309017577), Risques

Infectieux et Vecteurs en Occitanie (RIVOC) and Key Initiatives MUSE Risques

Infectieux et Vecteurs (KIM RIV) grants to H.M. and J.P., French Agence Nationale

pour la Recherche (ANR-20-CE15-0006 to J.P. and ANR-21-CE15-0041 to S.N.), EU

HORIZON-HLTH-2023-DISEASE-03-18 (#101137006) to J.P. and M.V.P., PHC Siam

grant (#49487UJ) from the French Embassy in Thailand to J.P. and clinical fellowship

from CIRM training grant (EDUC4-12822) to Y.L.

REFERENCES

1. Pierson, T.C., and Diamond, M.S. (2020). The continued threat of emerging flaviviruses. Nat. Microbiol. *5*, 796–812. https://doi.org/10.1038/s41564-020-0714-0.

2. Madere, F.S., Andrade da Silva, A.V., Okeze, E., Tilley, E., Grinev, A., Konduru, K., García, M., and Rios, M. (2025). Flavivirus infections and diagnostic challenges for dengue, West Nile and Zika Viruses. Npj Viruses *3*, 36. https://doi.org/10.1038/s44298-025-00114-z.

3. Santos-Peral, A., Luppa, F., Goresch, S., Nikolova, E., Zaucha, M., Lehmann, L., Dahlstroem, F., Karimzadeh, H., Thorn-Seshold, J., Winheim, E., et al. (2024). Prior flavivirus immunity skews the yellow fever vaccine response to cross-reactive antibodies with potential to enhance dengue virus infection. Nat. Commun. *15*, 1696. https://doi.org/10.1038/s41467-024-45806-x.

4. Halstead, S.B. (2021). Is Dengue Vaccine Protection Possible? Clin. Infect. Dis., ciab282. https://doi.org/10.1093/cid/ciab282.

5. Ferguson, N.M., Rodríguez-Barraquer, I., Dorigatti, I., Mier-y-Teran-Romero, L., Laydon, D.J., and Cummings, D.A.T. (2016). Benefits and risks of the Sanofi-Pasteur dengue vaccine: Modeling optimal deployment. Science *353*, 1033–1036. https://doi.org/10.1126/science.aaf9590.

6. WHO position paper on dengue vaccines, May 2024 https://www.who.int/publications/i/item/who-wer-9918-203-224.

7. Vegvari, C., Hadjichrysanthou, C., Cauët, E., Lawrence, E., Cori, A., Wolf, F. de, and Anderson, R.M. (2016). How Can Viral Dynamics Models Inform Endpoint Measures in Clinical Trials of Therapies for Acute Viral Infections? PLOS ONE *11*, e0158237. https://doi.org/10.1371/journal.pone.0158237.

8.   Obi, J.O., Gutiérrez-Barbosa, H., Chua, J.V., and Deredge, D.J. (2021). Current

Trends and Limitations in Dengue Antiviral Research. Trop. Med. Infect. Dis. *6*, 180.

https://doi.org/10.3390/tropicalmed6040180.

9.   Manning, J.E., Morens, D.M., Kamhawi, S., Valenzuela, J.G., and Memoli, M.

(2018). Mosquito Saliva: The Hope for a Universal Arbovirus Vaccine? J. Infect. Dis.

218, 7–15. https://doi.org/10.1093/infdis/jiy179.

10.   Marín-López, A., Raduwan, H., Chen, T.-Y., Utrilla-Trigo, S., Wolfhard, D.P.,

and Fikrig, E. (2023). Mosquito Salivary Proteins and Arbovirus Infection: From Viral

Enhancers to Potential Targets for Vaccines. Pathog. Basel Switz. *12*, 371.

https://doi.org/10.3390/pathogens12030371.

11.   Styer, L.M., Kent, K.A., Albright, R.G., Bennett, C.J., Kramer, L.D., and Bernard,

K.A. (2007). Mosquitoes Inoculate High Doses of West Nile Virus as They Probe and

Feed   on   Live   Hosts.   PLOS   Pathog.   *3*,   e132.

https://doi.org/10.1371/journal.ppat.0030132.

12.   Choumet, V., Attout, T., Chartier, L., Khun, H., Sautereau, J., Robbe-Vincent,

A., Brey, P., Huerre, M., and Bain, O. (2012). Visualizing Non Infectious and Infectious

Anopheles gambiae Blood Feedings in Naive and Saliva-Immunized Mice. PLOS ONE

7, e50464. https://doi.org/10.1371/journal.pone.0050464.

13.   Wei Xiang, B.W., Saron, W.A.A., Stewart, J.C., Hain, A., Walvekar, V., Missé,

D., Thomas, F., Kini, R.M., Roche, B., Claridge-Chang, A., et al. (2022). Dengue virus

infection modifies mosquito blood-feeding behavior to increase transmission to the

host.   Proc.   Natl.   Acad.   Sci.   U.   S.   A.   *119*,   e2117589119.

https://doi.org/10.1073/pnas.2117589119.

14.   Lefteri, D.A., Bryden, S.R., Pingen, M., Terry, S., McCafferty, A., Beswick, E.F.,

Georgiev, G., Van der Laan, M., Mastrullo, V., Campagnolo, P., et al. (2022).

Mosquito saliva enhances virus infection through sialokinin-dependent vascular

leakage. Proc. Natl. Acad. Sci. *119*, e2114309119.

https://doi.org/10.1073/pnas.2114309119.

15. Hastings, A.K., Uraki, R., Gaitsch, H., Dhaliwal, K., Stanley, S., Sproch, H.,

Williamson, E., MacNeil, T., Marin-Lopez, A., Hwang, J., et al. (2019). Aedes aegypti

NeSt1 Protein Enhances Zika Virus Pathogenesis by Activating Neutrophils. J. Virol.

93, e00395-19. https://doi.org/10.1128/JVI.00395-19.

16. Wang, Z., Nie, K., Liang, Y., Niu, J., Yu, X., Zhang, O., Liu, L., Shi, X., Wang,

Y., Feng, X., et al. (2024). A mosquito salivary protein-driven influx of myeloid cells

facilitates flavivirus transmission. EMBO J. *43*, 1690–1721.

https://doi.org/10.1038/s44318-024-00056-x.

17. Pingen, M., Bryden, S.R., Pondeville, E., Schnettler, E., Kohl, A., Merits, A.,

Fazakerley, J.K., Graham, G.J., and McKimmie, C.S. (2016). Host Inflammatory

Response to Mosquito Bites Enhances the Severity of Arbovirus Infection. Immunity

44, 1455–1469. https://doi.org/10.1016/j.immuni.2016.06.002.

18. Uraki, R., Hastings, A.K., Marin-Lopez, A., Sumida, T., Takahashi, T., Grover,

J.R., Iwasaki, A., Hafler, D., Montgomery, R.R., and Fikrig, E. (2019). Aedes aegypti

AgBR1 antibodies modulate early Zika virus infection of mice. Nat. Microbiol. *4*, 948–

955. https://doi.org/10.1038/s41564-019-0385-x.

19. Guerrero, D., Vo, H.T.M., Lon, C., Bohl, J.A., Nhik, S., Chea, S., Man, S., Sreng,

S., Pacheco, A.R., Ly, S., et al. (2022). Evaluation of cutaneous immune response in

a controlled human in vivo model of mosquito bites. Nat. Commun. *13*, 7036.

https://doi.org/10.1038/s41467-022-34534-9.

20. Duangkhae, P., Erdos, G., Ryman, K.D., Watkins, S.C., Falo, L.D., Marques,

E.T.A., and Barratt-Boyes, S.M. (2018). Interplay between Keratinocytes and Myeloid

Cells Drives Dengue Virus Spread in Human Skin. J. Invest. Dermatol. *138*, 618–626.

https://doi.org/10.1016/j.jid.2017.10.018.

21.   Schmid, M.A., Glasner, D.R., Shah, S., Michlmayr, D., Kramer, L.D., and Harris,

E. (2016). Mosquito Saliva Increases Endothelial Permeability in the Skin, Immune Cell

Migration, and Dengue Pathogenesis during Antibody-Dependent Enhancement.

PLoS Pathog. *12*, e1005676. https://doi.org/10.1371/journal.ppat.1005676.

22.   Wu, S.-J.L., Grouard-Vogel, G., Sun, W., Mascola, J.R., Brachtel, E., Putvatana,

R., Louder, M.K., Filgueira, L., Marovich, M.A., Wong, H.K., et al. (2000). Human skin

Langerhans cells are targets of dengue virus infection. Nat. Med. *6*, 816–820.

https://doi.org/10.1038/77553.

23.   Cerny, D., Haniffa, M., Shin, A., Bigliardi, P., Tan, B.K., Lee, B., Poidinger, M.,

Tan, E.Y., Ginhoux, F., and Fink, K. (2014). Selective Susceptibility of Human Skin

Antigen Presenting Cells to Productive Dengue Virus Infection. PLOS Pathog. *10*,

e1004548. https://doi.org/10.1371/journal.ppat.1004548.

24.   Martin, M.-F., Maarifi, G., Abiven, H., Seffals, M., Mouchet, N., Beck, C., Bodet,

C., Lévêque, N., Arhel, N.J., Blanchet, F.P., et al. (2022). Usutu Virus escapes

langerin-induced restriction to productively infect human Langerhans cells, unlike West

Nile   virus.   Emerg.   Microbes   Infect.   *11*,   761–774.

https://doi.org/10.1080/22221751.2022.2045875.

25.   Reynoso, G.V., Gordon, D.N., Kalia, A., Aguilar, C.C., Malo, C.S., Aleshnick,

M., Dowd, K.A., Cherry, C.R., Shannon, J.P., Vrba, S.M., et al. (2023). Zika virus

spreads   through   infection   of   lymph   node-resident   macrophages.   Cell   Rep.   *42*.

https://doi.org/10.1016/j.celrep.2023.112126.

26.   Kyle, J.L., Beatty, P.R., and Harris, E. (2007). Dengue Virus Infects

Macrophages and Dendritic Cells in a Mouse Model of Infection. J. Infect. Dis. *195*,

1808–1817. https://doi.org/10.1086/518007.

27.   Lim, P.-Y., Behr, M.J., Chadwick, C.M., Shi, P.-Y., and Bernard, K.A. (2011).

Keratinocytes Are Cell Targets of West Nile Virus In Vivo. J. Virol. *85*, 5197–5201.

https://doi.org/10.1128/jvi.02692-10.

28.   Bryden, S.R., Pingen, M., Lefteri, D.A., Miltenburg, J., Delang, L., Jacobs, S.,

Abdelnabi, R., Neyts, J., Pondeville, E., Major, J., et al. (2020). Pan-viral protection

against arboviruses by activating skin macrophages at the inoculation site. Sci. Transl.

Med. *12*. https://doi.org/10.1126/scitranslmed.aax2421.

29.   Mirzayev, M. (2025). ANATOMY OF THE SKIN: STRUCTURE AND

FUNCTIONS OF THE EPIDERMIS AND DERMIS. Int. J. Med. Sci. *1*, 302–306.

30.   Arnold, S.J., Osvath, S.R., Hall, R.A., King, N.J.C., and Sedger, L.M. (2004).

Regulation of antigen processing and presentation molecules in West Nile virus-

infected human skin fibroblasts. Virology *324*, 286–296.

https://doi.org/10.1016/j.virol.2004.03.036.

31.   Montes-Gómez, A.E., García-Cordero, J., Marcial-Juárez, E., Shrivastava, G.,

Visoso-Carvajal, G., Juárez-Delgado, F.J., Flores-Romo, L., Sanchez-Torres, M.C.,

Santos-Argumedo, L., Bustos-Arriaga, J., et al. (2020). Crosstalk Between Dermal

Fibroblasts and Dendritic Cells During Dengue Virus Infection. Front. Immunol. *11*,

538240. https://doi.org/10.3389/fimmu.2020.538240.

32.   Kim, J.-A., Seong, R.-K., Son, S.W., and Shin, O.S. (2019). Insights into ZIKV-

Mediated Innate Immune Responses in Human Dermal Fibroblasts and Epidermal

Keratinocytes. J. Invest. Dermatol. *139,* 391–399.

https://doi.org/10.1016/j.jid.2018.07.038.

33.   Hoover, L.I., and Fredericksen, B.L. (2014). IFN-Dependent and -Independent

Reduction in West Nile Virus Infectivity in Human Dermal Fibroblasts. Viruses *6*, 1424–

1441. https://doi.org/10.3390/v6031424.

34.   Hamel, R., Dejarnac, O., Wichit, S., Ekchariyawat, P., Neyret, A., Luplertlop, N.,

Perera-Lecoin, M., Surasombatpattana, P., Talignani, L., Thomas, F., et al. (2015).

Biology of Zika Virus Infection in Human Skin Cells. J. Virol. *89*, 8880–8896.

https://doi.org/10.1128/jvi.00354-15.

35.   Garcia, M., Alout, H., Diop, F., Damour, A., Bengue, M., Weill, M., Missé, D.,

Lévêque, N., and Bodet, C. (2018). Innate Immune Response of Primary Human

Keratinocytes to West Nile Virus Infection and Its Modulation by Mosquito Saliva. Front.

Cell. Infect. Microbiol. *8*. https://doi.org/10.3389/fcimb.2018.00387.

36.   Bustos-Arriaga, J., García-Machorro, J., León-Juárez, M., García-Cordero, J.,

Santos-Argumedo, L., Flores-Romo, L., Méndez-Cruz, A.R., Juárez-Delgado, F.J., and

Cedillo-Barrón, L. (2011). Activation of the Innate Immune Response against DENV in

Normal Non-Transformed Human Fibroblasts. PLoS Negl. Trop. Dis. *5*.

https://doi.org/10.1371/journal.pntd.0001420.

37.   López-González, M., Meza-Sánchez, D., García-Cordero, J., Bustos-Arriaga,

J., Vélez-Del Valle, C., Marsch-Moreno, M., Castro-Jiménez, T., Flores-Romo, L.,

Santos-Argumedo, L., Gutiérrez-Castañeda, B., et al. (2018). Human keratinocyte

cultures (HaCaT) can be infected by DENV, triggering innate immune responses that

include IFNλ and LL37. Immunobiology *223*, 608–617.

https://doi.org/10.1016/j.imbio.2018.07.006.

38.   Bustos-Arriaga, J., Mita-Mendoza, N.K., Lopez-Gonzalez, M., García-Cordero,

J., Juárez-Delgado, F.J., Gromowski, G.D., Méndez-Cruz, R.A., Fairhurst, R.M.,

Whitehead, S.S., and Cedillo-Barrón, L. (2016). Soluble mediators produced by the

crosstalk between microvascular endothelial cells and dengue-infected primary dermal fibroblasts inhibit dengue virus replication and increase leukocyte transmigration. Immunol. Res. *64*, 392–403. https://doi.org/10.1007/s12026-015-8675-8.

39.   Martí, M.M., Castanha, P.M.S., and Barratt-Boyes, S.M. (2024). The Dynamic Relationship between Dengue Virus and the Human Cutaneous Innate Immune Response. Viruses *16*, 727. https://doi.org/10.3390/v16050727.

40.   Garcia, M., Wehbe, M., Lévêque, N., and Bodet, C. (2017). Skin innate immune response to flaviviral infection. Eur. Cytokine Netw. *28*, 41–51. https://doi.org/10.1684/ecn.2017.0394.

41.   Pingen, M., Schmid, M.A., Harris, E., and McKimmie, C.S. (2017). Mosquito Biting Modulates Skin Response to Virus Infection. Trends Parasitol. *33*, 645–657. https://doi.org/10.1016/j.pt.2017.04.003.

42.   Dudley, D.M., Newman, C.M., Lalli, J., Stewart, L.M., Koenig, M.R., Weiler, A.M., Semler, M.R., Barry, G.L., Zarbock, K.R., Mohns, M.S., et al. (2017). Infection via mosquito bite alters Zika virus tissue tropism and replication kinetics in rhesus macaques. Nat. Commun. *8*, 2096. https://doi.org/10.1038/s41467-017-02222-8.

43.   Cox, J., Mota, J., Sukupolvi-Petty, S., Diamond, M.S., and Rico-Hesse, R. (2012). Mosquito Bite Delivery of Dengue Virus Enhances Immunogenicity and Pathogenesis in Humanized Mice. J. Virol. *86*, 7637–7649. https://doi.org/10.1128/JVI.00534-12.

44.   Medkour, H., Pruvost, L., Miot, E.F., Gong, X., Vaissayre, V., Tavadia, M., Boutinaud, P., Revel, J., Hitakarun, A., Sornjai, W., et al. (2025). Sphingomyelins in mosquito saliva reconfigure skin lipidome to promote viral protein levels and enhance transmission of flaviviruses. Cell Metab. *0*. https://doi.org/10.1016/j.cmet.2025.05.015.

45.   Barrows, N.J., Campos, R.K., Liao, K.-C., Prasanth, K.R., Soto-Acosta, R., Yeh,

S.-C., Schott-Lerner, G., Pompon, J., Sessions, O.M., Bradrick, S.S., et al. (2018).

Biochemistry and Molecular Biology of Flaviviruses. Chem. Rev. *118*, 4448–4482.

https://doi.org/10.1021/acs.chemrev.7b00719.

46.   Almet, A.A., Liu, Y., Nie, Q., and Plikus, M.V. (2025). Integrated Single-Cell

Analysis Reveals Spatially and Temporally Dynamic Heterogeneity in Fibroblast States

during   Wound   Healing.   J.   Invest.   Dermatol.   *145*,   645-659.e25.

https://doi.org/10.1016/j.jid.2024.06.1281.

47.   Zhang, Y., Liu, G., Zeng, Q., Wu, W., Lei, K., Zhang, C., Tang, M., Zhang, Y.,

Xiang, X., Tan, L., et al. (2024). CCL19-producing fibroblasts promote tertiary lymphoid

structure formation enhancing anti-tumor IgG response in colorectal cancer liver

metastasis. Cancer Cell *42*, 1370-1385.e9. https://doi.org/10.1016/j.ccell.2024.07.006.

48.   Onder, L., Papadopoulou, C., Lütge, A., Cheng, H.-W., Lütge, M., Perez-

Shibayama, C., Gil-Cruz, C., De Martin, A., Kurz, L., Cadosch, N., et al. (2025).

Fibroblastic reticular cells generate protective intratumoral T cell environments in lung

cancer. Cell *188*, 430-446.e20. https://doi.org/10.1016/j.cell.2024.10.042.

49.   Li, Y., Long, J., Zhang, Z., and Yin, W. (2024). Insights into the unique roles of

dermal white adipose tissue (dWAT) in wound healing. Front. Physiol. *15*, 1346612.

https://doi.org/10.3389/fphys.2024.1346612.

50.   Buechler, M.B., Pradhan, R.N., Krishnamurty, A.T., Cox, C., Calviello, A.K.,

Wang, A.W., Yang, Y.A., Tam, L., Caothien, R., Roose-Girma, M., et al. (2021). Cross-

tissue   organization   of   the   fibroblast   lineage.   Nature   *593*,   575–579.

https://doi.org/10.1038/s41586-021-03549-5.

51.   Garrity, R., Arora, N., Haque, M.A., Weis, D., Trinh, R.T., Neerukonda, S.V.,

Kumari, S., Cortez, I., Ubogu, E.E., Mahalingam, R., et al. (2023). Fibroblast-derived

PI16 sustains inflammatory pain via regulation of CD206+ myeloid cells. Brain. Behav.

Immun. *112*, 220–234. https://doi.org/10.1016/j.bbi.2023.06.011.

52.   Garrity, R., Haque, A., Kavelaars, A., Heijnen, C., and Shepherd, A. (2024).

Fibroblast-derived PI16 Drives the Physiological Process of Immune Cell Recruitment

and   Activation   During   Inflammatory   Response.   J.   Pain   *25*,   13.

https://doi.org/10.1016/j.jpain.2024.01.067.

53.   McCartney, E.E., Chung, Y., and Buechler, M.B. (2024). Life of Pi: Exploring

functions   of   Pi16+   fibroblasts.   F1000Research   *13*,   126.

https://doi.org/10.12688/f1000research.143511.2.

54.   Rahmani, W., Abbasi, S., Hagner, A., Raharjo, E., Kumar, R., Hotta, A.,

Magness, S., Metzger, D., and Biernaskie, J. (2014). Hair follicle dermal stem cells

regenerate the dermal sheath, repopulate the dermal papilla, and modulate hair type.

Dev. Cell *31*, 543–558. https://doi.org/10.1016/j.devcel.2014.10.022.

55.   Joost, S., Zeisel, A., Jacob, T., Sun, X., La Manno, G., Lönnerberg, P.,

Linnarsson, S., and Kasper, M. (2016). Single-Cell Transcriptomics Reveals that

Differentiation and Spatial Signatures Shape Epidermal and Hair Follicle

Heterogeneity. Cell Syst. *3*, 221-237.e9. https://doi.org/10.1016/j.cels.2016.08.010.

56.   Guerrero-Juarez, C.F., Dedhia, P.H., Jin, S., Ruiz-Vega, R., Ma, D., Liu, Y.,

Yamaga, K., Shestova, O., Gay, D.L., Yang, Z., et al. (2019). Single-cell analysis

reveals fibroblast heterogeneity and myeloid-derived adipocyte progenitors in murine

skin wounds. Nat. Commun. *10*, 650. https://doi.org/10.1038/s41467-018-08247-x.

57.   Philippeos, C., Telerman, S.B., Oulès, B., Pisco, A.O., Shaw, T.J., Elgueta, R.,

Lombardi, G., Driskell, R.R., Soldin, M., Lynch, M.D., et al. (2018). Spatial and Single-

Cell Transcriptional Profiling Identifies Functionally Distinct Human Dermal Fibroblast

Subpopulations. J. Invest. Dermatol. *138,* 811–825.

https://doi.org/10.1016/j.jid.2018.01.016.

58. Avigad Laron, E., Aamar, E., and Enshell-Seijffers, D. (2018). The

Mesenchymal Niche of the Hair Follicle Induces Regeneration by Releasing Primed

Progenitors from Inhibitory Effects of Quiescent Stem Cells. Cell Rep. *24,* 909-921.e3.

https://doi.org/10.1016/j.celrep.2018.06.084.

59. Plikus, M.V., Wang, X., Sinha, S., Forte, E., Thompson, S.M., Herzog, E.L.,

Driskell, R.R., Rosenthal, N., Biernaskie, J., and Horsley, V. (2021). Fibroblasts:

Origins, definitions, and functions in health and disease. Cell *184,* 3852–3872.

https://doi.org/10.1016/j.cell.2021.06.024.

60. Joost, S., Annusver, K., Jacob, T., Sun, X., Dalessandri, T., Sivan, U., Sequeira,

I., Sandberg, R., and Kasper, M. (2020). The Molecular Anatomy of Mouse Skin during

Hair Growth and Rest. Cell Stem Cell *26,* 441-457.e7.

https://doi.org/10.1016/j.stem.2020.01.012.

61. Blanpain, C., and Fuchs, E. (2009). Epidermal homeostasis: a balancing act of

stem cells in the skin. Nat. Rev. Mol. Cell Biol. *10,* 207–217.

https://doi.org/10.1038/nrm2636.

62. Watt, F.M. (2014). Mammalian skin cell biology: at the interface between

laboratory and clinic. Science *346,* 937–940. https://doi.org/10.1126/science.1253734.

63. Zouboulis, C.C., Picardo, M., Ju, Q., Kurokawa, I., Törőcsik, D., Bíró, T., and

Schneider, M.R. (2016). Beyond acne: Current aspects of sebaceous gland biology

and function. Rev. Endocr. Metab. Disord. *17,* 319–334.

https://doi.org/10.1007/s11154-016-9389-5.

64. Rogers, G.E. (2004). Hair follicle differentiation and regulation. Int. J. Dev. Biol.

48, 163–170.

65.   Reggio, A., Fuoco, C., Deodati, R., and Palma, A. (2025). SPP1 macrophages

across diseases: A call for reclassification? FASEB J. Off. Publ. Fed. Am. Soc. Exp.

Biol. *39*, e70448. https://doi.org/10.1096/fj.202403227R.

66.   Nalio Ramos, R., Missolo-Koussou, Y., Gerber-Ferder, Y., Bromley, C.P.,

Bugatti, M., Núñez, N.G., Tosello Boari, J., Richer, W., Menger, L., Denizeau, J., et al.

(2022). Tissue-resident FOLR2+ macrophages associate with CD8+ T cell infiltration

in   human   breast   cancer.   Cell   *185*,   1189-1207.e25.

https://doi.org/10.1016/j.cell.2022.02.021.

67.   Jenkins, S.J., and Allen, J.E. (2021). The expanding world of tissue-resident

macrophages. Eur. J. Immunol. *51*, 1882–1896. https://doi.org/10.1002/eji.202048881.

68.   Muntjewerff, E.M., Meesters, L.D., and van den Bogaart, G. (2020). Antigen

Cross-Presentation   by   Macrophages.   Front.   Immunol.   *11*,   1276.

https://doi.org/10.3389/fimmu.2020.01276.

69.   Guilliams, M., Ginhoux, F., Jakubzick, C., Naik, S.H., Onai, N., Schraml, B.U.,

Segura, E., Tussiwand, R., and Yona, S. (2014). Dendritic cells, monocytes and

macrophages: a unified nomenclature based on ontogeny. Nat. Rev. Immunol. *14*,

571–578. https://doi.org/10.1038/nri3712.

70.   Ohl, L., Mohaupt, M., Czeloth, N., Hintzen, G., Kiafard, Z., Zwirner, J.,

Blankenstein, T., Henning, G., and Förster, R. (2004). CCR7 governs skin dendritic

cell migration under inflammatory and steady-state conditions. Immunity *21*, 279–288.

https://doi.org/10.1016/j.immuni.2004.06.014.

71.   Randolph, G.J., Angeli, V., and Swartz, M.A. (2005). Dendritic-cell trafficking to

lymph   nodes   through   lymphatic   vessels.   Nat.   Rev.   Immunol.   *5*,   617–628.

https://doi.org/10.1038/nri1670.

72.   Shi, C., and Pamer, E.G. (2011). Monocyte recruitment during infection and

inflammation. Nat. Rev. Immunol. *11*, 762–774. https://doi.org/10.1038/nri3070.

73.   Nauseef, W.M., and Borregaard, N. (2014). Neutrophils at work. Nat. Immunol.

15, 602–611. https://doi.org/10.1038/ni.2921.

74.   Klose, C.S.N., and Artis, D. (2016). Innate lymphoid cells as regulators of

immunity, inflammation and tissue homeostasis. Nat. Immunol. *17*, 765–774.

https://doi.org/10.1038/ni.3489.

75.   Kim, B.S. (2015). Innate Lymphoid Cells in the Skin. J. Invest. Dermatol. *135*,

673–678. https://doi.org/10.1038/jid.2014.401.

76.   MacLeod, A.S., and Havran, W.L. (2011). Functions of skin-resident γδ T cells.

Cell. Mol. Life Sci. *68*, 2399–2408. https://doi.org/10.1007/s00018-011-0702-x.

77.   Bielecki, P., Riesenfeld, S.J., Hütter, J.-C., Torlai Triglia, E., Kowalczyk, M.S.,

Ricardo-Gonzalez, R.R., Lian, M., Amezcua Vesely, M.C., Kroehling, L., Xu, H., et al.

(2021). Skin-resident innate lymphoid cells converge on a pathogenic effector state.

Nature *592*, 128–132. https://doi.org/10.1038/s41586-021-03188-w.

78.   Kokkinou, E., Pandey, R.V., Mazzurana, L., Gutierrez-Perez, I., Tibbitt, C.A.,

Weigel, W., Soini, T., Carrasco, A., Rao, A., Nagasawa, M., et al. (2022).

CD45RA+CD62L− ILCs in human tissues represent a quiescent local reservoir for the

generation of differentiated ILCs. Sci. Immunol. *7*, eabj8301.

https://doi.org/10.1126/sciimmunol.abj8301.

79.   Maharaj, P.D., Widen, S.G., Huang, J., Wood, T.G., and Thangamani, S.

(2015). Discovery of Mosquito Saliva MicroRNAs during CHIKV Infection. PLoS Negl.

Trop. Dis. *9*. https://doi.org/10.1371/journal.pntd.0003386.

80.   Fiorillo, C., Yen, P.-S., Colantoni, A., Mariconti, M., Azevedo, N., Lombardo, F.,

Failloux, A.-B., and Arcà, B. (2022). MicroRNAs and other small RNAs in Aedes

aegypti saliva and salivary glands following chikungunya virus infection. Sci. Rep. *12*,

9536. https://doi.org/10.1038/s41598-022-13780-3.

81.   Galloway, C.J., Dean, G.E., Marsh, M., Rudnick, G., and Mellman, I. (1983).

Acidification of macrophage and fibroblast endocytic vesicles in vitro. Proc. Natl. Acad.

Sci. *80*, 3334–3338. https://doi.org/10.1073/pnas.80.11.3334.

82.   Yang, W., He, R., Qu, H., Lian, W., Xue, Y., Wang, T., Lin, W., Zhu, P., Xia, M.,

Lai, L., et al. (2023). FXYD3 enhances IL-17A signaling to promote psoriasis by

competitively binding TRAF3 in keratinocytes. Cell. Mol. Immunol. *20*, 292–304.

https://doi.org/10.1038/s41423-023-00973-7.

83.   Charles A Janeway, J., Travers, P., Walport, M., and Shlomchik, M.J. (2001).

The major histocompatibility complex and its functions. In Immunobiology: The

Immune System in Health and Disease. 5th edition (Garland Science).

84.   Li, J.-P., Fu, Y.-N., Chen, Y.-R., and Tan, T.-H. (2010). JNK Pathway-

associated Phosphatase Dephosphorylates Focal Adhesion Kinase and Suppresses

Cell   Migration.   J.   Biol.   Chem.   *285*,   5472–5478.

https://doi.org/10.1074/jbc.M109.060186.

85.   Xue, X., Bredell, B.X., Anderson, E.R., Martin, A., Mays, C., Nagao-Kitamoto,

H., Huang, S., Győrffy, B., Greenson, J.K., Hardiman, K., et al. (2017). Quantitative

proteomics identifies STEAP4 as a critical regulator of mitochondrial dysfunction

linking inflammation and colon cancer. Proc. Natl. Acad. Sci. U. S. A. *114*, E9608–

E9617. https://doi.org/10.1073/pnas.1712946114.

86.   Cheng, L., Xu, Y., Li, Z., Sun, D., Xu, Y., He, J., Chen, Y., and Yao, Q. (2025).

DHCR24 overexpression is involved in lipid metabolic reprogramming to drive cervical

cancer malignant progression and is associated with immune microenvironment. BMC

Cancer *25*, 1291. https://doi.org/10.1186/s12885-025-14663-2.

87.   Cai, D., Zhong, G.-C., Dai, X., Zhao, Z., Chen, M., Hu, J., Wu, Z., Cheng, L., Li,

S., and Gong, J. (2025). Targeting FDFT1 Reduces Cholesterol and Bile Acid

Production and Delays Hepatocellular Carcinoma Progression Through the

HNF4A/ALDOB/AKT1 Axis. Adv. Sci. Weinh. Baden-Wurtt. Ger. *12*, e2411719.

https://doi.org/10.1002/advs.202411719.

88.   El-Kurjieh, A., Al-Arab, R., Hachem, Q.A., Ibrahim, J.-N., and Kobeissy, P.H.

(2025). ACSS2 and metabolic diseases: from lipid metabolism to therapeutic target.

Lipids Health Dis. *24*, 74. https://doi.org/10.1186/s12944-025-02491-z.

89.   Houddane, A., Bultot, L., Novellasdemunt, L., Johanns, M., Gueuning, M.-A.,

Vertommen, D., Coulie, P.G., Bartrons, R., Hue, L., and Rider, M.H. (2017). Role of

Akt/PKB and PFKFB isoenzymes in the control of glycolysis, cell proliferation and

protein synthesis in mitogen-stimulated thymocytes. Cell. Signal. *34*, 23–37.

https://doi.org/10.1016/j.cellsig.2017.02.019.

90.   Leclerc, E.A., Huchenq, A., Kezic, S., Serre, G., and Jonca, N. (2014). Mice

deficient for the epidermal dermokine β and γ isoforms display transient cornification

defects. J. Cell Sci. *127*, 2862–2872. https://doi.org/10.1242/jcs.144808.

91.   Wang, H.U., Chen, Z.F., and Anderson, D.J. (1998). Molecular distinction and

angiogenic interaction between embryonic arteries and veins revealed by ephrin-B2

and its receptor Eph-B4. Cell *93*, 741–753. https://doi.org/10.1016/s0092-

8674(00)81436-1.

92.   Elkins, J.M., Amos, A., Niesen, F.H., Pike, A.C.W., Fedorov, O., and Knapp, S.

(2009). Structure of dystrophia myotonica protein kinase. Protein Sci. Publ. Protein

Soc. *18*, 782–791. https://doi.org/10.1002/pro.82.

93.   Cao, J.-M., Cheng, X.-N., Li, S.-Q., Heller, S., Xu, Z.-G., and Shi, D.-L. (2016).

Identification of novel MYO18A interaction partners required for myoblast adhesion

and muscle integrity. Sci. Rep. *6*, 36768. https://doi.org/10.1038/srep36768.

94.   Lin, J., He, Y., Chen, L., Chen, X., Zang, S., and Lin, W. (2018). MYLK promotes

hepatocellular carcinoma progression through regulating cytoskeleton to enhance

epithelial-mesenchymal transition. Clin. Exp. Med. *18*, 523–533.

https://doi.org/10.1007/s10238-018-0509-2.

95.   Yang, Z., Wei, W., Nie, D., Zhang, M., and Chen, Q. (2025). UCHL3 regulates

snail stability and promotes epithelial–mesenchymal transition in ovarian cancer.

Cytotechnology *77*, 152. https://doi.org/10.1007/s10616-025-00811-w.

96.   Feng, X., Wei, Z., Zhang, S., Zhou, J., Wu, J., Luan, B., Du, Y., and Zhao, H.

(2021). Overexpression of LVRN impedes the invasion of trophoblasts by inhibiting

epithelial-mesenchymal transition. Acta Biochim. Biophys. Sin. *53*, 249–257.

https://doi.org/10.1093/abbs/gmaa167.

97.   Zhang, S., Geng, Z., Sun, J., Chen, N., Liu, M., Wang, Y., Gu, X., and Pei, B.

(2025). TPD52 promotes the proliferation and metastasis of gastric cancer cells. BMC

Gastroenterol. *25*, 690. https://doi.org/10.1186/s12876-025-04295-y.

98.   Yu, J., Perri, M., Jones, J.W., Pierzchalski, K., Ceaicovscaia, N., Cione, E., and

Kane, M.A. (2022). Altered RBP1 Gene Expression Impacts Epithelial Cell Retinoic

Acid, Proliferation, and Microenvironment. Cells *11*, 792.

https://doi.org/10.3390/cells11050792.

99.   Li, H., Sun, C., Luo, B., Zhan, C., Li, W., Deng, L., Kang, K., and Gou, D. (2024).

Exploring the Spectrum of Long Non-Coding RNA CARMN in Physiological and

Pathological Contexts. Biomolecules *14*, 954. https://doi.org/10.3390/biom14080954.

100.   Bache, K.G., Slagsvold, T., Cabezas, A., Rosendal, K.R., Raiborg, C., and

Stenmark, H. (2004). The growth-regulatory protein HCRP1/hVps37A is a subunit of

mammalian ESCRT-I and mediates receptor down-regulation. Mol. Biol. Cell *15*, 4337–

4346. https://doi.org/10.1091/mbc.e04-03-0250.

101.   Ghuwalewala, S., Cao, J., Rezza, A., Rangl, M., Kwiatkowski, A., Brown, A.,

Grisanti, L., Heitman, N., Schober, M., Wang, Z., et al. (2025). Molecular signatures

and signaling interactions of the hair follicle stem cell niche. J. Invest. Dermatol.,

S0022-202X(25)03637-1. https://doi.org/10.1016/j.jid.2025.11.021.

102.   Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H.,

Myung, P., Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell

communication   using   CellChat.   Nat.   Commun.   *12*,   1088.

https://doi.org/10.1038/s41467-021-21246-9.

103.   Jin, S., Plikus, M.V., and Nie, Q. (2025). CellChat for systematic analysis of cell-

cell communication from single-cell transcriptomics. Nat. Protoc. *20*, 180–219.

https://doi.org/10.1038/s41596-024-01045-4.

104.   Kiseleva, E.P., and Rutto, K.V. (2022). Semaphorin 3A in the Immune System:

Twenty   Years   of   Study.   Biochem.   Mosc.   *87*,   640–657.

https://doi.org/10.1134/S0006297922070069.

105.   Sawada, Y., Mashima, E., Saito-Sasaki, N., and Nakamura, M. (2020). The Role

of Cell Adhesion Molecule 1 (CADM1) in Cutaneous Malignancies. Int. J. Mol. Sci. *21*.

https://doi.org/10.3390/ijms21249732.

106.   Laffranchi, M., Schioppa, T., Sozio, F., Piserà, A., Tiberio, L., Salvi, V., Bosisio,

D., Musso, T., Sozzani, S., and Del Prete, A. (2025). Chemerin in immunity. J. Leukoc.

Biol. *117*, qiae181. https://doi.org/10.1093/jleuko/qiae181.

107. Darlington, K.C., Shumway, A.J., Sona, C., Khaled, Z., Steinbach, E.C., Kikuchi,

Y., Fan, S., Gartner, V., Clough, K.M., Nishiyama, N.C., et al. (2026). Myeloid Cell

Adhesion Molecule 1 (CADM1) Promotes Proinflammatory Signaling in Human

Inflammatory Bowel Diseases. Cell. Mol. Gastroenterol. Hepatol. *20*, 101618.

https://doi.org/10.1016/j.jcmgh.2025.101618.

108. Zhu, X., Yang, Z., Wang, J., and Rosmarin, A.G. (2011). Consistent Activation

of Flt3 Promotes Hematopoietic Progenitors Toward Dendritic Cell Development in

Mouse. Blood *118*, 4789. https://doi.org/10.1182/blood.V118.21.4789.4789.

109. Hulpke, S., and Tampé, R. (2013). The MHC I loading complex: a multitasking

machinery in adaptive immunity. Trends Biochem. Sci. *38*, 412–420.

https://doi.org/10.1016/j.tibs.2013.06.003.

110. Suchitha, G.P., Devasahayam Arokia Balaya, R., Prasad, T.S.K., and

Dagamajalu, S. (2024). A signaling network map of Lipoxin (LXA4): an anti-

inflammatory molecule. Inflamm. Res. *73*, 1099–1106. https://doi.org/10.1007/s00011-

024-01885-6.

111. Li, J., Wang, Z., Qin, X., Zhong, M.-C., Tang, Z., Qian, J., Dou, J., Hussell, T.,

King, P.D., Nunès, J.A., et al. (2025). CD200R1-CD200 checkpoint inhibits

phagocytosis differently from SIRPα-CD47 to suppress tumor growth. Nat. Commun.

16, 5145. https://doi.org/10.1038/s41467-025-60456-3.

112. Worthington, J.J., Fenton, T.M., Czajkowska, B.I., Klementowicz, J.E., and

Travis, M.A. (2012). Regulation of TGFβ in the immune system: An emerging role for

integrins and dendritic cells. Immunobiology *217*, 1259–1265.

https://doi.org/10.1016/j.imbio.2012.06.009.

113. Barclay, A.N., and Brown, M.H. (2006). The SIRP family of receptors and

immune regulation. Nat. Rev. Immunol. *6*, 457–464. https://doi.org/10.1038/nri1859.

114.   Evans, D.T., Serra-Moreno, R., Singh, R.K., and Guatelli, J.C. (2010). BST-

2/tetherin: a new component of the innate immune response to enveloped viruses.

Trends Microbiol. *18*, 388–396. https://doi.org/10.1016/j.tim.2010.06.010.

115.   Pascoal Ramos, M.I., van der Vlist, M., and Meyaard, L. (2025). Inhibitory

pattern recognition receptors: lessons from LAIR1. Nat. Rev. Immunol. *25*, 711–724.

https://doi.org/10.1038/s41577-025-01181-2.

116.   Chitu, V., and Stanley, E.R. (2006). Colony-stimulating factor-1 in immunity and

inflammation.   Curr.   Opin.   Immunol.   *18*,   39–48.

https://doi.org/10.1016/j.coi.2005.11.006.

117.   Klaas, M., Dubock, S., Ferguson, D.J.P., and Crocker, P.R. (2023).

Sialoadhesin (CD169/Siglec-1) is an extended molecule that escapes inhibitory cis-

interactions and synergizes with other macrophage receptors to promote

phagocytosis. Glycoconj. J. *40*, 213–223. https://doi.org/10.1007/s10719-022-10097-

1.

118.   Sun, C., Mezzadra, R., and Schumacher, T.N. (2018). Regulation and Function

of   the   PD-L1   Checkpoint.   Immunity   *48*,   434–452.

https://doi.org/10.1016/j.immuni.2018.03.014.

119.   Yang, J., Yang, K., Wang, K., Zhou, D., Zhou, J., Du, X., Liu, S., and Cheng, Z.

(2023). Serum amyloid A regulates TLR2/4-mediated IFN-β signaling pathway against

Marek's   disease   virus.   Virus   Res.   *326*,   199044.

https://doi.org/10.1016/j.virusres.2023.199044.

120.   Loo, L., Waller, M.A., Moreno, C.L., Cole, A.J., Stella, A.O., Pop, O.-T., Jochum,

A.-K., Ali, O.H., Denes, C.E., Hamoudi, Z., et al. (2023). Fibroblast-expressed LRRC15

is a receptor for SARS-CoV-2 spike and controls antiviral and antifibrotic transcriptional

programs. PLoS Biol. *21*, e3001967. https://doi.org/10.1371/journal.pbio.3001967.

121. O'Prey, J., Wilkinson, S., and Ryan, K.M. (2008). Tumor antigen LRRC15

impedes adenoviral infection: implications for virus-based cancer therapy. J. Virol. *82*,

5933–5939. https://doi.org/10.1128/JVI.02273-07.

122. Yu, X., Zhu, Y., Yin, G., Wang, Y., Shi, X., and Cheng, G. (2024). Exploiting

hosts and vectors: viral strategies for facilitating transmission. EMBO Rep. *25*, 3187–

3201. https://doi.org/10.1038/s44319-024-00214-6.

123. Kolaczkowska, E., and Kubes, P. (2013). Neutrophil recruitment and function in

health and inflammation. Nat. Rev. Immunol. *13*, 159–175.

https://doi.org/10.1038/nri3399.

124. Greenlee-Wacker, M.C. (2016). Clearance of apoptotic neutrophils and

resolution of inflammation. Immunol. Rev. *273*, 357–370.

https://doi.org/10.1111/imr.12453.

125. Yunna, C., Mengru, H., Lei, W., and Weidong, C. (2020). Macrophage M1/M2

polarization. Eur. J. Pharmacol. *877*, 173090.

https://doi.org/10.1016/j.ejphar.2020.173090.

126. Ortega-Gómez, A., Perretti, M., and Soehnlein, O. (2013). Resolution of

inflammation: an integrated view. EMBO Mol. Med. *5*, 661–674.

https://doi.org/10.1002/emmm.201202382.

127. Reider, I.E., Lin, E., Krouse, T.E., Parekh, N.J., Nelson, A.M., and Norbury, C.C.

(2024). γδ T Cells Mediate a Requisite Portion of a Wound Healing Response

Triggered by Cutaneous Poxvirus Infection. Viruses *16*.

https://doi.org/10.3390/v16030425.

128. Lujan, R.A., Pei, L., Shannon, J.P., Dábilla, N., Dolan, P.T., and Hickman, H.D.

(2023). Widespread and dynamic expression of granzyme C by skin-resident antiviral

T cells. Front. Immunol. *14*. https://doi.org/10.3389/fimmu.2023.1236595.

129.   Simmons, J., and Gallo, R.L. (2024). The Central Roles of Keratinocytes in

Coordinating   Skin   Immunity.   J.   Invest.   Dermatol.   *144*,   2377–2398.

https://doi.org/10.1016/j.jid.2024.06.1280.

130.   Ding, J., Chou, Y.-Y., and Chang, T.L. (2009). Defensins in viral infections. J.

Innate Immun. *1*, 413–420. https://doi.org/10.1159/000226256.

131.   Charles A Janeway, J., Travers, P., Walport, M., and Shlomchik, M.J. (2001).

The complement system and innate immunity. In Immunobiology: The Immune System

in Health and Disease. 5th edition (Garland Science).

132.   Waskow, C., Liu, K., Darrasse-Jèze, G., Guermonprez, P., Ginhoux, F., Merad,

M., Shengelia, T., Yao, K., and Nussenzweig, M. (2008). The receptor tyrosine kinase

Flt3 is required for dendritic cell development in peripheral lymphoid tissues. Nat.

Immunol. *9*, 676–683. https://doi.org/10.1038/ni.1615.

133.   Takamatsu, H., and Kumanogoh, A. (2012). Diverse roles for semaphorin-plexin

signaling   in   the   immune   system.   Trends   Immunol.   *33*,   127–135.

https://doi.org/10.1016/j.it.2012.01.008.

134.   Suchitha,   G.P.,   Devasahayam   Arokia   Balaya,   R.,   Prasad,   T.S.K.,   and

Dagamajalu, S. (2024). A signaling network map of Lipoxin (LXA4): an anti-

inflammatory molecule. Inflamm. Res. Off. J. Eur. Histamine Res. Soc. Al *73*, 1099–

1106. https://doi.org/10.1007/s00011-024-01885-6.

135.   Vaine, C.A., and Soberman, R.J. (2014). The CD200-CD200R1 inhibitory

signaling pathway: immune regulation and host-pathogen interactions. Adv. Immunol.

121, 191–211. https://doi.org/10.1016/B978-0-12-800100-4.00005-2.

136.   Solé-Boldo, L., Raddatz, G., Schütz, S., Mallm, J.-P., Rippe, K., Lonsdorf, A.S.,

Rodríguez-Paredes, M., and Lyko, F. (2020). Single-cell transcriptomes of the human

skin reveal age-related loss of fibroblast priming. Commun. Biol. *3*, 188. https://doi.org/10.1038/s42003-020-0922-4.

137. Dolan, J., Walshe, K., Alsbury, S., Hokamp, K., O'Keeffe, S., Okafuji, T., Miller, S.F.C., Tear, G., and Mitchell, K.J. (2007). The extracellular leucine-rich repeat superfamily; a comparative survey and analysis of evolutionary relationships and expression patterns. BMC Genomics *8*, 320. https://doi.org/10.1186/1471-2164-8-320.

138. Imler, J.L., and Hoffmann, J.A. (2001). Toll receptors in innate immunity. Trends Cell Biol. *11*, 304–311. https://doi.org/10.1016/s0962-8924(01)02004-9.

139. Garg, H., Yeh, R., Watts, D.M., Mehmetoglu-Gurbuz, T., Resendes, R., Parsons, B., Gonzales, F., and Joshi, A. (2021). Enhancement of Zika virus infection by antibodies from West Nile virus seropositive individuals with no history of clinical infection. BMC Immunol. *22*, 5. https://doi.org/10.1186/s12865-020-00389-2.

140. Lucas, M., Frenkiel, M.-P., Mashimo, T., Guénet, J.-L., Deubel, V., Desprès, P., and Ceccaldi, P.-E. (2004). The Israeli strain IS-98-ST1 of West Nile virus as viral model for West Nile encephalitis in the Old World. Virol. J. *1*, 9. https://doi.org/10.1186/1743-422X-1-9.

141. Hamel, R., Narpon, Q., Serrato-Pomar, I., Gauliard, C., Berthomieu, A., Wichit, S., Missé, D., Sofonea, M.T., and Pompon, J. (2024). West Nile virus can be transmitted within mosquito populations through infectious mosquito excreta. iScience *27*, 111099. https://doi.org/10.1016/j.isci.2024.111099.

142. Kuno, G. (2010). Early history of laboratory breeding of Aedes aegypti (Diptera: Culicidae) focusing on the origins and use of selected strains. J. Med. Entomol. *47*, 957–971. https://doi.org/10.1603/me10152.

143. Pompon, J., Manuel, M., Ng, G.K., Wong, B., Shan, C., Manokaran, G., Soto-Acosta, R., Bradrick, S.S., Ooi, E.E., Missé, D., et al. (2017). Dengue subgenomic

flaviviral RNA disrupts immunity in mosquito salivary glands to increase virus

transmission. PLoS Pathog. *13*, e1006535.

https://doi.org/10.1371/journal.ppat.1006535.

144. Yeh, S.-C., Strilets, T., Tan, W.-L., Castillo, D., Medkour, H., Rey-Cadilhac, F.,

Serrato-Pomar, I.M., Rachenne, F., Chowdhury, A., Chuo, V., et al. (2023). The anti-

immune dengue subgenomic flaviviral RNA is present in vesicles in mosquito saliva

and is associated with increased infectivity. PLOS Pathog. *19*, e1011224.

https://doi.org/10.1371/journal.ppat.1011224.

145. Forootan, A., Sjöback, R., Björkman, J., Sjögreen, B., Linz, L., and Kubista, M.

(2017). Methods to determine limit of detection and limit of quantification in quantitative

real-time PCR (qPCR). Biomol. Detect. Quantif. *12*, 1–6.

https://doi.org/10.1016/j.bdq.2017.04.001.

146. Graham, J.B., Swarts, J.L., and Lund, J.M. (2017). A Mouse Model of West Nile

Virus Infection. Curr. Protoc. Mouse Biol. *7*, 221–235. https://doi.org/10.1002/cpmo.33.

147. Matthews, B.J., Dudchenko, O., Kingan, S.B., Koren, S., Antoshechkin, I.,

Crawford, J.E., Glassford, W.J., Herre, M., Redmond, S.N., Rose, N.H., et al. (2018).

Improved reference genome of Aedes aegypti informs arbovirus vector control. Nature

563, 501–507. https://doi.org/10.1038/s41586-018-0692-z.

148. West Nile virus lineage 1, complete genome (2019).

http://www.ncbi.nlm.nih.gov/nuccore/NC_009942.1

http://www.ncbi.nlm.nih.gov/nuccore/NC_009942.1.

149. Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A.,

Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., et al. (2024). Dictionary

learning for integrative, multimodal and scalable single-cell analysis. Nat. Biotechnol.

42, 293–304. https://doi.org/10.1038/s41587-023-01767-y.

150. Miller, S.A., Policastro, R.A., Sriramkumar, S., Lai, T., Huntington, T.D.,

Ladaika, C.A., Kim, D., Hao, C., Zentner, G.E., and O'Hagan, H.M. (2021). LSD1 and

Aberrant DNA Methylation Mediate Persistence of Enteroendocrine Progenitors That

Support BRAF-Mutant Colorectal Cancer. Cancer Res. *81*, 3791–3805.

https://doi.org/10.1158/0008-5472.CAN-20-3562.

151. Finak, G., McDavid, A., Yajima, M., Deng, J., Gersuk, V., Shalek, A.K., Slichter,

C.K., Miller, H.W., McElrath, M.J., Prlic, M., et al. (2015). MAST: a flexible statistical

framework for assessing transcriptional changes and characterizing heterogeneity in

single-cell RNA sequencing data. Genome Biol. *16*, 278.

https://doi.org/10.1186/s13059-015-0844-5.

152. Fang, Z., Liu, X., and Peltz, G. (2023). GSEApy: a comprehensive package for

performing gene set enrichment analysis in Python. Bioinformatics *39*, btac757.

https://doi.org/10.1093/bioinformatics/btac757.

153. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H.,

Myung, P., Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell

communication using CellChat. Nat. Commun. *12*, 1088.

https://doi.org/10.1038/s41467-021-21246-9.

154. Vora, A., Zhou, W., Londono-Renteria, B., Woodson, M., Sherman, M.B.,

Colpitts, T.M., Neelakanta, G., and Sultana, H. (2018). Arthropod EVs mediate dengue

virus transmission through interaction with a tetraspanin domain containing

glycoprotein Tsp29Fb. Proc. Natl. Acad. Sci. *115*, E6604–E6613.

https://doi.org/10.1073/pnas.1720125115.

155. Sun, P., Nie, K., Zhu, Y., Liu, Y., Wu, P., Liu, Z., Du, S., Fan, H., Chen, C.-H.,

Zhang, R., et al. (2020). A mosquito salivary protein promotes flavivirus transmission

by activation of autophagy. Nat. Commun. *11*, 260. https://doi.org/10.1038/s41467-

019-14115-z.

156.   Jones, D.C., Elz, A.E., Hadadianpour, A., Ryu, H., Glass, D.R., and Newell,

E.W. (2025). Cell Simulation as Cell Segmentation. Nat. Methods *22*, 1331–1342.

https://doi.org/10.1038/s41592-025-02697-0.

157.   Chen, X., Totty, M., and Hicks, S.C. (2025). Spotsweeper-py: spatially-aware

quality control metrics for spatial omics data in the Python ecosystem. bioRxiv,

2025.12.06.692760. https://doi.org/10.64898/2025.12.06.692760.

158.   Ergen, C., and Yosef, N. (2025). ResolVI - addressing noise and bias in spatial

transcriptomics. Preprint at bioRxiv, https://doi.org/10.1101/2025.01.20.634005

https://doi.org/10.1101/2025.01.20.634005.

159.   Kleshchevnikov, V., Shmatko, A., Dann, E., Aivazidis, A., King, H.W., Li, T.,

Elmentaite, R., Lomakin, A., Kedlian, V., Gayoso, A., et al. (2022). Cell2location maps

fine-grained cell types in spatial transcriptomics. Nat. Biotechnol. *40*, 661–671.

https://doi.org/10.1038/s41587-021-01139-4.

1506

1507

1508

Figure 1. scRNA-seq analysis on mouse skin at the onset of bite-initiated infection

(A, B) Early WNV replication kinetics in the skin (A) and inguinal LNs of mice (B) at 10 min, 1, 3, 6, 9 and 24 hours (h) post-bite. Bars show geometric means ± 95% C.I. Point

represents one skin biopsy (A) and the two inguinal LNs from the corresponding mouse

(B). N mice, 3-4 per time point. Different letters indicate statistical differences as

determined by post-hoc Fisher LSD's test.

(C) Experimental workflow for the scRNA-seq assay on bitten mouse skin.

(D) Numbers of high-quality single cells retained after quality control (QC) on scRNA-

seq data.

(E) UMAP dimensional reduction of sequenced skin cells revealing eight cell types.

Proportion of each cell cluster is provided in brackets. EC, endothelial cells; SMC,

smooth muscle cells.

(F) Heatmap of the top 20 enriched genes per cell cluster. Four marker genes for each

cluster are shown on the right.

(G-J) UMAP subclustering of fibroblasts (G), keratinocytes and ectodermal appendage

cells (H), myeloid cells (I), and lymphocytes (J). Relative cell proportions are indicated

in brackets. Fib, fibroblasts; Ker, keratinocytes; DP, dermal papilla cells; dWAT, dermal

white adipose tissue cells; HF, hair follicle cells; IRS, inner root sheath cells; HFSC,

hair follicle stem cells; DETC, dendritic epidermal T cells; ILC, innate lymphoid T cells.

See also **Figures S1-S6 and Datasets S1A-B, S2, S3.**

1530

1531

Figure 2. Cellular distribution of mosquito-derived RNAs at the bite site

(A) Gene identification of mosquito salivary RNA fragments.

(B) Distribution of mosquito RNA-positive mouse skin cells. Percent of mosquito RNA-

positive cells are indicated in brackets.

(C, D) Proportion of mosquito RNA-positive cells between cell types (C) and fibroblast

subtypes (D). Numbers above the bars indicate absolute number of positive mouse

skin cells over total cells.

(E) Differentially expressed genes (DEG) in mosquito RNA-positive fibroblasts.

Functional categories of DEGs are indicated by colored boxes.

See also **Figure S7 and Dataset S4.**

1542

Figure 3. Cellular landscape of mouse skin after infectious bite

(A-C) Distribution of cell types in unbitten skin (resting) (A) and in skin at 6 hours post-

infectious bite (infectious bite) (B), and percent comparisons and cell type fold change

between infectious *vs.* rest conditions (C).

(D-O) Distribution of cell subtypes among fibroblasts (D-F), keratinocytes and

ectodermal appendage cells (G-I), myeloid cells (J-L), and lymphocytes (M-O) in

resting skin and skin at 6 hours post-infectious bite.

(A, B) Proportions within all cells are shown in brackets. (D, E, G, H, J, K, M, N) Within-

cell type proportions are shown in brackets. (C, F, I, L, O) Changes in cell proportions

are expressed as log2 fold difference (Log2FD) means ± standard deviation in

infectious bite as compared to resting conditions. Dashed black lines indicate a fold

change of 1.5. Statistical significance indicated in red for p < 0.05 based on permutation

test. ECs, endothelial cells; SMCs, smooth muscle cells; DP, dermal papilla; dWAT,

dermal white adipose tissue; HF, hair follicle; IRS, inner root sheath; DCs, dendritic

cells; HFSC, hair follicle stem cells; DETC, dendritic epidermal T cells; ILC, innate

lymphoid T cells.

1560

Figure 4. Cell-cell communication changes in response to infectious bite

(A) Influence of infectious bite on the overall cell-cell communication (CCC) strength

between skin cell subtypes. NK, natural killer; F, fibroblasts; DP, dermal papilla; DS,

dermal sheath; migr, migratory; Prol., proliferative; Macro, macrophage; Neutro,

neutrophil; mono, monocytes; K, keratinocytes; Sup, suprabasal; Cor/cut,

cortex/cuticle; Germ, germinative; HFSC, hair follicle stem cells; IRS, inner root sheath;

S, sebaceous; UHF, upper hair follicle.

(B) Pathway information flow among active CCC pathways in resting and infectious

bite conditions. Pathway names in red indicate significant induction and in blue

significant inhibition by infectious bite (p < 0.001; as determined by Wilcoxon test[102]).

Functional categories of regulated pathways are indicated by colored boxes.

(C) Influence of infectious bite on signaling strength for the significantly induced and

inhibited pathways per cell subtype.

(D) Overview of the induced CCC immune pathways between cell subtypes.

See also **Figure S8.**

1577

1578

Figure 5. Cell-specific transcriptomic response to infectious bite

(A) Functional distribution of DEGs (adj. p-value < 0.001; |fold change| ≥ 4) within cell

subtypes. Bars indicate the numbers of regulated genes. Functional categories: SKR,

Skin Regeneration; METAB, Metabolism; RSMT, Redox-Stress-Mitochondria; RTT,

replication, transcription and translation; ncRNA, non-coding RNA; IMM/INF, immunity

and inflammation; DIVERS, other and unknown functions. N total DEGs, 2,481.

Number of up- and down-regulated DEGs are indicated in red and green, respectively.

(**B-E**) Gene Ontology (GO) biological process enrichment analysis for the top 4 most

abundant subtypes of fibroblasts (B), keratinocytes and ectodermal appendage cells

(C), myeloid cells (D) and lymphocytes (E). Subtypes are presented in an increasing

order of cell abundance. Significant GO terms encompass > 10% all DEGs within

subtypes and have p-value < 0.01. Functional categories are indicated by colored

boxes. dWAT, dermal white adipose tissue cells; IRS, inner root sheath cells; DC,

dendritic cells; DETC, dendritic epidermal T cells; ILC, innate lymphoid cells.

See also **Figure S9 and Dataset S5.**

1594

1595

Figure 6. Fibroblast response to infectious bite reduces transmission

(A) Expression of selected immune DEGs in fibroblasts.

(B) Infection of dermal fibroblast (HFF1) cells with or without mosquito saliva and

quantification of selected DEGs. hpt, hours post treatment.

(C) Regulation of *LRRC15* expression by either saliva, virus (WNV) or both in HFF1.

CTRL, mock infection. DEGs that were not regulated as in skin scRNA-seq dataset are

presented in Figure S10.

(D) Functional evaluation on WNV infection of LRRC15 in HFF1. CTRL, siRNA control.

(E) Infection levels in *LRRC15*-depleted HFF1 cells at 36 hours (h) post-infection.

(F) Evaluation of Lrrc15 function in transmission.

(G-I) Expression of Lrrc15 gene (G) and protein (H,I) in mouse skin at 48 h post-siRNA

injection. N mice, 2. Each lane represents the sample from one mouse.

(J, K) Infection level per skin biopsy (J) and pair of inguinal LNs (K) at 24 h post-

infectious bite. N mice, 4. Skin biopsies and LN from the same mouse are color-coded.

(L-N) Relative weight changes (L), clinical scores (M) and survival (N) over 14 days

post-infectious bite in mice intradermally injected with siRNA against *Lrrc15* or control

siRNA. N mice, 8.

(C, E, G, I-K) Bars and lines show means ± s.e.m. Dots indicate repeats. *, p < 0.05;

**, p < 0.01; ***, p < 0.001 by Dunett's test with CTRL or T-test. (L, M) Lines indicate

mean and shaded area ± s.e.m. p, as determined by mixed-effect ANOVA for time x

treatment interaction (L) and treatment effect (M). (N) p, as determined by Kaplan-

Meier Survival test.

See also **Figures S10-S11 and Datasets S1, S5.**

1619