**Develop a durable, memory-driven, CspZ-targeting Lyme disease vaccine by rationale**

**adjuvant selection**

Miranda McCarty[1], Sergio A. Hernández[1,2], Jill Malfetano[3], Ana Carolina Leão[4,5], Maria Jose

Villar[4,5], Xiuli Yang[6], Yi-Lin Chen[4,5], Jungsoon Lee[4,5], Zhuyun Liu[4,5], Ulrich Strych[4,5], Maria

Elena Bottazzi[4,5,7], Utpal Pal[6], Klemen, Strle[2], Wen-Hsiang Chen[4,5]*, Yi-Pin Lin[1,3]*

[1]Department of Infectious Disease and Global Health, Cummings School of Veterinary Medicine,

Tufts University, North Grafton, MA, USA; [2]Department of Molecular Biology and

Microbiology, School of Medicine, Tufts University, Boston; MA, USA. [3]Division of Infectious

Diseases, Wadsworth Center, NYSDOH, Albany, NY, USA; [4]Department of Pediatrics (Tropical

Medicine), National School of Tropical Medicine, Baylor College of Medicine, Houston, TX,

USA; [5]Texas Children's Hospital Center for Vaccine Development, Houston, TX, USA;

[6]Department of Veterinary Medicine, Virginia-Maryland Regional College of Veterinary

Medicine, University of Maryland, College Park, MD, United States; [7]Department of Biology,

Baylor University, Waco, TX, United States.

Short title: A durable, memory immunity-mediated Lyme disease vaccine

Keywords: Lyme disease, CspZ, *Borrelia*, Adjuvant, Vaccine

*correspondence: Wen-Hsiang Chen, Ph.D.
Departments of Pediatrics (Tropical Medicine), National School of Tropical
Medicine, Texas Children's Hospital Center for Vaccine Development, Baylor
College of Medicine, Houston, TX USA
Telephone: +1 832-824-0541
Email: Wen-Hsiang.Chen@bcm.edu
23
*correspondence: Yi-Pin Lin, Ph.D.
Department of Infectious Disease and Global health,
Cummings School of Veterinary Medicine, Tufts University
North Grafton, MA USA
Telephone: +1 508-887-4535
Email: Yi-Pin.Lin@tufts.edu

**ABSTRACT**

Rational adjuvant selection is a systematic approach based on adjuvant-mediated immunomodulation to identify safe vaccine regimens that enhance protective immunity. Transmitted through ticks and caused by the bacterium *Borrelia burgdorferi* (*Bb*), Lyme disease (LD) is the most common vector-borne disease in the Northern hemisphere. There are no effective vaccines, making it suitable for testing the concept of rational adjuvant selection. Here, we formulated our previously developed and effective LD vaccine antigen, CspZ-YA$_{C187S,}$ with different adjuvants suitable for human use; we analyzed the immune response by transcriptomics and tested the vaccine efficacy after *Bb* infection. We identified Alum-CpG and Alum-αGal to elicit the highest titers of CspZ-YA$_{C187S}$-dependent protective antibodies and robust levels of protection but through distinct mechanisms of immunomodulation. We demonstrated that immunization with Alum-CpG formulated CspZ-YA$_{C187S}$ provided up to nine months of protective bactericidal antibody titers, as well as recall-memory response to prevent LD after natural infection. Immunity was linked to elevated levels of IgG1 memory cells in the vaccine-triggered immune responses. This work thus identified a durable, memory immunity-driven LD vaccine, ultimately paving the road to understanding the mechanisms of rationale adjuvant selection for vaccine development.

## INTRODUCTION

Vaccines depend on the host's adaptive immune response to confer protection. An ideal vaccine offers safe, robust, and long-lasting protective immune responses. Each component of the vaccine, namely the antigen and the adjuvant, triggers distinct immune responses, but not all responses persist and/or confer protection [1, 2]. Some immune responses may even promote pathogen dissemination, exacerbate disease, and/or pose safety concerns [3]. Consequently, considerable efforts have focused on optimizing vaccine regimens to prolong protective immune responses while minimizing adverse effects [4]. One practical approach of vaccine regimen optimization is rational adjuvant selection whereby antigens and adjuvants are selected and combined based on their capacity to shape and elicit a favorable immune response [5, 6]. Broadly, adjuvants can act as either delivery systems or immune potentiators, according to their mechanism of action [7]. Delivery systems (e.g., oil-water emulsion, microparticles, or minerals such as aluminum hydroxide (Alum)) modulate antigen release to enhance durability of protection and promote antigen-specific responses [8]. Immune potentiators activate specific responses due to engagement of receptors and agonism of particular pathways (e.g., α-Galactosylceramide (αGal) for invariant natural killer T (iNKT) cells; PAM3CSK4 (PAM3) for Toll-like receptors (TLR) 1 and 2; CpG oligodeoxynucleotides (CpG) for TLR-9) [7]. Knowing which particular immune responses are required for pathogen survival or disease prevention allows for adjuvants to be selected individually or combinatorially to direct host immune responses toward pathogen clearance and mitigation of pathology. Understanding how specific adjuvant mechanisms translate into protective immunity is, therefore, central to rational adjuvant selection.

Lyme disease is the most common vector-borne infection in the Northern Hemisphere, with more than 476,000 human cases reported in the United States annually [9]. Transmitted by *Ixodes* ticks, this disease is caused by a variety of bacterial species within the *Borrelia burgdorferi* sensu lato complex (also known as *Borreliella burgdorferi* or Lyme borreliae) [1]. LYMErix, the only licensed human Lyme disease vaccine to date, was withdrawn from the market 20 years ago [10, 11]. This vaccine targeted a bacterial protein present in ticks only, OspA, or outer surface protein A, to block transmission to humans [10, 11]. A next-generation OspA-based vaccine, VLA15, is currently in clinical trials [12, 13, 14]. However, because Lyme borreliae express *ospA* predominantly in the tick midgut and downregulate *ospA* expression after transmission to mammals, to sustain high titers of anti-OspA antibodies required for protection, typically, higher vaccine doses and multiple boosters are necessary [11, 13, 15, 16, 17]. Efforts were taken to attempt in addressing such caveats through optimizing adjuvants and antigens in regimens [18, 19, 20, 21, 22, 23]. Although the tick-specific expression of *ospA* likely does not induce OspA-specific memory responses after natural infection in humans, the history of OspA vaccine development suggests Lyme disease as a tractable system for elucidating the immunological basis of how adjuvant formulations optimize protective and durable immunity.

The strategy we have pursued to overcome the challenges of Lyme disease vaccine development is to identify new vaccine antigens. Lyme borreliae produce outer surface proteins that bind host Regulators of Complement Activation (RCAs) to promote the degradation of the complement protein, C3b [24, 25, 26, 27]. This ultimately facilitates complement resistance and survival of the bacteria in circulation despite high concentrations of complement. Five of these Lyme borreliae proteins, so-called Complement Regulator Acquiring Surface Proteins (CRASPs), bind to a host RCA, Factor H (FH) [24]. Among them, CspZ (aka CRASP-2) is essential for

optimal bacterial dissemination after invasion of vertebrate hosts [20, 28], consistent with the fact

that CspZ is produced only when bacteria are in mammalian hosts, but not in ticks [29].

Furthermore, anti-CspZ IgG is constantly detectable in serologically confirmed Lyme disease

patients and infected murine models [30, 31, 32, 33], supporting CspZ as an antigen target for Lyme

disease vaccine development. However, because vaccination with wild-type CspZ has shown

limited protection in mice [20, 31, 34, 35], we have engineered a superior mutated antigen, CspZ-

$YA_{C187S}$ (CspZ-Y207A/Y211A/C187S)[20, 33, 35]. By exposing potential bactericidal epitopes near

FH-binding sites via Y207A/Y211A mutations and stabilizing overall folding via the C187S

mutation, $CspZ-YA_{C187S}$ has been shown to effectively prevent *B. burgdorferi* colonization and

joint inflammation with fewer immunization boosters[20, 33, 35]. These efforts enable us to apply

rational adjuvant selection by further identifying $CspZ-YA_{C187S}$-induced host immune responses

and linking these responses to durable and memory-mediated protection against Lyme disease.

In this study, we formulated $CspZ-YA_{C187S}$ with different adjuvants established for clinical

use and identified (i) the resulting immunomodulation of each formulation, (ii) the efficacy of

vaccination in a murine challenge model for Lyme borreliae, and (iii) the durability and memory-

inducing characteristics of the protective antibodies triggered by these formulations following

natural infection.

**RESULTS**

**CpG or αGal boost IgG titers and bactericidal activity triggered by Alum-CspZ-YA$_{C187S}$**

**vaccination**. Amongst the adjuvant used in humans HogenEsch, [36, 37], Alum(Aluminum

hydroxide/Alhydrogel) can trigger a Th2-biased immune response and antibody production and

IL-10, which can alliviate Lyme disease-associated arthritis preclinically [38]. PAM3(PAM3CSK4)

facilitates TLR2-mediated cytokines documented to promote *B. burgdorferi*-specific recognition

by the host for clearance [39]. To define vaccine formulations that elicit the highest levels of CspZ-

YA$_{C187S}$-specific IgG titers and protective antibodies, we thus immunized mice once or twice on

days 0 and 14 with CspZ-YA$_{C187S}$ formulated with Alum and PAM3. IgG titers after 14 and 28

days post initial immunization (dpii) were compared to those generated by formulating CspZ-

YA$_{C187S}$ with TiterMax Gold (TMG), an adjuvant previously used in preclinical mouse studies [20,

33, 35] (**Fig. 1A**). Anti-CspZ-YA$_{C187S}$ IgG titers were similar across all groups after one or two

immunizations (**Fig. 1B-C**). We then quantified bactericidal activity against *B. burgdorferi* strain

B31-A3 (OspC type A, the most common genotype from patients in North America) using BA$_{50}$

values, the serum dilution that kills 50% of spirochetes (**Fig. 1D-G, Table S1**). After a single

immunization, killing was detectable for the TMG formulation, but not for the Alum or PAM3

vaccines (**Fig. 1D-E, Table S1**). Following the second immunization, TMG adjuvanted CspZ-

YA$_{C187S}$ produced significantly higher BA$_{50}$ values than Alum or PAM3 formulations (**Fig. 1F-G,**

**Table S1**), indicating that Alum or PAM3 alone do not outperform the TMG benchmark.

Alum primarily functions as a delivery system for antigen formulations and has been

commonly co-formulated with other adjuvants considered "immune potentiators" to enhance

pathogen-killing efficacy [7]. We thus next evaluated whether adding immune potentiators could

enhance responses. As a one of the immune potentiators, αGal facilitates iNKT cells/B cells

production, the immune mechanisms documented to  promote Lyme borreliae clearance and

prevent Lyme disease-associated joint inflammation preclinically [40, 41]. The other immune

potentiator, CpG, is a agonist of TLR9, reported to be critical for Lyme borreliae clearance [42, 43].

We thus co-formulated Alum-CspZ-YA$_{C187S}$ with CpG or αGal  (**Fig. 1A**). Both combinations of

immune potentiators significantly increased anti-CspZ-YA$_{C187S}$ IgG titers relative to Alum alone

after one or two immunization doses (**Fig. 1B-C**). Importantly, bacterial killing was detectable

after a single immunization with Alum-CpG or Alum-αGal formulated CspZ-YA$_{C187S}$ and

indistinguishable from those with TMG formulated CspZ-YA$_{C187S}$ (**Fig. 1D-E, Table S1**) and

were significantly higher than Alum alone after the booster (**Fig. 1F-G, Table S1**). Together,

these data demonstrate that adding CpG or αGal as immune potentiators to Alum augments both

antibody titers and *B. burgdorferi*-killing .

**CpG or αGal formulations with Alum-CspZ-YA$_{C187S}$ prevent *B. burgdorferi* infection after**

**two immunizations**. After the mice were immunized with CspZ-YA$_{C187S}$ formulated with

various adjuvants, they were challenged by feeding of *B. burgdorferi* B31-A3-infected *Ixodes*

*scapularis* nymphal ticks at 35 dpii (**Fig. 1A**). Uninfected mice or TBS-inoculated and *B.*

*burgdorferi*-infected mice were included as controls. Bacterial burdens in replete nymphs were

comparable across all vaccine groups, agreeing with prior reports that mouse vaccination with

TMG-CspZ-YA$_{C187S}$ does not impact spirochetal burdens in fed nymphs (**Fig. 2A**) [33].

Seropositivity to the *B. burgdorferi* C6 peptide at 21 days post-nymph feeding (dpf) mirrored

protection: all TMG-CspZ-YA$_{C187S}$ animals remained C6-seronegative, whereas all mice in the

TBS control and Alum-formulated groups, and most of the PAM3-formulated group (5/5 mice

for TBS and Alum; 4/5 for PAM3), were C6-seropositive. Mice immunized with TBS, Alum, or

PAM3 formulations had significantly higher anti-C6 IgG titers than uninfected controls **(Fig. 2B)**.

In contrast, all mice receiving either Alum-CpG or Alum-αGal formulation were seronegative to

the C6 peptide, similar to uninfected animals **(Fig. 2B)**. Consistent with serology, bacterial

burdens at the tick bite site, bladder, heart, and knee joints were elevated in the TBS, Alum-, and

PAM3-formulated groups but were undetectable in uninfected controls, TMG, Alum-CpG, and

Alum-αGal groups (**Fig. 2C-F**). Thus, CpG and αGal adjuvants convert Alum-CspZ-YA$_{C187S}$

into an efficacious vaccine  that blocks *B. burgdorferi* tissue colonization and seroconversion

preclinically.

We next determined the severity of *B. burgdorferi*-triggered inflammation in the tibiotarsal

(ankle) joints of mice at 21 dpf (**Fig. 1A**). Histopathology revealed dense mononuclear infiltrates

in the muscles, tendons, and connective tissues from all five mice immunized with TBS or

Alum-, and most (4/5) PAM3-immunizeed mice (arrows in representative images, **Fig. 2G**).

Inflammation scores in these vaccination groups were significantly elevated versus uninfected

controls (**Fig. 2H**). Conversely, all five mice immunized with CspZ-YA$_{C187S}$ formulated with

Alum-CpG or Alum-αGal displayed no pathologic joint inflammation and had inflammatory

scores comparable to uninfected controls and the protective TMG adjuvant formulation (**Fig.**

**2G-H**). Overall, these data show that the addition of CpG or αGal to Alum-CspZ-YA$_{C187S}$

vaccination yields robust protection after two immunizations, preventing seroconversion, tissue

colonization, and Lyme disease-associated joint inflammation, whereas PAM3 or Alum does not.

**Alum-CpG  and Alum-αGal formulations drive distinct immune transcriptional programs**

**after CspZ-YA$_{C187S}$-vaccination.** We next compared the transcriptomic programs induced by

adding CpG or αGal to Alum-formulated CspZ-YA$_{C187S.}$ RNA was isolated from whole spleens

of mice 14 days after the second immunization (28 dpii) and profiled for RNA sequencing

(RNA-seq) (**Fig. 1A**). Relative to TBS controls, mice vaccinated with Alum-CpG-formulated

CspZ-YA$_{C187S}$ produced fewer differentially expressed genes (DEGs) whereas Alum-αGal-

formulated CspZ-YA$_{C187S}$ yielded more DEGs overall, suggesting distinct gene expression

profiles induced by CpG or αGal in Alum-CspZ-YA$_{C187S}$ immunized mice (**Fig. S1, Dataset 1**).

Using Alum-CspZ-YA$_{C187S}$ mice as the baseline, Alum-CpG vs. Alum identified 8 upregulated

DEGs, whereas Alum-αGal versus Alum identified 100 upregulated DEGs. Only three DEGs

(2.9%) were shared between these upregulated sets (**Fig. 3A**). In contrast, for downregulated

DEGs, Alum-CpG and Alum-αGal yielded 28 and 32 downregulated genes, respectively, with no

overlap (**Fig. 3B**). In the direct transcriptomic comparison of the Alum-CpG vs. Alum groups,

we observed the upregulation of genes involved mostly in immunoglobulin production and

selected inflammatory mediators (**Fig. 3C-D, Table S2**). In Alum-αGal vs. Alum, we observed

upregulation of immunoglobulin genes together with antigen presentation/recognition, Th-1 cell

modulation, and leukocyte trafficking, accompanied by reduced expression of several anti-

inflammatory genes (**Fig. 3E-F, Table S3**). Taken together, these results suggest that CpG and

αGal remodel splenic transcriptomics in qualitatively different ways when formulated with

Alum-CspZ-YA$_{C187S}$.

We next conducted pathway/over-representation analyses modulated by CpG or αGal when

added to Alum-CpG-CspZ-YA$_{C187S}$-vaccination. Given the modest DEG lists, pathway/term

over-representation tests were underpowered. We therefore utilized Gene Set Enrichment

Analysis (GSEA) on the complete ranked gene lists, enabling detection of modest but relevant

pathway changes with statistical control (False Discovery Rate (FDR)). In the comparisons vs.

Alum, immune-related functions were prominent among the top 10 significantly enriched

pathways in both formulations (3 of the top 10 in Alum-CpG; 4/10 in Alum- αGal) (bold

pathways in **Fig. S2**). We thus selected all immune-related pathways significantly upregulated in

the mice vaccinated with CspZ-YA$_{C187S}$ formulated with Alum-CpG or Alum-αGal (vs. Alum-

only) and characterized their functions. Most of the highest-ranked immune terms in both groups

(6/14 in Alum-CpG; 8/19 in Alum-αGal) mapped to adaptive B and T cell functions (**Fig. S3**),

aligning with the significantly greater levels of IgG titers and enhanced bactericidal activity seen for the Alum-CpG and Alum-αGal formulations of CspZ-YA$_{C187S}$ (**Fig. 1B-G**). To further differentiate the immune transcriptional programs induced by the formulations containing Alum-CpG and Alum-αGal, we directly compared the pathways upregulated by the two adjuvants relative to each other using GSEA. Alum-CpG (enriched relative to Alum-αGal) favored many interferon (IFN)-related pathways, including IFN-α responses, positive regulation of IFN-β production, NK-cell IFN-κ signatures, and an antiviral state overall (see bold pathways in **Fig. 3G**). In contrast, Alum-αGal (enriched relative to Alum-CpG) emphasized antigen processing and presentation, as well as antigen presenting cells (APC)/lymphocyte crosstalk pathways (see bold pathways in **Fig. 3H**). In sum, CpG skews Alum-CspZ-YA$_{C187S}$ towards an antiviral/interferon-modulated program, whereas Alum-αGal induces APC licensing, antigen processing, and lymphocyte priming processes. These results suggest two distinct routes of immunity for the two formulations that lead to a robust, protective humoral responses after CspZ-YA$_{C187S}$ vaccination.

**Alum-CpG vs. Alum-αGal impact cytokine secretion in *ex vivo* splenocytes from CspZ-YA$_{C187S}$ immunized mice but converge on memory B-cell activation.** We isolated splenocytes from mice at 7-days after the second inoculation with TBS (control), or CspZ-YA$_{C187S}$ formulated with Alum, Alum-αGal or Alum-CpG. Splenocytes were restimulated *ex vivo* with CspZ-YA$_{C187S}$, and cytokine supernatant concentrations were quantified. Across all measured mediators (IL-10, TNFα, IL-2, IL-6, IL-4, IFN-γ at two doses), splenocytes from Alum-αGal-vaccinated mice produced significantly higher levels than TBS controls (**Fig. 4A-G**). In contrast, vaccination with Alum-CpG adjuvant selectively increased only IFN-γ levels, while other

cytokines were not significantly different (**Fig. 4F and G**), supporting our transcriptomic

findings of Alum-CpG amplifying a marked IFN-specific immune program, whereas Alum-

αGal-CspZ-YA$_{C187S}$ elicits a broader cytokine profile.

To delineate the responders among splenocytes to the CspZ-YA$_{C187S}$ vaccination with

different tested adjuvants, we profiled splenocytes by flow cytometry after antigen (CspZ-

YA$_{C187S}$) restimulation. Germinal center (GC) B cells and T follicular helper (Tfh) cells were

significantly increased in mice immunized with Alum-CpG or Alum-αGal but not in mice

immunized with Alum alone, compared to TBS-inoculated mice (**Fig. 4H-I**). Total memory B

cell populations were similar among the tested groups of splenocytes, but IgG1+ memory B cells

were significantly enriched in both Alum-CpG and Alum-αGal cohorts compared to the TBS-

inoculated mice (**Fig. 4K**). These findings indicate that although CpG and αGal diverge in the

acute cytokine profile they elicit, both adjuvants promote GC B cells/Tfh engagement, and

expansion of IgG1+ memory B cells, consistent with the enhanced antibody titers and protective

phenotypes reported earlier.

**Vaccination with Alum-CpG-CspZ-YA$_{C187S}$ elicits durable, bactericidal antibodies for up**

**to nine months.** The abovementioned results identified Alum-CpG and Alum-αGal as the lead

adjuvants to formulate with CspZ-YA$_{C187S}$. However, CpG, but not αGal, has been FDA-

approved and used in licensed human vaccines [44, 45, 46, 47], we thus prioritized the Alum-CpG

formulation and sought to determine the durability of protective antibodies triggered by CspZ-

YA$_{C187S}$ vaccination. Mice received either one dose (at day 0) or two doses (at day 0 and 14dpii).

Serum was collected longitudinally at different days post last immunization (dpli) to quantify

antigen-specific IgG titers against OspA and CspZ-YA$_{C187S}$, as well as bactericidal activity (BA$_{50}$)

(**Fig. 5A**). TBS-inoculated mice and mice immunized three times with TMG-OspA were

included as controls (**Fig. 5A**). We detected anti-OspA IgG only in the TMG-OspA group, with

antibodies persisting until 182 dpli. No OspA reactivity was observed in any mice inoculated

with Alum-CpG-CspZ-YA$_{C187S}$ or TBS (**Fig. 5B and S4**). Conversely, TMG-OspA or TBS-

inoculated mice did not exhibit detectable anti-CspZ-YA$_{C187S}$ IgG, but these specific antibodies

were present in high titers in Alum-CpG-CspZ-YA$_{C187S}$ (**Fig. 5C and S5**). A two-dose

immunization regimen with CspZ-YA$_{C187S}$ yielded signifianctly higher titers at all time points

and extended seropositivity to 238 dpli, compared with 210 dpli after a single dose (**Fig. 5C and**

**S5**). No bactericidal activity was detected in TBS-inoculated mice. The TMG-OspA immunized

mice maintained measurable bactericidal antibody titers (BA$_{50}$) until 182 dpli (**Fig. 5D and S6**).

Importantly, sera from Alum-CpG-CspZ-YA$_{C187S}$ immunized mice showed durable bactericidal

activity up until 210 dpli after one dose and up to 238 dpli after two doses, mirroring the IgG

durability (**Fig. 5D and S6**). These results showed the capacity of Alum-CpG-CspZ-YA$_{C187S}$ to

induce dose-responsive, long-lived humoral immunity with bactericidal function lasting up to

nine months, supporting this formulation as the lead vaccine formulation.

**Natural infection rapidly recalled bactericidal anti-CspZ antibodies and conferred**

**protection in Alum-CpG-CspZ-YA$_{C187S}$-vaccinated mice.** We next determined whether

natural infection could recall CspZ-targeting immunity once vaccine-induced CspZ-YA$_{C187S}$ IgG

waned below detection. Mice previously immunized with Alum-CpG-CspZ-YA$_{C187S}$ (one or two

doses), TMG-OspA (three doses), or TBS were held until 273 dpli, when anti-CspZ-YA$_{C187S}$ IgG

and bactericidal antibody titers (BA$_{50}$) were undetectable (**Fig. 5A**). Animals were then exposed

to *B. burgdorferi* by infected nymphal tick feed and sera were collected at 4, 7, 10, 14, and 21

dpf to assess antibody recall (**Fig. 5A**). As expected, anti-OspA IgG remained negligible across

all immunization groups after tick feeding, including mice who received TMG-OspA (**Fig. 6A,**

**S7**), consistent with *ospA* downregulation in vertebrate hosts [15, 16]. In contrast, Alum-CpG-CspZ-

$YA_{C187S}$ vaccination primed a rapid recall response, as anti-$CspZ$-$YA_{C187S}$ IgG rose significantly

compared to uninfected controls by 4-7 dpf and remained elevated through all timepoints (**Fig.**

**6B, S8**). TMG-OspA and TBS groups lacked early anti-$CspZ$-$YA_{C187S}$ increases, and showed

only modest responses by 10 dpf, in line with detectable CspZ-targeting antibodies reported at

later stages during natural infection (*de novo* priming) [20, 29, 30, 32, 33](**Fig. 6B, S8**). Antibody

isotyping at 7 dpf, done to determine early protective responders, revealed a broad recall of IgG

subtypes (IgG1, IgG2a, IgG2b, IgG3) (**Fig. 6C**). Among these isotypes, IgG1 was found to be

the predominant subclass (**Fig. 6C**), mirroring the increase in IgG1+ memory B cells observed

post-vaccination (**Fig. 4K**). Further, early bactericidal function was unique to sera from Alum-

CpG-CspZ-$YA_{C187S}$ recipients, as $BA_{50}$ became detectable at 4-7 dpf and increased thereafter,

whereas other groups lacked bactericidal activity until $\geq$10 dpf (**Fig. 6D, S9, Table S4**). These

results suggest an early emergence of memory immunity-driven, bactericidal antibodies after

natural infection in the mice immunized with Alum-CpG-formulated CspZ-$YA_{C187S.}$

We next tested if this rapid recall via CspZ-$YA_{C187S}$ vaccination translated into protection by

determining anti-C6 seropositivity, *B. burgdorferi* burdens in tissues, and ankle joint

inflammation histologically at 21 dpf (**Fig. 5A**). Tick bacterial burdens in fed nymphs did not

differ among groups, as expected (**Fig. 6E**), but host outcomes differed sharply. At 21 dpf, all

TBS- or TMG-OspA inoculated mice were C6-seropositive and harbored significantly higher *B.*

*burgdorferi* burdens in skin (bite site), bladder, heart, and knee, compared to uninfected mice

(**Fig. 6F-J**). These mice also displayed extensive mononuclear infiltrates in ankle joints and

significantly elevated histologic inflammation scores relative to uninfected controls (**Fig. 6K-L**).

In contrast, one mouse out of five immunized with a single dose of Alum-CpG-CspZ-YA$_{C187S}$

and none in the two-dose group became C6-seropositive, with bacterial tissue burdens

indistinguishable from those of uninfected mice (**Fig. 6F-J**). Similarly, one animal (single dose)

and none from the two-dose group developed apparent joint inflammation, resulting in

histopathological scores comparable to those from uninfected mice (**Fig. 6J-K**). Taken together,

these results link a durable, memory B-cell antibody response primed by Alum-CpG formulated

CspZ-YA$_{C187S}$ vaccination and rapidly recalled by natural infection, to the prevention of C6-

seropositivity, bacterial colonization, and *B. burgdorferi*-associated joint pathology in the murine

Lyme disease model.

**DISCUSSION**

Adjuvants are critical in vaccines to enhance magnitude, quality, and durability of the

protective immunity against infectious diseases. For many vaccine prototypes, rational adjuvant

selections that utilize established immune potentiators or delivery systems to trigger desirable

immune pathways while avoiding unwanted host responses can improve immunization safety

and efficacy [5, 6]. Guided by this principle, we evaluated how different immune potentiators

layered onto a common delivery system (Alum) direct diverse immune responses elicited by a

structure-guided CspZ vaccine (CspZ-YA$_{C187S}$). Lyme borreliae are positioned as a tractable

model pathogen for vaccine development, given the numerous efforts invested to elucidate host

responses that enable pathogen clearance and alleviate the development of manifestations [25].

These data provide the foundation for applying the concept of rational adjuvant selection to elicit

disease-resolving immune responses in the development of Lyme disease vaccines. As the most

commonly used adjuvant in human vaccines, Alum activates the local innate immune system

through pattern recognition receptors (PRRs), which matures antigen presenting cells (APCs) for

appropriate presentation of the vaccine antigen. This process layered to immune potentiators

ultimately creates a unique inflammatory niche that produces CspZ-YA$_{C187S}$ antibodies to drive

more robust bactericidal responses, while reducing the number of boosters required [48, 49, 50].

However, CspZ-YA$_{C187S}$ formulated with Alum alone did not provide protection in our study.

Additionally, it is well-known that TLR1/2 recognize Lyme borreliae, triggering inflammatory

pathways favoring spirochete elimination [51, 52, 53, 54, 55] and that PAM3 is a TLR1/2 agonist [56, 57].

Yet, similar to Alum, PAM3-formulated CspZ-YA$_{C187S}$ also failed to protect mice from *B.*

*burgdorferi* colonization or joint inflammation after immunization. These findings likely reflect

insufficient priming of protective humoral and cellular immune programs by Alum or PAM3

adjuvants alone. We thus tested immune potentiators with defined mechanisms co-formulated

with Alum to bias an immune response towards protective pathways that drive beneficial

adaptive immune responses. One of the selected immune potentiators is CpG, a TLR9 agonist

known to trigger APC signaling, drives activation of immune cells and promotes Th1 immune

responses, and proliferation and maturation of B cells [58, 59, 60, 61]. Another immune potentiator, α-

galactosylceramide (αGal), is a synthetic glycolipid that can be presented by CD1d, which is

widely expressed on hematopoietic and non-hematopoietic cells, and subsequently recognized by

iNKT cells [62, 63]. Consistent with TLR9 ligation and iNKT cells in facilitating Lyme borreliae

clearance [40, 41, 42, 64, 65, 66], both adjuvant combinations (Alum-CpG and Alum-αGal) protected

against *B. burgdorferi* infection in murine models following vaccination with CspZ-YA$_{C187S}$.

Additionally, mice immunized with each of these two formulations developed significantly

higher levels of anti-CspZ-YA$_{C187S}$ IgG and borreliacidal activities than animals vaccinated with

Alum-CspZ-YA$_{C187S}$ alone, supporting well-documented engagement of downstream B cell

responses by TLR9 and iNKT cell pathways [67, 68, 69, 70]. Our results thus suggest similar favorable

immune responses (i.e., higher extents of bactericidal antibodies) could be boosted by distinct

pathways using diverse immune potentiators, underscoring the need to identify the unique

mechanisms of potentiator-mediated protective immunity.

In this study, transcriptomic profiling of splenocytes provided additional mechanistic insights

into specific adaptive immune responses of the immune simulators. Compared to TBS-inoculated

mice, both Alum-CpG and Alum-αGal-formulated CspZ-YA$_{C187S}$ upregulated genes associated

with B cell and T cell immunity, including the activation of signaling, immunoglobulin

production, and effector functions with associated cytokines. This is in line with previous

findings of the requirement of durable T-dependent B cell immune responses for effective Lyme

borreliae clearance [49, 71]. Furthermore, the enhanced expression of both T and B cell associated

genes can be directly attributed to CpG and αGal priming, as these genes remained significantly

upregulated relative to mice immunized with CspZ-YA$_{C187S}$ formulated with Alum alone.

However, Alum-αGal formulated CspZ-YA$_{C187S}$ uniquely permitted the enrichment of NK cell-

related pathways, agreeing with αGal as an iNKT cell agonist [60, 61, 62, 63]. Alum-αGal also

preferentially enriched antigen-processing and presentation programs (e.g., ER phagosome

pathway and MHC assembly/loading) in APCs (dendritic cells), together with the

aforementioned iNKT/NK-cell-linked terms. These results suggest the possibility for the role of

αGal in promoting more efficient APC licensing and crosstalk with T cells, thereby broadening

adaptive immune responses for Lyme borreliae clearance. Further, we found that immunization

with Alum-CpG formulated CspZ-YA$_{C187S}$ demonstrated an interferon-centric signature.

Activation of IFN-specific pathways was corroborated with targeted proteomics using

splenocytes from mice immunized with Alum-CpG formulated CspZ-YA$_{C187S}$. These consistent

results agree with published data on TLR9-triggered type IFN (IFN-α and IFN-β) signaling [72].

Together with type II IFNs (i.e., IFN-γ) and IFN-γ-induced cytokines being associated with

lower *B. burgdorferi* burdens in humans and the phenotypes of *B. burgdorferi* clearance in the

murine models [66, 73, 74, 75], our findings provide possible mechanisms for Alum-CpG-activated

IFN immunity in protection against Lyme borreliae colonization. Despite these upstream

differences between the formulations of Alum-CpG and Alum-αGal, both regimens converged

on expansion of germinal center B cells, T follicular helper cells, and IgG1+ memory B cells, all

aligning with increased antibody titers and serum bactericidal activity. Moreover, TLRs,

including TLR9, signal through MyD88, which can result in a broad spectrum of cytokine

secretion [76, 77]. However, we did not observe a lot of pathways other than those IFN-associated

be significantly enriched by Alum-CpG formulated CspZ-YA$_{C187S}$ when compared to Alum-

αGal-CspZ-YA$_{C187S}$, suggesting the possibility of antigen (CspZ-YA$_{C187S}$)-adjuvant interplay in

immunomodulation. This is consistent with many Lyme borreliae lipoproteins reported as having

immunomodulatory functions [34, 51, 78, 79]. Overall, the findings here provide mechanistic

information on how adjuvants can skew host protective immune responses.

The induction of immune memory is critical for vaccine development as the capacity and

durability of these responses impact the frequency of immunization needed to be effective [80]. For

Lyme disease vaccines, the ideal response would include the capability to trigger long-lasting

memory after immunization and rapid emergence of bactericidal antibodies upon natural

infection [81]. However, the currently developed OspA-based vaccine blocks the transmission of

the *B. burgdorferi* to the host but lack the ability to trigger persisting OspA-targeted antibodies

after several months (unless specific formulations were used [10, 11, 23]). Additionally, in support of

the downregulation of  OspA after *B. burgdorferi* invades vertebrate hosts[20, 28], we found no

induction of OspA-specific IgG in OspA-immunized mice after natural infection. Unlike OspA,

Alum☐CpG–CspZ☐YA$_{C187S}$ generated CspZ☐specific IgG and borreliacidal antibodies for up to

nine months in murine models, with higher magnitude and longer persistence after two doses

than one. This persistence of CspZ IgG exceeds what is typically achieved with OspA-based

vaccine, suggesting that CspZ-targeted vaccines may provide longer lasting immunity and reduce

the numbers of required booster doses. We also found that CspZ-targeted responses were readily

recalled, with Alum-CpG-primed animals mounting a rapid memory response detectable as early

as 4-days after natural infection. Such rapid responses were accompanied by a rise in functional

bactericidal activity, protection from tissue colonization, seroconversion, and joint inflammation.

Antibody isotype and flow cytometry analyses point to a mechanistic link between vaccine-

primed memory B cell responses and protection [82]. The dominance of IgG1 at recall and the

expansion of IgG1+ memory B cells after immunization with Alum-CpG formulated CspZ-

YA$_{C187S}$ support a model in which early reactivation of class-switched memory, likely aided by

Alum-CpG-driven interferon signaling and efficient T-cell help, delivers complement-fixing

antibodies. This posits IgG1 as the major contributor of CspZ-YA$_{C187S}$-primed memory

immunity which confers protection. IgG isotype differences against *B. burgdorferi* antigens have

been linked to infection vs. protection after *B. burgdorferi* invasion [49, 83]. For some antigens (i.e.,

OspC), IgG1 has been associated with protection, consistent with this subclass preferentially

triggering classical complement pathway-mediated pathogen killing, which has been documented

in Lyme borreliae clearance [84, 85]. Thus, our work supports the role of IgG1 in Lyme disease

protection, not only induced by vaccines but also in memory immune responses following

natural infection. Overall, IgG1-mediated protective responses from CspZ-YA$_{C187S,}$ or other

previously identified Lyme borreliae vaccine antigens, could be the determinant of protection

upon natural infection. Further examination of this possibility has the potential to aid

development of more precise antibody-based protection in memory immune responses induced

by Lyme disease vaccines.

We also showed that unvaccinated mice or mice immunized with other regimens that do not

confer protection to natural infection (i.e., TMG-OspA) also develop anti-CspZ-YA$_{C187S}$ IgG.

This result is consistent with the elevated titers of CspZ-specific IgG in Lyme disease patients

and mice colonized by Lyme borreliae [30, 31, 32, 33]. However, mice vaccinated with CspZ-YA$_{C187S}$

had an earlier onset of CspZ-specific IgG antibodies, compared to the unvaccinated or TMG-

OspA-immunized mice (4 dpf vs. 10 dpf). This result highlights the importance of a rapid recall

of anti-CspZ-YA$_{C187S}$ IgG for an efficacious CspZ-targeted vaccine. In fact, Lyme borreliae

infection kinetics begin with the pathogens transiently being in the bloodstream, followed by fast

migration to distal tissues [86, 87]. Such a unique progression during infection is largely due to the

vulnerability of Lyme borreliae while in the bloodstream, in which numerous host mechanisms

are present to eradicate spirochetes [25, 88]. Although Lyme borreliae develop several antibody-

evasion strategies immediately, those strategies do not mature until later in the infection [89, 90, 91].

Therefore, the ability to generate bactericidal antibodies within 4 days during natural infection, at

a time when spirochetes are transiently blood-borne and most susceptible to humoral clearance,

is expected to be critical in CspZ-YA$_{C187S}$ vaccine efficiency. Taken together, our study tested

the concept of various adjuvant formulations in eliciting beneficial memory B cell immune

responsesto CspZ-YA$_{C187S}$. The optimization of adjuvant selection identified a formulation that

can facilitate a long-lasting, memory-driven Lyme disease vaccine. Overall, our work examines

the concept of rational adjuvant selection for vaccine development and elucidates the vaccine-

triggered, adjuvant-influenced immune responses, which would benefit the development of other

infectious disease vaccines.

**MATERIALS AND METHODS**

**Ethics Statement.** All mouse experiments were performed in strict accordance with all

provisions of the Animal Welfare Act, the Guide for the Care and Use of Laboratory Animals,

and the PHS Policy on Humane Care and Use of Laboratory Animals. The protocols, docket

numbers 22-451 and G2024-07, were approved by the Institutional Animal Care and Use Agency

of Wadsworth Center, New York State Department of Health and Tufts University, respectively.

All efforts were made to minimize animal suffering.

**Mouse, ticks, and bacterial strains.** Four-week-old, female C3H/HeN mice were purchased

from Charles River (Wilmington, MA, USA). Although such an age of the mice has not reached

sexual maturity, the under development of immune system in this age of mice would allow such

mice to be more susceptible to Lyme borreliae infection, increasing the signal to noise ratio of

the readout. That will also provide more stringent criteria to define the protectivity. BALB/c C3-

deficient mice were from in-house breeding colonies [92] and *Ixodes scapularis* tick larvae were

obtained from BEI Resources (Manassas, VA). *Escherichia coli* strain BL21(DE3) and

derivatives were grown at 37°C or other appropriate temperatures in Luria-Bertani broth or agar,

supplemented with kanamycin (50μg/mL). *B. burgdorferi* strain B31-A3 were grown at 33°C in

BSK II complete medium [93]. Cultures of *B. burgdorferi* B31-A3 were tested with PCR to ensure

a full plasmid profile [94, 95].

**Purification of OspA and CspZ-YA$_{C187S}$.** The purification of hexahistidine-tagged lipidated OspA and untagged CspZ-YA$_{C187S}$ was executed as described [33, 96, 97]. The *E. coli* BL21(DE3) strains producing OspA and CspZ-YA$_{C187S}$ are listed in **Table S6**. OspA was lipidaded as lipidation is required for recombinant OspA proteins to protect mice from Lyme disease infection[97, 98].

**Mouse immunization and infection.** C3H/HeN mice were immunized as described, with slight modifications [33]. Twenty-five µg of untagged CspZ-YA$_{C187S}$ or hexahistine-tagged lipidated OspA (control) in 50 µl of Tris-buffered saline (TBS) buffer was thoroughly mixed with 50 µl TiterMax Gold adjuvant (Norcross, GA, USA), resulting 100µl inoculum. Twenty-five µg of tagged CspZ-YA$_{C187S}$ was also formulated with PAM3 (10 µg, PAM3CSK4, InvivoGen, CA, USA), Alum (400 µg, Alhydrogel, Corda, NJ, USA), Alum (400 µg) mixed with CpG (20 µg, CpG1826, InvivoGen, CA, USA), or Alum (400 µg) mixed with αGal (1 µg, α-galactosylceramide, Avanti, AL, USA) in 50 µl of **TBS** with 0.05% Pluronic F-127 and 7% sucrose (TBS). In addition, 50 µl of TBS only was included as a control. The inoculum was introduced into C3H/HeN mice subcutaneously once at 0- or twice at 0- and 14-days post initial immunization (**Fig. 1A and 4A**). Blood was collected via submandibular bleeding at 14-days post last immunization (dpli) (**Fig. 1A**) or 14, 42, 70, 98, 126, 154, 182, 210, 238, and 273 dpli (**Fig. 5A**). Sera obtained from the blood were used to determine the IgG titers of CspZ-YA$_{C187S}$ or OspA, and the bactericidal activities, described in the section "ELISAs" and "Bactericidal assays", respectively. At 21 dpli (35 days post initial immunization shown in **Fig. 1A**) or 273 dpli (**Fig. 5A**), *B. burgdorferi* B31-A3-infected flat nymphs were placed in a chamber on C3H/HeN mice as described [34]. Five nymphs were allowed to feed to repletion on each mouse,

and a subset of nymphs was collected pre- and post-feeding. At 21 days post nymph feeding

(dpf), tick bite sites of skin, bladder, knees, and heart were collected to determine the bacterial

burdens, and ankles were also collected at 21 dpli to determine the severity of arthritis described

in the section "Quantification of spirochete burdens and histological analysis of arthritis." At this

time point, blood was also collected via cardiac puncture bleeding to isolate sera for the

determination of seropositivity described in the section "ELISAs."

**ELISAs.** The determination of the titers of anti-OspA, CspZ-YA$_{C187S}$ IgG, C6 IgG or the IgG

isotypes (IgG1, IgG2a, IgG2b, and IgG3) in the serum samples was as described, with

modifications [20, 33, 81]. Specifically for anti-C6 IgG, this methodology has been commonly used

for human Lyme disease diagnosis [99]. One µg of untagged and lipidated OspA, untagged CspZ-

YA$_{C187S}$, or C6 peptides was coated on ELISA plate wells as described [20]. The procedures

following the protein coating are as described previously [20]. For each serum sample, the

maximum slope of optical density/minute of all the dilutions of the serum samples was

multiplied by the respective dilution factor, and the greatest value was used as arbitrary unit (AU)

to represent the antibody titers for the experiment to obtain anti-OspA, -CspZ-YA$_{C187S}$, or -C6

IgG, or IgG1, IgG2a, IgG2b, and IgG3 against CspZ-YA$_{C187S}$. For anti-C6 IgG, the seropositive

mice were defined as the mice with the serum samples yielding a value greater than the threshold,

the mean plus 1.5-fold standard deviation of the IgG values derived from the uninfected mice.

**Borreliacidal assays.** The ability of serum samples to eradicate *B. burgdorferi* B31-A3 was

determined as described with modifications [20, 33, 35]. Briefly, the sera collected from mice

immunized with lipidated OspA or CspZ-YA$_{C187S}$ under different adjuvant formulations at

different immunization frequency were heat-treated to inactivate complement. The rest

procedure was performed in the similar fashion as described [20, 35]. Surviving spirochetes at 24 h

of incubation with Guinea pig sera (Sigma) and different dilution rate of serum samples were

quantified by directly counting the motile spirochetes using dark-field microscopy and expressed

as the proportion of serum-treated to untreated Lyme borreliae. The 50% borreliacidal activities

($BA_{50}$), representing the serum dilution rate that kills 50% of spirochetes, was calculated using

dose-response stimulation fitting in GraphPad Prism 9.3.1.

**Quantification of spirochete burdens and histological analysis of arthritis.** DNA was

extracted from the indicated mouse tissues to determine the bacterial burdens using quantitative

PCR analysis as described [20, 33]. Basically, the forward and reverse primers with the sequences as

GTGGATCTATTGTATTAGATGAGGCTCTCG                                                                          and

GCCAAAGTTCTGCAACATTAACACCTAAAG, respectively, were used to amplify the *recA*

gene of *B. burgdorferi* strain B31-A3. The number of *recA* copies was calculated by establishing

a threshold cycle (Cq) standard curve of a known number of *recA* gene extracted from strain

B31-A3, and burdens were normalized to 100 ng of total DNA from mouse tissues or presented

as the number of bacteria per tick for *B. burgdorferi* in ticks. The histological analysis was

applied to ankles for Lyme disease-associated arthritis (**Fig. 3G**), as described [20, 33]. Basically,

images were scored based on the severity of the inflammation as 0 (no inflammation), 1 (mild

inflammation with less than two small foci of infiltration), 2 (moderate inflammation with two or

more foci of infiltration), or 3 (severe inflammation with focal and diffuse infiltration covering a

large area).

**mRNA Extractions for RNASeq analyses.** Spleens were placed immediately on dry ice and

stored at –80 C° and then submitted for RNA extraction and RNAseq analysis by GENEWIZ

(Waltham, MA). Turbo Capture mRNA Kits (Qiagen) were used for total RNA extraction of 50

mg of tissues. A drill and polypropylene pestles were used to lyse TRIzol® suspended tissues in

Eppendorf Tubes, followed by the treatment of DNAse I as described in the vendor's manuals.

Elutions were prepared in DNAase and RNAase free water, and the quality of RNA was

measured using the Nanodrop Eight Spectrophotometer (ThermoFisher Scientific), targeting a

260/280 ratio of 1.95-2.04. RINe (RNA Integrity Number) values greater than 6.5 with strong

indications of robust 18S and 28S (28S/18S) peaks, with minimal evidence of degraded small

RNA at the lower threshold range, were submitted for further processing. The RNA samples

were processed for mRNA enrichment using Dynabead's mRNA Purification Kit per vendor's

manual (ThermoFisher). The mRNA sequencing was then carried out by polyA selection using

Illumina HiSeq, PE 2x150 (150 bp paired end).

**RNASeq analysis** After the sequencing, the reads were trimmed via Trimmomatic v.0.36, and

mapped to the *Mus musculus* strain C3H/HeJ reference genome via ENSEMBL using the STAR

aligner v.2.5.2b. The gene hit counts (calculation of reads/gene/sample) were determined using

feature counts from the Subread package v.1.5.2. The results from Salmon were imported into R

v.4.2.3 with tximport and a DESeq2 object was created with the DESeq dataset from tximport

function. Differential expression analyses were performed by DESeq2 with minreplicates for

replace set to 30. The identification of DEGs was shown in **Dataset S1**. RNASeq results from

the spleens of mice immunized with CspZ-YA$_{C187S}$ formulated with Alum, Alum-CpG, or Alum-

αGal were first compared with those from the control (TBS-inoculated mice). Additionally, the

results from the spleens of mice vaccinated with Alum-CpG or Alum-αGal were compared with

those from mice injected with Alum. VolcaNoseR was used to plot and explore differentially

expressed genes (DEGs) in volcano plot format [100]. Adjusted p-values ≤ 0.05 and log2 fold

changes ≥ 0.58 or ≤ -0.58 were used as the initial cutoffs defining differentially expressed genes

(DEG). The number shared and unique DEGs among the mice immunized with CspZ-YA$_{C187S}$

formulated Alum, Alum-CpG, or Alum-αGal was visualized as Venn diagram with Venny 2.1 [101].

Heat map analysis was performed using Morpheus [102]. The Pearson correlation was used to

measure dissimilarities between individuals for the heat map's hierarchical clustering [102].

Complete ranked gene lists were analyzed with GSEA (v4.4.0;

http://software.broadinstitute.org/gsea/index.jsp), using 1,000 gene-set permutations,

minimum/maximum gene-set sizes of 15/500, respectively [103]. We queried the following mouse

MSigDB collections (v2025.1): M2 (curated pathways), M5 (ontology), M7 (immunologic

signatures), and M8 (cell-type signatures). Gene sets were considered enriched at FDR q <0.25

with normalized enrichment scores (NES) |NES| ≥ 1.5. The NES sign indicates direction of

enrichment (positive = comparator group, negative = baseline).

**Multiplex analysis of cytokines by Luminex.** Splenocytes from mice immunized with CspZ-

YA$_{C187S}$ formulated with each of the tested adjuvants were prepared and analyzed by Luminex,

as described previously [104]. Spleens were rinsed and homogenized in sterile PBS buffer, using a

gentleMACS Dissociator (Miltenyi Biotech, Charlestown, MA) and treated with ACK lysis

buffer (Lonza, Portsmouth, NH) to remove red blood cells. The suspension was then diluted 5-

fold with complete Roswell Park Memorial Institute (RPMI) 1640 medium (cRPMI, Sigma

Aldrich) supplemented with 10% fetal bovine serum (FBS, Sigma Aldrich), 1× penicillin-

streptomycin (Pen-Strep, Sigma Aldrich), and L-glutamine (Sigma Aldrich), followed by

centrifugation at 300 × g for 5 min. Splenocytes were resuspended in 5 mL of cRPMI, filtered

through 40 µm strainers (BD Biosciences, San Jose, CA) and counted using acridine orange–

propidium iodide (AOPI, Fisher Scientific) staining on a Cellometer Auto 2000 (Nexcelom

Bioscience, Lawrence, MA). For stimulation assays, viable splenocytes ($1 \times 10^6$) were incubated

for each sample in a 96-well non-tissue culture plate and re-stimulated with different

concentrations of CspZ-YA$_{C187S}$ or cRPMI (control). Splenocytes were incubated at 37°C in 5%

CO$_2$ and cell-free supernatants were collected after 48-h after re-stimulation. Cytokines levels of

IL-10, TNF-a, IL-2, IL-6, IL-4, and IFN-g in supernatants were measured by a Luminex-based

assay using Milliplex MAP Mouse Th 17 Magnetic Bead Panel (Millipore Sigma), as previously

described [105]. Cytokine responses were calculated by subtracting values obtained from

splenocytes restimulated with cRPMI alone from those restimulated with CspZ-YA$_{C187S}$.

**Flow Cytometry of splenocytes** Splenocytes from the section "Multiplex analysis of cytokines

by Luminex" were washed with 1X PBS, and stained with Live/Dead NIR fixable viability dye

(ThermoFisher). The viable population of the cells were incubated with purified rat anti-mouse

CD16/CD32 (Mouse BD Fc Block) and stained by the antibodies indicated below obtained from

ThermoFisher. Basically, the population of Tfh cells was stained with and evaluated using Alexa

Fluor 700 conjugated anti-CD4 IgG, fluorescein isothiocyanate (FITC) conjugated anti-CD44

IgG, BV421 Brilliant Violet 421 (BV-421) conjugated anti-CD185 (for CXCR5) IgG and PE

(phycoerythrin)-Cy7 (PE-Cy7) conjugated anti-CD279 (PD-1) IgG. To evaluate the population

of B cells, splenocytes were stained with FITC conjugated anti-CD4, anti-CD8 FITC, anti-Ly-

6G/Ly-6C (Gr-1), anti-F4/ 80, or anti-CD11c IgG, BV-421 or BV-711 conjugated anti-CD19

IgG, Cy5.5 or BV-711 conjugated anti-CD38 peridinin chlorophyll protein (PerCP) IgG, BV-

605-conjugated anti-mouse CD95 (Fas) IgG, allophycocyanin (APC) conjugated anti-CD138

IgG, Alexa Fluor 700 conjugated anti-IgD, and/or PE-Cy7 conjugated anti-IgM, and/or PerCP-

Cy5.5 conjugated anti-IgG1. Stained cell samples were then applied to an Attune Flowcytometer

(ThermoFisher), and at least 100,000 total events in a live gate were analyzed using FlowJo

software. Cells were gated on forward and side scatter for lymphocytes, exclusion of viability

dye, and singlet populations. B cells and Tfh cells in the spleen were determined based on

Fluorescence minus one (FMO).

**Statistical analyses.** Significant differences were determined with a Kruskal-Wallis test with the

two-stage step-up method of Benjamini, Krieger, and Yekutieli [106] and two-tailed Fisher test (for

seropositivity in **Fig. 2B and 5B**) [107] using GraphPad Prism 9.3.1. A p-value < 0.05 was used to

determine significance.

**AUTHOR CONTRIBUTIONS**

M.C.: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data

Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. S.H.:

Conceptualization, Software, Methodology, Validation, Formal analysis, Investigation, Data

Curation, Writing - Review & Editing, Visualization. J.M.: Methodology, Investigation. A.C.L.:

Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing - Review &

Editing, M.J.V.: Methodology, Validation, Formal analysis, Investigation, Data Curation,

Writing - Review & Editing, X.Y.: Methodology, Investigation, Data Curation,  Y.L.C.:

Investigation, Writing - Review & Editing, J.L.: Methodology, Investigation, Writing - Review

& Editing, Z.L.: Investigation, Writing - Review & Editing, U.S.: Methodology, Investigation,

Writing - Review & Editing, M.E.B.: Conceptualization, Methodology, Investigation, Writing -

Review & Editing, Supervision, Project administration, Funding acquisition. U.P.: Supervision,

Project administration, K.S.: Methodology, Investigation, Supervision, Project administration,

Funding acquisition, W.H.C.: Conceptualization, Methodology, Validation, Formal analysis,

Investigation, Data Curation, Writing - Original Draft, Writing - Review & Editing,

Visualization, Supervision, Project administration, Funding acquisition, Y.P.L.:

Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation,

Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Project

administration, Funding acquisition.

**ACKNOWLEDGEMENTS**

The authors thank Patricia Rosa and John Leong for providing *B. burgdorferi* strain B31-A3. We

thank Carly Fernandes and Connor McKaig's technical support. We are also grateful for Kate

Sulka in helping to review the manuscript and provide advice. We thank the Wadsworth Animal

Core for assistance with Animal Care and Tufts Laboratory Animal Medicine Service.

**FUNDING**

This work was supported by NIH grant R01AI181746 and R01AI81904 (for M.M., S.H., K.S.,

Y.L.), R01AI154542 (for X.Y., U.P.), R21AI144891 (J.M., Y.L.C., J.L., Z.L., U.S., M.E. B.,

W.H.C., Y.L.), and the U.S. Department of Defense, Congressionally Directed Medical Research

Programs, Grant Number W81XWH-20-1-0913 (J.M., Y.L.C., J.L., Z.L., U.S., M.E. B., W.H.C.,

Y.L.). The funders had no role in study design, data collection, interpretation, or the decision to

submit the work for publication.

**COMPETING INTERESTS**

Y. L. is the inventor on U.S. patent application no. US11771750B2 ("Composition and method

for generating immunity to *Borrelia burgdorferi*"). The remaining authors declare no competing

interests.

**DATA AVAILABILITY**

All data generated or analysed during this study are included in this published article and its

supplementary information files.

**REFERENCE**

1.    Cunningham AL*, et al.* Vaccine development: From concept to early clinical testing.

*Vaccine* **34**, 6655-6664 (2016).

2.    Roth GA, Picece V, Ou BS, Luo W, Pulendran B, Appel EA. Designing spatial and

temporal control of vaccine responses. *Nat Rev Mater* **7**, 174-195 (2022).

3.    Kennedy RB, Ovsyannikova IG, Palese P, Poland GA. Current Challenges in

Vaccinology. *Front Immunol* **11**, 1181 (2020).

4.    Gebre MS, Brito LA, Tostanoski LH, Edwards DK, Carfi A, Barouch DH. Novel

approaches for vaccine development. *Cell* **184**, 1589-1603 (2021).

5.    Schmidt ST, Pedersen GK, Christensen D. Rational Design and In Vivo Characterization

of Vaccine Adjuvants. *ILAR J* **59**, 309-322 (2018).

6.    O'Hagan DT, Fox CB. New generation adjuvants--from empiricism to rational design.

*Vaccine* **33 Suppl 2**, B14-20 (2015).

7.    Zhao T*, et al.* Vaccine adjuvants: mechanisms and platforms. *Signal Transduct Target*

*Ther* **8**, 283 (2023).

8.    Wang ZB, Xu J. Better Adjuvants for Better Vaccines: Progress in Adjuvant Delivery

Systems, Modifications, and Adjuvant-Antigen Codelivery. *Vaccines (Basel)* **8**,  (2020).

9.    Kugeler KJ, Schwartz, A.M., Delorey, M.J., Mead, P.S., Hinckley, A.F. Estimating the

Frequency of Lyme Disease Diagnoses, United States, 2010–2018. *Emerg Infect Dis* **27**,

(2021).

10.   Gomes-Solecki M*, et al.* Protective Immunity and New Vaccines for Lyme Disease. *Clin*

*Infect Dis*,  (2019).

11.    Dattwyler RJ, Gomes-Solecki M. The year that shaped the outcome of the OspA vaccine
694           for human Lyme disease. *NPJ Vaccines* **7**, 10 (2022).

696    12.    Wagner L*, et al.* Immunogenicity and safety of an 18-month booster dose of the VLA15
697           Lyme borreliosis vaccine candidate after primary immunisation in children, adolescents,
698           and adults in the USA: a randomised, observer-blind, placebo-controlled, phase 2 trial.
699           *Lancet Infect Dis*,  (2025).

701    13.    Lundberg U*, et al.* Preclinical Evidence for the Protective Capacity of Antibodies
702           Induced by Lyme Vaccine Candidate VLA15 in People. *Open Forum Infect Dis* **11**,
703           ofae467 (2024).

705    14.    Comstedt P, Schuler W, Meinke A, Lundberg U. The novel Lyme borreliosis vaccine
706           VLA15 shows broad protection against Borrelia species expressing six different OspA
707           serotypes. *PLoS One* **12**, e0184357 (2017).

709    15.    Caimano MJ*, et al.* The RpoS Gatekeeper in Borrelia burgdorferi: An Invariant
710           Regulatory Scheme That Promotes Spirochete Persistence in Reservoir Hosts and Niche
711           Diversity. *Front Microbiol* **10**, 1923 (2019).

713    16.    Ohnishi J, Piesman J, de Silva AM. Antigenic and genetic heterogeneity of Borrelia
714           burgdorferi populations transmitted by ticks. *Proceedings of the National Academy of*
715           *Sciences of the United States of America* **98**, 670-675 (2001).

17. Tahir D, *et al.* A Lyme disease mRNA vaccine targeting Borrelia burgdorferi OspA induces strong immune responses and prevents transmission in mice. *Mol Ther Nucleic Acids* **36**, 102514 (2025).

18. Gingerich MC, *et al.* Intranasal vaccine for Lyme disease provides protection against tick transmitted Borrelia burgdorferi beyond one year. *NPJ Vaccines* **9**, 33 (2024).

19. Pine M, *et al.* Development of an mRNA-lipid nanoparticle vaccine against Lyme disease. *Mol Ther* **31**, 2702-2714 (2023).

20. Federizon J, *et al.* Immunogenicity of the Lyme disease antigen OspA, particleized by cobalt porphyrin-phospholipid liposomes. *Vaccine* **38**, 942-950 (2020).

21. Kamp HD, *et al.* Design of a broadly reactive Lyme disease vaccine. *NPJ Vaccines* **5**, 33 (2020).

22. O'Bier NS, Camire AC, Patel DT, Billingsley JS, Hodges KR, Marconi RT. Development of novel multi-protein chimeric immunogens that protect against infection with the Lyme disease agent, Borreliella burgdorferi. *mBio* **15**, e0215924 (2024).

23. Rios S, *et al.* Rabies virus-vectored Lyme disease vaccine provides long-term protection against tick-transmitted Borrelia burgdorferi. *NPJ Vaccines* **10**, 231 (2025).

740    24.    Lin YP, Diuk-Wasser MA, Stevenson B, Kraiczy P. Complement Evasion Contributes to

741           Lyme Borreliae-Host Associations. *Trends Parasitol* **36**, 634-645 (2020).

743    25.    Bockenstedt LK, Wooten RM, Baumgarth N. Immune Response to Borrelia: Lessons

744           from Lyme Disease Spirochetes. *Curr Issues Mol Biol* **42**, 145-190 (2021).

746    26.    Skare JT, Garcia BL. Complement Evasion by Lyme Disease Spirochetes. *Trends

747           Microbiol* **28**, 889-899 (2020).

749    27.    Dulipati V, Meri S, Panelius J. Complement evasion strategies of Borrelia burgdorferi

750           sensu lato. *FEBS Lett* **594**, 2645-2656 (2020).

752    28.    Marcinkiewicz AL*, et al.* Structural evolution of an immune evasion determinant shapes

753           pathogen host tropism. *Proceedings of the National Academy of Sciences of the United

754           States of America* **120**, e2301549120 (2023).

756    29.    Bykowski T*, et al.* Coordinated expression of Borrelia burgdorferi complement regulator-

757           acquiring surface proteins during the Lyme disease spirochete's mammal-tick infection

758           cycle. *Infection and immunity* **75**, 4227-4236 (2007).

30.  Rogers EA, Marconi RT. Delineation of species-specific binding properties of the CspZ
protein (BBH06) of Lyme disease spirochetes: evidence for new contributions to the
pathogenesis of Borrelia spp. *Infection and immunity* **75**, 5272-5281 (2007).

31.  Rogers EA, Abdunnur SV, McDowell JV, Marconi RT. Comparative analysis of the
properties and ligand binding characteristics of CspZ, a factor H binding protein, derived
from Borrelia burgdorferi isolates of human origin. *Infection and immunity* **77**, 4396-
4405 (2009).

32.  Kraiczy P*, et al.* Borrelia burgdorferi complement regulator-acquiring surface protein 2
(CspZ) as a serological marker of human Lyme disease. *Clinical and vaccine*
*immunology : CVI* **15**, 484-491 (2008).

33.  Brangulis K*, et al.* Mechanistic insights into the structure-based design of a CspZ-
targeting Lyme disease vaccine. *Nat Commun* **16**, 2898 (2025).

34.  Coleman AS*, et al.* Borrelia burgdorferi complement regulator-acquiring surface protein
2 does not contribute to complement resistance or host infectivity. *PLoS One* **3**, 3010e
(2008).

35.  Marcinkiewicz AL*, et al.* Eliminating Factor H-Binding Activity of Borrelia burgdorferi
CspZ Combined with Virus-Like Particle Conjugation Enhances Its Efficacy as a Lyme
Disease Vaccine. *Front Immunol* **9**, 181 (2018).

36.    HogenEsch H, O'Hagan DT, Fox CB. Optimizing the utilization of aluminum adjuvants
785           in vaccines: you might just get what you want. *NPJ Vaccines* **3**, 51 (2018).

787    37.    Le Naour J, Kroemer G. Trial watch: Toll-like receptor ligands in cancer therapy.
788           *Oncoimmunology* **12**, 2180237 (2023).

790    38.    Brown JP, Zachary JF, Teuscher C, Weis JJ, Wooten RM. Dual role of interleukin-10 in
791           murine Lyme disease: regulation of arthritis severity and host defense. *Infection and*
792           *immunity* **67**, 5142-5150 (1999).

794    39.    Wang G, Ma Y, Buyuk A, McClain S, Weis JJ, Schwartz I. Impaired host defense to
795           infection and Toll-like receptor 2-independent killing of Borrelia burgdorferi clinical
796           isolates in TLR2-deficient C3H/HeJ mice. *FEMS Microbiol Lett* **231**, 219-225 (2004).

798    40.    Lee WY*, et al.* An intravascular immune response to Borrelia burgdorferi involves
799           Kupffer cells and iNKT cells. *Nat Immunol* **11**, 295-302 (2010).

801    41.    Tupin E*, et al.* NKT cells prevent chronic joint inflammation after infection with Borrelia
802           burgdorferi. *Proceedings of the National Academy of Sciences of the United States of*
803           *America* **105**, 19863-19868 (2008).

42.   Jahanbani S*, et al.* Increased macrophage phagocytic activity with TLR9 agonist

conjugation of an anti- Borrelia burgdorferi monoclonal antibody. *Clin Immunol* **246**,

109180 (2023).

43.   Bode C, Zhao G, Steinhagen F, Kinjo T, Klinman DM. CpG DNA as a vaccine adjuvant.

*Expert Rev Vaccines* **10**, 499-511 (2011).

44.   Thuluva S*, et al.* Immunogenicity and safety of Biological E's CORBEVAX vaccine

compared to COVISHIELD (ChAdOx1 nCoV-19) vaccine studied in a phase-3, single

blind, multicentre, randomized clinical trial. *Hum Vaccin Immunother* **19**, 2203632

(2023).

45.   Maddeppungeng M*, et al.* Safety and immunogenicity of a SARS-CoV-2 recombinant

protein subunit vaccine adjuvanted with Alum + CpG 1018 in healthy Indonesian adults:

A multicenter, randomized, comparative, observer-blind, placebo-controlled phase 2

study. *Hum Vaccin Immunother* **20**, 2429231 (2024).

46.   Tabarsi P*, et al.* Evaluating the efficacy and safety of SpikoGen(R), an Advax-CpG55.2-

adjuvanted severe acute respiratory syndrome coronavirus 2 spike protein vaccine: a

phase 3 randomized placebo-controlled trial. *Clin Microbiol Infect* **29**, 215-220 (2023).

47.   Lee GH, Lim SG. CpG-Adjuvanted Hepatitis B Vaccine (HEPLISAV-B(R)) Update.

*Expert Rev Vaccines* **20**, 487-495 (2021).

48.    Ulanova M, Tarkowski A, Hahn-Zoric M, Hanson LA. The Common vaccine adjuvant

830            aluminum hydroxide up-regulates accessory properties of human monocytes via an

831            interleukin-4-dependent mechanism. *Infection and immunity* **69**, 1151-1159 (2001).

833    49.    Hastey CJ, Elsner RA, Barthold SW, Baumgarth N. Delays and diversions mark the

834            development of B cell responses to Borrelia burgdorferi infection. *J Immunol* **188**, 5612-

835            5622 (2012).

837    50.    Blum LK*, et al.* Robust B Cell Responses Predict Rapid Resolution of Lyme Disease.

838            *Front Immunol* **9**, 1634 (2018).

840    51.    Hirschfeld M*, et al.* Cutting edge: inflammatory signaling by Borrelia burgdorferi

841            lipoproteins is mediated by toll-like receptor 2. *J Immunol* **163**, 2382-2386 (1999).

843    52.    Schenk M, Belisle JT, Modlin RL. TLR2 looks at lipoproteins. *Immunity* **31**, 847-849

844            (2009).

846    53.    Alexopoulou L*, et al.* Hyporesponsiveness to vaccination with Borrelia burgdorferi OspA

847            in humans and in TLR1- and TLR2-deficient mice. *Nat Med* **8**, 878-884 (2002).

54.    Fikrig E, Narasimhan S, Neelakanta G, Pal U, Chen M, Flavell R. Toll-like receptors 1

and 2 heterodimers alter Borrelia burgdorferi gene expression in mice and ticks. *J Infect*

*Dis* **200**, 1331-1340 (2009).

55.    Wooten RM*, et al.* Toll-like receptor 2 is required for innate, but not acquired, host

defense to Borrelia burgdorferi. *J Immunol* **168**, 348-355 (2002).

56.    Hoffmann P*, et al.* Stimulation of human and murine adherent cells by bacterial

lipoprotein and synthetic lipopeptide analogues. *Immunobiology* **177**, 158-170 (1988).

57.    Aliprantis AO*, et al.* Cell activation and apoptosis by bacterial lipoproteins through toll-

like receptor-2. *Science* **285**, 736-739 (1999).

58.    Cornelie S*, et al.* Direct evidence that toll-like receptor 9 (TLR9) functionally binds

plasmid DNA by specific cytosine-phosphate-guanine motif recognition. *J Biol Chem*

**279**, 15124-15129 (2004).

59.    Hemmi H*, et al.* A Toll-like receptor recognizes bacterial DNA. *Nature* **408**, 740-745

(2000).

60.    Krieg AM. Immune effects and mechanisms of action of CpG motifs. *Vaccine* **19**, 618-

622 (2000).

61.  Jiang W, Lederman MM, Harding CV, Rodriguez B, Mohner RJ, Sieg SF. TLR9

stimulation drives naive B cells to proliferate and to attain enhanced antigen presenting

function. *Eur J Immunol* **37**, 2205-2213 (2007).

62.  Brossay L*, et al.* CD1d-mediated recognition of an alpha-galactosylceramide by natural

killer T cells is highly conserved through mammalian evolution. *J Exp Med* **188**, 1521-

1528 (1998).

63.  Patel O*, et al.* NKT TCR recognition of CD1d-alpha-C-galactosylceramide. *J Immunol*

**187**, 4705-4713 (2011).

64.  Cervantes JL*, et al.* Phagosomal signaling by Borrelia burgdorferi in human monocytes

involves Toll-like receptor (TLR) 2 and TLR8 cooperativity and TLR8-mediated

induction of IFN-beta. *Proceedings of the National Academy of Sciences of the United*

*States of America* **108**, 3683-3688 (2011).

65.  Levi DT, Iablokova TB, Altynova MP, Mitinskaia LA, Bardina EE. [Activity and

specificity of tuberculins derived from BCG culture filtrates]. *Probl Tuberk*, 60-63 (1985).

66.  Lee WY*, et al.* Invariant natural killer T cells act as an extravascular cytotoxic barrier for

joint-invading Lyme Borrelia. *Proceedings of the National Academy of Sciences of the*

*United States of America* **111**, 13936-13941 (2014).

67.    Suthers AN, Sarantopoulos S. TLR7/TLR9- and B Cell Receptor-Signaling Crosstalk:

Promotion of Potentially Dangerous B Cells. *Front Immunol* **8**, 775 (2017).

68.    Hanten JA*, et al.* Comparison of human B cell activation by TLR7 and TLR9 agonists.

*BMC Immunol* **9**, 39 (2008).

69.    Doherty DG, Melo AM, Moreno-Olivera A, Solomos AC. Activation and Regulation of

B Cell Responses by Invariant Natural Killer T Cells. *Front Immunol* **9**, 1360 (2018).

70.    Leadbetter EA, Karlsson MCI. Invariant natural killer T cells balance B cell immunity.

*Immunol Rev* **299**, 93-107 (2021).

71.    Elsner RA, Hastey CJ, Olsen KJ, Baumgarth N. Suppression of Long-Lived Humoral

Immunity Following Borrelia burgdorferi Infection. *PLoS Pathog* **11**, e1004976 (2015).

72.    Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update

on Toll-like receptors. *Nat Immunol* **11**, 373-384 (2010).

73.    Lochhead RB, Arvikar SL, Aversa JM, Sadreyev RI, Strle K, Steere AC. Robust

interferon signature and suppressed tissue repair gene expression in synovial tissue from

patients with postinfectious, Borrelia burgdorferi-induced Lyme arthritis. *Cell Microbiol*

**21**, e12954 (2019).

74.    Strle K, Jones KL, Drouin EE, Li X, Steere AC. Borrelia burgdorferi RST1 (OspC type A)

genotype is associated with greater inflammation and more severe Lyme disease. *Am J*

*Pathol* **178**, 2726-2739 (2011).

75.    Mullegger RR, McHugh G, Ruthazer R, Binder B, Kerl H, Steere AC. Differential

expression of cytokine mRNA in skin specimens from patients with erythema migrans or

acrodermatitis chronica atrophicans. *J Invest Dermatol* **115**, 1115-1123 (2000).

76.    Honda K*, et al.* IRF-7 is the master regulator of type-I interferon-dependent immune

responses. *Nature* **434**, 772-777 (2005).

77.    Honda K, Taniguchi T. IRFs: master regulators of signalling by Toll-like receptors and

cytosolic pattern-recognition receptors. *Nat Rev Immunol* **6**, 644-658 (2006).

78.    Erdile LF, Guy B. OspA lipoprotein of Borrelia burgdorferi is a mucosal immunogen and

adjuvant. *Vaccine* **15**, 988-996 (1997).

79.    Giambartolomei GH, Dennis VA, Lasater BL, Philipp MT. Induction of pro- and anti-

inflammatory cytokines by Borrelia burgdorferi lipoproteins in monocytes is mediated by

CD14. *Infection and immunity* **67**, 140-147 (1999).

80.    Pollard AJ, Bijker EM. A guide to vaccinology: from basic principles to new

developments. *Nat Rev Immunol* **21**, 83-100 (2021).

81.     Chen WH, Strych U, Bottazzi ME, Lin YP. Past, present, and future of Lyme disease vaccines: antigen engineering approaches and mechanistic insights. *Expert Rev Vaccines*, 1-13 (2022).

82.     Zhang Y*, et al.* IgG1 Is Required for Optimal Protection after Immunization with the Purified Porin OmpD from Salmonella Typhimurium. *J Immunol* **199**, 4103-4109 (2017).

83.     Tunev SS, Hastey CJ, Hodzic E, Feng S, Barthold SW, Baumgarth N. Lymphoadenopathy during lyme borreliosis is caused by spirochete migration-induced specific B cell activation. *PLoS Pathog* **7**, e1002066 (2011).

84.     Frischauf N, Strasser J, Borg EGF, Labrijn AF, Beurskens FJ, Preiner J. Complement activation by IgG subclasses is governed by their ability to oligomerize upon antigen binding. *Proceedings of the National Academy of Sciences of the United States of America* **121**, e2406192121 (2024).

85.     Edmondson DG*, et al.* Enhanced Protective Immunogenicity of Homodimeric Borrelia burgdorferi Outer Surface Protein C. *Clinical and vaccine immunology : CVI* **24**,  (2017).

86.     Steere AC*, et al.* Lyme borreliosis. *Nature reviews Disease primers* **2**, 16090 (2016).

87.   Radolf JD, Caimano MJ, Stevenson B, Hu LT. Of ticks, mice and men: understanding the

dual-host lifestyle of Lyme disease spirochaetes. *Nature reviews Microbiology* **10**, 87-99

(2012).

88.   Tracy KE, Baumgarth N. Borrelia burgdorferi Manipulates Innate and Adaptive

Immunity to Establish Persistence in Rodent Reservoir Hosts. *Front Immunol* **8**, 116

(2017).

89.   Lone AG, Bankhead T. The Borrelia burgdorferi VlsE Lipoprotein Prevents Antibody

Binding to an Arthritis-Related Surface Antigen. *Cell Rep* **30**, 3663-3670 e3665 (2020).

90.   Zhang JR, Hardham JM, Barbour AG, Norris SJ. Antigenic variation in Lyme disease

borreliae by promiscuous recombination of VMP-like sequence cassettes. *Cell* **89**, 275-

285 (1997).

91.   Bankhead T, Chaconas G. The role of VlsE antigenic variation in the Lyme disease

spirochete: persistence through a mechanism that differs from other pathogens. *Mol*

*Microbiol* **65**, 1547-1558 (2007).

92.   Hart T*, et al.* Polymorphic factor H-binding activity of CspA protects Lyme borreliae

from the host complement in feeding ticks to facilitate tick-to-host transmission. *PLoS*

*Pathog* **14**, e1007106 (2018).

93.    Barbour AG, Burgdorfer W, Grunwaldt E, Steere AC. Antibodies of patients with Lyme

987           disease to components of the Ixodes dammini spirochete. *The Journal of clinical*

988           *investigation* **72**, 504-515 (1983).

990    94.    Purser JE, Norris SJ. Correlation between plasmid content and infectivity in Borrelia

991           burgdorferi. *Proceedings of the National Academy of Sciences of the United States of*

992           *America* **97**, 13865-13870 (2000).

994    95.    Bunikis I, Kutschan-Bunikis S, Bonde M, Bergstrom S. Multiplex PCR as a tool for

995           validating plasmid content of Borrelia burgdorferi. *Journal of microbiological methods*

996           **86**, 243-247 (2011).

998    96.    Chen YL*, et al.* Biophysical and biochemical characterization of a recombinant Lyme

999           disease vaccine antigen, CspZ-YA. *Int J Biol Macromol* **259**, 129295 (2024).

1000

1001   97.    Nayak A*, et al.* Broadly Protective Multivalent OspA Vaccine against Lyme Borreliosis,

1002          Developed Based on Surface Shaping of the C-Terminal Fragment. *Infection and*

1003          *immunity* **88**,  (2020).

1004

1005   98.    Erdile LF*, et al.* Role of attached lipid in immunogenicity of Borrelia burgdorferi OspA.

1006          *Infection and immunity* **61**, 81-90 (1993).

1007

1008    99.    Liang FT, Bowers LC, Philipp MT. C-terminal invariable domain of VlsE is
1009          immunodominant but its antigenicity is scarcely conserved among strains of Lyme
1010          disease spirochetes. *Infection and immunity* **69**, 3224-3231 (2001).

1011

1012    100.   Goedhart J, Luijsterburg MS. VolcaNoseR is a web app for creating, exploring, labeling
1013          and sharing volcano plots. *Sci Rep* **10**, 20560 (2020).

1014

1015    101.   Oliveros JC. Venny. An interactive tool for comparing lists with Venn's diagram.) (2007-
1016          2015).

1017

1018    102.   Morpheus.).

1019

1020    103.   Subramanian A*, et al.* Gene set enrichment analysis: a knowledge-based approach for
1021          interpreting genome-wide expression profiles. *Proceedings of the National Academy of*
1022          *Sciences of the United States of America* **102**, 15545-15550 (2005).

1023

1024    104.   Leao AC*, et al.* Different responses involving Tfh cells delay parasite-specific antibody
1025          production in Trypanosoma cruzi acute experimental models. *Front Immunol* **16**,
1026          1487317 (2025).

1027

1028    105.   Versteeg L*, et al.* Transferring Luminex(R) cytokine assays to a wall-less plate
1029          technology: Validation and comparison study with plasma and cell culture supernatants. *J*
1030          *Immunol Methods* **440**, 74-82 (2017).

1031

1032    106.    Benjamini YK, A. M. Yekutieli, D. . Adaptive linear step-up procedures that control the

1033            false discovery rate. *Biometrika* **93**, 491-507 (2006).

1034

1035    107.    Fisher RA. *Statistical methods for research workers*, 5th edn. Oliver&Boyd (1934).

1036

1037

**FIGURE LEGENDS**

**Figure 1. Vaccination with CspZ-YA$_{C187S}$ formulated with Alum-CpG or Alum-αGal**

**induced robust antibody  titers and bactericidal activities in mice. (A)** Pre-adolescent

C3H/HeN mice received two inoculations at 0- and 14-days post initial immunization (dpii) with

TBS (control) or CspZ-YA$_{C187S}$ formulated with the indicated adjuvants, followed by infection

via feeding of *I. scapularis* nymphs carrying *B. burgdorferi* strain B31-A3 at 35 dpii and

euthanasia at 56 dpii. Sera were collected at **(B, D, E)** 14 dpii for one immunization (Immun. 1)

and **(C, F, G)** 28 dpii for two immunizations (Immun. 2). **(B-C)** The levels of total IgG against

CspZ-YA$_{C187S}$ in the sera were determined using quantitative ELISA. Data shown are the

geometric mean ± geometric standard deviation of the titers of anti-CspZ-YA$_{C187S}$ antibodies

from five mice per group. **(D-G)** Sera were diluted as indicated, and mixed with guinea pig

complement and *B. burgdorferi* B31-A3 for 24 hours. Surviving spirochetes were quantified

microscopically from three fields of view in three independent experiments. **(D, F)** Survival

percentage was derived from the proportion of serum-treated to untreated spirochetes. Data

shown are the mean ± SEM of the survival percentage from three replicates in one representative

experiment. **(E, G)** The BA$_{50}$ value, representing the dilution rate that effectively killed 50% of

spirochetes, was obtained from curve-fitting and extrapolation of Panels C and F. Data shown are

the geometric mean ± geometric standard deviation of the borreliacidal titers from five mice per

group in three experiments per mouse sample and shown in **Table S1.** ("NK"), no killing.

Statistical significance ($p < 0.05$, Kruskal Wallis test with the two-stage step-up method of

Benjamini, Krieger, and Yekutieli) of IgG titers or borreliacidal titers from an indicated group of

mice was compared to the titers from TBS-inoculated mice ("*") or between indicated groups

were shown as "#".

1061

**Figure 2. CspZ-YA$_{C187S}$ formulated with Alum-CpG or Alum-αGal protected mice from**

**seroconversion, borrelial colonization and Lyme disease-associated joint inflammation.**

Five pre-adolescent C3H/HeN mice were inoculated with TBS or CspZ-YA$_{C187S}$ formulated with

indicated adjuvants, followed by infection using **(A)** nymphs carrying *B. burgdorferi* B31-A3 in

the fashion described in **Fig. 1A**. Mice inoculated with TBS that are not fed on by nymphs were

included as an uninfected control group (uninfect.). **(B)** Seropositivity was determined by

measuring the levels of IgG against C6 peptides in the sera of those mice at 21 dpf using ELISA.

A mouse was considered as seropositive if that mouse had IgG levels against C6 peptides greater

than the threshold, the mean plus 1.5-fold standard deviation of the IgG levels against C6

peptides from the TBS-inoculated, uninfected mice (dotted line). The number of mice in each

group with anti-C6 IgG levels greater than the threshold (seropositive) is shown. Data presented

here are the geometric mean ± geometric standard deviation of the titers of anti-C6 IgG.

Statistical significances ($p < 0.05$, Kruskal-Wallis test with the two-stage step-up method of

Benjamini, Krieger, and Yekutieli) of differences in IgG titers relative to (*) uninfected mice are

presented. **(A, C-F)** *B. burgdorferi* (*Bb*) burdens at **(A)** nymphs after feeding to repletion or **(C)**

the tick feeding site ("Bite Site"), **(D)** bladder, **(E)** heart, and **(F)** knees, were quantitatively measured at 21 dpf, shown as the number of **(A)** *Bb* per tick or **(C-F)** per 100 ng total DNA. Data shown are the geometric mean ± geometric standard deviation of the spirochete burdens from each group of mice. Asterisks indicate the statistical significance ($p < 0.05$, Kruskal Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli) of differences in bacterial burdens relative to uninfected mice. **(G-H)** Tibiotarsus joints at 42 dpli were collected to assess inflammation by staining these tissues using hematoxylin and eosin. **(G)** Representative images from one mouse per group are shown. Top panels are lower-resolution images (joint, ×10 [bar, 160 µm]); bottom panels are higher-resolution images (joint, 2×20 [bar, 80 µm]) of selected areas (highlighted in top panels). Arrows indicate infiltration of immune cells. **(H)** To quantitate inflammation of joint tissues, at least ten random sections of tibiotarsus joints from each mouse were scored on a scale of 0-3 for the levels of inflammation. Data shown are the mean inflammation score ± standard deviation of the inflammatory scores from each group of mice. Asterisks indicate the statistical significance ($p < 0.05$, Kruskal Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli) of differences in inflammation relative to uninfected mice.

**Figure 3. CpG or αGal skewed the Alum-CspZ-YA$_{C187S}$-induced immune pathways in different fashions.** Five pre-adolescent C3H/HeN mice per group were inoculated with TBS or CspZ-YA$_{C187S}$ formulated with Alum, Alum-CpG, and Alum-αGal twice in the fashion described in **Fig. 1A. (A-H)** Spleens were collected at 28dpii, and the expression levels of genes in the spleen from each group of the mice were determined. Differentially expressed genes (DEGs) are defined by an adjusted P values less than 0.05 and absolute (log$_2$fold change) ≥ 0.58 or ≤ -0.58.

(**A-B**) Venn diagram comparing the number of (**A**) up- or (**B**) down-regulated DEGs identified in the spleen from mice inoculated with CspZ-YA$_{C187S}$ formulated with Alum-CpG vs. Alum and Alum-αGal vs. Alum. (**C, E**) The expression levels of genes in the comparison of (**C**) Alum-CpG vs. Alum and (**E**) Alum-αGal vs. Alum are plotted as a Volcano plots. The up- and down-regulated DEGs are shown in red and blue, respectively. The immune-related DEGs in each comparison are highlighted. The details of all DEGs are shown in **Dataset S1**, and the details of these immune related DEGs are shown in **Table S2 to S3**. (**D, F**) Heat map exhibits the expression patterns of all DEGs in the comparison of (**D**) Alum-CpG vs. Alum and (**F**) Alum-αGal vs. Alum, and the gene IDs are shown in the right. (**G-H**) GSEA analysis to identify the enriched pathways triggered by the CspZ-YA$_{C187S}$ formulated with Alum-CpG or Alum-αGal when the enrichment analysis was performed in the comparison of the formulations under Alum-CpG vs. Alum-αGal. The DEGs enriched in the formulation of (**G**) Alum-CpG vs. (**H**) Alum-αGal are shown. The pathways shown were based on the statistical cutoff (FDR q-value <0.1 and NES >1.5).

**Figure 4. The splenocytes from vaccination with Alum-CpG or Alum-αGal formulated CspZ-YA$_{C187S}$ differed in the elevated cytokine species but were similar in the activated memory cell types.** Splenocytes isolated from the mouse spleens at 21 dpii were restimulated with CspZ-YA$_{C187S}$ for 48-h. (**A-G**) Multiplex Luminex analysis was used to measure the indicated secreted cytokines. (**H-K**) Flow cytometry analysis was applied to determine the percentage of (**H**) Germinal Center B cells (CD19+CD38-FAS+), (**I**) Follicular T helper cells (CD4$^+$CD44$^+$CXCR5hiPD$^-$1hi), (**J**) Memory B cells (CD19$^+$CD38$^+$IgD$^-$IgM$^-$), and (**K**) IgG1+ Memory B cells (CD19$^+$CD38$^+$IgD$^-$IgM$^-$IgG1$^+$). Data shown are the mean ± SEM of the (**A-G**)

1123 cytokine levels or **(H-K)** percentage of live/singlets cell populations from each group of mice.

1124 The statistical significance ($p < 0.05$, Kruskal Wallis test with the two-stage step-up method of

1125 Benjamini, Krieger, and Yekutieli) of differences in bacterial burdens relative to TBS-inoculated

1126 mice ("*") or between indicated groups of mice ("#") were indicated.

1127

**Figure 5. Titers and bactericidal activities of antibodies triggered by Alum-CpG**

**formulated CspZ-YA$_{C187S}$ were detectable for up to nine months. (A)** The schematic diagram

shows pre-adolescent C3H/HeN mice receiving one or two injections of CspZ-YA$_{C187S}$

formulated with Alum-CpG on 0 and 14 dpii.. Mice inoculated with TBS at 0 and 14 dpii, or

TMG-formulated OspA at 0, 14, and 28dpii were included as control. Sera were collected at 14,

42, 70, 98, 126, 154, 182, 210, 238, and 273 days post last immunization (dpli). **(B-C)** The levels

of total IgG against **(B)** OspA and **(C)** CspZ-YA$_{C187S}$ in the sera were determined using

quantitative ELISA. Data shown are the geometric mean ± geometric standard deviation of the

titers of **(B)** anti-OspA and **(C)** CspZ-YA$_{C187S}$ IgGs from five mice per group. **(D)** The sera were

diluted as indicated and mixed with guinea pig complement and *B. burgdorferi* B31-A3 for 24

hours. Surviving spirochetes were quantified from three fields of view microscopically in three

independent experiments. The survival percentage was derived from the proportion of serum-

treated to untreated spirochetes. Data shown are the mean ± SEM of the survival percentage from

three replicates in one representative experiment. The BA$_{50}$ value, representing the dilution rate

that effectively killed 50% of spirochetes, was obtained from curve-fitting and shown in **Table**

**S4**. Asterisks represent that no killing was detected in indicated groups of mice and time points.

1144

**Figure 6. Natural infection triggered CspZ-targeting bactericidal antibodies linked to prevention of seroconversion, borrelial colonization and joint inflammation in Alum-CpG-CspZ-YA$_{C187S}$ vaccinated mice.** As shown in **Fig. 5A**, five pre-adolescent C3H/HeN mice were inoculated with TBS or CspZ-YA$_{C187S}$ formulated with Alum-CpG. At 273 dpli, using *I. scapularis* nymphs carrying *B. burgdorferi* strain B31-A3, the mice were naturally infected. Sera were collected at 4, 7, 10, 14, and 21 dpf, and mice were euthanized at 21 dpf. Mice inoculated with TBS that were not fed on by nymphs were included as an uninfected control group (uninfect.). **(A-C)** The levels of total IgG against **(A)** OspA and **(B)** CspZ-YA$_{C187S}$ or indicated IgG isotypes against **(C)** CspZ-YA$_{C187S}$ in the sera at indicated time points were determined using quantitative ELISA. Data shown are the geometric mean ± geometric standard deviation of the titers of **(A)** anti-OspA and **(B-C)** CspZ-YA$_{C187S}$ IgGs from five mice per group. **(D)** The sera were diluted as indicated, and mixed with guinea pig complement and *B. burgdorferi* B31-A3 for 24 hours. Surviving spirochetes were quantified from three fields of view microscopically in three independent experiments. The survival percentage was derived from the proportion of serum-treated to untreated spirochetes. Data shown are the mean ± SEM of the survival percentage from three replicates in one representative experiment. The BA$_{50}$ value, representing the dilution rate that effectively killed 50% of spirochetes, was obtained from curve-fitting and shown in **Table S5**. Asterisks represent that no killing was detected in indicated groups of mice and time points. **(E)** The *B. burgdorferi burdens* of nymphs carrying *B. burgdorferi* B31-A3. Mice inoculated with TBS that were not fed on by nymphs were included as an uninfected control group (uninfect.). **(F)** Seropositivity was determined by measuring the levels of IgG against C6 peptides in the sera of those mice at 21 dpf using ELISA. The mouse was considered as seropositive if that mouse had IgG levels against C6 peptides greater than the threshold, the

mean plus 1.5-fold standard deviation of the IgG levels against C6 peptides from the TBS-

inoculated, uninfected mice (dotted line). The number of mice in each group with the anti-C6

IgG levels greater than the threshold (seropositive) is shown. Data shown are the geometric mean

± geometric standard deviation of the titers of anti-C6 IgG. **(G-J)** *B. burgdorferi* (*Bb*) burdens at

**(G)** the tick feeding site ("Bite Site"), **(H)** bladder, **(I)** heart, and **(J)** knees, were quantitatively

measured at 21 dpf, shown as the number of **(E)** *Bb* per tick or **(G-J)** per 100ng total DNA.  Data

shown are the geometric mean ± geometric standard deviation of the spirochete burdens from

each group of mice. **(K-L)** Tibiotarsus joints at 21 dpf were collected to assess inflammation by

staining these tissues using hematoxylin and eosin. **(K)** Representative images from one mouse

per group are shown. Top panels are lower-resolution images (joint, ×10 [bar, 160 µm]); bottom

panels are higher-resolution images (joint, 2×20 [bar, 80 µm]) of selected areas (highlighted in

top panels). Arrows indicate infiltration of immune cells. **(L)** To quantitate inflammation of joint

tissues, at least ten random sections of tibiotarsus joints from each mouse were scored on a scale

of 0-3 for the levels of inflammation.  Data shown are the mean inflammation score ± standard

deviation of the inflammatory scores from each group of mice. Asterisks indicate the statistical

significance ($p < 0.05$, Kruskal Wallis test with the two-stage step-up method of Benjamini,

Krieger, and Yekutieli) of differences in **(C, E, G-J)** bacterial burdens, **(F)** C6 IgG titers, **(L)**

inflammatory scores relative to uninfected mice.

**Upregulated DEGs**

A

Alum-CpG vs. Alum

Alum-αGal vs. Alum

5 (4.8%)  3 (2.9%)  97 (92.4%)

**Downregulated DEGs**

B

Alum-CpG vs. Alum

Alum-αGal vs. Alum

28 (46.7%)  0  32 (53.3%)

C

Alum-CpG vs. Alum Significance (-Log₁₀)

Fold Change (Log2)

Alum    Alum-CpG

Labels: 9330159F19Rik, Errfi1, Adamts1, Clec10a, Alox15, Pde1a, Klf9, Gdf6, Ccl11, Ebf3, Cd34, Cav2, Ighv8-12, Serpina3c, Maob, Mmp3, Ighv1-70

D

Alum / Alum-CpG / Gene name / ID

E

Alum-αGal vs. Alum Significance (-Log₁₀)

Fold Change (Log2)

Alum    Alum-αGal

Labels: Ifit1, Igkv11-125, Rock1, Siglec1, Chp2, Mmp27, Cpeb4, Cpeb3, Pla2g4c, Igkv2-109, Atoh8, H2-Q1, H2-Q2, H2-Q3, Iglc2, Tlr12, Igkv8-19, Socs1, Igkv12-46, Igkv7-33, Iglj3, Ighv8-12, Cd3d, Igkv3-8, H2-Q6

F

Alum / Alum-αGal / ID / Gene name

G

**Alum-CpG vs. Alum-αGal (Enriched in Alum-CpG)**

- NEGATIVE REGULATION OF VIRAL PROCESS
- PROTEIN K63 LINKED UBIQUITINATION
- IFNA1 RESPONSE UP
- ANTIVIRAL INNATE IMMUNE RESPONSE
- NK CELL IFNK RESPONSE UP
- POSITIVE REGULATION OF INTERFERON BETA PRODUCTION
- TYPE I INTERFERON PRODUCTION
- REGULATION OF VIRAL GENOME REPLICATION
- ILC IFNB RESPONSE UP
- CLATHRIN DEPENDENT ENDOCYTOSIS
- MYELOID CELL HOMEOSTASIS
- PROTEASOME AND UBIQUITIN DEPENDENT PROTEIN CATABOLISM

Normalized Enrichment Score (NES)

FDR q-value

H

**Alum-CpG vs. Alum-αGal (Enriched in Alum-αGal)**

- ER PHAGOSOME PATHWAY
- ANTIMICROBIAL HUMORAL IMMUNE RESPONSE
- T CELL CD4 TL1A RESPONSE UP
- ANTIGEN PRESENTATION AND LOADING OF CLASS I MHC
- MYELOID DENDRITIC CELL DIFFERENTIATION
- DENDRITIC CELL CHEMOTAXIS
- B CELL RECEPTOR SIGNALING PATHWAY
- T CELL CD8 TL1A RESPONSE UP
- PHOSPHORYLATION OF CD3 AND TCR ZETA CHAINS
- ENDOSOMAL VACUOLAR PATHWAY
- GRANULOCYTE ACTIVATION
- ANTIGEN PRESENTATION OF EXOGENOUS ANTIGEN
- NK CELL IL36A RESPONSE UP
- POSITIVE REGULATION OF ERK1 AND ERK2 CASCADE
- NEUTROPHIL DEGRANULATION

Normalized Enrichment Score (NES)

FDR q-value