Neuronal Toll-like Receptor-4 regulation of Matrix Metalloproteinase-9 Activity

Mediates Dentate Circuit Dysfunction after Traumatic Brain Injury

Deepak Subramanian[1#], Erick Contreras[1], Laura Dovek[2], Razieh Jaberi[1], Iryna M Ethell[2] and

Vijayalakshmi Santhakumar[1#]

[1]Department of Molecular, Cell and Systems Biology, [2]Biomedical Sciences, University of

California Riverside, Riverside, California 92521.

Keywords: Traumatic Brain Injury, Dentate gyrus, Toll-like Receptor 4, Matrix

Metalloproteinase-9, Neuro-immune signaling, Synaptic plasticity

Abstract: 273 words   Full Text Character count: 66,897   Number of pages: 47

Number of figures: 7   Supplementary figures: 4   Color figures: 10

Number of tables: 0   Number of supplementary tables: 3

Correspondence:
Deepak Subramanian, PhD
Department of Molecular, Cell and Systems Biology
University of California, Riverside
E-mail: dsubra@ucr.edu
19
Co-Correspondence:
Vijayalakshmi Santhakumar, PhD
Department of Molecular, Cell and Systems Biology
University of California, Riverside
E-mail: vijayas@ucr.edu

List of Abbreviations:

aCSF: Artificial Cerebrospinal Fluid

AMPAR: α-Amino-3-hydroxy-5-methyl-4-isoxazolepropionic Acid Receptor

ANOVA: Analysis of Variance

CA1 SR: CA1 Stratum Radiatum

CaMKIIa: Calcium/Calmodulin-Dependent Protein Kinase II Alpha

CreERT2: Cre Recombinase Estrogen Receptor T2

DG: Dentate Gyrus

DQ Gelatinase: Dye Quenched fluorogenic substrate for zymography

ECM: Extracellular Matrix

EPSC: Excitatory Postsynaptic Current

FPI: Fluid Percussion Injury

HMGB1: High-Mobility Group Box 1

IEI: Inter-Event Interval

IPSC: Inhibitory Postsynaptic Current

JSH-23: NF-κB Inhibitor

K-S test: Kolmogorov-Smirnov Test

LPS: Lipopolysaccharide

LTP: Long Term Potentiation

MMP-9: Matrix Metalloproteinase-9

NF-κB: Nuclear Factor Kappa B

TAK-242: TLR4 Antagonist

TBS: Theta Burst Stimulation

TBI: Traumatic Brain Injury

TTX: Tetrodotoxin

Abstract:

Neuroinflammatory pathways activated by traumatic brain injury (TBI) are critical mediators of long-term neurological dysfunction and represent promising therapeutic targets. Toll-like receptor 4 (TLR4), an innate immune receptor, was previously shown to contribute to seizure susceptibility and cognitive deficits in rats after lateral fluid percussion injury (FPI). However, the cellular and molecular mechanisms underlying TLR4-mediated circuit dysfunction early after brain injury is not fully understood. In this study, we define a cell- and circuit- specific neuroimmune-extracellular matrix signaling axis that mediates early post-TBI circuit dysfunction in the hippocampal Dentate Gyrus (DG). Using *ex vivo* electrophysiology in rat and mouse models one-week after brain injury, we demonstrate that neuronal TLR4 signaling regulates both excitatory and inhibitory synaptic inputs to dentate granule cells (DGC). Pharmacological inhibition of TLR4 in rats within 24 hours post-injury and cell-type-specific deletion of TLR4 in glutamatergic neurons in mice demonstrated that injury-driven excitatory circuit dysfunction is mediated by neuronal TLR4 and relies on downstream activation of Matrix Metalloproteinase-9 (MMP-9), an enzyme critical for extracellular matrix remodeling. In contrast, TLR4-dependent alterations in inhibitory inputs occurred independent of MMP-9, revealing a mechanistic divergence. Early inhibition of either TLR4 signaling or MMP-9 activity in rats within 24 hours post-injury reduced network hyperexcitability, improved long-term potentiation (LTP) in the dentate gyrus measured *in vivo* one-week after injury, and attenuated long-term deficits in spatial memory one-month post-injury. Furthermore, we demonstrate a context-dependent role for TLR4 signaling contributing to circuit homeostasis in the uninjured brain and driving

75   circuit dysfunction in the injured brain. Together, these results identify a novel TLR4 -MMP-9

76   axis as a key driver of early post-TBI dentate gyrus circuit dysfunction and behavioral

77   deficits.

INTRODUCTION:

Traumatic brain injury (TBI) initiates a robust and complex inflammatory cascade that exacerbates secondary structural and functional damage [1–4]. While cellular damage is widespread and diffuse, the hippocampus is highly susceptible to post-traumatic pathology. Specifically, the hippocampal dentate gyrus (DG) undergoes extensive loss of hilar neurons and circuit reorganization that critically alters hippocampal network physiology and contributes to the development of cognitive impairments and post-traumatic epilepsy [5–11]. In this context, we previously identified a pivotal role for the neuronally expressed innate immune receptor, Toll-like receptor 4 (TLR4), in enhancing neuronal excitability and driving epileptogenesis after injury [12–14]. Yet, the downstream mechanisms by which specific immune receptors influence circuit physiology remain poorly understood.

 Recent evidence indicates that canonical TLR4 activators, such as lipopolysaccharide (LPS) and high-mobility group box 1 (HMGB1), can upregulate Matrix Metalloproteinase-9 (MMP-9) activity in vitro [15,16]. MMP-9 , a zinc-dependent endopeptidase, is a critical regulator of synaptic communication and plasticity through the activity-dependent degradation of the extracellular matrix (ECM)[17–20]. Importantly, MMP-9 levels are elevated early in both animal models and clinical populations after brain injury [21–24] , driving pathological remodeling of synapses and circuits that promote epileptogenesis. [25–29]. Despite a strong temporal correlation between MMP-9 elevation and transient increases in TLR4 signaling after injury [12,30], a causal association between these two pathways has not been established. Based on the temporal correlation and shared

upstream activators, we hypothesized a mechanistic interaction between these two

pathways that contributes to the early circuit dysfunction after brain injury.

Here, we provide evidence for a novel mechanism involving TLR4-mediated upregulation of

MMP-9 activity following brain injury that drives maladaptive changes in synaptic

physiology, network excitability, and plasticity. Our findings reveal that the injury-induced

increase of MMP-9 in the dentate gyrus is specifically mediated by neuronal TLR4 signaling.

While TLR4-regulated MMP-9 activity was critical in enhancing excitatory synaptic inputs,

inhibitory input dysregulation was mediated by TLR4 signaling independent of MMP-9.

Furthermore, we demonstrate that this TLR4 and MMP-9 dependent pathway enhances

network excitability and impairs synaptic plasticity early after injury. Importantly, we show

that pharmacological inhibition of either TLR4 or MMP-9 effectively mitigates these early

pathological changes, highlighting the therapeutic potential of targeting this neuro-immune

signaling pathway to preserve circuit function after TBI.

METHODS:

All experiments were performed in accordance with IACUC protocols approved by the University of California-Riverside, CA and in keeping with the ARRIVE guidelines. Male and female Wistar rats (P23-P24) and mice (TLR4$^{f/f}$:CaMKIIa-CreERT2) were housed with littermates in a 12-hour light/dark cycle with access to water and food ad libidum. All mice were bred in-house. Animals were randomly assigned to either sham or FPI groups and treatment conditions.

Fluid Percussion Injury (FPI) and treatments

Juvenile male and female Wistar rats (24 days old) were subjected to moderate (1.8-2.0 atm) lateral FPI or sham injury using standard methods [12–14]. Briefly, a 2.7 mm hole was trephined on the left side of the skull 3 mm caudal to bregma and 3.5 mm lateral from the sagittal suture. A Luer-Lok syringe hub with a 3 mm inner diameter was placed over the exposed dura and bonded to the skull with cyanoacrylate adhesive. The animals were returned to their home cage following recovery. One day later, animals were briefly anesthetized with isoflurane and attached to a fluid percussion injury device (Virginia Commonwealth University, Richmond, VA, Model 01-B) directly at the metal nozzle. A pendulum was dropped to deliver a brief (20ms) 1.9–2.0 atm impact on the dura. For sham injury, the animals underwent surgical implantation of the hub and were anesthetized and attached to the FPI device, but the pendulum was not dropped. Only rats with >10 seconds apnea after injury were included in the injury group, and only the hemisphere ipsilateral to the injury was examined.

To study the specific role of neuronal TLR4 signaling, TLR4[f/f]:CaMKIIa-CreERT2 mice (8–10-

week-old male and female mice) were induced with tamoxifen and subjected to lateral FPI

(1.5 atm) or sham injury [31,32]. Tamoxifen (75mg/kg, i.p.) was prepared in 10% ethanol and

90% corn oil by sonicating at 37°C for 90 minutes. Mice received 3 doses of tamoxifen, 24

hours before injury, 30 minutes after injury and 24 hours after injury. TLR4 deletion in

glutamatergic neurons in dentate gyrus was confirmed through immunohistochemistry (see

immunohistochemistry methods below). Cre negative mice were used as genotype

controls. Animals were treated with vehicle (DMSO), TLR4 antagonist (TAK-242, 0.5mg/kg,

i.p., Invivogen, San Diego, CA) or MMP-9 antagonist (SB-3CT, 50mg/kg, i.p., SelleckChem,

Houston, TX) at 30 minutes, 8 hours, and 24 hours after injury.

Ex vivo electrophysiology

One-week after FPI or sham injury, animals were anesthetized with isoflurane and

decapitated. Coronal hippocampal slices (300 μm) were prepared in ice-cold sucrose

artificial cerebrospinal fluid (aCSF) containing (in mM): 85 NaCl, 75 sucrose, 24 NaHCO3,

25 glucose, 4 MgCl2, 2.5 KCl, 1.25 NaH2PO4, and 0.5 CaCl2 [33]. Slices were incubated at

32°C for 20 min in an interface holding chamber containing an equal volume of sucrose-

aCSF and recording aCSF containing (in mM) 126 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 1.25

NaH_2PO_4, 26 $NaHCO_3$ and 10 D-glucose. All solutions were saturated with 95% O_2 and 5%

CO_2 and maintained at a pH of 7.4. Slices were allowed to equilibrate for an hour at room

temperature before recordings. Whole-cell voltage- and current-clamp recordings from

dentate granule cells (DGCs) were obtained at 33-34°C under IR-DIC (Olympus BX51 with

40x water immersion objectives) using 3-6MΩ glass pipettes fabricated using dual stage

glass micropipette puller (Narishige PC-10, Amityville NY). Data was recorded using

MultiClamp 700B amplifiers, digitized using Digidata 1440A and acquired using pClamp10

(Molecular Devices, San Jose, CA) at 10-kHz sampling frequency. Spontaneous and

miniature excitatory/inhibitory currents from the same DGCs were obtained using a cesium-

based internal solution (in mM: Cesium methanesulfonate 140, NaCl 5, HEPES 10, EGTA

0.2, Mg2ATP 2, Na2GTP 0.2, Qx314 5, (pH 7.25, 270-285 mOsm). Biocytin (0.2%) was

included in the internal solution for post hoc cell identification. Spontaneous and miniature

excitatory postsynaptic potentials (sEPSC and mEPSC) were recorded from the same

neurons at a holding potential of -70 mV, and spontaneous and miniature inhibitory

postsynaptic potentials (sIPSC and mIPSC respectively) were recorded at a holding

potential of 0 mV. Action potential independent miniature currents were isolated using

tetrodotoxin (TTX, 1µM). In a subset of experiments TLR4 signaling was selectively activated

by incubating slices in LPS-B5 ultrapure (Escherichia Coli serotype O55:B5, 10 µg/ml,

Invivogen, San Diego, CA), a selective TLR4 agonist, for 60-90 minutes before recording [34].

Data from cell in which access resistance changes over 20% during recordings and

recordings with access resistance above 25 MΩ were excluded from analysis.

Spontaneous/miniature synaptic currents (100 events/cell) were detected using template

search feature in Easy Electrophysiology (version 2.6., London, UK.) by a blinded investigator

and interevent intervals and amplitudes were analyzed as cumulative distribution in

Graphpad Prism 10.0.1 (Boston, MA).

In vivo electrophysiology

One-week after FPI/Sham injury rats were anesthetized under urethane (1.5g/kg i.p.) and head-fixed onto a stereotaxic frame. Under aseptic conditions, rats were implanted with a 50 µm tungsten wire depth electrode (California Fine Wire company, Grover beach, CA) in the granule cell layer (AP: 3-4 mm, ML: 2 mm, DV: 3-3.5 mm) guided by periodic electrical stimulation of the angular bundle (ML:4-4.2 mm, DV: 1.8-2 mm; pulse width 150µs delivered at 0.1Hz). Two screw electrodes on the contralateral side served as reference and ground. The position of electrodes was secured using dental cement. Acute recordings from urethane anesthetized rats started 60 minutes after induction. A current-response curve was derived by electrical stimulation of the angular bundle (150 µs @ 0.1Hz). For assessing Long Term Potentiation (LTP), DG LFP responses were evoked every 30 seconds for a period of 30 minutes to derive baseline. LTP was induced by Theta Burst Protocol[35] and responses were recorded for an additional 180 minutes. Signals were amplified 100 times (Pinnacle systems, Lawrence, KS), sampled at 10kHz, digitized using PowerLab 16/35 and recorded using LabChart8 software (AD Instruments, Colorado Springs, CO). Analysis was conducted using LabChart modules. Dentate spikes were recorded from urethane anesthetized rats (1.5g/kg i.p.) using 16 Ch silicon probe (A1x16-5mm-100-703 CM16, Neuronexus, Ann Arbor, MI). Signals were amplified 100 times using Model 3600 amplifier (AM-Systems, Sequim, WA), digitized using PowerLab 16/35 and recorded using Labchart8 Software (AD instruments, Colorado Springs, CO). Dentate spikes were detected using Toothy analysis pipeline[36] (https://github.com/Farrell-Laboratory/Toothy). Briefly, large positive events were identified using a 4.5 S.D. threshold. Detected events were visually

confirmed by a blinded investigator and the total number of dentate spikes recorded in a 30-

minute period were counted. We did not classify dentate spikes as type 1 or type 2.

In Situ Zymography

MMP-9 activity was evaluated in hippocampal slices obtained from rats and mice 48 hours

after injury using a modified in situ zymography protocol [37]. Briefly, sham/injured rats or

TLR4$^{f/f}$:CaMKIIa-CreERT2 mice and Cre^{-ve} controls were transcardially perfused using an

alcohol-based formalin free tissue fixative (Accustain, A5472 Sigma-Aldrich, St.Louis, MO),

embedded in 2% agar and coronal slices (50μm, VF-310-0Z Compresstome, Precisionary

Instruments, Ashland, MA) were prepared at <4°C to preserve enzyme activity. Slices were

then incubated with a fluorogenic Dye Quenched gelatin substrate (1:50 dilution, Invitrogen,

Waltham MA) for 40 min at 37°C, washed with PBS, fixed in 4% paraformaldehyde and

mounted on slides in Vectashield (Vector Labs, Newark, CA). The specificity of reactions

was confirmed by adding MMP-9 inhibitor SB-3CT to control slices (see supplementary

figure 2J-K). Semiquantitative analysis of DQ fluorescence in DG molecular layer were

performed using ImageJ by a blinded investigator. Data was collected from 2-4 slices /

animals and averaged. Intensity measures were normalized to CA1 stratum radiatum since

we did not observe any effect of injury, treatment or genotype in this region (Supplementary

figure 2B-C).

Single molecule fluorescent in situ hybridization (RNAscope)

To visualize TLR4 in mice and MMP-9 mRNA in rat, we used RNAscope Multiplex Fluorescent

in situ hybridization on cryosectioned slices (20μm). RNAscope Multiplex Fluorescent

Reagent kit v2 assay (ACD, Newark, CA) was used following the protocol for fixed tissue [38].

The following kits were used for this study: RNAscope Multiplex Fluorescent Reagent kit

(Cat#323100), RNAscope Probe-MM-TLR4 (Cat#316801) and RNAscope probe-Rn-MMP-9

(Cat#532281). Slices were coverslipped with DABCO with DAPI mounting media and imaged

using a Zeiss Axioscope and analyzed using QuPath software by a blinded investigator.

Immunohistochemistry

Male and female TLR4$^{f/f}$:CaMKIIa-CreERT2 mice (8-10 weeks) were induced with tamoxifen

as described above and subjected to Sham or FPI. Three days after injury, brain tissue was

harvested following transcardial perfusion with PBS followed by 4% PFA. Brains were post-

fixed overnight and then transferred to 30% sucrose until fully equilibrated. Tissue was

cryosectioned at 20 µm thickness and mounted on Superfrost Plus glass slides. For

immunohistochemistry, slides were brought to room temperature to remove OCT and

washed three times in PBS. Sections were permeabilized with 0.2% Triton X-100 in PBS for

45 minutes at room temperature, followed by blocking with 20% goat serum for 45 minutes.

Sections were then incubated overnight at 4 °C with a primary antibody against TLR4 (mouse

anti-TLR4; Santa Cruz Biotechnology; 1:250). After three PBS washes, sections were

incubated with goat anti-mouse Alexa Fluor 546 secondary antibody (1:1000, Thermo Fisher,

S11227) for 1 hour at room temperature. Slides were washed three times with PBS and

mounted using an antifade mounting medium. Images were acquired using a Zeiss

Axioscope-5 with stereo investigator (MBF Bioscience) from three animals per group, with

six sections analyzed per animal. Quantification was performed using ImageJ software by

calculating the Corrected Total Cell Fluorescence (CTCF) in the granule cell layer and hilus

for each section (Supplementary figure 2C-E).

Behavioral Experiments

Rats were trained on Barnes Maze task to assess spatial learning 1-month post-injury. The

Barnes maze table (Maze Engineers, https://conductscience.com/maze/) was 122 cm in

diameter with 20 holes (10 cm each). The target hole was equipped with an escape box (22.5

cm x 14 cm x 9 cm LxWxH). The maze table was set up within black curtain walls with visual

cues and a camera installed above to record animal movement. Two bright floodlights and

a fan served as aversive stimuli to encourage animals reach the escape box. Animals were

held in their home cage outside the curtain in a dark room until their turn to run the trial. On

day 1, rats were habituated to the behavior room for at least 1 hour and then placed in the

middle of the table inside starter cup for 1 minute following which they were allowed to

explore the table for 180 seconds. At the end of the trial, animals were gently guided to their

designated escape hole. The escape hole locations were randomly assigned to each rat and

remained the same till the end of the experiment. The table was thoroughly wiped with

ethanol between tests to eliminate odor cues. Rats were subjected to a single 180-second

trial for 3 days. Task acquisition and retention were confirmed through tests on day 4 and

day 7. Four rats that fell from the table and exhibited signs of lethargy were removed from

the test and sacrificed. Behavioral analysis was performed by a blinded experimenter using

Any-maze software. Parameters such as latency to escape hole, distance travelled and

spatial search strategy were evaluated between sham and injured rats. Spatial search

strategies were scored based on 11-point scoring scheme[39]. Thigmotaxis was quantified

by defining a peripheral zone of the testing arena that was approximately 12 cm from the

apparatus's outer edges. The proportion of time the animals spent within this designated

zone was recorded and used as the metric for thigmotaxis.

Statistics

Data are presented as mean ± standard error of the mean (SEM) for parametric measures,

or as median and interquartile range (IQR) where non-parametric distributions were

observed. To ensure rigorous data interpretation and minimize investigator bias, all data

analysis, including quantification of in situ zymography, electrophysiological recordings,

and behavioral scoring, was performed by researchers blinded to the treatment groups and

genotypes.

Normality of data distributions was assessed using the Shapiro-Wilk test. For the analysis

of synaptic events (sEPSC, mEPSC, sIPSC, and mIPSC), cumulative probability distributions

of inter-event intervals (IEI) and amplitudes were compared across groups using equal

number of consecutive events with the non-parametric Kolmogorov-Smirnov (K-S) test.

Effect sizes for these distributions were calculated using Cohen's d. Differences between

two groups were evaluated using two-tailed unpaired t-tests. For experiments involving

multiple factors, such as injury and drug treatment or injury and genotype, two-way ANOVA

or two-way repeated measures (RM) ANOVA (for input/output curves and behavioral training

days) were employed, followed by Šídák's multiple comparisons test where appropriate.

Significance was defined at $P<0.05$. Statistical analyses were performed using Graphpad

Prism 10.0.1 (Boston, MA).

RESULTS:

Neuronal TLR4 signaling is necessary for early excitatory circuit changes after brain injury.

Prior studies have identified TLR4 contribution to early changes in glutamatergic synapse strength after brain injury [12–14]. However, the nature of early circuit reorganization after brain injury in the dorsal hippocampus and whether immune signaling regulates these changes has not been examined. Whole-cell voltage clamp recordings from DGCs in dorsal hippocampal slices one-week after FPI revealed an early decrease in the inter-event intervals (IEI) of spontaneous excitatory postsynaptic currents (sEPSCs), indicating an increase in sEPSC frequency (IEI in milliseconds: Sham$_{DMSO}$ mean – 336.9±13.93, median-216.4, IQR- 86.05-429.9; FPI$_{DMSO}$ mean – 225.3±8.26, median-144.8, IQR- 64.60-307.4., n = 8 cells/ group, $P<0.0001$ by K-S test, Cohen's d = 0.34, Figure 1A-E). Additionally, we observed a reduction in sEPSC amplitude in DGCs from rats after brain injury (Amplitude in pA: Sham$_{DMSO}$ mean – 13.16±0.17, median-12.20, IQR- 9.98-15.36; FPI$_{DMSO}$ mean – 11.23±0.16, median-10.31, IQR- 8.05-13.42, n = 8 cells/ group, $P<0.0001$ by K-S test, Cohen's d = 0.41). To assess potential post-traumatic synaptic reorganization, we recorded action potential-independent miniature EPSCs (mEPSCs) in DGCs the presence of tetrodotoxin (TTX, 1 μM). Similar to sEPSCs, DGCs in rats one week after FPI exhibited shorter mEPSC IEI and smaller mEPSC amplitudes compared to Sham controls (IEI in milliseconds: Sham$_{DMSO}$ mean: 824.8±34.17, median: 496.1, IQR:208.1-1096; FPI$_{DMSO}$ mean: 642.4±28.91, median: 396.4, IQR: 158.2- 858.6; Amplitude in pA: Sham$_{DMSO}$ mean: 14.16±0.25, median:12.32, IQR: 8.90-17.91; FPI$_{DMSO}$ mean: 9.28±0.16, median: 8.02, IQR:

6.55-10.55, n = 7 cells/ group, P <0.001 by K-S test, Cohen's d = 0.22 for IEI and 0.85 for

amplitude, Supplementary figure 1A-C). Together, these findings provide the first evidence

of an increase in excitatory synaptic inputs to DGCs in dorsal hippocampus of rats one

week after injury.

Significant changes in synaptic inhibition is often observed early after brain injury [5,7,40].

To evaluate whether inhibitory circuits are reorganized alongside the excitatory circuits, we

examined spontaneous inhibitory postsynaptic potentials (sIPSC) in the same neurons. In

contrast to changes in sEPSCs, sIPSC IEI was prolonged in DGCs from injured rats (IEI in

milliseconds: Sham$_{DMSO}$ mean: 190.3±6.64, median:128.1, IQR: 65.35- 252.4; FPI$_{DMSO}$ mean:

256.6±9.272, median:181.1, IQR:83.78-320.4; n= 8 cells/group, P <0.0001 by K-S test,

Cohen's d = 0.29 Figure 1F-H). These findings indicate that in addition to receiving more

frequent excitatory inputs, the DGCs receive fewer inhibitory inputs after brain injury. DGCs

also showed an increase in sIPSC amplitude compared to sham controls after brain injury

(amplitude in pA: Sham$_{DMSO}$ mean: 16.66±0.35, median:13.78, IQR:10.81-19.21; FPI$_{DMSO}$

mean: 18.24±0.35, median: 15.74, IQR: 11.94-21.70, n = 8 cells/ group, P <0.0001 by K-S

test, Cohen's d = 0.16). These findings differ from our findings in the ventral hippocampus

demonstrating an increase in sIPSC frequency in DGCs after FPI [33,41,42]. We also

observed an increase in mIPSC amplitudes with no change in mIPSC IEI (Supplemental

Figure 1D-F). Together, these findings indicate an early dysregulation in excitatory and

inhibitory inputs to DGC's one-week after FPI. Furthermore, these results identify early

circuit reorganization after FPI and reveal potential differences in synaptic inhibition and

post-traumatic plasticity in the dorsal vs ventral hippocampus.

To determine whether inhibition of TLR4 signaling early after injury modified DGC synaptic

currents, we treated sham and FPI rats with TLR4 inhibitor TAK-242 (0.5mg/kg, i.p, 30 min, 8

hours and 24 hours post-injury). TAK-242 treatment reduced changes in sEPSC and mEPSC

frequency and amplitude in injured rats (Figure.1A-E, Supplementary figure 1A-C, and

Supplementary table.1). In addition, TAK-242 reduced injury-driven changes in sIPSC

frequency and mIPSC amplitude (Figure.1F-H, Supplementary figure 1D-F).   These findings

demonstrate a role for TLR4 signaling in driving early synaptic dysregulation after FPI.

In contrast to injured animals, sham controls receiving TAK-242 showed paradoxically

opposing changes excitatory and inhibitory synaptic inputs to DGCs. In sham controls,

TAK-242 treatment reduced sEPSC IEI in DGCs, increasing frequency as observed after FPI

but failed to alter sEPSC amplitude (Figure 1B-E). Moreover, TAK-242 treated sham controls

showed a reduction in DGC sIPSC IEI and amplitude suggesting more frequent inhibitory

inputs (Figure 1F-H). Similar to sEPSCs, both mEPSC interval and amplitude were reduced

in TAK-242 treated sham controls (Supplementary figure 1A-C) consistent with synaptic

reorganization. However, Tak-242 treatment did not alter mIPSC frequency or amplitude.

Together, these findings identify a novel contribution of basal TLR4 signaling in maintaining

excitatory synaptic homeostasis in uninjured animals.

Our previous studies demonstrated neuronal TLR4 signaling as critical for post-traumatic

increase in DGCs AMPA currents [13]. To test whether the glutamatergic circuit changes

identified above require neuronal TLR4 signaling, we examined changes in sEPSC

frequency and amplitude in mice with selective deletion of TLR4 in CaMKII expressing

excitatory neurons (TLR4$^{fl/fl}$-CaMKIIa-CreERT2 mice). Care was taken to avoid constitutive

and developmental effects by inducing Cre recombinase just prior to injury.

Immunostaining for TLR4 in mice three-days after FPI, following the 3-day tamoxifen

induction strategy, revealed strong immunostaining for TLR4 in the granule cell layer and

hilus of Cre^{-ve} mice which was absent in Cre^{+ve} mice, confirming TLR4 deletion in DGCs (n =

3 mice/ group, $P<0.05$ by Nested t test, Supplementary figure 2A-C). Examination of active

and passive properties of DGCs revealed a significant decrease in capacitance and an

increase in firing frequency in Cre^{+ve} sham mice compared to Cre^{-ve} sham mice (n = 7 cells/

group, $P<0.05$ by two-way ANOVA, Supplementary figure 2E&I). We did not observe any

genotype effects on the resting membrane potential, input resistance or rheobase of DGCs

in sham controls (Supplementary figure 2F-H). Consistent with our previous reports in rat

and mice [31,43], there were no injury effects on DGC active properties in either genotype.

However, while there was no injury effect on DGC resting membrane potential in Cre^{-ve}

mice, DGCs from Cre^{+ve} mice with TLR4 deletion were more depolarized (Supplementary

figure 2F). Similar to our findings in rat, brain injury reduced DGC sEPSC IEI and amplitude

in Cre^{-ve} mice with intact TLR4 signaling (See Supplementary table 1; n = 7 cells/ group,

$P<0.0001$ by K-S test, Figure 1I-L). Additionally, DGC sEPSC IEI and amplitude were not

different between Cre^{-ve} and Cre^{+ve} sham injured mice. Importantly, although DGCs from

Cre^{+ve} mice with conditional deletion of TLR4 in excitatory neurons showed a small

decrease in sEPSC amplitude, they failed to exhibit injury-induced decrease in sEPSC IEI.

These findings demonstrate that TLR4 expression in glutamatergic neurons is necessary for

the early injury-driven changes in excitatory synaptic inputs to DGCs.

378 **MMP-9 activity is required for TLR4-dependent changes in DGC excitatory synapses**

379 **after brain injury**

380 We next sought to identify downstream molecular mechanisms through which TLR4

381 signaling dysregulates DGC synaptic inputs after brain injury. Several studies have reported

382 an upregulation in Matrix Metallproteinase-9 (MMP-9) expression in pre-clinical models of

383 TBI [27,30,44]. MMP-9 is a critical regulator of synaptic strength and plasticity [17,18], and

384 perturbations in MMP-9 activity is known to dysregulate synaptic physiology [19,37,45–48] .

385 Although extrinsic activation of TLR4 signaling has been shown to enhance MMP-9

386 expression in primary astrocyte/neuronal co-cultures and *in vivo* [15,16], whether TLR4

387 regulates MMP-9 activity and if this can induced synaptic and functional changes *in vivo*

388 and in the injured brain remains to be determined. Moreover, whether *neuronal* TLR4

389 regulates MMP-9 activity has not been examined.

390 To directly assess MMP-9 activity, we performed *in-situ* zymography on hippocampal slices

391 obtained from sham/FPI rats 48 hours after injury, focusing on the DG molecular layer

392 where the afferent glutamatergic synapses are located. DQ-fluorescence intensity values

393 were normalized to CA1 Stratum Radiatum (SR) since no changes were observed in this

394 region with injury, treatment or genotypes (Supplementary figure 2L-M, see methods). Our

395 experiments identified a significant increase in the MMP-9 activity in the DG molecular

396 layer in brain-injured rats compared to sham controls (MMP-activity normalized to CA1 SR:

397 Sham:14.56±2.15%, FPI:30.43±2.89%, *n* = 5 sham and 6 FPI rats, Figure 2A-C).

398 Pharmacologically blocking TLR4 early after FPI reduced MMP-9 activity in the DG of brain

399 injured rats (Sham$_{TAK-242}$: 17.22±2.68%, FPI$_{TAK-242}$: 16.36±2.97%, *n* = 5 Sham and 6 FPI rats;

Effect of Injury x Treatment F (1,18) = 9.13; P <0.05 by Two-way ANOVA). In contrast, MMP-9

activity in sham controls treated with TAK-242 was not different from vehicle treated sham

controls. These results identify TLR4 regulation of MMP-9 activity in the injured brain.

To isolate the role of neuronal TLR4 in regulating MMP-9 activity, we performed in-situ

zymography on mice with neuronal TLR4 deletion (TLR4[fl/fl]-CaMKIIa-CreERT2) and Cre[-ve]

controls (Supplementary figure 2A-C) 48 hours after Sham or FPI. We did not observe any

difference in MMP-9 activity between Cre[-ve] sham and Cre[+ve] sham mice, indicating that

TLR4 deletion does not alter normal MMP-9 levels without injury (Figure 2D-F). While Cre[-ve]

mice showed an increase in MMP-9 levels after injury, mice with selective TLR4 deletion in

glutamatergic neurons failed to develop elevated MMP-9 activity after FPI (MMP-activity

normalized to CA1 SR: Cre[-ve] Sham: 13.43±3.36%, Cre[-ve] FPI: 42.79±5.02%; Cre[+ve] SHAM:

10.59±3.15%, Cre[+ve] FPI: 12.20±1.16%., n=3-4 mice/group; Effect of Injury x Genotype, F

(1,11) = 14.70; P<0.05 by Two-way ANOVA, Figure 2D-F). These findings demonstrate that

neuronal TLR4 is necessary for the post-injury upregulation of MMP-9 activity.

Next, we tested the possibility that TLR4 signaling increased MMP-9 levels through

transcriptional upregulation. Naïve rats (P30) were injected with a selective TLR4 agonist,

LPS-B5 (5mg/kg., i.p.) or saline and euthanized after six hours for single molecule

fluorescent in situ hybridization (smFISH) to quantify MMP-9 mRNA levels. Sections from

LPS-B5 injected rats showed a significant increase in MMP-9 mRNA in DGCs, indicating

that TLR4 activation enhances MMP-9 via transcriptional upregulation (intensity A.U/μm^2:

saline controls 1.67±0.15; LPS-B5 treated 2.41±0.19; n= 5/group, Figure 2H-J).

Furthermore, inhibition of NF-κB, a key downstream effector of TLR4, using JSH-23 (10

mg/kg) administered one hour before LPS-B5 injection abolished the upregulation of MMP-

9 (intensity A.U/μm^2 : JSH-23 + LPS-B5 1.319±0.16, n = 5 rats, Main effect F(2,12) =11.20, *P*

= 0.002, Figure 2G-I) Together, these findings identify TLR4 and NF-κB dependent

transcriptional upregulation of MMP-9 as a candidate mechanism for increase in MMP9

activity in the DG following brain injury.

Extracellular matrix (ECM) remodeling by MMP-9 has been shown to play a crucial role in

plasticity of excitatory and inhibitory synapses [49]. Therefore, we reasoned that TLR4

mediated increase in MMP-9 activity may be required for the post-traumatic changes in

synaptic inputs to DGCs. To examine the acute effects of TLR4 signaling *ex vivo*, we

compared sEPSCs in hippocampal slices from sham rats incubated in LPS-B5 (10 μg/ml for

90min) or aCSF. Acute LPS-B5 incubation significantly reduced sEPSC IEI and increased

sEPSC amplitude in DGCs mimicking the FPI phenotype (IEI in milliseconds: LPS-B5

incubation mean: 265.3±10.99, median:157.3, IQR: 69.18-330.4; Amplitude in pA: mean –

17.86±0.21, median-17.29, IQR- 14.06-21.09; *n* = 8 cells from 4 rats, Sham vs LPS-B5

incubated *P* <0.0001 by K-S test, Figure 3A-D). Unlike sEPSCs, acute TLR4 activation failed

to alter sIPSC frequency or amplitude in DGCs (Supplementary figure 3A-C). Importantly,

inhibiting MMP-9 using SB-3CT (10 μM) mitigated LPS-B5-induced changes in sEPSC IEI

and amplitude, indicating that TLR4-mediated alterations in excitatory synaptic inputs to

DGCs require MMP-9 activity (IEI in milliseconds:SB-3CT+LPS-B5: mean: 339.7±12.40,

median:230.9, IQR: 94.53-434.6; amplitude in pA: mean: 15.90±0.19, median:14.66, IQR:

11.53-18.96; *n* = 9 cells from 4 rats, LPS-B5 incubated vs SB-3CT+LPS-B5 *P* <0.0001 by K-S

test, Figure 3A-D).

Since we find that TLR4-mediated alterations in DGC sEPSC frequency requires MMP-9

activity, we tested if systemic inhibition of MMP-9 using SB-3CT treatment (50 mg/kg, i.p.)

early after FPI would prevent post-traumatic changes in excitatory inputs to DGCs. SB-3CT

treatment effectively mitigated injury-induced the changes in sEPSC IEI and amplitude

one-week after FPI in rat (see Supplementary table.1, Figure 3E-H). Consistent with our

findings in rat, SB-3CT treatment reduced post-injury changes in sEPSC IEI in DGCs from

Cre^{-ve} -negative TLR4$^{fl/fl}$-CaMKIIa-CreERT2 mice (Figure 3I-L). Note that Cre^{+ve} TLR4$^{fl/fl}$-

CaMKIIa-CreERT2 mice did not show difference in sEPSC intervals or amplitudes after FPI.

SB-3CT treatment did not have any effect on sIPSC frequency or amplitude (Supplementary

figure 3G-I) after FPI but enhanced sIPSC amplitude in sham controls suggesting that TLR4

regulation of DGC inhibitory synaptic parameters after injury does not require MMP-9

activation. Together, these results indicate that injury-driven changes in excitatory inputs to

DGCs required both TLR4 signaling and MMP-9 activity.

Early activation of TLR4 - MMP-9 molecular axis contributes to DG network

dysfunction after brain injury

Having established that the TLR4- MMP-9 mechanistic axis increases glutamatergic inputs

to DGCs after brain injury, we examined the contribution of this molecular pathway to early

changes in DG circuit function in rats *in vivo* one-week after brain injury. Input/output plots

of the DG population spike amplitude elicited in response to perforant path stimulation at

increasing current intensities, recorded under urethane anesthesia and normalized to peak

amplitude, revealed a leftward shift after FPI indicating increased network excitability ($n = 6$

SHAM and 5 FPI rats; Effect of injury – $F_{(1,9)} = 6.462$, $P = 0.031$ by Two-way RM ANOVA,

Figure 4A-C). This increase in network excitability was mitigated by early inhibition of either

TLR4 signaling or MMP-9 activity, demonstrating their contribution in promoting post-

traumatic DG circuit dysfunction (n= 5-6 rats/group; Effect of Stimulus x Treatment: F

(6,166,40.08) = 4.752, P=0.0009 by Two-way RM ANOVA, Figure 4D). Paradoxically, as

previously reported in slices [12–14], TAK-242 treatment increased DG population spike

amplitude in sham controls (Figure 4E). However, consistent with its lack of effect on EPSC

frequency in sham controls, SB-3CT treatment did not alter DG network excitability in sham

rats (Figure 4E).

Enhanced network excitability combined with deficits in synaptic inhibition can increase

dentate throughput. We assessed DG throughput by examining dentate spikes, a distinct

DG population event driven by entorhinal cortex[50,51] . Linear 16- channel silicon probe

recordings of dentate spikes in anesthetized rats (Figure 4F-G) revealed a significant

increase in the frequency of dentate spikes one-week after FPI (in Hz: Sham: 0.11±0.019;

FPI 0.24±0.025, n = 4-5 rats/group, P= 0.006 by two tailed unpaired t-test), identifying

enhanced dentate throughput following FPI. To test if inhibition of either TLR4 signaling or

MMP-9 activity early after injury would reduce changes to DG throughput, we examined

dentate spikes in TAK-242 or SB-3CT treated sham/injured rats. Our experiments

demonstrate that early inhibition of either TLR4 signaling or MMP-9 activity effectively

prevented the post injury increase in dentate spike frequency without impacting it in sham

controls (n = 4-5 rats/group; Effect of Interaction F (2,21) = 9.975, P<0.001 by Two-way

ANOVA, Figure 4G).

We previously demonstrated that early post-injury increase in DGC calcium permeable

AMPA currents and deficits in working memory were mitigated by TLR4 inhibition [13]. To

assess whether the TLR4-MMP9 signaling impacts synaptic plasticity, long-term

potentiation (LTP) at perforant path-DGC synapses in anesthetized rats *in vivo* using theta-

burst stimulation (TBS) [35]. In the first 10 minutes after TBS, the field EPSP slope

normalized to the maximum slope at baseline trended lower but was not statistically

different between sham and FPI. However, there was a reduction in the field EPSP slope at

60 min, 120 min, and 180 min post-TBS indicating impairment in LTP maintenance (n = 6

rats/group, Two-way RM ANOVA with Šídák's multiple comparisons 10 min: P =0.065, 60

min: P = 0.003, 120 min: P = 0.002, 180 min: P = 0.125, Figure 5A-G). Crucially, early in vivo

antagonism of either TLR4 or MMP-9 early after FPI effectively prevented deficits in LTP in

injured rats, demonstrating contribution of TLR4-MMP-9 axis to LTP deficits after brain

injury (See Supplementary table 2, Figure 5A-G). Consistent with the paradoxical effects of

TAK-242 on sEPSC and network excitability in sham controls, TAK-242 treatment resulted in

deficits in LTP in sham rats (Figure 5F-G). Once again, unlike TAK242, SB-3CT treatment

failed to impact LTP Sham rats, paralleling on the differential effects of the treatments on

sEPSC frequency, population spike amplitude and dentate spike frequency in sham

animals. These findings demonstrate that TLR4-MMP-9 dependent post-traumatic circuit

changes compromise DG glutamatergic circuit function and plasticity.

Targeting TLR4 or MMP-9 axis early after brain injury limits long-term spatial memory

impairments.

Since we identify post injury increase in dentate spikes which can compromise hippocampal

information processing [52–55], and blocking TLR4 and MMP-9 early after injury prevent this

change, we sought to determine if TLR4 and MMP-9 antagonists could alleviate long term

cognitive deficits after brain injury. We adopted a Barnes maze spatial learning task which

has been shown to rely on intact DG function[56,57]. One month after sham/FPI, brain

injured rats exhibited a longer latency to find the escape hole and traveled a greater total

distance compared to sham controls, early during training but reached control levels of task

performance by Day 3 (See Supplementary table 3, n =10 rats / group, $P<0.05$ by TW-RM

ANOVA, Figure 6A-E). Additionally, there was no difference in task performance during trials

on Day 4 and Day 7, indicating that task retention is not impaired (Supplementary figure 4A-

D). Although there were no differences in speed of locomotion (Supplementary figure 4E),

injured rats showed increased thigmotaxis on Days 1 and 2 (Time spent in thigmotaxis zone

in seconds: $Sham_{DMSO}$ – 24.73±5.42, FPI_{DMSO} – 64.99±16.96; n= 10 rats/group, Effect of injury–

F (1,18) = 6.841, $P<0.05$ by Two-way -RM ANOVA, Figure 6, E, I). Since injured rats showed a

significant increase in both thigmotaxis and latency to find the escape on Day 2 of task

acquisition, we examined the use of spatial strategy for navigation using a 11-point scoring

system [39]. Analysis of search strategies on Day 2 showed that injured animals employed

random or serial strategies more often, demonstrating specific deficits in ability to recall

spatial cues for navigation (Search strategy score_Day2: $Sham_{DMSO}$ – 7.7±0.73, FPI_{DMSO} –

2.4±0.47; n= 10 rats/group, Effect of injury– F (1,18) = 17, $P<0.01$ by Two-way RM ANOVA,

Figure.6A-C). These results suggest a mild delay in task acquisition after brain injury which

is apparent on Day 2 of the task.

We focused our analysis on Day 2 of task acquisition when the injury-induced impairment is

most apparent. Early inhibition of TLR4 signaling or MMP-9 activity prevented post-injury

changes in latency to escape, and distance traveled observed on Day 2 (n = 6-10 rats/group;

P<0.05 by Two-way RM ANOVA, Supplementary table 3 and Figure 6F-G). Furthermore, the

search strategy scores significantly improved with MMP-9 inhibition. While TLR4 inhibition

trended to improve the strategy score, the difference did not reach statistical significance

(Figure 6H). Importantly, while MMP-9 did not have any adverse effects on sham controls,

TLR4 inhibition showed a trend towards increased latency, distance travelled and

thigmotaxis, and a lower spatial search strategy score (Figure 6F-I) which did not reach

statistical significance. Collectively, these findings demonstrate that early inhibition of the

TLR4-MMP-9 signaling axis effectively mitigates mild long term spatial learning deficits

observed after FPI.

DISCUSSION:

Understanding how inflammatory pathways modulate neuronal function after brain injury

is critical for developing effective therapeutic strategies to minimize the long-term

sequelae of TBI. While widespread secondary injury is driven by glial mediators of

neuroinflammation [1,2], we previously showed that the neuronal expression of innate

immune receptor TLR4 is upregulated in the hippocampus early after injury, contributing to

long-term seizure susceptibility and impaired working memory [12–14]. Here, we provide

mechanistic evidence that TLR4 signaling mediates rapid, circuit-level changes in both

excitatory and inhibitory synaptic transmission after injury. Using a combination of

pharmacology in rats and mouse genetics we demonstrate a central role for TLR4

expressed in CaMKII-positive excitatory neurons in mediating the observed circuit changes

through the upregulation of MMP-9. Importantly, we show that TLR4 or MMP-9 inhibition

early after brain injury not only reduces these pathological changes one-week after injury,

but improves deficits in spatial learning 1-month after injury.

TLR4 Signaling Augments Glutamatergic inputs to DGCs after brain injury.

TLR4 signaling is well established as a driver of neuronal excitability [3,58–62]. Our prior

work showed that TLR4 signaling enhances AMPA currents in granule cells contributing to

DG network hyperexcitability [12,13]. Examining circuit-level changes in injured animals

one week after injury, this study identifies an aberrant increase in excitatory inputs to DGCs

in the dorsal hippocampus. Considering that perforant path fibers are severed in coronal

hippocampal slice preparations [63], the post-injury increase in spontaneous EPSCs and

action potential-independent miniature events observed here indicate an increase in

synaptic contacts, consistent with early axonal sprouting [42], or elevated mossy cell

excitability previously reported after FPI [64]. Our findings highlight a critical role of TLR4

signaling in this process. First, we show that early pharmacological inhibition of TLR4

mitigates changes in both spontaneous and miniature EPSCs. We further demonstrate that

transient activation of TLR4 signaling in acute brain slices using LPS-B5 is sufficient to

augment sEPSCs frequency to levels comparable to those seen in injured animals.

Furthermore, by leveraging TLR4$^{fl/fl}$: CaMKIIa-CreERT2 mice, we demonstrate that selective

deletion of TLR4 in glutamatergic neurons alone is sufficient to mitigate the post-injury

increase in sEPSC frequency. Together, these findings show that neuronal TLR4 is both

necessary and sufficient to enhance glutamatergic drive to DGCs after brain injury.

A Novel Neuro-Immune-Extracellular Matrix Signaling Cascade Underlies Early Post-

Traumatic Glutamatergic Circuit Plasticity

To mechanistically elucidate how TLR4 enhances glutamatergic drive to DGCs after injury,

we focused on MMP-9 as a potential downstream effector. MMP-9 co-localizes with

excitatory synapses in an activity-dependent manner, and increased MMP-9 activity is

known to dysregulate glutamatergic synaptic transmission [37,49,65–67]. We previously

reported that neuronal TLR4 enhances calcium-permeable AMPA currents in DGCs post-

injury by increasing the surface expression of GluA1 subunits [13]. Notably, MMP-9 activity

has been shown to regulate AMPA receptor mobility and synaptic accumulation of GluA1

subunits following chemically induced LTP [68]. In addition, TLR4 ligands such as high-

mobility group box 1 (HMGB1) and lipopolysaccharide (LPS) are known to enhance MMP-9

expression in cultures and following ischemic insult [15,16].

Here, we show that pharmacologically inhibiting TLR4 signaling (rats) or deleting TLR4 in

excitatory neurons (mice) both limit post-injury increases in MMP-9 activity. By using LPS-

B5 to selectively activate TLR4, we demonstrate that activation of TLR4 alone increases

MMP-9 mRNA levels in the dentate gyrus of naïve rats in an NF-κB dependent manner.

Thus, as summarized in Figure 7, our findings identify a novel TLR4/NF-κB-dependent

transcriptional pathway that upregulates MMP-9 activity and promotes glutamatergic

circuit reorganization following brain injury. Consistent with this, inhibition of MMP-9 with

SB-3CT effectively reduced the glutamatergic changes driven by both direct activation of

TLR4 signaling in slices (LPS-B5) and in vivo following brain injury.

Mechanistic Divergence in TLR4 regulation of glutamatergic and GABAergic circuits

In contrast to the glutamatergic inputs, we observed a decrease in the frequency of

inhibitory inputs to DGCs one week after FPI. This stands in direct contrast to our previous

findings from the ventral hippocampus [33,41,42], highlighting potential regional

differences in functional plasticity between dorsal and ventral hippocampal inhibitory

circuits [69]. While TLR4 inhibition reduced post-traumatic changes in IPSCs, this effect

was independent of MMP-9 activity. Excessive MMP-9 activity was previously implicated in

PV hypofunction and perineuronal net remodeling underlying auditory hypersensitivity in

the mouse model of Fragile X Syndrome, a neurodevelopmental disorder associated with

autism [70]. However, our findings indicate a divergence of TLR4-dependent mechanisms

of circuit reorganization, with MMP-9-independent mechanisms contributing to changes in

inhibitory synapses (Figure 7). Furthermore, selective activation of TLR4 with LPS-B5 did

not alter sIPSC frequency in DGCs. Therefore, it is possible that injury-driven loss of hilar

interneurons contributes to the reduction in sIPSC after brain injury [7,43]. However, we

observed an increase in mIPSC frequency following LPS-B5 suggesting perturbations in

release probability or presynaptic calcium [40,71–73]. Future studies should determine

whether neuronal or glial TLR4 is involved in post-injury changes in inhibition and its

functional impact on dentate function and behaviors.

Network Consequences and behavioral impact of TLR4/MMP-9 pathway.

The dentate gyrus plays a central node in hippocampal memory processing, regulating the

flow of information through the hippocampal circuitry [53,74,75]. Thus, perturbations in DG

circuit function can directly impair cognitive processes. Our earlier studies showed

enhanced perforant path evoked EPSC amplitude suggesting stronger entorhinal-dentate

coupling in the injured brain and deficits in working memory [12–14]. Here we advance

these findings using in vivo approaches to identify increased dentate network excitability

early after injury. Furthermore, although the LTP induction was unaffected, LTP

maintenance was compromised in injured rats suggesting that the strengthening of

perforant path to granule cell synapses doesn't saturate synaptic strength and preclude

potentiation. Rather, we find selective deficits in LTP maintenance after brain injury

indicating altered transcriptional processes. Importantly, we report an increase in dentate

spike frequency early after brain injury indicating abnormal entorhinal-dentate throughput.

Since dentate spikes are critical in hippocampal information processing, memory

formation, and consolidation [50–54,76,77], abnormal dentate spikes could impair

memory processing after brain injury. Supporting this proposal, treatments that reduced

dentate spike frequency in injured animals also improved spatial learning deficits.

Crucially, dentate spikes are also increased in experimental models of epilepsy[78],

suggesting that these events may serve as biomarkers for post-traumatic epileptogenesis

alongside the pathological high frequency oscillations and altered sleep spindles identified

previously [79,80]. In summary, we find that inhibition of the novel TLR4-MMP9 signaling

axis after brain injury was effective in limiting post-traumatic deficits in dentate circuit

excitability, long term potentiation and spatial memory indicating therapeutic potential of

manipulating this pathway (Figure 7).

A Novel Constitutive Role for TLR4 Signaling in Basal Circuit Homeostasis

Our study reveals a fundamental role for basal TLR4 signaling in maintaining dentate circuit

homeostasis. We found that inhibition of TLR4 signaling in uninjured animals consistently

altered synaptic inputs, network excitability, and plasticity. Specifically, timed and

selective deletion of TLR4 in glutamatergic neurons in a mature circuit caused a genotype-

dependent reduction in granule cell capacitance and increased firing rates (Figure 7). This

indicates a constitutive role for neuronal TLR4 in limiting basal excitability. A reduction in

capacitance with increased firing in the absence of changes in action potential threshold

parallels observations in parvalbumin neurons in epilepsy models, where regulation of

perineuronal nets alters neuronal firing through effects on cell capacitance[38,81,82].

Indeed, our prior studies have found that ability of TLR4 blockers to increase dentate

excitability in uninjured controls is abolished when glial metabolic processes are

disrupted[14]. Our finding that TLR4 or MMP9 antagonism does not impact dentate spike

frequency in uninjured controls, is consistent with lack of TLR4 modulation of perforant

path evoked AMPA currents in the absence of injury. Nevertheless, the basal TLR4

regulation of network homeostasis appears critical as blocking TLR4 enhances network

excitability, impairs LTP maintenance and spatial memory (Figure 7). Future studies should

examine the mechanisms underlying basal TLR4 regulation of neuronal activity.

Collectively, our data establishes the TLR4-NFκB-MMP-9 signaling axis as a central

mechanism driving post-TBI pathology. By placing MMP-9 as the downstream mediator for

TLR4-induced glutamatergic synaptic and network dysfunction, these findings provide a

novel and targetable therapeutic avenue for mitigating epileptogenesis and cognitive

deficits following traumatic brain injury.

Declarations

Ethics approval and consent to participate

All experiments were performed in accordance with IACUC protocols approved by the University of California-Riverside, CA (Approval number #20210005/35) and in keeping with the ARRIVE guidelines.

Consent for publication

Not applicable.

Availability of data and materials

All data are available from the corresponding author upon reasonable request.

Competing interests

The authors declare no competing interests.

Funding

This work is supported by National Institutes of Health (NIH) NINDS R01 NS069861, R01NS097750 to V.S., NIH/NINDS F31NS124290 to L.D; CDMRP- Epilepsy Research Program grant W81XWH-21-1-0684 to D.S.

Authors' contributions

D.S, E.C and L.D performed experiments; D.S, E.C, L.D, and R.J analyzed data; D.S, I.E and V.S interpreted results of experiments; D.S prepared figures; D.S, V.S conception and design of research; D.S and V.S drafted manuscript; D.S, I.E and V.S finalized and approved manuscript.

Acknowledgements: We thank Dr. Leszek Kaczmarek (Nencki Institute, Warsaw, Poland) for his expert advice on *in situ* zymography. We also thank Dr. Amanda Schott and Dr. Jordan S Farrell (Harvard Medical School) for their expertise and help with Dentate spike detection algorithms.

REFERENCES:

1. Sharma R, Leung WL, Zamani A, O'brien TJ, Espinosa PMC, Semple BD. Neuroinflammation in post-traumatic epilepsy: Pathophysiology and tractable therapeutic targets. Brain Sciences. 2019 Nov 1;9(11).

2. Mukherjee S, Arisi GM, Mims K, Hollingsworth G, O'Neil K, Shapiro LA. Neuroinflammatory mechanisms of post-traumatic epilepsy. Journal of Neuroinflammation. 2020 June 17;17(1).

3. Webster KM, Sun M, Crack P, O'Brien TJ, Shultz SR, Semple BD. Inflammation in epileptogenesis after traumatic brain injury. Journal of Neuroinflammation. 2017 Jan 13;14(1).

4. Simon DW, McGeachy MJ, Baylr H, Clark RSB, Loane DJ, Kochanek PM. The far-reaching scope of neuroinflammation after traumatic brain injury. Nature Reviews Neurology. 2017 Mar 1;13(3):171–91.

5. Lowenstein D, Thomas M, Smith D, McIntosh T. Selective vulnerability of dentate hilar neurons following traumatic brain injury: a potential mechanistic link between head trauma and disorders of the hippocampus. J Neurosci. 1992 Dec 1;12(12):4846–53.

6. McIntosh, Saatman, Raghupathi, Graham, Smith, Lee, et al. The Dorothy Russell Memorial Lecture* The molecular and cellular sequelae of experimental traumatic brain injury: pathogenetic mechanisms. Neuropathology Appl Neurobio. 1998 Aug;24(4):251–67.

7. Toth Z, Hollrigel GS, Gorcs T, Soltesz I. Instantaneous Perturbation of Dentate Interneuronal Networks by a Pressure Wave-Transient Delivered to the Neocortex. 1997.

8. Atkins CM. Decoding Hippocampal Signaling Deficits After Traumatic Brain Injury. Transl Stroke Res. 2011 Dec;2(4):546–55.

9. Hunt RF, Boychuk JA, Smith BN. Neural circuit mechanisms of post–traumatic epilepsy. Front Cell Neurosci [Internet]. 2013 [cited 2025 Nov 13];7. Available from: http://journal.frontiersin.org/article/10.3389/fncel.2013.00089/abstract

10. Paterno R, Folweiler KA, Cohen AS. Pathophysiology and Treatment of Memory Dysfunction After Traumatic Brain Injury. Curr Neurol Neurosci Rep. 2017 July;17(7):52.

11. Neuberger EJ, Gupta A, Subramanian D, Korgaonkar AA, Santhakumar V. Converging early responses to brain injury pave the road to epileptogenesis. Journal of Neuroscience Research. 2019 Nov 1;97(11):1335–44.

12. Li Y, Korgaonkar AA, Swietek B, Wang J, Elgammal FS, Elkabes S, et al. Toll-like receptor
4 enhancement of non-NMDA synaptic currents increases dentate excitability after
brain injury. Neurobiology of Disease. 2015 Feb 1;74:240–53.

13. Korgaonkar AA, Li Y, Sekhar D, Subramanian D, Guevarra J, Swietek B, et al. Toll-like
Receptor 4 Signaling in Neurons Enhances Calcium-Permeable α-Amino-3-Hydroxy-5-
Methyl-4-Isoxazolepropionic Acid Receptor Currents and Drives Post-Traumatic
Epileptogenesis. Annals of Neurology. 2020 Apr 1;87(4):497–515.

14. Korgaonkar AA, Nguyen S, Li Y, Sekhar D, Subramanian D, Guevarra J, et al. Distinct
cellular mediators drive the Janus faces of toll-like receptor 4 regulation of network
excitability which impacts working memory performance after brain injury. Brain,
behavior, and immunity. 2020 Aug;88:381–95.

15. Yang CC, Lin CC, Hsiao LD, Kuo JM, Tseng HC, Yang CM. Lipopolysaccharide-induced
matrix metalloproteinase-9 expression associated with cell migration in rat brain
astrocytes. International Journal of Molecular Sciences. 2020 Jan 1;21(1).

16. Qiu J, Xu J, Zheng Y, Wei Y, Zhu X, Lo EH, et al. High-mobility group box 1 promotes
metalloproteinase-9 upregulation through toll-like receptor 4 after cerebral ischemia.
Stroke. 2010 Sept;41(9):2077–82.

17. Huntley GW. Synaptic circuit remodelling by matrix metalloproteinases in health and
disease. Nature Reviews Neuroscience. 2012 Nov;13(11):743–57.

18. Ethell IM, Ethell DW. Matrix metalloproteinases in brain development and remodeling:
synaptic functions and targets. Journal of neuroscience research. 2007
Oct;85(13):2813–23.

19. Szklarczyk A, Lapinska J, Rylski M, Mckay RDG, Kaczmarek L. Matrix Metalloproteinase-
9 Undergoes Expression and Activation during Dendritic Remodeling in Adult
Hippocampus. 2002.

20. Dityatev A, Schachner M. Extracellular matrix molecules and synaptic plasticity. Nature
reviews Neuroscience. 2003 June;4(6):456–68.

21. Minta K, Brinkmalm G, Al Nimer F, Thelin EP, Piehl F, Tullberg M, et al. Dynamics of
cerebrospinal fluid levels of matrix metalloproteinases in human traumatic brain
injury. Sci Rep. 2020 Oct 22;10(1):18075.

22. Wang X, Jung J, Asahi M, Chwang W, Russo L, Moskowitz MA, et al. Effects of Matrix
Metalloproteinase-9 Gene Knock-Out on Morphological and Motor Outcomes after
Traumatic Brain Injury. 2000.

23. Hayashi T, Kaneko Y, Yu S, Bae E, Stahl CE, Kawase T, et al. Quantitative analyses of
matrix metalloproteinase activity after traumatic brain injury in adult rats. Brain
Research. 2009 July;1280:172–7.

24. Grossetete M, Phelps J, Arko L, Yonas H, Rosenberg GA. ELEVATION OF MATRIX
METALLOPROTEINASES 3 AND 9 IN CEREBROSPINAL FLUID AND BLOOD IN PATIENTS
WITH SEVERE TRAUMATIC BRAIN INJURY. Neurosurgery. 2009 Oct;65(4):702–8.

25. Reinhard SM, Razak K, Ethell IM. A delicate balance: Role of MMP-9 in brain
development and pathophysiology of neurodevelopmental disorders. Frontiers in
Cellular Neuroscience. 2015 July 29;9(JULY).

26. Vafadari B, Salamian A, Kaczmarek L. MMP-9 in translation: from molecule to brain
physiology, pathology, and therapy. Journal of Neurochemistry. 2016 Oct 1;139:91–
114.

27. Pijet B, Stefaniuk M, Kostrzewska-Ksiezyk A, Tsilibary PE, Tzinia A, Kaczmarek L.
Elevation of MMP-9 Levels Promotes Epileptogenesis After Traumatic Brain Injury.
Molecular Neurobiology. 2018 Dec 1;55(12):9294–306.

28. Wilczynski GM, Konopacki FA, Wilczek E, Lasiecka Z, Gorlewicz A, Michaluk P, et al.
Important role of matrix metalloproteinase 9 in epileptogenesis. Journal of Cell
Biology. 2008 Mar 10;180(5):1021–35.

29. Dityatev A, Fellin T. Extracellular matrix in plasticity and epileptogenesis. Neuron Glia
Biol. 2008 Aug;4(3):235–47.

30. Jia F, Pan Y hua, Mao Q, Liang Y min, Jiang J yao. Matrix metalloproteinase-9 expression
and protein levels after fluid percussion injury in rats: The effect of injury severity and
brain temperature. Journal of neurotrauma. 2010 June;27(6):1059–68.

31. Corrubia L, Huang A, Nguyen S, Shiflett MW, Jones MV, Ewell LA, et al. Early deficits in
dentate circuit and behavioral pattern separation after concussive brain injury.
Experimental Neurology. 2023 Dec;370:114578.

32. Witgen BM, Lifshitz J, Smith ML, Schwarzbach E, Liang SL, Grady MS, et al. Regional
hippocampal alteration associated with cognitive deficit following experimental brain
injury: A systems, network and cellular evaluation. Neuroscience. 2005;133(1):1–15.

33. Gupta A, Elgammal FS, Proddutur A, Shah S, Santhakumar V. Decrease in Tonic
Inhibition Contributes to Increase in Dentate Semilunar Granule Cell Excitability after
Brain Injury. J Neurosci. 2012 Feb 15;32(7):2523–37.

34. Hirschfeld M, Ma Y, Weis JH, Vogel SN, Weis JJ. Cutting edge: repurification of
lipopolysaccharide eliminates signaling through both human and murine toll-like
receptor 2. J Immunol. 2000 July 15;165(2):618–22.

35. Cooke SF, Wu J, Plattner F, Errington M, Rowan M, Peters M, et al. Autophosphorylation
of αCaMKII is not a general requirement for NMDA receptor-dependent LTP in the adult
mouse. Journal of Physiology. 2006 Aug;574(3):805–18.

36. Schott AL, Esfahany KN, Grocott JM, Farrell JS. Toothy: an interactive platform for
dentate spike curation. bioRxiv. 2025 Oct 3;2025.10.03.680300.

37. Gawlak M, Górkiewicz T, Gorlewicz A, Konopacki FA, Kaczmarek L, Wilczynski GM. High
resolution in situ zymography reveals matrix metalloproteinase activity at
glutamatergic synapses. Neuroscience. 2009 Jan 12;158(1):167–76.

38. Proddutur A, Nguyen S, Yeh CW, Gupta A, Santhakumar V. Reclusive chandeliers:
Functional isolation of dentate axo-axonic cells after experimental status epilepticus.
Progress in Neurobiology. 2023 Dec;231:102542.

39. Rodríguez Peris L, Scheuber MI, Shan H, Braun M, Schwab ME. Barnes maze test for
spatial memory: A new, sensitive scoring system for mouse search strategies.
Behavioural Brain Research. 2024 Feb;458:114730.

40. Hunt RF, Scheff SW, Smith BN. Synaptic Reorganization of Inhibitory Hilar Interneuron
Circuitry after Traumatic Brain Injury in Mice. J Neurosci. 2011 May 4;31(18):6880–90.

41. Gupta A, Dovek L, Proddutur A, Elgammal FS, Santhakumar V. Long-Term Effects of
Moderate Concussive Brain Injury During Adolescence on Synaptic and Tonic GABA
Currents in Dentate Granule Cells and Semilunar Granule Cells. Front Neurosci.
2022;16:800733.

42. Santhakumar V, Ratzliff ADH, Jeng J, Toth Z, Soltesz I. Long-term hyperexcitability in the
hippocampus after experimental head trauma. Annals of Neurology. 2001
Dec;50(6):708–17.

43. Santhakumar V, Bender R, Frotscher M, Ross ST, Hollrigel GS, Toth Z, et al. Granule cell
hyperexcitability in the early post-traumatic rat dentate gyrus: the 'irritable mossy cell'
hypothesis. The Journal of Physiology. 2000 Apr;524(1):117–34.

44. Hadass O, Tomlinson BN, Gooyit M, Chen S, Purdy JJ, Walker JM, et al. Selective
Inhibition of Matrix Metalloproteinase-9 Attenuates Secondary Damage Resulting from
Severe Traumatic Brain Injury. PLoS ONE. 2013 Oct 23;8(10).

45. Michaluk P, Wawrzyniak M, Alot P, Szczot M, Wyrembek P, Mercik K, et al. Influence of
matrix metalloproteinase MMP-9 on dendritic spine morphology. J Cell Sci. 2011 Oct
1;124(Pt 19):3369–80.

46. Wiera G, Szczot M, Wojtowicz T, Lebida K, Koza P, Mozrzymas JW. Impact of matrix
metalloproteinase-9 overexpression on synaptic excitatory transmission and its

plasticity in rat CA3-CA1 hippocampal pathway. J Physiol Pharmacol. 2015
Apr;66(2):309–15.

47. Nagy V, Bozdagi O, Matynia A, Balcerzyk M, Okulski P, Dzwonek J, et al. Matrix
metalloproteinase-9 is required for hippocampal late-phase long-term potentiation
and memory. Journal of Neuroscience. 2006 Feb 15;26(7):1923–34.

48. Bozdagi O, Nagy V, Kwei KT, Huntley GW. In vivo roles for matrix metalloproteinase-9 in
mature hippocampal synaptic physiology and plasticity. Journal of Neurophysiology.
2007 July;98(1):334–44.

49. Wiera G, Mozrzymas JW. Extracellular Metalloproteinases in the Plasticity of Excitatory
and Inhibitory Synapses. Cells [Internet]. 2021 Aug 11;10(8). Available from:
http://www.ncbi.nlm.nih.gov/pubmed/34440823

50. Bragin A, Jand& G, Nadasdy Z, Van Landeghem M, Buzsaki G. Dentate EEG Spikes and
Associated Interneuronal Population Bursts in the Hippocampal Hilar Region of the
Rat. Vol. 73, JOURNALOF NEUROPHYSIOLOGY. 1995.

51. Penttonen M, Kamondi A, Sik A, Acsády L, Buzsáki G. Feed-forward and feed-back
activation of the dentate gyrus in vivo during dentate spikes and sharp wave bursts.
Hippocampus. 1997;7(4):437–50.

52. Lensu S, Waselius T, Penttonen M, Nokia MS. Dentate spikes and learning: disrupting
hippocampal function during memory consolidation can improve pattern separation. J
Neurophysiol. 2019;121:131–9.

53. Farrell JS, Soltesz I. Noncanonical circuits, states, and computations of the
hippocampus. Science. 2025 Sept 11;389(6765):eadv4420.

54. Nokia MS, Gureviciene I, Waselius T, Tanila H, Penttonen M. Hippocampal electrical
stimulation disrupts associative learning when targeted at dentate spikes. Journal of
Physiology. 2017 July 15;595(14):4961–71.

55. Farrell JS, Hwaun E, Dudok B, Soltesz I. Neural and behavioural state switching during
hippocampal dentate spikes. Nature. 2024 Apr 18;628(8008):590–5.

56. Bott JB, Muller MA, Jackson J, Aubert J, Cassel JC, Mathis C, et al. Spatial reference
memory is associated with modulation of theta-gamma coupling in the dentate Gyrus.
Cerebral Cortex. 2016 Sept 1;26(9):3744–53.

57. Dovek L, Ahmadi M, Marrero K, Zagha E, Santhakumar V. Cellular and circuit features
distinguish dentate gyrus semilunar granule cells and granule cells activated during
contextual memory formation. 2025 July 17 [cited 2026 Jan 1]; Available from:
https://elifesciences.org/reviewed-preprints/101428v2

58. Maroso M, Balosso S, Ravizza T, Liu J, Aronica E, Iyer AM, et al. Toll-like receptor 4 and
high-mobility group box-1 are involved in ictogenesis and can be targeted to reduce
seizures. Nature Medicine. 2010 Apr;16(4):413–9.

59. Kleen JK, Holmes GL. Taming TLR4 may ease seizures. Nature Medicine. 2010
Apr;16(4):369–70.

60. Vezzani A, Balosso S, Ravizza T. Neuroinflammatory pathways as treatment targets and
biomarkers in epilepsy. Nat Rev Neurol. 2019 Aug;15(8):459–72.

61. Okun E, Griffioen KJ, Mattson MP. Toll-like receptor signaling in neural plasticity and
disease. Trends in Neurosciences. 2011 May;34(5):269–81.

62. Ahmad A, Crupi R, Campolo M, Genovese T, Esposito E, Cuzzocrea S. Absence of TLR4
Reduces Neurovascular Unit and Secondary Inflammatory Process after Traumatic
Brain Injury in Mice. Dhandapani KM, editor. PLoS ONE. 2013 Mar 28;8(3):e57208.

63. Xiong G, Metheny H, Johnson BN, Cohen AS. A comparison of different slicing planes in
preservation of major hippocampal pathway fibers in the mouse. Frontiers in
Neuroanatomy. 2017 Nov 21;11.

64. Howard AL, Neu A, Morgan RJ, Echegoyen JC, Soltesz I. Opposing Modifications in
Intrinsic Currents and Synaptic Inputs in Post-Traumatic Mossy Cells: Evidence for
Single-Cell Homeostasis in a Hyperexcitable Network. Journal of Neurophysiology.
2007 Mar;97(3):2394–409.

65. Dziembowska M, Milek J, Janusz A, Rejmak E, Romanowska E, Gorkiewicz T, et al.
Activity-Dependent Local Translation of Matrix Metalloproteinase-9. J Neurosci. 2012
Oct 17;32(42):14538–47.

66. Dziembowska M, Wlodarczyk J. MMP9: A novel function in synaptic plasticity. The
International Journal of Biochemistry & Cell Biology. 2012 May;44(5):709–13.

67. Stefaniuk M, Beroun A, Lebitko T, Markina O, Leski S, Meyza K, et al. Matrix
Metalloproteinase-9 and Synaptic Plasticity in the Central Amygdala in Control of
Alcohol-Seeking Behavior. Biological Psychiatry. 2017 June;81(11):907–17.

68. Szepesi Z, Hosy E, Ruszczycki B, Bijata M, Pyskaty M, Bikbaev A, et al. Synaptically
released matrix metalloproteinase activity in control of structural plasticity and the
cell surface distribution of GluA1-AMPA receptors. PLoS ONE. 2014 May 22;9(5).

69. Fanselow EE, Kubota Y, Wehr M, Yavorska I. Somatostatin-Expressing Inhibitory
Interneurons in Cortical Circuits. Frontiers in Neural Circuits | www.frontiersin.org.
2016;10:76.

70. Pirbhoy PS, Rais M, Lovelace JW, Woodard W, Razak KA, Binder DK, et al. Acute
894   pharmacological inhibition of matrix metalloproteinase-9 activity during development
895   restores perineuronal net formation and normalizes auditory processing in Fmr1 KO
896   mice. Journal of Neurochemistry. 2020 Dec;155(5):538–58.

897   71. Sun DA, Deshpande LS, Sombati S, Baranova A, Wilson MS, Hamm RJ, et al. Traumatic
898   brain injury causes a long-lasting calcium (Ca2+)-plateau of elevated intracellular Ca
899   levels and altered Ca2+homeostatic mechanisms in hippocampal neurons surviving
900   brain injury. Eur J of Neuroscience. 2008 Apr;27(7):1659–72.

901   72. Weber JT. Altered Calcium Signaling Following Traumatic Brain Injury. Front Pharmacol
902   [Internet]. 2012 [cited 2025 Dec 22];3. Available from:
903   http://journal.frontiersin.org/article/10.3389/fphar.2012.00060/abstract

904   73. Hellstrom IC, Danik M, Luheshi GN, Williams S. Chronic LPS exposure produces
905   changes in intrinsic membrane properties and a sustained IL-β-dependent increase in
906   GABAergic inhibition in hippocampal CA1 pyramidal neurons. Hippocampus.
907   2005;15(5):656–64.

908   74. Hainmueller T, Bartos M. Dentate gyrus circuits for encoding, retrieval and
909   discrimination of episodic memories. Nature Reviews Neuroscience. 2020 Mar
910   1;21(3):153–68.

911   75. Hsu D. The dentate gyrus as a filter or gate: a look back and a look ahead. Prog Brain
912   Res. 2007;163:601–13.

913   76. Ewell LA. To sleep perchance to spike: a functional role for dentate spikes in memory.
914   Journal of Physiology. 2017 July 15;595(14):4565.

915   77. Farrell JS, Hwaun E, Dudok B, Soltesz I. Neural and behavioural state switching during
916   hippocampal dentate spikes. [cited 2024 Mar 20]; Available from:
917   https://doi.org/10.1038/s41586-024-07192-8

918   78. Flynn SP, Barrier S, Scott RC, Lenck-Santini PP, Holmes GL. Status epilepticus induced
919   spontaneous dentate gyrus spikes: In vivo current source density analysis. PLoS ONE.
920   2015 July 6;10(7).

921   79. Engel J, Bragin A, Staba R. Nonictal EEG biomarkers for diagnosis and treatment.
922   Epilepsia Open. 2018 Dec 1;3(S2):120–6.

923   80. Perucca P, Smith G, Santana-Gomez C, Bragin A, Staba R. Electrophysiological
924   biomarkers of epileptogenicity after traumatic brain injury. Neurobiology of Disease.
925   2019 Mar;123:69–74.

926   81. Tewari BP, Chaunsali L, Campbell SL, Patel DC, Goode AE, Sontheimer H. Perineuronal
927   nets decrease membrane capacitance of peritumoral fast spiking interneurons in a
928   model of epilepsy. Nature Communications. 2018 Dec 1;9(1).

929   82. Whitebirch AC, LaFrancois JJ, Jain S, Leary P, Santoro B, Siegelbaum SA, et al. Enhanced
930   excitability of the hippocampal CA2 region and its contribution to seizure activity in a
931   mouse model of temporal lobe epilepsy. Neuron. 2022 Oct;110(19):3121-3138.e8.

Figure legends:

Figure 1: TLR4 signaling drives early excitatory and inhibitory synaptic changes in DGCs after brain injury. A) Schematic of the experimental question. B) Experimental timeline for examining synaptic changes after FPI in rats. C) Representative sEPSC traces showing whole-cell voltage clamp recordings from granule cells in vehicle treated sham controls (black), vehicle treated FPI (red), TLR4 inhibited sham (grey) and TLR4 inhibited FPI (blue). D-E) Cumulative probability plots comparing sEPSC interevent intervals (D) and amplitude (E) between vehicle treated and TLR4 inhibited sham and injured rats. F) Representative sIPSC traces showing whole-cell voltage clamp recordings from granule cells in vehicle treated sham controls (black), vehicle treated FPI (red), TLR4 inhibited sham (grey) and TLR4 inhibited FPI (blue). G-H) Cumulative probability plots comparing sIPSC interevent intervals and amplitude between vehicle treated and TLR4 inhibited sham and injured rats. I) Experimental timeline for examining synaptic changes after FPI in TLR4$^{fl/fl}$: CaMKIIa- CreERT2 mice. J) Representative sEPSC traces from granule cells in Cre^{-ve} sham controls (black), Cre^{-ve} FPI (red), Cre^{+ve} sham (grey) and Cre^{+ve} FPI (blue). K-L) Cumulative probability plots comparing sEPSC interevent intervals and amplitude between granule cells in Cre^{-ve} and Cre^{+ve} mice. * indicates *P*<0.0001 by K-S test.

Figure 2: TLR4 signaling upregulates MMP-9 activity in the dentate gyrus after brain injury. A) Experimental timeline for in-situ zymography to quantify MMP-9 activity after brain injury in rat. B) Representative images showing MMP-9 activity in hippocampal sections from sham and injured rats. Images from vehicle treated (middle) and TLR inhibited mice (bottom) show MMP-9 activity in green. C) Quantification of MMP-9 activity,

normalized to CA1 stratum Radiatum, shows an increase in MMP-9 activity in vehicle

treated injured rats, but not in TAK-242 treated injured rats. Note that MMP-9 activity in

vehicle treated and TAK-242 treated sham rats were not different. D) Experimental timeline

for in-situ zymography to quantify MMP-9 activity after brain injury in mice. E)

Representative images showing MMP-9 activity in sham and injured mice. Images from

hippocampal sections from Cre[-ve] sham and injured mice (top) and Cre[+ve] sham and injured

mice (bottom) show MMP-9 activity in green.F) Quantification of MMP-9 activity, normalized

to CA1 stratum Radiatum, shows an increase in MMP-9 activity after brain injury in Cre[-ve]

mice, but not in Cre[+ve] mice. Note that MMP-9 activity in sham mice was not different. *

indicates $P<0.05$ by TW-ANOVA with Šídáks multiple comparisons post hoc tests. G)

Experimental timeline for RNAscope experiments in naïve rats. H) Representative images

showing increase in MMP-9 mRNA in sections from rats after LPS-B5 treatment. I)

Summary plot of mRNA levels in saline, LPS-B5 treated and LPS-B5+JSH-23 sections

shows an increase in mRNA levels after LPS-B5 treatment which is abolished by blocking

NF-κB. * indicates $P<0.05$ by OW-ANOVA with Šídáks multiple comparisons post hoc tests.

Figure 3: Blocking MMP-9 prevents early synaptic changes in excitatory inputs to DGCs

after brain injury. A) Experimental timeline for examining synaptic changes in acute rat

slices after LPS incubation. B) Representative sEPSC traces from granule celsl in LPS

incubated (orange), LPS+SB-3CT co-incubated (magenta) slices. C-D) Cumulative

probability plots comparing sEPSC interevent intervals and amplitude between LPS-B5

incubated and LPS-B5+SB-3CT co-incubated slices. Note that LPS incubation significantly

reduced interevent intervals. Co-incubation of slices in MMP-9 blocker SB-3CT prevented

this change and were not different from sham animals. MMP-9 inhibition did not rescue

changes in sEPSC amplitude. E) Experimental timeline for examining synaptic changes in

SB-3CT treated sham/injured rats. F) Representative granule cell sEPSC traces showing

from SB-3CT treated sham (grey) and injured rats (green). G-H) Cumulative probability plots

comparing sEPSC interevent intervals and amplitude between SB-3CT treated

sham/injured rats. I) Experimental timeline for examining synaptic changes in SB-3CT

treated mice after brain injury. J) Representative granule cell sEPSC traces from vehicle

treated (red) and SB-3CT treated mice(green). K-L) Cumulative probability plots comparing

sEPSC interevent intervals and amplitude between vehicle treated and SB-3CT treated

mice. * indicates *P*<0.0001 by K-S test.

Figure 4: TLR4 regulation of MMP-9 activity enhances dentate network excitability after

brain injury. A) Experimental timeline for *in vivo* experiments examining dentate excitability

and throughput. B) Representative traces showing DG population spike from sham and

injured animals in response to 800 µA stimulus (bold lines) and 1200 µA (dotted lines). C)

Input-output curve illustrating early saturation of population spike amplitude in injured

animals (red) compared to sham controls. D) Inhibition of TLR4 or MMP-9 effectively

mitigates these changes. E) MMP-9 inhibition (light grey) did not affect input-output curves

in sham animals, TLR4 inhibition increased network excitability of sham animals (grey). F)

Representative trace from 16 channel silicon probe recording local field potentials from

cortex to hippocampus. Current-source density plot shows the sink and source of an

individual dentate spike. inset- a single dentate spike recorded in the GC layer. G)

Quantification of dentate spikes shows a significant increase in frequency in injured

animals. Note that inhibition of TLR4 or MMP-9 effectively reduced changes in dentate

spike frequency in injured animals. * indicates<0.05 by TW-ANOVA with Šídáks multiple

comparisons post hoc tests.

Figure 5: Posttraumatic deficits in long-term potentiation is improved by early

inhibition of TLR4 or MMP-9. A) Experimental timeline for _in vivo_ assessment of long-term

potentiation in perforant path synapses to DGCS. B-C) Injured rats exhibit a significant

decrease in fEPSP slope values after 60 minutes(red) compared to sham controls (black).

D-E) Inhibition of TLR4 (blue) or MMP-9 (green) significantly improved LTP in injured

animals. F-G) MMP-9 inhibition did not affect LTP in SHAM controls. TLR4 inhibition

reduced LTP at 60 min and 180 min after induction. * indicates _P_ <0.05 by TW-RM ANOVA.

Figure 6: Early inhibition of TLR4 or MMP-9 improves spatial learning after brain injury.

A) Experimental timeline for testing spatial learning using Barnes maze. B) Representative

heat plot showing rats trajectory during the task on day 2. C-E) Injured animals showed

deficits in latency to escape, distance travelled, and search strategy on day 2. * indicates

P<0.05 by RM-ANOVA. F-H) Summary plots of effect of TLR4 or MMP-9 inhibition on latency

to escape (F), distance travelled (G), and search strategy (H) on day 2. I) Injured animals

showed increased thigmotaxis on day 2 which was reduced with TLR4 or MMP-9 inhibition.

* indicates P<0.05 by TW-ANOVA with Šídáks multiple comparisons post hoc tests.

1018

Figure 7: Graphical Summary of TLR4 regulation of dentate gyrus function in control

and injured brain.

In uninjured brain, pharmacological inhibition of TLR4 signaling increases the frequency of

both excitatory and inhibitory currents and enhances network excitability and impairs long-

term potentiation without altering MMP-9 activity. Genetic deletion of neuronal TLR4 alters

intrinsic properties without impacting synaptic physiology. Together these data are

consistent with glial TLR4 signaling contributing to basal synaptic homeostasis in line with

prior ex vivo studies [14].In the injured brain, augmented neuronal TLR4 signaling acts

through MMP-9 to enhance glutamatergic signaling, network excitability and dentate spikes

and impairs LTP maintenance. Mechanistically, inhibitory changes are driven by TLR4 in an

MMP-9 independent manner. Together, these findings demonstrate that TLR4 signaling

regulates DG function and seizure susceptibility [13] through distinct, context-dependent

cellular mechanisms in the healthy and injured brain.

1032

1033

FIGURE.1

FIGURE.2

FIGURE.3

FIGURE.4

FIGURE.5

FIGURE.6

FIGURE.7