Neuronal Toll-like Receptor-4 regulation of Matrix Metalloproteinase-9 Activity

Mediates Dentate Circuit Dysfunction after Traumatic Brain Injury

Deepak Subramanian[1#], Erick Contreras[1], Laura Dovek[1], Razieh Jaberi[1], Emmanuel Green[1],

Ysabelle K Lao[1], Iryna M Ethell[2] and Vijayalakshmi Santhakumar[1#]

[1]Department of Molecular, Cell and Systems Biology, University of California Riverside,

Riverside, California 92521, United States.

[2]Biomedical Sciences, University of California Riverside, Riverside, California 92521,

United States.

Keywords: Traumatic Brain Injury, Dentate gyrus, Toll-like Receptor 4, Matrix

Metalloproteinase-9, Neuro-immune signaling, Synaptic plasticity

Abstract: 287 words   Full Text Character count: 75,590   Number of pages: 51

Number of figures: 7   Supplementary figures: 6   Color figures: 7

Number of tables: 0   Number of supplementary tables: 5

Correspondence:
Deepak Subramanian, PhD
Department of Molecular, Cell and Systems Biology
University of California, Riverside
E-mail: dsubra@ucr.edu

Co-Correspondence:
Vijayalakshmi Santhakumar, PhD
Department of Molecular, Cell and Systems Biology
University of California, Riverside
E-mail: vijayas@ucr.edu

List of Abbreviations:

aCSF: Artificial Cerebrospinal Fluid

AMPAR: α-Amino-3-hydroxy-5-methyl-4-isoxazolepropionic Acid Receptor

ANOVA: Analysis of Variance

CA1 SR: CA1 Stratum Radiatum

CaMKIIa: Calcium/Calmodulin-Dependent Protein Kinase II Alpha

CreERT2: Cre Recombinase Estrogen Receptor T2

DG: Dentate Gyrus

DQ Gelatinase: Dye Quenched fluorogenic substrate for zymography

ECM: Extracellular Matrix

EPSC: Excitatory Postsynaptic Current

FPI: Fluid Percussion Injury

HMGB1: High-Mobility Group Box 1

IEI: Inter-Event Interval

IPSC: Inhibitory Postsynaptic Current

JSH-23: NF-κB Inhibitor

K-S test: Kolmogorov-Smirnov Test

LPS: Lipopolysaccharide

LTP: Long Term Potentiation

MMP-9: Matrix Metalloproteinase-9

NF-κB: Nuclear Factor Kappa B

TAK-242: TLR4 Antagonist

TBS: Theta Burst Stimulation

TBI: Traumatic Brain Injury

TTX: Tetrodotoxin

Abstract:

Neuroinflammatory pathways activated by traumatic brain injury (TBI) are critical mediators of long-term neurological dysfunction and represent promising therapeutic targets. Toll-like receptor 4 (TLR4), an innate immune receptor, was previously shown to contribute to increased seizure susceptibility and cognitive deficits in rats after lateral fluid percussion injury (FPI). However, the cellular and molecular mechanisms underlying TLR4-mediated circuit dysfunction early after brain injury are not fully understood. In this study, we define a cell- and circuit- specific neuroimmune-enzyme effector signaling axis that mediates early post-TBI circuit dysfunction in the hippocampal Dentate Gyrus (DG). Using *ex vivo* electrophysiology in rat and mouse models one-week after brain injury, we demonstrate that neuronal TLR4 signaling regulates both excitatory and inhibitory synaptic inputs to dentate granule cells (DGC). Collectively, pharmacological inhibition of TLR4 in rats and cell-type-specific deletion of TLR4 in mice show that neuronal TLR4 mediates injury-driven increase in DGC excitatory input frequency and relies on downstream activation of Matrix Metalloproteinase-9 (MMP-9). In contrast, TLR4 signaling contributed to decrease in inhibitory current frequency after injury, but independent of MMP-9, revealing a mechanistic divergence. Systemic inhibition of either TLR4 signaling or MMP-9 activity in rats within 24 hours after injury reduced network hyperexcitability and improved long-term potentiation (LTP) in the DG measured *in vivo* one-week after injury. Either TLR4 or MMP-9 inhibition early after injury effectively attenuated spatial memory deficits in a Barnes maze task one-month post-injury. Paradoxically, in sham controls, inhibition of TLR4 increased frequency of both excitatory and inhibitory inputs to DGCs and augmented network

77   excitability, without altering MMP-9 levels, identifying context-dependent roles for TLR4

78   signaling. Together, these results identify a novel TLR4 -MMP-9 axis as a key driver of early

79   post-TBI dentate gyrus circuit dysfunction and behavioral deficits.

INTRODUCTION:

Traumatic brain injury (TBI) initiates a robust and complex inflammatory cascade that exacerbates secondary structural and functional damage [1–4]. While cellular damage is widespread and diffuse, the hippocampus is highly susceptible to post-traumatic pathology. Specifically, the hippocampal dentate gyrus (DG) undergoes extensive loss of hilar neurons and circuit reorganization that critically alter hippocampal network physiology and contributes to the development of cognitive impairments and post-traumatic epilepsy [5–11]. In this context, we previously identified a pivotal role for the neuronally expressed innate immune receptor, Toll-like receptor 4 (TLR4), in enhancing neuronal excitability and driving epileptogenesis after injury [12–14]. Yet, the downstream mechanisms by which specific immune receptors influence circuit physiology remain poorly understood.

 Recent evidence indicates that canonical TLR4 activators, such as lipopolysaccharide (LPS) and high-mobility group box 1 (HMGB1), can upregulate Matrix Metalloproteinase-9 (MMP-9) activity in vitro [15,16]. MMP-9 , a zinc-dependent endopeptidase, is a critical regulator of synaptic communication and plasticity through the activity-dependent degradation of the extracellular matrix (ECM)[17–20]. Importantly, MMP-9 levels are elevated early in both animal models and clinical populations after brain injury [21–24] , driving pathological remodeling of synapses and circuits that promote epileptogenesis [25–29]. Despite a strong temporal correlation between MMP-9 elevation and transient increases in TLR4 signaling after injury [12,30], a causal association between these two pathways has not been established. Based on the temporal correlation and shared

upstream activators, we hypothesized that mechanistic interactions between these two

pathways contribute to early circuit dysfunction after brain injury.

Here, we provide evidence for a novel mechanism involving TLR4-mediated upregulation of

MMP-9 activity following brain injury that drives maladaptive changes in synaptic

physiology, network excitability, and plasticity. Our findings reveal that the injury-induced

increase of MMP-9 in the dentate gyrus is specifically mediated by neuronal TLR4 signaling.

While TLR4-regulated MMP-9 activity was critical in enhancing excitatory synaptic inputs,

inhibitory input dysregulation was mediated by TLR4 signaling independent of MMP-9.

Furthermore, we demonstrate that this TLR4 and MMP-9 dependent pathway enhances

network excitability and impairs synaptic plasticity early after injury. Importantly, we show

that pharmacological inhibition of either TLR4 or MMP-9 effectively mitigates these early

pathological changes, highlighting the therapeutic potential of targeting this neuro-immune

signaling pathway to preserve circuit function after TBI.

METHODS:

All experiments were performed in accordance with IACUC protocols approved by the University of California-Riverside, CA and in keeping with the ARRIVE guidelines. Male and female Wistar rats (P23-P24) and mice (TLR4$^{f/f}$:CaMKIIa-CreERT2) were housed with littermates in a 12-hour light/dark cycle with access to water and food ad libidum. All mice were bred in-house. Animals were randomly assigned to either sham or FPI groups and treatment conditions.

Fluid Percussion Injury (FPI) and treatments

Juvenile male and female Wistar rats (24 days old) were subjected to moderate (1.8-2.0 atm) lateral FPI or sham injury using standard methods [12–14]. Briefly, a 2.7 mm hole was trephined on the left side of the skull 3 mm caudal to bregma and 3.5 mm lateral from the sagittal suture. A Luer-Lok syringe hub with a 3 mm inner diameter was placed over the exposed dura and bonded to the skull with cyanoacrylate adhesive. The animals were returned to their home cage following recovery. One day later, animals were briefly anesthetized with isoflurane and attached to a fluid percussion injury device (Virginia Commonwealth University, Richmond, VA, Model 01-B) directly at the metal nozzle. A pendulum was dropped to deliver a brief (20ms) 1.9–2.0 atm impact on the dura. For sham injury, the animals underwent surgical implantation of the hub and were anesthetized and attached to the FPI device, but the pendulum was not dropped. Only rats with >10 seconds apnea after injury were included in the injury group, and only the hemisphere ipsilateral to the injury was examined.

To achieve neuron-specific deletion of TLR4, TLR4-floxed (TLR4$^{f/f}$)mice were crossed with

mice expressing inducible Cre recombinase under the control of the CamKIIa promoter

(CaMKIIa-CreERT2). Cre-positive (Cre^{+ve}) mice with TLR4 deletion and Cre-negative (Cre^{-ve})

littermate were used in experiments. TLR4$^{f/f}$:CaMKIIa-CreERT2 mice (8–10-week-old male

and female mice) were induced with tamoxifen and subjected to lateral FPI (1.5 atm) or

sham injury [31,32]. Tamoxifen (75mg/kg, i.p.) was prepared in 10% ethanol and 90% corn

oil by sonicating at 37°C for 90 minutes. Mice received 3 doses of tamoxifen, 24 hours before

injury, 30 minutes after injury and 24 hours after injury. TLR4 deletion in glutamatergic

neurons   in   dentate   gyrus   was   confirmed   through   immunohistochemistry   (see

immunohistochemistry methods below) and RNAscopy. Animals were treated with vehicle

(DMSO), TLR4 antagonist (TAK-242, 0.5mg/kg, i.p., Invivogen, San Diego, CA) or MMP-9

antagonist (SB-3CT, 50mg/kg, i.p., SelleckChem, Houston, TX) at 30 minutes, 8 hours, and

24 hours after injury. The dosing strategy for TAK-242 was based on the approach previously

adopted to identify functional outcomes [13,14,30,33]. Although SB3-CT has a shorter half

life than TAK-242 [34], prior studies have adopted 30 minutes - 24 hour dosing strategies

[30,33,35]. Therefore, a dosing interval similar to TAK-242 was adopted to ensure a

consistency in injections and animal handling across groups.

Ex vivo electrophysiology

One-week after FPI or sham injury, animals were anesthetized with isoflurane and

decapitated. Brain was quickly harvested, hemisected and coronal hippocampal slices

(300 μm) were prepared in ice-cold sucrose artificial cerebrospinal fluid (aCSF) containing

(in mM): 85 NaCl, 75 sucrose, 24 NaHCO3, 25 glucose, 4 MgCl2, 2.5 KCl, 1.25 NaH2PO4,

and 0.5 CaCl2 [36]. Slices were incubated at 32°C for 20 min in an interface holding chamber

containing an equal volume of sucrose-aCSF and recording aCSF containing (in mM) 126

NaCl, 2.5 KCl, 2 CaCl$_2$, 2 MgCl$_2$, 1.25 NaH$_2$PO$_4$, 26 NaHCO$_3$ and 10 D-glucose. All solutions

were saturated with 95% O$_2$ and 5% CO$_2$ and maintained at a pH of 7.4. Slices were allowed

to equilibrate for an hour at room temperature before recordings. Whole-cell voltage- and

current-clamp recordings were obtained from dentate granule cells (DGCs) in the

suprapyramidal blade at 33-34°C under IR-DIC (Olympus BX51 with 40x water immersion

objectives) using 3-6MΩ glass pipettes fabricated using dual stage glass micropipette puller

(Narishige PC-10, Amityville NY). Data was recorded using MultiClamp 700B amplifiers,

digitized using Digidata 1440A and acquired using pClamp10 (Molecular Devices, San Jose,

CA) at 10-kHz sampling frequency. Spontaneous and miniature excitatory/inhibitory

currents from the same DGCs were obtained using a cesium-based internal solution (in mM:

Cesium methanesulfonate 140, NaCl 5, HEPES 10, EGTA 0.2, Mg2ATP 2, Na2GTP 0.2, Qx314

5, (pH 7.25, 270-285 mOsm). Biocytin (0.2%) was included in the internal solution for post

hoc cell identification. Spontaneous and miniature excitatory postsynaptic potentials

(sEPSC and mEPSC) were recorded from the same neurons at a holding potential of -70 mV,

and spontaneous and miniature inhibitory postsynaptic potentials (sIPSC and mIPSC

respectively) were recorded at a holding potential of 0 mV. Action potential independent

miniature currents were isolated using tetrodotoxin (TTX, 1μM). In a subset of experiments

TLR4 signaling was selectively activated by incubating slices in LPS-B5 ultrapure

(Escherichia Coli serotype O55:B5, 10 μg/ml, Invivogen, San Diego, CA), a selective TLR4

agonist, for 60-90 minutes before recording [37]. Data from cell in which access resistance

changes over 20% during recordings and recordings with access resistance above 25 MΩ

were excluded from analysis. Spontaneous/miniature synaptic currents (100 events/cell)

were detected using template search feature in Easy Electrophysiology (version 2.6.,

London, UK.) by a blinded investigator and interevent intervals and amplitudes were

analyzed as cumulative distribution in Graphpad Prism 10.0.1 (Boston, MA).

In vivo electrophysiology

One-week after FPI/Sham injury rats were anesthetized under urethane (1.5g/kg i.p.) and

head-fixed onto a stereotaxic frame. Under aseptic conditions, rats were implanted with a

50 µm tungsten wire depth electrode (California Fine Wire company, Grover beach, CA) in

the granule cell layer (AP: 3-4 mm, ML: 2 mm, DV: 3-3.5 mm) guided by electrical stimulation

of the angular bundle (ML:4-4.2 mm, DV: 1.8-2 mm; pulse width 150µs delivered at 0.1Hz).

Two screw electrodes on the contralateral side served as reference and ground. The position

of electrodes was secured using dental cement. Acute recordings from urethane

anesthetized rats started 60 minutes after induction. Current-response curves were

obtained in response to electrical stimulation of the angular bundle (150 µs @ 0.1Hz). For

assessing Long Term Potentiation (LTP), DG LFP responses were evoked every 30 seconds

for a period of 30 minutes to establish a baseline. LTP was induced by Theta Burst Protocol

consisting six trains of six pulses at 400 Hz, 200 ms between trains, delivered six times at 20

seconds interval [38] and responses were recorded for an additional 180 minutes. At the end

of the experiment, animals were euthanized with a lethal dose of Euthasol (150 mg/kg, i.p.).

Signals were amplified 100 times (Pinnacle systems, Lawrence, KS), sampled at 10kHz,

digitized using PowerLab 16/35 and recorded using LabChart8 software (AD Instruments,

Colorado Springs, CO). Analysis was conducted using LabChart modules. Dentate spikes

were recorded from urethane anesthetized rats (1.5g/kg i.p.) using 16 Ch silicon probe

(A1x16-5mm-100-703 CM16, Neuronexus, Ann Arbor, MI). Signals were amplified 100 times

using Model 3600 amplifier (AM-Systems, Sequim, WA), digitized using PowerLab 16/35 and

recorded using Labchart8 Software (AD instruments, Colorado Springs, CO). Dentate spikes

were detected using Toothy analysis pipeline[39] (https://github.com/Farrell-

Laboratory/Toothy). Briefly, large positive events were identified using a 4.5 S.D. threshold.

Detected events were visually confirmed by a blinded investigator and the total number of

dentate spikes recorded in a 30-minute period were counted. We did not classify dentate

spikes as type 1 or type 2.

In Situ Zymography

MMP-9 activity was evaluated in hippocampal slices obtained from rats and mice 48 hours

after injury using a modified in situ zymography protocol [40]. Briefly, sham/injured rats or

TLR4$^{f/f}$:CaMKIIa-CreERT2 mice and Cre^{-ve} controls were transcardially perfused using an

alcohol-based formalin free tissue fixative (Accustain, A5472 Sigma-Aldrich, St.Louis, MO),

embedded in 2% agar and coronal slices (50µm, VF-310-0Z Compresstome, Precisionary

Instruments, Ashland, MA) were prepared at <4°C to preserve enzyme activity. Slices were

then incubated with a fluorogenic Dye Quenched gelatin substrate (1:50 dilution, Invitrogen,

Waltham MA) for 40 min at 37°C, washed with PBS, fixed in 4% paraformaldehyde and

mounted on slides in Vectashield (Vector Labs, Newark, CA). The specificity of reactions

was confirmed by adding MMP-9 inhibitor SB-3CT to control slices (see supplementary

figure 2J-K). Semiquantitative analysis of DQ fluorescence in DG molecular layer were

performed using ImageJ by a blinded investigator. Data was collected from 2-4 slices /

animals and averaged. Intensity measures were normalized to CA1 stratum radiatum since

we did not observe any effect of injury, treatment or genotype in this region (Supplementary

figure 4I-J).

Single molecule fluorescent in situ hybridization (RNAscope)

To visualize TLR4 in mice and MMP-9 mRNA in rat, we used RNAscope Multiplex Fluorescent

in situ hybridization on cryosectioned slices (20μm). RNAscope Multiplex Fluorescent

Reagent kit v2 assay (ACD, Newark, CA) was used following the protocol for fixed tissue [41].

The following kits were used for this study: RNAscope Multiplex Fluorescent Reagent kit

(Cat#323100), RNAscope Probe-MM-TLR4 (Cat#316801) and RNAscope probe-Rn-MMP-9

(Cat#532281). Slices were coverslipped with DABCO with DAPI mounting media and imaged

using a Zeiss Axioscope and analyzed using QuPath software by a blinded investigator.

Immunohistochemistry

Male and female TLR4[f/f]:CaMKIIa-CreERT2 mice (8-10 weeks) were induced with tamoxifen

as described above and subjected to Sham or FPI. Three days after injury, brain tissue was

harvested following transcardial perfusion with PBS followed by 4% PFA. Brains were post-

fixed overnight and then transferred to 30% sucrose until fully equilibrated. Tissue was

cryosectioned at 20 μm thickness and mounted on Superfrost Plus glass slides. For

immunohistochemistry, slides were brought to room temperature to remove OCT and

washed three times in PBS. Sections were permeabilized with 0.2% Triton X-100 in PBS for

45 minutes at room temperature, followed by blocking with 20% goat serum for 45 minutes.

Sections were then incubated overnight at 4 °C with a primary antibody against TLR4 (mouse

anti-TLR4; Santa Cruz Biotechnology; 1:250). After three PBS washes, sections were

incubated with goat anti-mouse Alexa Fluor 546 secondary antibody (1:1000, Thermo Fisher,

S11227) for 1 hour at room temperature. Slides were washed three times with PBS and

mounted using an antifade mounting medium. Images were acquired using a Zeiss

Axioscope-5 with stereo investigator (MBF Bioscience) from three animals per group, with

six sections analyzed per animal. Quantification was performed using ImageJ software by

calculating the Corrected Total Cell Fluorescence (CTCF) in the granule cell layer and hilus

for each section (Supplementary figure 3 A-C).

Stereological Estimation of Hilar Neurons

To quantify neuronal loss following injury, sham and FPI rats (n = 3 and 4 rats, respectively.,

1-month post-injury) were deeply anesthetized with Euthasol (150 mg/kg, i.p.) and

transcardially perfused with PBS followed by 4% PFA. Brains were post-fixed and coronally

sectioned (50 μm) using a compressotome (VF-310-Z, Precisionary Instruments). Sections

from the side of injury were processed for Nissl staining (0.1% cresyl violet acetate and

0.25% glacial acetic acid). Unbiased stereological estimates of hilar neuron populations

were obtained using the Optical Fractionator probe (Stereo Investigator; MBF Bioscience) on

a Zeiss Axioscope-5 microscope by blinded investigators (See Supplementary Figure 1). Cell

counts from three slices were averaged within each animal and used for statistical analysis.

Behavioral Experiments

Rats were trained on Barnes Maze task to assess spatial learning 1-month post-injury. The

Barnes maze table (Maze Engineers, https://conductscience.com/maze/) was 122 cm in

diameter with 20 holes (10 cm each). The target hole was equipped with an escape box (22.5

cm x 14 cm x 9 cm LxWxH). The maze table was set up within black curtain walls with visual

cues and a camera installed above to record animal movement. Two bright floodlights and

a fan served as aversive stimuli to encourage animals reach the escape box. Animals were

held in their home cage outside the curtain in a dark room until their turn to run the trial. On

day 1, rats were habituated to the behavior room for at least 1 hour and then placed in the

middle of the table inside starter cup for 1 minute following which they were allowed to

explore the table for 180 seconds. At the end of the trial, animals were gently guided to their

designated escape hole. The escape hole locations were randomly assigned to each rat and

remained the same till the end of the experiment. The table was thoroughly wiped with

ethanol between tests to eliminate odor cues. Rats were subjected to a single 180-second

trial for 3 days. Task acquisition and retention were confirmed through tests on day 4 and

day 7. Four rats that fell from the table and exhibited signs of lethargy were removed from

the test and sacrificed. Behavioral analysis was performed by a blinded experimenter using

Any-maze software. Parameters such as latency to escape hole, distance travelled and

spatial search strategy were evaluated between sham and injured rats. Spatial search

strategies were scored based on 11-point scoring scheme[42]. Thigmotaxis was quantified

by defining a peripheral zone of the testing arena that was approximately 12 cm from the

outer edges of the apparatus. The proportion of time the animals spent within this

designated zone was recorded and used as the metric for anxiety-like behavior [43].

Statistics

Data are presented as mean ± standard error of the mean (SEM) for parametric measures,

or as median and interquartile range (IQR) where non-parametric distributions were

observed. To ensure rigorous data interpretation and minimize investigator bias, all data

analysis, including quantification of in situ zymography, electrophysiological recordings,

and behavioral scoring, was performed by researchers blinded to the treatment groups and

genotypes.

Normality of data distributions was assessed using the Shapiro-Wilk test. For the analysis

of synaptic events (sEPSC, mEPSC, sIPSC, and mIPSC), cumulative probability distributions

of inter-event intervals (IEI) and amplitudes were compared across groups using equal

number of consecutive events with the non-parametric Kolmogorov-Smirnov (K-S) test.

Effect sizes for these distributions were calculated using Cohen's d. When multiple

comparisons were involved (Figures 1&3, Supplementary figure 2) Holms-Bonferroni correction

was adopted and corrected P values are reported (See Supplementary table 3). Differences

between two groups were evaluated using two-tailed unpaired t-tests. For experiments

involving multiple factors, such as injury and drug treatment or injury and genotype, two-way

ANOVA, two-way repeated measures (RM) ANOVA (for input/output curves and behavioral

training days) or three-way ANOVA were employed, followed by Šídák's multiple

comparisons test where appropriate. Significance was defined at $P<0.05$. Statistical outlier

tests were not applied. Statistical analyses were performed using Graphpad Prism 10.0.1

(Boston, MA).

RESULTS:

Neuronal TLR4 signaling is necessary for early excitatory circuit changes after brain injury.

Prior studies have identified TLR4 contribution to early changes in glutamatergic synapse strength after brain injury [12–14]. However, the nature of early circuit reorganization after brain injury in the dorsal hippocampus and whether immune signaling regulates these changes has not been examined. Whole-cell voltage clamp recordings from DGCs in dorsal hippocampal slices one-week after FPI revealed an early decrease in the inter-event intervals (IEI) of spontaneous excitatory postsynaptic currents (sEPSCs), indicating an increase in sEPSC frequency ($P<0.0001$ by K-S test with Holm-Šídák correction; Cohen's d = 0.34, n = 8 cells/ group, Figure 1A-E, Supplementary table1 & 3). Additionally, we observed a reduction in sEPSC amplitude in DGCs from rats after brain injury (n = 8 cells/ group, $P<0.0001$ by K-S test, Cohen's d = 0.41). To assess potential post-traumatic synaptic reorganization, we recorded action potential-independent miniature EPSCs (mEPSCs) in DGCs the presence of tetrodotoxin (TTX, 1 µM). Similar to sEPSCs, DGCs in rats one week after FPI exhibited shorter mEPSC IEI and smaller mEPSC amplitudes compared to Sham controls ($P <0.001$ by K-S test with Holm-Šídák correction; Cohen's d = 0.22 for IEI and 0.85 for amplitude, n = 7 cells/ group, Supplementary figure 2A-C, Supplementary table 2 & 3). Together, these findings provide the first evidence of an increase in excitatory synaptic inputs to DGCs in dorsal hippocampus of rats one week after injury.

Significant changes in synaptic inhibition are often observed early after brain injury

[5,7,44]. To evaluate whether inhibitory circuits are reorganized alongside the excitatory

circuits, we examined spontaneous inhibitory postsynaptic potentials (sIPSC) in the same

neurons. In contrast to changes in sEPSCs, sIPSC IEI was prolonged in DGCs from injured

rats (P <0.0001 by K-S test with Holm-Šídák correction; Cohen's d = 0.29, n= 8 cells/group,

Figure 1F-H, Supplementary tables 1 & 3). These findings indicate that in addition to

receiving more frequent excitatory inputs, the DGCs receive fewer inhibitory inputs after

brain injury. DGCs also showed an increase in sIPSC amplitude compared to sham

controls after brain injury (P <0.0001 by K-S test with Holm-Šídák correction; Cohen's d =

0.16, n = 8 cells/ group). These findings differ from our findings in the ventral hippocampus

demonstrating an increase in sIPSC frequency in DGCs after FPI [36,45,46]. We also

observed an increase in mIPSC amplitudes with no change in mIPSC IEI (Supplementary

Figure 2D-F, Supplementary tables 2 & 3). Together, these findings indicate an early

dysregulation in excitatory and inhibitory inputs to DGC's one-week after FPI. Furthermore,

these results identify early circuit reorganization after FPI and reveal potential differences

in synaptic inhibition and post-traumatic plasticity in the dorsal vs ventral hippocampus.

To determine whether inhibition of TLR4 signaling early after injury modified DGC synaptic

currents, we treated sham and FPI rats with TLR4 inhibitor TAK-242 (0.5mg/kg, i.p, 30 min, 8

hours and 24 hours post-injury). TAK-242 treatment reduced changes in sEPSC and mEPSC

frequency and amplitude in injured rats (P <0.0001 by K-S test with Holm-Šídák correction;

Cohen's d = 0.62 for sEPSC IEI, 0.50 for sEPSC amplitude, 0.54 for mEPSC IEI and 0.89 for

mEPSC amplitude, n = 8-9 cells/ group, Figure.1A-E, Supplementary figure 2A-C, and

Supplementary tables 1 & 3). In addition, TAK-242 reduced injury-driven changes in sIPSC

frequency and amplitude (P <0.0001 by K-S test with Holm-Šídák correction; Cohen's d =

0.22 for IEI, 0.11 for amplitude, n = 7-8 cells/ group, Figure.1F-H, Supplementary tables 2 &

3). While mIPSC frequency was unaltered, mIPSC amplitude was reduced by TAK-242

treatment in injured rats (P <0.0001 by K-S test with Holm-Šídák correction; Cohen's d =

0.47 for amplitude, n = 6-7 cells/ group, Supplementary figure 2D-F, Supplementary tables

2 & 3). These findings demonstrate a role for TLR4 signaling in driving early synaptic

dysregulation after FPI.

In contrast to injured animals, sham controls receiving TAK-242 showed paradoxically

opposing changes in excitatory and inhibitory synaptic inputs to DGCs. In sham controls,

TAK-242 treatment reduced sEPSC IEI in DGCs, increasing frequency as observed after FPI

but failed to alter sEPSC amplitude (P <0.0001 by K-S test with Holm-Šídák correction;

Cohen's d = 0.35 for IEI, n = 8-9 cells/ group, Figure 1B-E). Both mEPSC interval and

amplitude were reduced in TAK-242 treated sham controls (P <0.0001 by K-S test with

Holm-Šídák correction; Cohen's d = 0.24 for IEI and 0.30 for amplitude, n = 7-8 cells/

group, Supplementary figure 2A-C, Supplementary tables 2 & 3) consistent with synaptic

reorganization. Moreover, TAK-242 treated sham controls showed a reduction in DGC

sIPSC IEI and amplitude suggesting more frequent inhibitory inputs (P <0.01 by K-S test

with Holm-Šídák correction; Cohen's d = 0.07 for IEI and 0.21 for amplitude, n = 7-8 cells/

group, Figure 1F-H, Supplementary tables 1 & 3). However, TAK-242 treatment did not alter

mIPSC frequency or amplitude. Together, these findings identify a novel contribution of

basal TLR4 signaling in maintaining excitatory synaptic homeostasis in uninjured animals.

Our previous studies demonstrated neuronal TLR4 signaling as critical for post-traumatic

increase in DGCs AMPA currents [13]. To test whether the glutamatergic circuit changes

identified above require neuronal TLR4 signaling, we examined changes in sEPSC

frequency and amplitude in mice with selective deletion of TLR4 in CaMKII expressing

excitatory neurons (TLR4$^{fl/fl}$-CaMKIIa-CreERT2 mice). Care was taken to avoid constitutive

and developmental effects by inducing Cre recombinase just prior to injury.

Immunostaining for TLR4 in mice three-days after FPI, following the 3-day tamoxifen

induction strategy, revealed strong immunostaining for TLR4 in the granule cell layer and

hilus of Cre^{-ve} mice which was absent in Cre^{+ve} mice, confirming TLR4 deletion in DGCs

(Nested t-test: t(4) = 7.57, P = 0.0016, F(1,4) = 58.82, n = 5-6 slices from 3 mice/ group,

Supplementary figure 3A-C). In addition, RNAscope analysis of TLR4 mRNA confirmed the

lack of injury-driven TLR4 mRNA increase in Cre^{+ve} mice (F(2,9) = 8.353, P<0.05 by one-way

ANOVA, n = 3-5 mice/ group, Supplementary figure 3D-F). Validation studies to test for

effect on TLR4 deletion on neuronal intrinsic physiology failed to reveal genotype-

dependent effects on the resting membrane potential, input resistance or rheobase in

DGCs from sham controls (Supplementary figure 4C-E). However, there was a significant

decrease in capacitance and an increase in firing frequency in Cre^{+ve} sham mice compared

to Cre^{-ve} sham mice (Capacitance - Effect of Genotype- F(1,23) = 6.9, P = 0.015 by two-way

ANOVA; Firing frequency – Effect of Genotype – F(1,23) = 4.706, P =0.040 by three-way

ANOVA, n = 7 cells/ group, Supplementary figure 4B&F). Consistent with our previous

reports in rat and mice [31,47], there were no injury effects on DGC active properties in

either genotype. While there was no injury effect on DGC resting membrane potential in

Cre[-ve] mice, DGCs from Cre[+ve] mice with TLR4 deletion were more depolarized

(Supplementary figure 4C). Together these validation studies demonstrate the lack of

injury-induced TLR4 increase in Cre[+ve] mice and suggest a role for TLR4 in basal DGC

excitability and maintenance of RMP after injury.

Having established the lack of posttraumatic TLR4 upregulation in Cre[+ve] mice we

examined the contribution of neuronal TLR4 signaling to injury-induced changes in

synaptic inputs to DGCs. Similar to our findings in rat, brain injury reduced DGC sEPSC IEI

and amplitude in Cre[-ve] mice with intact TLR4 signaling (P <0.0001 by K-S test with Holm-

Šídák correction; Cohen's d = 0.42 for IEI and 0.31 for amplitude, n = 7 cells/ group, Figure

1I-L, Supplementary table 1). Importantly, DGCs from Cre[+ve] mice with conditional deletion

of TLR4 in excitatory neurons failed to exhibit injury-induced decrease in sEPSC IEI, but

showed a decrease in sEPSC amplitude after brain injury (P = 0.0008 by K-S test with Holm-

Šídák correction; Cohen's d = 0.16 for amplitude, n = 7 cells/ group. Interestingly, DGC

sEPSC IEI and amplitude were not different between Cre[-ve] and Cre[+ve] sham injured mice.

These findings demonstrate that TLR4 expression in glutamatergic neurons is necessary for

the early injury-driven changes in excitatory synaptic inputs to DGCs.

MMP-9 activity is required for TLR4-dependent changes in DGC excitatory synapses

after brain injury

We next sought to identify downstream molecular mechanisms through which TLR4

signaling dysregulates DGC synaptic inputs after brain injury. Several studies have reported

an upregulation in Matrix Metallproteinase-9 (MMP-9) expression in pre-clinical models of

TBI [27,30,35]. MMP-9 is a critical regulator of synaptic strength and plasticity [17,18], and

perturbations in MMP-9 activity is known to dysregulate synaptic physiology [19,40,48–51] .

Although extrinsic activation of TLR4 signaling has been shown to enhance MMP-9

expression in primary astrocyte/neuronal co-cultures and *in vivo* [15,16], whether TLR4

regulates MMP-9 activity and if this can induced synaptic and functional changes *in vivo*

and in the injured brain remains to be determined. Moreover, whether *neuronal* TLR4

regulates MMP-9 activity has not been examined.

To directly assess MMP-9 activity, we performed *in-situ* zymography on hippocampal slices

obtained from sham/FPI rats 48 hours after injury, focusing on the DG molecular layer

where the afferent glutamatergic synapses are located. DQ-fluorescence intensity values

were normalized to CA1 Stratum Radiatum (SR) since no changes were observed in this

region with injury, treatment or genotypes (Supplementary figure 4I-J, see methods). Our

experiments identified a significant increase in the MMP-9 activity in the DG molecular

layer in brain-injured rats compared to sham controls (MMP-activity normalized to CA1 SR:

Sham:14.56±2.15%, FPI:30.43±2.89%, *n* = 5 sham and 6 FPI rats, Figure 2A-C).

Pharmacologically blocking TLR4 early after FPI reduced MMP-9 activity in the DG of brain

injured rats (Sham$_{TAK-242}$: 17.22±2.68%, FPI $_{TAK-242}$: 16.36±2.97%, *n* = 5 Sham and 6 FPI rats;

Effect of Injury x Treatment F (1,18) = 9.13; *P* <0.05 by Two-way ANOVA). In contrast, MMP-9

activity in sham controls treated with TAK-242 was not different from vehicle treated sham

controls. These results identify TLR4 regulation of MMP-9 activity in the injured brain.

To isolate the role of neuronal TLR4 in regulating MMP-9 activity, we performed in-situ

zymography on mice with neuronal TLR4 deletion (TLR4$^{fl/fl}$-CaMKIIa-CreERT2) and Cre^{-ve}

controls (Supplementary figure 4) 48 hours after Sham or FPI. We did not observe any

difference in MMP-9 activity between Cre^{-ve} sham and Cre^{+ve} sham mice, indicating that

TLR4 deletion does not alter normal MMP-9 levels without injury (Figure 2D-F). While Cre^{-ve}

mice showed an increase in MMP-9 levels after injury, mice with selective TLR4 deletion in

glutamatergic neurons failed to develop elevated MMP-9 activity after FPI (MMP-activity

normalized to CA1 SR: Cre^{-ve} Sham: 13.43±3.36%, Cre^{-ve} FPI: 42.79±5.02%; Cre^{+ve} SHAM:

10.59±3.15%, Cre^{+ve} FPI: 12.20±1.16%., n=3-4 mice/group; Effect of Injury x Genotype, F

$(1,11) = 14.70$; $P<0.05$ by Two-way ANOVA, Figure 2D-F). These findings demonstrate that

neuronal TLR4 is necessary for the post-injury upregulation of MMP-9 activity.

Next, we tested the possibility that TLR4 signaling increased MMP-9 levels through

transcriptional upregulation. Naïve rats (P30) were injected with a selective TLR4 agonist,

LPS-B5 (5mg/kg., i.p.) or saline and euthanized after six hours for single molecule

fluorescent in situ hybridization (smFISH) to quantify MMP-9 mRNA levels. Sections from

LPS-B5 injected rats showed a significant increase in MMP-9 mRNA in DGCs, indicating

that TLR4 activation enhances MMP-9 via transcriptional upregulation (intensity A.U/µm^2:

saline controls 1.67±0.15; LPS-B5 treated 2.41±0.19; n= 5/group, Figure 2G-I). Furthermore,

inhibition of NF-κB, a key downstream effector of TLR4, using JSH-23 (10 mg/kg)

administered one hour before LPS-B5 injection abolished the upregulation of MMP-9

(intensity A.U/µm^2 : JSH-23 + LPS-B5 1.319±0.16, n = 5 rats, Main effect F$(2,12)$ =11.20, P =

0.002, Figure 2G-I) Together, these findings identify TLR4 and NF-κB dependent

transcriptional upregulation of MMP-9 as a candidate mechanism for increase in MMP9

activity in the DG following brain injury.

MMP-9 has been shown to play a crucial role in plasticity of excitatory and inhibitory

synapses [52]. Therefore, we reasoned that TLR4 mediated increase in MMP-9 activity may

be required for the post-traumatic changes in synaptic inputs to DGCs. To examine the

acute effects of TLR4 signaling *ex vivo*, we compared sEPSCs in hippocampal slices from

sham rats incubated in LPS-B5 (10 µg/ml for 90min) or aCSF. Acute LPS-B5 incubation

significantly reduced sEPSC IEI in DGCs, mimicking the increase in sEPSC frequency

observed in FPI animals (Sham vs LPS-B5 incubated: P <0.0001 by K-S test with Holm-

Šídák correction; Cohen's d = 0.20 for IEI, n = 8 cells from 4 rats, Figure 3A-B,C). However,

LPS-B5 incubation increased the sEPSC amplitude (Figure 3A-B,D). Importantly, inhibiting

MMP-9 using SB-3CT (10 µM) mitigated LPS-B5-induced changes in sEPSC IEI indicating

that TLR4-mediated alterations in excitatory synaptic inputs to DGCs require MMP-9

activity (LPS-B5 incubated vs SB-3CT+LPS-B5: P <0.0001 by K-S test with Holm-Šídák

correction; Cohen's d = 0.21 for IEI, n = 9 cells from 4 rats, Figure 3A-C). Conversely, SB-

3CT co-incubation failed to reduce, and further exacerbated, the LPS-B5 mediated

increase in sEPSC amplitude (Figure 3B, D). Acute TLR4 activation failed to alter sIPSC

frequency but increased mIPSC interevent intervals in DGCs (Supplementary figure 5A-B).

Since we find that TLR4-mediated alterations in DGC sEPSC frequency requires MMP-9

activity, we tested if systemic inhibition of MMP-9 using SB-3CT treatment (50 mg/kg, i.p.)

early after FPI would prevent post-traumatic changes in excitatory inputs to DGCs. SB-3CT

treatment effectively mitigated injury-induced the changes in sEPSC IEI and amplitude

one-week after FPI in rat (P <0.0001 by K-S test with Holm-Šídák correction; Cohen's d =

0.60 for IEI and 0.30 for amplitude, n = 7 cells / group, see Supplementary table1&3, Figure

3E-H). Since Cre[+ve] mice failed to show posttraumatic increase in sEPSC frequency (Figure

1K), we examined Cre[-ve] mice to assess the role of MMP9 in injury-induced changes in

sEPSC in mice. Consistent with our findings in rat, SB-3CT treatment reduced post-injury

changes in sEPSC IEI in DGCs from Cre[-ve] -negative TLR4[fl/fl]-CaMKIIa-CreERT2 mice (P

<0.0001 by K-S test with Holm-Šídák correction; Cohen's d = 0.283 for IEI, n = 9 cells

Figure 3I-L). SB-3CT treatment did not have any effect on sIPSC frequency or amplitude

(Supplementary figure 5G-I) after FPI suggesting that TLR4 regulation of DGC inhibitory

synaptic parameters after injury does not require MMP-9 activation. However, SB-3CT

caused a small increase in sIPSC amplitude in sham controls (Supplementary figure 5I).

Together, these results indicate that injury-driven changes in excitatory, but not inhibitory,

inputs to DGCs required both TLR4 signaling and MMP-9 activity.

Early activation of TLR4 - MMP-9 molecular axis contributes to DG network

dysfunction after brain injury

Having established that the TLR4- MMP-9 mechanistic axis increases glutamatergic inputs

to DGCs after brain injury, we examined the contribution of this molecular pathway to early

changes in DG circuit function in rats *in vivo* one-week after brain injury. Input/output plots

of the DG population spike amplitude in response to perforant path stimulation at

increasing current intensities was recorded under urethane anesthesia. DG population

spike responses normalized to peak amplitude, revealed a leftward shift after FPI

indicating increased network excitability (n = 6 SHAM and 5 FPI rats; Effect of injury – F (1,9)

= 6.462, *P* = 0.031 by Two-way RM ANOVA, Figure 4A-C). This increase in network

excitability was mitigated by early inhibition of either TLR4 signaling or MMP-9 activity,

demonstrating their contribution in promoting post-traumatic DG circuit dysfunction (n= 5-

6 rats/group; Effect of Stimulus x Treatment: F (6,166,40.08) = 4.752, *P*=0.0009 by Two-way

RM ANOVA, Figure 4D). Paradoxically, as previously reported in slices [12–14], TAK-242

treatment increased DG population spike amplitude in sham controls (Figure 4E). However,

consistent with its lack of effect on EPSC frequency in sham controls, SB-3CT treatment

did not alter DG network excitability in sham rats (Figure 4E).

Enhanced network excitability combined with deficits in synaptic inhibition can increase

dentate throughput. We assessed DG throughput by examining dentate spikes, a distinct

DG population event driven by entorhinal cortex [53,54]. Linear 16- channel silicon probe

recordings of dentate spikes in anesthetized rats (Figure 4F-G) revealed a significant

increase in the frequency of dentate spikes one-week after FPI (in Hz: Sham: 0.11±0.019;

FPI 0.24±0.025, n = 4-5 rats/group, *P*= 0.006 by two tailed unpaired t-test), identifying

enhanced dentate throughput following FPI. To test if inhibition of either TLR4 signaling or

MMP-9 activity early after injury would reduce changes to DG throughput, we examined

dentate spikes in TAK-242 or SB-3CT treated sham/injured rats. Our experiments

demonstrate that early inhibition of either TLR4 signaling or MMP-9 activity effectively

prevented the post injury increase in dentate spike frequency without impacting it in sham

controls (n = 4-5 rats/group; Effect of Interaction F (2,21) = 9.975, *P*<0.001 by Two-way

ANOVA, Figure 4G).

We previously demonstrated that early post-injury increase in DGC calcium permeable

AMPA currents and deficits in working memory were mitigated by TLR4 inhibition [13]. To

assess whether the TLR4-MMP9 signaling impacts synaptic plasticity, we examined long-

term potentiation (LTP) at perforant path-DGC synapses in anesthetized rats *in vivo* using

theta-burst stimulation (TBS) [38]. In the first 10 minutes after TBS, the field EPSP slope

normalized to the maximum slope at baseline trended lower but was not statistically

different between sham and FPI. However, there was a reduction in the field EPSP slope at

60 min, 120 min, and 180 min post-TBS indicating impairment in LTP maintenance (n = 6

rats/group, Effect of injury : $F_{(1,10)}$ = 32.79, *P*=0.0002 by Two-way RM ANOVA with Šídák's

multiple comparisons 10 min: *P* =0.065, 60 min: *P* = 0.003, 120 min: *P* = 0.002, 180 min: *P* =

0.125, Figure 5A-E). Crucially, systemic antagonism of either TLR4 or MMP-9 early after FPI

effectively prevented deficits in LTP in injured rats, demonstrating contribution of TLR4-

MMP-9 axis to LTP deficits after brain injury (Effect of treatment: $F_{(2,13)}$ = 18, *P*=0.0002; *n*=

5-6 rats/group; See Supplementary table 4, Figure 5A-E). Consistent with the paradoxical

effects of TAK-242 on sEPSC and network excitability in sham controls, TAK-242 treatment

resulted in deficits in LTP in sham rats (Effect of treatment: $F_{(2,13)}$ = 8.046, *P*=0.0053; *n*= 5-

6 rats/group; Figure 5E). Once again, unlike TAK242, SB-3CT treatment failed to impact LTP

Sham rats, paralleling on the differential effects of the treatments on sEPSC frequency,

population spike amplitude and dentate spike frequency in sham animals. These findings

demonstrate that TLR4-MMP-9 dependent post-traumatic circuit changes compromise DG

glutamatergic circuit function and plasticity.

Targeting TLR4 or MMP-9 axis early after brain injury limits long-term spatial memory impairments.

Since we identify post injury increase in dentate spikes which can compromise hippocampal information processing [55–58], and blocking TLR4 and MMP-9 early after injury prevent this change, we sought to determine if TLR4 and MMP-9 antagonists could alleviate long term cognitive deficits after brain injury. We adopted a Barnes maze spatial learning task which has been shown to rely on intact DG function[59,60]. One month after sham/FPI, brain injured rats exhibited a longer latency to find the escape hole and traveled a greater total distance compared to sham controls early during training but reached control levels of task performance by Day 3 (See Supplementary table 5, n =10 rats / group, P<0.05 by TW-RM ANOVA, Figure 6A-D). Additionally, there was no difference in task performance during trials on Day 4 and Day 7, indicating that task retention is not impaired (Supplementary figure 6B-C). Although there were no differences in speed of locomotion (Supplementary figure 6D), injured rats showed increased thigmotaxis on Days 1 and 2 (Time spent in thigmotaxis zone in seconds: $Sham_{DMSO}$ – 24.73±5.42, FPI_{DMSO} – 64.99±16.96; n= 10 rats/group, Effect of injury– F (1,18) = 6.841, P<0.05 by Two-way -RM ANOVA, Figure 6I). Since injured rats showed a significant increase in both thigmotaxis and latency to find the escape on Day 2 of task acquisition, we examined the use of spatial strategy for navigation using a 11-point scoring system [42]. Analysis of search strategies on Day 2 showed that injured animals employed random or serial strategies more often, demonstrating specific deficits in ability to recall spatial cues for navigation (Search strategy score_Day2: $Sham_{DMSO}$ – 7.7±0.73, FPI_{DMSO} – 2.4±0.47; n= 10 rats/group, Effect of injury– F (1,18) = 17, P<0.01 by Two-way RM ANOVA,

Figure.6E,H). These results suggest a mild delay in task acquisition after brain injury which

is apparent on Day 2 of the task.

We focused our analysis on Day 2 of task acquisition when the injury-induced impairment is

most apparent. Early inhibition of TLR4 signaling or MMP-9 activity prevented post-injury

increases in latency to escape, and distance traveled observed on Day 2 (n = 6-10 rats/group;

P<0.05 by Two-way RM ANOVA, Figure 6F-G, Supplementary table 3 and Supplementary

figure 6A). While the trend in ability of TLR4 inhibition to improve the strategy score in injured

animals did not reach statistical significance (Figure 6H), the search strategy scores in

injured animals significantly improved with MMP-9 inhibition (Figure 6H). Importantly, while

MMP-9 did not have any adverse effects on sham controls, TLR4 inhibition showed a trend

towards increased latency, distance travelled and thigmotaxis, and a lower spatial search

strategy score (Figure 6F-I) which did not reach statistical significance. Collectively, these

findings demonstrate that early inhibition of the TLR4-MMP-9 signaling axis effectively

mitigates mild long term spatial learning deficits observed after FPI.

DISCUSSION:

Understanding how immune signaling modulates neuronal function after brain injury is

critical for developing effective therapeutic strategies to minimize the long-term sequelae

of TBI. Aside from the widespread secondary injury driven by glial mediators of

neuroinflammation [1,2], we previously showed that the neuronal expression of innate

immune receptor TLR4 is upregulated in the hippocampus early after injury, contributing to

long-term seizure susceptibility and impaired working memory [12–14]. Here, we provide

mechanistic evidence that TLR4 signaling mediates rapid, circuit-level changes in both

excitatory and inhibitory synaptic transmission after injury. Using a combination of

pharmacology in rats and mouse genetics we demonstrate a central role for TLR4

expressed in CaMKII-positive excitatory neurons in mediating the posttraumatic circuit

changes through the upregulation of MMP-9. Importantly, we show that either TLR4 or

MMP-9 inhibition early after brain injury not only reduces these pathological changes one-

week after injury, but improves deficits in spatial learning 1-month after injury.

TLR4 Signaling Augments Glutamatergic inputs to DGCs after brain injury.

TLR4 signaling is well established as a driver of neuronal excitability [3,61–65]. Our prior

works showed that TLR4 signaling enhances AMPA currents in granule cells contributing to

DG network hyperexcitability [12,13]. Examining circuit-level changes in animals one week

after injury, this study identifies an aberrant increase in excitatory inputs to DGCs in the

dorsal hippocampus. Considering that perforant path fibers are severed in coronal

hippocampal slice preparations [66], the post-injury increase in spontaneous EPSCs and

action potential-independent miniature events observed here indicate an increase in

synaptic contacts, consistent with early axonal sprouting [46], or elevated mossy cell

excitability previously reported after FPI [67]. Our findings highlight a critical role of TLR4

signaling in this process. First, we show that early pharmacological inhibition of TLR4

mitigates changes in both spontaneous and miniature EPSC frequency. We further

demonstrate that transient activation of TLR4 signaling in acute brain slices using LPS-B5 is

sufficient to augment sEPSCs frequency to levels comparable to those seen in injured

animals. Furthermore, by leveraging TLR4$^{fl/fl}$: CaMKIIa-CreERT2 mice, we demonstrate that

selective deletion of TLR4 in glutamatergic neurons alone is sufficient to mitigate the post-

injury increase in sEPSC frequency. Together, these findings show that neuronal TLR4 is

necessary to enhance glutamatergic drive to DGCs after brain injury.

A Novel Neuro-Immune-enzyme Signaling Cascade Underlies Early Post-Traumatic

Glutamatergic Circuit Plasticity

To elucidate the mechanism by which TLR4 enhances glutamatergic drive to DGCs after

injury, we focused on MMP-9 as a potential downstream effector. MMP-9 co-localizes with

excitatory synapses in an activity-dependent manner, and increased MMP-9 activity is

known to dysregulate glutamatergic synaptic transmission [40,52,68–70]. We previously

reported that neuronal TLR4 enhances calcium-permeable AMPA currents in DGCs post-

injury by increasing the surface expression of GluA1 subunits [13]. Notably, MMP-9 activity

has been shown to regulate AMPA receptor mobility and synaptic accumulation of GluA1

subunits following chemically induced LTP [71]. In addition, TLR4 ligands such as high-

mobility group box 1 (HMGB1) and lipopolysaccharide (LPS) are known to enhance MMP-9

expression in cultures and following ischemic insult [15,16] making MMP9 an attractive

candidate effector of the synaptic effects of TLR4.

Here, we show that pharmacologically inhibiting TLR4 signaling (rats) or deleting TLR4 in

excitatory neurons (mice) both limit post-injury increases in MMP-9 activity identifying that

TLR4 signaling is necessary for injury-induced increase in MMP9. Furthermore, using LPS-

B5 to selectively activate TLR4, we demonstrate that activation of TLR4 is sufficient to

increase MMP-9 mRNA levels in the dentate gyrus of naïve rats in an NF-κB dependent

manner. Thus, as summarized in Figure 7, our findings identify a novel TLR4/NF-κB-

dependent transcriptional pathway that upregulates MMP-9 activity and promotes

glutamatergic circuit reorganization following brain injury. Consistent with this, inhibition of

MMP-9 with SB-3CT effectively reduced most of the glutamatergic circuit changes in the

DG resulting from direct activation of TLR4 signaling in slices (LPS-B5) and brain injury *in*

vivo.

Mechanistic Divergence in TLR4 regulation of glutamatergic and GABAergic circuits

In contrast to the glutamatergic inputs, we observed a decrease in the frequency of

inhibitory inputs to dorsal DGCs one week after FPI. This stands in direct contrast to our

previous findings from the ventral hippocampus [36,45,46], highlighting potential regional

differences in functional plasticity between dorsal and ventral hippocampal inhibitory

circuits [72]. While TLR4 inhibition reduced post-traumatic changes in IPSCs, this effect

was independent of MMP-9 activity. Although enhances MMP-9 activity is implicated in

hypofunction of parvalbumin expressing inhibitory neurons and perineuronal net

remodeling in neurodevelopmental disorders [73], we find that blocking MMP9 does not

impact changes in inhibitory synaptic inputs after brain injury. Notably, our findings

indicate a divergence of TLR4-dependent mechanisms for glutamatergic and GABAergic

circuit reorganization, with MMP-9-independent mechanisms contributing to changes in

inhibitory synapses (Figure 7). Interestingly, unlike after brain injury, selective activation of

TLR4 with LPS-B5 in slices from sham rats did not alter sIPSC frequency in DGCs raising the

possibility that injury-driven loss of hilar interneurons contributes to the reduction in sIPSC

after brain injury [7,47]. However, mIPSC frequency was increased following LPS-B5

incubation suggesting perturbations in release probability or presynaptic calcium [44,74–

76]. Curiously, mice with TLR4 deletion developed posttraumatic depolarization of RMP

not observed in cre[-ve] controls suggesting TLR4 signaling may contribute to cell intrinsic

compensatory processes. Future studies should determine whether neuronal or glial TLR4

signaling mediates the TLR4 dependent post-injury changes in inhibition and how this

impacts dentate function and behaviors.

Network Consequences and behavioral impact of TLR4/MMP-9 pathway.

The dentate gyrus is a central node in hippocampal memory processing, regulating the flow

of information through the hippocampal circuit [56,77,78]. Consequently, perturbations in

DG circuit function can directly impact cognitive processes. Our earlier studies showed

enhanced perforant path evoked EPSC amplitude in DGCs after FPI, suggesting stronger

entorhinal-dentate coupling in the injured brain, coupled with deficits in working memory

[12–14]. Here we advance these findings using in vivo approaches to identify increased

dentate network excitability early after injury. We further show that while LTP induction was

unaffected after injury, LTP maintenance was compromised in injured rats. Indeed the lack

of deficit in LTP induction argues against a role for synaptic saturation in mediating LTP

deficits. Rather, we find selective deficits in LTP maintenance after brain injury indicating

altered transcriptional processes. Importantly, we report an increase in dentate spike

frequency early after brain injury indicating abnormal entorhinal-dentate throughput. Since

dentate spikes are critical in hippocampal information processing, memory formation, and

consolidation [53–57,79,80], abnormal dentate spikes could impair memory processing

after brain injury. Supporting this proposal, treatments that reduced dentate spike

frequency in injured animals also improved spatial learning deficits. Crucially, dentate

spikes are also increased in experimental models of epilepsy [81], suggesting that these

events may serve as biomarkers for post-traumatic epileptogenesis alongside the

pathological high frequency oscillations and altered sleep spindles identified previously

[82,83]. In summary, inhibition of the novel TLR4-MMP9 signaling axis after brain injury

preserves dentate circuit excitability and long-term potentiation while rescuing spatial

memory deficits (Figure 7). These results highlight the manipulation of this pathway as a

promising therapeutic strategy for post-traumatic recovery.

A Novel Constitutive Role for TLR4 Signaling in Basal Circuit Homeostasis

Our study reveals a fundamental role for basal TLR4 signaling in maintaining dentate circuit

homeostasis. We found that inhibition of TLR4 signaling in uninjured animals consistently

altered synaptic inputs, network excitability, and plasticity. Specifically, timed and

selective deletion of TLR4 in glutamatergic neurons in a mature circuit caused a genotype-

dependent reduction in granule cell capacitance and increased firing rates (Figure 7). This

indicates a constitutive role for neuronal TLR4 in limiting basal excitability. A reduction in

capacitance with increased firing in the absence of changes in action potential threshold

parallels observations in parvalbumin neurons in epilepsy models, where regulation of

perineuronal nets alters neuronal firing through effects on cell capacitance[41,84,85].

Indeed, our prior studies have found that ability of TLR4 blockers to increase dentate

excitability in uninjured controls is abolished when glial metabolic processes are

disrupted[14]. Our finding that TLR4 or MMP9 antagonism does not impact dentate spike

frequency in uninjured controls, is consistent with lack of TLR4 modulation of perforant

path evoked AMPA currents in the absence of injury. Nevertheless, the basal TLR4

regulation of network homeostasis appears critical as blocking TLR4 enhances network

excitability, impairs LTP maintenance and spatial memory (Figure 7). The potential for TLR4

inhibitors to limit the progression of epilepsy and Alzheimer's disease [11,61,86,87],

highlights the need to understand non-canonical TLR4 signaling in the healthy brain. Future

research should focus on the underlying mechanisms of basal TLR4-mediated neuronal

regulation.

Collectively, our results position the TLR4-NFkB-MMP-9 axis as a central driver of TBI-

induced circuit dysfunction, providing a targetable molecular pathway to serve as a

therapeutic target for mitigating post-traumatic epilepsy and associated cognitive deficits.

Declarations

Ethics approval and consent to participate

All experiments were performed in accordance with IACUC protocols approved by the University of California-Riverside, CA (Approval number #20210005/35) and in keeping with the ARRIVE guidelines.

Consent for publication

Not applicable.

Availability of data and materials

All data are available from the corresponding author upon reasonable request.

Competing interests

The authors declare no competing interests.

Funding

This work is supported by National Institutes of Health (NIH) NINDS R01 NS069861, R01NS097750 to V.S., NIH/NINDS F31NS124290 to L.D; CDMRP- Epilepsy Research Program grant W81XWH-21-1-0684 to D.S.

Authors' contributions

D.S., E.C., and L.D., performed experiments; D.S., E.C., L.D., E.G., Y.L., and R.J., analyzed data; D.S., I.E., and V.S., interpreted results of experiments; D.S. prepared figures; D.S., V.S. conception and design of research; D.S. and V.S., drafted manuscript; D.S., I.E., and V.S. finalized and approved manuscript.

Acknowledgements: We thank Dr. Leszek Kaczmarek (Nencki Institute, Warsaw, Poland) for his expert advice on *in situ* zymography. We also thank Dr. Amanda Schott and Dr. Jordan S Farrell (Harvard Medical School) for their expertise and help with Dentate spike detection algorithms.

REFERENCES:

1. Sharma R, Leung WL, Zamani A, O'brien TJ, Espinosa PMC, Semple BD. Neuroinflammation in post-traumatic epilepsy: Pathophysiology and tractable therapeutic targets. Brain Sciences. 2019 Nov 1;9(11). doi:10.3390/brainsci9110318

2. Mukherjee S, Arisi GM, Mims K, Hollingsworth G, O'Neil K, Shapiro LA. Neuroinflammatory mechanisms of post-traumatic epilepsy. Journal of Neuroinflammation. 2020 Jun 17;17(1). doi:10.1186/s12974-020-01854-w PubMed PMID: 32552898.

3. Webster KM, Sun M, Crack P, O'Brien TJ, Shultz SR, Semple BD. Inflammation in epileptogenesis after traumatic brain injury. Journal of Neuroinflammation. 2017 Jan 13;14(1). doi:10.1186/s12974-016-0786-1 PubMed PMID: 28086980.

4. Simon DW, McGeachy MJ, Baylr H, Clark RSB, Loane DJ, Kochanek PM. The far-reaching scope of neuroinflammation after traumatic brain injury. Nature Reviews Neurology. 2017 Mar 1;13(3):171–91. doi:10.1038/nrneurol.2017.13 PubMed PMID: 28186177.

5. Lowenstein D, Thomas M, Smith D, McIntosh T. Selective vulnerability of dentate hilar neurons following traumatic brain injury: a potential mechanistic link between head trauma and disorders of the hippocampus. J Neurosci. 1992 Dec 1;12(12):4846–53. doi:10.1523/JNEUROSCI.12-12-04846.1992

6. McIntosh, Saatman, Raghupathi, Graham, Smith, Lee, et al. The Dorothy Russell Memorial Lecture* The molecular and cellular sequelae of experimental traumatic brain injury: pathogenetic mechanisms. Neuropathology Appl Neurobio. 1998 Aug;24(4):251–67. doi:10.1046/j.1365-2990.1998.00121.x

7. Toth Z, Hollrigel GS, Gorcs T, Soltesz I. Instantaneous Perturbation of Dentate Interneuronal Networks by a Pressure Wave-Transient Delivered to the Neocortex. 1997.

8. Atkins CM. Decoding Hippocampal Signaling Deficits After Traumatic Brain Injury. Transl Stroke Res. 2011 Dec;2(4):546–55. doi:10.1007/s12975-011-0123-z

9. Hunt RF, Boychuk JA, Smith BN. Neural circuit mechanisms of post–traumatic epilepsy. Front Cell Neurosci. 2013;7. doi:10.3389/fncel.2013.00089

10. Paterno R, Folweiler KA, Cohen AS. Pathophysiology and Treatment of Memory Dysfunction After Traumatic Brain Injury. Curr Neurol Neurosci Rep. 2017 Jul;17(7):52. doi:10.1007/s11910-017-0762-x

11. Neuberger EJ, Gupta A, Subramanian D, Korgaonkar AA, Santhakumar V. Converging early responses to brain injury pave the road to epileptogenesis. Journal of

774 Neuroscience Research. 2019 Nov 1;97(11):1335–44. doi:10.1002/jnr.24202 PubMed
775 PMID: 29193309.

776 12. Li Y, Korgaonkar AA, Swietek B, Wang J, Elgammal FS, Elkabes S, et al. Toll-like receptor
777 4 enhancement of non-NMDA synaptic currents increases dentate excitability after
778 brain injury. Neurobiology of Disease. 2015 Feb 1;74:240–53.
779 doi:10.1016/j.nbd.2014.11.021 PubMed PMID: 25497689.

780 13. Korgaonkar AA, Li Y, Sekhar D, Subramanian D, Guevarra J, Swietek B, et al. Toll-like
781 Receptor 4 Signaling in Neurons Enhances Calcium-Permeable α-Amino-3-Hydroxy-5-
782 Methyl-4-Isoxazolepropionic Acid Receptor Currents and Drives Post-Traumatic
783 Epileptogenesis. Annals of Neurology. 2020 Apr 1;87(4):497–515.
784 doi:10.1002/ana.25698 PubMed PMID: 32031699.

785 14. Korgaonkar AA, Nguyen S, Li Y, Sekhar D, Subramanian D, Guevarra J, et al. Distinct
786 cellular mediators drive the Janus faces of toll-like receptor 4 regulation of network
787 excitability which impacts working memory performance after brain injury. Brain,
788 behavior, and immunity. 2020 Aug;88:381–95. doi:10.1016/j.bbi.2020.03.035 PubMed
789 PMID: 32259563.

790 15. Yang CC, Lin CC, Hsiao LD, Kuo JM, Tseng HC, Yang CM. Lipopolysaccharide-induced
791 matrix metalloproteinase-9 expression associated with cell migration in rat brain
792 astrocytes. International Journal of Molecular Sciences. 2020 Jan 1;21(1).
793 doi:10.3390/ijms21010259 PubMed PMID: 31905967.

794 16. Qiu J, Xu J, Zheng Y, Wei Y, Zhu X, Lo EH, et al. High-mobility group box 1 promotes
795 metalloproteinase-9 upregulation through toll-like receptor 4 after cerebral ischemia.
796 Stroke. 2010 Sep;41(9):2077–82. doi:10.1161/STROKEAHA.110.590463 PubMed PMID:
797 20671243.

798 17. Huntley GW. Synaptic circuit remodelling by matrix metalloproteinases in health and
799 disease. Nature Reviews Neuroscience. 2012 Nov;13(11):743–57.
800 doi:10.1038/nrn3320 PubMed PMID: 23047773.

801 18. Ethell IM, Ethell DW. Matrix metalloproteinases in brain development and remodeling:
802 synaptic functions and targets. Journal of neuroscience research. 2007
803 Oct;85(13):2813–23. doi:10.1002/jnr.21273 PubMed PMID: 17387691.

804 19. Szklarczyk A, Lapinska J, Rylski M, Mckay RDG, Kaczmarek L. Matrix Metalloproteinase-
805 9 Undergoes Expression and Activation during Dendritic Remodeling in Adult
806 Hippocampus. 2002.

807 20. Dityatev A, Schachner M. Extracellular matrix molecules and synaptic plasticity. Nature
808 reviews Neuroscience. 2003 Jun;4(6):456–68. doi:10.1038/nrn1115 PubMed PMID:
809 12778118.

21. Minta K, Brinkmalm G, Al Nimer F, Thelin EP, Piehl F, Tullberg M, et al. Dynamics of
cerebrospinal fluid levels of matrix metalloproteinases in human traumatic brain
injury. Sci Rep. 2020 Oct 22;10(1):18075. doi:10.1038/s41598-020-75233-z

22. Wang X, Jung J, Asahi M, Chwang W, Russo L, Moskowitz MA, et al. Effects of Matrix
Metalloproteinase-9 Gene Knock-Out on Morphological and Motor Outcomes after
Traumatic Brain Injury. 2000.

23. Hayashi T, Kaneko Y, Yu S, Bae E, Stahl CE, Kawase T, et al. Quantitative analyses of
matrix metalloproteinase activity after traumatic brain injury in adult rats. Brain
Research. 2009 Jul;1280:172–7. doi:10.1016/j.brainres.2009.05.040

24. Grossetete M, Phelps J, Arko L, Yonas H, Rosenberg GA. ELEVATION OF MATRIX
METALLOPROTEINASES 3 AND 9 IN CEREBROSPINAL FLUID AND BLOOD IN PATIENTS
WITH SEVERE TRAUMATIC BRAIN INJURY. Neurosurgery. 2009 Oct;65(4):702–8.
doi:10.1227/01.NEU.0000351768.11363.48

25. Reinhard SM, Razak K, Ethell IM. A delicate balance: Role of MMP-9 in brain
development and pathophysiology of neurodevelopmental disorders. Frontiers in
Cellular Neuroscience. 2015 Jul 29;9(JULY). doi:10.3389/fncel.2015.00280

26. Vafadari B, Salamian A, Kaczmarek L. MMP-9 in translation: from molecule to brain
physiology, pathology, and therapy. Journal of Neurochemistry. 2016 Oct 1;139:91–
114. doi:10.1111/jnc.13415 PubMed PMID: 26525923.

27. Pijet B, Stefaniuk M, Kostrzewska-Ksiezyk A, Tsilibary PE, Tzinia A, Kaczmarek L.
Elevation of MMP-9 Levels Promotes Epileptogenesis After Traumatic Brain Injury.
Molecular Neurobiology. 2018 Dec 1;55(12):9294–306. doi:10.1007/s12035-018-1061-
5 PubMed PMID: 29667129.

28. Wilczynski GM, Konopacki FA, Wilczek E, Lasiecka Z, Gorlewicz A, Michaluk P, et al.
Important role of matrix metalloproteinase 9 in epileptogenesis. Journal of Cell
Biology. 2008 Mar 10;180(5):1021–35. doi:10.1083/jcb.200708213 PubMed PMID:
18332222.

29. Dityatev A, Fellin T. Extracellular matrix in plasticity and epileptogenesis. Neuron Glia
Biol. 2008 Aug;4(3):235–47. doi:10.1017/S1740925X09000118

30. Jia F, Pan Y hua, Mao Q, Liang Y min, Jiang J yao. Matrix metalloproteinase-9 expression
and protein levels after fluid percussion injury in rats: The effect of injury severity and
brain temperature. Journal of neurotrauma. 2010 Jun;27(6):1059–68.
doi:10.1089/neu.2009.1067 PubMed PMID: 20233042.

31. Corrubia L, Huang A, Nguyen S, Shiflett MW, Jones MV, Ewell LA, et al. Early deficits in
dentate circuit and behavioral pattern separation after concussive brain injury.
Experimental Neurology. 2023 Dec;370:114578. doi:10.1016/j.expneurol.2023.114578

32. Witgen BM, Lifshitz J, Smith ML, Schwarzbach E, Liang SL, Grady MS, et al. Regional
hippocampal alteration associated with cognitive deficit following experimental brain
injury: A systems, network and cellular evaluation. Neuroscience. 2005;133(1):1–15.
doi:10.1016/j.neuroscience.2005.01.052

33. Jia F, Yin YH, Gao GY, Wang Y, Cen L, Jiang JY. MMP-9 inhibitor SB-3CT attenuates
behavioral impairments and hippocampal loss after traumatic brain injury in rat.
Journal of Neurotrauma. 2014 Jul 1;31(13):1225–34. doi:10.1089/neu.2013.3230
PubMed PMID: 24661104.

34. Fumihiro J, Masaaki K, Toshiyuki T, Yoshihiko T, Takahiro K, Satoru A. Investigation of the
unique metabolic fate of ethyl (6R)-6-[N-(2-chloro-4-fluorophenyl)sulfamoyl]cyclohex-
1-ene-1-carboxylate (TAK-242) in rats and dogs using two types of 14C-labeled
compounds having different labeled positions. Arzneimittelforschung. 2011 Nov
27;61(08):458–71. doi:10.1055/s-0031-1296228

35. Hadass O, Tomlinson BN, Gooyit M, Chen S, Purdy JJ, Walker JM, et al. Selective
Inhibition of Matrix Metalloproteinase-9 Attenuates Secondary Damage Resulting from
Severe Traumatic Brain Injury. PLoS ONE. 2013 Oct 23;8(10).
doi:10.1371/journal.pone.0076904 PubMed PMID: 24194849.

36. Gupta A, Elgammal FS, Proddutur A, Shah S, Santhakumar V. Decrease in Tonic
Inhibition Contributes to Increase in Dentate Semilunar Granule Cell Excitability after
Brain Injury. J Neurosci. 2012 Feb 15;32(7):2523–37. doi:10.1523/JNEUROSCI.4141-
11.2012

37. Hirschfeld M, Ma Y, Weis JH, Vogel SN, Weis JJ. Cutting edge: repurification of
lipopolysaccharide eliminates signaling through both human and murine toll-like
receptor 2. J Immunol. 2000 Jul 15;165(2):618–22. doi:10.4049/jimmunol.165.2.618
PubMed PMID: 10878331.

38. Cooke SF, Wu J, Plattner F, Errington M, Rowan M, Peters M, et al. Autophosphorylation
of αCaMKII is not a general requirement for NMDA receptor-dependent LTP in the adult
mouse. Journal of Physiology. 2006 Aug;574(3):805–18.
doi:10.1113/jphysiol.2006.111559 PubMed PMID: 16728448.

39. Schott AL, Esfahany KN, Grocott JM, Farrell JS. Toothy: an interactive platform for
dentate spike curation. bioRxiv. 2025 Oct 3;2025.10.03.680300.
doi:10.1101/2025.10.03.680300 PubMed PMID: 41256680; PubMed Central PMCID:
PMC12621741.

40. Gawlak M, Górkiewicz T, Gorlewicz A, Konopacki FA, Kaczmarek L, Wilczynski GM. High
resolution in situ zymography reveals matrix metalloproteinase activity at
glutamatergic synapses. Neuroscience. 2009 Jan 12;158(1):167–76.
doi:10.1016/j.neuroscience.2008.05.045 PubMed PMID: 18588950.

41. Proddutur A, Nguyen S, Yeh CW, Gupta A, Santhakumar V. Reclusive chandeliers:
Functional isolation of dentate axo-axonic cells after experimental status epilepticus.
Progress in Neurobiology. 2023 Dec;231:102542.
doi:10.1016/j.pneurobio.2023.102542

42. Rodríguez Peris L, Scheuber MI, Shan H, Braun M, Schwab ME. Barnes maze test for
spatial memory: A new, sensitive scoring system for mouse search strategies.
Behavioural Brain Research. 2024 Feb;458:114730. doi:10.1016/j.bbr.2023.114730

43. Harrison FE, Reiserer RS, Tomarken AJ, McDonald MP. Spatial and nonspatial escape
strategies in the Barnes maze. Learn Mem. 2006 Nov;13(6):809–19.
doi:10.1101/lm.334306

44. Hunt RF, Scheff SW, Smith BN. Synaptic Reorganization of Inhibitory Hilar Interneuron
Circuitry after Traumatic Brain Injury in Mice. J Neurosci. 2011 May 4;31(18):6880–90.
doi:10.1523/JNEUROSCI.0032-11.2011

45. Gupta A, Dovek L, Proddutur A, Elgammal FS, Santhakumar V. Long-Term Effects of
Moderate Concussive Brain Injury During Adolescence on Synaptic and Tonic GABA
Currents in Dentate Granule Cells and Semilunar Granule Cells. Front Neurosci.
2022;16:800733. doi:10.3389/fnins.2022.800733 PubMed PMID: 35360164; PubMed
Central PMCID: PMC8964009.

46. Santhakumar V, Ratzliff ADH, Jeng J, Toth Z, Soltesz I. Long-term hyperexcitability in the
hippocampus after experimental head trauma. Annals of Neurology. 2001
Dec;50(6):708–17. doi:10.1002/ana.1230

47. Santhakumar V, Bender R, Frotscher M, Ross ST, Hollrigel GS, Toth Z, et al. Granule cell
hyperexcitability in the early post-traumatic rat dentate gyrus: the 'irritable mossy cell'
hypothesis. The Journal of Physiology. 2000 Apr;524(1):117–34. doi:10.1111/j.1469-
7793.2000.00117.x

48. Michaluk P, Wawrzyniak M, Alot P, Szczot M, Wyrembek P, Mercik K, et al. Influence of
matrix metalloproteinase MMP-9 on dendritic spine morphology. J Cell Sci. 2011 Oct
1;124(Pt 19):3369–80. doi:10.1242/jcs.090852 PubMed PMID: 21896646.

49. Wiera G, Szczot M, Wojtowicz T, Lebida K, Koza P, Mozrzymas JW. Impact of matrix
metalloproteinase-9 overexpression on synaptic excitatory transmission and its
plasticity in rat CA3-CA1 hippocampal pathway. J Physiol Pharmacol. 2015
Apr;66(2):309–15. PubMed PMID: 25903961.

50. Nagy V, Bozdagi O, Matynia A, Balcerzyk M, Okulski P, Dzwonek J, et al. Matrix
metalloproteinase-9 is required for hippocampal late-phase long-term potentiation
and memory. Journal of Neuroscience. 2006 Feb 15;26(7):1923–34.
doi:10.1523/JNEUROSCI.4359-05.2006 PubMed PMID: 16481424.

51. Bozdagi O, Nagy V, Kwei KT, Huntley GW. In vivo roles for matrix metalloproteinase-9 in
mature hippocampal synaptic physiology and plasticity. Journal of Neurophysiology.
2007 Jul;98(1):334–44. doi:10.1152/jn.00202.2007 PubMed PMID: 17493927.

52. Wiera G, Mozrzymas JW. Extracellular Metalloproteinases in the Plasticity of Excitatory
and Inhibitory Synapses. Cells. 2021 Aug 11;10(8). doi:10.3390/cells10082055
PubMed PMID: 34440823.

53. Bragin A, Jand& G, Nadasdy Z, Van Landeghem M, Buzsaki G. Dentate EEG Spikes and
Associated Interneuronal Population Bursts in the Hippocampal Hilar Region of the
Rat. JOURNALOF NEUROPHYSIOLOGY. 1995.

54. Penttonen M, Kamondi A, Sik A, Acsády L, Buzsáki G. Feed-forward and feed-back
activation of the dentate gyrus in vivo during dentate spikes and sharp wave bursts.
Hippocampus. 1997;7(4):437–50. doi:10.1002/(SICI)1098-
1063(1997)7:4%3C437::AID-HIPO9%3E3.0.CO;2-F PubMed PMID: 9287083.

55. Lensu S, Waselius T, Penttonen M, Nokia MS. Dentate spikes and learning: disrupting
hippocampal function during memory consolidation can improve pattern separation. J
Neurophysiol. 2019;121:131–9. doi:10.1152/jn.00696.2018.-Hippocampal

56. Farrell JS, Soltesz I. Noncanonical circuits, states, and computations of the
hippocampus. Science. 2025 Sep 11;389(6765):eadv4420.
doi:10.1126/science.adv4420

57. Nokia MS, Gureviciene I, Waselius T, Tanila H, Penttonen M. Hippocampal electrical
stimulation disrupts associative learning when targeted at dentate spikes. Journal of
Physiology. 2017 Jul 15;595(14):4961–71. doi:10.1113/JP274023 PubMed PMID:
28426128.

58. Farrell JS, Hwaun E, Dudok B, Soltesz I. Neural and behavioural state switching during
hippocampal dentate spikes. Nature. 2024 Apr 18;628(8008):590–5.
doi:10.1038/s41586-024-07192-8

59. Bott JB, Muller MA, Jackson J, Aubert J, Cassel JC, Mathis C, et al. Spatial reference
memory is associated with modulation of theta-gamma coupling in the dentate Gyrus.
Cerebral Cortex. 2016 Sep 1;26(9):3744–53. doi:10.1093/cercor/bhv177 PubMed
PMID: 26250776.

60. Dovek L, Ahmadi M, Marrero K, Zagha E, Santhakumar V. Cellular and circuit features
distinguish dentate gyrus semilunar granule cells and granule cells activated during
contextual memory formation [Internet]. 2025 Jul 17. doi:10.7554/eLife.101428.2

61. Maroso M, Balosso S, Ravizza T, Liu J, Aronica E, Iyer AM, et al. Toll-like receptor 4 and
high-mobility group box-1 are involved in ictogenesis and can be targeted to reduce

seizures. Nature Medicine. 2010 Apr;16(4):413–9. doi:10.1038/nm.2127 PubMed
PMID: 20348922.

62. Kleen JK, Holmes GL. Taming TLR4 may ease seizures. Nature Medicine. 2010
Apr;16(4):369–70. doi:10.1038/nm0410-369 PubMed PMID: 20376038.

63. Vezzani A, Balosso S, Ravizza T. Neuroinflammatory pathways as treatment targets and
biomarkers in epilepsy. Nat Rev Neurol. 2019 Aug;15(8):459–72. doi:10.1038/s41582-
019-0217-x

64. Okun E, Griffioen KJ, Mattson MP. Toll-like receptor signaling in neural plasticity and
disease. Trends in Neurosciences. 2011 May;34(5):269–81.
doi:10.1016/j.tins.2011.02.005 PubMed PMID: 21419501.

65. Ahmad A, Crupi R, Campolo M, Genovese T, Esposito E, Cuzzocrea S. Absence of TLR4
Reduces Neurovascular Unit and Secondary Inflammatory Process after Traumatic
Brain Injury in Mice. Dhandapani KM, editor. PLoS ONE. 2013 Mar 28;8(3):e57208.
doi:10.1371/journal.pone.0057208

66. Xiong G, Metheny H, Johnson BN, Cohen AS. A comparison of different slicing planes in
preservation of major hippocampal pathway fibers in the mouse. Frontiers in
Neuroanatomy. 2017 Nov 21;11. doi:10.3389/fnana.2017.00107

67. Howard AL, Neu A, Morgan RJ, Echegoyen JC, Soltesz I. Opposing Modifications in
Intrinsic Currents and Synaptic Inputs in Post-Traumatic Mossy Cells: Evidence for
Single-Cell Homeostasis in a Hyperexcitable Network. Journal of Neurophysiology.
2007 Mar;97(3):2394–409. doi:10.1152/jn.00509.2006

68. Dziembowska M, Milek J, Janusz A, Rejmak E, Romanowska E, Gorkiewicz T, et al.
Activity-Dependent Local Translation of Matrix Metalloproteinase-9. J Neurosci. 2012
Oct 17;32(42):14538–47. doi:10.1523/JNEUROSCI.6028-11.2012

69. Dziembowska M, Wlodarczyk J. MMP9: A novel function in synaptic plasticity. The
International Journal of Biochemistry & Cell Biology. 2012 May;44(5):709–13.
doi:10.1016/j.biocel.2012.01.023

70. Stefaniuk M, Beroun A, Lebitko T, Markina O, Leski S, Meyza K, et al. Matrix
Metalloproteinase-9 and Synaptic Plasticity in the Central Amygdala in Control of
Alcohol-Seeking Behavior. Biological Psychiatry. 2017 Jun;81(11):907–17.
doi:10.1016/j.biopsych.2016.12.026

71. Szepesi Z, Hosy E, Ruszczycki B, Bijata M, Pyskaty M, Bikbaev A, et al. Synaptically
released matrix metalloproteinase activity in control of structural plasticity and the
cell surface distribution of GluA1-AMPA receptors. PLoS ONE. 2014 May 22;9(5).
doi:10.1371/journal.pone.0098274 PubMed PMID: 24853857.

72. Fanselow EE, Kubota Y, Wehr M, Yavorska I. Somatostatin-Expressing Inhibitory
Interneurons in Cortical Circuits. Frontiers in Neural Circuits | www.frontiersin.org.
2016;10:76. doi:10.3389/fncir.2016.00076

73. Pirbhoy PS, Rais M, Lovelace JW, Woodard W, Razak KA, Binder DK, et al. Acute
pharmacological inhibition of matrix metalloproteinase-9 activity during development
restores perineuronal net formation and normalizes auditory processing in Fmr1 KO
mice. Journal of Neurochemistry. 2020 Dec;155(5):538–58. doi:10.1111/jnc.15037

74. Sun DA, Deshpande LS, Sombati S, Baranova A, Wilson MS, Hamm RJ, et al. Traumatic
brain injury causes a long-lasting calcium (Ca2+)-plateau of elevated intracellular Ca
levels and altered Ca2+homeostatic mechanisms in hippocampal neurons surviving
brain injury. Eur J of Neuroscience. 2008 Apr;27(7):1659–72. doi:10.1111/j.1460-
9568.2008.06156.x

75. Weber JT. Altered Calcium Signaling Following Traumatic Brain Injury. Front Pharmacol.
2012;3. doi:10.3389/fphar.2012.00060

76. Hellstrom IC, Danik M, Luheshi GN, Williams S. Chronic LPS exposure produces
changes in intrinsic membrane properties and a sustained IL-β-dependent increase in
GABAergic inhibition in hippocampal CA1 pyramidal neurons. Hippocampus.
2005;15(5):656–64. doi:10.1002/hipo.20086

77. Hainmueller T, Bartos M. Dentate gyrus circuits for encoding, retrieval and
discrimination of episodic memories. Nature Reviews Neuroscience. 2020 Mar
1;21(3):153–68. doi:10.1038/s41583-019-0260-z PubMed PMID: 32042144.

78. Hsu D. The dentate gyrus as a filter or gate: a look back and a look ahead. Prog Brain
Res. 2007;163:601–13. doi:10.1016/S0079-6123(07)63032-5 PubMed PMID:
17765740.

79. Ewell LA. To sleep perchance to spike: a functional role for dentate spikes in memory.
Journal of Physiology. 2017 Jul 15;595(14):4565. doi:10.1113/JP274502 PubMed PMID:
28485489.

80. Farrell JS, Hwaun E, Dudok B, Soltesz I. Neural and behavioural state switching during
hippocampal dentate spikes [Internet]. doi:10.1038/s41586-024-07192-8

81. Flynn SP, Barrier S, Scott RC, Lenck-Santini PP, Holmes GL. Status epilepticus induced
spontaneous dentate gyrus spikes: In vivo current source density analysis. PLoS ONE.
2015 Jul 6;10(7). doi:10.1371/journal.pone.0132630 PubMed PMID: 26148195.

82. Engel J, Bragin A, Staba R. Nonictal EEG biomarkers for diagnosis and treatment.
Epilepsia Open. 2018 Dec 1;3(S2):120–6. doi:10.1002/epi4.12233

83. Perucca P, Smith G, Santana-Gomez C, Bragin A, Staba R. Electrophysiological
biomarkers of epileptogenicity after traumatic brain injury. Neurobiology of Disease.
2019 Mar;123:69–74. doi:10.1016/j.nbd.2018.06.002

84. Tewari BP, Chaunsali L, Campbell SL, Patel DC, Goode AE, Sontheimer H. Perineuronal
nets decrease membrane capacitance of peritumoral fast spiking interneurons in a
model of epilepsy. Nature Communications. 2018 Dec 1;9(1). doi:10.1038/s41467-
018-07113-0 PubMed PMID: 30413686.

85. Whitebirch AC, LaFrancois JJ, Jain S, Leary P, Santoro B, Siegelbaum SA, et al. Enhanced
excitability of the hippocampal CA2 region and its contribution to seizure activity in a
mouse model of temporal lobe epilepsy. Neuron. 2022 Oct;110(19):3121-3138.e8.
doi:10.1016/j.neuron.2022.07.020

86. Zhou Y, Chen Y, Xu C, Zhang H, Lin C. TLR4 Targeting as a Promising Therapeutic
Strategy for Alzheimer Disease Treatment. Front Neurosci. 2020 Dec 18;14:602508.
doi:10.3389/fnins.2020.602508

87. Iori V, Iyer AM, Ravizza T, Beltrame L, Paracchini L, Marchini S, et al. Blockade of the IL-
1R1/TLR4 pathway mediates disease-modification therapeutic effects in a model of
acquired epilepsy. Neurobiology of Disease. 2017 Mar;99:12–23.
doi:10.1016/j.nbd.2016.12.007

1041

1042

Figure legends:

Figure 1: TLR4 signaling drives early excitatory and inhibitory synaptic changes in

DGCs after brain injury.   A) Schematic of the experimental question. B) Experimental

timeline for examining synaptic changes after FPI in rats. C) Representative sEPSC traces

showing whole-cell voltage clamp recordings from granule cells in vehicle treated sham

controls (black), vehicle treated FPI (red), TLR4 inhibited sham (grey) and TLR4 inhibited FPI

(blue). D-E) Cumulative probability plots of sEPSC interevent intervals (D) and amplitude

(E) in vehicle treated and TLR4 inhibited sham and injured rats. F) Representative sIPSC

traces showing whole-cell voltage clamp recordings from granule cells in vehicle treated

sham controls (black), vehicle treated FPI (red), TLR4 inhibited sham (grey) and TLR4

inhibited FPI (blue). G-H) Cumulative probability plots of sIPSC interevent intervals (G) and

amplitude (H) in vehicle treated and TLR4 inhibited sham and injured rats. I) Experimental

timeline for examining synaptic changes after FPI in TLR4$^{fl/fl}$: CaMKIIa- CreERT2 mice. J)

Representative sEPSC traces from granule cells in Cre^{-ve} sham controls (black), Cre^{-ve} FPI

(red), Cre^{+ve} sham (grey) and Cre^{+ve} FPI (blue). K-L) Cumulative probability plots of sEPSC

interevent intervals (K) and amplitude (L) in granule cells from Cre^{-ve} and Cre^{+ve} mice. n = 8-9

cells from 3 rats/group (A-H) and 7 cells from 3 mice/group (I-L). * indicates *P*<0.05 by K-S

test with Holm-Šídák correction.

Figure 2: TLR4 signaling upregulates MMP-9 activity in the dentate gyrus after brain

injury.   A) Experimental timeline for in-situ zymography to quantify MMP-9 activity after

brain injury in rat. B) Representative images showing MMP-9 activity in hippocampal

sections from sham and injured rats. Images from vehicle treated (middle) and TLR

inhibited mice (bottom) show MMP-9 activity in green. C) Quantification of MMP-9 activity,

normalized to CA1 stratum Radiatum, shows an increase in MMP-9 activity in vehicle

treated injured rats, but not in TAK-242 treated injured rats. Note that MMP-9 activity in

vehicle treated and TAK-242 treated sham rats were not different. D) Experimental timeline

for in-situ zymography to quantify MMP-9 activity after brain injury in mice. E)

Representative images showing MMP-9 activity in sham and injured mice. Images from

hippocampal sections from Cre^{-ve} sham and injured mice (top) and Cre^{+ve} sham and injured

mice (bottom) show MMP-9 activity in green .F) Quantification of MMP-9 activity,

normalized to CA1 stratum Radiatum, shows an increase in MMP-9 activity after brain

injury in Cre^{-ve} mice, but not in Cre^{+ve} mice. Note that MMP-9 activity in sham mice was not

different between genotypes. * indicates $P<0.05$, ** indicates $P<0.01$, *** indicates

$P<0.001$, **** indicates $P<0.0001$ by TW-ANOVA with Šídáks multiple comparisons post

hoc tests. G) Experimental timeline for RNAscope experiments in naïve rats. H)

Representative images showing increase in MMP-9 mRNA in sections from rats after LPS-

B5 treatment. I) Summary plot of mRNA levels in saline, LPS-B5 treated and LPS-B5+JSH-

23 sections shows an increase in mRNA levels after LPS-B5 treatment which is abolished

by blocking NF-κB. GCL- granule cell layer. n = 5-6 rats/group (A-C), 3-4 mice /genotype (D-

F), and 5 rats/group (G-I). * indicates $P<0.05$, ** indicates $P<0.01$, by OW-ANOVA with

Šídáks multiple comparisons post hoc tests. Data points in panel C, F and I represent

individual animals.

Figure 3: Blocking MMP-9 prevents early synaptic changes in excitatory inputs to DGCs

after brain injury. A) Experimental timeline for examining synaptic changes in acute rat

slices after LPS incubation. B) Representative sEPSC traces from granule cells in LPS

incubated (orange), LPS+SB-3CT co-incubated (magenta) slices. C-D) Cumulative

probability plots of sEPSC interevent intervals (C) and amplitude (D) between LPS-B5

incubated and LPS-B5+SB-3CT co-incubated slices. Note that LPS incubation significantly

reduced interevent intervals. Co-incubation of slices in MMP-9 blocker SB-3CT prevented

this change. E) Experimental timeline for examining synaptic changes in SB-3CT treated

sham/injured rats. F) Representative granule cell sEPSC traces showing from SB-3CT

treated sham (grey) and injured rats (green). G-H) Cumulative probability plots of sEPSC

interevent intervals (G) and amplitude (H) in SB-3CT treated sham/injured rats. I)

Experimental timeline for examining synaptic changes in SB-3CT treated Cre^{-ve} mice after

brain injury. J) Representative granule cell sEPSC traces from vehicle treated (red) and SB-

3CT treated mice(green). K-L) Cumulative probability plots of sEPSC interevent intervals (K)

and amplitude (L) in vehicle treated and SB-3CT treated Cre^{-ve} mice. n = 8-9 cells(A-D), 7-10

cells (E-H) from 3-4 rats/group and 7 cells from 3 mice /group (I-L). * indicates $P<0.05$ by K-

S test with Holm-Šídák correction.

Figure 4: TLR4 regulation of MMP-9 activity enhances dentate network excitability after

brain injury. A) Experimental timeline for *in vivo* experiments examining dentate excitability

and throughput. B) Representative traces showing DG population spike from sham and

injured animals in response to 800 µA stimulus (bold lines) and 1200 µA (dotted lines). C)

Input-output curve illustrating early saturation of population spike amplitude in injured

animals (red) compared to sham controls. D) Inhibition of TLR4 or MMP-9 effectively

mitigates these changes. E) MMP-9 inhibition (light grey) did not affect input-output curves

in sham animals, TLR4 inhibition increased network excitability of sham animals (grey).

Sham and FPI data from panel C are included as faded traces in D (FPI) and E (Sham) for

comparison. F) Representative trace from 16 channel silicon probe recording local field

potentials from cortex to hippocampus. Current-source density plot shows the sink and

source of an individual dentate spike. inset- a single dentate spike recorded in the GC layer.

G) Quantification of dentate spikes shows a significant increase in frequency in injured

animals. Inhibition of TLR4 or MMP-9 effectively reduced changes in dentate spike

frequency in injured animals. n = 5-6 rats/group (A-E) and 4-5 rats/ group (F-G). *

indicates<0.05, **** indicates P<0.0001   by TW-ANOVA with Šídáks multiple comparisons

post hoc tests. Data points in panels G represent individual animals.

1119

Figure 5: Posttraumatic deficits in long-term potentiation are improved by early

inhibition of TLR4 or MMP-9. A) Experimental timeline for *in vivo* assessment of long-term

potentiation in perforant path synapses to DGCs. B) Representative traces from Sham

(black) and FPI rats (red) at 10 minutes before (a) and 120 minutes after (b) LTP induction.

C) Injured rats exhibit a significant decrease in fEPSP slope values after 60 minutes(red)

compared to sham controls (black). D) Inhibition of TLR4 (blue) or MMP-9 (green)

significantly improved LTP in injured animals. E) MMP-9 inhibition did not affect LTP in

SHAM controls. TLR4 inhibition reduced LTP at 60 min and 180 min after induction. Note

that Sham and FPI data from panel C are included as faded traces in D (FPI) and E (Sham)

for comparison. n = 5-6 rats/group * indicates P <0.05 by TW-RM ANOVA.

Figure 6: Early inhibition of TLR4 or MMP-9 improves spatial learning after brain injury.

A) Experimental timeline for testing spatial learning using Barnes maze. B) Representative

heat plot showing rats trajectory during the task on day 2. C-E) Injured animals showed

deficits in latency to escape, distance travelled, and search strategy on day 2. * indicates

P<0.05 by RM-ANOVA. F-H) Summary plots of effect of TLR4 or MMP-9 inhibition on latency

to escape (F), distance travelled (G), and search strategy (H) on day 2. I) Injured animals

showed increased thigmotaxis on day 2 which was attenuated with TLR4 or MMP-9

inhibition. n = 6-10 rats/group. * indicates P<0.05, ** indicates *P*<0.01, *** indicates

P<0.001, **** indicates *P*<0.0001 by TW-ANOVA with Šídáks multiple comparisons post

hoc tests. Data points in panels F-I represent individual animals.

1140

1141

Figure 7: Graphical Summary of TLR4 regulation of dentate gyrus function in control

and injured brain.

In uninjured brain, pharmacological inhibition of TLR4 signaling increases the frequency of

both excitatory and inhibitory currents and enhances network excitability and impairs long-

term potentiation without altering MMP-9 activity. Genetic deletion of neuronal TLR4 alters

intrinsic properties without impacting synaptic physiology. Together these data are

consistent with non-neuronal TLR4 signaling contributing to basal synaptic homeostasis in

line with prior ex vivo studies [14]. In the injured brain, augmented neuronal TLR4 signaling

acts through MMP-9 to enhance glutamatergic signaling, network excitability and dentate

spikes and impairs LTP maintenance. Mechanistically, inhibitory changes are driven by

TLR4 in an MMP-9 independent manner. Together, these findings demonstrate that TLR4

signaling regulates DG function and seizure susceptibility [13] through distinct, context-

dependent cellular mechanisms in the healthy and injured brain.† refers to [14] ‡ refers to

[13].

1156

1157

A Wistar rat — P24

time post-injury: 30min, 8hrs, 24hrs → 2 DPI

Sham/FPI — DMSO (or) TAK-242 → in situ zymography

B MMP-9

SHAM_vehicle | FPI_vehicle
SHAM_TAK-242 | FPI_TAK-242

500 μm — TLR4 inhibition

C Molecular Layer

Normalized MMP-9 activity (%)

Vehicle | TAK-242

** **

■ Sham ■ FPI

D TLR4^fl/fl-CaMKIIa-CreERT2 mice

tamoxifen induction: -24hrs, 30min, 24hrs → 2 DPI

Sham/FPI → in situ zymography

E MMP-9

SHAM Cre^-ve | FPI Cre^-ve
SHAM Cre^+ve | FPI Cre^+ve

500 μm

F Molecular Layer

Normalized MMP-9 activity (%)

Cre -ve | Cre +ve

**** *** ns

○ Sham ○ FPI

G P30 (Wistar rats)

+ 1hr → + 6hrs

Saline/JSH-23 → Saline/LPS-B5 → Transcardial perfusion → RNAscope

H Saline | DAPI | MMP-9 | LPS-B5 | LPS-B5 + JSH-23

100 μm

I MMP-9 mRNA in GCL (a.u / μm²)

Saline | LPS-B5 | LPS+JSH-23

ns * **

A Wistar rat · Sham · P24 · 7 DPI · Slice incubation · aCSF (or) LPS-B5 (or) LPS-B5 + SB-3CT · + 90 min · Patch-Clamp recordings

B SHAM$_{LPS}$ · SHAM$_{LPS+SB-3CT}$ · 20 pA · 0.2 s

C sEPSC IEI · Cumulative Probability · SHAM vs SHAM$_{LPS}$ * · SHAM vs SHAM$_{LPS+SB-3CT}$ - ns · Interevent Interval (msec)

D sEPSC amplitude · Cumulative Probability · SHAM vs SHAM$_{LPS}$ * · SHAM$_{LPS}$ vs SHAM$_{LPS+SB-3CT}$ * · Amplitude (pA)

E Wistar rat · Sham/FPI · P24 · time post-injury · 30min · 8hrs · 24hrs · 7 DPI · DMSO (or) SB-3CT · Patch-Clamp recordings

F SHAM$_{SB-3CT}$ · FPI$_{SB-3CT}$ · 20 pA · 0.2 s

G sEPSC IEI · Cumulative Probability · SHAM$_{DMSO}$ vs SHAM$_{SB-3CT}$ ns · FPI$_{DMSO}$ vs FPI$_{SB-3CT}$ * · Interevent Interval (msec)

H sEPSC amplitude · Cumulative Probability · SHAM$_{DMSO}$ vs SHAM$_{SB-3CT}$ ns · FPI$_{DMSO}$ vs FPI$_{SB-3CT}$ * · Amplitude (pA)

I tamoxifen induction · -24hrs · 30min · 24hrs · 7 DPI · TLR4$^{fl/fl}$-CaMKIIa-CreERT2 mice · Sham/FPI · + 30min · DMSO (or) SB-3CT · Patch Clamp recordings

J Cre -ve FPI$_{DMSO}$ · Cre -ve FPI$_{SB-3CT}$ · 20 pA · 0.1 s

K sEPSC IEI · Cumulative Probability · Cre -ve SHAM vs Cre -ve FPI * · Cre -ve FPI vs Cre -ve FPI$_{SB-3CT}$ * · Interevent Interval (sec)

L sEPSC amplitude · Cumulative Probability · Cre -ve SHAM vs Cre -ve FPI * · Cre -ve FPI vs Cre -ve FPI$_{SB-3CT}$ * · Amplitude (pA)