Sex-stratified insights into the genetics of brain volumes in late adulthood

Édith Breton[a,b,c,*], Elleke Tissink[a], Gloria Matte Bon[d,e], Dennis van der Meer[a], Ole A. Andreassen[a], Tobias Kaufmann[a,d,f,*]

[a] Centre for Precision Psychiatry, Division of Mental Health and Addiction, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

[b] Centre intersectoriel en santé durable, département des sciences fondamentales, Université du Québec à Chicoutimi, Saguenay, Canada

[c] Centre intégré universitaire de santé et de services sociaux (CIUSSS) du Saguenay-Lac-Saint-Jean, Saguenay, QC, Canada

[d] Department of Psychiatry and Psychotherapy, Tübingen Center for Mental Health, University of Tübingen, Tübingen, Germany

[e] Department of Women's and Children's Health, Science for Life Laboratory, Uppsala University, Uppsala, Sweden

[f] German Center for Mental Health (DZPG), partner site Tübingen, Tübingen, Germany

* Correspondence:

Édith Breton, PhD. Département des sciences fondamentales, Université du Québec à Chicoutimi, 555 Bd de l'Université, G7H 2B1, Chicoutimi, QC, Canada. Email: ebreton@uqac.ca

Tobias Kaufmann, PhD. Department of Psychiatry and Psychotherapy, Tübingen Center for Mental Health, University of Tübingen, Calwerstr. 14, 72076 Tübingen, Germany. Email: Tobias.Kaufmann@med.uni-tuebingen.de

Abstract

Background. The prevalence, timing and disease course of mental and neurological disorders vary according to sex, yet the neurogenetic mechanisms underlying sex differences in these disorders remain poorly understood. *Methods.* Here, we explored the genetic architecture of the anatomical volumes of 257 regions of interest across the brain using multivariate genome-wide analyses in 15,740 male and 15,740 female participants from the UK Biobank, matched on age and scan site. *Results.* Our findings revealed that the genetics of brain volume is highly similar between females and males in late adulthood. Yet, we found evidence of possible autosomal sex heterogeneity, particularly in the number of brain-volume associated genes, which was higher in females than in males. Variability in the number of identified genes were marked in limbic regions such as the insula, the cingulate cortex, the hippocampus and the amygdala. *Conclusion.* Overall, our findings contribute to a better understanding of the genetic determinants of brain volumes in males and females. Because neurogenetics may also influence the risk for sex-prevalent brain disorders, the current findings have the potential to facilitate precision medicine approaches in improving prevention strategies and targeted treatments.

Plain English summary

Mental and neurobiological disorders often differ between females and males. Some disorders are more common in one sex than the other, and symptoms can present or evolve differently. Variation in brain biology, partly shaped by genetic factors, may contribute to these observed patterns. In this study, we investigated genetic factors linked to brain structure in females and males separately. We focused on the volumes of 257 brain regions and analyzed genetic data from more than 30,000 adults from the UK Biobank. Overall, we found that the patterns of

association between genetics and brain volumes are largely similar between females and males in

late adulthood. At the same time, we observed evidence of sex-dependent patterns, particularly in

the number of genes associated with brain volume, which tended to be higher in females than in

males. These differences were marked in brain regions involved in the limbic system. Together,

our findings improve our understanding of how genetics contribute to brain structure in females

and males.

Highlights

• Genetic influences on brain volumes are largely shared between females and males in late
adulthood.

• Gene-level variability was observed, with a higher number of brain volumes associated
genes identified in females than in male, in both multivariate and univariate analyses.

• This gene-level variability was most pronounced in limbic regions, such as the insular
and cingulate cortices.

Key words

63 Sex Differences; Brain Anatomy; Imaging Genetics

Background

Sex impacts the prevalence, clinical presentation, course and treatment response of several

mental and neurological disorders (1,2). Major depressive disorder and Alzheimer's disease are

twice as common in females as in males, and anorexia nervosa is 10 times more common in

females than in males (3–5). In contrast, autism spectrum disorder, alcohol use disorder and

attention deficit hyperactivity disorder are two to three times more common in males (6–8). The

main windows of vulnerability for mental disorders also differ by sex, with females more likely

to be affected by conditions that begin in adolescence or later, and males more likely to develop

neurodevelopmental disorders that express in childhood. Yet, the biological mechanisms

underlying these differences have not been elucidated (9,10). Such mechanistic understanding

could facilitate efforts towards precision medicine in the clinical neurosciences and contribute to

reducing sex inequalities in health care and research (11,12).

Neuroimaging studies have identified structural and functional brain differences between

males and females across the lifespan, including greater variability in brain structure in males

(13). There is evidence to suggest that these brain differences may be associated with mental

disorders (14), but more studies are needed to establish this association. Sex chromosomes and

hormones are obvious contributors to the sex differences in the brain (15), but they are not the

only ones. For example, genetic studies have identified sex-dependent associations between

autosomal gene variants and mood or psychotic disorders (16). There is also evidence that

autosomal gene variants interact with sex hormones to differentially affect the risk of conditions

such as dementia (17).

Yet, there is still little understanding of the interaction between sex, genetics and the

brain, and few large-scale studies have considered study designs that analyze females and males

separately (12). Existing evidence suggests that common genetic variants often impact gene

expression differently in the brains of males and females (18), which may be relevant for

disorder risk (19). Further studies have suggested that genes involved in neurotransmission,

immune function and mental disorders are differently targeted by transcriptional regulators in

both sexes (20). Throughout the life course, epigenetic and other regulatory factors,

inflammatory processes, neurodevelopmental processes and ageing all plausibly contribute to

shaping the genetics-brain interaction differently in males and females, potentially influencing

their vulnerability to common brain disorders.

Here, we investigated the genetic architecture of brain volume in 15,740 females and

15,740 males from the UK Biobank. Both groups were matched for age and scan site. Our

overarching aim was to gain further insight into how genetics are linked to brain structure in

females and males. We opted for sex-stratified analyses to characterize whether the overall

patterns of autosomal genetic associations observed in one sex also show up in the other. To this

end, we extracted volumes of 180 cortical and 77 subcortical regions across the entire brain for

genome-wide association studies (GWAS). We leveraged MOSTest (21), a multivariate GWAS

tool designed to enhance power and facilitate discovery in imaging genetics research, yielding

one multivariate summary statistic across the whole brain for females and males separately. In

addition, we assessed each brain region in females and males separately using a univariate

approach.

Methods
106

Sample

We selected participants from the UK Biobank with genetic and neuroimaging data and of

European descent, defined by self-identification as "White British" and similar genetic ancestry

based on genetic principal components analysis (accession code no. 27412). Study procedures

were approved by local institutional review boards and each sample was collected with

participants' written informed consent. Standardized brain MRI and genomic protocols are

described elsewhere (22–24).

Magnetic Resonance Imaging

The present study used T1-weighted magnetic imaging volumes extracted using FreeSurfer

(https://surfer.nmr.mgh.harvard.edu/). The whole brain volume measure correspond to the mean

of the right and left hemisphere volumes, and included a combination of 180 cortical ROIs based

on the Glasser Atlas (25), as well as subcortical structures (10 ROIs (26)) and further

segmentation of the amygdala (9 nuclei + whole amygdala (27)), the hippocampus (19

subregions + whole hippocampus (28)), the thalamus (25 subregions + whole thalamus (29)), the

brainstem (4 subregions + whole brainstem (30)) and the hypothalamus (5 subregions + whole

hypothalamus (31)), for a total of 257 ROIs.

Individual structures with volumes exceeding three standard deviations below the mean

of the Euler number were excluded from the analyses. This left us with 34,706 participants of

European descent with genetics and neuroimaging data. From these 34,704 participants, we

selected male-female pairs, ensuring matching for scan site and age (within a 6-months

difference) while minimizing discrepancies in estimated total intracranial volume by selecting

the pairs with the smallest differences, based on a matching procedure deployed by others (32).

We made sure that there was no significant difference ($p>0.05$) between males and females for

age and total intracranial volume, for each scan site. Participants who did not meet these match

criteria were excluded, leading to the inclusion of 15,740 males and 15,740 females with a mean

age of 64,4 years.

Genome-wide analyses

We deployed the Multivariate Omnibus Statistical Test (MOSTest) (21) to perform two

multivariate genome-wide association analyses of the whole brain volume, in females and males

separately, focusing on autosomes. Individuals with genotype missing rate >10%, variants with

genotype missing rate > 5%, variants failing Hardy-Weinberg equilibrium at $p=1\times10^{-9}$, variants

with minor allele frequency below 0.5%, and imputation info score below 0.5 were filtered out.

We controlled our analyses for covariates by pre-residualizing the ROI volumes for age, age^2,

scanning site, the first 20 genetic principal components, Euler number, and estimated total

intracranial volume (in R version 4.2).

In parallel, we extracted univariate GWASs from the univariate stream of MOSTest (21)

for each of the 257 ROIs, to gain a better understanding of the regional heterogeneity that may

occur between males and females in specific brain regions.

FUMA

The Functional Mapping and Annotation of Genome-Wide Association Studies (FUMA)

platform (33) was used to perform positional gene mapping for all candidate SNPs, using the

default settings of the SNP2GENE functionality. For the list of overlapping genes between

females and males, FUMA GENE2FUNC was used to perform enrichment analyses against 15

496 gene sets (curated gene sets: 5 500, Gene Ontology terms: 9 996) from MsigDB version

v7.0. We corrected for multiple testing using FDR and set the minimum overlapping genes to 2.

Heritability & Genetic correlations

The sex-stratified univariate GWASs summary statistics were used to compute heritability and

genetic correlations estimates for each ROI between females and males, using linkage

disequilibrium score regression (LDSC) (34,35) with default settings and LD Scores based on the

European 1000 Genomes reference panel. We filtered out phenotypes with low heritability z-

scores (h^2_{SNP} / SE < 1.96) from further analyses.

Results

The genetics of brain volumes is similar in females and males, with some variability at the gene level

Using a multivariate GWAS framework, we identified 231 genome-wide significant loci ($p < 5e-8$) associated with brain volumes in females and 241 loci associated with brain volumes in males, including 127 physically overlapping loci between both sexes (based on chromosomes, loci start positions and end positions). Genome-wide patterns of association appeared highly similar between sexes (**Figure 1A**). Among the identified loci, 199 (female) and 190 (male) loci could positionally be mapped to genes, leading to the identification of 1000 genes associated with brain volumes in females and 849 genes associated with brain volumes in males (since some loci map to many genes; **Figure 1B**). Approximately half of the identified genes (N=552) overlapped between males and females, while 448 genes were only found in females and 297 genes were only found in males (**Figure 1C**, **Supplementary Table 1**).

Figure 1: Sex-stratified multivariate analyses reveal shared and distinct genetic contributions to brain volume

Legend: (A) Miami plot displaying the multivariate genome-wide association statistics in females (top/pink) and males (bottom/blue). These statistics reflect the association across 257 brain regions of interest. (B) Comparison of the number of genes, lead single nucleotide polymorphisms (SNPs) and risk loci identified in females (pink) versus males (blue) based on the multivariate GWAS statistics. The number of risk loci correspond to the number of loci with at least one gene mapped. (C) Venn diagram showing the corresponding number of genes overlapping or not in females (pink) and males (blue).

The genes shared between females and males were significantly ($p_{FDR} < 0.05$) enriched in 123 gene-sets (see Methods, **Supplementary Figure 1**). 25 of these were related to neurodevelopmental pathways (e.g., neuron development, neurogenesis, regulation of neuron projection), 15 were related to developmental/growth processes, and 15 were related to immune pathways. Overlapping genes between males and females were significantly ($p_{FDR} < 0.05$) enriched for differentially expressed gene-sets in 21 tissue types. These included 11 key brain

regions such as the putamen, nucleus accumbens, caudate nucleus, amygdala, anterior cingulate

cortex, hypothalamus and hippocampus, frontal cortex, and substantia nigra (**Supplementary**

Figure 2).

The 448 genes only significantly associated with brain volumes in females were enriched

in 23 tissue-specific gene-sets with differential expression, of which 13 were brain tissues

(**Supplementary Figure 3**). Most of these (N=19) overlapped with the enrichment pattern of

genes shared between females and males described above, except for cerebellum, cerebellar

hemisphere, tibial artery and EBV-transformed lymphocytes.

On the other hand, genes only found as significant in males (N=297) were enriched for 34

differentially expressed tissue-specific gene-sets. These included 12 brain tissues that were also

identified in genes only found for females or genes shared between males and females

(**Supplementary Figure 4**).

Divergences in number of genes identified in males versus females are widespread across

the brain

Next, we sought to characterize patterns of sex-stratified associations across distinct brain

regions. To do this, we turned to a univariate framework and performed standard GWASs for

each of the 257 regions of interest (ROI) separately. We further assigned cortical ROIs to their

respective lobes (i.e., frontal, parietal, temporal, occipital, insula, cingulate). We performed sex-

stratified LDSC (34,35) to estimate SNP-based heritability across the 257 brain regions in

females and males, yielding highly similar mean heritability estimates of $h^2=0.19 \pm 0.07$ in

females and $h^2=0.18 \pm 0.06$ in males (**Supplementary Figure 5**). As expected, the genetic

correlations across the genome were very high between the sexes in all cortical and subcortical

brain regions (based on LDSC; all r >0.72; **Figure 2A-B; 4**). However, at the gene level, these

regions showed potential sex-related heterogeneity. (**Figure 2C**).

Figure 2: The genetic architecture of brain volume shows global similarity between sexes, yet

exhibits divergent gene-level patterns

Legend: (A) Genetic correlations for each cortical ROI between males and females. Note that LDSC

correlation estimates are not bound to 1 and can therefore exceed 1 in cases of high GWAS similarity and depending

on the estimation error. (B) Density plot showing the distribution of genetic correlations between males and females

for all cortical ROIs. (C) Differences between females and males (i.e., females – males) in the number of genes

identified through univariate GWASs, for each cortical regions of interest. Pink shades indicate a higher number of

genes in females, whereas blue shades indicate a higher number of genes in males. (D) Number of genes identified

in females (pink) versus males (blue) in the 180 univariate GWASs for cortical ROIs, presented by brain lobe. (E)

Venn diagrams showing the overlap of genes identified in females (pink) and males (blue) across brain lobes, based

on univariate GWASs. Full gene lists can be found in S2-S7.

Next, we examined sex-stratified patterns at the lobe level (**Figure 2D)**. Notably, the

insular and cingulate cortices showed indications of greater sex heterogeneity. Particularly, the

cingulate cortex was significantly associated to twice as many genes in males versus females,

with 38 genes identified compared to 14 in females, with only 2 genes overlapping (**Figure 2E)**.

In contrast, the insula was significantly associated with almost four times more genes in females

(31 genes) than in males (8 genes), with only two genes overlapping between the sexes (**Figure

2E**).

In addition, we examined the overlap of genes identified by univariate GWASs in the

different brain regions, both within and between sexes. Our analysis revealed limited gene

overlap across brain regions and within and between sexes, highlighting regional and potential

sex-dependent variability at the gene level (**Figure 3**).

Figure 3: Genes identified by univariate analyses vary across brain regions and sexes

Legend: Heatmap showing the percentage of overlapping genes out of the total number of genes identified for two given regions, for all cortical and subcortical brain regions, and within or between sexes. Color scale represents the percentage of overlapping genes between two regions whereas the absolute number of overlapping gene is displayed within each cell. The figure is based on univariate GWAS results. Full gene lists can be found in Supplementary Tables 2-12.

Subcortical structures appear to be associated with more genes in females than in males

In line with the cortical results, univariate GWAS for subcortical regions were also strongly

correlated between males and females (**Figure 4A**). All subcortical structures (except the

brainstem) showed a higher number of genes in females than in males (**Figure 4B**).

Different genes emerged in females and males within limbic structures such as the

amygdala and the hippocampus (**Figure 4C**; **Supplementary Tables 8-9)**. In the hippocampus,

we identified 69 genes only genome-wide significant ($p < 5e-8$) in females and 26 genes only

significant in males. In the amygdala, we found 47 genes in females, 37 genes in males and 4

overlapping genes. In the thalamus, only 32 genes were significant in both sex-stratified GWAS,

whereas we identified 166 reached genome-wide significance in the female-stratified GWAS

only, and 119 in the male-stratified GWAS only (**Figure 4C**, **Supplementary Table 10)**.

Lastly, there was no overlap between female and male genes for the hypothalamus

(**Figure 4C**, **Supplementary Table 11)**.

Figure 4: Limited overlap between females and males for genes associated with subcortical

structures

Legend: (A) Density plot showing the distribution of genetic correlations between males and females for all

subcortical ROIs. (B) Comparison of the number of genes identified in females (pink) versus males (blue) by the 77

univariate GWASs for each subcortical ROI, presented by subcortical structure. (C) Venn Diagrams showing the

number of genes identified by univariate GWASs in females (pink) and males (blue); for all ROIs located in

subcortical structures. Full gene lists can be found in S8-S12.

Discussion

The present study used a combination of multivariate and univariate genome-wide approaches to

quantify and overlap the genes significantly associated with human brain anatomy in females and

males separately. Genetic correlations between the sexes were high across all brain regions,

indicating largely shared genetic architecture. Yet, we observed sets of genes that reached

genome-wide significance in only one sex, possibly highlighting candidate regions for sex-

dependent genetic effects. Moreover, our results suggest that a larger set of genes shows

significant associations with brain volume anatomy in females than in males, at least in late

adulthood. This excess of genome-wide significant genes in females aligns with recent reports

demonstrating a higher number of genomic loci with significant local heritability in females

across a wide range of quantitative traits (36).

We found that, particularly in limbic structures, different genes were significant for

females versus males, with many structures being significantly associated with a higher number

of genes in females compared to males. Responsible for emotions, learning and memory (37), the

limbic system has been linked to variation between males and females, both in terms of brain

structure and association with neurological and psychiatric disorders. A previous study by our

group found that limbic structures show greater volume differences between sexes than non-

limbic structures, and that they are more heritable (38). In the present study, the cingulate cortex

was the only region where more genes were identified in males. Sex differences in the cingulate

cortex have been reported in healthy individuals (39), but also in people with autism spectrum

disorders (40) and depression (41). The cingulate cortex is also associated with impulsivity and

aggressive behaviors in childhood, and these associations have been shown to differ between

males and females (42,43). In line with these evidence, most of the cingulate genes only

identified as significant in males have previously been associated with male-prevalent mental

disorders, including autism spectrum disorders (e.g., *RPP25* (44)) and cocaine dependence (e.g.,

NSF (45)).

The present study identified genes associated with brain volumes that could potentially be

linked to variations in the clinical presentation of common brain disorders, although further

research is needed to confirm this hypothesis. As an example, in the hippocampus and thalamus

of females—but not males—we respectively found *CARF*, *NBEAL1*, and *LRP11*, as well as

CRHR1, *NSUN3*, and *NCKIPSD*. These genes have previously been implicated in eating

disorders or anxiety – conditions that disproportionately affect females (46,47). In contrast, the

genes that we identified as significant only in males were frequently involved in male-prevalent

conditions that typically onset in childhood, such as autism spectrum disorders, intellectual

disability and epilepsy (e.g., *AUTS2* (48), *RPP25* (44), *SLC1A2* (49)). Overall, our findings are a

first step in exploring how neurogenetic variation could relate to behavioural and clinical sex

differences in brain disorders. Sex-related variability has long been overlooked in research; yet, a

deeper understanding of the underlying biological mechanisms is crucial to translate

neuroscientific advances into clinical applications that can meaningfully reduce mental health

inequalities (1,9,50,51).

Some of the genes identified in the present study have previously been associated with

more than one disorder and are involved in neurotransmission. This includes *SLC6A4* (identified

in the female insula), the molecular target of selective serotonin reuptake inhibitors (SSRIs)

which has been associated with sex-dependent effects in depression (52), obsessive-compulsive

symptoms (53), and bipolar disorder (54). Another of the genes identified for females in the

insula is *GRID2*, a glutamate receptor possibly associated with antipsychotic response (55)

which has been associated with obsessive-compulsive symptoms in females but not in males

(56). In the hippocampus, a gene only identified in females was *GRIN2A,* a component of the

NMDA receptor complex previously associated with schizophrenia (57), bipolar disorder (58)

and depression (59). In contrast, one example of neurotransmission-related gene identified as

significant only in males is the GABA receptor *GABBR1*, previously associated with

schizophrenia, bipolar disorder, depression and autism spectrum disorder (60), and now linked to

male thalamic volume. These findings may indicate that sex could modulate neurotransmission

pathways, influencing specific brain structure in a sex-dependent manner.

On a different note, females and males may experience various environmental triggers or

events throughout their lives (e.g., trauma, pregnancy, parenting), during which their brains

undergo adaptations that may have long-lasting effects (61,62). Genetic regulatory factors may

modulate these adaptations (63). These include epigenetic modifications and other mechanisms

that regulate gene expression – factors that are now recognized as key players in sex variations in

human tissues (20,64,65). Our findings appear in line with such evidence since many genes only

significant in females or in males were involved in the regulation of gene expression, either

through transcriptional regulation (e.g., *FOXP2*, *TP53*, *RFX4*), or through chromatin

organization and epigenetic processes (e.g. *BLTP3A*, *ASH1L*, *PHF21A*). In the future, it will be

important to elucidate how these regulatory factors interact, respond to environmental triggers,

and contribute to the risk of common brain disorders in females and males.

While our results provide important new insights into sex patterns in neurogenetics, they

need to be interpreted in the light of some limitations. We studied a population of adults aged 40-

80, and the link between genetics, brain and disorder risks may not be the same at all stages of

life. Although we minimized the impact of age on our results by carefully matching males and

females on age, it will be interesting to explore, in the future, the degree of stability in the

observed associations across the lifespan. Lastly, our analyses comparing male and female

neurogenetics are descriptive and do not constitute a formal test of sex differences. Instead, they

provide a complementary perspective to interaction models by illustrating how genetic signals

are distributed across sexes, offering useful context while acknowledging that firm conclusions

would require dedicated inferential approaches. By mapping the here identified sex-dependent

genes to previous research, our study also highlights avenues for targeted follow-up work with

potential relevance for brain disorders.

Conclusion

In conclusion, while the overall genetic architecture of brain anatomy appears largely similar

between females and males, our gene-level results suggest the possibility of sex-dependent

features in the genetic underpinnings of brain structure. Future studies are needed to explore sex

sensitive neurogenetics across different life stages, and in relation to mental and neurological

disorders. Such evidence may improve our understanding of the relationship between sex,

genetics and the brain and their role in the onset, presentation, and course of brain disorders, and

could also help to identify periods or life events that exacerbate the potential of neurogenetic

factors to intensify risk differences between the sexes.

Declarations

Acknowledgements

This work was undertaken on the Tjeneste for Sensitive Data (TSD) facilities which is

operated/developed by University of Oslo TSD service group (IT-Department), and owned by

the University of Oslo as well as on resources provided by the National Infrastructure for High

Performance Computing and Data Storage in Norway (UNINETT Sigma2). We would like to

thank the UK Biobank team and their research participants, for making this work possible.

Funding

TK received funding from the Research Council of Norway (#323961, BrainGap), the German

Research Foundation (IRTG 2804), and the European Research Council (#101086793,

HealthyMom). TK is a member of the Machine Learning Cluster of Excellence, EXC number

2064/1 – Project number 39072764. EB was the recipient of postdoctoral training awards from

the Canadian Institutes of Health Research (#489915) and from the Fonds de Recherche du

Québec – Santé (#329028) when conducting this study. OAA has received funding from

Research Council of Norway (#324252, #324499) and Nordforsk (#164218).

Data availability

Data used in this study were obtained from the UK Biobank (22). GWAS summary statistics and

FUMA results will be made openly available upon acceptance.

Code availability

Codes and software needed to generate the results are available publicly.

379   MOSTest: https://github.com/precimed/mostest

380   FUMA: https://fuma.ctglab.nl/

381   LDSC: https://github.com/bulik/ldsc

Disclosures

384   OAA is a consultant to Precision Health and has received speaker's honoraria from Lilly, BMS,

385   Lundbeck and Janssen. The other authors have no conflict of interest to declare.

References

1. Franceschini A, Fattore L. Gender-specific approach in psychiatric diseases: Because sex matters. European Journal of Pharmacology. 2021;896:173895.

2. Mauvais-Jarvis F, Bairey Merz N, Barnes PJ, Brinton RD, Carrero JJ, DeMeo DL, et al. Sex and gender: modifiers of health, disease, and medicine. Lancet. 2020;396(10250):565–82.

3. Treasure J, Duarte TA, Schmidt U. Eating disorders. The Lancet. 2020;395(10227):899–911.

4. Kuehner C. Why is depression more common among women than among men? The Lancet Psychiatry. 2017;4(2):146–58.

5. Podcasy JL, Epperson CN. Considering sex and gender in Alzheimer disease and other dementias. Dialogues in Clinical Neuroscience. 2016;18(4):437–46.

6. Faheem M, Akram W, Akram H, Khan MA, Siddiqui FA, Majeed I. Gender-based differences in prevalence and effects of ADHD in adults: A systematic review. Asian Journal of Psychiatry. 2022;75:103205.

7. Loomes R, Hull L, Mandy WPL. What is the male-to-female ratio in autism spectrum cisorder? A systematic review and meta-analysis. J Am Acad Child Adolesc Psychiatry. 2017;56(6):466–74.

8. White AM. Gender differences in the epidemiology of alcohol use and related harms in the United States. Alcohol Res. 2020;40(2):01.

9. Shansky RM, Murphy AZ. Considering sex as a biological variable will require a global shift in science culture. Nat Neurosci. 2021;24(4):457–64.

10. Bölte S, Neufeld J, Marschik PB, Williams ZJ, Gallagher L, Lai MC. Sex and gender in neurodevelopmental conditions. Nat Rev Neurol. 2023;19(3):136–59.

11. Riecher-Rössler A. Sex and gender differences in mental disorders. The Lancet Psychiatry. 2017;4(1):8–9.

12. Rechlin RK, Splinter TFL, Hodges TE, Albert AY, Galea LAM. An analysis of neuroscience and psychiatry papers published from 2009 and 2019 outlines opportunities for increasing discovery of sex differences. Nat Commun. 2022;13(1):2137.

13. Wierenga LM, Doucet GE, Dima D, Agartz I, Aghajani M, Akudjedu TN, et al. Greater male than female variability in regional brain structure across the lifespan. Hum Brain Mapp. 2020;43(1):470–99.

14. Salminen LE, Tubi MA, Bright J, Thomopoulos SI, Wieand A, Thompson PM. Sex is a defining feature of neuroimaging phenotypes in major brain disorders. Hum Brain Mapp. 2021;43(1):500–42.

15. Bakker J. The sexual differentiation of the human brain: Role of sex hormones versus sex chromosomes. Curr Top Behav Neurosci. 2019;43:45–67.

16. Blokland GAM, Grove J, Chen CY, Cotsapas C, Tobet S, Handa R, et al. Sex-dependent shared and nonshared genetic architecture across mood and psychotic disorders. Biol Psychiatry. 2022;91(1):102–17.

17. Yaffe K, Haan M, Byers A, Tangen C, Kuller L. Estrogen use, APOE, and cognitive decline: evidence of gene-environment interaction. Neurology. 2000;54(10):1949–54.

18. Oliva M, Muñoz-Aguirre M, Kim-Hellmuth S, Wucher V, Gewirtz ADH, Cotter DJ, et al. The impact of sex on gene expression across human tissues. Science. 2020;

369(6509):eaba3066.
19.   Fass SB, Mulvey B, Chase R, Yang W, Selmanovic D, Chaturvedi SM, et al.
Relationship between sex biases in gene expression and sex biases in autism and Alzheimer's
disease. Biol Sex Differ. 2024;15:47.
20.   Lopes-Ramos CM, Chen CY, Kuijjer ML, Paulson JN, Sonawane AR, Fagny M,
et al. Sex differences in gene expression and regulatory networks across 29 human tissues. Cell
Rep. 2020;31(12):107795.
21.   van der Meer D, Frei O, Kaufmann T, Shadrin AA, Devor A, Smeland OB, et al.
Understanding the genetic determinants of the brain with MOSTest. Nat Commun.
2020;11(1):3512.
22.   Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK
Biobank resource with deep phenotyping and genomic data. Nature. 2018;562(7726):203–9.
23.   Alfaro-Almagro F, Jenkinson M, Bangerter NK, Andersson JLR, Griffanti L,
Douaud G, et al. Image processing and quality control for the first 10,000 brain imaging datasets
from UK Biobank. Neuroimage. 2018;166:400–24.
24.   Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J, et al.
Multimodal population brain imaging in the UK Biobank prospective epidemiological study. Nat
Neurosci. 2016;19(11):1523–36.
25.   Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, et al. A
multi-modal parcellation of human cerebral cortex. Nature. 2016;536(7615):171–8.
26.   Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole
brain segmentation: automated labeling of neuroanatomical structures in the human brain.
Neuron. 2002;33(3):341–55.
27.   Saygin ZM, Kliemann D, Iglesias JE, van der Kouwe AJW, Boyd E, Reuter M, et
al. High-resolution magnetic resonance imaging reveals nuclei of the human amygdala: manual
segmentation to automatic atlas. Neuroimage. 2017;155:370–82.
28.   Iglesias JE, Augustinack JC, Nguyen K, Player CM, Player A, Wright M, et al. A
computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI:
Application to adaptive segmentation of in vivo MRI. Neuroimage. 2015;115:117–37.
29.   Iglesias JE, Insausti R, Lerma-Usabiaga G, Bocchetta M, Van Leemput K, Greve
DN, et al. A probabilistic atlas of the human thalamic nuclei combining ex vivo MRI and
histology. Neuroimage. 2018;183:314–26.
30.   Iglesias JE, Van Leemput K, Bhatt P, Casillas C, Dutt S, Schuff N, et al. Bayesian
segmentation of brainstem structures in MRI. Neuroimage. 2015;113:184–95.
31.   Billot B, Bocchetta M, Todd E, Dalca AV, Rohrer JD, Iglesias JE. Automated
segmentation of the hypothalamus and associated subunits in brain MRI. Neuroimage.
2020;223:117287.
32.   Wiersch L, Hamdan S, Hoffstaedter F, Votinov M, Habel U, Clemens B, et al.
Accurate sex prediction of cisgender and transgender individuals without brain size bias. Sci
Rep. 2023;13(1):13868.
33.   Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and
annotation of genetic associations with FUMA. Nat Commun. 2017;8(1):1826.
34.   Bulik-Sullivan BK, Loh PR, Finucane HK, Ripke S, Yang J, Schizophrenia
Working Group of the Psychiatric Genomics Consortium, et al. LD Score regression
distinguishes confounding from polygenicity in genome-wide association studies. Nat Genet.

2015;47(3):291–5.

35. Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh PR, et al. An
atlas of genetic correlations across human diseases and traits. Nat Genet. 2015;47(11):1236–41.

36. Uffelmann E, de Leeuw C, Schipper M, Posthuma D. Local genetic sex
differences in quantitative traits. Nat Commun. 2025;16(1):7232.

37. Rolls ET. The cingulate cortex and limbic systems for emotion, action, and
memory. Brain Struct Funct. 2019;224(9):3001–18.

38. Matte Bon G, Kraft D, Comasco E, Derntl B, Kaufmann T. Modeling brain sex in
the limbic system as phenotype for female-prevalent mental disorders. Biol Sex Differ.
2024;15(1):42.

39. Ge R, Liu X, Long D, Frangou S, Vila-Rodriguez F. Sex effects on cortical
morphological networks in healthy young adults. NeuroImage. 2021;233:117945.

40. Floris DL, Filho JOA, Lai MC, Giavasis S, Oldehinkel M, Mennes M, et al.
Towards robust and replicable sex differences in the intrinsic brain function of autism. Mol
Autism. 2021;12:19.

41. Mohammadi S, Seyedmirzaei H, Salehi MA, Jahanshahi A, Zakavi SS, Dehghani
Firouzabadi F, et al. Brain-based sex differences in depression: A systematic review of
neuroimaging studies. Brain Imaging and Behavior. 2023;17(5):541–69.

42. Boes AD, Tranel D, Anderson SW, Nopoulos P. Right anterior cingulate: A
neuroanatomical correlate of aggression and defiance in boys. Behavioral Neuroscience.
2008;122(3):677–84.

43. Chen Y, Ide JS, Li CS, Chaudhary S, Le TM, Wang W, et al. Gray matter
volumetric correlates of dimensional impulsivity traits in children: Sex differences and
heritability. Hum Brain Mapp. 2022;43(8):2634–52.

44. Huang HS, Cheung I, Akbarian S. RPP25 is developmentally regulated in
prefrontal cortex and expressed at decreased levels in autism spectrum disorder. Autism Res.
2010;3(4):153–61.

45. Cabana-Domínguez J, Roncero C, Grau-López L, Rodríguez-Cintas L, Barral C,
Abad AC, et al. A highly polymorphic copy number variant in the NSF gene is associated with
cocaine dependence. Sci Rep. 2016;6:31033.

46. Hinney A, Kesselmeier M, Jall S, Volckmar AL, Föcker M, Antel J, et al.
Evidence for three genetic loci involved in both anorexia nervosa risk and variation of body mass
index. Mol Psychiatry. 2017;22(2):192–201.

47. Burstein D, Griffen TC, Therrien K, Bendl J, Venkatesh S, Dong P, et al.
Genome-wide analysis of a model-derived binge eating disorder phenotype identifies risk loci
and implicates iron metabolism. Nat Genet. 2023;55(9):1462–70.

48. Hori K, Shimaoka K, Hoshino M. AUTS2 gene: Keys to understanding the
pathogenesis of neurodevelopmental disorders. Cells. 2021;11(1):11.

49. Green JL, dos Santos WF, Fontana ACK. Role of glutamate excitotoxicity and
glutamate transporter EAAT2 in Epilepsy: opportunities for novel therapeutics development.
Biochem Pharmacol. 2021;193:114786.

50. Tannenbaum C, Greaves L, Graham ID. Why sex and gender matter in
implementation research. BMC Med Res Methodol. 2016;16(1):145, s12874-016-0247–7.

51. Bhargava A, Arnold AP, Bangasser DA, Denton KM, Gupta A, Hilliard Krause
LM, et al. Considering sex as a biological variable in basic and clinical studies: An Endocrine

Society Scientific Statement. Endocr Rev. 2021;42(3):219–58.

52. Baune BT, Hohoff C, Mortensen LS, Deckert J, Arolt V, Domschke K. Serotonin
transporter polymorphism (5-HTTLPR) association with melancholic depression: a female
specific effect? Depress Anxiety. 2008;25(11):920–5.

53. Lei X, Chen C, He Q, Chen C, Moyzis RK, Xue G, et al. Sex determines which
section of the *SLC6A4* gene is linked to obsessive–compulsive symptoms in normal Chinese
college students. Journal of Psychiatric Research. 2012;46(9):1153–60.

54. Wang TY, Lee SY, Chen SL, Chang YH, Chen SH, Huang SY, et al. Gender-
specific association of the SLC6A4 and DRD2 gene variants in bipolar disorder. International
Journal of Neuropsychopharmacology. 2014;17(2):211–22.

55. Allen JD, Bishop JR. A systematic review of genome-wide association studies of
antipsychotic response. Pharmacogenomics. 2019;20(4):291–306.

56. Khramtsova EA, Heldman R, Derks EM, Yu D, Tourette Syndrome/Obsessive-
Compulsive Disorder Working Group of the Psychiatric Genomics Consortium, Davis LK, et al.
Sex differences in the genetic architecture of obsessive-compulsive disorder. Am J Med Genet B
Neuropsychiatr Genet. 2019;180(6):351–64.

57. Harrison PJ, Bannerman DM. GRIN2A (NR2A): a gene contributing to
glutamatergic involvement in schizophrenia. Mol Psychiatry. 2023;28(9):3568–72.

58. Zandi PP, Jaffe AE, Goes FS, Burke EE, Collado-Torres L, Huuki-Myers L, et al.
Amygdala and anterior cingulate transcriptomes from individuals with bipolar disorder reveal
downregulated neuroimmune and synaptic pathways. Nat Neurosci. 2022;25(3):381–9.

59. Brown SJ, Brown AM, Purves-Tyson TD, Huang XF, Shannon Weickert C,
Newell KA. GRIN2B gene expression is increased in the anterior cingulate cortex in major
depression. J Psychiatr Res. 2023;160:204–9.

60. Wu Y, Cao H, Baranova A, Huang H, Li S, Cai L, et al. Multi-trait analysis for
genome-wide association study of five psychiatric disorders. Transl Psychiatry. 2020;10:209.

61. Barha CK, Galea LAM. The maternal 'baby brain' revisited. Nat Neurosci.
2017;20(2):134–5.

62. Helpman L, Zhu X, Suarez-Jimenez B, Lazarov A, Monk C, Neria Y. Sex
differences in trauma-related psychopathology: a critical review of neuroimaging literature
(2014–2017). Curr Psychiatry Rep. 2017;19(12):104.

63. Kundakovic M, Tickerhoof M. Epigenetic mechanisms underlying sex differences
in the brain and behavior. Trends in Neurosciences. 2024;47(1):18–35.

64. McCarthy MM. A new view of sexual differentiation of mammalian brain. J
Comp Physiol A Neuroethol Sens Neural Behav Physiol. 2020;206(3):369–78.

65. Glass K, Quackenbush J, Silverman EK, Celli B, Rennard SI, Yuan GC, et al.
Sexually-dimorphic targeting of functionally-related genes in COPD. BMC systems biology.
2014;8:118.

	Cingulate males	Frontal males	Insula males	Occipital males	Parietal males	Temporal males	Amygdala males	Hippocampus males	Thalamus males	Hypothalamus males	Brainstem males	Cingulate females	Frontal females	Insula females	Occipital females	Parietal females	Temporal females	Amygdala females	Hippocampus females	Thalamus females	Hypothalamus females	Brainstem females
Brainstem females	0	0	0	0	0	0	0	1	10	0	25	0	0	0	0	0	2	0	1	1	1	49
Hypothalamus females	16	0	0	16	0	15	0	0	1	0	1	0	0	0	16	0	15	0	0	17	22	1
Thalamus females	18	2	0	19	4	24	4	6	32	1	2	2	3	0	18	2	16	2	14	198	17	1
Hippocampus females	2	3	2	2	8	19	5	32	15	1	1	2	0	2	3	8	7	28	101	14	0	1
Amygdala females	0	1	0	0	0	4	4	8	2	1	0	0	0	1	1	1	6	51	28	2	0	0
Temporal females	15	0	0	15	3	20	1	8	6	0	2	1	15	0	18	2	84	6	7	16	15	2
Parietal females	3	3	0	2	14	5	1	10	0	0	0	2	3	0	4	79	2	1	8	2	0	0
Occipital females	18	4	0	26	10	15	0	2	0	0	0	2	3	0	46	4	18	1	3	18	16	0
Insula females	1	0	2	0	1	0	0	0	2	0	0	0	0	31	0	0	0	1	2	0	0	0
Frontal females	0	9	0	1	4	0	0	0	7	0	0	0	49	0	3	3	15	0	0	3	0	0
Cingulate females	2	0	0	2	2	0	0	2	0	0	0	14	0	0	2	2	1	0	2	2	0	0
Brainstem males	0	0	0	0	0	0	0	1	9	0	49	0	0	0	0	0	2	0	1	2	1	25
Hypothalamus males	0	0	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0	1	1	1	0	0
Thalamus males	0	1	2	1	2	9	3	3	151	1	9	0	7	2	0	0	6	2	15	32	1	10
Hippocampus males	2	3	0	2	10	9	9	58	3	1	1	2	0	0	2	10	8	8	32	6	0	1
Amygdala males	0	0	0	0	1	1	41	9	3	1	0	0	0	0	0	1	1	4	5	4	0	0
Temporal males	15	0	0	15	6	63	1	9	9	0	0	0	0	0	15	5	20	4	19	24	15	0
Parietal males	2	6	0	9	74	6	1	10	2	0	0	2	4	1	10	14	3	0	8	4	0	0
Occipital males	18	1	0	54	9	15	0	2	1	0	0	2	1	0	26	2	15	0	2	19	16	0
Insula males	0	1	8	0	0	0	0	0	2	0	0	0	0	2	0	0	0	0	2	0	0	0
Frontal males	0	64	1	1	6	0	0	3	1	0	0	0	9	0	4	3	0	1	3	2	0	0
Cingulate males	38	0	0	18	2	15	0	2	0	0	0	2	0	1	18	3	15	0	2	18	16	0

% of overlapping genes