Kidney disease reprograms microbiome-host signaling to promote heart failure

Alex Yarritu[1,2,3,4,*], Wibke Anders[1,2,3,4,5,*], Arne Thiele[1,2,3,4], Olena Potapenko[1,2,3,4], Fabian Schumacher[5], István A. Szijartó[1,2,3,4], Ariana Matz-Rauch[1,2,3,4], Jakob Versnjak[6], Natnael Gebremedhin[1,2,3,4], Victoria McParland[1,2,3,4], Simon Heckscher[7], Sakshi Kamboj[7], Giuseppe Trimarchi[3,8], Harithaa Anandakumar[1,2,3,4], Franziska Fuckert[1,2,3,4], Carina Hoffmann[1], Sara A. Hassan[1,5], Paul M. Bonnekoh[1,2,3,4], Moritz I. Wimmer[1,2,3,4], Felix Behrens[1,2,3,4], Jakob Voelkl[3,4,9], Rafael Kramann[10], Ulrich Kintscher[3,8], Titus Kuehne[3,6,11], Burkhard Kleuser[5], Alma Zernecke[12], Peter J. Oefner[7], Wolfram Gronwald[7], Katja Dettmer[7], Kai-Uwe Eckardt[4], Dominik N. Müller[1,2,3,13], Marcus Kelm[3,6,11], Johannes Holle[1,2,3,14], Hendrik Bartolomaeus[2,3,12,†], Nicola Wilck[1,2,3,4,†]

[1]Experimental and Clinical Research Center (ECRC), a cooperation of Charité – Universitätsmedizin Berlin and Max Delbrück Center for Molecular Medicine, Berlin, Germany

[2]Max-Delbrück-Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany

[3]DZHK (German Centre for Cardiovascular Research), partner site Berlin, Germany.

[4]Department of Nephrology and Medical Intensive Care, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

[5]Freie Universität Berlin, Department of Biology, Chemistry and Pharmacy, Institute of Pharmacy, Berlin, Germany

[6]Institute of Computer-assisted Cardiovascular Medicine, Deutsches Herzzentrum der Charité, Berlin, Germany

[7]Institute of Functional Genomics, University of Regensburg, Regensburg, Germany

[8]Max Rubner Center for Cardiovascular Metabolic Renal Research, Institute of Pharmacology, Charité – Universitätsmedizin Berlin, Berlin, Germany

[9]Institute for Physiology and Pathophysiology, Johannes Kepler University of Linz, Linz, Austria

[10]Division of Nephrology and Clinical Immunology, University Hospital RWTH Aachen,

Aachen, Germany

[11]Department of Congenital Heart Disease – Pediatric Cardiology, Deutsches

Herzzentrum der Charité, Berlin, Germany

[12]Institute of Experimental Biomedicine, University Hospital Würzburg, Würzburg,

Germany

[13]Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin

and Humboldt-Universität zu Berlin, Berlin, Germany

[14]Department of General Pediatrics and Hematology/Oncology, University Children's

Hospital, University Hospital Tübingen, Tübingen, Germany

*These authors contributed equally to this work.

†These authors contributed equally to this work.

Correspondence: nicola.wilck@charite.de

Summary

Gut microbiome-derived metabolites regulate host physiology, with systemic levels controlled by renal excretion. In kidney disease, impaired clearance leads to metabolite accumulation, inflammation, and secondary cardiovascular damage. We show that adverse cardiac remodeling following kidney disease critically depends on the microbiome. We identify microbiome-derived indoxyl sulfate-mediated activation of the aryl hydrocarbon receptor (AhR) as a key mechanism promoting IL-17A-producing T cells and cardiac fibrosis. AhR inhibition attenuates inflammation and cardiac remodeling. On population level, IL-17A is elevated in chronic kidney disease, particularly with coexisting heart failure. Mechanistically, AhR and IL-17A signaling induce a pro-fibrotic phenotype in cardiac fibroblasts, with conserved responses in human cells. This work identifies a microbiome-AhR-IL-17A axis as a mediator and druggable target of cardiovascular damage in kidney disease.

Introduction

Multimorbidity, the coexistence of two or more chronic conditions, is a major challenge for future medicine[1]. Secondary organ damage arises from a complex interplay between intrinsic physiological processes and environmental exposures. The crosstalk between the kidney, a key regulator of internal homeostasis, and the gut microbiome[2], a metabolically active interface between host and environment is particularly relevant. Chronic kidney disease (CKD), a growing global health burden[3], impairs the clearance of host- and microbiome-derived metabolites, leading to their accumulation and toxic effects[4,5]. Cardiovascular disease (CVD) is the most common comorbidity and the primary cause of mortality in CKD[6], with persistent low-grade inflammation recognized as a major but unaddressed driver of this pathology[7]. Although the gut microbiome strongly influences immunity[8], it is not yet targeted by standard CKD therapies. CKD is associated with an altered gut microbiome composition and production of metabolites entering the host circulation, including ligands of the aryl hydrocarbon receptor (AhR), a widely expressed transcription factor[9]. Indoxyl sulfate (IxS), a prototypical uremic toxin and agonistic AhR ligand, is derived from dietary tryptophan utilisation by gut microbes and hepatic sulfation. Upon ligation and activation, the AhR translocates to the nucleus to regulate gene expression and immune cell differentiation[10]. AhR signaling contributes to inflammation[11], but its effects vary with ligand type, concentration, and context[12].

Here, we examine how gut microbiome-immune system interactions drive adverse cardiac remodeling in kidney disease. We show that microbiome depletion confers protection, while IL-17A emerges as a microbiome-dependent driver of cardiac fibrosis and dysfunction in CKD. IL-17A is increased in CKD patients and even more so with coexisting heart failure. Bacterial metabolite IxS promotes T_H17 differentiation via the AhR, and in heart failure, induces T_H17 and aggravates cardiac remodeling. AhR inhibition suppresses T_H17 induction and cardiac fibrosis despite metabolite accumulation. We identify cardiac murine and human fibroblasts as AhR-modulated, IL-17A-responsive effectors. These findings reveal microbiome-dependent, pro-fibrotic pathways and position the AhR as a therapeutic target for cardiovascular protection in CKD.

Results

Gut microbiota accelerates cardiac remodeling in kidney disease

To assess the contribution of the gut microbiota to the host's response to kidney disease, we compared colonized and microbiome-depleted (oral antibiotics, Abx) 129/Sv mice subjected to subtotal nephrectomy (STNx), with Sham-operated mice as controls (Fig.1a). Disease progression was monitored over a 13-week period. Reduced 16S rRNA gene copies in cecal content confirmed successful microbiota depletion over the course of the experiment (Fig.S1a) and cecal weight was increased in Abx-treated mice, reminiscent of the germ-free (GF) mouse cecum (Fig.S1b)[13]. 16S rRNA gene sequencing of cecal samples from the non-microbiome-depleted groups showed a minor reduction in alpha diversity (Fig.S1c-e) in STNx mice while phyla composition remained similar (Fig.S1f). Principal component analysis (PCA) indicated a separation of bacterial communities (Fig.S1g). We found this to be due to an increase in known CKD-associated genera such as *Bacteroides* and *Ruminococcus* (Fig.S1h)[14]. Notably, the taxonomic shifts observed after STNx were less pronounced than those reported in patients with CKD[15], likely because factors such as diet and medication – major drivers of microbial variation in humans – are standardized or absent in our mouse model[16]. Functionally, bacterial metabolites are particularly relevant in kidney disease, as they can accumulate as a function of microbial production and insufficient renal excretion[5]. Thus, we performed plasma metabolomics by nuclear magnetic resonance (NMR), covering a large spectrum of metabolites (Fig.1b). Antibiotic depletion led to a substantial reduction of metabolite accumulation in STNx mice, with most pronounced changes in the aromatic region of the NMR spectrum (Fig.1c). We therefore quantified a targeted panel of aromatic metabolites in plasma by mass spectrometry (MS, Fig.1d). Bacterially derived toxins such as indoxyl sulfate (IxS), p-cresol sulfate and phenyl sulfate were significantly increased in STNx (Fig.1e) and consequently reduced by Abx treatment. Thus, kidney disease induces taxonomic alterations in the microbiome and elevates bacterial metabolite concentrations in the plasma that were abrogated by microbiome depletion.

To assess the impact of microbiome depletion on the degree of kidney dysfunction, we measured glomerular filtration rate (GFR) longitudinally. While both STNx groups displayed a comparable initial GFR reduction until week 8 (Fig.1f), Abx-treated STNx

mice preserved higher GFR at the endpoint (Fig.1g) despite similar remnant kidney

mass (Fig.S2a). Plasma cystatin C (CysC) confirmed this observation (Fig.S2b).

Focusing on cardiovascular comorbidities as the leading cause of mortality in CKD[6,17],

we next assessed cardiac and vascular damage. STNx induced cardiac remodeling

as reflected by increased heart weight, which was abolished by microbiome depletion

(Fig.1h). Longitudinal echocardiographic assessment of cardiac dimensions

corroborated this effect, becoming evident from week 8 onward (Fig.1i-k; Table S1)

despite similar GFR at this time point. While systolic function (ejection fraction, EF)

was preserved in all groups (Fig. 1l; Table S1), STNx mice displayed an increase in

E/e' (Fig.1i, m; Table S1) indicative of diastolic dysfunction, and a reduction in global

longitudinal strain (GLS, Fig.1n; Table S1), reflecting ventricular stiffening – both

rescued by microbiome depletion. Adverse cardiac remodeling is characterized by

cardiomyocyte hypertrophy, fibrosis and inflammation[18]. Interestingly, we observed a

comparable cardiomyocyte size in STNx (Fig.S2c-d), suggesting that the increased

heart weight is driven by extracellular matrix (ECM) deposition. Sirius red staining

revealed increased collagen deposition in STNx hearts, which was mitigated in

microbiome-depleted STNx mice (Fig.1o-p). In addition, STNx mice exhibited

increased cardiac leukocyte infiltration (Fig.1q), comprising mostly neutrophils and T

cells (Fig.S2e), which was ameliorated in Abx-treated STNx mice. In agreement,

plasma C-reactive protein (CRP) – a routine clinical marker of systemic inflammation

associated with cardiovascular risk[19] – was elevated in STNx mice and reduced by

Abx treatment (Fig.1r). These findings demonstrate that microbiome depletion

prevents hallmarks of adverse cardiac remodeling and diastolic dysfunction induced

by kidney disease.

Impaired kidney function typically leads to hypertension, a common cause for adverse

cardiac remodeling. Further evidence suggests that the gut microbiota influences

blood pressure (BP) and cardiac remodeling[20,21]. We therefore measured BP

longitudinally by radiotelemetry in both control and Abx-treated STNx mice (Fig.1s).

Both groups exhibited a similar increase in both systolic (SBP) and diastolic (DBP)

throughout the experiment, which was sustained towards the endpoint (Fig.1t-u, S2f-

i), suggesting that the cardioprotective effects observed in microbiota-depleted STNx

mice are independent of BP. To assess vascular function, we measured

vasorelaxation in isolated mesenteric arteries. Arteries from Abx-treated STNx mice

displayed an improved endothelium-dependent relaxation compared to those from

STNx mice (Fig.1v-w). Taken together, microbiome depletion protected mice with

reduced kidney function from further GFR loss, inflammation, cardiac and vascular

dysfunction despite a similar blood pressure increase, suggesting these benefits are

linked to immunological mechanisms.

Microbiome-Dependent T_H17 Immunity and AhR Activation

Since cardiac leukocyte influx was attenuated by microbiome depletion, we

hypothesized that microbiome-immune interactions are critical for the pathological

cardiac response to STNx. To assess immune cell composition, we performed flow

cytometry on splenic leukocytes (Fig.S3a). STNx induced an increased abundance of

neutrophils (Fig.2a) and a shift of T cells towards $CD4^+$ T helper subsets (Fig.2b), both

of which were prevented by Abx treatment. Flow cytometric analysis further revealed

an expansion of $CD44^+CD62L^-$ effector memory phenotypes (Fig.2c) for both $FoxP3^-$

conventional T helper cells (T_{conv}, Fig.2d) and $FoxP3^+$ regulatory T cells (T_{reg}, Fig.2e).

These alterations were abrogated in Abx-treated mice. Moreover, splenic T_{conv} from

STNx mice showed an increased abundance of $ROR\gamma t^+$ T_{conv} (T_H17, Fig.2f), which was

prevented by Abx. Similar patterns were detected in circulating and intestinal lamina

propria T cells (Fig.2g), indicating a microbiome-dependent T_H17-type inflammatory

response.

In parallel, we observed an expansion of $ROR\gamma t^+$ T_{reg}, hereafter referred to as $T_{reg}17$

cells (Fig.2h). While $T_{reg}17$ are known to limit inflammation in mucosal tissues[22], they

can also display T_H17-like effector functions, including IL-17A secretion[23]. To further

characterize $T_{reg}17$ in this inflammatory setting, we compared frequencies of $PD-1^+$,

$CD69^+$ and $CD103^+$ cells within the $T_{reg}17$ and T_{reg} compartment. $T_{reg}17$ cells showed

enrichment for these markers relative to total T_{reg} populations in Sham and Abx-treated

STNx mice; however, this pattern was not observed in STNx mice (Fig. S3b–d).

Additionally, $IL-10^+$ T_{reg} were reduced in STNx, but restored under Abx treatment

(Fig.S3e), altogether suggesting a loss of suppressive capacity favoring T_H17-like

effector functions. In line with these findings, STNx induced a sustained T_H17 and

$T_{reg}17$ increase over time (Fig.2i), while T_H17 of Sham and Abx did not show this

response. The parallel expansion of $T_{reg}17$ and T_H17 cells suggests ineffective

regulatory control and persistence of an inflammatory response.

Microbiota-derived metabolites can activate the aryl hydrocarbon receptor (AhR)[24],

which is known to modulate T cell function, including T_H17 differentiation[25,26].

Consistent with the observed increase in microbiome-derived tryptophan catabolites,

predominantly IxS, a known AhR ligand, plasma from STNx mice induced significantly

higher AhR activation in a cell-based reporter assay compared with Abx-treated and

Sham controls (Fig.2j). Moreover, *Cyp1a1*, a canonical AhR target gene, was

upregulated in splenic leukocytes from STNx mice compared to Abx-treated STNx

mice (Fig.2k), suggesting immune cells are affected by bacterial AhR ligands.

Next, we evaluated RORγt+ T cells in fibrotic hearts. We found an increase in $T_{reg}17$,

but not T_H17 cells, in whole hearts of STNx mice after 13 weeks (Fig.2l-m, Fig.S3f),

consistent with the observation that the cardiac microenvironment promotes T cell

transdifferentiation towards regulatory phenotypes[27]. Additionally, infiltrated

neutrophils were reduced in STNx mice treated with Abx (Fig.S3g). To further

characterize the inflammatory microenvironment of the heart, we performed bulk

mRNA sequencing from cardiac tissue. A substantial part of the STNx-associated

differentially expressed genes (DEG) were reversed by Abx treatment (Fig.2n).

Enrichment analysis on DEG identified Reactome pathways and Gene Ontology (GO)

terms involving tissue remodeling (e.g. fibronectin matrix formation, collagen

formation, extracellular matrix organization) and immunity (e.g. T cell receptor

signaling, interleukin 17 signaling, Fig.2o, S3h). Variation analysis of IL-17A-

associated gene sets revealed a significant increase in enrichment scores in STNx,

an effect normalized by Abx (Fig.2p). Taken together, our findings suggest that

microbiome-derived, AhR-activating metabolites drive a systemic T_H17-skewed

inflammatory response that impairs $T_{reg}17$ function and promotes adverse cardiac

remodeling in kidney disease.

IxS exacerbates heart failure

IxS has been shown to modulate T_H17 immunity in neuroinflammation[28]. *In vitro* T_H17

polarization of murine naïve CD4+ T cells was indeed stimulated dose-dependently by

IxS (Fig.3a-b). To test whether IxS stimulates a T_H17 response *in vivo* and aggravates

cardiac remodeling independent of advanced kidney dysfunction, we examined the

effects of IxS supplementation in a mouse model of heart failure with preserved

ejection fraction[29] (HFpEF, Fig.3c), the form of heart failure most frequently associated

with CKD[30]. We confirmed elevated IxS levels in mice exposed to IxS (Fig.3d),

although to a lesser extent compared to STNx mice and patients with CKD[31], likely

due to preserved renal clearance (Fig.3e). Still, IxS levels were sufficient to increase

splenic *Cyp1a1* expression (Fig.3f), but not liver *Cyp1a1* (Fig.S4a)*,* indicating that the

immune system is particularly susceptible to AhR activation by bacterial metabolites.

Similar to STNx mice, splenic IL-17A$^+$ T$_{conv}$ (Fig.3g), IL-17A$^+$ T$_{reg}$ (Fig.3h) and

neutrophils (Fig.S4b) frequencies were elevated in response to IxS. IxS exposure

exacerbated heart failure, evidenced by increased heart weights (Fig.3i) and

pulmonary congestion (Fig.S4c). Echocardiography confirmed aggravation of diastolic

dysfunction (E/e', E/A, Fig.S4d-e; Table S2) with a preserved left ventricular EF

(Fig.S4f; Table S2). Next, we analyzed cardiac fibrosis, the most prominent feature of

STNx-associated cardiac remodeling. IxS-exposed hearts displayed higher collagen I

and fibronectin content, indicating increased interstitial cardiac fibrosis (Fig.3j-l). Thus,

IxS exacerbates heart failure and elicits a similar inflammatory response as observed

in experimental kidney disease.

Cardiac fibroblasts respond to IL-17A and IxS to produce ECM

In addition to the T cell-driven increase in IL-17A production, *Il17ra* expression was

upregulated in STNx hearts (Fig.4a), and likewise in IxS-treated HFpEF mice (Fig.4b).

To identify IL-17A-responsive cells among cardiac cells, we re-analyzed published

single nuclei RNA (sn-RNA) sequencing data from cardiac tissue of STNx and control

mice[32]. Among cardiac cell types, fibroblasts showed the highest *Il17ra* expression in

STNx (Fig.4c, Fig.S5a). Notably, fibroblasts also showed higher expression of AhR-

dependent genes (Fig.S5b) Interestingly, *Il17ra*$^+$ fibroblasts displayed higher ECM

gene set scores[33] than *Il17ra*$^-$ fibroblasts in STNx mice (Fig.4d). In line, the DEG were

characterized by a dominant representation of ECM components (Fig.S5c). To further

characterize *Il17ra*$^+$ fibroblast transcriptomes, we compared their DEG with different

fibroblast signatures known in both mouse and human cardiac tissue[34]. DEG were

enriched in myofibroblast-related and, to a lesser degree, cyclic and WNT-related

fibroblast-specific genes (Fig.4e), known for their pro-fibrotic and activated status.

Cytokine enrichment analysis revealed IL-17A as a prominent factor shaping DEG in

cardiac STNx lymphocytes (Fig.S5d). To test the response of cardiac fibroblasts to

AhR ligand IxS and IL-17A, respectively, we isolated fibroblasts from healthy murine

hearts (Fig.4f). Cell identity and purity were confirmed by immunocytochemistry and

flow cytometry (Fig.4g-h). As suggested by the sn-RNA data, IxS upregulated *Il17ra*

mRNA expression (Fig.4i), indicating IxS primes fibroblasts to sense IL-17A.
Consistent with this, coincubation with IL-17A and IxS resulted in a synergistic
increase in *Col1a1*, *Col1a2*, and *Acta2* expression (Fig.4j-l)*.* In line with *in vivo* data,
IL17A downregulated *Abi3bp* expression, an ECM protein shown to be reduced in
heart failure[35] (Fig.4m). Immunocytochemistry confirmed the presence of COL1 and
IL17RA at the protein level (Fig.4n). Our data demonstrate that IL-17A signaling in
cardiac fibroblasts primed by IxS exposure leads to increased ECM production,
ultimately exacerbating cardiac fibrosis.

260 AhR inhibition ameliorates cardiac remodeling in kidney disease

Since STNx-associated cardiac remodeling occurs in an AhR-activating milieu that in
turn promotes IL-17A production and cardiac fibrosis, we investigated whether AhR
inhibition confers therapeutic benefit. For optimal translational potential, we re-
purposed the orally bioavailable AhR inhibitor BAY 2416964 (AhRi), currently under
investigation in solid malignancies[36]. *In vitro*, AhRi dose-dependently reduced the IxS-
induced upregulation of *Cyp1a1* expression in murine splenocytes (Fig.5a-b).
Furthermore, AhRi dose-dependently suppressed IxS-driven T_H17 polarization
(Fig.5a,c). In both assays, AhRi was effective at doses of $10^{-7.5}$-$10^{-6.5}$ M. Next, we
aimed to test the efficacy of AhRi *in vivo* in STNx mice (Fig.5d). Therefore, mice were
randomized based on serum CysC levels (Fig.S6a) to receive either AhRi or vehicle
starting two weeks after complete STNx through to week 13 by daily oral gavage.
Sham-operated vehicle-treated mice served as additional control. At the endpoint, we
detected similar levels of CysC in AhRi- and vehicle-treated mice (Fig.S6b). AhRi
treatment improved survival of STNx mice (Fig.5e) and lowered systemic
inflammation, as measured by plasma CRP (Fig.5f). Upregulation of canonical AhR
target gene *Cyp1a1* in immune cells was prevented by AhRi (Fig.5g), as well as the
STNx-associated increase in both T_H17 (Fig.5h) and $T_{reg}17$(Fig.5i). In line, IL-17A+
T_{conv} and T_{reg} were reduced by AhRi (Fig.S6c-d). We observed an improvement of the
cardiac phenotype upon AhRi reminiscent of the Abx effect, with lower left ventricular
mass (Fig.5j), and improved cardiac function (Fig.5k-l, Table S3) and fibrosis (Fig.5m-
n), together with lower cardiac *Il17ra* expression (Fig.5o). Bulk transcriptomic analyzes
of cardiac tissue showed similar effects as observed with Abx treatment, with a
reduction in GO terms and pathways involved in ECM formation and inflammatory
processes (Fig.5p, Fig.S6e). Targeted analysis of T_H17-related signaling pathways

revealed significantly higher scores upon STNx, which were rescued by AhRi (Fig.5q).

These findings identify the AhR as a key mediator of inflammation and cardiac

remodeling in kidney disease and suggest AhRi as a therapeutic strategy.

288   Conserved pro-fibrotic IxS-AhR-T$_H$17 axis in patients with CKD

We further aimed to explore the translational potential of AhRi by investigating the

efficacy in human cells. AhRi dose-dependently blocked the IxS-stimulated AhR

activity in human AhR reporter cells (Fig.6a). Likewise, AhRi reduced AhR activity of

serum collected from patients with kidney failure requiring hemodialysis (HD) to levels

observed with serum from healthy donors (HC, Fig.6b-c; Table S4). Moreover, addition

of HD serum to human T$_H$17 polarization assays increased T$_H$17 cells compared to

HC (Fig.6b,d). Hence, the kidney disease-responsive AhR-T$_H$17 axis identified in mice

is conserved in humans.

Next, we analyzed plasma IL-17A levels in patients with CKD and healthy controls

from the UK Biobank (Table S5). Circulating IL-17A concentrations increased

progressively with worsening kidney function (Fig.6e, n= 15 120 with available eGFR

and albuminuria). IL-17A concentrations were further elevated in patients with eGFR

below 60 (CKD) and HFpEF (n = 5 497) compared to patients with only CKD

(n = 15 981, Fig.6f). Furthermore, higher plasma IL-17A in individuals with CKD

(n = 2 785) was associated with all-cause mortality, even after adjustment for

conventional risk factors (Fig.6g-h). These findings substantiate kidney disease-

dependent IL-17A response at population scale.

To verify the role of AhR signaling and IL-17A in fibroblast activation in humans, we

used publicly available single cell RNA (sc-RNA) sequencing data. Since cardiac sc-

RNA data from patients with CKD were not available, we explored data from kidney

tissue of patients with CKD and healthy controls[37]. Fibroblasts of patients with CKD

displayed the highest AhR enrichment scores compared to other kidney cell types

(Fig.S7a). Moreover, myofibroblasts expressed the highest levels of *IL17RA* (Fig.S7b),

and *IL17RA* expressing myofibroblasts exhibited markedly higher ECM module scores

(Fig.S7c). In line, pro-fibrotic and ECM-associated genes were among the major DEG

(Fig.S7d), indicating that IL-17A-responsive fibroblasts represent a pro-fibrotic subset.

Next, we stimulated primary human cardiac fibroblasts with serum from hemodialysis

patients (HD) or healthy controls (HC) (Fig.6i). HD serum induced upregulation of

IL17RA (Fig.6j, Fig.S8a) together with ECM-associated transcripts such as *COL1A2*

and *ACTA2* (Fig.6k-m, Fig.S8b-d). We tested whether these effects could be
prevented by AhRi or secukinumab, a human monoclonal antibody used in clinical
routine targeting IL-17A (αIL-17A). Both AhRi and αIL-17A mitigated the pro-fibrotic
effects (Fig.6k-m, Fig.S8b-d). However, only AhRi reduced *IL-17RA* expression
(Fig.6j, Fig.S8a). Similar to murine cardiac fibroblasts, IxS upregulated *IL17RA*
expression in primary human cardiac fibroblasts, an effect abrogated by AhRi
(Fig.S8e). Both IxS and IL-17A increased *COL1A2* and *ACTA2* expression, whereas
co-stimulation with both ligands again had a synergistic effect, leading to the highest
pro-fibrotic transcriptional response (Fig.S8f-h). Proliferation was not altered with IL-
17A and IxS co-incubation, as determined by Ki-67 staining (Fig.S8i-j). Taken
together, translation of our findings to patients with CKD using data from UK Biobank,
human sc-RNA sequencing and *in vitro* assays using patient-derived serum show a
conserved pro-fibrotic IxS-AhR-T_H17 axis.

Discussion

This study demonstrates that kidney disease recalibrates microbiome-host interactions to potentiate inflammation-mediated cardiac fibrosis. We identify a microbial AhR-activating metabolite driving a pathogenic T_H17 response that engages IL-17A-responsive cardiac fibroblasts, positioning the metabolite-AhR-IL-17A axis as a therapeutic target to prevent cardiac fibrosis in kidney disease.

Inflammation is a key driver of CVD. In CKD, inflammation is substantially elevated, paralleling a markedly increased risk for CVD[38]. Evidence suggests that CVD development in CKD extends beyond traditional risk factors[39] such as hypertension, hyperlipidemia, and diabetes, yet underlying mechanisms remain less characterized. Adverse cardiac remodeling and fibrosis are prevalent in CKD[40], with heart failure representing a frequent clinical manifestation[41]. The microbiome plays a central role in immune modulation, in part via action of bacterial metabolites. In CKD, these metabolites accumulate and contribute to a pro-inflammatory internal milieu. Such metabolites were shown to be of bacterial colonic origin and can reach toxic levels in CKD[5]. To date, therapeutic strategies aimed at improving outcomes by reducing bacterial metabolite levels through unspecific intestinal adsorption showed limited success[42], underscoring the need for a refined understanding of microbiome-host interaction in CKD.

Our mechanistic investigations were prompted by the observation that the depletion of the gut microbiome in a state of reduced kidney function drastically reduced the development of cardiac fibrosis as well as diastolic dysfunction, and suppressed systemic inflammation, as evidenced by a lower plasma CRP. Remarkably, cardioprotection occurred despite persistently high blood pressure, in line with a clinical observation showing that cardiac fibrosis in CKD occurs independent of hypertension[40], which has so far not been linked to the gut microbiome. The profound inflammation encouraged us to investigate cellular immunity further, in which a microbiome-dependent T_H17 response to STNx became evident. While IL-17A has previously been associated to ischemic heart disease[43] and kidney disease progression after acute kidney injury[44], it has not been highlighted as a cytokine linking heart disease and CKD. Our analyzes of UK Biobank data suggest that this may indeed be the case, demonstrating elevated IL-17A levels in patients with CKD and

HFpEF. IL-17A plasma concentrations increased in a CKD stage-dependent manner

and were associated with excess mortality.

We previously hypothesized that CKD-linked inflammation is mediated by the AhR[26].

Plasma AhR activation potential increased following STNx, with splenocytes exhibiting

AhR-dependent gene expression; both effects were abrogated by microbiome

depletion. We present evidence that AhR activation and T_H17 response are mediated

via the AhR by increased levels of IxS, a known microbiome-derived AhR-activating

ligand. *In vitro*, IxS induced T_H17 polarization dose-dependently. *In vivo*, oral

administration of IxS in an experimental heart failure model increased AhR activation

and AhR-dependent gene expression in immune cells, promoted T_H17 expansion, and

aggravated cardiac fibrosis, despite preserved kidney function. Further *in vivo*

evidence is provided by AhR inhibition in STNx mice, which abrogated AhR-dependent

gene expression in immune cells, T_H17 expansion and systemic inflammation, as well

as cardiac fibrosis and diastolic dysfunction. Thus, IxS-dependent AhR activation

appears to be central to cardiac remodeling in kidney disease.

As an environmental sensor integrating various external and endogenous cues, the

AhR is known to regulate T_H17 differentiation[45]. AhR-deficient naïve T cells exhibit

reduced differentiation[45], while AhR activation by high affinity ligands has been shown

to promote T_H17 and autoimmune disease[46]. However, AhR-mediated effects on T_H17

differentiation depend on the ligand and cytokine milieu[47]. IxS is a known endogenous

AhR ligand[28] and has been linked with inflammatory milieus[48] as well as T_H17-

mediated autoimmunity[28]. We present evidence that IxS activates AhR transcriptional

programs in CD4+ cells to promote a pathogenic T_H17 response in kidney disease.

We show that IxS drives RORγt expression and IL-17A production in CD4+ T cells *in*

vitro and *in vivo*. Intriguingly, we also detected increased IL-17A production by Foxp3+

CD4+ T_{reg}, as well as increased RORγt expression in CD4+ FoxP3+ cells ($T_{reg}17$),

especially in cardiac tissue. While $T_{reg}17$ cells typically restrain IL-17A-driven

inflammation through IL-10[48], their regulatory capacity appears diminished in the

present uremic setting, where IL-17A-mediated inflammation is less effectively

controlled. Consistently, $T_{reg}17$ cells from uremic STNx mice exhibited reduced

expression of PD-1, CD103, and CD69 – proteins implicated in the maintenance of

suppressive function[49-51] – compared with microbiome-depleted and kidney-healthy

controls. This phenotype suggests a functional shift in $T_{reg}17$ cells toward RORγt-

driven effector activity at the expense of FoxP3-dependent regulatory functions.

Moreover, IL-10 production was lowest in T_{reg}s from STNx mice. The STNx-induced
IL-17A response was accompanied by neutrophilia and enhanced cardiac neutrophil
infiltration. IL-17A is a known regulator of neutrophils, promoting granulopoiesis and
inducing neutrophil-attracting chemokines in stromal and cardiac cells[52]. Circulating
neutrophils correlate with IL-17A levels[53]. Once recruited, neutrophils maintain IL-17A-
driven inflammation through release of inflammatory mediators, including IL-1β and
neutrophil extracellular traps[54]. Together, these findings reveal a microbiome-
dependent, AhR-mediated and IL-17A-driven inflammatory program in kidney disease.
This study identifies cardiac fibroblasts as key targets of IL-17A, characterized by
enhanced IL-17RA expression *in vivo*, which in turn drives ECM production and
promotes cardiac fibrosis. It has been reported that IL-17A enhances fibroblast
proliferation and migration[55] and induces a proinflammatory fibroblast phenotype
involving GM-CSF[56]. We demonstrate that IxS effectively primes cardiac fibroblasts
via AhR-dependent upregulation of IL-17RA in murine and human cells, suggesting
that the same bacterial metabolite that drives IL-17A-mediated inflammation also
sensitizes fibroblasts to IL-17A signaling and ECM production. Thus, IxS orchestrates
inflammation and tissue remodeling through AhR signaling, providing a mechanistic
explanation for the pronounced cardiac fibrosis observed in CKD and a strong
rationale for AhR inhibition as a therapeutic strategy. Notably, pharmacological AhR
targeting with a small-molecule inhibitor currently under clinical investigation for cancer
treatment[36] (NCT04999202) proved effective both *in vivo* in STNx mice and *in vitro* in
human cardiac fibroblasts, highlighting its translational potential. Targeting host
sensing of uremic toxins may uncouple disease-driving microbiome-derived
metabolites from downstream inflammatory responses. Integrating AhR inhibition with
existing therapies could therefore provide a complementary strategy to mitigate
adverse cardiovascular comorbidities in CKD.
Taken together, our findings identify an IxS-AhR-IL-17A axis as a central driver of
cardiac inflammation and fibrosis in kidney disease. We demonstrate how kidney
dysfunction promotes cardiac pathology through specific microbiome-derived
metabolites. Our study positions IxS not merely as a biomarker of uremia but as an
active pathogenic effector linking microbial metabolite accumulation to maladaptive
immunity and tissue fibrosis via AhR signaling. AhR inhibition may offer a means to
selectively target uremic downstream signaling while preserving potentially beneficial

microbiome functions. More broadly, these findings refine our understanding of

disease-to-multimorbidity transition in chronic kidney disease.

Methods

Animal studies

All experiments were performed in accordance with the German/European law for
animal protection. Experimental protocols were approved by the local ethic authorities
under the G0190/21 and G0019/21 licenses. For all *in vivo* experiments, male 129/Sv
mice (wild type [WT]; strain 129S2/SvPasOrlRj) were obtained from Janvier Labs,
France, and housed at 12-hour day:night cycle with free access to water and food
(Chow diet, Ssniff, V1124-300) in the Preclinical Research Center (PRC) of the Max-
Delbrück-Center for Molecular Medicine (MDC). Mice were habituated for 2 weeks
before use. Individual mice served as experimental units, and all animals within a cage
received the same treatment.

For the Abx trial, mice were distributed according to body weight (BW) to four groups
prior to treatment (STNx ± Abx and Sham ± Abx). The Abx experiment consisted of
two independent cohorts. A longitudinal cohort was designed to evaluate dynamics in
kidney function, metabolomics and immunity in STNx groups (n=11 per group) and
Shams (n=7 per group) across timepoints (w1, w3, w8, w13). Each timepoint had its
own independent cohort to collect biomaterial. A second cohort was designed for
terminal assessments, focused on organ damage and tissue level responses in STNx
groups (n=12) and Shams (n=7). Both cohorts were exploratory, thereby no prior
sample size calculation was performed. Endpoint analyzes were performed combining
both cohorts if applicable. Animals were excluded in cases of surgical failure or failure
to reach minimum kidney function thresholds. Animals were sacrificed during the
course of the study if BW dropped under 20% or they showed clear signs of pain or
discomfort. The longitudinal cohort resulted in: w1 - Sham n=7; Abx n=6; STNx n= 6;
STNx Abx n= 6. w3 - Sham n=6; Abx n= 7; STNx n=10; STNx Abx n=7. w8 - Sham
n=5; Abx n=5; STNx n=6; STNx Abx n=6. w13 - Sham n=6; Abx n=4; STNx n=6; STNx
Abx n=6. The endpoint cohort resulted in: Sham n=7; Abx n=7; STNx n=11; STNx Abx
n=10.

For the HFD/L-NAME trial, mice were assigned according to BW and cage to receive
either potassium indoxyl sulfate (IxS) exposure (n=16) or control (KCl; n=16) over the
drinking water *ad libitum.* Sample size was calculated using G Power according to a
partial effect size of the alleviation of the STNx-induced diastolic dysfunction by Abx.

During the experiments n=3 from the control group were taken out of the experiment
due to prespecified criteria.

For the AhRi trial, mice underwent STNx (n=24) or Sham (n=6) surgery. Sample size
was calculated using G Power according to the effect size of the alleviation of the
STNx-induced cardiac fibrosis by Abx. The same exclusion criteria were followed as
in the Abx trial. To ensure equal kidney dysfunction before treatment, mice were
equally distributed into 2 groups according to BW and plasma CysC two weeks after
completed STNx surgery, and assigned to receive either AhRi (n=10) or vehicle (n=9).
During the experiments n=4 vehicle and n=2 AhRi-treated animals were taken out of
the experiment due to prespecified criteria and analyzed at an earlier time point.

Across all experiments, blinding during *in vivo* procedures was not possible due to
model-specific features; however, all outcome measurements and data analysis were
performed blindly.

Subtotal nephrectomy

Mice between 10 and 12 weeks old were anaesthetized with isoflurane (2.5%
isoflurane-air) and subjected to subtotal nephrectomy operation (STNx). Briefly, the
left kidney was exposed, and the lower and upper poles were excised using a
cauterizer (Surtron 160, LED SpA). The adrenal gland was separated from the kidney
to preserve its function. 2 weeks later, a total nephrectomy of the right kidney was
performed. Sham-operated mice underwent a simple flank incision and sutures. All
animals received pre-emptive buprenorphine (30 mg/kg, intraperitoneal (i.p.) injection)
30 min before surgery. Post surgery, analgesia consisted of metamizole (800
mg/kg/24h) administered over the drinking water for 3 days.

Antibiotic treatment

Mice were given an antibiotic cocktail (all obtained individually from Thermofisher)
containing ampicillin (0.5 g/L), vancomycin hydrochloride (0.5 g/L), neomycin sulfate
(1 g/L) and metronidazole (1 g/L) in the drinking water throughout the entire study
period. 1% glucose was added into the mixture for organoleptic purposes. Control
animals received only glucose in the drinking water. Treatment was started one week
prior to the first operation and antibiotics were refreshed every 3 days.

<u>AhR inhibitor treatment</u>

Mice were stratified and randomized as described above to receive either small-
molecule AhR inhibitor BAY2416964 (GlpBio) or vehicle, starting two weeks after
completed STNx surgery. AhRi was formulated in a mixture of water, Solutol® HS 15
(polyethylene glycol 12-hydroxystearate; Merck) and ethanol at a 50:40:10 ratio
(v/v)[36]. The vehicle control consisted of the solvent mixture alone. A daily AhRi dose
of 30 mg/kg was selected based on a previously reported safe and effective regimen[36].

<u>HFpEF model</u>

12-week-old mice were switched to a high fat diet (HFD; 60% kcal fat, Research diets,
D19032801-1.5V) and administered Nω-nitro-L-arginine methyl ester hydrochloride
(L-NAME; 1 g/L, Merck) in the drinking water for 15 weeks. Mice received either
potassium indoxyl sulfate (IxS; 1 g/L, Merck) or equimolar potassium chloride (KCl;
0.3 g/L, Merck) in the drinking water throughout the study. This model has been
published previously[29].

Microbiome

<u>Cecal DNA isolation</u>

At different timepoints (w1, w3, w8, w13), cecal contents were collected at the time of
euthanasia. Samples were immediately fresh-frozen on dry ice and kept at -80 °C.
DNA was isolated from cecal samples using the Quick-DNA Miniprep Kit (Zymo
Research) according to manufacturer's protocol. Absolute bacterial load was
determined by qPCR for the *16S* gene using SYBR Green as described below.

<u>Microbiome analysis</u>

Samples were stored at -20°C and sent for 16S rRNA gene amplicon sequencing
(Novogene, Cambridge, UK). The V3-V4 hypervariable region of the 16S rRNA gene
was amplified using universal primers 341F (5′-CCTAYGGGRBGCASCAG-3′) and
806R (5′-GGACTACNNGGGTATCTAAT-3′), which were appended with sample-
specific barcodes and Illumina sequencing adapters. Amplicons were purified using
AMPure XP magnetic beads and quantified for library preparation.

Libraries were constructed using the NEBNext® Ultra™ II DNA Library Prep Kit and
sequenced on an Illumina NovaSeq 6000 platform (paired-end 2×250 bp), targeting

approximately 30,000 raw reads per sample. Annotation was performed in R (v4.0.3).

Demultiplexed reads were processed through DADA2 pipeline (v1.25.2) and

taxonomic assignment was based on the rRNA database SILVA (v138.1) with a 80%

boodstrapped confidence. Only ASVs that were present in at least 20% of the samples

were considered for further downstream analysis.

Nuclear magnetic resonance (NMR) spectroscopy

<u>Sample preparation</u>

Plasma samples were prepared using 3 mm SampleJet NMR tubes. 100 µL of plasma

was mixed thoroughly with 78 µL of 0.1 mol/L phosphate buffer (pH 7.4) and 19.5 µL

of a 0.75% (w) solution of 3-trimethylsilyl-2,2,3,3-tetradeuteropropionate (TSP; Sigma-

Aldrich, Taufkirchen, Germany) in deuterium oxide, serving as an internal standard for

NMR measurements. As an additional internal standard 2.5 µL of formic acid (240

mM) was added. Prepared samples were measured as described below.

<u>Data acquisition</u>

All NMR experiments were performed on a 600 MHz Bruker Avance III HD

spectrometer, using a triple resonance (^{1}H, ^{13}C, ^{15}N, ^{2}H lock) helium cooled cryoprobe

with z-gradient. Samples were handled by an automatic Bruker SampleJet sample

changer (Bruker Biospin GmbH, Ettlingen, Germany). Tuning and matching of the

probe as well as locking and shimming of the sample were performed automatically.

For each sample one dimensional ^{1}H NMR spectra were obtained at 310 K using Carr-

Purcell-Meiboom-Gill (CPMG) pulse sequence. For each spectrum, 512 scans were

collected into 72k datapoints over a spectral width of 12019 Hz. An acquisition time

and T_1 relaxation delay of 3.07 and 4 seconds per scan were used, respectively. All

NMR spectra were processed with TopSpin 4.5.0. Free induction decays (FIDs) were

Fourier transformed, phase corrected and subjected to exponential line broadening

function of 0.3 Hz. All the chemical shifts were manually referenced to formic acid at

8.46 ppm.

To compensate for chemical shifts perturbations bucket tables were generated from

the processed 1D CPMG spectra using AMIX software (version 3.9.13, Bruker,

Ettlingen, Germany). The spectra over a range from δ0.11 to δ 9.5 ppm were divided

into the buckets at an equal width of 0.01 ppm with the regions at $\delta 4.59$ - $\delta 4.77$ ppm

being excluded as the water peak. The spectra were not normalized.

Bucket analysis

Only buckets within the aromatic region (6.75-8.00 ppm) were retained for subsequent

analyzes. For each treatment background, sham animals were used to calculate the

mean intensity for every aromatic bucket, providing treatment-matched reference

values. Each STNx (± Abx) sample was then compared with its corresponding sham

reference by determining, for each bucket, whether its intensity exceeded the sham

mean. The proportion and number of aromatic buckets classified as increased were

calculated per sample and statistically compared between groups using Mann Whitney

U test.

Liquid chromatography-mass spectrometry

Tryptophan panel

Tryptophan and its downstream metabolites were quantified in serum and plasma

samples. If not stated otherwise, the sample preparation was applied as stated

previously[57]. Briefly, 10 µL of an aqueous stable-isotope labelled internal standard

mixture was added to 50 µL of serum/ plasma. For protein precipitation samples were

diluted in a ratio of 1:5 with 100% cold methanol, vortexed, and stored at -80 °C for

24 h. Samples were centrifugated at 10,000×g at 4 °C for 5 min. The supernatant was

collected. The protein pellet was washed twice by addition of 200 µL of cold 80%

MeOH and subsequent centrifugation at 10,000×g (washing step 1) and 12,000×g

(washing step 2) at 4 °C for 5 min. For each sample, all supernatants were combined

and dried using a vacuum evaporator (CombiDancer, Hettich AG, Bäch, Switzerland).

All samples were reconstituted in 100 µL in water.

An ExionLC-30AD HPLC system (AB Sciex, Germany, Darmstadt) was used for the

chromatographic separation of metabolites on an ACQUITY Premier HSS T3, 1.8 µm,

2.1 × 150 mm reversed-phase column (Waters, Eschborn, Germany) reversed-phase

column. Gradient elution was performed with mobile phase A consisting of 0.1% formic

acid in water and mobile phase B of 0.1% formic acid in acetonitrile. Metabolite

detection was performed with a TripleQuad6500$^+$ (AB Sciex, Germany, Darmstadt)

performing positive/ negative-switching for the ionization. Peak integration and data

evaluation were carried out using SciexOS-MQ Software (Version 2.1.6, AB Sciex,

Germany, Darmstadt). Results were normalized to sample volumes.

3-Indoxyl sulfate quantification

For the HFD/L-NAME trial, frozen plasma samples were thawed at RT and then 70 µL

of each sample was mixed with 210 µL of a cold methanolic solution of the internal

standard d_5-IxS (Toronto Research Chemicals, Toronto, Canada), concentrated to

1 µmol/L . Proteins were precipitated by vortexing for 1 min and subsequent storage

for 60 min at -20°C. Supernatants were obtained by centrifugation at 9,400 g at 4°C

for 10 min and subjected to LC-MS/MS IxS quantification applying the multiple reaction

monitoring (MRM) method. Chromatographic separation was achieved on a 1290

Infinity II HPLC (Agilent Technologies, Waldbronn, Germany) equipped with a

Poroshell 120 EC-C18 column (3.0 x 150 mm, 2.7 µm; Agilent Technologies) guarded

by a pre-column (3.0 x 5 mm, 2.7 µm) of identical material. Water (eluent A) and

acetonitrile (eluent B), both acidified with 0.1% formic acid, were pumped with

0.4 mL/min. Elution of IxS and its internal standard was achieved with a 10-min linear

gradient from 5% to 90% eluent B. Total run-time was 16 min including re-equilibration

of the LC system. MS/MS analyzes were carried out using an Ultivo (6465B) triple-

quadrupole mass spectrometer (Agilent Technologies) operating in the negative

electrospray ionization mode (ESI-). The following ion source parameters were set:

sheath gas temperature, 400°C; sheath gas flow, 12 l/min of nitrogen; nebulizer

pressure, 50 psi; drying gas temperature, 100°C; drying gas flow, 9 l/min of nitrogen;

capillary voltage, 2.0 kV; nozzle voltage, 0.4 kV. The following mass transitions were

recorded (fragmentor voltage [FV] and collision energies [CE] in parentheses): IxS:

m/z 212.0 → 79.8 (FV: 102 V, CE: 28 eV, quantifier), m/z 212.0 → 80.9 (FV: 102 V,

CE: 16 eV), m/z 212.0 → 131.9 (FV: 102 V, CE: 20 eV), d_5-IxS: m/z 217.0 → 79.8 (FV:

82 V, CE: 28 eV, quantifier), m/z 217.0 → 80.9 (FV: 82 V, CE: 16 eV), m/z 217.0 →

137.0 (FV: 82 V, CE: 20 eV). Peak areas were determined with MassHunter

Quantitative Analysis software (version 10.1, Agilent Technologies) and IxS was

directly quantified via its internal standard d_5-IxS that was concentrated to 750 nmol/L

in the samples.

Transdermal measurement of glomerular filtration rate

FITC-sinistrin was reconstituted with PBS (40 mg/mL) and stored at -20 °C. Mice were anaesthetized with isoflurane and prone-placed on a heat pad. After shaving, the GFR monitor was attached on the right side of the back below the ribs. The measurement was recorded for 2-3 minutes prior to intravenous FITC-sinistrin injection (60 mg/kg) via either the tail vein or retro-orbital injection to set a steady baseline. Mice were then allowed to recover from anesthesia for the measurement of FITC decay (1-2 hours). Transcutaneous GFR was calculated from FITC-sinistrin half-time using a 3-compartment model (MBLab2, Medibeacon).

Clinical chemistry

Clinical parameters (plasma CRP, Cystatin C) were analyzed using an AU480 clinical chemistry analyzer (Beckman Coulter).

Echocardiography

Echocardiography was performed according to a standard operation protocol[58]. Briefly, mice were examined on a Vevo 3100 Imaging System equipped with a 30 MHz linear transducer (MX400; FUJIFILM VisualSonics Inc., Canada). Anaesthesia was induced by 3% isoflurane (in 80% oxygen). For image acquisition, isoflurane concentration was reduced to 1-1.5%, and adjusted to maintain comparable heart rates. All images were analyzed using Vevo LAB analysis software (FUJIFILM VisualSonics Inc.). Systolic function and cardiac dimensions were assessed using B-mode and M-mode echocardiography obtained from the parasternal long-axis view and the parasternal short-axis view at the mid-papillary level. Diastolic function was evaluated using conventional pulsed-wave Doppler to measure transmitral flow in the apical four-chamber view. Tissue Doppler recordings from the septal mitral annulus were used to assess myocardial relaxation velocity. The following parameters were analyzed: E/A (early diastolic filling velocity to atrial contraction), E/e′ (early filling velocity to early diastolic annular motion), and IVRT (isovolumetric relaxation time). Strain and M-mode measurements were derived from three images per parameter, with three cardiac cycles per image, and subsequently averaged. Diastolic Doppler measurements were obtained from three independent images, each with three

650 repeated measurements. Speckle-tracking analysis was performed using parasternal
651 long-axis and short-axis images, with epicardial and endocardial contours used to
652 quantify longitudinal, circumferential, and radial deformation[59]. Echocardiographic
653 parameters for each trial are summarized in Tables S1–S3.

Cardiac histology

Hearts were immediately collected after sacrifice. Cardiac tips were snap-frozen at
-80 °C for RNA isolation. The remaining tissue was either fixed in 10% neutral buffered
formalin solution (Merck) for 36 h at room temperature (RT) and then switched to 1%
until further use, or fixed in 4% paraformaldehyde (PFA) in PBS for 5 h at 4 °C.

For Picrosirius Red staining (PSR), formalin-fixed heart tissues were embedded in
paraffin and transversally sectioned (2 μm) using an HM 340E Rotary Microtome
(Epredia). Prior to staining with PSR, tissue sections were deparaffinated, rehydrated,
washed and incubated in PSR working solution (Morphisto) according to
manufacturer's protocol, with the exception of using ROTIClear (Carl Roth) as a xylol
substitute. After washing and dehydration, samples were mounted in water-free
medium (ROTIMount, Carl Roth, HP68.1).

For IHC stainings, PFA-fixed cardiac tissue was washed in PBS and incubated in 15%
sucrose in PBS for 6-12h and 30% sucrose for 6-12h under constant rotation at 4°C
for cryoprotection. Cardiac tissue was frozen embedded in Tissue-Tek O.C.T.
Compound (Sakura Finetek) in -40°C methyl butane and stored at -80% until further
use. Hearts were cut into 7 μm cryosections on a Cryostat Microtome (CMC3050 S,
Leica Biosystems). Prior to staining, cryosections were re-hydrated in Tris-buffered
saline (TBS) for 10 min at RT and subsequently permeabilized in TBS with 0.1%
Tween 20 (TBST) for 5 min at RT. Blocking was performed with 5% bovine serum
albumin (BSA) and 10% normal donkey serum (NDS) in TBST for 20 min at RT.

For wheat germ agglutinin (WGA) staining, cryosections were incubated with WGA
Fluorescein (1:800, Vector Laboratories) and DAPI staining solution (1:1000, Biotium)
in TBST + 5% NDS for 2 h at RT. For extracellular matrix (ECM) protein staining,
cryosections were incubated with primary antibodies, rabbit anti-fibronectin (1:100,
polyclonal, abcam) and goat anti-type I collagen (1:50, polyclonal, Southern Biotech),
in TBST + 5% NDS overnight at 4 °C. After washing in TBST, cryosections were

incubated with secondary antibodies, donkey anti-rabbit IgG Alexa Fluor Plus 488
(1:1000, Thermo Fisher) and anti-goat IgG Alexa Fluor Plus 555 (1:1000, Thermo
Fisher), together with DAPI (1:1000, Biotium) in TBST + 5% NDS for 1 h at RT. After
washing in TBS, slides were mounted with DAKO fluorescent mounting medium.

Slides were imaged using the PANNORAMIC™ MIDI II digital slide scanner
(3DHistech) with a 20x objective. For each heart, six representative snapshots were
taken at 20x magnification using CaseViewer (v2.2.1, 3DHistech) for downstream
analysis.

For PSR and ECM stainings, the percentage of positively stained area was quantified.
Snapshots were pre-processed using Fiji to generate a binary image of stained versus
unstained areas, and quantification was performed using CellProfiler. For WGA
staining, the size of the unstained area was quantified. After pre-processing in Fiji,
each snapshot was segmented in Ilastik by classifying pixels as stained or unstained
using an interactive machine-learning algorithm. The size of segmented objects
(unstained areas) was subsequently quantified using CellProfiler. The mean value
across all snapshots was calculated per animal.

Telemetry

In a subset of mice (n= 7 STNx, n= 7 STNx Abx) subcutaneous radiotelemetry
transmitters (PA-C10 BP, DSI data science international, St. Paul, MN, USA) were
implanted one week prior to the administration of antibiotics. Due to the length of the
study, we measured blood pressure on a weekly 3-day period (except w2, which had
a 2-day measurement). Means and SD per each 3-day measurement period were
calculated and values with an absolute SD > 5 were excluded from downstream
analysis. Longitudinal curves were generated by a non-parametric, locally weighted
scatterplot smoothing (LOESS) regression. For statistical analysis, systolic and
diastolic blood pressure AUCs were computed and divided by the total elapsed
recording time (in minutes) to yield average blood pressure per minute. Group
differences were assessed using a Wilcoxon test. Of the 14 mice initially implanted
with radiotelemetry transmitters, 5 mice died perioperatively and were excluded. An
additional 3 mice were removed prior to analysis due to telemetry signal instability and
inconsistent blood pressure recordings. Thus, only animals that survived surgery and

maintained stable telemetry signals throughout the study (n=3 per group) were
included in the final analysis.

Wire myography

From a subset of animals (n=5 STNx, n=8 STNx + Abx), mesenteric arteries were
removed, immediately transferred to cold (4°C) oxygenated (95% O_2/ 5% CO_2)
physiological salt solution (PSS: 119 NaCl, 4.7 KCl, 1.2 KH_2PO_4, 25 $NaHCO_3$, 1.2
Mg_2SO_4, 11.1 glucose, and 1.6 $CaCl_2$ in mmol/L), and dissected into 2-mm rings (n=2
per animal). Perivascular fat and connective tissues were removed. Each ring was
positioned between two stainless steel wires (0.0394-mm dia.) in a 5-mL PSS-filled
organ bath of Small Vessel Myograph (DMT 630MA, Danish Myo Technology, Aarhus,
Denmark). The PSS was continuously oxygenated and kept at 37°C (pH 7.4).
Mesenteric rings were placed under a tension equivalent to that generated at 0.9 times
the diameter of the vessel at 100 mm Hg. This normalization procedure was performed
to obtain the passive diameter of the vessel at 100 mm Hg. The software LabChart 8
was used for data acquisition and display. The rings were precontracted with the KCl
buffer (119 NaCl, 60 KCl, 1.2 KH_2PO_4, 25 $NaHCO_3$, 1.2 Mg_2SO_4, 11.1 glucose, and
1.6 $CaCl_2$ in mmol/L) and equilibrated until a stable resting tension was acquired. To
test the endothelial function, vessels were precontracted with 1 µmol/L phenylephrine.
After all vessel segments had reached a stable contraction plateau, increasing doses
of acetylcholine were administered to the organ baths (3 to 1000 nmol/L). Tension is
expressed as a percentage of the steady-state tension (100%) obtained with
phenylephrine.

Immunophenotyping

<u>Plasma collection, immune cell isolation and flow cytometry measurements</u>

Whole blood was collected in lithium-heparin syringes (80 IU/3 mL, BD Vacutainer A-
Line) for anticoagulation and centrifuged at 1,500 g for 15 min at RT. Plasma was
collected and frozen at -80 °C, and the remaining blood cell pellet was resuspended
in erythrocyte lysis buffer (83 g/L NH_4Cl, 8.5 g/L $NaHCO_3$, 10 mM EDTA) and
incubated at 37 °C for 6 min for peripheral blood cell (PBC) isolation. After
centrifugation at 400 g for 5 min (RT), supernatant was discarded and cells underwent

a second erythrocyte lysis step (37 °C, 4-6 min). Reaction was stopped using ice-cold

PBS + 0.5 % BSA + 2 mM EDTA (referred to as FACS buffer) and cell suspension was

filtered through a 30 µm strainer.

Spleens were excised and kept in FACS buffer at 4 °C. Single cell suspensions were

obtained by pressing tissue through 70 µm EasyStrainer, followed by erythrocyte lysis

and filtration through 40 µm EasyStrainer.

Hearts were excised, atria were removed and tissue was kept in HBSS without

Ca^{2+}/Mg^{2+} + 10 mM HEPES (referred to HBSS w/o) at 4 °C. Ventricles were cut into 2-

3 mm pieces before transfer into gentleMACS C Tubes (Miltenyi Biotec). A fresh

digestion mix was prepared with DNase I (30 U/mL), Hyaluronidase I-S (60 U/mL),

Collagenase XI (125 U/mL) in 1 mL HBSS with Ca^{2+}/Mg^{2+}. Samples were incubated

for 1 h at 700 rpm and 37 °C, followed by mechanical dissociation using the

gentleMACS Octo Dissociator (Miltenyi Biotec, program m_heart_02). Digestion was

quenched using ice-cold PBS + 10 % FBS. Cells were pelleted (400 g, 10 min, 4 °C),

treated with erythrocyte lysis buffer for 3 min at RT, neutralized with FACS buffer, and

filtered sequentially through 70 µm and 40 µm EasyStrainers. An aliquot was collected

for absolute counting (1:100 dilution).

Intestines were excised, flushed with HBSS w/o to remove all contents, divided into

small intestine (duodenum, jejunum, ileum) and large intestine (cecum, colon) and

stored in HBSS w/o at 4 °C. Lamina propria cells were isolated using the Lamina

Propria Dissociation Kit, mouse (Miltenyi Biotec).

Heart and lamina propria cell suspensions were subjected to a 40%/80% Percoll

density gradient (centrifugation at 800 g, 20 min, RT, no brake). Immune cells were

harvested at the interphase and washed in FACS buffer.

Splenocyte and lamina propria cell suspensions were counted on a LUNA-FL Dual

Fluorescence Cell Counter (Logos Biosystems), centrifuged and resuspended at

$1x10^6$ cells per 100 µL medium (RPMI 1640 + 10% FBS + 1% penicillin/streptomycin

[P/S, 10,000 U/mL / 10,000 µg/mL]) for further analysis. PBC and heart cell

suspensions were directly centrifuged and resuspended in 100 µL medium per flow

cytometry panel.

Cells were either immediately stained for flow cytometry or restimulated with 50 ng/mL

PMA, 500 ng/mL ionomycin, and 0.75 µL/mL GolgiStop in medium for 4 h at 37 °C, 5 %

CO_2. Viability was assessed with Live/Dead Fixable Aqua for 405 nm (Thermo Fisher),

surface markers were stained in Brilliant Stain Buffer (BD Biosciences) with Fc block

(Miltenyi Biotec), then cells were fixed/permeabilized (eBioscience

FoxP3/Transcription Factor Buffer Kit) for intracellular staining. All antibody

incubations were 30 min at 4 °C (Table S7). Data were acquired on a BD LSRFortessa

with high throughput sampler (HTS) using BD FACSDiva and analyzed in

FlowJo 10.8.1.

<u>Flow cytometry analysis</u>

Immune cell populations were identified by flow cytometry using the gating strategies

in Figure S9. For the immune cell abundance analysis, batch effects arising from

different flow cytometry acquisition days were corrected using ComBat from the *sva* R

package (v.3.20). Raw abundance values were imported into R and the matrix of cell-

type frequencies was extracted. Batch identifiers were supplied as the batch variable,

and group assignment was included as a covariate in the ComBat model matrix.

ComBat was run with parametric priors and without generating prior plots, yielding

batch-adjusted abundance values that were re-combined with sample metadata for

downstream analyzes. All subsequent analyzes were performed on the batch-

corrected dataset using the same gating-based cell populations described above.

UMAP visualizations were generated from the T-cell panel using two representative

samples per group. $CD3^+$ T cells were isolated using the hierarchical gating strategy,

and all gated events (10,000 $CD3^+$ T cells per sample) were used for dimensionality

reduction. UMAP embeddings were computed in FlowJo using the UMAP_R plugin

(version 4.1.1) with Euclidean distance, minimum distance = 0.1, and 15 nearest

neighbours. FlowJo's default logical transformation was applied to all fluorescence

channels. Group-level embedding intensity plots were created in FlowJo by overlaying

all cells from the same group and visualizing them as smoothed density distributions.

Mouse AhR reporter assay

Mouse AhR-responsive reporter cell line (H1L1.1c2), derived from mouse hepatoma

(Hepa1c1c7) cells and stably transfected with the pGudLuc1.1 vector containing AhR-

responsive dioxin response elements (DRE) upstream of a luciferase gene, was kindly

provided by Prof. Michael S. Denison, University of California, Davis, CA, USA.
H1L1.1c2 cells were cultured in Opti-MEM (Thermo Fisher, 31985070) supplemented
with 1% FBS (Merck, F7524), 1% P/S, and 0.2 mg/mL G418 (Sigma, A1720-1G), and
maintained at 37 °C with 5% CO_2. Subculture was performed twice a week using
TrypLE™ Express Enzyme (Gibco, 12604-02).

For the assay, cells were seeded in 96-well plates at a density of $7.5x10^4$ cells/well
and allowed to attach for 24 h. Cells were then incubated with 10% mouse plasma in
the same medium for 24 h. Assays were performed in triplicate, and 6-formylindolo[2-
b]carbazole (FICZ) was included as a positive control at a concentration of 10 nM.
After treatment, luciferase activity and cell viability were quantified using the One-
Glo™ + Tox Luciferase Reporter and Cell Viability Assay (Promega, E7120) according
to the manufacturer's instructions. Luminescence and fluorescence were measured
using a microplate reader (Infinite 200 plate reader, Tecan). Luminescence values
were normalized to cell viability (fluorescence signal) and subsequently expressed
relative to the FICZ control within each plate.

T_H17 polarization assay, murine

Mesenteric and peripheral lymph nodes (LN) were harvested from 129S2/SvPasOrlRj
mice and kept in cold FACS Flow buffer. LN were mechanically dissociated through a
70 μm cell strainer using cold FACS Flow buffer. The cell suspension was
subsequently filtered through a 40 μm strainer, washed with 20 mL FACS Flow buffer,
and centrifuged at 400 × g for 10 minutes at 4°C. Cells were resuspended in FACS
Flow buffer at $1 × 10^7$ cells per 100 μL.

CD4+ T cells were enriched via magnetic-activated cell sorting (MACS) using the
MojoSort Mouse CD4 T Cell Isolation Kit (BioLegend) with minor modifications for the
use of magnetic columns. Briefly, cells were incubated with a biotinylated antibody
cocktail (2 μL per 10^7 cells in 100 μL) for 15 minutes at 4°C, followed by Streptavidin
Nanobeads (2 μL per 10^7 cells) for an additional 15 minutes at 4°C. Cells were
separated using LS columns (Miltenyi Biotec) and the unlabelled fraction was used for
further analysis. For naïve CD4+ T cell isolation, cells were stained with CD4
(VioGreen), CD25 (PE), CD44 (BB700), CD62L (BV421), and a live/dead stain

(Zombie Green) and CD4$^+$CD25$^-$CD44^{low}CD62L^{high} cells were sorted on a FACS
Melody (BD Bioscience, 100 μm nozzle).

96-well flat-bottom plates were coated overnight at 4°C with anti-CD3 antibody (2
μg/mL in PBS, 50 μL per well). Naïve CD4+ T cells were resuspended in T$_H$17
differentiation medium (RPMI-1640 supplemented with 10% FCS, 1% penicillin-
streptomycin, 1% L-glutamine, 1% non-essential amino acids, 1 mM sodium pyruvate,
and 0.01 mM β-mercaptoethanol). Cells were plated at 5×10^4 cells per well in a final
volume of 200 μL with anti-CD28 (2 μg/mL), mIL-6 (80 000 U/mL, Miltenyi Biotec),
rhTGFβ1 (40 U/mL, Miltenyi Biotec) and mIL-1β (33 600 U/mL, Miltenyi Biotec) and
test compounds (IxS, KCl, AhRi at concentrations indicated in the respective figure) or
DMSO vehicle control, as appropriate. Cells were cultured at 37°C with 5% CO$_2$ for 4
to 5 days and subsequently restimulated for 4 hours with PMA (50 ng/ mL), ionomycin
(750 ng/mL), and GolgiPlug (0.5 μL/mL). Following restimulation, cells were washed
with FACS Flow buffer, stained with BD Horizon Fixable Viability Stain 700 (1:500
dilution) for 30 minutes cold, fixed, and permeabilized using Fix/Perm buffer
(eBioscience), and stained with the antibodies indicated in Table S7. Cells were
analyzed using a BD LSRFortessa with high throughput sampler (HTS). Data were
processed using FlowJo.

RNA isolation

Splenocytes were isolated as described above, enriched for CD45+ using mouse
CD45 MicroBeads (Miltenyi Biotec). Hearts were excised and apex was snap frozen
at -80°C. Spleens were excised and a small piece of tissue was placed into 300 μL
RNAprotect Tissue Reagent (Qiagen) at 4 °C for a minimum of 24 h. RNAprotect
Tissue Reagent was then discarded, and tissues were frozen at -80 °C.

RNA isolation from splenocytes and spleen was performed using the RNeasy Mini Kit
(Qiagen). RNA isolation from heart was performed using the RNeasy Fibrous Tissue
Mini Kit (Qiagen). For cell homogenisation, enriched CD45+ splenocytes were
resuspended in buffer RLT + β-mercaptoethanol and vortexed vigorously. Tissue
pieces were immersed in RLT buffer + β-mercaptoethanol with 1.5 mm ceramic beads
and placed into a Precellys 24 tissue homogeniser. RNA was subsequently isolated
from both sample types according to the instructions of the respective kits.

Fibroblasts on well plates were washed in PBS and incubated with 700 µL Qiazol Lysis

Reagent (Qiagen) per well at -80 °C over night. Then, lysed cells were harvested,

transferred to a tube and vortexed vigorously. 200 µL of chloroform was added and

samples were centrifuged at 12,000 g for 15 min at 4 °C. Afterwards, the aqueous

phase was collected and RNA was isolated with the RNeasy Mini Kit according to the

manufacturer's protocol.

Quantitative polymerase chain reaction (qPCR)

All qPCR was performed on a QuantStudio™ 5 Real-Time PCR System (Thermo

Fisher Scientific) using a 384-well plate format. For each 5-µL reaction mix, 4 ng of

DNA or cDNA was used, and all detections were performed in triplicate. All primer and

probe sequences and concentrations are listed in Table S6.

For all SYBR-Green based detection, the PowerUP™ SYBR™ Green Master Mix

(Thermo Fisher Scientific) was used with a fast cycling protocol of 50 °C for 2 min, 95

°C for 2 min, followed by 40 cycles of 95 °C for 1 s and 60 °C for 30 s. For all TaqMan-

based detection, the TaqMan® Multiplex Master Mix (Thermo Fisher Scientific,

4461882) was used with a fast cycling protocol of 95 °C for 20 s, followed by 40 cycles

of 95 °C for 2 s and 60 °C for 20 s.

For *16S* quantification, cecal DNA was amplified and detected using SYBR Green and

the standard curve method. Standard curves were generated using 10-fold serial

dilutions ranging from 10^1 to 10^9 copies of the *16S* rRNA gene of *E. coli* (Invitrogen)

amplified with primers 27F (5′-GTTTGATCCTGGCTCAG-3′) and

1492R (5′-CGGCTACCTTGTTACGAC-3′). Using the resulting curve, *16S* rDNA copy

numbers per gram content were calculated.

For all other genes, cDNA was synthesized prior to qPCR using the High Capacity

cDNA Reverse Transcription Kit (Thermo Fisher Scientific). Murine *Col1a2*, *Acta2*,

Cyp1a1 and *18s* were detected using TaqMan. Gene expression was normalized to

the *18s* signal using the ΔΔCt method. All other murine genes were quantified using

SYBR Green, and *Hprt* or *Gapdh* as the housekeeping genes. All human genes were

quantified using SYBR Green and normalized to *HPRT1* using the ΔΔCt method. All

primer and probes are shown in Table S6.

Bulk mRNA sequencing

Illumina stranded mRNA library preparation and bulk mRNA sequencing were performed at the BIH/MDC Genomics Technology Platform. Paired-end sequencing was performed on an Illumina NovaSeq X Plus platform, targeting ~30 million reads per sample. Transcript quantification of raw sequencing data to the *Mus musculus* reference transcriptome (GRCm38) was performed using Salmon (v1.10.1). Quality assessment of the preprocessing was performed using MultiQC (v1.6). Following quality control, 26.4M-34.7M paired-end reads per sample were retained and included in downstream analysis. Gene symbols were generated from Ensembl gene IDs using biomaRt (v2.52.0, Ensembl release 102). Post-processing quality control and downstream data analysis were conducted using DESeq2 (v1.42.1).

For the Abx trial, separate DeSeq2 datasets were generated for each reference condition (Sham, STNx). Lowly expressed genes were removed by filtering for transcripts with at least 1 count per million in a minimum of two samples using edgeR (v4.0.16). Differential expression analysis was performed using the DeSeq2 pipeline. Log$_2$ fold changes were adjusted using lfcShrink with the ashr method (v2.2.63), and p-values were corrected using a false discovery threshold of 10%. Variance-stabilized expression values were obtained using the rlog transformation.

Gene set enrichment analysis (GSEA) was performed on the overlap of significantly regulated genes from the STNx vs. Sham and STNx Abx vs. STNx comparisons (adjusted p-value < 0.1). Curated *Mus musculus* gene sets were downloaded from MSigDB. Enrichment was calculated with Fisher's exact test and FDR correction, using all Ensembl gene IDs present in the complete MSigDB *Mus musculus* dataset as the background. Pathways and GO terms not relevant to the study were excluded from visualisation using targeted keyword filtering.

For the AhRi trial, differential expression analysis with DeSeq2 was restricted to the set of DEGs in STNx that were Abx responsive. Batch effects between experimental rounds were corrected using ComBat-seq (sva, v3.50.0). Genes significantly regulated by AhRi-treatment in the same direction as the Abx-dependent response (p < 0.1) were subjected to GSEA.

934 For both trials, targeted gene set variation analysis (GSVA) was performed on rlog-
935 transformed expression values using curated gene sets relevant to the T_H17 axis (gsva
936 package, v1.50.5).

Single-cell RNA-sequencing

We analyzed publicly available mouse cardiac single-nucleus and human single-cell
RNA-seq datasets[32,37]. Processed Seurat objects were imported into R (v4.2.2) using
Seurat (v5.0). No reclustering or reannotation was performed. For each dataset, cells
belonging to diseased samples (mouse CKD or human CKD) and appropriate controls
were subset for downstream analyzes. Resident structural cell types were selected
based on the original annotations, and expression of IL17ra was quantified from log-
normalized RNA values to identify positive and negative stromal populations.
Extracellular matrix (ECM) activity was assessed using an extended NABA matrisome
gene list[33] and compared using Wilcoxon tests. AhR pathway activity was similarly
scored using a recently published AhR-responsive gene set[60]. Differential expression
was performed using the MAST framework[61], including only genes expressed in more
than 10% of nuclei. Genes were considered differentially expressed when meeting:
adjusted $p < 0.1$, absolute $\log_2$ fold change > 0.3, and average normalized expression
> 3 in Il17ra$^+$ cells. Differentially expressed genes were compared with curated
fibroblast transcriptional signatures derived from human cardiac and stromal
datasets[34] to identify enriched transcriptional programs. Briefly, DEG were intersected
with each fibroblast signature and the number of overlapping genes, proportion of
signature genes present among upregulated genes, and average FC and expression
of the overlapping genes were used to find similar transcriptional programs. Cytokine-
response enrichment was conducted from DEG of cardiac lymphocytes as described
above using the Immune Response Enrichment Analysis (IREA)[62] software.

Murine cardiac fibroblasts

Hearts were freshly harvested from 7-10 weeks old male and female C57BL/6J mice
and rinsed twice in PBS. Aortic and atrial tissues were removed, and ventricles were
minced and transferred into 1 mL of enzymatic digestion solution containing DNase I
(300 U/mL), Hyaluronidase I-S (60 U/mL), Collagenase XI (125 U/mL), and

Collagenase I (450 U/mL) in DMEM. Digestion was performed on a shaking incubator
(800 rpm, 37 °C) for 45 minutes.

Following digestion, the cell suspension was filtered through a 70 µm strainer using
PBS and centrifuged at 400 g for 10 minutes at 37 °C. The pellet was resuspended in
1 mL erythrocyte lysis buffer (Erylysis), incubated for 3 minutes at RT, and quenched
with PBS. Cells were centrifuged again under the same conditions, supernatant was
discarded, and the pellet was resuspended in 4 mL fibroblast culture medium (DMEM
supplemented with 20% FBS, 2% penicillin-streptomycin, 1 ng/mL endothelial cell
growth factor [ECGF], and 10 µg/mL heparin). Cells were maintained at 37 °C and 5%
CO_2. Media was changed after 2.5 h and further replaced every 2-3 days. For
subculture, fibroblasts were detached using Trypsin-EDTA (1mL, 2-3 min incubation).
Reaction was stopped with culture media, fibroblasts were transferred to a 15 mL tube
and centrifuged at 400 g for 10 min. Pellet was resuspended in culture media, counted
and plated into tissue flasks or well plates depending on cell counts (6.000-10.000
cells/cm^2).

Assay was performed on a 6-well plate. In brief, fibroblasts were starved in culture
media containing 1% FBS and incubated for 4 h at 37 °C, followed by a 24 h incubation
with the test substances. Murine recombinant IL-17A was used at 1 ng/mL in culture
media. Potassium 3-indoxyl sulfate was used at 50 µM and 100 µM on its own, and at
100 µM in combination with IL-17A in culture media. After 24 hours incubation,
fibroblasts were washed in PBS and incubated with 700 µL Qiazol Lysis Reagent per
well at -80 °C over night. Lysed fibroblasts were then harvested, RNA was isolated
and RT-qPCR was performed as described above.

For immunofluorescence staining, 18x18 mm coverslips were sterilized with ethanol
70% and placed into 6-well plates. Fibroblasts were grown on the coverslips in culture
media until desired confluence. Cells were then washed with PBS and fixed with 4%
Paraformaldehyde, followed by permeabilization with 0.1% Triton X-100 and blocking
with 10% normal donkey serum (NDS) in PBS.

For PDGFRα and VIM staining, cells were incubated with conjugated antibodies (anti-
mouse CD140a PE, 1:100, clone APA5, Miltenyi, 130-102-502 and AF647 anti-
vimentin, 1:500, clone EPR3776, ab194719) for 2 h at RT in PBS + 10% NDS. For
COL-1 and IL17RA staining, cells were incubated with primary antibodies (goat anti-

type I collagen, 1:500, polyclonal, Southern Biotech, 1310-01 and rabbit anti-IL17RA, 1:500, polyclonal, abcam, ab180904) for 2 h at RT in PBS + 10% NDS. After washing with PBS + 1% NDS, secondary antibodies, donkey anti-rabbit IgG Alexa Fluor Plus 488 (1:1000, Thermo Fisher) and anti-goat IgG Alexa Plus 555 (1:1000, Thermo Fisher), were incubated together with DAPI (1:1000, Biotium) in PBS + 10% NDS for 1 h at RT. After washing, coverslips were taken out of the well plates and mounted onto microscopy slides using DAKO fluorescent mounting medium. Slides were imaged on the Keyence Compact Fluorescence microscope with the Plan Apochromat 20x objective lens.

For flow cytometry staining, cells were stained for PDGFRα, CD45 and CD31 (antibody concentrations in Table S7) together with SYTOX^{TM} Green Dead Cell Stain (1:500, Invitrogen, S34860) and Fc blocking reagent (Miltenyi Biotec, 130-092-575) in FACS buffer. Data were acquired on a BD LSRFortessa using BD FACSDiva and analyzed in FlowJo 10.8.1.

Splenocyte incubation assay

Spleens from 129S2/SvPasOrlRj mice were processed to obtain spleen single-cell suspensions as described above. For *ex vivo* stimulation, splenocytes were incubated for 4 h at 37 °C, 5 % CO_2 in RPMI-1640 in the presence of IxS and AhRi at the indicated concentrations. After incubation, cells were resuspended in buffer RLT + β-mercaptoethanol and frozen at −80 °C for subsequent RNA isolation as described above.

Patient serum collection

Patients with CKD stage 5 dependent on hemodialysis were recruited Department of Nephrology and Medical Intensive Care at Charité-Universitätsmedizin Berlin. Healthy volunteers were recruited at the Experimental and Clinical Research Center (ECRC) of Charité-Universitätsmedizin Berlin. Ethical approval was obtained from the institutional review board of Charité–Universitätsmedizin Berlin (EA2/162/17) and all participants provided written informed consent before biomaterial was collected. Blood was obtained from patients at the start of hemodialysis. Serum was collected by

1029 centrifugation at 1500 g for 15 min at RT, and stored at -80 °C until further use.

1030 Demographic and clinical characteristics of the study cohort are summarized in Table

1031 S4.

1032

Human luciferase reporter assay

Human AhR reporter cell line (HT29-Lucia^{TM} AhR, InvivoGen, ht21-ahr), containing

AhR-responsive DRE upstream of a Lucia luciferase gene, was cultured in Dulbecco's

Modified Eagle Medium (DMEM) supplemented with 10% FBS and 100 µg/mL Zeocin

(InvivoGen, ant-zn-05), maintained at 37 °C with 5% CO_2. Cells were subcultured once

per week using a standard Trypsin-EDTA protocol and partially replenished with fresh

medium mid-week between passages.

For the assay, 180 µL of cell suspension per well was seeded on a 96-well plate at a

density of 2.8×10^5 cells/mL in medium without Zeocin. For the dose-response curve of

AhRi, cells were incubated with 100 µM IxS and AhRi concentrations between 1 nM

and 10 µM or VEH (vehicle, DMSO content of the highest AhRi concentration) for 24

h in quadruples. To assess the AhR activation potential of human biomaterial, cells

were incubated with 10% patient serum and AhRi in a final concentration of 1 or 0.1

µM, or VEH for 24 h in triplicates. For quantification, 20 µL of culture supernatant was

used to determine luciferase activity using the QUANTI-Luc^{TM} Luciferase Detection

Reagent (InvivoGen, rep-qlc4lg1). Luminescence was measured using a microplate

reader (Infinite 200 plate reader, Tecan).

1050

Human T$_H$17 polarization assay

Peripheral blood mononuclear cells (PBMC) were collected together with serum from

healthy volunteers under the ethics approval described above (EA2/162/17). PBMC

were isolated from EDTA-anticoagulated blood of a healthy donor by diluting samples

1:1 in PBS + 2% FBS and layering them into 50 mL SepMate tubes (Stemcell) prefilled

with 15 mL Pancoll (PAN Biotech). After centrifugation at 1200 g for 10 min at RT, the

mononuclear cell layer was collected, transferred into a fresh 50 mL tube, and washed

with PBS + 2% FBS, followed by erythrocyte lysis.

T cells were isolated from PBMC using MojoSort Human CD4 T Cell Isolation Kit (Biolegend) with LS columns (Miltenyi) following the manufacturer's protocol. To obtain naïve CD4 cells, the enriched T helper cell fraction was sorted using a BD FACSAria III (BD Bioscience, 70 µm nozzle) by staining for CD4, CD25, CD127, CD45RO (Table S8), and dead cells exclusion with DRAQ7 (1:250, Miltenyi Biotec).

Naïve CD4 T cells were resuspended in X-VIVO 15 serum-free hematopoietic media (Lonza Bioscience) and plated at 5×10^4 cells per well in anti-CD3-coated (10 µg/mL, BD) 96-well U-bottom plates and 1 µg/mL anti-CD28 (BD). To induce T_H17 polarization, naïve CD4 T cells were cultured with 12.5 ng/mL IL-1β, 25 ng/mL IL-6, 25 ng/mL IL-21, 25 ng/mL IL-23, and 5 ng/mL TGF-β (all Miltenyi Biotec), while T_H0 control wells were left without cytokine supplementation. Cells were treated with 2.5% serum from either hemodialysis patients or healthy controls in triplicates and maintained for 7 days at 37°C and 5% CO_2.

Prior to analysis, cells were restimulated for 4 h with 10 ng/mL PMA (Merck), 500 ng/mL ionomycin (Merck), and 1 µL/mL GolgiPlug (BD), stained with LIVE/DEAD fixable aqua dye (Invitrogen), fixed and permeabilized using the Foxp3/Transcription Factor Staining Buffer Set (Invitrogen), and finally stained with antibody panels with Fc Block, CD4 and IL-17A (antibodies listed in Table S8).

Cells were analyzed using a BD LSRFortessa with HTS and data were processed using FlowJo (v10.10).

UK Biobank

Study participants

The UK Biobank is a population-based cohort of 502,366 participants aged 40-69 years recruited between 2006 and 2010 across 22 assessment centers in England, Scotland, and Wales. Ethical approval was granted by the North West Multi-Centre Research Ethics Committee (reference 11/NW/0382), and all participants provided written informed consent. Data available as of 16 November 2023 were used for this study.

Participants with impaired kidney function were defined as those with an estimated glomerular filtration rate (eGFR) below 60 mL/min/1.73 m², corresponding to KDIGO

categories G3-G5. Demographic and clinical characteristics of the study cohort are
summarized in Table S5.

Identification of individuals with heart failure with preserved ejection fraction (HFpEF)
followed a previously published multi-stage algorithm[63]. All HFpEF cases were
required to exhibit at least one clinical symptom of heart failure in accordance with
European Society of Cardiology (ESC) guidelines, supplemented by additional criteria
depending on data availability. When cardiac magnetic resonance (CMR) imaging was
available, participants with a left ventricular ejection fraction above 49% were
considered to have preserved systolic function. When NT-proBNP measurements
were available, patients exceeding the 90th percentile were classified as HFpEF,
whereas those with lower values or evidence of reduced ejection fraction were
excluded. For participants lacking imaging or proteomic data, HFpEF was assigned
when the pre-test probability exceeded 90%.

<u>Data modalities</u>

All variables used in this study were assessed prior to or at the baseline visit. eGFR
was calculated from plasma cystatin C concentrations using the equation eGFR =
$74.835 / (cystatin\ C^{1.333})$. Albuminuria was quantified as the urine albumin-to-creatinine
ratio. Essential hypertension was defined by the presence of an ICD-10 I10 diagnosis
code recorded before baseline. Proteomics profiling was available for approximately
10% of the UK Biobank participants and was performed using Olink proximity
extension assays, with protein concentrations reported as log_2-normalized protein
expression (NPX) values. For the present analysis, we focused on NT-proBNP and IL-
17A, which were available for 51,578 and 50,573 participants, respectively. Smoking
exposure was quantified using the UK Biobank variable "pack years of smoking"
(Field-ID 20161), calculated from baseline questionnaire responses using the formula:
(cigarettes per day / 20) * (age stopped smoking - age started smoking). This measure
reflects cumulative smoking intensity, with higher values reflecting greater lifetime
exposure.

<u>Statistical Analysis</u>

Time-to-event outcomes were evaluated using Kaplan-Meier estimators implemented
in the Python package lifelines (version 0.30.0). Participants were stratified into three
IL-17A groups according to the plasma IL-17A concentrations: below the 25th

percentile, between the 25th and 75th percentiles, and above the 75th percentile. Survival time was defined as the interval from baseline to the occurrence of the event of interest, with censoring at the end of follow-up. Differences across IL-17A strata were assessed using the log-rank test.

To evaluate the association between IL-17A and clinical outcomes, Cox proportional hazards models were fitted. All covariates were standardized before model fitting to facilitate direct comparison of effect sizes. Cox proportional hazards models were estimated using an unpenalized framework with robust variance estimation. To ensure numerical stability, a limited grid of ridge penalisation terms (λ = 0 to 0.01) was evaluated, and the model yielding stable convergence without materially influencing effect estimates was selected. Hazard ratios (HRs) and 95% confidence intervals (CIs) were derived from the final model and displayed using forest plots.

Normality of continuous variables was assessed using the Shapiro-Wilk test. When both groups satisfied the normality assumption ($p > 0.05$), differences were evaluated using Welch's *t* test; otherwise, comparisons were performed using the non-parametric Mann-Whitney U test. For analyzes involving multiple statistical tests, p-values were adjusted using the Benjamini-Hochberg false discovery rate (FDR) procedure. All statistical analyzes were performed in Python 3.11.

Human cardiac fibroblasts

Human cardiac fibroblasts were obtained from PromoCell (Germany). Cells were seeded (4000 cells per cm^2 seeding density) with 18 mL of Fibroblast Growth Medium 3 (PromoCell, Germany) into a tissue flask T75. Media contained supplements (PromoCell, Germany), P/S (1%) and Amphotericin B (Gibco, 0.2%). Cells were maintained at 37 °C and 5% CO_2 with medium exchanged every 2-3 days and passaged at 70-90% confluence as described above. Subculture was performed using the standard Trypsin-EDTA protocol.

For all experiments, cells were seeded at manufacturer recommended densities and allowed to attach overnight. The following day, cultures were starved for 4h in medium containing 1% FBS and subsequently stimulated for 24h under the appropriate conditions. Treatments included IL-17A (1 ng/mL), IxS (100 µM), AhRi (1 µM), Secukinumab (Cosentyx, 10 nM), human serum (HD/HC; 10%), or the corresponding

combinations. After stimulation, cells were washed once with PBS and lysed in QIAzol
as descrived above. Lysed cells were then harvested, RNA was isolated and RT-
qPCR was performed as described above.

To assess viability and proliferation, fibroblasts were co-incubated of IL-17A and IxS
in starved medium for 24 h. After detaching, cells were stained with Zombie Green
Fixable Viability Kit (1:500, BioLegend), fixed/permeabilized (eBioscience
FoxP3/Transcription Factor Buffer Kit) for intracellular staining, and stained for Ki-67
(Table S8.)

1162

1163 **Data and code availability**

1164 All code, data, and scripts used in this study will be made publicly available. The code
1165 and scripts will be made accessible on Zenodo. The raw sequencing data will be
1166 deposited in the NCBI Sequence Read Archive.

Author contribution

CRediT: Conceptualization: AY, WA, AT, JH, HB, NW; Data curation: AY, WA, JVe, HA, MK; Formal Analysis: AY, WA, OP; Funding acquisition: VM, UK, AZ, PO, WG, KUE, DNM, MK, HB, NW; Investigation: AY, WA, AT, OP, FS, IAS, AMR, JVe, NG, VM, SH, SK, GT, HA, FF, CH, SAH, PMB, MIW, FB; Methodology: AY, WA, OP, FS, IAS, JVe, JVo, PO, WG, KD, MK, HB, NW; Project administration: AY, WA; Resources: UK, TK, BK, AZ, PO, WG, KD, KUE, DNM, MK, JH, HB, NW; Software: AY, WA, JVe, HA; Supervision: JVo, UK, TK, BK, AZ, PO, WG, KD, KUE, DNM, MK, JH, HB, NW; Validation: AY, WA, RK, MK; Visualization: AY, WA, JVe, HB, NW; Writing - original draft: AY, WA, HB, NW; Writing - review & editing: all authors.

Acknowledgment

We thank Melanie Röhr and Gudrun Koch for their technical assistance, Ilona Kamer for the implantation of the blood pressure telemetry devices and Heike Schwede for support with animal husbandry. We also thank Laura Mossier for contributing to the HFpEF trial, and Daniel Herrmann for his technical assistance with the IxS quantification. Clinical chemistry and echocardiography measurements were performed at the Preclinical Research Center of the MDC. Bulk mRNA-seq was performed at the BIH/MDC Genomics Technology Platform. Human T cells were sorted using the flow cytometry platform of the MDC. We thank the MDC Max Cluster for providing us with the scientific compute cluster resources and Ulrike Löber for bioinformatics support.

Funding

FACSMelody used in this study was funded by German Federal Ministry of Research, Technology and Space (BMFTR), project-ID: 01EJ2202A. V.M., D.N.M., W.G., P.J.O., H.B., and N.W. were supported by the BMFTR, TAhRget consortium (project-ID 01EJ2502A (N.W.), 01EJ2502B (W.G., P.J.O.), 01EJ2502E (V.M., D.N.M.), and 01EJ2502G (H.B.)), as well the QEED consortium (project-ID 13N16386, to H.B.). U.K., D.N.M., A.Z. and N.W. were supported by the German Research Foundation (DFG), CRC1470 (project-ID 437531118, subproject A10 to N.W., A06 to D.N.M., and A09 to U.K.), and CRC1525 (project-ID 453989101, to A.Z.) and project-ID 432915089

(to A.Z). D.N.M. and N.W. were supported by the DZGIF (DZG Innovation Fund), Topic

"Microbiome". N.W. was supported from the European Research Council (ERC) under

the European Union's Horizon 2020 research and innovation program (852796). N.W.

was supported by the Corona Foundation in the German Stifterverband

(S199/10080/2019). K.U.E. received funding from the European Union's Horizon 2020

research and innovation program under a Marie Skłodowska-Curie Innovative Training

Networks (H2020-MSCA-ITN-2019) grant (TrainCKDis, project-ID 860977).

1206

Conflict of interest

The authors declare no conflict of interest.

References

1   Langenberg, C., Hingorani, A. D. & Whitty, C. J. M. Biological and functional
multimorbidity—from mechanisms to management. *Nature Medicine* **29**, 1649-1657
(2023). https://doi.org/10.1038/s41591-023-02420-6

2   Evenepoel, P., Poesen, R. & Meijers, B. The gut-kidney axis. *Pediatr Nephrol* **32**,
2005-2014 (2017). https://doi.org/10.1007/s00467-016-3527-x

3   Eckardt, K.-U. *et al.* Evolving importance of kidney disease: from subspecialty to
global health burden. *The Lancet* **382**, 158-169 (2013).
https://doi.org/10.1016/S0140-6736(13)60439-0

4   Mishima, E. *et al.* Evaluation of the impact of gut microbiota on uremic solute
accumulation by a CE-TOFMS–based metabolomics approach. *Kidney International*
92, 634-645 (2017). https://doi.org/https://doi.org/10.1016/j.kint.2017.02.011

5   Mair, R. D., Sirich, T. L., Plummer, N. S. & Meyer, T. W. Characteristics of Colon-
Derived Uremic Solutes. *Clin J Am Soc Nephrol* **13**, 1398-1404 (2018).
https://doi.org/10.2215/cjn.03150318

6   Jankowski, J., Floege, J., Fliser, D., Böhm, M. & Marx, N. Cardiovascular Disease in
Chronic Kidney Disease. *Circulation* **143**, 1157-1172 (2021).
https://doi.org/doi:10.1161/CIRCULATIONAHA.120.050686

7   Speer, T., Dimmeler, S., Schunk, S. J., Fliser, D. & Ridker, P. M. Targeting innate
immunity-driven inflammation in CKD and cardiovascular disease. *Nature Reviews*
Nephrology **18**, 762-778 (2022). https://doi.org/10.1038/s41581-022-00621-9

8   Wilck, N. *et al.* Salt-responsive gut commensal modulates T(H)17 axis and disease.
Nature **551**, 585-589 (2017). https://doi.org/10.1038/nature24628

9   Granados, J. C. *et al.* AHR is a master regulator of diverse pathways in endogenous
metabolism. *Scientific Reports* **12**, 16625 (2022). https://doi.org/10.1038/s41598-
022-20572-2

10   Rothhammer, V. & Quintana, F. J. The aryl hydrocarbon receptor: an environmental
sensor integrating immune responses in health and disease. *Nature Reviews*
Immunology **19**, 184-197 (2019). https://doi.org/10.1038/s41577-019-0125-8

11   Dong, F. & Perdew, G. H. The aryl hydrocarbon receptor as a mediator of host-
microbiota interplay. *Gut Microbes* **12**, 1859812 (2020).
https://doi.org/10.1080/19490976.2020.1859812

12   Gutiérrez-Vázquez, C. & Quintana, F. J. Regulation of the Immune Response by the
Aryl Hydrocarbon Receptor. *Immunity* **48**, 19-33 (2018).
https://doi.org/10.1016/j.immuni.2017.12.012

13   Anandakumar, H. *et al.* Segmental patterning of microbiota and immune cells in the
murine intestinal tract. *Gut microbes* **16**, 2398126 (2024).
https://doi.org/10.1080/19490976.2024.2398126

14   Voroneanu, L. *et al.* Gut Microbiota in Chronic Kidney Disease: From Composition to
Modulation towards Better Outcomes-A Systematic Review. *J Clin Med* **12** (2023).
https://doi.org/10.3390/jcm12051948

15   Chaves, L. D. *et al.* Chronic kidney disease, uremic milieu, and its effects on gut
bacterial microbiota dysbiosis. *Am J Physiol Renal Physiol* **315**, F487-f502 (2018).
https://doi.org/10.1152/ajprenal.00092.2018

16   Bloom, P. P. *et al.* Microbiota and kidney disease: the road ahead. *Nature Reviews*
Nephrology **21**, 702-716 (2025). https://doi.org/10.1038/s41581-025-00988-5

17 Chen, T. K., Knicely, D. H. & Grams, M. E. Chronic Kidney Disease Diagnosis and
Management: A Review. *Jama* **322**, 1294-1304 (2019).
https://doi.org/10.1001/jama.2019.14745
18 Patel, N., Yaqoob, M. M. & Aksentijevic, D. Cardiac metabolic remodelling in chronic
kidney disease. *Nature Reviews Nephrology* **18**, 524-537 (2022).
https://doi.org/10.1038/s41581-022-00576-x
19 Kaptoge, S. *et al.* C-reactive protein concentration and risk of coronary heart disease,
stroke, and mortality: an individual participant meta-analysis. *Lancet* **375**, 132-140
(2010). https://doi.org/10.1016/s0140-6736(09)61717-7
20 Marques, F. Z., Mackay, C. R. & Kaye, D. M. Beyond gut feelings: how the gut
microbiota regulates blood pressure. *Nature Reviews Cardiology* **15**, 20-32 (2018).
https://doi.org/10.1038/nrcardio.2017.120
21 Avery, E. G. *et al.* Quantifying the impact of gut microbiota on inflammation and
hypertensive organ damage. *Cardiovasc Res* **119**, 1441-1452 (2023).
https://doi.org/10.1093/cvr/cvac121
22 Yang, B. H. *et al.* Foxp3+ T cells expressing RORγt represent a stable regulatory T-cell
effector lineage with enhanced suppressive capacity during intestinal inflammation.
Mucosal Immunology **9**, 444-457 (2016). https://doi.org/10.1038/mi.2015.74
23 Blatner, N. R. *et al.* Expression of RORγt marks a pathogenic regulatory T cell subset
in human colon cancer. *Sci Transl Med* **4**, 164ra159 (2012).
https://doi.org/10.1126/scitranslmed.3004566
24 Rothhammer, V. & Quintana, F. J. The aryl hydrocarbon receptor: an environmental
sensor integrating immune responses in health and disease. *Nature reviews.*
Immunology **19**, 184-197 (2019). https://doi.org/10.1038/s41577-019-0125-8
25 Quintana, F. J. *et al.* Control of T(reg) and T(H)17 cell differentiation by the aryl
hydrocarbon receptor. *Nature* **453**, 65-71 (2008).
https://doi.org/10.1038/nature06880
26 Veldhoen, M. *et al.* The aryl hydrocarbon receptor links TH17-cell-mediated
autoimmunity to environmental toxins. *Nature* **453**, 106-109 (2008).
https://doi.org/10.1038/nature06881
27 Delgobo, M. *et al.* Myocardial Milieu Favors Local Differentiation of Regulatory T
Cells. *Circulation research* **132**, 565-582 (2023).
https://doi.org/10.1161/CIRCRESAHA.122.322183
28 Hwang, S. J. *et al.* Indoxyl 3-sulfate stimulates Th17 differentiation enhancing
phosphorylation of c-Src and STAT3 to worsen experimental autoimmune
encephalomyelitis. *Toxicol Lett* **220**, 109-117 (2013).
https://doi.org/10.1016/j.toxlet.2013.04.016
29 Schiattarella, G. G. *et al.* Nitrosative stress drives heart failure with preserved
ejection fraction. *Nature* **568**, 351-356 (2019). https://doi.org/10.1038/s41586-019-
1100-z
30 Bansal, N., Zelnick, L. R., Scherzer, R., Estrella, M. & Shlipak, M. G. Risk Factors and
Outcomes Associated With Heart Failure With Preserved and Reduced Ejection
Fraction in People With Chronic Kidney Disease. *Circulation. Heart failure* **17**,
e011173 (2024). https://doi.org/10.1161/CIRCHEARTFAILURE.123.011173
31 Meijers, B. K. *et al.* p-Cresyl sulfate and indoxyl sulfate in hemodialysis patients. *Clin J*
Am Soc Nephrol **4**, 1932-1938 (2009). https://doi.org/10.2215/cjn.02940509

32   Kaesler, N. *et al.* Mapping cardiac remodeling in chronic kidney disease. *Sci Adv* **9**,
eadj4846 (2023). https://doi.org/10.1126/sciadv.adj4846

33   Naba, A. *et al.* The matrisome: in silico definition and in vivo characterization by
proteomics of normal and tumor extracellular matrices. *Mol Cell Proteomics* **11**,
M111.014647 (2012). https://doi.org/10.1074/mcp.M111.014647

34   Patrick, R. *et al.* Integration mapping of cardiac fibroblast single-cell transcriptomes
elucidates cellular principles of fibrosis in diverse pathologies. *Science Advances* **10**,
eadk8501 (2024). https://doi.org/10.1126/sciadv.adk8501

35   Barallobre-Barreiro, J. *et al.* Glycoproteomics Reveals Decorin Peptides With Anti-
Myostatin Activity in Human Atrial Fibrillation. *Circulation* **134**, 817-832 (2016).
https://doi.org/10.1161/CIRCULATIONAHA.115.016423

36   Kober, C. *et al.* Targeting the aryl hydrocarbon receptor (AhR) with BAY 2416964: a
selective small molecule inhibitor for cancer immunotherapy. *J Immunother Cancer*
11 (2023). https://doi.org/10.1136/jitc-2023-007495

37   Kuppe, C. *et al.* Decoding myofibroblast origins in human kidney fibrosis. *Nature* **589**,
281-286 (2021). https://doi.org/10.1038/s41586-020-2941-1

38   Batra, G. *et al.* Interleukin 6 and Cardiovascular Outcomes in Patients With Chronic
Kidney Disease and Chronic Coronary Syndrome. *JAMA Cardiol* **6**, 1440-1445 (2021).
https://doi.org/10.1001/jamacardio.2021.3079

39   Tonelli, M., Karumanchi, S. A. & Thadhani, R. Epidemiology and Mechanisms of
Uremia-Related Cardiovascular Disease. *Circulation* **133**, 518-536 (2016).
https://doi.org/doi:10.1161/CIRCULATIONAHA.115.018713

40   Hayer, M. K. *et al.* Defining Myocardial Abnormalities Across the Stages of Chronic
Kidney Disease: A Cardiac Magnetic Resonance Imaging Study. *JACC Cardiovasc*
Imaging **13**, 2357-2367 (2020). https://doi.org/10.1016/j.jcmg.2020.04.021

41   Bansal, N. *et al.* Burden and Outcomes of Heart Failure Hospitalizations in Adults
With Chronic Kidney Disease. *Journal of the American College of Cardiology* **73**, 2691-
2700 (2019). https://doi.org/10.1016/j.jacc.2019.02.071

42   Schulman, G. *et al.* Randomized Placebo-Controlled EPPIC Trials of AST-120 in CKD.
Journal of the American Society of Nephrology : JASN **26**, 1732-1746 (2015).
https://doi.org/10.1681/ASN.2014010042

43   Holle, J. *et al.* Inflammation in Children with CKD Linked to Gut Dysbiosis and
Metabolite Imbalance. *J Am Soc Nephrol* **33**, 2259-2275 (2022).
https://doi.org/10.1681/asn.2022030378

44   Collett, J. A. *et al.* IL-17A Levels and Progression of Kidney Disease Following
Hospitalization with and without Acute Kidney Injury. *Kidney360* **5**, 1623-1632
(2024). https://doi.org/10.34067/kid.0000000000000559

45   Kimura, A., Naka, T., Nohara, K., Fujii-Kuriyama, Y. & Kishimoto, T. Aryl hydrocarbon
receptor regulates Stat1 activation and participates in the development of Th17
cells. *Proceedings of the National Academy of Sciences* **105**, 9721-9726 (2008).
https://doi.org/doi:10.1073/pnas.0804231105

46   Schroeder, J. C. *et al.* The uremic toxin 3-indoxyl sulfate is a potent endogenous
agonist for the human aryl hydrocarbon receptor. *Biochemistry* **49**, 393-400 (2010).
https://doi.org/10.1021/bi901786x

47   Kaminski, T. W., Pawlak, K., Karbowska, M., Mysliwiec, M. & Pawlak, D. Indoxyl
sulfate - the uremic toxin linking hemostatic system disturbances with the

prevalence of cardiovascular disease in patients with chronic kidney disease. *BMC*
Nephrol **18**, 35 (2017). https://doi.org/10.1186/s12882-017-0457-1

48   Lochner, M. *et al.* In vivo equilibrium of proinflammatory IL-17+ and regulatory IL-
10+ Foxp3+ RORgamma t+ T cells. *The Journal of experimental medicine* **205**, 1381-
1393 (2008). https://doi.org/10.1084/jem.20080034

49   Kim, M. J. *et al.* Deletion of PD-1 destabilizes the lineage identity and metabolic
fitness of tumor-infiltrating regulatory T cells. *Nature immunology* **24**, 148-161
(2023). https://doi.org/10.1038/s41590-022-01373-1

50   Cortes, J. R. *et al.* Maintenance of immune tolerance by Foxp3+ regulatory T cells
requires CD69 expression. *Journal of autoimmunity* **55**, 51-62 (2014).
https://doi.org/10.1016/j.jaut.2014.05.007

51   Ichikawa, T. *et al.* CD103(hi) T(reg) cells constrain lung fibrosis induced by CD103(lo)
tissue-resident pathogenic CD4 T cells. *Nature immunology* **20**, 1469-1480 (2019).
https://doi.org/10.1038/s41590-019-0494-y

52   Mantovani, A., Cassatella, M. A., Costantini, C. & Jaillon, S. Neutrophils in the
activation and regulation of innate and adaptive immunity. *Nature Reviews*
Immunology **11**, 519-531 (2011). https://doi.org/10.1038/nri3024

53   Stark, M. A. *et al.* Phagocytosis of Apoptotic Neutrophils Regulates Granulopoiesis
via IL-23 and IL-17. *Immunity* **22**, 285-294 (2005).
https://doi.org/https://doi.org/10.1016/j.immuni.2005.01.011

54   Wilson, A. S. *et al.* Neutrophil extracellular traps and their histones promote Th17
cell differentiation directly via TLR2. *Nature Communications* **13**, 528 (2022).
https://doi.org/10.1038/s41467-022-28172-4

55   Valente, A. J. *et al.* Interleukin-17A stimulates cardiac fibroblast proliferation and
migration via negative regulation of the dual-specificity phosphatase MKP-1/DUSP-1.
Cellular Signalling **24**, 560-568 (2012).
https://doi.org/https://doi.org/10.1016/j.cellsig.2011.10.010

56   Wu, L. *et al.* Cardiac fibroblasts mediate IL-17A-driven inflammatory dilated
cardiomyopathy. *The Journal of experimental medicine* **211**, 1449-1464 (2014).
https://doi.org/10.1084/jem.20132126

57   Träger, C. *et al.* A probiotic approach identifies a Treg-centred immunoregulation via
modulation of gut microbiota metabolites in people with multiple sclerosis and
healthy individuals. *EBioMedicine* **116**, 105743 (2025).
https://doi.org/10.1016/j.ebiom.2025.105743

58   Thiele, A. *et al.* Pharmacological inhibition of adipose tissue adipose triglyceride
lipase by Atglistatin prevents catecholamine-induced myocardial damage. *Cardiovasc*
Res **118**, 2488-2505 (2022). https://doi.org/10.1093/cvr/cvab182

59   Beyhoff, N. *et al.* Application of Speckle-Tracking Echocardiography in an
Experimental Model of Isolated Subendocardial Damage. *J Am Soc Echocardiogr* **30**,
1239-1250.e1232 (2017). https://doi.org/10.1016/j.echo.2017.08.006

60   Sadik, A. *et al.* IL4I1 Is a Metabolic Immune Checkpoint that Activates the AHR and
Promotes Tumor Progression. *Cell* **182**, 1252-1270.e1234 (2020).
https://doi.org/10.1016/j.cell.2020.07.038

61   Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional
changes and characterizing heterogeneity in single-cell RNA sequencing data.
Genome Biology **16**, 278 (2015). https://doi.org/10.1186/s13059-015-0844-5

62   Cui, A. *et al.* Dictionary of immune responses to cytokines at single-cell resolution.
Nature **625**, 377-384 (2024). https://doi.org/10.1038/s41586-023-06816-9
63   Versnjak, J. *et al.* Deep phenotyping of heart failure with preserved ejection fraction
through multi-omics integration. *Eur J Heart Fail* (2025).
https://doi.org/10.1002/ejhf.70041

1398

 Figure Legends

 Figure 1. Depletion of microbiome-derived metabolites attenuates cardiac
remodeling in mice with impaired kidney function. (a) 129/Sv mice were
randomized to receive a broad-spectrum, oral antibiotic cocktail (Abx) or control
drinking water. After one week, mice underwent two-stage subtotal nephrectomy
(STNx) or sham surgery. **(b)** Representative ^{1}H-NMR plasma spectra 13 weeks after
surgery and **(c)** quantification of the aromatic region, normalized to the respective
sham group. **(d)** Heatmap of plasma metabolites quantified by liquid chromatography-
mass spectrometry (LC-MS). **(e)** Plasma concentrations of sulfated microbiome-
derived metabolites quantified by LC-MS. Transdermal glomerular filtration rate (GFR)
normalized measured **(f)** longitudinally and **(g)** at endpoint. **(h)** Heart weight
normalized to tibia length at the endpoint. **(i)** Representative echocardiographic
images at the endpoint. **(j)** Left ventricular posterior wall thickness at end-diastole
(LVPWd). **(k)** Longitudinal changes in normalized left ventricular (LV) mass. **(l)**
Ejection fraction (EF), **(m)** E/e′ ratio, **(n)** global longitudinal strain (GLS) at the
endpoint. **(o)** Histological assessment of left ventricular fibrosis by Sirius Red staining
(scale bars: upper, 1 mm; lower, 100 µm) and **(p)** quantification of fibrotic area. **(q)**
Quantification of leukocytes in the heart at the endpoint by flow cytometry. **(r)** Plasma
C-reactive protein (CRP) levels at the endpoint. **(s)** Radiotelemetric blood pressure
(BP) measurement. **(t)** Representative average systolic BP (SBP) plots at baseline
(week 0) and week 12, and **(u)** quantification. **(v)** Acetylcholine (ACh)-induced
endothelial-dependent vasorelaxation of mesenteric arteries assessed by wire
myography, and **(w)** corresponding EC_{50} values. Symbols in **(f)** and **(k)** indicate the
following comparisons: Sham vs. STNx (‡), Sham Abx vs. STNx Abx (#), and STNx
vs. STNx Abx (*). Data are shown as mean ± SEM. Each dot represents one mouse.
Statistical analyses were performed using one-way ANOVA or Kruskal-Wallis test
followed by Tukey's or Dunn's *post hoc* correction, as appropriate, unless otherwise
indicated. Mann Whitney U-tests with false discovery rate correction according to
Benjamini-Hochberg were used for metabolomics data. Two-group comparisons for
the aromatic NMR region **(c)** and EC_{50} values **(w)** were performed using unpaired two-
tailed Student's t test. Two-group comparisons of systolic blood pressure in **(u)** were
assessed by Mann–Whitney U test.

Figure 2. Kidney disease induces microbiota-dependent Th17-driven systemic
and cardiac inflammation. Immune cells were isolated from spleens and analyzed
by flow cytometry. **(a)** Abundance of neutrophils (CD45$^+$ CD3$^-$ NK1.1$^-$ CD11b$^+$ Ly6G$^+$)
among CD45$^+$ cells. **(b)** Ratio of T helper (T$_H$, CD45$^+$ CD3$^+$ TCRγδ$^-$ CD4$^+$ CD8$^-$) to
cytotoxic T cells (T$_C$, CD45$^+$ CD3$^+$ TCRγδ$^-$ CD4$^+$ CD8$^-$). **(c)** UMAP projection of splenic
T cell populations based on flow cytometry. **(d)** Frequencies of effector memory (EM,
CD44$^+$ CD62L$^-$) conventional T cells (T$_{conv}$, FoxP3$^-$ T$_H$). **(e)** Frequencies of EM
regulatory T cells (T$_{reg}$, FoxP3$^+$ T$_H$). **(f)** Frequencies of RORγt$^+$ T$_{conv}$. **(g)** Changes in
RORγt$^+$ T$_{conv}$ abundance between experimental groups
(* p-adj. < 0.1). **(h)** Frequencies of RORγt$^+$ T$_{reg}$. **(i)** Longitudinal frequencies of RORγt$^+$
T$_{conv}$ (left) and T$_{reg}$ (right). **(j)** Aryl hydrocarbon receptor (AhR) activation of plasma in
reporter cell line. **(k)** *Cyp1a1* expression in splenocytes. **(l)** Representative flow
cytometric plots of cardiac RORγt$^+$ T$_{reg}$ (T$_{reg}$17) and **(m)** quantification. **(n)** Overlap of
differentially expressed genes (DEG) in cardiac bulk mRNA-sequencing between the
two indicated comparisons. **(o)** Significantly enriched biological pathways in the
cardiac transcriptome. **(p)** Targeted gene ontology (GO) and pathway analysis of IL-
17A-related differentially enriched gene sets. Data are shown as mean ± SEM. Each
dot represents one mouse. Statistical analyses were performed using one-way
ANOVA or Kruskal-Wallis test followed by Tukey's or Dunn's *post hoc* correction, as
appropriate, unless otherwise indicated. For (g), Benjamini-Hochberg false discovery
rate (FDR) correction was performed. For RNA sequencing, DEG were generated
using the DeSeq2 pipeline with appropriate FDR correction. Gene set enrichment
analyses were performed using Fisher's exact test followed by FDR correction.

1455

Figure 3. Indoxyl sulfate induces inflammation and cardiac fibrosis independent
of kidney function. (a, b) *In vitro* T$_H$17 differentiation of naïve T helper cells from
129/Sv mice in the presence and absence of indoxyl sulfate (IxS). Two biological
replicates, three technical replicates each. **(c)** Heart failure with preserved ejection
fraction was induced in 129/Sv mice using high-fat diet (HFD) and L-NAME in the
drinking water for 15 weeks. Mice received either IxS or potassium chloride (KCl,
control) in the drinking water throughout the experiment. **(d)** Plasma IxS measured by
liquid chromatography-mass spectrometry (LC-MS). **(e)** Plasma cystatin C levels. (f)
Cyp1a1 expression in spleen. Flow cytometric quantification of IL17A in splenic **(g)**

T_{conv} (CD45$^+$ CD3$^+$ TCRγδ$^-$ CD4$^+$ CD8$^-$ FoxP3$^-$) and **(h)** T_{reg} (CD45$^+$ CD3$^+$ TCRγδ$^-$

CD4$^+$ CD8$^-$ FoxP3$^+$). **(i)** Heart weight normalized to tibia length. **(j)** Cardiac interstitial

fibrosis, assessed by histological staining for collagen type I (COL-1) and fibronectin

(FN-1) (scale bar: 100 µm), with quantification of **(k)** FN-1 and **(l)** COL-1. Data are

shown as mean ± SEM. Each dot represents one mouse. Statistical analyses were

performed using one-tailed Student's t test or Mann-Whitney U-test, as appropriate.

(b) Statistical analysis was performed using linear regression.

1472

Figure 4. IL-17A-signaling in IxS-primed cardiac fibroblasts promotes

extracellular matrix production. Cardiac *Il17ra* expression in **(a)** 129/Sv mice

subjected to subtotal nephrectomy (STNx) or sham operation and treated with either

an antibiotic cocktail (Abx) or control drinking water, and in **(b)** 129/Sv mice subjected

to high-fat diet (HFD) plus L-NAME and indoxyl sulfate (IxS) or potassium chloride

(KCl, control) in the drinking water. Re-analysis of previously published cardiac single

nuclei (sn) RNA-sequencing data (Kaesler *et al.*) from STNx mice shows **(c)** highest

Il17ra expression in fibroblasts and **(d)** higher extracellular matrix (ECM) gene set

score in *Il17ra* expressing cardiac fibroblasts. **(e)** Enrichment analysis for known

cardiac fibroblasts subtypes within differentially expressed genes. **(f)** Cardiac

fibroblasts were isolated from C57BL/6J mice for *ex vivo* treatment with IxS and IL-

17A. Purity validation **(g)** by histology using platelet-derived growth factor receptor

alpha (PDGFRA) and vimentin (VIM) (scale bar: 100 µm), and **(h)** by flow cytometry

showing CD45$^-$ CD31$^-$ PDGFRA$^+$ fibroblasts. **(i)** *Il17ra* expression following *in vitro*

incubation with IxS. Isolated fibroblasts were incubated with 100µM IxS, 1ng/mL IL-

17A, or both and expression of **(j)** *Col1a1*, **(k)** *Col1a2*, **(l)** *Acta2*, **(m)** *Abi3bp* were

quantified. **(n)** Histological confirmation of IL-17RA expression in cardiac fibroblasts

(scale bar: 100 µm). **(a, b)** Data are shown as mean ± SEM. Each dot represents one

mouse. **(i-m)** Data are combined from four biological replicates, with three technical

replicates each. Statistical analyses were performed using one-way ANOVA or

Kruskal-Wallis test followed by Tukey's or Dunn's *post hoc* correction, as appropriate.

(b) Statistical analysis was performed using one-tailed Student's t test. **(d)** Comparison

by two-sided Mann-Whitney U-test.

1496

Figure 5. Pharmacological blockade of the AhR ameliorates inflammation and
cardiac remodeling in kidney disease. (a) *Ex vivo* aryl hydrocarbon receptor (AhR)
inhibition in immune cells from 129/Sv mice with BAY 2416964 (AhRi). **(b)** AhRi blocks
indoxyl sulfate (IxS)-induced *Cyp1a1* gene expression in splenocytes. **(c)** AhRi
suppresses IxS-enhanced TH17 differentiation of naïve T cells. **(d)** 129/Sv mice
underwent two stage subtotal nephrectomy (STNx) or sham surgery. Two weeks after
completion of surgery, STNx mice were randomized to AhRi or vehicle (VEH)
treatment based on plasma cystatin C. **(e)** Survival analysis according to pre-specified
humane endpoints. **(f)** Plasma C-reactive protein (CRP) levels. **(g)** *Cyp1a1* expression
in splenocytes. RORγt in splenic **(h)** T_{conv} (CD45$^+$ CD3$^+$ TCRγδ$^-$ CD4$^+$ CD8$^-$ FoxP3$^-$)
and **(i)** T_{reg} (CD45$^+$ CD3$^+$ TCRγδ$^-$ CD4$^+$ CD8$^-$ FoxP3$^+$) by flow cytometry. **(j)** Left
ventricular (LV) mass normalized to tibia length, **(i)** E/e' ratio and **(l)** global longitudinal
strain (GLS) from endpoint echocardiography. **(m)** Cardiac fibrosis assessed by Sirius
Red staining (scale bar top: 1mm, bottom: 100µm). **(n)** Quantification of Sirius Red-
positive interstitial fibrosis. **(o)** Cardiac *Il17ra* expression. **(p)** Enriched biological
pathways among differentially expressed genes significant in STNx rescued by AhRi.
(q) Targeted gene ontology (GO) analysis of IL-17A-related differentially expressed
gene sets. **(b-c)** Data are combined from three technical replicates. **(c)** Data were
analyzed by nonlinear sigmoidal regression. **(e-o)** Data are shown as mean ± SEM.
Each dot represents one mouse. Statistical analyses were performed using one-way
ANOVA or Kruskal-Wallis test followed by Tukey's or Dunn's *post hoc* correction, as
appropriate, unless otherwise indicated. Gene set enrichment analyses were
performed using Fisher's exact test.

1520

Figure 6. Validation of the pro-fibrotic AhR-IL-17A axis in patients with chronic
kidney disease. (a) Activation of aryl hydrocarbon receptor (AhR) in HT29-Lucia AhR
reporter cells with 100µM indoxyl sulfate (IxS) is blockable by the AhR inhibitor BAY
2416964 (AhRi). **(b)** Serum was obtained from healthy controls (HC) or patients with
chronic kidney disease treated with hemodialysis (HD) for *in vitro* testing on AhR
reporter and naïve T cells. **(c)** Activation of the AhR in response to HC or HD serum
in the presence or absence of AhRi. **(d)** *In vitro* T_H17 differentiation of naïve T cells
from healthy volunteers in the presence of HC and HD serum. **(e)** Plasma IL-17A
derived from OLINK Explore HT in the UK Biobank (UKBB) stratified by estimated

glomerular filtration rate (eGFR) and albuminuria in patients with available data.

Statistical comparisons are made between the reference cell (G1, A1) and each

corresponding cell. Adjusted p-values indicate significance as follows: *** < 0.001,

** < 0.01, * < 0.05. **(f)** Plasma IL-17A in patients with CKD defined by an eGFR < 60

mL/min/1.73 m². with and without heart failure with persevered ejection fraction

(HFpEF). **(g)** Kaplan-Meier survival curves for all-cause mortality stratified by IL-17A

levels (<25th, 25th -75th, >75th percentile) among participants with eGFR < 60

mL/min/1.73 m². **(h)** Corresponding hazard ratios and 95% confidence intervals from

a Cox proportional hazards model for all-cause mortality. Model 1 only IL-17A. Model

2 additionally adjusted for age, sex, and eGFR. Model 3 further adjusted for

hypertension, diabetes, pack years, BMI, and NT-proBNP. **(i)** Human primary cardiac

fibroblasts were incubated *in vitro* with HC and HD serum in addition of anti-IL-17A

(Secukinumab, αIL17A) or AhRi and analyzed for **(j)** *IL17RA*, **(k)** *COL1A1*, **(l)** *COL1A2*,

and **(m)** *ACTA2* expression. **(a, c)** Data are combined from three biological replicates,

with three technical replicates each and **(h-k)** data combined from two primary cell

lines from different donors treated with HC serum (n = 4 donors) or HD serum (n = 9

donors) in triplicates. Statistical analyses were performed using one-way ANOVA or

Kruskal-Wallis test followed by Tukey's or Dunn's *post hoc* correction, as appropriate.

Figure 1

Figure 2

Figure 3

a

Naïve T$_H$ cell isolation from lymph nodes → T$_N$

+ α-CD3ε, α-CD28
+ TGF-β, IL-1β, IL-6
± IxS
4 d incubation

T$_H$17

Analysis by flow cytometry

129/Sv

b

T$_H$17 polarisation [%]

p < 0.0001
R^2 = 0.4022

T$_N$ T$_H$17 0.1 0.5 1 5 10 50 100 200

T$_H$17 + IxS [μM]

c

HFD / L-NAME | KCl
              | IxS

W0   W15

129/Sv

d

Plasma indoxyl sulfate [μmol/L]

0.0003

KCl IxS
HFD / L-NAME

e

Plasma cystatin C [mg/L]

0.4186

KCl IxS
HFD / L-NAME

f

Splenic *Cyp1a1* Relative mRNA expression

0.0889

KCl IxS
HFD / L-NAME

g

T$_H$17 [% of T$_{conv}$]

0.0029

KCl IxS
HFD / L-NAME

h

T$_{reg}$17 [% of T$_{reg}$]

<0.0001

KCl IxS
HFD / L-NAME

i

Heart weight / tibia length [g/m]

0.0260

KCl IxS
HFD / L-NAME

j

FN-1 COL-1

FN-1

COL-1

HFD/L-NAME KCl   HFD/L-NAME IxS

k

Fibronectin positive area [%]

0.0355

KCl IxS
HFD / L-NAME

l

Collagen I positive area [%]

0.0552

KCl IxS
HFD / L-NAME

Figure 4

Figure 5

Figure 6