Type I and III interferons synergize with TNF to promote virally-triggered damage to the intestinal epithelium

Lucie Bernard-Raichon[1,2,3,4]*, Jessica A Neil[1,2,5]*, Kayla Kim[3], Thomas Heaney[1,2], Brittany M Miller[3], Damee Moon[2,6], Ashira Lubkin[2,7,8], Ashley L DuMont[2,9], Victor J Torres[2,9], Jordan Axelrad[10], Yu Matsuzawa-Ishimoto[1,2,11], and Ken Cadwell[1,2,3,12]

Affiliations:

1. Kimmel Center for Biology and Medicine at the Skirball Institute, New York University Grossman School of Medicine, New York, NY, USA.
2. Department of Microbiology, New York University Grossman School of Medicine, New York, NY, USA.
3. Division of Gastroenterology and Hepatology, Department of Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA.
4. Institut de Génomique Fonctionnelle de Lyon, École Normale Supérieure de Lyon, CNRS UMR 5242, UCBL Lyon-1, Lyon F-69007, France.
5. Department of Microbiology and Immunology, Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, VIC, Australia
6. Institute for Systems Genetics, New York University Grossman School of Medicine, New York, New York, United States
7. Fungal Pathogenesis Section, Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Bethesda, MD, USA.
8. Postdoctoral Research Associate program, National Institute of General Medical Sciences, National Institutes of Health (NIH), Bethesda, MD, USA.
9. Department of Host-Microbe Interactions, St. Jude Children's Research Hospital, Memphis, TN, USA.
10. Division of Gastroenterology, Department of Medicine, NYU Grossman School of Medicine, New York, USA.

11. Department of Clinical and Diagnostic Laboratory Science Graduate School of Medical
and Dental Science Institute of Science Tokyo (Science Tokyo), Tokyo, Japan
12. Department of Pathobiology, University of Pennsylvania School of Veterinary Medicine,
Philadelphia, PA, USA.
* Authors contributed equally to this work

Correspondence:

Ken Cadwell: ken.cadwell@pennmedicine.upenn.edu, (215) 573-1076, 421 Curie Blvd,

Philadelphia, PA 19104

Yu Matsuzawa-Ishimoto: matsuzawa.yu@tmd.ac.jp, +81-3-5803-5375, 1-5-45 Yushima,

Bunkyo-ku, Tokyo,113-8510 Japan

Competing interests
K.C. has received research funding from Pfizer, Takeda, Pacific Biosciences, Genentech and

Abbvie. K.C. has consulted for or received an honorarium for speaking engagement from

Puretech Health, Genentech, Moderna, Gentibio, Pioneering Medicines, Prologue Medicines,

and Abbvie. K.C. is an inventor on US patent 10,722,600 and provisional patent 63/157,225.

J.A. has received research grants from BioFire Diagnostics, Genentech, Takeda, and Johnson

& Johnson; consultancy fees, honorarium, or advisory board fees from Abbvie, Abviax, Adiso,

BioFire Diagnostics, Biomerieux, Bristol-Myers Squibb, Celltrion, Ferring, Fresenius, Johnson &

Johnson, Johnson & Johnson, Merck, Pfizer, Sanofi, Takeda, and Vedanta. V.J.T. is an inventor

on patents and patent applications filed by New York University, which are currently under

commercial license to Janssen Biotech Inc. Janssen Biotech Inc. provides research funding and

other payments associated with the licensing agreement.

Abstract

Excessive cell death in the epithelium due to prolonged immune activation is associated with intestinal diseases such as Crohn's disease. Mice with mutations in the Crohn's disease susceptibility gene *Atg16l1* are susceptible to inflammation associated with intestinal epithelial cell (IEC) death, including loss of antimicrobial Paneth cells triggered by infection with murine norovirus (MNV). Here, we show that intestinal disease downstream of MNV depends on IFN-α/β and IFN-λ signaling in IECs. In mouse organoids, IFNs synergize with TNF to induce RIPK1-dependent cell death, amplified by ATG16L1 deficiency. We further show that human intestinal organoids harboring the *ATG16L1* risk allele exhibit heightened sensitivity to TNF and IFN co-stimulation and to serum from severe COVID-19 patients. Our findings reveal that virally triggered cytokines including IFNs exacerbate epithelial vulnerability in genetically predisposed hosts.

Keywords

Norovirus, interferons, autophagy, intestinal inflammation, necroptosis, Paneth cells.

Introduction

A single layer of epithelial cells in the intestine forms the first line of defense against diverse microbial agents ranging from beneficial members of the microbiota to pathogens that cause life-threatening disease[1]. Intestinal epithelial cells (IECs) include enterocytes that absorb nutrients, chemosensory tuft cells, mucus-secreting goblet cells, and enteroendocrine cells that produce hormones. The small intestinal crypt also harbors Paneth cells, IECs that secrete antimicrobial factors such as lysozyme and defensins. The proper turnover of IECs maintains anatomical integrity and contributes to resolving infections, such as through sloughing of damaged or infected cells[2]. However, inflammatory or excessive cell death triggered by prolonged immune activation is a hallmark of inflammatory bowel diseases (IBDs) such as Crohn's disease (CD) and ulcerative colitis[3]. The cell death modality, timing, and cell type specificity are likely important variables that contribute to outcomes in infectious and inflammatory diseases of the gut.

The etiology of IBDs is complex and associated with genetic and environmental factors[4]. Although sustaining remission continues to be a challenge in many patients, the success of therapies that target cytokines and their signaling activities support a disease model in which these soluble immune mediators disrupt the epithelial barrier in a susceptible host. Consistent with this possibility, we demonstrated that mutations in the CD susceptibility gene *ATG16L1* are associated with loss of function and viability of Paneth cells as well as susceptibility to intestinal injury downstream of TNF[5-8]. ATG16L1 is necessary for the cell biological process of autophagy in which cytosolic material, such as protein aggregates and damaged organelles, is targeted to the lysosome for degradation and recycling[9]. ATG16L1 and other autophagy genes have an epithelial-intrinsic role in promoting intestinal barrier function[7, 10-16]. Autophagy and related processes may be particularly important in Paneth cells because of their longevity (2-3 months)

and high secretory burden that make these cells susceptible to ER and mitochondrial stress[5, 7,

17, 18].

We previously developed a model in which *Atg16l1*-mutant mice, including mice with an

IEC-specific deletion of *Atg16l1* (*Atg16l1*[fl/fl];villin-cre, hereafter referred to as *Atg16l1*[ΔIEC]),

persistently infected with murine norovirus (MNV) are susceptible to chemical injury to the gut

by dextran sodium sulfate (DSS) and develop Paneth cell alterations similar to those observed

in patients with CD homozygous for the *ATG16L1 T300A* risk variant[6, 7]. Virally-triggered

disease outcomes are dependent on TNF[6, 7]. This IEC-intrinsic role for ATG16L1 in mitigating

damaged caused by TNF is reproduced in small intestinal organoids, a 3D cell culture model

differentiated from intestinal stem cells. We and others showed that organoids generated from

Atg16l1-mutant mice or patients with *ATG16L1*[T300A] undergo necroptosis (programmed

necrosis) and loss of Paneth cells upon TNF treatment[7, 13]. Given that MNV can promote

beneficial immune development under certain settings and is otherwise tolerated[19-21], the virally-

triggered intestinal damage observed in *Atg16l1*[ΔIEC] mice provides an opportunity to identify the

immune factors that regulate IEC-viability in response to environmental perturbations.

Type I Interferons (IFN-α/β) and type III interferons (IFN-λs) are best known for their

antiviral functions downstream of Janus kinases (JAK) and signal transducer and activator of

transcription proteins (STAT) upon receptor engagement. Although type I and III IFNs induce

similar interferon stimulated genes (ISGs), they have distinct activities mediated in part by

differential expression of their receptors. In the intestine, the type I IFN receptor (IFNAR)

primarily functions in immune cells and is downregulated in IECs in adult mice whereas the type

III IFN receptor (IFNLR) is highly expressed in IECs and neutrophils[22-24]. In the context of MNV

infection, IFN-α/β prevents extraintestinal dissemination while IFN-λ restricts local replication

and fecal shedding[25-29]. Although homeostatic IFN-λ produced by lymphocytes in response to

commensal bacteria limits viral replication during early infection[30], the viral non-structural protein

NS1 encoded by persistent strains mediates evasion of IFN-λ to establish prolonged infection of tuft cells[31, 32]. In addition to their antiviral role, IFNs impact intestinal homeostasis. A recent study reported that mice lacking both IFNAR and IFNLR do not survive even low levels of DSS exposure, which was associated with extensive destruction of the colonic epithelium[33]. By contrast, IFN-λ levels are high in serum and small intestinal tissue of patients with CD, and over-expression or administration of IFN-λ induces inflammatory forms of epithelial cell death, necroptosis and pyroptosis[34, 35].

Although MNV can activate cell death proteins directly in infected IECs to promote its reproduction and immune evasion[36-41], persistent strains display specific tropism for tuft cells due to the restricted expression of the viral entry receptor *Cd300lf* [31, 42, 43]. Thus, in the *Atg16l1* mutant setting, in which cell death is observed in Paneth cells that do not express *Cd300lf*, extracellular factors mediate IEC defects. MNV increases TNF and IFN-γ producing intra-epithelial lymphocytes and inhibits protective γδ T cells upstream of Paneth cell death and DSS-induced disease[6-8]. Here, we identify a critical cytotoxic role of type I and III IFNs in mediating these effects. We report that deletion of IFNAR or IFNLR in IECs rescues intestinal defects in *Atg16l1* deficient mice and identify a role for IFNs in potentiating TNF-induced cell death. Using organoids and serum from patients with COVID, we provide evidence that the general epithelial cytotoxic effect of cytokines produced in response to viral infections is conserved in humans. Thus, our results suggest that a component of viral sequelae includes production of soluble immune mediators such as TNF and IFNs that compromise the viability of IECs, which is especially detrimental in genetically susceptible hosts.

Results

Epithelial type I and III interferon signaling mediate intestinal damage downstream of viral infection.

Atg16l1-mutant mice display spontaneous ISG expression in the gut dependent on the microbiota[44, 45]. To determine whether ISG expression is further increased in the presence of MNV, we examined the small intestine (ileum) and colon of WT and Atg16l1$^{\Delta IEC}$ mice 10 days post intragastric infection with MNV strain CR6. The levels of three representative ISGs (*Mx2*, *Oasl2*, and *Zbp1*) in the ileum and colon were increased by MNV infection, and for the ileum, highest in MNV-infected Atg16l1$^{\Delta IEC}$ mice (Sup. Fig. 1A). IFN-λ is primarily responsible for ISG expression in IECs[46]. To confirm that IFN-λ is responsible for the ISG expression, we crossed *Ifnlr* floxed mice to Atg16l1$^{\Delta IEC}$ mice to obtain Atg16l1;Ifnlr$^{\Delta IEC}$ mice that are deficient in ATG16L1 and IFNLR in IECs. ISG expression in MNV-infected Atg16l1;Ifnlr$^{\Delta IEC}$ mice was not significantly different from uninfected Cre-negative controls, hereafter referred to as wild-type (WT) (Sup. Fig. 1A).

To determine whether IFN-λ signaling contributes to virally triggered intestinal defects, we examined the susceptibility of Atg16l1;Ifnlr$^{\Delta IEC}$ mice to DSS following MNV infection. Atg16l1$^{\Delta IEC}$ mice showed increased lethality, weight loss, and disease score compared to WT mice upon MNV + DSS treatment (Fig. 1A-C), similar to our previous study[7]. In contrast, these disease signs were abrogated in MNV + DSS treated Atg16l1;Ifnlr$^{\Delta IEC}$ mice and were similar to Ifnlr$^{\Delta IEC}$ (*Ifnlr*$^{fl/fl}$villin-cre) and WT controls (Fig. 1A-C). We also generated Atg16l1;Ifnar$^{\Delta IEC}$ (*Atg16l1*$^{fl/fl}$*Ifnar*$^{fl/fl}$villin-cre) mice for additional comparison and unexpectedly found that disease was ameliorated. Consistent with a dominant antiviral role of IFNLR at the epithelial barrier during MNV infection[26], Atg16l1;Ifnlr$^{\Delta IEC}$ mice, but not Atg16l1;Ifnar$^{\Delta IEC}$ mice, displayed increased shedding of virus in stool compared to Atg16l1$^{\Delta IEC}$ or WT mice (Fig. 1D). None of the

mice examined displayed mortality or exacerbated disease in response to DSS in the absence

of MNV, except *Atg16l1;Ifnlr*[ΔIEC] mice (Sup. Fig 1B). These results indicate that IFNLR and

IFNAR in IECs are necessary for the exacerbated intestinal injury response in *Atg16l1* [ΔIEC] mice

infected with MNV and may have a different role in uninfected *Atg16l1* mutant mice.

We previously showed that Paneth cells are less frequent and structurally altered in the

ileum and that colons are shortened in MNV + DSS treated *Atg16l1*[ΔIEC] mice[7]. *Atg16l1*[ΔIEC] mice

also displayed a lower number of Paneth cells compared to similarly-treated WT mice, and the

remaining Paneth cells showed diffuse lysozyme staining and/or reduced number of lysozyme[+]

granules that are a characteristic of CD[6]. Deleting *Ifnlr* or *Ifnar* in *Atg16l1* [ΔIEC] mice led to partial

reversal of Paneth cell abnormalities. The number of Paneth cells in *Atg16l1;Ifnar*[ΔIEC] mice were

increased compared to *Atg16l1*[ΔIEC] mice and the proportion of abnormal Paneth cells were

reduced in *Atg16l1;Ifnlr*[ΔIEC] mice (Fig. 1E-G). To test potential redundancy, we crossed

Atg16l1;Ifnlr[ΔIEC] mice with *Atg16l1;Ifnar*[ΔIEC] mice to generate *Atg16l1;Ifnlr;Ifnar* [ΔIEC] mice deficient

in both receptors. However, *Atg16l1;Ifnlr;Ifnar* [ΔIEC] mice still displayed reductions in Paneth cells

and a high proportion showed abnormal lysozyme staining compared with WT mice (Fig. 1E-G).

We cannot rule out the possibility that the above analyses would miss Paneth cells that lose

their granules, which can happen in the *Atg16l1* mutant setting[5]. To more broadly capture

abnormalities, we used terminal deoxynucleotidyl transferase–mediated dUTP nick-end labeling

(TUNEL) to detect cell death. TUNEL-positive cells were rarely detected in the crypts of WT

mice whereas a subset of crypts displayed multiple TUNEL-positive cells in *Atg16l1* [ΔIEC] mice

(Fig. 1E, H). Rather than a decrease, the number of TUNEL-positive cells displayed a non-

significant increase in *Atg16l1;Ifnar* [ΔIEC] mice compared with *Atg16l1* [ΔIEC] mice. However, the

Atg16l1;Ifnlr [ΔIEC] mice were similar to WT controls and *Atg16l1;Ifnlr;Ifnar* [ΔIEC] mice were similar to

Atg16l1 [ΔIEC] mice (Fig. 1E, H). *Atg16l1*[ΔIEC] mice also exhibited increased colon shortening

compared to WT and *Atg16l1;Ifnlr* [ΔIEC] mice following MNV + DSS (Fig. 1I).

These results indicate that IFNLR and IFNAR signaling mediate lethality in MNV-infected

Atg16l1$^{\Delta IEC}$ mice. IFNLR and IFNAR contribute to Paneth cell abnormalities and cell death, but

deleting these receptors individually or together do not fully reverse these IEC defects.

IFNβ and IFNλ exacerbate TNF-mediated cell death of intestinal organoids.

We previously found that organoids generated from *Atg16l1$^{\Delta IEC}$* mice (*Atg16l1$^{-/-}$*

organoids) display a striking spontaneous ISG signature and that treatment with the JAK

inhibitor ruxolitinib prevents TNF-induced cell death of these *Atg16l1$^{-/-}$* organoids[10]. Although

these results are consistent with a role for signaling downstream of cytokine receptors that use

the JAK/STAT pathway, it was unclear whether IFNs can cooperate with TNF to induce loss of

organoid viability. Since the majority of *Atg16l1$^{-/-}$* organoids die within the first 2-3 days of

seeding when TNF is administered on day 0[7], we modified our protocol to increase the dynamic

range of the assay and assess the synergistic effect of TNF and IFNs. We found that >50% of

Atg16l1$^{-/-}$ organoids and ~80% of *Atg16L1$^{+/+}$* organoids (derived from *Atg16l1$^{fl/fl}$* mice that were

negative for Cre) remained viable 72 hours post-stimulation when TNF was added 3 days after

seeding (Fig. 2A, B). Treatment with IFN-β and IFN-λ2 led to similar or slightly greater loss of

viability as TNF for both *Atg16L1$^{+/+}$* and *Atg16l1$^{-/-}$* organoids (Fig. 2A, B). In contrast, the

combination of TNF and IFN led to substantial organoid death. After only 24 h, *Atg16l1$^{-/-}$*

organoids stimulated with TNF + IFN-β or TNF + IFN-λ2 showed complete loss of viability (Fig.

2A, B). *Atg16L1$^{+/+}$* organoids started to display loss of viability at 24 h and were completely dead

by 48-72 h post-treatment with TNF + IFN-β or TNF + IFN-λ2 (Fig. 2A, B). Titration of IFN-β and

IFN-λ2 in the presence of TNF showed dose responsiveness and that *Atg16l1$^{-/-}$* organoids are

more sensitive to IFN since 10x less IFN induced same level of cell death in this genotype

compared with *Atg16L1$^{+/+}$* organoids (Fig. 2C). Therefore, IFN-β and IFN-λ2 increase the toxicity

of TNF, and ATG16L1 deficiency further sensitizes organoids to the combination of these

cytokines.

IFN-β can induce TNF production[13], which may explain why IFN treatment alone can

induce some degree of cytotoxicity to organoids (Fig. 2B). We observed that *Tnf* gene

expression and protein secretion were increased in *Atg16l1*$^{-/-}$ organoids but not *Atg16L1*$^{+/+}$

organoids treated with IFN-β (Fig. 2D and Sup. Fig. 2A). Antibody-mediated blockade of TNF

rescued viability of *Atg16l1*$^{-/-}$ organoids stimulated with TNF as expected and also greatly

improved viability of *Atg16l1*$^{-/-}$ organoids treated with IFN-β (Fig. 2E). Anti-TNF improved

viability of *Atg16L1*$^{+/+}$ and *Atg16l1*$^{-/-}$ organoids treated with TNF + IFN-λ2 but not IFN-λ2 alone

(Sup. Fig. 2B). These results indicate that IFN-β increases the production of TNF in *Atg16l1*$^{-/-}$

organoids, which contributes to its toxicity.

We confirmed the specificity of IFN-β and IFN-λ2 for their cognate receptors. *Atg16l1*$^{-/-}$

Ifnar$^{-/-}$ organoids were resistant to TNF + IFN-β and *Atg16l1*$^{-/-}$*Ifnlr*$^{-/-}$ organoids were resistant to

TNF + IFN-λ2 (Fig. 2F). Also, IFN-β and IFN-λ2 did not induce representative ISGs in the

absence of their respective receptors (Sup. Fig. 2C). However, the spontaneous ISG expression

observed in untreated *Atg16l1*$^{-/-}$ organoids was absent in *Atg16l1*$^{-/-}$*Ifnlr*$^{-/-}$ but not *Atg16l1*$^{-/-}$*Ifnar*$^{-/-}$

organoids (Sup. Fig. 2C), suggesting a dominant role for IFNLR, possibly due to the

overexpression of *Ifnl2* in absence of IFNAR as previously described[47]. The necroptosis gene

Mlkl is also an ISG and displayed similar IFNAR and IFNLR dependent expression patterns

(Sup. Fig. 2C). Given our prior finding that ATG16L1-deficient organoids display spontaneous

ISG expression[10], we determined whether IFN receptor signaling contributed to cell death

induced by TNF alone. For these experiments, we added TNF to organoids on day 0, a

condition that allows us to examine toxicity of TNF treatment alone. Although the effect of

deleting IFNAR or IFNLR individually (*Atg16l1*$^{-/-}$*Ifnar*$^{-/-}$ and *Atg16l1*$^{-/-}$*Ifnlr*$^{-/-}$) was modest, deleting

both IFN receptors improved the viability of *Atg16l1*$^{-/-}$ organoids (*Atg16l1*$^{-/-}$*Ifnar*$^{-/-}$*Ifnlr*$^{-/-}$) (Fig. 2G).

When these experiments using exogenous cytokines and receptor deletions are taken together,

they show that IFNs synergize with TNF to mediate cell death, especially in the ATG16L1-

deficient setting.

Exacerbation of TNF-induced toxicity in organoids by IFNs is partially dependent on necroptosis

signaling.

TNF receptor engagement in IECs leads to activation of receptor-interacting

serine/threonine protein kinase 1 (RIPK1) to mediate either apoptosis or necroptosis in IECs[48,

49]. RIPK1 mediates apoptosis through caspases, and when apoptosis or caspases are inhibited,

necroptosis occurs through forming a complex with RIPK3 to phosphorylate the pore-forming

molecule MLKL that disrupts cell membrane integrity. We and others have shown that TNF-

induced cell death in *Atg16l1* deficient intestinal organoids and macrophages is dependent on

RIPK1, RIPK3, and MLKL[7, 13, 50]. To investigate how the combination of TNF and IFN reduced

viability, we treated organoids with ruxolitinib as a control, a pan-caspase inhibitor (QVD), and a

RIPK1-inhibitor (Nec1s). Ruxolitinib and Nec1s improved viability of both WT and *Atg16l1*-

deficient organoids treated with TNF + IFN-λ2 and TNF + IFN-β (Fig. 3A, B, Sup. Fig. 3A). By

contrast, QVD did not improve the viability of TNF + IFN stimulated organoids.

We first examined protein levels of cell death mediators at an early time point of cytokine

treatment before *Atg16l1*[-/-] organoids displayed signs of cytotoxicity. Even at 2 h post-

stimulation, phosphorylated RIPK3 (pRIPK3) and cleaved caspases, especially caspase-3, were

visible in TNF or TNF + IFN stimulated *Atg16l1*[-/-] organoids and not WT organoids (Fig. 3C-F,

Sup. Fig. 3B). This result is consistent with the delayed onset of cell death in WT organoids.

MLKL and cleaved caspase-3 were detected in WT organoids treated with IFN-λ2 for 24 h, with

or without TNF; pMLKL was faintly visible when both cytokines were present (Fig. 3F, G, Sup.

Fig. 3B). Although our earlier observation implicated necroptosis because RIPK1 inhibition and

not caspase inhibition protected organoids, we noted the prominent cleavage of caspase-3.

Therefore, we tested whether cell death induced by TNF and IFN treatment were dependent on

MLKL. $Mlkl^{-/-}$ organoids displayed increased viability when treated with TNF + IFN-λ2 or TNF +

IFN-β at 24 h, but this protection was less consistent at 48 h (Fig. 3I, H). These results indicate

that necroptosis contributes to the loss of viability induced by TNF + IFN treatment of organoids.

Virally-triggered intestinal disease in Atg16l1 mutant mice is RIPK3-dependent.

Paneth cell abnormalities and mortality are ameliorated by RIPK1 inhibition in MNV +

DSS-treated $Atg16l1^{ΔIEC}$ mice[7]. As a kinase, RIPK3 may also be a targetable molecule. We

found that $Atg16l1^{ΔIEC};Ripk3^{-/-}$ mice showed substantially reduced lethality and weight loss

compared to $Atg16l1^{ΔIEC}$ mice following MNV + DSS treatment similar to WT and $Ripk3^{-/-}$ control

mice (Fig. 4A). Although not a complete reversal, disease score and colon length of

$Atg16l1^{ΔIEC};Ripk3^{-/-}$ mice were not significantly different from WT and $Ripk3^{-/-}$ mice (Fig. 4A, B),

indicating a general improvement in disease outcomes. Additionally, Paneth cell numbers and

lysozyme staining were restored in $Atg16l1^{ΔIEC};Ripk3^{-/-}$ mice (Fig. 4C-E). These results show

that RIPK3 is necessary for susceptibility of $Atg16l1^{ΔIEC}$ mice to MNV.

Human organoids are susceptible to cell death induced by antiviral factors.

The combination of TNF and IFNs is toxic to colonic organoids derived from humans,

although the concentration of cytokines necessary for cell death is high[51]. Based on our data

with mouse organoids, we hypothesized that human small intestinal organoids will be sensitive

to TNF + IFN treatment, especially in the presence of the *ATG16L1* risk allele. To test this

hypothesis, we utilized organoids that we previously generated from endoscopic biopsy

specimens that were genotyped for the *ATG16L1^T300A* variant (Sup Table 1)[10]. As in the mouse

model, the combination of TNF + IFN-λ2 induced cell death in human organoids, which was

exacerbated in those derived from individuals homozygous for *ATG16L1^T300A* (Fig. 5A-C).

TNF and IFNs are produced in response to most viral infections. We previously showed

that SARS-CoV-2 infected mice display a reduction in Paneth cell numbers and abnormal

lysozyme staining, reminiscent of *Atg16l1* mutant mice infected with MNV[52]. We did not detect

replication competent SARS-CoV-2 particles in the gut of these mice, raising the possibility that

circulating antiviral cytokines can mediate long-range cytotoxic effects on IECs in situations in

which they reach high levels, such as observed in severe COVID[53, 54]. Indeed, gastrointestinal

symptoms are common in patients with COVID-19 and many hospitalized patients display blood

stream infections (BSI) caused by microbes originating from the gut, suggesting intestinal

barrier defects[52]. To test whether soluble immune mediators associated with severe COVID-19

are cytotoxic to human IECs, we incubated the above organoids with serum samples pooled

from hospitalized patients during the height of the pandemic when individuals were screened for

SARS-CoV-2 infection. These samples were compared with serum from healthy volunteers and

hospitalized individuals who were COVID-negative but BSI-positive. Compared with serum from

healthy volunteers, addition of serum from patients with severe COVID led to increased loss of

viability, which was worse in organoids homozygous for *ATG16L1^T300A* (Fig. 5D-F). Serum from

BSI-positive control also worsened organoid death but not as severely as the COVID-positive

samples (Fig. 5 D-F). Therefore, circulating factors associated with severe COVID-19 display

cytotoxicity to IECs, and the *ATG16L1* risk variant further sensitizes cells to this activity.

Discussion

Our study demonstrates that type I and III IFNs are central mediators of intestinal epithelial damage following viral infection in the context of ATG16L1 deficiency. Antiviral IFNs and ISG expression have been observed in IBD patients but their contribution to the pathology is incompletely understood[34, 55-57]. We show that IFN signaling in IECs drives epithelial cell death and disease exacerbation in MNV-infected $Atg16l1^{\Delta IEC}$ mice, while organoid models reveal a synergistic cytotoxicity between IFNs and TNF associated with necroptosis signaling. Notably, deletion of IFN receptors rescued overall survival but had partial effects on Paneth cells. This may reflect a confounding role for IFNs in controlling the virus, or alternatively, redundancy with other cytokines such as IFN-γ, which is known to be upregulated during MNV infection and can also trigger Paneth cell dysfunction[8, 58, 59]. IFN-γ and type I/III IFNs display redundancy in a model of chronic murine astrovirus infection[60]. Also, the role of IFNAR was unexpected given the dominance of IFNLR signaling in IECs and may reflect selective functions in a subset of IECs or a role specific to the $Atg16l1$ mutant setting. Future studies should explore whether each IFN has a distinct IEC target and timing of activity that controls their cytotoxic effects.

These findings strengthen the link between infectious agents such as noroviruses and CD[4, 61, 62]. Although noroviruses are associated with CD in some patients, they most often cause self-resolving gastroenteritis and are detected in many asymptomatic individuals, suggesting that as in our mouse model, the consequences of their presence depend on host factors and may be immune mediated[63, 64]. A recent study showed that Epstein-Bar virus (EBV) infection precedes CD[65]. Thus, triggers of intestinal inflammation are unlikely to be exclusive to noroviruses but there may be some specificity. Although many types of gut microbes including commensal bacteria evoke TNF and IFNs, our finding that serum from patients with severe COVID-19 displays higher toxicity to organoids compared with serum from patients with non-viral BSIs raises the possibility that certain types of infections are more likely to trigger

responses that cause IEC death. These observations also raise the intriguing possibility that

viral sequelae and chronic diseases like IBD share underlying mechanisms that include IEC

death. However, we acknowledge that our experimental system does not allow for such

comparisons, and that IBD is much more complex with a prolonged disease development and

cycles of healing and recurrence.

Therapeutically targeting signaling molecules upstream of cell death may offer a strategy

to protect the epithelium in at-risk individuals. Although RIPK3 deletion ameliorated disease in

Atg16l1$^{\Delta IEC}$ mice *in vivo*, RIPK1 inhibition may provide broader protection by blocking multiple

death pathways. Crosstalk between necroptosis, apoptosis, and pyroptosis and their co-

occurrence[66-68] may be especially relevant to organoids that harbor multiple IEC types. Our time

course experiments show that necroptosis inhibition delays but does not completely prevent

loss of viability in organoids. The partial rescue of viability is reminiscent of findings showing that

deleting *Tnfaip3* and *Tnip1* (genes that are also linked to IBD and autoimmunity) sensitizes to

TNF toxicity in the presence of microbial and viral ligands[69, 70]. Interventions that can be

personalized to restore function of a cell type that is specifically affected in a given susceptible

host, such as Paneth cells in the ATG16L1-deficient background, may be ideal. In this sense,

our observation that RIPK3 deletion better restores Paneth cells compared with IFN receptor

deletion supports targeting nodes at the intersection of multiple cytokine signaling pathways.

Indeed, clinically used JAK inhibitors effectively protect patient-derived colonic organoids from

cytokine cocktails[51].

In summary, this work positions IFNs as key players in virally triggered intestinal

pathology associated with TNF toxicity, particularly in the setting of ATG16L1 deficiency. It

underscores the need to consider synergistic cytokine responses in the management of CD and

other barrier-associated disorders.

Methods

Mice

Age- and gender-matched 8–12-week-old mice on the C57BL/6J (B6) background were used.

Atg16l1^{f/f};VillinCre (*Atg16l1^{ΔIEC}*) mice were provided by S. Virgin (Washington University School

of Medicine, St. Louis, MO) and crossed to *Ifnar1^{f/f}* (Jax, 028256) and *Ifnlr1^{f/f}* mice (provided by

S. Virgin) to obtain *Atg16l1;Ifnar^{ΔIEC}* and *Atg16l1;Ifnlr^{ΔIEC}* mice, respectively. To generate

littermate VillinCre-negative and *Atg16l1^{ΔIEC}* controls, mice were bred heterozygous for either

the *Ifnar^f* or *Ifnlr^f* genes and crossed VillinCre-positive with VillinCre-negative. Ifnar^{ΔIEC} and

Ifnlr^{ΔIEC} mice were bred separately by crossing VillinCre-positive with VillinCre-negative mice.

Atg16l1;Ifnlr;Ifnar^{ΔIEC} mice were obtained by crossing *Atg16l1;Ifnar1^{ΔIEC}* and *Atg16l1;Ifnlr^{ΔIEC}*

mice. WT mice are a pool of VillinCre-negative mice from different crosses. *Rip3^{−/−}* mice were

provided by G. Miller (NYU School of Medicine, New York, NY). *Mlkl^{-/-}* mice were provided by J.

Murphy (WEHI, Parkville, Australia). All animal studies were performed according to approved

protocols by the New York University School of Medicine Institutional Animal Care and Use

Committees.

MNV-triggered disease model

MNV.CR6 stock was prepared as previously described and as detailed in the Supplementary

method section[71]. Mice were infected orally with 3×10^6 PFUs resuspended in PBS. Viral

burden in stool at day 10 post infection was determined by plaque assay. On day 10 post-

infection, mice were administered 5% Dextran Sulfate Sodium (DSS) (TdB Consultancy) in their

drinking water for 6 d after which it was replaced by regular drinking water. *Ripk3^{-/-}* mice showed

increased lethality when given 5% DSS. When these mice were used, 3% DSS was used

instead of 5% as indicated in the figure legend. Clinical disease score and mouse weight were

assessed daily. Disease score was quantified on the basis of five parameters as described

previously, including diarrhea (0–2), hunched posture (0–2), fur ruffling (0 or 1), mobility (0–2),

and blood stool (0 or 1), in which 8 was the maximum score for the pathology[7]. For survival

experiments, mice were euthanized if they reached the humane end point. In other experiments,

mice were sacrificed before receiving DSS or after receiving 5 days of DSS (which occurred

before clinical disease signs).

Intestinal organoid culture

Murine and human organoids (generated during previous studies[10, 72]) were cultured as

described in the Supplementary methods. Murine organoids were treated with 20 ng/ml mTNFα

(Peprotech) and indicated concentrations of mIFNβ (PBL assay science) and mIFNλ2

(Peprotech). In some experiments, organoids were pre-treated with 100nM Ruxolitinib, 10µM Q-

VD-OPh (Millipore Sigma), 10 µM Necrostatin-1s (Nec-1s) (Biovision), 20 µg/ml anti-TNFα

antibody (XT3.11, BioXcell) or 20 µg/ml Rat IgG1 isotype control (HRPN, BioXcell). Human

organoids were treated with 50 ng/mL hTNFα (Peprotech), 100 ng/mL hIFNλ2 (R&D systems).

In another experiment, half of the DMEM/F-12 in the differentiation medium was replaced by

human serum (growth factors were maintained at same concentrations). Serum from healthy

donors from the United States was purchased from SeraCare. The serum from patients with BSI

and SARS-CoV-2 infection was collected at the NYU Langone hospital in 2020 during a

previous study (Table S3)[73]. Each organoid donor was stimulated with the same pool of serum

obtained from healthy donors, patients with BSI and negative for SARS-CoV-2 by qPCR or with

BSI and positive for SARS-CoV-2. Percentages of viable organoids were determined based on

morphology by daily quantification as before[7]. Dead organoids were also marked by staining

with 100 µg/ml PI (Sigma-Aldrich) and 100 µg/ml Hoechst 33342 (Invitrogen) to confirm

phenotypes.

Statistical analysis

GraphPad Prism version 10 was used for statistical analysis unless otherwise specified.

Survival data were analyzed by Kaplan-Meier with Log-rank test followed by Holm Sidak's

multiple comparison test (Figure 1A, 4A). For other parameters, ordinary or lognormal ordinary

(PFU, Figure 1D) one-way ANOVA and Tukey's multiple comparison were used when data

followed a normal distribution or a Kruskal-Wallis test followed by Dunn's multiple comparison

test when they did not (disease score, Figure 1B). For histological measurements, the effect of

mouse genotype was determined on R using individual measurements (e.g. number of Paneth

cells in 50 crypts per mouse) while adjusting for batch effect caused by individual mouse and

experiments. ANOVA type III followed by Tukey's multiple comparison Generalized Linear

Mixed Model (GLMM) was used for Paneth cell counts. A beta-binomial analysis followed by

Tukey's multiple comparison was used for Paneth cell abnormality. Negative Binomial GLMM

was   used   to   compare   the   number   of   TUNEL+   cells   per   crypt.

In Figure 2 and 3, in each experiment, organoids derived from a single mouse per genotype

were stimulated with different cytokines and inhibitors (experiments were then repeated in

independent experiments using organoids derived from different mice). Ordinary one-way

ANOVA were used to compare the effect of the treatment within a genotype, two-way ANOVA

were used when comparing the effect of the treatment and of the genotypes, followed by

Sidak's multiple comparison test. Comparison of area under the curve was performed to

compare time-course curves. For human organoids, an ANOVA followed by Sidak's multiple

comparison test was used to compare the effect of the stimulation on organoid viability

depending on the genotype.

Data availability

All data generated or analysed during this study are included in this published article and its supplementary information files.

Acknowledgements

We thank the NYU Langone's Experimental Pathology Research Laboratory (RRID:SCR_017928) and the NYU Langone's Microscopy Laboratory (RRID: SCR_017934), supported in part by NYU Langone Health's Laura and Isaac Perlmutter Cancer Center Support (grant P30CA016087), for the use of their instruments and technical assistance. We thank the following University of Pennsylvania core facilities for their expert technical support: the Cell and Developmental Biology (CDB) Microscopy Core (RRID SCR_022373) as well as the Molecular Pathology & Imaging Core (MPIC) (RRID: SCR_022420) and Genetically Engineered Mouse Core supported by the UPENN Digestive and Liver Center (P30DK050306). We also thank M. B. da Silva, Y. Shono and M. R. M. van den Brink for their support with TUNEL staining and N Desvignes and A Lambert for their help with lysozyme staining. We also thank Filipe De Vadder for his kind help with statistical analyses. This research was funded in part by NIH grants DK093668 (K.C.), AI121244 (K.C.), AI140754 (K.C.), AI179896 (K.C.), and DK050306 (K.C.); Shanahan Family Fund for Inflammatory Bowel Disease Research (K.C.); pilot grant from King Center for Lynch Syndrome (K.C.); pilot grant from Abramson Cancer Center (K.C.); and an industry partnership with AbbVie Inc. (K.C.).

Author contributions

LBR, JAN, YMI and KC designed the study. LBR, JAN and YMI performed most of the experiments with assistance from KK, TH, and DM and BMM. YMI and JA provided the human organoids lines. AL, AD and VJT provided the human serum samples. LBR, JAN and KC

458   drafted the manuscript. All authors contributed to refinement of the study protocol and approved

459   the final manuscript.

References

1.   Ramanan D, Cadwell K. Intrinsic Defense Mechanisms of the Intestinal Epithelium. *Cell Host*
Microbe 2016; **19**(4)**:** 434-441.

2.   Churchill MJ, Mitchell PS, Rauch I. Epithelial Pyroptosis in Host Defense. *J Mol Biol* 2022; **434**(4)**:**
167278.

3.   Patankar JV, Becker C. Cell death in the gut epithelium and implications for chronic
inflammation. *Nat Rev Gastroenterol Hepatol* 2020; **17**(9)**:** 543-556.

4.   Cadwell K, Loke P. Gene-environment interactions shape the host-microbial interface in
inflammatory bowel disease. *Nat Immunol* 2025; **26**(7)**:** 1023-1035.

5.   Cadwell K, Liu JY, Brown SL, Miyoshi H, Loh J, Lennerz JK *et al.* A key role for autophagy and the
autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature* 2008; **456**(7219)**:**
259-263.

6.   Cadwell K, Patel KK, Maloney NS, Liu TC, Ng AC, Storer CE *et al.* Virus-plus-susceptibility gene
interaction determines Crohn's disease gene Atg16L1 phenotypes in intestine. *Cell* 2010; **141**(7)**:**
1135-1145.

7.   Matsuzawa-Ishimoto Y, Shono Y, Gomez LE, Hubbard-Lucey VM, Cammer M, Neil J *et al.*
Autophagy protein ATG16L1 prevents necroptosis in the intestinal epithelium. *J Exp Med* 2017;
214(12)**:** 3687-3705.

8.   Matsuzawa-Ishimoto Y, Yao X, Koide A, Ueberheide BM, Axelrad JE, Reis BS *et al.* The
gammadelta IEL effector API5 masks genetic susceptibility to Paneth cell death. *Nature* 2022;
610(7932)**:** 547-554.

9.   Matsuzawa-Ishimoto Y, Hwang S, Cadwell K. Autophagy and Inflammation. *Annu Rev Immunol*
2018; **36:** 73-101.

10.   Matsuzawa-Ishimoto Y, Hine A, Shono Y, Rudensky E, Lazrak A, Yeung F *et al.* An intestinal
organoid-based platform that recreates susceptibility to T-cell-mediated tissue injury. *Blood*
2020; **135**(26)**:** 2388-2401.

11.   Jackson DN, Panopoulos M, Neumann WL, Turner K, Cantarel BL, Thompson-Snipes L *et al.*
Mitochondrial dysfunction during loss of prohibitin 1 triggers Paneth cell defects and ileitis. *Gut*
2020; **69**(11)**:** 1928-1938.

12.   Foerster EG, Tsang DKL, Goyal S, Robertson SJ, Robert LM, Maughan H *et al.* ATG16L1 protects
from interferon-gamma-induced cell death in the small intestinal crypt. *Mucosal Immunol* 2023;
16(2)**:** 135-152.

13.   Aden K, Tran F, Ito G, Sheibani-Tezerji R, Lipinski S, Kuiper JW *et al.* ATG16L1 orchestrates
interleukin-22 signaling in the intestinal epithelium via cGAS-STING. *J Exp Med* 2018; **215**(11)**:**
2868-2886.

14.   Pott J, Kabat AM, Maloy KJ. Intestinal Epithelial Cell Autophagy Is Required to Protect against
TNF-Induced Apoptosis during Chronic Colitis in Mice. *Cell Host Microbe* 2018; **23**(2)**:** 191-202
e194.

15.   Jung H, Leal-Ekman JS, Lu Q, Stappenbeck TS. Atg14 protects the intestinal epithelium from TNF-
triggered villus atrophy. *Autophagy* 2019; **15**(11)**:** 1990-2001.

16.   Asano J, Sato T, Ichinose S, Kajita M, Onai N, Shimizu S *et al.* Intrinsic Autophagy Is Required for
the Maintenance of Intestinal Stem Cells and for Irradiation-Induced Intestinal Regeneration.
Cell Rep 2017; **20**(5)**:** 1050-1060.

17.   Bel S, Pendse M, Wang Y, Li Y, Ruhn KA, Hassell B *et al.* Paneth cells secrete lysozyme via
secretory autophagy during bacterial infection of the intestine. *Science* 2017; **357**(6355)**:** 1047-
1052.

18.   Adolph TE, Tomczak MF, Niederreiter L, Ko HJ, Bock J, Martinez-Naves E *et al.* Paneth cells as a
site of origin for intestinal inflammation. *Nature* 2013; **503**(7475)**:** 272-276.

19.   Kernbauer E, Ding Y, Cadwell K. An enteric virus can replace the beneficial function of
commensal bacteria. *Nature* 2014; **516**(7529)**:** 94-98.

20.   Neil JA, Matsuzawa-Ishimoto Y, Kernbauer-Holzl E, Schuster SL, Sota S, Venzon M *et al.* IFN-I and
IL-22 mediate protective effects of intestinal viral infection. *Nat Microbiol* 2019; **4**(10)**:** 1737-
1749.

21.   Abt MC, Buffie CG, Susac B, Becattini S, Carter RA, Leiner I *et al.* TLR-7 activation enhances IL-22-
mediated colonization resistance against vancomycin-resistant enterococcus. *Sci Transl Med*
2016; **8**(327)**:** 327ra325.

22.   Ingle H, Peterson ST, Baldridge MT. Distinct Effects of Type I and III Interferons on Enteric
Viruses. *Viruses* 2018; **10**(1).

23.　Lin JD, Feng N, Sen A, Balan M, Tseng HC, McElrath C *et al.* Distinct Roles of Type I and Type III
Interferons in Intestinal Immunity to Homologous and Heterologous Rotavirus Infections. *PLoS*
Pathog 2016; **12**(4)**:** e1005600.

24.　Mahlakoiv T, Hernandez P, Gronke K, Diefenbach A, Staeheli P. Leukocyte-derived IFN-
alpha/beta and epithelial IFN-lambda constitute a compartmentalized mucosal defense system
that restricts enteric virus infections. *PLoS Pathog* 2015; **11**(4)**:** e1004782.

25.　Rocha-Pereira J, Jacobs S, Noppen S, Verbeken E, Michiels T, Neyts J. Interferon lambda (IFN-
lambda) efficiently blocks norovirus transmission in a mouse model. *Antiviral Res* 2018; **149:** 7-
15.

26.　Nice TJ, Baldridge MT, McCune BT, Norman JM, Lazear HM, Artyomov M *et al.* Interferon-
lambda cures persistent murine norovirus infection in the absence of adaptive immunity.
Science 2015; **347**(6219)**:** 269-273.

27.　Thackray LB, Duan E, Lazear HM, Kambal A, Schreiber RD, Diamond MS *et al.* Critical role for
interferon regulatory factor 3 (IRF-3) and IRF-7 in type I interferon-mediated control of murine
norovirus replication. *J Virol* 2012; **86**(24)**:** 13515-13523.

28.　Baldridge MT, Lee S, Brown JJ, McAllister N, Urbanek K, Dermody TS *et al.* Expression of Ifnlr1 on
Intestinal Epithelial Cells Is Critical to the Antiviral Effects of Interferon Lambda against
Norovirus and Reovirus. *J Virol* 2017; **91**(7).

29.　Nice TJ, Osborne LC, Tomov VT, Artis D, Wherry EJ, Virgin HW. Type I Interferon Receptor
Deficiency in Dendritic Cells Facilitates Systemic Murine Norovirus Persistence Despite Enhanced
Adaptive Immunity. *PLoS Pathog* 2016; **12**(6)**:** e1005684.

30.　Van Winkle JA, Peterson ST, Kennedy EA, Wheadon MJ, Ingle H, Desai C *et al.* Homeostatic
interferon-lambda response to bacterial microbiota stimulates preemptive antiviral defense
within discrete pockets of intestinal epithelium. *Elife* 2022; **11**.

31.　Wilen CB, Lee S, Hsieh LL, Orchard RC, Desai C, Hykes BL, Jr. *et al.* Tropism for tuft cells
determines immune promotion of norovirus pathogenesis. *Science* 2018; **360**(6385)**:** 204-208.

32.　Lee S, Wilen CB, Orvedahl A, McCune BT, Kim KW, Orchard RC *et al.* Norovirus Cell Tropism Is
Determined by Combinatorial Action of a Viral Non-structural Protein and Host Cytokine. *Cell*
Host Microbe 2017; **22**(4)**:** 449-459 e444.

33.　McElrath C, Espinosa V, Lin JD, Peng J, Sridhar R, Dutta O *et al.* Critical role of interferons in
gastrointestinal injury repair. *Nat Commun* 2021; **12**(1)**:** 2624.

34.   Gunther C, Ruder B, Stolzer I, Dorner H, He GW, Chiriac MT *et al.* Interferon Lambda Promotes
590   Paneth Cell Death Via STAT1 Signaling in Mice and Is Increased in Inflamed Ileal Tissues of
591   Patients With Crohn's Disease. *Gastroenterology* 2019; **157**(5)**:** 1310-1322 e1313.

593   35.   Jena KK, Mambu J, Boehmer D, Sposito B, Millet V, de Sousa Casal J *et al.* Type III interferons
594   induce pyroptosis in gut epithelial cells and impair mucosal repair. *Cell* 2024; **187**(26)**:** 7533-
595   7550 e7523.

597   36.   Robinson BA, Van Winkle JA, McCune BT, Peters AM, Nice TJ. Caspase-mediated cleavage of
598   murine norovirus NS1/2 potentiates apoptosis and is required for persistent infection of
599   intestinal epithelial cells. *PLoS Pathog* 2019; **15**(7)**:** e1007940.

601   37.   Wang G, Zhang D, Orchard RC, Hancks DC, Reese TA. Norovirus MLKL-like protein initiates cell
602   death to induce viral egress. *Nature* 2023; **616**(7955)**:** 152-158.

604   38.   Song J, Zhang L, Moon S, Fang A, Wang G, Gheshm N *et al.* Norovirus co-opts NINJ1 for selective
605   protein secretion. *Sci Adv* 2025; **11**(9)**:** eadu7985.

607   39.   Deerain JM, Aktepe TE, Trenerry AM, Ebert G, Hyde JL, Charry K *et al.* Murine norovirus infection
608   of macrophages induces intrinsic apoptosis as the major form of programmed cell death.
609   *Virology* 2024; **589:** 109921.

611   40.   Bok K, Prikhodko VG, Green KY, Sosnovtsev SV. Apoptosis in murine norovirus-infected
612   RAW264.7 cells is associated with downregulation of survivin. *J Virol* 2009; **83**(8)**:** 3647-3656.

614   41.   Aktepe TE, Deerain JM, Hyde JL, Fritzlar S, Mead EM, Carrera Montoya J *et al.* Norovirus-
615   mediated translation repression promotes macrophage cell death. *PLoS Pathog* 2024; **20**(9)**:**
616   e1012480.

618   42.   Orchard RC, Wilen CB, Doench JG, Baldridge MT, McCune BT, Lee YC *et al.* Discovery of a
619   proteinaceous cellular receptor for a norovirus. *Science* 2016; **353**(6302)**:** 933-936.

621   43.   Graziano VR, Alfajaro MM, Schmitz CO, Filler RB, Strine MS, Wei J *et al.* CD300lf Conditional
622   Knockout Mouse Reveals Strain-Specific Cellular Tropism of Murine Norovirus. *J Virol* 2021;
623   **95**(3).

625   44.   Martin PK, Marchiando A, Xu R, Rudensky E, Yeung F, Schuster SL *et al.* Autophagy proteins
626   suppress protective type I interferon signalling in response to the murine gut microbiota. *Nat*
627   *Microbiol* 2018; **3**(10)**:** 1131-1141.

45.   Marchiando AM, Ramanan D, Ding Y, Gomez LE, Hubbard-Lucey VM, Maurer K *et al.* A deficiency
in the autophagy gene Atg16L1 enhances resistance to enteric bacterial infection. *Cell Host*
Microbe 2013; **14**(2)**:** 216-224.

46.   Odendall C, Dixit E, Stavru F, Bierne H, Franz KM, Durbin AF *et al.* Diverse intracellular pathogens
activate type III interferon expression from peroxisomes. *Nat Immunol* 2014; **15**(8)**:** 717-726.

47.   Ardanuy J, Scanlon K, Skerry C, Fuchs SY, Carbonetti NH. Age-Dependent Effects of Type I and
Type III IFNs in the Pathogenesis of Bordetella pertussis Infection and Disease. *J Immunol* 2020;
204(8)**:** 2192-2202.

48.   Dannappel M, Vlantis K, Kumari S, Polykratis A, Kim C, Wachsmuth L *et al.* RIPK1 maintains
epithelial homeostasis by inhibiting apoptosis and necroptosis. *Nature* 2014; **513**(7516)**:** 90-94.

49.   Kattah MG, Shao L, Rosli YY, Shimizu H, Whang MI, Advincula R *et al.* A20 and ABIN-1
synergistically preserve intestinal epithelial cell survival. *J Exp Med* 2018; **215**(7)**:** 1839-1852.

50.   Lim J, Park H, Heisler J, Maculins T, Roose-Girma M, Xu M *et al.* Autophagy regulates
inflammatory programmed cell death via turnover of RHIM-domain proteins. *Elife* 2019; **8**.

51.   Jang KK, Hudesman D, Jones DR, Loke P, Axelrad JE, Cadwell K *et al.* Tofacitinib Uptake by
Patient-Derived   Intestinal   Organoids   Predicts   Individual   Clinical   Responsiveness.
Gastroenterology 2024; **167**(7)**:** 1453-1456 e1455.

52.   Bernard-Raichon L, Venzon M, Klein J, Axelrad JE, Zhang C, Sullivan AP *et al.* Gut microbiome
dysbiosis in antibiotic-treated COVID-19 patients is associated with microbial translocation and
bacteremia. *Nat Commun* 2022; **13**(1)**:** 5926.

53.   Ghaffarpour S, Ghazanfari T, Ardestani SK, Naghizadeh MM, Vaez Mahdavi MR, Salehi M *et al.*
Cytokine profiles dynamics in COVID-19 patients: a longitudinal analysis of disease severity and
outcomes. *Sci Rep* 2025; **15**(1)**:** 14209.

54.   Araujo R, Von Rekowski CP, Fonseca TAH, Calado CRC, Ramalhete L, Bento L. Multiplex Targeted
Proteomic Analysis of Cytokine Ratios for ICU Mortality in Severe COVID-19. *Proteomes* 2025;
13(3).

55.   Mavragani CP, Nezos A, Dovrolis N, Andreou NP, Legaki E, Sechi LA *et al.* Type I and II Interferon
Signatures Can Predict the Response to Anti-TNF Agents in Inflammatory Bowel Disease
Patients: Involvement of the Microbiota. *Inflamm Bowel Dis* 2020; **26**(10)**:** 1543-1553.

56.   Veyssiere M, Hammoudi N, Le Bourhis L, Hassid D, Bonnet J, Tran Minh ML *et al.* Blood
670   proteomic signatures associated with disease activity in inflammatory bowel diseases. *J Crohns*
671   *Colitis* 2025; **19**(9).

672
673   57.   Ostvik AE, Svendsen TD, Granlund AVB, Doseth B, Skovdahl HK, Bakke I *et al.* Intestinal Epithelial
674   Cells Express Immunomodulatory ISG15 During Active Ulcerative Colitis and Crohn's Disease. *J*
675   *Crohns Colitis* 2020; **14**(7)**:** 920-934.

676
677   58.   Farin HF, Karthaus WR, Kujala P, Rakhshandehroo M, Schwank G, Vries RG *et al.* Paneth cell
678   extrusion and release of antimicrobial products is directly controlled by immune cell-derived
679   IFN-gamma. *J Exp Med* 2014; **211**(7)**:** 1393-1405.

680
681   59.   Tomov VT, Palko O, Lau CW, Pattekar A, Sun Y, Tacheva R *et al.* Differentiation and Protective
682   Capacity of Virus-Specific CD8(+) T Cells Suggest Murine Norovirus Persistence in an Immune-
683   Privileged Enteric Niche. *Immunity* 2017; **47**(4)**:** 723-738 e725.

684
685   60.   Herrmann C, Zaldana K, Lustig AM, Bee GCW, Agostino EL, Koralov SB *et al.* Environmental stress
686   drives clearance of a persistent enteric virus in mice. *Nat Microbiol* 2025; **10**(8)**:** 1975-1988.

687
688   61.   Axelrad JE, Chen Z, Devlin J, Ruggles KV, Cadwell K. Pathogen-Specific Alterations in the Gut
689   Microbiota Predict Outcomes in Flare of Inflammatory Bowel Disease Complicated by
690   Gastrointestinal Infection. *Clin Transl Gastroenterol* 2023; **14**(2)**:** e00550.

691
692   62.   Axelrad JE, Olen O, Askling J, Lebwohl B, Khalili H, Sachs MC *et al.* Gastrointestinal Infection
693   Increases Odds of Inflammatory Bowel Disease in a Nationwide Case-Control Study. *Clin*
694   *Gastroenterol Hepatol* 2019; **17**(7)**:** 1311-1322 e1317.

695
696   63.   Bernard-Raichon L, Cadwell K. Immunomodulation by Enteric Viruses. *Annu Rev Virol* 2023;
697   **10**(1)**:** 477-502.

698
699   64.   Lindesmith LC, Donaldson EF, Lobue AD, Cannon JL, Zheng DP, Vinje J *et al.* Mechanisms of GII.4
700   norovirus persistence in human populations. *PLoS Med* 2008; **5**(2)**:** e31.

701
702   65.   Nandy A, Petralia F, Porter CK, Elledge S, Viruses, Group IBDS *et al.* Epstein-Barr Virus Exposure
703   Precedes Crohn's Disease Development. *Gastroenterology* 2025; **169**(1)**:** 150-153.

704
705   66.   Araujo A, Safronova A, Burger E, Lopez-Yglesias A, Giri S, Camanzo ET *et al.* IFN-gamma mediates
706   Paneth cell death via suppression of mTOR. *Elife* 2021; **10**.

67. Newton K, Dugger DL, Wickliffe KE, Kapoor N, de Almagro MC, Vucic D *et al.* Activity of protein kinase RIPK3 determines whether cells die by necroptosis or apoptosis. *Science* 2014; **343**(6177)**:** 1357-1360.

68. Malireddi RKS, Kanneganti TD. Three decades of caspases and RIPKs in life and death. *Hum Mol Genet* 2025; **34**(R1)**:** R121-R136.

69. Su Z, Dziedzic SA, Hu D, Barrett VJ, Broekema N, Li W *et al.* ABIN-1 heterozygosity sensitizes to innate immune response in both RIPK1-dependent and RIPK1-independent manner. *Cell Death Differ* 2019; **26**(6)**:** 1077-1088.

70. Rusu I, Mennillo E, Bain JL, Li Z, Sun X, Ly KM *et al.* Microbial signals, MyD88, and lymphotoxin drive TNF-independent intestinal epithelial tissue damage. *J Clin Invest* 2022; **132**(5).

71. Strong DW, Thackray LB, Smith TJ, Virgin HW. Protruding domain of capsid protein is necessary and sufficient to determine murine norovirus replication and pathogenesis in vivo. *J Virol* 2012; **86**(6)**:** 2950-2958.

72. Jang KK, Kaczmarek ME, Dallari S, Chen YH, Tada T, Axelrad J *et al.* Variable susceptibility of intestinal organoid-derived monolayers to SARS-CoV-2 infection. *PLoS Biol* 2022; **20**(3)**:** e3001592.

73. Lubkin A, Bernard-Raichon L, DuMont AL, Valero Jimenez AM, Putzel GG, Gago J *et al.* SARS-CoV-2 infection predisposes patients to coinfection with Staphylococcus aureus. *mBio* 2024; **15**(8)**:** e0166724.

Figure legends

Figure 1: *Epithelial type I and III interferon signaling mediate intestinal damage downstream of viral infection.* (A) Survival of wildtype (WT; villin-cre⁻ littermate controls), *Atg16l1^{ΔIEC}*, *Atg16l1;Ifnlr^{ΔIEC}*, *Ifnlr^{ΔIEC}*, *Atg16l1;Ifnar^{ΔIEC}*, and *Ifnar^{ΔIEC}* mice receiving 5% DSS in drinking water for 6 days starting 10 days post inoculation (dpi) with MNV. **(B-C)** Cumulative disease score (B) and weight loss (C) at day 6 post DSS from mice in (A). **(D)** Plaque forming units (PFUs) of MNV per gram of stool from mice in (A) at 10 dpi. **(E-F)** Representative images of ileal crypts stained for lysozyme (red) and DAPI (blue) (top) or for deoxynucleotidyl transferase–mediated dUTP nick-end labeling (TUNEL) (bottom) of WT, *Atg16l1^{ΔIEC}*, *Atg16l1;Ifnar^{ΔIEC}*, *Atg16l1;Ifnlr^{ΔIEC}*, and *Atg16l1;Ifnlr;Ifnar^{ΔIEC}* mice infected with MNV on day 5 post DSS treatment as in (A) (scale bars 20μm). **(F)** Violin plots depicting the repartition of the number of Paneth cells observed in individual crypts, overlayed with symbols representing the average number of Paneth cells per crypt observed in each mouse based on the lysozyme staining shown in (E). **(G)** Average proportion of abnormal Paneth cells per crypt observed in each mouse as determined on the basis of whether each Paneth cell displayed a typical staining pattern with distinguishable granules (normal, orange arrow) or depleted and/or diffuse lysozyme staining (abnormal, white arrow) as indicated in (E). **(H)** Proportion of crypts with 1, 2 or 3-5 TUNEL+ cells per mouse of each genotype. Orange arrowheads indicate TUNEL⁺ cells in (E). At least 50 villi and crypts were quantified per mouse. **(I)** Measurement of colon length from mice in (E-H). Each symbol represents individual mice (or the mean in A) with bars representing mean ± SEM from at least two independent experiments. Comparison between all genotypes was performed, not significant comparisons are not indicated, *p≤0.05, **p≤0.01, ***p≤0.001, ****p≤0.0001, See also Figure S1.

Figure 2: IFNβ and IFNλ2 exacerbate TNF-mediated cell death of intestinal organoids. (A)

Representative images of small intestinal organoids derived from WT ($Atg16l1^{+/+}$)

and $Atg16l1^{\Delta IEC}$ mice ($Atg16l1^{-/-}$) cultivated in ENR medium for 3 days and stimulated for another

24h ± 20 ng/ml TNF, ± 100 U/ml IFNβ and ± 1ng/ml IFNλ2, then stained with Hoechst (blue) for

nuclei and propidium iodide (PI) (red) for cell death. Scale bars represent 1000 μm. **(B)** Viability

of $Atg16l1^{+/+}$ and $Atg16l1^{-/-}$ organoids stimulated with indicated cytokines over time as

determined by PI staining and organoid morphology. **(C)** Viability of organoids 48h post

stimulation with 20 ng/ml TNF and 1 (low), 10 (med), 100 (high) U/ml IFNβ (left panel) or 0.01

(low), 0.1 (med), 1 (high) ng/ml IFNλ2 (right panel). **(D)** Quantification of TNF in culture

supernatants by ELISA from indicated organoids stimulated or not with 100 U/ml IFNβ for 48h.

(E) Viability of $Atg16l1^{+/+}$ (left) or $Atg16l1^{-/-}$ (right) organoids treated with 20 μg/ml anti-TNFα

antibody or isotype control for at least 30min before stimulation with cytokines. **(F)** Viability of

$Atg16l1^{+/+}$, $Atg16l1^{-/-}$, $Atg16l1^{-/-}Ifnar^{-/-}$ (derived from $Atg16l1;Ifnar^{\Delta IEC}$ mice), and $Atg16l1^{-/-}Ifnlr^{-/-}$

organoids (derived from $Atg16l1;Ifnlr^{\Delta IEC}$ mice) stimulated on day 3 for 48h. **(G)** Viability of

organoids from the indicated genotype stimulated 2h post seeding with 20 ng/ml TNF for 24 and

48h. For each condition, at least 20 organoids were seeded in 3 different wells (technical

triplicate), which was repeated in at least 3 independent experiments with organoids derived

from different mice each repeat. Each symbol represents the average of one technical triplicate

(or the mean for each stimulation in B) with bars representing mean ± SEM. ns = not significant,

*p≤0.05, **p≤0.01, ****p≤0.0001. See also Figure S2.

Figure 3: TNF and IFNλ synergize to promote RIPK1-dependent cell death in small

intestinal organoids. (A) Representative images of Hoechst (blue) and PI (red) staining of

small intestinal organoids from WT (*Atg16l1$^{+/+}$*) and *Atg16l1^{ΔIEC}* mice (*Atg16l1$^{-/-}$*) cultivated in

ENR medium for 3 days and stimulated with ± 20 ng/ml TNF, ± 1ng/ml IFNλ2 for 48h.

Organoids were pre-treated with vehicle control, 100nM Ruxolitinib, 10μM Q-VD-OPh, or 10 μM

Nec-1s for at least 30min before TNF+IFNλ2 stimulation. Scale bars represent 1000μm. **(B)**

Quantification of organoid viability from (A) at 24 and 48h post-stimulation. For each condition,

at least 20 organoids were seeded in 3 different wells (technical triplicate), which was repeated

in 3 independent experiments with organoids derived from different mice). **(C-E)** Representative

western blot of indicated proteins (C) and quantification of phosphorylated RIPK3 (pRIPK3) (D)

and cleaved caspase-3 normalized to β-actin and unstimulated control (E) in lysates from

Atg16l1$^{+/+}$ and *Atg16l1$^{-/-}$* organoids stimulated for 2h with 20 ng/ml TNF alone or combined with

1ng/ml IFNλ2. C and FL designate cleaved and full length, respectively. **(F-G)** Representative

western blot of indicated proteins (F) and quantification of MLKL normalized to β-actin and

unstimulated control (G) in lysates from *Atg16l1$^{+/+}$* organoids stimulated for 24h with 20 ng/ml

TNF alone, 1ng/ml IFNλ2 alone or the combination of both. **(H-I)** Viability of organoids derived

from WT and *Mlkl$^{-/-}$* mice treated at day 3 with the indicated combinations of TNF, IFNλ2, and

IFNβ for 24h (H) or 48h (I). Each symbol represents the average of one technical triplicate (B, I-

J) or protein expression observed from a pool of different wells in an independent experiment

(F-H) with bars representing mean ± SEM. ns = not significant, *p≤0.05, **p≤0.01, ****p≤0.0001.

See also Figure S2 and S3.

Figure 4: *Virally-triggered intestinal disease in Atg16l1 mutant mice is RIPK3-dependent*.

(A) Survival over time, cumulative disease score, and weight loss in *Atg16L1*^{f/f}, *Atg16L1*^{ΔIEC},

Atg16L1^{f/f};*Ripk3*^{-/-}, and *Atg16L1*^{ΔIEC};*Ripk3*^{-/-} mice infected for 10 days with MNV before receiving

3% DSS in their drinking water for 6 days. **(B)** Colon length on day 5 post DSS treatment of

indicated mice infected with MNV as in (A). **(C)** Representative images of H&E staining (top,

scale bar 100μm) and immunofluorescence of ileal crypts from mice in (B) stained for lysozyme

(red) and DAPI (blue) (bottom, scale bar 20μm). **(D)** Number of Paneth cells observed in each

crypt in mice from each genotype. **(E)** Average proportion of abnormal Paneth cells per crypt

observed in each mouse as determined on the basis of whether each Paneth cell displayed a

typical staining pattern with distinguishable granules (normal, orange arrow) or depleted and/or

diffuse staining (abnormal, white arrow). At least 50 crypts were quantified per mouse. Each

symbol represents individual mice with bars representing mean ± SEM from at least two

independent experiments (except for Paneth cell abnormality, n=1). ns = not significant,

*p≤0.05, **p≤0.01, ***p≤0.001, ****p≤0.0001.

Figure 5: Intestinal organoids derived from *ATG16L1T300A* homozygous individuals exhibit heightened susceptibility to inflammatory factors. (A-B) Representative images at 24h post-stimulation with ± 50 ng/mL hTNFα, ± 100 ng/mL hIFNλ2 (A) and viability over time (B) of human organoids from controls (n = 6 donors homozygous or heterozygous for the non-risk allele) and *ATG16L1^{T300A}* homozygous individuals (n=3 donors). **(C)** Bar graph representation of data from (B) comparing non-risk control and *ATG16L1^{T300A}* homozygous organoids. **(D-E)** Representative pictures at 48h (D) and viability over time (E) of non-risk (n=3) and *ATG16L1T300A* homozygous (n=3) organoids post incubation with pooled serum from healthy donors, patients with a blood stream infection (BSI+ SARS-CoV-2-) or patients with a BSI and infected with SARS-CoV-2 (BSI+ SARS-CoV-2+). **(F)** Bar graph representation of data from (E) comparing non-risk control and *ATG16L1^{T300A}* homozygous organoids. For each condition, at least 20 organoids were seeded in 3 different wells (technical triplicate). Each symbol represents the average viability per donor (calculated from technical triplicates) with bars representing mean ± SEM for each group. For (A-C) 2-5 independent experiments were performed for each organoid donor. For (D-E), one experiment was performed for each organoid donor due to the limited quantity of serum available. ns = not significant, *p≤0.05, **p≤0.01, ****p≤0.0001.

Figure 1

Figure 3

Figure 4

Figure 5