The STA1–DOT2 interaction promotes nuclear speckle formation and

splicing robustness in growth and heat stress responses

Heejin Kim[1], Kyoung-jae Yu[1], So Young Park[2], Dong Hye Seo[3, 4, 5], Dong-Hoon Jeong[2], Woo

Taek Kim[5], Dae-Jin Yun[6], and Byeong-ha Lee[1*]

[1]Department of Life Science, Sogang University, Mapo-gu, Seoul 04107, Republic of Korea

[2]Department of Life Science and and Multidisciplinary Genome Institute, Hallym University,
Chuncheon 24252, Republic of Korea

[3]Plant Biotechnology Research Center, Ghent University, Global Campus, Incheon 21985,
Republic of Korea

[4]Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent,
Belgium

[5]Department of Systems Biology and Institute of Life Science and Biotechnology, Yonsei
University, Seoul 03722, Republic of Korea

[6]Department of Advanced Biotechnology and Global Plant Stress Research Center, Konkuk
University, 05029, Seoul, Republic of Korea

Contact information

Heejin Kim: gmlwls7686@sogang.ac.kr / 0009-0000-2341-5581

Kyoung-jae Yu: kyoungjae@sogang.ac.kr / 0009-0003-1125-572X

Dong Hye Seo: Donghye.Seo@ghent.ac.kr / 0000-0003-1243-4466

So Young Park: soyoungp@hallym.ac.kr / 0000-0001-9202-0874

Dong-Hoon Jeong: dhjeong73@hallym.ac.kr / 0000-0002-2378-1706

Woo Taek Kim: wtkim@yonsei.ac.kr / 0000-0001-5027-4622

Dae-Jin Yun: djyun@konkuk.ac.kr / 0000-0002-3638-6043

*Correspondence: Byeong-ha Lee: byeongha@sogang.ac.kr / 0000-0001-7633-3264

ABSTRACT

Pre-mRNA splicing is carried out by the spliceosome, a large and dynamic ribonucleoprotein complex. The spliceosome is known to be stored in nuclear speckles (NS), which are now recognized as active subnuclear organelles for splicing. However, it remains poorly understood how spliceosomal protein-protein interactions are functionally coupled to NS organization to maintain splicing robustness in plants. Here, we report the functional significance of a specific interaction between two U4/U6·U5 tri-snRNP components of the spliceosome, STA1 and DOT2, in regulating NS organization, pre-mRNA splicing, and heat stress responses in *Arabidopsis*. We identified a missense mutation in DOT2 (a Snu66/SART1 homolog) from a genetic suppressor of the PRP6 homolog mutant *sta1-1* (named *S307*). This mutation restored the weakened interaction between STA1 and DOT2 in the *sta1-1* mutant background. Genetic, biochemical, and cell biological analyses showed that variation in the strength of the STA1-DOT2 interaction was closely associated with changes in NS formation, splicing efficiency, as well as growth and heat tolerance. Pharmacological inhibition of STA1-associated NS formation by tubercidin recapitulated *sta1-1*-like phenotypes and splicing defects, supporting a functional link between NS organization and splicing outcomes. In addition, heat-induced weakening of the STA1-DOT2 interaction was accompanied by reduced NS formation and increased intron retention at the transcriptome-wide level including key heat-responsive transcripts. Based on these observations, we propose that the STA1-DOT2 interaction, likely reflecting the assembly state of the U4/U6·U5 tri-snRNP, functions as a heat-sensitive interaction node that couples spliceosome assembly to NS organization and splicing robustness under stress conditions.

Keywords: *STA1*, *DOT2*, U4/U6·U5 tri-snRNP, spliceosomal protein interaction, splicing, nuclear speckle

INTRODUCTION

Pre-mRNA splicing is an essential step in eukaryotic gene expression and is carried out by the spliceosome, a large and dynamic ribonucleoprotein complex composed of five small nuclear ribonucleoproteins (snRNPs) and numerous auxiliary factors (Will and Lührmann, 2011). Among spliceosomal subcomplexes, the U4/U6·U5 tri-snRNP plays a pivotal role in spliceosome activation, and undergoes extensive structural rearrangements during the splicing cycle (Sander et al., 2006; Will and Lührmann, 2011). Defects in tri-snRNP assembly or stability result in splicing failure, underscoring the importance of precise and coordinated interactions among its components (Klimesová et al., 2021; Schneider et al., 2010).

Arabidopsis STABILIZED1 (STA1) is homologous to the pre-mRNA splicing factors PRP1, Prp6p, and PRPF6 (hereafter collectively referred to as PRP6, unless otherwise indicated) in *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae*, and *Homo sapiens*, respectively. Within the tri-snRNP, PRP6 resides in the U5 snRNP and interacts with components of the U4/U6 di-snRNP, thereby physically bridging the U5 snRNP and the U4/U6 di-snRNP (Bertram et al., 2017; Liu et al., 2006; Makarov et al., 2000; Nguyen et al., 2016; Zhan et al., 2018). Consistent with this role, depletion or mutation of PRP6 in yeast and human cells disrupts tri-snRNP formation and leads to the accumulation of U4/U6 di-snRNP, indicating that PRP6 is essential for tri-snRNP assembly (Galisson and Legrain, 1993; Schaffert et al., 2004).

The *Arabidopsis sta1-1* mutant carries a 6-bp deletion within the half-a-tetratricopeptide (HAT) repeat domain of STA1 (Lee et al., 2006). HAT domains are known to mediate protein–protein interactions and promote the assembly of multiprotein complexes (Zeytuni and Zarivach, 2012). Supporting this, yeast two-hybrid assays using human tri-snRNP components have shown that PRPF6 interacts with several tri-snRNP components through its HAT domain (Liu *et al.*, 2006). These findings suggest that the PRP6 homolog STA1 contributes to pre-mRNA splicing in plants primarily by promoting proper spliceosome assembly through its interactions with other tri-snRNP components.

In the nucleus, nuclear speckles (NS) are prominent subnuclear membraneless compartments enriched in spliceosomal proteins and RNAs (Ilik et al., 2020; Smith et al., 2020). NS serve as sites for the storage, assembly, and dynamic exchange of splicing factors and are closely associated with transcriptionally active chromatin, and are increasingly recognized as sites that actively contribute to efficient pre-mRNA processing (Daguenet et al., 2012; Wu et al., 2024). NS are thought to form through the self-assembly of resident proteins and RNAs and to be maintained by transient protein-protein and protein-RNA interactions (Choi et al., 2025; Day et al., 2012; Kim et al., 2016; Paul et al., 2024). Importantly, localization of spliceosomal components to NS enhances the splicing efficiency of nearby genes, whereas impaired speckle association leads to splicing defects (Liu et al., 2021; Wang et al., 2024).

Several lines of evidence suggest that tri-snRNP components contribute directly to NS organization. In human cells, depletion of PRPF6 disrupts the NS localization of another tri-snRNP component PRPF31 and causes splicing defects (Yildirim et al., 2021). In plants, NS localization of splicing factors has also been implicated in stress adaptation. For example, two alternatively spliced isoforms of RDM16 (a tri-snRNP component PRPF3 homolog) cooperatively promote nuclear condensation and enhance heat tolerance in *Arabidopsis* (Ma et al., 2025). These studies support the idea that NS organization is functionally linked to spliceosome integrity and splicing outcomes, particularly under stress. However, it remains unclear how the interaction-dependent assembly state of the tri-snRNP contributes to NS organization in plants.

In this study, from a *sta1-1* suppressor line (*S307*), we identified a missense mutation in DOT2 (*dot2^S*), a U4/U6·U5 tri-snRNP component homologous to yeast Snu66p and human SART1, which are known to bind to and stabilize the PRP6-involved bridge within the tri-snRNP (Bertram *et al.*, 2017; Zhan *et al.*, 2018). The *dot2^S* suppressor allele restores the weakened interaction between STA1 and DOT2 in the *sta1-1* background, leading to recovery of NS formation, improved pre-mRNA splicing, and restoration of growth and heat tolerance. Our results support the idea that the STA1-DOT2 interaction functions as a critical and heat-sensitive interaction node within the tri-snRNP, contributing to spliceosome assembly to NS organization, and splicing robustness under stress conditions.

RESULTS

A mutation in *DOT2*, a U4/U6·U5 tri-snRNP-associated protein, suppresses the *sta1-1* defects

The *sta1-1* mutants are small in size and hypersensitive to heat (Kim et al., 2017; Lee *et al.*, 2006). To elucidate the molecular mechanism behind it, we employed a forward genetic approach to identify *sta1-1* modifiers, which could reveal the genetic interactors of *STA1*. From an ethyl methanesulfonate (EMS)-treated *sta1-1* mutant pool, we isolated a *sta1-1* suppressor *S307* based on its size recovery. *S307* suppressed the small size phenotype of *sta1-1*, exhibiting the phenotype similar to wild-type (WT, Columbia harboring a *RD29A* promoter-driven luciferase transgene) (Figure 1A). In addition to the restored size, the splicing defects observed in *sta1-1* were also restored to wild-type levels in *S307* (Figures 1B and 1C). Intron retention (IR) profiles of *S307* were more similar to those of WT rather than to those of *sta1-1*, as shown by heatmaps (Figure 1B). Only 16 transcripts (16 increased, 0 decreased) displayed differential IR between *S307* and WT, in contrast to 2,876 transcripts (2,860 increased, 16 decreased) between s*ta1-1* and WT (Figure 1C, Supplemental Table 2).

Whole-genome sequencing combined with map-based positional cloning, using the F2 suppressor seedlings from a cross between *S307* and Landsberg *erecta* identified a G-to-A point mutation in the 8th exon of the *DEFECTIVELY ORGANIZED TRIBUTARIES 2* (*DOT2*, At5g16780) gene (herein, called *dot2^S*). This mutation results in an amino acid change from aspartate to asparagine (Supplemental Figure 1). The *DOT2* gene encodes a protein homologous to Snu66p in *Saccharomyces cerevisiae* and SART1 in *Homo sapiens*, both of which are components of the U4/U6·U5 tri-snRNP (Supplemental Figure 2A) together with STA1. Indeed, *DOT2* promoter-driven GUS activity was detected across all tissues, as found in the *STA1::GUS* plants (Lee et al., 2006) (Supplemental Figure 2B). This widespread and overlapping expression pattern suggests constant and cooperative roles for DOT2 and STA1.

To further examine whether DOT2 functions together with STA1, we characterized two independent *dot2* single mutant alleles: a *dot2* single mutant carrying the *S307* allele and a *dot2* T-DNA insertion mutant (SAIL_775_F10, also known as *mdf-2*) (de Luxan-Hernandez et al., 2022), hereafter referred to as *dot2^S* and *dot2^T*, respectively. *dot2^S* plants were comparable to WT in both growth and splicing patterns, whereas *dot2^T* plants showed growth and splicing

defects resembling those of *sta1-1* (Supplemental Figure 3A and 3B). Consistent with previous reports (de Luxan-Hernandez *et al.*, 2022), $dot2^T$ plants displayed cell death in the root apical meristem (RAM) as revealed by propidium iodide (PI)-stained confocal imaging. However, no detectable RAM defects were observed in the $dot2^S$ plants (Supplemental Figure 3C). Together, these results indicate that $dot2^S$ represents a weak loss-of-function allele and that *dot2* loss-of-function mutants display *sta1-1*-like defects, suggesting a coordinated function for DOT2 and STA1 in growth and pre-mRNA splicing.

Genetic and molecular assays confirm $dot2^S$ as a semi-dominant suppressor of *sta1-1*

To confirm that the $dot2^S$ mutation is the causal mutation, a genomic fragment of *DOT2* including its endogenous promoter was introduced into *S307* via *Agrobacterium*-mediated transformation. Unexpectedly, transgenic plants expressing this genomic fragment of *DOT2* ($S307^{DOT2END}$) did not revert the WT-like phenotype of *S307* to the small-sized *sta1-1* phenotypes (Figure 1D). However, when the *DOT2* gene was overexpressed by the Cauliflower Mosaic Virus (CaMV) 35S promoter in *S307* ($S307^{DOT2OX}$), the overexpressing plants exhibited the *sta1-1*-like small size (Figure 1D). In addition, compared to $S307^{DOT2END}$, $S307^{DOT2OX}$ exhibited higher accumulation of intron-retained transcripts for several marker genes that show IR in *sta1-1*, showing stronger phenotypic reversion by the overexpression (Figure 1E). We speculated that phenotypic reversion only by *DOT2* overexpression might be due to the dominant or semi-dominant effects of the dot2^S mutant protein on sta1-1, relative to those of the wild-type DOT2 protein; the dot2^S protein may interact more effectively with the sta1-1 protein than the DOT2 protein. Thus, we performed a dominance test on $dot2^S$ by genetically combining $dot2^S$ with the *STA1* homozygotes (WT) or the *sta1-1* homozygotes. While no clear size defects were observed in the combinations of $dot2^S$ mutation in the wild-type background, the $dot2^S$ heterozygote in the *sta1-1* background showed an intermediate size, supporting the notion of semi-dominant effects of dot2^S on sta1-1 (Figure 1F). Consistently, the $dot2^S$ heterozygote in the *sta1-1* background exhibited intermediate levels of splicing defects between the *DOT2* (i.e., *sta1-1*) and the $dot2^S$ homozygotes in *sta1-1* (i.e., *S307*) (Figure 1G). Taken together, these results confirm that the mutation in *DOT2* is responsible for the *sta1-1* phenotypic suppression in *S307*.

The *dot2^S* mutation restores physical interaction with sta1-1, rescuing *sta1-1* defects

Homologs of STA1 and DOT2 are known to interact, as supported by structural and

experimental evidence (Häcker et al., 2008; Liu *et al.*, 2006; Sun et al., 2018; Zhan *et al.*, 2018).

Our genetic data suggested that STA1 and DOT2 likely interact physically. In support of this,

our structural modeling predicted an interaction between *Arabidopsis* STA1 and DOT2, with

the *sta1-1* and *dot2^S* mutations positioned near the predicted interface (Supplemental Figure

4A). The *sta1-1* mutation resides within the HAT domain, which mediates protein-protein

interactions (Zeytuni and Zarivach, 2012), and the *dot2^S* mutation maps to the predicted STA1-

interacting surface identified by InterProSurf and PDBePISA (Krissinel and Henrick, 2007;

Negi et al., 2007) (Supplemental Figure 4B). These findings suggest that both mutations may

alter the STA1-DOT2 interaction. We examined interactions between the wild type and mutant

proteins using yeast two-hybrid (Y2H) assays. As expected, STA1 and DOT2 showed

substantial interaction (Figure 1H), whereas interactions between a wild-type protein (STA1 or

DOT2) and a mutant protein (sta1-1 or dot2^S) were not detectable. Interestingly, and consistent

with our expectation, sta1-1 and dot2^S showed interactions comparable to those between STA1

and DOT2 (Figure 1H). Consistent with these observations, bimolecular fluorescence

complementation (BiFC) analysis further confirmed the sta1-1-dot2^S interaction similar to the

STA1-DOT2 interaction (Figure 1I and 1J).

To determine whether the suppressive effect is allele-specific, we generated an

additional *dot2* allele using CRISPR/Cas9 (herein, called *dot2^C*). The *dot2^C* mutation is a 3-bp

deletion located 44 bp upstream of the *dot2^S* mutation, outside the predicted STA1-DOT2

interaction interface (Supplemental Figure 4A-4C). Unlike dot2^S, the dot2^C protein did not alter

interaction strength with STA1 or sta1-1 proteins (Figure 1I and 1J). Consistent with this, *dot2^C*

mutation did not alter the defects in size or intron retention observed in *sta1-1* (Supplemental

Figure 4D and 4E). These results indicate that *dot2^S* is a specific suppressor allele whose

product interacts with the sta1-1 mutant protein, thereby likely restoring the U4/U6·U5 tri-

snRNP assembly in *S307*.

Interaction between STA1 and DOT2 promotes NS formation

We noticed diverse nuclear localization signals in our BiFC assays. Thus, we examined the localization of each GFP-tagged protein (STA1/sta1-1 and DOT2/dot2^S/dot2^C) and classified nuclear signals into five subclasses depending on their subcompartments (Figure 2A). Our observation revealed that STA1-GFP formed nuclear condensates (Class IV and V) at a higher frequency than sta1-1-GFP, whereas GFP-DOT2/dot2^S/dot2^C were mainly distributed throughout the nucleoplasm and nucleolus (Class II) (Figure 2B). Interestingly, co-expression of STA1-GFP and mRFP-DOT2 significantly increased condensate formation of both proteins compared to expression of each protein alone (from ~20%/0% to ~90% for STA1/DOT2) (Figure 2C-2E). This suggests that the STA1-DOT2 interaction contributes to the localization of each protein to nuclear condensates. This idea is supported by the following observations: sta1-1-GFP showed limited condensate formation when co-expressed with mRFP-DOT2/dot2^C, whereas mRFP-dot2^S strongly enhanced sta1-1-GFP condensate formation (Figure 2C-2E), consistent with the interaction strengths of these protein combinations. To identify the nature of the nuclear condensates observed above, we co-expressed each protein with the NS marker mRFP-U1-70K. STA1-GFP condensates showed partial co-localization with mRFP-U1-70K (Figure 2F), whereas the EYFP signals from the BiFC assay, indicating the STA1-DOT2 interaction, showed complete co-localization with mRFP-U1-70K (Figure 2G). In addition, the partial co-localization of STA1-GFP with mRFP-U1-70K became nearly complete in the presence of DOT2 (Figure 2H). These results reconfirm that the STA1-DOT2 interaction promotes nuclear condensate formation and reveal that the condensates correspond to NS. NS are a specific type of nuclear condensates that are enriched in the spliceosome components and associated with mRNA splicing and mRNA metabolism (Bhat et al., 2024; Daguenet *et al.*, 2012). Therefore, our results indicate that the STA1-DOT2 interaction contributes to their NS formation and probably to proper mRNA splicing. It should be noted that STA1-GFP condensates that did not overlap with mRFP-U1-70K suggest that STA1-GFP may form non-functional or non-speckle-like condensates (herein called "aggregates") (see below).

The *dot2^S* mutation restores sta1-1 localization to NS, enabling proper splicing

We next examined the functional significance of NS formation of STA1 and DOT2 for mRNA splicing. To this end, we generated the STA1-GFP/sta1-1-GFP overexpressing plants

in *Salk007933*, a *sta1* T-DNA insertion knock-out line (each named *Salk007933*[STA1-GFP] and

Salk007933[sta1-1-GFP]). The homozygous *sta1* knockouts are lethal. However, introduction of

either STA1-GFP or sta1-1-GFP rescued this lethality and the resulting plants phenocopied the

wild-type and *sta1-1* plants, respectively (Supplemental Figure 5). Using protoplasts isolated

from these GFP-expressing plants, we also confirmed that the interaction between STA1 and

DOT2 (or sta1-1 and dot2[S]) promoted the localization of the GFP-tagged protein to NS (Figure

3A). Previously, we developed a protoplast-based splicing activity assay using a luciferase

reporter gene fused with a genomic fragment of *HSFA3* (*gHSFA3-LUC*) which undergoes

STA1-dependent splicing (Kim *et al.*, 2017). After transfection of the alleles to be tested,

splicing activity was measured using protoplasts from *Salk007933*[sta1-1-GFP]. As expected, the

introduction of STA1 into the protoplasts successfully restored full splicing activity of *gHSFA3-*

LUC, while sta1-1 did not (Figure 3B). Among the tested DOT2 alleles, only the dot2[S] allele

showed the full splicing activity in *Salk007933*[sta1-1-GFP] (Figure 3B). Taken together, these

results indicate that the STA1-DOT2 or sta1-1-dot2[S] interaction is critical for NS formation

and splicing activity, thereby linking NS formation to proper splicing activity.

We then examined the effect of disrupting NS formation by applying tubercidin, an

inhibitor of speckle formation (Kurogi et al., 2014) to *Salk007933*[STA1-GFP] plants. STA1-GFP in

Salk007933[STA1-GFP] is localized predominantly to NS (Figure 3C). However, tubercidin

treatment dispersed the GFP signals into the nucleoplasm (Figure 3C). Tubercidin treatment

also eliminated the differences in splicing activity and growth patterns among WT, *sta1-1*, and

S307, resulting in comparable splicing and growth phenotypes across all genotypes (Figure 3D

and 3E). Altogether, these results demonstrate that the interaction-dependent localization of

STA1 and DOT2 is functionally critical for proper mRNA splicing and that the *S307* suppressor

phenotypes are due to the rescued interaction between sta1-1 and dot2[S].

Heat-induced weakening of the STA1–DOT2 interaction reduces nuclear speckle
formation

To further assess the functional connection between the STA1-DOT2 interaction and

speckle formation, we employed heat as an additional perturbation to weaken protein-protein

interactions, because elevated temperatures can weaken molecular interactions. We also

hypothesized that the heat-hypersensitive phenotype of *sta1-1* might result from heat-induced

weakening of the sta1-1 interaction. Consistent with this idea, BiFC analysis revealed a

reduction in EYFP signals corresponding to the STA1-DOT2 interaction after heat treatment

(Figure 4A-4C).

Unexpectedly, despite the reduction in BiFC signals, heat treatment increased the

numbers of STA1-GFP condensates when STA1-GFP was expressed alone (Supplemental

Figure 6A). We hypothesized that the increase of heat-induced condensates might result from

STA1 self-interaction (i.e., STA1 aggregation). To test this possibility, we performed BiFC

analysis using STA1-nEYFP and STA1-cEYFP. Heat treatment indeed increased BiFC signals,

indicating enhanced STA1 self-interaction under heat (Supplemental Figure 6B). Most EYFP

signals derived from STA1 self-interaction did not overlap with signals from the speckle

marker mRFP-U1-70K under either normal or heat conditions (Supplemental Figure 6C and

6D), suggesting that the STA1 self-interacting condensates are STA1 aggregates. This idea is

further supported by the presence of a predicted intrinsically disordered regions (IDRs) within

STA1 (residues 109-216, 298-318, 910-918, and 1022-1029), as identified by MolPhase (Liang

et al., 2024) (Supplemental Figure 6E). Therefore, the increase of STA1-GFP condensates

observed under the STA1-GFP single expression is likely the result of enhanced STA1-GFP

self-aggregation.

To distinguish the NS from STA1 aggregates, we used BiFC analysis to quantify NS

induced by the STA1-DOT2 interaction. The number of these speckle-representing BiFC

signals in nuclei decreased after heat treatment and speckle formation was reduced in

combinations of the weakly interacting STA1 and DOT2 allele products (Figures 4D-4G).

These results reaffirm that the STA1-DOT2 interaction is required for efficient NS formation.

The *dot2^S* mutation suppresses the heat sensitivity of *sta1-1*

285   Next, we sought to determine whether the strength of the STA1-DOT2 interaction also

286 plays an important role in the heat responses. The *sta1-1* mutants have previously been shown

287 to display heat-sensitive phenotypes (Kim *et al.*, 2017; Kim et al., 2018). At both the seedling

and reproductive stages, the heat-sensitive phenotypes of *sta1-1* were almost fully rescued in *S307* (Figure 5A–5C). These results along with the partial or full reversion to *sta1-1* heat sensitivity observed in $S307^{DOT2OX}$ indicated that DOT2 functions together with STA1 under heat stress (Figure 5A and 5C). In addition, the $dot2^C$ allele did not suppress the heat sensitivity of s*ta1-1*, supporting the allele-specific suppressive effect of $dot2^S$ (Figure 5B and 5C). These findings also suggest that the restored heat tolerance in *S307* results from the rescued interaction between sta1-1 and dot2^S.

The $dot2^S$ mutation mitigates transcriptome-wide intron retention defects in *sta1-1* under heat stress

Under heat stress, IR profiles in *S307* were partially restored toward WT levels (Figure 6A). Although the global splicing patterns in *S307* did not fully return to wild-type levels, the IR defects of *sta1-1* were markedly improved in *S307*. Among 67,103 transcripts with significant IR changes across genotypes, *S307* showed a reduced median IR ratio (0.14) compared with *sta1-1* (0.34) (Figure 6B). Across the three genotypes under heat, 1,114 introns (from 940 genes) showed increased IR in WT, whereas 58,522 introns (12,140 genes) and 36,617 introns (10,654 genes) showed increased IR in *sta1-1* and *S307*, respectively (Figure 6C, Supplemental Table 3). Venn diagram analysis further revealed that 27,514 introns with elevated IR in *sta1-1* did not show increased IR in *S307* (Figure 6D).

Among the genes with restored splicing patterns, *HEAT SHOCK TRANSCRIPTION FACTOR A1*s (*HSFA1*s), key upstream regulators of heat-responsive genes (Li et al., 2023; Nishizawa-Yokoi et al., 2011; Röth et al., 2017), showed reduced intron retention in *S307* compared with *sta1-1* (Figure 6E, Supplemental Figure 7). This recovery was further supported by decreased levels of spliced transcripts and increased levels of intron-retained transcripts of *HSFA1*s in $S307^{DOT2END}$ and $S307^{DOT2OX}$ relative to *S307* (Figure 6F). The $dot2^C$ allele failed to rescue splicing defects of *HSFA1s* in *sta1-1*, consistent with its heat-sensitive phenotypes (Figure 5B and 5C, Supplemental Figure 8). Proper splicing of *HSFA1s* in *S307* enabled normal expression of downstream heat-responsive genes (Figure 6G). These findings indicate that $dot2^S$ suppresses intron retention defects in heat-responsive genes in *sta1-1*, thereby

contributing to the restoration of heat tolerance. Taken together with the previous results, these

findings highlight the importance of the STA1-DOT2 interaction for proper heat stress

responses through NS formation and accurate mRNA splicing under heat conditions. It should

be noted that a partial rescue of splicing defects and heat sensitivity in *sta1-1*, particularly at

the pollen stage, under heat stress indicates that the sta1-1-dot2^S interaction is weaker than the

original STA1-DOT2 interaction.

DISCUSSION

Nuclear speckles (NS) are dynamic, membraneless nuclear compartments enriched in spliceosomal components and RNAs (Mintz et al., 1999; Smith *et al.*, 2020), and are increasingly recognized as active hubs that facilitate efficient pre-mRNA processing rather than acting merely as passive storage sites (Wu *et al.*, 2024). Previous studies have emphasized the contribution of intrinsically disordered regions, low-complexity domains, and multivalent RNA–protein interactions to NS organization (Ku et al., 2025; Paul *et al.*, 2024; Wang *et al.*, 2024; Zhang et al., 2024). To what extent spliceosomal protein-protein interactions involved in spliceosome assembly contribute to NS formation remains poorly understood. In this study, we provide evidence that NS organization in plants is tightly associated with the integrity of a specific interaction between two core U4/U6·U5 tri-snRNP components, STA1 and DOT2, thereby linking spliceosome assembly with subnuclear organelle organization.

Our genetic and cell biological analyses show that the weakening of the STA1-DOT2 interaction by the *sta1-1* mutation is accompanied by reduced NS formation, compromised splicing efficiency, and growth defects, whereas restoration of this interaction by the *dot2^S* suppressor allele is associated with recovery of these phenotypes. Importantly, this suppression is achieved through reinforcement of the STA1-DOT2 interaction probably within the tri-snRNP. These observations suggest that NS organization is not independent of spliceosome assembly status and is closely linked to the stability of specific spliceosomal interactions.

The spliceosome is a highly dynamic macromolecular complex, and formation of the U4/U6·U5 tri-snRNP represents a critical step in each splicing cycle (Will and Lührmann, 2011). PRP6/STA1 functions as a central bridging factor between the U4/U6 di-snRNP and the U5 snRNP, with its interactions stabilized by tri-snRNP–specific proteins such as SART1/DOT2 (Bertram *et al.*, 2017; Nguyen *et al.*, 2016). Our observation of a positive correlation between STA1–DOT2 interaction strength and NS abundance supports a model in which proper tri-snRNP assembly facilitates efficient recruitment or retention of mature spliceosomal complexes within NS. Consistent with this view, quantitative subnuclear localization analyses revealed that co-expression of STA1-GFP and mRFP-DOT2 significantly increased NS formation (types IV and V), compared with expression of each protein alone

(Figure 2B and 2D). In contrast, co-expression of sta1-1-GFP and mRFP-DOT2 resulted in

reduced NS-associated localization patterns (types IV and V) and increased localization to

Cajal bodies (type III) compared with the wild-type protein (Figure 2D). Tri-snRNPs are

assembled in Cajal bodies and subsequently redistributed to NS, which contain high levels of

mature, splicing-competent complexes (Schaffert *et al.*, 2004; Yildirim *et al.*, 2021). Thus, our

results suggest that the *sta1-1* mutation impairs later steps associated with NS organization due

to defects in the STA1-DOT2 interaction, likely leading to impaired formation of splicing-

competent spliceosome. In agreement with our results and speculation, when SART1,

homologous to DOT2, was immunologically depleted from HeLa cells, pre-mRNA splicing

was inhibited but the integrity of the U4/U6·U5 tri-snRNP was not affected, indicating the

importance of SART1 in active spliceosome formation and recruitment of the tri-snRNP to the

pre-spliceosome (Makarova et al., 2001; Utans et al., 1992). Thus, the STA1–DOT2 interaction

is likely to promote DOT2-dependent localization of STA1 to NS, concomitant with the

delivery of the tri-snRNP to the pre-spliceosome for active spliceosome assembly.

The functional relevance of our notion becomes particularly apparent under heat stress

conditions. Elevated temperatures are known to destabilize spliceosomal assemblies and

broadly impair splicing fidelity (Bond, 1988; Bond and James, 2000; Park et al., 2024). The

most sensitive component of the spliceosome is the U4/U6·U5 tri-snRNP (Bond, 1988). In this

context, the STA1–DOT2 interaction appears to function as a heat-sensitive interaction node,

since the heat-induced weakening of this interaction correlates with reduced NS formation,

increased intron retention, and weakened heat tolerance (Figure 4, Figure 5, Figure 6).

Reinforcement of the interaction by the dot2^S allele in sta1-1 (i.e., the *S307* suppressor) is

associated with improved NS formation and splicing performance, as well as partial recovery

of survival under heat stress (Figure 4, Figure 5, Figure 6). Although it is still debated whether

NS primarily function as storage sites or as active hubs for RNA metabolism, our findings

support the latter view by linking speckle organization to both defined molecular interactions

and functional splicing outcomes. It should also be noted that STA1 itself appeared to form

non-functional aggregates under heat conditions due to its IDRs (Supplemental Figure 6).

DOT2 seemed to inhibit the aggregation of STA1 and direct STA1 to NS, presumably

facilitating formation of an active spliceosome. Thus, STA1-DOT2 interaction contributes to

the robustness of pre-mRNA splicing under heat stress.

Taken together, our study expands current understanding of NS biology by

demonstrating that NS organization in plants is closely associated with the assembly state of

the spliceosome, mediated by specific protein–protein interactions within the U4/U6·U5 tri-

snRNP. On this basis, we propose that heat-sensitive interaction nodes such as the STA1-DOT2

bridge represent "molecular weak points" or "interaction-level sensors" that integrate

spliceosome architecture, subnuclear organelle formation, and environmental responsiveness.

This interaction-based perspective provides insight into how plants maintain splicing

robustness under stress conditions and suggests that targeted modulation of spliceosomal

interactions may offer a potential strategy for improving stress tolerance.

METHODS

Plant growth conditions

Arabidopsis thaliana seeds were planted on Murashige and Skoog (MS) medium containing MS salts with vitamins (Caisson Laboratories Inc, USA), 2% sucrose, and 0.3% gelite (Duchefa, the Netherlands), adjusted to pH 5.7-5.8. For vertical growth, 0.6% gelrite was supplemented. Seeds were surface-sterilized with commercial bleach supplemented with 0.01% Tween 20 (USB, USA), stratified at 4°C for 2 days in the dark, and then transferred to a growth chamber under long-day conditions (16 h light/8 h dark) at 22°C. For soil-grown plants, seeds were sown on Sun Gro propagation mix (Sun Gro, Canada) and maintained under long-day conditions (16 h light/8 h dark) at 22°C.

Positional cloning and mutation identification

To map the *S307* mutation locus, *S307* (Columbia background) was crossed with Landsberg *erecta* (L*er*) to generate an F2 mapping population. WT-sized F2 plants homozygous for the *sta1-1* mutation were selected, and recovery of the size phenotype was confirmed in the F3 population pool. Genomic DNA was extracted from pooled leaves of 94 selected F2 mutant individuals using the RNeasy Plant Mini Kit (QIAGEN, Germany), and whole-genome sequencing was performed at Macrogen Inc. The genomic region between the SSLP markers Ch5_5.46 Mb and Ch5_5.71Mb, identified by initial mapping, was compared with the WT genomic sequence, identifying a candidate point mutation at position 1,651 bp of *DOT2* (At5g16780) coding sequence. The candidate mutation was validated by complementation using genomic constructs driven by either the native *DOT2* promoter (DOT2::DOT2) or the Cauliflower mosaic virus 35S promoter (35S::DOT2). For native *DOT2* promoter complementation lines (*S307^{DOT2END}*), a 3,783 bp genomic fragment spanning 1,695 bp upstream of the DOT2 start codon to 294 bp downstream of the stop codon was amplified and cloned into pCAMBIA1300 between the *Kpn*I and *Bam*HI sites (Takara, Japan). For 35S-driven complementation lines (*S307^{DOT2OX}*), the 2,463 bp *DOT2* coding sequence was amplified, cloned into pDONR221 entry vector, and subsequently recombined into the pMDC32 destination vector using the Gateway cloning system. All primers used to generate complementation constructs are listed in Supplemental Table S1.

Generation of different alleles of *dot2* mutants

The *dot2* single mutant carrying the *S307* allele (*dot2^S^*) was obtained from the F2 progeny of a cross between WT and *S307*. A derived cleaved amplified polymorphic sequence (dCAPS) marker detecting the single-nucleotide substitution in the *dot2^S^* allele using *Eco*RI (Takara, Japan), together with a marker detecting the 6-nucleotide deletion in *sta1-1* (used to exclude plants carrying the *sta1-1* mutation), was employed to isolate the *dot2^S^* single mutant. The *dot2* T-DNA insertion mutant *SAIL_775_F10* (*dot2^T^*) was obtained from the Arabidopsis Biological Resource Center (ABRC). Because homozygous *dot2^T^* plants are infertile, only heterozygotes were maintained, and homozygotes were identified by their dwarf phenotype (Petricka et al., 2008).

A CRISPR/Cas9-generated *dot2* mutant allele (*dot2^C^*) was produced in the *sta1-1* background using the CRISPR/Cas9 system as previously described (Cho et al., 2017; Liu et al., 2015; Yun et al., 2022). A single guide RNA (sgRNA) targeting *DOT2* was designed using the CRISPR web toolbox CHOPCHOP (https://chopchop.cbu.uib.no) (Labun et al., 2019). The annealed oligonucleotides (DOT2sgRNA3_infu_CF and DOT2sgRNA3_infu_CR) for sgRNA were cloned into the psgR-Cas9-At to generate *DOT2*-targeting construct. The resultant DOT2sgRNA3-Cas9 cassette, excised with *Hin*dIII and *Eco*RI (Takara, Japan), was subsequently subcloned into the *Hin*dIII-*Eco*RI sites of pCAMBIA1300. *Agrobacterium tumefaciens* strain GV3101 carrying these constructs was used to transform *sta1-1* plants by the floral dip method to obtain T1 seeds. CRISPR/Cas9-edited elite lines were identified by Sanger sequencing of the DOT2sgRNA3-targeted region in the T1 generation, and homozygous lines were subsequently isolated. All primers used for the isolation of different *dot2* mutant allels are listed in Supplemental Table S1.

RNA extraction and cDNA synthesis

Total RNA was extracted from seedlings at the designated developmental stage using RNAiso Plus reagent (Takara, Japan) and treated with RNase-free DNase (New England BioLabs, USA). cDNA was synthesized from total RNA using the TOPscript^TM^ Reverse Transcriptase kit (Enzynomics, Korea). Semi-quantitative RT-PCR and realtime PCR analyses were performed using 2 μL of the resulting cDNA (corresponding to 50 ng of input RNA). All

450   primers used for intron retention analyses and gene expression quantification are listed in

451   Supplemental Table S1.

GUS staining assay

For the *STA1* promoter-driven *GUS* (*STA1::GUS*) and *DOT2* promoter-driven *GUS*

(*DOT2::GUS*) constructs, 1,475 bp and 1,695 bp promoter fragments upstream of the start

codon of each gene were amplified, respectively. These fragments were cloned into

pCAMBIA1381 between the *Bam*HI and *Hind*III sites for *STA1::GUS*, or between the *Bam*HI

and *Nco*I sites for *DOT2::GUS* (Takara, Japan). *Agrobacterium tumefaciens* strain GV3101

carrying these constructs was used to transform Col-0 plants by the floral dip method.

Tissues of transgenic *Arabidopsis* seedlings carrying the *STA1::GUS* or *DOT2::GUS*

constructs were collected in 90% acetone (Duksan, Korea) in conical tubes on ice and incubated

for 10-20 min at room temperature. After removal of acetone, samples were washed twice for

5 min with washing buffer (50 mM potassium phosphate buffer (pH = 7.0), 0.2% Triton X-100,

2 mM potassium ferricyanide, and 2 mM potassium ferrocyanide). The washing buffer was

then replaced with staining buffer (washing buffer supplemented with 2 mM X-Gluc (Sigma,

Japan), and tissues were incubated overnight at 37°C. Subsequently, tissues were incubated in

70% ethanol to remove chlorophyll. GUS-staining was observed using a light microscope

(Leica ICC50, Germany).

Nuclear localization, co-localization, and BiFC analyses

For nuclear localization analyses, the full-length coding regions of *STA1* (*sta1-1*)

lacking a stop codon and *DOT2* (*dot2^S* and *dot2^C*) containing a stop codon were amplified,

cloned into pDONR221 entry vector, and subsequently recombined into the pGWB551 or

pGWB552 destination vectors, respectively, using the Gateway cloning system. *Agrobacterium*

tumefaciens cells harboring STA1 (sta1-1)-GFP or GFP-DOT2 (dot2^S and dot2^C) constructs

were infiltrated into *Nicotiana benthamiana* leaves in expression buffer (50 mM MES (pH =

5.7), 10 mM MgSO$_4$, 5% D-Glucose, and 0.1 mM of acetosyringone) together with P19, each

at a final concentration of OD$_{600}$ of 0.2. After 36 h of incubation of infiltrated *N. benthamiana*

at 22°C, fluorescence images were acquired using a confocal microscope (TCS SP5 or

STELLARIS; Leica, Germany). GFP signals were excited at 488 nm and detected at 500-530

nm. For heat treatment, samples were incubated in liquid MS medium (0.5x MS mineral salts,

1% sucrose, 5 mM 2-(N-Morpholino) ethanesulfonic acid (MES; Sigma, Japan), pH adjusted

to 5.7 with KOH) for 1 h at 37°C following the initial 36 h incubation.

For co-localization analyses, the full-length DOT2 (dot2^S and dot2^C) coding region

containing a stop codon was cloned downstream of the CaMV 35S promoter and fused in frame

with mRFP in the p35S-FAST-mRFP vector. *Agrobacterium* cultures carrying STA1 (sta1-1)-

GFP and mRFP-DOT2 (dot2^S and dot2^C) constructs were co-infiltrated into *N. benthamiana*

leaves and fluorescent signals were observed as described above. RFP signals were excited at

532 nm and detected at 580-620 nm.

For BiFC analyses, the same entry clones used for nuclear localization analyses were

used. STA1 (sta1-1) and DOT2 (dot2^S and dot2^C) entry clones were recombined downstream

of the CaMV 35S promoter and fused in frame with nEYFP or cEYFP in the pSITE-nEYFP-

N1 (GU734648) or pSITE-cEYFP-C1 (GU734652) Gateway vectors, respectively.

Agrobacterium cells carrying pSITE-STA1 (sta1-1)-nEYFP or pSITE-cEYFP-DOT2 (dot2^S

and dot2^C) constructs were co-infiltrated in combination into *N. benthamiana* leaves in the

indicated combinations, and fluorescent signals were observed as described above. EYFP

signals were excited at 488 nm and detected at 500-530 nm. As a nuclear speckle marker, the

full-length coding region of *U1-70K* (*At3g50670*) was amplified and cloned into p35S-FAST-

mRFP between the *Bam*HI and *Sal*I (Enzynomics, Korea) sites. As a nucleolus marker,

FIBRILLARIN 2 (FIB2, At4g25630)-mRFP, provided by Dr. Hyun-Sook Pai, was used (Park

et al., 2021). All primers used to generate constructs are listed in Supplemental Table S1.

Yeast two-hybrid assay

The full-length coding region of *STA1* or *sta1-1* was inserted into the pGADT7 vector,

and *DOT2* or *dot2^S* was inserted into the pGBKT7 vector. The resulting constructs, pGADT7-

STA1 (sta1-1) as prey and pGBKT7-DOT2 (dot2^S) as bait, were co-transformed into the

Saccharomyces cerevisiae strain AH109 (Takara, Japan) and incubated at 30°C for 3 days.

Transformants were selected on SD/-Leu/-Trp or SD/-Leu/-Trp/-His medium supplemented

with 3-amino-1,2,4-triazole (3-AT; Sigma, Japan), a competitive inhibitor of the *HIS3* gene

product. All primers used to generate constructs are listed in Supplemental Table S1.

Arabidopsis mesophyll protoplast transient expression assay

Protoplast isolation and transient expression assays were performed as previously described (Kim *et al.*, 2017). Effector constructs (STA1, sta1-1, DOT2, dot2^S, and dot2^C) and reporter constructs (g*HSFA3* fused to NanoLuc luciferase (nLUC) were generated by inserting the corresponding coding sequences between the HBT promoter (a modified CaMV 35S promoter) and the NOS terminator in a plant expression vector. Firefly luciferase (fLUC), driven by the *UBQ10* promoter, was included as an internal control.

For nuclear localization analysis in protoplasts, HBT-DOT2 (dot2^S and dot2^C) constructs were transfected to *Salk007933*$^{STA1-GFP}$ or *Salk007933*$^{sta1-1-GFP}$. Fluorescence images were acquired using a confocal microscope (TCS SP5; Leica, Germany). GFP signals were excited at 488 nm and detected at 500-530 nm.

For functional splicing assays, HBT-DOT2 (dot2^S and dot2^C) constructs were transfected to *Salk007933*$^{sta1-1-GFP}$ and reporter activities were calculated as the nLUC/fLUC ratio. Transfected protoplasts were lysed in passive lysis buffer (Promega, USA) containing 1% Triton X-100 (USB, USA) and briefly vortexed. Lysates were incubated at room temperature for 10 minutes and then centrifuged at 13,000 rpm for 30 s. The supernatant was transferred for measurement of fLUC and nLUC activities using the Nano-Glo® Dual-Luciferase® Reporter Assay System (Promega, USA) and a GloMax 20/20 Luminometer (Promega, USA). All primers used to generate constructs are listed in Supplemental Table S1.

Pollen viability

To assess pollen viability, 2-month-old plants were subjected to heat treatment (37°C for 1 day) or maintained under control conditions. Pollen grains were stained using the Alexander staining method as previously described (Peterson, 2010). Flower buds prior to anthesis were collected and fixed in Carnoy's fixative (6 ethanol : 3 chloroform : 1 acetic acid) for at least 2 h. Fixed flower buds were then stained in the final staining solution at 50°C for 2 min. Stained pollen grains were observed using a light microscope (Leica ICC50, Germany).

Propidium iodide staining

Seedlings from 3-day-old plants were collected and incubated in a 10 μg/mL

propidium iodide (PI) solution (Sigma, Japan) for 1 min. Seedlings were then washed with
distilled water and observed using a Leica TCS SP5 laser scanning confocal microscope (Leica,
Germany). PI fluorescence was excited at 532 nm and detected at 590–630 nm.

RNA-seq analysis

Total RNA was extracted from 7-day-old WT, *sta1-1*, and *S307* seedlings using
RNAiso Plus (Takara, Japan). Heat stress (37°C for 3 h) was applied to 7-day-old plants, and
heat-treated samples were collected. Three independent biological replicates were prepared for
each condition. Sequencing libraries were constructed using the TruSeq Stranded mRNA
Library Prep Kit with indexed adapters and RNA sequencing was performed using the Illumina
NovaSeq platform (MacroGen, Korea). Paired-end raw sequencing reads were pre-processed
to trim adaptors and low-quality reads using CLC Genomics Workbench v20.0.4 with default
parameters (quality scores, 0.05; maximum ambiguous nucleotides, 2). Clean reads were
mapped to *Arabidopsis* reference genome (TAIR10). Intron retention ratios at the whole-
transcript level were estimated using IRFinder (v.1.3.0) (Middleton et al., 2017). Intron
retention events were defined as those showing a change in IR ratio (ΔIR ratio) greater than 0.1
with an adjusted *p*-value (p_{adj}) < 0.05. IR ratios were transformed into z-scores and visualized
as heatmaps. Volcano plots were used to assess the overall distribution of IR changes in each
comparison. RNA sequencing data have been deposited in the Gene Expression Omnibus (GEO)
under accession number XXX.

FUNDING

This work was supported by the National Research Foundation of Korea grant funded by the
Korean government (MSIT) (RS-2024-00355816 to D.-H.J., RS-2024-00407469 for D.J.Y. and
B.-h.L. and RS-2025-00564152 for B.-h.L.) and G-LAMP grant (NRF) funded by the Korean
government (MOE) (RS-2024-00441954 for B.-h.L.), Korea Basic Science Institute (National
Research Facilities and Equipment Center) grant funded by the Korean government (MOE)
(RS-2020-NF000322 for B.-h.L.), and the Basic Science Research Program (NRF) funded by
the Korean government MOE (2018R1A6A1A03025607 for W.T.K.).

ACKNOWLEDGMENTS

We would like to thank our lab members, as well as Inki Kim (Asan Medical Center, Korea), Dr. Hyun-Sook Pai (Yonsei University, Korea) and Dr. Jae-Hoon Jung (Sungkyunkwan University, Korea), for technical assistance, materials and helpful discussion. The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

H.K., K.-j.Y., S.P., and D.S. conducted the experiments and performed bioinformatic analyses.
W.T.K., D.-H.J., D.J.Y., and B.-h.L. advised on the research. All authors discussed the results.
H.K. and B.-h.L. wrote the manuscript.

REFERENCES

Bertram, K., Agafonov, D.E., Dybkov, O., Haselbach, D., Leelaram, M.N., Will, C.L., Urlaub, H., Kastner, B., Lührmann, R., and Stark, H. (2017). Cryo-EM Structure of a Pre-catalytic Human Spliceosome Primed for Activation. Cell. **170**:701-713.

Bhat, P., Chow, A., Emert, B., Ettlin, O., Quinodoz, S.A., Strehle, M., Takei, Y., Burr, A., Goronzy, I.N., Chen, A.W., et al. (2024). Genome organization around nuclear speckles drives mRNA splicing efficiency. Nature. **629**

Bond, U. (1988). Heat shock but not other stress inducers leads to the disruption of a sub-set of snRNPs and inhibition of in vitro splicing in HeLa cells. EMBO J. **7**:3509-3518.

Bond, U., and James, T.C. (2000). Dynamic changes in small nuclear ribonucleoproteins of heat-stressed and thermotolerant HeLa cells. Int J Biochem Cell Biol. **32**:643-656.

Cho, S., Yu, S.I., Park, J., Mao, Y.F., Zhu, J.K., Yun, D.J., and Lee, B.H. (2017). Accession-Dependent CBF Gene Deletion by CRISPR/Cas System in Arabidopsis. Front. Plant Sci. **8**

Choi, S., Ha, S., Wolfgeher, D.J., Kim, J.W., Go, Y.H., Cha, H.J., Bae, G.U., Kron, S.J., and Kim, W.Y. (2025). N-Terminal deleted isoforms of E3 ligase RNF220 are ubiquitously expressed and required for mouse muscle differentiation. Mol. Cell. **48**

Daguenet, E., Baguet, A., Degot, S., Schmidt, U., Alpy, F., Wendling, C., Spiegelhalter, C., Kessler, P., Rio, M.C., Le Hir, H., et al. (2012). Perispeckles are major assembly sites for the exon junction core complex. Mol Biol Cell. **23**:1765-1782.

Day, I.S., Golovkin, M., Palusa, S.G., Link, A., Ali, G.S., Thomas, J., Richardson, D.N., and Reddy, A.S.N. (2012). Interactions of SR45, an SR-like protein, with spliceosomal proteins and an intronic sequence: insights into regulated splicing. Plant J. **71**:936-947.

de Luxan-Hernandez, C., Lohmann, J., Tranque, E., Chumova, J., Binarova, P., Salinas, J., and Weingartner, M. (2022). MDF is a conserved splicing factor and modulates cell division and stress response in Arabidopsis. Life Sci. Alliance. **6**

Galisson, F., and Legrain, P. (1993). THE BIOCHEMICAL DEFECTS OF PRP4-1 AND PRP6-1 YEAST SPLICING MUTANTS REVEAL THAT THE PRP6 PROTEIN IS REQUIRED FOR THE ACCUMULATION OF THE [U4/U6.U5] TRI-SNRNP. Nucl Acids Res. **21**:1555-1562.

Häcker, I., Sander, B., Golas, M.M., Wolf, E., Karagoez, E., Kastner, B., Stark, H., Fabrizio, P., and Lührmann, R. (2008). Localization of Prp8, Brr2, Snu114 and U4/U6 proteins in the yeast tri-snRNP by electron microscopy. Nat. Struct. Mol. Biol. **15**:1206-1212.

Ilik, I.A., Malszycki, M., Lübke, A.K., Schade, C., Meierhofer, D., and Aktas, T. (2020). SON and SRRM2 are essential for nuclear speckle formation. eLife. **9**

Kim, C.H., Kim, Y.D., Choi, E.K., Kim, H.R., Na, B.R., Im, S.H., and Jun, C.D. (2016). Nuclear Speckle-related Protein 70 Binds to Serine/Arginine-rich Splicing Factors 1 and 2 via an Arginine/Serine-like Region and Counteracts Their Alternative Splicing Activity. J. Biol. Chem. **291**:6169-6181.

Kim, G.D., Yoo, S.D., and Cho, Y.H. (2018). STABILIZED1 as a heat stress-specific splicing factor
in Arabidopsis thaliana. Plant Signal Behav. **13**

Kim, G.D., Cho, Y.H., Lee, B.H., and Yoo, S.D. (2017). STABILIZED1 Modulates Pre-mRNA
Splicing for Thermotolerance. Plant Physiol. **173**:2370-2382.

Klimesová, K., Vojácková, J., Radivojevic, N., Vandermoere, F., Bertrand, E., Verheggen, C., and
Stanek, D. (2021). TSSC4 is a component of U5 snRNP that promotes tri-snRNP formation. Nat.
Commun. **12**

Krissinel, E., and Henrick, K. (2007). Inference of macromolecular assemblies from crystalline state.
J. Mol. Biol. **372**:774-797.

Ku, D., Yang, Y., and Kim, Y. (2025). RNA-associated nuclear condensates: Where the nucleus keeps
its RNAs in check. Mol. Cells. **48**

Kurogi, Y., Matsuo, Y., Mihara, Y., Yagi, H., Shigaki-Miyamoto, K., Toyota, S., Azuma, Y.,
Igarashi, M., and Tani, T. (2014). Identification of a chemical inhibitor for nuclear speckle
formation: Implications for the function of nuclear speckles in regulation of alternative pre-
mRNA splicing. Biochem. Biophys. Res. Commun. **446**:119-124.

Labun, K., Montague, T.G., Krause, M., Cleuren, Y.N.T., Tjeldnes, H., and Valen, E. (2019).
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. Nucl Acids Res.
47:W171-W174.

Lee, B.H., Kapoor, A., Zhu, J.H., and Zhu, J.K. (2006). STABILIZED1, a stress-upregulated nuclear
protein, is required for pre-mRNA splicing, mRNA turnover, and stress tolerance in Arabidopsis.
Plant Cell. **18**:1736-1749.

Li, J., Cao, Y.M., Zhang, J.X., Zhu, C.J., Tang, G.L., and Yan, J. (2023). The miR165/166-
PHABULOSA module promotes thermotolerance by transcriptionally and posttranslationally
regulating HSFA1. Plant Cell. **35**:2952-2971.

Liang, Q.Y., Peng, N.N., Xie, Y., Kumar, N., Gao, W.B., and Miao, Y.S. (2024). MolPhase, an
advanced prediction algorithm for protein phase separation. EMBO J. **43**:1898-1918.

Liu, S.B., Rauhut, R., Vornlocher, H.P., and Lührmann, R. (2006). The network of protein-protein
interactions within the human U4/U6.U5 tri-snRNP. RNA. **12**:1418-1430.

Liu, S.Y., Wang, T.S., Shi, Y.L., Bai, L., Wang, S.S., Guo, D., Zhang, Y., Qi, Y.F., Chen, C.Q., Zhang,
J.R., et al. (2021). USP42 drives nuclear speckle mRNA splicing via directing dynamic phase
separation to promote tumorigenesis. Cell Death Differ. **28**:2482-2498.

Liu, W.S., Zhu, X.H., Lei, M.G., Xia, Q.Y., Botella, J.R., Zhu, J.K., and Mao, Y.F. (2015). A detailed
procedure for CRISPR/Cas9-mediated gene editing in Arabidopsis thaliana. Sci. Bull. **60**:1332-
1347.

Ma, J., Li, S., Wang, T.Y., Tao, Z., Huang, S.J., Lin, N., Zhao, Y.B., Wang, C.H., and Li, P.J. (2025).
Cooperative condensation of RNA-DIRECTED DNA METHYLATION 16 splicing isoforms
enhances heat tolerance in Arabidopsis. Nat Commun. **16**

Makarov, E.M., Makarova, O.V., Achsel, T., and Lührmann, R. (2000). The human homologue of
the yeast splicing factor Prp6p contains multiple TPR elements and is stably associated with the
U5 snRNP via protein-protein interactions. J. Mol. Biol. **298**:567-575.

Makarova, O.V., Makarov, E.M., and Lührmann, R. (2001). The 65 and 110 kDa SR-related proteins
of the U4/U6•U5 tri-snRNP are essential for the assembly of mature spliceosomes. EMBO J.
20:2553-2563.

Middleton, R., Gao, D.D., Thomas, A., Singh, B., Au, A., Wong, J.J.L., Bomane, A., Cosson, B.,
Eyras, E., Rasko, J.E.J., et al. (2017). IRFinder: assessing the impact of intron retention on
mammalian gene expression. Genome Biol. **18**

Mintz, P.J., Patterson, S.D., Neuwald, A.F., Spahr, C.S., and Spector, D.L. (1999). Purification and
biochemical characterization of interchromatin granule clusters. EMBO J. **18**:4308-4320.

Negi, S.S., Schein, C.H., Oezguen, N., Power, T.D., and Braun, W. (2007). InterProSurf: a web server
for predicting interacting sites on protein surfaces. Bioinform. **23**:3397-3399.

Nguyen, T.H.D., Galej, W.P., Bai, X.C., Oubridge, C., Newman, A.J., Scheres, S.H.W., and Nagai,
K. (2016). Cryo-EM structure of the yeast U4/U6.U5 tri-snRNP at 3.7 Å resolution. Nature.
530:298-302.

Nishizawa-Yokoi, A., Nosaka, R., Hayashi, H., Tainaka, H., Maruta, T., Tamoi, M., Ikeda, M.,
Ohme-Takagi, M., Yoshimura, K., Yabuta, Y., et al. (2011). HsfA1d and HsfA1e Involved in
the Transcriptional Regulation of HsfA2 Function as Key Regulators for the Hsf Signaling
Network in Response to Environmental Stress. Plant Cell Physiol. **52**:933-945.

Park, S.J., Choi, S.W., Kim, G.M., Moller, C., Pai, H.S., and Yang, S.W. (2021). Light-stabilized
FHA2 suppresses miRNA biogenesis through interactions with DCL1 and HYL1. Mol. Plant.
14:647-663.

Park, S.Y., Chang, H.S., Lim, S.H., Jeong, D.H., and Kim, Y.J. (2024). Functional Characterization
of PRP43a, a Component for Splicing Disassembly in Arabidopsis. J. Plant Biol. **67**:427-435.

Paul, S., Arias, M.A., Wen, L., Liao, S.E., Zhang, J.C., Wang, X.S., Regev, O., and Fei, J.Y. (2024).
RNA molecules display distinctive organization at nuclear speckles. iScience. **27**

Peterson, R.S., J. P. Chen, C. (2010). A simplified method for differential staining of aborted and non-
aborted pollen grains. Int. J. Plant Biol. **1**:e13-e13.

Petricka, J.J., Clay, N.K., and Nelson, T.M. (2008). Vein patterning screens and the defectively
organized tributaries mutants in Arabidopsis thaliana. Plant J. **56**:251-263.

Röth, S., Mirus, O., Bublak, D., Scharf, K.D., and Schleiff, E. (2017). DNA-binding and repressor
function are prerequisites for the turnover of the tomato heat stress transcription factor HsfB1.
Plant J. **89**:31-44.

Sander, B., Golas, M.M., Makarov, E.M., Brahms, H., Kastner, B., Luehrmann, R., and Stark, H.
(2006). Organization of core spliceosomal components U5 snRNA loop I and U4/U6 Di-snRNP
within U4/U6.U5 tri-snRNP as revealed by electron cryomicroscopy. Mol Cell. **24**:267-278.

Schaffert, N., Hossbach, M., Heintzmann, R., Achsel, T., and Lührmann, R. (2004). RNAi
knockdown of hPrp31 leads to an accumulation of U4/U6 di-snRNPs in Cajal bodies. EMBO J.
23:3000-3009.

Schneider, M., Hsiao, H.H., Will, C.L., Giet, R., Urlaub, H., and Lührmann, R. (2010). Human
PRP4 kinase is required for stable tri-snRNP association during spliceosomal B complex
formation. Nat. Struct. Mol. Biol. **17**:216-U212.

Smith, K.P., Hall, L.L., and Lawrence, J.B. (2020). Nuclear hubs built on RNAs and clustered
organization of the genome. Curr. Opin. Cell Biol. **64**:67-76.

Sun, M.L., Zhang, Y.M., Wang, Q.H., Wu, C.L., Jiang, C., and Xu, J.R. (2018). The tri-snRNP
specific protein FgSnu66 is functionally related to FgPrp4 kinase in Fusarium graminearum. Mol.
Microbiol. **109**:494-508.

Utans, U., Behrens, S.E., Luhrmann, R., Kole, R., and Kramer, A. (1992). A SPLICING FACTOR
THAT IS INACTIVATED DURING INVIVO HEAT-SHOCK IS FUNCTIONALLY
EQUIVALENT TO THE [U4/U6.U5] TRIPLE SNRNP-SPECIFIC PROTEINS. Genes Dev.
6:631-641.

Wang, B., Li, J., Song, Y.Y., Qin, X.H., Lu, X., Huang, W., Peng, C.T., Wei, J.X., Huang, D.H., and
Wang, W. (2024). CLK2 Condensates Reorganize Nuclear Speckles and Induce Intron Retention.
Adv. Sci. **11**

Will, C.L., and Lührmann, R. (2011). Spliceosome Structure and Function. CSH Perspect. Biol. **3**

Wu, J.J., Xiao, Y., Liu, Y.Z., Wen, L., Jin, C.Y., Liu, S., Paul, S., He, C., Regev, O., and Fei, J.Y.
(2024). Dynamics of RNA localization to nuclear speckles are connected to splicing efficiency.
Sci. Adv. **10**

Yildirim, A., Mozaffari-Jovin, S., Wallisch, A.K., Schaefer, J., Ludwig, S.E.J., Urlaub, H.,
Luehrmann, R., and Wolfrum, U. (2021). SANS (USH1G) regulates pre-mRNA splicing by
mediating the intra-nuclear transfer of tri-snRNP complexes. Nucl Acids Res. **49**:5845-5866.

Yun, J.Y., Yu, S.I., Bang, S.E., Kim, J.Y., Lee, S.H., and Lee, B.H. (2022). Identification of CRISPR-
Induced Mutations in Plants: with a Focus on the Next-Generation Sequencing Assay. J. Plant
Biol. **65**:435-443.

Zeytuni, N., and Zarivach, R. (2012). Structural and Functional Discussion of the Tetra-Trico-Peptide
Repeat, a Protein Interaction Module. Structure. **20**:397-405.

Zhan, X.C., Yan, C.Y., Zhang, X.F., Lei, J.L., and Shi, Y.G. (2018). Structures of the human pre-
catalytic spliceosome and its precursor spliceosome. Cell Res. **28**:1129-1140.

Zhang, M.J., Gu, Z., Guo, S.H., Sun, Y.T., Ma, S.B., Yang, S., Guo, J.R., Fang, C.X., Shu, L., Ge,
Y.F., et al. (2024). SRRM2 phase separation drives assembly of nuclear speckle
subcompartments. Cell Rep. **43**

Figure Legends

Figure 1. The *dot2^S* mutation in *S307* rescues *sta1-1* defects and restores interaction with the sta1-1 protein

(A) Growth phenotypes and fresh weights of 3-week-old WT, *sta1-1*, and *S307* plants (scale bar = 5 cm; n = 22).

(B) Heatmap showing IR ratios in WT, *sta1-1*, and *S307*. Three independent biological replicates were analyzed for each genotype.

(C) Volcano plots showing significantly repressed and induced IR events. Blue and red dots represent significantly decreased and increased IR events, respectively.

(D) Growth phenotypes of 3-week-old WT, *sta1-1*, *S307*, *S307^{DOT2END}*, and *S307^{DOT2OX}* plants (scale bar = 5 cm).

(E) Experimental validation of IR levels for representative genes (*Leaf curly and small 1* (*FCS1*, At1g78910), *BOI-RELATED GENE 1* (*BRG1*, At5g45100), and *BETA GLUCOSIDASE 6* (*BGLU6*, At1g60270). Box diagrams show spliced and unspliced isoforms. *Protein phosphatase 2A* (*PP2A*, At1g13320) is used as a loading control.

(F and G) Semi-dominant effects of the *dot2^S* mutation on plant growth and splicing in the *sta1-1* mutant background (scale bar = 1 cm). Three-week-old plants derived from an F2 population segregating for both the *sta1-1* and *dot2^S* alleles (WT x *S307* cross). Genotypes are indicated as follows; S, *STA1*; s, *sta1-1*; D, *DOT2*; d^S, *dot2^S*. *PP2A* is used as a loading control. Box diagrams showed spliced and unspliced isoforms.

(H) Y2H assays demonstrating the interaction between STA1/sta1-1 and DOT2/dot2^S. Yeast cells co-transformed with the fusion constructs of the GAL4 DNA-binding domain and activation domain were grown overnight and spotted onto selection medium (-L/-W), and interaction medium (-L/-W/-H) supplemented with 0.5 mM 3-amino-1,2,4-triazole (3-AT).

(I and J) BiFC analysis for detecting the interaction between STA1/sta1-1 and DOT2/dot2^S/dot2^C in leaves of *N. benthamiana* (scale bar = 75 μm). Red arrows indicate EYFP signals representing protein-protein interactions in the nucleus. EYFP signals were quantified from identical areas of leaf disks (2 x 2 cm). Data are presented as mean ± SE (n = 5). Statistical analysis is performed using one-way ANOVA with Tukey's HSD test (**$P < 0.01$; n.s., not significant).

Figure 2. The disrupted nuclear localization of sta1-1 protein is re-localized to NS by the dot2^S protein

(A) Classification of nuclear compartments showing the localization of gene of interest (GOI)-GFP driven by the CaMV 35S promoter into five subclasses. The nucleolus marker FIBRILLARIN 2 (FIB2, At4g25630)-mRFP was co-infiltrated (scale bar = 10 μm).

(B) Subcellular localization of each protein tagged with GFP in leaves of *N. benthamiana*. Nuclear localization patterns of each GFP-tagged protein were classified into the five distinct categories defined in **(A)**. For each set, 100 GFP-expressing cells were analyzed per replicate, and data are presented as mean ± SE (n = 4). Statistical analysis is performed using one-way ANOVA with Tukey's HSD test (**$**P < 0.01$; n.s., not significant).

(C-E) Co-localization of GFP-tagged STA1/sta1-1 with mRFP-tagged DOT2/dot2^S/dot2^C in leaves of *N. benthamiana*. Nuclear localization categories are shown in **(D)**, classified according to the groups defined in **(A)**. The number of NS per nucleus is shown in **(E)**, based on the STA1/sta1-1-GFP signals upon co-expression with mRFP-DOT2/dot2^S/dot2^C. For each analysis, 100 cells **(D)** or 50 nuclei **(E)** were randomly selected and analyzed (scale bar = 10 μm).

(F-H) Co-localization of GFP-tagged STA1 **(F and H)** or EYFP signals from BiFC analysis of STA1 and DOT2 **(G)** with the NS marker U1 SMALL NUCLEAR RIBONUCLEOPROTEIN-70K (U1-70K, At3g50670) tagged with mRFP (mRFP-U1-70K) in leaves of *N. benthamiana*. Red boxes indicate regions of overlap between STA1-GFP and mRFP-U1-70K (scale bar = 10 μm).

Figure 3. The localization of STA1 to NS is closely associated with proper pre-mRNA splicing

(A) Nuclear localization patterns of GFP-tagged STA1/sta1-1 upon transfection with DOT2/dot2^S/dot2^C in *Salk007933$^{STA1-GFP}$* or *Salk007933$^{sta1-1-GFP}$* protoplasts. Cells with nuclear-localized GFP signals of two distinct patterns (nucleoplasm and NS) were quantified as a ratio. A total of 15 cells were analyzed for each combination (scale bar = 5 μm).

(B) Splicing activity of DOT2/dot2^S/dot2^C on the *gHSFA3* DNA in *Salk007933$^{sta1-1-GFP}$*

protoplasts. Splicing activity on the *gHSFA3* DNA is represented by the relative expression of

NanoLuc luciferase (nLUC) compared with firefly luciferase (fLUC). *Clathrin* (At4g24550)

was used as a loading control. Data are presented as mean ± SE (n = 2). Statistical analysis is

performed using one-way ANOVA with Tukey's HSD test (**P < 0.01; n.s., not significant).

(C) Tubercidin-induced delocalization of STA1 from NS to nucleoplasm. Ten-day-old

Salk007933$^{STA1-GFP}$ plants were treated with DMSO or 100 μM tubercidin for 1 h (scale bar =

25 μm).

(D) Tubercidin-induced disruption of pre-mRNA splicing. Ten-day-old WT, *sta1-1*, and *S307*

plants were treated with DMSO or 100 μM tubercidin for 12 h. The box diagrams show spliced

and unspliced isoforms. *PP2A* is used as a loading control.

(E) Abolished size differences between WT, *sta1-1*, and *S307* by tubercidin treatment (scale

bar = 1 cm). Five-day-old seedlings were transferred to medium containing either DMSO or

10 μM tubercidin for 6 d. Data are presented as mean ± SE (n = 4).

**Figure 4. The dot2^S protein facilitates NS localization of the sta1-1 protein under heat
conditions**

(A-C) BiFC analysis to detect interaction between STA1 and DOT2 under heat stress

conditions in leaves of *N. benthamiana* (scale bar = 75 μm). Red arrows indicate EYFP signals

representing protein-protein interactions in the nucleus. The number of EYFP signals per unit

area (2 x 2 cm) or per nucleus (n = 43) was quantified in **(B)** and **(C)**.

(D and E) BiFC analysis to detect interaction between STA1/sta1-1 and DOT2/dot2^S/dot2^C

under heat stress conditions in leaves of *N. benthamiana* (scale bar = 75 μm). Red arrows

indicate EYFP signals. EYFP signals were quantified from the same area of leaf disks (1 x 1

cm). Data are presented as means from three independent biological replicates. Statistical

analysis is performed using one-way ANOVA with Tukey's HSD test. Significant differences

are denoted by letters above the plots.

(F and G) Co-localization of EYFP signals from BiFC analysis of STA1/sta1-1 and

DOT2/dot2^S/dot2^C with the NS marker mRFP-U1-70K under heat stress conditions in leaves

of *N. benthamiana* (scale bar = 10 μm). Box plots show the number of NS exhibiting EYFP

signals. Center lines indicate medians; boxes represent interquartile range, and whiskers extend

to minimum and maximum values. For heat treatment, samples were incubated at 37°C for 1 h

in liquid MS medium following a 36 h post-infiltration incubation.

Figure 5. Heat-sensitivity of *sta1-1* is restored to wild type levels in the *S307* suppressor

(A and B) Heat survival test and quantification of survival rate of each plant genotype. Ten-

day-old plants of each genotype were exposed to heat stress (37°C for 48 h), and survived

plants were scored. Statistical analysis is performed using one-way ANOVA with Tukey's HSD

test (**$P < 0.01$; *$P < 0.05$; n.s., not significant).

(C) Assessment of pollen viability using Alexander staining (scale bar = 50 μm). Viable pollen

grains appear purple, while non-viable pollen grains are transparent. Red arrows indicate non-

viable pollen grains. Data are presented as means ± SE from three independent biological

replicates. Statistical analysis is performed using one-way ANOVA with Tukey's HSD test.

Significant differences are denoted by letters above the plots.

Figure 6. Defective splicing of *HSFA1s* in *sta1-1* is rescued in *S307*, leading to proper

expression of downstream target genes

(A) Heatmap showing the IR ratios in WT under heat stress (WT(H)), *sta1-1* under heat stress

(*sta1-1*(H)), and *S307* under heat stress (*S307*(H)). Each genotype is represented by three

independent biological replicates.

(B) Boxplot analysis showing IR ratio trends across genotypes for 67,103 transcripts exhibiting

significant changes in intron retention. Each box represents the IR ratio distribution for each

genotype, with lines connecting genotypes to indicate intron-specific changes. The ribbon

highlights the median and variability for visual summary, and the bold red line indicates the

overall trend of median IR ratios across genotypes. Median IR ratios for each genotype are

shown in parentheses. Statistical significance among genotypes was assessed using the

Kruskal–Wallis test followed by a post hoc Dunn's test with Benjamini–Hochberg correction.

Significant differences are denoted by letters above the plots.

(C) Changes in intron retention levels of transcripts in response to heat stress across different

genotypes. Paired notation denotes (number of introns/number of genes).

(D) Venn diagram showing transcripts with increased intron retention under heat stress in each

genotype.

(E and F) Quantification and validation of splicing defects in the master heat stress regulators

*HSFA1*s. Box diagrams show spliced and unspliced isoforms. *PP2A* is used as a loading control.

(G) qRT-PCR analysis of heat stress-related genes regulated by *HSFA1*s. The relative

expression levels of each gene are shown in the graphs. Data are presented as mean ± SE from

three independent biological replicates. Statistical analysis is performed using one-way

ANOVA with Tukey's HSD test (**$P < 0.01$; *$P < 0.05$; n.s., not significant).

Figure 7. A proposed model for STA1-DOT2 interaction-mediated NS formation and pre-

mRNA splicing in growth and heat stress responses

The STA1-DOT2 interaction, presumably reflecting the U4/U6·U5 tri-snRNP assembly, is

important for NS formation. Proper NS formation ensures the production of correctly spliced

transcripts required for normal growth and heat stress responses. The *sta1-1* mutation or heat

stress weakens the STA1-DOT2 interaction, thereby reducing NS formation. This disrupted

subnuclear localization of splicing components increases intron-retained transcripts, resulting

in defective growth and heat stress responses.

Supplemental Figure 1. Positional cloning and the identification of the *dot2^S* mutation in

S307

(A) Map of the *DOT2* locus on chromosome 5. Ninety-four F2 seedlings derived from an *S307*

x L*er* cross were used for mapping. The Simple sequence length polymorphism (SSLP)

markers used for mapping, Ch5_5.46Mb and Ch5_5.71Mb, are shown on the 184A1 and 35B3

BAC clones, respectively. Recombination numbers are shown under each marker locus.

(B) The mutation information of *dot2^S* mutation, including nucleotide and corresponding amino

acid changes.

Supplemental Figure 2. Homologs and expression patterns of *DOT2*, a U4/U6·U5 tri-

snRNP associated protein

(A) Multiple sequence alignment of DOT2 homologs.

(B) Expression patterns of *DOT2* and *STA1::GUS* in *Arabidopsis*. GUS signals are observed in

whole seedling, flower, silique, and trichome from left to right.

Supplemental Figure 3. Phenotypes of the *dot2* single mutant alleles

(A) Growth phenotypes of 3-week-old WT, *sta1-1*, *S307*, *dot2^S*, and *dot2^T* (SAIL_775_F10)

plants (scale bar = 5 cm).

(B) Splicing defects in marker genes across genotypes. Box diagrams show spliced and

unspliced isoforms. *PP2A* is used as a loading control.

(C) Representative confocal images of PI-stained root tips of 3-day-old WT, *sta1-1*, *S307*, *dot2^S*,

and *dot2^T* seedlings (scale bar = 25 μm). Arrowheads indicate dead cells, and white bars mark

the division zone. Division zone length in 3-day-old seedlings is quantified (n = 10). Statistical

analysis is performed using one-way ANOVA with Tukey's HSD test (**$P < 0.01$; n.s., not

significant).

Supplemental Figure 4. Residue positions of mutations in sta1-1 and dot2 mutant proteins

(A) Predicted protein structures of *Arabidopsis* STA1 and DOT2. In the model, STA1 protein

is shown in green and DOT2 protein in blue, with their respective mutations highlighted as

follows: the *sta1-1* mutation in magenta, the *dot2^S* mutation in red, and the *dot2^C* mutation in

orange.

(B) Domain structure of STA1 and DOT2. Predicted STA1-DOT2 interaction regions identified using InterProSurf and PDBePISA are indicated by red lines (STA1: 113−123 aa, 185−202 aa; DOT2: 548−569 aa). Domains are labeled as follows: UBQ, ubiquitin-like; PRP1P, pre−mRNA processing protein 1; HAT, half-A-tetratricopeptide repeat; NLS, nuclear localization signal; TPR, tetratricopeptide repeat; HIND, hub1−interacting domain.

(C) The $dot2^S$ and $dot2^C$ mutations, showing nucleotides and the corresponding amino acid changes.

(D) Growth phenotypes of 3-week-old WT, *sta1-1*, *S307*, and the *sta1-1 dot2^C* double mutant (*ssd^{C}d^C*) plants (scale bar = 5 cm).

(E) Splicing defects in the marker genes across genotypes. The box diagrams show spliced and unspliced isoforms. *PP2A* is used as a loading control.

Supplemental Figure 5. Rescue of *Salk007933* lethality through introduction of STA1-GFP or sta1-1-GFP, resulting wild-type- and *sta1-1*-like phenotypes, respectively
Growth phenotypes of 3-week-old $Salk007933^{STA1-GFP}$ and $Salk007933^{sta1-1-GFP}$ plants (scale bar = 5 cm).

Supplemental Figure 6. Increased STA1-aggregates under heat conditions
(A and B) Subcellular localization of GFP-tagged STA1 and EYFP signals from BiFC analysis of STA1-STA1 interaction under heat conditions (scale bar = 75 μm).
(C and D) Co-localization of EYFP signals from BiFC analysis of STA1-STA1 interaction with the mRFP-U1-70K in leaves of *N. benthamiana* (scale bar = 10 μm).
(E) MolPhase prediction analysis of IDR region within STA1 (Liang *et al.*, 2024).
For heat treatment, samples were incubated at 37°C for 1 h in liquid MS medium following a 36 h post-infiltration incubation.

Supplemental Figure 7. Intron-retained *HSFA1* transcripts in *sta1-1*
(A) Increased IR events of *HSFA1s* in *sta1-1* (H). IR events in *WT*(H), *sta1-1*(H), and *S307*(H)

are visualized using the Integrative Genomics Viewer (IGV).

(B) Protein domain structures of premature termination codon (PTC)-containing intron-

retained pre-mRNA isoforms in *sta1-1* (H). Domains are labeled as follows: DB, DNA-binding

domain; OD, oligomerization domain; NLS, nuclear localization signal; AHA1/2, aromatic and

large hydrophobic amino acid residues embedded in acidic surroundings. PTCs are labeled

with a red symbol.

Supplemental Figure 8. Splicing defects of *HSFA1*s in *sta1-1 dot2^C* double mutants

Splicing defects of the master heat stress regulators *HSFA1*s in the *sta1-1 dot2^C* double mutant

(*ssd^{C}d^C*) plants. Box diagrams show spliced and unspliced isoforms. *PP2A* is used as a loading

control.

STA1-DOT2 nuclear speckle
sta1-DOT2 nuclear speckle
STA1-DOT2 aggregates

sta1-DOT2 weak interaction

Heat

Normally spliced transcripts

Intron retained transcripts

Intron retained transcripts

Growth and Heat Tolerance

A Heat 48h

WT

sta1-1

S307

S307^(R85S)

sta1-1 csa1-1 csa2-1 csa3-1

Survival rate (%)

B Heat 48h

WT

sta1-1

S307

sta1^(ΔF)

Survival rate (%)

C

WT   *sta1-1*   *S307*   *S307^(R85S)*   *sta1^(ΔF)*

Heat 0h

Heat 24h

Pollen viability (%)

■ Non-viable □ Viable

WT   *sta1-1*   *S307*   *S307^(R85S)*   *sta1^(ΔF)*