The STA1–DOT2 interaction promotes nuclear speckle formation and splicing robustness in growth and heat stress responses

Heejin Kim[1], Kyoung-jae Yu[1], So Young Park[2], Dong Hye Seo[3, 4, 5], Woo Taek Kim[5], Dae-Jin Yun[6], Jae-Hoon Jung[7], Dong-Hoon Jeong[2], and Byeong-ha Lee[1*]

[1]Department of Life Science, Sogang University, Mapo-gu, Seoul 04107, Republic of Korea

[2]Department of Life Science and Multidisciplinary Genome Institute, Hallym University, Chuncheon 24252, Republic of Korea

[3]Plant Biotechnology Research Center, Ghent University, Global Campus, Incheon 21985, Republic of Korea

[4]Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent, Belgium

[5]Department of Systems Biology and Institute of Life Science and Biotechnology, Yonsei University, Seoul 03722, Republic of Korea

[6]Department of Advanced Biotechnology and Global Plant Stress Research Center, Konkuk University, Seoul 05029, Republic of Korea

[7]Department of Biological Sciences, Sungkyunkwan University, Suwon 16419, Republic of Korea

Contact information

Heejin Kim: gmlwls7686@sogang.ac.kr / 0009-0000-2341-5581

Kyoung-jae Yu: kyoungjae@sogang.ac.kr / 0009-0003-1125-572X

So Young Park: soyoungp@hallym.ac.kr / 0000-0001-9202-0874

Dong Hye Seo: Donghye.Seo@ghent.ac.kr / 0000-0003-1243-4466

Woo Taek Kim: wtkim@yonsei.ac.kr / 0000-0001-5027-4622

Dae-Jin Yun: djyun@konkuk.ac.kr / 0000-0002-3638-6043

Jae-Hoon Jung: jhjung19@skku.edu / 0009-0001-9562-526X

Dong-Hoon Jeong: dhjeong73@hallym.ac.kr / 0000-0002-2378-1706

*Correspondence: Byeong-ha Lee: byeongha@sogang.ac.kr / 0000-0001-7633-3264

ABSTRACT

Pre-mRNA splicing is carried out by the spliceosome, a large and dynamic ribonucleoprotein complex. The spliceosome is known to be stored in nuclear speckles (NS), which are now recognized as active subnuclear organelles for splicing. However, it remains poorly understood how spliceosomal protein-protein interactions are functionally coupled to NS organization to maintain splicing robustness in plants. Here, we report the functional significance of a specific interaction between two U4/U6·U5 tri-snRNP components of the spliceosome, STA1 and DOT2, in regulating NS organization, pre-mRNA splicing, and heat stress responses in *Arabidopsis*. We identified a missense mutation in *DOT2* (a Snu66/SART1 homolog) from a genetic suppressor of the PRP6 homolog mutant *sta1-1* (named *S307*). This mutation restored the weakened interaction between STA1 and DOT2 in the *sta1-1* mutant background. Genetic, biochemical, and cell biological analyses showed that variation in the strength of the STA1-DOT2 interaction was closely associated with changes in NS formation, splicing efficiency, as well as growth and heat tolerance. Pharmacological inhibition of STA1-associated NS formation by tubercidin recapitulated *sta1-1*-like phenotypes and splicing defects, supporting a functional link between NS organization and splicing outcomes. In addition, heat-induced weakening of the STA1-DOT2 interaction was accompanied by reduced NS formation and increased intron retention at the transcriptome-wide level including key heat-responsive transcripts. Based on these observations, we propose that the STA1-DOT2 interaction, likely reflecting the assembly state of the U4/U6·U5 tri-snRNP, functions as a heat-sensitive interaction node that couples spliceosome assembly to NS organization and splicing robustness under stress conditions.

Keywords: *STA1*, *DOT2*, U4/U6·U5 tri-snRNP, spliceosomal protein interaction, splicing, nuclear speckle

INTRODUCTION

Pre-mRNA splicing is an essential step in eukaryotic gene expression and is carried out by the spliceosome, a large and dynamic ribonucleoprotein complex composed of five small nuclear ribonucleoproteins (snRNPs) and numerous auxiliary factors (Albakri et al., 2025; Will and Lührmann, 2011). Among spliceosomal subcomplexes, the U4/U6·U5 tri-snRNP plays a pivotal role in spliceosome activation, and undergoes extensive structural rearrangements during the splicing cycle (Sander et al., 2006; Will and Lührmann, 2011). Defects in tri-snRNP assembly or stability result in splicing failure, underscoring the importance of precise and coordinated interactions among its components (Klimesová et al., 2021; Schneider et al., 2010).

Arabidopsis STABILIZED1 (STA1) is homologous to the pre-mRNA splicing factors PRP1, Prp6p, and PRPF6 (hereafter collectively referred to as PRP6, unless otherwise indicated) in *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae*, and *Homo sapiens*, respectively. Within the tri-snRNP, PRP6 resides in the U5 snRNP and interacts with components of the U4/U6 di-snRNP, thereby physically bridging the U5 snRNP and the U4/U6 di-snRNP (Bertram et al., 2017; Liu et al., 2006; Makarov et al., 2000; Nguyen et al., 2016; Zhan et al., 2018). Consistent with this role, depletion or mutation of PRP6 in yeast and human cells disrupts tri-snRNP formation and leads to the accumulation of U4/U6 di-snRNP, indicating that PRP6 is essential for tri-snRNP assembly (Galisson and Legrain, 1993; Schaffert et al., 2004).

The *Arabidopsis sta1-1* mutant carries a 6-bp deletion within the half-a-tetratricopeptide (HAT) repeat domain of STA1 (Lee et al., 2006). HAT domains are known to mediate protein–protein interactions and promote the assembly of multiprotein complexes (Zeytuni and Zarivach, 2012). Supporting this, yeast two-hybrid assays using human tri-snRNP components have shown that PRPF6 interacts with several tri-snRNP components through its HAT domain (Liu *et al.*, 2006). These findings suggest that the PRP6 homolog STA1 contributes to pre-mRNA splicing in plants primarily by promoting proper spliceosome assembly through its interactions with other tri-snRNP components.

In the nucleus, nuclear speckles (NS) are prominent subnuclear membraneless compartments enriched in spliceosomal proteins and RNAs (Ilik et al., 2020; Smith et al., 2020). NS serve as sites for the storage, assembly, and dynamic exchange of splicing factors and are closely associated with transcriptionally active chromatin, and are increasingly recognized as sites that actively contribute to efficient pre-mRNA processing (Daguenet et al., 2012; Wu et al., 2024). NS are thought to form through the self-assembly of resident proteins and RNAs and to be maintained by transient protein-protein and protein-RNA interactions (Choi et al., 2025; Day et al., 2012; Kim et al., 2016; Paul et al., 2024). Importantly, localization of spliceosomal components to NS enhances the splicing efficiency of nearby genes, whereas impaired speckle association leads to splicing defects (Liu et al., 2021; Wang et al., 2024).

Several lines of evidence suggest that tri-snRNP components contribute directly to NS organization. In human cells, depletion of PRPF6 disrupts the NS localization of another tri-snRNP component PRPF31 and causes splicing defects (Yildirim et al., 2021). In plants, NS localization of splicing factors has also been implicated in stress adaptation. For example, two alternatively spliced isoforms of RDM16 (a tri-snRNP component PRPF3 homolog) cooperatively promote nuclear condensation and enhance heat tolerance in *Arabidopsis* (Ma et al., 2025). These studies support the idea that NS organization is functionally linked to spliceosome integrity and splicing outcomes, particularly under stress. However, it remains unclear how the interaction-dependent assembly state of the tri-snRNP contributes to NS organization in plants.

In this study, from a *sta1-1* suppressor line (*S307*), we identified a missense mutation in *DEFECTIVELY ORGANIZED TRIBUTARIES2* (*DOT2*; also known as *MERISTEM-DEFECTIVE*, *MDF* (de Luxan-Hernandez et al., 2022; Petricka et al., 2008)), hereafter referred to as *dot2^S*. DOT2 is a U4/U6·U5 tri-snRNP component homologous to yeast Snu66p and human SART1, which are known to bind to and stabilize the PRP6-involved bridge within the tri-snRNP (Bertram *et al.*, 2017; Zhan *et al.*, 2018). The *dot2^S* suppressor allele restores the weakened interaction between STA1 and DOT2 in the *sta1-1* background, leading to recovery of NS formation, improved pre-mRNA splicing, and restoration of growth and heat tolerance. Our results support the idea that the STA1-DOT2 interaction

functions as a critical and heat-sensitive interaction node within the tri-snRNP, contributing

to spliceosome assembly to NS organization, and splicing robustness under stress conditions.

RESULTS

A mutation in DOT2, a U4/U6·U5 tri-snRNP-associated protein, suppresses the *sta1-1*

defects

The *sta1-1* mutants, defective in the PRP6 homolog *STA1* gene, are small in size and

hypersensitive to heat (Kim et al., 2017; Lee *et al.*, 2006). To elucidate the molecular

mechanism underlying these phenotypes, we employed a forward genetic approach to

identify *sta1-1* modifiers, aiming to uncover genetic interactors of *STA1*. From an ethyl

methanesulfonate (EMS)-treated *sta1-1* mutant pool, we isolated a *sta1-1* suppressor *S307*

based on its size recovery. *S307* suppressed the small size phenotype of *sta1-1*, exhibiting the

phenotype similar to wild-type (WT, Columbia harboring a *RD29A* promoter-driven

luciferase transgene) (Figure 1A). In addition to the restored size, the splicing defects

observed in *sta1-1* were also restored to wild-type levels in *S307* (Figures 1B and 1C). Intron

retention (IR) profiles of *S307* were more similar to those of WT rather than to those of *sta1-*

1, as shown by heatmaps (Figure 1B). Only 16 transcripts (16 increased, 0 decreased)

displayed differential IR between *S307* and WT, in contrast to 2,876 transcripts (2,860

increased, 16 decreased) between s*ta1-1* and WT (Figure 1C, Supplemental Table 1).

Whole-genome sequencing combined with map-based positional cloning using the F2

suppressor seedlings from a cross between *S307* and Landsberg *erecta* identified a G-to-A

point mutation in the 8th exon of the *DEFECTIVELY ORGANIZED TRIBUTARIES 2* (*DOT2*,

At5g16780) gene. This mutation results in an amino acid change from aspartate to asparagine

(Supplemental Figure 1). The *DOT2* gene encodes a protein homologous to Snu66p in

Saccharomyces cerevisiae and SART1 in *Homo sapiens* (Supplemental Figure 2A), both of

which are components of the U4/U6·U5 tri-snRNP together with STA1. Indeed, *DOT2*

promoter-driven GUS activity was detected across all tissues, as found in the *STA1::GUS*

plants (Lee et al., 2006) (Supplemental Figure 2B). This widespread and overlapping

expression pattern suggests constant and cooperative roles for DOT2 and STA1.

To further examine whether DOT2 functions together with STA1, we characterized

two independent *dot2* single mutant alleles: a *dot2* single mutant carrying the *S307* allele and

a *dot2* T-DNA insertion mutant (SAIL_775_F10, also known as *mdf-2*) (de Luxan-Hernandez

et al., 2022), hereafter referred to as *dot2^S* and *dot2^T*, respectively. *dot2^S* plants were

comparable to WT in both growth and splicing patterns, whereas *dot2^T* plants showed growth

and splicing defects resembling those of *sta1-1* (Supplemental Figures 3A and 3B).

Consistent with previous reports (de Luxan-Hernandez *et al.*, 2022), *dot2^T* plants displayed

cell death in the root apical meristem (RAM) as revealed by propidium iodide (PI)-stained

confocal imaging. However, no detectable RAM defects were observed in the *dot2^S* plants

(Supplemental Figure 3C). Together, these results indicate that *dot2^S* represents a weak loss-

of-function allele and that *dot2* loss-of-function mutants display *sta1-1*-like defects,

suggesting a coordinated function for DOT2 and STA1 in growth and pre-mRNA splicing.

157 Genetic and molecular assays confirm *dot2^S* as a semi-dominant suppressor of *sta1-1*

To confirm that the *dot2^S* mutation is the causal mutation, a genomic fragment of

DOT2 including its endogenous promoter was introduced into *S307* via *Agrobacterium*-

mediated transformation. Unexpectedly, transgenic plants expressing this genomic fragment

of *DOT2* (*S307^{DOT2END}*) did not revert the WT-like phenotype of *S307* to the small-sized *sta1-

1* phenotypes (Figure 1D). However, when the *DOT2* gene was overexpressed by the

Cauliflower Mosaic Virus (CaMV) 35S promoter in *S307* (*S307^{DOT2OX}*), the overexpressing

plants exhibited the *sta1-1*-like small size (Figure 1D). In addition, compared to *S307^{DOT2END},*

S307^{DOT2OX} exhibited higher accumulation of intron-retained transcripts for several marker

genes that show IR in *sta1-1*, showing stronger phenotypic reversion by the overexpression

(Figure 1E). We speculated that phenotypic reversion only by *DOT2* overexpression might be

due to the dominant or semi-dominant effects of the dot2^S mutant protein on sta1-1, relative

to those of the wild-type DOT2 protein; the dot2^S protein may interact more effectively with

the sta1-1 protein than the DOT2 protein. Thus, we performed a dominance test on *dot2^S* by

genetically combining *dot2^S* with the *STA1* homozygotes (WT) or the *sta1-1* homozygotes.

No clear size defects were observed in the combinations of *dot2^S* mutation in the wild-type

background (Figure 1F). In contrast, the *dot2^S* heterozygote in the *sta1-1* background showed

an intermediate size, supporting the notion of semi-dominant effects of dot2^S on sta1-1

(Figure 1F). Consistently, the *dot2^S* heterozygote in the *sta1-1* background exhibited

intermediate levels of splicing defects between the *DOT2* (i.e., *sta1-1*) and the *dot2^S*

homozygotes in *sta1-1* (i.e., *S307*) (Figure 1G). Taken together, these results confirm that the

mutation in *DOT2* is responsible for the *sta1-1* phenotypic suppression in *S307*.

The *dot2^S* mutation restores physical interaction with sta1-1, rescuing *sta1-1* defects

Homologs of STA1 and DOT2, as well as STA1 and DOT2 themselves, are known to

interact, as supported by structural and experimental evidence (de Luxan-Hernandez *et al.*,

2022; Häcker et al., 2008; Liu *et al.*, 2006; Sun et al., 2018; Zhan *et al.*, 2018). Our structural

modeling predicted that the *sta1-1* and *dot2^S* mutations are positioned near the STA1-DOT2

interaction interface (Supplemental Figure 4A). The *sta1-1* mutation resides within the HAT

domain which mediates protein-protein interactions (Zeytuni and Zarivach, 2012), and the

dot2^S mutation maps to the predicted STA1-interacting surface identified by InterProSurf and

PDBePISA (Krissinel and Henrick, 2007; Negi et al., 2007) (Supplemental Figure 4B). These

findings suggest that both mutations may alter the STA1-DOT2 interaction. We examined

interactions between the wild type and mutant proteins using yeast two-hybrid (Y2H) assays.

As expected, STA1 and DOT2 showed substantial interaction (Figure 1H), whereas

interactions between a wild-type protein (STA1 or DOT2) and a mutant protein (sta1-1 or

dot2^S) were not detectable. Interestingly, and consistent with our expectation, sta1-1 and

dot2^S showed interactions comparable to those between STA1 and DOT2 (Figure 1H).

Consistent with these observations, bimolecular fluorescence complementation (BiFC)

analysis further confirmed the sta1-1-dot2^S interaction similar to the STA1-DOT2 interaction

(Figures 1I and 1J).

To determine whether the suppressive effect is allele-specific, we generated an

additional *dot2* allele using CRISPR/Cas9 (herein, called *dot2^C*). The *dot2^C* mutation is a 3-

bp deletion located 44 bp upstream of the *dot2^S* mutation, outside the predicted STA1-DOT2

interaction interface (Supplemental Figures 4A-4C). Unlike dot2^S, the dot2^C protein did not

alter interaction strength with STA1 or sta1-1 proteins (Figures 1I and 1J). Consistent with

this, *dot2^C* mutation did not alter the defects in size or intron retention observed in *sta1-1*

(Supplemental Figures 4D and 4E). These results indicate that $dot2^S$ is a specific suppressor

allele whose product interacts with the sta1-1 mutant protein, thereby likely restoring the

U4/U6·U5 tri-snRNP assembly in *S307*.

Interaction between STA1 and DOT2 promotes NS formation

We noticed diverse nuclear localization signals in our BiFC assays. Thus, we

examined the localization of each GFP-tagged protein (STA1/sta1-1 and DOT2/dot2^S/dot2^C)

and classified the nuclear signals into five subclasses based on their predominant subnuclear

compartments (Figure 2A). Our observation revealed that STA1-GFP formed nuclear

condensates (Class IV and V) at a higher frequency than sta1-1-GFP, whereas GFP-

DOT2/dot2^S/dot2^C were predominantly localized to the nucleolus (Class II) (Figure 2B).

Interestingly, co-expression of STA1-GFP and mRFP-DOT2 significantly increased

condensate formation of both proteins compared to expression of each protein alone (from

~15%/0% to ~92% for STA1/DOT2) (Figures 2A-2E). This suggests that the STA1-DOT2

interaction contributes to the localization of each protein to nuclear condensates. This idea

was supported by the following observations: sta1-1-GFP showed limited condensate

formation when co-expressed with mRFP-DOT2/dot2^C, whereas mRFP-dot2^S strongly

enhanced sta1-1-GFP condensate formation (Figures 2C-2E), consistent with the interaction

strengths of these protein combinations. To identify the nature of the nuclear condensates

observed above, we co-expressed each protein with the NS marker mRFP-U1-70K (Ali et al.,

2008). STA1-GFP condensates formed upon single expression of STA1-GFP showed partial

co-localization with mRFP-U1-70K (Figure 2F). Upon co-expression with DOT2, STA1-GFP

became nearly completely co-localized with mRFP-U1-70K (Figure 2G). Furthermore, the

EYFP signals from the BiFC assay (i.e., co-expression of STA1 and DOT2), indicating the

STA1-DOT2 interaction, showed complete co-localization with mRFP-U1-70K (Figure 2H).

These results confirm that the STA1-DOT2 interaction promotes the formation of nuclear

condensates corresponding to NS. NS are a specific type of nuclear condensates that are

enriched in the spliceosome components and associated with mRNA splicing and mRNA

metabolism (Bhat et al., 2024; Daguenet *et al.*, 2012). Therefore, our results indicate that the

STA1-DOT2 interaction contributes to their NS localization and probably to proper mRNA

splicing. It should be noted that STA1-GFP condensates that did not overlap with mRFP-U1-
70K suggest that STA1-GFP may form non-functional or NS-like condensates (herein called
"aggregates") (see below).

The $dot2^S$ mutation restores sta1-1 localization to NS, enabling proper splicing

We next examined the functional significance of NS formation of STA1 and DOT2
for mRNA splicing. To this end, we generated the STA1-GFP/sta1-1-GFP overexpressing
plants in *Salk007933*, a *sta1* T-DNA insertion knock-out line (hereafter referred to as
Salk007933$^{STA1-GFP}$ and *Salk007933$^{sta1-1-GFP}$*, respectively). Homozygous *sta1* knockouts are
lethal. However, introduction of either STA1-GFP or sta1-1-GFP rescued this lethality and
the resulting plants phenocopied the wild-type and *sta1-1* plants, respectively (Supplemental
Figure 5). Using protoplasts isolated from these GFP-expressing plants, we confirmed again
that the interaction between STA1 and DOT2 (or sta1-1 and dot2^S) promoted the localization
of the GFP-tagged protein to NS (Figure 3A).

Previously, we developed a protoplast-based splicing activity assay using a
luciferase reporter gene fused with a genomic fragment of *HSFA3* (*gHSFA3-LUC*) which
undergoes STA1-dependent splicing (Kim *et al.*, 2017). After transfection with the alleles to
be tested, splicing activity was measured in protoplasts derived from *Salk007933$^{sta1-1-GFP}$*. As
expected, the introduction of *STA1* into the protoplasts successfully restored full splicing
activity of *gHSFA3-LUC*, while *sta1-1* did not (Figure 3B). Among the tested *DOT2* alleles,
only the *dot2^S* allele showed the full splicing activity in *Salk007933$^{sta1-1-GFP}$* (Figure 3B).
Taken together, these results indicate that the STA1-DOT2 or sta1-1-dot2^S interaction is
critical for NS formation and splicing activity, thereby linking NS formation to proper
splicing activity.

We then examined the effect of disrupting NS formation by applying tubercidin, an
inhibitor of speckle formation (Kurogi et al., 2014) to *Salk007933$^{STA1-GFP}$* plants. We observed
that STA1-GFP in *Salk007933$^{STA1-GFP}$* was localized predominantly to NS (Figure 3C).
However, tubercidin treatment dispersed the GFP signals into the nucleoplasm (Figure 3C).
Tubercidin treatment also eliminated the differences in splicing activity and growth patterns

among WT, *sta1-1*, and *S307*, resulting in comparable splicing and growth phenotypes across

all genotypes (Figures 3D and 3E). Altogether, these results demonstrate that the interaction-

dependent localization of STA1 and DOT2 is functionally critical for proper mRNA splicing

and that the *S307* suppressor phenotypes are due to the rescued interaction between sta1-1

and dot2^S.

Heat-induced weakening of the STA1–DOT2 interaction reduces nuclear speckle
269
270   **formation**

271   To further assess the functional connection between the STA1-DOT2 interaction and

speckle formation, we employed heat as an additional perturbation to weaken protein-protein

interactions, because elevated temperatures can weaken molecular interactions. We also

hypothesized that the heat-hypersensitive phenotype of *sta1-1* might result from heat-induced

weakening of the sta1-1 interaction. Consistent with this idea, BiFC analysis revealed a

reduction in EYFP signals corresponding to the STA1-DOT2 interaction after heat treatment

(Figures 4A-4C).

278   Unexpectedly, despite the reduction in BiFC signals, heat treatment increased the

numbers of STA1-GFP condensates when STA1-GFP was expressed alone (Supplemental

Figure 6A). We hypothesized that the increase of heat-induced condensates might result from

STA1 self-interaction (i.e., STA1 aggregation). To test this possibility, we performed BiFC

analysis using STA1-nEYFP and STA1-cEYFP. Heat treatment indeed increased BiFC signals,

indicating enhanced STA1 self-interaction under heat (Supplemental Figure 6B). Most EYFP

signals derived from STA1 self-interaction did not overlap with signals from the speckle

marker mRFP-U1-70K under either normal or heat conditions (Supplemental Figures 6C and

6D), suggesting that the STA1 self-interacting condensates represent STA1 aggregates. This

interpretation is further supported by the presence of predicted intrinsically disordered

regions (IDRs) within STA1 (residues 109-216, 298-318, 910-918, and 1022-1029), as

identified by MolPhase (Liang et al., 2024) (Supplemental Figure 6E). Therefore, the increase

in STA1-GFP condensates observed under STA1-GFP single expression is likely the result of

STA1-GFP self-aggregation.

To distinguish the NS from STA1 aggregates, we used BiFC analysis to quantify NS
induced by the STA1-DOT2 interaction. The number of these speckle-representing BiFC
signals in nuclei decreased after heat treatment and speckle formation was constantly lower in
combinations of the weakly interacting STA1 and DOT2 allele products (Figures 4D-4G).
These results reaffirm that the STA1-DOT2 interaction is required for efficient NS formation.

The *dot2^S* mutation suppresses the heat sensitivity of *sta1-1*

Next, we sought to determine whether the strength of the STA1-DOT2 interaction
also plays an important role in the heat responses. The *sta1-1* mutants have previously been
shown to display heat-sensitive phenotypes (Kim *et al.*, 2017; Kim et al., 2018). At both the
seedling and reproductive stages, the heat-sensitive phenotypes of *sta1-1* were almost fully
rescued in *S307* (Figures 5A–5C). Together, these results, along with the partial or full
reversion to *sta1-1* heat sensitivity observed in *S307^{DOT2OX}*, indicate that DOT2 functions
together with STA1 under heat stress (Figures 5A and 5C). In addition, the *dot2^C* allele did
not suppress the heat sensitivity of s*ta1-1*, supporting the allele-specific suppressive effect of
dot2^S (Figures 5B and 5C). These findings also suggest that the restored heat tolerance in
S307 results from the rescued interaction between sta1-1 and dot2^S.

The *dot2^S* mutation mitigates transcriptome-wide intron retention defects in *sta1-1*
under heat stress

Under heat stress, IR profiles in *S307* were partially restored toward WT levels
(Figure 6A). Although the global splicing patterns in *S307* did not fully return to wild-type
levels, the IR defects of *sta1-1* were markedly improved in *S307*. Among 67,103 transcripts
with significant IR changes across genotypes, *S307* showed a reduced median IR ratio (0.14)
compared with *sta1-1* (0.34) (Figure 6B). Across the three genotypes under heat, 1,114
introns (from 940 genes) showed increased IR in WT, whereas 58,522 introns (12,140 genes)
and 36,617 introns (10,654 genes) showed increased IR in *sta1-1* and *S307*, respectively
(Figure 6C, Supplemental Table 2). Venn diagram analysis further revealed that 27,514 of the
58,522 introns (~47%) with elevated IR in *sta1-1* did not exhibit increased IR in *S307* (Figure

6D). Together, these results indicate a partial restoration of *sta1-1* splicing defects in *S307* under heat stress.

Among the genes with restored splicing patterns, *HSFA1*s (*HEAT SHOCK TRANSCRIPTION FACTOR A1*s), key upstream regulators of heat-responsive genes (Li et al., 2023; Nishizawa-Yokoi et al., 2011; Röth et al., 2017), showed reduced intron retention in *S307* compared with *sta1-1* (Figure 6E, Supplemental Figure 7). This recovery was further supported by decreased levels of spliced transcripts and increased levels of intron-retained transcripts of *HSFA1*s in $S307^{DOT2END}$ and $S307^{DOT2OX}$ relative to *S307* (Figure 6F). The $dot2^{C}$ allele failed to rescue splicing defects of *HSFA1s* in *sta1-1*, consistent with its heat-sensitive phenotypes (Figures 5B and 5C, Supplemental Figure 8). Proper splicing of *HSFA1s* in *S307* enabled normal expression of downstream heat-responsive genes (Figure 6G). These findings indicate that $dot2^{S}$ suppresses intron retention defects in heat-responsive genes in *sta1-1*, thereby contributing to the restoration of heat tolerance. Taken together with the previous results, these findings highlight the importance of the STA1-DOT2 interaction for proper heat stress responses through NS formation and accurate mRNA splicing under heat conditions. The incomplete rescue of splicing defects and heat sensitivity in *S307* under heat stress, particularly at the pollen stage (Figure 5C), is likely attributable to the weaker sta1-1-dot2^{S} interaction compared with the wild-type STA1-DOT2 interaction.

DISCUSSION

Nuclear speckles (NS) are dynamic, membraneless nuclear compartments enriched in spliceosomal components and RNAs (Mintz et al., 1999; Smith *et al.*, 2020), and are increasingly recognized as active hubs that facilitate efficient pre-mRNA processing rather than acting merely as passive storage sites (Wu *et al.*, 2024). Previous studies have emphasized the contribution of intrinsically disordered regions, low-complexity domains, and multivalent RNA–protein interactions to NS organization (Ku et al., 2025; Paul *et al.*, 2024; Wang *et al.*, 2024; Zhang et al., 2024). To what extent spliceosomal protein-protein interactions involved in spliceosome assembly contribute to NS formation remains poorly understood. In this study, we provide evidence that NS organization in plants is tightly associated with the integrity of a specific interaction between two core U4/U6·U5 tri-snRNP components, STA1 and DOT2, thereby linking spliceosome assembly with subnuclear organelle organization.

Our genetic and cell biological analyses show that the weakening of the STA1-DOT2 interaction by the *sta1-1* mutation is accompanied by reduced NS formation, compromised splicing efficiency, and growth defects, whereas restoration of this interaction by the $dot2^S$ suppressor allele is associated with recovery of these phenotypes. Importantly, this suppression is achieved through reinforcement of the STA1-DOT2 interaction probably within the tri-snRNP. These observations suggest that NS organization is not independent of spliceosome assembly status and is closely linked to the stability of specific spliceosomal interactions.

The spliceosome is a highly dynamic macromolecular complex, and formation of the U4/U6·U5 tri-snRNP represents a critical step in each splicing cycle (Will and Lührmann, 2011). PRP6/STA1 functions as a central bridging factor between the U4/U6 di-snRNP and the U5 snRNP, with its interactions stabilized by tri-snRNP–specific proteins such as SART1/DOT2 (Bertram *et al.*, 2017; Nguyen *et al.*, 2016). Our observation of a positive correlation between STA1–DOT2 interaction strength and NS abundance supports a model in which proper tri-snRNP assembly facilitates efficient recruitment or retention of mature spliceosomal complexes within NS. Consistent with this view, quantitative subnuclear

localization analyses revealed that co-expression of STA1-GFP and mRFP-DOT2 significantly increased NS formation (types IV and V), compared with expression of each protein alone (Figures 2B and 2D). In contrast, co-expression of sta1-1-GFP and mRFP-DOT2 resulted in reduced NS-associated localization patterns (types IV and V) and increased localization to Cajal bodies (type III) compared with the wild-type protein (Figure 2D). Tri-snRNPs are assembled in Cajal bodies and subsequently redistributed to NS, which contain high levels of mature, splicing-competent complexes (Schaffert *et al.*, 2004; Yildirim *et al.*, 2021). Thus, our results suggest that the *sta1-1* mutation impairs later steps associated with NS organization due to defects in the STA1–DOT2 interaction, likely leading to impaired formation of splicing-competent spliceosome. In agreement with our results and speculation, when SART1, homologous to DOT2, was immunologically depleted from HeLa cells, pre-mRNA splicing was inhibited but the integrity of the U4/U6·U5 tri-snRNP was not affected, indicating the importance of SART1 in active spliceosome formation and recruitment of the tri-snRNP to the pre-spliceosome (Makarova et al., 2001; Utans et al., 1992). Thus, the STA1–DOT2 interaction is likely to promote DOT2-dependent localization of STA1 to NS, concomitant with the delivery of the tri-snRNP to the pre-spliceosome for active spliceosome assembly.

The functional relevance of our notion becomes particularly apparent under heat stress conditions. Elevated temperatures are known to destabilize spliceosomal assemblies and broadly impair splicing fidelity (Bond, 1988; Bond and James, 2000; Park et al., 2024). The most sensitive component of the spliceosome is the U4/U6·U5 tri-snRNP (Bond, 1988). In this context, the STA1–DOT2 interaction appears to function as a heat-sensitive interaction node, since the heat-induced weakening of this interaction correlates with reduced NS formation, increased intron retention, and weakened heat tolerance (Figure 4, Figure 5, Figure 6). Reinforcement of the interaction by the dot2[S] allele in sta1-1 (i.e., the *S307* suppressor) is associated with improved NS formation and splicing performance, as well as partial recovery of survival under heat stress (Figure 4, Figure 5, Figure 6). Although it is still debated whether NS primarily function as storage sites or as active hubs for RNA metabolism, our findings support the latter view by linking speckle organization to both defined molecular interactions and functional splicing outcomes. It should also be noted that STA1 itself

appeared to form non-functional aggregates under heat conditions due to its IDRs (Supplemental Figure 6). DOT2 seemed to inhibit the aggregation of STA1 and direct STA1 to NS, presumably facilitating formation of an active spliceosome. Thus, STA1–DOT2 interaction contributes to the robustness of pre-mRNA splicing under heat stress.

Taken together, our study expands current understanding of NS biology by demonstrating that NS organization in plants is closely associated with the assembly state of the spliceosome, mediated by specific protein–protein interactions within the U4/U6·U5 tri-snRNP. On this basis, we propose that heat-sensitive interaction nodes such as the STA1–DOT2 bridge represent "molecular weak points" or "interaction-level sensors" that integrate spliceosome architecture, subnuclear organelle formation, and environmental responsiveness (Figure 7). This interaction-based perspective provides insight into how plants maintain splicing robustness under stress conditions and suggests that targeted modulation of spliceosomal interactions may offer a potential strategy for improving stress tolerance.

METHODS

Plant growth conditions

Arabidopsis thaliana seeds were planted on Murashige and Skoog (MS) medium containing MS salts with vitamins (Caisson Laboratories Inc, USA), 2% sucrose, and 0.3% gelite (Duchefa, the Netherlands), adjusted to pH 5.7–5.8. For vertical growth, 0.6% gelrite was supplemented. Seeds were surface-sterilized with commercial bleach supplemented with 0.01% Tween 20 (USB, USA), stratified at 4°C for 2 days in the dark, and then transferred to a growth chamber under long-day conditions (16 h light/8 h dark) at 22°C. For soil-grown plants, seeds were sown on Sun Gro propagation mix (Sun Gro, Canada) and maintained under long-day conditions (16 h light/8 h dark) at 22°C.

Positional cloning and mutation identification

To map the *S307* mutation locus, *S307* (Columbia background) was crossed with Landsberg *erecta* (L*er*) to generate an F2 mapping population. WT-sized F2 plants homozygous for the *sta1-1* mutation were selected, and recovery of the size phenotype was confirmed in the F3 population pool. Genomic DNA was extracted from pooled leaves of 94 selected F2 mutant individuals using the RNeasy Plant Mini Kit (QIAGEN, Germany), and whole-genome sequencing was performed at Macrogen Inc. The genomic region between the SSLP markers Ch5_5.46 Mb and Ch5_5.71 Mb, identified by initial mapping, was compared with the WT genomic sequence, identifying a candidate point mutation at position 1,651 bp of *DOT2* (At5g16780) coding sequence. The candidate mutation was validated by complementation using genomic constructs driven by either the native *DOT2* promoter (DOT2::DOT2) or the Cauliflower mosaic virus 35S promoter (35S::DOT2). For native *DOT2* promoter complementation lines (*S307^{DOT2END}*), a 3,783 bp genomic fragment spanning 1,695 bp upstream of the *DOT2* start codon to 294 bp downstream of the stop codon was amplified and cloned into pCAMBIA1300 between the *Kpn*I and *Bam*HI sites (Takara, Japan). For 35S-driven complementation lines (*S307^{DOT2OX}*), the 2,463 bp *DOT2* coding sequence was amplified, cloned into pDONR221 entry vector, and subsequently recombined into the pMDC32 destination vector using the Gateway cloning system. All primers used to generate complementation constructs are listed in Supplemental Table 3.

Generation of different alleles of *dot2* mutants

The *dot2* single mutant carrying the *S307* allele (*dot2^S^*) was obtained from the F2

progeny of a cross between WT and *S307*. A derived cleaved amplified polymorphic

sequence (dCAPS) marker detecting the single-nucleotide substitution in the *dot2^S^* allele

using *Eco*RI (Takara, Japan), together with a marker detecting the 6-nucleotide deletion in

sta1-1 (used to exclude plants carrying the *sta1-1* mutation), was employed to isolate the

*dot2^S^* single mutant. The *dot2* T-DNA insertion mutant *SAIL_775_F10* (*dot2^T^*) was obtained

from the Arabidopsis Biological Resource Center (ABRC). Because homozygous *dot2^T^*

plants are infertile, only heterozygotes were maintained, and homozygotes were identified by

their dwarf phenotype (Petricka *et al.*, 2008).

A CRISPR/Cas9-generated *dot2* mutant allele (*dot2^C^*) was produced in the *sta1-1*

background using the CRISPR/Cas9 system as previously described (Cho et al., 2017; Liu et

al., 2015; Yun et al., 2022). A single guide RNA (sgRNA) targeting *DOT2* was designed

using the CRISPR web toolbox CHOPCHOP (https://chopchop.cbu.uib.no) (Labun et al.,

2019). The annealed oligonucleotides (DOT2sgRNA3_infu_CF and DOT2sgRNA3_infu_CR)

for sgRNA were cloned into the psgR-Cas9-At to generate *DOT2*-targeting construct. The

resultant DOT2sgRNA3-Cas9 cassette, excised with *Hin*dIII and *Eco*RI (Takara, Japan), was

subsequently subcloned into the *Hin*dIII-*Eco*RI sites of pCAMBIA1300. *Agrobacterium*

tumefaciens strain GV3101 carrying these constructs was used to transform *sta1-1* plants by

the floral dip method to obtain T1 seeds. CRISPR/Cas9-edited elite lines were identified by

Sanger sequencing of the DOT2sgRNA3-targeted region in the T1 generation, and

homozygous lines were subsequently isolated. All primers used for the isolation of different

dot2 mutant allels are listed in Supplemental Table 3.

RNA extraction and cDNA synthesis

Total RNA was extracted from seedlings at the designated developmental stage using

RNAiso Plus reagent (Takara, Japan) and treated with RNase-free DNase (New England

BioLabs, USA). cDNA was synthesized from total RNA using the TOPscript^TM^ Reverse

Transcriptase kit (Enzynomics, Korea). Semi-quantitative RT-PCR and realtime PCR

analyses were performed using 2 µL of the resulting cDNA (corresponding to 50 ng of input

RNA). All primers used for intron retention analyses and gene expression quantification are

listed in Supplemental Table 3.

GUS staining assay

For the *STA1* promoter-driven *GUS* (*STA1::GUS*) and *DOT2* promoter-driven *GUS*

(*DOT2::GUS*) constructs, 1,475 bp and 1,695 bp promoter fragments upstream of the start

codon of each gene were amplified, respectively. These fragments were cloned into

pCAMBIA1381 between the *Bam*HI and *Hind*III sites for *STA1::GUS*, or between the *Bam*HI

and *Nco*I sites for *DOT2::GUS* (Takara, Japan). *Agrobacterium tumefaciens* strain GV3101

carrying these constructs was used to transform Col-0 plants by the floral dip method.

Tissues of transgenic *Arabidopsis* seedlings carrying the *STA1::GUS* or *DOT2::GUS*

constructs were collected in 90% acetone (Duksan, Korea) in conical tubes on ice and

incubated for 10–20 min at room temperature. After removal of acetone, samples were

washed twice for 5 min with washing buffer (50 mM potassium phosphate buffer (pH = 7.0),

0.2% Triton X-100, 2 mM potassium ferricyanide, and 2 mM potassium ferrocyanide). The

washing buffer was then replaced with staining buffer (washing buffer supplemented with 2

mM X-Gluc (Sigma, Japan)), and tissues were incubated overnight at 37°C. Subsequently,

tissues were incubated in 70% ethanol to remove chlorophyll. GUS-staining was observed

using a light microscope (Leica ICC50, Germany).

Nuclear localization, co-localization, and BiFC analyses

For nuclear localization analyses, the full-length coding regions of *STA1* (*sta1-1*)

lacking a stop codon and *DOT2* ($dot2^S$ and $dot2^C$) containing a stop codon were amplified,

cloned into pDONR221 entry vector, and subsequently recombined into the pGWB551 or

pGWB552 destination vectors, respectively, using the Gateway cloning system.

Agrobacterium tumefaciens cells harboring *STA1 (sta1-1)-GFP* or *GFP-DOT2* ($dot2^S$ and

$dot2^C$) constructs were infiltrated into *Nicotiana benthamiana* leaves in expression buffer (50

mM MES (pH = 5.7), 10 mM $MgSO_4$, 5% D-Glucose, and 0.1 mM of acetosyringone)

together with P19, each at a final concentration of OD_{600} of 0.2. After 36 h of incubation of

infiltrated *N. benthamiana* at 22°C, fluorescence images were acquired using a confocal

microscope (TCS SP5 or STELLARIS; Leica, Germany). GFP signals were excited at 488

nm and detected at 500–530 nm. For heat treatment, samples were incubated in liquid MS

medium (0.5x MS mineral salts, 1% sucrose, 5 mM 2-(N-Morpholino) ethanesulfonic acid

(MES; Sigma, Japan), pH adjusted to 5.7 with KOH) for 1 h at 37°C following the initial 36 h

incubation.

For co-localization analyses, the full-length *DOT2* (*dot2^S* and *dot2^C*) coding region

containing a stop codon was cloned downstream of the CaMV 35S promoter and fused in

frame with mRFP in the p35S-FAST-mRFP vector. *Agrobacterium* cultures carrying *STA1*

(sta1-1)-GFP and *mRFP-DOT2* (*dot2^S* and *dot2^C*) constructs were co-infiltrated into *N.*

benthamiana leaves and fluorescent signals were observed as described above. RFP signals

were excited at 532 nm and detected at 580–620 nm.

For BiFC analyses, the same entry clones used for nuclear localization analyses were

used. *STA1 (sta1-1)* and *DOT2* (*dot2^S* and *dot2^C*) entry clones were recombined downstream

of the CaMV 35S promoter and fused in frame with nEYFP or cEYFP in the pSITE-nEYFP-

N1 (GU734648) or pSITE-cEYFP-C1 (GU734652) Gateway vectors, respectively.

Agrobacterium cells carrying *pSITE-STA1 (sta1-1)-nEYFP* or *pSITE-cEYFP-DOT2* (*dot2^S*

and *dot2^C*) constructs were co-infiltrated in combination into *N. benthamiana* leaves in the

indicated combinations, and fluorescent signals were observed as described above. EYFP

signals were excited at 488 nm and detected at 500–530 nm. As a nuclear speckle marker, the

full-length coding region of *U1-70K* (*U1 SMALL NUCLEAR RIBONUCLEOPROTEIN-70K*,

At3g50670) was amplified and cloned into p35S-FAST-mRFP between the *Bam*HI and *Sal*I

(Enzynomics, Korea) sites. As a nucleolus marker, *FIB2* (*FIBRILLARIN* 2, At4g25630)-

mRFP, provided by Dr. Hyun-Sook Pai, was used (Park et al., 2021). All primers used to

generate constructs are listed in Supplemental Table 3.

Yeast two-hybrid assay

The full-length coding region of *STA1* or *sta1-1* was inserted into the pGADT7

vector, and *DOT2* or *dot2^S* was inserted into the pGBKT7 vector. The resulting constructs,

pGADT7-STA1 (sta1-1) as prey and *pGBKT7-DOT2* (*dot2^S*) as bait, were co-transformed into

the *Saccharomyces cerevisiae* strain AH109 (Takara, Japan) and incubated at 30°C for 3 days.

Transformants were selected on SD/-Leu/-Trp or SD/-Leu/-Trp/-His medium supplemented

with 3-amino-1,2,4-triazole (3-AT; Sigma, Japan), a competitive inhibitor of the *HIS3* gene

product. All primers used to generate constructs are listed in Supplemental Table 3.

Arabidopsis mesophyll protoplast transient expression assay

Protoplast isolation and transient expression assays were performed as previously

described (Kim *et al.*, 2017). Effector constructs (*STA1*, *sta1-1*, *DOT2*, *dot2^S*, and *dot2^C*) and

reporter constructs (g*HSFA3* fused to NanoLuc luciferase (nLUC)) were generated by

inserting the corresponding coding sequences between the HBT promoter (a modified CaMV

35S promoter) and the NOS terminator in a plant expression vector. Firefly luciferase (fLUC),

driven by the UBQ10 promoter, was included as an internal control.

For nuclear localization analysis in protoplasts, *HBT-DOT2* (*dot2^S* and *dot2^C*)

constructs were transfected to *Salk007933$^{STA1-GFP}$* or *Salk007933$^{sta1-1-GFP}$*. Fluorescence

images were acquired using a confocal microscope (TCS SP5; Leica, Germany). GFP signals

were excited at 488 nm and detected at 500–530 nm.

For functional splicing assays, *HBT-DOT2* (*dot2^S* and *dot2^C*) constructs were

transfected to *Salk007933$^{sta1-1-GFP}$* and reporter activities were calculated as the nLUC/fLUC

ratio. Transfected protoplasts were lysed in passive lysis buffer (Promega, USA) containing 1%

Triton X-100 (USB, USA) and briefly vortexed. Lysates were incubated at room temperature

for 10 minutes and then centrifuged at 13,000 rpm for 30 s. The supernatant was transferred

for measurement of fLUC and nLUC activities using the Nano-Glo® Dual-Luciferase®

Reporter Assay System (Promega, USA) and a GloMax 20/20 Luminometer (Promega, USA).

All primers used to generate constructs are listed in Supplemental Table 3.

Pollen viability

To assess pollen viability, 2-month-old plants were subjected to heat treatment (37°C

for 1 day) or maintained under control conditions. Pollen grains were stained using the

Alexander staining method as previously described (Peterson, 2010). Flower buds prior to

anthesis were collected and fixed in Carnoy's fixative (6 ethanol : 3 chloroform : 1 acetic acid)

for at least 2 h. Fixed flower buds were then stained in the final staining solution at 50°C for

2 min. Stained pollen grains were observed using a light microscope (Leica ICC50,

Germany).

Propidium iodide staining

557 Seedlings from 3-day-old plants were collected and incubated in a 10 μg/mL
558 propidium iodide (PI) solution (Sigma, Japan) for 1 min. Seedlings were then washed with
559 distilled water and observed using a Leica TCS SP5 laser scanning confocal microscope
560 (Leica, Germany). PI fluorescence was excited at 532 nm and detected at 590–630 nm.

RNA-seq analysis

Total RNA was extracted from 7-day-old WT, *sta1-1*, and *S307* seedlings using
RNAiso Plus (Takara, Japan). Heat stress (37°C for 3 h) was applied to 7-day-old plants, and
heat-treated samples were collected. Three independent biological replicates were prepared
for each condition. Sequencing libraries were constructed using the TruSeq Stranded mRNA
Library Prep Kit with indexed adapters and RNA sequencing was performed using the
Illumina NovaSeq platform (MacroGen, Korea). Paired-end raw sequencing reads were pre-
processed to trim adaptors and low-quality reads using CLC Genomics Workbench v20.0.4
with default parameters (quality scores, 0.05; maximum ambiguous nucleotides, 2). Clean
reads were mapped to *Arabidopsis* reference genome (TAIR10). Intron retention ratios at the
whole-transcript level were estimated using IRFinder (v.1.3.0) (Middleton et al., 2017).
Intron retention events were defined as those showing a change in IR ratio (ΔIR ratio) greater
than 0.1 with an adjusted p-value (p_{adj}) < 0.05. IR ratios were transformed into z-scores and
visualized as heatmaps. Volcano plots were used to assess the overall distribution of IR
changes in each comparison. RNA sequencing data have been deposited in the Gene
Expression Omnibus (GEO) under accession number GSE316463.

FUNDING

This work was supported by the National Research Foundation of Korea grant funded by the
Korean government (MSIT) (RS-2024-00355816 to D.-H.J., RS-2024-00407469 for D.J.Y.
and B.-h.L. and RS-2025-00564152 for B.-h.L.) and G-LAMP grant (NRF) funded by the
Korean government (MOE) (RS-2024-00441954 for B.-h.L.), Korea Basic Science Institute

(National Research Facilities and Equipment Center) grant funded by the Korean government

(MOE) (RS-2020-NF000322 for B.-h.L.), and the Basic Science Research Program (NRF)

funded by the Korean government MOE (2018R1A6A1A03025607 for W.T.K.).

ACKNOWLEDGMENTS

588 We thank our lab members, as well as Dr. Inki Kim (Asan Medical Center, Korea) and Dr.

589 Hyun-Sook Pai (Yonsei University, Korea), for technical assistance and materials. The

590 authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

593 H.K., K.-j.Y., S.P., and D.S. conducted the experiments and performed bioinformatic analyses.
594 J.-H.J., D.-H.J., W.T.K., D.J.Y., and B.-h.L. advised on the research. All authors discussed the
595 results. H.K. S.P., D.-H.J., and B.-h.L. wrote the manuscript.

REFERENCES

Albakri, N., Bader, Z.E., Park, K.S., Bae, M.J., Zareen, S., Yun, H., Chin, H.G., Pradhan, S., Yun, D.J., and Park, J. (2025). HOS15 Modulates Alternative Splicing Homeostasis Under Cold Stress in Arabidopsis Thaliana. J. Plant Biol. **68**:203-212.

Ali, G.S., Prasad, K., Hanumappa, M., and Reddy, A.S.N. (2008). Analyses of In Vivo Interaction and Mobility of Two Spliceosomal Proteins Using FRAP and BiFC. PLoS ONE. **3**:e1953.

Bertram, K., Agafonov, D.E., Dybkov, O., Haselbach, D., Leelaram, M.N., Will, C.L., Urlaub, H., Kastner, B., Lührmann, R., and Stark, H. (2017). Cryo-EM Structure of a Pre-catalytic Human Spliceosome Primed for Activation. Cell. **170**:701-713.

Bhat, P., Chow, A., Emert, B., Ettlin, O., Quinodoz, S.A., Strehle, M., Takei, Y., Burr, A., Goronzy, I.N., Chen, A.W., et al. (2024). Genome organization around nuclear speckles drives mRNA splicing efficiency. Nature. **629**:1165-1173.

Bond, U. (1988). Heat shock but not other stress inducers leads to the disruption of a sub□set of snRNPs and inhibition of in vitro splicing in HeLa cells. EMBO J. **7**:3509-3518.

Bond, U., and James, T.C. (2000). Dynamic changes in small nuclear ribonucleoproteins of heat-stressed and thermotolerant HeLa cells. Int J Biochem Cell Biol. **32**:643-656.

Cho, S., Yu, S.I., Park, J., Mao, Y.F., Zhu, J.K., Yun, D.J., and Lee, B.H. (2017). Accession-Dependent CBF Gene Deletion by CRISPR/Cas System in Arabidopsis. Front. Plant Sci. **8**:1910.

Choi, S., Ha, S., Wolfgeher, D.J., Kim, J.W., Go, Y.H., Cha, H.J., Bae, G.U., Kron, S.J., and Kim, W.Y. (2025). N-Terminal deleted isoforms of E3 ligase RNF220 are ubiquitously expressed and required for mouse muscle differentiation. Mol. Cells. **48**:100250.

Daguenet, E., Baguet, A., Degot, S., Schmidt, U., Alpy, F., Wendling, C., Spiegelhalter, C., Kessler, P., Rio, M.C., Le Hir, H., et al. (2012). Perispeckles are major assembly sites for the exon junction core complex. Mol Biol Cell. **23**:1765-1782.

Day, I.S., Golovkin, M., Palusa, S.G., Link, A., Ali, G.S., Thomas, J., Richardson, D.N., and Reddy, A.S.N. (2012). Interactions of SR45, an SR-like protein, with spliceosomal proteins and an intronic sequence: insights into regulated splicing. Plant J. **71**:936-947.

de Luxan-Hernandez, C., Lohmann, J., Tranque, E., Chumova, J., Binarova, P., Salinas, J., and Weingartner, M. (2022). MDF is a conserved splicing factor and modulates cell division and stress response in Arabidopsis. Life Sci. Alliance. **6**:e202201507.

Galisson, F., and Legrain, P. (1993). THE BIOCHEMICAL DEFECTS OF PRP4-1 AND PRP6-1 YEAST SPLICING MUTANTS REVEAL THAT THE PRP6 PROTEIN IS REQUIRED FOR THE ACCUMULATION OF THE [U4/U6.U5] TRI-SNRNP. Nucl Acids Res. **21**:1555-1562.

Häcker, I., Sander, B., Golas, M.M., Wolf, E., Karagoez, E., Kastner, B., Stark, H., Fabrizio, P., and Lührmann, R. (2008). Localization of Prp8, Brr2, Snu114 and U4/U6 proteins in the yeast tri-snRNP by electron microscopy. Nat. Struct. Mol. Biol. **15**:1206-1212.

Ilik, I.A., Malszycki, M., Lübke, A.K., Schade, C., Meierhofer, D., and Aktas, T. (2020). SON and

SRRM2 are essential for nuclear speckle formation. eLife. **9**:e60579.

Kim, C.H., Kim, Y.D., Choi, E.K., Kim, H.R., Na, B.R., Im, S.H., and Jun, C.D. (2016). Nuclear
Speckle-related Protein 70 Binds to Serine/Arginine-rich Splicing Factors 1 and 2 via an
Arginine/Serine-like Region and Counteracts Their Alternative Splicing Activity. J. Biol. Chem.
291:6169-6181.

Kim, G.D., Yoo, S.D., and Cho, Y.H. (2018). STABILIZED1 as a heat stress-specific splicing factor
in Arabidopsis thaliana. Plant Signal Behav. **13**:e1432955.

Kim, G.D., Cho, Y.H., Lee, B.H., and Yoo, S.D. (2017). STABILIZED1 Modulates Pre-mRNA
Splicing for Thermotolerance. Plant Physiol. **173**:2370-2382.

Klimesová, K., Vojácková, J., Radivojevic, N., Vandermoere, F., Bertrand, E., Verheggen, C.,
and Stanek, D. (2021). TSSC4 is a component of U5 snRNP that promotes tri-snRNP formation. Nat.
Commun. **12**:3646.

Krissinel, E., and Henrick, K. (2007). Inference of macromolecular assemblies from crystalline state.
J. Mol. Biol. **372**:774-797.

Ku, D., Yang, Y., and Kim, Y. (2025). RNA-associated nuclear condensates: Where the nucleus
keeps its RNAs in check. Mol. Cells. **48**:100240.

Kurogi, Y., Matsuo, Y., Mihara, Y., Yagi, H., Shigaki-Miyamoto, K., Toyota, S., Azuma, Y.,
Igarashi, M., and Tani, T. (2014). Identification of a chemical inhibitor for nuclear speckle formation:
Implications for the function of nuclear speckles in regulation of alternative pre-mRNA splicing.
Biochem. Biophys. Res. Commun. **446**:119-124.

Labun, K., Montague, T.G., Krause, M., Cleuren, Y.N.T., Tjeldnes, H., and Valen, E. (2019).
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. Nucl Acids Res.
47:W171-W174.

Lee, B.H., Kapoor, A., Zhu, J.H., and Zhu, J.K. (2006). STABILIZED1, a stress-upregulated
nuclear protein, is required for pre-mRNA splicing, mRNA turnover, and stress tolerance in
Arabidopsis. Plant Cell. **18**:1736-1749.

Li, J., Cao, Y.M., Zhang, J.X., Zhu, C.J., Tang, G.L., and Yan, J. (2023). The miR165/166-
PHABULOSA module promotes thermotolerance by transcriptionally and posttranslationally
regulating HSFA1. Plant Cell. **35**:2952-2971.

Liang, Q.Y., Peng, N.N., Xie, Y., Kumar, N., Gao, W.B., and Miao, Y.S. (2024). MolPhase, an
advanced prediction algorithm for protein phase separation. EMBO J. **43**:1898-1918.

Liu, S.B., Rauhut, R., Vornlocher, H.P., and Lührmann, R. (2006). The network of protein-protein
interactions within the human U4/U6.U5 tri-snRNP. RNA. **12**:1418-1430.

Liu, S.Y., Wang, T.S., Shi, Y.L., Bai, L., Wang, S.S., Guo, D., Zhang, Y., Qi, Y.F., Chen, C.Q.,
Zhang, J.R., et al. (2021). USP42 drives nuclear speckle mRNA splicing via directing dynamic phase
separation to promote tumorigenesis. Cell Death Differ. **28**:2482-2498.

Liu, W.S., Zhu, X.H., Lei, M.G., Xia, Q.Y., Botella, J.R., Zhu, J.K., and Mao, Y.F. (2015). A

detailed procedure for CRISPR/Cas9-mediated gene editing in Arabidopsis thaliana. Sci. Bull.
60:1332-1347.

Ma, J., Li, S., Wang, T.Y., Tao, Z., Huang, S.J., Lin, N., Zhao, Y.B., Wang, C.H., and Li, P.J.
(2025). Cooperative condensation of RNA-DIRECTED DNA METHYLATION 16 splicing isoforms
enhances heat tolerance in Arabidopsis. Nat Commun. **16**:433.

Makarov, E.M., Makarova, O.V., Achsel, T., and Lührmann, R. (2000). The human homologue of
the yeast splicing factor Prp6p contains multiple TPR elements and is stably associated with the U5
snRNP via protein-protein interactions. J. Mol. Biol. **298**:567-575.

Makarova, O.V., Makarov, E.M., and Lührmann, R. (2001). The 65 and 110 kDa SR-related
proteins of the U4/U6•U5 tri-snRNP are essential for the assembly of mature spliceosomes. EMBO J.
20:2553-2563.

Middleton, R., Gao, D.D., Thomas, A., Singh, B., Au, A., Wong, J.J.L., Bomane, A., Cosson, B.,
Eyras, E., Rasko, J.E.J., et al. (2017). IRFinder: assessing the impact of intron retention on
mammalian gene expression. Genome Biol. **18**:51.

Mintz, P.J., Patterson, S.D., Neuwald, A.F., Spahr, C.S., and Spector, D.L. (1999). Purification and
biochemical characterization of interchromatin granule clusters. EMBO J. **18**:4308-4320.

Negi, S.S., Schein, C.H., Oezguen, N., Power, T.D., and Braun, W. (2007). InterProSurf: a web
server for predicting interacting sites on protein surfaces. Bioinform. **23**:3397-3399.

Nguyen, T.H.D., Galej, W.P., Bai, X.C., Oubridge, C., Newman, A.J., Scheres, S.H.W., and Nagai,
K. (2016). Cryo-EM structure of the yeast U4/U6.U5 tri-snRNP at 3.7 Å resolution. Nature. **530**:298-
302.

Nishizawa-Yokoi, A., Nosaka, R., Hayashi, H., Tainaka, H., Maruta, T., Tamoi, M., Ikeda, M.,
Ohme-Takagi, M., Yoshimura, K., Yabuta, Y., et al. (2011). HsfA1d and HsfA1e Involved in the
Transcriptional Regulation of HsfA2 Function as Key Regulators for the Hsf Signaling Network in
Response to Environmental Stress. Plant Cell Physiol. **52**:933-945.

Park, S.J., Choi, S.W., Kim, G.M., Moller, C., Pai, H.S., and Yang, S.W. (2021). Light-stabilized
FHA2 suppresses miRNA biogenesis through interactions with DCL1 and HYL1. Mol. Plant. **14**:647-
663.

Park, S.Y., Chang, H.S., Lim, S.H., Jeong, D.H., and Kim, Y.J. (2024). Functional Characterization
of PRP43a, a Component for Splicing Disassembly in Arabidopsis. J. Plant Biol. **67**:427-435.

Paul, S., Arias, M.A., Wen, L., Liao, S.E., Zhang, J.C., Wang, X.S., Regev, O., and Fei, J.Y.
(2024). RNA molecules display distinctive organization at nuclear speckles. iScience. **27**:109603.

Peterson, R.S., J. P. Chen, C. (2010). A simplified method for differential staining of aborted and
non-aborted pollen grains. Int. J. Plant Biol. **1**:e13.

Petricka, J.J., Clay, N.K., and Nelson, T.M. (2008). Vein patterning screens and the defectively
organized tributaries mutants in Arabidopsis thaliana. Plant J. **56**:251-263.

Röth, S., Mirus, O., Bublak, D., Scharf, K.D., and Schleiff, E. (2017). DNA-binding and repressor

function are prerequisites for the turnover of the tomato heat stress transcription factor HsfB1. Plant J.
89:31-44.

Sander, B., Golas, M.M., Makarov, E.M., Brahms, H., Kastner, B., Luehrmann, R., and Stark, H.
(2006). Organization of core spliceosomal components U5 snRNA loop I and U4/U6 Di-snRNP
within U4/U6.U5 tri-snRNP as revealed by electron cryomicroscopy. Mol Cell. **24**:267-278.

Schaffert, N., Hossbach, M., Heintzmann, R., Achsel, T., and Lührmann, R. (2004). RNAi
knockdown of hPrp31 leads to an accumulation of U4/U6 di-snRNPs in Cajal bodies. EMBO J.
23:3000-3009.

Schneider, M., Hsiao, H.H., Will, C.L., Giet, R., Urlaub, H., and Lührmann, R. (2010). Human
PRP4 kinase is required for stable tri-snRNP association during spliceosomal B complex formation.
Nat. Struct. Mol. Biol. **17**:216-221.

Smith, K.P., Hall, L.L., and Lawrence, J.B. (2020). Nuclear hubs built on RNAs and clustered
organization of the genome. Curr. Opin. Cell Biol. **64**:67-76.

Sun, M.L., Zhang, Y.M., Wang, Q.H., Wu, C.L., Jiang, C., and Xu, J.R. (2018). The tri-snRNP
specific protein FgSnu66 is functionally related to FgPrp4 kinase in Fusarium graminearum. Mol.
Microbiol. **109**:494-508.

Utans, U., Behrens, S.E., Luhrmann, R., Kole, R., and Kramer, A. (1992). A SPLICING FACTOR
THAT IS INACTIVATED DURING INVIVO HEAT-SHOCK IS FUNCTIONALLY EQUIVALENT
TO THE [U4/U6.U5] TRIPLE SNRNP-SPECIFIC PROTEINS. Genes Dev. **6**:631-641.

Wang, B., Li, J., Song, Y.Y., Qin, X.H., Lu, X., Huang, W., Peng, C.T., Wei, J.X., Huang, D.H.,
and Wang, W. (2024). CLK2 Condensates Reorganize Nuclear Speckles and Induce Intron Retention.
Adv. Sci. **11**:2309588.

Will, C.L., and Lührmann, R. (2011). Spliceosome Structure and Function. CSH Perspect. Biol.
3:a003707.

Wu, J.J., Xiao, Y., Liu, Y.Z., Wen, L., Jin, C.Y., Liu, S., Paul, S., He, C., Regev, O., and Fei, J.Y.
(2024). Dynamics of RNA localization to nuclear speckles are connected to splicing efficiency. Sci.
Adv. **10**:eadp7727.

Yildirim, A., Mozaffari-Jovin, S., Wallisch, A.K., Schaefer, J., Ludwig, S.E.J., Urlaub, H.,
Luehrmann, R., and Wolfrum, U. (2021). SANS (USH1G) regulates pre-mRNA splicing by
mediating the intra-nuclear transfer of tri-snRNP complexes. Nucl Acids Res. **49**:5845-5866.

Yun, J.Y., Yu, S.I., Bang, S.E., Kim, J.Y., Lee, S.H., and Lee, B.H. (2022). Identification of
CRISPR-Induced Mutations in Plants: with a Focus on the Next-Generation Sequencing Assay. J.
Plant Biol. **65**:435-443.

Zeytuni, N., and Zarivach, R. (2012). Structural and Functional Discussion of the Tetra-Trico-
Peptide Repeat, a Protein Interaction Module. Structure. **20**:397-405.

Zhan, X.C., Yan, C.Y., Zhang, X.F., Lei, J.L., and Shi, Y.G. (2018). Structures of the human pre-
catalytic spliceosome and its precursor spliceosome. Cell Res. **28**:1129-1140.

Zhang, M.J., Gu, Z., Guo, S.H., Sun, Y.T., Ma, S.B., Yang, S., Guo, J.R., Fang, C.X., Shu, L., Ge,
Y.F., et al. (2024). SRRM2 phase separation drives assembly of nuclear speckle subcompartments.
Cell Rep. **43**:113827.

 Figure Legends

Figure 1. The *dot2^S* mutation in *S307* rescues *sta1-1* defects and restores interaction with

the sta1-1 protein

(A) Growth phenotypes and fresh weights of 3-week-old WT, *sta1-1*, and *S307* plants (scale

bar = 5 cm; n = 22).

(B) Heatmap showing IR ratios in WT, *sta1-1*, and *S307*. Three independent biological

replicates were analyzed for each genotype.

(C) Volcano plots showing significantly altered IR events. Blue and red dots represent

significantly decreased and increased IR events, respectively.

(D) Growth phenotypes of 3-week-old WT, *sta1-1*, *S307*, *S307^{DOT2END},* and *S307^{DOT2OX}* plants

(scale bar = 5 cm).

(E) Experimental validation of IR levels for representative genes *FCS1* (*LEAFY CURLY AND*

SMALL1, At1g78910), *BRG1* (*BOI-RELATED GENE1*, At5g45100), and *BGLU6* (*BETA*

GLUCOSIDASE6, At1g60270). Asterisks indicate unspliced transcripts. Box diagrams show

spliced and unspliced isoforms. *Clathrin* (At4g24550) is used as a loading control.

(F and G) Semi-dominant effects of the *dot2^S* mutation on plant growth and splicing in the

sta1-1 mutant background (scale bar = 1 cm). Three-week-old plants derived from an F2

population segregating for both the *sta1-1* and *dot2^S* alleles (WT x *S307* cross). Genotypes

are indicated as follows; S, *STA1*; s, *sta1-1*; D, *DOT2*; *d^S*, *dot2^S*. PP2A (*Protein phosphatase*

2A, At1g13320) is used as a loading control. Box diagrams showed spliced and unspliced

isoforms.

(H) Y2H assays demonstrating the interaction between STA1/sta1-1 and DOT2/dot2^S. Yeast

cells co-transformed with the fusion constructs of the GAL4 DNA-binding domain and

activation domain were grown overnight and spotted onto selection medium (-L/-W), and

interaction medium (-L/-W/-H) supplemented with 0.5 mM 3-AT (3-amino-1,2,4-triazole).

(I and J) BiFC analysis for detecting the interaction between STA1/sta1-1 and

DOT2/dot2^S/dot2^C in leaves of *N. benthamiana* (scale bar = 75 μm). Red arrows indicate

EYFP signals representing protein-protein interactions in the nucleus. EYFP signals were

quantified from identical areas of leaf disks (2 x 2 cm). Data are presented as mean ± SE (n =

5). Statistical analysis is performed using one-way ANOVA with Tukey's HSD test (**P <

0.01; n.s., not significant).

Figure 2. **The disrupted nuclear localization of sta1-1 protein is re-localized to NS by the**

782   **dot2^S protein**

(A) Classification of nuclear compartments showing the localization of GOI (gene of interest)-GFP driven by the CaMV 35S promoter into five subclasses. The nucleolus marker FIB2 (FIBRILLARIN 2, At4g25630)-mRFP was co-infiltrated (scale bar = 10 μm).

(B) Subcellular localization of each protein tagged with GFP in leaves of *N. benthamiana*. Nuclear localization patterns of each GFP-tagged protein were classified into the five distinct categories defined in **(A)**. For each set, 100 GFP-expressing cells were analyzed per replicate, and data are presented as mean ± SE (n = 4). Statistical analysis is performed using one-way ANOVA with Tukey's HSD test (**$P < 0.01$; n.s., not significant).

(C-E) Co-localization of GFP-tagged STA1/sta1-1 with mRFP-tagged DOT2/dot2^S/dot2^C in leaves of *N. benthamiana*. Nuclear localization categories are shown in **(D)**, classified according to the groups defined in **(A)**. The number of NS per nucleus is shown in **(E)**, based on the STA1/sta1-1-GFP signals upon co-expression with mRFP-DOT2/dot2^S/dot2^C. For each analysis, 100 cells **(D)** or 50 nuclei **(E)** were randomly selected and analyzed (scale bar = 10 μm).

(F-H) Localization of GFP-tagged STA1 or BiFC signals of STA1 and DOT2 in leaves of *N. benthamiana*. Co-localization of GFP-tagged STA1 **(F and G),** upon single or co-expression with DOT2, and EYFP signals from BiFC analysis of the STA1 and DOT2 interaction **(H)** with the NS marker U1-70K (U1 SMALL NUCLEAR RIBONUCLEOPROTEIN-70K, At3g50670) tagged with mRFP (mRFP-U1-70K). Red boxes indicate regions of overlap between STA1-GFP/BiFC signals and mRFP-U1-70K (scale bar = 10 μm). No red boxes are indicated in **(H)** as the signals overlap completely.

Figure 3. **The localization of STA1 to NS is closely associated with proper pre-mRNA**

806   **splicing**

(A) Nuclear localization patterns of GFP-tagged *STA1/sta1-1* upon transfection with

$DOT2/dot2^S/dot2^C$ in $Salk007933^{STA1\text{-}GFP}$ or $Salk007933^{sta1\text{-}1\text{-}GFP}$ protoplasts. Cells with

nuclear-localized GFP signals of two distinct patterns (nucleoplasm and NS) were quantified

as a ratio. A total of 15 cells were analyzed for each combination (scale bar = 5 μm).

(B) Splicing activity of $DOT2/dot2^S/dot2^C$ on the *gHSFA3* DNA in $Salk007933^{sta1\text{-}1\text{-}GFP}$

protoplasts. Splicing activity on the *gHSFA3* DNA is represented as the relative luminescence

intensity by NanoLuc luciferase (nLUC) normalized to firefly luciferase (fLUC). *Clathrin* is

used as a loading control. Data are presented as mean ± SE (n = 2). Statistical analysis is

performed using one-way ANOVA with Tukey's HSD test (**$P < 0.01$; n.s., not significant).

(C) Tubercidin-induced delocalization of STA1 from NS to nucleoplasm. Ten-day-old

$Salk007933^{STA1\text{-}GFP}$ plants were treated with DMSO or 100 μM tubercidin for 1 h (scale bar =

25 μm).

(D) Tubercidin-induced disruption of pre-mRNA splicing. Ten-day-old WT, *sta1-1*, and *S307*

plants were treated with DMSO or 100 μM tubercidin for 12 h. The box diagrams show

spliced and unspliced isoforms. *PP2A* is used as a loading control.

(E) Abolished size differences between WT, *sta1-1*, and *S307* by tubercidin treatment (scale

bar = 1 cm). Five-day-old seedlings were transferred to medium containing either DMSO or

10 μM tubercidin for 6 d. Data are presented as mean ± SE (n = 4).

Figure 4. The dot2^S protein facilitates NS localization of the sta1-1 protein under heat

conditions

(A-C) BiFC analysis to detect interaction between STA1 and DOT2 under heat conditions in

leaves of *N. benthamiana* (scale bar = 75 μm). Red arrows indicate EYFP signals

representing STA1-DOT2 interactions in the nucleus. The number of EYFP signals per unit

area (2 x 2 cm) and per nucleus (n = 43) was quantified in **(B)** and **(C)**, respectively.

(D and E) BiFC analysis to detect interaction between STA1/sta1-1 and $DOT2/dot2^S/dot2^C$

under heat stress conditions in leaves of *N. benthamiana* (scale bar = 75 μm). Red arrows

indicate EYFP signals. EYFP signals were quantified from the same area of leaf disks (1 x 1

cm). Data are presented as means from three independent biological replicates. Statistical

analysis is performed using one-way ANOVA with Tukey's HSD test. Significant differences

are denoted by letters above the plots.

(F and G) Co-localization of EYFP signals from BiFC analysis of STA1/sta1-1 and

DOT2/dot2^S/dot2^C with the NS marker mRFP-U1-70K under heat stress conditions in leaves

of *N. benthamiana* (scale bar = 10 μm). Box plots show the number of NS exhibiting EYFP

signals. Center lines indicate medians; boxes represent interquartile range, and whiskers

extend to minimum and maximum values. For heat treatment, samples were incubated at

37°C for 1 h in liquid MS medium following a 36 h post-infiltration incubation.

Figure 5. **Heat-sensitivity of *sta1-1* is restored to wild type levels in the *S307* suppressor**

(A and B) Heat survival test and quantification of survival rate of each plant genotype. Ten-

day-old plants of each genotype were exposed to heat stress (37°C for 48 h), and survived

plants were scored. Statistical analysis is performed using one-way ANOVA with Tukey's

HSD test (**P < 0.01; *P < 0.05; n.s., not significant).

(C) Assessment of pollen viability using Alexander staining (scale bar = 50 μm). Viable

pollen grains appear purple, while non-viable pollen grains are transparent. Red arrows

indicate non-viable pollen grains. Data are presented as means ± SE from three independent

biological replicates. Statistical analysis is performed using one-way ANOVA with Tukey's

HSD test. Significant differences are denoted by letters above the plots.

Figure 6. **Defective splicing of *HSFA1s* in *sta1-1* is rescued in *S307*, leading to proper**

expression of downstream target genes

(A) Heatmap showing the IR ratios in WT under heat stress (WT(H)), *sta1-1* under heat stress

(*sta1-1*(H)), and *S307* under heat stress (*S307*(H)). Each genotype is represented by three

independent biological replicates.

(B) Boxplot analysis showing IR ratio trends across genotypes for 67,103 transcripts

exhibiting significant changes in intron retention. Each box represents the IR ratio

distribution for each genotype, with lines connecting genotypes to indicate intron-specific

changes. The ribbon highlights the median and variability for visual summary, and the bold

red line indicates the overall trend of median IR ratios across genotypes. Median IR ratios for

each genotype are shown in parentheses. Statistical significance among genotypes was

assessed using the Kruskal–Wallis test followed by a post hoc Dunn's test with Benjamini–

Hochberg correction. Significant differences are denoted by letters above the plots.

(C) Changes in intron retention levels of transcripts in response to heat stress across different genotypes. Paired notation denotes (number of introns/number of genes).

(D) Venn diagram showing transcripts with increased intron retention under heat stress in each genotype.

(E and F) Quantification and validation of splicing defects in the master heat stress regulators *HSFA1*s. Box diagrams show spliced and unspliced isoforms. *PP2A* is used as a loading control.

(G) qRT-PCR analysis of heat stress-related genes regulated by *HSFA1*s. The relative expression levels of each gene are shown in the graphs. Data are presented as mean ± SE from three independent biological replicates. Statistical analysis is performed using one-way ANOVA with Tukey's HSD test (**$P < 0.01$; *$P < 0.05$; n.s., not significant).

Figure 7. A proposed model for STA1-DOT2 interaction-mediated NS formation and pre-mRNA splicing in growth and heat stress responses

The STA1-DOT2 interaction, presumably reflecting the U4/U6·U5 tri-snRNP assembly, is important for NS formation. Proper NS formation ensures the production of correctly spliced transcripts required for normal growth and heat stress responses. The *sta1-1* mutation or heat stress weakens the STA1-DOT2 interaction, thereby reducing NS formation. This disrupted subnuclear localization of splicing components increases intron-retained transcripts, resulting in defective growth and heat stress responses.

Supplemental Figure 1. **Positional cloning and the identification of the $dot2^S$ mutation in**

S307

(A) Map of the *DOT2* locus on chromosome 5. Ninety-four F2 seedlings derived from an

S307 x L*er* cross were used for mapping. The Simple sequence length polymorphism (SSLP)

markers used for mapping, Ch5_5.46Mb and Ch5_5.71Mb, are shown on the 184A1 and

35B3 BAC clones, respectively. Recombination numbers are shown under each marker locus.

(B) The mutation information of $dot2^S$ mutation, including nucleotide and corresponding

amino acid changes.

Supplemental Figure 2. **Homologs and expression patterns of *DOT2*, a U4/U6·U5 tri-**

snRNP associated protein

(A) Multiple sequence alignments of DOT2 homologs.

(B) Expression patterns of *DOT2* and *STA1::GUS* in *Arabidopsis*. GUS signals are observed

in whole seedling, flower, silique, and trichome from left to right.

Supplemental Figure 3. Phenotypes of the *dot2* single mutant alleles

(A) Growth phenotypes of 3-week-old WT, *sta1-1*, *S307*, $dot2^S$, and $dot2^T$ (SAIL_775_F10)

plants (scale bar = 5 cm).

(B) Splicing defects in marker genes across genotypes. Box diagrams show spliced and

unspliced isoforms. *PP2A* is used as a loading control.

(C) Representative confocal images of PI-stained root tips of 3-day-old WT, *sta1-1*, *S307*,

$dot2^S$, and $dot2^T$ seedlings (scale bar = 25 μm). Arrowheads indicate dead cells, and white bars

mark the division zone. Division zone length in 3-day-old seedlings is quantified (n = 10).

Statistical analysis is performed using one-way ANOVA with Tukey's HSD test (**$P < 0.01$;

n.s., not significant).

Supplemental Figure 4. Residue positions of mutations in sta1-1 and dot2 mutant

proteins

(A) Predicted protein structures of *Arabidopsis* STA1 and DOT2. In the model, STA1 protein

is shown in green and DOT2 protein in blue, with their respective mutations highlighted as

follows: the *sta1-1* mutation in magenta, the $dot2^S$ mutation in red, and the $dot2^C$ mutation in

orange.

(**B**) Domain structure of STA1 and DOT2. Predicted STA1-DOT2 interaction regions

identified using InterProSurf and PDBePISA are indicated by red lines (STA1: 113–123 aa,

185–202 aa; DOT2: 548–569 aa). Domains are labeled as follows: UBQ, ubiquitin-like;

PRP1P, pre–mRNA processing protein 1; HAT, half-A-tetratricopeptide repeat; NLS, nuclear

localization signal; TPR, tetratricopeptide repeat; HIND, hub1–interacting domain.

(**C**) The $dot2^S$ and $dot2^C$ mutations, showing nucleotides and the corresponding amino acid

changes.

(**D**) Growth phenotypes of 3-week-old WT, *sta1-1*, *S307*, and the *sta1-1 $dot2^C$* double mutant

($ssd^C d^C$) plants (scale bar = 5 cm).

(**E**) Splicing defects in the marker genes across genotypes. The box diagrams show spliced

and unspliced isoforms. *PP2A* is used as a loading control.

Supplemental Figure 5. Rescue of *Salk007933* lethality through introduction of STA1-

GFP or sta1-1-GFP, resulting wild-type- and *sta1-1*-like phenotypes, respectively

Growth phenotypes of 3-week-old $Salk007933^{STA1\text{-}GFP}$ and $Salk007933^{sta1\text{-}1\text{-}GFP}$ plants (scale

bar = 5 cm).

Supplemental Figure 6. Increased STA1-aggregates under heat conditions

(**A and B**) Subcellular localization of GFP-tagged STA1 and EYFP signals from BiFC

analysis of STA1-STA1 interaction under heat conditions (scale bar = 75 μm).

(**C and D**) Co-localization of EYFP signals from BiFC analysis of STA1-STA1 interaction

with the mRFP-U1-70K in leaves of *N. benthamiana* (scale bar = 10 μm).

(**E**) MolPhase prediction analysis of IDR region within STA1 (Liang *et al.*, 2024).

For heat treatment, samples were incubated at 37°C for 1 h in liquid MS medium following a

36 h post-infiltration incubation.

Supplemental Figure 7. Intron-retained *HSFA1* transcripts in *sta1-1*

(A) Increased IR events of *HSFA1s* in *sta1-1* (H). IR events in *WT*(H), *sta1-1*(H), and *S307*(H)

are visualized using the Integrative Genomics Viewer (IGV).

(B) Protein domain structures of premature termination codon (PTC)-containing intron-

retained pre-mRNA isoforms in *sta1-1* (H). Domains are labeled as follows: DB, DNA-

binding domain; OD, oligomerization domain; NLS, nuclear localization signal; AHA1/2,

aromatic and large hydrophobic amino acid residues embedded in acidic surroundings. PTCs

are labeled with a red symbol.

Supplemental Figure 8. Splicing defects of *HSFA1*s in *sta1-1 dot2^C* double mutants

Splicing defects of the master heat stress regulators *HSFA1*s in the *sta1-1 dot2^C* double

mutant (*ssd^{C}d^C*) plants. Box diagrams show spliced and unspliced isoforms. *PP2A* is used as

a loading control.

Supplemental Table 1. Intron retained transcripts under normal conditions

Supplemental Table 2. Intron retained transcripts under heat conditions

Supplemental Table 3. Primer sets used in this study

Fig2

Fig3

Fig4

Fig6