Functional Annotation Workflow for Fungal Transcriptomes

Nagisa Morihara [1] and **Hidemasa Bono** [1,2,*]

[1] Graduate School of Integrated Sciences for Life, Hiroshima University, 1-4-4 Kagamiyama, Higashi-Hiro-shima City, Hiroshima 739-8528, Japan; d235537@hiroshima-u.ac.jp

[2] Genome Editing Innovation Center, Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima City 739-0046, Japan; bonohu@hiroshima-u.ac.jp

* Correspondence: bonohu@hiroshima-u.ac.jp; Tel.: +81-82-424-4013

Abstract

Although RNA sequencing (RNA-seq) enables rapid transcriptome profiling, functional annotation of fungal transcriptomes remains challenging. Existing tools prioritize broad taxonomic coverage, and reference genomes are scarce for non-model species. This study aimed to develop a fungal-specific functional annotation workflow to support rapid and accurate functional analyses downstream of RNA-seq, independent of reference genome availability. To evaluate the workflow, RNA-seq data from 57 samples of *Lentinula edodes* strain H600 (shiitake mushroom) were retrieved, along with full-length transcript sequencing (Iso-Seq) data and corresponding RNA-seq data from 20 samples of *Phakopsora pachyrhizi* (Asian soybean rust) from public databases. The workflow successfully annotated over 96% of protein-coding transcripts and demonstrated applicability to Iso-Seq data. Functional enrichment analyses revealed higher-resolution functional detection than existing annotation tools. Furthermore, integrating homology searches against fungal-specific databases with expression pattern-based annotations highlighted the workflow's utility for target identification in genome editing and other applications. Overall, the results of this study highlight the potential of the developed workflow in facilitating the discovery of functionally important transcripts and their translation into biotechnological applications.

Keywords: RNA sequencing; functional annotation; shiitake mushroom; soybean rust; full-length transcript sequencing; transcriptome profiling

1. Introduction

Fungi represent one of nature's most diverse organismal groups, with their exceptional functional diversity and ecological importance increasingly recognized in biotechnology, agriculture, environmental conservation, and human health [1]. Next-generation sequencing (NGS) technologies have enabled rapid, cost-effective genome sequencing, driving an exponential increase in sequenced fungal genomes. These resources offer opportunities to identify novel genes as targets for genome editing and functional characterization. RNA sequencing (RNA-seq), a key NGS application, is widely used to obtain transcript sequences and perform comparative transcriptome analyses across various conditions, time points, and treatments. This approach reveals actively expressed genes and their expression dynamics, providing direct insights into regulatory mechanisms and aiding research target prioritization. However, genome or transcriptome sequencing alone cannot elucidate gene function, as accurate functional annotation is essential for biological interpretation. Inaccurate or incomplete annotations hinder downstream analyses, such

as functional enrichment studies and genome editing target identification [2]. Fungal functional annotation presents multiple inherent challenges compared with that in well-studied model organisms. First, many fungal species lack reference genomes, requiring reliance on phylogenetically distant or poorly annotated species; intraspecific diversity further necessitates comparative analyses across multiple strains. Second, fungi exhibit distinctive features, including biosynthetic gene clusters and species-specific adaptations, that are often absent in species-agnostic protein or domain databases, resulting in failure to capture. Finally, many predicted fungal genes lack experimental validation and uncharacterized with uncertain functional predictions. Consequently, annotation based on these resources often results in failure when analyzing closely related homologs lacking functional annotation [3, 4]. For these reasons, large-scale fungal sequence data have not yet been fully utilized.

Gene annotation involves identifying loci in genome sequences and assigning structural and functional information, including gene structure prediction, coding/non-coding region delineation, and functional inference based on sequence homology and existing functional data. Existing annotation workflows, such as BRAKER3 [5] and MAKER2 [6] for eukaryotic genomes, the Prokaryotic Genome Annotation Pipeline [7] for bacterial genomes, and funannotate [8] and FunGAP [9] for fungi, are often designed for submission to public repositories, including the National Center for Biotechnology Information (NCBI) [10], and therefore emphasize genome DNA-based structural prediction and gene determination. Meanwhile, several analytical tools are available for functional annotation of RNA-seq-assembled transcripts, including Trinotate [11], which provides multi-database annotation for model/non-model organisms; Blast2GO [12], which enables Gene Ontology [GO] enrichment *via* graphical user interface; and Fanflow4Insects [13], an insect-specific annotation tool. However, no dedicated tools exist for fungal transcriptomes.

In this study, a fungal functional annotation workflow applicable to transcriptomes with or without reference genomes was developed. By integrating fungus-optimized homology search databases and condition-specific annotations (e.g., tissue type and developmental stage), it aimed to address fungal-specific challenges, improving the quality and efficiency of downstream analyses, such as functional enrichment and genome editing target identification. Additionally, the applicability of this workflow was evaluated for full-length transcript sequencing (Iso-Seq) [14]. Iso-Seq enables high-resolution identification of splicing isoform structures, and data submissions to public repositories are expected to increase. The findings of this study demonstrate the workflow compatibility with both short-read RNA-seq and long-read transcripts, which highlight its versatility.

2. Materials and Methods

2.1. Acquisition of Expression Data

RNA-seq data for 57 samples of *Lentinula edodes* strain H600 were obtained from the Sequence Read Archive (SRA) (accession numbers SRR21185407–SRR21185463). Filtered, error-corrected Iso-Seq transcripts for *Phakopsora pachyrhizi* were retrieved from GenBank (accession numbers GHWK00000000 and GHWL00000000) [15]. Corresponding RNA-seq data, collected at days 3, 7, 10, and 14 post-infection, were obtained from the SRA (accession numbers SRR10130097–SRR10130116) [15].

2.2. Assembly and Coding Region Prediction

For the *L. edodes* dataset, raw reads from all 57 samples were merged and trimmed using TrimGalore (version 0.6.10) [16], followed by *de novo* transcriptome assembly with the rnaSPAdes module of SPAdes (version 3.15.5) [17]. Assembly quality was assessed using Benchmarking Universal Single-Copy Orthologs (version 5.8.0) [18], revealing

97.3% completeness. Coding regions were predicted using TransDecoder (version 5.7.1) [19]. For the *P. pachyrhizi* dataset, pre-assembled Iso-Seq transcripts were used directly, and short-read RNA-seq data were processed with TrimGalore.

2.3. Expression Quantification

Expression quantification was performed using Salmon (version 1.10.3) [20]. Treating each transcript as an independent unit, we used the '—keepDuplicates' option during index creation to retain all variants. Transcripts Per Million (TPM) values were used for principal component analysis (PCA) in R (version 4.3.2) [21] with the stats and ggplot2 (version 3.5.2) [22] packages.

2.4. Functional Annotation

Workflow scripts are available in the fungifunate GitHub repository [23], built on the Systematic Analysis for Quantification of Everything framework [24]. Homology searches of retrieved coding region protein sequences used ggsearch36 from the FASTA package (version 36.3.8g) [25] with parameters '-d 1 -m10 -E 0.1'. Reference protein sequences for comparison were from human (*Homo sapiens*) [26], mouse (*Mus musculus*) [27], and budding yeast (*Saccharomyces cerevisiae*) [28], obtained from Ensembl [29], UniProtKB/SwissProt [30], and FungiDB [31, 32]. Release68 protein files were concatenated into a single FASTA file, and protein domain searches followed using InterProScan (version 5.67-99.0) [33]. GO terms [34, 35] were retrieved for human, mouse, budding yeast, and UniProtKB protein IDs *via* the biomaRt package (version 2.58.2) [36] in R, and for InterProScan results using the InterPro2GO mapping file [37]. For the *L. edodes* dataset, condition-specific annotations were generated by averaging transcript-level TPM across the following three groups: mycelia (n = 19), primordia (n = 12), and fruiting bodies (n = 26). Stage-specific transcripts were defined as those expressed (mean TPM ≥ 1) in only one developmental stage, with low variability (coefficient of variation [CV] < 1) and maximum mean TPM ≥ 2. These were consolidated into a single table using an R script.

2.5. Differential Expression Analysis

Differential gene expression analysis for *L. edodes* data used DESeq2 package (version 1.42.1) [38] in R, with pairwise comparisons across developmental stages. Wald tests identified differentially expressed transcripts, with statistical significance, using Benjamini–Hochberg adjusted *p*-values (padj) < 1×10^{-11}.

For *P. pachyrhizi*, time-course expression patterns were analyzed with maSigPro package (version 1.74.0) [39] in R using third-degree polynomial regression. Significant genes (Benjamini–Hochberg adjusted Q-value < 0.05, ≥10 observations) underwent backward stepwise regression; those with $R^2 \geq 0.6$ were clustered into four groups *via* hierarchical clustering.

2.6. Functional Analysis

Functional enrichment analysis targeted differentially expressed transcripts using web tools Metascape (version 3.5) [40] and gProfiler [41] with human and budding yeast gene identifiers. Additionally, the topGO package (version 2.54.0) [42] in R analyzed all assigned GO terms and duplicates were removed. For topGO analysis, GO biological process (BP) ontology was examined *via* elim algorithms and Fisher's exact test.

2.7. Comparative Annotation Method

Protein sequences were searched against the NCBI nr database [43] using DIAMOND (version 2.1.8.162) [44] in the blastp mode. GO terms were assigned with Blast2GO (version 6.0.3), followed by topGO functional enrichment analysis using the abovementioned parameters (section 2.6).

3. Results

3.1. Overview of the Functional Annotation Workflow

This workflow generated an annotation table from RNA-seq reads for functional enrichment analysis, integrating both functional annotations and differential expression results for transcript filtering. The functional annotation comprised four main components. First, homology-based annotation using a global alignment tool (ggsearch) against well-annotated protein databases, including human, mouse, budding yeast, UniProtKB/Swiss-Prot, and FungiDB. Second, GO term assignment for protein IDs from human, mouse, yeast, and UniProtKB. Third, protein domain annotation using InterProScan. Finally, condition-specific expression annotation.

Figure 1. Overview of the annotation table generation including functional annotation. RNA-seq, RNA sequencing; GO, Gene Ontology.

Transcript sequences can be derived from *de novo* RNA-seq assembly and coding sequence prediction or from public databases. The ggsearch tool performs global alignments, making it suitable for detecting distant homologs in genetically diverse fungi GO terms provide a standardized framework for gene function across BPs, molecular functions, and cellular components, enabling functional enrichment analysis of significantly overrepresented functions. InterProScan identifies conserved protein domains and functional motifs, annotating transcripts missed by homology searches against protein databases and improving overall coverage. For condition-specific expression annotation, we applied custom binary criteria rather than continuous metrics, such as the Tau index [45], which are widely used to score tissue-specific expression on a 0–1 scale. A transcript was classified as condition-specific if it met all three criteria: (1) expression (TPM ≥ 1) in only one

group, (2) CV < 1 across all groups, and (3) maximum mean TPM ≥ 2 in the expressing 162
group. This binary TRUE/FALSE classification facilitates evaluation of expression pat- 163
terns under specific biological conditions, such as developmental stages. 164

3.2. Application to L. edodes 165

To evaluate workflow utility, 57 RNA-seq samples from *L. edodes* strain H600 under- 166
went *de novo* assembly and coding region prediction, followed by workflow application. 167
These samples, originally classified into 20 groups by developmental stage and tissue type 168
(Table S1) were consolidated into three broad groups (namely mycelia, primordia, and 169
fruiting bodies) based on PCA (Figure S1) for condition-specific annotation and differen- 170
tial expression analysis. Annotation and expression data for each transcript were then in- 171
tegrated into a comprehensive table (Table S2). Of 227,580 analyzed transcripts, 98.2% re- 172
ceived functional annotations. Contributions by database were as follows: human (42%), 173
mouse (38%), budding yeast (30%), UniProtKB (55%), FungiDB (81%), InterPro (43%), con- 174
dition-specific (0.13%), and GO terms (74%). In contrast, the NCBI nr database method 175
annotated 66% of transcripts with only 12% assigned GO terms. Differential expression 176
analysis identified 1,926 transcripts between mycelia and primordia, 1,739 between pri- 177
mordia and fruiting bodies, and 3,801 between fruiting bodies and mycelia. A Venn dia- 178
gram (Figure 2a) was constructed, and functional enrichment analysis was performed for 179
non-overlapping differentially expressed genes, including 492 specific to the mycelium 180
versus primordium comparison and 711 specific to the primordium vs fruiting body com- 181
parison. Differential analysis between mycelia and primordia was performed through 182
Metascape (Figure 2b) and g:Profiler (Figure 2c) analyses using human gene identifiers. 183
This enabled the use of web-based tools that were previously inaccessible for non-model 184
organisms. The results revealed significantly enriched metabolic processes (carboxylic 185
acid, xenobiotic, and lipid metabolism) and functions related to oxidoreductase activity, 186
cell membrane, and cytoskeleton. These results align with metabolic demands for primor- 187
dium initiation and membrane/cell wall remodeling during mycelium-to-primordia tran- 188
sition [46-48]. Subsequently, GO enrichment analysis was performed using topGO with 189
GO terms assigned by both approaches. As a result, the comparative method detected 190
basic metabolic functions, such as carbohydrate metabolism and aromatic amino acid bi- 191
osynthesis (Figure 2e), the other hand, our workflow detected more detailed functions 192
related to cellular dynamics and specific pathways including NAD-cap decapping, fatty 193
acid α-oxidation, microtubule-based peroxisome localization, and lamellipodium assem- 194
bly regulation (Figure 2d). These differences reflect the differing GO term assignment 195
rates between the two approaches. 196

Between fruiting bodies and primordium, Metascape and g:Profiler analyses de- 197
tected common enrichments in olefinic compound metabolism, xenobiotic metabolism, 198
lysosomal lumen pH regulation, and iron ion-related functions (Figure S2a, b). Olefinic 199
acid metabolism relates to unsaturated fatty acid metabolism, consistent with previously 200
reported elevated expression during fruiting body development [49], along with poten- 201
tially novel functions. Similarly, topGO analysis using our workflow identified mitosis, 202
cell wall remodeling, and autophagy among top-ranked functions (Figure S2c), aligning 203
with prior reports [50,51]. In contrast, the comparison method prioritized phospholipid 204
biosynthesis, endoplasmic reticulum unfolded protein response, and microtubule-based 205
nuclear migration (Figure S2d). Although these appear related to cell proliferation, they 206
represented more ambiguous functional categories. 207

Figure 2. Functional enrichment analysis of *Lentinula edodes* transcriptome data. (a) Venn diagram of differentially expressed transcripts. The analyses shown in (b–e) were performed on the 492 transcripts uniquely differentially expressed between primordia and mycelia. (b) Metascape results using human gene identifiers. (c) g:Profiler results using human gene identifiers. (d) topGO dot plot based on GO terms aggregated from all annotation databases. (e) topGO dot plot based on GO terms assigned by the nr database-based method. GO, Gene Ontology; BP, biological process; CC, cellular component; MF, molecular function.

Notably, approximately 13% of transcripts were annotated exclusively *via* FungiDB, with many from uncharacterized basidiomycete genes, such as *Coprinopsis*, *Pleurotus*, and *Lentinus* (Table S2). Some of these appeared in differential expression analyses and considered priority candidates for genome-editing targets. Additionally, four transcripts received unique developmental stage-specific annotations. Among primordium-specific

transcripts, a highly expressed homolog (TPM = 44) of yeast *SWS2*, involved in sporulation and oxidative stress responses, was identified (Table S2). This highlights the workflow's ability to uncover functionally important genes beyond differential expression analysis.

3.3. Application to P. pachyrhizi

To evaluate workflow applicability to Iso-Seq data, filtered, error-corrected *P. pachyrhizi* Iso-Seq transcripts from GenBank were analyzed using the same pipeline. Annotation and expression data were consolidated into a single file (Table S3). Of 9,680 protein-coding transcripts, 96.1% received functional annotations, with contributions from each database as follows: human (56%), mouse (50%), budding yeast (44%), UniProtKB (55%), FungiDB (87%), InterPro (52%), and GO terms (79%). The comparative method annotated 80% of transcripts with only 19% GO terms assigned.

PCA confirmed separation of transcriptome data at days 3, 7, 10, and 14 post-infection (Figure S2). Time-course analysis identified 3,038 upregulated transcripts (Clusters 1–3) and 40 downregulated transcripts (Cluster 4) (Figure 3a). Functional enrichment was performed as described for *L. edodes*. For upregulated transcripts, both methods detected relatively broad functional categories (data not shown), which was attributed to their high proportion of upregulated transcripts compared with that in the total transcriptome. For downregulated transcripts, Metascape and g:Profiler identified CENP-A (histone H3 variant) and RNA-related processes. Although histone modifications regulate pathogenicity [52] and host-derived histones exhibit antimicrobial activity [53], CENP-A associations with infection remain unreported. Subsequent topGO analysis showed that the comparative method prioritized nucleosome/ribosomal assembly and host-related functions (reductive pentose-phosphate cycle and photorespiration) as top enriched GO terms. In contrast, the developed workflow detected specific, biologically relevant terms, including negative regulation of K48-linked ubiquitination, cell proliferation, and histone deacetylation. These results confirm improved annotation accuracy for Iso-Seq data and applicability of the developed workflow.

Several differentially expressed transcripts received FungiDB-exclusive annotations, including uncharacterized genes from rust fungi and other pathogens (Table S3). These results suggest novel functions in rust fungi and represent future genome editing targets. Overall, the workflow delivers detailed, fungus-specific insights from both RNA-seq and Iso-Seq datasets.

Figure 3. Functional enrichment analysis of *Phakopsora pachyrhizi* transcriptome data. (a) Cluster analysis of time-course differentially expressed transcripts. The analyses shown in (b–e) were performed on transcripts in Cluster 4. (b) Metascape results using human gene identifiers. (c) g:Profiler results using human gene identifiers. (e) topGO dot plot based on GO terms aggregated from all annotation databases. (e) topGO dot plot based on GO terms assigned by the nr database-based method. GO, Gene Ontology; BP, biological process; CC, cellular component; MF, molecular function.

4. Discussion

Publicly available RNA-seq data remain underutilized, particularly in fungi lacking comprehensive functional annotation. Reanalysis with the proposed workflow can uncover novel functional genes and response pathways. Even species with reference genomes may harbor unidentified splicing variants [54], making it valuable to utilize RNA-seq data-assembled transcript sequences. For *L. edodes*, the NCBI reference genome (GCF_021015755.1) contained 14,078 transcripts compared with our assembly result of 227,580 transcripts. Although redundant and non-translated sequences may exist, they may be filtered out during low-expression transcript removal or differential expression

analysis. Moreover, the benefit of detecting novel transcript variants outweighs these limitations. In this regard, Iso-Seq analysis represents an effective approach. However, it should be noted that read number limitations may challenge complete coverage of all transcripts.

The developed workflow achieved superior annotation coverage and biological validity, particularly in GO term assignment. This was partially attributed to the use of NCBI nr database for BLAST results input to Blast2GO for comparison, which contains uncharacterized genes or entries lacking GO term associations. These results further emphasize the importance of incorporating databases with identifiers compatible with integrative functional analyses. Functional enrichment analysis using topGO revealed notable differences in biological insights between the two approaches. Differences originating from web-based tools, such as Metascape and topGO, may be partly attributed to the algorithm choice used in topGO. Herein, the elim algorithm was employed that preferentially detects more specific GO terms located at lower hierarchical levels. In contrast, Metascape and similar tools typically produce results closer to those generated by the classic algorithm, resulting in the enrichment of broader, higher-level parent terms. Additionally, web-based tools perform enrichment using gene identifiers from a specific model organism (e.g., human), whereas topGO integrates GO terms aggregated from all databases, further generating differences in the outputs. Results obtained for *L. edodes* and *P. pachyrhizi* showed that the proposed workflow identified more specific and narrowly defined functions, which were directly linked to core cellular dynamics, including transcript remodeling, the G2/M transition of the mitotic cell cycle, and ubiquitination, thereby efficiently annotating candidate genes for downstream functional validation. The workflow also detected lower-level GO categories involving relatively small numbers of genes; for example, the NAD-cap decapping category contained 37 genes, and the negative regulation of protein K48-linked ubiquitination category contained 12 genes, representing a practically manageable number of candidates for gene-focused analyses. In the topGO results for *P. pachyrhizi*, the detected broad host-associated functions included photorespiration and photosynthesis by nr-based annotation approaches, requiring careful interpretation. Although host derived sequences were removed at the transcript level, these annotations may arise from assignments to plant-related genes in the nr database or artifacts inherent to homology-based annotation. In contrast, the developed workflow utilized databases with explicit taxonomic constraints, including fungal-specific resources, model organism datasets, and UniProt, thereby reducing such artifacts. Notably, integrating GO terms from multiple databases reduces annotation bias, increases the detection of smaller gene-set categories, and supports more accurate biological interpretation. FungiDB contributed the most to annotation efficiency owing to its continuous integration of published and unpublished fungal data, including functionally unknown genes. This further suggests that several fungal genes remain undercharacterized and unregistered in species-agnostic databases. Annotations of differentially expressed transcripts exclusively by FungiDB were largely derived from closely related species, suggesting that FungiDB complements important fungus-specific information not captured by conventional annotation approaches. Developmental stage-specific annotation in *L. edodes* exclusively annotated four transcripts through this method. A high-expression homolog of a yeast oxidative stress response gene was detected among primordium-specific transcripts; however, it was excluded from the differentially expressed transcript list because of the applied q-value cutoff. This approach allowed prioritization of potentially important genes independent of other annotation strategies and differential expression analyses. Despite the low developmental stage-specific annotation rate (0.13%), analysis based on the original 20 classification categories may reveal additional stage- or tissue-specific transcripts.

The workflow developed in this study relies on homology-based annotation, making it dependent on reference database quality and imperfect cross-species functional inference. This underscores the need for experimental validation. Despite these limitations, opportunities exist for optimizing annotation strategies, refining database selection, and expanding the analysis. In a multi-database integration strategy, annotation order across these databases is important, as prioritizing well-annotated species may yield more useful functional information. This observed higher annotation rate from human sequences compared with those from phylogenetically closer yeast species may be attributable to the more extensive characterization of human transcript variants. Although this workflow integrates GO terms for facilitating easy functional interpretation and widespread use, its flexibility allows for the incorporation of additional database information per requirements, such as enzyme commission numbers, Clusters of Orthologous Groups/Eukaryotic Clusters of Orthologous Groups classifications, or secondary metabolite biosynthetic gene cluster annotations from resources, such as antiSMASH database. Although focused on protein-coding transcripts, expanding to non-coding RNAs (ncRNAs), such as long ncRNAs and micro RNAs, as key regulators and genome editing targets [55] would benefit from rRNA depletion-based library preparation, instead of mRNA-seq. Future directions of this study include machine learning-based structural annotation for both coding and non-coding transcripts, which is particularly valuable considering fungi's high genetic diversity and low sequence homology.

In conclusion, the proposed workflow provides a practical, reference-genome-independent framework for fungal transcriptome functional annotation. It enables mechanistic investigation of underexplored, complex biological processes underlying developmental programs and environmental responses even in fungal species, capturing genes and functions missed by conventional approaches. Furthermore, prioritized candidates serve as rational CRISPR-based functional validation, facilitating the translation of annotated transcripts from basic to applied research.

Supplementary Materials: The following supporting information can be downloaded from: https://doi.org/10.6084/m9.figshare.30944468, Figure S1: PCA plot of *L. edodes* RNA-seq datasets; Figure S2: PCA plot of *P. pachyrhizi* RNA-seq datasets; Table S1: *L. edodes* dataset metadata; Table S2: *L. edodes* annotation table; Table S3: *P. pachyrhizi* annotation table.

Author Contributions: Conceptualization, N.M. and H.B.; methodology, N.M. and H.B.; software, N.M. and H.B.; validation, N.M..; formal analysis, N.M.; investigation, N.M.; resources, H.B.; data curation, N.M.; writing—original draft preparation, N.M.; writing—review and editing, N.M. and H.B.; visualization, N.M.; supervision, H.B.; project administration, H.B.; funding acquisition, H.B. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the Center of Innovation for Bio-Digital Transformation (Bio-DX), an open innovation platform for industry-academia co-creation (COI-NEXT), and the Japan Science and Technology Agency (JST), grant number JPMJPF2010.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original data presented in this study are openly available on GitHub [23].

Acknowledgments: Experiments were performed using computers at the Hiroshima University Genome Editing Innovation Center.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Abbreviations

The following abbreviations are used in this manuscript:

TPM	Transcripts Per Million
RNA-seq	RNA Sequencing
Iso-Seq	Full-Length Transcript Sequencing
NGS	Next-Generation Sequencing
NCBI	National Center for Biotechnology Information
SRA	Sequence Read Archive
PCA	Principal Component Analysis
CV	Coefficient of Variation

References

1. Corbu, V.M.; Gheorghe-Barbu, I.; Dumbravă, A. Ştefania; Vrâncianu, C.O.; Şesan, T.E. Current Insights in Fungal Importance—a Comprehensive Review. *Microorganisms* **2023**, *11*, 1384, doi:10.3390/microorganisms11061384.

2. Griesemer, M.; Kimbrel, J.A.; Zhou, C.E.; Navid, A.; D'haeseleer, P. Combining Multiple Functional Annotation Tools Increases Coverage of Metabolic Annotation. *BMC Genomics* **2018**, *19*, 948, doi:10.1186/s12864-018-5221-9.

3. Hyde, K.D.; Baldrian, P.; Chen, Y.; Thilini Chethana, K.W.; De Hoog, S.; Doilom, M.; De Farias, A.R.G.; Gonçalves, M.F.M.; Gonkhom, D.; Gui, H.; et al. Current Trends, Limitations and Future Research in the Fungi? *Fungal Diversity* **2024**, *125*, 1–71, doi:10.1007/s13225-023-00532-5.

4. Mohanta, T.K.; Al-Harrasi, A. Fungal Genomes: Suffering with Functional Annotation Errors. *IMA Fungus* **2021**, *12*, 32, doi:10.1186/s43008-021-00083-x.

5. Gabriel, L.; Brůna, T.; Hoff, K.J.; Ebel, M.; Lomsadze, A.; Borodovsky, M.; Stanke, M. BRAKER3: Fully Automated Genome Annotation Using RNA-Seq and Protein Evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res.* **2024**, *34*, 769–777, doi:10.1101/gr.278090.123.

6. Holt, C.; Yandell, M. MAKER2: An Annotation Pipeline and Genome-Database Management Tool for Second-Generation Genome Projects. *BMC Bioinformatics* **2011**, *12*, 491, doi:10.1186/1471-2105-12-491.

7. Tatusova, T.; DiCuccio, M.; Badretdin, A.; Chetvernin, V.; Nawrocki, E.P.; Zaslavsky, L.; Lomsadze, A.; Pruitt, K.D.; Borodovsky, M.; Ostell, J. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* **2016**, *44*, 6614–6624, doi:10.1093/nar/gkw569.

8. Palmer, J.M.; Stajich, J. Funannotate v1.8.1: Eukaryotic Genome Annotation 2020. https://doi.org/10.5281/zenodo.4054262

9. Min, B.; Grigoriev, I.V.; Choi, I.-G. FunGAP: Fungal Genome Annotation Pipeline Using Evidence-Based Gene Model Evaluation. *Bioinformatics* **2017**, *33*, 2936–2937, doi:10.1093/bioinformatics/btx353.

10. Wheeler, D.L.; Barrett, T.; Benson, D.A.; Bryant, S.H.; Canese, K.; Chetvernin, V.; Church, D.M.; DiCuccio, M.; Edgar, R.; Federhen, S.; et al. Database Resources of the National Center for Biotechnology Information. *Nucleic Acids Research* **2007**, *35*, D5–D12, doi:10.1093/nar/gkl1031.

11. Bryant, D.M.; Johnson, K.; DiTommaso, T.; Tickle, T.; Couger, M.B.; Payzin-Dogru, D.; Lee, T.J.; Leigh, N.D.; Kuo, T.-H.; Davis, F.G.; et al. A Tissue-Mapped Axolotl de Novo Transcriptome Enables Identification of Limb Regeneration Factors. *Cell Reports* **2017**, *18*, 762–776, doi:10.1016/j.celrep.2016.12.063.

12. Conesa, A.; Götz, S.; García-Gómez, J.M.; Terol, J.; Talón, M.; Robles, M. Blast2GO: A Universal Tool for Annotation, Visualization and Analysis in Functional Genomics Research. *Bioinformatics* **2005**, *21*, 3674–3676, doi:10.1093/bioinformatics/bti610.

13. Bono, H.; Sakamoto, T.; Kasukawa, T.; Tabunoki, H. Systematic Functional Annotation Workflow for Insects. *Insects* **2022**, *13*, 586, doi:10.3390/insects13070586.

14. Rhoads, A.; Au, K.F. Pacbio Sequencing and Its Applications. *Genomics, Proteomics & Bioinformatics* **2015**, *13*, 278–289, doi:10.1016/j.gpb.2015.08.002.

15. Elmore, M.G.; Banerjee, S.; Pedley, K.F.; Ruck, A.; Whitham, S.A. De Novo Transcriptome of Phakopsora Pachyrhizi Uncovers Putative Effector Repertoire during Infection. *Physiological and Molecular Plant Pathology* **2020**, *110*, 101464, doi:10.1016/j.pmpp.2020.101464.

16. Krueger, F. Felixkrueger/Trimgalore 2025. https://github.com/FelixKrueger/TrimGalore

17. Prjibelski, A.; Antipov, D.; Meleshko, D.; Lapidus, A.; Korobeynikov, A. Using Spades de Novo Assembler. *CP in Bioinformatics* **2020**, *70*, e102, doi:10.1002/cpbi.102.

18. Tegenfeldt, F.; Kuznetsov, D.; Manni, M.; Berkeley, M.; Zdobnov, E.M.; Kriventseva, E.V. OrthoDB and BUSCO Update: Annotation of Orthologs with Wider Sampling of Genomes. *Nucleic Acids Research* **2025**, *53*, D516–D522, doi:10.1093/nar/gkae987.

19. Haas, B.J.; Papanicolaou, A.; Yassour, M.; Grabherr, M.; Blood, P.D.; Bowden, J.; Couger, M.B.; Eccles, D.; Li, B.; Lieber, M.; et al. De Novo Transcript Sequence Reconstruction from RNA-Seq Using the Trinity Platform for Reference Generation and Analysis. *Nat Protoc* **2013**, *8*, 1494–1512, doi:10.1038/nprot.2013.084.

20. Patro, R.; Duggal, G.; Love, M.I.; Irizarry, R.A.; Kingsford, C. Salmon Provides Fast and Bias-Aware Quantification of Transcript Expression. *Nat Methods* **2017**, *14*, 417–419, doi:10.1038/nmeth.4197.

21. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*; Use R!; Second edition.; Springer international publishing: Cham, 2016; ISBN 9783319242774.

22. R Development Core Team. *R a Language and Environment for Statistical Computing: Reference Index*; R Foundation for Statistical Computing: [Vienna], 2010; ISBN 9783900051075.

23. Github Repository: moriharanagisa/fungifunate. Available online: https://github.com/moriharanagisa/fungifunate (accessed on 8 Sep 2025).

24. Github Repository: Bonohu/SAQE. Available online: https://github.com/bonohu/SAQE (accessed on 8 Sep 2025).

25. Pearson, W.R. FASTA Search Programs. In *Encyclopedia of Life Sciences*; Wiley, 2014 ISBN 9780470016176.

26. Ensembl FTP: Homo sapiens GRCh38 protein sequences. Available online: https://ftp.ensembl.org/pub/current_fasta/homo_sapiens/pep/Homo_sapiens.GRCh38.pep.all.fa.gz (accessed on 22 Apr 2023).

27. Ensembl FTP: Mus musculus GRCm39 protein sequences. Available online: https://ftp.ensembl.org/pub/current_fasta/mus_musculus/pep/Mus_musculus.GRCm39.pep.all.fa.gz (accessed on 22 Apr 2023).

28. Ensembl FTP: Saccharomyces cerevisiae R64-1-1 protein sequences. Available online: https://ftp.ensembl.org/pub/current_fasta/saccharomyces_cerevisiae/pep/Saccharomyces_cerevisiae.R64-1-1.pep.all.fa.gz (accessed on 5 Oct 2023).

29. Martin, F.J.; Amode, M.R.; Aneja, A.; Austine-Orimoloye, O.; Azov, A.G.; Barnes, I.; Becker, A.; Bennett, R.; Berry, A.; Bhai, J.; et al. Ensembl 2023. *Nucleic Acids Research* **2023**, *51*, D933–D941, doi:10.1093/nar/gkac958.

30. Uniprot FTP: UniprotKB/swiss-prot protein sequences. Available online: https://ftp.uniprot.org/pub/databases/uniprot/knowledgebase/complete/uniprot_sprot.fasta.gz (accessed on 24 Jan 2024).

31. Basenko, E.Y.; Pulman, J.A.; Shanmugasundram, A.; Harb, O.S.; Crouch, K.; Starns, D.; Warrenfeltz, S.; Aurrecoechea, C.; Stoeckert, C.J.; Kissinger, J.C.; et al. FungiDB: An Integrated Bioinformatic Resource for Fungi and Oomycetes. *JoF* **2018**, *4*, 39, doi:10.3390/jof4010039.

32. FungiDB. Available online: https://fungidb.org/fungidb/app/downloads (accessed on 30 May 2024).

33. Jones, P.; Binns, D.; Chang, H.-Y.; Fraser, M.; Li, W.; McAnulla, C.; McWilliam, H.; Maslen, J.; Mitchell, A.; Nuka, G.; et al. InterProScan 5: Genome-Scale Protein Function Classification. *Bioinformatics* **2014**, *30*, 1236–1240, doi:10.1093/bioinformatics/btu031.

34. Ashburner, M.; Ball, C.A.; Blake, J.A.; Botstein, D.; Butler, H.; Cherry, J.M.; Davis, A.P.; Dolinski, K.; Dwight, S.S.; Eppig, J.T.; et al. Gene Ontology: Tool for the Unification of Biology. *Nat Genet* **2000**, *25*, 25–29, doi:10.1038/75556.

35. The Gene Ontology Consortium; Aleksander, S.A.; Balhoff, J.; Carbon, S.; Cherry, J.M.; Drabkin, H.J.; Ebert, D.; Feuermann, M.; Gaudet, P.; Harris, N.L.; et al. The Gene Ontology Knowledgebase in 2023. *GENETICS* **2023**, *224*, iyad031, doi:10.1093/genetics/iyad031.

36. Durinck, S.; Moreau, Y.; Kasprzyk, A.; Davis, S.; De Moor, B.; Brazma, A.; Huber, W. BioMart and Bioconductor: A Powerful Link between Biological Databases and Microarray Data Analysis. *Bioinformatics* **2005**, *21*, 3439–3440, doi:10.1093/bioinformatics/bti525.

37. InterPro2GO. Available online: https://current.geneontology.org/ontology/external2go/interpro2go (accessed on)

38. Love, M.I.; Huber, W.; Anders, S. Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2. *Genome Biol* **2014**, *15*, 550, doi:10.1186/s13059-014-0550-8.

39. Conesa, A.; Nueda, M.J.; Ferrer, A.; Talón, M. maSigPro: A Method to Identify Significantly Differential Expression Profiles in Time-Course Microarray Experiments. *Bioinformatics* **2006**, *22*, 1096–1102, doi:10.1093/bioinformatics/btl056.

40. Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A.H.; Tanaseichuk, O.; Benner, C.; Chanda, S.K. Metascape Provides a Biologist-Oriented Resource for the Analysis of Systems-Level Datasets. *Nat Commun* **2019**, *10*, 1523, doi:10.1038/s41467-019-09234-6.

41. Raudvere, U.; Kolberg, L.; Kuzmin, I.; Arak, T.; Adler, P.; Peterson, H.; Vilo, J. G:Profiler: A Web Server for Functional Enrichment Analysis and Conversions of Gene Lists (2019 Update). *Nucleic Acids Research* **2019**, *47*, W191–W198, doi:10.1093/nar/gkz369.

42. topGO. Available online: http://bioconductor.org/packages/topGO/ (accessed on 17 September 2025).

43. NCBI nr database. Available online: https://ftp.ncbi.nlm.nih.gov/blast/db/FASTA/nr.gz (accessed on 17 May 2024)

44. Buchfink, B.; Reuter, K.; Drost, H.-G. Sensitive Protein Alignments at Tree-of-Life Scale Using DIAMOND. *Nat Methods* **2021**, *18*, 366–368, doi:10.1038/s41592-021-01101-x.

45. Yanai, I.; Benjamin, H.; Shmoish, M.; Chalifa-Caspi, V.; Shklar, M.; Ophir, R.; Bar-Even, A.; Horn-Saban, S.; Safran, M.; Domany, E.; et al. Genome-Wide Midrange Transcription Profiles Reveal Expression Level Relationships in Human Tissue Specification. *Bioinformatics* **2005**, *21*, 650–659, doi:10.1093/bioinformatics/bti042.

46. Huang, X.; Zhang, R.; Qiu, Y.; Wu, H.; Xiang, Q.; Yu, X.; Zhao, K.; Zhang, X.; Chen, Q.; Penttinen, P.; et al. RNA-Seq Profiling Showed Divergent Carbohydrate-Active Enzymes (CAZymes) Expression Patterns in Lentinula Edodes at Brown Film Formation Stage Under Blue Light Induction. *Front. Microbiol.* **2020**, *11*, 1044, doi:10.3389/fmicb.2020.01044.

47. Li, Y.; Wang, H.; Zhang, Y.; Xiang, Q.; Chen, Q.; Yu, X.; Zhang, L.; Peng, W.; Penttinen, P.; Gu, Y. Hydrated Lime Promoted the Polysaccharide Content and Affected the Transcriptomes of Lentinula Edodes during Brown Film Formation. *Front. Microbiol.* **2023**, *14*, 1290180, doi:10.3389/fmicb.2023.1290180.

48. Krizsán, K.; Almási, É.; Merényi, Z.; Sahu, N.; Virágh, M.; Kószó, T.; Mondo, S.; Kiss, B.; Bálint, B.; Kües, U.; et al. Transcriptomic Atlas of Mushroom Development Reveals Conserved Genes behind Complex Multicellularity in Fungi. *Proc. Natl. Acad. Sci. U.S.A.* **2019**, *116*, 7409–7418, doi:10.1073/pnas.1817822116.

49. Wang, Y.; Zeng, X.; Liu, W. De Novo Transcriptomic Analysis during Lentinula Edodes Fruiting Body Growth. *Gene* **2018**, *641*, 326–334, doi:10.1016/j.gene.2017.10.061.

50. Tang, L.; Chu, T.; Shang, J.; Yang, R.; Song, C.; Bao, D.; Tan, Q.; Jian, H. Oxidative Stress and Autophagy Are Important Processes in Post Ripeness and Brown Film Formation in Mycelium of Lentinula Edodes. *Front. Microbiol.* **2022**, *13*, 811673, doi:10.3389/fmicb.2022.811673.

51. Shen, N.; Xie, H.; Liu, K.; Li, X.; Wang, L.; Deng, Y.; Chen, L.; Bian, Y.; Xiao, Y. Near-Gapless Genome and Transcriptome Analyses Provide Insights into Fruiting Body Development in Lentinula Edodes. *International Journal of Biological Macromolecules* **2024**, *263*, 130610, doi:10.1016/j.ijbiomac.2024.130610.

52. Winter, C.; Fehr, M.; Craig, I.R.; Grammenos, W.; Wiebe, C.; Terteryan-Seiser, V.; Rudolf, G.; Mentzel, T.; Quintero Palomar, M.A. Trifluoromethyloxadiazoles: Inhibitors of Histone Deacetylases for Control of Asian Soybean Rust. *Pest Management Science* **2020**, *76*, 3357–3368, doi:10.1002/ps.5874.

53. Hoeksema, M.; Van Eijk, M.; Haagsman, H.P.; Hartshorn, K.L. Histones as Mediators of Host Defense, Inflammation and Thrombosis. *Future Microbiol.* **2016**, *11*, 441–453, doi:10.2217/fmb.15.151.

54. Sieber, P.; Voigt, K.; Kämmer, P.; Brunke, S.; Schuster, S.; Linde, J. Comparative Study on Alternative Splicing in Human Fungal Pathogens Suggests Its Involvement During Host Invasion. *Front. Microbiol.* **2018**, *9*, 2313, doi:10.3389/fmicb.2018.02313.

55. Gervais, N.C.; Shapiro, R.S. Discovering the Hidden Function in Fungal Genomes. *Nat Commun* **2024**, *15*, 8219, doi:10.1038/s41467-024-52568-z.